

Connections on Parahoric Torsors over Curves

by

Vikraman BALAJI, Indranil BISWAS and Yashonidhi PANDEY

Abstract

We define parahoric \mathcal{G} -torsors for a certain class of Bruhat–Tits group schemes \mathcal{G} on a smooth complex projective curve X when the weights are real, and also define connections on them. We prove that a \mathcal{G} -torsor is given by a homomorphism from $\pi_1(X \setminus D)$ to a maximal compact subgroup of G , where the finite subset $D \subset X$ is the parabolic divisor, if and only if the \mathcal{G} -torsor is polystable.

2010 Mathematics Subject Classification: Primary 14F22, 14D23; Secondary 14D20.

Keywords: Bruhat–Tits group scheme, parahoric torsor, connection, polystability.

§1. Introduction

Let X be an irreducible smooth projective curve defined over the field of complex numbers. Let G be a simple and simply connected affine algebraic group defined over \mathbb{C} . Fix a finite subset $D \subset X$. Let \mathcal{G} be a Bruhat–Tits group scheme with parabolic points at D (see Definition 3.3).

In [BS], an analogue of the Mehta–Seshadri theorem in [MS] was proved relating stable parahoric torsors under Bruhat–Tits group schemes with irreducible homomorphisms of certain Fuchsian groups into a maximal compact subgroup of G . This was done under the assumption that the parabolic weights are rational, or equivalently, the fixed points of the Fuchsian group are all *elliptic*. Recall that in [BS] it is shown that if the weights are chosen to be *rational* then one can recover the \mathcal{G} -torsors as invariant direct images of orbifold principal bundles with respect to suitable ramified covers of X ramified over the parabolic points.

Communicated by T. Mochizuki. Received February 7, 2017. Revised May 3, 2017.

V. Balaji: Chennai Mathematical Institute, Sipcot IT Park, Siruseri 603103, India;
e-mail: balaji@cmi.ac.in

I. Biswas: School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005, India;
e-mail: indranil@math.tifr.res.in

Y. Pandey: Indian Institute of Science Education and Research, Mohali Knowledge city, Sector 81, SAS Nagar, Manauli PO 140306, India;
e-mail: ypandey@iisermohali.ac.in

An obstruction to covering real weights in the setting of parahoric torsors is that the classical Bruhat–Tits group scheme is defined on spectra of discrete valuation rings, while the phenomenon of parabolic bundles with real weights naturally lies in the setting of an analytic neighborhood of the origin. Further, the notion of invariant direct image fails to generalize directly to the analytic setting when the covering map is no longer algebraic. We address this issue by working with a natural analogue of Deligne extensions to the parahoric torsor setting: to a pair (E, \widehat{d}) of a principal G -bundle E equipped with a flat connection \widehat{d} on the punctured disc, we give a canonical extension to a torsor under the Bruhat–Tits group scheme on the compact Riemann surface (see Section 10.2). We use this to construct the parahoric torsor associated to a representation of the fundamental group of the punctured Riemann surface.

Another obstruction in the case of real weights is the notion of stability for parahoric torsors. Filling the lacuna in [BS], we have a definition of stability for parahoric torsors (6.2) that covers real weights as well. Then, following the approach in the paper of Mehta–Seshadri ([MS, p. 217]), in Section 7 we develop the theme of *variation of weights* in the setting of parahoric torsors, where the notions of (semi)stability for the case of real weights are shown to coincide for “nearby” rational weights. Although the broad lines are the same, this generalization is not entirely straightforward.

We then go on to define the notion of a connection on a \mathcal{G} -torsor over a smooth projective curve over \mathbb{C} and prove the analogue of the Donaldson–Uhlenbeck–Yau correspondence in the parahoric setting when the weights are *real*. The basic idea is to use the Tannakian formalism and the argument reduces to the case of the associated parabolic Lie algebra bundle. The more general reductive group case is then an easy generalization.

§1.1. Origins

In the early 1960s, Mumford defined the notion of stability for vector bundles on curves as a tool to get Hausdorff moduli spaces; using geometric-invariant theory, he then constructed the moduli space of stable vector bundles. In [NS], Narasimhan and Seshadri gave an alternative characterization of stability using flat connections; more precisely, they proved that a holomorphic vector bundle E on a compact Riemann surface is stable if and only if E arises from an irreducible projective unitary representation of the fundamental group of the Riemann surface. This correspondence between flat projective unitary connections and stable vector bundles has been generalized in several directions. Ramanathan ([Ra]) extended the correspondence to the case of holomorphic principal G -bundles, where G is a complex reductive affine algebraic group ([Ra]). On the other hand, Mehta and

Seshadri ([MS]) generalized the Narasimhan–Seshadri construction to include unitary logarithmic connections; more precisely, they identified the irreducible unitary representations of general Fuchsian groups with fixed conjugacy classes as stable parabolic bundles with fixed parabolic data determined by the conjugacy classes. These logarithmic connections are those that have regular singularity at finitely many points and were already apparent in the early work of Weil ([We]); their importance was emphasized by Deligne in [De]. The objects replacing stable bundles in the Mehta–Seshadri correspondence are stable parabolic vector bundles. Following Donaldson’s reinterpretation of the Narasimhan–Seshadri correspondence, Biquard ([Biq]) gave a differential geometric interpretation of this Mehta–Seshadri correspondence.

It is a very natural problem to generalize the Mehta–Seshadri correspondence from the setting of parabolic vector bundles to that of principal G -bundles, where G is a complex reductive affine algebraic group. On the side of representations, the objects were easy to define: they were homomorphisms of Fuchsian groups taking values in a maximal compact subgroup of G such that, for each puncture of the Riemann surface, the associated conjugacy class in the fundamental group of the surface is mapped to a fixed conjugacy class of the maximal compact subgroup of G . Similarly, on the side of connections, the corresponding objects have been quite well understood since the work of Deligne. The central problem was to generalize the notion of a stable parabolic vector bundle to the setting of principal G -bundles. From a Tannakian perspective ([BBN1]) it became apparent that a naive generalization in terms of *parabolic G -bundles*, i.e., *principal G bundles with parabolic structures*, was insufficient for setting up a comprehensive analogue of the Mehta–Seshadri correspondence.

At a technical level, it was not clear what the correct notion of weight should be in the general setting. A breakthrough came in the work of Boalch ([Bo]) and Balaji–Seshadri ([BS]). These papers introduced the correct and very natural notion of weight, namely, as a point in the affine apartment of G . As a consequence of this viewpoint, the reason for the inadequacy of parabolic G -bundles also became clear. It was realized that instead of parabolic G -bundles, one should consider torsors or principal homogeneous spaces under parahoric group schemes in the sense of Bruhat and Tits. Balaji and Seshadri ([BS]) extended the Mehta–Seshadri theorem to the case of parahoric torsors with rational weights.

Prior to these works, there were at least two partial approaches to generalize the Mehta–Seshadri theorem, both around the turn of the millennium. The approach in [BBN1] was Tannakian in spirit and followed the method of Nori ([No]); this Tannakian approach identified the basic problem, namely that the object associated to a representation of a Fuchsian group into the maximal compact of G ,

such that for each puncture of the Riemann surface the associated conjugacy class in the fundamental group of the surface is mapped to a fixed conjugacy class of the maximal compact subgroup of G , in general will not be a principal G -bundle on the Riemann surface. The approach in [TW] again gave a partial solution to the problem; in the language of Weyl alcoves, the solution was for weights in the interior of the Weyl alcove that corresponds to the subclass of *parabolic G -bundles*.

Parahoric group schemes \mathcal{G} and \mathcal{G} -torsors over smooth projective curves were first defined and studied by Pappas–Rapoport ([PR1]). Subsequently, in [PR2], they made several conjectures on the moduli stacks of \mathcal{G} -torsors. Most of these conjectures were answered by Heinloth ([He1]) providing a precise setting for the study of these moduli stacks. The paper of Seshadri ([Se2]) takes up the question of the analogue of the Mehta–Seshadri theorem; the main emphasis in [Se2] was again to point out that for a solution to the problem of obtaining analogues of the Mehta–Seshadri theorem, one has to go beyond the category of principal G -bundles. Evidence of the role of Bruhat–Tits theory was also given in a few illustrative examples. The paper of Boalch ([Bo]) studies logarithmic connections on G -bundles and the notion of parahoric torsors comes along with the first appearance of the notion of weights for these torsors. Around the same time and independently in [BS], a similar notion of weights was defined towards providing a satisfactory analogue of the Mehta–Seshadri theorem in the general setting for semisimple groups G , thereby completing the broad picture outlined in [Se2]. The notion of “invariant direct images” of torsors in [BS] plays a key role analogous to the one in [MS] for the case of vector bundles.

§2. Preliminaries

In this section we collect together some standard notions and notation that will be used throughout this paper. See [BT1], [BT2], [BS], [He1] for this section and the next one. The base field will always be \mathbb{C} .

Define

$$(2.1) \quad A := \mathbb{C}[[t]] \quad \text{and} \quad K := \mathbb{C}((t)),$$

where t denotes a uniformizing parameter. Let G be a *semisimple simply connected affine algebraic group* defined over \mathbb{C} . The Lie algebra of G will be denoted by \mathfrak{g} . Fix a maximal torus $T \subset G$; let $Y(T) = \text{Hom}(\mathbb{G}_m, T)$ be the group of all holomorphic one-parameter subgroups of T .

§2.1. Apartment of G

For each maximal torus T of G , we have the *standard affine apartment* \mathcal{A}_T . It is an affine space under $Y(T) \otimes_{\mathbb{Z}} \mathbb{R}$. In general, there is no origin in the apartment

(cf. [BT1]), but for purposes of this paper, we shall identify \mathcal{A}_T with $Y(T) \otimes_{\mathbb{Z}} \mathbb{R}$ (see [BS, §2]).

Let \mathcal{V} be a real affine space. A function $f : \mathcal{V} \rightarrow \mathbb{R}$ is said to be an *affine functional* if

$$f(rx + (1 - r)y) = rf(x) + (1 - r)f(y)$$

for all $x, y \in \mathcal{V}$ and $r \in \mathbb{R}$. Thus, for a root α of G and an integer $n \in \mathbb{Z}$ there is the affine functional

$$\alpha + n : \mathcal{A}_T \rightarrow \mathbb{R}, \quad x \mapsto \alpha(x - 0) + n.$$

We note that these are called the *affine roots of G* . The zero locus of $\alpha + n$ will be denoted by $H_{\alpha+n}$; it is called an *affine wall*. The set of affine walls is known to be *locally finite*, meaning any compact subset of \mathcal{A}_T intersects only finitely many affine walls. For any point $x \in \mathcal{A}_T$, let Z_x denote the set of affine functionals vanishing on x . For an integer $n \geq 0$, define

$$\mathcal{H}_n = \{x \in \mathcal{A} \mid |Z_x| = n\},$$

which is the set of all points where n of the affine functionals vanish. A *facet* Ω of \mathcal{A}_T is defined to be a connected component of \mathcal{H}_n for some n . The *dimension* of a facet is its dimension as a real manifold. We then have a decomposition of the apartment,

$$(2.2) \quad \mathcal{A}_T = \bigsqcup_n \mathcal{H}_n.$$

Although, as mentioned above, almost always Θ will be a point of \mathcal{A}_T , sometimes Θ will also be a facet of \mathcal{A}_T .

§3. Parahoric group scheme and torsors

§3.1. Invariant direct image

The base field is \mathbb{C} . Let $p : W \rightarrow U$ be a finite flat surjective morphism of normal, integral Noetherian schemes that is Galois. So the Galois group $\text{Gal}(p)$, which we will denote by Γ , acts on W with $U = W/\Gamma$ being the quotient. Such a morphism p is called a *Galois covering* with Galois group Γ .

Let \mathcal{G} be an *affine group scheme* over W . For the above Galois covering map p , the direct image $p_*\mathcal{G}$ is defined as follows: for each U -scheme S , set

$$(3.1) \quad (p_*\mathcal{G})(S) = \text{Hom}_W(S \times_U W, \mathcal{G});$$

this is representable by a group scheme ([BLR, Thm. 4 and Prop. 6]). Assume that the group scheme \mathcal{G} is equipped with an action of the Galois group Γ that lifts the

action of Γ on W ; in particular, the “multiplication map” and the “inverse map” on \mathcal{G} are Γ -equivariant. Such a \mathcal{G} will be called a Γ -group scheme over W .

There is a left action of Γ on $S \times_U W$ induced by the action of Γ on W . This and the left action of Γ on \mathcal{G} together induce the following right action of Γ on $(p_*\mathcal{G})(S)$:

$$(3.2) \quad (f \cdot \gamma)([s, w]) := \gamma^{-1} \cdot f(\gamma \cdot [s, w]), \quad [s, w] \in S \times_U W, \quad \gamma \in \Gamma.$$

Consider the *fixed point* subscheme under the above action of Γ on $p_*\mathcal{G}$. The general results on fixed point subschemes given in [Ed, Sect. 3] can be applied to our situation *since the characteristic is 0*. Consequently, a canonically defined *smooth* closed X -subgroup scheme

$$p_*^\Gamma \mathcal{G} := (p_*\mathcal{G})^\Gamma \subset p_*\mathcal{G}$$

is obtained. This $p_*^\Gamma \mathcal{G}$ is representable because $p_*\mathcal{G}$ is representable.

Definition 3.1 (Invariant direct image). Let $p : W \rightarrow U$ be as above, and let $\Gamma = \text{Gal}(W/U)$. Let \mathcal{G} be a smooth affine Γ -group scheme over W . Define the *invariant direct image* of \mathcal{G} to be

$$(3.3) \quad p_*^\Gamma(\mathcal{G}) := (p_*\mathcal{G})^\Gamma,$$

so $(p_*^\Gamma \mathcal{G})(S) = \mathcal{G}(S \times_U W)^\Gamma$ for any U -scheme S .

More generally, if E is any affine scheme over W with a lift of the Γ -action on W , then define the *invariant direct image* of E to be

$$p_*^\Gamma E := (p_*E)^\Gamma.$$

§3.2. Parahoric torsors

The notation of Section 2.1 will be followed. Let $R = R(T, G)$ denote the root system of G (cf. [Sp, p. 125]). Thus, for every $r \in R$, there is the root homomorphism $u_r : \mathbb{G}_a \rightarrow G$ ([Sp, Prop. 8.1.1]).

For any nonempty subset $\Theta \subset \mathcal{A}_T$, the *parahoric subgroup* $\mathcal{P}_\Theta \subset G(K)$,

$$(3.4) \quad \mathcal{P}_\Theta := \langle T(A), \{u_r(t^{m_r(\Theta)} A)\}_{r \in R} \rangle,$$

is the subgroup generated by $T(A)$ and $\{u_r(t^{m_r(\Theta)} A)\}_{r \in R}$, where

$$m_r = m_r(\Theta) = - \left\lceil \inf_{\theta \in \Theta} (\theta, r) \right\rceil,$$

and A is defined in (2.1) ([BS, p. 8]). Moreover, by [BT2, Sect. 1.7] we have an affine flat smooth group scheme $\mathcal{G}_\Theta \rightarrow \text{Spec}(A)$ corresponding to Θ . The set of K -valued

(respectively, A -valued) points of \mathcal{G}_Θ is identified with $G(K)$ (respectively, \mathcal{P}_Θ). The group scheme \mathcal{G}_Θ is uniquely determined by its A -valued points. Here we will often take Θ to be just a point of \mathcal{A}_T .

Remark 3.2. We remark that the notion of a parahoric subgroup is defined in the greatest generality in the basic papers of Bruhat and Tits. For our purposes, the definition given above is sufficient.

Let X be a smooth complex projective curve. We fix once and for all a nonempty finite set of closed points

$$(3.5) \quad D = \{x_j\}_{j=1}^m \subset X.$$

These will play the role of parabolic points. Let $A_j := \widehat{\mathcal{O}_{X,x_j}}$ be the complete discrete valuation ring with function field $K_j \simeq \mathbb{C}((t))$ and residue field \mathbb{C} , obtained by completing the local rings \mathcal{O}_{X,x_j} . We shall denote $\text{Spec}(A_j)$ by D_j .

Definition 3.3. Let \mathcal{G} be a flat, affine group scheme on X of finite type. We call \mathcal{G} a *Bruhat–Tits group scheme* with parabolic points D , if

- (1) restricted to $X \setminus D$, it is isomorphic to the split group scheme $G \times (X \setminus D)$, and
- (2) $\mathcal{G}|_{D_j} \rightarrow D_j$ is a Bruhat–Tits group scheme for each j .

We shall denote \mathcal{G} by \mathcal{G}_Ω , where $\Omega = \{\Omega_j\}_{j=1}^m$ is a collection of facets of the Bruhat–Tits building with $\mathcal{G}|_{D_j}$ corresponding to Ω_j .

By the general theory due to Bruhat and Tits, one has an affine flat smooth group scheme \mathcal{G}_Θ on $\text{Spec}(A)$ corresponding to Θ . The set of K -valued (respectively, A -valued) points of \mathcal{G}_Θ is identified with $G(K)$ (respectively, \mathcal{P}_Θ). The group scheme \mathcal{G}_Θ is uniquely determined by its A -valued points. These notions were defined and the moduli stack of \mathcal{G}_Θ -torsors studied extensively in the papers of Pappas–Rapoport (cf. [PR1], [PR2]) and Heinloth ([He1]). To construct one on the whole of X one can proceed as in [BS, 5.2]. The existence of such group schemes also follows from the invariant direct images constructed above (see also [BS, Thm. 5.2.7]).

Definition 3.4 ([BS, Sect. 6]). A *quasi-parahoric torsor* \mathcal{E} is a $\mathcal{G}_{\Omega,X}$ -torsor on X .

Definition 3.5 ([BS, Sect. 6], [Bo]). A *parahoric torsor* is a pair (\mathcal{E}, θ) consisting of

- (1) a $\mathcal{G}_{\Omega,X}$ -torsor $\mathcal{E} \rightarrow X$, and

(2) weights, meaning elements $\theta = \{\theta_i\}_{i=1}^m \in (Y(T) \otimes \mathbb{R})^m$ in the interior of Ω_i .

Remark 3.6. The above notion of weight is the precise analogue of the classical weight for a parabolic vector bundle with *multiplicity* (cf. [MS, p. 211, Def. 1.5]).

Remark 3.7. It should be noted that as in [BS], the theory of Bruhat–Tits group schemes that are used here assumes that the group G is semisimple and simply connected. On the other hand, for the case of $\mathrm{GL}(V)$, which satisfies neither of these conditions, the parabolic bundles are classic (cf. [Se1], [MS], [Bis]). In [BS, Exa. 2.3.4] and [BS, Rem. 6.1.5], it is noted that the torsors under Bruhat–Tits group schemes for $\mathrm{GL}(V)$ are the same as the parabolic vector bundles. Let us spell this out for the convenience of the reader: Let $\mathcal{GL}(V)$ be a Bruhat–Tits group scheme on $\mathrm{Spec}(A)$ with generic fiber $\mathrm{GL}(V)$. Fix a maximal torus $T(V) \subset \mathrm{GL}(V)$. Let \mathcal{E} be a $\mathcal{GL}(V)$ -torsor, and let $\theta(V) \in \mathrm{Al}(T(V))_{\mathbb{R}}$ be a weight as a point in the so-called Weyl alcove (see [BS, p. 9]). Then, the associated vector bundle gets a canonical parabolic structure with quasi-parabolic type determined by the group scheme $\mathcal{GL}(V)$ while the parabolic weights are given by $\theta(V)$. We observe that in the case of $\mathrm{GL}(n, \mathbb{C})$, or $\mathrm{SL}(n, \mathbb{C})$, giving a point $\theta(V) \in \mathrm{Al}(T(V))_{\mathbb{R}}$ in the Weyl alcove is equivalent to giving n -tuples $(\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{R}^n$, such that every $\alpha_i \geq 0$ and $\alpha_i \leq \alpha_{i+1}$, for $1 \leq i \leq n-1$; in the case of $\mathrm{SL}(n, \mathbb{C})$, the condition of the determinant translates to the further condition

$$\sum_{i=1}^n \alpha_i \in \mathbb{Z}.$$

By a Weyl group conjugation, we can also arrange the α_i 's in increasing order, i.e., $0 \leq \alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_n < 1$. Once the parahoric subgroup is chosen, this determines a flag type and hence we may even order the parabolic weights in a strictly increasing sequence.

§3.3. Rational weights

Starting with a m -tuple of weights $\theta \in (Y(T) \otimes \mathbb{Q})^m$, following the proof of the converse in [BS, Thm. 2.3.1], we get positive integers d_1, d_2, \dots, d_m such that $d_i \cdot \theta_i \in Y(T)$. By choosing these d_i to be *smallest with this property*, we see that a choice of θ entails a choice of ramification index d_x at each point $x \in D$.

There exists a ramified Galois cover of curves $p : Y \rightarrow X$ that is

- unramified over $X \setminus D$, and
- the ramification index over $x \in D$ is d_x ,

if and only if exactly one of the following conditions holds:

- (1) the genus of X is nonzero;
- (2) $X = \mathbb{P}^1$ and $\#D \geq 3$;
- (3) $X = \mathbb{P}^1$, $\#D = 2$ and $d_x = d_y$, where $D = \{x, y\}$.

(See [Na, p. 26, Prop. 1.2.12].)

When we consider parahoric torsors as invariant direct images of (Γ, G) bundles, it will always be with respect to such a Galois cover. See also [Bis].

§3.4. Rational weights and parahoric \mathcal{G} -torsors as (Γ, G) -bundles

Let (X, D) be as above, and let $\mathcal{G} \rightarrow X$ be a Bruhat–Tits group scheme over X . By [BS, Thm. 5.3.1], there exists a (possibly ramified) finite Galois cover $p : Y \rightarrow X$ branched over D , and a principal G -bundle F over Y (cf. [BS, Notn. 5.1.0.1]) equipped with a lift of the action of the Galois group $\Gamma := \text{Gal}(p)$, such that the following statements hold:

- (1) Let ${}_F G := \text{Isom}_Y(F, F)$ be the twisting of the constant group scheme $G \rightarrow Y$ by F . The invariant direct image satisfies the condition

$$(3.6) \quad p_*^\Gamma {}_F G = \mathcal{G}.$$

- (2) Let $D_Y := p^{-1}(D) \subset Y$ denote the ramification points of the covering p . For each $y \in D_Y$, let $\Gamma_y \subset \Gamma$ be the isotropy subgroup that fixes y . Let $\tau_y : \Gamma_y \rightarrow \text{Aut}(F_y)$ be the action of Γ_y on the fiber F_y . Its conjugacy class is called the local type of F at y ; this local type will be denoted by $[\tau_y]$. By the type τ of F , we mean the set of conjugacy classes $[\tau_y]$ of τ_y :

$$\tau := \{[\tau_y] \mid y \in D_Y\}.$$

Let $\mathcal{M}_Y^\tau(\Gamma, G)$ denote the moduli stack of (Γ, G) bundles over Y of type τ , and let $\mathcal{M}_X(\mathcal{G})$ denote the moduli stack of \mathcal{G} torsors on X . Then there is an isomorphism of moduli stacks

$$(3.7) \quad \alpha_F : \mathcal{M}_Y^\tau(\Gamma, G) \rightarrow \mathcal{M}_X(\mathcal{G}),$$

given by the (Γ, G) bundle F as follows: Denote by F^{op} the left G -bundle defined by $gf := fg^{-1}$, where $g \in G$ and $f \in F$. The above group scheme ${}_F G$ acts on the right of F^{op} . The isomorphism in (3.7) is

$$(3.8) \quad E \mapsto p_*^\Gamma(E \times_{Y, G} F^{\text{op}}) = p_*^\Gamma(\text{Isom}_Y(E, F)).$$

The inverse of the map in (3.7) is given by

$$(3.9) \quad \mathcal{E} \mapsto p^* \mathcal{E} \times_{p^* \mathcal{G}} F.$$

(3) Let $y \in D_Y$, and $x := p(y)$. Let $N_y = \text{Spec}(B)$, where $B = \widehat{\mathcal{O}_{Y,y}}$, and also $D_x = \text{Spec}(A)$ with $A = \widehat{\mathcal{O}_{X,x}}$. Let U_y denote the group of local Γ_y - G automorphisms of $F|_{N_y}$ (cf. [BS, Def. 2.2.7]). Then by [BS, Prop. 5.1.2] we have

$$(3.10) \quad \mathcal{G}|_{D_x}(A) = U_y.$$

For the equality in (3.10), we need the existence of the (Γ, G) bundle F only locally on N_y and not on the entire Y .

A different (Γ, G) bundle F' of type τ will, in general, provide a different (cf. (3.7)) isomorphism $\alpha_{F'} : \mathcal{M}_Y(\Gamma, G) \rightarrow \mathcal{M}_X(\mathcal{G})$.

§4. Change of weights under a homomorphism

§4.1. The local homomorphism problem

Consider the following problem. Let A be an arbitrary discrete valuation ring with quotient field K . Let G be a semisimple and simply connected group. Let $\rho : G \rightarrow H$ be a homomorphism. Fix a maximal torus $T \subset G$. Then fix a weight θ in the affine apartment \mathcal{A}_T for T . It should be emphasized that θ may not be a rational point, meaning it may not lie in the image of $Y(T) \otimes_{\mathbb{Z}} \mathbb{Q}$. Fix a maximal torus $T_H \subset H$ such that $\rho(T) \subset T_H$. Via the identification between \mathcal{A}_T and $Y(T) \otimes_{\mathbb{Z}} \mathbb{R}$ mentioned in Section 2.1, we have a linear map

$$(4.1) \quad \mathcal{A}_T \rightarrow \mathcal{A}_{T_H}$$

between the apartments. Let θ_H denote the image of θ under this map. We wish to construct, through ρ , a canonical homomorphism of group schemes over $\text{Spec}(A)$: from \mathcal{G}_θ corresponding to θ to \mathcal{G}_{θ_H} corresponding to θ_H .

§4.2. Facets of a homomorphism

Let $\rho : G \rightarrow H$ be a homomorphism. The affine roots of H give affine functionals on \mathcal{A}_{T_H} (cf. Section 2). These functionals are defined over the rationals. By (4.1) it follows that the linear map of apartments $\mathcal{A}_T \rightarrow \mathcal{A}_{T_H}$ corresponding to ρ is also defined over the rationals. Indeed, it is induced by the algebraic map $\rho|_T : T \rightarrow T_H$. Via $\mathcal{A}_T \rightarrow \mathcal{A}_{T_H}$, we view the above affine functionals on \mathcal{A}_{T_H} , associated to the affine roots of H , as affine functionals on \mathcal{A}_T . Take the union of these affine functionals with the usual affine functionals on \mathcal{A}_T corresponding to the affine roots of G . Note that all these functionals are defined over rationals. We shall call them ρ -functionals.

Definition 4.1. An *affine wall* of ρ is the zero locus of a ρ -functional. For any $x \in \mathcal{A}_T$, define Z_x to be the set of all affine functionals on \mathcal{A}_T vanishing at x . Define

$$\mathcal{H}_n := \{x \in \mathcal{A} \mid |Z_x| = n\}.$$

A *facet* of ρ is a connected component of \mathcal{H}_n for some $n \geq 0$.

Notice that for the identity homomorphism $G \rightarrow G$, we have just got the usual facets. By the theory of buildings, one knows that the usual affine walls of G provide a decomposition of \mathcal{A}_T (cf. (2.2)).

We claim that more generally, the facets of a homomorphism $\rho : G \rightarrow H$ provide an even finer decomposition of

$$\mathcal{A}_T = \bigsqcup_n \mathcal{H}_n;$$

here n varies over a finite set. To prove this claim, first note that the set of all ρ -affine walls corresponding to ρ -functionals forms a locally finite set. Indeed, this follows because any compact set C of \mathcal{A}_T meets finitely many usual affine walls in \mathcal{A}_T . Now under the map $\mathcal{A}_T \rightarrow \mathcal{A}_{T_H}$, the image of C being compact, meets finitely many affine walls of \mathcal{A}_{T_H} . Thus, C meets only finitely many walls of ρ .

Proposition 4.2. *Let $\rho : G \rightarrow H$ be a homomorphism of algebraic groups. Given a weight $\theta \in \mathcal{A}_T$, there exists a rational weight $\eta \in \mathcal{A}_T$ lying in the same facet as θ such that η_H and θ_H also lie in a common facet of \mathcal{A}_{T_H} .*

Proof. It can be shown that if the element $\theta \in \mathcal{A}_T$ lies in a zero-dimensional facet, then it must be a rational point. Indeed, this follows from that fact that we are looking at a common zero locus of a set of rational functionals. Taking the contrapositive of the last statement, if θ is not rational then a ρ -facet of θ cannot be zero-dimensional. So we can find a rational weight η in it.

Now, by construction, both η and θ lie in the same facet of \mathcal{A}_T , and also θ_H and η_H lie in a common facet of \mathcal{A}_{T_H} . □

Remark 4.3. Henceforth, when we say that a *rational weight η is close to θ with respect to $\rho : G \rightarrow H$* , we mean a *rational weight* in the sense of Proposition 4.2.

The constructions in the proof of Proposition 4.2 can be extended to the context of a finite set of representations $\{\rho_i : G \rightarrow \text{GL}(V_i)\}_{i \leq n}$ of G .

Now further assume that η and θ are actually interior points of one ρ -facet. This assumption implies that they define isomorphic Bruhat–Tits group schemes with generic fiber G and also group schemes with generic fiber H . There are canonical homomorphisms between them by the following proposition.

Proposition 4.4. *Let $\rho : G \rightarrow H$ be given. Let $\theta \in \mathcal{A}_T$ be a weight with image θ_H under the map of apartments $\mathcal{A}_T \rightarrow \mathcal{A}_{T_H}$. Then there is a canonical homomorphism of group schemes $\mathcal{G}_\theta \rightarrow \mathcal{G}_{\theta_H}$ over $\text{Spec}(A)$.*

Proof. Let us recall a characterization of parahoric groups when the weight η lies in $\mathcal{A}_\mathbb{Q}$ ([BS, Thm. 2.3.1]). In this case, by (3.10), there exists a ramified Galois cover

$$p : \text{Spec}(B) \rightarrow \text{Spec}(A)$$

with Galois group say Γ , and a (Γ, G) bundle $F \rightarrow \text{Spec}(B)$, such that

$$(4.2) \quad \mathcal{G}_{\Omega(A)} = \mathcal{G}_\eta(A) = \text{Aut}_{(\Gamma, G)}(F).$$

Choose a rational weight η close to θ with respect to ρ . This gives an element η_H close to θ_H . Further, let

$$F(H) := F \times_G H$$

be the (Γ, H) bundle obtained by extending the structure group using ρ . Via the natural homomorphism $\text{Aut}_{(\Gamma, G)}(F) \rightarrow \text{Aut}_{(\Gamma, H)}(F(H))$ and the equalities $\mathcal{G}_\eta(A) = \text{Aut}_{(\Gamma, G)}(F)$ and $\mathcal{G}_{\eta_H}(A) = \text{Aut}_{(\Gamma, H)}(F(H))$, we obtain a map of parahoric groups

$$\rho_K : \mathcal{G}_\eta(A) \rightarrow \mathcal{G}_{\eta_H}(A).$$

Now the characterizing property of the Bruhat–Tits group schemes is that they are *étouffé*, which means that any morphism at the level of group schemes over $\text{Spec}(A)$ is determined completely by the A -valued points alone (cf. [BT2, Def. 1.7.1]).

Thus, we get a morphism over $\text{Spec}(A)$ of group schemes

$$(4.3) \quad \mathcal{G}_\eta \rightarrow \mathcal{G}_{\eta_H}$$

extending (4.2). Since η_H is close to θ_H , it follows that $\mathcal{G}_{\eta_H}(A) = \mathcal{G}_{\theta_H}(A)$, and hence we have an induced isomorphism of group schemes $\mathcal{G}_{\eta_H} \rightarrow \mathcal{G}_{\theta_H}$ on $\text{Spec}(A)$, which gives the homomorphism $\rho_\theta : \mathcal{G}_\theta \rightarrow \mathcal{G}_{\theta_H}$. \square

To the best of our knowledge, Proposition 4.4 is not available in the papers of Bruhat and Tits. Alternatively it can be proved using the general framework of functoriality of buildings as in [La, Thm. 2.1.8] and [La, Thm. 2.2.1].

§5. Associated constructions

As before, G is semisimple and simply connected. Let $\rho : G \rightarrow \text{GL}(V)$ be a rational representation. Fix a maximal torus $T \subset G$, and also fix a maximal torus $T_V \subset \text{GL}(V)$ containing $\rho(T)$. Let \mathcal{E} be a parahoric $\mathcal{G}_{\theta, X}$ -torsor on X with weights θ .

§5.1. The construction

As in (3.5), fix a finite subset $D = \{x_j\}_{j=1}^m \subset X$. For each parahoric group scheme $\mathcal{G}_j \rightarrow D_j = \text{Spec}(A_j)$, $1 \leq j \leq m$, we get a facet Ω_j in the Bruhat–Tits building $\mathcal{B}(G)$. Any weight $\theta \in \mathcal{A}_T^m$ induces a weight θ_V in $\mathcal{A}_{T(V)}^m$ (see Section 4.2).

The group scheme $\mathcal{G}_{\theta, X}$ on X is obtained by gluing \mathcal{G}_{θ_j} , for every $1 \leq j \leq m$, with

$$\mathcal{G}|_{X \setminus D} \simeq (X \setminus D) \times G.$$

Using the same gluing data via the representation ρ one immediately gets a Bruhat–Tits group scheme $\mathcal{GL}_{\theta_V, X}$, and using locally Proposition 4.4, we see that ρ gives a natural global homomorphism of group schemes over X :

$$(5.1) \quad \rho_X : \mathcal{G}_{\theta, X} \rightarrow \mathcal{GL}_{\theta_V, X}.$$

Using ρ_X one gets the standard construction of extension of structure groups.

By Remark 3.7, via ρ one obtains an associated parabolic vector bundle

$$\mathcal{E}_{V, \theta_V} := (\mathcal{E}(V), \theta_V)$$

with parabolic weights θ_V . We will mostly need to apply associated constructions under the adjoint homomorphism $\text{Ad} : G \rightarrow \text{GL}(\mathfrak{g})$.

§5.2. Tensor product of parabolic vector bundles

Assume that we have homomorphisms $\rho_1 : G \rightarrow \text{GL}(V)$ and $\rho_2 : G \rightarrow \text{GL}(W)$, and let

$$\rho_1 \otimes \rho_2 : G \rightarrow \text{GL}(V \otimes W)$$

be their tensor product. Therefore, for a parahoric torsor (\mathcal{E}, θ) we get parabolic vector bundles $\mathcal{E}_{V, \theta_V}$, $\mathcal{E}_{W, \theta_W}$ and $\mathcal{E}_{V \otimes W, \theta_{V \otimes W}}$.

When the weights θ are rational numbers then, by (Γ, G) bundle theory, we have a canonical isomorphism of parabolic vector bundles ([MY]):

$$\mathcal{E}_{V, \theta_V} \otimes^p \mathcal{E}_{W, \theta_W} \simeq \mathcal{E}_{V \otimes W, \theta_{V \otimes W}},$$

where \otimes^p is a parabolic tensor product. The following proposition, which extends this isomorphism to the case of *real* weights, is immediate by observing that the quasi-parabolic bundle for a parabolic tensor product does not change under a sufficiently small change of the parabolic weights of the factors, while the parabolic weights of the parabolic tensor product are given by the parabolic weights of the factors using a standard algebraic formula.

Proposition 5.1. *Let $\theta \in \mathcal{A}_T$ be arbitrary. Then there is a canonical isomorphism*

$$\mathcal{E}_{V, \theta_V} \otimes^p \mathcal{E}_{W, \theta_W} \simeq \mathcal{E}_{V \otimes W, \theta_{V \otimes W}}$$

of parabolic vector bundles.

Let V and W be two representations of G and $\psi : V \rightarrow W$ a G -equivariant homomorphism. Let (\mathcal{E}, θ) be a parahoric torsor. By identifying the G -module $\text{Hom}(V, W)$ with $W \otimes V^*$ it follows that ψ induces a homomorphism

$$(5.2) \quad (\mathcal{E}, \theta)(\psi) : \mathcal{E}_{V, \theta_V} \rightarrow \mathcal{E}_{W, \theta_W}.$$

§5.3. Lie algebra bundle of a \mathcal{G} -torsor

Consider a parahoric \mathcal{G}_θ -torsor (\mathcal{E}, θ) , where θ is a system of real weights. We define a parabolic Lie bracket operation on $\mathcal{E}(\mathfrak{g})$ as follows. The Lie bracket $[\cdot, \cdot] : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ is a G -equivariant homomorphism for the adjoint action of G , and hence by (5.2) we have a homomorphism of parabolic vector bundles

$$\mathcal{E}_{[\cdot, \cdot]} : \mathcal{E}(\mathfrak{g} \otimes \mathfrak{g}) \rightarrow \mathcal{E}(\mathfrak{g}).$$

Now Proposition 5.1 gives an isomorphism

$$\mathcal{E}(\mathfrak{g}) \otimes^p \mathcal{E}(\mathfrak{g}) \xrightarrow{\sim} \mathcal{E}(\mathfrak{g} \otimes \mathfrak{g})$$

of parabolic vector bundles. Combining with $\mathcal{E}_{[\cdot, \cdot]}$, the Lie bracket can be defined as the parabolic homomorphism

$$(5.3) \quad [\cdot, \cdot] : \mathcal{E}(\mathfrak{g}) \otimes_X^p \mathcal{E}(\mathfrak{g}) \rightarrow \mathcal{E}(\mathfrak{g}).$$

§6. Semistability and stability of torsors

Let $\mathcal{G}_{\Omega, X}$ be a Bruhat–Tits group scheme on X as in Definition 3.3, and let θ be such that $\mathcal{G}_{\Omega, X} \simeq \mathcal{G}_{\theta, X}$. Let (\mathcal{E}, θ) be a parahoric $\mathcal{G}_{\theta, X}$ -torsor with arbitrary real weights $\theta \in \mathcal{A}_T^m$.

Remark 6.1. Let G_K be a split group scheme over a field K . Let E_K be a G_K -torsor. Consider the twisted group scheme $E(G)_K$. Then giving a parabolic subgroup scheme

$$P_K \subset E(G)_K$$

(of fiber type P) is equivalent to giving a reduction of structure group of E_K to P (cf. [DG, Exp. XXVI, Cor. 3.6]).

§6.1. First definition

Let $\mathcal{E}(\mathcal{G})$ denote the group scheme of automorphisms of \mathcal{E} obtained by taking the quotient $\mathcal{E} \times_X \mathcal{G}$ by the left \mathcal{G} -action on \mathcal{E} and the right \mathcal{G} -action on itself by conjugation. Let $\text{Lie}(\mathcal{E}(\mathcal{G}))$ denote the Lie algebra bundle of $\mathcal{E}(\mathcal{G})$. One has the following well-known identification of Lie algebra bundles:

$$(6.1) \quad \text{Lie}(\mathcal{E}(\mathcal{G})) = \mathcal{E}(\mathfrak{g}).$$

Since (\mathcal{E}, θ) is a parahoric torsor, by Section 5.3, the above Lie algebra bundle $\mathcal{E}(\mathfrak{g})$ gets a natural parabolic Lie algebra bundle structure $(\mathcal{E}(\mathfrak{g}), \theta_{\mathfrak{g}})$. Thus, via the isomorphism (6.1) the bundle $\text{Lie}(\mathcal{E}(\mathcal{G}))$ gets a parabolic vector bundle structure with a Lie bracket operation compatible with the parabolic structure:

$$(6.2) \quad (\text{Lie}(\mathcal{E}(\mathcal{G})), \theta_{\mathfrak{g}}, [\cdot, \cdot]).$$

By Remark 6.1, giving a generic parabolic reduction is equivalent to giving a parabolic subgroup scheme

$$P_K \subset \mathcal{E}(\mathcal{G}_K).$$

For each $1 \leq j \leq m$, we take the flat closure of P_K in $\mathcal{E}(\mathcal{G}_{A_j})$. This will give a subgroup scheme $\mathcal{P} \subset \mathcal{E}(\mathcal{G})$.

The extended subgroup scheme \mathcal{P} also gives a Lie subalgebra bundle

$$\text{Lie}(\mathcal{P}) \subset \text{Lie}(\mathcal{E}(\mathcal{G})) \simeq \mathcal{E}(\mathfrak{g}).$$

Then endow the bundle $\text{Lie}(\mathcal{P})$ with the *canonical induced parabolic structure* on the divisor D and denote this parabolic subbundle by $\text{Lie}(\mathcal{P})_*$.

Definition 6.2. We say that the parahoric torsor

$$(\mathcal{E}, \theta)$$

with arbitrary real weights $\theta \in \mathcal{A}_T^m$ is semistable (respectively, stable) as a $\mathcal{G}_{\theta, X}$ -torsor if for every generic parabolic reduction datum as above,

$$\text{par.deg}(\text{Lie}(\mathcal{P})_*) \leq 0 \quad (\text{respectively, } \text{par.deg}(\text{Lie}(\mathcal{P})_*) < 0).$$

§6.2. Second definition

Now assume that (\mathcal{E}, θ) is a parahoric torsor with *rational weights*. Let $P_K \subset \mathcal{G}_K$ be a maximal parabolic subgroup of the generic fiber \mathcal{G}_K of $\mathcal{G}_{\Omega, X}$. Let $\chi : P_K \rightarrow \mathbb{G}_{m, K}$ be a strictly antidominant character of the parabolic subgroup P_K . Therefore, the associated line bundle on \mathcal{G}_K/P_K is ample. Since the quotient map $\mathcal{E}_K \rightarrow \mathcal{E}_K/P_K$ defines a principal P_K -bundle, it follows that χ defines a line bundle L_χ on $\mathcal{E}_K/P_K = \mathcal{E}_K(\mathcal{G}_K/P_K)$. For any reduction of structure group

$$s_K : X \setminus D \rightarrow \mathcal{E}_K(\mathcal{G}_K/P_K),$$

we have the pulled-back line bundle $s_K^*(L_\chi)$ on $X \setminus D$. This line bundle $s_K^*(L_\chi)$ extends X by the following proposition.

Proposition 6.3 ([BS, Prop. 6.3.1]). *Let (\mathcal{E}, θ) be a parahoric torsor with rational weights. Let s_K be a generic reduction of structure group of \mathcal{E}_K to P_K . Then*

the line bundle $s_K^*(L_x)$ on $X \setminus D$ has a canonical extension L_x^θ to X as a parabolic line bundle.

Definition 6.4 ([BS, Def. 6.3.4]). A parahoric torsor (\mathcal{E}, θ) with rational weights is called *stable* (respectively, *semistable*) if for every maximal parabolic $\mathcal{P}_K \subset \mathcal{G}_K$, for all strictly antidominant characters χ of \mathcal{P}_K and for every reduction of structure group s_K as above,

$$\text{par.deg}(L_x^\theta) < 0 \quad (\text{respectively, } \text{par.deg}(L_x^\theta) \leq 0).$$

We observe that the two definitions, namely Definitions 6.2 and 6.4, are equivalent when the weights are rational; the proof is identical to the proof of [Ra, Lem. 2.1].

Remark 6.5. In a recent paper, Heinloth ([He2]) studies the Hilbert–Mumford criterion in terms of algebraic stacks. The viewpoint developed there allows natural choices of test objects for the verification of stability, leading to criteria for the existence of separated coarse moduli spaces.

Let \mathcal{E} be a \mathcal{G} -torsor for a parahoric group scheme \mathcal{G} and let $\mathcal{B} \subset \text{Aut}_\mathfrak{g}(\mathcal{E})$. Let $\mathcal{E}_\mathcal{B}$ be a reduction of structure group to \mathcal{B} . To a character χ of \mathcal{B} , one can then associate a line bundle $\mathcal{E}_\mathcal{B}(\chi)$ on X . In the language of Section 6.2 this will be a parabolic line bundle. In [He2, Sect. 3.5] the classical notion of a parabolic degree of $\mathcal{E}_\mathcal{B}(\chi)$ is recovered in the new setting, which leads to the definition of stability as in Section 6.2. An advantage with this new definition is that it works in positive characteristics as well and moreover, the parahoric group scheme \mathcal{G} need not be assumed to be generically split as in the present paper.

§7. (Semi)stability and polystability under variation of weights

Let \mathcal{V} be the category of all quasi-parahoric $\mathcal{G}_{\theta, X}$ -torsors along D with weights θ varying in \mathcal{A}_T^m . For convenience, we shall work with a *single* parabolic point $P \in X$. The generalization to finitely many points follows without any difficulty.

Let r be the rank of $\mathcal{E}(\mathfrak{g})_*$. The degree of the underlying vector bundle is denoted by d_1 . Note that its parabolic degree is 0 because $\mathfrak{g} = \mathfrak{g}^*$ as G -modules. Let $\mathcal{V}_\mathfrak{g}$ denote the space of all parabolic vector bundles such that

- the rank is the fixed integer r ,
- the quasi-parabolic structure at P is given by that of $\mathcal{E}(\mathfrak{g})_*$,
- the degree of the underlying vector bundle is d_1 ,
- the parabolic degree is zero, and

- the parabolic weights $0 \leq \alpha_1 < \alpha_2 < \dots < \alpha_{r'} < 1$ are not fixed, but the length r' and the multiplicities $m_1, \dots, m_{r'}$ are fixed.

Take any $V_* \in \mathcal{V}_{\mathfrak{g}}$. For a subbundle W of V , if n_k for each $1 \leq k \leq r'$ denotes the multiplicity of α_k for the parabolic structure on W induced by $\mathcal{V}_{\mathfrak{g}}$, the condition for α_* -stability of V_* is

$$(7.1) \quad \text{degree}(W) + \sum_{k \leq r'} n_{i_k} \alpha_{i_k} < 0$$

for all subbundles W of V ; for semistability the strict inequality is replaced by inequality. Let $\chi(V, W, \alpha)$ denote the left-hand side of (7.1).

There exists a constant $C_2 \geq 0$ such that if $\text{degree}(W) > C_2$, then

$$\chi(V, W, \alpha) \geq 0$$

for all α . Thus, for any $V \in \mathcal{V}_{\mathfrak{g}}$, if the underlying vector bundle V admits a subbundle W with $\text{degree}(W) \geq C_2$, then V_* can never be parabolic semistable for any choice of weights α_* . We note that the quasi-parabolic structure of $V \in \mathcal{V}_{\mathfrak{g}}$ alone determines such bundles.

§7.1. Facets of a quasi-parahoric torsor

As before, G is simple and simply connected.

In this subsection we shall consider only parahoric torsors $\mathcal{E} \in \mathcal{V}$ such that the quasi-parabolic bundle $\mathcal{E}(\mathfrak{g})$ admits no subbundle of degree greater than C_2 :

$$(7.2) \quad \mathcal{V}_{\mathfrak{g}}(C_2) := \{V_* \in \mathcal{V}_{\mathfrak{g}} \mid V \text{ has no subbundle of degree greater than } C_2\}.$$

Proposition 7.1. *The set of inequalities required to verify the (semi)stability of any bundle in $\mathcal{V}_{\mathfrak{g}}(C_2)$ has finite cardinality.*

Proof. From (7.1) we see that there exists a constant $C_1 \leq 0$ such that

$$\text{degree}(W) \leq d_1 + C_1 \implies \chi(V, W, \alpha) < 0.$$

In other words, subbundles of degree at most C_1 will never be destabilizing with respect to any inequality. Thus, to check (semi)stability of $(\mathcal{E}(\mathfrak{g}), \theta_{\mathfrak{g}})$ we may restrict ourselves to subbundles W of $\mathcal{E}(\mathfrak{g})$ such that

$$(7.3) \quad C_1 \leq \text{degree}(W) \leq C_2.$$

The ranks of subbundles W vary between 1 and $r-1$. Let $m_1, \dots, m_{r'}$ be the multiplicities of the quasi-parabolic structure on $\mathcal{V}_{\mathfrak{g}}$. The multiplicities n_{i_1}, \dots, n_{i_k} of the induced parabolic structure are positive integers. Thus, as one varies over $\mathcal{V}_{\mathfrak{g}}$, only finitely many inequalities appear. □

If $\mathcal{E}(\mathfrak{g}) \in \mathcal{V}_{\mathfrak{g}}(C_2)$, it follows that to check the (semi)stability of (\mathcal{E}, θ) we need to consider only finitely many inequalities corresponding to a (possibly proper) subset of the set of inequalities seen in Proposition 7.1. This is because we need to check these inequalities for subbundles that are Lie algebra bundles of certain subgroup schemes (see Definition 6.2).

We fix a maximal torus T of G and also fix a maximal torus $T_{\mathfrak{g}}$ of $GL(\mathfrak{g})$ such that $Ad(T_G) \subset T_{\mathfrak{g}}$.

For every inequality,

$$\text{degree}(W) + \sum_{k \leq r'} n_{i_k} \alpha_{i_k} < 0 \quad (\text{respectively, } \leq 0)$$

in (7.1) and for every integer c between C_1 (as in the proof of Proposition 7.1) and C_2 , we associate a functional

$$\ell_c : \mathcal{A}_{T_{\mathfrak{g}}} \longrightarrow \mathbb{R}$$

as follows: for any $\eta = (\alpha_1, \dots, \alpha_r) \in \mathcal{A}_{T_{\mathfrak{g}}}$,

$$\ell_c(\eta) = \frac{c}{\text{rank}(W)} + \frac{\sum_{k \leq r'} n_{i_k} \alpha_{i_k}}{\text{rank}(W)}.$$

Define $f : \mathcal{A}_T \longrightarrow \mathbb{R}$ by $f(\theta) = \ell_c(\theta(\mathfrak{g}))$. These are finitely many in number. We denote the set of these functionals by $S_T^{\mathcal{E}}$ (or S_T for notational convenience). Further, they are defined over rationals, since clearly the definition of f involves only rational numbers and the map $\mathcal{A}_T \longrightarrow \mathcal{A}_{T_{\mathfrak{g}}}$ is defined over rationals and is linear. For any functional f in S_T , define the f -wall in \mathcal{A}_T as

$$W_f := \{x \in \mathcal{A}_T \mid f(x) = 0\}.$$

The collection $\{W_f\}_{f \in S_T^{\mathcal{E}}}$ will be called the *walls* of \mathcal{E} .

Definition 7.2. Fix a quasi-parahoric torsor \mathcal{E} . For any $\theta \in \mathcal{A}_T$, let

$$S_1^{\theta} = \{f \in S_T^{\mathcal{E}} \mid x \in W_f\}.$$

Let $\mathcal{H}_n = \{x \in \mathcal{A}_T \mid |S_1^{\theta}| = n\}$. Define a *facet* of \mathcal{E} to be a connected component of \mathcal{H}_n for some $n \geq 0$.

Thus, the facets of \mathcal{E} provide a decomposition

$$\mathcal{A}_T = \bigsqcup_n \mathcal{H}_n$$

of \mathcal{A}_T . Note that only finitely many n appear here.

For any weight $\theta \in \mathcal{A}_T$ there is a unique $n \geq 0$ such that $\theta \in \mathcal{H}_n$. Thus, for any $\theta \in \mathcal{A}_T$, the following three conditions are equivalent:

- (1) θ does not belong to any \mathcal{E} -wall,
- (2) S_1^θ is empty, and
- (3) $\theta \in \mathcal{H}_0$.

The *facet* of θ is the unique facet of \mathcal{E} containing it.

The following propositions generalize [MS, Sect. 2, p. 217]. We note that the quasi-parabolic bundles $\mathcal{E}(\mathfrak{g})$ cannot admit any subbundle of degree greater than C_2 (cf. (7.2)).

Proposition 7.3. *Let θ lie in $\mathcal{A}_T = Y(T) \otimes \mathbb{R}$. Then there exists an element θ' in the rational apartment $\mathcal{A}_{\mathbb{Q}} = Y(T) \otimes \mathbb{Q}$ such that for all $\mathcal{E} \in \mathcal{V}$, the pair (\mathcal{E}, θ) is semistable (respectively, stable) if and only if (\mathcal{E}, θ') is semistable (respectively, stable).*

Proof. Let S_2^θ be the complement of S_1^θ in S_T .

We note that θ has a strictly positive distance from each W_f , where $f \in S_2^\theta$. Let d be the minimum distance if S_2^θ is nonempty and set d to be ∞ if S_2^θ is empty. Thus, in all cases $d > 0$. Let U be the ball in the alcove of radius d around θ .

Let I denote the \mathcal{E} -facet of θ . Let I_1 be a connected component of $I \cap U$ containing θ , where U is the above ball. Now if I_1 is not reduced to a single point, then we can take a rational weight θ' in it. If I_1 is just a point, then θ must be rational because $d > 0$ and all the functionals are defined over \mathbb{Q} . In this case, we take θ' to be θ itself.

Let us check that the (semi)stability conditions for θ and θ' coincide. For each functional J in S_T ,

- (1) if $J \in S_1^\theta$ then $J(\theta') = J(\theta)$, because $\theta' \in I_1 \subset I$, and
- 2. if $J \in S_2$ then $\text{sign}(J(\theta')) = \text{sign}(J(\theta))$, because $\theta' \in I_1 \subset U$.

So for \mathcal{E} , one has that θ' -(semi)stability is equivalent to θ -(semi)stability. □

We return to the setting of m -marked points on X noting that the above discussion immediately goes through for multiple marked points.

Lemma 7.4. *Let $\theta \in \mathcal{A}_T^m$. Let $\rho_i : G \rightarrow \text{GL}(V_i)$ for $i \leq m$ be finitely many representations. Then there exists $\theta' \in \mathcal{A}_{\mathbb{Q}}^m$ such that for any $\mathcal{E} \in \mathcal{V}$, and any $i \leq m$, the parahoric torsor $(\mathcal{E}, \theta(V_i))$ is stable (respectively, semistable) if and only if $(\mathcal{E}, \theta'(V_i))$ is stable (respectively, semistable).*

Proof. This is immediate from Proposition 7.3. □

The following proposition is a generalization of [MS, Prop. 2.1].

Proposition 7.5. *Given any $\theta_0 \in \mathcal{A}_T^m$, there exists a neighborhood U of θ_0 in \mathcal{A}_T^m with the property that for all $\mathcal{E} \in \mathcal{V}$ such that (\mathcal{E}, θ_0) is stable, the pair (\mathcal{E}, θ) is stable for all $\theta \in U$.*

Proof. Now θ may as well be rational. Owing to the stability condition, $f(\theta) < 0$ for all $f \in S_T$. Thus, we have $S_2^\theta = S_T$. Let d be the minimum distance between θ and any f -wall. Now we take U to be the ball around θ with radius d . □

Definition 7.6. A parahoric torsor (\mathcal{E}, θ) for the linear parahoric group scheme $\mathcal{GL}(V)$ is called *polystable* if the associated parabolic vector bundle $\mathcal{E}(V)_*$ is polystable (i.e., a direct sum of stable parabolic bundles of parabolic degree 0).

It is straightforward to check that Lemma 7.4 remains valid if stability in the lemma is substituted by polystability.

Corollary 7.7. *Let $\rho : G \rightarrow \mathrm{GL}(V)$ be a representation and $\rho_x : \mathcal{G} \rightarrow \mathcal{GL}(V)$ the induced homomorphism of parahoric group schemes as in (5.1). Let \mathcal{E} be a \mathcal{G} -torsor. Then, for a weight $\theta \in \mathcal{A}_T^m$ such that (\mathcal{E}, θ) is stable, the pair $(\mathcal{E}(V)_*, \theta(V))$ is polystable.*

Proof. By Proposition (7.5), we can assume θ is rational. For a stable equivariant principal G -bundle, the associated bundles are polystable. Consequently, in view of the equivalence of categories in Section 3.4, The stability of (\mathcal{E}, θ) implies that $(\mathcal{E}(V), \theta(V))$ is polystable. □

§8. Connections on parahoric \mathcal{G} -torsors

The main aim of this section is to define connections on a parahoric $\mathcal{G}_{\theta,x}$ -torsor.

§8.1. D_X -modules

We briefly recall the definition of D_X -modules.

Definition 8.1. Let $X \rightarrow S$ be an S -scheme. Let dx denote the image of x under the canonical de Rham differentiation map $d : \mathcal{O}_X \rightarrow \Omega_{X/S}^1$. Let \mathcal{F} be a coherent sheaf of \mathcal{O}_X -modules over X . By a *D_X -module structure* on \mathcal{F} we mean an \mathcal{O}_S -linear homomorphism of a sheaf of abelian groups $\nabla : \mathcal{F} \rightarrow \mathcal{F} \otimes_{\mathcal{O}_X} \Omega_{X/S}^1$ satisfying the Leibniz rule which says that

$$(8.1) \quad \nabla(xf) = f \otimes dx + x\nabla(f),$$

where f and x are local sections of \mathcal{F} and \mathcal{O}_X respectively.

Definition 8.2. Let $\nabla_{\mathcal{F}} : \mathcal{F} \rightarrow \mathcal{F} \otimes \Omega_{X/S}^1$ and $\nabla_{\mathcal{E}} : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_{X/S}^1$ be two connections over \mathcal{F} and \mathcal{E} respectively. Define their tensor product $\nabla_{\mathcal{F}} \otimes \nabla_{\mathcal{E}} : \mathcal{F} \otimes \mathcal{E} \rightarrow \mathcal{F} \otimes \mathcal{E} \otimes \Omega_{X/S}^1$ to be

$$(8.2) \quad \nabla_{\mathcal{F} \otimes \mathcal{E}}(f \otimes e) = \nabla_{\mathcal{F}}(f) \otimes e + f \otimes \nabla_{\mathcal{E}}(e),$$

where f and e are local sections of \mathcal{F} and \mathcal{E} respectively.

Similarly define $\nabla_{\text{Hom}} : \text{Hom}(\mathcal{E}, \mathcal{F}) \rightarrow \text{Hom}(\mathcal{E}, \mathcal{F}) \otimes \Omega_{X/S}^1$ to be

$$(8.3) \quad \nabla_{\text{Hom}(\mathcal{E}, \mathcal{F})}(\Phi)(e) = \nabla_{\mathcal{F}}(\Phi(e)) - \Phi(\nabla_{\mathcal{E}}(e)),$$

where Φ and e are local sections of $\text{Hom}(\mathcal{E}, \mathcal{F})$ and \mathcal{E} respectively.

§8.2. Logarithmic connections on curves

The canonical line bundle of the smooth complex projective curve X will be denoted by K_X . Fix a finite subset $D = \{x_i\}_{1 \leq i \leq m} \subset X$; define

$$K_X(\log D) = K_X \otimes \mathcal{O}_X(D).$$

A logarithmic connection on a vector bundle $V \rightarrow X$ singular on D is a first-order algebraic differential operator $\nabla : V \rightarrow V \otimes K_X(\log D)$ satisfying the Leibniz rule.

For a point $x \in D$, the fiber $K_X(\log D)_x$ is identified with \mathbb{C} using the Poincaré adjunction formula. For a logarithmic connection (V, ∇) , the composition

$$V \xrightarrow{\nabla} V \otimes K_X(\log D) \rightarrow (V \otimes K_X(\log D))_x \rightarrow V_x,$$

which is a \mathbb{C} -linear endomorphism of V_x , is called the *residue of ∇ at x* ([De, p. 53]), and it is denoted by $\text{Res}(\nabla, x)$. The monodromy of ∇ around x is conjugate to

$$(8.4) \quad \exp(-2\pi\sqrt{-1} \text{Res}(\nabla, x))$$

([De, p. 53, Thm. 1.17]).

§8.3. Connection on parabolic vector bundles

Let $V \rightarrow X$ be a vector bundle on X . A *quasi-parabolic* structure on V over D is a filtration, for each $x \in D$, of subspaces

$$(8.5) \quad V_x = F_1^x \supseteq F_2^x \supseteq \dots \supseteq F_{a_x}^x \supseteq F_{a_x+1}^x = \{0\}.$$

A *parabolic structure* on V over D is a quasi-parabolic structure as above together with real numbers

$$0 \leq \alpha_1^x < \dots < \alpha_i^x < \dots < \alpha_{a_x}^x < 1$$

associated to the quasi-parabolic flags. We shall often abbreviate a parabolic vector bundle $(V, \{F_*^x, \alpha_*^x\}_{x \in D})$ by V_* .

Definition 8.3. A connection on V_* is a logarithmic connection ∇ on V such that for all $x \in D$,

- the residue $\text{Res}(\nabla, x)$ is semisimple and preserves the quasi-parabolic flag at x , meaning $\text{Res}(\nabla, x)(F_i^x) \subseteq F_i^x$ for all i , and
- $\text{Res}(\nabla, x)(F_i^x/F_{i+1}^x) = \alpha_i^x \text{Id}_{F_i^x/F_{i+1}^x}$.

A connection on V_* induces a connection on the dual parabolic vector bundle V_*^* . To see this, given a logarithmic connection ∇ on V defining a connection on V_* , consider the logarithmic connection on $V^* \otimes \mathcal{O}_X(D)$ induced by ∇ . This logarithmic connection preserves the subsheaf of $V^* \otimes \mathcal{O}_X(D)$ identified with the vector bundle underlying the parabolic vector bundle V_*^* . The logarithmic connection on this subsheaf obtained this way defines a connection on V_*^* .

Let V_*^1 and V_*^2 be parabolic vector bundles with underlying vector bundles V^1 and V^2 respectively. Let ∇^1 and ∇^2 be connections on V_*^1 and V_*^2 respectively. Consider the logarithmic connection on $V^1 \otimes V^2 \otimes \mathcal{O}_X(D)$ induced by ∇^1 and ∇^2 . It preserves the subsheaf of $V^1 \otimes V^2 \otimes \mathcal{O}_X(D)$ corresponding to the parabolic tensor product $V_*^1 \otimes^p V_*^2$. The logarithmic connection on this subsheaf obtained this way defines a connection on $V_*^1 \otimes^p V_*^2$.

§8.4. Lie connection on a principal G -bundle

For a principal G -bundle $E \rightarrow X$, let $E(\mathfrak{g}) = E_G \times^G \mathfrak{g}$ be its adjoint bundle. The fibers of $E(\mathfrak{g})$ are equipped with a Lie bracket structure $[\cdot, \cdot] : E(\mathfrak{g}) \otimes E(\mathfrak{g}) \rightarrow E(\mathfrak{g})$ induced by the Lie algebra structure of \mathfrak{g} .

Definition 8.4. A Lie connection on E is a connection

$$\nabla : E(\mathfrak{g}) \rightarrow E(\mathfrak{g}) \otimes \Omega_X^1$$

such that following diagram is commutative:

$$(8.6) \quad \begin{array}{ccc} E(\mathfrak{g}) \otimes E(\mathfrak{g}) & \xrightarrow{[\cdot, \cdot]} & E(\mathfrak{g}) \\ \nabla_{\otimes} \downarrow & & \downarrow \nabla \\ E(\mathfrak{g}) \otimes E(\mathfrak{g}) \otimes \Omega_X^1 & \xrightarrow{[\cdot, \cdot] \otimes \text{Id}_{\Omega_X^1}} & E(\mathfrak{g}) \otimes \Omega_X^1 \end{array}$$

where ∇_{\otimes} is the connection on $E(\mathfrak{g}) \otimes E(\mathfrak{g})$ induced by ∇ .

The above commutativity condition means that the section of $E(\mathfrak{g}) \otimes (E(\mathfrak{g}) \otimes E(\mathfrak{g}))^*$ given by the Lie bracket operation on $E(\mathfrak{g})$ is flat with respect to the connection on $E(\mathfrak{g}) \otimes (E(\mathfrak{g}) \otimes E(\mathfrak{g}))^*$ induced by ∇ .

A connection on E_G produces a Lie connection on E_G . Therefore, we get a map from the space of all connections on E_G to the space of all Lie connections on E_G . Since G is semisimple, the adjoint homomorphism $G \rightarrow \mathrm{GL}(\mathfrak{g})$ has finite kernel and its image is the connected component, containing the identity element, of the group of all automorphisms of the Lie algebra \mathfrak{g} . From this it follows that the above map from the space of all connections on E_G to the space of all Lie connections on E_G is a bijection. This fact motivates the definition of a connection on a parahoric torsor.

§8.5. Connection on parahoric \mathcal{G} -torsors

We refer to Section 5.3 for the notation. Take any $\mathcal{G} = \mathcal{G}_{\theta,x}$.

Definition 8.5. A *connection* on a \mathcal{G} -torsor \mathcal{E} is a logarithmic parabolic connection (see Definition 8.3) ∇ on $\mathcal{E}(\mathfrak{g})$ satisfying the condition that the section of $\mathrm{Hom}^p(\mathcal{E}(\mathfrak{g}) \otimes^p \mathcal{E}(\mathfrak{g}), \mathcal{E}(\mathfrak{g}))$ given by the homomorphism in (5.3) is flat with respect to the connection ∇_{hom} on the parabolic vector bundle $\mathrm{Hom}^p(\mathcal{E}(\mathfrak{g}) \otimes^p \mathcal{E}(\mathfrak{g}), \mathcal{E}(\mathfrak{g}))$ induced by ∇ .

§8.6. A Tannakian description of connection

Let M be a smooth complex variety. Let G be a complex reductive algebraic group and $E_G \rightarrow M$ a principal G -bundle. Take any pair (H, f) , where H is a complex algebraic group and $f : G \rightarrow H$ an algebraic homomorphism such that the corresponding homomorphism of Lie algebras $df : \mathrm{Lie}(G) \rightarrow \mathrm{Lie}(H)$ is injective. Let $E_H := E_H \times^f H \rightarrow M$ be the principal H -bundle obtained by extending the structure group of E_G using f . Let

$$\tilde{f} : E_G \rightarrow E_H$$

be the natural morphism. A connection on E_G induces a connection on E_H . The converse is also true. To see this, fix a G -equivariant splitting

$$\sigma : \mathrm{Lie}(H) \rightarrow \mathrm{Lie}(G),$$

meaning $\sigma \circ df = \mathrm{Id}_{\mathrm{Lie}(G)}$ (such a splitting exists because G is reductive). If D is a $\mathrm{Lie}(H)$ -valued 1-form on E_H defining a connection on H , then $\sigma \circ \tilde{f}^*D$ is a connection on E_G . If D_0 is a connection on E_G and D the connection on E_H induced by D_0 , then the connection $\sigma \circ \tilde{f}^*D$ on E_G coincides with D_0 . Indeed, this follows immediately from the fact that $\sigma \circ df = \mathrm{Id}_{\mathrm{Lie}(G)}$.

Therefore, the map from connections on E_G to connections on E_H is injective. The image of this map from connections on E_G admits a group-theoretic description. This will be explained below.

We remove the assumption that the algebraic group G is reductive. As before, $f : G \rightarrow H$ is any algebraic homomorphism such that df is injective.

A theorem of Chevalley (see [Hu, p. 80]) says that there is a finite-dimensional left H -module

$$\rho : H \rightarrow \mathrm{GL}(W)$$

and a complex line $\ell \subset W$ such that $f(G)$ is exactly the isotropy subgroup, of the point in the projective space $P(W)$ representing the line ℓ , for the action of H on the projective space $P(W)$ of lines in W induced by the action of H on W . Let

$$E_W := E_H \times^H W \rightarrow M$$

be the vector bundle associated to E_H for the H -module W . For a connection D on E_H , the connection on E_W induced by D will be denoted by D_W . Note that E_W is identified with the vector bundle associated to E_G for the action $\rho \circ f$ of G on W . The condition on ℓ implies that the action of G on W preserves it. Let $E_\ell \subset E_W$ be the line subbundle associated to the G -submodule $\ell \subset W$.

A connection D on E_H is induced by a connection on E_G if and only if the corresponding connection D_W on E_W preserves the above-line subbundle $E_\ell \subset E_W$. This characterizes the connections on E_H that are induced by connections on E_G .

We recall that Tannakian theory involves describing properties of principal bundles in terms of properties of associated vector bundles. For a Tannakian description of connections on E_G , take $H = \mathrm{GL}(V)$, so V is a finite-dimensional G -module. Let

$$E_V := E_G \times^f V \rightarrow M$$

be the vector bundle associated to E_G for the G -module V . From the above observation we know that a connection on E_G is a connection D on the vector bundle E_V such that the connection on the vector bundle E_W induced by D preserves the line subbundle $E_\ell \subset E_W$.

§9. Connections on (Γ, G) -bundles and rational weights

Let F be a principal G -bundle on a curve Y with adjoint bundle

$$(9.1) \quad \mathrm{Ad}(F) = F(G) = F \times^G G,$$

where G acts on itself by conjugation. Given a principal G -bundle E on Y , define the “twisting” by F ,

$$E \times^G F^{\mathrm{op}} := (E \times_Y F) / \sim,$$

where the equivalence relation identifies all pairs $(e, f), (ez, fz) \in E \times_Y F$, with $z \in G$. Consider the map

$$\xi : E \times_Y F \times_Y F \times G \longrightarrow E \times_Y F, \quad (e, f, fz, z_1) \longmapsto (ezz_1, fz),$$

where $(e, f) \in E \times_Y F$ and $z, z_1 \in G$. There is a unique map

$$\widehat{\xi} : (E \times^G F^{\text{op}}) \times_Y F(G) \longrightarrow E \times^G F^{\text{op}}$$

such that the following diagram is commutative:

$$\begin{array}{ccc} (E \times_Y F) \times_Y (F \times G) & \xrightarrow{\xi} & E \times_Y F \\ \downarrow & & \downarrow \\ (E \times^G F^{\text{op}}) \times_Y F(G) & \xrightarrow{\widehat{\xi}} & E \times^G F^{\text{op}}; \end{array}$$

recall that $E \times^G F^{\text{op}}$ and $F(G)$ are quotients of $E \times_Y F$ and $F \times G$ respectively. This map $\widehat{\xi}$ makes $E \times^G F^{\text{op}}$ an $F(G)$ -torsor on Y . Let α_F be the map from the space of principal G -bundles to the space of $F(G)$ -torsors on Y defined by $E \longmapsto E \times^G F^{\text{op}}$. This α_F is an equivalence of categories.

Consider the adjoint action of G on $\text{Lie}(G) = \mathfrak{g}$. Let $\text{ad}(F) = F \times^G \mathfrak{g} \longrightarrow Y$ be the associated adjoint vector bundle. We note that $\text{ad}(F)$ is the Lie algebra bundle associated to the group scheme $F(G)$.

Let ∇_0 be a connection F . Using ∇_0 we will define connections on an $F(G)$ -torsor.

The connection ∇_0 induces a connection on every fiber bundle associated to F . In particular, it produces a connection on $F(G)$; this connection on $F(G)$ given by ∇_0 will be denoted by ∇_0^G . The kernel of the differential $d\pi : TF(G) \longrightarrow \pi^*TY$ of the map π in (9.1) is identified with $\pi^* \text{ad}(F)$. So the above connection ∇_0^G on $F(G)$ gives a homomorphism

$$(9.2) \quad \nabla_0^G : TF(G) \longrightarrow \pi^* \text{ad}(F).$$

Take any $F(G)$ -torsor $\varphi : E \longrightarrow Y$. Consider the action $E \times F(G) \longrightarrow E$ of $F(G)$ on E . Let

$$(9.3) \quad \delta : TE \oplus \varphi^*TF(G) \longrightarrow TE$$

be the differential of this map giving the action. Consider the differential of φ ,

$$d\varphi : TE \longrightarrow \varphi^*TY.$$

Let

$$T_\varphi := \text{kernel}(d\varphi) \subset TE$$

be the relative tangent bundle for the projection φ . The action of $F(G)$ on E identifies T_φ with $\varphi^* \text{ad}(F)$.

A connection on an $F(G)$ -torsor $\varphi : E \rightarrow Y$ is a holomorphic homomorphism of vector bundles over Y ,

$$\beta : TE \rightarrow \varphi^* \text{ad}(F) = \text{ad}(\varphi^* F)$$

such that

- (1) the restriction of β to T_φ coincides with the above identification of T_φ with $\varphi^* \text{ad}(F)$, and
- (2) for the homomorphism δ in (9.3),

$$\delta(\text{kernel}(\beta) \oplus \varphi^* \text{kernel}(\nabla_0^G)) \subset \text{kernel}(\beta),$$

where ∇_0^G is the homomorphism in (9.2).

Note that the above definition of a connection on E depends on ∇_0 .

If F is the trivial principal G -bundle $Y \times G$, then $F(G) = Y \times G$, and an $F(G)$ -torsor is in fact a principal G -bundle on Y . If we choose ∇_0 to be the trivial connection on $Y \times G$, then connections on an $F(G)$ -torsor are the same as connections on the corresponding principal G -bundle.

The following lemma is straightforward to check.

Lemma 9.1. *Given an $F(G)$ -torsor $\varphi : E \rightarrow Y$, a homomorphism*

$$\beta : TE \rightarrow \varphi^* \text{ad}(F) = \text{ad}(\varphi^* F)$$

defines a connection on E if and only if

$$\delta^* \beta = \beta \oplus \nabla_0^G$$

on $TE \oplus \varphi^ TF(G)$, where δ is constructed in (9.3) and ∇_0^G is the homomorphism in (9.2).*

Proposition 9.2. *Twisting by F defines an equivalence between principal G -bundles equipped with a connection and $F(G)$ -torsors equipped with a connection.*

Proof. Let E be a principal G -bundle on Y . Let D be a connection on E . Consider D as a \mathfrak{g} -valued 1-form on E . Let D' denote the \mathfrak{g} -valued 1-form on F corresponding to the connection ∇_0 on F . So (D, D') is a \mathfrak{g} -valued 1-form on the fiber product $E \times_Y F$. The pullback of $\text{ad}(F)$ to F is identified with the trivial vector bundle $F \times \mathfrak{g} \rightarrow F$. Therefore, (D, D') defines a 1-form with values in the pullback of $\text{ad}(F)$ to $E \times_Y F$. This form on $E \times_Y F$ descends to the quotient $F(G)$ -torsor

$E \times^G F^{\text{op}}$ as a 1-form with values in the pullback of $\text{ad}(F)$ to $E \times^G F^{\text{op}}$. It is straightforward to check that this form defines a connection on the $F(G)$ -torsor corresponding to E .

Conversely, let β be a connection on an $F(G)$ -torsor $\varphi : E \rightarrow Y$. Consider the pullback β' of β to $E \times_Y F$ as a 1-form with values in the pullback of $\text{ad}(F)$. As noted above, the pullback of $\text{ad}(F)$ to $E \times_Y F$ is identified with the trivial vector bundle with fiber \mathfrak{g} . So β' is a 1-form on $E \times_Y F$ with values in \mathfrak{g} . Let D' be the pullback of the connection form D to $E \times_Y F$. Then $\beta' - D'$ descends to E by the projection $E \times_Y F \rightarrow E$, and this descended form defines a connection on the principal G -bundle E .

The above two constructions are evidently inverses of each other. □

Assume that Y is equipped with the action of a finite group Γ . A Γ -connection on a (Γ, G) -bundle E on Y is a connection on E that is preserved by the action of Γ .

Proposition 9.3. *Let $E \rightarrow Y$ be a (Γ, G) -bundle on some Galois cover $p : Y \rightarrow X$ with Galois group Γ . Let \mathcal{E} be the parahoric torsor on X with rational weights corresponding to E . Then there is a natural bijection between the connections on \mathcal{E} and the Γ -connections on E .*

Proof. This follows from the fact that the connections on a Γ -equivariant vector bundle on Y are in bijection with the connections on the corresponding parabolic vector bundle on X . □

§10. Flat unitary connections on parahoric torsors and stability

§10.1. Polystable parahoric torsors

Lemma 10.1. *Let V_* be a polystable parabolic vector bundle of parabolic degree 0 with real weights θ . Then the parabolic vector bundle $(V_*)^{\otimes m} \otimes^p ((V_*)^*)^{\otimes n}$ is also polystable.*

Proof. A parabolic vector bundle of parabolic degree 0 is polystable if and only if it is given by a unitary representation of $\pi_1(X \setminus D)$, where D is the parabolic divisor ([MS], [Biq]). Since V_* is polystable, it is given by a representation ρ of $\pi_1(X \setminus D)$. The parabolic vector bundle $(V_*)^{\otimes m} \otimes^p ((V_*)^*)^{\otimes n}$ is given by the representation $\rho^{\otimes m} \otimes (\bar{\rho})^{\otimes n}$. This implies that $(V_*)^{\otimes m} \otimes^p ((V_*)^*)^{\otimes n}$ is polystable. □

Corollary 10.2. *Take V_* as in Lemma 10.1. Take any homomorphism*

$$\rho : \text{GL}(r, \mathbb{C}) \rightarrow \text{GL}(N, \mathbb{C}),$$

where r is the rank of V_* . Let W_* be the parabolic vector bundle associated to V_* for ρ . Then W_* is also polystable.

Proof. Consider \mathbb{C}^N as a $\mathrm{GL}(r, \mathbb{C})$ -module using ρ and the standard representation of $\mathrm{GL}(N, \mathbb{C})$. This $\mathrm{GL}(r, \mathbb{C})$ -module \mathbb{C}^N is a direct summand of a direct sum of $\mathrm{GL}(r, \mathbb{C})$ -modules of the form $(\mathbb{C}^r)^{\otimes m_i} ((\mathbb{C}^r)^*)^{\otimes n_i}$ ([DMOS, p. 40, Prop. 3.1(a)]). Therefore, from Lemma 10.1 we conclude that W_* is a direct summand of a polystable parabolic vector bundle of parabolic degree 0. Hence W_* is polystable. \square

We define polystability for parahoric torsors.

Definition 10.3. Let $\mathcal{G}_{\theta,x} \rightarrow X$ be a Bruhat–Tits group scheme with generic fiber G . A parahoric $\mathcal{G}_{\theta,x}$ -torsor \mathcal{E} with real weights θ is said to be polystable if for every representation $\rho : G \rightarrow \mathrm{GL}(V)$, the corresponding parabolic vector bundle $(\mathcal{E}, \theta(V))$ is polystable in the sense of Definition 7.6.

§10.2. Polystable parahoric torsors from unitary representations

In this subsection we will first assume that $D = \{x\}$ is a single point. The multi-point case is actually a straightforward generalization.

The complement $X \setminus \{x\}$ will be denoted by Y . For a base point $y_0 \in Y$, set $\Gamma = \pi_1(Y, y_0)$. Choose an analytic disc $U \subset X$ around x such that $y_0 \in U$. The inclusion of $U \setminus \{x\}$ in Y produces an inclusion $\pi_1(U \setminus \{x\}, y_0) \hookrightarrow \Gamma$. Using the orientation of $U \setminus \{x\}$, the group $\pi_1(U \setminus \{x\}, y_0)$ gets identified with \mathbb{Z} . The element of $\pi_1(U \setminus \{x\}, y_0)$ corresponding to $1 \in \mathbb{Z}$ will be denoted by γ .

We now recall a description of the set of conjugacy classes in a compact semisimple and simply connected group in terms of the Weyl alcove (see [BS, p. 9]). Let $K_G \subset G$ be a fixed maximal compact subgroup and T a fixed maximal torus in K_G . The corresponding Weyl group is the quotient $N_G(T)/T$. The set of conjugacy classes of elements in K_G gets identified with the T/W , which is in fact the Weyl alcove because any element of K_G is conjugate to an element in the maximal torus up to an element of the Weyl group (cf. [Mor, p. 151]). Given any $t \in K_G$, let θ_t denote the point in the Weyl alcove corresponding to t .

Given any homomorphism

$$\rho : \Gamma \rightarrow K_G,$$

let E_ρ be the flat principal G -bundle on Y associated to it. To construct E_ρ , let (\tilde{Y}, \tilde{y}_0) be the pointed universal cover of Y corresponding to the base point y_0 ; note that Γ acts on \tilde{Y} . Identify two points $(y_1, g_1), (y_2, g_2) \in \tilde{Y} \times G$ if there is an element $\gamma \in \Gamma$ such that $(y_2, g_2) = (y_1\gamma, \rho(\gamma^{-1})g_2)$. The quotient of $\tilde{Y} \times G$ is

a principal G -bundle on Y , and is denoted by E_ρ ; the right-translation action of G on $\tilde{Y} \times G$ produces an action of G on E_ρ . The flat connection on the trivial principal G -bundle $\tilde{Y} \times G \rightarrow \tilde{Y}$ descends to a flat connection on E_ρ . For any $h \in K_G$, the map

$$\tilde{Y} \times G \rightarrow \tilde{Y} \times G, \quad (y, z) \mapsto (y, \rho(h)z)$$

descends to an isomorphism

$$(10.1) \quad E_{h\rho h^{-1}} \xrightarrow{\sim} E_\rho$$

as flat principal G -bundles on Y .

Let

$$t = \rho(\gamma) \in K_G$$

be the image of γ . Since $h.\rho.h^{-1}(\gamma) = hth^{-1}$ for all $h \in K_G$, there is the map

$$\text{Hom}(\Gamma, K_G)/K_G \rightarrow \mathcal{A}, \quad \rho \mapsto \theta_t.$$

After conjugating ρ by an element of K_G , we may assume that t belongs to a fixed maximal torus T of K_G .

Let $\mathfrak{t} \in \text{Lie}(T)$ be such that

$$(10.2) \quad \exp(-2\pi\sqrt{-1}\mathfrak{t}) = t,$$

where \exp denotes the exponential map on the Lie algebra $\text{Lie}(T)$. Consider the trivial principal G -bundle $(U \setminus \{x\}) \times G$ over $U \setminus \{x\}$. The trivial connection on it given by this trivialization will be denoted by d_0 . On $(U \setminus \{x\}) \times G$, we now have the flat connection

$$(10.3) \quad \hat{d} = d_0 + \frac{\mathfrak{t} dz}{z},$$

where z is a holomorphic coordinate function on U with $z(x) = 0$.

Restrict the representation ρ to the subgroup $\pi_1(U \setminus \{x\}, y_0)$. This produces a flat principal G -bundle $E_\rho(\infty) \rightarrow U \setminus \{x\}$. Note that

$$(10.4) \quad E_\rho|_{U \setminus \{x\}} \simeq E_\rho(\infty)$$

as flat principal G -bundles. Both the flat principal G -bundles in (10.4) are isomorphic to the flat principal G -bundle $((U \setminus \{x\}) \times G, \hat{d})$ constructed in (10.3). This is because all of them have the same residue, namely \mathfrak{t} , at x . Recall that the residue determines the conjugacy class of the monodromy (see (8.4)).

To the element $t \in K_G \cap T$, we have the associated conjugacy class $\theta_t \in \mathcal{A}$, and hence by Bruhat–Tits theory have a group scheme \mathcal{G}_{θ_t} on a formal neighborhood $\hat{U} = \text{Spec } \mathbb{C}[[t]]$ of x . This group scheme \mathcal{G}_{θ_t} is the trivial group scheme

$\text{Spec } \mathbb{C}((t)) \times G$ over $\widehat{U} \setminus x = \text{Spec } \mathbb{C}((t))$. Therefore, we can extend \mathcal{G}_{θ_t} uniquely to X by setting it to be the trivial group scheme $Y \times G$ over Y ([BS, Sect. 5.2, p. 28]). We denote this group scheme on X by $\mathcal{G}_{t,X}$.

We observe that there is a morphism $\widehat{U} \hookrightarrow U$, where one identifies the formal power series ring with the completion of the convergent power series ring. Consider the trivial \mathcal{G}_{θ_t} -torsor on \widehat{U} , which we denote by P_t . Note that the connection on $E_\rho(\infty)$ restricts to a natural connection on $P_t|_{\widehat{U} \setminus \{x\}}$.

We now patch together (using, e.g., [BS, 5.2.3]) the trivial \mathcal{G}_{θ_t} -torsor P_t on \widehat{U} and the principal G -bundle E_ρ over Y along the intersection

$$\widehat{U} \setminus \{x\} = \text{Spec } \mathbb{C}((t))$$

such that the patching is connection preserving; as noted above, on $\text{Spec } \mathbb{C}((t))$, both the principal G -bundles are the trivial principal G -bundle $\text{Spec } \mathbb{C}((t)) \times G$ equipped with the connection \widehat{d} .

The above construction is summarized in the following proposition.

Proposition 10.4. *Given any homomorphism $\rho : \Gamma \rightarrow K_G$, and any $\mathfrak{t} \in \text{Lie}(T)$ satisfying (10.2), the flat principal G -bundle on Y has a canonical extension to a $\mathcal{G}_{t,X}$ -torsor over X .*

It should be clarified that the $\mathcal{G}_{t,X}$ -torsor in Proposition 10.4 depends on the choice of \mathfrak{t} (a branch of the logarithm), while the isomorphism class of the Bruhat–Tits group scheme $\mathcal{G}_{t,X}$ depends only on the conjugacy class $[t]$ of t . For $G = \text{GL}(r, \mathbb{C})$, if the logarithm \mathfrak{t} is chosen as in [MS] (meaning \mathfrak{t} is semisimple and eigenvalues are nonnegative and less than 1), then the construction in Proposition 10.4 coincides with the construction in [MS] of a parabolic vector bundle from a homomorphism $\Gamma \rightarrow \text{U}(r)$. This follows by comparing the two constructions.

The $\mathcal{G}_{t,X}$ -torsor in Proposition 10.4 will be denoted by $E_\rho(t)$.

We mention that we may restrict the connection \widehat{d} in (10.3) to $\widehat{U} \setminus \{x\} \subset U \setminus \{x\}$, where \widehat{U} as before is the formal completion along x . A $\mathcal{G}_{t,X}$ -torsor on X can be trivialized over both Y and \widehat{U} . Conversely, given $\mathcal{G}_{t,X}$ -torsors on Y and \widehat{U} , and an isomorphism between them over $\widehat{U} \setminus \{x\}$, we get a $\mathcal{G}_{t,X}$ -torsor on X . Therefore, the connection over $\widehat{U} \setminus \{x\}$ is enough to construct the $\mathcal{G}_{t,X}$ -torsor $E_\rho(t)$.

§10.3. Polystable parahoric torsors and unitary representations

As before, fix a maximal compact subgroup K_G of G .

Theorem 10.5. *Let (\mathcal{E}, θ) be a parahoric $\mathcal{G}_{\theta,X}$ -torsor on X with arbitrary real weights $\theta \in \mathcal{A}_T^m$. Then (\mathcal{E}, θ) is polystable if and only if (\mathcal{E}, θ) is given by a homomorphism from $\pi_1(X \setminus D)$ to K_G as described in Section 10.2.*

Proof. First assume that (\mathcal{E}, θ) is given by a homomorphism

$$\beta : \pi_1(X \setminus D) \longrightarrow K_G.$$

Let $\rho : G \longrightarrow \mathrm{GL}(V)$ be any homomorphism. Fix a maximal compact subgroup $K_{\mathrm{GL}(V)}$ of $\mathrm{GL}(V)$ such that $\beta(K_G) \subset K_{\mathrm{GL}(V)}$. Then the parabolic vector bundle W_* associated to (\mathcal{E}, θ) for ρ is given by the homomorphism

$$\rho \circ \beta : \pi_1(X \setminus D) \longrightarrow K_{\mathrm{GL}(V)}.$$

Therefore, this associated parabolic vector bundle W_* is polystable.

To prove the converse, assume that (\mathcal{E}, θ) is polystable. Let $\mathcal{E}(\mathfrak{g})$ and $\mathcal{E}(\mathfrak{g} \otimes \mathfrak{g})$ be the parabolic vector bundles associated to (\mathcal{E}, θ) for the G -modules \mathfrak{g} and $\mathfrak{g} \otimes \mathfrak{g}$ respectively. From Definition 10.3 we know that both $\mathcal{E}(\mathfrak{g})$ and $\mathcal{E}(\mathfrak{g} \otimes \mathfrak{g})$ are polystable of parabolic degree 0. If $\mathcal{E}(\mathfrak{g})$ is given by a homomorphism β from $\pi_1(X \setminus D)$ to a maximal compact subgroup of $\mathrm{GL}(\mathfrak{g})$, then $\mathcal{E}(\mathfrak{g} \otimes \mathfrak{g})$ is given by $\beta \otimes \beta$.

For any two parabolic vector bundles given by unitary representations of $\pi_1(X \setminus D)$, any homomorphism between them is given by a homomorphism of $\pi_1(X \setminus D)$ -modules. Let

$$\gamma : \mathcal{E}(\mathfrak{g} \otimes \mathfrak{g}) \longrightarrow \mathcal{E}(\mathfrak{g})$$

be the homomorphism of parabolic vector bundles given by the Lie bracket $\mathfrak{g} \otimes \mathfrak{g} \longrightarrow \mathfrak{g}$. From the above statement we conclude that γ is given by a homomorphism of $\pi_1(X \setminus D)$ -modules. This implies that the connection on $\mathcal{E}(\mathfrak{g})$ is induced by a connection on (\mathcal{E}, θ) (see Definition 8.5). Therefore, (\mathcal{E}, θ) is given by a homomorphism from $\pi_1(X \setminus D)$ to K_G . \square

A homomorphism $\rho : \pi_1(X \setminus D) \longrightarrow K_G$ is called irreducible if $\rho(\pi_1(X \setminus D))$ is not contained in some proper parabolic subgroup of G . A homomorphism ρ is irreducible if and only if the space of invariants in \mathfrak{g} for the adjoint action of $\rho(\pi_1(X \setminus D))$ is the zero element.

Corollary 10.6. *Let (\mathcal{E}, θ) be a parahoric $\mathcal{G}_{\theta, X}$ -torsor on X with arbitrary real weights $\theta \in \mathcal{A}_T^m$. Then (\mathcal{E}, θ) is stable if and only if (\mathcal{E}, θ) is given by an irreducible homomorphism from $\pi_1(X \setminus D)$ to K_G as described in Section 10.2.*

Proof. Assume that (\mathcal{E}, θ) is stable. Therefore, (\mathcal{E}, θ) is polystable. If the homomorphism $\rho : \pi_1(X \setminus D) \longrightarrow K_G$ corresponding to (\mathcal{E}, θ) has the property that $\rho(\pi_1(X \setminus D))$ is contained in a proper parabolic subgroup P of G , then the reduction of \mathcal{E} to P over $X \setminus D$ contradicts the stability of (\mathcal{E}, θ) . Therefore, ρ is irreducible.

Conversely, for a polystable (\mathcal{E}, θ) , if the corresponding homomorphism $\rho : \pi_1(X \setminus D) \rightarrow K_G$ is irreducible, then the polystable parabolic vector bundle $\mathcal{E}(\mathfrak{g})$ does not admit any holomorphic section ([Si, p. 744, Thm. 3]). Consequently, the polystable torsor (\mathcal{E}, θ) is stable. \square

Remark 10.7. We note that the Corlette–Donaldson–Hitchin–Simpson correspondence between flat G -bundles and G -Higgs bundles also extends to the parahoric case. When $G = \mathrm{GL}(n, \mathbb{C})$ this was proved by Simpson in [Si]. This result of [Si] is the key ingredient in this extension for general G . Using this result the question for G is reduced to one on vector bundles using the adjoint representation of G . The approach in the present paper then goes through without any essential difficulty.

Remark 10.8. The paper [BGM] considers the problem of parabolic Higgs G -bundle and the Corlette–Donaldson–Hitchin–Simpson correspondence on curves from a somewhat different perspective and also consider real representations.

Remark 10.9. The Atiyah–Weil criterion ([At], [We], [AB]) for the existence of a holomorphic connection on a holomorphic principal G -bundle generalizes to \mathcal{G} -torsors. The proof in [AB] has a straightforward generalization. Similarly, the Atiyah–Krull–Schmidt reduction of a holomorphic principal G -bundle ([BBN2]) generalizes to \mathcal{G} -torsors.

Remark 10.10. Theorem 10.5 evidently generalizes to the situation where G is a product of simple and simply connected groups. The more general case of semisimple groups G that are not simply connected is covered by using twisted bundles as in [BLS]. For a reductive group G , the natural map $G \rightarrow G/Z_0(G) \times (G/[G, G])$ is surjective with finite kernel, where $Z_0(G)$ is the connected component of the center of G containing the identity element. Since $G/Z_0(G)$ is semisimple and $/[G, G]$ is a product of copies of \mathbb{G}_m , to prove Theorem 10.5 it suffices to prove it for \mathbb{G}_m . But this was done in [Si].

§10.4. The reductive case

We now indicate briefly how to extend the consideration of (semi)stability of torsors in the case when the structure group G is a connected reductive algebraic group and identify it with the space of homomorphisms from π to K_G . However, the corresponding relationship with parahoric torsors for reductive G needs a closer analysis of Bruhat–Tits theory for reductive groups.

Let $S = [G, G]$ be the derived group, i.e., the *maximal connected semisimple subgroup* of G . Let Z_0 be the connected component of the center of G (which is a

torus) and one knows that S and Z_0 together generate G . Let $H = Z_0 \times S$. Then in fact, $H \rightarrow G$ is a finite covering map. It is easy to see (following [Ra, p. 145]) that (Γ, H) -bundles give rise to (Γ, G) -bundles and the stability and semistability of the associated (Γ, G) -bundles follow immediately from those of the (Γ, H) -bundles.

Thus, the problem of handling the reductive group H reduces to the problem of handling the semisimple group G that is *not simply connected*. On the side of Bruhat–Tits group schemes and parahoric group schemes, for a general connected reductive G the existence of Bruhat–Tits group schemes are well known and would give the existence of similar group schemes on the whole of X . There are several technical issues, which we have avoided, in the setting of the Bruhat–Tits buildings. Canonical choices of apartments and alcoves that give a transparent meaning to the association of conjugacy classes with weights in the alcove would need technical modifications leading us too far afield; future considerations would therefore need a careful discussion of a “canonical” choice of apartment as, for example, indicated in [Tits, p. 32].

Acknowledgements

We are very grateful to the referee for very detailed comments. A portion of Section 1.1 was formulated by the referee. The first two authors acknowledge partial support from the J. C. Bose Fellowship. The first author thanks the Department of Mathematics, Pondicherry University for its hospitality where this work was completed.

References

- [At] M.F. Atiyah, Complex analytic connections in fibre bundles, *Trans. Amer. Math. Soc.* **85** (1957) 181–207. [Zbl 0078.16002](#) [MR 0086359](#)
- [AB] H. Azad and I. Biswas, On holomorphic principal bundles over a compact Riemann surface admitting a flat connection, *Math. Ann.* **322** (2002), 333–346. [Zbl 0999.32010](#) [MR 1893920](#)
- [BBN1] V. Balaji, I. Biswas and D. S. Nagaraj, Principal bundles over projective manifolds with parabolic structure over a divisor, *Tohoku Math. J.* **53** (2001), 337–368. [Zbl 1070.14506](#) [MR 1844373](#)
- [BBN2] V. Balaji, I. Biswas and D. S. Nagaraj, Krull-Schmidt reduction for principal bundles, *J. reine angew. Math.* **578** (2005), 225–234. [Zbl 1084.14041](#) [MR 2113896](#)
- [BS] V. Balaji and C. S. Seshadri, Moduli of parahoric \mathcal{G} -torsors on a compact Riemann surface, *J. Alg. Geom.* **24** (2015), 1–49. [arXiv:1009.3485](#) [[math.AG](#)] [Zbl 1330.14059](#) [MR 3275653](#)
- [BLS] A. Beauville, Y. Laszlo and C. Sorger, The Picard group of the moduli of G -bundles on a curve, *Compos. Math.* **112** (1998), 183–216. [Zbl 0976.14024](#) [MR 1626025](#)
- [Biq] O. Biquard, Fibrés paraboliques stables et connexions singulières plates, *Bull. Soc. Math. France* **119** (1991), 231–257. [Zbl 0769.53013](#) [MR 1116847](#)

- [BGM] O. Biquard, O. Garcia-Prada, I. Mundet i Riera, Parabolic Higgs bundles and representations of the fundamental group of a punctured surface into a real group. [arXiv:1510.04207v1](#)
- [Bis] I. Biswas, Parabolic bundles as orbifold bundles, *Duke Math. J.* **88** (1997), 305–325. [Zbl 0955.14010](#) [MR 1455522](#)
- [Bo] P. Boalch, Riemann-Hilbert for tame complex parahoric connections, *Transform. Groups* **16** (2011), 27–50. [arXiv:1003.3177](#) [[math.DG](#)] [Zbl 1232.34117](#) [MR 2785493](#)
- [BLR] S. Bosch, W. Lutkebohmert and M. Raynaud, *Néron models*, *Ergebnisse der Mathematik und ihrer Grenzgebiete* 21, Springer, Berlin-New York, 1990. [Zbl 0705.14001](#) [MR 1045822](#)
- [BT1] F. Bruhat and J. Tits, Groupes réductifs sur un corps local, I: Données radicielles valuées, *Publ. Math. Inst. Hautes Études Sci.* **41** (1972), 5–251. [Zbl 0254.14017](#) [MR 0327923](#)
- [BT2] F. Bruhat and J. Tits, Groupes réductifs sur un corps local II, Schémas en groupes, *Publ. Math. Inst. Hautes Études Sci.* **60** (1984), 5–184. [Zbl 0597.14041](#) [MR 0756316](#)
- [De] P. Deligne, *Équations différentielles à points singuliers réguliers*, *Lecture Notes in Mathematics* 163, Springer, Berlin-New York, 1970. [Zbl 0244.14004](#) [MR 0417174](#)
- [DMOS] P. Deligne, J. S. Milne, A. Ogus and K.-y. Shih, *Hodge cycles, motives, and Shimura varieties*, *Lecture Notes in Mathematics* 900, Springer, Berlin-New York, 1982. [Zbl 0465.00010](#) [MR 0654325](#)
- [DG] M. Demazure and A. Grothendieck (eds.), *Séminaire de géométrie algébrique du Bois-Marie 1962/1964 (SGA 3), Schémas en groupes I*, *Lecture Notes in Mathematics* 151 (in French) Springer, Berlin-New York, 1970. [Zbl 0207.51401](#)
- [Ed] B. Edixhoven, Néron models and tame ramification, *Compos. Math.* **81** (1992), 291–306. [Zbl 0759.14033](#) [MR 1149171](#)
- [He1] J. Heinloth, Uniformization of \mathcal{G} -bundles, *Math. Ann.* **347** (2010), 499–528. [Zbl 1193.14014](#) [MR 2640041](#)
- [He2] J. Heinloth, Hilbert-Mumford stability on algebraic stacks and applications to \mathcal{G} -bundles on curves. [arXiv:1609.06058](#)
- [Hu] J. E. Humphreys, *Linear algebraic groups*, *Graduate Texts in Mathematics* 21, Springer, New York, Heidelberg, Berlin, 1987. [Zbl 0471.20029](#) [MR 0396773](#)
- [La] E. Landvogt, Some functorial properties of the Bruhat-Tits building, *J. reine angew. Math.* **518** (2000), 213–241. [Zbl 0937.20026](#) [MR 1739403](#)
- [MY] M. Maruyama and K. Yokogawa, Moduli of parabolic stable sheaves, *Math. Ann.* **293** (1992), 77–99. [Zbl 0735.14008](#) [MR 1162674](#)
- [MS] V. Mehta and C. S. Seshadri, Moduli of vector bundles on curves with parabolic structures, *Math. Ann.* **248** (1980), 205–239. [Zbl 0454.14006](#) [MR 0575939](#)
- [Mor] J. W. Morgan, Holomorphic bundles over elliptic manifolds, in *School on algebraic geometry (Trieste 1999)*, *ICTP Lecture Notes* 1 (2000), 135–203. [Zbl 0995.32014](#) [MR 1795863](#)
- [Na] M. Namba, *Branched coverings and algebraic functions*, *Pitman Research Notes in Mathematics Series* 161, Longman Scientific & Technical, Harlow; John Wiley & Sons, New York, 1987. [Zbl 0706.14017](#) [MR 0933557](#)
- [NS] M. S. Narasimhan and C. S. Seshadri, Stable and unitary vector bundles on a compact Riemann surface, *Ann. of Math.* **82** (1965), 540–567. [Zbl 0171.04803](#) [MR 0184252](#)
- [No] M. V. Nori, The fundamental group scheme, *Proc. Indian Acad. Sci. Math. Sci.* **91** (1982), 73–122. [Zbl 0586.14006](#) [MR 0682517](#)

- [PR1] G. Pappas and M. Rapoport, Twisted loop groups and their affine flag varieties, *Adv. Math.* **219** (2008), 118–198. [Zbl 1159.22010](#) [MR 2435422](#)
- [PR2] G. Pappas and M. Rapoport, Some questions about G -bundles on curves, in *Algebraic and arithmetic structures of moduli spaces (Sapporo 2007)*, *Advanced Studies in Pure Mathematics* 58, Math. Soc. Japan, Tokyo, 2010, 159–171. [Zbl 1213.14028](#) [MR 2676160](#)
- [Ra] A. Ramanathan, Stable principal bundles on a compact Riemann surface, *Math. Ann.* **213** (1975), 129–152. [Zbl 0284.32019](#) [MR 0369747](#)
- [Se1] C. S. Seshadri, Moduli of π -vector bundles over an algebraic curve, *Questions on algebraic varieties, C.I.M.E (Varenna 1969)*, 1970, 139–260. (For a new corrected version see <http://www.cmi.ac.in/balaji/Publications/CSS1.19.pdf>). [Zbl 0209.24503](#) [MR 0280496](#)
- [Se2] C. S. Seshadri, Remarks on parabolic structures, in *Vector bundles and complex geometry. Conference in honor of S. Ramanan (Madrid 2008)*, *Contemporary Mathematics* 522, 2010, 171–182. [Zbl 1267.14044](#) [MR 2681729](#)
- [Si] C. T. Simpson, Harmonic bundles on noncompact curves, *J. Amer. Math. Soc.* **3** (1990), 713–770. [Zbl 0713.58012](#) [MR 1040197](#)
- [Sp] T. A. Springer, *Linear algebraic groups*, 2nd ed., *Progress in Mathematics* 9, Birkhäuser Boston, Boston, MA, 1998. [Zbl 0927.20024](#) [MR 1642713](#)
- [TW] C. Teleman and C. Woodward, Parabolic bundles, products of conjugacy classes, and Gromov-Witten invariants, *Ann. Inst. Fourier* **3** (2003), 713–748. [Zbl 1041.14025](#) [MR 2008438](#)
- [Tits] J. Tits, Reductive groups over local fields, in *Automorphic forms, representations and L-functions*, *Proc. Symp. Pure Math.* 33, Amer. Math. Soc., Providence, RI, 1979, 29–69. [Zbl 0415.20035](#) [MR 0546588](#)
- [We] A. Weil, Généralisation des fonctions abéliennes, *J. Math. Pures Appl.* **17** (1938), 47–87. [Zbl 64.0361.02](#)