

Hydra groups

Will Dison and Timothy R. Riley

Abstract. We give examples of $CAT(0)$, biautomatic, free-by-cyclic, one-relator groups which have finite-rank free subgroups of huge (Ackermannian) distortion. This leads to elementary examples of groups whose Dehn functions are similarly extravagant. This behaviour originates in manifestations of *Hercules-versus-the-hydra battles* in string-rewriting.

Mathematics Subject Classification (2010). 20F65, 20F10, 20F67.

Keywords. Free-by-cyclic group, subgroup distortion, Dehn function, hydra.

1. Introduction

1.1. Hercules versus the hydra. Hercules' second labour was to fight the Lernaean hydra, a beast with multiple serpentine heads enjoying magical regenerative powers: whenever a head was severed, two grew in its place. Hercules succeeded with the help of his nephew, Iolaus, who stopped the regrowth by searing the stumps with a burning torch after each decapitation. The extraordinarily fast-growing functions we will encounter in this article stem from a re-imagining of this battle.

For us, a *hydra* will be a finite-length *positive* word on the alphabet a_1, a_2, a_3, \dots – that is, it includes no inverse letters $a_1^{-1}, a_2^{-1}, a_3^{-1}, \dots$. Hercules fights a hydra by striking off its first letter. The hydra then regenerates as follows: each remaining letter a_i , where $i > 1$, becomes $a_i a_{i-1}$ and the a_1 are unchanged. This process – removal of the first letter and then regeneration – repeats, with Hercules victorious *when* (not *if*!) the hydra is reduced to the empty word ε .

For example, Hercules defeats the hydra $a_2 a_3 a_1$ in five strikes:

$$a_2 a_3 a_1 \rightarrow a_3 a_2 a_1 \rightarrow a_2 a_1 a_1 \rightarrow a_1 a_1 \rightarrow a_1 \rightarrow \varepsilon.$$

(Each arrow represents the removal of the first letter and then regeneration.)

Proposition 1.1. *Hercules defeats all hydra.*

Proof. When fighting a hydra in which the highest index present is k , no a_i with $i > k$ will ever appear, and nor will any new a_k . The prefix before the first a_k is itself

a hydra, which, by induction, we can assume Hercules defeats. Hercules will then remove that a_k , decreasing the total number of a_k present. It follows that Hercules eventually wins. \square

However these battles are of extreme duration. Define $\mathcal{H}(w)$ to be the number of strikes it takes Hercules to vanquish the hydra w , and for integers $k \geq 1, n \geq 0$, define $\mathcal{H}_k(n) := \mathcal{H}(a_k^n)$. We call the \mathcal{H}_k *hydra functions*. Here are some values of $\mathcal{H}_k(n)$.

	1	2	3	4	...	n	...
1	1	2	3	4	...	n	...
2	1	3	7	15	...	$2^n - 1$...
3	1	4	46	$3(2^{46}) - 2$
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
k	1	$k + 1$	⋮	⋮	⋮	⋮	⋮
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮

To see that $\mathcal{H}_2(n) = 2^n - 1$ for all n , note that

$$\mathcal{H}(a_2^{n+1}) = \mathcal{H}(a_2^n) + \mathcal{H}(a_2 a_1^{\mathcal{H}(a_2^n)}) = 2\mathcal{H}(a_2^n) + 1.$$

And $\mathcal{H}_3(n)$ is essentially an n -fold iterated exponential function because, for all $n > 0$,

$$\mathcal{H}_3(n + 1) = 3(2^{\mathcal{H}_3(n)}) - 2,$$

by the calculations

$$\mathcal{H}(a_3^{n+1}) = \mathcal{H}(a_3^n) + 1 + \mathcal{H}(a_2 a_1 a_2 a_1^2 \dots a_2 a_1^{\mathcal{H}(a_3^n)}),$$

$$\mathcal{H}(a_2 a_1 a_2 a_1^2 \dots a_2 a_1^m) = 3(2^m) - m - 3.$$

Extending this line of reasoning, we will derive relationships (15) and (19) in Section 3 from which it will follow, for example, that

$$\mathcal{H}_4(3) = 3 \left(2 \left(3 \left(2 \left(3 \left(2^{3(2^{25})-1} \right) - 1 \right) - 1 \right) - 1 \right) - 1 \right) - 1.$$

So these functions are extremely wild. The reason behind the fast growth is a nested recursion. What we have is a variation on Ackermann’s functions $A_k: \mathbb{N} \rightarrow \mathbb{N}$, defined for integers $k, n \geq 0$ by

$$A_0(n) = n + 2 \quad \text{for } n \geq 0,$$

$$A_k(0) = \begin{cases} 0 & \text{for } k = 1, \\ 1 & \text{for } k \geq 2, \end{cases}$$

and

$$A_{k+1}(n+1) = A_k(A_{k+1}(n)) \quad \text{for } k, n \geq 0.$$

So, in particular, $A_1(n) = 2n$, $A_2(n) = 2^n$ and $A_3(n) = \exp_2^{(n)}(1)$, the n -fold iterated power of 2. (Definitions of Ackermann's functions occur with minor variations in the literature.) Ackermann's functions are representatives of the successive levels of the Grzegorzcyk hierarchy, which is a grading of all primitive recursive functions – see, for example, [37].

We will prove the following relationship in Section 3. Our notation in this proposition and henceforth is that for $f, g: \mathbb{N} \rightarrow \mathbb{N}$, we write $f \preceq g$ when there exists $C > 0$ such that for all n we have $f(n) \leq Cg(Cn + C) + Cn + C$. This gives an equivalence relation capturing qualitative agreement of growth rates: $f \simeq g$ if and only if $f \preceq g$ and $g \preceq f$.

Proposition 1.2. *For all $k \geq 1$, $\mathcal{H}_k \simeq A_k$.*

Other hydra dwell in the mathematical literature, particularly in the context of results concerning independence from Peano arithmetic and other logical systems. The hydra of Kirby and Paris [27], based on finite rooted trees, are particularly celebrated. Similar, but yet more extreme hydra were later constructed by Buchholz [14]. And creatures that, like ours, are finite strings that regenerate on *decapitation* were defined by Hamano and Okada [25] and then independently by Beklemishev [7]. They go by the name of *worms*, are descended from Buchholz's hydra, involve more complex regeneration rules, and withstand Hercules even longer.

1.2. Wild subgroup distortion. The distortion function $\text{Dist}_H^G: \mathbb{N} \rightarrow \mathbb{N}$ for a subgroup H with finite generating set T inside a group G with finite generating set S compares the intrinsic word metric d_T on H with the extrinsic word metric d_S :

$$\text{Dist}_H^G(n) := \max \{ d_T(1, g) \mid g \in H \text{ with } d_S(1, g) \leq n \}.$$

Up to \simeq it does not depend on the particular finite generating sets used.

A manifestation of our Hercules-versus-the-hydra battle leads to the result that even for apparently benign G and H , distortion can be wild.

Theorem 1.3. *For each integer $k \geq 1$, there is a finitely generated group G_k that*

- *is free-by-cyclic,*
- *can be presented with only one defining relator,*
- *is CAT(0),*
- *and is biautomatic,*

and yet has a rank- k free subgroup H_k that is distorted like the k -th of Ackermann's functions – that is, $\text{Dist}_{H_k}^{G_k} \simeq A_k$.

This distortion of a free subgroup of a CAT(0) group stands in stark contrast to that of any abelian subgroup – they are always quasi-isometrically embedded (see Theorem 4.10 of Chapter III.Γ in [13], for example) and so no more than linearly distorted.

The distortion we achieve exceeds that found in the hyperbolic groups of Mitra [32] and the subsequent 2-dimensional CAT(−1) groups of Barnard, Brady and Dani [2]. They give families of groups that have free subgroups distorted like the iterated exponential function $\exp^{(k)}(n)$, and examples with faster growing distortion like $\exp^{(\lfloor \log_4 n \rfloor)}(1)$. Their approach is to iterate the exponential distortion of the subgroup F in certain free-by-cyclic groups $F \rtimes \mathbb{Z}$.

In contrast to those of Mitra and of Barnard, Brady and Dani, our examples contain \mathbb{Z}^2 subgroups and so are not hyperbolic. However, in a subsequent article [10] with N. Brady we will give an elaboration of G_k that is hyperbolic and has a free subgroup distorted $\geq A_k$.

Explicitly, our examples here are

$$G_k = \langle a_1, \dots, a_k, t \mid t^{-1}a_1t = a_1, t^{-1}a_it = a_ia_{i-1} \text{ (for all } i > 1) \rangle \quad (1)$$

and their subgroups

$$H_k := \langle a_1t, \dots, a_kt \rangle.$$

So G_k is the free-by-cyclic group $F(a_1, \dots, a_k) \rtimes \mathbb{Z}$ where $\mathbb{Z} = \langle t \rangle$ and t acts by the automorphism of $F(a_1, \dots, a_k)$ that is the restriction of the automorphism θ of $F(a_1, a_2, \dots)$ defined by

$$\theta(a_i) = \begin{cases} a_1, & i = 1, \\ a_ia_{i-1}, & i > 1. \end{cases} \quad (2)$$

This automorphism of $F(a_1, \dots, a_k)$ is polynomial growing and of the type studied by Bestvina, Feighn and Handel in [9]. Indeed, our technique in Section 6 and following of using *pieces* to analyze its affect on words is also employed in [9].

For $i \leq j$, the canonical homomorphism $G_i \rightarrow G_j$ is an inclusion as the free-by-cyclic normal forms of an element of G_i and its image in G_j are the same. So the direct limit of the G_i under these inclusions is

$$G = \langle t, a_1, a_2, \dots \mid t^{-1}a_1t = a_1, t^{-1}a_it = a_ia_{i-1} \text{ (for all } i > 1) \rangle.$$

Also, the subgroup $H := \langle a_1t, a_2t, \dots \rangle$ of G is $\varinjlim H_i$ and $H_k = G_k \cap H$.

Our convention is that $[a, b] = a^{-1}b^{-1}ab$. By re-expressing the original relations as $[a_1, t] = 1$ and $a_{i-1} = [a_i, t]$ for $i > 1$ and then eliminating a_1, \dots, a_{k-1} and

defining $a := a_k$, one can present G_k with one relation, a nested commutator, known as an *Engel* relation:

$$G_k \cong \langle a, t \mid \underbrace{[a, t, \dots, t]}_k = 1 \rangle.$$

That is, the relation is $v_k = 1$ where v_k is the word defined recursively by $v_0 = a$ and $v_{i+1} = [v_i, t]$ for $i \geq 0$.

Recursively define a family of words by $u_0 = a$ and $u_{i+1} = u_i^{-1} s u_i$ for $i \geq 0$. By inducting on i , one can verify that after substituting $t^{\pm 1}$ for every $s^{\mp 1}$ in u_i , the words $t^{-(i-1)} u_i t^i$ and v_i become freely equal for all $i \geq 1$. So the relation $v_k = 1$ can be replaced by $u_k = s$ to give an alternative one-relator presentation for G_k :

$$G_k \cong \left\langle a, s \mid \underbrace{s \overset{s}{\cdot} \overset{s}{\cdot} \dots \overset{s}{\cdot}}_k = s \right\rangle.$$

That the groups G_k are CAT(0) was proved by Samuelson: set $\kappa = 1$ in Lemma 5.2 of [38]. We explain the result by re-expressing the presentation via $\alpha_i := u_{k-i}$ for $1 \leq i \leq k$ as

$$G_k \cong \langle \alpha_1, \dots, \alpha_k, s \mid \alpha_1^{-1} s \alpha_1 = s, \alpha_i^{-1} s \alpha_i = \alpha_{i-1} \ (i > 1) \rangle.$$

By checking the link condition (see, for example, [13], II.5.24) one finds that the Cayley 2-complex of this presentation (that is, the universal cover of the associated presentation 2-complex), metrized so that each 2-cell is a Euclidean square, is CAT(0). Gersten & Short [23] proved that all such groups are automatic, and later Niblo & Reeves [33] proved that a more general class of groups, those acting geometrically on CAT(0) cube complexes, are biautomatic.

The groups G_k are well-behaved in a couple of senses not mentioned in Theorem 1.3. They are residually torsion-free nilpotent by Baumslag [4]¹ and enjoy the property of *rapid decay* by Jolissaint [26], Corollary 2.1.10. We thank Gilbert Baumslag and Indira Chatterji, respectively, for these observations.

We remark that a corollary of our recursive upper bound on $\text{Dist}_{H_k}^{G_k}$ is that the membership problem for H_k in G_k is decidable.

The family G_k have received attention elsewhere. From a geometric point-of-view, it is natural to see G_k as the fundamental group of a mapping torus, and indeed G_2 is a 3-manifold group. In [22] Gersten showed the group G_2 to be CAT(0) with quadratic divergence function. He gave the free-by-cyclic, the one-relator, and the CAT(0) presentations of G_2 we have described. In [30] Macura shows G_3 to be CAT(0) and proves that an associated CAT(0) complex has a cubic divergence

¹*Added in proof:* Baumslag adds that Mikhailov should also be credited for this result and a proof is in their recent article *On residual properties of generalized Hydra groups*, [arXiv:1301.4629](https://arxiv.org/abs/1301.4629) [math.GR].

function. Results in [30] imply that the divergence function of the universal cover of the mapping torus associated to the free-by-cyclic presentation of G_k is polynomial of degree k (up to \simeq) and in [28] Macura proves the same result for CAT(0) spaces associated to each G_k . Macura also mentions G_2 and G_3 in [29] as examples in the context of Kolchin maps and quadratic isoperimetric functions, and she and Cashen use G_k as examples in [15] when studying novel quasi-isometry invariants they call *line patterns*. It is stated in Example 4 of [5] that G_3 is biautomatic. Bridson uses G_k in [12] as a starting point to construct free-by-free groups with Dehn functions that are polynomial of degree $k + 1$ and he shows them to be subgroups of $\text{Out}(F_n)$ for suitable n . Additionally, he shows his examples are asynchronously automatic via normal forms which have length $\simeq n^k$, but by no shorter normal form. En route he shows (Section 4.1 (3)) that free-by-cyclic $F_k \rtimes \mathbb{Z}$ groups, such as G_k , embed in $\text{Aut}(F_k)$.

Examples of yet more extreme distortion are known, even for subgroups of hyperbolic groups. Arzhantseva & Osin [1], §3.4, and Pittet [35] explain an argument attributed to Sela in §3, 3. K_3'' of [24]: the Rips construction, applied to a finitely presentable group with unsolvable word problem yields a hyperbolic (indeed, $C'(1/6)$ small-cancellation) group G with a finitely generated subgroup N such that Dist_N^G is not bounded above by any recursive function. The reason is that when N is a finitely generated normal subgroup of a finitely presented group G , there is an upper bound for the Dehn function of G/N in terms of the Dehn function of G and the distortion of N in G – see Corollary 8.2 in [19], [35]. Ol’shanskii & Sapir in [34], Theorem 2, provide another source of extreme examples – using Mikhailova’s construction as their starting point, they show that the set of distortion functions of finitely generated subgroups of $F_2 \times F_2$ coincides (up to \simeq) with the set of Dehn functions of finitely presented groups. As for finitely presented subgroups, Baumslag, Bridson, Miller and Short [6] explain how to construct groups Γ that are both CAT(0) and hyperbolic and yet such that $\Gamma \times \Gamma$ has a finitely presented subgroup whose distortion is not bounded above by any recursive function.

We are not aware of any systematic study of subgroup distortion in one-relator groups. It seems natural to ask whether our examples are best-possible – that is, whether there is a one-relator group with a finite-rank free subgroup of distortion $\geq A_k$ for every k .

1.3. Extreme Dehn functions. The Dehn function $\text{Area}(n)$ of a finitely presented group $\langle A \mid R \rangle$ is related to the group’s word problem in that $\text{Area}(n)$ is the minimal N such that given any word w of length at most n that represents the identity, w freely equals some product $\prod_{i=1}^{N'} u_i^{-1} r_i u_i$ of $N' \leq N$ conjugates of relators $r_i \in R^{\pm 1}$, or, equivalently, one can reduce w to the empty word by applying defining relations at most N times and removing or inserting inverse pairs of letters. At the same time, the Dehn function is a natural geometric invariant (in fact, a quasi-isometry invariant

up to \simeq) of a group: $\text{Area}(n)$ is the minimal N such that any edge-loop of length at most n in the Cayley 2-complex of $\langle A \mid R \rangle$ can be spanned by a combinatorial filling disc (a van Kampen diagram) with area (that is, number of 2-cells) at most N . This geometric perspective is related to the classical notion of an isoperimetric function in Riemannian geometry in that if $\langle A \mid R \rangle$ is the fundamental group of a closed Riemannian manifold M , then its Dehn function is \simeq -equivalent to the minimal isoperimetric function of the universal cover of M .

Theorem 1.3 leads to strikingly simple examples of finitely presented groups with huge Dehn functions, namely the HNN-extensions of G_k with stable letter commuting with all elements of the subgroup H_k .

Theorem 1.4. *For $k \geq 2$, the Dehn function of the group*

$$\Gamma_k := \langle a_1, \dots, a_k, t, p \mid t^{-1}a_1t = a_1, t^{-1}a_it = a_ia_{i-1} \ (i > 1), [p, a_it] = 1 \ (i > 0) \rangle.$$

is \simeq -equivalent to A_k .

So, together with Γ_1 , which has Dehn function \simeq -equivalent to $n \mapsto n^2$ (see Proposition 9.1), these groups have Dehn functions that are representative of each graduation of the Grzegorzczuk hierarchy of primitive recursive functions. Details of the proof are in Section 9.

These are not the only such examples (but we believe they are the first that are explicit and elementary): Cohen, Madlener and Otto [17], [18], [31] embedded algorithms (*modular Turing machines*, in fact) with running times like $n \mapsto A_k(n)$ in groups so that the running of the algorithm is displayed in van Kampen diagrams so as to make the Dehn function reflect the time-complexity of the algorithms. They state that their techniques produce yet more extreme examples as they also apply to an algorithm with running time like $n \mapsto A_n(n)$, and so yield a group with Dehn function that is recursive but not primitive recursive. More extreme still, any finitely presentable group with undecidable word problem is not bounded above by any recursive function.

Elementary examples of groups with large Dehn function are described by Gromov in [24], §4, but their behaviour is not so extreme. There is the family

$$\langle x_0, \dots, x_k \mid x_{i+1}^{-1}x_ix_{i+1} = x_i^2 \ (i < k) \rangle,$$

which has Dehn function \simeq -equivalent to $n \mapsto \exp_2^{(k)}(n)$. [We write $\exp_2(n)$ to denote 2^n .] And Baumslag's group [3]

$$\langle a, b \mid (b^{-1}a^{-1}b)a(b^{-1}ab) = a^2 \rangle, \tag{3}$$

which contains $\langle x_0, \dots, x_k \mid x_{i+1}^{-1}x_ix_{i+1} = x_i^2 \ (i \geq 0) \rangle$ as a normal subgroup, was shown by Platonov [36] to have Dehn function \simeq -equivalent to $n \mapsto \exp^{(\lceil \log_2 n \rceil)}(1)$. (Prior partial results in this direction are in [8], [20], [21].)

1.4. The organisation of the article. We believe the most compelling assertion of Theorem 1.3 to be the existence of groups H_k and G_k with H_k free of rank k , G_k enjoying the bulleted list of properties, and $\text{Dist}_{H_k}^{G_k}$ bounded below by A_k . In particular, this shows that there is no uniform upper bound on the level in the Grzegorzcyk hierarchy at which the functions $\text{Dist}_{H_k}^{G_k}$ appear. The reader who is primarily interested in these components of Theorem 1.3 need only read up to the end of Section 5. In Section 2 we derive a collection of elementary properties of the Ackermann functions that will be used elsewhere in the paper. Section 3 contains a proof of Proposition 1.2 comparing the hydra functions to Ackermann's functions. In Section 4 we prove that the subgroups H_k are free. And in Section 5 we prove that each function $\text{Dist}_{H_k}^{G_k}$ is bounded below by \mathcal{H}_k – combining this result with Proposition 1.2 gives the lower bound A_k .

Our proof that each function $\text{Dist}_{H_k}^{G_k}$ lies in the same \simeq -equivalence class of functions as A_k – i.e. that A_k is an upper bound for $\text{Dist}_{H_k}^{G_k}$ – is considerably more involved than that of the lower bound and occupies most of the second half of the article: Sections 6, 7 and 8. In deriving the upper bound, a key notion will be that of passing a power of t through a word w on the letters a_i . We explain this idea in Section 6, where we also identify recursive structure that will be crucial in facilitating an inductive analysis. In Section 7 we focus on the situation where w is of the form $\theta^n(a_k^{\pm 1})$ and derive preliminary result that will feed into the main proof, presented in Section 8, that $\text{Dist}_{H_k}^{G_k} \preceq A_k$.

Finally, in Section 9, we prove Theorem 1.4, which gives the Dehn functions of the groups Γ_k .

We illustrate some of our arguments using van Kampen diagrams, particularly observing their *corridors* (also known as *bands*). For an introduction see, for example, I.8A.4 and the proof of Proposition 6.16 in III.Γ of [13].

We denote the length of a word w by $\ell(w)$. We write $w = w(a_1, \dots, a_k)$ when w is a word on $a_1^{\pm 1}, \dots, a_k^{\pm 1}$.

1.5. Acknowledgements. We are grateful to Martin Bridson for a number of conversations on this work, to Volker Diekert for a discussion of Ackermann's functions, to Arye Juhasz for background on one-relator groups, and to John McCammond for help with some computer explorations of Hercules' battle with the hydra. We also thank an anonymous referee for a careful reading, for bringing to our attention the connections between this work and [9], and for simplifying our proof of Proposition 5.2.

2. Ackermann's functions

Throughout this article we will frequently compare functions to Ackermann's functions and will find the following relationships useful.

Lemma 2.1. *For integers k, l, m, n , the following relations hold within the given domains:*

$$A_k(A_{k+1}(n)) = A_{k+1}(n + 1), \quad k, n \geq 0, \tag{4}$$

$$A_k(1) = 2, \quad k \geq 1, \tag{5}$$

$$A_k(2) = 4, \quad k \geq 0, \tag{6}$$

$$A_k(n) \leq A_{k+1}(n), \quad k \geq 1; n \geq 0, \tag{7}$$

$$A_k(n) < A_k(n + 1), \quad k, n \geq 0, \tag{8}$$

$$n \leq A_k(n), \quad k, n \geq 0, \tag{9}$$

(with equality holding in (9) if and only if $(k, n) = (1, 0)$)

$$mA_k(n) \leq A_k(nm), \quad k, n \geq 1; m \geq 0, \tag{10}$$

$$mA_k^{(l)}(n) \leq A_k^{(l+m)}(n), \quad k \geq 1; l, m, n \geq 0, \tag{11}$$

$$A_k(n) + A_k(m) \leq A_k(n + m), \quad k, n, m \geq 1, \tag{12}$$

$$A_k(n) + m \leq A_k(n + m), \quad k, n, m \geq 0, \tag{13}$$

$$(A_k(n))^m \leq A_k(nm), \quad k \geq 2; n, m \geq 0. \tag{14}$$

Proof. Equation (4) follows immediately from the definition of the Ackermann functions. Equations (5) and (6) follow from (4) by an easy induction on k .

Before proving (7), (8) and (9), we first prove that non-strict versions of these inequalities hold. The proof is by induction on k and n . It is easy to check that (7) holds if $k = 1$ or if $n = 0$ and that (8) and (9) hold if $k = 0$, if $k = 1$ or if $n = 0$. Now let $k' > 1$ and $n' > 0$ and suppose, as an inductive hypothesis, that (7), (8) and (9) hold (not necessarily strictly) if $k < k'$ or if $k = k'$ and $n < n'$. We prove that the inequalities hold if $k = k'$ and $n = n'$. For (7), we calculate that $A_{k'}(n') = A_{k'-1}(A_{k'}(n' - 1)) \leq A_{k'-1}(A_{k'+1}(n' - 1)) \leq A_{k'}(A_{k'+1}(n' - 1)) = A_{k'+1}(n')$, where we have applied (4) and the inductive hypothesis versions of (7) and (8). For (8), we calculate that $A_{k'}(n') \leq A_{k'-1}(A_{k'}(n')) = A_{k'}(n' + 1)$, where we have used (4) and the inductive hypothesis version of (9). For (9), we calculate that $n' \leq 2n' = A_1(n') \leq A_{k'}(n')$, where we have used the inductive hypothesis version of (7). This completes the proof that (7), (8) and (9) hold in non-strict form. Now observe that equality in (9) at $(k, n) = (k', n')$ requires $n' = 2n'$, whence $n' = 0$. Since $A_k(0) = 1$ for all $k \geq 2$, equality in (9) holds if and only if $(k, n) = (1, 0)$. It follows that equality in (8) at $(k, n) = (k', n')$ would require that $A_{k'}(n') = 0$ and $k' - 1 = 1$, whence $A_2(n') = 0$. But $A_2(n) = 2^n > 0$ for all n and so the inequality (8) is strict.

We now prove inequality (10). This clearly holds if $m = 0$, so suppose that $m \geq 1$. The proof is by induction on k and n . It is clear that (10) holds if $k = 1$. The inequality also holds if $n = 1$ since, applying (5) and (7), we calculate that

$m A_k(1) = 2m = A_1(m) \leq A_k(m)$. Now let $k', n' > 1$ and suppose, as an inductive hypothesis, that (10) holds if $k < k'$ or if $k = k'$ and $n < n'$. We calculate that

$$\begin{aligned} m A_{k'}(n') &= m A_{k'-1}(A_{k'}(n' - 1)) \leq A_{k'-1}(m A_{k'}(n' - 1)) \\ &\leq A_{k'-1}(A_{k'}(m n' - m)) \leq A_{k'-1}(A_{k'}(m n' - 1)) = A_{k'}(m n'), \end{aligned}$$

where we have used (4) and (8). Thus the inequality holds if $(k, n) = (k', n')$, completing the proof of (10).

For inequality (11) observe that, by (9), $m A_k^{(l)}(n) \leq A_{k+1}(m) A_k^{(l)}(n) = A_k^{(m)}(1) A_k^{(l)}(n)$. It also follows from (9) that $A_k^{(i)}(1) \geq 1$ for all $i \geq 0$. We can thus apply (10), together with (8), to show that

$$A_k^{(m)}(1) A_k^{(l)}(n) \leq A_k^{(m)}(A_k^{(l)}(n)) = A_k^{(l+m)}(n).$$

We prove (12) by induction on k . We will make repeated use of the identity $A_k(m) = A_{k-1}^{(m)}(1)$. It is clear that the inequality holds if $k = 1$, so suppose that $k > 1$ and that the result is true for smaller values of k . Without loss of generality suppose that $n \leq m$. It follows from (9) that $A_{k-1}^{(i)} \geq 1$ for all $i \geq 0$, and so we can apply the induction hypothesis to calculate that $A_k(n) + A_k(m) = A_{k-1}^{(n)}(1) + A_{k-1}^{(m)}(1) \leq A_{k-1}^{(n)}(1 + A_{k-1}^{(m-n)}(1)) = A_{k-1}^{(n)}(1 + A_k(m - n))$. Applying (8) gives that this quantity is at most $A_{k-1}^{(n)}(A_k(m - n + 1)) = A_k(m + 1) \leq A_k(m + n)$.

We now prove inequality (13). This clearly holds if $k = 0, k = 1$ or $m = 0$. If $k \geq 2$ and $n = 0$, then $A_k(n) + m = m + 1 \leq A_k(m) = A_k(n + m)$ by (9). It remains to prove (13) if $k, n, m \geq 1$. But in this case $A_k(n) + m \leq A_k(n) + A_k(m) \leq A_k(n + m)$ by (9) and (12).

Finally, we prove (14) by induction on k . It is clear that the inequality holds if $k = 2$, so suppose that $k \geq 3$ and that the result holds for smaller values of k . It is also clear that the inequality holds if $n = 0$ or if $m = 0$; suppose that $n, m \geq 1$. Applying the induction hypothesis, together with (4), we calculate that $A_k(n)^m = A_{k-1}(A_k(n - 1))^m \leq A_{k-1}(m A_k(n - 1))$. Applying (4), (8) and (10), we see that this quantity is at most $A_{k-1}(A_k(n m - m)) \leq A_{k-1}(A_k(n m - 1)) = A_k(n m)$. \square

3. Comparing the hydra functions to Ackermann’s functions

In this section we prove Proposition 1.2 comparing Ackermann’s functions to the hydra functions. The proof will proceed via a third family of functions ϕ_k . In this section $\phi_k(n)$ will be defined for $n \geq 0$; subsequently we will give a more general definition with an expanded domain.

For integers $k \geq 1$ and $n \geq 0$, define $\phi_k(n) := \mathcal{H}(\theta^n(a_k))$. The functions \mathcal{H}_k satisfy

$$\mathcal{H}_k(n + 1) = \mathcal{H}_k(n) + \phi_k(\mathcal{H}_k(n)) \tag{15}$$

since after $\mathcal{H}_k(n)$ strikes the word a_k^{n+1} has become $\theta^{\mathcal{H}_k(n)}(a_k)$. We will need the following elementary properties of the functions ϕ_k .

Lemma 3.1. *For integers $k \geq 1$ and $n \geq 0$,*

$$\phi_k(0) = 1, \tag{16}$$

$$\phi_2(n) = n + 1, \tag{17}$$

$$\phi_k(n) \geq 1, \tag{18}$$

$$\phi_{k+1}(n + 1) = \phi_{k+1}(n) + \phi_k(\phi_{k+1}(n) + n). \tag{19}$$

For integers $k \geq 2$ and $n \geq 0$,

$$\phi_k(n) < \phi_k(n + 1), \tag{20}$$

$$\phi_k(n) \geq n. \tag{21}$$

Proof. Assertions (16), (17), (18) are straightforward. For (19), note that, by induction on n , $\theta^{n+1}(a_{k+1}) = a_{k+1}a_k\theta(a_k) \dots \theta^n(a_k)$ and hence

$$\theta^{n+1}(a_{k+1}) = \theta^n(a_{k+1})\theta^n(a_k).$$

Thus, after $\phi_{k+1}(n)$ strikes, $\theta^{n+1}(a_{k+1})$ has become

$$\theta^{\phi_{k+1}(n)}(\theta^n(a_k)) = \theta^{\phi_{k+1}(n)+n}(a_k).$$

Inequality (20) follows immediately from (18) and (19) and inequality (21) follows from (18) and (20). □

It is easy to check that $\phi_1 \simeq A_0$ and $\phi_2 \simeq A_1$. As such, the next result is sufficient to establish that $\phi_k \simeq A_{k-1}$ for $k \geq 1$.

Lemma 3.2. (i) *For integers $k \geq 3$ and $n \geq 0$, $\phi_k(n) \geq A_{k-1}(n)$.*

(ii) *For integers $k \geq 2$ and $n \geq 0$, $\phi_k(n) \leq A_{k-1}(n + k) - n - k$.*

Proof. We prove (i) by simultaneous induction on k and n . It is immediate from (16) that the inequality holds if $n = 0$. By (17) and (19), $\phi_3(n) = 2\phi_3(n - 1) + n$, which, combined with (16), gives $\phi_3(n) = 3(2^n) - n - 2$. Since $A_2(n) = 2^n$, it is easy to check that (i) holds if $k = 3$. Now let $k' > 3$ and $n' > 0$ and suppose, as an inductive hypothesis, that the result is true if $k < k'$ or if $k = k'$ and $n < n'$. Applying (4), (18) and (20), we calculate that $\phi_{k'}(n') = \phi_{k'}(n' - 1) + \phi_{k'-1}(\phi_{k'}(n' - 1) + n' - 1) \geq \phi_{k'-1}(\phi_{k'}(n' - 1)) \geq \phi_{k'-1}(A_{k'-1}(n' - 1)) \geq A_{k'-2}(A_{k'-1}(n' - 1)) = A_{k'-1}(n')$. Thus the result holds at $(k, n) = (k', n')$, completing the proof of (i).

We now make the following claim: for all $k \geq 2, n \geq 0$ and $c \geq k$,

$$\phi_k(n) \leq A_{k-1}(n + c) - n + k - 2c. \tag{22}$$

Assertion (ii) will follow by setting $c = k$. The proof of this inequality is by simultaneous induction on k and n . Since $A_1(n) = 2n$ and, by (17), $\phi_2(n) = n + 1$, it is straightforward to check that (22) holds if $k = 2$. The inequality also holds for $n = 0$ since, by (7) and (16), $\phi_k(0) = 1 \leq k = A_1(c) + k - 2c \leq A_{k-1}(c) + k - 2c$. Now let $c \geq k' > 2$ and $n' > 0$ and suppose, as an induction hypothesis, that (22) holds if $k < k'$ or if $k = k'$ and $n < n'$. We calculate that

$$\begin{aligned} \phi_{k'}(n') &= \phi_{k'}(n' - 1) + \phi_{k'-1}(\phi_{k'}(n' - 1) + n' - 1) && \text{by (19)} \\ &\leq \phi_{k'}(n' - 1) + A_{k'-2}(\phi_{k'}(n' - 1) + n' + c - 1) \\ &\quad - \phi_{k'}(n' - 1) - n' + k' - 2c \\ &= A_{k'-2}(\phi_{k'}(n' - 1) + n' + c - 1) - n' + k' - 2c \\ &\leq A_{k'-2}(A_{k'-1}(n' + c - 1) + k' - c) - n' + k' - 2c && \text{by (8)} \\ &\leq A_{k'-2}(A_{k'-1}(n' + c - 1)) - n' + k' - 2c && \text{by (8)} \\ &= A_{k'-1}(n' + c) - n' + k' - 2c. && \text{by (4)} \end{aligned}$$

Thus the inequality holds if $(k, n) = (k', n')$, completing the proof of (22). □

Since $A_1(n) = 2n$, $\mathcal{H}_1(n) = n$, $A_2(n) = 2^n$ and $H_2(n) = 2^n - 1$, the next result is sufficient to establish Proposition 1.2.

Proposition 3.3. (i) For integers $k \geq 3$ and $n \geq 2$, $\mathcal{H}_k(n) \geq A_k(n)$.

(ii) For integers $k \geq 1$ and $n \geq 0$, $\mathcal{H}_k(n) \leq A_k(n + k)$.

Proof. We prove (i) by induction on n . The inequality certainly holds for $n = 2$ since, by (6), $\mathcal{H}_k(2) = \mathcal{H}(a_k a_{k-1} a_{k-2}) \geq 4 = A_k(2)$. Now let $n' > 2$ and suppose that (i) holds for $n < n'$. Applying (4), (15) and (20), together with Lemma 3.2 (i), we calculate that $\mathcal{H}_k(n') = \mathcal{H}_k(n' - 1) + \phi_k(\mathcal{H}_k(n' - 1)) \geq \phi_k(\mathcal{H}_k(n' - 1)) \geq \phi_k(A_k(n' - 1)) \geq A_{k-1}(A_k(n' - 1)) = A_k(n')$. Thus the inequality holds for $n = n'$, completing the proof of (i).

For (ii), we prove the stronger claim that, for all $k \geq 1, n \geq 0$,

$$\mathcal{H}_k(n) \leq A_k(n + k) - k. \tag{23}$$

The proof is by simultaneous induction on k and n . Since $A_1(n) = 2n$ and $\mathcal{H}_1(n) = n$, it is straightforward to check that (23) holds if $k = 1$. The inequality holds if $n = 0$ since, by (7), $\mathcal{H}_k(0) = 0 \leq k = A_1(k) - k \leq A_k(k) - k$. Now let $k' > 1$ and $n' > 0$ and suppose, as an inductive hypothesis, that (23) holds if $k < k'$ or if $k = k'$ and $n < n'$. We calculate that

$$\begin{aligned} \mathcal{H}_{k'}(n') &= \mathcal{H}_{k'}(n' - 1) + \phi_{k'}(\mathcal{H}_{k'}(n' - 1)) && \text{by (15)} \\ &\leq \mathcal{H}_{k'}(n' - 1) + A_{k'-1}(\mathcal{H}_{k'}(n' - 1) + k') \\ &\quad - \mathcal{H}_{k'}(n' - 1) - k' && \text{by Lemma 3.2 (ii)} \end{aligned}$$

$$\begin{aligned}
 &= A_{k'-1}(\mathcal{H}_{k'}(n' - 1) + k') - k' \\
 &\leq A_{k'-1}(A_{k'}(n' + k' - 1)) - k' && \text{by (8)} \\
 &= A_{k'}(n' + k') - k' && \text{by (4)}.
 \end{aligned}$$

Thus the inequality holds if $(k, n) = (k', n')$, completing the proof of (23). □

4. Freeness of the subgroups H and H_k

In this section we prove:

Proposition 4.1. *The subgroup H_k of G_k is free with free basis a_1t, \dots, a_kt , and the subgroup H of G is free with free basis a_1t, a_2t, \dots .*

To facilitate an induction argument, we will prove the following more elaborate proposition. Proposition 4.1 will follow because if $w = w(a_1t, \dots, a_kt)$ is freely reduced and represents 1 in G_k (or, equivalently, in G), then $w = \varepsilon$ by conclusion (i), and so a_1t, \dots, a_kt are each not the identity and satisfy no non-trivial relations.

Proposition 4.2. *Let $u = u(a_1t, \dots, a_kt)$ be a freely reduced word with free-by-cyclic normal form vt^r – that is, $u = vt^r$ in G_k , $v = v(a_1, \dots, a_k)$ is reduced, and $r \in \mathbb{Z}$.*

- (i) *If $v = \varepsilon$, then $u = \varepsilon$.*
- (ii) *If $v = \theta(a_{k+1}^{-1})\theta^{1-r}(a_{k+1})$ in $F(a_1, a_2, \dots)$, then $u = \varepsilon$.*
- (iii) *If v is positive, then u is positive.*

We emphasise that we are considering u as a word on the a_it – it is freely reduced if and only if it contains no subword $(a_it)^{\pm 1}(a_it)^{\mp 1}$.

Proof of Proposition 4.2. We first show that for all fixed $k \geq 1$, if (iii) holds, then so do (i) and (ii).

For (i), note that if $u = t^r$ in G , then $u^{-1} = t^{-r}$. Thus (iii) implies that both of the freely reduced words u and u^{-1} are positive. Hence $u = \varepsilon$.

For (ii), we will separately consider the cases $r = 0$, $r < 0$, and $r > 0$. If $r = 0$, then $u = 1$ in G and hence $u = \varepsilon$ by (i). If $r < 0$, then $1 - r \geq 1$ and so $\theta^{1-r}(a_{k+1}) = a_{k+1}a_k w$ in $F(a_1, a_2, \dots)$ for some positive word $w = w(a_1, \dots, a_k)$. It follows that v is positive and therefore (iii) implies that u is positive. Thus $r \geq 0$, giving a contradiction. If $r > 0$, one calculates that $u^{-1} = t^{-r}\theta^{1-r}(a_{k+1}^{-1})\theta(a_{k+1}) = \theta(a_{k+1}^{-1})\theta^{1+r}(a_{k+1})t^{-r}$ in $F(a_1, a_2, \dots)$. Since $1 + r \geq 1$, the reduced form of $\theta(a_{k+1}^{-1})\theta^{1+r}(a_{k+1})$ is positive, and so (iii) implies that u^{-1} is positive. Thus $-r \geq 0$, giving a contradiction.

We now prove (iii) by induction on k . Since G_1 is free abelian with basis a_1, t , it is easy to check that (iii) holds in the case $k = 1$. As an inductive hypothesis, assume that assertions (i), (ii) and (iii) all hold for smaller values of k . If u contains no occurrence of an $(a_k t)^{\pm 1}$, then we are done. Otherwise, write $u = \sigma_0 (a_k t)^{\epsilon_1} \sigma_1 (a_k t)^{\epsilon_2} \dots (a_k t)^{\epsilon_m} \sigma_m$, where each $\sigma_i = \sigma_i(a_1 t, \dots, a_{k-1} t)$ and each $\epsilon_i \in \{\pm 1\}$.

Each σ_i has free-by-cyclic normal form $\tau_i t^{s_i}$ for some $\tau_i = \tau_i(a_1, \dots, a_{k-1})$ and some $s_i \in \mathbb{Z}$. Direct calculation of the normal form of u – moving all the $t^{\pm 1}$ to the right-hand end and applying the automorphism $\theta^{\mp 1}$ whenever a $t^{\pm 1}$ is moved past a letter a_i – gives that v freely equals

$$v' := \tau_0 \theta^{\lambda_1} (a_k^{\epsilon_1}) \theta^{\mu_1} (\tau_1) \theta^{\lambda_2} (a_k^{\epsilon_2}) \dots \theta^{\lambda_m} (a_k^{\epsilon_m}) \theta^{\mu_m} (\tau_m),$$

where

$$\lambda_i = \begin{cases} -(s_0 + \dots + s_{i-1} + \epsilon_1 + \dots + \epsilon_{i-1}) & \text{if } \epsilon_i = 1, \\ -(s_0 + \dots + s_{i-1} + \epsilon_1 + \dots + \epsilon_i) & \text{if } \epsilon_i = -1, \end{cases}$$

$$\mu_i = -(s_0 + \dots + s_{i-1} + \epsilon_1 + \dots + \epsilon_i).$$

We claim that $\epsilon_i = 1$ for all i . For a contradiction, suppose otherwise. Observe that, for each $s \in \mathbb{Z}$, there are words $w_s = w_s(a_1, \dots, a_{k-1})$ and $w'_s = w'_s(a_1, \dots, a_{k-1})$ such that $\theta^s(a_k) = a_k w_s$ and $\theta^s(a_k^{-1}) = w'_s a_k^{-1}$. Since v is positive, there must be a subword $a_k^{\pm 1} \chi a_k^{\mp 1}$ in v' which freely equals the empty word and in which $\chi = \chi(a_1, \dots, a_{k-1})$. The way this subword must arise is that for some i , either

- (a) $\epsilon_i = -1$, $\epsilon_{i+1} = 1$ and $\theta^{\mu_i}(\tau_i) = 1$, or
- (b) $\epsilon_i = 1$, $\epsilon_{i+1} = -1$ and $\theta^{\lambda_i}(a_k) \theta^{\mu_i}(\tau_i) \theta^{\lambda_{i+1}}(a_k^{-1}) = 1$.

In the first case $\tau_i = 1$ and hence the induction hypothesis (assertion (i)) gives that $\sigma_i = \varepsilon$. But this contradicts the supposition that u is freely reduced. In the second case, one calculates that $\lambda_i - \mu_i = 1$ and $\lambda_{i+1} - \mu_i = 1 - s_i$, and so $\tau_i = \theta(a_k^{-1}) \theta^{1-s_i}(a_k)$. The induction hypothesis (assertion (ii)) implies that $\sigma_i = \varepsilon$, but again this contradicts the supposition that u is freely reduced.

To complete our proof of (iii), we will show that all the σ_i are positive. Since v is positive and each $\epsilon_i = 1$, we have that τ_0 is positive and each $\theta^{\lambda_i}(a_k) \theta^{\mu_i}(\tau_i)$ is positive. The inductive hypothesis (assertion (iii)) immediately gives that σ_0 is positive. Suppose we have shown that $\sigma_0, \dots, \sigma_{j-1}$ are positive, for some j . It follows that $s_0, \dots, s_{j-1} \geq 0$, whence $\lambda_j \leq 0$. Note that if $w = w(a_1, \dots, a_k)$ is positive and $s \geq 0$, then $\theta^s(w)$ is positive. Hence $a_k \theta^{\mu_j - \lambda_j}(\tau_j) = a_k \theta^{-1}(\tau_j)$ is positive. Since $\theta^{-1}(\tau_j)$ is a word on $a_1^{\pm 1}, \dots, a_{k-1}^{\pm 1}$, it follows that $\theta^{-1}(\tau_j)$ is positive, and hence that τ_j is positive. Applying the induction hypothesis (assertion (iii)) gives that σ_j is positive. \square

5. A lower bound on the distortion of H_k in G_k

In the following lemma we see the battle between Hercules and the hydra manifest in G_k .

Lemma 5.1. *For all $k, n \geq 1$, there is a positive word $u_{k,n} = u_{k,n}(a_1t, \dots, a_kt)$ of length $\mathcal{H}_k(n)$ that equals $a_k^n t^{\mathcal{H}_k(n)}$ in G_k .*

Proof. Consider the following calculation in which successive t are moved to the front and paired off with the a_i . [We illustrate the calculation in the case $k \geq 3$ and $n \geq 2$ – for $k = 2$, the letters a_{k-2} would not appear and for $k = 1$, neither would the a_{k-1} .]

$$\begin{aligned}
 a_k^n t^{\mathcal{H}_k(n)} &= (a_k t) t^{-1} a_k^{n-1} t^{\mathcal{H}_k(n)-1} \\
 &= (a_k t) (a_k a_{k-1})^{n-1} t^{\mathcal{H}_k(n)-1} \\
 &= (a_k t) (a_k t) t^{-1} a_{k-1} (a_k a_{k-1})^{n-2} t^{\mathcal{H}_k(n)-2} \\
 &= (a_k t) (a_k t) a_{k-1} a_{k-2} (a_k a_{k-1} a_{k-1} a_{k-2})^{n-2} t^{\mathcal{H}_k(n)-2} \\
 &\vdots
 \end{aligned}$$

A van Kampen diagram displaying this calculation in the case $k = 2$ and $n = 4$ is shown in Figure 1.




Figure 1. A van Kampen diagram showing that $a_2^4 t^{15} = u_{2,4}$ in G_2 where $u_{2,4} = a_2 t a_2 t a_1 t a_2 t (a_1 t)^3 a_2 t (a_1 t)^7$.

One sees the Hercules-versus-the-hydra battle

$$a_k^n \rightarrow (a_k a_{k-1})^{n-1} \rightarrow a_{k-1} a_{k-2} (a_k a_{k-1} a_{k-1} a_{k-2})^{n-2} \rightarrow \dots$$

being played out in this calculation. The pairing off of a t with an a_i corresponds to a decapitation, and the conjugation by t that moves that t into place from the right-hand end causes a hydra-regeneration for the intervening subword. So by Proposition 1.1, after $\mathcal{H}_k(n)$ steps we have a positive word on $u_{k,n} = u_{k,n}(a_1t, \dots, a_k t)$, and its length is $\mathcal{H}_k(n)$. \square

Our next proposition establishes that $\text{Dist}_{H_k}^{G_k} \geq \mathcal{H}_k$ for all $k \geq 2$. The case $k = 1$ is straightforward: $H_1 \cong \mathbb{Z}$ is undistorted in $G_1 \cong \mathbb{Z}^2$ and $\mathcal{H}_1(n) = n$. The calculation in the proof of the proposition is illustrated by a van Kampen diagram in Figure 2 in the case $k = 2$ and $n = 4$ – the idea is that a copy of the diagram from Figure 1 fits together with its mirror image along intervening a_1 - and a_2 -corridors to make a diagram demonstrating the equality of a freely reduced word of extreme length on $a_1t, \dots, a_k t$ with a short word on a_1, \dots, a_k, t .

Proposition 5.2. *For all $k \geq 2$ and $n \geq 1$, there is a reduced word of length $2\mathcal{H}_k(n) + 3$ on the free basis $a_1t, \dots, a_k t$ for H_k which, in G_k , equals a word of length $2n + 4$ on a_1, \dots, a_k, t .*

Proof. As t commutes with a_1 in G_k , it also commutes with $a_2ta_1a_2^{-1}$. So

$$t^{-\mathcal{H}_k(n)} a_2 t a_1 a_2^{-1} t^{\mathcal{H}_k(n)} = a_2 t a_1 a_2^{-1} = (a_2 t) (a_1 t) (a_2 t)^{-1},$$

and then by Lemma 5.1,

$$a_k^n a_2 t a_1 a_2^{-1} a_k^{-n} = u_{k,n} (a_2 t) (a_1 t) (a_2 t)^{-1} u_{k,n}^{-1}.$$

The word on the left has length $2n + 4$. The word on the right, viewed as a word on $a_1t, \dots, a_k t$, is freely reduced and has length $2\mathcal{H}_k(n) + 3$, since $u_{k,n}$ is a positive word. \square




Figure 2. A van Kampen diagram demonstrating the equality $a_2^4 a_2 t a_1 a_2^{-1} a_2^{-4} = u_{2,4} (a_2 t) (a_1 t) (a_2 t)^{-1} u_{2,4}^{-1}$ in G_2 .

6. Recursive structure of words

This section contains preliminaries that will feed into the proof, presented in Section 8, that $\text{Dist}_{H_k}^{G_k} \preceq A_k$. Here is an outline of how we will bound the distortion of H_k in G_k . We will first suppose $u = u(t, a_1, \dots, a_k)$ represents an element of H_k . We will shuffle all the $t^{\pm 1}$ in u to the start, with the effect of applying $\theta^{\pm 1}$ to each $a_i^{\pm 1}$ they pass. After freely reducing, we will have a word $t^r w$ where $w = w(a_1, \dots, a_k)$. We will then look to carry the t^r back through w from left to right, converting all it passes to a word on $a_1 t, \dots, a_k t$. Estimating the length of this word will give an upper bound on $\text{Dist}_{H_k}^{G_k}$.

For convenience, we work with the group G and its subgroup H defined in Section 1.2.

When carrying the power of t through w we will face the problem of whether a word $t^r w$, where $w = w(a_1, a_2, \dots)$, represents an element of a coset Ht^s in G for some $s \in \mathbb{Z}$. We will see that the answer is not always affirmative – these cosets do not cover G . However, if $t^r w = \sigma t^s$ for some $\sigma = \sigma(a_1 t, a_2 t, \dots)$ and some $s \in \mathbb{Z}$, then σ is unique up to free-equivalence since H is free (Proposition 4.1) and s is unique by our next lemma. Indeed, we learn that Ht^{s_1} and Ht^{s_2} are equal precisely when $s_1 = s_2$.

Lemma 6.1. *If $\ell \in \mathbb{Z}$ and $t^\ell \in H$, then $\ell = 0$.*

Proof. Were $t^\ell \in H$ for some integer $\ell \neq 0$, then $\mathbb{Z}^2 \cong \langle a_1 t, t^\ell \rangle$ would be a subgroup of H contrary to the freeness of H established in Proposition 4.1. \square

Our next lemma will be the crux of our method for establishing an upper bound on distortion. It identifies recursive structure that will allow us to analyse the process of passing a power of t through a word $w = w(a_1, a_2, \dots)$, so as to leave behind a word on $a_1 t, a_2 t, \dots$

For a non-empty freely-reduced word $w = w(a_1, a_2, \dots)$, define the *rank of w* to be the highest k such that $a_k^{\pm 1}$ occurs in w . We define the empty word to have rank 0. For an integer $k \geq 1$, define a *piece of rank k* to be a freely-reduced word $a_k^{\epsilon_1} \pi a_k^{-\epsilon_2}$ where $\pi = \pi(a_1, \dots, a_{k-1})$ and $\epsilon_1, \epsilon_2 \in \{0, 1\}$. Notice that a piece of rank k will always also be a piece of rank $k + 1$ and that the empty word is a piece of rank k for every k .

For a non-empty freely-reduced word w of rank k , define the *number of pieces in w* to be the least integer m such that w can be expressed as a concatenation $w_1 \dots w_m$ of subwords w_i each of which is a piece of rank k . (We say the empty word is composed of 0 pieces.) Observe that

- (i) each a_k and a_k^{-1} in w is the first or last letter of some w_i , respectively;
- (ii) for $i = 1, \dots, m - 1$, either the final letter of w_i is a_k^{-1} or the first of w_{i+1} is a_k , but never both; and

(iii) if $a_k^{-1}\chi a_k$ is a subword of w and $\chi = \chi(a_1, \dots, a_{k-1})$, then $\chi = w_i$ for some i .

In particular, w_1, \dots, w_m are uniquely determined by the locations of the $a_k^{\pm 1}$ in w , and so we call the list of subwords w_1, \dots, w_m *the partition of w into pieces*.

For example, $w := a_3^{-1}a_1a_2a_3a_2^{-1}a_3a_1^{-1}a_3^{-1}$ has rank 3 and its partition into pieces is $w = w_1w_2w_3w_4$ where $w_1 = a_3^{-1}$, $w_2 = a_1a_2$, $w_3 = a_3a_2^{-1}$, and $w_4 = a_3a_1^{-1}a_3^{-1}$.

Lemma 6.2. *Suppose $w = w(a_1, \dots, a_k)$ is a non-empty freely-reduced word of rank k and r and s are integers such that $t^r w \in Ht^s$. Let $w = w_1 \dots w_n$ be the partition of w into pieces. Then there exist integers $r = r_0, r_1, \dots, r_n = s$ such that $t^{r_i} w_{i+1} \in Ht^{r_{i+1}}$ for each i .*

Proof. As $t^r w \in Ht^s$, there is some reduced word $v = v(a_1t, \dots, a_k t)$ such that $t^r w = vt^s$. Form the analogue of a *partition into pieces* for v – that is, express v as a concatenation $v_1 \dots v_m$ of subwords v_i each of the form $(a_k t)^{\epsilon_1} \tau (a_k t)^{-\epsilon_2}$ where $\tau = \tau(a_1t, \dots, a_{k-1}t)$ and $\epsilon_1, \epsilon_2 \in \{0, 1\}$ and m is minimal.

Note that v is non-empty as otherwise w would equal t^{s-r} in G and so be the empty word by the free-by-cyclic structure of G . Note also that no v_i is the empty word since m is minimal.

One can obtain $t^r w$ from vt^s by carrying all the $t^{\pm 1}$ to the left and freely reducing. More particularly, the t^s at the end of vt^s and all the $t^{\pm 1}$ in v_m can be collected immediately to the left of v_m , and then those $t^{\pm 1}$ and the $t^{\pm 1}$ in v_{m-1} can be carried to the left of v_{m-1} , and so on. Accordingly, inductively define w'_m, \dots, w'_1 and r_m, \dots, r_0 by setting $r_m := s$ and then, for $i = m, \dots, 1$, taking r_{i-1} and $w'_i = w'_i(a_1, \dots, a_k)$ to be the unique integer and reduced word such that $v_i t^{r_i} = t^{r_{i-1}} w'_i$. Then $r_0 = r$ and w is (a priori) the freely reduced form of $w'_1 \dots w'_m$. We claim that, in fact, $w'_1 \dots w'_m$ is the partition of w into pieces of rank k – that is, $m = n$ and $w'_i = w_i$ for all i . This will suffice to establish the lemma.

To prove this claim, we will show that for all i , if $v_i = (a_k t)^{\epsilon_1} \tau (a_k t)^{-\epsilon_2}$ where $\tau = \tau(a_1t, \dots, a_{k-1}t)$ and $\epsilon_1, \epsilon_2 \in \{0, 1\}$, then w'_i is a reduced word $a_k^{\epsilon_1} \pi a_k^{-\epsilon_2}$ for some $\pi = \pi(a_1, \dots, a_{k-1})$. Moreover, if $\epsilon_1 = \epsilon_2 = 0$, then π is not the empty word. In particular, no w'_i is the empty word.

Well, $v_i t^{r_i} = t^{r_{i-1}} w'_i$. Consider the process of carrying each $t^{\pm 1}$ in $v_i t^{r_i}$ to the front of the word, applying $\theta^{\pm 1}$ to each a_j they pass and then freely reducing, to give $t^{r_{i-1}} w'_i$. Throughout this process, no new $a_k^{\pm 1}$ are produced and, such is θ , no a_l appears to the left of the a_k in v_i (if present) or to the right of the a_k^{-1} (if present) – see (2) and Lemma 7.1. This means that the only way w'_i could fail to be a reduced word of the form $a_k^{\epsilon_1} \pi a_k^{-\epsilon_2}$ where $\pi = \pi(a_1, \dots, a_{k-1})$, would be for ϵ_1 and ϵ_2 to both be 1 and π be the empty word. But in that case, w'_i would be the empty word and so v_i would equal $t^{r_{i-1}-r_i}$ in G_k and $r_{i-1} - r_i$ would be 0 by Lemma 6.1. But then v_i would be the empty word by Proposition 4.1 which, as we observed, is not the

case. Likewise, when $\epsilon_1 = \epsilon_2 = 0$, it cannot be the case that $\pi = w'_i$ is the empty word, as otherwise v_i would again be the empty word.

So properties (i), (ii) and (iii) all apply to w'_1, \dots, w'_m as they are inherited the corresponding properties for v_1, \dots, v_m . It follows from these properties together with the fact that each w'_i is reduced, that $w'_1 \dots w'_m$ is reduced and is the partition of w into pieces of rank k . □

7. Passing powers of t through $\theta^n(a_k^{\pm 1})$

The words $\theta^n(a_k^{\pm 1})$ will play a crucial role in our proof that $\text{Dist}_{H^k}^{G^k} \leq A_k$. The next lemma reveals their recursive structure. The first part is proved by an induction on n . The second part is then an immediate consequence.

Lemma 7.1.

$$\theta^n(a_k) = \begin{cases} a_k \theta^0(a_{k-1}) \theta^1(a_{k-1}) \dots \theta^{n-1}(a_{k-1}), & n > 0, \\ a_k, & n = 0, \\ a_k \theta^{-1}(a_{k-1}^{-1}) \theta^{-2}(a_{k-1}^{-1}) \dots \theta^n(a_{k-1}^{-1}), & n < 0, \end{cases}$$

$$\theta^n(a_k^{-1}) = \begin{cases} \theta^{n-1}(a_{k-1}^{-1}) \theta^{n-2}(a_{k-1}^{-1}) \dots \theta^0(a_{k-1}^{-1}) a_k^{-1}, & n > 0, \\ a_k^{-1}, & n = 0, \\ \theta^n(a_{k-1}) \theta^{n+1}(a_{k-1}) \dots \theta^{-1}(a_{k-1}) a_k^{-1}, & n < 0. \end{cases}$$

When attempting to carry a power of t through a word $w = w(a_1, a_2, \dots)$, we will frequently be faced with the special case where w is of the form $\theta^n(a_k^{\pm 1})$. We now focus on this situation.

Definition 7.2. Define

$$\Lambda = \bigcup_{i \in \mathbb{Z}} Ht^i.$$

For each integer $k \geq 1$, define

$$S_k = \{n \in \mathbb{Z} : \theta^n(a_k) \in \Lambda\}$$

and define the function $\phi_k: S_k \rightarrow \mathbb{Z}$ by setting $\phi_k(n)$ to be the unique integer satisfying

$$\theta^n(a_k)t^{\phi_k(n)} \in H.$$

Note that this extends the previous definition of the functions ϕ_k given in Section 3 since $\phi_k(n) = \mathcal{H}(\theta^n(a_k))$ for $n \geq 0$.

Lemma 7.3. (i) $S_1 = \mathbb{Z}$ and $\phi_1(n) = 1$ for all $n \in S_1$.

(ii) $S_2 = \mathbb{Z}$ and $\phi_2(n) = n + 1$ for all $n \in S_2$.

(iii) If $k \geq 3$, then $S_k = \mathbb{N}$.

Proof. It is easy to check that $S_1 = S_2 = \mathbb{Z}$, $\phi_1(n) = 1$, $\phi_2(n) = n + 1$ and that $\mathbb{N} \subseteq S_k$ for all k .

Let $k \geq 3$ and suppose that $n < 0$ lies in S_k . Since $\theta^n(a_k)t^{\phi_k(n)}$ lies in H , so does $(a_k t)^{-1}\theta^n(a_k)t^{\phi_k(n)} = a_{k-1}^{-1}\theta^{-1}(a_{k-1}^{-1}) \dots \theta^{n+1}(a_{k-1}^{-1})t^{\phi_k(n)-1}$, and hence, by Lemma 6.2, a_{k-1}^{-1} lies in Ht^r for some r . It follows that $\theta^{-r}(a_{k-1})t^r \in H$ and so $r = \phi_{k-1}(-r)$. If $k = 3$, this is a contradiction, since it implies $r = -r + 1$. If $k > 3$, then $-r \in S_{k-1}$, and so, by the induction hypothesis, $r \leq 0$. But then $\phi_{k-1}(-r) \geq 1$, by (18), and hence $r \geq 1$, a contradiction. \square

Let d_H denote the word metric on H with respect to the generating set $a_1 t, a_2 t, \dots$

Lemma 7.4. If $n \in S_k$ and $h = \theta^n(a_k)t^{\phi_k(n)}$, then $d_H(1, h) = \phi_k(|n|)$.

Proof. If $k = 1$, then the result is obvious. If $k = 2$, then $h = a_2 a_1^n t^{n+1} = (a_2 t)(a_1 t)^n$ so $d_H(1, h) = 1 + |n| = \phi_k(|n|)$. If $k \geq 3$, then $n \geq 0$. Thus the word $\theta^n(a_k)$ is positive and hence $d_H(1, h) = \phi_k(n) = \phi_k(|n|)$. \square

Lemma 7.5. (i) Let $h = t^r \theta^i(a_k)t^{-s}$. Then $h \in H$ if and only if $i - r \in S_k$ and $s = r - \phi_k(i - r)$.

(ii) Let $h = t^r \theta^i(a_k^{-1})t^{-s}$. Then $h \in H$ if and only if $i - s \in S_k$ and $r = s - \phi_k(i - s)$.

Proof. For (i), note that $h = \theta^{i-r}(a_k)t^{r-s}$ and apply Definition 7.2. For (ii), note that $h^{-1} = t^s \theta^i(a_k)t^{-r}$ and apply (i). \square

Lemma 7.6. If $k \geq 3$ and $t^r \theta^i(a_k^{-1}) \in \Lambda$, then $r < i$.

Proof. If $t^r \theta^i(a_k^{-1}) \in Ht^s$, then Lemmas 7.3 and 7.5 give that $i - s \geq 0$ and $s - r = \phi_k(i - s) \geq 1$. Thus $i - r \geq 1$. \square

The exceptional nature of S_1 and S_2 highlighted by Lemma 7.3 means that small values of k will have to be treated separately in our proof. This motivates the inclusion of the following result, a special case of Lemma 7.5. Note in particular that (ii) implies that $t^r \theta^i(a_2^{-1}) \in \Lambda$ if and only if $r + i$ is odd.

Lemma 7.7. (i) Let $h = t^r \theta^i(a_2)t^{-s}$. Then $h \in H$ if and only if $s = 2r - i - 1$.

(ii) Let $h = t^r \theta^i(a_2^{-1})t^{-s}$. Then $h \in H$ if and only if $s = \frac{1}{2}(r + i + 1)$.

Proof. This follows immediately from Lemma 7.5 and the fact, given in Lemma 7.3, that $\phi_2(n) = n + 1$. □

The following result concerns passing a power of t through a sequence of terms of the form $\theta^i(a_2^{\pm 1})$. The statement is made neater by the use of the following formula, which is a consequence of Lemma 7.1:

$$\theta^a(a_3^{-1})\theta^b(a_3) = \begin{cases} \theta^a(a_2) \dots \theta^{b-1}(a_2), & a < b, \\ 1, & a = b, \\ \theta^{a-1}(a_2^{-1}) \dots \theta^b(a_2^{-1}), & a > b. \end{cases}$$

Lemma 7.8. *Let $\sigma = t^r \theta^a(a_3^{-1})\theta^b(a_3)$ and $s = 2^{b-a}(r - a - 2) + b + 2$ for some integers r, a, b . Then $\sigma \in \Lambda$ if and only if s is an integer. Furthermore, in this case, $\sigma \in Ht^s$.*

Proof. We split the proof into two claims. The first claim is that if $\sigma \in Ht^{s'}$ for some integer s' , then $s = s'$. In particular, this implies that if $\sigma \in \Lambda$, then s is an integer. If $a = b$, then clearly $s' = r = s$. If $a < b$, then $\theta^a(a_3^{-1})\theta^b(a_3) = \theta^a(a_2) \dots \theta^{b-1}(a_2)$. By the Lemma 6.2, there exist integers $r = r_0, r_1, \dots, r_{b-a} = s'$ such that $t^{r_i} \theta^{a+i}(a_2) \in Ht^{r_{i+1}}$. By Lemma 7.7, $r_{i+1} = 2r_i - a - i - 1$, which solves to give $r_i = 2^i(r - a - 2) + i + a + 2$. Substituting $i = b - a$ gives $s' = s$. On the other hand, suppose that $a > b$. Note that $t^r \theta^a(a_3^{-1})\theta^b(a_3) \in Ht^{s'}$ implies that $t^{s'} \theta^b(a_3^{-1})\theta^a(a_3) \in Ht^r$. Since $b < a$, we can substitute into the above solution to obtain $r = 2^{a-b}(s' - b - 2) + a + 2$, which rearranges to give $s' = s$. This completes the proof of our first claim.

The second claim is that if s is an integer, then $\sigma \in Ht^s$. If $a = b$, then this clearly holds. Suppose that $a < b$. Then $\sigma = t^r \theta^a(a_2) \dots \theta^{b-1}(a_2)$, so certainly $\sigma \in \Lambda$ since all the letters $a_2^{\pm 1}$ that appear are positive. Therefore $\sigma \in Ht^s$ by the first claim. Now suppose that $a > b$. Since s is an integer, we can define $\tau = t^s \theta^b(a_3^{-1})\theta^a(a_3) = t^s \theta^b(a_2) \dots \theta^{a-1}(a_2)$. Then certainly $\tau \in \Lambda$ – say $\tau \in Ht^{r'}$. By the first claim, $r' = 2^{a-b}(s - b - 2) + a + 2 = r$. Therefore $t^s \theta^b(a_3^{-1})\theta^a(a_3) \in t^r$, whence $t^r \theta^a(a_3^{-1})\theta^b(a_3) \in Ht^s$, and the second claim is proved. □

8. An upper bound on the distortion of H_k in G_k

Next we turn to estimates associated with pushing a power of t from left to right through a word $w = w(a_1, \dots, a_k)$ or through a piece of w , so as to leave a word on a_1t, \dots, a_kt times a power of t . We will need to keep track of both the length of that word on the a_1t, \dots, a_kt and the power of t that emerges to its right. Accordingly, let us define four families of functions, $\psi_{k,l}(n)$, $\Psi_{k,l,p}(n)$, $\kappa_{k,l}(n)$, $K_{k,l,p}(n)$ for integers $k \geq 1$ and $l, p, n \geq 0$.

- $\psi_{k,l}(n)$ is the least integer N such that if $h \in H$ is represented by a word $t^r \pi t^{-s}$ with π a piece of rank k , with $\ell(\pi) \leq l$, and with $|r| \leq n$, then $d_H(1, h) \leq N$.
- $\Psi_{k,l,p}(n)$ is the least integer N such that if $h \in H$ is represented by a word $t^r w t^{-s}$ with $w = w(a_1, \dots, a_k)$ a word of at most p pieces, with $\ell(w) \leq l$, and with $|r| \leq n$, then $d_H(1, h) \leq N$.
- $\kappa_{k,l}(n)$ is the least integer N such that if π is a piece of rank k with $\ell(\pi) \leq l$ and r is an integer with $|r| \leq n$ and $t^r \pi \in \Lambda$, then $t^r \pi \in H t^s$ for some s with $|s| \leq N$.
- $K_{k,l,p}(n)$ is the least integer N such that if w is a word of rank at most k with at most p pieces and with $\ell(w) \leq l$ and r is an integer with $|r| \leq n$ and $t^r w \in \Lambda$, then $t^r w \in H t^s$ for some s with $|s| \leq N$.

We will frequently make use, without further comment, of the fact that each of these functions is increasing in k, l, p and n .

The main technical result of this section is the following proposition. In the corollary that follows it we explain how the upper bound it gives on $\Psi_{k,l,p}(n)$ leads to our desired bound $\text{Dist}_{H_k}^{G_k} \leq A_k$.

Proposition 8.1. *For all $k \geq 1$, there exist integers $C_k \geq 1$ such that for all $l, p, n \geq 0$,*

$$\begin{aligned} \kappa_{k,l}(n) &\leq A_{k-1}(C_k n + C_k l), \\ \Psi_{k,l,p}(n) &\leq A_{k-1}^{(p)}(C_k n + C_k l), \\ \psi_{k,l}(n) &\leq A_{k-1}(C_k n + C_k l), \\ \Psi_{k,l,p}(n) &\leq A_{k-1}^{(3p)}(C_k n + C_k l). \end{aligned}$$

Corollary 8.2. *For all $k \geq 1$, the distortion function of H_k in G_k satisfies*

$$\text{Dist}_{H_k}^{G_k} \leq A_k.$$

Proof of Corollary 8.2. Since $G_1 \cong \mathbb{Z}^2$ and $H_1 \cong \mathbb{Z}$, H_1 is undistorted in G_1 and $\text{Dist}_{H_1}^{G_1} \leq A_1$. Now suppose that $k \geq 2$ and that $u = u(a_1, \dots, a_k, t)$ is a word of length at most n representing an element of H . By carrying each $t^{\pm 1}$ to the front, we see that u is equal in G_k to $t^r w$ for some integer r and some freely reduced word $w = w(a_1, \dots, a_k)$. These satisfy $|r| \leq n$ and $\ell(w) \leq C n^k$ for some integer $C > 0$ depending only on k – see, for example, Section 3.3 of [12].

We first show that the number of pieces of w is at most $n + 1$. Indeed, the process of carrying each $t^{\pm 1}$ to the front of u has the effect of applying $\theta^{\pm 1}$ to each a_i it passes.

The form of the automorphism θ ensures that no new $a_k^{\pm 1}$ are created by this process. The number of occurrences of $a_k^{\pm 1}$ in w , which we denote by $\ell_k(w)$, is therefore at most n . Let $w = w_1 \dots w_p$ be the partition of w into pieces. Say $w_i = a_k^{\epsilon_i^-} \pi_i a_k^{-\epsilon_i^+}$ where $\epsilon_i^-, \epsilon_i^+ \in \{0, 1\}$ and $\pi_i = \pi_i(a_1, \dots, a_{k-1})$. Observe that, for each i , precisely one of ϵ_i^+ and ϵ_{i+1}^- is equal to 1. Indeed, if $\epsilon_i^+ = \epsilon_{i+1}^- = 0$, then the pieces w_i and w_{i+1} could be concatenated to form a single piece, contradicting the minimality of p , and if $\epsilon_i^+ = \epsilon_{i+1}^- = 1$, then w would not be freely reduced. So

$$\ell_k(w) = \sum_{i=1}^p (\epsilon_i^- + \epsilon_i^+) = \epsilon_1^- + \sum_{i=1}^{p-1} (\epsilon_i^+ + \epsilon_{i+1}^-) + \epsilon_n^+ = \epsilon_1^- + p - 1 + \epsilon_n^+,$$

whence $p \leq \ell_k(w) + 1 \leq n + 1$.

Now,

$$d_H(1, u) = d_H(1, t^r w) \leq \Psi_{k, \ell(w), p}(|r|) \leq \Psi_{k, Cn^k, n+1}(n),$$

which is at most

$$A_{k-1}^{(3n+3)}(C_k C n^k + C_k n)$$

by Proposition 8.1. Choose an integer N large enough that $n^k \leq 2^n$ for $n \geq N$. Then, for $n \geq \max\{N, 1\}$,

$$\begin{aligned} d_H(1, u) &\leq A_{k-1}^{(3n+3)}(C_k C A_2(n) + C_k n) && \text{by (8)} \\ &\leq A_{k-1}^{(3n+3)}(C_k C A_k(n) + C_k n) && \text{by (7), (8)} \\ &\leq A_{k-1}^{(3n+3)}(A_k(C_k C n) + C_k n) && \text{by (8), (10)} \\ &\leq A_{k-1}^{(3n+3)}(A_k((C_k C + C_k)n)) && \text{by (8), (13)} \\ &= A_k((C_k C + C_k + 3)n + 3) && \text{by (4)}. \quad \square \end{aligned}$$

Proposition 8.1 will follow from the relationships between $\psi_{k,l}(n)$, $\Psi_{k,l,p}(n)$, $\kappa_{k,l}(n)$ and $K_{k,l,p}(n)$ set out in the next proposition. Of its claims, (26) and (29) are the most challenging to establish; we postpone their proof to Proposition 8.4, which itself will draw on Lemmas 8.5, 8.6 and 8.7.

Proposition 8.3. *For integers $k \geq 1$ and $l, p, n \geq 0$,*

$$\kappa_{1,l}(n) \leq n + 1, \tag{24}$$

$$K_{k,l,p}(n) \leq \max_{\substack{q \leq p \\ l_1 + \dots + l_q \leq l}} \{\kappa_{k,l_1}(\dots \kappa_{k,l_{q-1}}(\kappa_{k,l_q}(n)) \dots)\}, \tag{25}$$

$$\kappa_{k+1,l}(n) \leq 2K_{k,l,l}(2\phi_{k+1}(n)), \tag{26}$$

$$\psi_{1,l}(n) \leq 1, \tag{27}$$

$$\Psi_{k,l,p}(n) \leq p\psi_{k,l}(K_{k,l,p}(n)), \tag{28}$$

$$\psi_{k+1,l}(n) \leq 3K_{k,l,l}(2\phi_{k+1}(n)) + \Psi_{k,l,l}(2\phi_{k+1}(n)). \tag{29}$$

Proof. We first establish (24) and (27). Consideration of the empty word gives that $\kappa_{k,0}(n) = n$ and $\psi_{k,0} = 0$. Now suppose that $l \geq 1$ and note that the only pieces of rank 1 are $a_1^{\pm 1}$. If $h = t^r a_1^{\pm 1} t^{-s}$ lies in H , then $d_H(1, h) = 1$ and $r - s = \pm 1$, whence $|s| \leq |r| + 1$. Thus $\kappa_{1,l}(n) \leq n + 1$ and $\psi_{1,l}(n) = 1$.

For (25) and (28), let $h = t^r w t^{-s}$ where $w = w(a_1, \dots, a_k)$ is a word of length at most l with at most p pieces and $|r| \leq n$. Let $w = w_1 \dots w_q$ be the partition of w into pieces, where $q \leq p$. If $h \in H$, then Lemma 6.2 implies that there exist integers $r = r_0, r_1, \dots, r_q = s$ and elements h_1, \dots, h_q in H such that $t^{r_{i-1}} w_i = h_i t^{r_i}$. Thus $|r_i| \leq \kappa_{k,\ell(w_i)}(|r_{i-1}|)$, whence

$$|s| \leq \kappa_{k,\ell(w_q)}(\dots(\kappa_{k,\ell(w_1)}(|r|))\dots) \leq \kappa_{k,\ell(w_q)}(\dots(\kappa_{k,\ell(w_1)}(n))\dots)$$

and we obtain inequality (25). For inequality (28), note that

$$|r_i| \leq K_{k,\ell(w_1 \dots w_i),i}(|r|) \leq K_{k,l,p}(n),$$

whence

$$d_H(1, h) \leq \sum_{i=1}^q d_H(1, h_i) \leq \sum_{i=1}^q \psi_{k,\ell(w_i)}(|r_{i-1}|) \leq p \psi_{k,l}(K_{k,l,p}(n)).$$

Finally, (26) and (29) will follow from Proposition 8.4. □

We now derive Proposition 8.1 from Proposition 8.3. We first use (24), (25) and (26) to obtain bounds on $\kappa_{k,l}(n)$ and $K_{k,l,p}(n)$ in terms of Ackermann's functions. We then derive bounds on $\psi_{k,l}(n)$ and $\Psi_{k,l,p}(n)$ from (27), (28) and (29), having fed in our bounds on $\kappa_{k,l}(n)$ and $K_{k,l,p}(n)$.

Proof of Proposition 8.1. We will need the inequality, established in Lemma 3.2, that for $n \geq 0$ and $k \geq 2$,

$$\phi_k(n) \leq A_{k-1}(n + k). \tag{30}$$

We first prove that there exist integers $D_k \geq 1$ such that

$$\kappa_{k,l}(n) \leq A_{k-1}(D_k n + D_k l), \tag{31}$$

$$K_{k,l,p}(n) \leq A_{k-1}^{(p)}(D_k n + D_k l). \tag{32}$$

Inequalities (24) and (25) together imply that $K_{1,l,p}(n) \leq n + p$. Thus (31) and (32) hold in the case $k = 1$ with $D_1 = 1$. Now suppose that $k \geq 2$ and that (31) and (32) hold for smaller values of k . If $l = 0$, then, using (9), we calculate that $\kappa_{k,l}(n) = n \leq A_{k-1}(n)$. If $l \geq 1$, then

$$\begin{aligned} \kappa_{k,l}(n) &\leq 2K_{k-1,l,l}(2\phi_k(n)) && \text{by (26)} \\ &\leq 2K_{k-1,l,l}(2A_{k-1}(n + k)) && \text{by (30)} \\ &\leq 2A_{k-2}^{(l)}(2D_{k-1}A_{k-1}(n + k) + D_{k-1}l) \end{aligned}$$

$$\begin{aligned}
 &\leq 2A_{k-2}^{(l)}(A_{k-1}(2D_{k-1}n + D_{k-1}l + 2D_{k-1}k)) && \text{by (8), (10), (13)} \\
 &= 2A_{k-1}(2D_{k-1}n + (D_{k-1} + 1)l + 2D_{k-1}k) && \text{by (4)} \\
 &\leq A_{k-1}(4D_{k-1}n + 2(D_{k-1} + 1)l + 4D_{k-1}k) && \text{by (10)} \\
 &\leq A_{k-1}(4D_{k-1}n + [2(D_{k-1} + 1) + 4D_{k-1}k]l) && \text{by (8)}.
 \end{aligned}$$

Taking $D_k = \max\{2(D_{k-1} + 1) + 4D_{k-1}k, 1\}$, we obtain (31).

For (32) we calculate that

$$\begin{aligned}
 K_{k,l,p}(n) &\leq \max_{\substack{q \leq p \\ l_1 + \dots + l_q \leq l}} \{ \kappa_{k,l_1}(\dots \kappa_{k,l_{q-1}}(\kappa_{k,l_q}(n)) \dots) \} && \text{by (25)} \\
 &\leq \max_{l_1 + \dots + l_q \leq l} \{ A_{k-1}(\dots A_{k-1}(A_{k-1}(D_k n + D_k l_q) + D_k l_{q-1}) \dots) \} \\
 &&& \text{by (8)} \\
 &\leq \max_{\substack{q \leq p \\ l_1 + \dots + l_q \leq l}} \left\{ A_{k-1}^{(q)} \left(D_k n + D_k \sum_{i=1}^q l_i \right) \right\} && \text{by (8), (13)} \\
 &\leq \max_{q \leq p} \{ A_{k-1}^{(q)}(D_k n + D_k l) \} && \text{by (8)} \\
 &\leq A_{k-1}^{(p)}(D_k n + D_k l) && \text{by (9)}.
 \end{aligned}$$

Next, we combine (27), (28) and (29) with (31) and (32) to deduce that there exist integers $E_k, F_k \geq 1$ such that

$$\psi_{k,l}(n) \leq A_{k-1}(E_k n + E_k l), \tag{33}$$

$$\Psi_{k,l,p}(n) \leq A_{k-1}^{(3p)}(F_k n + F_k l). \tag{34}$$

It follows from (27) and (28) that $\Psi_{1,l,p}(n) \leq p$. Thus (33) and (34) hold in the case $k = 1$ with $E_k = F_k = 1$. Now suppose that $k \geq 2$ and that (33) and (34) hold for smaller values of k . If $l = 0$, then $\psi_{k,l}(n) = 0 \leq A_{k-1}(0)$. If $l \geq 1$, then

$$\begin{aligned}
 \psi_{k,l}(n) &\leq 3K_{k-1,l,l}(2\phi_k(n)) + \Psi_{k-1,l,l}(2\phi_k(n)) && \text{by (29)} \\
 &\leq 3K_{k-1,l,l}(2A_{k-1}(n+k)) + \Psi_{k-1,l,l}(2A_{k-1}(n+k)) && \text{by (30)} \\
 &\leq 3A_{k-2}^{(l)}(2D_{k-1}A_{k-1}(n+k) + D_{k-1}l) \\
 &\quad + A_{k-2}^{(3l)}(2F_{k-1}A_{k-1}(n+k) + F_{k-1}l) && \text{by (32)} \\
 &\leq 3A_{k-2}^{(l)}(A_{k-1}(2D_{k-1}(n+k) + D_{k-1}l)) \\
 &\quad + A_{k-2}^{(3l)}(A_{k-1}(2F_{k-1}(n+k) + F_{k-1}l)) && \text{by (8), (10), (13)}
 \end{aligned}$$

$$\begin{aligned}
 &= 3A_{k-1}(2D_{k-1}(n+k) + (D_{k-1} + 1)l) \\
 &\quad + A_{k-1}(2F_{k-1}(n+k) + (F_{k-1} + 3)l) && \text{by (4)} \\
 &\leq A_{k-1}(6D_{k-1}(n+k) + 3(D_{k-1} + 1)l) \\
 &\quad + A_{k-1}(2F_{k-1}(n+k) + (F_{k-1} + 3)l) && \text{by (10)} \\
 &\leq A_{k-1}(2(3D_{k-1} + F_{k-1})(n+k) + (3D_{k-1} + F_{k-1} + 4)l) && \text{by (12)} \\
 &\leq A_{k-1}(2(3D_{k-1} + F_{k-1})n + (3(2k + 1)D_{k-1} + (2k + 1)F_{k-1} + 4)l).
 \end{aligned}$$

Taking $E_k = 3(2k + 1)D_{k-1} + (2k + 1)F_{k-1} + 4$, we obtain (33).

If $p = 0$ or $l = 0$, then, using (9), we calculate that $\Psi_{k,l,p}(n) = 0 \leq A_{k-1}^{(3p)}(0)$.
 If $l, p \geq 1$, then

$$\begin{aligned}
 \Psi_{k,l,p}(n) &\leq p\psi_{k,l}(K_{k,l,p}(n)) && \text{by (28)} \\
 &\leq p\psi_{k,l}(A_{k-1}^{(p)}(D_k n + D_k l)) && \text{by (32)} \\
 &\leq pA_{k-1}(E_k A_{k-1}^{(p)}(D_k n + D_k l) + E_k l) \\
 &\leq pA_{k-1}^{(p+1)}(D_k E_k n + (D_k + 1)E_k l) && \text{by (8), (9), (10), (13)} \\
 &\leq A_{k-1}^{(2p+1)}(D_k E_k n + (D_k + 1)E_k l) && \text{by (11),} \\
 &\leq A_{k-1}^{(3p)}(D_k E_k n + (D_k + 1)E_k l) && \text{by (9).}
 \end{aligned}$$

Taking $F_k = (D_k + 1)E_k$, we obtain (34).

Finally, the proof is completed by taking $C_k = \max\{D_k, E_k, F_k\}$ and applying (8). □

The remainder of this section is devoted to establishing (26) and (29). This is done in Proposition 8.4, which draws on Lemmas 8.5, 8.6 and 8.7 that follow. We now outline our strategy.

Suppose that $t^r a_k^{\epsilon_1} w a_k^{-\epsilon_2} t^{-s}$, where $r, s \in \mathbb{Z}$, $\epsilon_1, \epsilon_2 \in \{0, 1\}$ and $w = w(a_1, \dots, a_{k-1})$, represents an element $h \in H$. Our approach will be to find elements $h_1, h_2 \in H$, integers r', s' and a word $w' = w'(a_1, \dots, a_{k-1})$ such that h is represented by $h_1 t^{r'} w' t^{-s'} h_2$. The functions $K_{k-1,*,*}$ and $\Psi_{k-1,*,*}$ will then control the behaviour of the subword $t^{r'} w' t^{-s'}$. Together with estimates for $d_H(1, h_i)$, $|r'|$, $|s'|$ and $\ell(w')$, this will allow us to derive bounds on $|s|$ and $d_H(1, h)$.

As indicated by Lemma 7.3, the case $k = 2$ is exceptional and so will be treated separately. For $k \geq 3$, the h_1, h_2, r', s' and w will be produced by Lemma 8.5. This lemma takes integers k, n and ϵ , with $k \geq 3$ and $\epsilon \in \{0, 1\}$, and gives an integer n' , an element $h \in H$ and a word $u = u(a_1, \dots, a_{k-1})$ such that $t^n a_k^\epsilon = h t^{n'} u$ in G . Applying Lemma 8.5 to k, r and ϵ_1 will produce r', h_1 and a word u_1 . Applying Lemma 8.5 to k, s and ϵ_2 will produce s', h_2^{-1} and a word u_2 . The word w' will then be defined to be the free reduction of $\tilde{w} := u_1 w u_2^{-1}$.

The relationship between the input and output of Lemma 8.5 is determined by which of the following holds:

- (i) $\epsilon = 0$,
- (ii) $\epsilon = 1$ and $n \leq 0$, or
- (iii) $\epsilon = 1$ and $n > 0$.

A priori, this would lead to us having to consider nine distinct cases, depending on the values of ϵ_1 and ϵ_2 and the signs of r and s . To streamline the process, Lemma 8.5 packages (i) and (ii) together: it considers the cases that either $n\epsilon \leq 0$ or $n\epsilon > 0$. As such, we need now only consider four cases, depending on the signs of $r\epsilon_1$ and $s\epsilon_2$.

The form of \tilde{w} will depend on which of (i), (ii) or (iii) applies to r and ϵ_1 and to s and ϵ_2 . Lemmas 8.6 and 8.7 will be brought to bear to ensure that enough cancellation occurs to obtain a sufficiently strong bound on $\ell(w')$.

Proposition 8.4. *Let $h = t^r a_k^{\epsilon_1} w a_k^{-\epsilon_2} t^{-s}$ where $k \geq 2$, $\epsilon_1, \epsilon_2 \in \{0, 1\}$, and $w = w(a_1, \dots, a_{k-1})$. Let n and l be integers with $|r| \leq n$ and $\ell(w) \leq l$. If $h \in H$, then*

$$|s| \leq 2K_{k-1,l,l}(2\phi_k(n)),$$

$$d_H(1, h) \leq 3K_{k-1,l,l}(2\phi_k(n)) + \Psi_{k-1,l,l}(2\phi_k(n)).$$

Proof. We claim that there exist $h_1, h_2 \in H$, $r', s' \in \mathbb{Z}$ and $w' = w'(a_1, \dots, a_{k-1})$ such that $h = h_1 t^{r'} w' t^{-s'} h_2$ in G and

$$|r'| \leq 2\phi_k(n), \tag{35}$$

$$|s| \leq |s'| + 1, \tag{36}$$

$$d_H(1, h_1) \leq |r'| + 1, \tag{37}$$

$$d_H(1, h_2) \leq |s'| + 1, \tag{38}$$

$$\ell(w') \leq l. \tag{39}$$

The result follows from the claim by direct calculation. Indeed, since the number of pieces of a word is bounded by its length,

$$|s'| \leq K_{k-1,\ell(w'),\ell(w')}(|r'|), \tag{40}$$

$$d_H(1, t^{r'} w' t^{-s'}) \leq \Psi_{k-1,\ell(w'),\ell(w')}(|r'|). \tag{41}$$

We will also need the inequality

$$K_{k,l,p}(n) \geq n, \tag{42}$$

which follows immediately from consideration of the empty word. We can now calculate that

$$\begin{aligned} |s| &\leq |s'| + 1 && \text{by (36)} \\ &\leq K_{k-1,\ell(w'),\ell(w')}(|r'|) + 1 && \text{by (40)} \\ &\leq K_{k-1,l,l}(2\phi_k(n)) + 1 && \text{by (35), (39)} \\ &\leq 2K_{k-1,l,l}(2\phi_k(n)) && \text{by (18), (42)} \end{aligned}$$

and

$$\begin{aligned}
 d_H(1, h) &\leq d_H(1, h_1) + d_H(1, t^{r'} w' t^{-s'}) + d_H(1, h_2) \\
 &\leq |r'| + 1 + \Psi_{k-1, \ell(w'), \ell(w')}(|r'|) + |s'| + 1 && \text{by (37), (38), (41)} \\
 &\leq 2\phi_k(n) + 1 + \Psi_{k-1, l, l}(2\phi_k(n)) + K_{k-1, \ell(w'), \ell(w')}(|r'|) + 1 \\
 &&& \text{by (35), (39), (40)} \\
 &\leq 4\phi_k(n) + \Psi_{k-1, l, l}(2\phi_k(n)) + K_{k-1, l, l}(2\phi_k(n)) && \text{by (18), (35), (39)} \\
 &\leq 3K_{k-1, l, l}(2\phi_k(n)) + \Psi_{k-1, l, l}(2\phi_k(n)) && \text{by (42)}.
 \end{aligned}$$

We first prove the claim for $k = 2$. Since $t^q a_2 = (a_2 t)(a_1 t)^{-q} t^{2q-1}$, we can take w' to be w and define h_1, h_2, r' and s' by

$$\begin{aligned}
 h_1 &= \begin{cases} 1, & \epsilon_1 = 0, \\ (a_2 t)(a_1 t)^{-r}, & \epsilon_1 = 1, \end{cases} & r' &= \begin{cases} r, & \epsilon_1 = 0, \\ 2r - 1, & \epsilon_1 = 1, \end{cases} \\
 h_2 &= \begin{cases} 1, & \epsilon_2 = 0, \\ (a_1 t)^{-s}(a_2 t)^{-1}, & \epsilon_2 = 1, \end{cases} & s' &= \begin{cases} s, & \epsilon_2 = 0, \\ 2s - 1, & \epsilon_2 = 1. \end{cases}
 \end{aligned}$$

Inequalities (36) and (39) are immediate. For (35), use the fact, from Lemma 7.3, that $\phi_2(n) = n + 1$. Inequality (37) is immediate if $\epsilon_1 = 0$. If $\epsilon_1 = 1$, then $r = \frac{1}{2}(r' + 1)$, whence $|r| \leq \frac{1}{2}(|r'| + 1)$. But $r' \neq 0$, so $|r| \leq |r'|$ and $d_H(1, h_1) = |r| + 1 \leq |r'| + 1$. Inequality (38) is derived similarly.

We now prove the claim for $k \geq 3$. First apply Lemma 8.5 to k, r, ϵ_1 to produce r', h_1 and a word u_1 . Then apply it to k, s, ϵ_2 to produce s', h_2^{-1} and a word u_2 . Defining $\tilde{w} := u_1 w u_2^{-1}$, we have that h is represented by $h_1 t^{r'} \tilde{w} t^{-s'} h_2$ and hence that $t^{r'} \tilde{w} t^{-s'} \in H$. It is immediate from the bounds given in Lemma 8.5 that (35)–(38) hold. Finally, we define w' to be the free reduction of \tilde{w} . To establish (39), we consider four cases.

Case $r\epsilon_1 \leq 0, s\epsilon_2 \leq 0$. We have that $\tilde{w} = w$ and so it is immediate that $\ell(w') \leq \ell(w)$.

Case $r\epsilon_1 > 0, s\epsilon_2 \leq 0$. We have that $\tilde{w} = \theta^{r-1}(a_{k-1}^{-1}) \dots \theta^0(a_{k-1}^{-1})w$. Since $t^{r'} \theta^{r-1}(a_{k-1}^{-1})$ does not lie in Λ , applying Lemma 6.2 to $t^{r'} \tilde{w} t^{-s'}$ shows that, when \tilde{w} is freely reduced, each a_{k-1}^{-1} in $\theta^{r-1}(a_{k-1}^{-1}) \dots \theta^0(a_{k-1}^{-1})$ cancels into w . It follows from Lemma 8.7 that $\ell(w') \leq \ell(w)$.

Case $r\epsilon_1 \leq 0, s\epsilon_2 > 0$. We have that $\tilde{w} = w\theta^0(a_{k-1}) \dots \theta^{s-1}(a_{k-1})$. Since $t^{s'} \theta^{s-1}(a_{k-1}^{-1})$ does not lie in Λ , applying Lemma 6.2 to $t^{s'} \tilde{w}^{-1} t^{-r'}$ shows that, when \tilde{w} is freely reduced, each a_{k-1} in $\theta^0(a_{k-1}) \dots \theta^{s-1}(a_{k-1})$ cancels into w . It follows from Lemma 8.7 that $\ell(w') \leq \ell(w)$.

Case $r\epsilon_1 > 0, s\epsilon_2 > 0$. We have that

$$\tilde{w} = \theta^{r-1}(a_{k-1}^{-1}) \dots \theta^0(a_{k-1}^{-1})w\theta^0(a_{k-1}) \dots \theta^{s-1}(a_{k-1}).$$

Neither $t^{r'}\theta^{r-1}(a_{k-1}^{-1})$ nor $t^{s'}\theta^{s-1}(a_{k-1}^{-1})$ lies in Λ , so we are in a position to apply Lemma 8.6. If case (i) of Lemma 8.6 occurs, then, when \tilde{w} is freely reduced, each a_{k-1}^{-1} in $\theta^{r-1}(a_{k-1}^{-1}) \dots \theta^0(a_{k-1}^{-1})$ and each a_{k-1} in $\theta^0(a_{k-1}) \dots \theta^{s-1}(a_{k-1})$ cancels into w . Applying Lemma 8.7 gives that $\ell(w') \leq \ell(w)$. On the other hand, suppose that case (ii) of Lemma 8.6 occurs, so w' is the free reduction of $\theta^{r-1}(a_{k-1}^{-1})\theta^{s-1}(a_{k-1})$. We will show that $r = s$, whence w' is the empty word and trivially $\ell(w') \leq l$. If $k = 3$, then $t^{r'}w't^{-s'} = t^{r-1}\theta^{r-1}(a_2^{-1})\theta^{s-1}(a_2)t^{1-s} = t^{r-s}a_1^{s-r}$ in G . Since this element lies in H , $r - s = s - r$, whence $r = s$. If $k = 4$, then $t^{r'}w't^{-s'}$ is freely equal to $t^{r-1}\theta^{r-1}(a_3^{-1})\theta^{s-1}(a_3)t^{1-s}$. Since this lies in H , applying Lemma 7.8 and solving the resulting equation gives $r = s$. Finally, suppose that $k > 4$. Lemma 7.1 gives that

$$t^{r'}w't^{-s'} \stackrel{\text{fr}}{=} \begin{cases} t^{r-1}\theta^{r-1}(a_{k-2}) \dots \theta^{s-2}(a_{k-2})t^{1-s}, & r < s, \\ t^{r-s}, & r = s, \\ t^{r-1}\theta^{r-2}(a_{k-2}^{-1}) \dots \theta^{s-1}(a_{k-2}^{-1})t^{1-s}, & r > s. \end{cases}$$

By Lemma 7.6, neither $t^{r-1}\theta^{r-2}(a_{k-2}^{-1})$ nor $t^{s-1}\theta^{s-2}(a_{k-2}^{-1})$ lies in Λ , since $k - 2 \geq 3$. Thus, by Lemma 6.2, both $r < s$ and $s > r$ lead to a contradiction. Hence $r = s$ as required. □

Lemma 8.5. *Given integers k, n, ϵ , with $k \geq 3$ and $\epsilon \in \{0, 1\}$, there exists an integer n' , an element $h \in H$ and a word $u = u(a_1, \dots, a_{k-1})$ such that $t^n a_k^\epsilon = h t^{n'} u$ in G ,*

$$|n| - 1 \leq |n'| \leq 2\phi_k(|n|) \quad \text{and} \quad d_H(1, h) \leq \max\{|n'|, 1\}.$$

Furthermore,

- (i) if $n\epsilon \leq 0$, then u is the empty word;
- (ii) if $n\epsilon > 0$, then $n' = n - 1$,

$$u = \theta^{n-1}(a_{k-1}^{-1}) \dots \theta^0(a_{k-1}^{-1}) \quad \text{and} \quad t^{n'}\theta^{n-1}(a_{k-1}^{-1}) \notin \Lambda.$$

Proof. We consider three cases.

Case $\epsilon = 0$. We trivially obtain an instance of conclusion (i) by taking $n' = n$, $h = 1$ and u to be the empty word. The upper bound on $|n'|$ follows from (18) and (21).

Case $\epsilon = 1$ and $n \leq 0$. Following the calculation

$$t^n a_k = \theta^{-n}(a_k)t^n = \theta^{-n}(a_k)t^{\phi_k(|n|)}t^{n-\phi_k(|n|)},$$

we obtain an instance of conclusion (i) by taking $n' = n - \phi_k(|n|)$, $h = \theta^{-n}(a_k)t^{\phi_k(|n|)}$ and u to be the empty word. It follows immediately from the definition of the function

ϕ_k that $h \in H$ and from Lemma 7.4 that $d_H(1, h) = \phi_k(|n|)$. By (18), $\phi_k(|n|)$ is positive whence $|n'| = |n| + \phi_k(|n|)$ and $d_h(1, h) \leq |n'|$. Applying (18) and (21) gives $|n| + 1 \leq |n'| \leq 2\phi_k(|n|)$.

Case $\epsilon = 1$ and $n > 0$. Following the calculation

$$t^n a_k = a_k a_k^{-1} t^n a_k = a_k t^n \theta^n(a_k^{-1}) a_k = (a_k t) t^{n-1} \theta^{n-1}(a_{k-1}^{-1}) \dots \theta^0(a_{k-1}^{-1})$$

we obtain an instance of conclusion (ii) by taking $n' = n - 1$, $h = (a_k t)$ and

$$u = \theta^{n-1}(a_{k-1}^{-1}) \dots \theta^0(a_{k-1}^{-1}).$$

The upper bound on $|n'|$ follows from (18) and (21). The fact that $t^{n'} \theta^{n-1}(a_{k-1}^{-1})$ does not lie in Λ follows from Lemmas 7.6 and 7.7. \square

Lemma 8.6. *Let $\sigma = t^r \theta^a(a_k^{-1}) \dots \theta^0(a_k^{-1}) w \theta^0(a_k) \dots \theta^b(a_k) t^{-s}$ where $w = w(a_1, \dots, a_k)$ is freely reduced and $a, b \geq 0$. Suppose σ represents an element of H but $t^r \theta^a(a_k^{-1}) \notin \Lambda$ and $t^s \theta^b(a_k^{-1}) \notin \Lambda$. Then either*

(i) *w has a prefix $\theta^0(a_k) \dots \theta^{a-1}(a_k) a_k$ and suffix $a_k^{-1} \theta^{b-1}(a_k^{-1}) \dots \theta^0(a_k^{-1})$,
or*

(ii) *$w = \theta^0(a_k) \dots \theta^{a-1}(a_k) \theta^{b-1}(a_k^{-1}) \dots \theta^0(a_k^{-1})$.*

Proof. Write l_1 for the letter a_k^{-1} of the term $\theta^a(a_k^{-1})$ of σ and write l_2 for the letter a_k of the term $\theta^b(a_k)$ of σ . Lemma 6.2 implies that, when σ is freely reduced, both l_1 and l_2 cancel. Let l' be the letter a_k that cancels with l_1

If l' lies in w , then l_2 must cancel with a letter to the right of l' in w , and we have case (i).

On the other hand, suppose that l' lies in the subword $\theta^0(a_k) \dots \theta^b(a_k)$. If l' is distinct from l_2 , then l_2 must cancel with an a_k^{-1} lying to the right of l' . But this is a contradiction, since all the occurrences of $a_k^{\pm 1}$ in $\theta^0(a_k) \dots \theta^b(a_k)$ are positive. Thus $l' = l_2$. Now $\theta^{a-1}(a_k^{-1}) \dots \theta^0(a_k^{-1}) w \theta^0(a_k) \dots \theta^{b-1}(a_k)$ must be freely trivial and we have case (ii). \square

Lemma 8.7. *Let $w = \theta^0(a_k) \dots \theta^r(a_k)$ where $r \geq 0$. Let l be the last a_k appearing in w and partition w as $w = uv$ where u is the prefix of w ending with l and v is the suffix of w coming after l . Then $\ell(u) \geq \ell(v)$.*

Proof. Note that $u = \theta^0(a_k) \dots \theta^{r-1}(a_k) a_k$, and $v = \theta^0(a_{k-1}) \dots \theta^{r-1}(a_{k-1})$ by Lemma 7.1. It thus suffices to prove that $\ell(\theta^i(a_k)) \geq \ell(\theta^i(a_{k-1}))$ for $i \geq 0$. But this follows by an easy induction on k from the structures of $\theta^i(a_k)$ and $\theta^i(a_{k-1})$ respectively given by Lemma 7.1. \square

9. Groups with Ackermannian Dehn functions

Recall that Γ_k is the HNN extension of G_k over H_k in which the stable letter commutes with all elements of H_k :

$$\Gamma_k := \langle a_1, \dots, a_k, t, p \mid t^{-1}a_1t = a_1, t^{-1}a_it = a_ia_{i-1} \ (i > 1), [p, a_it] = 1 \ (i > 0) \rangle.$$

Proposition 9.1. *The group Γ_1 has Dehn function \simeq -equivalent to $n \mapsto n^2$.*

Proof. Making the substitution $\alpha = a_1t$ shows that Γ_1 is a right-angled Artin group with presentation $\langle \alpha, t, p \mid [t, \alpha], [p, \alpha] \rangle$. It follows that Γ_1 is CAT(0) [16] whence it has Dehn function \simeq -equivalent to n^2 by [13], Proposition 1.6.III.Γ. □

Proposition 9.2. *For all $k \geq 2$, the group Γ_k has Dehn function \simeq -equivalent to A_k .*

Proof. Let $k \geq 2$. The Dehn function of a CAT(0) group is either linear or quadratic [11], Theorem 6.2.1, with the linear case occurring precisely when the group is hyperbolic [11], Theorem 6.1.5. By Theorem 1.3, the group G_k is CAT(0). However, since it contains an embedded copy of \mathbb{Z}^2 it is not hyperbolic [11], Theorem 6.1.10. The Dehn function of G_k is therefore quadratic. By Theorem 1.3, the distortion function of H_k in G_k is \simeq -equivalent to A_k . Plugging these two functions into Theorem 6.20.III.Γ of [13] gives lower and upper bounds for the Dehn function of Γ_k of $\max\{n^2, nA_k(n)\}$ and $nA_k(n)^2$ respectively, up to \simeq -equivalence. So, by (9), the Dehn function of Γ_k is between $A_k(n)$ and $A_k(n)^3$. But (14) implies that, for any $C \geq 1$, the function $n \mapsto A_k(n)^C$ is \simeq -equivalent to A_k . □

The ideas behind [13], Theorem 6.20.III.Γ, used here are most transparent via the tools of van Kampen diagrams and corridors. For example, towards the lower bound, consider the words

$$v_{k,n} := a_k^n a_2 t a_1 a_2^{-1} a_k^{-n}$$

of Section 5, which equal

$$w_{k,n} := u_{k,n} (a_2t) (a_1t) (a_2t)^{-1} u_{k,n}^{-1}$$

in G_k . Observe that $[v_{k,n}, p] = 1$ in Γ_k and that in any van Kampen diagram for $[v_{k,n}, p]$, there must be a p -corridor connecting the two boundary edges labelled by p . (Figure 3 is an example of such a diagram when $k = 2$ and $n = 4$.) The word on a_1t, \dots, a_kt written along each side of this corridor must equal $v_{k,n}$ in G_k and so freely equals $w_{k,n}$. It follows that any van Kampen diagram for $[v_{k,n}, p]$ has area at least the length of $w_{k,n}$, which is $2\mathcal{H}_k(n) + 3$. So, as the length of $[v_{k,n}, p]$ is $4n + 10$, this leads to a lower bound of $A_k(n) \simeq \mathcal{H}_k$ on the Dehn function of G_k .




Figure 3. A van Kampen diagram for $[v_{2,4}, p]$ – an example of a word which represents the identity in Γ_k but can only be filled by a large area diagram.

References

- [1] G. Arzhantseva and D. Osin, Solvable groups with polynomial Dehn functions. *Trans. Amer. Math. Soc.* **354** (2002), no. 8, 3329–3348. [Zbl 0998.20040](#) [MR 1897402](#)
- [2] J. Barnard, N. Brady, and P. Dani, Super-exponential distortion of subgroups of $\text{CAT}(-1)$ groups. *Algebr. Geom. Topol.* **7** (2007), 301–308. [Zbl 1187.20055](#) [MR 2308946](#)
- [3] G. Baumslag, A non-cyclic one-relator group all of whose finite quotients are cyclic. *J. Austral. Math. Soc.* **10** (1969), 497–498. [Zbl 0214.27402](#) [MR 0254127](#)
- [4] G. Baumslag, Recognizing powers in nilpotent groups and nilpotent images of free groups. *J. Austral. Math. Soc.* **83** (2007), no. 2, 149–155. [Zbl 1146.20030](#) [MR 2396860](#)
- [5] G. Baumslag, M. R. Bridson, C. F. Miller, and H. Short, Finitely presented subgroups of automatic groups and their isoperimetric functions. *J. London Math. Soc.* (2) **56** (1997), no. 2, 292–304. [Zbl 0910.20023](#) [MR 1489138](#)
- [6] G. Baumslag, M. R. Bridson, C. F. Miller, III, and H. Short, Fibre products, non-positive curvature, and decision problems. *Comment. Math. Helv.* **75** (2000), no. 3, 457–477. [Zbl 0973.20034](#) [MR 1793798](#)
- [7] L. D. Beklemishev, The worm principle. In *Logic Colloquium '02*, Lect. Notes Log. 27, Assoc. Symbol. Logic, La Jolla, CA, 2006, 75–95. [Zbl 1108.03055](#) [MR 2258703](#)
- [8] A. A. Bernasconi, On HNN-extensions and the complexity of the word problem for one-relator groups. PhD thesis, University of Utah, 1994. www.math.utah.edu/~sg/Papers/bernasconi-thesis.pdf.

- [9] M. Bestvina, M. Feighn, and M. Handel, The Tits alternative for $\text{Out}(F_n)$. II. A Kolchin type theorem. *Ann. of Math. (2)* **161** (2005), no. 1, 1–59. [Zbl 1139.20026](#) [MR 2150382](#)
- [10] N. Brady, W. Dison, and T. R. Riley, Hyperbolic hydra. *Groups Geom. Dyn.*, to appear; preprint [arXiv:1105.1535](#).
- [11] M. R. Bridson, The geometry of the word problem. In *Invitations to geometry and topology*, M. R. Bridson and S. M. Salamon, eds., Oxf. Grad. Texts Math. 7, Oxford University Press, Oxford 2002, 29–91. [Zbl 0996.54507](#) [MR 1967746](#)
- [12] M. R. Bridson, Polynomial Dehn functions and the length of asynchronously automatic structures. *Proc. London Math. Soc.* **85** (2002), no. 2, 441–465. [Zbl 1046.20027](#) [MR 1912057](#)
- [13] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*. Grundlehren Math. Wiss. 319, Springer-Verlag, Berlin 1999. [Zbl 0988.53001](#) [MR 1744486](#)
- [14] W. Buchholz, An independence result for $(\Pi_1^1\text{-CA})+\text{BI}$. *Ann. Pure Appl. Logic* **33** (1987), no. 2, 131–155. [Zbl 0636.03052](#) [MR 0874022](#)
- [15] C. H. Cashen and N. Macura, Line patterns in free groups. *Geom. Topol.* **15** (2011), no. 3, 1419–1475. [Zbl 05934844](#) [MR 2825316](#)
- [16] R. Charney and M. W. Davis, Finite $K(\pi, 1)$ s for Artin groups. In *Prospects in topology* (Princeton, NJ, 1994), Ann. of Math. Stud. 138, Princeton University Press, Princeton, NJ, 1995, 110–124. [Zbl 0930.55006](#) [MR 1368655](#)
- [17] D. E. Cohen, The mathematician who had little wisdom: a story and some mathematics. In *Combinatorial and geometric group theory* (Edinburgh, 1993), London Math. Soc. Lecture Note Ser. 204, Cambridge University Press, Cambridge 1995, 56–62. [Zbl 0962.20500](#) [MR 1320274](#)
- [18] D. E. Cohen, K. Madlener, and F. Otto, Separating the intrinsic complexity and the derivational complexity of the word problem for finitely presented groups. *Math. Logic Quart.* **39** (1993), no. 2, 143–157. [Zbl 0799.03049](#) [MR 1269904](#)
- [19] B. Farb, The extrinsic geometry of subgroups and the generalised word problem. *Proc. London Math. Soc.* (3) **68** (1994), no. 3, 577–593. [Zbl 0816.20032](#) [MR 1262309](#)
- [20] S. M. Gersten, Isodiametric and isoperimetric inequalities in group extensions. Preprint, University of Utah, 1991.
- [21] S. M. Gersten, Isoperimetric and isodiametric functions of finite presentations. In *Geometric group theory I*, G. Niblo and M. Roller, eds., London Math. Soc. Lecture Note Ser. 181, Cambridge University Press, Cambridge 1993, 79–96. [Zbl 0829.20054](#) [MR 1238517](#)
- [22] S. M. Gersten, Quadratic divergence of geodesics in $\text{CAT}(0)$ spaces. *Geom. Funct. Anal.* **4** (1994), no. 1, 37–51. [Zbl 0809.53054](#) [MR 1254309](#)
- [23] S. M. Gersten and H. Short, Small cancellation theory and automatic groups. *Invent. Math.* **102** (1990), no. 2, 305–334. [Zbl 0714.20016](#) [MR 1074477](#)
- [24] M. Gromov, Asymptotic invariants of infinite groups. In *Geometric group theory II*, G. Niblo and M. Roller, eds., London Math. Soc. Lecture Note Ser. 182, Cambridge University Press, Cambridge 1993, 1–295. [Zbl 0841.20039](#) [MR 1253544](#)
- [25] M. Hamano and M. Okada, A relationship among Gentzen’s proof-reduction, Kirby-Paris’ hydra game and Buchholz’s hydra game. *Math. Logic Quart.* **43** (1997), no. 1, 103–120. [Zbl 0872.03038](#) [MR 1429324](#)

- [26] P. Jolissaint, Rapidly decreasing functions in reduced C^* -algebras of groups. *Trans. Amer. Math. Soc.* **317** (1990), no. 1, 167–196. [Zbl 0711.46054](#) [MR 0943303](#)
- [27] L. Kirby and J. Paris, Accessible independence results for Peano arithmetic. *Bull. London Math. Soc.* **14** (1982), no. 4, 285–293. [Zbl 0501.03017](#) [MR 0663480](#)
- [28] N. Macura, CAT(0) spaces with polynomial divergence of geodesics. *Geom. Dedicata* **163** (2013), no. 1, 361–378. [MR 3032700](#)
- [29] N. Macura, Quadratic isoperimetric inequality for mapping tori of polynomially growing automorphisms of free groups. *Geom. Funct. Anal.* **10** (2000), no. 4, 874–901. [Zbl 0980.20031](#) [MR 1791144](#)
- [30] N. Macura, Detour functions and quasi-isometries. *Q. J. Math.* **53** (2002), no. 2, 207–239. [Zbl 1036.20033](#) [MR 1909513](#)
- [31] K. Madlener and F. Otto, Pseudonatural algorithms for the word problem for finitely presented monoids and groups. *J. Symbolic Comput.* **1** (1985), no. 4, 383–418. [Zbl 0591.20038](#) [MR 0849044](#)
- [32] M. Mitra, Cannon–Thurston maps for trees of hyperbolic metric spaces. *J. Differential Geom.* **48** (1998), no. 1, 135–164. [Zbl 0906.20023](#) [MR 1622603](#)
- [33] G. A. Niblo and L. D. Reeves, The geometry of cube complexes and the complexity of their fundamental groups. *Topology* **37** (1998), no. 3, 621–633. [Zbl 0911.57002](#) [MR 1604899](#)
- [34] A. Yu. Ol’shanskii and M. V. Sapir, Length and area functions on groups and quasi-isometric Higman embeddings. *Internat. J. Algebra Comput.* **11** (2001), no. 2, 137–170. [Zbl 1025.20030](#) [MR 1829048](#)
- [35] Ch. Pittet, Géométrie des groupes, inégalités isopérimétriques de dimension 2 et distortions. PhD thesis, Université de Genève, 1992.
- [36] A. N. Platonov, An isoperimetric function of the Baumslag–Gersten group. *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* **2004**, no. 3, 12–17, 70; English trans. *Moscow Univ. Math. Bull.* **59** (2004), no. 3, 12–17 (2005). [Zbl 1084.20022](#) [MR 2127449](#)
- [37] H. E. Rose, *Subrecursion: functions and hierarchies*. Oxford Logic Guides 9, The Clarendon Press/Oxford University Press, New York 1984. [Zbl 0539.03018](#) [MR 0752696](#)
- [38] P. Samuelson, On CAT(0) structures for free-by-cyclic groups. *Topology Appl.* **153** (2006), no. 15, 2823–2833. [Zbl 1140.20034](#) [MR 2248386](#)

Received May 14, 2010

Will Dison, Department of Mathematics, University Walk, Bristol, BS8 1TW, U.K.

E-mail: w.dison@bristol.ac.uk

<http://www.maths.bris.ac.uk/~mawjd/>

Timothy R. Riley, Department of Mathematics, 310 Malott Hall, Cornell University, Ithaca, NY 14853, U.S.A.

E-mail: tim.riley@math.cornell.edu

<http://www.math.cornell.edu/~riley/>