

Counting embedded curves in symplectic 6-manifolds

Aleksander Doan and Thomas Walpuski

Abstract. Based on computations of Pandharipande (1999), Zinger (2011) proved that the Gopakumar–Vafa BPS invariants $\text{BPS}_{A,g}(X, \omega)$ for primitive Calabi–Yau classes and arbitrary Fano classes A on a symplectic 6-manifold (X, ω) agree with the signed count $n_{A,g}(X, \omega)$ of embedded J -holomorphic curves representing A and of genus g for a generic almost complex structure J compatible with ω . Zinger’s proof of the invariance of $n_{A,g}(X, \omega)$ is indirect, as it relies on Gromov–Witten theory. In this article we give a direct proof of the invariance of $n_{A,g}(X, \omega)$. Furthermore, we prove that $n_{A,g}(X, \omega) = 0$ for $g \gg 1$, thus proving the Gopakumar–Vafa finiteness conjecture for primitive Calabi–Yau classes and arbitrary Fano classes.

1. Introduction

Are there invariants of symplectic manifolds which count embedded pseudo-holomorphic curves? Such counts can fail to be invariants for two reasons:

- (a) pseudo-holomorphic embeddings can degenerate to multiple covers, and
- (b) they can undergo bubbling and their domains can degenerate.

In the following we consider two situations in which both of these can be ruled out.

Let (X, ω) be a closed symplectic 6-manifold equipped with an almost complex structure J compatible with ω . Denote by $\mathcal{M}_{A,g}^*(X, J)$ the moduli space of simple J -holomorphic maps representing a homology class $A \in H_2(X, \mathbf{Z})$ and of genus g . For a generic choice of J the moduli space $\mathcal{M}_{A,g}^*(X, J)$ is an oriented smooth manifold of dimension

$$\dim \mathcal{M}_{A,g}^*(X, J) = 2\langle c_1(X, \omega), A \rangle.$$

If A is a *Calabi–Yau class*, that is, $\langle c_1(X, \omega), A \rangle = 0$, then $\mathcal{M}_{A,g}^*(X, J)$ is a finite set of signed points and can be counted. If A is primitive in $H_2(X, \mathbf{Z})$, then multiple cover phenomena can be ruled out, and it will be proved that this count defines an invariant $n_{A,g}(X, \omega)$. If A is a *Fano class*, that is, $\langle c_1(X, \omega), A \rangle > 0$, then $\mathcal{M}_{A,g}^*(X, J)$

can be cut-down to a finite set of signed points by imposing incidence conditions governed by suitable cohomology classes $\gamma, \dots, \gamma_\Lambda \in H^{\text{even}}(X, \mathbf{Z})$. In this case, multiple cover phenomena can be ruled out regardless of whether A is primitive or not, and it will be proved that counting the cut-down moduli space defines an invariant $n_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda)$.

These invariants are not new. They were considered by Zinger [39, Theorem 1.5 and footnote 11], who proved that they agree with Gopakumar and Vafa’s BPS invariants. The proof of the invariance of $n_{A,g}(X, \omega)$ and $n_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda)$ in [39] is indirect: it relies on these numbers satisfying the Gopakumar–Vafa formula and the invariance of Gromov–Witten invariants. The novelty in the present work is that we give a much simpler direct proof of invariance. Furthermore, we prove that the invariants vanish for g sufficiently large; thus establishing the Gopakumar–Vafa finiteness conjecture for primitive Calabi–Yau classes and arbitrary Fano classes.

1.1. Ghost components

The main technical result of this paper allows us to rule out, in certain situations, degenerations in which the limiting nodal pseudo-holomorphic map has a *ghost component*, that is, a component on which it is constant. The precise definitions used in the following statement are given in Section 2 and Section 3.

Theorem 1.1. *Let (X, g_∞, J_∞) be an almost Hermitian manifold and let $(J_k)_{k \in \mathbb{N}}$ be a sequence of almost complex structure on X converging to J_∞ in the C^1 topology. If $(u_k: (\Sigma_k, j_k) \rightarrow (X, J_k))_{k \in \mathbb{N}}$ is a sequence of pseudo-holomorphic maps from smooth, closed Riemann surfaces which Gromov converges to the nodal J_∞ -holomorphic map $u_\infty: (\Sigma_\infty, j_\infty, \nu_\infty) \rightarrow (X, J_\infty)$, then one of the following holds:*

- (1) $(\Sigma_\infty, j_\infty, \nu_\infty)$ has no ghost components;
- (2) $(\Sigma_\infty, j_\infty, \nu_\infty)$ has a ghost component C with at least two non-ghost components attached to C ;
- (3) $(\Sigma_\infty, j_\infty, \nu_\infty)$ has a ghost component C with a non-ghost component attached to C in at least two nodes;
- (4) $(\Sigma_\infty, j_\infty, \nu_\infty)$ has a ghost component C with precisely one non-ghost component attached to C at a single node $n \in C$; in that case, $d_{\nu_\infty(n)} u_\infty = 0$, that is, the corresponding node $\nu_\infty(n)$ in the non-ghost component is a critical point of u_∞ .

Remark 1.2. Zinger [38, Theorem 1.2] has analyzed in detail when a nodal pseudo-holomorphic map whose domain has arithmetic genus one appears as a Gromov limit of pseudo-holomorphic maps with smooth domain. Jingchen Niu’s PhD thesis [25] extends Zinger’s analysis to genus two. Their results are based on analyzing the

obstruction map of a Kuranishi model of a neighborhood of the limiting pseudo-holomorphic map. The proof of Theorem 1.1 in Section 5 uses similar methods. This idea goes back to Ionel [18, Proposition 1.20] and Pandharipande [27, Lemma 1]. Recently, a different proof of a result similar to Theorem 1.1 has appeared in the work of Ekholm and Shende [9, Lemma 4.9].

Given a symplectic manifold (X, ω) of dimension at least 6, denote by $\mathcal{J}(X, \omega)$ the set of almost complex structures J compatible with ω and denote by $\mathcal{J}_{\text{emb}}(X, \omega)$ the subset of those J for which the following hold:

- (a) there are no simple J -holomorphic maps of negative index,
- (b) every simple J -holomorphic map is an embedding, and
- (c) every two simple J -holomorphic maps of index zero either have disjoint images or are related by a reparametrization;

see Definition 2.34. The complement of $\mathcal{J}_{\text{emb}}(X, \omega)$ in $\mathcal{J}(X, \omega)$ has codimension two; in particular: $\mathcal{J}_{\text{emb}}(X, \omega)$ is open and dense, and every path $(J_t)_{t \in [0,1]}$ in $\mathcal{J}(X, \omega)$ with end points in $\mathcal{J}_{\text{emb}}(X, \omega)$ is homotopic relative to the end points to a path in $\mathcal{J}_{\text{emb}}(X, \omega)$.

Theorem 1.3. *Let (X, ω) be a compact symplectic 6-manifold, let $(J_k)_{k \in \mathbf{N}}$ be a sequence of almost complex structures compatible with ω converging to J_∞ , and let $(u_k: (\Sigma_k, j_k) \rightarrow (X, J_k))_{k \in \mathbf{N}}$ be a sequence of pseudo-holomorphic maps which Gromov converges to the nodal J_∞ -holomorphic map $u_\infty: (\Sigma_\infty, j_\infty, \nu_\infty) \rightarrow (X, J_\infty)$. Set $A := (u_\infty)_*[\Sigma_\infty] \in H_2(X, \mathbf{Z})$. If A is primitive, satisfies $\langle c_1(X, \omega), A \rangle = 0$, and $J_\infty \in \mathcal{J}_{\text{emb}}(X, \omega)$, then $(\Sigma_\infty, j_\infty, \nu_\infty)$ is smooth and u_∞ is an embedding.*

There is a variant of the definition of $\mathcal{J}(X, \omega)$ adapted to pseudo-holomorphic maps with Λ marked points constrained by pseudo-cycles f_1, \dots, f_Λ . (See Section B for a review of the theory of pseudo-cycles.) The precise definition of this subspace $\mathcal{J}(X, \omega; f_1, \dots, f_\Lambda)$ is rather lengthy and deferred to Definition 2.42.

Theorem 1.4. *Let (X, ω) be a compact symplectic 6-manifold, let $(J_k)_{k \in \mathbf{N}}$ be a sequence of almost complex structures compatible with ω converging to J_∞ , and let $(u_k: (\Sigma_k, j_k) \rightarrow (X, J_k))_{k \in \mathbf{N}}$ be a sequence of pseudo-holomorphic maps which Gromov converges to the nodal J_∞ -holomorphic map $u_\infty: (\Sigma_\infty, j_\infty, \nu_\infty) \rightarrow (X, J_\infty)$. Set $A := (u_\infty)_*[\Sigma_\infty] \in H_2(X, \mathbf{Z})$. Let f_1, \dots, f_Λ be even-dimensional pseudo-cycles of positive codimension in general position. If*

- (1) $\text{im } u_k \cap \text{im } f_\lambda \neq \emptyset$ for every $\lambda = 1, \dots, \Lambda$,
- (2) $2\langle c_1(X, \omega), A \rangle = \sum_{\lambda=1}^\Lambda (\text{codim } f_\lambda - 2) > 0$, and
- (3) $J_\infty \in \mathcal{J}_{\text{emb}}(X, \omega; f_1, \dots, f_\Lambda)$,

then $(\Sigma_\infty, j_\infty, \nu_\infty)$ is smooth and u_∞ is an embedding with $\text{im } u_\infty \cap \text{im } f_\lambda \neq \emptyset$ for every $\lambda = 1, \dots, \Lambda$.

1.2. Embedded curve counts

Denote by $\mathcal{J}_{\text{emb}}^*(X, \omega)$ the subset of those $J \in \mathcal{J}_{\text{emb}}(X, \omega)$ for which every simple J -holomorphic map is unobstructed; see Definition 2.34.

Theorem 1.5. *Let (X, ω) be a symplectic 6-manifold. Let $A \in H_2(X, \mathbf{Z})$ be a primitive class such that $\langle c_1(X, \omega), A \rangle = 0$.*

- (1) *For every $g \in \mathbf{N}_0$ and $J \in \mathcal{J}_{\text{emb}}^*(X, \omega)$, the moduli space $\mathcal{M}_{A,g}^*(X, J)$ of simple J -holomorphic maps representing the class A and of genus g is a compact oriented zero-dimensional manifold, and the signed count*

$$n_{A,g}(X, \omega) := \#\mathcal{M}_{A,g}^*(X, J)$$

is independent on the choice of J .

- (2) *There is a $g_0 \in \mathbf{N}_0$, depending on (X, ω) and A , such that*

$$n_{A,g}(X, \omega) = 0 \quad \text{for every } g \geq g_0.$$

Remark 1.6. In fact, $n_{A,g}(X, \omega)$ depends on ω only up to deformation.

Again, there is a variant $\mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda)$ of $\mathcal{J}_{\text{emb}}^*(X, \omega)$ adapted to pseudo-holomorphic maps with Λ marked points constrained by pseudo-cycles f_1, \dots, f_Λ ; see Definition 2.43.

Theorem 1.7. *Let (X, ω) be a symplectic 6-manifold, let $A \in H_2(X, \mathbf{Z})$, and let $\gamma_1, \dots, \gamma_\Lambda \in H^{\text{even}}(X, \mathbf{Z})$ be such that $\text{deg}(\gamma_\lambda) > 0$ and*

$$2\langle c_1(X, \omega), A \rangle = \sum_{\lambda=1}^{\Lambda} (\text{deg}(\gamma_\lambda) - 2) > 0.$$

- (1) *Let f_1, \dots, f_Λ be pseudo-cycles in X which are Poincaré dual to $\gamma_1, \dots, \gamma_\Lambda$ and in general position. For every $g \in \mathbf{N}_0$ and $J \in \mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda)$ the moduli space $\mathcal{M}_{A,g}^*(X, J; f_1, \dots, f_\Lambda)$ of simple J -holomorphic maps representing the class A , of genus g , and intersecting f_1, \dots, f_Λ is a compact oriented zero-dimensional manifold, and the signed count*

$$n_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda) := \#\mathcal{M}_{A,g}^*(X, J; f_1, \dots, f_\Lambda)$$

is independent on the choice of f_1, \dots, f_Λ and J .

(2) *There exists a $g_0 \in \mathbf{N}_0$, depending on (X, ω) , A , and $\gamma_1, \dots, \gamma_\Lambda$, such that*

$$n_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda) = 0 \quad \text{for all } g \geq g_0.$$

Remark 1.8. Remark 1.6 applies mutatis mutandis.

1.3. Gopakumar and Vafa’s BPS invariants

Using ideas from M -theory, Gopakumar and Vafa [10, 11] predicted that there are integer invariants $\text{BPS}_{A,g}(X, \omega)$ associated with every closed symplectic 6-manifold (X, ω) , a class $A \in H_2(X, \mathbf{Z})$ with $\langle c_1(X, \omega), A \rangle = 0$, and $g \in \mathbf{N}_0$, which count BPS states supported on embedded J -holomorphic curves representing A and of genus g . Gopakumar and Vafa did not give a direct mathematical definition of $\text{BPS}_{A,g}(X, \omega)$; however, they conjectured that their invariants are related to the Gromov–Witten invariants $\text{GW}_{A,g}(X, \omega)$ by the marvelous formula

$$\begin{aligned} & \sum_A \sum_{g=0}^{\infty} \text{GW}_{A,g}(X, \omega) \cdot t^{2g-2} q^A \\ &= \sum_A \sum_{g=0}^{\infty} \text{BPS}_{A,g}(X, \omega) \cdot \sum_{k=1}^{\infty} \frac{1}{k} (2 \sin(kt/2))^{2g-2} q^{kA} \end{aligned} \tag{1.9}$$

with the sum taken over all non-zero Calabi–Yau classes A and, moreover, that

$$\text{BPS}_{A,g}(X, \omega) = 0 \quad \text{for } g \gg 1.$$

In algebraic geometry, there are approaches to defining the BPS invariants for projective Calabi–Yau three-folds [15, 21, 23, 29, 30]. These satisfy the Gopakumar–Vafa formula (1.9) in some cases, but it is not currently known whether the formula holds in general.

An alternative approach is to take (1.9) as the *definition* of $\text{BPS}_{A,g}(X, \omega)$; see [3, Section 2]. This approach leads to the following conjecture.

Conjecture 1.10 (Gopakumar and Vafa [10, 11]; see also [3, Conjecture 1.2]). *The numbers $\text{BPS}_{A,g}(X, \omega)$ defined by (1.9) satisfy*

- (integrality) $\text{BPS}_{A,g}(X, \omega) \in \mathbf{Z}$, and
- (finiteness) $\text{BPS}_{A,g}(X, \omega) = 0$ for $g \gg 1$.

The Gopakumar–Vafa integrality conjecture has been proved by Ionel–Parker [19]. Zinger [39, footnote 11] has proved that for primitive Calabi–Yau classes

$$\text{BPS}_{A,g}(X, \omega) = n_{A,g}(X, \omega);$$

see also Section C. Therefore, Theorem 1.5 implies the following.

Corollary 1.11. *The Gopakumar–Vafa finiteness conjecture holds for primitive Calabi–Yau classes; that is, for every closed symplectic 6-manifold (X, ω) and every primitive Calabi–Yau class $A \in H_2(X, \mathbf{Z})$, there is a $g_0(\omega, A)$ such that for every $g \geq g_0(\omega, A)$,*

$$\text{BPS}_{A,g}(X, \omega) = 0.$$

Remark 1.12. The finiteness conjecture for general Calabi–Yau classes has been resolved recently [6].

The genus bound in Corollary 1.11 is not effective; therefore, it is natural to ask the following.

Definition 1.13. Let (X, ω) be a closed symplectic 6-manifold and $A \in H_2(X, \mathbf{Z})$ a Calabi–Yau class. Define the *BPS Castelnuovo number* $\gamma_A(X, \omega)$ by

$$\gamma_A(X, \omega) := \inf\{g \in \mathbf{N} : \text{BPS}_{A,g}(X, \omega) = 0\} \in \mathbf{N}_0$$

Question 1.14. Is there an bound on $\gamma_A(X, \omega)$ analogous to Castelnuovo’s bound for the genus of an irreducible degree d curve in \mathbf{P}^n (see [4] and [2, Chapter III, Section 2]); that is, a bound of $\gamma_A(X, \omega)$ by a formula involving A and the geometry of X ? (See [16, 22] for some work in this direction.)

There is an analogue of the Gopakumar–Vafa formula for Fano classes. Given $A \in H_2(X, \mathbf{Z})$, $g \in \mathbf{N}_0$, and $\gamma_1, \dots, \gamma_\Lambda \in H^{\text{even}}(X, \mathbf{Z})$ satisfying $\deg(\gamma_\lambda) > 0$ and

$$2\langle c_1(X, \omega), A \rangle = \sum_{\lambda=1}^{\Lambda} (\deg(\gamma_\lambda) - 2) > 0, \tag{1.15}$$

denote by $\text{GW}_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda)$ be the corresponding Gromov–Witten invariant. The analogue of (1.9) is

$$\begin{aligned} & \sum_A \sum_{g=0}^{\infty} \text{GW}_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda) \cdot t^{2g-2} q^A \\ &= \sum_A \sum_{g=0}^{\infty} \text{BPS}_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda) \cdot (2 \sin(t/2))^{2g-2+\langle c_1(X, \omega), A \rangle} q^A \end{aligned}$$

with the sum taken over all $A \in H_2(X, \mathbf{Z})$ satisfying (1.15). Zinger [39, Theorem 1.5] has proved that

$$\text{BPS}_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda) = n_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda);$$

thus establishing the analogue of the Gopakumar–Vafa integrality conjecture. Furthermore, Theorem 1.7 implies the following.

Corollary 1.16. *The analogue of the Gopakumar–Vafa finiteness conjecture holds for all Fano classes.*

Of course, there is an analogue of Question 1.14 in the Fano case.

2. Nodal pseudo-holomorphic maps

This section reviews a few definitions and results regarding nodal pseudo-holomorphic maps.

2.1. Nodal manifolds

Definition 2.1. Let X be a manifold, possibly disconnected. A *nodal structure* on X is an involution $\nu: X \rightarrow X$ whose fixed-point set has a discrete complement. (This involution is discontinuous unless $\nu = \text{id}$.) The set of points not fixed by ν is called the *nodal set*. A *nodal manifold* is a manifold together with a nodal structure.

The quotient X/ν should be considered as the topological space underlying the nodal manifold (X, ν) . The atlas of X induces a “nodal atlas” for X/ν consisting of “charts” mapping either to \mathbf{R}^n or $\mathbf{R}^n \times \{0\} \cup \{0\} \times \mathbf{R}^n \subset \mathbf{R}^{2n}$. The nodes of X/ν are precisely the points mapping to $(0, 0) \in \mathbf{R}^{2n}$ in some chart or, equivalently, the images of the points in the nodal set.

Definition 2.2. Let (X_1, ν_1) and (X_2, ν_2) be nodal manifolds. A *nodal map*

$$f: (X_1, \nu_1) \rightarrow (X_2, \nu_2)$$

is a smooth map $f: X_1 \rightarrow X_2$ such that

$$f \circ \nu_1 = \nu_2 \circ f.$$

Definition 2.3. Let (X, ν) be a nodal manifold. A *diffeomorphism* of (X, ν) is an element of

$$\text{Diff}(X, \nu) := \{\phi \in \text{Diff}(X) : \phi \circ \nu = \nu \circ \phi\}.$$

Every manifold X canonically is a nodal manifold with $\nu = \text{id}_X$ and a smooth map between manifolds, trivially, is a nodal map. In other words, the category of manifolds is a full subcategory of the category of nodal manifolds.

Definition 2.4. Let (X, ν) be a nodal manifold, Y be a manifold, and $f: (X, \nu) \rightarrow Y$ be a nodal map. For a vector bundle $E \rightarrow Y$, set

$$\Gamma(X, \nu; f^*E) := \{\xi \in \Gamma(X, f^*E) : \xi \circ \nu = \xi\}.$$

Remark 2.5. In the situation of the preceding definition, set $n := \dim X$ and let $p > n$. Given a Riemannian metric on X , a Euclidean metric on E , and a metric covariant derivative on E , denote by $W^{1,p}\Gamma(X, f^*E)$ the completion of $\Gamma(X, f^*E)$ with respect to the corresponding $W^{1,p}$ norm. By Morrey’s embedding theorem,

$$W^{1,p} \hookrightarrow C^{0,1-n/p}.$$

Therefore, the evaluations maps $\text{ev}_x: \Gamma(X, f^*E) \rightarrow E_{f(x)}$ extend to $W^{1,p}\Gamma(X, f^*E)$ and

$$W^{1,p}\Gamma(X, \nu; f^*E) = \{\xi \in W^{1,p}\Gamma(X; f^*E) : \xi(\nu(x)) = \xi(x) \text{ for every } x \in X\}.$$

For $p < n$, it can be shown that the $W^{1,p}$ completion of $\Gamma(X, \nu; f^*E)$ agrees with the $W^{1,p}$ completion of $\Gamma(X; f^*E)$.

2.2. Nodal Riemann surfaces

Definition 2.6. A nodal Riemann surface is a Riemann surface (Σ, j) together with a nodal structure ν .

Definition 2.7. Let C be a complex analytic curve. A point of C is a *node* if it has a neighborhood which is isomorphic to a neighborhood of the point $(0, 0)$ in the curve

$$\{(z, w) \in \mathbb{C}^2 : zw = 0\}.$$

A *nodal curve* is a complex analytic curve all of whose points are either smooth or a node.

Let C be a nodal curve and denote by $\pi: \tilde{C} \rightarrow C$ its normalization. The complex analytic curve \tilde{C} is smooth and, hence, equivalent to a closed Riemann surface (Σ, j) . Since \tilde{C} is obtained from C by replacing every node with a pair of points, Σ inherits a canonical nodal structure ν . This sets up an equivalence between complete, nodal curves C and closed, nodal Riemann surfaces (Σ, j, ν) .

Definition 2.8. The *automorphism group* of a nodal Riemann surface (Σ, j, ν) is

$$\text{Aut}(\Sigma, j, \nu) := \{\phi \in \text{Diff}(\Sigma, \nu) : \phi_*j = j\}.$$

A nodal Riemann surface (Σ, j, ν) is *stable* if $\text{Aut}(\Sigma, j, \nu)$ is finite.

Definition 2.9. Let (Σ, ν) be a nodal surface with nodal set S . The *arithmetic genus* of (Σ, ν) is

$$p_a(\Sigma, \nu) := 1 - \frac{1}{2}(\chi(\Sigma) - \#S). \tag{2.10}$$

Remark 2.11. If $(\tilde{\Sigma}, \tilde{\nu})$ denotes a nodal surface obtained from (Σ, ν) by attaching a 1-handle at some pairs of nodes $\{n, \nu(n)\}$, then

$$p_a(\Sigma, \nu) = p_a(\tilde{\Sigma}, \tilde{\nu}).$$

2.3. Nodal J -holomorphic maps

Throughout the next four subsections, let (X, J) be an almost complex manifold of dimension $2n$.

Definition 2.12. A nodal J -holomorphic map $u: (\Sigma, j, \nu) \rightarrow (X, J)$ is a nodal Riemann surface (Σ, j, ν) together with a nodal map $u: (\Sigma, \nu) \rightarrow X$ which is J -holomorphic; that is,

$$\bar{\partial}_J(u, j) := \frac{1}{2}(du + J(u) \circ du \circ j) = 0.$$

Definition 2.13. If $u: (\Sigma, j, \nu) \rightarrow (X, J)$ is a nodal J -holomorphic map and $\phi \in \text{Diff}(\Sigma, \nu)$, then the *reparametrization*

$$\phi_*u := u \circ \phi^{-1}: (\Sigma, \phi_*j, \nu) \rightarrow (X, J)$$

is a nodal J -holomorphic map as well. The *automorphism group* of a nodal J -holomorphic map $u: (\Sigma, j, \nu) \rightarrow (X, J)$ is

$$\text{Aut}(u) := \{\phi \in \text{Aut}(\Sigma, j, \nu) : u \circ \phi = u\}.$$

The map u is said to be *stable* if $\text{Aut}(u)$ is finite.

Definition 2.14. Let (Σ, j) and $(\tilde{\Sigma}, \tilde{j})$ be smooth Riemann surfaces. Let $u: (\Sigma, j) \rightarrow (X, J)$ be a J -holomorphic map and let $\pi: (\tilde{\Sigma}, \tilde{j}) \rightarrow (\Sigma, j)$ be a holomorphic map of degree $\text{deg}(\pi) \geq 2$. The composition $u \circ \pi: (\tilde{\Sigma}, \tilde{j}) \rightarrow (X, J)$ is said to be a *multiple cover* of u . A J -holomorphic map is *simple* if it is not constant and not a multiple cover.

2.4. Ghost components

Let $u: (\Sigma, j, \nu) \rightarrow (X, J)$ be a nodal J -holomorphic map. Let S be the nodal set of (Σ, ν) .

Definition 2.15. Suppose $C \subset \Sigma$ is a union of connected components of Σ . Set

$$S_C^{\text{int}} := \{n \in S : n \in C \text{ and } \nu(n) \in C\} \quad \text{and} \quad S_C^{\text{ext}} := \{n \in S : n \in C \text{ and } \nu(n) \notin C\},$$

and denote by ν_C the nodal structure on C which agrees with ν_0 on S_C^{int} and the identity on the complement of S_C^{int} . Denote by \check{C} the nodal curve associated with (C, j, ν) .

Definition 2.16. A ghost component of u is a union C of connected components of Σ such that $u|_C$ is constant, \check{C} is connected, and which is a maximal subset satisfying these properties.

Proposition 2.18 below, which will be used in the proof of Theorem 1.1 (specifically in Section 5.7), concerns the dualizing sheaf of a nodal curve C . The dualizing sheaf is a generalization of the canonical sheaf of a smooth curve; for the reader’s convenience, we describe its construction in the proof of Proposition 2.18.

Definition 2.17. Let C be a nodal curve. The dual graph of C is the weighted graph whose set of vertices is the set of irreducible components of C with the genus as the weight function, and edges between two vertices if and only if the corresponding irreducible components of C intersect.

Proposition 2.18. Let C be a nodal curve. Denote the dual graph of C by Γ . Denote by ω_C the dualizing sheaf of C and by B its base-locus:

$$B := \{x \in C : \zeta(x) = 0 \text{ for every } \zeta \in H^0(C, \omega_C)\}.$$

The base-locus has the following description:

- (1) B is a union of irreducible rational components of C .
- (2) The dual graph of B is the subgraph $\Delta \subset \Gamma$ obtained by
 - (a) removing every vertex of non-zero weight, and
 - (b) removing every simple cycle in Γ .

In particular, Δ is a forest with weight zero. Moreover, if e_1, e_2 are distinct vertices of a tree $T \subset \Delta$, then they cannot be connected by a path in $(\Gamma \setminus T) \cup \{e_1, e_2\}$.

The proof relies on the following.

Proposition 2.19. Let Σ be a connected smooth curve. For every three p, q, r distinct points on Σ there is a $\zeta \in H^0(K_\Sigma(p + q))$ with

$$\text{Res}_p \zeta = -\text{Res}_q \zeta \neq 0 \quad \text{and} \quad \zeta(r) \neq 0.$$

Here $\text{Res}_p \zeta$ denotes the residue at p of the meromorphic 1-form ζ .

Proof. For $\Sigma = \mathbf{P}^1$ without loss of generality $p = 0$ and $q = \infty$; hence, the meromorphic 1-form can be taken to be $z^{-1}dz$.

Suppose $\Sigma \neq \mathbf{P}^1$. Consider the exact sequence

$$H^0(K_\Sigma) \hookrightarrow H^0(K_\Sigma(p + q)) \xrightarrow{\rho} H^0(\mathcal{O}_p \oplus \mathcal{O}_q) \cong \mathbf{C} \oplus \mathbf{C} \xrightarrow{\delta} H^1(K_\Sigma) \cong \mathbf{C}$$

with $\rho(\zeta) := (\text{Res}_p \zeta, \text{Res}_q \zeta)$ and $\delta(a, b) := a - b$. This implies that there is a $\zeta \in H^0(K_\Sigma(p + q))$ with non-vanishing residues at p and q . Since K_Σ is base-point-free, ζ can be arranged not to vanish at r . ■

Proof of Proposition 2.18. The dualizing sheaf of C is constructed as follows; see [1, p. 91]. Denote by $\pi: \Sigma \rightarrow C$ the normalization map. Denote by S the set of nodal points of C . Denote by $\tilde{\omega}_C$ the subsheaf of $K_\Sigma(S)$ whose sections ζ satisfy

$$\text{Res}_n \zeta + \text{Res}_{\nu(n)} \zeta = 0$$

for every $n \in S$. Here ν denotes the obvious involution on $\pi^{-1}(S)$. The dualizing sheaf ω_C then is

$$\omega_C = \pi_* \tilde{\omega}_C.$$

The base-locus of K_Σ are precisely the rational connected components of Σ . This implies Proposition 2.18 (1). It follows from the above Proposition 2.19 that the dual graph of B is contained in Δ . By the Residue theorem any meromorphic 1-form with simple poles must have at least two poles. This implies that the dual graph of B agrees with Δ . ■

2.5. Moduli spaces of nodal pseudo-holomorphic maps

Definition 2.20. Given $A \in H_2(X, \mathbf{Z})$ and $g \in \mathbf{N}_0$, the *moduli space of stable nodal J -holomorphic maps* representing A and of genus g is the set

$$\bar{\mathcal{M}}_{A,g}(X, J)$$

of equivalence classes of stable nodal J -holomorphic maps $u: (\Sigma, j, \nu) \rightarrow (X, J)$ up to reparametrization with

$$u_*[\Sigma] = A \quad \text{and} \quad p_a(\Sigma, \nu) = g.$$

The subset of $\bar{\mathcal{M}}_{A,g}(X, J)$ parametrizing simple J -holomorphic maps is denoted by

$$\mathcal{M}_{A,g}^*(X, J).$$

At this stage, $\bar{\mathcal{M}}_{A,g}(X, J)$ is just a set. In Section 3.2, it will be equipped with the *Gromov topology*. This topology induces the C^∞ topology on $\mathcal{M}_{A,g}^*(X, J)$.

Definition 2.21. Let (X, ω) be a symplectic manifold. Denote by $\mathcal{J}(X, \omega)$ the space of almost complex structures on X which are *compatible with ω* ; that is,

$$g(\cdot, \cdot) := \omega(\cdot, J\cdot)$$

defines a Riemannian metric on X . Equip $\mathcal{J}(X, \omega)$ with the C^∞ topology.

Definition 2.22. Given $A \in H_2(X, \mathbf{Z})$ and $g \in \mathbf{N}$, set

$$\bar{\mathcal{M}}_{A,g}(X, \omega) := \coprod_{J \in \mathcal{J}(X, \omega)} \bar{\mathcal{M}}_{A,g}(X, J) \quad \text{and} \quad \mathcal{M}_{A,g}^*(X, \omega) := \coprod_{J \in \mathcal{J}(X, \omega)} \mathcal{M}_{A,g}^*(X, J).$$

Denote by $\pi: \bar{\mathcal{M}}_{A,g}(X, \omega) \rightarrow \mathcal{J}(X, \omega)$ the canonical projection.

2.6. Linearization of the J -holomorphic map equation

Let $u: (\Sigma, j, \nu) \rightarrow (X, J)$ be a nodal J -holomorphic map. Let h be a Hermitian metric on (X, J) and let ∇ be a torsion-free connection on TX . Throughout the remainder of this article, let $p > 2$.

Definition 2.23. Given $\xi \in W^{1,p}\Gamma(\Sigma, \nu; u^*TX)$, set

$$u_\xi := \exp_u(\xi)$$

and denote by

$$\Psi_\xi: L^p\Omega^{0,1}(\Sigma, u^*TX) \rightarrow L^p\Omega^{0,1}(\Sigma, u_\xi^*TX)$$

the map induced by parallel transport along the geodesics $t \mapsto \exp_u(t\xi)$. Define

$$\mathfrak{F}_{u,j,\nu;J}: W^{1,p}\Gamma(\Sigma, \nu; u^*TX) \rightarrow L^p\Omega^{0,1}(\Sigma, u^*TX)$$

by

$$\mathfrak{F}_{u,j,\nu;J}(\xi) := \Psi_\xi^{-1} \bar{\partial}_J(u_\xi, j).$$

Definition 2.24. Define the linear operator

$$\mathfrak{d}_{u,j,\nu;J}: W^{1,p}\Gamma(\Sigma, \nu; u^*TX) \rightarrow L^p\Omega^{0,1}(\Sigma, u^*TX)$$

by

$$\mathfrak{d}_{u,j,\nu;J}\xi := \mathfrak{d}_0\mathfrak{F}_{u,j,\nu;J}\xi = \frac{1}{2}(\nabla\xi + J(u) \circ (\nabla\xi) \circ j + (\nabla_\xi J) \circ du \circ j).$$

Remark 2.25. If u is J -holomorphic, then $\mathfrak{d}_{u,j,\nu;J}$ does not depend on the choice of torsion-free connection ∇ on TX ; see [24, Proposition 3.1.1].

The operator $\mathfrak{d}_{u,j,\nu;J}$ is the restriction to $W^{1,p}\Gamma(\Sigma, \nu; u^*TX)$ of the operator

$$\mathfrak{d}_{u,j;J}: W^{1,p}\Gamma(\Sigma, u^*TX) \rightarrow L^p\Omega^{0,1}(\Sigma, u^*TX)$$

given by the same formula. The former controls the deformation theory of u as a nodal J -holomorphic map from the nodal Riemann surface (Σ, j, ν) whereas the latter controls the deformation theory of u as a smooth J -holomorphic map from the smooth Riemann surface (Σ, j) , ignoring the nodal structure.

Proposition 2.26. *The index of $\delta_{u,j,v;J}$ is given by*

$$\text{index } \delta_{u,j,v;J} = 2\langle [\Sigma], u^*c_1(X, J) \rangle + 2n(1 - p_a(\Sigma, v)). \tag{2.27}$$

Proof. The inclusion

$$W^{1,p}\Gamma(\Sigma, v; u^*TX) \rightarrow W^{1,p}\Gamma(\Sigma, u^*TX)$$

has index $-n\#S$. By the Riemann–Roch theorem,

$$\text{index } \delta_{u,j;J} = 2\langle [\Sigma], u^*c_1(X, J) \rangle + n\chi(\Sigma).$$

These together with (2.10) imply the index formula. ■

Remark 2.28. For our discussion in Section 5.7, which establishes the key technical result of this article, the following detailed description of the kernel and cokernel of $\delta_{u,j,v;J}$ will be important. Denote by

$$V_- \subset \bigoplus_{n \in S} T_{u(n)}X$$

the subspace of those $(v_n)_{n \in S}$ satisfying

$$v_{v(n)} = -v_n.$$

Define $\text{diff}: \ker \delta_{u,j;J} \rightarrow V_-$ by

$$\text{diff}k := (\kappa(n) - \kappa(v(n)))_{n \in S}.$$

Evidently,

$$\ker \delta_{u,j,v;J} = \ker \text{diff}.$$

The map diff is induced by the analogously defined map $W^{1,p}(\Sigma, u^*TX) \rightarrow V_-$, which fits in to the following commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & W^{1,p}(\Sigma, v; u^*TX) & \longrightarrow & W^{1,p}(\Sigma, u^*TX) & \longrightarrow & V_- \longrightarrow 0 \\ & & \downarrow \delta_{u,j,v;J} & & \downarrow \delta_{u,j;J} & & \downarrow \\ 0 & \longrightarrow & L^p\Omega^{0,1}(\Sigma, u^*TX) & \xrightarrow{=} & L^p\Omega^{0,1}(\Sigma, u^*TX) & \longrightarrow & 0 \longrightarrow 0. \end{array}$$

Therefore, the Snake lemma yields the short exact sequence

$$0 \longrightarrow \text{coker } \text{diff} \longrightarrow \text{coker } \delta_{u,j,v;J} \longrightarrow \text{coker } \delta_{u,j;J} \longrightarrow 0.$$

The dual sequence

$$0 \longrightarrow (\text{coker } \delta_{u,j;J})^* \longrightarrow (\text{coker } \delta_{u,j,v;J})^* \longrightarrow (\text{coker } \text{diff})^* \longrightarrow 0$$

can be understood as follows. Let $q \in (1, 2)$ be such that $1/p + 1/q = 1$. The dual space $(\text{coker } \mathfrak{d}_{u,j,v;J})^*$ can be identified via the pairing between L^p and L^q with the space \mathcal{H} consisting of those $\zeta \in L^q \Omega^{0,1}(\Sigma, u^*TX)$ which satisfy a distributional equation of the form

$$\mathfrak{d}_{u,j,J}^* \zeta = \sum_{n \in S} v_n \delta_n.$$

with $v = (v_n)_{n \in S} \in (\text{im diff})^\perp \cong (\text{coker diff})^*$ and δ_n denoting the Dirac δ distribution at n . The map $(\text{coker } \mathfrak{d}_{u,j,v;J})^* \rightarrow (\text{coker diff})^*$ maps ζ to v .

Definition 2.29. Define the map

$$\mathfrak{n}_{u,j,v;J}: W^{1,p} \Gamma(\Sigma, \nu; u^*TX) \rightarrow L^p \Omega^{0,1}(\Sigma, u^*TX)$$

by

$$\mathfrak{n}_{u,j,v;J}(\xi) := \mathfrak{F}_{u,j,v;J}(\xi) - \bar{\partial}_J(u, j) - \mathfrak{d}_{u,j,v;J} \xi.$$

Proposition 2.30 ([24, Proposition 3.5.3 and Remark 3.5.5]). *Denote by $c_S > 0$ an upper bound for the norm of the embedding $W^{1,p}(\Sigma) \hookrightarrow C^{0,1-2/p}(\Sigma)$ and let $c_\xi > 0$. For every ξ_1, ξ_2 with $\|\xi_1\|_{W^{1,p}} \leq c_\xi$ and $\|\xi_2\|_{W^{1,p}} \leq c_\xi$,*

$$\begin{aligned} & \|\mathfrak{n}_{u,j,v;J}(\xi_1) - \mathfrak{n}_{u,j,v;J}(\xi_2)\|_{L^p} \\ & \leq c(c_S, c_\xi, \|\text{du}\|_{L^p}) \cdot (\|\xi_1\|_{W^{1,p}} + \|\xi_2\|_{W^{1,p}}) \cdot \|\xi_1 - \xi_2\|_{W^{1,p}}. \end{aligned}$$

So far, the complex structure j has been held fixed. Denote by $\mathcal{J}(\Sigma)$ the space of complex structures on Σ and by $\text{Diff}_0(\Sigma, \nu)$ the group of diffeomorphism of Σ which are isotopic to the identity and commute with ν . Denote by

$$\mathcal{T} := \mathcal{J}(\Sigma) / \text{Diff}_0(\Sigma, \nu)$$

the corresponding Teichmüller space. This is a complex manifold whose real dimension satisfies

$$\dim \mathcal{T} - \dim \text{aut}(\Sigma, j, \nu) + \#S = 6(p_a(\Sigma, \nu) - 1).$$

For every $j \in \mathcal{J}(\Sigma)$, there is a *Teichmüller slice through j* ; that is, an open neighborhood Δ of $0 \in \mathbf{C}^{\dim \mathcal{T}}$ together with a $\text{Aut}(\Sigma, j, \nu)$ -equivariant map $J: \Delta \rightarrow \mathcal{J}(\Sigma)$ such that $J(0) = j$.

Definition 2.31. Consider the bundle over Δ whose fiber over $\sigma \in \Delta$ is the Banach space

$$L^p \Omega^{0,1}(\Sigma, u^*TX).$$

Here the space of $(0, 1)$ forms on Σ is defined with respect to complex structure $J(\sigma)$. A choice of a trivialization of this bundle gives rise to the map

$$\begin{aligned} W^{1,p}\Gamma(\Sigma, \nu; u^*TX) \times \Delta &\rightarrow L^p\Omega^{0,1}(\Sigma, u^*TX), \\ (\xi, \sigma) &\mapsto \mathcal{F}_{u,J(\sigma),\nu;J}(\xi). \end{aligned} \tag{2.32}$$

Define $d_{u,j}\bar{\partial}_{\nu;J}: W^{1,p}\Gamma(\Sigma, \nu; u^*TX) \oplus T_0\Delta \rightarrow L^p\Omega^{0,1}(\Sigma, u^*TX)$ to be the derivative of the map (2.32) at $(0, 0)$.

Definition 2.33. The *index* of u is

$$\begin{aligned} \text{index}(u) &:= \text{index}(d_{u,j}\bar{\partial}_{\nu;J}) - \dim \text{aut}(\Sigma, j, \nu) + \#S \\ &= 2\langle [\Sigma], u^*c_1(X, J) \rangle + 2(n - 3)(1 - p_a(\Sigma, \nu)). \end{aligned}$$

The map $u: (\Sigma, j, \nu) \rightarrow (X, J)$ is said to be *unobstructed* if $d_{u,j}\bar{\partial}_{\nu;J}$ is surjective.

Henceforth, to simplify notation, we will often drop some or all of the subscripts j, ν, J from the maps defined above.

2.7. Transversality for simple maps

Throughout the remainder of this section, (X, ω) is a compact symplectic manifold of dimension $2n \geq 6$ and we only consider pseudo-holomorphic maps from smooth Riemann surfaces.

Definition 2.34. Denote by $\mathcal{J}_{\text{emb}}(X, \omega) \subset \mathcal{J}(X, \omega)$ the subspace of those almost complex structures compatible with ω for which the following hold:

- (1) there are no simple J -holomorphic maps with negative index,
- (2) every simple J -holomorphic map with $\text{index}(u) < 2n - 4$ is an embedding, and
- (3) every pair of simple J -holomorphic maps u_1, u_2 satisfying

$$\text{index}(u_1) + \text{index}(u_2) < 2n - 4$$

either have disjoint images or are related by a reparametrization.

Denote by $\mathcal{J}_{\text{emb}}^*(X, \omega) \subset \mathcal{J}_{\text{emb}}(X, \omega)$ the subset of those J for which, moreover,

- (4) every simple J -holomorphic map is unobstructed.

Definition 2.35. Given $J_0, J_1 \in \mathcal{J}(X, \omega)$, denote by $\mathcal{J}(X, \omega; J_0, J_1)$ the space of smooth paths $(J_t)_{t \in [0,1]}$ in $\mathcal{J}(X, \omega)$ from J_0 and J_1 . Given $J_0, J_1 \in \mathcal{J}_{\text{emb}}^*(X, \omega)$, denote by $\mathcal{J}_{\text{emb}}^*(X, \omega, J_0, J_1)$ the subset of those $(J_t)_{t \in [0,1]} \in \mathcal{J}(X, \omega; J_0, J_1)$ such that for every $t \in [0, 1]$:

- (1) $J_t \in \mathcal{J}_{\text{emb}}(X, \omega)$, and
- (2) if $u: (\Sigma, j) \rightarrow (X, J_t)$ is a simple J_t -holomorphic map, then either
 - (a) $\text{coker } d_{u,j} \bar{\partial}_{J_t} = \{0\}$, or
 - (b) $\dim \text{coker } d_{u,j} \bar{\partial}_{J_t} = 1$ and the map $\ker d_{u,j} \bar{\partial}_{J_t} \rightarrow \text{coker } d_{u,j} \bar{\partial}_{J_t}$ defined by

$$\xi \mapsto \text{pr} \left(\frac{d}{ds} \Big|_{s=t} d_{u,j} \bar{\partial}_{J_s} \xi \right),$$

with $\text{pr}: \Omega^{0,1}(\Sigma, u^*TX) \rightarrow \text{coker } d_{u,j} \bar{\partial}_{J_t}$ denoting the canonical projection, is surjective.

Proposition 2.36. *Let $A \in H_2(X, \mathbf{Z})$ and $g \in \mathbf{N}_0$.*

- (1) *For every $J \in \mathcal{J}_{\text{emb}}^*(X, \omega)$, the moduli space $\mathcal{M}_{A,g}^*(X, J)$ is an oriented smooth manifold of dimension*

$$2\langle c_1(X, \omega), A \rangle + 2(n - 3)(1 - g).$$

- (2) *For every pair $J_0, J_1 \in \mathcal{J}_{\text{emb}}^*(X, \omega)$ and $(J_t)_{t \in [0,1]} \in \mathcal{J}_{\text{emb}}^*(X, \omega; J_0, J_1)$, the moduli space*

$$\mathcal{M}_{A,g}^*(X, (J_t)_{t \in [0,1]}) := \coprod_{t \in [0,1]} \mathcal{M}_{A,g}^*(X, J_t),$$

is an oriented smooth manifold with boundary

$$\mathcal{M}_{A,g}^*(X, J_1) \amalg -\mathcal{M}_{A,g}^*(X, J_0).$$

This is a consequence of the Implicit function theorem; see [24, Theorem 3.1.6 and Theorem 3.1.7]. The orientation on the moduli spaces is obtained by trivializing the determinant line bundle of the family of operators $d_{u,j} \bar{\partial}_J$; see [24, Proof of Theorem 3.1.6, Remark 3.2.5, Appendix A.2]. If the moduli space is zero-dimensional, that is, a discrete set, then every $[u] \in \mathcal{M}_{A,g}^*(X, J)$ is assigned a sign

$$\text{sign}[u] \in \{+1, -1\}.$$

The signed count of $\mathcal{M}_{A,g}^*(X, J)$ is then

$$\#\mathcal{M}_{A,g}^*(X, J) := \sum_{[u] \in \mathcal{M}_{A,g}^*(X, J)} \text{sign}[u].$$

Proposition 2.37. *The following hold:*

- (1) $\mathcal{J}_{\text{emb}}^*(X, \omega) \subset \mathcal{J}(X, \omega)$ is residual.

(2) For every pair

$$J_0, J_1 \in \mathcal{J}_{\text{emb}}^*(X, \omega), \quad \mathcal{J}_{\text{emb}}^*(X, \omega; J_0, J_1) \subset \mathcal{J}(X, \omega; J_0, J_1)$$

is residual.

The proof is a standard application of the Sard–Smale theorem; cf. [26, Theorem 1.2], [19, Proposition A.4], and [24, Sections 3.2 and 6.3]. Some details of the proof will be reviewed in the proof of Proposition 2.45.

2.8. J -holomorphic maps with constraints

Definition 2.38. Let $\Lambda \in \mathbf{N}$. A J -holomorphic map with Λ marked points is a J -holomorphic map $u: (\Sigma, j) \rightarrow (X, J)$ together with Λ distinct labeled points $z_1, \dots, z_\Lambda \in \Sigma$.

The reparametrization of $(u; z_1, \dots, z_\Lambda)$ by $\phi \in \text{Diff}(\Sigma)$ is the J -holomorphic map with Λ marked points $\phi_*(u; z_1, \dots, z_\Lambda) := (u \circ \phi^{-1}; \phi(z_1), \dots, \phi(z_\Lambda))$.

A J -holomorphic map $(u; z_1, \dots, z_\Lambda)$ with Λ marked points is said to be simple if u is simple.

Definition 2.39. Given $A \in H_2(X, \mathbf{Z})$, $g \in \mathbf{N}_0$, $\Lambda \in \mathbf{N}$, and $J \in \mathcal{J}(X, \omega)$, the moduli space of simple J -holomorphic maps with Λ marked points representing A and of genus g is the set

$$\mathcal{M}_{A,g,\Lambda}^*(X, J)$$

of equivalence classes J -holomorphic maps $u: (\Sigma, j) \rightarrow (X, J)$ with Λ marked points z_1, \dots, z_Λ up to reparametrization with

$$u_*[\Sigma] = A \quad \text{and} \quad g(\Sigma) = g.$$

Define the evaluation map $\text{ev}: \mathcal{M}_{A,g,\Lambda}^*(X, J) \rightarrow X^\Lambda$ by

$$\text{ev}([u; z_1, \dots, z_\Lambda]) := (u(z_1), \dots, u(z_\Lambda)).$$

Remark 2.40. Given two maps $f: X \rightarrow Z$ and $g: Y \rightarrow Z$, the fiber product is

$$X \times_f \times_g Y := (f \times g)^{-1}(\Delta)$$

with $\Delta \subset Z \times Z$ denoting the diagonal. If X, Y, Z are smooth manifolds and f and g are transverse smooth maps, then $X \times_f \times_g Y$ is a submanifold of $X \times Y$ of dimension $\dim(X) + \dim(Y) - \dim(Z)$.

Let $(f_\lambda: V_\lambda \rightarrow X)_{\lambda=1}^\Lambda$ be a Λ -tuple of pseudo-cycles in general position such that

$$\text{codim}(f_\lambda) := \dim X - \dim V_\lambda$$

is even and positive for every λ . The following discussion assumes some familiarity with the notions of pseudo-cycle, pseudo-cycle cobordism, and pseudo-cycle transversality. In particular, we make use of the following facts, which are discussed in Section B:

- (a) For every $\lambda \in \{1, \dots, \Lambda\}$, there is a manifold V_λ^∂ of dimension $\dim(V_\lambda) - 2$ and a smooth map $f_\lambda^\partial: V_\lambda^\partial \rightarrow X$ whose image contains the pseudo-cycle boundary $\text{bd}(f_\lambda)$.
- (b) A smooth map $g: M \rightarrow X$ is said to be transverse to the pseudo-cycle f_λ if it is transverse to both f_λ and f_λ^∂ in the usual sense.
- (c) For every $I \subset \{1, \dots, \Lambda\}$ the product $\prod_{\lambda \in I} f_\lambda$ is a pseudo-cycle and f_λ^∂ induces in a natural way a map from a smooth manifold whose image contains $\text{bd}(\prod_{\lambda \in I} f_\lambda)$.

In the following, $f_\lambda^\bullet: V_\lambda^\bullet \rightarrow X$ stands for either $f_\lambda: V_\lambda \rightarrow X$ or $f_\lambda^\partial: V_\lambda^\partial \rightarrow X$.

Definition 2.41. Given $A \in H_2(X, \mathbf{Z})$, $g \in \mathbf{N}_0$, and $J \in \mathcal{J}(X, \omega)$, set

$$\mathcal{M}_{A,g}^\star(X, J; f_1^\bullet, \dots, f_\Lambda^\bullet) := \mathcal{M}_{A,g,\Lambda}^\star(X, J)_{\text{ev} \times f_1^\bullet \times \dots \times f_\Lambda^\bullet} V_1^\bullet \times \dots \times V_\Lambda^\bullet.$$

The *expected dimension* of $\mathcal{M}_{A,g}^\star(X, J; f_1, \dots, f_\Lambda)$ is defined to be

$$\begin{aligned} \text{vdim } \mathcal{M}_{A,g}^\star(X, J; f_1, \dots, f_\Lambda) &:= 2\langle c_1(X, \omega), A \rangle \\ &\quad + (n - 3)(2 - 2g) + \sum_{\lambda=1}^\Lambda (2 - \text{codim}(f_\lambda)). \end{aligned}$$

The following are analogues of Definition 2.34 and Definition 2.35 in the setting of J -holomorphic maps with constraints.

Definition 2.42. Denote by

$$\mathcal{J}_{\text{emb}}(X, \omega; f_1, \dots, f_\Lambda) \subset \mathcal{J}(X, \omega)$$

the subset of those almost complex structures J compatible with ω for which the following conditions hold for every $A, A_1, A_2 \in H_2(X, \mathbf{Z})$, $g, g_1, g_2 \in \mathbf{N}_0$, and $I, I_1, I_2 \subset \{1, \dots, \Lambda\}$ with $I_1 \cap I_2 = \emptyset$:

- (1) if $\text{vdim } \mathcal{M}_{A,g}^\star(X, J; (f_\lambda^\bullet)_{\lambda \in I}) < 0$, then $\mathcal{M}_{A,g}^\star(X, J; (f_\lambda^\bullet)_{\lambda \in I}) = \emptyset$;
- (2) if $\text{vdim } \mathcal{M}_{A,g}^\star(X, J; (f_\lambda^\bullet)_{\lambda \in I}) < 2n - 4$, then every J -holomorphic map underlying an element of $\mathcal{M}_{A,g}^\star(X, J; (f_\lambda^\bullet)_{\lambda \in I})$ is an embedding; and
- (3) if $\text{vdim } \mathcal{M}_{A_1,g_1}^\star(X, J; (f_\lambda^\bullet)_{\lambda \in I_1}) + \text{vdim } \mathcal{M}_{A_2,g_2}^\star(X, J; (f_\lambda^\bullet)_{\lambda \in I_2}) < 2n - 4$, then every pair of every J -holomorphic maps underlying elements of

$$\mathcal{M}_{A_1,g_1}^\star(X, J; (f_\lambda^\bullet)_{\lambda \in I_1}) \quad \text{and} \quad \mathcal{M}_{A_2,g_2}^\star(X, J; (f_\lambda^\bullet)_{\lambda \in I_2})$$

either have disjoint images or are related by a reparametrization.

Denote by

$$\mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda)$$

the subset of those elements of $\mathcal{J}_{\text{emb}}(X, \omega; f_1, \dots, f_\Lambda)$ for which, moreover:

- (4) every simple J -holomorphic map is unobstructed, and
- (5) for every $A \in H_2(X, \mathbf{Z})$, $g \in \mathbf{N}$, and $I \subset \{1, \dots, \Lambda\}$, the pseudo-cycle $\prod_{\lambda \in I} f_\lambda$ is transverse to

$$\text{ev}: \mathcal{M}_{A,g,|I|}^*(X, J) \rightarrow X^{|I|}$$

in the sense of Definition B.3.

Definition 2.43. Given $J_0, J_1 \in \mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda)$, denote by

$$\mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda; J_0, J_1)$$

the space of smooth paths $(J_t)_{t \in [0,1]}$ in $\mathcal{J}(X, \omega)$ from J_0 and J_1 such that for every $t \in [0, 1]$:

- (1) $J_t \in \mathcal{J}_{\text{emb}}(X, \omega; f_1, \dots, f_\Lambda)$,
- (2) if $u: (\Sigma, j) \rightarrow (X, J_t)$ is a simple J_t -holomorphic map, then either
 - (a) $\text{coker } d_{u,j} \bar{\partial}_{J_t} = \{0\}$, or
 - (b) $\dim \text{coker } d_{u,j} \bar{\partial}_{J_t} = 1$ and the map $\ker d_{u,j} \bar{\partial}_{J_t} \rightarrow \text{coker } d_{u,j} \bar{\partial}_{J_t}$ defined by

$$\xi \mapsto \text{pr} \left(\frac{d}{ds} \Big|_{s=t} d_{u,j} \bar{\partial}_{J_s} \xi \right),$$

with $\text{pr}: \Omega^{0,1}(\Sigma, u^*TX) \rightarrow \text{coker } d_{u,j} \bar{\partial}_{J_t}$ denoting the canonical projection, is surjective; in particular, for every $A \in H_2(X, \mathbf{Z})$, $g \in \mathbf{N}$, and $k \in \mathbf{N}$ the moduli space

$$\mathcal{M}_{A,g,k}^*(X, (J_t)_{t \in [0,1]}) := \bigsqcup_{t \in [0,1]} \mathcal{M}_{A,g,k}^*(X, J_t)$$

is an oriented smooth manifold with boundary

$$\mathcal{M}_{A,g,k}^*(X, J_1) \amalg -\mathcal{M}_{A,g,k}^*(X, J_0),$$

and

- (3) for every $A \in H_2(X, \mathbf{Z})$, $g \in \mathbf{N}$, and $I \subset \{1, \dots, \Lambda\}$ the pseudo-cycle $\prod_{\lambda \in I} f_\lambda$ is transverse to the evaluation map

$$\text{ev}: \mathcal{M}_{A,g,|I|}^*(X, A; (J_t)_{t \in [0,1]}) \rightarrow X^{|I|}$$

in the sense of Definition B.3.

The next two results are analogues of Proposition 2.36 and Proposition 2.37.

Proposition 2.44. *Let $A \in H_2(X, \mathbf{Z})$ and $g \in \mathbf{N}_0$.*

(1) *For every $J \in \mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda)$, the moduli space*

$$\mathcal{M}_{A,g}^*(X, J; f_1^\bullet, \dots, f_\Lambda^\bullet)$$

is an oriented smooth manifold of dimension

$$\text{vdim } \mathcal{M}_{A,g}^*(X, J; f_1^\bullet, \dots, f_\Lambda^\bullet).$$

(2) *For every pair*

$$J_0, J_1 \in \mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda), \quad (J_t)_{t \in [0,1]} \in \mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda; J_0, J_1),$$

the moduli space

$$\mathcal{M}_{A,g}^*(X, (J_t)_{t \in [0,1]}; f_1^\bullet, \dots, f_\Lambda^\bullet) := \coprod_{t \in [0,1]} \mathcal{M}_{A,g}^*(X, J_t; f_1^\bullet, \dots, f_\Lambda^\bullet)$$

is an oriented smooth manifold with boundary

$$\mathcal{M}_{A,g}^*(X, J; f_1^\bullet, \dots, f_\Lambda^\bullet) \amalg -\mathcal{M}_{A,g}^*(X, J; f_1^\bullet, \dots, f_\Lambda^\bullet).$$

Proposition 2.45. *The following hold:*

(1) $\mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda) \subset \mathcal{J}(X, \omega)$ *is residual.*

(2) *For every pair $J_0, J_1 \in \mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda)$,*

$$\mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda; J_0, J_1) \subset \mathcal{J}(X, \omega; J_0, J_1)$$

is residual.

Proof. We will prove the first part; the proof of the second part is similar. It follows from Proposition A.2, proved in Section A, that the set of $J \in \mathcal{J}(X, \omega)$ satisfying conditions (4) and (5) from Definition 2.42 is residual. Note that condition (4) implies condition (1). To prove that condition (2) is also satisfied by a generic J , consider the evaluation map

$$\text{ev}: \mathcal{M}_{A,g,2}^*(X, J; (f_\lambda^\bullet)_{\lambda \in I}) \rightarrow X^2.$$

If ev is transverse to the diagonal $X = \Delta \subset X^2$, then $\text{ev}^{-1}(\Delta)$ is a submanifold of codimension

$$\dim \mathcal{M}_{A,g,2}^*(X, J; (f_\lambda^\bullet)_{\lambda \in I}) - 2n = \dim \mathcal{M}_{A,g}^*(X, J; (f_\lambda^\bullet)_{\lambda \in I}) - (2n - 4).$$

Therefore, if $\dim \mathcal{M}_{A,g}^*(X, J; (f_\lambda^\bullet)_{\lambda \in I}) < 2n - 4$, then $\text{ev}^{-1}(\Delta)$ is empty and two distinct maps in $\mathcal{M}_{A,g}^*(X, J; (f_\lambda^\bullet)_{\lambda \in I})$ have disjoint images. By Proposition A.2, the

set of J for which the map ev is transverse to the diagonal $X \hookrightarrow X^2$ is residual. This shows that the set of J satisfying condition (2) from Definition 2.42 is residual. In the same way, we conclude that the set of J satisfying condition (3) is residual. ■

The following will be important for relating moduli spaces defined using cobordant pseudo-cycles. Let $F: W \rightarrow X$ be a cobordism between two pseudo-cycles f_1^0 and f_1^1 in X , and let $F^\partial: W^\partial \rightarrow X$ be such that $\text{bd}(F)$ is contained in the image of F^∂ ; see Definition B.1 for the notation and the definition of a pseudo-cycle cobordism. In what follows, F^\bullet denotes either F or F^∂ . Let f_2, \dots, f_Λ be pseudo-cycles in X such that F, f_2, \dots, f_Λ are in general position, as in Definition B.4.

Given $J \in \mathcal{J}(X, \omega)$ and a subset $I \subset \{2, \dots, \Lambda\}$, set

$$\mathcal{M}_{A,g}^*(X, J; F^\bullet, (f_\lambda^\bullet)_{\lambda \in I}) := \mathcal{M}_{A,g,|I|+1}^*(X, J)_{\text{ev} \times F^\bullet \times \prod_{\lambda \in I} f_\lambda^\bullet} W^\bullet \times \prod_{\lambda \in I} V_\lambda^\bullet.$$

Definition 2.46. Let

$$\mathcal{J}_{\text{emb}}^*(X, \omega; F, f_2, \dots, f_\Lambda) \subset \mathcal{J}_{\text{emb}}^*(X, \omega; f_1^0, f_2, \dots, f_\Lambda) \cap \mathcal{J}_{\text{emb}}^*(X, \omega; f_1^1, f_2, \dots, f_\Lambda)$$

be the subset of those J for which the following conditions hold for every $A, A_1, A_2 \in H_2(X, \mathbf{Z})$, $g, g_1, g_2 \in \mathbf{N}_0$, and $I, I_1, I_2 \subset \{2, \dots, \Lambda\}$ with $I_1 \cap I_2 = \emptyset$:

- (1) if $\text{vdim} \mathcal{M}_{A,g}^*(X, J; F^\bullet, (f_\lambda^\bullet)_{\lambda \in I}) < 2n - 4$, then every J -holomorphic map underlying an element of $\mathcal{M}_{A,g}^*(X, J; F^\bullet, (f_\lambda^\bullet)_{\lambda \in I})$ is an embedding;
- (2) if $\text{vdim} \mathcal{M}_{A_1,g_1}^*(X, J; F^\bullet, (f_\lambda^\bullet)_{\lambda \in I_1}) + \text{vdim} \mathcal{M}_{A_2,g_2}^*(X, J; (f_\lambda^\bullet)_{\lambda \in I_2}) < 2n - 4$, then every pair of J -holomorphic maps underlying elements of

$$\mathcal{M}_{A_1,g_1}^*(X, J; F^\bullet, (f_\lambda^\bullet)_{\lambda \in I_1}) \quad \text{and} \quad \mathcal{M}_{A_2,g_2}^*(X, J; F^\bullet, (f_\lambda^\bullet)_{\lambda \in I_2})$$

either have disjoint images or are related by a reparametrization; and

- (3) for every $A \in H_2(X, \mathbf{Z})$, $g \in \mathbf{N}$, and $I \subset \{2, \dots, \Lambda\}$, the pseudo-cycle $F \times \prod_{\lambda \in I} f_\lambda$ is transverse as pseudo-cycle with boundary to

$$\text{ev}: \mathcal{M}_{A,g,|I|+1}^*(X, J) \rightarrow X^{|I|+1}$$

in the sense of Definition B.3.

It follows from this definition that for every $J \in \mathcal{J}_{\text{emb}}^*(X, \omega; F, f_2, \dots, f_\Lambda)$, the pseudo-cycle $F \times f_2 \times \dots \times f_\Lambda$ is transverse as a pseudo-cycle cobordism to

$$\text{ev}: \mathcal{M}_{A,g,\Lambda}^*(X, J) \rightarrow X^\Lambda.$$

In this case, $\mathcal{M}_{A,g}^*(X, J; F, f_2, \dots, f_\Lambda)$ is an oriented cobordism from

$$\mathcal{M}_{A,g}^*(X, J; f_1^0, f_2, \dots, f_\Lambda)$$

to $\mathcal{M}_{A,g}^*(X, J; f_1^1, f_2, \dots, f_\Lambda)$.

Proposition 2.47. $\mathcal{J}_{\text{emb}}^*(X, \omega; F, f_2, \dots, f_\Lambda)$ is residual in $\mathcal{J}(X, \omega)$.

The proof is almost identical to that of Proposition 2.45.

3. Gromov compactness

3.1. Deformations of nodal Riemann surfaces

Definition 3.1. Let \mathcal{X} and A be complex manifolds and let $\pi: \mathcal{X} \rightarrow A$ be a holomorphic map. Set $n := \dim_{\mathbb{C}} A$ and suppose that $\dim_{\mathbb{C}} \mathcal{X} = n + 1$. A critical point $x \in \mathcal{X}$ of π is called *nodal* if there are holomorphic coordinates at x and holomorphic coordinates at $\pi(x)$ with respect to which

$$\pi(z, w, t_2, \dots, t_n) = (zw, t_2, \dots, t_n).$$

A *nodal family* is a surjective, proper, holomorphic map $\pi: \mathcal{X} \rightarrow A$ between complex manifolds of dimension $\dim_{\mathbb{C}} \mathcal{X} = \dim_{\mathbb{C}} A + 1$ such that every critical point of π is nodal. The *fiber* over $a \in A$ is the nodal Riemann surface (Σ, j, ν) associated with the nodal curve $\pi^{-1}(a)$. Henceforth, we engage in the abuse of notation of identifying $\pi^{-1}(a)$ with (Σ, j, ν) .

Definition 3.2. Let (Σ, j, ν) be a nodal Riemann surface. A *deformation* of (Σ, j, ν) is a nodal family $\pi: \mathcal{X} \rightarrow A$, together with a base-point $\star \in A$, and a nodal, biholomorphic map $\iota: (\Sigma, j, \nu) \rightarrow \pi^{-1}(\star)$.

Definition 3.3. Let (Σ, j, ν) be a nodal Riemann surface, and let $(\pi: \mathcal{X} \rightarrow A, \star, \iota)$ and $(\rho: \mathcal{Y} \rightarrow B, \dagger, \kappa)$ be two deformations of (Σ, j, ν) . A pair of holomorphic maps $\Phi: \mathcal{X} \rightarrow \mathcal{Y}$ and $\phi: A \rightarrow B$ forms a *morphism* $(\Phi, \phi): (\rho, \star, \iota) \rightarrow (\mathcal{Y}, \dagger, \kappa)$ of deformations if

$$\phi(\star) = \dagger, \quad \rho \circ \Phi = \phi \circ \pi, \quad \Phi \circ \iota = \kappa$$

and for every $a \in A$ the restriction $\Phi: \pi^{-1}(a) \rightarrow \rho^{-1}(\phi(a))$ induces a nodal, biholomorphic map.

Definition 3.4. A deformation $(\rho: \mathcal{Y} \rightarrow B, \dagger, \kappa)$ of (Σ, j, ν) is *(uni)versal* if for every deformation $(\pi: \mathcal{X} \rightarrow A, \star, \iota)$ of (Σ, j, ν) there exists an open neighborhood U of $\star \in A$ and a (unique) morphism of deformations $(\pi: \pi^{-1}(U) \rightarrow U, \star, \iota) \rightarrow (\rho, \dagger, \kappa)$.

A nodal Riemann surface (Σ, j, ν) admits a universal deformation if and only if it is stable (see [5], [1, Chapter XI, Theorem 4.3], and [33, Theorem A]). However, every nodal Riemann surface (Σ, j, ν) admits a versal deformation. This will be discussed in detail in Section 4.

Definition 3.5. Let $(\pi: \mathcal{X} \rightarrow A, \star, \iota)$ be a deformation of a nodal Riemann surface (Σ, j, ν) . Denote by S the nodal set of ν . A *framing* of (π, \star, ι) is a smooth embedding $\Psi: (\Sigma \setminus S) \times A \rightarrow \mathcal{X}$ such that

$$\pi \circ \Psi = \text{pr}_A \quad \text{and} \quad \Psi(\cdot, \star) = \iota.$$

3.2. The Gromov topology

Let X be a manifold and denote by $\mathcal{H}(X)$ the set of almost Hermitian structures (J, h) on X equipped with the C^∞ topology. The following defines a topology on

$$\bar{\mathcal{M}}_{A,g}(X) := \coprod_{(J,h) \in \mathcal{H}(X)} \bar{\mathcal{M}}_{A,g}(X, J).$$

Definition 3.6. Let (X, J, h) be an almost Hermitian manifold. Let (Σ, j, ν) be a closed, nodal Riemann surface. The *energy* of a nodal map $u: (\Sigma, \nu) \rightarrow X$ is

$$E(u) := \frac{1}{2} \int_{\Sigma} |du|^2 \text{ vol.}$$

Implicit in this definition is a choice of Riemannian metric in the conformal class determined by j . The right-hand side, however, is independent of this choice.

Definition 3.7. Let $(J_0, h_0) \in \mathcal{H}(X)$. Let $[u_0: (\Sigma_0, j_0, \nu_0) \rightarrow (X, J_0)] \in \bar{\mathcal{M}}_{A,g}(X, J_0)$, let $(\pi: \mathcal{X} \rightarrow A, \star, \iota)$ be a versal deformation of (Σ_0, j_0, ν_0) , let Ψ be a framing of (π, \star, ι) , and let $\varepsilon > 0$. Let $U_0 \subset C^\infty(\Sigma_0 \setminus S, X)$ be an open neighborhood of $u_\infty|_{\Sigma_0 \setminus S}$ in the C_{loc}^∞ topology, and let $U_{\mathcal{H}}$ be an open neighborhood of (J_0, h_0) in $\mathcal{H}(X)$. Define

$$\mathcal{U}(u_0, \varepsilon, U_0, U_{\mathcal{H}}) \subset \bar{\mathcal{M}}_{A,g}(X)$$

to be the subset of the equivalence classes of nodal J -holomorphic maps $u: (\Sigma, j, \nu) \rightarrow (X, J)$ satisfying the following:

- (1) $(J, h) \in U_{\mathcal{H}}$,
- (2) $|E(u) - E(u_0)| < \varepsilon$,
- (3) $(\Sigma, j, \nu) = \pi^{-1}(a)$ for some $a \in A$, and
- (4) $\tilde{u} := u \circ \Psi(\cdot, a) \in U_0$,

The *Gromov topology* on $\bar{\mathcal{M}}_{A,g}(X)$ is the coarsest topology with respect to which every subset of the form $\mathcal{U}(u_0, \varepsilon, U_0, U_{\mathcal{H}})$ is open.

In practice, it is more convenient to use the notion of *Gromov convergence* defined on the level of nodal maps.

Definition 3.8. Let (X, J_∞, h_∞) be an almost Hermitian manifold and let $(J_k, h_k)_{k \in \mathbf{N}}$ be a sequence of almost Hermitian structures on X converging to (J_∞, h_∞) in the C^∞ topology. For every $k \in \mathbf{N} \cup \{\infty\}$ let $u_k: (\Sigma_k, j_k, \nu_k) \rightarrow (X, J_k)$ be a nodal J_k -holomorphic map. Denote by S the nodal set of $(\Sigma_\infty, \nu_\infty)$. The sequence $(u_k, j_k)_{k \in \mathbf{N}}$ Gromov converges to (u_∞, j_∞) if

- (1) $\lim_{k \rightarrow \infty} E(u_k) = E(u_\infty)$, and
- (2) there are:
 - (a) a deformation $(\pi: \mathcal{X} \rightarrow A, a_\infty, \iota_\infty)$ of $(\Sigma_\infty, j_\infty, \nu_\infty)$ together with a framing Ψ ,
 - (b) a sequence $(a_k)_{k \in \mathbf{N}}$ in A converging to a_∞ , and
 - (c) a nodal, biholomorphic map $\iota_k: (\Sigma_k, j_k, \nu_k) \rightarrow \pi^{-1}(a_k)$ for every sufficiently large $k \in \mathbf{N}$,

such that the sequence of maps

$$\tilde{u}_k := u_k \circ \iota_k^{-1} \circ \Psi(\cdot, a_k) \circ \iota_\infty: \Sigma_\infty \setminus S \rightarrow X$$

converges to $u_\infty|_{\Sigma_\infty \setminus S}$ in the C_{loc}^∞ topology.

Remark 3.9. If (π, \star, ι) is a versal deformation of $(\Sigma_\infty, j_\infty, \nu_\infty)$ and Ψ is a framing of this deformation, then for every sequence $(u_k, j_k)_{k \in \mathbf{N}}$ which Gromov converges to (u_∞, j_∞) the deformation in Definition 3.8 can be assumed to be (π, \star, ι) and the framing can be assumed to be Ψ . This is an immediate consequence of the definition of a versal deformation.

Theorem 3.10 (Gromov [13]; see also [17, 32, 36] and [24, Chapters 4 and 5]). *Let (X, J_∞, h_∞) be a closed almost Hermitian manifold and let $(J_k, h_k)_{k \in \mathbf{N}}$ be a sequence of almost Hermitian structures on X converging to (J_∞, h_∞) in the C^∞ topology. For every $k \in \mathbf{N}$, let*

$$u_k: (\Sigma_k, j_k, \nu_k) \rightarrow (X, J_k)$$

be a stable nodal J -holomorphic map. Denote by $\#\pi_0(\Sigma_k)$ the number of connected components of Σ_k . If

$$\limsup_{k \rightarrow \infty} \#\pi_0(\Sigma_k) < \infty, \quad \limsup_{k \rightarrow \infty} p_a(\Sigma_k, \nu_k) < \infty, \quad \text{and} \quad \limsup_{k \rightarrow \infty} E(u_k) < \infty,$$

then there exists a stable nodal J_∞ -holomorphic map $u_\infty: (\Sigma_\infty, j_\infty, \nu_\infty) \rightarrow (X, J_\infty)$ and a subsequence of $(u_k, j_k)_{k \in \mathbf{N}}$, which Gromov converges to (u_∞, j_∞) . The limit $(u_\infty, j_\infty, \mu_\infty)$ is unique up to automorphism.

Remark 3.11. The Gromov topology on $\bar{\mathcal{M}}_{A,g}(X)$ is metrizable, which can be seen as follows. Theorem 3.10 implies that it is Hausdorff and the projection map

$$\bar{\mathcal{M}}_{A,g}(X) \rightarrow \mathcal{H}(X) \times \mathbf{R}$$

is proper and closed. This implies, in particular, that $\bar{\mathcal{M}}_{A,g}(X)$ is a regular topological space. (In general, if A is a Hausdorff space, B is a regular space, and $f: A \rightarrow B$ is a proper, closed map, then A is a regular space.) Urysohn’s metrization theorem says that a second countable, Hausdorff, regular space is metrizable.

Henceforth, let (X, ω) be a symplectic manifold. The set $\mathcal{J}(X, \omega)$ of almost complex structures compatible with ω injects into $\mathcal{H}(X)$.

Proposition 3.12 (Gromov [13]; see also [24, Lemma 2.2.1]). *Let (X, ω) be a symplectic manifold and $J \in \mathcal{J}(X, \omega)$. Let (Σ, ν, j) be a closed, nodal Riemann surface. For every nodal map $u: (\Sigma, \nu) \rightarrow X$,*

$$E(u) \geq \langle u^*[\omega], [\Sigma] \rangle,$$

and the equality holds if and only if u is J -holomorphic. Here E is to be understood with respect to the Riemannian metric $h = \omega(\cdot, J\cdot)$ on X .

Set

$$\bar{\mathcal{M}}_{A,g}(X, \omega) := \coprod_{J \in \mathcal{J}(X, \omega)} \bar{\mathcal{M}}_{A,g}(X, J).$$

By the above energy identity, in the symplectic context, Theorem 3.10 is equivalent to the map

$$\pi: \bar{\mathcal{M}}_{A,g}(X, \omega) \rightarrow \mathcal{J}(X, \omega)$$

being proper.

3.3. Behavior near the vanishing cycles

The results of this subsection will be important for proving the surjectivity of the gluing construction in Section 5.6. Assume the situation of Definition 3.8. By condition Definition 1 for every $\delta > 0$ there are $K \in \mathbf{N}_0$ and $r > 0$ such that for every $k \geq K$,

$$E(u_k|_{N_k^r}) \leq \delta$$

with

$$N_k^r := \Sigma_k \setminus \{\Psi(z, a_k) : z \in \Sigma_0 \text{ with } d(z, S) \geq r\}. \tag{3.13}$$

The subset N_k^r can be partitioned into regions $N_{k,n}^r$ corresponding to the nodes $n \in S$. If n is not smoothed out in Σ_k , then the corresponding region is biholomorphic to

$$B_1(0) \amalg B_1(0)$$

with ν_k identifying the origins. If n is smoothed out in Σ_k , then the corresponding region is biholomorphic to

$$S^1 \times (-L_k, L_k)$$

with $\lim_{k \rightarrow \infty} L_k = \infty$.

The behavior of J -holomorphic maps from such domains and with small energy can be understood through the following two results.

Lemma 3.14 ([24, Lemma 4.3.1]). *Let (X, J, h) be an almost Hermitian manifold. There is a constant $\delta = \delta(X, J, h) > 0$, depending continuously on (J, h) , such that for every $r > 0$ the following holds. If $u: (B_{2r}(0), i) \rightarrow (X, J)$ is a J -holomorphic map with*

$$E(u) \leq \delta,$$

then

$$\|du\|_{L^\infty(B_r(0))} \leq cr^{-1} E(u)^{1/2}.$$

Lemma 3.15 ([24, Lemma 4.7.3]). *Let (X, J, h) be an almost Hermitian manifold. For every $\mu \in (0, 1)$ there are constants: $\delta = \delta(X, J, h, \mu) > 0$, depending continuously on (J, h) , and $c = c(\mu) > 0$ such that for every $L > 0$ the following holds. If $u: (S^1 \times (-L, L), j_{\text{cyl}}) \rightarrow (X, J)$ is a J -holomorphic map with*

$$E(u) \leq \delta,$$

then for every $\ell \in (0, L)$,

$$E(u|_{S^1 \times (L+\ell, L-\ell)}) \leq ce^{-2\mu(L-\ell)} E(u),$$

and for every $\theta \in S^1$ and $\ell \in [-L + 1, L - 1]$,

$$|du|(\theta, \ell) \leq ce^{-\mu(L-|\ell|)} E(u)^{1/2}.$$

Proof. The first assertion is [24, Lemma 4.7.3]. The second assertion follows from the first by Lemma 3.14. ■

The following is an important consequence of the previous two lemmas.

Proposition 3.16. *Let $(u_k: (\Sigma_k, j_k, \nu_k) \rightarrow (X, J_k))_{k \in \mathbb{N}}$ be a sequence of nodal pseudo-holomorphic maps which Gromov converges to*

$$u_\infty: (\Sigma_\infty, j_\infty, \nu_\infty) \rightarrow (X, J_\infty).$$

Denote by S the nodal set of $(\Sigma_\infty, \nu_\infty)$ and let N_k^r be as in (3.13). For every $\delta > 0$, there are $r > 0$ and $K \in \mathbb{N}$ such that for every $k \geq K$ and $n \in S$,

$$u_k(N_{k,n}^r) \subset B_\delta(u_\infty(n));$$

in particular, provided δ is sufficiently small,

$$(u_k)_*[\Sigma_k] = (u_\infty)_*[\Sigma_\infty].$$

4. Versal deformations of nodal Riemann surfaces

The purpose of this section is to construct a versal deformation of a nodal Riemann surface in a rather explicit manner.

4.1. Deformations of nodal curves

Let us briefly review parts of the deformation theory of nodal curves in the complex analytic category. For further details and proofs we refer the reader to [1, Chapter XI, Section 3]. A thorough discussion of deformation theory in the algebraic category can be found in [14].

Definition 4.1. Let C be a nodal curve. A *deformation* of C consists of

- (1) a proper flat¹ morphism $\pi: \mathcal{X} \rightarrow A$ between analytic spaces such that every fiber of π is a nodal curve,
- (2) a base-point $\star \in A$, and
- (3) an isomorphism $\iota: C \rightarrow \pi^{-1}(\star)$.

Proposition 4.2. *Every nodal family $\pi: \mathcal{X} \rightarrow A$ is flat. In particular, a deformation of a nodal Riemann surface (Σ, j, ν) is also a deformation of the associated nodal curve C .*

Definition 4.3. Let C be a nodal Riemann surface and let $(\pi: \mathcal{X} \rightarrow A, \star, \iota)$ and $(\rho: \mathcal{Y} \rightarrow B, \dagger, \kappa)$ be two deformations of C . A pair of analytic maps $\Phi: \mathcal{X} \rightarrow \mathcal{Y}$ and $\phi: A \rightarrow B$ forms a *morphism* $(\Phi, \phi): (\rho, \star, \iota) \rightarrow (\mathcal{Y}, \dagger, \kappa)$ of deformations if

$$\phi(\star) = \dagger, \quad \rho \circ \Phi = \phi \circ \pi, \quad \Phi \circ \iota = \kappa,$$

and for every $a \in A$ the restriction $\Phi: \pi^{-1}(a) \rightarrow \rho^{-1}(\phi(a))$ induces an analytic isomorphism.

Definition 4.4. A deformation $(\rho: \mathcal{Y} \rightarrow B, \dagger, \kappa)$ of C is *(uni)versal* if for every deformation $(\pi: \mathcal{X} \rightarrow A, \star, \iota)$ of (Σ, j, ν) , there exists an open neighborhood U of $\star \in A$ and a (unique) morphism of deformations $(\pi: \pi^{-1}(U) \rightarrow A, \star, \iota) \rightarrow (\rho, \dagger, \kappa)$.

Definition 4.5. Denote by $\mathbf{C}[\varepsilon]/\varepsilon^2$ the ring of dual numbers and set

$$D := \text{Spec}(\mathbf{C}[\varepsilon]/\varepsilon^2).$$

A *first order deformation* is a deformation over D .

¹A morphism $f: A \rightarrow B$ between two analytic spaces is *flat* if it makes the stalk $\mathcal{O}_{A,a}$ into a flat $\mathcal{O}_{B,f(a)}$ -module for every a , that is, tensoring by $\mathcal{O}_{A,a}$ preserves short exact sequences of $\mathcal{O}_{B,f(a)}$ -modules.

Let C be a nodal curve. Every first order deformation $(\pi: \mathcal{X} \rightarrow D, 0, \iota)$ of C induces a short exact sequence

$$0 \rightarrow \mathcal{O}_C \cong \pi^* \Omega_D^1 \rightarrow \Omega_{\mathcal{X}}^1 \otimes \mathcal{O}_C \xrightarrow{\iota^*} \Omega_C^1 \rightarrow 0.$$

The map $\pi^* \Omega_D^1 \rightarrow \Omega_{\mathcal{X}}^1 \otimes \mathcal{O}_C$ is given by pulling-back forms from D to \mathcal{X} and $\Omega_{\mathcal{X}}^1 \otimes \mathcal{O}_C \rightarrow \Omega_C^1$ is given by restricting forms on \mathcal{X} to C . The extension class $\delta \in \text{Ext}^1(\Omega_C^1, \mathcal{O}_C)$ of this sequence depends on the first order deformation only up to isomorphism of deformations. Indeed, two first order deformation of C are isomorphic if and only if they yield the same extension class δ .

Definition 4.6. Let C be a nodal curve and let $(\pi: \mathcal{X} \rightarrow A, \star, \iota)$ be a deformation of C . Every $v \in T_{\star} A$ corresponds to an analytic map $\phi: D \rightarrow A$ mapping 0 to \star . The pullback of (π, \star, ι) via ϕ is a first order deformation. Denote by $\delta(v) \in \text{Ext}^1(\Omega_C^1, \mathcal{O}_C)$ the corresponding extension class. The map $\delta: T_{\star} A \rightarrow \text{Ext}^1(\Omega_C^1, \mathcal{O}_C)$ thus defined is called the *Kodaira–Spencer map*.

It is instructive to analyze $\text{Ext}^1(\Omega_C^1, \mathcal{O}_C)$ more closely. The local-to-global Ext spectral sequence yields a short exact sequence

$$0 \rightarrow H^1(C, \mathcal{H}om(\Omega_C^1, \mathcal{O}_C)) \rightarrow \text{Ext}^1(\Omega_C^1, \mathcal{O}_C) \rightarrow H^0(C, \mathcal{E}xt^1(\Omega_C^1, \mathcal{O}_C)) \rightarrow 0.$$

This can be interpreted in terms of the normalization $\pi: \tilde{C} \rightarrow C$ as follows. Denote by S the set of nodes of C and set $\tilde{S} := \pi^{-1}(S)$. It can be shown that

$$\mathcal{H}om(\Omega_C^1, \mathcal{O}_C) = \pi_* \mathcal{T}_{\tilde{C}}(-\tilde{S});$$

hence,

$$H^1(C, \mathcal{H}om(\Omega_C^1, \mathcal{O}_C)) = H^1(\tilde{C}, \mathcal{T}_{\tilde{C}}(-\tilde{S})).$$

The space $H^1(\tilde{C}, \mathcal{T}_{\tilde{C}}(-\tilde{S}))$ parametrizes the deformations of the marked curve (\tilde{C}, \tilde{S}) ; that is, deformations of \tilde{C} which fix \tilde{S} point-wise. The sheaf $\mathcal{E}xt^1(\Omega_C^1, \mathcal{O}_C)$ is supported on the nodes of C :

$$\mathcal{E}xt^1(\Omega_C^1, \mathcal{O}_C) = \bigoplus_{n \in S} \text{Ext}^1(\Omega_{C,n}^1, \mathcal{O}_{C,n}).$$

For every $n \in \tilde{S}$ and $\{n_1, n_2\} = \pi^{-1}(n)$,

$$\text{Ext}^1(\Omega_{C,n}^1, \mathcal{O}_{C,n}) = T_{n_1} \tilde{C} \otimes T_{n_2} \tilde{C}.$$

If we consider the deformation $\{zw = \varepsilon\}$ of the node $\{zw = 0\}$, then the space $\text{Ext}^1(\Omega_{C,n}^1, \mathcal{O}_{C,n})$ can be seen to parametrize smoothings of the node n . The above discussion show that to first order all deformations of C arise from smoothing nodes and deforming its normalization while fixing the points mapping to the nodes. In the following we construct a deformation of C which induces all of these deformations to first order.

4.2. Smoothing nodal Riemann surfaces

Let (Σ_0, j_0, ν_0) be a closed, nodal Riemann surface with nodal set S . Let g_0 be a Riemannian metric on Σ_0 in the conformal class determined by j_0 and such that there is a constant $R_0 > 0$ such that for every $n \in S$ the restriction of g_0 to $B_{4R_0}(n)$ is flat and for every $n_1, n_2 \in S$ the balls $B_{4R_0}(n_1)$ and $B_{4R_0}(n_2)$ are disjoint. For every $n \in S$, define the holomorphic charts $\phi_n: B_{4R_0}(n) \subset T_n \Sigma_0 \rightarrow \Sigma_0$ by

$$\phi_n(v) := \exp_n(v)$$

and define $r_n: \Sigma_0 \rightarrow [0, \infty)$ by

$$r_n(z) := \max\{d(n, z), 4R_0\}.$$

Given a pair of complex vector spaces V and W , denote by $\sigma: V \otimes W \rightarrow W \otimes V$ the isomorphism defined by $\sigma(v \otimes w) := w \otimes v$.

Definition 4.7. A *smoothing parameter* for (Σ_0, j_0, ν_0) is an element

$$\tau = (\tau_n)_{n \in S} \in \prod_{n \in S} T_n \Sigma_0 \otimes_{\mathbb{C}} T_{\nu(n)} \Sigma_0$$

such that for every $n \in S$, we have $\tau_{\nu(n)} = \sigma(\tau_n)$ and $|\tau_n| < R_0^2$. Given a smoothing parameter τ , for every $n \in S$, set

$$\varepsilon_n := |\tau_n| \quad \text{and} \quad \hat{\tau}_n := \tau_n / |\tau_n| \quad \text{if } \tau_n \neq 0;$$

furthermore, set

$$\varepsilon := \max\{\varepsilon_n : n \in S\}.$$

Henceforth, let $\tau = (\tau_n)_{n \in S}$ be a smoothing parameter for (Σ_0, j_0, ν_0) .

Definition 4.8. Set

$$A_\tau := \{w \in \Sigma_0 : \varepsilon_n / R_0 < r_n(w) < R_0 \text{ for some } n \in S \text{ with } \varepsilon_n \neq 0\}$$

and denote by $\iota_\tau: A_\tau \rightarrow A_\tau$ the biholomorphic map characterized by

$$\phi_{\nu(n)}^{-1} \circ \iota_\tau \circ \phi_n(v) \otimes v = \tau_n$$

for every $n \in S$ and $v \in T_n \Sigma_0$ with $\varepsilon_n / R_0 < |v| < R_0$.

Definition 4.9. Consider the Riemann surface with boundary

$$\Sigma_\tau^\circ := \{z \in \Sigma_0 : r_n(z) \geq \varepsilon_n^{1/2} \text{ for every } n \in S\}.$$

Denote by \sim_τ the equivalence relation on Σ_τ° generated by identifying the boundary components via ι_τ . The quotient

$$\Sigma_\tau := \Sigma_\tau^\circ / \sim_\tau$$

is a closed surface. The restrictions of the complex structure j_0 and the nodal structure ν_0 to Σ_τ° descends to a complex structure j_τ and a nodal structure ν_τ on Σ_τ . The nodal Riemann surface $(\Sigma_\tau, j_\tau, \nu_\tau)$ is called the *partial smoothing* of (Σ_0, j_0, ν_0) associated with τ .

Remark 4.10. The above construction smooths out every node with $\varepsilon_n > 0$. In particular, if all of the ε_n are positive, then ν_τ is the trivial nodal structure and $(\Sigma_\tau, j_\tau, \nu_\tau)$ is simply the Riemann surface (Σ_τ, j_τ) .

Definition 4.11. Denote by Δ the space of smoothing parameters for (Σ_0, j_0, ν_0) . Set

$$\mathcal{X} := \{(z, \tau) \in \Sigma_0 \times \Delta : z \in \Sigma_\tau^\circ\} / \sim$$

with $(z_1, \tau_1) \sim (z_2, \tau_2)$ if and only if $\tau_1 = \tau_2$ and $z_1 \sim_{\tau_1} z_2$ or $z_1, z_2 \in S, \nu(z_1) = z_2$, and $\varepsilon_{z_1} = \varepsilon_{z_2} = 0$. Denote by $\pi: \mathcal{X} \rightarrow \Delta$ the canonical projection.

The following example will be important in the proof of Theorem 1.1 in Section 5.8.

Example 4.12. Let (Σ_1, j_1, ν_1) and (Σ_2, j_2, ν_2) be two nodal Riemann surfaces with nodal sets S_1 and S_2 . Given $x_i \in \Sigma_i \setminus S_i$ for $i = 1, 2$, we define a new nodal Riemann surface $(\Sigma_\clubsuit, j_\clubsuit, \nu_\clubsuit)$ by setting $\Sigma_\clubsuit = \Sigma_1 \amalg \Sigma_2$ and $\nu_\clubsuit(x_1) = x_2$ (and otherwise agreeing with ν_1 and ν_2). The nodal set of $(\Sigma_\clubsuit, j_\clubsuit, \nu_\clubsuit)$ is

$$S_\clubsuit = \{x_1, x_2\} \amalg S_1 \amalg S_2.$$

Accordingly, the space of smoothing parameters for $(\Sigma_\clubsuit, j_\clubsuit, \nu_\clubsuit)$ is

$$\Delta_\clubsuit = \Delta_0 \times \Delta_1 \times \Delta_2,$$

where Δ_0 is an open neighborhood of zero in $T_{x_1} \Sigma_1 \otimes_{\mathbb{C}} T_{x_2} \Sigma_2$.

Suppose now that (Σ_1, j_1, ν_1) is a tree of spheres. It is easy to see that for every smoothing parameter $\tau_\clubsuit = (\tau_0, \tau_1, \tau_2)$ such that $\tau_0 \neq 0$ and $\tau_{1,n} \neq 0$ for every node $n \in S_1$, there is a biholomorphism

$$(\Sigma_{\clubsuit, \tau_\clubsuit}, j_{\tau_\clubsuit}, \nu_{\tau_\clubsuit}) \cong (\Sigma_{2, \tau_2}, j_{\tau_2}, \nu_{\tau_2}).$$

In particular, if $\tau_0 \neq 0, \tau_{1,n} \neq 0$ for every $n \in S_1$, and $\tau_2 = 0$, there is a biholomorphism

$$(\Sigma_{\clubsuit, \tau_\clubsuit}, j_{\tau_\clubsuit}, \nu_{\tau_\clubsuit}) \cong (\Sigma_2, j_2, \nu_2).$$

Proposition 4.13. *The space \mathcal{X} defined in Definition 4.11 is a smooth manifold and the complex structure on $\Sigma_0 \times \Delta$ induces a complex structure on \mathcal{X} such that π is a nodal family and for every $\tau \in \Delta$ the canonical map $\Sigma_\tau \rightarrow \pi^{-1}(\tau)$ induces a nodal, biholomorphic map*

$$\iota_\tau: (\Sigma_\tau, j_\tau, \nu_\tau) \rightarrow \pi^{-1}(\tau).$$

Proof. It suffices to consider the local model of a node $C_0 := \{(z, w) \in \mathbb{C}^2 : zw = 0\}$. $\tilde{\mathcal{X}} := \{(z, w, \tau) \in \mathbb{C}^2 \times \mathbb{C} : zw = \tau\}$ is a complex manifold. The map $\tilde{\pi}: \tilde{\mathcal{X}} \rightarrow \mathbb{C}$ defined by $\tilde{\pi}(z, w, \tau) := \tau$ has only nodal critical points and its fiber over 0 is C_0 . The nodal Riemann surface associated with C_0 is $\Sigma_0 = \mathbb{C} \amalg \mathbb{C}$ with the complex structure i on both components and the nodal structure which interchanges the origins of the components. The partial smoothing defined in Definition 4.9 is

$$\Sigma_\tau = (\{z \in \mathbb{C} : |z| \geq |\tau|^{1/2}\} \amalg \{w \in \mathbb{C} : |w| \geq |\tau|^{1/2}\}) / \sim_\tau.$$

The map $\Phi: \mathcal{X} \rightarrow \tilde{\mathcal{X}}$ defined by $\Phi([z], \tau) := (z, \tau/z, \tau)$ and $\Phi([w], \tau) := (\tau/z, z, \tau)$ is biholomorphic. This implies the assertion. ■

4.3. Construction of a versal deformation

Let (Σ_0, j_0, ν_0) be a nodal Riemann surface with nodal set S . Denote by $\mathcal{J}(\Sigma_0)$ the space of almost complex structures on Σ_0 and by $\text{Diff}_0(\Sigma_0, \nu_0)$ the group of diffeomorphism of Σ_0 which are isotopic to the identity and commute with ν_0 . Denote by

$$\mathcal{T} := \mathcal{J}(\Sigma_0) / \text{Diff}_0(\Sigma_0, \nu_0)$$

the corresponding Teichmüller space. This is a complex manifold and there is an open neighborhood Δ_1 of $0 \in \mathbb{C}^{\dim_{\mathbb{C}} \mathcal{T}}$ together with a map $J: \Delta_1 \rightarrow \mathcal{J}(\Sigma_0)$ such that:

- (1) $J(0) = j_0$,
- (2) for every $\sigma \in \Delta_1$ the almost complex structure $J(\sigma)$ agrees with j_0 in some neighborhood U of S , and
- (3) the map $[J]: \Delta_1 \rightarrow \mathcal{T}$ is an embedding.

For every $\sigma \in \Delta_1$, set

$$\Sigma_{\sigma,0} := \Sigma_0, \quad j_{\sigma,0} := J(\sigma), \quad \text{and} \quad \nu_{\sigma,0} := \nu_0.$$

Choose a family of metrics $(g_{\sigma,0})_{\sigma \in \Delta_1}$ whose restriction to the neighborhood U of S is independent of σ and such that $g_{\sigma,0}$ is in the conformal class determined by $j_{\sigma,0}$ for every $\sigma \in \Delta_1$. Let $R_0 > 0$ be such that the conditions at the beginning of Section 4.2 hold for every $\sigma \in \Delta_1$ and $B_{4R_0}(S) \subset U$.

Denote by Δ_2 the space of elements

$$\tau = (\tau_n)_{n \in S} \in \prod_{n \in S} T_n \Sigma_0 \otimes_{\mathbb{C}} T_{\nu(n)} \Sigma_0$$

such that for every $n \in S$, we have $\tau_{\nu(n)} = \sigma(\tau_n)$ and $|\tau_n| < R_0^2$.

Definition 4.14. Set $\Delta := \Delta_1 \times \Delta_2$. Set

$$\mathcal{X} := \{(z; \sigma, \tau) \in \Sigma_0 \times \Delta_1 \times \Delta_2 : z \in \Sigma_{\sigma, \tau}^{\circ}\} / \sim$$

with $(z_1; \sigma_1, \tau_1) \sim (z_2; \sigma_2, \tau_2)$ if and only if $\sigma_1 = \sigma_2$, $\tau_1 = \tau_2$ and $z_1 \sim_{\tau_1} z_2$ or $\tau_1 = \tau_2$ and $z_1 \sim_{\tau_1} z_2$ or $z_1, z_2 \in S$, $\nu(z_1) = z_2$, and $\varepsilon_{z_1} = \varepsilon_{z_2} = 0$. Denote by $\pi: \mathcal{X} \rightarrow \Delta$ the canonical projection.

Proposition 4.15. *The space \mathcal{X} defined in Definition 4.14 is a smooth manifold and the complex structure on $\Sigma_0 \times \Delta$ induces a complex structure on \mathcal{X} such that π is a nodal family and for every $(\sigma, \tau) \in \Delta$ the canonical map $\Sigma_{\sigma, \tau} \rightarrow \pi^{-1}(\sigma, \tau)$ induces a nodal, biholomorphic map*

$$\iota_{\sigma, \tau}: (\Sigma_{\sigma, \tau}, j_{\sigma, \tau}, \nu_{\sigma, \tau}) \rightarrow \pi^{-1}(\sigma, \tau).$$

Theorem 4.16 (cf. [1, Chapter XI, Theorem 3.17 and Section 4]). *Set $\star := (0, 0)$ and $\iota := \iota_{0, 0}$. The deformation (π, \star, ι) of (Σ_0, j_0, ν_0) is versal.*

Proof. Denote by C the nodal curve associated with (Σ_0, j_0, ν_0) . It is proved in [1, Chapter XI, Theorem 3.17] that the Kodaira–Spencer map

$$\delta: T_0 \Delta_1 \times T_0 \Delta_2 \rightarrow \text{Ext}^1(\Omega_C^1, \mathcal{O}_C)$$

is an isomorphism. This implies that the deformation is versal. Indeed, C has some versal family $(\rho: \mathcal{Y} \rightarrow B, \dagger, \kappa)$ for which the Kodaira–Spencer map is an isomorphism. Therefore, after possibly shrinking Δ , there exists a morphism of deformations

$$(\Phi, \phi): (\pi, \star, \iota) \rightarrow (\rho, \dagger, \kappa).$$

Since both Kodaira–Spencer maps are isomorphism, after possibly shrinking Δ , ϕ is a holomorphic embedding. Therefore, after possibly shrinking both Δ and A , both deformations become isomorphic. ■

To define a framing of the deformation (π, \star, ι) , choose an increasing, smooth function $\eta: [0, 2] \rightarrow [1, 2]$ such that

$$\eta(0) = 1 \quad \text{and} \quad \eta(r) = r$$

for every $3/2 \leq r \leq 2$.

Definition 4.17. Define the framing $\Psi: \Sigma_0 \setminus S \times \Delta \rightarrow \mathcal{X}$ of (π, \star, ι) by

$$\Psi(z; \sigma, \tau) := \begin{cases} \psi_n(z) & \text{if } r_n(z) \leq 2\varepsilon_n^{1/2} \text{ for some } n \in S, \\ (z; \sigma, \tau) & \text{otherwise,} \end{cases}$$

with $\psi_n(z)$ defined by

$$\psi_n(z) = \phi_n \left(\eta(r_n(z)/\varepsilon_n^{1/2}) \cdot \frac{\phi_n^{-1}(z)}{r_n(z)/\varepsilon_n^{1/2}} \right).$$

Observe that ψ_n is defined so that:

- (1) $r_n(\psi_n(z)) \geq \varepsilon_n^{1/2}$, so that indeed $\psi_n(z)$ corresponds to a point in \mathcal{X} ,
- (2) ψ_n defines an embedding from a punctured neighborhood of n in Σ_0 to \mathcal{X} ,
- (3) $\psi_n(z) = z$ when $r_n(z) \geq 3/2\varepsilon_n^{1/2}$, so that Ψ is continuous.

Remark 4.18. Let $(\sigma, \tau) \in \Delta$ and $r \in (2\varepsilon^{1/2}, R_0)$. Set

$$\Sigma_0^r := \{z \in \Sigma_0 : r_n(z) \geq r \text{ for every } n \in S\}.$$

Denote by

$$N_{\sigma,\tau}^r := \Sigma_{\sigma,\tau} \setminus \Psi(\Sigma_0^r \times \{(\sigma, \tau)\})$$

the part of $\Sigma_{\sigma,\tau}$ not covered by Σ_0^r under the framing Ψ , cf. Section 3.3. By construction,

$$N_{\sigma,\tau}^r = \bigcup_{n \in S} N_{\sigma,\tau;n}^r$$

with

$$N_{\sigma,\tau;n}^r = N_{\sigma,\tau;\nu(n)}^r := \{z \in \Sigma_{\sigma,\tau}^\circ : r_n(z) < r \text{ or } r_{\nu(n)}(z) < r\} / \sim_\tau.$$

If $\varepsilon_n = 0$, then $N_{\sigma,\tau;n}^r$ is biholomorphic to $B_r(0) \amalg B_r(0)$ and the nodal structure $\nu_{\sigma,\tau}$ identifies the two origins. If $\varepsilon_n \neq 0$, then $N_{\sigma,\tau;n}^r$ is biholomorphic to

$$\{z \in \mathbf{C} : \varepsilon_n/r < |z| < r\} \cong S^1 \times (-\log(r\varepsilon_n^{-1/2}), \log(r\varepsilon_n^{-1/2})).$$

5. Smoothing nodal J -holomorphic maps

The purpose of this section is to prove Theorem 1.1. The strategy is to construct a Kuranishi model for a Gromov neighborhood of u_∞ and analyze the obstruction map. This idea goes back to Ionel [18] and has been used by Zinger [38] and Niu [25] to give a sharp compactness results for genus one and two pseudo-holomorphic maps.

Throughout this section, fix a smooth function $\chi: [0, \infty) \rightarrow [0, 1]$ with

$$\chi|_{[0,1]} = 1 \quad \text{and} \quad \chi|_{[2,\infty)} = 0 \tag{5.1}$$

and, moreover, fix $p \in (2, \infty)$.

5.1. Riemannian metrics on smoothings

Let (Σ_0, j_0, ν_0) be a nodal Riemann surface with nodal set S . Denote by g_0 a Riemannian metric on Σ_0 as at the beginning of Section 4.2. In Section 4.2 we discussed the construction of a smoothing Σ_τ of Σ_0 for every smoothing parameter τ . In this section we construct a Riemannian metric g_τ on Σ_τ which is uniformly equivalent to the metric g_0 on Σ_0 in the smoothing region. This property will be useful for proving estimates in the construction of a smoothing of a nodal pseudo-holomorphic map from Σ_0 .

Definition 5.2. Given a smoothing parameter τ , let Σ_τ° be as in Definition 4.9. Recall that for every node $n \in S$ we have the corresponding number $\varepsilon_n = |\tau_n|$, the size of the smoothing parameter at n , and local radial coordinate $r_n: \Sigma_0 \rightarrow [0, \infty)$.

Define the Riemannian metric g_τ° on Σ_τ° by

$$g_\tau^\circ := g_0 + \sum_{n \in S} \chi\left(\frac{r_n}{2\varepsilon_n^{1/2}}\right) \cdot (\varepsilon_n \cdot (\phi_n)_*(r^{-2}dr \otimes dr + \theta \otimes \theta) - g_0)$$

with r denoting the distance from origin in $T_n \Sigma_0 \cong \mathbf{C}$ and $\theta = -dr \circ j_0$. Since the Riemannian metric $r^{-2}dr \otimes dr + \theta \otimes \theta$ on \mathbf{C}^* is invariant under the involution $z \mapsto \varepsilon/z$, g_τ° descends to a Riemannian metric g_τ on Σ_τ .

Proposition 5.3. *There is a constant $c > 1$ such that for every nodal Riemann surface and every smoothing parameter τ ,*

$$c^{-1}g_0 < g_\tau^\circ < cg_0.$$

Proof. Let $n \in S$ and let r and θ be as in Definition 5.2. On the annulus

$$\{z \in \Sigma_0 : \varepsilon_n^{1/2} \leq r_n(z) \leq 4\varepsilon_n^{1/2}\},$$

we have $\phi_n^*g_0 = dr \otimes dr + r^2\theta \otimes \theta$ and, therefore,

$$g_\tau^\circ = (F_{\varepsilon_n} \circ r_n) \cdot g_0 \quad \text{with} \quad F_{\varepsilon_n}(r) := 1 + \chi\left(\frac{r}{2\varepsilon_n^{1/2}}\right) \cdot (\varepsilon_n r^{-2} - 1).$$

This implies the assertion because $c^{-1} < F_{\varepsilon_n}(r) < c$ for $\varepsilon_n^{1/2} \leq r \leq 4\varepsilon_n^{1/2}$. ■

Henceforth, the L^p and $W^{1,p}$ norms of all sections and differential forms on Σ_τ are understood with respect to the metric g_τ . The above proposition will often be implicitly used to bound these norms by estimating various expressions with respect to g_0 over the corresponding region in Σ_0 .

5.2. Approximate smoothing of nodal J -holomorphic maps

Throughout the next four sections, let (X, J, h) be an almost Hermitian manifold, let $c_u > 0$, let $u_0: (\Sigma_0, j_0, \nu_0) \rightarrow (X, J)$ be a nodal map, and let τ be a smoothing parameter. Furthermore, choose g_0 and R_0 as at the beginning of Section 4.2.

Definition 5.4. For every point $x \in X$, denote by $\tilde{U}_x \subset T_x X$ the segment/injectivity domain and set $U_x := \exp_x(\tilde{U}_x)$ and $\frac{1}{2}U_x := \exp_x(\frac{1}{2}\tilde{U}_x)$. The map $\exp_x: \tilde{U}_x \rightarrow U_x$ is a diffeomorphism and its inverse is denoted by $\exp_x^{-1}: U_x \rightarrow \tilde{U}_x$.

Furthermore, we assume the following.

Hypothesis 5.5. *The map u_0 and $R_0 > 0$ satisfy*

$$\|u_0\|_{C^2} \leq c_u \quad \text{and} \quad u_0(B_{4R_0}(n)) \subset U_{u_0(n)} \quad \text{for every } n \in S.$$

Convention 5.6. Henceforth, constants may depend on $p, (\Sigma_0, j_0, \nu_0), (X, J, h), c_u,$ and R_0 , but not on τ .

Definition 5.7. For $n \in S$, define $\chi_\tau^n: \Sigma_\tau \rightarrow [0, 1]$ by

$$\chi_\tau^n(z) := \chi\left(\frac{r_n(z)}{R_0}\right).$$

Let ι_τ be the map defined in Definition 4.8. Define $\tilde{u}_\tau^\circ: \Sigma_\tau^\circ \rightarrow X$ by

$$\tilde{u}_\tau^\circ(z) := \begin{cases} \exp_{u_0(n)}(\exp_{u_0(n)}^{-1} \circ u_0(z) + \chi_\tau^n(z) \cdot \exp_{u_0(n)}^{-1} \circ u_0(\iota_\tau(z))) & \text{if } r_n(z) \leq 2R_0, \\ u_0(z) & \text{otherwise.} \end{cases}$$

Since $u_0(\nu_0(n)) = u_0(n)$, the restriction of \tilde{u}_τ° to

$$\{z \in \Sigma_\tau^\circ : r_n(z) \leq R \text{ for some } n \in S\}$$

is invariant under ι_τ . Therefore, \tilde{u}_τ° descends to a smooth map

$$\tilde{u}_\tau: \Sigma_\tau \rightarrow X.$$

This map is called the *approximate smoothing* of u associated with τ .

Remark 5.8. This construction differs from that found, for example, in [24, Section 10.2] and [31, Section B.3] in which the approximate smoothing is constant in the middle of the neck region. The above construction is very similar to that in [12, Section 2.1]. It leads to a smaller error term and significantly simplifies the discussions in Section 5.7. Morally, this section analyzes how the interaction between the different components of u_0 affects whether u_0 can be smoothed or not. The constructions in [24, Section 10.2] and [31, Section B.3] make it difficult to see such interactions.

Proposition 5.9. *For every nodal map $u_0 : \Sigma_0 \rightarrow X$ (not necessarily J -holomorphic) the map \tilde{u}_τ satisfies*

$$\|\bar{\partial}_J(\tilde{u}_\tau, j_\tau)\|_{L^p} \leq c \|\bar{\partial}_J(u_0, j_0)\|_{L^p} + c\varepsilon^{\frac{1}{2} + \frac{1}{p}},$$

with $\varepsilon = \max\{\varepsilon_n : n \in S\}$ as in Definition 4.7.

For the proof of this result and for future reference let us observe that for every $k \geq 1$,

$$\left(\int_{\varepsilon_n^{1/2} \leq r_n \leq 2R_0} r_n^{-kp} \right)^{\frac{1}{p}} \leq \left(\frac{2\pi}{kp - 2} \right)^{\frac{1}{p}} \varepsilon_n^{\frac{1}{p} - \frac{k}{2}}. \tag{5.10}$$

Proof of Proposition 5.9. The map \tilde{u}_τ° agrees with u_0 in the region where $r_n \geq 2R_0$ for every $n \in S$. Therefore, it suffices to consider the regions where $r_n \leq 2R_0$ for some $n \in S$. To simplify notation, identify U_x with \tilde{U}_x via \exp_x for $x := u(n)$. Here U_x and \tilde{U}_x are as in Definition 5.4. Having made this identification, in such a region, \tilde{u}_τ° is given by

$$\tilde{u}_\tau^\circ = u_0 + \chi_\tau^n \cdot u_0 \circ \iota_\tau.$$

Note that addition is well defined since, with respect to the above identification, u_0 takes values in an sufficiently small open subset of $T_x X$. Therefore,

$$\begin{aligned} \bar{\partial}_J(\tilde{u}_\tau^\circ, j_\tau) &= \frac{1}{2} (d\tilde{u}_\tau^\circ + J(\tilde{u}_\tau^\circ) \circ d\tilde{u}_\tau^\circ \circ j) \\ &= \underbrace{\bar{\partial}_J(u_0, j_0) + \chi_\tau^n \cdot \bar{\partial}_J(u_0 \circ \iota_\tau, j_0)}_{=: \text{I}} + \underbrace{\frac{1}{2} (J(\tilde{u}_\tau^\circ) - J(u_0)) \circ du_0 \circ j_0}_{=: \text{II}_1} \\ &\quad + \underbrace{\chi_\tau^n \cdot \frac{1}{2} (J(\tilde{u}_\tau^\circ) - J(u_0 \circ \iota_\tau)) \circ d(u_0 \circ \iota_\tau) \circ j_0}_{=: \text{II}_2} + \underbrace{\bar{\partial} \chi_\tau^n \cdot u_0 \circ \iota_\tau}_{=: \text{III}}. \end{aligned}$$

The L^p norm of the term I is controlled by the L^p norm of $\bar{\partial}_J(u_0, j_0)$ over the regions of Σ_0 , where $\varepsilon_n/2R_0 \leq r_n \leq 2R_0$ for some $n \in S$. By Taylor expansion,

$$|\text{II}_1| \leq c \|J\|_{C^1} \cdot |u_0 \circ \iota_\tau| \cdot |du_0| \leq c\varepsilon_n/r_n$$

and

$$\begin{aligned} |\text{II}_2| &\leq c \|J\|_{C^1} \cdot (|u_0| + (1 - \chi_\tau^n) \cdot |u_0 \circ \iota|) \cdot |d(u_0 \circ \iota_\tau)| \\ &\leq c \cdot (r_n + (1 - \chi_\tau^n) \varepsilon_n / r_n) \cdot \varepsilon_n / r_n^2 \leq c \varepsilon_n / r_n. \end{aligned}$$

On Σ_τ° , by definition, $r_n \geq \varepsilon_n^{1/2}$. Therefore and by (5.10),

$$\|\text{II}_1 + \text{II}_2\|_{L^p} \leq c \varepsilon_n^{\frac{1}{2} + \frac{1}{p}}.$$

The term III is supported in the region where $R_0 \leq r_n \leq 2R_0$, whose area is independent of ε_n , and satisfies

$$|\text{III}| \leq c |d\chi_\tau^n| |u_0 \circ \iota_\tau| \leq c r_n \cdot (\varepsilon_n / r_n) = c \varepsilon_n,$$

where in the last inequality we use that, with respect to the identifications introduced earlier, $u_0(0) = 0$ so $|u_0(z)| \leq c|z|$ in a neighborhood of 0. Therefore,

$$\|\text{III}\|_{L^p} \leq c \varepsilon_n. \quad \blacksquare$$

5.3. Fusing nodal vector fields

The purpose of this section is to introduce the fusing operator. This operator assigns to every vector field ξ along u_0 a vector field $\text{fuse}_\tau(\xi)$ along \tilde{u}_τ , which agrees with ξ outside the gluing region. The construction of the fusing operator makes use of the following local trivialisations of TX .

Definition 5.11. For every $x \in X$ and $y \in U_x$, define an isomorphism

$$\Phi_y = \Phi_y^x: T_x X \rightarrow T_y X$$

by

$$\Phi_y^x(v) := d_{\exp_x^{-1}(y)} \exp_x(v)$$

As y varies in U_x , these maps define a trivialisation $\Phi = \Phi^x: U_x \times T_x X \rightarrow TX|_{U_x}$.

The definition of the fusing operator uses a different cutoff function than the definition of the approximate smoothing u_τ .

Definition 5.12. For $n \in S$, define $\rho_\tau^n: \Sigma_\tau \rightarrow [0, 1]$ by

$$\rho_\tau^n(z) := \chi\left(\frac{r_n(z)}{\varepsilon_n^{1/4}}\right).$$

Here χ is the cutoff function (5.1); that is, $\rho_\tau^n = 1$ in the region where $r_n \leq \varepsilon_n^{1/4}$, and $\rho_\tau^n = 0$ in the region where $r_n \geq 2\varepsilon_n^{1/4}$.

Definition 5.13. Define $\text{fuse}_\tau^\circ: W^{1,p}\Gamma(\Sigma_0, \nu_0; u_0^*TX) \rightarrow W^{1,p}\Gamma(\Sigma_\tau^\circ, \nu_0; (\tilde{u}_\tau^\circ)^*TX)$ by

$$\text{fuse}_\tau^\circ(\xi)(z) := \begin{cases} \Phi_{\tilde{u}_\tau}([\!|z|]) (\Phi_{u_0(z)}^{-1}\xi(z) + \rho_\tau^n(z) \cdot (\Phi_{u_0 \circ \iota_\tau(z)}^{-1}\xi(\iota_\tau(z)) - \Phi_{u_0}^{-1}\xi(n))) & \text{if } r_n(z) \leq 2\varepsilon_n^{1/4}, \\ \xi(z) & \text{otherwise.} \end{cases}$$

In the above formula, $\Phi = \Phi^x$ with $x = u_0(n)$. For every $n \in S$ the restriction of $\text{fuse}_\tau^\circ(\xi)$ to

$$\{z \in \Sigma_\tau^\circ : r_n(z) \leq \varepsilon_n^{1/4} \text{ for some } n \in S\}$$

is invariant under ι_τ . Therefore, fuse_τ° induces a map

$$\text{fuse}_\tau: W^{1,p}\Gamma(\Sigma_0, \nu_0; u_0^*TX) \rightarrow W^{1,p}\Gamma(\Sigma_\tau, \nu_\tau; \tilde{u}_\tau^*TX).$$

The following is a counterpart of Proposition 5.9.

Proposition 5.14. For every $\xi \in W^{1,p}\Gamma(\Sigma_0, \nu_0; u_0^*TX)$,

$$\|\text{d}_{\tilde{u}_\tau} \text{fuse}_\tau(\xi)\|_{L^p} \leq c \|\text{d}_{u_0}\xi\|_{L^p} + c \sum_{n \in S} (\varepsilon_n^{\frac{1}{2p}} + \varepsilon_n^{\frac{1}{2} - \frac{1}{p}}) \|\xi\|_{W^{1,p}}.$$

The proof requires the following results as a preparation.

Proposition 5.15. For every $n \in S$ and $\xi \in W^{1,p}\Gamma(\Sigma_0, \nu; u_0^*TX)$,

$$\|\text{d}\rho_\tau^n \cdot (\xi \circ \iota_\tau - \xi(n))\|_{L^p} \leq c\varepsilon_n^{\frac{1}{2} - \frac{1}{p}} \|\xi\|_{W^{1,p}}.$$

Proof. Morrey’s embedding theorem asserts that $W^{1,p} \hookrightarrow C^{0,1-2/p}$. Hence,

$$|\xi \circ \iota_\tau(z) - \xi(n)| \leq c(\varepsilon_n/r_n(z))^{1-2/p} \|\xi\|_{W^{1,p}}.$$

The term $\text{d}\rho_\tau^n$ is supported in the annulus $P_\tau^n = \{\varepsilon_n^{1/4} \leq r_n \leq 2\varepsilon_n^{1/4}\}$ and satisfies

$$|\text{d}\rho_\tau^n| \leq c\varepsilon_n^{-1/4}.$$

Since the area of P_τ^n is proportional to $\varepsilon_n^{1/2}$,

$$\begin{aligned} \|\text{d}\rho_\tau^n \cdot (\xi \circ \iota_\tau - \xi(n))\|_{L^p} &\leq c\varepsilon_n^{-1/4} \varepsilon_n^{3/4(1-2/p)} \varepsilon_n^{1/2p} \|\xi\|_{W^{1,p}} \\ &= c\varepsilon_n^{\frac{1}{2} - \frac{1}{p}} \|\xi\|_{W^{1,p}}. \end{aligned}$$

■

Proposition 5.16. Let $U \subset \Sigma_0$ be an open subset. Let $u_1, u_2: U \rightarrow U_x$ and set

$$v := \exp_x^{-1} \circ u_2 - \exp_x^{-1} \circ u_1.$$

For every $\xi \in C^\infty(U, T_x X)$,

$$|(\Phi_{u_1} \circ \text{d}_{u_1} \circ \Phi_{u_1}^{-1} - \Phi_{u_2} \circ \text{d}_{u_2} \circ \Phi_{u_2}^{-1})\xi| \leq c(|v| |\text{d}\xi| + |\text{d}v| |\xi| + |\text{d}u_1| |\xi| |v|).$$

Proof. To simplify notation, identify U_x with \tilde{U}_x via \exp_x . Having made this identification, Φ becomes the identity map and $v = u_2 - u_1$. Therefore,

$$\begin{aligned} \delta_{u_1}\xi - \delta_{u_2}\xi &= \frac{1}{2}(J(u_1) - J(u_2)) \circ \nabla\xi \circ j \\ &\quad + \frac{1}{2}((\nabla_\xi J)(u_1) - (\nabla_\xi J)(u_2)) \circ du_1 \circ j \\ &\quad + \frac{1}{2}(\nabla_\xi J)(u_2) \circ (du_1 - du_2) \circ j. \end{aligned}$$

This implies the asserted inequality. ■

Proof of Proposition 5.14. Outside the regions where $r_n \leq 2R_0$ for some $n \in S$, the operators δ_{u_0} and $\delta_{\tilde{u}_\tau}$ agree. Within such a region and with the usual identifications

$$\begin{aligned} \delta_{\tilde{u}_\tau^\circ} \text{fuse}_\tau^\circ(\xi) &= \delta_{u_0}\xi + \underbrace{(\delta_{\tilde{u}_\tau^\circ} - \delta_{u_0})\xi}_{=: \text{I}} + \underbrace{\bar{\partial}\rho_\tau^n \cdot (\xi \circ \iota_\tau - \xi(n))}_{=: \text{II}} \\ &\quad + \underbrace{\rho_\tau^n \cdot \delta_{\tilde{u}_\tau^\circ}(\xi \circ \iota_\tau)}_{=: \text{III}} - \underbrace{\rho_\tau^n \cdot \delta_{\tilde{u}_\tau^\circ}\xi(n)}_{=: \text{IV}}. \end{aligned}$$

The difference $v := \tilde{u}_\tau^\circ - u_0 = \chi_\tau^n \cdot u_0 \circ \iota_\tau$ satisfies

$$\begin{aligned} |v| &\leq c\varepsilon_n/r_n \leq c\varepsilon_n^{1/2}, \\ |dv| &\leq |d\chi_\tau^n \cdot u_0 \circ \iota_\tau| + |\chi_\tau^n d(u_0 \circ \iota_\tau)| \leq c\varepsilon_n/r_n^2. \end{aligned}$$

Therefore, by Proposition 5.16 and (5.10),

$$\|\text{I}\|_{L^p} \leq c\varepsilon_n^{\frac{1}{p}} \|\xi\|_{W^{1,p}}.$$

By Proposition 5.15,

$$\|\text{II}\|_{L^p} \leq c\varepsilon_n^{\frac{1}{2} - \frac{1}{p}} \|\xi\|_{W^{1,p}}.$$

The term III can be written as

$$\text{III} = \rho_\tau^n \cdot (\delta_{\tilde{u}_\tau^\circ}\xi) \circ \iota_\tau + \rho_\tau^n \cdot (\delta_{\tilde{u}_\tau^\circ} - \delta_{\tilde{u}_\tau^\circ \circ \iota_\tau})(\xi \circ \iota_\tau).$$

The first term in this sum satisfies

$$\|\rho_\tau^n \cdot (\delta_{\tilde{u}_\tau^\circ}\xi) \circ \iota_\tau\|_{L^p} \leq \|\delta_{u_0}\xi\|_{L^p} + \|\text{I}\|_{L^p}.$$

To estimate the second term, consider the difference

$$w := \tilde{u}_\tau^\circ - \tilde{u}_\tau^\circ \circ \iota_\tau = (1 - \chi_\tau^n)(u_0 - u_0 \circ \iota_\tau).$$

It satisfies

$$|w| \leq cr_n \quad \text{and} \quad |dw| \leq c.$$

Since ρ_τ^n is supported in the region where $\varepsilon_n^{1/4} \leq r_n \leq 2\varepsilon_n^{1/4}$, Proposition 5.16 implies that

$$\|\rho_\tau^n \cdot (\delta_{\tilde{u}_\tau^\circ} - \delta_{\tilde{u}_\tau^\circ \circ \iota_\tau})(\xi \circ \iota_\tau)\|_{L^p} \leq c\varepsilon_n^{1/2p} \|\xi\|_{W^{1,p}}.$$

Therefore,

$$\|\text{III}\|_{L^p} \leq \|\delta_{u_0}\xi\|_{L^p} + c\varepsilon_n^{1/2p} \|\xi\|_{W^{1,p}}.$$

To estimate the term IV, write it in the form

$$\text{IV} = \rho_\tau^n \cdot (\delta_{\tilde{u}_\tau^\circ} - \delta_{u(n)})\xi(n),$$

with $\delta_{u(n)}$ denoting the operator associated with the constant map with value $u(n)$. Since the difference $u_\tau^\circ - u(n)$ and its derivative are bounded, again from Proposition 5.16 we conclude that

$$\|\text{IV}\|_{L^p} \leq c\varepsilon_n^{1/2p} \|\xi\|_{W^{1,p}}. \quad \blacksquare$$

5.4. Construction of right inverses

Throughout this subsection, let $\mathcal{O} \subset L^p\Omega^{0,1}(\Sigma_0, u_0^*TX)$ be a finite-dimensional subspace such that

$$\text{im } \delta_{u_0} + \mathcal{O} = L^p\Omega^{0,1}(\Sigma_0, u_0^*TX). \tag{5.17}$$

In particular, \mathcal{O} surjects onto $\text{coker } \delta_{u_0}$.

Definition 5.18. Define $\text{pull}_\tau: L^p\Omega^{0,1}(\Sigma_0, u_0^*TX) \rightarrow L^p\Omega^{0,1}(\Sigma_\tau, \tilde{u}_\tau^*TX)$ by

$$\text{pull}_\tau(\eta)([z]) := \begin{cases} \Phi_{\tilde{u}_\tau([z])} \Phi_{u_0(z)}^{-1} \eta(z) & \text{if } \varepsilon_n^{1/2} \leq r_n(z) \leq 2R_0, \\ \eta(z) & \text{otherwise.} \end{cases}$$

Recall that Σ_τ is defined in Definition 4.9 by identifying the boundary components of $\{r_n \geq \varepsilon_n^{1/2}\}$ and $\{r_{v(n)} \geq \varepsilon_n^{1/2}\}$ using ι_τ . The operator pull_τ is obtained by simply restricting $(0, 1)$ -forms to these regions. The resulting $(0, 1)$ -form on Σ_τ is typically not continuous but it is still in L^p . In particular, the ambiguity at $r_n = \varepsilon_n^{1/2}$ in Definition 5.18 is immaterial. The reader should contrast Definition 5.18 with the definition of fuse_τ , cf. Definition 5.13, which produces sections of class $W^{1,p}$, and therefore continuous.

Definition 5.19. Define $\bar{\delta}_{u_0}: W^{1,p}\Gamma(\Sigma_0, \nu_0; u_0^*TX) \oplus \mathcal{O} \rightarrow L^p\Omega^{0,1}(\Sigma_0, u_0^*TX)$ by

$$\bar{\delta}_{u_0}(\xi, o) := \delta_{u_0}\xi + o.$$

Define $\bar{\delta}_{\tilde{u}_\tau}: W^{1,p}\Gamma(\Sigma_\tau, \nu_\tau; \tilde{u}_\tau^*TX) \oplus \mathcal{O} \rightarrow L^p\Omega^{0,1}(\Sigma_\tau, \tilde{u}_\tau^*TX)$ by

$$\bar{\delta}_{\tilde{u}_\tau}(\xi, o) := \delta_{\tilde{u}_\tau}\xi + \text{pull}_\tau(o).$$

By construction, $\bar{\delta}_{u_0}$ is surjective and, hence, has a right inverse

$$r_{u_0}: L^p \Omega^{0,1}(\Sigma_0, u_0^*TX) \rightarrow W^{1,p} \Gamma(\Sigma_0, \nu_0; u_0^*TX) \oplus \mathcal{O}$$

of $\bar{\delta}_{u_0}$. Henceforth, fix a choice of r_{u_0} . The purpose of this subsection is to construct a right inverse $r_{\tilde{u}_\tau}$ to $\bar{\delta}_{\tilde{u}_\tau}$ for sufficiently small ε .

Definition 5.20. Define $\text{push}_\tau: L^p \Omega^{0,1}(\Sigma_\tau, \tilde{u}_\tau^*TX) \rightarrow L^p \Omega^{0,1}(\Sigma_0, u_0^*TX)$ by

$$\text{push}_\tau(\eta)(z) := \begin{cases} 0 & \text{if } r_n(z) < \varepsilon_n^{1/2}, \\ \Phi_{u_0(z)} \Phi_{\tilde{u}_\tau^{-1}([z])}^{-1} \eta([z]) & \text{if } \varepsilon_n^{1/2} \leq r_n(z) \leq 2R_0, \\ \eta([z]) & \text{otherwise.} \end{cases}$$

Definition 5.21. Define $\tilde{r}_{\tilde{u}_\tau}: L^p \Omega^{0,1}(\Sigma_\tau, \tilde{u}_\tau^*TX) \rightarrow W^{1,p} \Gamma(\Sigma_\tau, \nu_\tau; \tilde{u}_\tau^*TX) \oplus \mathcal{O}$ by

$$\tilde{r}_{\tilde{u}_\tau} := (\text{fuse}_\tau \oplus \text{id}_{\mathcal{O}}) \circ r_{u_0} \circ \text{push}_\tau.$$

Proposition 5.22. *The linear operator $\tilde{r}_{\tilde{u}_\tau}$ satisfies*

$$\begin{aligned} \|\bar{\delta}_{\tilde{u}_\tau} \circ \tilde{r}_{\tilde{u}_\tau} - \text{id}\| &\leq c \sum_{n \in \mathcal{S}} (\varepsilon_n^{\frac{1}{2p}} + \varepsilon_n^{\frac{1}{2} - \frac{1}{p}}) \|r_{u_0}\|, \\ \|\tilde{r}_{\tilde{u}_\tau}\| &\leq c \|r_{u_0}\|. \end{aligned} \tag{5.23}$$

Proof. The map push_τ is bounded by a constant independent of τ and, by Proposition 5.15, so is fuse_τ . This implies the estimate on $\|\tilde{r}_{\tilde{u}_\tau}\|$.

Let $\eta \in L^p \Omega^{0,1}(\Sigma_\tau, \tilde{u}_\tau^*TX)$. In order to prove Proposition 5.23, we estimate $\|\bar{\delta}_{\tilde{u}_\tau} \tilde{r}_{\tilde{u}_\tau} \eta - \eta\|_{L^p}$ as follows. Set

$$(\xi, o) := r_{u_0} \circ \text{push}_\tau(\eta),$$

so that

$$\delta_{u_0} \xi + o = \text{push}_\tau(\eta).$$

By Proposition 5.16 applied to \tilde{u}_τ° and u_0 and using (5.10) and the fact that on Σ_τ° we have $\text{pull}_\tau(o) = o$,

$$\|\delta_{\tilde{u}_\tau^\circ} \xi + \text{pull}_\tau(o) - \eta\|_{L^p} \leq c \varepsilon^{\frac{1}{p}} \|\xi\|_{W^{1,p}} \leq c \varepsilon^{\frac{1}{p}} \|r_{u_0}\| \|\eta\|_{L^p}.$$

Therefore, it remains to estimate

$$\begin{aligned} \delta_{\tilde{u}_\tau^\circ}(\rho_\tau^n \cdot (\xi \circ \iota_\tau - \xi(n))) &= \underbrace{\bar{\partial} \rho_\tau^n \cdot (\xi \circ \iota_\tau - \xi(n))}_{=: \text{I}} + \underbrace{\rho_\tau^n \cdot \delta_{\tilde{u}_\tau^\circ}(\xi \circ \iota_\tau)}_{=: \text{II}} \\ &\quad - \underbrace{\rho_\tau^n \cdot \delta_{\tilde{u}_\tau^\circ} \xi(n)}_{=: \text{III}}. \end{aligned}$$

By Proposition 5.15,

$$\|I\|_{L^p} \leq c\varepsilon^{\frac{1}{2}-\frac{1}{p}} \|\xi\|_{W^{1,p}} \leq c\varepsilon^{\frac{1}{2}-\frac{1}{p}} \|r_{u_0}\| \|\eta\|_{L^p}.$$

To estimate the second term, observe that in the region where $r_n \geq \varepsilon_n^{1/2}$,

$$\delta_{u_0 \circ \iota_\tau}(\xi \circ \iota_\tau) = \iota_\tau^*(\delta_{u_0}\xi) = \iota_\tau^*(\text{push}_\tau(\eta)) = 0.$$

To understand the last identity, observe that $r_n(\iota_\tau(z)) = \varepsilon_n r_{v(n)}^{-1}(z)$ and $\text{push}_\tau(\eta)$ is defined to vanish in the region of Σ_0 where $r_{v(n)} \leq \varepsilon_n^{1/2}$. Thus, by Proposition 5.16 applied to \tilde{u}_τ and $u_0 \circ \iota_\tau$, and using the fact that ρ_τ^n is supported in the region where $\varepsilon_n^{1/4} \leq r_n \leq 2\varepsilon_n^{1/4}$ whose area is proportional to $\varepsilon_n^{1/2}$,

$$\|II\|_{L^p} \leq c\varepsilon^{\frac{1}{2p}} \|\xi\|_{W^{1,p}} \leq c\varepsilon^{\frac{1}{2p}} \|r_{u_0}\| \|\eta\|_{L^p}.$$

The vector field $\xi(n)$ is constant with respect the chosen trivialization. Since the operator $\delta_{u_0(n)}$ associated with the constant map agrees with the standard $\bar{\partial}$ -operator,

$$\delta_{u_0(n)}\xi(n) = 0.$$

Therefore, using Proposition 5.16 applied to \tilde{u}_τ and the constant map $u(n)$, and the estimate on the area of the support of ρ_τ^n , we arrive at

$$\|III\|_{L^p} \leq c\varepsilon^{\frac{1}{2p}} \|r_{u_0}\| \|\eta\|_{L^p}. \quad \blacksquare$$

Throughout the remainder of this subsection, suppose the following.

Hypothesis 5.24. *The smoothing parameter τ is such that the right-hand side of Proposition 5.23 is at most $1/2$.*

Definition 5.25. Define the *right inverse*

$$r_{\tilde{u}_\tau} : L^p \Omega^{0,1}(\Sigma_\tau, \tilde{u}_\tau^* TX) \rightarrow W^{1,p} \Gamma(\Sigma_\tau, \nu_\tau; \tilde{u}_\tau^* TX) \oplus \mathcal{O}$$

associated with r_{u_0} by

$$r_{\tilde{u}_\tau} := \tilde{r}_{\tilde{u}_\tau} (\bar{\delta}_{\tilde{u}_\tau} \tilde{r}_{\tilde{u}_\tau})^{-1} = \tilde{r}_{\tilde{u}_\tau} \sum_{k=0}^{\infty} (\text{id} - \bar{\delta}_{\tilde{u}_\tau} \tilde{r}_{\tilde{u}_\tau})^k.$$

The following is an immediate consequence of the definition.

Proposition 5.26. *The right inverse*

$$r_{\tilde{u}_\tau} : L^p \Omega^{0,1}(\Sigma_\tau, \tilde{u}_\tau^* TX) \rightarrow W^{1,p} \Gamma(\Sigma_\tau, \nu_\tau; \tilde{u}_\tau^* TX) \oplus \mathcal{O}$$

satisfies

$$\bar{\delta}_{u_\tau} r_{\tilde{u}_\tau} = \text{id} \quad \text{and} \quad \|r_{\tilde{u}_\tau}\| \leq c \|r_{u_0}\|;$$

furthermore,

$$\text{im } r_{\tilde{u}_\tau} = \text{im } \tilde{r}_{\tilde{u}_\tau}.$$

5.5. Complements of the image of $r_{\tilde{u}_\tau}$

Proposition 5.27. *Given $c_f > 0$ there is a constant $\delta = \delta(c_f) > 0$ such that the following holds. If τ satisfies $\varepsilon < \delta$ and $K \subset W^{1,p}\Gamma(\Sigma_0, \nu_0; u_0^*TX)$ is a subspace with $\dim K = \dim \ker \bar{d}_{u_0}$ and such that for every $\kappa \in K$,*

$$\|d_{u_0}\kappa\|_{L^p} \leq \delta\|\kappa\|_{W^{1,p}} \quad \text{and} \quad \|\kappa\|_{W^{1,p}} \leq c_f\|\text{fuse}_\tau(\kappa)\|_{W^{1,p}},$$

then every $(\xi, o) \in W^{1,p}\Gamma(\Sigma_\tau, \nu_\tau; \tilde{u}_\tau^*TX) \oplus \mathcal{O}$ can be uniquely written as

$$(\xi, o) = r_{\tilde{u}_\tau}\eta + (\kappa, 0)$$

with $\eta \in L^p\Omega^{0,1}(\Sigma_\tau, \tilde{u}_\tau^*TX)$ and $\kappa \in K$; moreover,

$$\|\eta\|_{L^p} + \|\kappa\|_{W^{1,p}} \leq c(c_f)(\|\xi\|_{W^{1,p}} + |o|).$$

Here $|o| = \|o\|_{L^p}$ is the norm of o induced by the inclusion $\mathcal{O} \subset L^p\Omega^{0,1}(\Sigma_0, u_0^*TX)$.

Proof. Because $r_{\tilde{u}_\tau}$ and $\text{fuse}_\tau|_K$ are injective and given the hypothesis on $\text{fuse}_\tau|_K$, it suffices to show that $W^{1,p}\Gamma(\Sigma_\tau, \nu_\tau; \tilde{u}_\tau^*TX) \oplus \mathcal{O}$ is the direct sum of $\text{im}(r_{\tilde{u}_\tau})$ and $\text{im}(\text{fuse}_\tau|_K) \oplus 0$.

By the index formula (2.27), Remark 2.11, and Proposition 3.16,

$$\begin{aligned} \text{index } d_{u_0} &= 2\langle (u_0^*c_1(X, J), [\Sigma_0]) \rangle + 2n(1 - p_a(\Sigma_0, \nu_0)) \\ &= 2\langle (\tilde{u}_\tau^*c_1(X, J), [\Sigma_\tau]) \rangle + 2n(1 - p_a(\Sigma_\tau, \nu_\tau)) \\ &= \text{index } d_{\tilde{u}_\tau}. \end{aligned}$$

Therefore and because \bar{d}_{u_0} is surjective and $r_{\tilde{u}_\tau}$ is injective,

$$\text{codim } \text{im}(r_{\tilde{u}_\tau}) = \text{index } \bar{d}_{\tilde{u}_\tau} = \text{index } \bar{d}_{u_0} = \dim \ker \bar{d}_{u_0}.$$

Hence, it remains to prove that $\text{im}(r_{\tilde{u}_\tau})$ and $\text{im}(\text{fuse}_\tau|_K) \oplus 0$ intersect trivially.

Suppose that $\eta \in L^p\Omega^{0,1}(\Sigma_{\sigma,\tau}, \tilde{u}_\tau^*TX)$ and $\kappa \in K$ satisfy

$$r_{\tilde{u}_\tau}(\eta) = (\text{fuse}_\tau(\kappa), 0).$$

By Proposition 5.14 as well as the hypothesis on fuse_τ and for sufficiently small δ ,

$$\begin{aligned} \|\eta\|_{L^p} &= \|d_{\tilde{u}_\tau}\text{fuse}_\tau(\kappa)\|_{L^p} \\ &\leq c[\delta + \#S \cdot (\delta^{\frac{1}{2p}} + \delta^{\frac{1}{2} - \frac{1}{p}})]\|\kappa\|_{W^{1,p}} \\ &\leq cc_f[\delta + \#S \cdot (\delta^{\frac{1}{2p}} + \delta^{\frac{1}{2} - \frac{1}{p}})]\|\eta\|_{L^p} \\ &\leq \frac{1}{2}\|\eta\|_{L^p}. \end{aligned}$$

Therefore, η vanishes. ■

5.6. Kuranishi model for a neighborhood of nodal maps

Throughout, let (Σ_0, j_0, ν_0) be a nodal Riemann surface with nodal set S , let (X, J_0, h) be an almost Hermitian manifold, and let $u_0: (\Sigma_0, j_0, \nu_0) \rightarrow (X, J_0)$ be a nodal J_0 -holomorphic map. Let $(\pi: \mathcal{X} \rightarrow \Delta, \star = (0, 0), \iota)$ be the versal deformation of (Σ_0, j_0, ν_0) constructed in Section 4.3 with fibers

$$(\Sigma_{\sigma, \tau}, j_{\sigma, \tau}, \nu_{\sigma, \tau}) = \pi^{-1}(\sigma, \tau).$$

Let $\delta_j > 0$ and let

$$\mathcal{U} \subset \{J \in \mathcal{J}(X) : \|J - J_0\|_{C^1} < \delta_j\}$$

be such that for every $k \in \mathbb{N}$,

$$\sup_{J \in \mathcal{U}} \|J - J_0\|_{C^k} < \infty.$$

In the upcoming discussion we may implicitly shrink Δ and δ_j , in order to ensure that Hypothesis 5.5 and Hypothesis 5.24 hold and various expressions involving $|\sigma|$, $\varepsilon := \max\{\varepsilon_n : n \in S\}$ with $\varepsilon_n := |\tau_n|$, and $\|J - J_0\|_{C^1}$ are sufficiently small.

The purpose of this subsection is to analyze whether u_0 can be slightly deformed to a J -holomorphic map $u_{\sigma, \tau}: (\Sigma_{\sigma, \tau}, j_{\sigma, \tau}, \nu_{\sigma, \tau}) \rightarrow (X, J)$ with $J \in \mathcal{U}$. More precisely, we show that a Gromov neighborhood of u_0 in the space of nodal J -holomorphic maps with $J \in \mathcal{U}$ is homeomorphic to the zero set of a continuous map

$$\text{ob}: \Delta \times \mathcal{U} \times \mathcal{J} \rightarrow \mathcal{O},$$

where \mathcal{J} an open subset of the deformation space $\ker \delta_{u_0, J_0}$ and $\mathcal{O} \cong \text{coker } \delta_{u_0, J_0}$ is the obstruction space. This is a local Kuranishi model at u_0 for the universal moduli space of pseudo-holomorphic nodal maps.

Remark 5.28. Since we are not interested here in the global properties of the universal moduli space, we do not require that u_0 is a stable map. A local Kuranishi model can be constructed around any pseudo-holomorphic map. However, the Gromov limit, as defined in Definition 3.8, is not necessarily unique for unstable maps, so the universal moduli space of all nodal pseudo-holomorphic maps is not a Hausdorff space.

To facilitate the discussion in Section 5.7 (and although it makes the present discussion somewhat more awkward than it needs to be) the construction of the Kuranishi model proceeds in two steps. Choose a partition

$$S = S_1 \amalg S_2 \quad \text{with } \nu_0(S_1) = S_1 \text{ and } \nu_0(S_2) = S_2$$

and write every smoothing parameter τ as

$$\tau = (\tau_1, \tau_2) \quad \text{with } \tau_1 = (\tau_{1,n})_{n \in S_1} \text{ and } \tau_2 = (\tau_{2,n})_{n \in S_2}.$$

The first step of our construction varies σ and τ_1 but $\tau_2 = 0$ is fixed. The second step holds σ and τ_1 fixed and varies τ_2 .

Denote by $u_{\sigma,0}: \Sigma_{\sigma,0} \rightarrow X$ the smooth map underlying u_0 . Denote by

$$\delta_{u_0;J_0}: W^{1,p}\Gamma(\Sigma_0, \nu_0; u_0^*TX) \rightarrow L^p\Omega^{0,1}(\Sigma_0, u_0^*TX)$$

the linear operator associated with u_0 defined in Definition 2.24. Let

$$\mathcal{O} \subset \Omega^{0,1}(\Sigma_0, u_0^*TX) \subset L^p\Omega^{0,1}(\Sigma_0, u_0^*TX)$$

be a lift of coker $\delta_{u_0;J_0}$; that is, $\dim \mathcal{O} = \dim \text{coker } \delta_{u_0}$ and (5.17) holds. We will assume that all 1-forms in \mathcal{O} are smooth on each component of Σ_0 . (The canonical choice is $\mathcal{O} = \ker \delta_{u_0;J_0}^*$, but this choice is not always the most convenient.) Let Δ_1 parametrize complex structures on (Σ_0, ν_0) as at the beginning of Section 4.3, and let \mathcal{U} be an open neighborhood of J_0 as above. Trivialize the bundle over $\Delta_1 \times \mathcal{U}$ whose fiber over $(\sigma, J) \in \Delta_1 \times \mathcal{U}$ is $\Omega^{0,1}(\Sigma_{\sigma,0}, u_{\sigma,0}^*TX)$ with the $(0, 1)$ -part taken with respect to $j_{\sigma,0}$ and J . This identifies $\Omega^{0,1}(\Sigma_0, u_0^*TX)$ and $\Omega^{0,1}(\Sigma_{\sigma,0}, u_{\sigma,0}^*TX)$, and thus exhibits \mathcal{O} as a subset of $L^p\Omega^{0,1}(\Sigma_{\sigma,0}, u_{\sigma,0}^*TX)$ for which (5.17) holds for $\delta_{\tilde{u}_{\sigma,0};J}$ instead of δ_{u_0} . Define

$$\bar{\delta}_{\tilde{u}_{\sigma,\tau_1,0};J}: W^{1,p}\Gamma(\Sigma_{\sigma,\tau_1,0}, \nu_{\sigma,0}; \tilde{u}_{\sigma,\tau_1,0}^*TX) \oplus \mathcal{O} \rightarrow L^p\Omega^{0,1}(\Sigma_{\sigma,\tau_1,0}, \tilde{u}_{\sigma,\tau_1,0}^*TX)$$

as in Definition 5.19. The construction in Section 5.4 yields a right inverse

$$r_{\tilde{u}_{\sigma,\tau_1,0};J}: L^p\Omega^{0,1}(\Sigma_{\sigma,\tau_1,0}, \tilde{u}_{\sigma,\tau_1,0}^*TX) \rightarrow W^{1,p}\Gamma(\Sigma_{\sigma,\tau_1,0}, \nu_{\sigma,\tau_1,0}; \tilde{u}_{\sigma,\tau_1,0}^*TX) \oplus \mathcal{O}.$$

of $\bar{\delta}_{\tilde{u}_{\sigma,\tau_1,0};J}$.

The next two propositions build a Kuranishi model for a neighborhood of u_0 in the Gromov compactification of the moduli space of pseudo-holomorphic maps. In essence, they assert that for every smoothing parameter τ and an infinitesimal deformation κ the pseudo-holomorphic map equation can be solved modulo obstructions. The construction of the Kuranishi model proceeds in two steps, described by Proposition 5.29 and Proposition 5.33, in order to obtain better control of the obstruction map. The first step is to smooth the nodes in S_1 .

Proposition 5.29. *There are constants $\delta_\kappa, \Lambda > 0$ such that for every $(\sigma, \tau_1, 0) \in \Delta$ and $\kappa \in \ker \delta_{u_0}$ with $|\kappa| < \delta_\kappa$, there exists a unique pair*

$$\begin{aligned} (\xi(\sigma, \tau_1; J; \kappa), o(\sigma, \tau_1; J; \kappa)) &\in \text{im } r_{\tilde{u}_{\sigma,\tau_1,0};J} \\ &\subset W^{1,p}\Gamma(\Sigma_{\sigma,\tau_1,0}, \nu_{\sigma,\tau_1,0}; \tilde{u}_{\sigma,\tau_1,0}^*TX) \oplus \mathcal{O} \end{aligned}$$

with

$$\|\xi(\sigma, \tau_1; J; \kappa)\|_{W^{1,p}} + |o(\sigma, \tau_1; J; \kappa)| \leq \Lambda$$

satisfying

$$\mathfrak{F}\tilde{u}_{\sigma,\tau_1,0;J}(\text{fuse}_{\tau_1,0}\kappa + \xi(\sigma, \tau_1; J; \kappa)) + \text{pull}_{\tau_1,0}(o(\sigma, \tau_1; J; \kappa)) = 0, \tag{5.30}$$

with \mathfrak{F} as in Definition 2.23. Furthermore,

$$\begin{aligned} & \|\xi(\sigma, \tau_1; J; \kappa)\|_{W^{1,p}} + |o(\sigma, \tau_1; J; \kappa)| \\ & \leq c(|\sigma| + |\tau_1|^{\frac{1}{2} + \frac{1}{p}} + \|J - J_0\|_{C^0} + |\kappa|). \end{aligned} \tag{5.31}$$

Proof. Since $r_{\tilde{u}_{\sigma,\tau_1,0;J}}$ is injective, (5.30) is equivalent to the fixed-point equation

$$\eta = \mathbf{F}(\eta) := \eta - \mathfrak{F}\tilde{u}_{\sigma,\tau_1,0;J}(\text{fuse}_{\tau_1,0}\kappa + \text{pr}_1 r_{\tilde{u}_{\sigma,\tau_1,0;J}}\eta) - \text{pull}_{\tau_1,0}(\text{pr}_2 r_{\tilde{u}_{\sigma,\tau_1,0;J}}\eta).$$

Here pr_1 and pr_2 denote the projections to the first and second summand of

$$W^{1,p}\Gamma(\Sigma_{\sigma,\tau_1,0}, \nu_{\sigma,\tau_1,0}; \tilde{u}_{\sigma,\tau_1,0}^*TX) \oplus \mathcal{O},$$

respectively. By Proposition 2.30,

$$\begin{aligned} \mathbf{F}(\eta) = & -\bar{\partial}_J(\tilde{u}_{\sigma,\tau_1,0}, j_{\sigma,\tau_1,0}) - \delta_{\tilde{u}_{\sigma,\tau_1,0;J}}\text{fuse}_{\tau_1,0}\kappa \\ & - \pi_{\tilde{u}_{\sigma,\tau_1,0;J}}(\text{fuse}_{\tau_1,0}\kappa + \text{pr}_1 \circ r_{\tilde{u}_{\sigma,\tau_1,0;J}}\eta). \end{aligned}$$

By Proposition 5.9 and Proposition 2.30,

$$\|\mathbf{F}(0)\|_{L^p} \leq c(|\sigma| + \|J - J_0\|_{C^0} + |\tau_1|^{\frac{1}{2} + \frac{1}{p}} + |\kappa| + |\kappa|^2).$$

Moreover, by Proposition 5.26 and Proposition 2.30,

$$\|\mathbf{F}(\eta_1) - \mathbf{F}(\eta_2)\|_{L^p} \leq c(|\kappa| + \|\eta_1\|_{L^p} + \|\eta_2\|_{L^p})\|\eta_1 - \eta_2\|_{L^p}.$$

Therefore, provided δ_κ is sufficiently small, there is an $R > 0$ such that $\|\mathbf{F}(0)\|_{L^p} \leq R/2$ and for every $\eta_1, \eta_2 \in \bar{B}_R(0) \subset L^p\Omega^{0,1}(\Sigma_{\sigma,\tau}, \tilde{u}_{\sigma,\tau}^*TX)$,

$$\|\mathbf{F}(\eta_1) - \mathbf{F}(\eta_2)\|_{L^p} \leq \frac{1}{2}\|\eta_1 - \eta_2\|_{L^p}.$$

This shows that \mathbf{F} maps $\bar{B}_R(0)$ into $\bar{B}_R(0)$ and $\mathbf{F}: \bar{B}_R(0) \rightarrow \bar{B}_R(0)$ is a contraction. Thus, the first assertion follows from Banach’s fixed-point theorem. The second follows from the above and Proposition 5.9. ■

This completes the first step. In the second step, we smooth the nodes in S_2 . This step is analogous to the first one, with u_0 being replaced by the maps obtained from Proposition 5.29. For $(\sigma, \tau) \in \Delta$ and $\kappa \in \ker \delta_{u_0}$ with $\|\kappa\|_{W^{1,p}} < \delta_\kappa$ set

$$u_{\sigma,\tau_1,0;J;\kappa} := \exp_{\tilde{u}_{\sigma,\tau_1,0}}(\text{fuse}_{\tau_1,0}\kappa + \xi(\sigma, \tau_1; J; \kappa)), \quad \tilde{u}_{\sigma,\tau;J;\kappa} := \widetilde{(u_{\sigma,\tau_1,0;J;\kappa})_\tau};$$

that is, $\tilde{u}_{\sigma,\tau;J;\kappa}$ is obtained from $u_{\sigma,\tau_1,0;J;\kappa}$ by the construction in Definition 5.7.

Definition 5.32. Define

$$\text{pull}_{\sigma, \tau_1, 0; J; \kappa}: L^p \Omega^{0,1}(\Sigma_0, u_0^* TX) \rightarrow L^p \Omega^{0,1}(\Sigma_{\sigma, \tau_1, 0}, u_{\sigma, \tau_1, 0}^* TX)$$

to be the composition of $\text{pull}_{\tau_1, 0}$ with the map induced by parallel transport along the geodesics

$$t \mapsto \exp_{\tilde{u}_{\sigma, \tau_1, 0}}(t(\text{fuse}_{\tau_1, 0} \kappa + \xi(\sigma, \tau_1; J; \kappa))).$$

Furthermore, denote by

$$\text{pull}_{\sigma, \tau; J; \kappa}: L^p \Omega^{0,1}(\Sigma_0, u_0^* TX) \rightarrow L^p \Omega^{0,1}(\Sigma_{\sigma, \tau}, \tilde{u}_{\sigma, \tau}^* TX)$$

the composition of $\text{pull}_{\sigma, \tau_1, 0; J; \kappa}$ with

$$\text{pull}_{\tau_2}: L^p \Omega^{0,1}(\Sigma_{\sigma, \tau_1, 0}, u_{\sigma, \tau_1, 0}^* TX) \rightarrow L^p \Omega^{0,1}(\Sigma_{\sigma, \tau}, \tilde{u}_{\sigma, \tau}^* TX)$$

defined in Definition 5.18.

The subspace $\text{pull}_{\sigma, \tau_1, 0; J; \kappa}(\mathcal{O})$ satisfies (5.17) for $u_{\sigma, \tau_1, 0; J; \kappa}$ instead of u_0 . Define

$$\bar{\delta}_{\tilde{u}_{\sigma, \tau; J; \kappa}}: W^{1,p} \Gamma(\Sigma_{\sigma, \tau}, \nu_{\sigma, \tau}; \tilde{u}_{\sigma, \tau}^* TX) \oplus \mathcal{O} \rightarrow L^p \Omega^{0,1}(\Sigma_{\sigma, \tau}, \tilde{u}_{\sigma, \tau}^* TX)$$

as in Definition 5.19. The construction in Section 5.4 yields a right inverse

$$r_{\tilde{u}_{\sigma, \tau; J; \kappa}}: L^p \Omega^{0,1}(\Sigma_{\sigma, \tau}, \tilde{u}_{\sigma, \tau}^* TX) \rightarrow W^{1,p} \Gamma(\Sigma_{\sigma, \tau}, \nu_{\sigma, \tau}; \tilde{u}_{\sigma, \tau}^* TX) \oplus \mathcal{O}$$

of $\bar{\delta}_{\tilde{u}_{\sigma, \tau; J; \kappa}}$.

Proposition 5.33. *There are constants $\delta_\kappa, \Lambda > 0$ such that for every $(\sigma, \tau; J) \in \Delta \times \mathcal{U}$ and $\kappa \in \ker \delta_{u_0}$ with $\|\kappa\|_{W^{1,p}} < \delta_\kappa$, there exists a unique pair*

$$(\hat{\xi}(\sigma, \tau; J; \kappa), \hat{\delta}(\sigma, \tau; J; \kappa)) \in \text{im } r_{\tilde{u}_{\sigma, \tau; J; \kappa}} \subset W^{1,p} \Gamma(\Sigma_{\sigma, \tau}, \nu_{\sigma, \tau}; \tilde{u}_{\sigma, \tau}^* TX) \oplus \mathcal{O}$$

with

$$\|\hat{\xi}(\sigma, \tau; J; \kappa)\|_{W^{1,p}} + |\hat{\delta}(\sigma, \tau; J; \kappa)| \leq \Lambda$$

satisfying

$$\mathfrak{F}_{\tilde{u}_{\sigma, \tau; J; \kappa}}(\hat{\xi}(\sigma, \tau; J; \kappa)) + \text{pull}_{\sigma, \tau; J; \kappa}(o(\sigma, \tau_1; J; \kappa) + \hat{\delta}(\sigma, \tau; J; \kappa)) = 0.$$

Furthermore,

$$\begin{aligned} & \|\hat{\xi}(\sigma, \tau; J; \kappa)\|_{W^{1,p}} + |\hat{\delta}(\sigma, \tau; J; \kappa)| \\ & \leq c \|\bar{\partial}_J(\tilde{u}_{\sigma, \tau; J; \kappa}, j_{\sigma, \tau}) + \text{pull}_{\sigma, \tau; J; \kappa}(o(\sigma, \tau_1; J; \kappa))\|_{L^p}. \end{aligned} \tag{5.34}$$

Proof. This is similar to the proof of Proposition 5.29. ■

Definition 5.35. Set $\mathcal{J} := B_{\delta_\kappa}(0) \subset \ker \delta_{u_0}$. The *Kuranishi map* $\text{ob}: \Delta \times \mathcal{U} \times \mathcal{J} \rightarrow \mathcal{O}$ is defined by

$$\text{ob}(\sigma, \tau; J; \kappa) := o(\sigma, \tau_1; J; \kappa) + \widehat{o}(\sigma, \tau; J; \kappa),$$

with o and \widehat{o} as in Proposition 5.29 and Proposition 5.33.

The upshot of the preceding discussion is that u_0 can be slightly deformed to a J -holomorphic map

$$u_{\sigma, \tau}: (\Sigma_{\sigma, \tau}, j_{\sigma, \tau}, \nu_{\sigma, \tau}) \rightarrow (X, J)$$

if and only if there is a $\kappa \in \mathcal{J}$ with $\text{ob}(\sigma, \tau; J; \kappa) = 0$. The following shows that this Kuranishi model indeed describes a Gromov neighborhood of $u_0: (\Sigma_0, j_0, \nu_0) \rightarrow (X, J_0)$.

Proposition 5.36. *Let $(\sigma_k, \tau_k)_{k \in \mathbb{N}}$ be a sequence in Δ converging to $(0, 0)$ and let $(J_k)_{k \in \mathbb{N}}$ be a sequence in \mathcal{U} converging to J_0 . If*

$$(u_k: (\Sigma_{\sigma_k, \tau_k}, j_{\sigma_k, \tau_k}, \nu_{\sigma_k, \tau_k}) \rightarrow (X, J_k))_{k \in \mathbb{N}}$$

is a sequence of nodal pseudo-holomorphic maps which Gromov converges to

$$u_0: (\Sigma_0, j_0, \nu_0) \rightarrow (X, J_0),$$

then there is a $K \in \mathbb{N}$ such that for every $k \geq K$ there are $\kappa_k \in \ker \delta_{u_0}$ and $(\xi_k, 0) \in \text{im } \tilde{r}_{\tilde{u}_{\sigma_k, \tau_k; \kappa_k}; J_k}$ with

$$u_k = \exp_{\tilde{u}_{\sigma_k, \tau_k; \kappa_k}}(\xi_k);$$

moreover,

$$\lim_{k \rightarrow \infty} |\kappa_k| = 0 \quad \text{and} \quad \lim_{k \rightarrow \infty} \|\xi_k\|_{W^{1,p}} = 0.$$

In particular,

$$\text{ob}(\sigma_k, \tau_k; J_k; \kappa_k) = 0.$$

The proof of this proposition relies on the following result.

Proposition 5.37. *Assume the situation of Proposition 5.36. There are $K \in \mathbb{N}$, $\delta_\kappa > 0$, and $c > 0$ such that for every $k \geq K$ and $\kappa \in \ker \delta_{u_0}$ with $\|\kappa\|_{W^{1,p}} < \delta_\kappa$, there is a $\zeta_{k; \kappa} \in \Gamma(\Sigma_{\sigma_k, \tau_k}, \tilde{u}_{\sigma_k, \tau_k; J_k}^* TX)$ with*

$$u_k = \exp_{\tilde{u}_{\sigma_k, \tau_k; J_k; \kappa}}(\zeta_{k; \kappa});$$

moreover,

$$\limsup_{k \rightarrow \infty} \|\zeta_{k; \kappa}\|_{W^{1,p}} \leq c|\kappa|.$$

Proof. The proof has two steps: the construction of $\zeta_{k; \kappa}$ and the proof of the estimate. Recall the definition of U_x from Definition 5.4.

Step 1. There are $K \in \mathbb{N}$ and $\delta_\kappa > 0$ such that for every $k \geq K$, $\kappa \in \ker \delta_{u_0}$ with $\|\kappa\|_{W^{1,p}} < \delta_\kappa$ and $z \in \Sigma_{\sigma_k, \tau_k}$,

$$u_k(z) \in U_{\tilde{u}_{\sigma_k, \tau_k; J_k; \kappa}(z)};$$

in particular, there is a section $\zeta_{k; \kappa} \in \Gamma(\Sigma_{\sigma_k, \tau_k}, \tilde{u}_{\sigma_k, \tau_k; J_k}^* TX)$ given by

$$\zeta_{k; \kappa} := \exp_{\tilde{u}_{\sigma_k, \tau_k; J_k; \kappa}}^{-1} \circ u_k.$$

By (5.31), (5.34), and Proposition 5.9,

$$d(\tilde{u}_{\sigma_k, \tau_k; J_k; \kappa}, \tilde{u}_{\sigma_k, \tau_k; J_k; 0}) \leq c(|\sigma_k| + \varepsilon_k^{\frac{1}{2} + \frac{1}{p}} + \|J_k - J_0\|_{C^0} + |\kappa|). \tag{5.38}$$

Therefore, it suffices to consider $\kappa = 0$ and prove that there exists a $K \in \mathbb{N}$ such that for every $k \geq K$,

$$u_k(z) \in \frac{1}{2} U_{\tilde{u}_{\sigma_k, \tau_k; J_k; 0}(z)}.$$

Using the framing Ψ from Definition 4.17, define $v_k: \Sigma_0 \setminus S \rightarrow X$, $\tilde{v}_{k; \kappa}: \Sigma_0 \setminus S \rightarrow X$ by

$$\begin{aligned} v_k &:= u_k \circ \iota_k^{-1} \circ \Psi(\cdot; \sigma_k, \tau_k) \circ \iota_0, \\ \tilde{v}_k &:= \tilde{u}_{\sigma_k, \tau_k; J_k; 0} \circ \iota_k^{-1} \circ \Psi(\cdot; \sigma_k, \tau_k) \circ \iota_0, \end{aligned}$$

respectively, cf. Definition 3.8. Both of the sequences $(v_k)_{k \in \mathbb{N}}$ and $(\tilde{v}_k)_{k \in \mathbb{N}}$ converge to $u_0: \Sigma_0 \setminus S \rightarrow X$ in the C_{loc}^∞ topology – the former by Definition 3.8 and the latter by construction.

With the notation of Remark 4.18 for $r > 0$ and $n \in S$, set

$$N_{k,n}^r := N_{\sigma_k, \tau_k; n}^r.$$

Choose $r > 0$ as in Proposition 3.16 with $\delta := \frac{1}{8} \text{inj}_g(X)$. By the preceding paragraph, the assertion holds for sufficiently large k and $z \notin N_{k,n}^r$. By Proposition 3.16 and by construction of $\tilde{u}_{\sigma, \tau}$, for sufficiently large k

$$u_k(N_{k,n}^r) \subset B_\delta(u_0(n)) \quad \text{and} \quad \tilde{u}_{\sigma_k, \tau_k; J_k; 0}(N_{k,n}^r) \subset B_\delta(u_0(n));$$

hence, for every $z \in N_{k,n}^r$

$$u_k(z) \in \frac{1}{2} U_{\tilde{u}_{\sigma_k, \tau_k; J_k; 0}(z)}.$$

Step 2. There is a constant $c > 0$ such that the sections $\zeta_{k; \kappa}$ defined in the preceding step satisfy

$$\limsup_{k \rightarrow \infty} \|\zeta_{k; \kappa}\|_{W^{1,p}} \leq c|\kappa|.$$

By (5.31), (5.34), and Proposition 5.9, we can restrict to $\kappa = 0$. Furthermore, it suffices to prove that for every $n \in S$

$$\lim_{s \downarrow 0} \limsup_{k \rightarrow \infty} \|\zeta_{k;0}\|_{W^{1,p}(N_{k;n}^s)} = 0. \tag{5.39}$$

The case when n is not smoothed is straightforward. The framing extends to identify a neighborhood of n in $\Sigma_{\sigma_k, \tau_k}$ with a neighborhood of n in Σ_0 . It follows from Lemma 3.14 and elliptic regularity that, on this subset, the maps u_k converge to u_0 in the C_{loc}^∞ topology. Let us therefore assume that n is smoothed out; that is, $\varepsilon_{k;n} \neq 0$ for sufficiently large k .

Define $\rho_k \in C^\infty(N_{k;n}^r, T_{u_0(n)}X)$ and $\tilde{\rho}_k \in C^\infty(N_{k;n}^r, T_{u_0(n)}X)$ by

$$\rho_k := \exp_{u_0(n)}^{-1} \circ u_k \quad \text{and} \quad \tilde{\rho}_k := \exp_{u_0(n)}^{-1} \circ \tilde{u}_{\sigma_k, \tau_k; J_k; 0}.$$

By construction,

$$\lim_{k \rightarrow \infty} \|\tilde{\rho}_k\|_{W^{1,p}} = 0.$$

Therefore, it suffices to prove that

$$\lim_{s \downarrow 0} \limsup_{k \rightarrow \infty} \|\rho_k\|_{W^{1,p}(N_{k;n}^s)} = 0.$$

As explained in Remark 4.18, the subset $N_{k;n}^r$ is biholomorphic to the cylinder

$$S^1 \times (-L_k, L_k) \quad \text{with} \quad L_k := \log(r\varepsilon_{n;k}^{-1/2}).$$

Hence, ρ_k can be thought of as a map $\rho_k^{\text{cyl}}: S^1 \times (-L_k, L_k) \rightarrow T_{u_0(n)}X$. More concretely, the canonical chart ϕ_n defines a holomorphic embedding

$$\phi_n: \{v \in T_n \Sigma_0 : \varepsilon_{n;k}^{1/2} \leq |v| < r\} \rightarrow N_{k;n}^r$$

which glues via ι_τ with the embedding $\phi_{v(n)}$ to a biholomorphic map

$$B_r(0) \setminus \bar{B}_{\varepsilon_{n;k}/r}(0) \cong N_{k;n}^r.$$

Choose identifications $T_n \Sigma_0 \cong \mathbf{C} \cong T_{v(n)} \Sigma_0$ such that $\iota_\tau(z) = \varepsilon_n/z$. The map ρ_k^{cyl} is then defined by

$$\rho_k^{\text{cyl}}(\theta, \ell) := \begin{cases} \rho_k \circ \phi_n(\varepsilon_{n;k}^{1/2} e^{\ell+i\theta}) & \text{if } t \geq 0, \\ \rho_k \circ \phi_{v(n)}(\varepsilon_{n;k}^{1/2} e^{-\ell-i\theta}) & \text{if } t \leq 0. \end{cases}$$

Since u_k is J_k -holomorphic, ρ_k^{cyl} is $\exp_{u_0(n)}^*(J_k)$ -holomorphic. Since the energy is conformally invariant,

$$E(\rho_k^{\text{cyl}}) = E(u_k|_{N_{k;n}^r}).$$

Choose $\mu \in (1 - 2/p, 1)$. By Lemma 3.15,

$$|\nabla \rho_k^{\text{cyl}}(\theta, \ell)| \leq c e^{-\mu(L_k - |\ell|)} E(u_k |_{N_{k;n}^r})^{1/2}.$$

By the above and Proposition 5.3, for $z \in \Sigma_{\sigma, \tau}^\circ$ with $r_n(z) < r$

$$|\nabla \rho_k(z)| \leq c r^{-\mu} r_n(z)^{\mu-1} E(u_k |_{N_{k;n}^r})^{1/2}.$$

There is a corresponding estimate with n replaced with $\nu(n)$. Hence,

$$\|\nabla \rho_k(z)\|_{L^p(N_{k;n}^s)}^p \leq \frac{c r^{-\mu p}}{(\mu - 1)p + 2} s^{(\mu-1)p+2} E(u_k |_{N_{k;n}^r})^{p/2}.$$

Since $(\mu - 1)p + 2 > 0$, the right-hand converges to zero as s converges to zero. ■

Proof of Proposition 5.36. Let $k \geq K$ and $\kappa \in \ker \delta_{u_0}$ with $|\kappa| < \delta_\kappa$. Let $\zeta_{k;\kappa}$ be as in Proposition 5.37. By Proposition 5.27 the latter can be uniquely written as

$$\zeta_{k,\kappa} = \text{fuse}_{\tau_k}(\lambda_{k;\kappa}) + \text{pr}_1 \circ r \tilde{u}_{\sigma_k, \tau_k; J_k; \kappa} \eta_{k;\kappa} \quad \text{with } \lambda_{k;\kappa} \in \ker \delta_{u_0}.$$

It remains to be proved that after possibly increasing K for every $k \geq K$ there exists a $\kappa \in \ker \delta_{u_0}$ with $|\kappa| < \delta_\kappa$ and $\lambda_{k;\kappa} = 0$. The following statement is a consequence of (5.38), (5.39), and the fact that $r \tilde{u}_{\sigma_k, \tau_k; J_k; \kappa}$ depends smoothly on κ when interpreted as a family of operators on a fixed Banach space $L^p \Omega^{0,1}(\Sigma_{\sigma_k, \tau_k}, \tilde{u}_{\sigma_k, \tau_k; J_k; 0}^* TX) \oplus \mathcal{O}$ using parallel transport along geodesics. If δ_κ is sufficiently small, then for every κ , u_k can be written in the form

$$u_k = \exp_{\tilde{u}_{\sigma_k, \tau_k; J_k; 0}}(\text{fuse}_{\tau_k}(\kappa + \lambda_{k;\kappa}) + r \tilde{u}_{\sigma_k, \tau_k; J_k; 0} \hat{\eta}_{k;\kappa} + e_{k;\kappa}),$$

with $e_{k;\kappa}$ satisfying

$$\limsup_{k \rightarrow \infty} \|e_{k;0}\|_{W^{1,p}} = 0$$

and a quadratic estimate

$$\|e_{k;\kappa_1} - e_{k;\kappa_2}\|_{W^{1,p}} \leq c(|\kappa_1| + |\kappa_2|)|\kappa_1 - \kappa_2| \tag{5.40}$$

It follows from Proposition 5.27 that for $|\kappa| \leq \delta_\kappa$,

$$\kappa + \lambda_{k;\kappa} + \pi(e_{k;\kappa}) = 0,$$

where π denotes the projection on $\text{fuse}_{\tau_k}(\ker \delta_{u_0})$ precomposed $\text{fuse}_{\tau_k}^{-1}$; the latter is defined because fuse_{τ_k} is injective on $\ker \delta_{u_0}$ provided k is sufficiently large. Thus, the existence of a unique small κ such that $\lambda_{k;\kappa} = 0$ is a consequence of (5.40) and the Banach fixed point theorem applied to the map $\kappa \mapsto -\pi(e_{k;\kappa})$. ■

5.7. The leading order term of the obstruction on ghost components

Assume the situation of Section 5.6. The purpose of this subsection is to analyze the leading order term of part of the obstruction map ob constructed in Section 5.6. This construction requires a choice of partition of S and a choice of lift

$$\mathcal{O} \subset L^p \Omega^{0,1}(\Sigma_0, u_0^* TX)$$

of coker δ_{u_0} . The following paragraphs introduce a particular choice tailored to the upcoming discussion.

Let $C \subset \Sigma_0$ be a ghost component of u_0 ; see Section 2.4 for the definitions of a ghost component and related notation. Denote by $x_0 \in X$ the constant value which u_0 takes on C .

To simplify the upcoming discussion, we will make the following assumption, which will be satisfied in the situation considered in the proof of Theorem 1.1.

Hypothesis 5.41. S_C^{ext} consists of one point, that is, C and $\Sigma_0 \setminus C$ meet at one node.

Denote by $B \subset C$ the base-locus of the dualizing sheaf of \check{C} , cf. Proposition 2.18. If B does not contain the node at which C and $\Sigma_0 \setminus C$ meet, then set $B_0 := \emptyset$; otherwise, denote by B_0 the connected component of B containing the node. Set

$$C_\bullet := C \setminus B_0 \quad \text{and} \quad \Sigma_\clubsuit := \Sigma_0 \setminus C_\bullet$$

and abbreviate

$$\nu_\bullet := \nu_{C_\bullet} \quad \text{and} \quad \nu_\clubsuit := \nu_{\Sigma_\clubsuit}.$$

The significance of this construction is as follows. By Proposition 2.18, every connected component of C_\bullet attaches to Σ_\clubsuit at a unique node; moreover: these nodes are not contained in the base-locus of the dualizing sheaf of C_\bullet .

The partition of the set of nodes we choose is, with the notation from Section 2.4,

$$S = S_1 \amalg S_2 \quad \text{with} \quad S_1 := S_{\Sigma_\clubsuit}^{\text{int}} \amalg S_{C_\bullet}^{\text{int}} \quad \text{and} \quad S_2 := S_{\Sigma_\clubsuit}^{\text{ext}} \amalg S_{C_\bullet}^{\text{ext}}, \tag{5.42}$$

that is, first, we will smooth the interior nodes of Σ_\clubsuit and C_\bullet and then the exterior nodes connecting them.

Next we discuss the choice of the obstruction space \mathcal{O} . Set $\delta_{u_0, \clubsuit} := \delta_{u_0|_{\Sigma_\clubsuit}}$ and let

$$\mathcal{O}_\clubsuit \subset \Omega^{0,1}(\Sigma_\clubsuit, u_0^* TX) \subset L^p \Omega^{0,1}(\Sigma_\clubsuit, u_0^* TX)$$

be a lift of coker $\delta_{u_0, \clubsuit}$ such that every $o \in \mathcal{O}_\clubsuit$ vanishes in a neighborhood of $S_{\Sigma_\clubsuit}^{\text{ext}}$ if B_0 is empty, and over all of B_0 if B_0 is non-empty. Furthermore, let

$$\mathcal{O}_\bullet \subset \Omega^{0,1}(C_\bullet, \mathbb{C}) \otimes_{\mathbb{C}} T_{x_0} X \subset L^p \Omega^{0,1}(C_\bullet, u_0^* TX)$$

be a lift of coker $(\bar{\partial} \otimes_{\mathbb{C}} \mathbf{1})$.

Every $\xi \in W^{1,p}\Gamma(\Sigma_{\clubsuit}, \nu_{\clubsuit}; u_0^*TX)$ can be extended to Σ_0 in the following way. Given $n \in S_{\Sigma_{\clubsuit}}^{\text{ext}}$, extend ξ to a constant section taking value $\xi(n)$ over the connected component of the nodal curve $(C_{\bullet}, \nu_{\bullet})$ containing $\nu(n)$. This defines an inclusion

$$W^{1,p}\Gamma(\Sigma_{\clubsuit}, \nu_{\clubsuit}; u_0^*TX) \subset W^{1,p}\Gamma(\Sigma_0, \nu_0; u_0^*TX). \tag{5.43}$$

Furthermore, extension by zero defines inclusions

$$\begin{aligned} L^p\Omega^{0,1}(\Sigma_{\clubsuit}, u_0^*TX) &\subset L^p\Omega^{0,1}(\Sigma_0, u_0^*TX), \\ L^p\Omega^{0,1}(C_{\bullet}, u_0^*TX) &\subset L^p\Omega^{0,1}(\Sigma_0, u_0^*TX). \end{aligned}$$

Set $\mathcal{O} := \mathcal{O}_{\clubsuit} \oplus \mathcal{O}_{\bullet}$.

Proposition 5.44. *The map (5.43) induces an isomorphism $\ker \delta_{u_0, \clubsuit} \cong \ker \delta_{u_0}$ and \mathcal{O} is a lift of coker δ_{u_0} .*

Proof. Denote by ν_{\amalg} the nodal structure on Σ_0 which agrees with ν_0 on the complement of S_2 and is the identity on S_2 . This nodal structure disconnects Σ_{\clubsuit} and C_{\bullet} . Denote by

$$\delta_{u_0, \amalg}: W^{1,p}\Gamma(\Sigma_0, \nu_{\amalg}; u_0^*TX) \rightarrow L^p\Omega^{0,1}(\Sigma_0, u_0^*TX)$$

the operator induced by δ_{u_0} . Define V_- and $\text{diff}: \ker \delta_{u_0, \amalg} \rightarrow V_-$ as in Remark 2.28 with S_2 instead of S . As is explained in Remark 2.28,

$$\ker \delta_{u_0} = \ker \text{diff}$$

and there is a short exact sequence

$$0 \longrightarrow \text{coker diff} \longrightarrow \text{coker } \delta_{u_0} \longrightarrow \text{coker } \delta_{u_0, \amalg} \longrightarrow 0.$$

The domain and codomain of $\delta_{u_0, \amalg}$ decompose as

$$\begin{aligned} W^{1,p}\Gamma(\Sigma_0, \nu_{\amalg}; u_0^*TX) &= W^{1,p}\Gamma(\Sigma_{\clubsuit}, \nu_{\clubsuit}; u_0^*TX) \oplus W^{1,p}\Gamma(C_{\bullet}, \nu_{\bullet}; \mathbf{C}) \otimes_{\mathbf{C}} T_{x_0}X, \\ L^p\Omega^{0,1}(\Sigma_0, u_0^*TX) &= L^p\Omega^{0,1}(\Sigma_{\clubsuit}, u_0^*TX) \oplus L^p\Omega^{0,1}(C_{\bullet}, \mathbf{C}) \otimes_{\mathbf{C}} T_{x_0}X. \end{aligned}$$

With respect to these decompositions

$$\delta_{u_0, \amalg} = \begin{pmatrix} \delta_{u_0, \clubsuit} & 0 \\ 0 & \bar{\partial} \otimes_{\mathbf{C}} \mathbf{1} \end{pmatrix}$$

with $\delta_{u_0, \clubsuit} = \delta_{u_0|_{\Sigma_{\clubsuit}}}$ and $\bar{\partial} \otimes_{\mathbf{C}} \mathbf{1} = \delta_{u_0|_{C_{\bullet}}}$ is the standard Cauchy–Riemann operator since u_0 is constant on C_{\bullet} . Therefore,

$$\begin{aligned} \ker \delta_{u_0, \amalg} &= \ker \delta_{u_0, \clubsuit} \oplus \ker(\bar{\partial} \otimes_{\mathbf{C}} \mathbf{1}), \\ \text{coker } \delta_{u_0, \amalg} &= \text{coker } \delta_{u_0, \clubsuit} \oplus \text{coker}(\bar{\partial} \otimes_{\mathbf{C}} \mathbf{1}). \end{aligned}$$

The task at hand is to understand $\ker \delta_{u_0}$ and $\text{coker } \delta_{u_0}$ in terms of the above.

Since elements of $\ker(\bar{\partial} \otimes_{\mathbf{C}} \mathbf{1})$ are locally constant, $\ker(\bar{\partial} \otimes_{\mathbf{C}} \mathbf{1})$ has a direct summand $T_{x_0}X$ for every connected component of $(C_{\bullet}, \nu_{\bullet})$. Hypothesis 5.41 and Proposition 2.18 imply that there is one connected component for each node in $S_{\Sigma_{\clubsuit}}^{\text{ext}}$. Therefore,

$$\ker(\bar{\partial} \otimes_{\mathbf{C}} \mathbf{1}) = V_- = \text{Map}(S_{\Sigma_{\clubsuit}}^{\text{ext}}, T_{x_0}X).$$

With respect to this identification the map $\text{diff}: \ker \delta_{u_0, \clubsuit} \oplus \ker(\bar{\partial} \otimes_{\mathbf{C}} \mathbf{1}) \rightarrow V_-$ is given by

$$\text{diff}(\kappa, v)(n) = \kappa(n) - v(n).$$

Therefore,

$$\ker \text{diff} \cong \ker \delta_{u_0, \clubsuit} \quad \text{and} \quad \text{coker diff} = \{0\},$$

which, by Remark 2.28, completes the proof of the proposition. ■

Construct the Kuranishi model as in Section 5.6 for the above choices of $S = S_1 \amalg S_2$ and \mathcal{O} .

As a final piece of preparation, let us make the following observation, which by Remark 2.28, in particular, gives an explicit description of $\mathcal{O}_{\bullet}^* = \text{coker}(\bar{\partial} \otimes_{\mathbf{C}} \mathbf{1})^*$.

Proposition 5.45. *Let (C, ν) be a nodal Riemann surface with nodal set S . Denote the corresponding nodal curve by \check{C} and its dualizing sheaf by $\omega_{\check{C}}$. Let $q \in (1, 2)$ be such that $1/p + 1/q = 1$. Define*

$$\mathcal{H} \subset L^q \Omega^{0,1}(C, \mathbf{C})$$

to be the subspace of solutions $\bar{\zeta}$ of the distributional equation

$$\bar{\partial}^* \bar{\zeta} = \sum_{n \in S} f(n) \delta(n) \tag{5.46}$$

for some weight function $f: S \rightarrow \mathbf{C}$ with $f \circ \nu = -f$. Here $\delta(n)$ is the Dirac delta distribution at n . The subspace \mathcal{H} satisfies

$$\bar{\mathcal{H}} = H^0(\check{C}, \omega_{\check{C}}).$$

Proof. If \check{C} is smooth, then $\check{C} = C$ and the dualizing sheaf ω_C is simply the canonical sheaf K_C . By the Kähler identities,

$$\begin{aligned} \bar{\mathcal{H}} &= \overline{\ker(\bar{\partial}^*: \Omega^{0,1}(C, \mathbf{C}) \rightarrow \Omega^0(C, \mathbf{C}))} \\ &= \overline{\ker(\partial: \Omega^{0,1}(C, \mathbf{C}) \rightarrow \Omega^{1,1}(C, \mathbf{C}))} \\ &\cong \overline{\ker(\bar{\partial}: \Omega^{1,0}(C, \mathbf{C}) \rightarrow \Omega^{1,1}(C, \mathbf{C}))} \\ &\cong H^0(C, K_C). \end{aligned}$$

Recall from the proof of Proposition 2.18, that the dualizing sheaf of \check{C} is constructed as follows; Denote by $\pi: C \rightarrow \check{C}$ the normalization map. Denote by $\tilde{\omega}_{\check{C}}$ the subsheaf of $K_C(S)$ whose sections ζ satisfy

$$\text{Res}_n \zeta + \text{Res}_{\nu(n)} \zeta = 0$$

for every $n \in S$, with $\text{Res}_n \eta$ being the residue of the meromorphic 1-form η at n . The dualizing sheaf $\omega_{\check{C}}$ is then

$$\omega_{\check{C}} = \pi_* \tilde{\omega}_{\check{C}}.$$

Therefore, $H^0(\check{C}, \omega_{\check{C}}) = H^0(C, \tilde{\omega}_{\check{C}})$. By definition every $\zeta \in H^0(C, \tilde{\omega}_{\check{C}})$ is smooth away from S and blows-up at most like $1/\text{dist}(n, \cdot)$ at n for $n \in S$; hence, $\zeta \in L^q \Omega^{0,1}(C, \mathbb{C})$. The residue condition amounts to (5.46). This shows that

$$H^0(\check{C}, \omega_{\check{C}}) \subset \mathcal{H}.$$

Conversely, by elliptic regularity every $\zeta \in \mathcal{H}$ defines an element of $H^0(\check{C}, \omega_{\check{C}})$. ■

The following is the technical backbone of the proof of Theorem 1.1. The reader is advised to recall Definition 5.35 and Proposition 5.36 because these are the main ingredients of the proof.

Lemma 5.47. *Denote by \check{C}_\bullet the nodal curve corresponding to (C_\bullet, ν_\bullet) . There is a constant $c > 0$ such that the obstruction map defined in Definition 5.35 satisfies the following. For every $(\sigma, \tau; J; \kappa) \in \Delta \times \mathcal{U} \times \mathcal{J}$, $\zeta \in H^0(\check{C}_\bullet, \omega_{\check{C}_\bullet})$, and $v \in T_{x_0} X$,*

$$\begin{aligned} & \langle \text{pull}_{\sigma, \tau; J; \kappa}(\text{ob}(\sigma, \tau; J; \kappa)), \text{pull}_{\sigma, \tau; J; \kappa}(\bar{\zeta} \otimes_{\mathbb{C}} v) \rangle_{L^2} \\ &= \sum_{n \in S_{C_\bullet}^{\text{ext}}} \pi \langle (\zeta(n) \otimes_{\mathbb{C}} d_{\nu_0(n)} u_{\sigma, \tau_1, 0; J; \kappa})(\tau_n), v \rangle + e \end{aligned}$$

with

$$|e| \leq c |\zeta| |v| \varepsilon_\bullet^{1/2} \sum_{n \in S_{C_\bullet}^{\text{ext}}} \varepsilon_n \quad \text{and} \quad \varepsilon_\bullet := \max\{\varepsilon_n : n \in S_{C_\bullet}^{\text{int}} \cup S_{C_\bullet}^{\text{ext}}\}.$$

Example 5.48. To understand better the significance of Lemma 5.47, it is helpful to consider the following example. Suppose that (Σ_0, j_0, ν_0) consists of two components: the higher genus ghost $C = C_\bullet$ and a spherical bubble $\Sigma_\bullet = CP^1$ meeting at points $n \in C$ and $\nu(n) \in CP^1$. (In that case, $B = B_0 = \emptyset$.) In that case, there is only one smoothing parameter τ , which, after trivializing the tangent spaces of C and Σ_\bullet at the nodes, can be thought of as a complex number, and $\varepsilon = |\tau|$. By Lemma 5.47, for every holomorphic 1-form $\zeta \in H^0(C, \omega_C)$,

$$\begin{aligned} & \langle \text{pull}_{\sigma, \tau; J; \kappa}(\text{ob}(\sigma, \tau; J; \kappa)), \text{pull}_{\sigma, \tau; J; \kappa}(\bar{\zeta} \otimes_{\mathbb{C}} v) \rangle_{L^2} \\ &= \pi \langle (\zeta(n) \otimes_{\mathbb{C}} d_{\nu_0(n)} u_{\sigma, 0; J; \kappa})(\tau), v \rangle + e \end{aligned} \tag{5.49}$$

with

$$|e| \leq c|\zeta||v|\varepsilon^{3/2}.$$

Since C has positive genus, there exists $\zeta \in H^0(C, \omega_C)$ such that $\zeta(n) \neq 0$. If the restriction of u_0 to the bubble $\mathbf{C}P^1$ is unobstructed, then $u_{\sigma,0;J;\kappa} = u_0$ on $\mathbf{C}P^1$ for all σ and κ . If, moreover, $d_{v(n)}u_0 \neq 0$, it follows that the right-hand side of (5.49) is never zero unless $\varepsilon = 0$, and so $\text{ob}(\sigma, \tau, J; \kappa) \neq 0$ for $\tau \neq 0$. We conclude that in that case u_0 cannot be smoothed.

Proof of Lemma 5.47. The proof is based on analyzing the expression

$$0 = \langle \tilde{\mathfrak{D}}_{\tilde{u}_{\sigma,\tau;J;\kappa}}(\hat{\xi}(\sigma, \tau; J; \kappa)) + \text{pull}_{\sigma,\tau;J;\kappa}(o(\sigma, \tau_1; J; \kappa) + \hat{o}(\sigma, \tau; J; \kappa)), \text{pull}_{\sigma,\tau;J;\kappa}(\bar{\zeta} \otimes_C v) \rangle_{L^2}$$

and the identity

$$\text{ob}(\sigma, \tau; J; \kappa) := o(\sigma, \tau_1; J; \kappa) + \hat{o}(\sigma, \tau; J; \kappa).$$

Step 1. The vector field $\hat{\xi}(\sigma, \tau_1; J; \kappa)$ is constant on C_\bullet and $o(\sigma, \tau_1; J; \kappa)$ is supported on Σ_\bullet ; in particular,

$$\langle \text{pull}_{\sigma,\tau;J;\kappa}(o(\sigma, \tau_1; J; \kappa)), \text{pull}_{\sigma,\tau;J;\kappa}(\bar{\zeta} \otimes_C v) \rangle_{L^2} = 0.$$

The construction in Proposition 5.29 can be carried out for $u_0|_{\Sigma_\bullet}$ with the choice of $\mathcal{O} = \mathcal{O}_\bullet$ and $S_2 = \emptyset$. For every $(\sigma, \tau_1, 0; J) \in \Delta \times \mathcal{U}$ and $\kappa \in \ker d_{u_0, \star}$ with $|\kappa| < \delta_\kappa$ denote by $\hat{\xi}(\sigma, \tau_1; J; \kappa)$ and $o(\sigma, \tau_1; J; \kappa)$ the solution of (5.30) obtained in this way.

Henceforth, regard $\hat{\xi}(\sigma, \tau_1; J; \kappa)$ as an element of $W^{1,p}\Gamma(\Sigma_0 \nu_0; u_0^*TX)$ and $o(\sigma, \tau_1; J; \kappa)$ as an element of \mathcal{O} . By construction these satisfy (5.30) for u_0 and with the choices of \mathcal{O} and $S = S_1 \amalg S_2$ made in the discussion preceding Lemma 5.47. Therefore, and since $\ker d_{u_0, \star} = \ker d_{u_0}$, $\hat{\xi}(\sigma, \tau_1; J; \kappa)$ and $o(\sigma, \tau_1; J; \kappa)$ are precisely the output produced by Proposition 5.29. The first part of the assertion thus holds by construction.

Step 2. The term

$$\bar{\partial}_J(\tilde{u}_{\sigma,\tau;J;\kappa}, j_{\sigma,\tau}) + \text{pull}_{\sigma,\tau;J;\kappa}(o(\sigma, \tau_1; J; \kappa))$$

is supported in the regions where $r_n \leq 2R_0$ for some $n \in S_{C_\bullet}^{\text{ext}}$. Set $x_n := u_{\sigma,\tau_1,0;J;\kappa}(n)$. Identifying U_{x_n} with \tilde{U}_{x_n} via \exp_{x_n} in the region where $r_n \leq 2R_0$, the error term can be written as

$$\bar{\partial} \chi_{\tau_2}^n \cdot u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2} + \mathfrak{q}$$

with

$$|\bar{\partial} \chi_{\tau_2}^n \cdot u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2}| \leq c\varepsilon_n \quad \text{and} \quad |\mathfrak{q}| \leq c\varepsilon_n^2. \tag{5.50}$$

The proof is a refinement of that of Proposition 5.9. A priori, the error term $\bar{\partial}_J(\tilde{u}_{\sigma,\tau;J;\kappa}, j_{\sigma,\tau}) + \text{pull}_{\sigma,\tau;J;\kappa}(o(\sigma, \tau_1; J; \kappa))$ is supported in the in the regions where $r_n \leq 2R_0$ for some $n \in S_2$. If $n \in S_{\Sigma_\bullet}^{\text{ext}}$, then it is immediate from Definition 5.7 that $\tilde{u}_{\sigma,\tau;J;\kappa}$ agrees with $u_{\sigma,\tau_1,0;J;\kappa}$ in the region under consideration; hence, the error term vanishes. For $n \in S_{C_\bullet}^{\text{ext}}$, in the region under consideration and with the identifications having been made,

$$\tilde{u}_{\sigma,\tau;J;\kappa}^\circ = \chi_{\tau_2}^n \cdot u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2}.$$

Therefore,

$$\begin{aligned} \bar{\partial}_J(\tilde{u}_{\sigma,\tau;J;\kappa}^\circ, j_{\sigma,\tau}) &= \bar{\partial}\chi_{\tau_2}^n \cdot u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2} \\ &+ \underbrace{\chi_{\tau_2}^n \cdot \frac{1}{2}(J(\tilde{u}_{\sigma,\tau;J;\kappa}^\circ) - J(u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2})) \circ d(u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2}) \circ j_{\sigma,\tau_1,0}}_{=:I} \\ &+ \underbrace{\chi_{\tau_2}^n \cdot \bar{\partial}_J(u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2}, j_{\sigma,\tau_1,0})}_{=:II}. \end{aligned}$$

(Observe that by elliptic regularity and (5.30), the map $u_{\sigma,\tau;J;\kappa}$ is smooth in the region in question, so we can take its derivative. We will use this fact in the remaining part of the proof.) The term I is supported in the region where $R_0 \leq r_n \leq 2R_0$. By Taylor expansion at $v_0(n)$, in this region

$$\begin{aligned} |u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2}| &\leq c\varepsilon_n/r_n, \\ |d(u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2})| &\leq c\varepsilon_n/r_n^2. \end{aligned}$$

Therefore,

$$|I| \leq c\varepsilon_n^2 \quad \text{and} \quad |\bar{\partial}\chi_{\tau_2}^n \cdot u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2}| \leq c\varepsilon_n.$$

Since ι_{τ_2} is holomorphic and $o(\sigma, \tau_1; \kappa)$ is defined by (5.30),

$$II = \chi_{\tau_2}^n \cdot \iota_{\tau_2}^* \bar{\partial}_J(u_{\sigma,\tau_1,0;J;\kappa}, j_{\sigma,\tau_1,0}) = -\chi_{\tau_2}^n \cdot \iota_{\tau_2}^* o(\sigma, \tau_1; J; \kappa),$$

and thus II vanishes by our choice of \mathcal{O} .

Step 3. For every $n \in S_{C_\bullet}^{\text{ext}}$,

$$\begin{aligned} &\langle \bar{\partial}\chi_{\tau_2}^n \cdot u_{\sigma,\tau_1,0;J;\kappa} \circ \iota_{\tau_2}, \text{pull}_{\sigma,\tau;J;\kappa}(\bar{\zeta} \otimes_C v) \rangle_{L^2} \\ &= -\pi \langle (\bar{\zeta} \otimes_C d_{v_0(n)} u_{\sigma,\tau_1,0;J;\kappa})(\tau_n), v \rangle + e_2 \end{aligned}$$

with

$$|e_2| \leq c\varepsilon_n^2 |\zeta| |v|.$$

To simplify the notation, we choose an identification $T_n C = \mathbf{C}$ and work in the canonical holomorphic coordinate z on C at n and the coordinate system at $v_0(n)$ with respect to which $w = \iota_{\tau_2}(z) = \varepsilon_n/z$. In particular, with respect to the induced identification $T_{v_0(n)} C = \mathbf{C}$ the gluing parameter is simply $\tau_n = \varepsilon_n \cdot 1 \otimes_{\mathbf{C}} 1$.

Since $u_{\sigma, \tau_1, 0; J; \kappa}$ is J -holomorphic, by Taylor expansion,

$$u_{\sigma, \tau_1, 0; J; \kappa}(\varepsilon_n/z) = \partial_w u_{\sigma, \tau_1, 0; J; \kappa}(0) \cdot \varepsilon_n/z + r \quad \text{with} \quad |r| \leq c\varepsilon_n^2/|z|^2.$$

The term $e'_2 := \langle \bar{\partial} \chi_{\tau_2}^n \cdot r, \text{pull}_{\sigma, \tau; J; \kappa}(\zeta \otimes_{\mathbf{C}} v) \rangle_{L^2}$ satisfies

$$|e'_2| \leq c\varepsilon_n^2 |\zeta| |v|.$$

Since ζ is holomorphic,

$$\int_{S^1} \zeta(re^{i\alpha}) \, d\alpha = 2\pi \cdot \zeta(0).$$

Therefore,

$$\begin{aligned} \langle \bar{\partial} \chi_{\tau_2}^n \cdot z^{-1}, \bar{\zeta} \rangle_{L^2} &= \int_{R_0 \leq |z| \leq 2R_0} \frac{1}{2} \chi' \left(\frac{|z|}{R_0} \right) \frac{\zeta(z)}{R_0 |z|} \, \text{vol} \\ &= \int_{R_0}^{2R_0} \frac{1}{2} \chi' \left(\frac{r}{R_0} \right) \frac{1}{R_0} \cdot \left(\int_{S^1} \zeta(re^{i\alpha}) \, d\alpha \right) \, dr \\ &= \int_{R_0}^{2R_0} \chi' \left(\frac{r}{R_0} \right) \frac{1}{R_0} \, dr \cdot \pi \zeta(re^{i\alpha}). \end{aligned}$$

The integral evaluates to -1 . Thus, the assertion follows because the term

$$\langle \zeta(0) \cdot \partial_w u_{\sigma, \tau_1, 0; J; \kappa}(0), v \rangle$$

can be written in coordinate-free form as

$$\pi \langle (\zeta \otimes_{\mathbf{C}} d_{v_0(n)} u_{\sigma, \tau_1, 0; J; \kappa})(\tau_n), v \rangle.$$

Step 4. The term

$$e_3 := \langle d_{\tilde{u}_{\sigma, \tau; J; \kappa}} \widehat{\xi}(\sigma, \tau; J; \kappa) + n_{\tilde{u}_{\sigma, \tau; J; \kappa}}(\widehat{\xi}(\sigma, \tau; J; \kappa)), \text{pull}_{\sigma, \tau; J; \kappa}(\bar{\zeta} \otimes_{\mathbf{C}} v) \rangle_{L^2}$$

satisfies

$$|e_3| \leq c\varepsilon_n^{1/2} \sum_{n \in S_{\mathbf{C}}^{\text{ext}}} \varepsilon_n |\zeta| |v|.$$

By Step 2 and Proposition 5.33,

$$\|\widehat{\xi}(\sigma, \tau; J; \kappa)\|_{W^{1,p}} \leq c\varepsilon_n.$$

This immediately implies that

$$e'_3 := \langle n_{\tilde{u}_{\sigma,\tau;J;\kappa}}(\widehat{\xi}(\sigma, \tau; J; \kappa)), \text{pull}_{\sigma,\tau;J;\kappa}(\bar{\zeta} \otimes_C v) \rangle_{L^2}$$

satisfies

$$|e'_3| \leq c \sum_{n \in S_{C_\bullet}^{\text{ext}}} \varepsilon_n^2 |\zeta| |v|.$$

It remains to estimate

$$e''_3 := \langle d_{\tilde{u}_{\sigma,\tau;J;\kappa}} \widehat{\xi}(\sigma, \tau; J; \kappa), \text{pull}_{\sigma,\tau;J;\kappa}(\bar{\zeta} \otimes_C v) \rangle_{L^2}.$$

Set

$$C_{\sigma,\tau}^\circ := C_\bullet \cap \Sigma_{\sigma,\tau}^\circ.$$

Since κ and $\xi(\sigma, \tau_1; \kappa)$ are constant on C , Proposition 5.16 implies that the term

$$d_{\tilde{u}_{\sigma,\tau;J;\kappa}} \widehat{\xi}(\sigma, \tau; J; \kappa) - \bar{\partial} \widehat{\xi}(\sigma, \tau; J; \kappa),$$

defined over $C_{\sigma,\tau}^\circ$, is supported in the regions where $\varepsilon_n^{1/2} \leq r_n \leq 2R_0$ for some $n \in S_{C_\bullet}^{\text{ext}}$ and satisfies

$$\begin{aligned} & \sum_{n \in S_{C_\bullet}^{\text{ext}}} \int_{\varepsilon_n^{1/2} \leq r_n \leq 2R_0} |d_{\tilde{u}_{\sigma,\tau;J;\kappa}} \widehat{\xi}(\sigma, \tau; J; \kappa) - \bar{\partial} \widehat{\xi}(\sigma, \tau; J; \kappa)| \\ & \leq c \sum_{n \in S_{C_\bullet}^{\text{ext}}} \int_{\varepsilon_n^{1/2} \leq r_n \leq 2R_0} |\widehat{\xi}(\sigma, \tau; J; \kappa)| |\nabla \widehat{\xi}(\sigma, \tau; J; \kappa)| + |\widehat{\xi}(\sigma, \tau; J; \kappa)|^2 \\ & \leq c \|\widehat{\xi}(\sigma, \tau; J; \kappa)\|_{W^{1,p}}^2 \\ & \leq c \sum_{n \in S_{C_\bullet}^{\text{ext}}} \varepsilon_n^2. \end{aligned}$$

Therefore,

$$|e''_3| \leq c \sum_{n \in S_{C_\bullet}^{\text{ext}}} \varepsilon_n^2 + |\langle \bar{\partial} \widehat{\xi}(\sigma, \tau; J; \kappa), \bar{\zeta} \otimes_C v \rangle_{L^2(C_{\sigma,\tau}^\circ)}|.$$

Since $\bar{\partial}^* \zeta = 0$ on $C_{\sigma,\tau}^\circ$, integration by parts yields

$$\begin{aligned} |\langle \bar{\partial} \widehat{\xi}(\sigma, \tau; J; \kappa), \bar{\zeta} \otimes_C v \rangle_{L^2(C_{\sigma,\tau}^\circ)}| & \leq c \varepsilon_\bullet^{1/2} \sum_{n \in S_{C_\bullet}^{\text{ext}}} \|\widehat{\xi}(\sigma, \tau; J; \kappa)\|_{W^{1,p}} |\zeta| |v| \\ & \leq c \varepsilon_\bullet^{1/2} \sum_{n \in S_{C_\bullet}^{\text{ext}}} \varepsilon_n. \end{aligned}$$

Combining the above estimates yields the asserted estimate on e_3 .

Step 5. Conclusion of the proof.

By Step 1, Step 2, Step 3, and Step 4 the term

$$o := \langle \text{pull}_{\sigma,\tau;J;\kappa}(\text{ob}(\sigma, \tau; J; \kappa)), \text{pull}_{\sigma,\tau;J;\kappa}(\bar{\zeta} \otimes_C v) \rangle_{L^2}$$

satisfies

$$\begin{aligned} o &= -\langle \mathfrak{F}\tilde{u}_{\sigma,\tau;J;\kappa}(\hat{\xi}(\sigma, \tau; J; \kappa)), \text{pull}_{\sigma,\tau;J;\kappa}(\bar{\zeta} \otimes_C v) \rangle_{L^2} \\ &= \sum_{n \in S_{C_\bullet}^{\text{ext}}} \pi \langle (\zeta \otimes_C d_{\nu_0(n)} u_{\sigma,\tau_1,0;J;\kappa})(\tau_n), v \rangle + e_1 + e_2 + e_3 \end{aligned}$$

with the error terms arising from the corresponding terms in the preceding steps. The term e_1 arises from the L^2 inner product of the sum, over all nodes $n \in S_{C_\bullet}^{\text{ext}}$, of error terms from Step 2 and $\text{pull}_{\sigma,\tau;J;\kappa}(\bar{\zeta} \otimes_C v)$, which multiplies the estimate (5.50) by $|\xi||v|$. The preceding steps thus yield the asserted estimate on $e = e_1 + e_2 + e_3$. ■

5.8. Proof of Theorem 1.1

Without loss of generality, suppose that

$$(\Sigma_k, j_k) = (\Sigma_{\sigma_k, \tau_k}, j_{\sigma_k, \tau_k})$$

with $\tau_{k;n} \neq 0$ for every $k \in \mathbb{N}$ and $n \in S$; that is, all nodes are smoothed and ν_{σ_k, τ_k} is the trivial nodal structure. By Proposition 5.37 there is a sequence $(\kappa_k)_{k \in \mathbb{N}}$ in \mathcal{J} corresponding to $(u_k)_{k \in \mathbb{N}}$.

Again, without loss of generality, $u_\infty: (\Sigma_\infty, j_\infty, \nu_\infty) \rightarrow (X, J_\infty)$ has at least one ghost component C with one non-ghost component

$$\Sigma_{\text{bubble}} = \Sigma_\infty \setminus C$$

attached at a single node, so that Hypothesis 5.41 is satisfied. Denote by \check{C} the nodal curve corresponding to C and let B be the base-locus of the dualizing sheaf of \check{C} . Define $B_0 \subset B$ and

$$C_\bullet := C \setminus B_0 \quad \text{and} \quad \Sigma_\clubsuit := \Sigma_0 \setminus C_\bullet$$

as at the beginning of Section 5.7. Observe that

$$\Sigma_\clubsuit = B_0 \amalg \Sigma_{\text{bubble}}$$

so that Σ_∞ decomposes into

$$\Sigma_\infty = C_\bullet \amalg \underbrace{B_0 \amalg \Sigma_{\text{bubble}}}_{\Sigma_\clubsuit}.$$

Write $\tau_k = (\tau_{k,1}, \tau_{k,2})$ with $\tau_{k,1}$ and $\tau_{k,2}$ denoting the smoothing parameters corresponding to the sets of nodes S_1 and S_2 defined in (5.42). Since B_0 is a tree of spheres, the partial smoothing $\Sigma_{\sigma_k, \tau_{k,1}, 0}$ contains a component biholomorphic to Σ_{bubble} , as discussed in Example 4.12. Let

$$b_k: \Sigma_{\text{bubble}} \rightarrow X$$


be the restriction of $u_{\sigma, \tau_{k,1}, 0; \kappa_k}$ to this component. After reparametrizing b_k by biholomorphisms of Σ_{bubble} we can guarantee that b_k converges to $u_0|_{\Sigma_{\text{bubble}}}$ in the C^∞ topology. Let x be any node in $S_{C_\bullet}^{\text{ext}}$; the point $v_k(x)$ under the above biholomorphisms it is mapped to a point $x_k \in \Sigma_{\text{bubble}}$ and the sequence (x_k) satisfies

$$\lim_{k \rightarrow \infty} x_k = v_\infty(n).$$

Let \check{C}_\bullet be the nodal curve corresponding to C_\bullet . By the construction of C_\bullet , for every node $n \in C_\bullet$ such that $v(n) \in \Sigma_\bullet$, there is a holomorphic section $\zeta \in H^0(\check{C}_\bullet, \omega_{\check{C}_\bullet})$ with $\zeta(n) \neq 0$. Since $\text{ob}(\sigma_k, \tau_k; J_k; \kappa_k) = 0$, it follows from Lemma 5.47 that

$$|d_{x_k} b_k| \leq \varepsilon_{k; \text{ghost}}^{1/2}.$$

(Observe that reparametrizing b_k by biholomorphisms of Σ_{bubble} does not affect the estimate in Lemma 5.47, which is independent of the choice of a Riemannian metric in the given conformal class.) Passing to the limit $k \rightarrow \infty$ yields that $d_{v_\infty(n)} u_0 = 0$.



6. Calabi–Yau classes in symplectic 6-manifolds

6.1. Proof of Theorem 1.3

Denote by C_1, \dots, C_I the connected components of Σ_∞ on which u_∞ is non-constant and set $u_\infty^i := u_\infty|_{C_i}$ and $A_i := (u_\infty^i)_* [C_i]$. By the index formula (2.27),

$$\sum_{i=1}^I \text{index}(u_\infty^i) = \sum_{i=1}^I 2\langle c_1(X, \omega), A_i \rangle = 2\langle c_1(X, \omega), A \rangle = 0$$

Since $J_\infty \in \mathcal{J}_{\text{emb}}(X, \omega)$, for every $i = 1, \dots, I$, we have $\text{index}(u_\infty^i) \geq 0$, and thus $\text{index}(u_\infty^i) = 0$. Consequently, the images of the simple maps underlying u_∞^i and u_∞^j either agree or are disjoint. However,

$$\text{im } u_\infty = \bigcup_{i=1}^I \text{im } u_\infty^i$$

is connected. Therefore and since A is primitive, $I = 1$ and u_∞^1 is simple and, hence, an embedding because $J \in \mathcal{J}_{\text{emb}}(X, \omega)$. Given the above, it follows from Theorem 1.1 that $(\Sigma_\infty, j_\infty, \nu_\infty)$ is smooth.

6.2. Proof of Theorem 1.5

The proof of Theorem 1.5 (1) is completely standard and straightforward. Nevertheless, let us spell it out. Let $J \in \mathcal{J}_{\text{emb}}^*(X, \omega)$. By Proposition 2.36 and Theorem 1.3, $\mathcal{M}_{A,g}^*(X, J)$ is a compact oriented zero-dimensional manifold; that is, a finite set of points with signs. The signed count $\#\mathcal{M}_{A,g}^*(X, J)$ is independent of the choice of J . To see this, let $J_0, J_1 \in \mathcal{J}_{\text{emb}}^*(X, \omega)$ and $(J_t)_{t \in [0,1]} \in \mathcal{J}_{\text{emb}}^*(X, \omega; J_0, J_1)$. By Proposition 2.36 and Theorem 1.3, $\mathcal{M}_{A,g}^*(X, (J_t)_{t \in [0,1]})$ is a compact oriented manifold with boundary

$$\mathcal{M}_{A,g}^*(X, J_1) \amalg -\mathcal{M}_{A,g}^*(X, J_0).$$

Therefore,

$$\#\mathcal{M}_{A,g}^*(X, J_1) = \#\mathcal{M}_{A,g}^*(X, J_0).$$

Theorem 1.5 (2) follows from [7, Theorem 1.6]. Indeed, the latter asserts that for every $J \in \mathcal{J}_*(X, \omega)$, the set

$$\coprod_{g=0}^\infty \mathcal{M}_{A,g}^*(X, J)$$

is a finite set. Therefore, there exists a $g_0 \in \mathbf{N}_0$ such that for every $g \geq g_0$ the moduli space $\mathcal{M}_{A,g}^*(X, A; J)$ is empty; in particular, $n_{A,g}(X, \omega) = 0$ for $g \geq g_0$. ■

7. Fano classes in symplectic 6-manifolds

The proofs in this section make use of definitions and results from Section 2.8.

7.1. Proof of Theorem 1.4

Denote by $\tilde{C}_1, \dots, \tilde{C}_I$ the connected components of Σ_∞ on which u_∞ is non-constant, set $\tilde{u}_\infty^i := u_\infty|_{\tilde{C}_i}$, denote by $u_\infty^i: C_i \rightarrow X$ the simple map underlying \tilde{u}_∞^i , let $d_i \in \mathbf{N}$ be the degree of the covering map relating \tilde{u}_∞^i and u_∞^i , set $A_i := (u_\infty^i)_*[C_i]$, and set $g_i := g(C_i)$. Since $(u_k)_{k \in \mathbf{N}}$ Gromov converges to u_∞ ,

- (1) $u_k(\Sigma_k)$ converges to $u_\infty(\Sigma_\infty)$ in the Hausdorff topology, and
- (2) $\sum_{i=1}^I d_i A_i = A$.

These are the only two consequences of Gromov convergence that will be used in the following argument.

Denote by I_0 the subset of those $i \in \{1, \dots, I\}$ with $\langle c_1(X, \omega), A_i \rangle = 0$ and set $I_+ := \{1, \dots, I\} \setminus I_0$. Without loss of generality all of the pseudo-cycles f_λ have $\text{codim}(f_\lambda) \geq 4$. For every $i \in I_+$, denote by Λ_i the subset of those $\lambda \in \{1, \dots, \Lambda\}$ such that

$$\text{im } u_\infty^i \cap \overline{\text{im } f_\lambda} \neq \emptyset. \tag{7.1}$$

Since $J_\infty \in \mathcal{J}_{\text{emb}}(X, \omega; f_1, \dots, f_\Lambda)$, for every $i \in I_0$ and $\lambda \in \{1, \dots, \Lambda\}$ we have $\text{im } u_\infty^i \cap \overline{\text{im } f_\lambda} = \emptyset$. Therefore, and since $u_k(\Sigma_k)$ converges to $u_\infty(\Sigma_\infty)$ in the Hausdorff topology, for every $\lambda \in \{1, \dots, \Lambda\}$ there exists at least one $i \in I_+$ such that (7.1) holds. For every $i \in \{1, \dots, I\}$ and $\lambda \in \Lambda_i$ set

$$f_\lambda^i := \begin{cases} f_\lambda & \text{if } \text{im } u_\infty^i \cap \text{im } f_\lambda \neq \emptyset, \\ f_\lambda^\partial & \text{otherwise,} \end{cases}$$

with f_λ^∂ as in Section 2.8; in particular: $\text{codim } f_\lambda \leq \text{codim } f_\lambda^i$ with equality if and only if

$$\text{im } u_\infty^i \cap \text{im } f_\lambda \neq \emptyset.$$

By definition, u_∞^i represents an element of $\mathcal{M}_{A,g}^*(X, J; (f_\lambda^i)_{\lambda \in \Lambda_i})$. Therefore, and since $J \in \mathcal{J}_{\text{emb}}(X, \omega, f_1, \dots, f_\Lambda)$,

$$2\langle c_1(X, \omega), A_i \rangle - \sum_{\lambda \in \Lambda_i} (\text{codim}(f_\lambda^i) - 2) \geq 0.$$

On the one hand, multiplying by d_i and summing yields

$$\begin{aligned} \sum_{i \in I_+} \sum_{\lambda \in \Lambda_i} (\text{codim}(f_\lambda^i) - 2) &\leq \sum_{i \in I_+} \sum_{\lambda \in \Lambda_i} d_i (\text{codim}(f_\lambda^i) - 2) \\ &\leq \sum_{i=1}^I 2\langle c_1(X, \omega), d_i A_i \rangle \\ &= 2\langle c_1(X, \omega), A \rangle \\ &= \sum_{\lambda=1}^\Lambda (\text{codim}(f_\lambda) - 2). \end{aligned}$$

On the other hand, by the preceding discussion, the reverse inequality also holds. Therefore, equality holds and this implies that

- (1) $d_i = 1$ for every $i \in I_+$,
- (2) $2\langle c_1(X, \omega), A_i \rangle = \sum_{\lambda \in \Lambda_i} (\text{codim}(f_\lambda^i) - 2)$,
- (3) $f_\lambda^i = f_\lambda$, and
- (4) the subsets Λ_i are non-empty and pairwise disjoint.

This implies that for every $i \in I_+$ the map \tilde{u}_∞^i agrees with u_∞^i and thus is simple; moreover, this map has index zero (in the sense of (2.27)) and its image intersects f_λ for every $\lambda \in \Lambda_i$. Furthermore, every f_λ intersects the image of precisely one map u_∞^i with $i \in I_+$. Therefore, the images of the maps u_∞^i with $i \in I_+$ are pairwise disjoint.

Since $2\langle c_1(X, \omega), A \rangle > 0$, I_+ is non-empty. For $i \in I_+$ and $j \in I_0$, the images of u_∞^i and u_∞^j must also be disjoint, because otherwise they would have to agree – contradicting $A_i \neq A_j$. However,

$$\text{im } u_\infty = \bigcup_{i=1}^I \text{im } u_\infty^i$$

is connected. Therefore, if $I_0 \neq \emptyset$, then there are $i \in I_0$ and $j \in I_+$ such that the images of u_∞^i and u_∞^j intersect. The preceding discussion shows this to be impossible; hence: $I_0 = \emptyset$. Similarly, if I_+ were to contain more than one element, then there are $i, j \in I_+$ with such that the images of u_∞^i and u_∞^j intersect – which is impossible. Therefore, $I = 1$ and $\tilde{u}_\infty^1 = u_\infty^1$ is an embedding.

Given the above, it follows from Theorem 1.1 that $(\Sigma_\infty, j_\infty, \nu_\infty)$ is smooth and $\text{im } u_\infty \cap \text{im } f_\lambda \neq \emptyset$ every $\lambda = 1, \dots, \Lambda$. ■

7.2. Proof of Theorem 1.7

Given Gromov compactness, Theorem 1.4, and Proposition 2.44, the proof that

$$n_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda)$$

is well defined and independent of the choice of J is identical to that of Theorem 1.5 up to changes in notation.

To prove that $n_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda)$ is independent of the choice of pseudo-cycle representatives, suppose that f_1^0 and f_1^1 are two representatives of $\text{PD}[\gamma_1]$ such that $f_1^i, f_2, \dots, f_\Lambda$ are in general position for $i = 0, 1$. Let $F: W \rightarrow X$ be a pseudo-cycle cobordism between f_1^0 and f_1^1 such that F, f_2, \dots, f_Λ are in general position. Let J be an element of the set $\mathcal{J}_{\text{emb}}^*(X, \omega; F, f_2, \dots, f_\Lambda)$ defined in Definition 2.46, which is residual by Proposition 2.47. It follows that $\mathcal{M}_{A,g}^*(X, J; f_1^0, \dots, f_\Lambda)$ and $\mathcal{M}_{A,g}^*(X, J; f_1^1, \dots, f_\Lambda)$ are finite sets of points with orientations and

$$\mathcal{M}_{A,g}^*(X, J; F, f_2, \dots, f_\Lambda)$$

is an oriented 1-dimensional cobordism between them. This cobordism is compact by Gromov compactness and the argument used in the proof in Theorem 1.4. Thus,

$$\#\mathcal{M}_{A,g}^*(X, J; f_1^0, \dots, f_\Lambda) = \#\mathcal{M}_{A,g}^*(X, J; f_1^1, \dots, f_\Lambda).$$

The fact that $n_{A,g}(X, \omega; \gamma_1, \dots, \gamma_\Lambda) = 0$ for $g \gg 1$ is a consequence of the following analogue of [7, Theorem 1.6] for Fano classes.

Theorem 7.2. *Let (X, ω) be a compact symplectic 6-manifold, f_1, \dots, f_Λ be a collection of even-dimensional pseudo-cycles in general position, and let $A \in H_2(X, \mathbf{Z})$ be such that*

$$2\langle c_1(X, \omega), A \rangle = \sum_{\lambda=1}^{\Lambda} (\text{codim } f_\lambda - 2) > 0.$$

For every $J \in \mathcal{J}(X, \omega; f_1, \dots, f_\Lambda)$ there are only finitely many simple J -holomorphic maps representing A and passing through $\text{im } f_\lambda$ for every $\lambda = 1, \dots, \Lambda$.

Proof. The proof is a minor variation of the proof of [7, Theorem 1.6]. Suppose, by contradiction, that there are infinitely many distinct J -holomorphic curves C_k representing A and passing through $\text{im } f_\lambda$ for all $\lambda = 1, \dots, \Lambda$. Here, by a J -holomorphic curve we mean the image of a simple J -holomorphic map. Considering C_k as J -holomorphic cycles, we can pass to a subsequence which converges geometrically to a J -holomorphic cycle $C_\infty = \sum_{i=1}^I d_i C_\infty^i$, see [7, Definitions 4.1, 4.2, Lemma 1.9]. Here $d_i > 0$ are integers and each C_∞^i is a J -holomorphic curve. Geometric convergence implies that

$$\sum_{i=1}^I d_i [C_\infty^i] = [C_\infty] = A$$

and that $(C_k)_{k \in \mathbf{N}}$ converges to C_∞ in the Hausdorff topology. Since these were the only two conditions needed for the argument in the proof Theorem 1.4, the same argument shows that:

- (1) $d_i = 1$ for every $i \in I$,
- (2) C_∞ has only one connected component,
- (3) C_∞ intersects every $\text{im } f_\lambda$, and consequently
- (4) C_∞ is embedded and unobstructed by the condition $J \in \mathcal{J}_{\text{emb}}^*(X, \omega; f_1, \dots, f_\Lambda)$.

We will now adapt the rescaling argument from the proof of [7, Proposition 5.1] – originally due to Taubes in the 4-dimensional setting [35] – to the present situation. Let $N \rightarrow C_\infty$ be the normal bundle of C_∞ in X . Identify a neighborhood of C_∞ with a neighborhood of the zero section in N using the exponential map. For sufficiently large k , C_k is contained in that neighborhood and by abuse of notation we will consider C_k as an $\text{exp}^* J$ -holomorphic curve in N and f_λ as maps to N .

Since the C_k are distinct, $C_k \neq C_\infty$. For $\varepsilon > 0$ denote by $\sigma_\varepsilon: N \rightarrow N$ the map which rescales the fibers by ε . Let $(\varepsilon_k)_{k \in \mathbf{N}}$ be the sequence of positive numbers such that the rescaled sequence

$$\tilde{C}_k := (\sigma_{\varepsilon_k})^{-1}(C_k)$$

satisfies

$$d_H(\tilde{C}_k, C_\infty) = 1,$$

where d_H is the Hausdorff distance. The sequence $(\varepsilon_k)_{k \in \mathbb{N}}$ converges to zero. The curves \tilde{C}_k are J_k -holomorphic where $J_k := \sigma_{\varepsilon_k}^* \exp^* J$. The sequence of rescaled almost complex structures $(J_k)_{k \in \mathbb{N}}$ converges to an almost complex structure J_∞ which is tamed by a symplectic form [7, Proposition 3.10]. In the same way as in the proof of [7, Proposition 5.1] we conclude that the sequence $(\tilde{C}_k)_{k \in \mathbb{N}}$ converges geometrically to a J_∞ -holomorphic cycle whose support is a union of J_∞ -holomorphic curves $\tilde{C}_\infty \subset N$ satisfying

$$d_H(\tilde{C}_\infty, C_\infty) = 1.$$

Since $[\tilde{C}_k] = [C_\infty] = A$ for all k , and the bundle projection $\pi: N \rightarrow C_\infty$ is J_∞ -holomorphic, π induces an isomorphism $\tilde{C}_\infty \cong C_\infty$. Let $\iota: C_\infty \rightarrow X$ be the inclusion map and denote by δ_ι the deformation operator corresponding to ι , as in Definition 2.24. By [7, Proposition 3.12], \tilde{C}_∞ is the graph of a non-zero section

$$\xi \in \Gamma(C_\infty, N) \subset \Gamma(C_\infty, \iota^*TX)$$

satisfying $\delta_\iota \xi = 0$. In Proposition 7.3 below we show that there is an algebraic constraint for the values of ξ at the points of intersection of C_∞ with each pseudo-cycle.

For every $\lambda = 1, \dots, \Lambda$, denote by V_λ the domain of f_λ , and let $z_{\lambda,k} \in C_k$ and $x_{\lambda,k} \in V_\lambda$ be such that

$$z_{\lambda,k} = f_\lambda(x_{\lambda,k}).$$

After passing to a subsequence, we may assume that

$$\lim_{k \rightarrow \infty} z_{\lambda,k} = z_\lambda \in C_\infty \quad \text{and} \quad \lim_{k \rightarrow \infty} x_{\lambda,k} = x_\lambda \in V_\lambda,$$

and $z_\lambda = f_\lambda(x_\lambda)$.

Proposition 7.3. *For $\lambda = 1, \dots, \Lambda$, there exist $v_\lambda \in T_{z_\lambda} C_\infty$ and $w_\lambda \in T_{x_\lambda} V_\lambda$ such that*

$$\xi(z_\lambda) + d_{z_\lambda} \iota \cdot v_\lambda = d_{x_\lambda} f_\lambda \cdot w_\lambda. \tag{7.4}$$

Equation (7.4) can be understood as the limit as $k \rightarrow \infty$ of the condition that C_k intersects each of the $\text{im } f_\lambda$. The proof is deferred to the end of this section. We will now show that Proposition 7.3 implies Theorem 7.2. Let g be the genus of C_∞ , so that the embedding $\iota: C_\infty \rightarrow X$ corresponds to an element in $\mathcal{M}_{A,g,\Lambda}^*(X, J)$. Since $J \in \mathcal{J}_{\text{emb}}^*(X, \omega, f_1, \dots, f_\Lambda)$,

- (1) the derivative of $\text{ev}_\Lambda: \mathcal{M}_{A,g,\Lambda}^*(X, J) \rightarrow X^\Lambda$ at $[l, z_1, \dots, z_\Lambda]$, and
- (2) the derivative of $\prod_{\lambda=1}^\Lambda f_\lambda: \prod_{\lambda=1}^\Lambda V_\lambda \rightarrow X^\Lambda$ at $\prod_{\lambda=1}^\Lambda x_\lambda$

are transverse to each other. Since

$$\dim \mathcal{M}_{A,g,\Lambda}^*(X, J) + \sum_{\lambda=1}^{\Lambda} \dim V_{\lambda} = \Lambda \dim X,$$

the images of these two maps intersect trivially. Since $\xi \neq 0$, this contradicts the existence of v_{λ} and w_{λ} satisfying (7.4). The contradiction shows that the sequence (C_k) cannot exist. ■

Proof of Proposition 7.3. Set $\tilde{z}_{\lambda,k} := \sigma_{\varepsilon_k}^{-1}(z_{\lambda,k})$. After possibly passing to a further subsequence,

$$\lim_{k \rightarrow \infty} \tilde{z}_{\lambda,k} = \xi(z_{\lambda}). \tag{7.5}$$

Let $\text{pr}_N d_{x_{\lambda}} f_{\lambda} : T_{x_{\lambda}} V_{\lambda} \rightarrow N_{z_{\lambda}}$ be the projection of the derivative of f_{λ} at x_{λ} on $N_{z_{\lambda}} \subset T_{z_{\lambda}} X$. We will show that for every λ there exists $w_{\lambda} \in T_{x_{\lambda}} V_{\lambda}$ such that

$$\lim_{k \rightarrow \infty} \tilde{z}_{\lambda,k} = \text{pr}_N d_{x_{\lambda}} f_{\lambda} \cdot w_{\lambda}.$$

The fact that the images of the maps (1) and (2) introduced above intersect trivially implies that $\text{pr}_N d_{x_{\lambda}} f_{\lambda}$ is injective for every λ . Indeed, otherwise there would exist $v \in T_{z_{\lambda}} C_{\infty}$ and $w \in T_{x_{\lambda}} V_{\lambda}$ for some λ such that

$$d_{z_{\lambda}} \iota \cdot v = d_{x_{\lambda}} f_{\lambda} \cdot w,$$

violating the above transversality condition. Fix a trivialization of N in a neighborhood of z_{λ} and a chart centered at x_{λ} in V_{λ} . Denoting by pr_N the projection on the fiber $N_{z_{\lambda}}$ in the given trivialization, Taylor expansion gives us

$$\text{pr}_N z_{\lambda,k} = \text{pr}_N f_{\lambda}(x_{\lambda,k}) = \text{pr}_N d_{x_{\lambda}} f_{\lambda}(x_{\lambda,k} - x_{\lambda}) + O(|x_{\lambda,k} - x_{\lambda}|^2).$$

Since $\text{pr}_N d_{x_{\lambda}} f_{\lambda}$ is injective, there is a constant $c > 0$ such that

$$|x_{\lambda,k} - x_{\lambda}| \leq c |\text{pr}_N z_{\lambda,k}| \leq c \varepsilon_k.$$

Thus, after passing to a subsequence, we may assume that the sequence $\varepsilon_k^{-1}(x_{\lambda,k} - x_{\lambda})$ converges to a limit $w_{\lambda} \in T_{x_{\lambda}} V_{\lambda}$. By construction,

$$\lim_{k \rightarrow \infty} \tilde{z}_{\lambda,k} = \lim_{k \rightarrow \infty} \text{pr}_N \tilde{z}_{\lambda,k} = \text{pr}_N d_{x_{\lambda}} f_{\lambda} \cdot w_{\lambda}.$$

Comparing this with (7.5), we see that for every λ there exists $v_{\lambda} \in T_{z_{\lambda}} C_{\infty}$ such that (7.4) holds. ■

A. Transversality for evaluation maps

Throughout this section, (X, ω) is a symplectic manifold of dimension $\dim X \geq 6$ and $\mathcal{J}(X, \omega)$ denotes the space of almost complex structures on X compatible with ω .

Definition A.1. Let $\Lambda \in \mathbf{N}$. Given a partition into non-empty pairwise disjoint subsets

$$\{1, 2, \dots, \Lambda\} = I_1 \sqcup \dots \sqcup I_k \quad \text{with } k < \Lambda,$$

the *generalized diagonal* associated with the partition is the submanifold $\Delta \subset X^\Lambda$ consisting of the points (x_1, \dots, x_Λ) such that for every pair of indices $\alpha, \beta \in I_i$ we have $x_\alpha = x_\beta$.

Generalized diagonals are partially ordered by inclusion and each point of X^Λ which belongs to a generalized diagonal belongs to a unique one which is minimal with respect to the partial order.

Proposition A.2. Let V be a manifold and let $f: V \rightarrow X^\Lambda$ be a map which is transverse to every generalized diagonal. Denote by $\mathcal{J}^*(X, \omega; f)$ the set of all $J \in \mathcal{J}^*(X, \omega)$ such that:

- (1) every simple J -holomorphic map is unobstructed, and
- (2) for every $A \in H_2(X, \mathbf{Z})$ and $g \in \mathbf{N}_0$, the evaluation map from the Λ -pointed moduli space (cf. Definition 2.39)

$$\text{ev}: \mathcal{M}_{A,g,\Lambda}^*(X, J) \rightarrow X^\Lambda$$

is transverse to f .

The set $\mathcal{J}^*(X, \omega; f)$ is residual in $\mathcal{J}^*(X, \omega)$.

Proof. The proof that condition (1) is generic is a standard application of the Sard–Smale theorem [26, Theorem 1.2], [19, Proposition A.4], [24, Sections 3.2 and 6.3]. Below we outline this proof and adapt it to show that condition (2) is generic.

Let (Σ, j_0) be a closed Riemann surface of genus g , and let $A \in H_2(X, \mathbf{Z})$. Denote by $W_{\text{inj}}^{1,p}(\Sigma, X; A)$ the subset of $W^{1,p}(\Sigma, X)$ consisting of functions $u: \Sigma \rightarrow X$ which represent A and are *somewhere injective* in the sense that there exist $z_0 \in \Sigma$ and $\delta > 0$ such that for all $z \in \Sigma$

$$\text{dist}_X(u(z_0), u(z)) \geq \delta \text{dist}_\Sigma(z_0, z).$$

A J -holomorphic map is somewhere injective if and only if it is simple [24, Proposition 2.5.1]. Given a slice $\mathcal{S} \subset \mathcal{J}(\Sigma)$ for the action of $\text{Diff}_0(\Sigma)$ on $\mathcal{J}(\Sigma)$ passing through j_0 , set

$$\mathcal{X} = W_{\text{inj}}^{1,p}(\Sigma, X; A) \times \mathcal{S}$$

and let $\mathcal{E} \rightarrow \mathcal{X}$ be a Banach vector bundle whose fiber over (u, j) is the space

$$L^p \Omega^{0,1}(\Sigma, u^*TX)$$

defined using the complex structure j .

Let $s: \mathcal{J}(X, \omega) \times \mathcal{X} \rightarrow \mathcal{E}$ be a section given by $s(J, u, j) = \bar{\partial}_J(u, j)$. The following hold; see, for example, [24, Section 3.2]:

- this section is Fredholm,
- it is transverse to the zero section, therefore
- $s^{-1}(0)$ is a submanifold of \mathcal{X} ; in particular, it is a Banach manifold, and
- the universal moduli space $\mathcal{M}_{A,g}^*(X, \omega)$ can be covered by a countable number of submanifolds of the form $s^{-1}(0)$, for different choices of (Σ, j_0) .

The projection $\pi: s^{-1}(0) \rightarrow \mathcal{J}(X, \omega)$ is a Fredholm map of index $\text{vdim } \mathcal{M}_{A,g}^*(X, J)$; in fact, the kernel and cokernel of $d\pi_{u,j}$ are isomorphic to the kernel and cokernel of $d_{u,j} \bar{\partial}_J$, and therefore finite-dimensional. It follows from the implicit function theorem that if J is a regular value of this map, the preimage

$$\pi^{-1}(J) = \mathcal{M}_{A,g}^*(X, J)$$

is a manifold of dimension $\text{vdim } \mathcal{M}_{A,g}^*(X, J)$ and every map in $\mathcal{M}_{A,g}^*(X, J)$ is unobstructed. Since $\pi: s^{-1}(0) \rightarrow \mathcal{J}(X, \omega)$ is a Fredholm map between separable Banach manifolds, the Sard–Smale theorem implies that the set of regular values of π is residual in $\mathcal{J}(X, \omega)$. This shows that condition (1) holds for a generic J .

Using a similar argument, we will show for a generic J , the evaluation map

$$\text{ev}: \mathcal{M}_{A,g,\Lambda}^*(X, J) \rightarrow X^\Lambda$$

is transverse to f . With the notation introduced above, consider the Fredholm map

$$S: \mathcal{J}(X, \omega) \times \mathcal{X} \times \Sigma^\Lambda \rightarrow \mathcal{E} \times X^\Lambda, \\ S(J, (u, j), z_1, \dots, z_\Lambda) = (s(J, u, j), u(z_1), \dots, u(z_\Lambda)).$$

We will show that S is transverse to the map

$$\text{zero section} \times f: \mathcal{X} \times V \rightarrow \mathcal{E} \times X^\Lambda. \tag{A.3}$$

Since s is transverse to the zero section $\mathcal{X} \rightarrow \mathcal{E}$, it suffices to show that whenever $(J, (u, j), z_1, \dots, z_\Lambda)$ and $x \in V$ satisfy

$$s(J, u, j) = 0 \quad \text{and} \quad (u(z_1), \dots, u(z_\Lambda)) = f(x),$$

then

$$\text{im } dS + \text{im } d_x f = T_{f(x)} X^\Lambda = \bigoplus_{i=1}^\Lambda T_{u(z_i)} X.$$

Here we have used dS to denote the projection on $T_{f(x)} X^\Lambda$ of the derivative of S at $(J, (u, j), z_1, \dots, z_\Lambda)$ and $d_x f$ is the derivative of f at x . The variation of S in the direction of a vector field

$$\xi \in W^{1,p} \Gamma(\Sigma, u^* TX)$$

is

$$dS(\xi) = (\xi(z_1), \dots, \xi(z_\Lambda)). \tag{A.4}$$

If $(u(z_1), \dots, u(z_\Lambda))$ does not lie on any generalized diagonal in X^Λ , we can find ξ with any prescribed values at z_1, \dots, z_Λ , and

$$\text{im } dS = T_{f(x)} X^\Lambda.$$

Suppose, on the other hand, that $(u(z_1), \dots, u(z_\Lambda))$ belongs to a generalized diagonal $\Delta \subset X^\Lambda$, and let Δ be the minimal such diagonal. In that case, (A.4) implies that

$$\text{im } dS = T_{f(x)} \Delta.$$

Since f is transverse to Δ , we have

$$\text{im } dS + \text{im } d_x f = T_{f(x)} \Delta + \text{im } d_x f = T_{f(x)} X^\Lambda,$$

as desired. This shows that S is transverse to the map (A.3). It follows from the Sard–Smale theorem that the set of J such that $S(J, \cdot)$ is transverse to (A.3) is residual in $\mathcal{J}(X, \omega)$. This completes the proof that condition (2) is generic. ■

B. Pseudo-cycles

Given a collection of homology classes, we are interested in counting J -holomorphic maps passing through cycles representing these classes. Since not every homology class is represented by a map from a manifold, it is convenient to use the language of pseudo-cycles. We briefly review the theory of pseudo-cycles below; for details, see [24, Section 6.5] and [20, 34, 37].

Definition B.1. We make the following definitions:

- (1) A subset of a smooth manifold X is said to have *dimension at most k* if it is contained in the image of a smooth map from a smooth k -dimensional manifold.

(2) A k -pseudo-cycle is a smooth map $f: V \rightarrow X$ from an oriented k -dimensional manifold V such that the closure $\overline{f(V)}$ is compact and the *boundary of f* , defined by

$$\text{bd}(f) := \bigcap_{K \subset V \text{ compact}} \overline{f(V - K)},$$

has dimension at most $k - 2$. We will use notation

$$\text{codim}(f) := \dim(X) - \dim(V).$$

(3) Two k -pseudo-cycles $f_i: V_i \rightarrow X$, for $i = 0, 1$, are *cobordant* if there exists a smooth, oriented $(k + 1)$ -dimensional manifold with boundary W and a smooth map $F: W \rightarrow X$ such that $\overline{F(W)}$ is compact, $\text{bd}(F)$ has dimension at most $k - 1$, and

$$\partial W = V_1 \amalg -V_0 \quad \text{and} \quad F|_{V_1} = f_1, \quad F|_{V_0} = f_0.$$

(4) Denote by $H_k^{\text{pseudo}}(X)$ the set of equivalence classes of k -pseudo-cycles up to cobordism. The disjoint union operation endows $H_k^{\text{pseudo}}(X)$ with the structure of an abelian group.

A smooth map $g: X \rightarrow Y$ between two smooth manifolds induces a group homomorphism $g_*: H_*^{\text{pseudo}}(X) \rightarrow H_*^{\text{pseudo}}(Y)$ by composing pseudo-cycles with g . Thus, $H_*^{\text{pseudo}}(\cdot)$ is a functor from the category of smooth manifolds to the category of \mathbf{Z} -graded abelian groups.

Theorem B.2 ([20,34,37]). *There exists a natural isomorphism $H_*^{\text{pseudo}}(\cdot) \cong H_*(\cdot, \mathbf{Z})$ as functors from the category of smooth manifolds to the category of \mathbf{Z} -graded abelian groups.*

In what follows we will use this isomorphism to identify these two homology theories and represent any class in $H_*(X, \mathbf{Z})$ by a pseudo-cycle.

Definition B.3. Let M be a smooth manifold and let $g: M \rightarrow X$ be a smooth map. We say that a k -pseudo-cycle $f: V \rightarrow X$ is *transverse² as a pseudo-cycle to g* if

- (1) there exists a smooth manifold V^∂ of dimension $\dim V^\partial \leq \dim V - 2$ and a smooth map $f^\partial: V^\partial \rightarrow X$ such that $\text{bd}(f) \subset \text{im } f^\partial$, and
- (2) f and f^∂ are transverse to g as smooth maps from manifolds.

²McDuff and Salamon [24, Definition 6.5.10] use the term *weakly transverse*, which we prefer to avoid, regarding that this notion of transversality is stronger than the transversality of f and g as smooth maps in the usual sense.

If W is a manifold with boundary ∂W , we require additionally that f is transverse as a pseudo-cycle to $g|_{\partial W}: \partial W \rightarrow X$.

Similarly, if M is a manifold without boundary and $F: W \rightarrow X$ is a cobordism between two pseudo-cycles f_0 and f_1 , we say that F is *transverse as a pseudo-cycle cobordism to g* if

- (1) there exists a smooth manifold with boundary W^∂ of dimension $\dim W^\partial \leq \dim W - 2$ and a smooth map $F^\partial: W^\partial \rightarrow X$ such that $\text{bd}(F) \subset \text{im } F^\partial$ and $\text{bd}(f_i) \subset \text{im } F^\partial|_{\partial W^\partial}$ for $i = 0, 1$,
- (2) F and F^∂ are transverse to g as smooth maps from manifolds with boundary.

Note that if $f: V \rightarrow X$ be a k -pseudo-cycle and $g: W \rightarrow X$ is an ℓ -pseudo-cycle, then $f \times g: V \times W \rightarrow X^2$ is a $(k + \ell)$ -pseudo-cycle.

Definition B.4. Let $(f_\lambda: V_\lambda \rightarrow X)$ be a collection of pseudo-cycles indexed by a finite set I . We say that $(f_\lambda)_{\lambda \in I}$ are *in general position* if the pseudo-cycle

$$\prod_{\lambda \in I} f_\lambda: \prod_{\lambda \in I} V_\lambda \rightarrow X^{|I|}$$

is transverse as a pseudo-cycle to all generalized diagonals in $X^{|I|}$; see Definition A.1 for the definition of a generalized diagonal. This is equivalent to the following condition: for every subset $S \subset I$, the pseudo-cycle $\prod_{\lambda \in S} f_\lambda$ is transverse as a pseudo-cycle to the diagonal $X \hookrightarrow X^{|S|}$.

Similarly, if one of f_λ is a cobordism between two pseudo-cycles, then so is $\prod_{\lambda \in I} f_\lambda$ and we require that it is transverse to all generalized diagonals as a pseudo-cycle cobordism.

Proposition B.5. *Given a finite collection of pseudo-cycles $(f_\lambda: V_\lambda \rightarrow X)_{\lambda \in I}$, the set*

$$\{(\phi_\lambda)_{\lambda \in I} \in \text{Diff}(X)^{|I|} : (\phi_\lambda \circ f_\lambda)_{\lambda \in I} \text{ are in general position}\}$$

is residual in $\text{Diff}(X)^{|I|}$.

Proof. The proof is similar to that of [24, Lemma 6.5.5]. Let us work with the group $\text{Diff}_k(X)$ of C^k diffeomorphism for any integer $k \geq 1$; the corresponding statement for $\text{Diff}(X)$ follows then using standard arguments [24, Remark 3.2.7, pp. 52–54]. A countable intersection of residual sets is residual; therefore, without loss of generality, consider the case $S = I$ in Definition B.4. Define the map

$$\mathcal{F}: \text{Diff}_k(X)^{|I|} \times \prod_{\lambda \in I} V_\lambda \rightarrow X^{|I|}$$

by

$$\mathcal{F}((\phi_\lambda)_{\lambda \in I}, (x_\lambda)_{\lambda \in I}) := (\phi_\lambda \circ f_\lambda(x_\lambda))_{\lambda \in I}.$$

Let $\Delta \subset X^{|I|}$ be the diagonal. If we show that \mathcal{F} is transverse to Δ , then it follows from the Sard–Smale theorem that for all $(\phi_\lambda)_{\lambda \in I}$ from a residual subset of $\text{Diff}_k(X)$ the maps $\prod \phi_\lambda \circ f_\lambda$ is transverse to Δ . (The same argument can be applied to f_λ^∂ to conclude transversality as pseudo-cycles.) In fact, the derivative of \mathcal{F} is surjective at every point $\mathbf{x} = ((\phi_\lambda)_{\lambda \in I}, (x_\lambda)_{\lambda \in I})$. Without loss of generality suppose that $\phi_\lambda = \text{id}$ for all $\lambda \in I$. Let $\text{Vect}_k(X)$ denote the space of C^k vector fields on X . Given

$$\xi = (\xi_\lambda)_{\lambda \in I} \in \prod_{\lambda \in I} T_{\text{id}} \text{Diff}_k(X) = \prod_{\lambda \in I} \text{Vect}_k(X),$$

we have

$$d_{\mathbf{x}}\mathcal{F}(\xi) = (\xi_\lambda(f_\lambda(x_\lambda)))_{\lambda \in I} \in \prod_{\lambda \in I} T_{f_\lambda(x_\lambda)}X.$$

Since for every $p \in X$ the evaluation map $\text{Vect}(X) \rightarrow T_p X$ is surjective, the map $d_{\mathbf{x}}\mathcal{F}$ is surjective, which finishes the proof. ■

C. Proof of $n_{A,g} = \text{BPS}_{A,g}$

In this section, we outline Zinger’s proof that for a primitive Calabi–Yau class

$$n_{A,g}(X, \omega) = \text{BPS}_{A,g}(X, \omega),$$

where $\text{BPS}_{A,g}(X, \omega)$ is the Gopakumar–Vafa invariant, which is defined in terms of the Gromov–Witten invariants via (1.9). We use the same notation as in the proof of Theorem 1.5.

Given $J \in \mathcal{J}_{\text{emb}}^*(X, \omega)$, every stable J -holomorphic map of arithmetic genus h factors through a J -holomorphic embedding from a smooth domain of genus $g \leq h$. In other words, every element of $\tilde{\mathcal{M}}_{A,h}(X, J)$ is of the form $[u \circ \varphi]$ for some $[u] \in \mathcal{M}_{A,g}^*(X, J)$ with $g \leq h$, and $[\varphi] \in \tilde{\mathcal{M}}_{[\Sigma],h}(\Sigma, j)$. Here (Σ, j) is the domain of u . Denote by $(\tilde{\Sigma}, \tilde{\nu}, \tilde{j})$ the domain of φ . Given such J -holomorphic maps, let N be the normal bundle of $u(\Sigma)$, and let

$$\delta_u^N: W^{1,p} \Gamma(\Sigma, u^* N) \rightarrow L^p \Omega^{0,1}(\Sigma, u^* N)$$

be the restriction of the operator $\delta_u = \delta_{u,j;J}$ to the subbundle $u^* N \subset u^* TX$ followed by the projection on $\tilde{u}^* N$. Similarly, we define

$$\delta_{\tilde{u}}^N: W^{1,p} \Gamma(\tilde{\Sigma}, \tilde{\nu}; \tilde{u}^* N) \rightarrow L^p \Omega^{0,1}(\tilde{\Sigma}, \tilde{u}^* N).$$

The spaces $\text{coker } \delta_{\tilde{u}}^N$, as φ varies, play an important role in computing the contribution of maps factoring through u to the Gromov–Witten invariant of (X, ω) . In this case, there is a simple description of these spaces.

First, we will see that $\ker \delta_u^N = \{0\}$ and $\text{coker } \delta_u^N = \{0\}$. Indeed, the Hermitian metric on u^*TX induced from X gives us a splitting $u^*TX = T\Sigma \oplus N_u$, with respect to which

$$\delta_u = \begin{pmatrix} \bar{\partial}_{T\Sigma} & * \\ 0 & \delta_u^N \end{pmatrix};$$

see, for example, [8, Appendix A]. Since u is unobstructed, i.e., $\text{coker } \delta_u = \{0\}$, and $\text{index}(u) = 0$, we have $\ker \delta_u^N = \{0\}$ and $\text{coker } \delta_u^N = \{0\}$.

Second, since $\varphi: (\tilde{\Sigma}, \tilde{\nu}, \tilde{j}) \rightarrow (\Sigma, j)$ has degree one, $(\tilde{\Sigma}, \tilde{\nu}, \tilde{j})$ has a unique irreducible component which is mapped by φ biholomorphically to (Σ, j) , and φ is constant on the other components. In particular, \tilde{u}^*N is trivial over these components. It follows that $\ker \delta_u^N \cong \{0\}$ and $\text{coker } \delta_u^N$ is the direct sum of the corresponding spaces for the standard $\bar{\partial}$ -operator with values in the trivial bundle \tilde{u}^*N over the components which are mapped to a point by φ .

In this situation, the following is a special instance of [39, Theorem 1.2].

Proposition C.1. *The following hold:*

- (1) *The family of vector spaces $\text{coker } \delta_{u \circ \varphi}^N$, as $[\tilde{\Sigma}, \tilde{\nu}, \tilde{j}, \varphi] \in \bar{\mathcal{M}}_{[\Sigma],h}(\Sigma, j)$ varies, forms an oriented orbundle $\mathfrak{D}_h(\Sigma, j, u) \rightarrow \bar{\mathcal{M}}_{[\Sigma],h}(\Sigma, j)$, called the obstruction bundle.*
- (2) *Denoting by $[\bar{\mathcal{M}}_{[\Sigma],h}(\Sigma, j)]^{\text{vir}}$ the virtual fundamental class and denoting by $e(\mathfrak{D}_h(\Sigma, j, u))$ the Euler class of the obstruction bundle, we have*

$$\begin{aligned} & \text{GW}_{A,h}(X, \omega) \\ &= \sum_{g=0}^h \sum_{[u] \in \mathcal{M}_{A,g}^*(X, J)} \text{sign}(\Sigma, j, u) \langle e(\mathfrak{D}_h(\Sigma, j, u)), [\bar{\mathcal{M}}_{[\Sigma],h}(\Sigma, j)]^{\text{vir}} \rangle. \end{aligned}$$

Pandharipande [28, Section 2.3] proved that for $g := g(\Sigma)$,

$$\sum_{h=g}^{\infty} \langle e(\mathfrak{D}_h(\Sigma, j, u)), [\bar{\mathcal{M}}_{[\Sigma],h}(\Sigma, j)]^{\text{vir}} \rangle t^{2h-2} = t^{2g-2} \left(\frac{\sin(t/2)}{t/2} \right)^{2g-2}$$

Therefore, after changing the order of summation $\sum_{h=0}^{\infty} \sum_{g=0}^h = \sum_{g=0}^{\infty} \sum_{h=g}^{\infty}$, we obtain

$$\sum_{h=0}^{\infty} \text{GW}_{A,h}(X, \omega) t^{2h-2} = \sum_{g=0}^{\infty} n_{A,g}(X, \omega) t^{2g-2} \left(\frac{\sin(t/2)}{t/2} \right)^{2g-2}.$$

Since the numbers $\text{BPS}_{A,g}(X, \omega)$ are uniquely determined by the Gopakumar–Vafa formula (1.9) (see [3, Section 2]), $n_{A,g}(X, \omega) = \text{BPS}_{A,g}(X, \omega)$.

Acknowledgments. We thank Aleksei Zinger for several discussions about [38] and Eleny Ionel for answering our questions about [19]. We thank the referees for their meticulous work and, especially, for alerting us to an issue in an earlier version of Section 2.4 and its implications for Section 5.7.

Funding. This material is based upon work supported by the National Science Foundation under grant no. 1754967, an Alfred P. Sloan Research Fellowship, the Simons Collaboration on Special Holonomy in Geometry, Analysis, and Physics, and the Simons Society of Fellows.

References

- [1] E. Arbarello, M. Cornalba, and P. A. Griffiths, *Geometry of algebraic curves. Volume II*. Grundlehren Math. Wiss. 268, Springer, Heidelberg, 2011 Zbl 1235.14002 MR 2807457
- [2] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris, *Geometry of algebraic curves. Volume I*. Grundlehren Math. Wiss. 267, Springer, New York, 1985 Zbl 0559.14017 MR 770932
- [3] J. Bryan and R. Pandharipande, BPS states of curves in Calabi–Yau 3-folds. *Geom. Topol.* **5** (2001), 287–318 Zbl 1063.14068 MR 1825668
- [4] G. Castelnuovo, Ricerche di geometria sulle curve algebriche. *Atti della Accademia delle Scienze di Torino* **24** (1889), 346–373 Zbl 21.0669.01
- [5] P. Deligne and D. Mumford, The irreducibility of the space of curves of given genus. *Inst. Hautes Études Sci. Publ. Math.* (1969), no. 36, 75–109 Zbl 0181.48803 MR 262240
- [6] A. Doan, E.-N. Ionel, and T. Walpuski, The Gopakumar–Vafa finiteness conjecture. 2021, arXiv:2103.08221
- [7] A. Doan and T. Walpuski, Castelnuovo’s bound and rigidity in almost complex geometry. *Adv. Math.* **379** (2021), article no. 107550 Zbl 1460.53033 MR 4199270
- [8] A. Doan and T. Walpuski, Equivariant Brill–Noether theory for elliptic operators and super-rigidity of J -holomorphic maps. *Forum Math. Sigma* **11** (2023), article no. e3 Zbl 07643021 MR 4537309
- [9] T. Ekhholm and V. Shende, Skeins on branes. 2019, arXiv:1901.08027
- [10] R. Gopakumar and C. Vafa, M-theory and topological strings. I. 1998, arXiv:9809187
- [11] R. Gopakumar and C. Vafa, M-theory and topological strings. II. 1998, arXiv:9812127
- [12] A. Gournay, Complex surfaces and interpolation on pseudo-holomorphic cylinders. 2009, arXiv:1006.1775
- [13] M. Gromov, Pseudo holomorphic curves in symplectic manifolds. *Invent. Math.* **82** (1985), no. 2, 307–347 Zbl 0592.53025 MR 809718
- [14] R. Hartshorne, *Deformation theory*. Grad. Texts in Math. 257, Springer, New York, 2010 Zbl 1186.14004 MR 2583634
- [15] S. Hosono, M.-H. Saito, and A. Takahashi, Relative Lefschetz action and BPS state counting. *Internat. Math. Res. Notices* (2001), no. 15, 783–816 Zbl 1060.14017 MR 1849482

- [16] M.-x. Huang, S. Katz, and A. Klemm, [Topological string on elliptic CY 3-folds and the ring of Jacobi forms](#). *J. High Energy Phys.* (2015), no. 10, article no. 125
Zbl [1388.81219](#) MR [3435514](#)
- [17] C. Hummel, [Gromov’s compactness theorem for pseudo-holomorphic curves](#). *Prog. Math.* 151, Birkhäuser, Basel, 1997 Zbl [0870.53002](#) MR [1451624](#)
- [18] E.-N. Ionel, [Genus 1 enumerative invariants in \$\mathbf{P}^n\$ with fixed \$j\$ invariant](#). *Duke Math. J.* **94** (1998), no. 2, 279–324 Zbl [0974.14038](#) MR [1638587](#)
- [19] E.-N. Ionel and T. H. Parker, [The Gopakumar–Vafa formula for symplectic manifolds](#). *Ann. of Math. (2)* **187** (2018), no. 1, 1–64 Zbl [1459.53078](#) MR [3739228](#)
- [20] P. Kahn, [Pseudohomology and homology](#). 2001, arXiv:[0111223](#)
- [21] Y. Kiem and J. Li, [Categorification of Donaldson–Thomas invariants via perverse sheaves](#). 2012, arXiv:[1212.6444](#)
- [22] J. Knapp, E. Scheidegger, and T. Schimannek, [On genus one fibered Calabi–Yau threefolds with 5-sections](#). 2021, arXiv:[2107.05647](#)
- [23] D. Maulik and Y. Toda, [Gopakumar–Vafa invariants via vanishing cycles](#). *Invent. Math.* **213** (2018), no. 3, 1017–1097 Zbl [1400.14141](#) MR [3842061](#)
- [24] D. McDuff and D. Salamon, [J-holomorphic curves and symplectic topology](#). Second edn., Amer. Math. Soc. Colloq. Publ. 52, American Mathematical Society, Providence, RI, 2012 Zbl [1272.53002](#) MR [2954391](#)
- [25] J. Niu, [Refined convergence for genus-two pseudo-holomorphic maps](#). PhD thesis, State University of New York at Stony Brook, 2016
- [26] Y.-G. Oh and K. Zhu, [Embedding property of J-holomorphic curves in Calabi–Yau manifolds for generic J](#). *Asian J. Math.* **13** (2009), no. 3, 323–340 Zbl [1193.53180](#) MR [2570442](#)
- [27] R. Pandharipande, [A note on elliptic plane curves with fixed j-invariant](#). 1995, arXiv:[9505023](#)
- [28] R. Pandharipande, [Hodge integrals and degenerate contributions](#). *Comm. Math. Phys.* **208** (1999), no. 2, 489–506 Zbl [0953.14036](#) MR [1729095](#)
- [29] R. Pandharipande and R. P. Thomas, [Curve counting via stable pairs in the derived category](#). *Invent. Math.* **178** (2009), no. 2, 407–447 Zbl [1204.14026](#) MR [2545686](#)
- [30] R. Pandharipande and R. P. Thomas, [Stable pairs and BPS invariants](#). *J. Amer. Math. Soc.* **23** (2010), no. 1, 267–297 Zbl [1250.14035](#) MR [2552254](#)
- [31] J. Pardon, [An algebraic approach to virtual fundamental cycles on moduli spaces of pseudo-holomorphic curves](#). *Geom. Topol.* **20** (2016), no. 2, 779–1034 Zbl [1342.53109](#) MR [3493097](#)
- [32] T. H. Parker and J. G. Wolfson, [Pseudo-holomorphic maps and bubble trees](#). *J. Geom. Anal.* **3** (1993), no. 1, 63–98 Zbl [0759.53023](#) MR [1197017](#)
- [33] J. W. Robbin and D. A. Salamon, [A construction of the Deligne–Mumford orbifold](#). *J. Eur. Math. Soc. (JEMS)* **8** (2006), no. 4, 611–699 Zbl [1105.32011](#) MR [2262197](#)
- [34] M. Schwarz, [Equivalences for Morse homology](#). In *Geometry and topology in dynamics (Winston-Salem, NC, 1998/San Antonio, TX, 1999)*, pp. 197–216, Contemp. Math. 246, Amer. Math. Soc., Providence, RI, 1999 Zbl [0951.55009](#) MR [1732382](#)

- [35] C. H. Taubes, *SW \Rightarrow Gr: from the Seiberg–Witten equations to pseudo-holomorphic curves*. *J. Amer. Math. Soc.* **9** (1996), no. 3, 845–918 Zbl 0867.53025 MR 1362874
- [36] R. Ye, *Gromov’s compactness theorem for pseudo holomorphic curves*. *Trans. Amer. Math. Soc.* **342** (1994), no. 2, 671–694 Zbl 0810.53024 MR 1176088
- [37] A. Zinger, *Pseudocycles and integral homology*. *Trans. Amer. Math. Soc.* **360** (2008), no. 5, 2741–2765 Zbl 1213.57031 MR 2373332
- [38] A. Zinger, *A sharp compactness theorem for genus-one pseudo-holomorphic maps*. *Geom. Topol.* **13** (2009), no. 5, 2427–2522 Zbl 1174.14012 MR 2529940
- [39] A. Zinger, *A comparison theorem for Gromov–Witten invariants in the symplectic category*. *Adv. Math.* **228** (2011), no. 1, 535–574 Zbl 1225.14046 MR 2822239

Received 17 January 2022.

Aleksander Doan

Department of Mathematics, University College London, London WC1E 6BT; and Trinity College, Cambridge CB2 1TQ, UK; a.doan@ucl.ac.uk

Thomas Walpuski

Institut für Mathematik, Humboldt-Universität, Unter den Linden 6, 10099 Berlin, Germany; walpuski@math.hu-berlin.de