# Duality of orthogonal and symplectic random tensor models

Razvan Gurau and Hannes Keppler

**Abstract.** The groups O(N) and Sp(N) are related by an analytic continuation to negative values of N,  $O(-N) \simeq Sp(N)$ . This duality has been studied for vector models, SO(N) and Sp(N) gauge theories, as well as some random matrix ensembles. We extend this duality to real random tensor models of arbitrary order D with no symmetry under permutation of the indices and with quartic interactions. The N to -N duality is shown to hold graph by graph to all orders in perturbation theory for the partition function, the free energy and the connected two-point function.

# 1. Introduction and conclusion

Dualities are non-trivial relations between seemingly different models and therefore of great use in physics and mathematics. It has been known for some time [41] that, for even N, SO(N) and Sp(N) gauge theories are related by changing N to -N and that one can make sense of the relation SO(-N)  $\simeq$  Sp(N) for the representations of the respective groups [14]. This duality has furthermore been shown to hold between orthogonal and symplectic matrix ensembles [43]. (These correspond to the O(N)  $\otimes$ O(N) and Sp(N)  $\otimes$  Sp(N) matrix models of Section 3.)

The N to -N duality inspired in part the conjectured holographic duality between Vasiliev's higher spin gravity [50] in four-dimensional de Sitter space and the threedimensional Euclidean Sp(N) vector model with anticommuting scalars [2]. This dS/CFT correspondence is in turn based on the conjectured Giombi–Klebanov–Polyakov–Yin duality [22, 35] relating the three-dimensional O(N) vector model in the large N limit to Vasiliev gravity in four-dimensional anti-de Sitter space. In this context,  $N \sim (\Lambda G_N)^{-1}$  so that the sign change of the cosmological constant  $\Lambda$  (holding  $G_N$  fixed) is accompanied by a change  $N \rightarrow -N$ .

The perturbative expansion of random matrix models is a sum over *ribbon graphs* representing topological surfaces. The weight of each graph is fixed by the Feynman rules and the perturbative series can be organized [48] as a topological expan-

<sup>2020</sup> Mathematics Subject Classification. Primary 81T32; Secondary 57M15.

Keywords. Tensor models, random tensors, symplectic group, matrix models, ribbon graphs.

sion in 1/N. Random matrices yield a theory of random two-dimensional topological surfaces relevant for the study of conformal field theories (CFTs) coupled to two-dimensional Liouville gravity [6, 16, 17, 34, 37] and two-dimensional Jackiw– Teitelboim gravity [33, 45, 47]. They have applications as combinatorial generating functions to several counting problems [4, 49, 53] and to the intersection theory on the moduli space of Riemann surfaces [38, 44, 51].

Random matrices generalize to random tensor models [1, 27, 28, 30] of higher order<sup>1</sup> D which are probability measures of the type

$$d\mu[T] = e^{-S[T]} \prod_{(a_1,\dots,a_D)} \frac{dT^{a_1\dots a_D}}{\sqrt{2\pi}}$$

where the action S[T] is build out of invariants under some symmetry transformation. These models can also be viewed as 0-dimensional quantum field theories. The Feynman graphs of such models can be interpreted as higher-dimensional cellular complexes and the perturbative series can be reorganized as a series in 1/N [3, 5, 9, 10, 12, 25] which is not topological for  $D \ge 3$ . Zero-dimensional random tensors yield a framework for the study random topological spaces; in one-dimension tensor models provide an alternative to the Sachdev–Ye–Kitaev model without quenched disorder [52]; in higher dimensions, they lead to tensor field theories and a new class of large N melonic conformal field theories [8, 20, 21, 31, 36].

**Main result.** In this paper, we deal with tensors with D indices (i.e., of order D) with no symmetry under their permutations. The position of an index is called its color c, with c = 1, 2, ..., D. The tensors transform in the tensor product of D fundamental representations of O(N) and/or Sp(N), i.e., each tensor index is transformed by a different O(N) or Sp(N) matrix. The tensor components are real Grassmann valued (anticommuting, odd) if the number of Sp(N) factors is odd and real bosonic (commuting, even) if this number is even.<sup>2</sup> We assign a parity to the tensor indices: |c| = 0 or |c| = 1 if the index transforms under  $O(N_c)$  or  $Sp(N_c)$ , respectively. We consider actions consisting in invariants up to quartic order (see Section 2 for more details).

<sup>&</sup>lt;sup>1</sup>In the physics literature, one often uses "rank" instead of order, but this may lead to confusion with the many notions of tensor rank in abstract algebra.

<sup>&</sup>lt;sup>2</sup>The tensors are even multilinear maps on  $\mathbb{R}^{m|n}$ , the real graded supervector space with *m* even and *n* odd directions. This is natural because the orthosymplectic super Lie group OSp(m, n) contains both O(m) and Sp(n) and acts on  $\mathbb{R}^{m|n}$ . This will be our guideline in constructing the models of interest.

**Definition 1.1.** The real quartic graded tensor model, where "graded" refers to symmetry under

$$\boldsymbol{O}_1(N_1) \otimes \boldsymbol{O}_2(N_2) \otimes \cdots \otimes \boldsymbol{O}_D(N_D), \quad \boldsymbol{O}_c(N_c) = \begin{cases} O(N_c), & |c| = 0, \\ Sp(N_c), & |c| = 1 \end{cases}$$

is defined by the measure

$$d\mu[T] \simeq e^{-S[T]} \prod_{a_1,...,a_D} dT^{a_1...a_D},$$
  
$$S[T] = \frac{1}{2} \left( T^{a_1...a_D} T^{b_1...b_D} \prod_{c=1}^D g^c_{a_c b_c} \right) + \sum_{q \in \Omega} \frac{\lambda_q}{4} I^q(T).$$

where  $g_{a_cb_c}^c$  is the Kronecker delta  $\delta_{a_cb_c}$  for |c| = 0 or the canonical symplectic form  $\omega_{a_cb_c}$  for |c| = 1 and the sum over  $\Omega$  runs over all the independent quartic trace invariants  $I^q(T)$ .

The partition function Z and the connected two-point function  $G_2$  of the model are defined by

$$Z(\lambda) = \int d\mu[T] \text{ and } G_2(\lambda) = \frac{1}{Z} \int d\mu[T] T^{a_1...a_D} T^{b_1...b_D} \prod_{c=1}^D g^c_{a_c b_c}$$

and can be evaluated in a perturbative expansion. Our main theorem is the following.

**Theorem 1.2.** The perturbative series of the free energy  $\ln Z$  and of the connected two-point function  $G_2$  can be expressed as formal sums over connected, colored multiribbon graphs

$$\ln Z(\lambda) = \sum_{\substack{[\mathbb{G}] \text{ connected, rooted,} \\ \text{at least one } E_q > 0}} \frac{1}{2^{C(\mathbb{G} - \mathcal{E}^{\varrho}) + 1} \sum_{q \in \mathcal{Q}} E_q}} \mathcal{A}(\mathbb{G}),$$
$$G_2(\lambda) = \sum_{\substack{[\mathbb{G}] \text{ connected, rooted}}} \frac{1}{2^{C(\mathbb{G} - \mathcal{E}^{\varrho}) - 1}} \mathcal{A}(\mathbb{G})$$

with amplitude

$$\mathcal{A}(\mathbb{G}) = 2^{E^{\varrho}(\mathbb{G})} \prod_{q \in \Omega} (-\lambda_q)^{E_q(\mathbb{G})} \prod_{c=1}^{D} ((-1)^{|c|} N_c)^{F_c(\mathbb{G})},$$
(1.1)

where  $E_q$ ,  $E^{\varrho}$ ,  $F_c$ ,  $C(\mathbb{G} - \mathcal{E}^{\varrho})$  are some combinatorial numbers associated to the multi-ribbon graph  $\mathbb{G}$  (see Section 4.2 for the relevant definitions).

*Proof.* The theorem follows from equations (4.3), (4.5) and (4.6).

The crucial remark is that all the factors  $N_c$  come in the form  $(-1)^{|c|}N_c$ , hence each term is mapped into itself by exchanging  $O(N_c) \leftrightarrow Sp(N_c)$  and  $N_c \leftrightarrow -N_c$ .

**Conclusion and outlook.** We list some comments on, and possible generalizations of, our result:

- In order to prove our main theorem, we will use in this paper an intermediate field representation adapted to quartic interactions. It should however be possible to extend this result to more general interactions [40].
- While more general models with OSp(*m*, *n*) symmetry could be considered, the construction of super tensor actions is complicated because of the abundance of sign factors [46].
- For D = 2 (matrices), the contributions of ribbon graphs and their duals cancel exactly in the fermionic case (see Remark 3.1). It would be interesting to understand similar cancellations in the graded tensor models. This should be related to Poincaré duality between lower-dimensional colored subgraphs.
- One should explore the implications of the N → -N duality for tensor field theories. The sign changes may generate new renormalization group fixed points, and the duality may not hold for all the physical properties [39]. Quantum mechanical models of order three tensors with Sp(N) symmetry have been studied in [11,24].

**Outline of the paper.** This paper is organized as follows. In Section 2, the quartic graded tensor model is defined, the relation between directed edge colored graphs and quartic trace invariants is explained, and we collect some definitions and notations on ribbon graphs. Section 3 deals in detail with the order 2 (matrix) case. Section 4 continues with the general case of arbitrary order *D* tensors. Appendix A contains the calculation of the sign of each ribbon graph amplitude and Appendix B gives details on the calculation of the symmetry factors of the Feynman graphs.

# 2. Definitions

In this section, we define the models we will be studying. We also give some standard definitions about ribbon graphs and combinatorial maps.

### 2.1. The real quartic graded tensor models

The orthosymplectic super Lie group OSp(m, n) is the isometry group of the canonical graded-symmetric bilinear form on the supervector space  $\mathbb{R}^{m|n}$ ,

$$\eta: \mathbb{R}^{m|n} \times \mathbb{R}^{m|n} \to \Lambda_{\infty}, \quad (\eta_{ij}) = \left(\frac{\delta \mid 0}{0 \mid \omega}\right),$$

where  $\Lambda_{\infty}$  is the Grassmann algebra generated by an infinite number of anticommuting generators. The space  $\mathbb{R}^{m|n}$  is a free module over  $\Lambda_{\infty}$  with *m* even (commuting) and *n* odd (anticommuting) basis vectors. Note that non-singularity of  $\eta$  demands that *n* is an even integer. For later comparison, *m* is also taken to be even.

Since we are only interested in O(N) and Sp(N), and not the whole OSp(m, n), we restrict to supervector spaces that are either purely odd or purely even, and thus have either O(N) or Sp(N) as their isometry group. This information can be encoded as a parity of the index color  $|c| \in \{0, 1\}$ , with |c| = 0 corresponding to orthogonal, and |c| = 1 to symplectic symmetry. The tensor components are commuting bosonic or anticommuting Grassmannian, depending on whether the number of indices with |c| = 1 is even or odd. Suitable invariants are defined to construct the actions of the models.

**Vector spaces.** Let  $\mathcal{H}_c = \mathbb{R}^{N_c|0}$  for |c| = 0,  $\mathcal{H}_c = \mathbb{R}^{0|N_c}$  for |c| = 1, respectively, be a real supervector space of dimension  $N_c$ , that is, either purely even or purely odd and is endowed with a non-degenerate graded symmetric inner product  $g^c: \mathcal{H}_c \times \mathcal{H}_c \to \Lambda_\infty$ ,

$$g^{c}(u,v) = (-1)^{|c|} g^{c}(v,u) \quad \forall u,v \in \mathcal{H}_{c}, \quad g^{c}(\cdot,v) = 0 \Leftrightarrow v = 0.$$

In the standard basis  $g^c$  agrees with the standard symmetric or symplectic form, that is,  $g^c_{a_cb_c} = \delta_{a_cb_c}$  for |c| = 0,  $g^c_{a_cb_c} = \omega_{a_cb_c}$  for |c| = 1, respectively. We denote by  $(g^c)^{a_cb_c}$  the matrix element of the inverse  $(g^c)^{-1}$ . The isometry group preserving  $g^c$  is either  $O(N_c)$  in the |c| = 0 case or  $Sp(N_c)$  in the |c| = 1 case, denoted collectively by  $\boldsymbol{O}_c(N_c) := \{O_c \mid g^c_{a_cb_c} = O^{a'_c}_{a_c} O^{b'_c}_{b'_c} g^c_{a'_cb'_c} = (Og^c O^{\mathsf{T}})_{a_cb_c}\}.$ 

**Tensors.** Tensors are *even* elements of the tensor product space  $T \in \bigotimes_{c=1}^{D} \mathcal{H}_c$ . Choosing a basis  $\{[\psi^c]_{a_c}\}_{a_c=1,...,N_c}$  in each  $\mathcal{H}_c$  and denoting by  $\{[\psi_c^{\vee}]_{a_c}\}_{a_c=1,...,N_c}$  the dual basis, the components of a tensor are

$$T^{a_1\dots a_D} \equiv T([\psi_1^{\vee}]^{a_1},\dots,[\psi_D^{\vee}]^{a_D}),$$
  
$$T = \sum_{a_c=1,\dots,N_c \ \forall c} T^{a_1\dots a_D} \ [\psi^1]_{a_1} \otimes \dots \otimes [\psi^D]_{a_D}.$$

A generic tensor has no symmetry properties under permutation of its indices  $a^1, \ldots, a^D$ , hence the indices have a well-defined position c called their *color*. The set of colors is denoted by  $\mathcal{D} = \{1, \ldots, D\}$ . We sometimes call the colors with |c| = 0 even and the ones with |c| = 1 odd. As the tensors are taken to be even elements of the tensor product space, the tensor components are bosonic (even) if the number of colors with |c| = 1 (i.e., odd colors) is even and fermionic (odd) otherwise: the Grassmann number  $T^{a_1...a_D}$  has the same parity as  $\sum_{c \in \mathcal{D}} |c|$ .

The tensors transform in the tensor product representation of several orthogonal and symplectic groups according to the type of the individual  $\mathcal{H}_c$ 's:

$$T^{a_1\dots a_D} \to (O_1)^{a_1}_{b_1} \cdots (O_D)^{a_D}_{b_D} T^{b_1\dots b_D}, \quad O_1 \otimes \cdots \otimes O_D \in \bigotimes_{c \in \mathcal{D}} \mathcal{O}_c(N_c).$$

A tensor can be viewed as a multilinear map  $T: \bigotimes_{c \in \mathcal{C}} \mathcal{H}_c^{\vee} \to \bigotimes_{c \in \mathcal{D} \setminus \mathcal{C}} \mathcal{H}_c$  for any subset of colors  $\mathcal{C} \subset \mathcal{D}$ . As the inner product,  $g^c$  induces an isomorphism between  $\mathcal{H}_c$  and its dual, denoting by  $a_{\mathcal{C}} = (a_c, c \in \mathcal{C})$ , the matrix elements of this linear map in the tensor product basis are  $T^{a_{\mathcal{D} \setminus \mathcal{C}} a_{\mathcal{C}}} \equiv T^{a_1 \dots a_D}$ .

Edge colored graphs. Invariant polynomials in the tensor components can be constructed by contracting the indices of color c with the inner product  $g^c$ . The unique quadratic invariant is

$$g^{\otimes \mathcal{D}}(T,T) := T^{a_{\mathcal{D}}} T^{b_{\mathcal{D}}} \prod_{c \in \mathcal{D}} g^{c}_{a_{c}b_{c}}.$$

General *trace invariants* are polynomials in the  $T^{a_{\mathcal{D}}}$ 's build by contracting pairs of indices of the same color. These invariants form an algebraic complete set for all invariant polynomials and admit a straightforward graphical representation as *edge colored graphs*.

**Definition 2.1** (Edge colored graphs [28]). A closed edge *D*-colored graph is a graph  $\mathcal{B} = (\mathcal{V}(\mathcal{B}), \mathcal{E}(\mathcal{B}))$  with vertex set  $\mathcal{V}(\mathcal{B})$  and edge set  $\mathcal{E}(\mathcal{B})$  such that

- The edge set is partitioned into *D* disjoint subsets  $\mathscr{E}(\mathscr{B}) = \bigsqcup_{c=1}^{D} \mathscr{E}^{c}(\mathscr{B})$ , where  $\mathscr{E}^{c}(\mathscr{B}) \ni e^{c} = (v, w), v, w \in \mathcal{V}(\mathscr{B})$ , is the subset of edges of color *c*.
- All vertices are *D*-valent with all the edges incident to a vertex having distinct colors.

In order to incorporate the odd colors appropriately, one needs to consider *directed graphs*, that is, graphs with an additional arrow for every edge (see Figure 1 for an example). Two graphs which are identical up to reorienting one edge of an odd color represent the same invariant up to a global "—" sign. We will fix the global sign in the case of quartic invariants below.

**Quartic invariants.** Quartic invariants are represented by *D*-colored graphs with four vertices (see Figure 1) and directed edges. Due to the sign ambiguity induced by reversing the edges corresponding to the odd colors, we need to give a prescription to fix the global sign of an invariant. Every directed quartic *D*-colored graph can be canonically oriented as follows (see again Figure 1):

• The color 1 edges give a pairing of the vertices. We denote by  $a^1$  and  $b^1$  the source vertices of the oriented edges 1 and  $a^2$  and  $b^2$  their targets.



Figure 1. Left: Quartic 5-colored graph. Right: Schematic representation of a general quartic invariant.

- We orient all the edges that connect (a<sup>1</sup>, a<sup>2</sup>) (resp. (b<sup>1</sup>, b<sup>2</sup>)) parallel to the edges 1.
   We denote their colors by c ∈ D \ C.
- All the edges of colors  $c \in \mathcal{C}$  connect the *a* pair with the *b* pair. We orient all of them from the *a* pair to the *b* pair. These edges fall into two classes:
  - either they connect  $a^1$  with  $b^1$  and  $a^2$  with  $b^2$  in which case we say they run in the *parallel* channel,
  - or they connect  $a^1$  with  $b^2$  and  $a^2$  with  $b^1$  in which case we say they run in the *cross* channel.

A canonically oriented graph is indexed by a subset of colors  $\mathcal{C} \subset \mathcal{D}$ ,  $1 \notin \mathcal{C}$  and permutations of two elements  $\pi^c \in \mathfrak{S}_2 = \{id, (12)\}, c \in \mathcal{C}$ . The associated invariant is

$$I(T) = \sum_{a_{\mathcal{D}}^{1}, a_{\mathcal{D}}^{2}, b_{\mathcal{D}}^{1}, b_{\mathcal{D}}^{2}} \left( T^{a_{\mathcal{D}}^{1}} T^{a_{\mathcal{D}}^{2}} \prod_{c \in \mathcal{D} \setminus \mathcal{C}} g^{c}_{a_{c}^{1} a_{c}^{2}} \right) \left( T^{b_{\mathcal{D}}^{1}} T^{b_{\mathcal{D}}^{2}} \prod_{c \in \mathcal{D} \setminus \mathcal{C}} g^{c}_{b_{c}^{1} b_{c}^{2}} \right) \\ \times \left( \prod_{c \in \mathcal{C}} (-\operatorname{sgn}(\pi^{c}))^{|c|} g^{c}_{a_{c}^{1} b_{c}^{\pi^{c}}(1)} g^{c}_{a_{c}^{2} b_{c}^{\pi^{c}}(2)} \right) \\ = \sum_{a_{\mathcal{C}}^{1}, a_{\mathcal{C}}^{2}, b_{\mathcal{C}}^{1}, b_{\mathcal{C}}^{2}} (g^{\otimes \mathcal{D} \setminus \mathcal{C}} (T, T))^{a_{\mathcal{C}}^{1} a_{\mathcal{C}}^{2}} K_{a_{\mathcal{C}}^{1} a_{\mathcal{C}}^{2}, b_{\mathcal{C}}^{1} b_{\mathcal{C}}^{2}} (g^{\otimes \mathcal{D} \setminus \mathcal{C}} (T, T))^{b_{\mathcal{C}}^{1} b_{\mathcal{C}}^{2}}, \quad (2.1)$$

where we introduced the shorthand notation K for the contractions of the indices transmitted between the pairs. Note that this is invariant by exchanging the b vertices and that  $(-\operatorname{sgn}(\pi^c))^{|c|}$  is the signature of the permutation  $(a^1a^2)(b^1b^2)$  to  $(a^1b^{\pi(1)})(a^2b^{\pi(2)})$  for the odd colors.

**Lemma 2.2.** There are  $\frac{1+3^{D-1}}{2}$  different quartic trace invariants (see Figure 2 for the D = 3 case).

*Proof.* There is only one invariant corresponding to  $\mathcal{C} = \emptyset$ . If  $\mathcal{C}$  has q elements, there are  $2^q$  choices for the channels and an overall  $\frac{1}{2}$  for the relabeling of the b vertices. Thus the total number of invariants is

$$1 + \frac{1}{2} \sum_{q=1}^{D-1} {D \choose q} 2^q = \frac{3^{D-1} + 1}{2}.$$



Figure 2. The 5 quartic invariants at order 3 known as double trace, pillow and tetrahedron.

Denote the set of distinct quartic *D*-colored graphs and the associate trace invariants by  $\mathfrak{Q} \ni q$  and  $I^q(T)$ , respectively.

**Definition 2.3** (Real quartic graded tensor model). The *real quartic "graded" tensor model* is the measure

$$d\mu[T] = e^{-S[T]} [dT],$$
  

$$[dT] = \prod_{a_{\mathcal{D}}} dT^{a_1...a_D} \cdot \begin{cases} \frac{1}{(2\pi)^{\prod_c N_c/2}}, & \sum_{c=1}^D |c| = 0 \mod 2, \\ 1, & \sum_{c=1}^D |c| = 1 \mod 2 \end{cases}$$

with

$$S[T] = \frac{1}{2}g^{\otimes \mathcal{D}}(T,T) + \sum_{q \in \mathcal{Q}} \frac{\lambda_q}{4}I^q(T),$$

where the normalization is such that

$$\int d\mu[T] = 1 \quad \text{for } \lambda_q = 0 \ \forall q \in \Omega.$$

**Convergence issues.** Throughout this paper, we treat the measures  $d\mu[T]$  as perturbed Gaussian measures. As such we do not concern ourselves with the convergence of the various tensor and matrix integrals. The integrals are always convergent if T is fermionic. If T is bosonic, the integrals converge if |c| = 0 for all c, but not necessarily in the other cases. As we treat the Gaussian integrals as generating functions of graphs, we will not worry about such issues.

#### 2.2. Ribbon graphs and combinatorial maps

As ribbon graphs [19,23] and combinatorial maps play a significant role in the derivation of our results, we review here some of their properties.

Ribbon graphs, see Figure 3 for some examples, are cellularly embedded graphs on topological surfaces, and thus can be viewed as 2-cell-complexes. Due to the embedding, each vertex carries an orientation and the order of edges around a vertex is fixed. A vertex can be re-embedded with the opposite orientation: this amounts to reversing the order of the incident edges and giving them a twist, see Figure 4.



**Figure 3.** Ribbon graphs, which we denote by  $\mathscr{G}_{T^2}$  and  $\mathscr{G}_{\mathbb{RP}^2}$ , and their cellular embeddings. The rightmost surface is the hemisphere representation of the real projective plane where opposite points along the equator are identified.



**Figure 4.** Re-embedding a vertex: the order of halfedges is reversed and they gain additional twists. This is an equivalence relation of ribbon graphs.

**Definition 2.4** (Ribbon graph [19]). A *ribbon graph*  $\mathcal{G} = (\mathcal{V}(\mathcal{G}), \mathcal{E}(\mathcal{G}))$  is a (possibly non-orientable) surface with boundary, represented as the union of two sets of topological discs, a set of *vertices*  $\mathcal{V}(\mathcal{G})$ , and a set of *edges*  $\mathcal{E}(\mathcal{G})$  such that

- (1) The vertices and edges intersect in disjoint line segments.
- (2) Each such line segment lies on the boundary of precisely one vertex and precisely one edge.
- (3) Every edge contains exactly two such line segments.

The boundary components of  $\mathscr{G}$  are called *faces*. The two disjoint boundary segments of an edge that are not connected to a vertex (i.e., the two sides of the edge) are called *strands*. We denote the set of faces of  $\mathscr{G}$  by  $\mathscr{F}(\mathscr{G})$ . A ribbon graph becomes a two-dimensional CW complex by sewing two-dimensional patches along its faces.

The numbers of vertices, edges and faces of  $\mathcal{G}$  are denoted by  $V(\mathcal{G})$ ,  $E(\mathcal{G})$  and  $F(\mathcal{G})$ , respectively.

Several remarks are in order:

- The strands of an edge can run parallel, in which case the edge is called untwisted, or cross, in which case the edge is called twisted.
- If  $V(\mathcal{G}) = 1$ , the graph is called a *rosette graph*. A rosette graph with only one face is called a *superrosette graph*.
- A self-loop in 𝔅 is an edge e = {h<sub>v</sub>, h'<sub>v</sub>} ∈ 𝔅(𝔅) connected to just one vertex v ∈ 𝒱(𝔅). A simple self-loop is a self-loop such that its halfedges are direct neighbors in the cyclic ordering around v, thus v has a corner of the form (h<sub>v</sub>, h'<sub>v</sub>). If e is (un-)twisted, the simple self-loop is called likewise.

We denote the ribbon graph consisting in only one vertex with no edge by 𝔅₀. By definition, this graph has one face. We denote the ribbon graph with one vertex and one twisted self-loop edge by 𝔅<sub>ℝP<sup>2</sup></sub> and the ribbon graph with one vertex, two untwisted self-loop edges but no simple self-loop by 𝔅<sub>T<sup>2</sup></sub>.<sup>3</sup> The last two graphs are depicted in Figure 3. As a topological surface with boundary 𝔅<sub>ℝP<sup>2</sup></sub> is homeomorphic to a Möbius strip.

Every ribbon graph has a dual ribbon graph with the same number of edges, but with the roles of the vertices and the faces interchanged.

**Definition 2.5** (Dual ribbon graph [19]). Let  $\mathscr{G}$  be a ribbon graph. The *dual* ribbon graph  $\mathscr{G}^*$  is obtained by sewing discs along the faces of  $\mathscr{G}$  and deleting the original vertex discs of  $\mathscr{G}$ . The new discs make up the dual vertex set  $\mathcal{V}(\mathscr{G}^*)$ , and the new boundary components created by the deletion are the faces of  $\mathscr{G}^*$ . See Figure 5 for an illustration.



Figure 5. The dual graphs.

Besides ribbon graphs, we will encounter combinatorial maps below.

**Definition 2.6** (Combinatorial map). A *combinatorial map*  $\mathcal{M} = (\mathcal{S}, \pi, \alpha)$  is a finite set  $\mathcal{S}$  of halfedges (or darts) of even cardinality, together with a couple of permutations  $(\pi, \alpha)$  on  $\mathcal{S}$ , where  $\alpha$  is an involution with no fixed points (a "pairing" of halfedges).

The map  $\mathcal{M}$  is called connected if the group freely generated by  $\pi$  and  $\alpha$  acts transitively on  $\mathcal{S}$ . The dual of  $\mathcal{M}$  is the combinatorial map  $\mathcal{M}^* = (\mathcal{S}, \alpha \circ \pi, \alpha)$ .

Combinatorial maps can be represented as graphs embedded in orientable surfaces. The cycles of  $\pi$  represent vertices with a cyclic order of their halfedges (chosen to be counter-clockwise), and  $\alpha$  encodes pairings of halfedges into edges. The faces of a combinatorial map are the cycles of the permutation  $\alpha \circ \pi$ . In the dual combinatorial map, the role of vertices and faces is reversed.

The definition of combinatorial maps and ribbon graphs can be extended to include a second kind of edges.

<sup>&</sup>lt;sup>3</sup>As their names suggest, these graphs can be cellularly embedded into  $\mathbb{RP}^2$  or  $T^2$ , respectively.

Definition 2.7 (Combinatorial map with *Q*-edges). A combinatorial map with *Q*-edges

$$\mathcal{M}^{\varrho} = (\mathcal{S} \sqcup \mathcal{S}^{\varrho}, \pi, \alpha, \alpha^{\varrho})$$

is a finite set  $\overline{S} = S \sqcup S^{\varrho}$  that is the disjoint union of two sets of halfedges, both of even cardinality, together with a triple of permutations  $(\pi, \alpha, \alpha^{\varrho})$  on  $\overline{S}$ . Here  $\alpha$  and  $\alpha^{\varrho}$  are fixed-point free involutions on S and  $S^{\varrho}$ , respectively, and extended to the whole of  $\overline{S}$  by setting  $\alpha(h) = h \forall h \in S^{\varrho}$  and analogous for  $\alpha^{\varrho}$ .

The cycles of  $\alpha^{\varrho}$  are pairs of halfedges in  $\mathcal{S}^{\varrho}$  which we will call  $\varrho$ -edges. The combinatorial map  $\mathcal{M}^{\varrho}$  is connected if the group freely generated by  $\pi$ ,  $\alpha$  and  $\alpha^{\varrho}$  acts transitively on  $\overline{\mathcal{S}}$ . The cycles of  $\pi$  are the vertices and the cycles of  $\pi \circ \alpha$  are the faces of  $\mathcal{M}^{\varrho}$ . The dual map is defined by changing the role of vertices and faces but not touching the  $\varrho$ -edges  $\mathcal{M}^{\varrho*} = (\mathcal{S} \sqcup \mathcal{S}^{\varrho}, \alpha \circ \pi, \alpha, \alpha^{\varrho})$ .

Deleting all the  $\rho$ -edges, one obtains an ordinary combinatorial map.

Ribbon graphs can be obtained from combinatorial maps by replacing their edges by twisted or untwisted ribbon edges. The same holds true for combinatorial maps with  $\rho$ -edges and *ribbon graphs with*  $\rho$ -edges.

**Definition 2.8** (Ribbon graph with *Q*-edges). A ribbon graph with *Q*-edges

$$\mathscr{G}^{\varrho} = (\mathcal{V}, \mathcal{E}, \mathcal{E}^{\varrho})$$

is a ribbon graph  $\mathscr{G} = (\mathcal{V}, \mathscr{E})$ , together with a set of line segments  $\mathscr{E}^{\varrho}$ , called  $\varrho$ -edges, such that their endpoints are connected to the corners of the ribbon graph. The ribbon graph  $\mathscr{G}^{\varrho}$  is called connected if it is connected as a topological space. The notions of faces, corners and edges of  $\mathscr{G}^{\varrho}$  refer to the ones of the ribbon graph  $\mathscr{G} = \mathscr{G}^{\varrho} - \mathscr{E}^{\varrho}$ , which is obtained by deleting the  $\varrho$ -edges.<sup>4</sup>

This dual of a ribbon graph with  $\rho$ -edges is obtained by performing the partial dual [13, 18, 19] with respect to the ribbon edges. This is the dual of the underlying ribbon graph obtained by ignoring the  $\rho$ -edges, where we keep track of the corners to which the  $\rho$ -edges are hooked.

### 3. Matrix models

We first deal with the case of matrices (order D = 2 tensors) in Definition 2.3. In particular,

$$M^{a_1a_2}M^{b_1b_2} = (-1)^{|1|+|2|}M^{b_1b_2}M^{a_1a_2}$$

<sup>&</sup>lt;sup>4</sup>Ribbon graphs with  $\rho$ -edges are embedded in nodal surfaces, that is, Riemann surfaces glued at marked points. The ribbon graphs encode closed topological surfaces and by identifying points that are connected by a  $\rho$ -edge, a gluing prescription is given.



Figure 6. Graphical representation of the matrix model invariants up to quartic order.

i.e., the models with mixed symmetry are fermionic. We show that, for each ribbon graph in the perturbative expansion of the free energy and the two-point function of the model, changing one (or both) of the symmetry group factors in the  $O(N_1) \otimes O(N_2)$ -model from O(N) to Sp(N) amounts to changing the sign accompanying the corresponding N factor.

Complex random matrix models in the intermediate field representation have been studied in [32]. The sign changes between the  $O(N_1) \otimes O(N_2)$ - and the  $Sp(N_1) \otimes Sp(N_2)$ -models have also been studied in [43] by different methods.

Denoting by the superscript T the transpose, the action of the real quartic graded matrix model is written as  $^5$ 

$$S[M] = \frac{1}{2} M^{a_1 a_2} g^1_{a_1 b_1} g^2_{a_2 b_2} M^{b_1 b_2} + \frac{\kappa}{4} (M^{a_1 a_2} g^1_{a_1 b_1} g^2_{a_2 b_2} M^{b_1 b_2})^2 + \frac{\lambda}{4} (-1)^{|2|} (M^{a_1^1 a_2^1} g^1_{a_1^1 a_1^2} M^{a_1^2 a_2^2}) g^2_{a_2^1 b_2^1} g^2_{a_2^2 b_2^2} (M^{b_1^1 b_2^1} g^1_{b_1^1 b_1^2} M^{b_1^2 b_2^2}) = \frac{1}{2} \operatorname{Tr}[M g^2 M^{\mathsf{T}}(g^1)^{\mathsf{T}}] + \frac{\kappa}{4} (\operatorname{Tr}[M g^2 M^{\mathsf{T}}(g^1)^{\mathsf{T}}])^2 + \frac{\lambda}{4} (-1)^{|1|} \operatorname{Tr}[(M g^2 M^{\mathsf{T}} g^1)^2],$$
(3.1)

where we note that the trace is  $Tr[A] = A_a^a = A^{ab}g_{ba}$ . This action is invariant under the transformation  $M \to O_1 X O_2^{\mathsf{T}}$  with  $O_1 \in \mathcal{O}_1(N_1)$ ,  $O_2 \in \mathcal{O}_2(N_2)$ . The three terms in (3.1) can be represented by 2-colored graphs or alternatively ribbon graphs, as depicted in Figure 6.

Whereas all terms in the action of the  $O(N_1) \otimes O(N_2)$ -model are positive for  $\kappa, \lambda \in \mathbb{R}_+$ , in the Sp( $N_1$ )  $\otimes$  Sp( $N_2$ )-model, this is only true for the  $\kappa$  term: the quadratic and the  $\lambda$  terms are in general indefinite.

<sup>&</sup>lt;sup>5</sup>In the pure Sp(N) case with  $g^1 = g^2 = \omega$ , the convergence of (3.1) is not clear, since the quadratic part has negative modes.

### 3.1. Intermediate field representation

The intermediate field (Hubbard–Stratonovich) representation is obtained by introducing an auxiliary field per quartic interaction and integrating out the original field. To be precise, we use that

$$\exp\left\{-\frac{\kappa}{4}(\operatorname{Tr}[Mg^{2}M^{\mathsf{T}}(g^{1})^{\mathsf{T}}])^{2} - \frac{\lambda}{4}(-1)^{|1|}\operatorname{Tr}[Mg^{2}M^{\mathsf{T}}g^{1}Mg^{2}M^{\mathsf{T}}g^{1}]\right\}$$
$$= \left[e^{\frac{1}{2}(\frac{\partial}{\partial\sigma}P\frac{\partial}{\partial\sigma})}e^{\frac{1}{2}\frac{\partial}{\partial\rho}\frac{\partial}{\partial\rho}}\exp\left\{-\iota\sqrt{\frac{\kappa}{2}}\operatorname{Tr}[Mg^{2}M^{\mathsf{T}}(g^{1})^{\mathsf{T}}]\rho\right]$$
$$-\iota\sqrt{\frac{\lambda}{2}}\operatorname{Tr}[Mg^{2}M^{\mathsf{T}}(g^{1}\sigma g^{1})^{\mathsf{T}}]\right\}_{\varrho,\sigma=0},$$
(3.2)

where  $\rho$  is a real commuting (bosonic) scalar field and  $\sigma = (-1)^{|1|} \sigma^{\mathsf{T}}$  is a (bosonic) real graded-symmetric matrix, and we introduce the shorthand notation:

$$\left(\frac{\partial}{\partial\sigma}P\frac{\partial}{\partial\sigma}\right) := \frac{\partial}{\partial\sigma^{ab}}P^{ab,dc}\frac{\partial}{\partial\sigma^{cd}}, \quad P^{ab,dc} = \frac{1}{2}(g_1^{ad}g_1^{bc} + (-1)^{|1|}g_1^{ac}g_1^{bd})$$

with *P* the (anti-)symmetric projector, taking into account the symmetry of the  $\sigma$  field. Note that  $g^1 M g^2 M^{\mathsf{T}} g^1$  has the same graded symmetry as  $\sigma$ .

Equation (3.2) is just a Gaussian integral over the intermediate fields  $\rho$  and  $\sigma$ . We favor here the notation of the Gaussian integral as a differential operator (see, for instance, [7]) for two reasons. First, the Gaussian integral is formal in some cases (that is, the covariance is not necessarily positively defined). Second, in this form the perturbative expansion of the Gaussian integral is straightforward. In order to prove (3.2), we expand the exponentials and commute the sum and the derivatives:

$$\begin{split} & \left[ e^{\frac{1}{2} \left(\frac{\partial}{\partial \sigma} P \frac{\partial}{\partial \sigma}\right)} e^{\frac{1}{2} \frac{\partial}{\partial \varrho} \frac{\partial}{\partial \varrho}} \sum_{n,p=0}^{\infty} \frac{\left(-\frac{\kappa}{2}\right)^n \left(-\frac{\lambda}{2}\right)^p}{(2n)!(2p)!} \right]_{\varrho,\sigma=0} \\ & \times \left( \operatorname{Tr}[Mg^2 M^{\mathsf{T}}(g^1)^{\mathsf{T}}] \varrho \right)^{2n} \left( \operatorname{Tr}[Mg^2 M^{\mathsf{T}}(g^1 \sigma g^1)^{\mathsf{T}}] \right)^{2p} \right]_{\varrho,\sigma=0} \\ & = \left[ \sum_{n,p=0}^{\infty} \frac{\left(-\frac{\kappa}{2}\right)^n \left(-\frac{\lambda}{2}\right)^p}{2^n n! 2^p p!(2n)!(2p)!} \left(\frac{\partial}{\partial \varrho} \frac{\partial}{\partial \varrho} \right)^n \left( \operatorname{Tr}[Mg^2 M^{\mathsf{T}}(g^1)^{\mathsf{T}}] \varrho \right)^{2n} \right. \\ & \times \left( \frac{\partial}{\partial \sigma} P \frac{\partial}{\partial \sigma} \right)^p \left( \operatorname{Tr}[Mg^2 M^{\mathsf{T}}(g^1 \sigma g^1)^{\mathsf{T}}] \right)^{2p} \right]_{\varrho,\sigma=0} \\ & = \left[ \sum_{n,p=0}^{\infty} \frac{\left(-\frac{\kappa}{4}\right)^n \left(-\frac{\lambda}{4}\right)^p}{n! p!} \left( \operatorname{Tr}[Mg^2 M^{\mathsf{T}}(g^1)^{\mathsf{T}}] \right)^{2n} \right. \\ & \times \left( \left(-1\right)^{|1|} \operatorname{Tr}[Mg^2 M^{\mathsf{T}}g^1 Mg^2 M^{\mathsf{T}}g^1] \right)^p \right]_{\varrho,\sigma=0}, \end{split}$$

where we used

$$[g^{1}Mg^{2}M^{\mathsf{T}}g^{1}]_{ab}g^{ad}g^{bc}[g^{1}Mg^{2}M^{\mathsf{T}}g^{1}]_{cd}$$
  
= (-1)<sup>|1|</sup> Tr[g^{1}Mg^{2}M^{\mathsf{T}}g^{1}Mg^{2}M^{\mathsf{T}}g^{1}].

The partition function now reads

$$Z(\kappa,\lambda) = \int [dM] e^{-\frac{1}{2}\operatorname{Tr}[Mg^2M^{\mathsf{T}}(g^1)^{\mathsf{T}}]} \Big[ e^{\frac{1}{2}(\frac{\partial}{\partial\sigma}P\frac{\partial}{\partial\sigma})} e^{\frac{1}{2}\frac{\partial}{\partial\varrho}\frac{\partial}{\partial\varrho}} \\ \times e^{-i\sqrt{\frac{\kappa}{2}}\operatorname{Tr}[Mg^2M^{\mathsf{T}}(g^1)^{\mathsf{T}}]} e^{-i\sqrt{\frac{\lambda}{2}}\operatorname{Tr}[Mg^2M^{\mathsf{T}}(g^1\sigma g^1)^{\mathsf{T}}]} \Big]_{\varrho,\sigma=0},$$

and all the terms containing M can be collected in a quadratic form using

$$\operatorname{Tr}(MAM^{\mathsf{T}}B^{\mathsf{T}}) = M(B \otimes A)M.$$

The exponent can be written as  $-\frac{1}{2}M(R^{-1}\otimes g^2)M$  with the *resolvent operator* R:

$$[R^{-1}(\kappa,\lambda)]^a{}_b = (1+\iota\sqrt{2\kappa}\varrho)\delta^a{}_b + \iota\sqrt{2\lambda}(\sigma g^1)^a{}_b.$$

As the resolvent and its inverse are operators, we write them with a covariant and a contravariant index. These indices are lowered with  $g^1$  and raised with  $(g^1)^{-1}$ .

Commuting the integral and derivative operators, the integral M is Gaussian and can be performed leading to the *intermediate field representation*:

$$Z(\kappa,\lambda) = \left[e^{\frac{1}{2}\left(\frac{\partial}{\partial\sigma}P\frac{\partial}{\partial\sigma}\right)}e^{\frac{1}{2}\frac{\partial}{\partial\varrho}\frac{\partial}{\partial\varrho}}e^{(-1)^{|1|+|2|}\frac{N_2}{2}\operatorname{Tr}\ln R(\kappa,\lambda)}\right]_{\varrho,\sigma=0}.$$
(3.3)

Now  $N_2$  is an explicit parameter in the integral, while  $N_1$  is hidden in the remaining traces. The sign  $(-1)^{|1|+|2|}$  tracks the bosonic/fermionic character of the original matrix. The sign  $(-1)^{|1|}$  tracks the symmetry of the intermediate matrix field  $\sigma$  (which agrees with that of  $g^1$ ). Both indices of  $\sigma$  have color 1 which reflects the fact that  $\sigma$ transforms in the (anti-)symmetric tensor representation of  $O_1(N_1)$ , that is,  $\sigma \rightarrow O_1 \sigma O_1^{\mathsf{T}}$  for  $O_1 \in O_1(N_1)$ . This is to be contrasted with the field M which transforms in the tensor product of the fundamental representations of  $O_1(N_1)$  and  $O_2(N_2)$ .

### 3.2. Perturbative expansion

The perturbative expansion of Z is obtained by Taylor expanding the interaction:

$$Z(\kappa,\lambda) = \left[ e^{\frac{1}{2} \left(\frac{\partial}{\partial\sigma} P \frac{\partial}{\partial\sigma}\right)} e^{\frac{1}{2} \frac{\partial}{\partial\varrho} \frac{\partial}{\partial\varrho}} \sum_{V=0}^{\infty} \frac{1}{V!} \left( \frac{(-1)^{|1|+|2|} N_2}{2} \operatorname{Tr} \ln R(\kappa,\lambda;\varrho,\sigma) \right)^V \right]_{\varrho,\sigma=0},$$

and commuting the Gaussian integration with the sum. Note that R denotes the resolvent operator, hence it naturally has a covariant and a contravariant index. Taking into account that

$$\ln R = -\sum_{p \ge 1} \frac{(-1)^{p+1}}{p} (\iota \sqrt{2\kappa} \varrho + \iota \sqrt{2\lambda} \sigma g^1)^p,$$
$$R = \sum_{p \ge 0} (-1)^p (\iota \sqrt{2\kappa} \varrho + \iota \sqrt{2\lambda} \sigma g^1)^p,$$

the derivatives of the resolvent and its logarithm are

$$\frac{\partial}{\partial \sigma^{ab}} \operatorname{Tr} \ln R = -\iota \sqrt{2\lambda} P_{ab,cd} R^{dc},$$

$$\frac{\partial}{\partial \sigma^{ab}} R^{cd} = -\iota \sqrt{2\lambda} P_{ab,ef} R^{ce} R^{fd},$$

$$\frac{\partial}{\partial \rho} \operatorname{Tr} \ln R = -\iota \sqrt{2\kappa} \operatorname{Tr}[R],$$

$$\frac{\partial}{\partial \rho} R^{ab} = -\iota \sqrt{2\kappa} R^{ac} g^{1}_{cd} R^{db} = -\iota \sqrt{2\kappa} (R^{2})^{ab}$$

where  $R^2$  denotes the square of the operator R.

Each term in the perturbative series can be represented as a ribbon graph with  $\rho$ -edges (see Section 2.2) as depicted in Figure 7:

- We represent each  $Tr \ln R$  as a disc with boundary oriented counter-clockwise.
- The derivatives with respect to σ create ribbon halfedges representing the free indices of *R*. The first derivative acting on a vertex creates a halfedge and an *R* associated to the *corner* (region between two consecutive halfedges) of the vertex. Subsequent derivatives split the existing corners creating new *R*'s. The indices *ab* of the resolvent *R<sup>ab</sup>* are associated to the ends of the corner: *a* for the source and *b* for the target in the sense of the arrow.
- The ribbon halfedges are connected into ribbon edges corresponding to the projectors P inside the  $\frac{\partial}{\partial\sigma^{ab}}P^{ab,dc}\frac{\partial}{\partial\sigma^{cd}}$  operators. The edges have an orientation represented by arrows on the strands bounding an edge: corresponding to  $P^{ab,dc}$ , we orient the strands from (ab) to (dc). Note that



**Figure 7.** *Left:* A ribbon graph with  $\rho$ -edges in the priori orientation: corners counter-clockwise and strands parallel. *Right:* Coherent orientation of arrows along every face. Five arrows had to be reoriented.

The projector generates two terms. The first one  $g_{a'd'}^1 g_{b'c'}^1$  corresponds to an edge with parallel strands. The second one  $(-1)^{|1|} g_{a'c'}^1 g_{b'd'}^1$  corresponds to a twisted edge.

A derivative with respect to ρ splits corner of a vertex also, but connects these two halves by g<sup>1</sup>. We represent this by a new type of halfedge, called ρ-halfedge. The ρ-halfedges are connected into ρ-edges corresponding to the ∂/∂ρ ∂/∂ρ operators. We represent these edges as dashed lines.

In the end, all intermediate fields are set to zero thus the resolvents are set to the identity R = 1. A corner that has been split by  $\rho$ -halfedges behaves like a single ordinary corner of a ribbon graph: for this reason corner will always refer to the region between two ribbon halfedges only.

Ignoring the twisting of the edges, a ribbon graph is a combinatorial map with  $\rho$ edges  $\mathcal{M}^{\rho}$ . We denote by  $h_v$  the ribbon-halfedges of the vertex v, each of which comes equipped with a pair of indices  $(b_{h_v}, a_{h_v})$ : b is the target of an arrow and a the source of another one. If  $h_v$  and  $h'_v$  are two neighboring ribbon-halfedges with  $h_v < h'_v$  in the cyclic order around v, the corner between them is denoted by  $(h_v, h'_v)$ . A ribbonedge connecting two vertices  $v, w \in \mathcal{M}^{\rho}$  is denoted by its halfedges  $e = \{h_v, h_w\}$ . Furthermore, we denote by  $V(\mathcal{M}^{\rho})$ ,  $E(\mathcal{M}^{\rho})$  and  $E^{\rho}(\mathcal{M}^{\rho})$  the numbers of vertices, ribbon-edges and  $\rho$ -edges of  $\mathcal{M}^{\rho}$  and by deg v and deg<sup> $\rho$ </sup> v the number of ribbonand  $\rho$ -halfedges at v. The perturbative series can be written as a sum over labeled combinatorial maps with  $\rho$ -edges:

$$Z(\lambda) = \sum_{\mathcal{M}^{\varrho}} \frac{1}{V(\mathcal{M}^{\varrho})! \, 2^{V(\mathcal{M}^{\varrho})}} \Big(\prod_{\substack{v \in \mathcal{M}^{\varrho} \\ v \in \mathcal{M}^{\varrho}}} \frac{1}{\deg v! \deg^{\varrho} v!} \Big)$$
$$\times ((-1)^{|1|+|2|} N_2)^{V(\mathcal{M}^{\varrho})} (-\lambda)^{E(\mathcal{M}^{\varrho})} (-2\kappa)^{E^{\varrho}(\mathcal{M}^{\varrho})}$$
$$\times \Big(\prod_{\substack{v \in \mathcal{M}^{\varrho} \\ v \in \mathcal{M}^{\varrho}}} \prod_{\substack{(h_v, h'_v) \\ comer of v}} g^1_{a_{h_v} b_{h'_v}} \Big) \Big(\prod_{\substack{e = \{h_v, h_w\} \\ ribbon-edge}} 2P^{b_{h_v} a_{h_v}, a_{h_w} b_{h_w}} \Big)$$

We expand the two terms in each edge projector to sum over ribbon graphs with (twisted) edges and  $\rho$ -edge. This is because the amplitude depends on the twisting: every face (closed strand) of the ribbon graph contributes a factor of  $N_1$  because along a face an even number of  $g^1$ 's concatenate into a trace. However, it might be necessary to transpose several  $g^1$ 's in order to get this trace: we represent these transpositions by reversing the corresponding arrows along the edge strands and the corners of the ribbon graph, see Figure 7. Overall we get (we explain the notation below)

$$Z(\lambda) = \sum_{\mathcal{M}^{\varrho}} \frac{1}{V(\mathcal{M}^{\varrho})! \, 2^{V(\mathcal{M}^{\varrho})}} \Big( \prod_{v \in \mathcal{M}^{\varrho}} \frac{1}{\deg v! \deg^{\varrho} v!} \Big) \\ \times ((-1)^{|1|+|2|} N_2)^{V(\mathcal{M}^{\varrho})} (-\lambda)^{E(\mathcal{M}^{\varrho})} (-2\kappa)^{E^{\varrho}(\mathcal{M}^{\varrho})} \\ \times \sum_{[\mathscr{G}^{\varrho}] \in \operatorname{Orb}_{\mathfrak{T}}(\mathcal{M}^{\varrho})} |\operatorname{Stab}_{\mathfrak{T}}(\mathscr{G}^{\varrho})| (N_1)^{F(\mathscr{G}^{\varrho})} ((-1)^{|1|})^{\#\operatorname{ranspositions} + \#\operatorname{twists}} \\ = \sum_{\mathcal{M}^{\varrho}} \frac{|\operatorname{Stab}_{\mathfrak{T}}(\mathcal{M}^{\varrho})|}{V(\mathcal{M}^{\varrho})! \, 2^{V(\mathcal{M}^{\varrho})}} \Big( \prod_{v \in \mathcal{M}^{\varrho}} \frac{1}{\deg v! \deg^{\varrho} v!} \Big) \sum_{[\mathscr{G}^{\varrho}] \in \operatorname{Orb}_{\mathfrak{T}}(\mathcal{M}^{\varrho})} \mathcal{A}(\mathscr{G}^{\varrho}) \quad (3.4)$$

with the amplitude  $\mathcal{A}(\mathscr{G}^{\varrho})$  of the ribbon graph. Some notation has been introduced in this equation. Because every ribbon-edge can be twisted or not, there are naively  $2^E$ ribbon graphs, associated to  $\mathcal{M}^{\varrho}$ . But ribbon graphs are in fact equivalence classes, emphasized by  $[\mathscr{G}^{\varrho}]$ . Two graphs are equivalent if one can be obtained from the other by successively reversing the order of halfedges of a subset of its vertices and—for each vertex separately—twisting all ribbon-edges connected to these vertices (edges with two twists are again untwisted). As proven in Appendix B, this degeneracy is counted by the cardinal of the stabilizer  $|\operatorname{Stab}_{\mathfrak{T}}(\mathscr{G}^{\varrho})|$  of the action of a finite group<sup>6</sup>  $\mathfrak{T}$ whose elements twist a subset of the ribbon-edges of  $\mathscr{G}^{\varrho}$  ( $\mathfrak{T}$  acts trivially on the  $\varrho$ edges). In the last step leading to (3.4), we used the fact that, as  $\mathfrak{T}$  is abelian,

$$\operatorname{Stab}_{\mathfrak{T}}(\mathscr{G}) = \operatorname{Stab}_{\mathfrak{T}}(\mathscr{M})$$

for any  $\mathscr{G} \in \operatorname{Orb}_{\mathfrak{T}}(\mathcal{M})$ , where  $\operatorname{Orb}_{\mathfrak{T}}(\mathcal{M})$  is the orbit of the combinatorial map  $\mathcal{M}$  under the action of  $\mathfrak{T}$ .

For example, the amplitude of the ribbon graph in Figure 7 is

$$\underbrace{((-1)^{|1|+|2|}N_2)^3}_{\text{vertices}}\underbrace{(-\lambda)^4}_{\text{edges}}\underbrace{(-2\kappa)^2}_{\varrho\text{-edges}}\underbrace{((-1)^{|1|})}_{\text{twists}}\underbrace{(N_1)}_{\text{faces}}\underbrace{((-1)^{|1|})^5}_{\text{reorientations}}$$
$$= (-\lambda)^4 (-2\kappa)^2 ((-1)^{|1|}N_1) ((-1)^{|2|}N_2)^3.$$

<sup>6</sup>The group  $\mathfrak{T}$  is a subgroup of the so called ribbon group, introduced in [18], which also includes the operation of taking the partial dual of a ribbon graph with respect to a subset of its edges.

**Amplitudes.** The amplitude  $\mathcal{A}(\mathcal{G}^{\varrho})$  can be further computed.

In Proposition A.1, we prove that any ribbon graph can be deformed into a connected sum of

- a graph without twisted edges embeddable into a closed orientable surface Σ<sub>g</sub> of genus g;
- either no, one or two graphs with a single twisted edge, embeddable into the projective plane ℝP<sup>2</sup>.

This is the ribbon graph equivalent of the classification theorem of closed twodimensional surfaces. The crucial observation is that one can track the power of -1 in the amplitude under these deformations. In Theorem A.11, we show that for a ribbon graph with twists and which requires transpositions in order to coherently orient the faces:

$$(-1)^{V(\mathscr{G})} \Big(\prod_{e \in \mathscr{E}(\mathscr{G})} (-1)^{\tau(e)} \Big) \Big(\prod_{f \in \mathscr{F}(\mathscr{G})} (-1)^{t(f)} \Big) = (-1)^{F(\mathscr{G})},$$

where  $\tau(e) = 0$  if the edge *e* is untwisted (straight) and  $\tau(e) = 1$  if the edge is twisted; t(f) is the number of reorientations of arrows required to coherently orient the face *f*.

The graph  $\mathcal{G}^{\varrho}$  can be seen as the union of two ribbon graphs:

- One ribbon graph has color 2 and is trivial. It consisting in all the vertices of *β<sup>ρ</sup>*, each bounded by one face and has no edges. The graph has no twists (as it has no edges) and all its faces are coherently oriented.
- The second one is the graph of color 1. It has twisted edges, and some transpositions are needed in order to coherently orient its faces.

Then the amplitude of a graph in (3.4) can be written as

$$\mathcal{A}(\mathscr{G}^{\varrho}) = (-2\kappa)^{E^{\varrho}(\mathscr{G}^{\varrho})} (-\lambda)^{E(\mathscr{G}^{\varrho})} ((-1)^{|1|} N_1)^{F(\mathscr{G}^{\varrho})} ((-1)^{|2|} N_2)^{V(\mathscr{G}^{\varrho})},$$

proving equation (1.1) in Theorem 1.2 for D = 2.

**Combinatorial weights.** The combinatorial weights in equation (3.4) simplify by gathering the labeled graphs corresponding to the same unlabeled ribbon graph with  $\rho$ -edges:

$$Z(\kappa,\lambda) = \sum_{[\mathscr{G}^{\varrho}]} \mathcal{W}(\mathscr{G}^{\varrho}) \,\mathcal{A}(\mathscr{G}^{\varrho}),$$

where the (positive) weights  $\mathcal{W}(\mathcal{G}^{\varrho})$  include all the combinatorial factors coming from partially resuming (3.4) to a sum over equivalence classes.

**Remark 3.1** (Dual graphs). The amplitudes of a ribbon graph and its dual are related by

$$\begin{aligned} \mathcal{A}(\mathcal{G}^*) &= (-\lambda)^{E(\mathcal{G}^*)} ((-1)^{|1|} N_1)^{F(\mathcal{G}^*)} ((-1)^{|2|} N_2)^{V(\mathcal{G}^*)} \\ &= (-\lambda)^{E(\mathcal{G})} ((-1)^{|2|} N_2)^{F(\mathcal{G})} ((-1)^{|1|} N_1)^{V(\mathcal{G})}. \end{aligned}$$

We will see below that  $\mathcal{W}(\mathcal{G}^{\varrho}) = \mathcal{W}(\mathcal{G}^{\varrho*})$ . In particular, for the mixed  $O(N) \otimes$ Sp(N)-models, we get  $\mathcal{A}(\mathcal{G}^{\varrho*}) = (-1)^{V(\mathcal{G}^{\varrho}) + F(\mathcal{G}^{\varrho})} \mathcal{A}(\mathcal{G}^{\varrho})$ , hence the contributions of a graph and its dual cancel if  $V(\mathcal{G}^{\varrho}) + F(\mathcal{G}^{\varrho})$  is odd.

A heuristic argument why  $\mathcal{W}(\mathcal{G}^{\varrho}) = \mathcal{W}(\mathcal{G}^{\varrho*})$  goes as follows. We split the quartic interactions using an intermediate field  $\sigma_1$  with indices of color 1 coupling to M via  $\propto M((g^1\sigma_1g^1) \otimes g^2)M$ . But one can choose the intermediate field to have indices of color 2 and coupling  $\propto M(g^1 \otimes (g^2\sigma_2g^2))M$ . The vertices now contribute factors of  $N_1$  and the faces  $N_2$ . For any graph, contracting the intermediate field  $\sigma_1$  and introducing  $\sigma_2$  in the orthogonal channel passes to the dual graph.

As the combinatorics is insensitive to the symmetry, we focus on the  $O(N_1) \otimes O(N_2)$ -model. The connected two-point function of this model

$$G_2(\kappa,\lambda) = Z^{-1}(\kappa,\lambda) \int d\mu[M] \operatorname{Tr}[M\delta M^{\mathsf{T}}\delta],$$

obeys a Dyson-Schwinger equation (DSE). Using

$$0 = Z^{-1}(\kappa, \lambda) \int [dM] \\ \times \frac{\partial}{\partial M^{a_1 a_2}} \left( M^{a_1 a_2} e^{-\frac{1}{2} \operatorname{Tr} \left[ M \delta M^{\mathsf{T}} \delta \right] - \frac{\kappa}{4} (\operatorname{Tr} [M \delta M^{\mathsf{T}} \delta])^2 - \frac{\lambda}{4} \operatorname{Tr} [(M \delta M^{\mathsf{T}} \delta)^2]} \right),$$

we conclude that

$$G_2(\kappa,\lambda) = N_1 N_2 + (4\kappa \partial_\kappa + 4\lambda \partial_\lambda) \ln Z(\kappa,\lambda).$$
(3.5)

The free energy  $\ln Z$  expands in connected graphs. The derivative operator  $2\kappa \partial_{\kappa} + 2\lambda \partial_{\lambda}$  generates a rooting of the graph, that we get a sum over graphs with a marked  $\rho$ -or ribbon-halfedge. Rooted graphs are simpler to count. In Proposition B.1, we show that the perturbative series of  $G_2$  can be written as

$$G_2(\kappa,\lambda) = \sum_{[\mathscr{G}^{\varrho}] \text{ connected, rooted}} \frac{1}{2^{C(\mathscr{G}^{\varrho} - \mathscr{E}^{\varrho}) - 1}} \mathcal{A}(\mathscr{G}^{\varrho}), \qquad (3.6)$$

where  $\mathscr{G}^{\varrho} - \mathscr{E}^{\varrho}$  is the graph obtained from  $\mathscr{G}^{\varrho}$  by deleting all the  $\varrho$ -edges and  $C(\mathscr{G}^{\varrho} - \mathscr{E}^{\varrho})$  denotes the number of its connected components. Note that even if  $\mathscr{G}^{\varrho}$  is connected as a ribbon graph with  $\varrho$ -edges, the graph  $\mathscr{G}^{\varrho} - \mathscr{E}^{\varrho}$  may be disconnected.

It is well known that rooting trivializes the symmetry factors in ordinary combinatorial maps. What is non-trivial is that it also simplifies the factor  $2^{-V} |\text{Stab}_{\mathfrak{T}}(\mathcal{M}^{\varrho})|$ in (3.4) to  $2^{-(C(\mathscr{G}^{\varrho} - \mathscr{E}^{\varrho}) - 1)}$ . The combinatorial weight in (3.6) is manifestly invariant under duality. Rooted ribbon graphs can be embedded into two-dimensional surfaces with one boundary component corresponding to the rooted face.

The Dyson–Schwinger equation for the connected two-point function can be integrated in the sense of formal power series to yield the perturbative expansion of the free energy:

$$\ln Z(\kappa, \lambda) = \sum_{\substack{[\mathscr{G}^{\varrho}] \text{ connected, rooted,} \\ E \text{ or } E^{\varrho} > 0}} \frac{1}{2^{C(\mathscr{G}^{\varrho} - \mathscr{E}^{\varrho}) + 1}(E + E^{\varrho})} \mathcal{A}(\mathscr{G}^{\varrho}).$$

where E and  $E^{\varrho}$  denote the number of ribbon edges and  $\varrho$ -edges in  $\mathscr{G}^{\varrho}$ , respectively. The integration does not spoil the symmetry under duality because the powers of the coupling constants in the amplitude only depend on the number of edges. Finally, the partition function Z can then be obtained by exponentiating  $\ln Z$ .

### 4. Tensor models

The case  $D \ge 3$  is treated similarly to the case D = 2. However, as the number of available quartic invariants grows exponentially with D (recall Lemma 2.2), the number of intermediate fields grows also. Moreover, the intermediate fields are matrices with different dimensions. At most one of the  $N_c$  factors can be rendered explicit as a parameter in the integral, and one must rely on graphical methods to track the other  $N_c$ 's.

In  $D \ge 3$  the perturbative expansion is an expansion in *colored multi-ribbon* graphs which can be understood intuitively as stacked ribbon graphs. The  $N_c$  to  $-N_c$  duality holds graph by graph because only the combination  $(-1)^{|c|}N_c$  appears in the amplitude of a graph.

If one aims to study tensor (or matrix) models with a sensible large N limit, one needs to rescale the coupling constants with powers of N. Care has to be taken if one wants to preserve the manifest N to -N duality: this can sometimes require a flip of the sign of some of the coupling constants.

#### 4.1. Intermediate field representation

Complex random tensor models in the intermediate field representation were, for example, studied in [15, 26]. We introduce an intermediate field per quartic interaction. For a subset of the colors  $\mathcal{C}$ , we denote by  $\Sigma^{\mathcal{C}}$  the set of  $N^{|\mathcal{C}|} \times N^{|\mathcal{C}|}$  matrices

(where  $|\mathcal{C}|$  denotes the cardinality of  $\mathcal{C}$ ) taken to be symmetric if the sum of the parities of the indices in  $\mathcal{C}$  is even and anti-symmetric if it is odd:

$$\Sigma^{\mathcal{C}} = \begin{cases} \operatorname{Sym}^{2}(\bigotimes_{c \in \mathcal{C}} \mathcal{H}_{c}), & \sum_{c \in \mathcal{C}} |c| = 0 \mod 2, \\ \Lambda^{2}(\bigotimes_{c \in \mathcal{C}} \mathcal{H}_{c}), & \sum_{c \in \mathcal{C}} |c| = 1 \mod 2, \end{cases}$$
$$\sigma^{a_{\mathcal{C}}^{1} a_{\mathcal{C}}^{2}} = (-1)^{\sum_{c \in \mathcal{C}} |c|} \sigma^{a_{\mathcal{C}}^{2} a_{\mathcal{C}}^{1}} \quad \forall \sigma \in \Sigma^{\mathcal{C}}, \end{cases}$$

where we recall that  $a_{\mathcal{C}}$  denotes a multi-index  $(a_c \mid c \in \mathcal{C})$ . Note that  $\sigma$  is always commuting (bosonic) because  $\bigotimes_{c \in \mathcal{C}} \mathcal{H}_c$  is either purely odd or even. For  $\mathcal{C} = \emptyset$ , set  $\sigma \in \Lambda^0_{\infty}$  the commuting scalars. Since  $\sigma$  are (anti-)symmetric under exchange of their two multi-indices, it is useful to introduce the (anti-)symmetric projector

$$P_{\mathcal{C}}: \Sigma^{\mathcal{C}} \to \Sigma^{\mathcal{C}},$$

$$(P_{\mathcal{C}})^{a_{\mathcal{C}}^{1}a_{\mathcal{C}}^{2}}_{,b_{\mathcal{C}}^{1}b_{\mathcal{C}}^{2}} := \frac{1}{2} \Big( \prod_{c \in \mathcal{C}} \delta^{a_{c}^{1}}_{b_{c}^{1}} \delta^{a_{c}^{2}}_{b_{c}^{2}} + (-1)^{\sum_{c \in \mathcal{C}} |c|} \prod_{c \in \mathcal{C}} \delta^{a_{c}^{1}}_{b_{c}^{2}} \delta^{a_{c}^{2}}_{b_{c}^{1}} \Big),$$

and  $P_{\mathcal{C}}$  is the identity for  $\mathcal{C} = \emptyset$ . The projector is such that

**Lemma 4.1** (Hubbard–Stratonovich transformation). *Every quartic tensor invariant* I(T) *in equation* (2.1)

$$\begin{split} I(T) &= \sum_{\substack{a_{\mathcal{D}}^{1}, a_{\mathcal{D}}^{2}, b_{\mathcal{D}}^{1}, b_{\mathcal{D}}^{2}}} \left( T^{a_{\mathcal{D}}^{1}} T^{a_{\mathcal{D}}^{2}} \prod_{c \in \mathcal{D} \setminus \mathcal{C}} g^{c}_{a_{c}^{1} a_{c}^{c}} \right) \left( T^{b_{\mathcal{D}}^{1}} T^{b_{\mathcal{D}}^{2}} \prod_{c \in \mathcal{D} \setminus \mathcal{C}} g^{c}_{b_{c}^{1} b_{c}^{c}} \right) \\ &\times \left( \prod_{c \in \mathcal{C}} (-\operatorname{sgn}(\pi^{c}))^{|c|} g^{c}_{a_{c}^{1} b_{c}^{\pi^{c}(1)}} g^{c}_{a_{c}^{2} b_{c}^{\pi^{c}(2)}} \right) \\ &= \sum_{a_{\mathcal{C}}^{1}, a_{\mathcal{C}}^{2}, b_{\mathcal{C}}^{1}, b_{\mathcal{C}}^{2}} (g^{\otimes \mathcal{D} \setminus \mathcal{C}} (T, T))^{a_{\mathcal{C}}^{1} a_{\mathcal{C}}^{2}} K_{a_{\mathcal{C}}^{1} a_{\mathcal{C}}^{2}, b_{\mathcal{C}}^{1} b_{\mathcal{C}}^{2}} (g^{\otimes \mathcal{D} \setminus \mathcal{C}} (T, T))^{b_{\mathcal{C}}^{1} b_{\mathcal{C}}^{2}} \end{split}$$

with  $\pi^{c}$  fixed permutations of two elements, can (formally) be expressed as a Gaussian integral

$$e^{-\frac{\lambda}{4}I(T)} = \left[e^{\frac{1}{2}\left(\frac{\partial}{\partial\sigma}, P_{\mathcal{C}}K\frac{\partial}{\partial\sigma}\right)}e^{-i\sqrt{\frac{\lambda}{2}}(\sigma, g^{\otimes \mathcal{D} \setminus \mathcal{C}}(T, T))}\right]_{\sigma=0}$$

with

$$(P_{\mathcal{C}}K)^{a_{\mathcal{C}}^1a_{\mathcal{C}}^2, b_{\mathcal{C}}^1b_{\mathcal{C}}^2} = (P_{\mathcal{C}})^{a_{\mathcal{C}}^1a_{\mathcal{C}}^2}_{,c_{\mathcal{C}}^1c_{\mathcal{C}}^2}K^{c_{\mathcal{C}}^1c_{\mathcal{C}}^2, b_{\mathcal{C}}^1b_{\mathcal{C}}^2}_{,c_{\mathcal{C}}^1c_{\mathcal{C}}^2}$$

and

$$(A, B) = g^{a_1 a_2} g^{b_1 b_2} A_{a_1 b_1} B_{a_2 b_2},$$

the standard pairing between a vector space and its dual.

*Proof.* The indices of color c of the kernel K are connected as

$$K_{a_{\mathcal{C}}^{1}a_{\mathcal{C}}^{2},b_{\mathcal{C}}^{1}b_{\mathcal{C}}^{2}} \sim \begin{cases} (-1)^{|c|} g_{a_{c}^{1}b_{c}^{1}}^{c} g_{a_{c}^{2}b_{c}^{2}}^{c}, & \pi^{c} = (1)(2), \\ g_{a_{c}^{1}b_{c}^{2}}^{c} g_{a_{c}^{2}b_{c}^{1}}^{c}, & \pi^{c} = (12), \end{cases}$$

hence, as an operator,  $K^2 = 1$  and  $P_{\mathcal{C}}K = KP_{\mathcal{C}}$ . Taking into account that

$$\begin{split} g^{c}_{b^{1}_{c}b^{2}_{c}} &= (-1)^{|c|} g^{c}_{b^{2}_{c}b^{1}_{c}}, \quad K^{a^{1}_{\mathcal{C}}a^{2}_{\mathcal{C}}}_{,b^{1}_{\mathcal{C}}b^{2}_{\mathcal{C}}} = K^{a^{2}_{\mathcal{C}}a^{1}_{\mathcal{C}}}_{,b^{2}_{\mathcal{C}}b^{2}_{\mathcal{C}}}, \\ T^{b^{1}_{\mathcal{D}}} T^{b^{2}_{\mathcal{D}}} &= (-1)^{\sum_{c \in \mathcal{D}} |c|} T^{b^{2}_{\mathcal{D}}} T^{b^{1}_{\mathcal{D}}}, \end{split}$$

we have

$$\begin{split} K^{a_{\mathcal{C}}^{l}a_{\mathcal{C}}^{2}} & \underset{b_{\mathcal{C}}^{l}b_{\mathcal{C}}^{2}}{\overset{1}{\mathcal{E}}} \left( T^{b_{\mathcal{D}}^{1}}T^{b_{\mathcal{D}}^{2}} \prod_{c \in \mathcal{D} \setminus \mathcal{C}} g^{c}_{b_{c}^{1}b_{c}^{2}} \right) \\ &= (-1)^{\sum_{c \in \mathcal{C}}|c|} K^{a_{\mathcal{C}}^{2}a_{\mathcal{C}}^{1}}_{,b_{\mathcal{C}}^{2}b_{\mathcal{C}}^{1}} \left( T^{b_{\mathcal{D}}^{2}}T^{b_{\mathcal{D}}^{1}} \prod_{c \in \mathcal{D} \setminus \mathcal{C}} g^{c}_{b_{c}^{2}b_{c}^{1}} \right) \end{split}$$

that is,  $P_{\mathcal{C}}Kg^{\otimes \mathcal{D} \setminus \mathcal{C}}(T,T) = Kg^{\otimes \mathcal{D} \setminus \mathcal{C}}(T,T)$ , hence  $Kg^{\otimes \mathcal{D} \setminus \mathcal{C}}(T,T)$  is a matrix with the same symmetry type as  $\sigma$ . It follows that

$$(g^{\otimes \mathcal{D} \setminus \mathcal{C}}(T,T), (P_{\mathcal{C}}KP_{\mathcal{C}})g^{\otimes \mathcal{D} \setminus \mathcal{C}}(T,T)) = (g^{\otimes \mathcal{D} \setminus \mathcal{C}}(T,T), Kg^{\otimes \mathcal{D} \setminus \mathcal{C}}(T,T)),$$

hence expanding the exponentials and commuting the sum and the derivative operator we get

$$\begin{split} \left[\sum_{n=0}^{\infty} \frac{(-\frac{\lambda}{4})^n}{n!(2n)!} \left(\frac{\partial}{\partial\sigma}, P_{\mathcal{C}} K \frac{\partial}{\partial\sigma}\right)^n (\sigma, g^{\otimes \mathcal{D} \setminus \mathcal{C}} (T, T))^{2n} \right]_{\sigma=0} &= \left[\sum_{n=0}^{\infty} \frac{(-\frac{\lambda}{4})^n}{n!} \times \left( (g^{\otimes \mathcal{D} \setminus \mathcal{C}} (T, T) P_{\mathcal{C}})_{a_{\mathcal{C}}^1 a_{\mathcal{C}}^2} (P_{\mathcal{C}} K)^{a_{\mathcal{C}}^1 a_{\mathcal{C}}^2, b_{\mathcal{C}}^1 b_{\mathcal{C}}^2} (P_{\mathcal{C}} g^{\otimes \mathcal{D} \setminus \mathcal{C}} (T, T))_{b_{\mathcal{C}}^1 b_{\mathcal{C}}^2} \right)^n \right]_{\sigma=0} \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \left( -\frac{\lambda}{4} I(T) \right)^n. \quad \blacksquare$$

When dealing with several quartic invariants, we will label them q and the corresponding subset of colors  $C_q$ . In order to simplify the notation, we sometimes drop this subscript. Using the intermediate fields, the partition function of the graded quadratic tensor model of Definition 2.3 becomes

$$Z(\lambda) = \int \mu[T] = \int [dT] e^{-\frac{1}{2}g^{\otimes \mathcal{D}}(T,T)} \times \left[ e^{\sum_{q \in \Omega} \frac{1}{2}(\frac{\partial}{\partial \sigma_q}, P_{\mathcal{C}_q} K_q \frac{\partial}{\partial \sigma_q})} \cdot e^{-\sum_q i \sqrt{\frac{\lambda q}{2}}(\sigma_q, g^{\otimes \mathcal{D} \setminus \mathcal{C}_q}(T,T))} \right]_{\sigma_q = 0},$$

where we denoted the coupling constants generically by  $\lambda$ . We denote by  $1^{\otimes \mathcal{C}}$  the identity operator acting on  $\bigotimes_{c \in \mathcal{C}} \mathcal{H}_c$ . We define the operator acting on  $\bigotimes_{c=1}^{D} \mathcal{H}_c$ 

$$A(\lambda) = \sum_{q \in \Omega} i \sqrt{2\lambda_q} 1^{\otimes \mathcal{D} \setminus \mathcal{C}_q} \otimes \sigma_q,$$
  
$$A^{a^1_{\mathcal{D}}}_{a^2_{\mathcal{D}}} = \sum_{q \in \Omega} i \sqrt{2\lambda_q} \left( \prod_{c \in \mathcal{D} \setminus \mathcal{C}_q} \delta^{a^1_c}_{a^2_c} \right) (\sigma_q)^{a^2_{\mathcal{C}_q}}_{a^1_{\mathcal{C}_q}}$$

and perform the Gaussian integral over T to obtain the partition function in the *inter*mediate field representation:

$$Z(\lambda) = \left[ e^{\sum_{q \in \mathcal{Q}} \frac{1}{2} \left( \frac{\partial}{\partial \sigma_q}, P_{\mathcal{C}_q} K_q \frac{\partial}{\partial \sigma_q} \right)} e^{-\frac{(-1)\sum_{c \in \mathcal{D}} |c|}{2} \operatorname{Tr} \ln(1^{\otimes \mathcal{D}} + A(\lambda))} \right]_{\sigma_q = 0}.$$
 (4.1)

This is the generalization of equation (3.3) to D > 3. The *resolvent operator* for tensors is  $R = (1^{\otimes \mathcal{D}} + A(\lambda))^{-1}$ . The field  $\varrho$  we encountered in D = 2 corresponds to the unique disconnected quartic invariant  $\mathcal{C}_q = \emptyset$ . For now, we keep all factors  $N_c$  in the trace: the trace over the color 1 space can be performed explicitly because  $1 \notin \mathcal{C}_q$  for all  $q \in Q$ . In strict generalization of the matrix cases, the sign  $(-1)^{\sum_{c \in \mathcal{D}} |c|}$  accounts for fermionic/bosonic nature of the tensor field T. Each intermediate field  $\sigma_q$  has its own symmetry captured by the sign  $(-1)^{\sum_{c \in \mathcal{C}_q} |c|}$ . The effect of the Hubbard–Stratonovich transformation on the Feynman diagrams is depicted schematically in Figure 8.



Figure 8. The Hubbard–Stratonovich transformation.

#### 4.2. Perturbative expansion

Because of the tensor products, the Feynman graphs of the perturbative expansion of (4.1) are *D*-colored multi-ribbon graphs. Intuitively, they can be understood as *D* stacked ribbon graphs. Ribbon graphs are obtained from combinatorial maps by replacing their edges by ribbon edges which can then be twisted or not. Similarly, *D*-colored multi-ribbon graphs are obtained from *edge multicolored combinatorial maps*. These, in turn, are combinatorial maps with edges labeled by subsets of colors  $C \subset D$ .



**Figure 9.** *Left:* Edge multicolored combinatorial map with D = 3. Each edge carries a subset of colors. *Center:* A multi-ribbon graph obtained from this multicolored combinatorial map. *Right:* Multi-ribbon edge corresponding to the quartic invariant of Figure 1 in its untwisted (*top*) and twisted (*down*) state.

**Definition 4.2** (Edge multicolored combinatorial map [28]). An *edge multicolored combinatorial map*  $\mathcal{M}$ , depicted in Figure 9 on the left, is composed of

- a finite set S that is the disjoint union of sets S<sup>C</sup> of halfedges of the colors C ∈ D, all of even cardinality S = ∐<sub>C⊂D</sub> S<sup>C</sup>;
- a permutation  $\pi$  on S;
- for every  $\mathcal{C} \in \mathcal{D}$  an involution  $\alpha^{\mathcal{C}}$  on  $\mathcal{S}^{\mathcal{C}}$  with no fixed points. The involution  $\alpha^{\mathcal{C}}$  can be extended to the whole of  $\mathcal{S}$  by setting  $\alpha^{\mathcal{C}}(h) = h \forall h \in \mathcal{S} \setminus \mathcal{S}^{\mathcal{C}}$ .

The set of cycles of  $\pi$  is the set of vertices of the map  $\mathcal{V}(\mathcal{M})$ . The set of cycles of  $\alpha^{\mathcal{C}}$  is the set of edges of colors  $\mathcal{C}$ ,  $\mathcal{E}_{\mathcal{C}}(\mathcal{M})$ , and  $\mathcal{E}(\mathcal{M}) = \bigcup_{\mathcal{C} \subset \mathcal{D}} \mathcal{E}_{\mathcal{C}}(\mathcal{M})$  is the set of all the edges of the map. The cardinalities of these sets are denoted by  $V(\mathcal{M})$ ,  $E_{\mathcal{C}}(\mathcal{M})$ ,  $E(\mathcal{M})$ , respectively.

An edge multicolored combinatorial map is connected if and only if the group freely generated by  $\pi$  and the  $\alpha^{\mathcal{C}}$  acts transitively on  $\mathcal{S}$ .

The following definition of multi-ribbon graphs is a generalization of signed rotation systems [19] which are equivalent to ribbon graphs.

**Definition 4.3** (*D*-colored multi-ribbon graph). A *D*-colored multi-ribbon graph  $\mathbb{G}$ , depicted in Figure 9 in the center, is an edge multicolored combinatorial map  $\mathcal{M}$  equipped with binary variables  $|\mathcal{C}|$  taking values 0 or 1 on each edge with colors  $\mathcal{C}$  (for each edge, we have either a 0 or a 1 for each of its colors):

for 
$$e \in \mathscr{E}_{\mathcal{C}}(\mathcal{M})$$
,  $e \mapsto \tau(e) = \{\tau^{c}(e) \in \{0, 1\} \mid c \in \mathcal{C}\}$ 

These edges are called (twisted) multi-ribbon edges. Twisting a multi-ribbon edge *e* amounts to flipping all the variables  $\tau(e)$ , that is,  $\tau^c(e) \rightarrow \tau^c(e) + 1 \mod 2$ .

Two *D*-colored multi-ribbon graphs are equivalent if they differ by reversing the order of halfedges around a vertex and simultaneously twisting every incident multi-ribbon edge (self-loops are twisted twice) at a finite number of vertices.

The following graphical representation is depicted in Figure 9. The vertices of a multi-ribbon graph are represented by D concentric discs with colors ordered from the innermost to the outermost circle. A multi-ribbon edge  $e \in \mathcal{E}_{\mathcal{C}}(\mathcal{M})$  connects the discs with colors in  $\mathcal{C}$  of its end vertices by ribbon edges. Only discs of the same color can be connected and the ribbons carry the color of the discs they are connecting. A 0/1 value of  $\tau^c(e)$  indicates that the ribbon with the color c of the edge e is un-/twisted. The whole multi-ribbon edge is called untwisted if the ribbon of biggest color in  $\mathcal{C}$  is untwisted. The  $\rho$ -edges encountered in Section 3 are the edges with colors  $\mathcal{C} = \emptyset$ . They can be represented as dashed.

The *faces* of color *c* of  $\mathbb{G}$  are the closed circuits obtained by going along the sides of the ribbon edges and along the discs of the vertices of color *c*. The set of faces of color *c* of  $\mathbb{G}$  is denoted  $\mathcal{F}_c(\mathbb{G})$  and its cardinality is denoted  $F_c(\mathbb{G})$ . The restriction of  $\mathbb{G}$  to a single color  $\mathbb{G}_c$  is obtained by deleting all the discs and ribbon with other colors. The graph  $\mathbb{G}_c$  is an ordinary ribbon graph, possibly disconnected. Observe that  $F_c(\mathbb{G})$  is also the number of faces of the ribbon graph  $\mathbb{G}_c$ .

The perturbative expansion of (4.1) is obtained by Taylor expanding and commuting the sum and the Gaussian integral:

$$Z(\lambda) = \sum_{V=1}^{\infty} \frac{(-1)^{V \sum_{c=1}^{D} |c|}}{V! 2^{V}} \Big[ e^{\sum_{q \in \mathcal{Q}} \frac{1}{2} (\frac{\partial}{\partial \sigma_{q}}, P_{\mathcal{C}_{q}} K_{q} \frac{\partial}{\partial \sigma_{q}})} (-\operatorname{Tr} \ln(1^{\otimes \mathcal{D}} + A))^{V} \Big]_{\sigma_{q} = 0},$$

where we suppressed the argument of A. Each Tr  $\ln(1^{\otimes D} + A)^{-1}$  represents a multiribbon vertex. The derivatives

$$\frac{\partial}{\partial \sigma_q} (1^{\otimes \mathcal{D}} + A)^{-1} = (1^{\otimes \mathcal{D}} + A)^{-1} \left( -\frac{\partial A}{\partial \sigma_q} \right) (1^{\otimes \mathcal{D}} + A)^{-1}$$
$$\frac{\partial}{\partial \sigma_q} \operatorname{Tr} \ln(1^{\otimes \mathcal{D}} + A)^{-1} = \operatorname{Tr} \left[ (1^{\otimes \mathcal{D}} + A)^{-1} \left( -\frac{\partial A}{\partial \sigma_q} \right) \right],$$
$$\frac{\partial A}{\partial \sigma_q} = \iota \sqrt{2\lambda_q} 1^{\otimes \mathcal{D} \setminus \mathcal{C}_q} \otimes P_{\mathcal{C}_q}$$

create multi-ribbon halfedges which, because of the projector, are joined in a twisted or untwisted way. The possible types of multi-ribbon edges depend on the quartic invariants  $q \in \Omega$ : for brevity the multi-ribbon edges associated to the quartic invariant q are called q-edges. The trace induces a cyclic ordering around the vertex which by convention we take to be counter-clockwise. Following an index of color c, it goes around the vertex until it encounters a multi-ribbon halfedge with  $c \in C_q$ . As in the matrix case, the order of indices is important if |c| = 1. This is accounted for by



**Figure 10.** *Left:* A 3-colored multi-ribbon graph. The arrows indicate the order of indices of the  $g_{a_cb_c}^c$ . In the a priori orientation, arrows point counter-clockwise around vertices and parallel along edges. *Right:* Ribbon graphs obtained by restricting to a single color. The black arrows had to be flipped to arrive at a coherent orientation along each face. Compare to Figure 7.

orienting the strands of a vertex in a counter-clockwise manner (Figure 10). Denoting  $R = (1^{\otimes \mathcal{D}} + A)^{-1}$ , the contribution of an edge can be written as

$$(-\iota\sqrt{2\lambda_q})R_{.b^1}R_{a^1.}(P_{\mathcal{C}_q}K_q)^{b^1a^1,b^2a^2}R_{.b^2}R_{a^2.}$$

and upon setting  $\sigma_q = 0$  all the resolvents reduce to the identity operator.

We denote by  $\mathcal{M}$  the edge multicolored maps and by  $\deg^q v$  the number of edges of type q incident to the vertex v. As in the matrix case, the edges with  $\mathcal{C} = \emptyset$  are special. We call them  $\varrho$ -edges and denote sometimes the number of such edges  $E^{\varrho}(\mathcal{M})$ . However, note that the  $\varrho$ -edges are also counted as a particular case q-edges for  $q \in Q$ .

The halfedges incident at a vertex have colors  $\mathcal{C}_q$ , and we denote them  $h_v^{\mathcal{C}_q}$ ,  $f_v^{\mathcal{C}_{q'}}$ and so on. Each halfedge is composed of  $|\mathcal{C}_q|$  ribbon halfedges, one for each color in  $\mathcal{C}_q$ . The corners<sup>7</sup> of the map  $\mathcal{M}$  are the pieces of vertices comprised between two consecutive halfedges, and we denote them by  $(h_v^{\mathcal{C}_q}, f_v^{\mathcal{C}_{q'}})$ , with the successor  $f_v^{\mathcal{C}_{q'}}$ of  $h_v^{\mathcal{C}_q}$  when turning around v. The partition function becomes

$$Z(\lambda) = \sum_{\mathcal{M}} \frac{1}{V(\mathcal{M})! 2^{V(\mathcal{M})}} \Big( \prod_{v \in \mathcal{M}} \frac{1}{\prod_{q} \deg^{q} v!} \Big) ((-1)^{\sum_{c \in \mathcal{D}} |c|})^{V(\mathcal{M})} 2^{E^{\rho}(\mathcal{M})}$$

$$\times \prod_{q \in \mathcal{Q}} (-\lambda_{q})^{E_{q}(\mathcal{M})} \prod_{v \in \mathcal{V}(\mathcal{M})} \Big( \prod_{(h_{v}^{\mathcal{C}_{q}}, f_{v}^{\mathcal{C}_{q'}}) \text{ corner of } v} \prod_{c \in \mathcal{D}} g_{(a_{c})}^{c} \Big|_{h_{v}^{\mathcal{C}_{q}}(b_{c})} \Big|_{f_{v}^{\mathcal{C}_{q'}}} \Big)$$

$$\times \prod_{e = \{h_{v}^{\mathcal{C}_{q}}, h_{w}^{\mathcal{C}_{q}}\} \in \mathcal{E}(\mathcal{M})} \Big( (2P_{\mathcal{C}_{q}}K_{q})^{(b_{\mathcal{C}_{q}})} \Big|_{h_{v}^{\mathcal{C}_{q}}(a_{\mathcal{C}_{q}})} \Big|_{h_{v}^{\mathcal{C}_{q}}, (b_{\mathcal{C}_{q}})} \Big|_{h_{w}^{\mathcal{C}_{q}}(a_{\mathcal{C}_{q}})} \Big|_{h_{w}^{\mathcal{C}_{q}}} \Big|_{h_{w}^{\mathcal{C}_{$$

<sup>&</sup>lt;sup>7</sup>We exclude the  $\rho$  halfedges when identifying the corners.

with the convention such that if  $C_q = \emptyset$ , then there is no corner and the edge  $(2P_{C_q}K_q)$  equals 1.

An index of color *c* is insensitive to the halfedges with colors different from *c*: an index follows a face and closes in a trace when the face closes. As in the matrix case, we obtain either straight edges or twisted ones coming from the two terms in  $P_{\mathcal{C}_q}$ . In turn, the edges contract on  $K_p$  kernels that send the color *c* either in a parallel channel or in a cross one. Overall, the ribbon of color *c* of the edge  $(2P_{\mathcal{C}_q}K_q)$  can either be straight, which we denote  $\tau^c(e) = 0$ , or twisted, which we denote  $\tau^c(e) = 1$ . Let us track the indices of color *c* coming from a term in  $P_{\mathcal{C}_q}$  and one possible  $K_q$ , for instance,

$$(2P_{\mathcal{C}_q}K_q) \sim \delta^{b_c^1}_{\ i} \delta^{a_c^1}_{\ j} (-1)^{|c|} g^{i b_c^2} g^{j a_c^2} = (-1)^{|c|} g^{b_c^1 b_c^2} g^{a_c^1 a_c^2}.$$

As this term contracts the indices b together and the indices a together, it corresponds to a ribbon of color c which is twisted. Proceeding similarly for all the edges and recalling that some g's need to be transposed in order to orient coherently the faces, we conclude that

$$Z(\lambda) = \sum_{\mathcal{M}} \frac{|\operatorname{Stab}_{\mathfrak{T}}(\mathcal{M})|}{V(\mathcal{M})! \, 2^{V(\mathcal{M})}} \Big( \prod_{v \in \mathcal{M}} \frac{1}{\prod_{q} \deg^{q} v!} \Big) \sum_{[\mathbb{G}] \in \operatorname{Orb}_{\mathfrak{T}}(\mathcal{M})} \mathcal{A}(\mathbb{G}),$$
  

$$\mathcal{A}(\mathbb{G}) = ((-1)^{\sum_{c \in \mathcal{D}} |c|})^{V(\mathbb{G})} 2^{E^{\varrho}(\mathbb{G})} \prod_{q \in Q} (-\lambda_{q})^{E_{q}(\mathbb{G})}$$

$$\times \Big( \prod_{C_{q} \neq \emptyset} \prod_{e \in \mathcal{E}_{\mathcal{C}_{q}}(\mathbb{G})} (-1)^{\sum_{c \in \mathcal{C}_{q}} \tau^{c}(e)|c|} \Big) \prod_{c} \prod_{f \in \mathcal{F}_{c}(\mathbb{G})} (-1)^{t(f)|c|} N_{c},$$

$$(4.2)$$

where t(f) denotes the number of transpositions needed to orient the face f coherently.

**Amplitudes.** Up to the overall coupling constants, the amplitude of a graph factors over the ribbon graphs  $\mathbb{G}_c$ 

$$\begin{aligned} \mathcal{A}(\mathbb{G}) &= 2^{E^{\varrho}(\mathbb{G})} \prod_{q \in \mathcal{Q}} (-\lambda_q)^{E_q(\mathbb{G})} \prod_{c \in \mathcal{D}} \left[ (-1)^{V(\mathbb{G}_c)|c|} \Big( \prod_{e \in \mathcal{E}(\mathbb{G}_c)} (-1)^{\tau^c(e)|c|} \Big) \right. \\ & \times \Big( \prod_{f \in \mathcal{F}_c(\mathbb{G})} (-1)^{t(f)|c|} N_c \Big) \Big], \end{aligned}$$

and using Theorem A.11, this is

$$\mathcal{A}(\mathbb{G}) = 2^{E^{\varrho}(\mathbb{G})} \prod_{q \in \mathcal{Q}} (-\lambda_q)^{E_q(\mathbb{G})} \prod_{c=1}^{D} ((-1)^{|c|} N_c)^{F_c(\mathbb{G})},$$
(4.3)

and thus obeys the  $N_c \rightarrow -N_c$  duality. The  $\rho$ -edges associated to the unique disconnected invariant  $\mathcal{C}_q = \emptyset$  do not have a twisted or untwisted state and bring a relative factor of two compared to the other multi-ribbon edges.

The two-point function. The connected two-point function of the tensor model

$$G_2(\lambda) := Z^{-1}(\lambda) \int d\mu[T] T^{a_{\mathcal{D}}} T^{b_{\mathcal{D}}} g^1_{a_1b_1} \cdots g^D_{a_Db_D}$$

can be expressed as a perturbative series over rooted multi-ribbon graphs. As in the matrix case, rooting drastically simplifies the combinatorial factors. The DSE for  $G_2$  follows from

$$0 = \int dT \frac{\partial}{\partial T^{a_{\mathcal{D}}}} (T^{a_{\mathcal{D}}} e^{-S[T]}) \Rightarrow G_2(\lambda) = \prod_{c \in \mathcal{D}} N_c + \sum_{q \in Q} 4\lambda_q \partial_{\lambda_q} \ln Z.$$
(4.4)

Graphically, the derivatives select an edge of a multi-ribbon graph and because every edge has two halfedges,  $\sum_{q \in \Omega} 2\lambda_q \partial_{\lambda_q}$  generates a sum over all possible rootings. Rooted unlabeled multi-ribbon graphs are equivalence classes of labeled multi-ribbon graphs that differ only by relabeling of their halfedges, but keeping the root halfedge fixed. The calculation of  $|\text{Stab}_{\mathfrak{T}}(\mathbb{G})|$  is a straightforward generalization of the ordinary ribbon graph case, and in Proposition B.1 we show that

$$G_2(\lambda) = \sum_{[\mathbb{G}] \text{ connected, rooted}} \frac{1}{2^{C(\mathbb{G} - \mathcal{E}^{\varrho}) - 1}} \mathcal{A}(\mathbb{G}), \tag{4.5}$$

where  $C(\mathbb{G} - \mathcal{E}^{\varrho})$  counts the number of connected components of the multi-ribbon graph obtained after deletion of the  $\varrho$ -edges. The free energy  $\ln Z(\lambda)$  can be obtained by integrating the DSE,

$$\ln Z(\lambda) = \sum_{\substack{[\mathbb{G}] \text{ connected, rooted,}\\ \text{at least one } E_q > 0}} \frac{1}{2^{C(\mathbb{G} - \mathcal{E}^{\varrho}) + 1} \sum_{q \in \mathcal{Q}} E_q(\mathbb{G})} \mathcal{A}(\mathbb{G}).$$
(4.6)

**Rescaled theories.** Models which admit a good 1/N expansion involve couplings rescaled by various powers of N. In order to maintain the N to -N duality of the amplitudes, one needs sometimes to flip the sign of the couplings. For instance, for D = 2, in order to get a sensible large N limit, one needs to rescale the coupling by a factor N. If one rescales  $\lambda \to \lambda/N$  in the  $O(N) \otimes O(N)$ -model and  $\lambda \to -\lambda/N$  in the  $Sp(N) \otimes Sp(N)$ -model, the amplitudes graphs differ by  $(-1)^{\chi(\mathcal{G})}$ .<sup>8</sup> The equality is reestablished if one sends at the same time  $\lambda \to -\lambda$ .

<sup>&</sup>lt;sup>8</sup>This was also found in [43].

# A. Classification of ribbon graphs

#### A.1. Canonical form

We prove in this subsection that a ribbon graph can be brought into a canonical form obtained by first separating the oriented and unoriented parts of the graph (Proposition A.1) and then simplifying the oriented part (Proposition A.3).

**Proposition A.1.** Every connected ribbon graph  $\mathcal{G}$  is homeomorphic as a topological surface (two-dimensional CW complex) to a ribbon graph  $\mathcal{G}'$  such that

- *G'* has only one vertex;
- *G'* has either none, or one or two twisted simple self-loops;
- all the remaining edges of  $\mathcal{G}'$  are untwisted.

Equivalently,

$$\mathscr{G} \cong \mathscr{G}' \cong \begin{cases} \mathscr{G}_{\Sigma_g}, & \text{orientable with } k = 2g, \\ \mathscr{G}_{\Sigma_g} \lor \mathscr{G}_{\mathbb{RP}^2}, & \text{unorientable with } k = 2g+1, \\ \mathscr{G}_{\Sigma_g} \lor \mathscr{G}_{\mathbb{RP}^2} \lor \mathscr{G}_{\mathbb{RP}^2}, & \text{unorientable with } k = 2g+2, \end{cases}$$

where  $\mathcal{G}_{\Sigma_g}$  a ribbon subgraph of  $\mathcal{G}'$  containing only untwisted edges and is cellularly embedded into a closed orientable surface  $\Sigma_g$  with orientable genus g (we reserve the notation g for the orientable genus) and k is the non-orientable genus of  $\mathcal{G}'$ .

Proposition A.1 is illustrated in Figure 11.

In order to state our second proposition, we need the notion of *clean nice crossing*.

**Definition A.2** (Nice crossing and clean nice crossing). Let  $e = \{e_1, e_2\}$  and  $f = \{f_1, f_2\}$  be two untwisted self-loop edges connected to the same vertex v of a ribbon graph. Assume  $f_1 < f_2$  and  $e_1 < e_2$  in the cyclic order around v.

- The pair (e, f) is a *nice crossing* [29] if and only if  $e_2$  is the successor of  $f_1$ .
- A nice crossing (e, f) is called *clean nice crossing* if there is no other halfedge h of v distinct from  $e_2$ ,  $f_1$  satisfying  $e_1 < h < f_2$ , i.e., along v the halfedges are encountered in the order  $\ldots e_1 f_1 e_2 f_2 \ldots$

**Proposition A.3.** Every ribbon graph  $\mathcal{G}$  composed of only untwisted edges is homeomorphic as a topological surface (two-dimensional CW complex) to a ribbon graph  $\mathcal{G}'$ with one vertex, one face and 2g edges forming g clean nice crossings, where g is the orientable genus of  $\mathcal{G}$ . Equivalently,

$$\mathscr{G} \simeq \mathscr{G}' \cong \mathscr{G}_{\circ} \bigvee_{g} \mathscr{G}_{T^2} \quad with \ \chi = 2 - 2g.$$



**Figure 11.** Illustration of Proposition A.1. In the first line the orientable part was further simplified using Proposition A.3. We call the right-hand side the canonical form.

Note that Proposition A.3 can be applied to  $\mathscr{G}_{\Sigma_g}$  in Proposition A.1, yielding

$$\mathscr{G} \simeq \mathscr{G}_{\circ} \bigvee_{g} \mathscr{G}_{T^2} \bigvee_{0, 1 \text{ or } 2} \mathscr{G}_{\mathbb{RP}^2}.$$

We call the right-hand side of this equation the *canonical form* of  $\mathcal{G}$ , see Figure 11. This is the ribbon graph version of classification theorem of closed surfaces, stating that every such surface is homeomorphic to the connected sum of a sphere, some number of tori, and either no, one or two real projective planes.

**Contraction and sliding of edges.** We introduce two homeomorphisms of ribbon graphs, viewed as a topological surface with boundary. Similar moves are known in the literature [23, 42].

**Definition A.4** (Contraction of an edge, see Figure 12). Let  $\mathscr{G}$  be a ribbon graph and  $e \in \mathscr{E}(\mathscr{G})$  an edge connecting two distinct vertices  $v, w \in \mathcal{V}(\mathscr{G})$  of coordination p and q. Remember that v, w and e are all topological discs.

If e is untwisted, we define  $\mathscr{G}/e$  to be the ribbon graph obtained from  $\mathscr{G}$  by replacing v, w and e by the single vertex  $u = v \cup e \cup w$  (which is again a topological disc) of coordination p + q - 2 such that in the cyclic ordering around this vertex the halfedges of v proceed the halfedges of w. The ribbon graph  $\mathscr{G}/e$  has one vertex and one edge fewer than  $\mathscr{G}$ , but the same number of faces.

If e is twisted, we first push the twist along the graph by reembedding the vertex w such that e is untwisted and proceed as before.



Figure 12. Contraction of an untwisted (*first line*) and twisted (*second line*) edge in a ribbon graph.



**Figure 13.** Sliding of edges Ia and Ib. The horizontal line is the vertex with ordering from right to left.

The contraction preserves the Euler characteristic and the orientability, and is thus a homeomorphism of surfaces.

A spanning tree of  $\mathcal{G}$ , that is, a connected acyclic subgraph  $\mathcal{T} \subset \mathcal{G}$ , has  $E(\mathcal{T}) = V(\mathcal{G}) - 1$  edges. One can contract all the edges in a spanning tree and decrease the numbers of vertices and edges of  $\mathcal{G}$  to

$$V(\mathcal{G}) \to V(\mathcal{G}) - (V(\mathcal{G}) - 1) = 1$$
 and  $E(\mathcal{G}) \to E(\mathcal{G}) - (V(\mathcal{G}) - 1)$ 

The resulting graph is a rosette graph homeomorphic to  $\mathcal{G}$ .

**Definition A.5** (Sliding of edges Ia, see Figure 13). Let  $e = \{e_1, e_2\}$  be a twisted self-loop edge on the vertex v of a ribbon graph. In the cyclic ordering of halfedges around v, let  $e_1 < e_2$  and denote by  $e_1 < h_1 < h_2 < \cdots < h_n < e_2$  all the halfedges of v that are between  $e_1$  and  $e_2$ .

Sliding of the halfedges  $h_1, \ldots, h_n$  out of the twisted edge e is defined as

- (1) Reordering the halfedges to  $h_n < \cdots < h_2 < h_1 < e_1 < e_2$ .
- (2) Adding a twist (recall that two twists on the same edge cancel) to all the edges to which  $h_1, \ldots, h_n$  belong.



Figure 14. Sliding of edges IIa and IIb.

Note that the order of the  $h_i$ 's has been reversed. Also, note that and after the sliding, e is a simple twisted self-loop.

**Definition A.6** (Sliding of edges Ib, see Figure 13). Let  $e = \{e_1, e_2\}$  be a simple twisted self-loop on the vertex v of a ribbon graph. In the cyclic ordering of halfedges around v, let  $e < 1 < e_2 < h_1 < h_2 < \cdots < h_n$  a with  $h_i$  a collection of consecutive halfedges preceding  $e_1$  on v. As e is a simple self-loop, there is no halfedge between  $e_1$  and  $e_2$ .

Sliding of the halfedges  $h_1, \ldots, h_n$  past the twisted edge e is defined as

(1) Reordering the halfedges to  $e_1 < e_2 < h_1 < h_2 < \cdots < h_n$ .

Note that the relative order of the  $h_i$ 's has not changed, no additional twists were introduced and e remains a simple twisted self-loop.

Both sliding operation (Ia) and (Ib) preserve the number of faces, do not change the numbers of vertices and edges and do not alter the orientability. Thus these operations are homeomorphisms of two-dimensional surfaces.

**Definition A.7** (Sliding of edges IIa, see Figure 14). Let  $(e = \{e_1, e_2\}, f = \{f_1, f_2\})$  be a nice crossing at the vertex v of a ribbon graph. In the cyclic ordering of halfedges around v, let us denote by

$$e_1 < h_1 < \dots < h_n < f_1 < e_2 < k_1 < \dots < k_m < f_2$$

the halfedges located between  $e_1$  and  $f_2$ .

Sliding of the halfedges  $h_1, \ldots, h_n, k_1, \ldots, k_m$  out of the nice crossing (e, f) is defined as

(1) Reordering the halfedges to  $k_1 < \cdots < k_m < h_1 < \cdots < h_n < e_1 < f_1 < e_2 < f_2$ .

Note that the order of the set of  $h_i$ 's and  $k_j$ 's was interchanged, but the relative order in each set remained unchanged. After sliding, (e, f) is a clean nice crossing.

**Definition A.8** (Sliding of edges IIb, see Figure 14). Let  $(e = \{e_1, e_2\}, f = \{f_1, f_2\})$  be a clean nice crossing at the vertex v of a ribbon graph. In the cyclic ordering of halfedges around v, let us denote by

$$h_1 < \cdots < h_n < e_1 < f_1 < e_2 < f_2$$

a collection of consecutive halfedges preceding  $e_1$  on v.

Sliding of the halfedges  $h_1, \ldots, h_n$  past the clean nice crossing (e, f) is defined as

(1) Reordering the halfedges to  $e_1 < f_1 < e_2 < f_2 < h_1 < \cdots < h_n$ .

Note that the relative order of the  $h_i$ 's is unchanged; (e, f) remains a clean nice crossing.

Like the sliding along twisted edges, the sliding along a nice crossing (IIa, IIb) is a homeomorphism of two-dimensional surfaces.

*Proof of Proposition* A.1. Let  $\mathcal{G}$  be a connected ribbon graph.

*First.* Contract a spanning tree  $\mathcal{T} \subset \mathcal{G}$ . This decreases the number of edges and vertices by  $V(\mathcal{G}) - 1$  and the resulting ribbon graph  $\mathcal{G}/\mathcal{T}$  is a rosette graph, that is, a graph with only one vertex.

Second. If  $\mathscr{G}/\mathcal{T}$  does not contain any twisted edges, then it can be embedded into an orientable surface  $\Sigma_g$  of genus g.

Otherwise, use sliding out of twisted self-loop edges (Ia) to create simple twisted self-loops. This operation may create new twists in the halfedges. Once a twisted self-loop is created, use the slide (Ib) to move it "to the right" on the vertex.

Proceed until all the twisted edges of the rosette graph belong to simple twisted self-loops. The resulting graph is a connected sum of an orientable graph  $\mathcal{O}$  containing only untwisted edges and p copies of  $\mathscr{G}_{\mathbb{RP}^2}$ , i.e., ribbon graphs with only one simple twisted loop:

$$\mathcal{O} \vee \underbrace{\mathscr{G}_{\mathbb{RP}^2} \vee \cdots \vee \mathscr{G}_{\mathbb{RP}^2}}_{p\text{-times}}.$$

*Third.* By sliding as depicted in Figure 15, three neighboring simple twisted self-loops can be reduced to one simple twisted self-loop and a clean nice crossing:

$$\mathscr{G}_{\mathbb{RP}^2} \vee \mathscr{G}_{\mathbb{RP}^2} \vee \mathscr{G}_{\mathbb{RP}^2} \cong \mathscr{G}_{\mathbb{RP}^2} \vee \mathscr{G}_{T^2},$$

hence it is possible to reduce the number of simple twisted self-loops (and twisted edges in total) to zero, one or two. Slide (Ib) the clean nice crossings to the left of the twisted self-loops.



**Figure 15.** Deforming three neighboring simple twisted self-loops into a graph with only one twisted edge. (a) and (b): By inverting (Ia), slide a halfedge of the left and right twisted simple loop into the central one. This creates a nice crossing. (c): Use (IIa) to slide the central twisted loop out of the nice crossing. (d): The result has only a single simple twisted loop.

*Finally.* One arrives at a graph  $\mathcal{G}'$  of the form

$$\mathscr{G}' \cong \begin{cases} \mathscr{G}_{\Sigma_g}, & \text{orientable with } k = 2g, \\ \mathscr{G}_{\Sigma_g} \lor \mathscr{G}_{\mathbb{RP}^2}, & \text{unorientable with } k = 2g+1, \\ \mathscr{G}_{\Sigma_g} \lor \mathscr{G}_{\mathbb{RP}^2} \lor \mathscr{G}_{\mathbb{RP}^2}, & \text{unorientable with } k = 2g+2 \end{cases}$$

with  $\chi(\mathscr{G}) = \chi(\mathscr{G}') = 2 - k$ .

*Proof of Proposition* A.3. Let  $\mathscr{G}$  be a connected ribbon graph with only untwisted edges. Such a graph can be embedded into an orientable surface.

*First.* Contract a spanning tree  $\mathcal{T}_1 \subset \mathcal{G}$  to arrive at a rosette graph  $\mathcal{G}/\mathcal{T}_1$ .

Second. Contract a spanning tree in the dual graph  $\mathcal{T}_2 \subset (\mathcal{G}/\mathcal{T}_1)^*$ . This corresponds to deleting edges in  $\mathcal{G}$  in a way that preserves the Euler characteristic, the orientability and the connectivity.

This reduces the number of faces and edges by  $F(\mathcal{G}) - 1$  and gives a superrosette graph  $\mathcal{R}$ , that is, a graph with one vertex, one face and only untwisted edges. A superrosette always contains at least one nice crossing.

*Third.* Choose a nice crossing (e, f) in  $\mathcal{R}$  and slide (IIa) all the halfedges encompassed by the nice crossing out of (e, f). The result has the structure

$$\mathcal{R} \cong \mathcal{R}/(e, f) \vee \mathcal{G}_{T^2},$$

where  $\mathcal{R}/(e, f)$  is again a superrosette graph with genus decreased by one. Iterating one arrives at

$$\mathscr{R} \cong \mathscr{G}_{\circ} \vee \underbrace{\mathscr{G}_{T^2} \vee \cdots \vee \mathscr{G}_{T^2}}_{g\text{-times}}.$$

#### A.2. Sign of a ribbon graph

Let  $\mathscr{G}$  be a connected ribbon graph. An *a priori arrow orientation*<sup>9</sup> of a  $\mathscr{G}$  (which has nothing to do with the orientability of the embedding surface) is defined by the following:

- (1) counter-clockwise pointing arrows at the corners of each vertex;
- (2) parallel pointing arrows on the strands of each edge.

We denote  $\tau(e) = 0$  if the edge *e* is untwisted (straight) and  $\tau(e) = 1$  if the edge *e* is twisted. Furthermore, we denote by t(f) the number of reorientations of arrows required to coherently orient the face *f* with all the arrows pointing in the same direction along its boundary. The *sign* of  $\mathcal{G}$  is defined as

$$\operatorname{sgn}(\mathscr{G}) = (-1)^{V(\mathscr{G})} \Big(\prod_{e \in \mathscr{E}(\mathscr{G})} (-1)^{\tau(e)} \Big) \Big(\prod_{f \in \mathscr{F}(\mathscr{G})} (-1)^{t(f)} \Big).$$

This is well defined. In order to determine the sign of  $\mathcal{G}$ , one needs to determine the number of arrow flips that are necessary to go from an a priori orientation of  $\mathcal{G}$ to an orientation where all arrows point coherently along the faces of  $\mathcal{G}$  (such an orientation will be called *coherent*). Since every face consists of as many corners as edge strands, the total number of arrows along a face is even and switching between two coherent orientations requires an even number of arrow flips. Also, as any two a priori orientations differ by an even number of arrow flips (pairs of arrows along the edge strands), switching between a priori orientations at fixed coherent orientation does not change the sign of the graph.

#### **Lemma A.9.** The sign of a graph is

- invariant under reembedding of the vertices;
- invariant under contraction of a tree edge.

*Proof.* Consider an a priori arrow orientation of  $\mathcal{G}$ . Re-embedding a vertex of degree d brings d new twists, but one needs to reverse d vertex corners in order to orient the re-embedded vertex counter-clockwise.

Consider now a tree edge e connecting two vertices v and w in a graph with a priori orientation (which by the first item we can assume to be untwisted). A flip of an arrow coherently orients the disc  $u = v \cup e \cup w$ , but this is canceled by the fact that under contraction the number of vertices of the graph goes down by 1.

**Lemma A.10.** The sign of a graph is invariant under the sliding moves.

*Proof.* We consider a graph  $\mathscr{G}$  having a twisted self-loop as in Figure 16 (a).

<sup>&</sup>lt;sup>9</sup>This is the arrow orientation encountered in Section 4.



**Figure 16.** Sliding I at a twisted self-loop in a coherently oriented graph. The red and blue corners and strands belong to the red and blue face, respectively. The number of reversed arrows and additional twists is always even.



Figure 17. Sliding II at a nice crossing in a coherently oriented graph. No arrows are reversed, nor are halfedges twisted.

We denote by  $\mathscr{G}'$  the graph obtained from  $\mathscr{G}$  by the sliding Ia. All else being equal, in order to pass from an a priori orientation of  $\mathscr{G}$  to the coherent orientation depicted in Figure 16 (a) two corner arrows had to be reversed, while for  $\mathscr{G}'$  only one. However,  $\mathscr{G}'$  has one twist more than  $\mathscr{G}$ . As the graphs are otherwise identical, they have the same sign. For Ib sliding, there is no extra twist, but both graphs need only one local arrow reorientation.

We now consider a graph  $\mathcal{G}$  having a nice crossing as in Figure 17 (a) and denote by  $\mathcal{G}'$  the graph obtained from  $\mathcal{G}$  after sliding. In all the cases, the same number of arrow flips is needed in order to pass locally from an a priori to the coherent orientations depicted. As  $\mathcal{G}$  and  $\mathcal{G}'$  are identical elsewhere, they have the same sign.

**Theorem A.11** (Sign of a ribbon graph). For any connected ribbon graph G, we have

$$\operatorname{sgn}(\mathscr{G}) = (-1)^{V(\mathscr{G})} \Big(\prod_{e \in \mathscr{E}(\mathscr{G})} (-1)^{\tau(e)} \Big) \Big(\prod_{f \in \mathscr{F}(\mathscr{G})} (-1)^{t(f)} \Big) = (-1)^{F(\mathscr{G})} \Big)$$

where  $F(\mathcal{G})$  is the number of faces of  $\mathcal{G}$ .

*Proof.* From Lemmas A.9 and A.10, the sign of a graph is invariant under the reduction moves used in Proposition A.1. It follows that, not only

$$\mathcal{G} \cong \mathcal{G}' \cong \begin{cases} \mathcal{G}_{\Sigma_g}, & \text{orientable with } k = 2g, \\ \mathcal{G}_{\Sigma_g} \vee \mathcal{G}_{\mathbb{RP}^2}, & \text{unorientable with } k = 2g+1, \\ \mathcal{G}_{\Sigma_g} \vee \mathcal{G}_{\mathbb{RP}^2} \vee \mathcal{G}_{\mathbb{RP}^2}, & \text{unorientable with } k = 2g+2, \end{cases}$$

but also  $sgn(\mathcal{G}) = sgn(\mathcal{G}')$ . The sign of  $\mathcal{G}'$  is easy to compute using the following:

- $\mathcal{G}'$  has one vertex;
- each simple twisted self-loop brings a (-1) for the twist and another (-1) in order to change from an a priori arrow orientation to a coherent one;
- the number of untwisted edges of S' is the number of edges of S<sub>Σg</sub>, that is, *E*(S<sub>Σg</sub>). Exactly one arrow for each such edge needs to be flipped in order to pass from an a priori to a coherent arrow orientation of S'.

Therefore,  $\operatorname{sgn}(\mathscr{G}') = (-1)^{1+E(\mathscr{G}_{\Sigma_g})}$ . The theorem follows by observing that the Euler relation for  $\mathscr{G}_{\Sigma_g}$  reads  $1 - E(\mathscr{G}_{\Sigma_g}) + F(\mathscr{G}_{\Sigma_g}) = 2 - 2g(\mathscr{G}_{\Sigma_g})$ , hence  $F(\mathscr{G}_{\Sigma_g}) = 1 + E(\mathscr{G}_{\Sigma_g}) \mod 2$ , and the number of faces is invariant under contraction and sliding  $F(\mathscr{G}_{\Sigma_g}) = F(\mathscr{G}') = F(\mathscr{G})$ .

# **B.** Symmetry factors

The aim of this section is to prove the following proposition.

Proposition B.1. The perturbative series of the two-point function

$$G_2(\lambda) := Z^{-1}(\lambda) \int d\mu[T] T^{a_{\mathcal{D}}} T^{b_{\mathcal{D}}} g^1_{a_1b_1} \cdots g^D_{a_Db_D}$$

can be written as the sum

$$G_2(\kappa, \lambda) = \sum_{[\mathscr{G}^{\varrho}] \text{ connected, rooted}} \frac{1}{2^{C(\mathscr{G}^{\varrho} - \mathscr{E}^{\varrho}) - 1}} \mathcal{A}(\mathscr{G}^{\varrho}).$$

Before proving this proposition, we discuss some useful facts. The symmetry factor of a ribbon graph in the perturbative series (3.4) of  $Z(\lambda)$  is obtained as

- a factor  $\frac{1}{2^V n_{\text{IVP}}}$ , where  $n_{\text{IVP}}$  is the number of permutations of vertex labels, that give the same labeled map;
- a factor  $\frac{1}{\deg v! \deg^{\varrho} v!}$  for every vertex;
- a factor counting the number of ways to connect labeled halfedges to form the same combinatorial map *M<sup>Q</sup>* underlying *S<sup>Q</sup>*, taking into account the different ways to label the halfedges;

 a factor |Stab<sub>𝔅</sub>| and the number of combinatorial maps such that 𝔅<sup>𝔅</sup> is contained in their orbits under 𝔅.

For example, the weight of the ribbon graph in Figure 7 is

$$\underbrace{\frac{1}{2^{3} \cdot 1}}_{2^{V} n_{\text{IVP}}} \underbrace{\frac{2^{2}}{5! \cdot 1! \cdot 2! \cdot 0! \cdot 1! \cdot 3}}_{\prod_{v} \log v! \deg^{\varrho} v!} \cdot \underbrace{\frac{1}{2 \cdot 5! \cdot 3!}}_{\text{Interpretation}} = \frac{1}{2}.$$

Stabilizer of rooted ribbon graphs with respect to  $\mathfrak{T}$ . Rooting simplifies the calculation of  $|\operatorname{Stab}_{\mathfrak{T}}(\mathcal{M})|$ . The finite group  $\mathfrak{T}$  that twists the ribbon edges is defined on graphs with a fixed but arbitrary labeling of their edges. The rooting can be used to induce such a labeling: Fix a spanning tree and enumerate all edges as they are encountered on a counter-clockwise walk following the unique face of the tree, starting at the root.

We first focus on ordinary combinatorial maps and ribbon graphs. The results can be generalized to graphs with  $\rho$ -edges, by considering the ordinary ribbon graph that is obtained by deleting the  $\rho$ -edges.

**Lemma B.2.** Let  $\mathcal{G}$  be a rooted, connected ribbon graph. We denote by  $V_1$  and  $V_2$  the numbers of non-root vertices of degree one and two, respectively. Then

$$|\operatorname{Stab}_{\mathfrak{T}}(\mathscr{G})| = 2^{V_1 + V_2}$$

*Proof.* The orientation of the root vertex is held fixed. If a non-root vertex has degree one, twisting the edge incident to it does not change the ribbon graph—the twist is "reducible". If a non-root vertex has degree two, twisting both incident edges again does not change the ribbon graph. If both halfedges of a degree two vertex belong to the same edge, it is necessarily the root vertex, since  $\mathscr{G}$  is assumed to be connected. This is depicted in Figure 18.

It follows that

$$\sum_{\substack{\mathcal{M} \\ \text{connected,} \\ \text{rooted}}} \frac{|\text{Stab}_{\mathfrak{T}}(\mathcal{M})|}{2^{V}} \sum_{[\mathscr{G}] \in \text{Orb}_{\mathfrak{T}}(\mathcal{M})} \mathcal{A}(\mathscr{G}) = \sum_{\substack{\mathcal{M} \\ \text{connected,} \\ \text{rooted}}} \frac{1}{2^{V_{\geq 3}}} \sum_{[\mathscr{G}] \in \text{Orb}_{\mathfrak{T}}(\mathcal{M})} \mathcal{A}(\mathscr{G}), \quad (B.1)$$

where  $V_{\geq 3}$  denotes the number of non-root vertices of degree  $\geq 3$ . In order to reshuffle this expression into a sum over rooted ribbon graphs, we recall that two ribbon graphs are equivalent if one can be obtained from the other by vertex re-embeddings. This implies that if two combinatorial maps  $\mathcal{M}_1$  and  $\mathcal{M}_2$  differ only by reversing the order of halfedges around some of their vertices, then  $\operatorname{Orb}_{\mathfrak{T}}(\mathcal{M}_1) = \operatorname{Orb}_{\mathfrak{T}}(\mathcal{M}_2)$ . Reversing the order of halfedges at a vertex of degree lower than three is trivial, hence for rooted



Figure 18. Reducible twists at vertices of degree one and two.

ribbon graphs the multiplicity is  $2^{V \ge 3}$ . As a result, the perturbative series of the twopoint function  $G_2$  (3.5) for  $\kappa = 0$  is

$$G_2(0,\lambda) = \sum_{[\mathscr{G}] \text{ connected, rooted}} \mathcal{A}(\mathscr{G}).$$

When taking the  $\rho$ -edges into account, we recall that  $\mathfrak{T}$  acts trivially on them. Thus, it is sufficient to consider the ribbon graph  $\mathscr{G} = \mathscr{G}^{\rho} - \mathscr{E}^{\rho}$  obtained by deleting all the  $\rho$ -edges of  $\mathscr{G}^{\rho}$ . However, when calculating  $|\operatorname{Stab}_{\mathfrak{T}}(\mathscr{G}^{\rho})|$  a subtlety arises:  $\mathscr{G} = \mathscr{G}^{\rho} - \mathscr{E}^{\rho}$  is not necessarily connected. The  $\rho$ -edges can be used to induce a rooting at every connected component  $\mathscr{G}_c \subset \mathscr{G}$ : Consider the connected components as effective vertices in a graph with only  $\rho$ -edges; pick a spanning tree in that graph; from every  $\mathscr{G}_c$ there is a unique path in the tree to the original root; let the halfedge of  $\mathscr{G}_c$ , belonging to that path, be another root. The stabilizer  $\operatorname{Stab}_{\mathfrak{T}}(\mathscr{G})$  factors over the  $\mathscr{G}_c$  and using Lemma B.2 for each rooted connected component, one obtains

$$|\operatorname{Stab}_{\mathfrak{T}}(\mathscr{G})| = \prod_{\mathscr{G}_{\mathcal{C}} \subset \mathscr{G}} 2^{V_1(\mathscr{G}_{\mathcal{C}}) + V_2(\mathscr{G}_{\mathcal{C}})}$$

One has to partially resume the double sum over combinatorial maps and ribbon graphs with  $\rho$ -edges analogous to (B.1) into a sum over rooted ribbon graphs with  $\rho$ -edges. The multiplicity of a ribbon graph with multiple rooted connected components is  $\prod_{\mathscr{G}_C \subset \mathscr{G}} 2^{V_{\geq 3}(\mathscr{G}_C)}$  and one arrives at

$$\prod_{\mathscr{G}_C \subset \mathscr{G}} 2^{V(\mathscr{G}_C) - 1} = 2^{V(\mathscr{G}) - C(\mathscr{G})},$$

where C denotes the number of connected components: the -1 in the exponent appears because  $V_1$ ,  $V_2$  and  $V_{\geq 3}$  count only non-root vertices, hence sum up to  $V(\mathcal{G}_c) - 1$  in each connected component.

*Proof of Proposition* B.1. The discussion above goes through mutatis mutandis for multi-ribbon graphs. Combining (4.2) with (4.4), the perturbative series of the two-point function can be written as

$$G_{2}(\lambda) = \sum_{\substack{\mathcal{M} \\ \text{connected, rooted}}} \frac{|\text{Stab}_{\mathfrak{T}}(\mathcal{M})|}{V(\mathcal{M})! \, 2^{V(\mathcal{M})-1}} \Big(\prod_{v \in \mathcal{M}} \frac{1}{\prod_{q} \deg^{q} v!} \Big) \sum_{[\mathbb{G}] \in \text{Orb}_{\mathfrak{T}}(\mathcal{M})} \mathcal{A}(\mathbb{G})$$

with  $\mathcal{M}$  and edge multicolored combinatorial map. All objects in the above expression are fully labeled. Rooting prevents non-trivial symmetry factors, thus it is sufficient to count the ways to assign labels to a multi-ribbon graph:

- (1) Pick a spanning tree.
- (2) There are V! ways to label the vertices.
- (3) At the root vertex  $v_0$ , the root breaks the cyclicity of halfedges, thus there are  $\prod_q \deg^q v_0!$  ways to label the different types of multi-ribbon halfedges.
- (4) At each non-root vertex one halfedge is part of the unique path in the tree towards the root.

This again breaks cyclicity and there are  $\prod_q \deg^q v!$  ways to label the halfedges. The amplitudes do not depend on the labeling, thus, in terms of unlabeled but rooted objects,

$$G_{2}(\lambda) = \sum_{\substack{\mathcal{M} \text{ connected,} \\ \text{unlabeled}}} \frac{|\text{Stab}_{\mathfrak{T}}(\mathcal{M})|}{2^{V(\mathcal{M})-1}} \sum_{[\mathbb{G}] \in \text{Orb}_{\mathfrak{T}}(\mathcal{M})} \mathcal{A}(\mathbb{G}) = \sum_{\substack{[\mathbb{G}] \\ \text{connected,} \\ \text{roted}}} \frac{1}{2^{C(\mathbb{G}-\mathcal{E}^{\varrho})-1}} \mathcal{A}(\mathbb{G}),$$

where  $C(\mathbb{G} - \mathcal{E}^{\varrho})$  counts the number of connected components of the multi-ribbon graph after deletion of the  $\varrho$ -edges.

For example,  $G_2$  up to quadratic order in the coupling constants for D = 2 is

Take, for example, the last three graphs. After deleting the  $\rho$ -edges, each graph splits into two connected components, this gives a factor  $\frac{1}{2}$  and in addition there are 4 distinct ways of rooting these graphs.

Acknowledgments. The authors would like to thank Dario Benedetti for comments and discussions at the early stages of this project.

**Funding.** This work was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No 818066) and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC-2181/1 – 390900948 (the Heidelberg STRUCTURES Cluster of Excellence).

## References

- J. Ambjørn, B. Durhuus, and T. Jónsson, Three-dimensional simplicial quantum gravity and generalized matrix models. *Modern Phys. Lett. A* 6 (1991), no. 12, 1133–1146 Zbl 1020.83537 MR 1115607
- [2] D. Anninos, T. Hartman, and A. Strominger, Higher spin realization of the DS/CFT correspondence. *Classical Quantum Gravity* 34 (2017), no. 1, article no. 015009
   Zbl 1354.83043 MR 3596141
- [3] D. Benedetti, S. Carrozza, R. Gurau, and M. Kolanowski, The 1/N expansion of the symmetric traceless and the antisymmetric tensor models in rank three. *Comm. Math. Phys.* 371 (2019), no. 1, 55–97 Zbl 1425.81071 MR 4015340
- [4] D. Bessis, C. Itzykson, and J. B. Zuber, Quantum field theory techniques in graphical enumeration. Adv. in Appl. Math. 1 (1980), no. 2, 109–157 Zbl 0453.05035 MR 603127
- [5] V. Bonzom, R. Gurau, and V. Rivasseau, Random tensor models in the large N limit: Uncoloring the colored tensor models. *Phys. Rev. D* 85 (2012), no. 2, article no. 084037
- [6] E. Brézin and V. A. Kazakov, Exactly solvable field theories of closed strings. *Phys. Lett. B* 236 (1990), no. 2, 144–150 MR 1040213
- [7] D. C. Brydges, Lectures on the renormalisation group. In *Statistical mechanics*, pp. 7–93, IAS/Park City Math. Ser. 16, American Mathematical Society, Providence, RI, 2009 Zbl 1186.82033 MR 2523458
- [8] K. Bulycheva, I. R. Klebanov, A. Milekhin, and G. Tarnopolsky, Spectra of operators in large N tensor models. Phys. Rev. D 97 (2018), no. 2, article no. 026016 MR 3877831
- [9] S. Carrozza, Large N limit of irreducible tensor models: O(N) rank-3 tensors with mixed permutation symmetry. J. High Energy Phys. 2018 (2018), no. 6, article no. 039 Zbl 1395.81155 MR 3850828
- [10] S. Carrozza and S. Harribey, Melonic large N limit of 5-index irreducible random tensors. Comm. Math. Phys. 390 (2022), no. 3, 1219–1270 Zbl 1487.81125 MR 4389081

- S. Carrozza and V. Pozsgay, SYK-like tensor quantum mechanics with Sp(N) symmetry. Nuclear Phys. B 941 (2019), 28–52 Zbl 1415.81131 MR 3916553
- [12] S. Carrozza and A. Tanasă, O(N) random tensor models. Lett. Math. Phys. 106 (2016), no. 11, 1531–1559 Zbl 1362.83010 MR 3555413
- [13] S. Chmutov, Generalized duality for graphs on surfaces and the signed Bollobás–Riordan polynomial. J. Combin. Theory Ser. B 99 (2009), no. 3, 617–638 Zbl 1172.05015 MR 2507944
- [14] P. Cvitanović and A. D. Kennedy, Spinors in negative dimensions. *Phys. Scripta* 26 (1982), no. 1, 5–14 Zbl 1063.22500 MR 669170
- [15] T. Delepouve, R. Gurau, and V. Rivasseau, Universality and Borel summability of arbitrary quartic tensor models. Ann. Inst. Henri Poincaré Probab. Stat. 52 (2016), no. 2, 821–848 Zbl 1341.81045 MR 3498011
- [16] P. Di Francesco, P. Ginsparg, and J. Zinn-Justin, 2D gravity and random matrices. *Phys. Rep.* 254 (1995), no. 1–2, 1–133 MR 1320471
- [17] M. R. Douglas and S. H. Shenker, Strings in less than one dimension. *Nuclear Phys. B* 335 (1990), no. 3, 635–654 MR 1059822
- [18] J. A. Ellis-Monaghan and I. Moffatt, Twisted duality for embedded graphs. *Trans. Amer. Math. Soc.* 364 (2012), no. 3, 1529–1569 Zbl 1238.05067 MR 2869185
- [19] J. A. Ellis-Monaghan and I. Moffatt, *Graphs on surfaces: Dualities, polynomials, and knots.* SpringerBriefs Math., Springer, New York, 2013 Zbl 1283.57001 MR 3086663
- [20] S. Giombi, I. R. Klebanov, F. Popov, S. Prakash, and G. Tarnopolsky, Prismatic large N models for bosonic tensors. *Phys. Rev. D* 98 (2018), no. 10, article no. 105005 MR 3954724
- [21] S. Giombi, I. R. Klebanov, and G. Tarnopolsky, Bosonic tensor models at large N and small ε. Phys. Rev. D 96 (2017), no. 10, article no. 106014 MR 3867822
- [22] S. Giombi and X. Yin, Higher spin gauge theory and holography: the three-point functions. J. High Energy Phys. 2010 (2010), no. 9, article no. 115 Zbl 1291.83107 MR 2776932
- [23] J. L. Gross and T. W. Tucker, *Topological graph theory*. Wiley-Intersci. Ser. Discrete Math. Optim., John Wiley & Sons, Inc., New York, 1987 Zbl 0621.05013 MR 898434
- [24] S. S. Gubser, M. Heydeman, C. Jepsen, S. Parikh, I. Saberi, B. Stoica, and B. Trundy, Melonic theories over diverse number systems. *Phys. Rev. D* 98 (2018), no. 12, article no. 126007 MR 3974315
- [25] R. Gurau, The complete 1/N expansion of colored tensor models in arbitrary dimension. Ann. Henri Poincaré 13 (2012), no. 3, 399–423 Zbl 1245.81118 MR 2909101
- [26] R. Gurau, The 1/N expansion of tensor models beyond perturbation theory. Comm. Math. Phys. 330 (2014), no. 3, 973–1019 Zbl 1297.81126 MR 3227505
- [27] R. Gurau, Invitation to random tensors. SIGMA Symmetry Integrability Geom. Methods Appl. 12 (2016), article no. 094 Zbl 1346.83030 MR 3550395
- [28] R. Gurau, Random tensors. Oxford University Press, Oxford, 2017 Zbl 1371.81007 MR 3616422
- [29] R. Gurau and V. Rivasseau, Parametric representation of noncommutative field theory. Comm. Math. Phys. 272 (2007), no. 3, 811–835 Zbl 1156.81465 MR 2304476

- [30] R. Gurau and J. P. Ryan, Colored tensor models a review. SIGMA Symmetry Integrability Geom. Methods Appl. 8 (2012), article no. 020 Zbl 1242.05094 MR 2942819
- [31] R. G. Gurau, Notes on tensor models and tensor field theories. Ann. Inst. Henri Poincaré D 9 (2022), no. 1, 159–218 Zbl 07509420 MR 4408001
- [32] R. G. Gurau and T. Krajewski, Analyticity results for the cumulants in a random matrix model. Ann. Inst. Henri Poincaré D 2 (2015), no. 2, 169–228 Zbl 1353.60009 MR 3354330
- [33] C. V. Johnson, The microstate physics of JT gravity and supergravity. 2022, arXiv:2201.11942
- [34] V. A. Kazakov, Ising model on a dynamical planar random lattice: Exact solution. *Phys. Lett. A* **119** (1986), no. 3, 140–144 MR 871244
- [35] I. R. Klebanov and A. M. Polyakov, AdS dual of the critical O(N) vector model. Phys. Lett. B 550 (2002), no. 3–4, 213–219 Zbl 1001.81057 MR 1948547
- [36] I. R. Klebanov, F. Popov, and G. Tarnopolsky, TASI lectures on large N tensor models. Proc. of Sci. 305 (2018), article no. PoS(TASI2017)004
- [37] V. G. Knizhnik, A. M. Polyakov, and A. B. Zamolodchikov, Fractal structure of 2Dquantum gravity. *Modern Phys. Lett. A* 3 (1988), no. 8, 819–826 MR 947880
- [38] M. Kontsevich, Intersection theory on the moduli space of curves and the matrix Airy function. *Comm. Math. Phys.* 147 (1992), no. 1, 1–23 Zbl 0756.35081 MR 1171758
- [39] A. LeClair and M. Neubert, Semi-Lorentz invariance, unitarity, and critical exponents of symplectic fermion models. J. High Energy Phys. 2007 (2007), no. 10, article no. 027 MR 2357951
- [40] L. Lionni and V. Rivasseau, Intermediate field representation for positive matrix and tensor interactions. Ann. Henri Poincaré 20 (2019), no. 10, 3265–3311 Zbl 1425.81074 MR 4009682
- [41] R. L. Mkrtchian, The equivalence of Sp(2N) and SO(-2N) gauge theories. *Phys. Lett. B* 105 (1981), no. 2–3, 174–176
- [42] I. Moffatt and E. Mphako-Banda, Handle slides for delta-matroids. *European J. Combin.* 59 (2017), 23–33 Zbl 1348.05109 MR 3546900
- [43] M. Mulase and A. Waldron, Duality of orthogonal and symplectic matrix integrals and quaternionic Feynman graphs. *Comm. Math. Phys.* 240 (2003), no. 3, 553–586 Zbl 1033.81062 MR 2005857
- [44] R. C. Penner, Perturbative series and the moduli space of Riemann surfaces. J. Differential Geom. 27 (1988), no. 1, 35–53 Zbl 0608.30046 MR 918455
- [45] P. Saad, S. H. Shenker, and D. Stanford, JT gravity as a matrix integral. 2019, arXiv:1903.11115
- [46] N. Sasakura, Super tensor models, super fuzzy spaces and super *n*-ary transformations. *Internat. J. Modern Phys. A* 26 (2011), no. 24, 4203–4216 Zbl 1247.83058
- [47] D. Stanford and E. Witten, JT gravity and the ensembles of random matrix theory. Adv. Theor. Math. Phys. 24 (2020), no. 6, 1475–1680 Zbl 07433590 MR 4285616
- [48] G. 't Hooft, A planar diagram theory for strong interactions. Nuclear Phys. B 72 (1974), no. 3, 461–473

- [49] A. Tanasă, Combinatorial physics: Combinatorics, quantum field theory, and quantum gravity models. Oxford University Press, Oxford, 2021 Zbl 1480.81004 MR 4516827
- [50] M. A. Vasiliev, Consistent equations for interacting gauge fields of all spins in 3 + 1 dimensions. *Phys. Lett. B* 243 (1990), no. 4, 378–382 Zbl 1332.81084 MR 1062824
- [51] E. Witten, Two-dimensional gravity and intersection theory on moduli space. In Surveys in differential geometry (Cambridge, MA, 1990), pp. 243–310, Lehigh University, Bethlehem, PA, 1991 Zbl 0757.53049 MR 1144529
- [52] E. Witten, An SYK-like model without disorder. J. Phys. A 52 (2019), no. 47, article no. 474002 Zbl 1509.81564 MR 4028950
- [53] P. Zinn-Justin, The general O(n) quartic matrix model and its application to counting tangles and links. *Comm. Math. Phys.* 238 (2003), no. 1–2, 287–304 Zbl 1033.57004 MR 1990878

Communicated by Adrian Tanasă

Received 13 July 2022; revised 13 January 2023.

#### Razvan Gurau

Institut für Theoretische Physik, Heidelberg University, Philosophenweg 19, 69120 Heidelberg, Germany; Centre de Physique Théorique (CPHT), Ecole Polytechnique, Route de Saclay, 91128 Palaiseau, France; Perimeter Institute for Theoretical Physics, 31 Caroline St. N, Waterloo, ON N2L 2Y5, Canada; gurau@thphys.uni-heidelberg.de

#### Hannes Keppler

Institut für Theoretische Physik, Heidelberg University, Philosophenweg 19, 69120 Heidelberg, Germany; keppler@thphys.uni-heidelberg.de