

Localizations and completions of stable ∞ -categories

LORENZO MANTOVANI (*)

ABSTRACT – We extend some classical results of Bousfield on homology localizations and nilpotent completions to a presentably symmetric monoidal stable ∞ -category \mathcal{M} admitting a multiplicative left-complete t -structure. If E is a homotopy commutative algebra in \mathcal{M} , we show that E -nilpotent completion, E -localization, and a suitable formal completion agree on bounded-below objects when E satisfies some reasonable conditions.

MATHEMATICS SUBJECT CLASSIFICATION (2020) – Primary 18N60; Secondary 55P60, 14F42.

KEYWORDS – Bousfield localization, nilpotent completion.

CONTENTS

1. Introduction	2
2. Preliminaries	5
3. Moore objects	12
4. Localization at some homotopy commutative algebras	23
5. Examples and applications	31
6. The E -based Adams–Novikov spectral sequence	35
7. Nilpotent resolutions	37
A. Appendices	54
References	60

(*) *Indirizzo dell’A.*: Institut für Mathematik, Johannes Gutenberg-Universität Mainz, Staudingerweg 9, 55128 Mainz, Germany; lorenzo.mantovani@uni-mainz.de

1. Introduction

1.1 – Motivation

Let \mathcal{M} be a presentably symmetric monoidal ∞ -category, with monoidal product \wedge and unit $\mathbf{1}$, and let E be an object of \mathcal{M} . One can construct a homology localization of \mathcal{M} by inverting all the maps ϕ in \mathcal{M} such that $\phi \wedge E$ is an equivalence. This construction was first introduced for the topological stable category \mathcal{SH} in [8]. The associated localization functor $X \mapsto X_E$ is called E -homology localization, and has a particularly simple universal property: X_E is the initial E -local object with a map $\lambda(X) : X \rightarrow X_E$. Unfortunately very little can be said about X_E for a general E , even in the case $X = \mathbf{1}$.

Assume now that E is a commutative algebra object in \mathcal{M} . In this situation, given any object $X \in \mathcal{M}$, we can perform a second construction X_E^\wedge , by setting

$$X_E^\wedge := \lim_{\Delta} X \wedge E^{\wedge \bullet+1}.$$

By construction, there is a natural map $\alpha(X) : X \rightarrow X_E^\wedge$ that factors through λ . The object X_E^\wedge is called *nilpotent completion of X at E* .

Instances of this construction have appeared in a wide range of contexts. When \mathcal{M} is (the nerve of) the category of modules over a ring A , and E is a commutative A -algebra, the map $\alpha(X)$ can be interpreted as the obstruction to recover X from its associated descent datum along $A \rightarrow E$. In algebraic topology, nilpotent completions were used by Adams and many others in relation to computing homotopy groups of spectra. Indeed, starting with the cosimplicial object

$$X \wedge E^{\wedge \bullet+1},$$

one can construct the tower of its partial totalizations and the inverse limit of such a tower recovers X_E^\wedge . In addition, the tower of partial totalizations gives rise to a Bousfield–Kan spectral sequence conditionally converging to X_E^\wedge . In some particularly favorable situation, the page of this spectral sequence is amenable to computations, and sometimes a good deal of information on X_E^\wedge can be understood via this spectral sequence.

Here is a crucial fact that follows from combining several parts of [8].

THEOREM 1.1.1 (Bousfield). *When E is a (-1) -connected commutative algebra in \mathcal{SH} , with $\pi_0(E) \simeq \mathbb{Z}/n\mathbb{Z}$, then for every bounded-below spectrum X the natural map $X_E \rightarrow X_E^\wedge$ is an equivalence. Furthermore, under the above assumptions, X_E is naturally identified with the derived completion $X_n^\wedge = \lim_k X/n^k$.*

1.2 – Actual content

In this paper, we axiomatize some of the techniques used by Bousfield and adapt them to work in a presentably symmetric monoidal stable ∞ -category \mathcal{M} . Our aim is to reach a formal analogue for \mathcal{M} of the above theorem of Bousfield. The main assumption we need on \mathcal{M} is that it comes endowed with a t -structure which has the following properties:

- (1) the t -structure is left-complete, i.e., $X \simeq \lim_n P^n(X)$ for all $X \in \mathcal{M}$;
- (2) the t -structure is multiplicative, i.e., $\mathbf{1} \in \mathcal{M}_{\geq 0}$ and $\mathcal{M}_{\geq p} \wedge \mathcal{M}_{\geq q} \subseteq \mathcal{M}_{\geq p+q}$;

The main application we have in mind being motivic homotopy theory, we have decided to dedicate Section 2 to recollecting some well-known facts about the motivic stable category $\mathcal{SH}(S)$ and about the categories of modules $\text{Mod}_A(S)$ over a commutative algebra $A \in \mathcal{SH}(S)_{\geq 0}$. In particular, we review how the t -structure that $\text{Mod}_A(S)$ inherits from $\mathcal{SH}(S)$ has the above two properties when S is a Noetherian scheme of finite Krull dimension.

In order to work with an abstract symmetric monoidal ∞ -category \mathcal{M} , we have chosen to axiomatize the elements of $\pi_0(\mathbb{S}) \simeq \mathbb{Z}$ in terms of maps $L \rightarrow \mathbf{1}$ where L is a \wedge -invertible object of \mathcal{M} such that $L \wedge -$ respects both $\mathcal{M}_{\geq 0}$ and $\mathcal{M}_{\leq 0}$; objects satisfying this property are called *tif* objects (*tif* stands for “tensor invertible and flat”).

Another choice we have made is to work with localizations at homotopy commutative algebras of \mathcal{M} , i.e., with a commutative algebra of the homotopy category $h\mathcal{M}$. We refer to Section 2.1.4 for a clearer definition.

In this framework, the main assumptions on E is essentially the following. E is a homotopy commutative algebra of $\mathcal{M}_{\geq 0}$. Furthermore, there exist a finite set of *tif* objects $\{L_i\}_{i=1}^r$ and maps $f_i : L_i \rightarrow \mathbf{1}$ such that the unit $\mathbf{1} \rightarrow E$ induces an isomorphism $\tau_0(\mathbf{1})/(f_1, \dots, f_r) \simeq \tau_0(E)$. For a more precise statement of this technical assumption we direct the reader to Assumption 4.2.1.

The main results we obtain in this general framework are then condensed in the following.

THEOREM 1.2.1 (Theorem 4.3.7 and Proposition 3.2.15). *Let E be a homotopy commutative algebra in \mathcal{M} satisfying Assumption 4.2.1 in the special case of $J = \emptyset$. Then for every bounded-below object X in \mathcal{M} there is a canonical equivalence $X_E \simeq X_{f_1}^{\wedge} \cdots^{\wedge} X_{f_r}^{\wedge}$ compatible with the localization map $\lambda_E(X) : X \rightarrow X_E$ and the formal completion map $\chi_{\underline{f}}(X) : X \rightarrow X_{f_1}^{\wedge} \cdots^{\wedge} X_{f_r}^{\wedge}$.*

The proof combines two main steps, where one compares formal completions and homology localization with some other intermediate object. This involves a sort of axiomatization of Moore spectra which is performed in Section 3. In the context of

Theorem 1.2.1, the relevant Moore object is $M = C(f_1) \wedge \cdots \wedge C(f_r)$, where $C(f_i)$ denotes the cofiber of f_i . The construction of M is what dictates the rather strong assumptions we have on $\tau_0(E)$. With this notation the first main step consists in proving the well-known statement that $X_M \simeq X_{f_1} \wedge \cdots \wedge X_{f_r}$ for every $X \in \mathcal{M}$. The second step, performed in Section 4, consists in showing that when X is bounded below, we have $X_M \simeq X_E$. This is done with a careful use of Bousfield classes, and uses crucially the left-completeness of the t -structure.

Section 5 contains a list of relevant examples and applications in the motivic setting. We mention here that, as an application, we partially recover a conservativity result for motives of Bachmann (cf. [3]). For this we work with $\mathcal{M} = \mathcal{SH}(K)[\frac{1}{p}]$ where K is a field and p is the exponential characteristic of K . We have thus functors $M : \mathcal{SH}(K)[\frac{1}{p}] \rightarrow \mathcal{DM}(K)[\frac{1}{p}]$ and $\tilde{M} : \mathcal{SH}(K)[\frac{1}{p}] \rightarrow \widetilde{\mathcal{DM}}(K)[\frac{1}{p}]$ associating with every spectrum its motive $M(X)$ and its Chow–Witt motive $\tilde{M}(X)$.

COROLLARY 1.2.2 (Corollary 5.4.1). *Let K be a perfect field of exponential characteristic $p \neq 2$, and let X be a bounded-below spectrum.*

- (1) *Assume that -1 is a sum of squares in K . If $M(X[\frac{1}{2}]) = 0$, then $X[\frac{1}{2}] = 0$.*
- (2) *Assume that K has finite étale 2-cohomological dimension and that X is strongly dualizable. If $M(X) = 0$, then $X = 0$.*
- (3) *Assume that k is infinite. If $\tilde{M}(X) = 0$, then $X = 0$.*

Actually analogous results hold for categories of motives that arise as categories of modules over a homotopy commutative algebra E in $\mathcal{SH}(K)$ whose $\tau_0(E)$ is Milnor (or Milnor–Witt) K -theory. We direct the reader to Remark 5.4.2 for a precise description of the relation with the work of Bachmann.

Section 6 contains a construction of E -nilpotent completions using an axiomatized version of the Adams tower, and a construction of the associated spectral sequence. The Adams tower is an alternative to the tower of partial totalizations mentioned above, and allows to avoid the assumption that E be a commutative algebra of \mathcal{M} .

Section 7 contains the axiomatization of nilpotent resolutions and a general proof of some of their properties. Using these, we can provide a universal property for the Adams tower as a pro-object. Comparing the pro-objects associated with the Adams tower and with the formal completion, we obtain the following result.

THEOREM 1.2.3 (Theorems 7.3.5 and 7.3.9). *Let E be a homotopy commutative algebra in \mathcal{M} satisfying Assumption 4.2.1 in the special case where either $J = \emptyset$ or $I = \emptyset$. Then for every bounded-below object X in \mathcal{M} the natural map $X_E \rightarrow X_E^\wedge$ is an equivalence in \mathcal{M} .*

This result has already appeared in [16], although in a very special case, and with a different proof. It was the reading of this specific work that stimulated our interest in the topic. Our approach to the problem is in fact very different in spirit from that of [16].

In their recent work [5], Bachmann and Østvær have generalized and streamlined the arguments of a previous version of our results which appeared in [24]. Although our results of [24] are phrased for the motivic stable homotopy category $h\mathcal{SH}(K)$ of a perfect field K , the structure and the arguments of the present paper are essentially the same as those of [24]. As a consequence the present paper and the second section of [5] present similar results with similar techniques, although [5] has a more direct and a simpler approach. The present paper was written, independently of [5], during the spring of 2020. I am grateful to T. Bachmann and P. A. Østvær for allowing me to publish the present work despite the overlap with theirs.

2. Preliminaries

We begin this section by introducing the categorical framework that we will be working with in order to fix some ideas and notation. This will happen in Section 2.1. In Section 2.2, we review some well-known facts about motivic stable categories, and in Section 2.3 we review Morel's homotopy t -structure, which is by far the most important tool we need. In Section 2.4, we introduce E -homology localizations associated to an object E , and recall some formal properties of these constructions. We conclude with a review of the formalism of Bousfield classes in Section 2.5, which turns out to be very useful for keeping track of the mutual relations between various localization functors appearing at the same time.

2.1 – Categorical framework

2.1.1. Along this paper we will be working with a presentably symmetric monoidal stable ∞ -category \mathcal{M} . We will denote the monoidal product by $- \wedge -$ and by $\mathbf{1}$ the unit. We also introduce the symbol $\text{Map}(-, -)$ to denote the mapping space between two objects. Finally, we denote by $h\mathcal{M}$ the homotopy category of \mathcal{M} , and by $[-, -]$ the Hom groups of the homotopy category, so that for every non-negative integer k and every pair of objects M and N in \mathcal{M} we have

$$\pi_k \text{Map}(M, N) \simeq \pi_0 \Omega^k \text{Map}(M, N) \simeq \pi_0 \text{Map}(\Sigma^k M, N) \simeq [\Sigma^k M, N].$$

We will assume that \mathcal{M} is endowed with a t -structure: the objects of $\mathcal{M}_{\geq n}$ will be referred to as $(n - 1)$ -connected. We will denote by $P^n(-)$ (resp. $P_n(-)$, resp. $P_n^n(-)$)

the $\leq n$ (resp. $\geq n$, resp. $= n$) truncation functors. For every $n \in \mathbb{Z}$ we then have a fiber sequence

$$P_n(X) \xrightarrow{\delta_n} X \xrightarrow{\pi_{n-1}} P^{n-1}(X).$$

The heart

$$\mathcal{M}^\heartsuit := \mathcal{M}_{\geq 0} \cap \mathcal{M}_{\leq 0}$$

is the nerve of an abelian category (see [22, Remark 1.2.1.12]), and for every object X of \mathcal{M} , $\tau_k(X) = \Sigma^{-k} P_k^k(X)$ will be referred to as the k -th homotopy object of X . Most of this paper works under the following list of assumptions on the t -structure on \mathcal{M} :

- (1) the t -structure is left-complete, i.e., $X \simeq \lim_n P^n(X)$ for all $X \in \mathcal{M}$;
- (2) the t -structure is multiplicative, i.e., $\mathbf{1} \in \mathcal{M}_{\geq 0}$ and $\mathcal{M}_{\geq p} \wedge \mathcal{M}_{\geq q} \subseteq \mathcal{M}_{\geq p+q}$;

Some sections actually work with less assumptions: for this we direct the reader to the introduction of each section and to the specific statements. The only additional assumption that appears on the t -structure of \mathcal{M} is the following:

- (3) the t -structure is compatible with filtered colimits, i.e., the truncation functors P^k commute with filtered colimits.

This last assumption is needed, in our opinion, to ensure that inverting homotopy elements is a t -exact functor (cf. Corollary 3.4.6). This assumption is used only in Corollary 4.3.6 when $J \neq \emptyset$. In any case, this assumption is not needed for the main theorems of the paper.

LEMMA 2.1.2. *If the t -structure on \mathcal{M} is multiplicative, the monoidal product $- \otimes^\heartsuit -$ induced on \mathcal{M}^\heartsuit is right exact.*

PROOF. Let

$$F \xrightarrow{f} G \xrightarrow{g} H \rightarrow 0$$

be an exact sequence of objects of \mathcal{M}^\heartsuit . Let $C := \text{cofib}(f)$, so that $H \simeq P^0(\text{cofib}(f))$. Thus given any $D \in \mathcal{M}^\heartsuit$, we have an induced fiber sequence

$$D \wedge F \rightarrow D \wedge G \rightarrow D \wedge C,$$

and we only need to check that the natural map

$$P^0(D \wedge C) \rightarrow P^0(D \wedge P^0(C))$$

is an equivalence. This follows from the fact that $D \wedge P_1(C) \in \mathcal{M}_{\geq 1}$. ■

2.1.3. In this situation, $\mathcal{M}_{\geq 0}$ has a natural structure of presentably symmetric monoidal ∞ -category induced via restriction along $\mathcal{M}_{\geq 0} \subseteq \mathcal{M}$. The inclusion $\mathcal{M}_{\geq 0} \subseteq \mathcal{M}$ is a symmetric monoidal functor, while its left adjoint $P_0 : \mathcal{M} \rightarrow \mathcal{M}_{\geq 0}$ is lax symmetric monoidal, and hence preserves algebra and module categories. All the claims follow at once from [22, Proposition 2.2.1.1].

The category $(\mathcal{M}_{\geq 0})_{\leq n}$ inherits a symmetric monoidal structure via the inclusion functor in $\mathcal{M}_{\geq 0}$ which, on its turn, inherits the structure of a lax symmetric monoidal functor. The left adjoint $P^n : \mathcal{M}_{\geq 0} \rightarrow (\mathcal{M}_{\geq 0})_{\leq n}$ of the inclusion inherits the structure of a symmetric monoidal functor, while the projection maps π_n inherit the structure of monoidal natural transformation. All the claims follow at once from [22, Proposition 2.2.1.8, Proposition 2.2.1.9, Example 2.2.1.10].

In particular, the inclusion

$$\mathcal{M}^\heartsuit \subseteq \mathcal{M}_{\geq 0}$$

induces on the heart of the t -structure the structure of a symmetric monoidal ∞ -category with monoidal product

$$- \otimes^\heartsuit - \simeq \tau_0(- \wedge -)$$

and monoidal unit

$$\mathbf{1}^\heartsuit \simeq P^0(\mathbf{1}) \simeq \tau_0(\mathbf{1}).$$

2.1.4. We will also use the notion of *homotopy commutative algebra* in \mathcal{M} . With this expression we mean an object E of \mathcal{M} together with maps $e : \mathbf{1} \rightarrow E$ and $\mu : E \wedge E \rightarrow E$ and suitable homotopies, making (E, e, μ) a commutative monoid in the homotopy category $h\mathcal{M}$. We will similarly use the notion of homotopy E -module, defined in an analogous way.

2.1.5. If E is a homotopy commutative algebra in \mathcal{M} and X is a homotopy E -module, then $\tau_0(E)$ has the structure of a commutative monoid in \mathcal{M}^\heartsuit and each object $\tau_k(X)$ inherits the structure of a $\tau_0(E)$ -module.

2.2 – Motivic stable categories

2.2.1. A *base scheme* is a Noetherian scheme of finite Krull dimension. Given a base scheme S , we denote by $\mathcal{SH}(S)$ the Morel–Voevodsky \mathbb{P}^1 -stable ∞ -category. We recall that $\mathcal{SH}(S)$ is a presentably symmetric monoidal stable ∞ -category; its monoidal product is denoted by \wedge and the unit, the motivic sphere spectrum, is denoted by \mathbb{S} . We redirect the reader to Appendix A.2 for a reference to the previous claim and a quick review of the higher categorical terminology.

The ∞ -category $\mathcal{SH}(S)$ is actually compactly generated by the set

$$\{\Sigma^{p+q\alpha} \Sigma^\infty X_+ : X \in \mathbf{Sm}_S, p, q \in \mathbb{Z}\},$$

where $\Sigma^{p+q\alpha}$ is defined as $\Sigma_{S^1}^p \Sigma_{\mathbb{G}_m}^q$. Here \mathbf{Sm}_S denotes the category of smooth schemes of finite type over S with S -morphisms as arrows.

2.2.2. To any commutative algebra A of $\mathcal{SH}(S)$ (cf. Definition A.2.7) we associate a category $\mathcal{M}od_A(S)$ whose objects are called A -module spectra, or simply A -modules. $\mathcal{M}od_A(S)$ inherits from $\mathcal{SH}(S)$ the property of being a presentably symmetric monoidal stable ∞ -category. Once again references and definitions are postponed to Section A.2.8. The monoidal product is denoted by $- \wedge_A -$, or simply by $- \wedge -$ when no confusion arises; the monoidal unit is denoted by $\mathbf{1}_A$. Similarly $\text{Map}_A(-, -)$ will denote the mapping space of the ∞ -category $\mathcal{M}od_A(S)$.

In addition, we have a free-forget adjunction

$$F_A : \mathcal{SH}(S) \rightleftarrows \mathcal{M}od_A(S) : U_A.$$

The forgetful functor U_A is right adjoint of F_A : it commutes with all small limits and colimits, and it is conservative. The functor F_A is symmetric monoidal and commutes with all small colimits. Since U_A is conservative and commutes with colimits, the category $\mathcal{M}od_A(S)$ is compactly generated by the set

$$\{F_A(\Sigma^{p+q\alpha} \Sigma^\infty X_+) : X \in \mathbf{Sm}_S, p, q \in \mathbb{Z}\}.$$

We conclude by observing that the composition $U_A \circ F_A \simeq A \wedge -$, and thus the monoidal unit $\mathbf{1}_A \in \mathcal{M}od_A(S)$ is mapped to A in $\mathcal{SH}(S)$. We will abuse the language and confuse $\mathbf{1}_A$ and A . We give a reference for all these facts in Section A.2.8.

2.3 – Homotopy t -structure

2.3.1. We define $\mathcal{M}od_A(S)_{\geq n}$ as the smallest full sub- ∞ -category of $\mathcal{M}od_A(S)$ closed under small colimits and extensions that contains the collection

$$\{F_A(\Sigma^{p+q\alpha} \Sigma^\infty X_+) : X \in \mathbf{Sm}_S, p \geq n, q \in \mathbb{Z}\}.$$

Furthermore, we denote by $\mathcal{M}od_A(S)_{\leq n}$ the full sub- ∞ -category spanned by those A -modules Z such that $\text{Map}_A(X, Z) \simeq *$ for every $X \in \mathcal{M}od_A(S)_{\geq n+1}$.

By [22, Proposition 1.4.4.11], the pair of subcategories

$$(\mathcal{M}od_A(S)_{\geq 0}, \mathcal{M}od_A(S)_{\leq -1})$$

defines an accessible t -structure on $\mathcal{M}od_A(S)$ (cf. [22, Definition 1.4.4.12]), called *homotopy t -structure*. It follows that for every $n \in \mathbb{Z}$ we have a cofiber sequence in $\mathcal{M}od_A(S)$,

$$(2.1) \quad P_n(X) \xrightarrow{\delta_n} X \xrightarrow{\pi_{n-1}} P^{n-1}(X),$$

which is functorial in the A -module X , with $P_n(X) \in \mathcal{M}od_A(S)_{\geq n}$ and $P^{n-1}(X) \in \mathcal{M}od_A(S)_{\leq n-1}$. It is clear from the choice of the generators that the homotopy t -structure on $\mathcal{M}od_A(S)$ is multiplicative.

THEOREM 2.3.2. *Let S be a Noetherian scheme of finite Krull dimension and A be a commutative algebra in $\mathcal{S}\mathcal{H}(S)_{\geq 0}$. The homotopy t -structure on $\mathcal{M}od_A(S)$ is left-complete, right-complete and compatible with filtered colimits.*

PROOF. For the left-completeness we need to show that $\lim_n P_n(X) \simeq 0$, and for the right-completeness that $\operatorname{colim}_n P^n(X) \simeq 0$. The forgetful functor U_A commutes with small limits and colimits (cf. Appendix A.2.9). Moreover, since $A \in \mathcal{S}\mathcal{H}(S)_{\geq 0}$, it is easy to see that an A -module Y is in $\mathcal{M}od_A(S)_{\geq 0}$ (resp. $\mathcal{M}od_A(S)_{\leq 0}$) if and only if $U_A(Y)$ is in $\mathcal{S}\mathcal{H}(S)_{\geq 0}$ (resp. $\mathcal{S}\mathcal{H}(S)_{\leq 0}$). As a consequence U_A commutes with the Postnikov truncations of $\mathcal{M}od_A(S)$ and of $\mathcal{S}\mathcal{H}(S)$. We are thus reduced to the case of $A \simeq \mathbb{S}$. Left-completeness then follows from [35, Corollary 3.8], while right completeness follows from [35, Remark 1.29]. The subcategories $\mathcal{M}od_A(S)_{\leq n}$ are closed under filtered colimits, since the generators of $\mathcal{M}od_A(S)_{\geq n+1}$ are compact. In particular, the truncation functors P^n commute with filtered colimits. ■

COROLLARY 2.3.3. *Let S be a Noetherian scheme of finite Krull dimension, and A be a commutative algebra in $\mathcal{S}\mathcal{H}(S)_{\geq 0}$. Then the ∞ -category $\mathcal{M}od_A(S)$ is a presentably symmetric monoidal stable ∞ -category and it is compactly generated. The homotopy t -structure is accessible, left-complete, right-complete, multiplicative and compatible with filtered colimits.*

2.3.4. When K is a perfect field of characteristic not 2, Morel [25, Section 6.1] constructs a map

$$\sigma : K_*^{MW}(K) \rightarrow [\mathbb{S}, \mathbb{G}_m^{\wedge *}]_{\mathbb{S}}$$

by defining it on the generators and checking that it passes to the quotient through the defining relations of $K_*^{MW}(K)$ (cf. [26, p. 49, Definition 3.1] for precise formulas).

In [25, Theorem 6.2.1], Morel shows that σ is actually an isomorphism, and in [25, Corollary 6.4.1] he shows that σ extends uniquely to an isomorphism of homotopy modules $\sigma : \mathcal{K}_*^{MW} \xrightarrow{\cong} \underline{\pi}_0 \mathbb{S}$.

2.4 – Homology localizations

DEFINITION 2.4.1. Let \mathcal{M} be a presentably symmetric monoidal stable ∞ -category and let E be an object of \mathcal{M} . We say that an arrow $f : X \rightarrow Y$ in \mathcal{M} is an E -homology equivalence (or, shortly, an E -equivalence) if the induced map $f \wedge \operatorname{id} : X \wedge E \rightarrow Y \wedge E$

is an equivalence in \mathcal{M} . We say that an object C is *E-acyclic* if $E \wedge C \simeq 0$ in \mathcal{M} . Finally, we say that an object $Z \in \mathcal{M}$ is *E-local* if for every *E*-homology equivalence $X \rightarrow Y$, the induced map on mapping spaces $\text{Map}(Y, Z) \rightarrow \text{Map}(X, Z)$ is an equivalence of spaces.

REMARK 2.4.2. One sees immediately that the full sub- ∞ -category $\text{Ac}(E)$ spanned by *E*-acyclic objects is closed under arbitrary (small) colimits and retracts, and that *E*-acyclic objects have the 2-out-of-3 property in fiber sequences in \mathcal{M} . More precisely if K is any simplicial set and $p : K \rightarrow \text{Ac}(E)$ is a diagram whose composition with the inclusion $\text{Ac}(E) \subseteq \mathcal{M}$ extends to a colimit diagram $\bar{p} : K^\triangleright \rightarrow \mathcal{M}$, then there exists a unique lift $\bar{p} : K^\triangleright \rightarrow \text{Ac}(E)$ and such lift is again a colimit diagram. Furthermore, since the inclusion functor $\text{Ac}(E) \subseteq \mathcal{M}$ has a right adjoint (cf. Proposition 2.4.5), then whenever $\bar{p} : K^\triangleright \rightarrow \text{Ac}(E)$ is a colimit diagram, its composition $K^\triangleright \rightarrow \mathcal{M}$ is again a colimit diagram; the converse to this statement is instead implied by the previous observation. Note that $\text{Ac}(E)$ is also closed under smashing with an arbitrary object.

REMARK 2.4.3. Similarly, it is immediate to see that the full sub- ∞ -category $\text{Loc}(E)$ spanned by *E*-local objects is closed under arbitrary (small) limits and retracts, and that *E*-local objects have the 2-out-of-3 property in fiber sequences. More precisely if K is any simplicial set and $p : K \rightarrow \text{Loc}(E)$ is a diagram whose composition with the inclusion $\text{Loc}(E) \subseteq \mathcal{M}$ extends to a limit diagram $\bar{p} : K^\triangleleft \rightarrow \mathcal{M}$, then there exists a unique lift of $\bar{p} : K^\triangleleft \rightarrow \text{Loc}(E)$ and such lift is again a limit diagram. We finally note that an object Z is *E*-local if and only if for every *E*-acyclic object C the space $\text{Map}(C, Z)$ is contractible, and this happens if and only if for every *E*-acyclic object C , the group $[C, Z]$ is zero.

REMARK 2.4.4. Note that we could define, a priori, a more general notion of equivalence: we call a map $M \rightarrow N$ an *E-local equivalence* if for every *E*-local object X the natural map $\text{Map}(N, X) \rightarrow \text{Map}(M, X)$ is an equivalence of spaces. Clearly all *E*-equivalences are *E*-local equivalences. The reverse holds if and only if the class $\text{Ac}(E)$ of *E*-acyclic objects coincides with its double orthogonal

$${}^\perp(\text{Ac}(E)^\perp) := \{X \in \mathcal{M} : \forall C \in \text{Loc}(E), \text{Map}(X, C) \simeq *\}.$$

This follows immediately from the existence of a left adjoint to the inclusion $\text{Loc}(E) \subseteq \mathcal{M}$.

PROPOSITION 2.4.5. *Let \mathcal{M} be a presentably symmetric monoidal stable ∞ -category and let E be an object of \mathcal{M} . Then:*

- (ℓ1) *The inclusion $i_{\text{Loc}} : \text{Loc}(E) \subseteq \mathcal{M}$ has a left adjoint $L_E : \mathcal{M} \rightarrow \text{Loc}(E)$.*
- (ℓ2) *For every X in \mathcal{M} there is an *E*-equivalence $X \rightarrow X'$ with target $X' \in \text{Loc}(E)$.*

($\ell 3$) The ∞ -category $\text{Loc}(E)$ is presentable.

($\ell 4$) A map $f : X \rightarrow Y$ in \mathcal{M} is an E -equivalence if and only if $L_E(f)$ is an equivalence.

Moreover:

(a1) The inclusion $i_{\text{Ac}} : \text{Ac}(E) \subseteq \mathcal{M}$ has a right adjoint $A_E : \mathcal{M} \rightarrow \text{Ac}(E)$.

(a2) For every X in \mathcal{M} there is a co-local equivalence $X'' \rightarrow X$ with source $X'' \in \text{Ac}(E)$.

(a3) The ∞ -category $\text{Ac}(E)$ is presentable.

PROOF. Let us concentrate on E -local objects first. The class of E -equivalences in \mathcal{M} is a strongly saturated class of morphisms according to [21, Definition 5.5.4.5]. Moreover, the collection of E -equivalences is a strongly saturated class of small generation: this can be seen combining [21, Proposition 5.5.4.16 and Remark 5.5.4.6], given that the functor $- \wedge E$ is presentable. Properties ($\ell 1$)–($\ell 4$) follow immediately from [21, Proposition 5.5.4.15].

Let us now turn to E -acyclic objects. Applying [21, Proposition 5.2.7.8] to \mathcal{M}^{op} , we get that (a2) is equivalent to (a1). In order to prove (a1), we take for every $X \in \mathcal{M}$ an E -equivalence to an E -local object $\lambda : X \rightarrow X'$ and define $X'' \rightarrow X$ as the fiber of λ . Then clearly X'' is E -acyclic and for every E -acyclic object C composing with λ induces an equivalence $\text{Map}(C, X'') \rightarrow \text{Map}(C, X)$, given that $\text{Map}(C, X')$ is contractible by assumption. In order to prove (a3), we use [22, Proposition 1.4.4.13]. ■

DEFINITION 2.4.6. A choice of composition $(-)_E := L_E \circ i_{\text{Loc}}$ is called E -localization functor, while a unit transformation $\lambda_E : \text{id} \rightarrow (-)_E$ is called E -localization map. Similarly, a choice of composition ${}_E(-) : i_{\text{Ac}} \circ A_E$ is called E -acyclicization functor and a co-unit transformation ${}_E(-) \rightarrow \text{id}$ is called E -acyclicization map.

PROPOSITION 2.4.7 (cf. [22, Proposition 2.2.1.9]). The functor L_E is symmetric monoidal and the natural transformation $\text{id} \rightarrow \iota_E \circ L_E$ is monoidal.

2.5 – Bousfield classes

2.5.1. Let \mathcal{M} be a presentably symmetric monoidal stable ∞ -category. We introduce an equivalence relation on the class of equivalence classes of objects of \mathcal{M} following [7]. We set $E \sim_B F$ if, for every $X \in \mathcal{M}$, we have that $E \wedge X = 0$ if and only if $F \wedge X = 0$. By Proposition 2.4.5, localization functors at E and F exist and they are equivalent exactly when $E \sim_B F$. We denote by $\mathcal{A}(\mathcal{M})$ the class of Bousfield classes in \mathcal{M} and by $\langle E \rangle$ the element in $\mathcal{A}(\mathcal{M})$ represented by an object E . On $\mathcal{A}(\mathcal{M})$ we introduce a partial ordering by setting $\langle E \rangle \leq \langle F \rangle$ if every F -acyclic object is E -acyclic.

2.5.2. Given a possibly infinite collection of Bousfield classes $\langle E_i \rangle_{i \in I}$ we have a *join* operation which is defined as $\bigoplus_{i \in I} \langle E_i \rangle := \langle \bigoplus_{i \in I} E_i \rangle$. We note that the join is always the minimal upper bound of its summands.

Similarly, given a finite collection of Bousfield classes $\langle E_i \rangle_{i \in I}$ we have a *meet* operation which is defined as $\bigwedge_{i \in I} \langle E_i \rangle := \langle \bigwedge_{i \in I} E_i \rangle$. We note that the meet operation is a lower bound for its factors, but in general does not need to be the maximal lower bound.

2.5.3. Following [7], we denote by $\mathcal{DL}(\mathcal{M})$ the subclass of $\mathcal{A}(\mathcal{M})$ of those Bousfield classes $\langle E \rangle$ such that $\langle E \rangle \wedge \langle E \rangle = \langle E \rangle$. The operations of meet and join restrict to $\mathcal{DL}(\mathcal{M})$ and it is elementary to check that $\mathcal{DL}(\mathcal{M})$ satisfies the axioms of a distributive lattice. The partial ordering \leq on $\mathcal{A}(\mathcal{M})$ restricts to a partial ordering on $\mathcal{DL}(\mathcal{M})$. We wish to observe that for given $\langle E \rangle, \langle F \rangle \in \mathcal{DL}(\mathcal{M})$ their meet $\langle E \wedge F \rangle$ is actually their maximal lower bound. Most of the objects we will consider later actually belong to this subclass: for instance, every homotopy algebra E belongs to $\mathcal{DL}(\mathcal{M})$, since E is a retract of $E \wedge E$.

2.5.4. We say that a Bousfield class $\langle E \rangle \in \mathcal{A}(\mathcal{M})$ has a complement if there is another Bousfield class $\langle F \rangle$ such that $\langle E \rangle \wedge \langle F \rangle = \langle 0 \rangle$ and $\langle E \rangle \oplus \langle F \rangle = \langle \mathbf{1} \rangle$. If $\langle E \rangle$ has a complement, then such complement is unique, and we denote it by $\langle E \rangle^c$. Moreover, when $\langle E \rangle$ has a complement, $\langle E \rangle \in \mathcal{DL}(\mathcal{M})$. We denote by $\mathcal{BA}(\mathcal{M})$ the sub-lattice of $\mathcal{DL}(\mathcal{M})$ of those Bousfield classes admitting a complement. Lemma 2.7 of [7] shows that the inclusion $\mathcal{BA}(\mathcal{M}) \subseteq \mathcal{DL}(\mathcal{M})$ is in general strict. Assume that both $\langle E \rangle, \langle F \rangle$ have complements, then the following equalities are satisfied:

$$(2.2) \quad \begin{cases} \langle E \rangle^{cc} = \langle E \rangle, \\ \langle E \oplus F \rangle^c = \langle E \rangle^c \wedge \langle F \rangle^c, \\ \langle E \wedge F \rangle^c = \langle E \rangle^c \oplus \langle F \rangle^c. \end{cases}$$

3. Moore objects

In the topological stable category \mathcal{SH} , one can associate to every abelian group A a Moore spectrum $\mathbb{S}A$. Up to equivalence $\mathbb{S}A$ is characterized by the following properties: $\mathbb{S}A \in \mathcal{SH}_{\geq 0}$, $\pi_0(\mathbb{S}A) \simeq A$, and $\mathbb{H}\mathbb{Z} \wedge \mathbb{S}A \simeq \mathbb{H}A$. Moore spectra are fundamental in the study of Bousfield classes for two reasons. One is that $\langle \mathbb{S}A \rangle$ depends only on torsion and divisibility properties of A (cf. [8, Proposition 2.3]). The other reason is that for a spectrum $X \in \mathcal{SH}_{\geq k}$, and a homotopy commutative ring spectrum $E \in \mathcal{SH}_{\geq 0}$ we have $X_E \simeq X_{\mathbb{S}\pi_0 E}$. Given a set J , a finite set I , and collections of maps $\{f_i : L_i \rightarrow \mathbf{1}\}_{i \in I}$,

$\mathcal{J} := \{g_j : L_j \rightarrow \mathbf{1}\}_{j \in J}$, we introduce a weak version of Moore object

$$C(f_1) \wedge \cdots \wedge C(f_r) \wedge \mathbf{1}[\mathcal{J}^{-1}],$$

that in general depends on the choice of the f_i and g_j . We start the section with reviewing some technical tools for dealing with towers in Section 3.1. We then treat the f_i and g_j separately in Sections 3.2 and 3.4; in Section 3.3 we make an example on η -completions in $\mathcal{SH}(S)$.

In this section, \mathcal{M} is a presentably symmetric monoidal stable ∞ -category, endowed with a left-complete multiplicative t -structure.

3.1 – Towers

Let K^n be the sub-simplicial set

$$\Delta^{\{0,1\}} \coprod_{\Delta^0} \Delta^{\{1,2\}} \cdots \Delta^{\{n-2,n-1\}} \coprod_{\Delta^0} \Delta^{\{n-1,n\}} \subseteq \Delta^n,$$

where the pushouts are taken with respect to the maps

$$\Delta^{\{k-1,k\}} \xleftarrow{k} \Delta^0 \xrightarrow{k} \Delta^{\{k,k+1\}}.$$

In other words, K^n is the sub-simplicial set of Δ^n generated by its non-degenerate 1-simplices. We also define $K = \bigcup_n K^n$. A diagram in \mathcal{M} indexed by $K^{n\text{op}}$ (resp. K^{op}) is thus a collection of n composable arrows of \mathcal{M} :

$$X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_0$$

(resp. a countable collection of composable arrows $\cdots \rightarrow X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_0$). A composition of n composable arrows $p : K^{n\text{op}} \rightarrow \mathcal{M}$ is an extension of p to $\Delta^{n\text{op}}$, and is essentially unique in the following sense.

- LEMMA 3.1.1. (1) *The inclusion $i_n : K^n \subseteq \Delta^n$ is an inner anodyne map.*
 (2) *If \mathcal{C} is an ∞ -category, the natural restriction map*

$$i_n^* : \text{Fun}(\Delta^{n\text{op}}, \mathcal{C}) \rightarrow \text{Fun}(K^{n\text{op}}, \mathcal{C})$$

is a trivial Kan fibration between ∞ -categories, and in particular a categorical equivalence.

PROOF. For (1) observe that when $n = 0, 1$, there is nothing to do, while for $n = 2$ we have just rewritten [21, Corollary 2.3.2.2]. Assume now that the inclusion $i_n : K^n \subseteq \Delta^n$

is inner anodyne. The inclusion i_{n+1} is the composition

$$K^{n+1} = K^n \coprod_{\Delta^{\{n\}}} \Delta^{\{n,n+1\}} \subseteq \Delta^n \coprod_{\Delta^{\{n\}}} \Delta^{\{n,n+1\}} \subseteq \Delta^{n+1},$$

and we claim that both of the above inclusions are inner anodyne. Indeed, the central inclusion is $i_n \coprod_{\Delta^{\{n\}}} \Delta^{\{n,n+1\}}$, and this is inner anodyne because the collection of inner anodyne maps is closed under push-outs (being a weakly saturated collection of maps). The rightmost inclusion is identified with the natural inclusion

$$\Delta^{n-1} * \Delta^0 \coprod_{\emptyset * \Delta^0} \emptyset * \Delta^1 \subseteq \Delta^{n-1} * \Delta^1,$$

which is inner anodyne after [21, Lemma 2.1.2.3].

For (2) one combines [21, Propositions 2.3.2.1 and 1.2.7.3] with (1) and with the fact that a map is inner anodyne if and only if the opposite is. ■

LEMMA 3.1.2. *Let \mathbb{N} be the poset of natural numbers. Then the natural inclusion*

$$i = \bigcup_n i_n : K = \bigcup_n K^n \xrightarrow{i_n} \bigcup_n \Delta^n = \mathbb{N}(\mathbb{N})$$

is inner anodyne. In particular, for every ∞ -category \mathcal{C} we have a trivial Kan fibration of ∞ -categories

$$\mathrm{Map}(\mathbb{N}(\mathbb{N})^{\mathrm{op}}, \mathcal{C}) \xrightarrow{i^*} \mathrm{Map}(K^{\mathrm{op}}, \mathcal{C}).$$

PROOF. The map i can be realized as well as the transfinite composition of the natural inclusions

$$j_n : \Delta^n \coprod_{K^n} K \subseteq \Delta^{n+1} \coprod_{K^{n+1}} K,$$

which on their turn are push-outs along $K^{n+1} \subseteq K$ of the inclusions

$$\Delta^n \coprod_{\Delta^{\{n\}}} \Delta^{\{n,n+1\}} \simeq \Delta^n \coprod_{K^n} K^{n+1} \subseteq \Delta^{n+1}$$

appearing in the proof of Lemma 3.1.1. Each j_n is thus inner anodyne, and so is their transfinite composition. The last part of the statement follows directly by [21, Proposition 2.3.2.5]. ■

3.1.3. As a consequence, if we have a countable family of composable arrows $f_n : X_n \rightarrow X_{n-1}$, we can essentially uniquely extend this collection to a diagram $X_\bullet : \mathbb{N}(\mathbb{N}^{\mathrm{op}}) \rightarrow \mathcal{M}$. If $\mathbf{c}X_0$ is the constant diagram on X_0 , we can thus upgrade the datum of the f_n to a map

of towers $f^\bullet : X_\bullet \rightarrow \mathbf{c}X_0$ which is a 1-simplex of $\text{Fun}(\mathbb{N}(\mathbb{N}^{\text{op}}), \mathcal{M})$. Finally, taking the cofiber of f^\bullet yields a commutative ladder of fiber sequences:

$$\begin{array}{ccccc} X_n & \xrightarrow{f^n} & X_0 & \longrightarrow & C(f^n) \\ f_n \downarrow & & \parallel & & p_n \downarrow \\ X_{n-1} & \xrightarrow{f^{n-1}} & X_0 & \longrightarrow & C(f^{n-1}), \end{array}$$

where f^n denotes a composition of the f_i for $i \leq n$.

3.2 – Quotients

DEFINITION 3.2.1. Let $r \geq 1$ be an integer and for $i \in \{1, \dots, r\}$ let $f_i : L_i \rightarrow \mathbf{1}$ be a map in \mathcal{M} . We denote by $C(f_i)$ the cofiber of f_i and by $M(\underline{f})$ the iterated cofiber $C(f_1) \wedge \cdots \wedge C(f_r)$. We call $M(\underline{f})$ the Moore object associated with the collection f_1, \dots, f_r .

REMARK 3.2.2. If $\mathcal{M} = \mathcal{SH}$, $r = 1$, and $n : \mathbb{S} \rightarrow \mathbb{S}$ represents the multiplication by $n \in \mathbb{Z}$, then $C(n)$ is the usual modulo n Moore spectrum. Note that when $r \geq 2$, the spectrum $M(n_1, n_2) = C(n_1) \otimes C(n_2)$ is not always a Moore spectrum in the classical sense: take $n_1 = n_2 = 2$ for instance. However, for every pair of integers n_1, n_2 ,

$$\langle M(n_1, n_2) \rangle = \langle C(\text{g.c.d.}(n_1, n_2)) \rangle,$$

which follows from [7, Proposition 2.13]. This observation points out that our definition of Moore object M really depends on choices, and not only on its $\tau_0(M)$.

3.2.3. We introduce some further notation. If $f : L \rightarrow \mathbf{1}$ is a map in \mathcal{M} , we will denote by $l_f(X)$ the left multiplication by f on $X \in \mathcal{M}$, i.e., the composition

$$L \wedge X \xrightarrow{f \wedge X} \mathbf{1} \wedge X \xrightarrow{\cong} X.$$

Similarly, $r_f(X)$ will denote the right multiplication by f , given by the composition

$$X \wedge L \xrightarrow{X \wedge f} X \wedge \mathbf{1} \xrightarrow{\cong} X.$$

Note that left multiplication commutes with any map in $\phi : X \rightarrow Y$ in \mathcal{M} , in the sense that the square

$$\begin{array}{ccc} L \wedge X & \xrightarrow{l_f(X)} & X \\ L \wedge \phi \downarrow & & \downarrow \phi \\ L \wedge Y & \xrightarrow{l_f(Y)} & Y \end{array}$$

commutes; right multiplication behaves similarly.

DEFINITION 3.2.4. An object L of \mathcal{M} is called *flat* if the functor $L \wedge -$ respects $\mathcal{M}_{\geq 0}$ and $\mathcal{M}_{\leq 0}$. L is called *sdf* (resp. *tif*) if it is strongly dualizable (resp. \wedge -invertible) and flat.

3.2.5. In the motivic setting, the main example of a tif object in $\mathcal{SH}(S)$ is of course $\Sigma^{-\mathrm{rk}(V)}\mathrm{Th}(V) \in \mathcal{SH}(S)$ where V is a virtual vector bundle on the base scheme S , and where $\mathrm{rk}(V)$ is interpreted as a locally constant function on S . For instance \mathbb{G}_m is a tif object in $\mathcal{SH}(S)$.

REMARK 3.2.6. Let $f : L \rightarrow \mathbf{1}$ be a morphism of \mathcal{M} with tensor-invertible source. The map $f^{\wedge 3} : L^{\wedge 3} \rightarrow \mathbf{1}$ gives an object of $\mathcal{M}/_1$ and the cyclic permutation σ of the factors of $L^{\wedge 3}$ gives a loop in the space $\mathrm{Map}_{\mathcal{M}}(L^{\wedge 3}, \mathbf{1})$ based at the point $f^{\wedge 3}$. As noted in [10], this loop is trivial. Indeed the permutation action of the factors of $L^{\wedge 3}$ induces a map $B\Sigma_3 \rightarrow \mathrm{Map}_{\mathcal{M}}(L^{\wedge 3}, \mathbf{1})$ sending the base point of $B\Sigma_3$ to $f^{\wedge 3}$. The induced map on π_1 has thus σ in its image, but at the same time factors through the abelianization of Σ_3 , since \mathcal{M} is stable. As a consequence the object $f \in \mathcal{M}/_1$ is 3-symmetric in the sense of the discussion carried over in [14, Section 3].

LEMMA 3.2.7. Let $f : L \rightarrow \mathbf{1}$ be a map with flat source and let X be any object of \mathcal{M} . A canonical zig-zag of natural maps induces an equivalence $\tau_n(L \wedge X) \simeq L \wedge \tau_n(X)$. Under this equivalence the maps $\tau_n(l_f(X)) : \tau_n(L \wedge X) \rightarrow \tau_n(X)$ and $l_f(\tau_n(X)) : L \wedge \tau_n(X) \rightarrow \tau_n(X)$ are naturally identified. Similar statements hold for right multiplication.

PROOF. The lemma follows from an easy diagram chase using the fiber sequences

$$P_n(-) \rightarrow (-) \rightarrow P^{n-1}(-) \quad \text{and} \quad P_{n+1}(-) \rightarrow P_n(-) \rightarrow \Sigma^n \tau_n(-)$$

for X and $L \wedge X$. ■

LEMMA 3.2.8. For every $i \in \{1, \dots, r\}$ let $f_i : L_i \rightarrow \mathbf{1}$ be a map in \mathcal{M} , where L_i is a flat object, and let (f_1, \dots, f_r) be the sub- $\tau_0(\mathbf{1})$ -module of $\tau_0(\mathbf{1})$ obtained as the image of the map

$$\Sigma_i l_{f_i}(\tau_0(\mathbf{1})) : \bigoplus_i L_i \wedge \tau_0(\mathbf{1}) \rightarrow \tau_0(\mathbf{1})$$

in \mathcal{M}^\heartsuit . Then $M(\underline{f}) \in \mathcal{M}_{\geq 0}$,

$$\tau_0 M(\underline{f}) \simeq \otimes_i^\heartsuit \tau_0(C(f_i)) \simeq \tau_0(\mathbf{1})/(f_1, \dots, f_r),$$

and the canonical map $\mathbf{1} \rightarrow M(\underline{f})$ induces on τ_0 the quotient map

$$\tau_0(\mathbf{1}) \rightarrow \tau_0(\mathbf{1})/(f_1, \dots, f_r).$$

PROOF. An easy diagram chase allows to reduce to the following claim: If \mathbf{A} is an abelian category, M, N, P are objects of \mathbf{A} , and we have a diagram $f : N \rightarrow M \leftarrow P : g$, then $(M/N)/\overline{\text{Im}(g)} \simeq M/\text{Im}(f + g)$, where $f + g : N \oplus P \rightarrow M$ is the natural map induced by f and g . On its turn this claim is easily proved using that colimits commute with each other. \blacksquare

3.2.9. Let $f : L \rightarrow \mathbf{1}$ be a map in \mathcal{M} and consider the collection of composable maps $r_f(L^{\wedge n-1}) : L^{\wedge n} \rightarrow L^{\wedge(n-1)}$. By applying the argument of Section 3.1.3, we obtain a commutative ladder of fiber sequences

$$(3.1) \quad \begin{array}{ccccc} L^{\wedge n} & \xrightarrow{f^n} & \mathbf{1} & \longrightarrow & C(f^n) \\ r_f(L^{\wedge(n-1)}) \downarrow & & \parallel & & p_n \downarrow \\ L^{\wedge(n-1)} & \xrightarrow{f^{n-1}} & \mathbf{1} & \longrightarrow & C(f^{n-1}). \end{array}$$

Note that we have equivalences

$$\begin{aligned} \text{fib}(C(f^n) \xrightarrow{p_n} C(f^{n-1})) &\simeq \text{cofib}(L^{\wedge n} \xrightarrow{r_f(L^{\wedge(n-1)})} L^{\wedge(n-1)}) \\ &\simeq L^{\wedge(n-1)} \wedge C(f) \end{aligned}$$

yielding a fiber sequence

$$(3.2) \quad L^{\wedge(n-1)} \wedge C(f) \rightarrow C(f^n) \xrightarrow{p_n} C(f^{n-1}).$$

DEFINITION 3.2.10. Let $f : L \rightarrow \mathbf{1}$ be a map in \mathcal{M} . In view of Section 3.2.9, we define the f -adic completion of $X \in \mathcal{M}$ as the object

$$X_f^\wedge := \lim_{\mathbf{N}(\mathbf{N}^{\text{op}})} (X \wedge C(f^\bullet)).$$

The (essentially unique) map $\chi_f(X) : X \rightarrow X_f^\wedge$ induced by $X \simeq \lim(X \wedge \mathbf{1}) \rightarrow \lim(X \wedge C(f^\bullet))$ is called f -adic completion map. An object X is f -complete if $\chi_f(X)$ is an equivalence.

REMARK 3.2.11. Since the operations of taking inverse limits and of smashing with an object X commute with finite limits, the fiber sequences of towers introduced in Section 3.2.9 yields the fiber sequence

$$\lim(X \wedge L^{\wedge \bullet}) \rightarrow X \xrightarrow{\chi_f(X)} X_f^\wedge.$$

In particular, when X is k -connected and $L \in \mathcal{M}_A(S)_{\geq 1}$, X is f -complete.

DEFINITION 3.2.12. If $r \geq 1$, then for every $X \in \mathcal{M}$ the object

$$X_{\underline{f}}^{\wedge} := X_{f_1 f_2 \cdots f_r}^{\wedge} \in \mathcal{M}$$

is called \underline{f} -adic completion of X . The \underline{f} -adic completion map $\chi_{\underline{f}}(X)$ is defined as a composition of the natural maps

$$\chi_{f_r}(X_{f_1, \dots, f_{r-1}}^{\wedge}) \circ \cdots \circ \chi_{f_2}(X_{f_1}^{\wedge}) \circ \chi_{f_1}(X).$$

LEMMA 3.2.13. Let L be a \wedge -invertible object of \mathcal{M} and $f : L \rightarrow \mathbf{1}$ be a map in \mathcal{M} . Then the cofiber $C(f)$ is strongly dualizable in \mathcal{M} and

$$D(C(f)) \simeq \text{fib}(D(f)) \simeq \Sigma^{-1}D(L) \wedge C(f),$$

where $D(-) = \text{Hom}(-, \mathbf{1})$ and $\text{Hom}(-, -)$ denotes the right adjoint of $- \wedge -$.

PROOF. We have that

$$D(C(f)) \simeq \text{fib}(D(f)) \simeq \text{fib}(D(L) \wedge f) \simeq D(L) \wedge \text{fib}(f) \simeq D(L) \wedge \Sigma^{-1}C(f).$$

The fact that $\text{fib}(D(f)) \simeq \text{fib}(D(L) \wedge f)$ follows directly from the definition of dual map via the duality adjunction. \blacksquare

PROPOSITION 3.2.14. Let L be a \wedge -invertible object of \mathcal{M} and $f : L \rightarrow \mathbf{1}$ be a map in \mathcal{M} . Then for every object X the natural map $\chi_f(X) : X \rightarrow X_f^{\wedge}$ is a $C(f)$ -localization of X in \mathcal{M} .

PROOF. We need to check that $\chi_f(X)$ is a $C(f)$ -equivalence and that X_f^{\wedge} is $C(f)$ -local. Let $\phi : M \rightarrow N$ be a map in \mathcal{M} . For every $F \in \mathcal{M}$ we get an induced map

$$\text{Map}(C(f) \wedge N, F) \rightarrow \text{Map}(C(f) \wedge M, F),$$

and since $C(f)$ has a strong dual $D(C(f))$, we get a natural map

$$(3.3) \quad \text{Map}(N, D(C(f)) \wedge F) \rightarrow \text{Map}(M, D(C(f)) \wedge F).$$

ϕ is a $C(f)$ -equivalence if and only if the map (3.3) is an equivalence for every F . In particular, for every F in \mathcal{M} , $D(C(f)) \wedge F$ is $C(f)$ -local. Furthermore, Lemma 3.2.13 implies that $D(C(f)) \simeq \Sigma^{-1}D(L) \wedge C(f)$. Since local objects are stable under smashing with invertible objects, we conclude that for every $F \in \mathcal{M}$ the object $C(f) \wedge F$ is $C(f)$ -local.

Now we show that X_f^{\wedge} is $C(f)$ -local: since local objects are closed under inverse limits, we reduce to showing that $X \wedge C(f^n)$ is $C(f)$ -local. This easily follows by

induction. The base case is that $X \wedge C(f)$ is $C(f)$ -local, which was observed above. Assume we know that $X \wedge C(f^{n-1})$ is $C(f)$ -local. We conclude using the fiber sequence

$$(3.4) \quad L^{\wedge n-1} \wedge C(f) \wedge X \rightarrow C(f^n) \wedge X \rightarrow C(f^{n-1}) \wedge X$$

deduced from (3.2) and the 2-out-of-3 property of $C(f)$ -local objects in fiber sequences.

In order to show that the canonical map $X \rightarrow X_f^\wedge$ is a $C(f)$ -local equivalence, it suffices to show that $C(f) \wedge Y \simeq 0$ in \mathcal{M} , where $Y := \text{fib}(X \rightarrow X_f^\wedge)$. For this note that

$$Y \simeq \lim(X \wedge L^{\wedge*}) = \lim(\cdots \rightarrow X \wedge L^{\wedge n} \xrightarrow{r_f(L^{\wedge n-1})} X \wedge L^{\wedge(n-1)} \rightarrow \cdots)$$

and that $C(f) \wedge Y \simeq \text{cofib}(r_f(Y) : Y \wedge L \rightarrow Y)$. However it is easily checked that the multiplication by f on Y is induced by the multiplication by f on each component of the tower $X \wedge L^{\wedge*}$. The inverse limit of this tower only depends on its associated pro-object (see Appendix A.1), so we only need to show that the multiplication by f induces an equivalence of the pro-object associated to $X \wedge L^{\wedge*}$. On its turn this follows easily from [17, Lemma 3.6]. \blacksquare

PROPOSITION 3.2.15. *Let $r \geq 2$ be an integer, for every $i = 1, \dots, r$ let $f_i : L_i \rightarrow \mathbf{1}$ be a map in \mathcal{M} , and assume that for all i , L_i is \wedge -invertible. Then for every $X \in \mathcal{M}$ the natural map $\chi_{\underline{f}}(X) : X \rightarrow X_{f_1}^\wedge \cdots \wedge_{f_r}^\wedge$ is an $M(\underline{f})$ -localization of X in \mathcal{M} .*

PROOF. By Lemma 3.2.13, $M(\underline{f}) = C(f_1) \wedge \cdots \wedge C(f_r)$ has a strong dual

$$D(M(\underline{f})) \simeq \Sigma^{-r} (\wedge_{i=1}^r D(L_i)) \wedge M(\underline{f}).$$

Hence by running the same argument as in the proof of Proposition 3.2.14, we deduce that $F \wedge M(\underline{f})$ is $M(\underline{f})$ -local for every $F \in \mathcal{M}$. In order to show that $X_{f_1}^\wedge \cdots \wedge_{f_n}^\wedge$ is $M(\underline{f})$ -local, thanks to the identification

$$X_{f_1}^\wedge \cdots \wedge_{f_r}^\wedge \simeq \lim_{\mathbf{N}(\mathbf{N}^{\text{op}})} (X \wedge C(f_1^*) \wedge \cdots \wedge C(f_r^*)),$$

we only need to prove that each of the objects $X \wedge C(f_1^n) \wedge \cdots \wedge C(f_r^n)$ is $M(\underline{f})$ -local. This can be done by induction using iteratively the fiber sequence (3.4) and the fact that $M(\underline{f})$ -local objects satisfy the 2-out-of-3 property in fiber sequences.

The natural map

$$X \xrightarrow{\chi_{f_1}} X_{f_1}^\wedge \xrightarrow{\chi_{f_2}} X_{f_1 f_2}^\wedge \xrightarrow{\chi_{f_3}} \cdots \xrightarrow{\chi_{f_r}} X_{f_1}^\wedge \cdots \wedge_{f_r}^\wedge$$

is a composition of $M(\underline{f})$ -equivalences since $\langle M(\underline{f}) \rangle \leq \langle C(f_i) \rangle$ and since χ_{f_i} is a $C(f_i)$ -equivalence for every $i = 1, \dots, r$ by Proposition 3.2.14. \blacksquare

3.3 – A remark on η -completions

Let K be a perfect field of characteristic $\neq 2$, and consider the case where $\mathcal{M} = \mathcal{SH}(K)$. In this setting, let $\eta \in \pi_0(\mathbb{S})_{-1}(K)$ be the algebraic Hopf map. We have proved above in Proposition 3.2.14 that for every spectrum X the η -completion map $\chi_\eta(X) : X \rightarrow X_\eta^\wedge$ is the $M(\eta)$ -localization map in $\mathcal{SH}(K)$. We want to bring the discussion on η -completions a bit further.

LEMMA 3.3.1. *Assume that the base field K is not formally real. Then for every spectrum $X \in \mathcal{SH}(K)$ the spectrum $X[\frac{1}{2}]$ is η -complete.*

PROOF. It follows from [34, Chapter 2, Theorem 7.9] that there exists an integer n such that 2^n acts as 0 on the Witt ring of K . In particular, we deduce that in the Grothendieck–Witt ring $\text{GW}(K)$ the relation $2^n = h\omega$ holds, where h is the rank 2 hyperbolic space and ω is some element of $\text{GW}(K)$. It follows that $2^n\eta = h\omega\eta = 0$ in $\mathbb{K}_*^{MW}(K)$. It follows that on $X[\frac{1}{2}]$ the multiplication by η is the zero map, which in view of Section 3.2.9 is enough to conclude. ■

LEMMA 3.3.2. *Assume that the base field has finite étale 2-cohomological dimension. Then every strongly dualizable object of $\mathcal{SH}(K)$ is η -complete.*

PROOF. If C is a dualizable object of $\mathcal{SH}(K)$, then the operation of smashing with C commutes with inverse limits. In particular, $C_\eta^\wedge \simeq C \wedge \mathbb{S}_\eta^\wedge$ and thus we reduce to showing that \mathbb{S} is η -complete. In view of Proposition 4.1.1, we just need to show that the spectra $\mathbb{S}[\frac{1}{2}]$, $(\mathbb{S}_2^\wedge)[\frac{1}{2}]$ and \mathbb{S}_2^\wedge are η -complete. For the two former spectra the previous claim follows from Lemma 3.3.1, while for \mathbb{S}_2^\wedge the claim follows from the combination of [16, Lemma 21] and [13, Lemma A.7]. ■

3.4 – Inversions

3.4.1. Let J be a set and $\mathcal{B} = \{B_j\}_{j \in J}$ be a collection of objects of \mathcal{M} . Since \mathcal{M} is presentable, we can choose a set G of κ -compact objects that generate \mathcal{M} under κ -filtered colimits. Note that we can assume that κ has been chosen so that $\mathbf{1}$ is a κ -compact object of \mathcal{M} . Without loss of generality we can and will assume that the set G is stable under desuspension and that $\mathbf{1} \in G$. We define now \mathcal{B} as the smallest full sub- ∞ -category of \mathcal{M} which contains the objects of the form

$$V_1 \wedge \cdots \wedge V_k \wedge B_j \quad \text{where } k \in \mathbb{N}, V_1, \dots, V_k \in G, j \in J,$$

and which is closed under small colimits and extensions. Using [22, Proposition 1.4.1.11], we deduce that \mathcal{B} is a stable presentable ∞ -category. We set \mathcal{B}' to be the

full sub- ∞ -category of \mathcal{M} spanned by those objects X such that $\text{Map}(C, X) \simeq *$ for every $C \in \mathcal{B}$. The pair $\{\mathcal{B}, \mathcal{B}'\}$ forms then an accessible t -structure on \mathcal{M} (see [22, Proposition 1.4.4.11]). We denote by $\tau^{\mathcal{B}} : \mathcal{B}(-) \rightarrow \text{id}$ the associated co-localization map and by $\lambda^{\mathcal{B}} : \text{id} \rightarrow (-)^{\mathcal{B}}$ the associated localization map. The notation we use here is inspired by [7]. In particular, we have a natural fiber sequence of functors

$$\mathcal{B}(-) \xrightarrow{\tau^{\mathcal{B}}} \text{id} \xrightarrow{\lambda^{\mathcal{B}}} (-)^{\mathcal{B}}.$$

LEMMA 3.4.2. *The ∞ -category \mathcal{B}' is stable and presentably symmetric monoidal. The functor $(-)^{\mathcal{B}}$ is exact and symmetric monoidal, and the natural transformation $\lambda^{\mathcal{B}}$ is monoidal; finally the functor $\mathcal{B}(-)$ is exact.*

PROOF. The part of the statement about multiplicative structures follows essentially from [22, Proposition 2.2.1.9]. Indeed, according to [22, Example 2.2.1.7] we only need to check that \mathcal{B} is closed under smashing with arbitrary objects. This follows immediately from the fact that $- \wedge -$ commutes with colimits in both variables (since \mathcal{M} is presentably symmetric monoidal) and the definition of \mathcal{B} . The ∞ -categories \mathcal{B} and \mathcal{B}' are stable by [22, Corollary 1.4.2.27], so the functors $\mathcal{B}(-)$ and $(-)^{\mathcal{B}}$ are exact. \blacksquare

PROPOSITION 3.4.3. *Let $\mathcal{B} := \{B_j\}_{j \in J}$ be a set of strongly dualizable objects of \mathcal{M} . Then:*

(t1) *For every pair of objects X and Y of \mathcal{M} we have $\mathcal{B}X \wedge Y^{\mathcal{B}} \simeq 0$.*

(t2) *For every $X \in \mathcal{M}$ we have $X^{\mathcal{B}} \simeq \mathbf{1}^{\mathcal{B}} \wedge X$ and $\mathcal{B}X \simeq \mathcal{B}\mathbf{1} \wedge X$.*

(t3) *The following are equivalent for $X \in \mathcal{M}$:*

(a) *X is \mathcal{B} -local;*

(b) *$X \simeq \mathbf{1}^{\mathcal{B}} \wedge X$;*

(c) *$\mathcal{B}\mathbf{1} \wedge X \simeq 0$;*

(d) *$B_j \wedge V \wedge X \simeq 0$ for all $j \in J$ and all $V \in G$.*

(t4) *$\langle \bigoplus_{j \in J} B_j \rangle = \langle \bigoplus_{V \in G, j \in J} V \wedge B_j \rangle = \langle \mathcal{B}\mathbf{1} \rangle$.*

(t5) *$\langle \bigoplus_{j \in J} B_j \rangle^c = \langle \bigoplus_{V \in G, j \in J} V \wedge B_j \rangle^c = \langle \mathbf{1}^{\mathcal{B}} \rangle$.*

(t6) *The multiplication map $\mathbf{1}^{\mathcal{B}} \wedge \mathbf{1}^{\mathcal{B}} \rightarrow \mathbf{1}^{\mathcal{B}}$ is an equivalence.*

(t7) *An object X is $\bigoplus_{j \in J} B_j$ -acyclic if and only if X is a $\mathbf{1}^{\mathcal{B}}$ -module if and only if X is $\mathbf{1}^{\mathcal{B}}$ -local.*

PROOF. We start with (t1). Note that if X is in \mathcal{B}' , so is $X \wedge Y$ for every $Y \in \mathcal{M}$. Indeed, an object Z is in \mathcal{B}' if and only if $\text{Map}(V_1 \wedge \cdots \wedge V_k \wedge B_j, Z) \simeq *$ for every $j \in J$, every $k \in \mathbb{N}$ and every choice of $V_1, \dots, V_k \in G$. On its turn this is equivalent

to the condition that $D(B_j) \wedge Z \simeq 0$ for every $j \in J$. The latter condition is clearly stable under smashing over with an arbitrary object of \mathcal{M} . As a consequence ${}^{\mathcal{B}}X \wedge Y^{\mathcal{B}}$ is both local and co-local (for every $X, Y \in \mathcal{M}$), and thus equivalent to 0.

Now the rest easily follows. Property (t6) follows from (t1) and the fact that $\lambda^{\mathcal{B}} : \mathbf{1} \rightarrow \mathbf{1}^{\mathcal{B}}$ is a unit for the algebra $\mathbf{1}^{\mathcal{B}}$. Property (t2) follows from the facts that by design the localization and co-localization functors are exact, and the fact that both \mathcal{B} and \mathcal{B}' are tensor-ideals. Property (t3) is a direct consequence of (t1), (t2) and the fact that $- \wedge -$ commutes with colimits. Property (t4) follows from (t3). Property (t5) follows from (t1) and (t4).

The first implication of (t7) follows from (t3) and (t4). For the second implication note that $\mathbf{1}^{\mathcal{B}}$ -modules are obviously $\mathbf{1}^{\mathcal{B}}$ -local; on the other hand, $\mathbf{1}^{\mathcal{B}} \wedge X$ is a $\mathbf{1}^{\mathcal{B}}$ -localization, since the multiplication map of $\mathbf{1}^{\mathcal{B}}$ is an equivalence. ■

DEFINITION 3.4.4. Let $\mathcal{J} = \{g_j\}_{j \in J}$ be a collection of maps in \mathcal{M} of the form $g_j : L_j \rightarrow \mathbf{1}$. Let $B_j := C(g_j)$ and consider the set of objects $\mathcal{B} = \{B_j\}_{j \in J}$. In this case, the functor $(-)^{\mathcal{B}}$ (resp. the map $\lambda^{\mathcal{B}}$) is called \mathcal{J} -inversion functor (resp. map). In this special case, we use the following notational convention: $(-)^{\mathcal{B}} = (-)[\mathcal{J}^{-1}]$ and $\lambda^{\mathcal{B}} = \iota_{\mathcal{J}}$. The object $\mathbf{1}[\mathcal{J}^{-1}]$ is called \mathcal{J} -inverted Moore object.

3.4.5. The topic of inverting homotopy elements has been treated already in the language of ∞ -categories, for instance in [14] and [10, Appendix C]. In [14], Hoyois carries over a precise discussion about the possibility of describing $X[\mathcal{J}^{-1}]$ in terms of a telescope construction. This is relevant to our discussion, since his discussion directly implies the following.

COROLLARY 3.4.6. *Assume that \mathcal{M} is a presentably symmetric monoidal stable ∞ -category equipped with a multiplicative t -structure. Let J be a set and let $\mathcal{J} = \{g_j\}_{j \in J}$ be a collection of maps in \mathcal{M} of the form $g_j : L_j \rightarrow \mathbf{1}$, where for every $j \in J$, L_j is a t if object of \mathcal{M} . The functor $(-)[\mathcal{J}^{-1}]$ is right t -exact. If, in addition, the t -structure on \mathcal{M} is compatible with filtered colimits, then $(-)[\mathcal{J}^{-1}]$ is t -exact.*

PROOF. The objects L_j are \wedge -invertible, and by Remark 3.2.6 also 3-symmetric. We can thus combine [14, Theorem 3.8] and the discussion therein before Lemma 3.3 to ensure that $X[\mathcal{J}^{-1}]$ can be expressed via a filtered colimit, whose terms are of the form

$$\left(\bigwedge_{j \in H} D(L_j)^{\wedge a_j} \right) \wedge X,$$

where H varies among the collection of finite subsets of J . In addition, the flatness of the objects L_j , and thus of $D(L_j)$, implies that each term of the diagram is at least as connected as X ; in particular $(-)[\mathcal{J}^{-1}]$ respects $\mathcal{M}_{\geq 0}$. On the other hand $(-)[\mathcal{J}^{-1}]$ respects $\mathcal{M}_{\leq 0}$ when the t -structure is compatible with filtered colimits. ■

COROLLARY 3.4.7. *Let I be a finite set and J be any set. Assume that we have a pair of collections*

$$\{L_i\}_{i \in I} \quad \text{and} \quad \{L_j\}_{j \in J}$$

of strongly dualizable objects of \mathcal{M} and a pair of collections of maps

$$\{f_i : L_i \rightarrow \mathbf{1}\}_{i \in I} \quad \text{and} \quad \mathcal{J} = \{g_j : L_j \rightarrow \mathbf{1}\}_{j \in J}.$$

Denote by M the object $M(\underline{f}) \wedge \mathbf{1}[\mathcal{J}^{-1}]$. Then $\langle M \rangle$ has a complement, given explicitly by

$$\langle M \rangle^c = \left\langle \bigoplus_{i \in I} \mathbf{1}[f_i^{-1}] \right\rangle \oplus \left\langle \bigoplus_{j \in J} C(g_j) \right\rangle.$$

PROOF. It is an immediate consequence of equalities (2.2) together with Proposition 3.4.3 (t5). ■

REMARK 3.4.8. In the discussion carried out around [7, Proposition 2.13], Bousfield observes that Moore spectra span a subset of $\mathcal{B}\mathcal{A}(\mathcal{S}\mathcal{H})$ isomorphic to the power set of $\text{Spec}(\mathbb{Z})$. We believe it would be interesting to investigate a better (than our ad hoc) notion of Moore objects in the motivic category $\mathcal{S}\mathcal{H}(K)$ which involved Thornton's computation of the homogeneous spectrum of $K_*^{MW}(K)$ (see [37]).

4. Localization at some homotopy commutative algebras

In this section, we prove our main results about homology localizations. We dedicate Section 4.1 to some preliminary results and Section 4.2 to the statement of our main technical assumption. These are later used along Section 4.3 and in the proof of Theorem 4.3.7, which is the main result of this section. Throughout this section \mathcal{M} is a presentably symmetric monoidal stable ∞ -category, endowed with a left-complete multiplicative t -structure.

4.1 – Fracture squares

PROPOSITION 4.1.1. *Let E, F, G be objects of \mathcal{M} such that $\langle E \rangle = \langle F \rangle \oplus \langle G \rangle$ and such that every G -local object is F -acyclic. Consider the square*

$$\begin{array}{ccc} X_E & \xrightarrow{\lambda_F^E} & X_F \\ \downarrow \lambda_G^E & & \downarrow i \\ X_G & \xrightarrow{c} & (X_F)_G \end{array}$$

where the maps λ_F^E and λ_G^E are induced by $\langle F \rangle \leq \langle E \rangle \geq \langle G \rangle$, $c = \lambda_G(\lambda_F(X))$ is the G -localization of the map $\lambda_F(X) : X \rightarrow X_F$, and finally $i = \lambda_G(X_F)$. In this situation, the above square is cartesian.

In particular, let $E \in \mathcal{M}$ and $f : L \rightarrow \mathbf{1}$ be a map in \mathcal{M} with strongly dualizable source L . Set $E/f := E \wedge C(f)$. Then for every object $X \in \mathcal{M}$ the square

$$(4.1) \quad \begin{array}{ccc} X_E & \xrightarrow{\lambda_{E/f}^E} & X_{E/f} \\ \downarrow \lambda_{E[f^{-1}]}^E & & \downarrow i \\ X_{E[f^{-1}]} & \xrightarrow{c} & (X_{E/f})_{E[f^{-1}]} \end{array}$$

is cartesian.

PROOF. Let us denote by $P(X)$ the pull-back of the diagram

$$(4.2) \quad \begin{array}{ccc} & & X_F \\ & & \downarrow i \\ X_G & \xrightarrow{c} & (X_F)_G \end{array}$$

Since $\langle F \rangle \leq \langle E \rangle \geq \langle G \rangle$ all the objects appearing in diagram (4.2) are E -local, and thus $P(X)$ is E -local as well. We are thus left to prove that the natural map $u : X \rightarrow P(X)$, which is induced by the localization maps $\lambda_F(X) : X \rightarrow X_F$ and $\lambda_G(X) : X \rightarrow X_G$, is an E -equivalence. Since $\langle E \rangle = \langle F \rangle \oplus \langle G \rangle$, it suffices to show that u is both an F -equivalence and a G -equivalence. In particular, by the 2-out-of-3 property of local equivalences, we reduce to showing that

- the natural map $P(X) \xrightarrow{\alpha} X_G$ is a G -equivalence, which is obvious;
- the natural map $P(X) \xrightarrow{\beta} X_F$ is an F -equivalence, which follows from the assumption that G -local objects are F -acyclic.

The second part of the statement follows easily: set $F = E/f$, $G = E[f^{-1}]$, and notice that we have $\langle \mathbf{1} \rangle = \langle C(f) \rangle \oplus \langle \mathbf{1}[f^{-1}] \rangle$ by Proposition 3.4.3 (t5), so that $\langle E \rangle = \langle E/f \rangle \oplus \langle E[f^{-1}] \rangle$ by Proposition 3.4.3 (t2). Finally, use the chain of inclusions

$$\mathrm{Loc}_{E[f^{-1}]} \subseteq \mathrm{Loc}_{\mathbf{1}[f^{-1}]} = \mathrm{Ac}_{C(f)} \subseteq (Ac)_{E[f^{-1}]};$$

the only non-obvious part is the central equality: it follows from Proposition 3.4.3 (t7). This concludes the proof. \blacksquare

COROLLARY 4.1.2. *Let E , X and f be as in the statement of Proposition 4.1.1. Then $X_{E \wedge C(f)}$ is equivalent to $(X_E)_{C(f)}$.*

PROOF. We start by considering the following commutative diagram in \mathcal{M} :

$$\begin{array}{ccc} X_E & \xrightarrow{\lambda_1} & (X_E)_{C(f)} \\ \downarrow \lambda_3 & & \downarrow \lambda_4 \\ X_{E/f} & \xrightarrow{\lambda_2} & (X_{E/f})_{C(f)} \end{array}$$

where $\lambda_1 = \lambda_{C(f)}(X_E)$, $\lambda_3 = \lambda_{E/f}^E$ is the map introduced above, $\lambda_4 = \lambda_{C(f)}(\lambda_3)$, and finally $\lambda_2 = \lambda_{C(f)}(X_{E/f})$.

Since $\langle E/f \rangle \leq \langle C(f) \rangle$, λ_2 is actually an equivalence in \mathcal{M} and we are left to prove that λ_4 is too. For this we apply the $C(f)$ -localization functor to the square (4.1) and we use Proposition 4.1.1 to reduce the proof to checking that

$$X_{E[f^{-1}]} \xrightarrow{c} (X_{E/f})_{E[f^{-1}]}$$

is a $C(f)$ -equivalence. Now both the source and target of c are $\mathbf{1}[f^{-1}]$ -local, being in fact $E[f^{-1}]$ -local. Thus, after $C(f)$ -localization both source and target of c become zero. \blacksquare

COROLLARY 4.1.3. *Let r be a positive integer. For every $i = 1, \dots, r$ consider strongly dualizable objects $L_i \in \mathcal{M}$ and maps $f_i : L_i \rightarrow \mathbf{1}$ in \mathcal{M} . Let $M := C(f_1) \wedge \dots \wedge C(f_r)$ be the Moore object associated to the maps f_1, \dots, f_r . Then for every pair of objects E and X of \mathcal{M} we have*

$$X_{E \wedge M} \simeq (\dots((X_E)_{C(f_1)})_{C(f_2)} \dots)_{C(f_r)} \simeq (X_E)_M.$$

PROOF. Apply inductively Corollary 4.1.2. \blacksquare

4.2 – Technical assumption

We now formulate a technical assumption, which we need for the proofs of Theorems 4.3.7, 7.3.5 and 7.3.9.

ASSUMPTION 4.2.1. E is an object of \mathcal{M} satisfying the following properties:

- (1) E is a homotopy commutative algebra.
- (2) $E \in \mathcal{M}_{\geq 0}$.
- (3) There are a finite set of tif objects $\{L_i\}_{i \in I}$ in \mathcal{M} and maps $f_i : L_i \rightarrow \mathbf{1}$, a set of tif objects $\{L_j\}_{j \in J}$ and maps $g_j : L_j \rightarrow \mathbf{1}$, and a morphism of $\tau_0(\mathbf{1})$ -algebras

$$\phi : \tau_0((\tau_0(\mathbf{1})/\mathcal{I})[\mathcal{J}^{-1}]) \rightarrow \tau_0 E,$$

where \mathcal{I} is the image of $\sum_{i \in I} \tau_0 f_i : \bigoplus_{i \in I} \tau_0(L_i) \rightarrow \tau_0 \mathbf{1}$ and \mathcal{J} is the collection of elements $\{g_j\}_{j \in J}$.

(4) The map ϕ given in (3) is an isomorphism.

In this situation, we will denote by $M_{\tau_0} E$ the object $C(f_1) \wedge \cdots \wedge C(f_r) \wedge \mathbf{1}[\mathcal{J}^{-1}]$, and sloppily refer to it as the Moore object associated with $\tau_0(E)$, rather than with the maps f_i and g_j .

Note that the natural map of commutative algebras

$$(\tau_0(\mathbf{1})/\mathcal{I})[\mathcal{J}^{-1}] \rightarrow \tau_0((\tau_0(\mathbf{1})/\mathcal{I})[\mathcal{J}^{-1}])$$

is an equivalence as soon as $J = \emptyset$ or when the t -structure on \mathcal{M} is compatible with filtered colimits.

4.3 – Comparison of E - and $\tau_0 E$ -localization

LEMMA 4.3.1. *Let $E \in \mathcal{M}_{\geq k}$ be a homotopy commutative algebra in \mathcal{M} and let $f : L \rightarrow \mathbf{1}$ be a map in \mathcal{M} whose source L is an sdf object. Then left multiplication by f on E , $l_f(E)$, is an equivalence if and only if left multiplication by f on $\tau_0(E)$, $l_f(\tau_0(E))$, is an equivalence.*

PROOF. If $l_f(E)$ is an equivalence, then the induced map $\tau_0(l_f(E))$ is too, and by Lemma 3.2.7 $\tau_0(l_f(E)) = l_f(\tau_0(E))$, as morphisms of \mathcal{M}^\heartsuit . For the other implication we argue as follows. By left-completeness and the fact that L is a strongly dualizable object, the map $l_f(E) : L \wedge E \rightarrow E$ is naturally identified with the limit of $l_f(P^n(E)) : L \wedge P^n(E) \rightarrow P^n(E)$, and for checking that it is an equivalence one only needs to check that, for every integer n , multiplication by f on $\tau_n(E)$ is an equivalence. Now $\tau_n(E)$ is a $\tau_0(E)$ -module and thus is endowed with an action map $a : \tau_0(E) \otimes^\heartsuit \tau_n(E) \rightarrow \tau_n(E)$. Multiplication by f commutes with such action map by Section 3.2.3, in the sense that we have a commutative square

$$\begin{array}{ccc} L \wedge \tau_n(E) & \xrightarrow{l_f(\tau_n(E))} & \tau_n(E) \\ \uparrow a & & \uparrow a \\ L \wedge \tau_0(E) \otimes^\heartsuit \tau_n(E) & \longrightarrow & \tau_0(E) \otimes^\heartsuit \tau_n(E), \end{array}$$

where the lower horizontal map is $l_f(\tau_0(E) \otimes^\heartsuit \tau_n(E))$. However, such map is homotopic to $l_f(\tau_0(E)) \otimes^\heartsuit \tau_n(E)$, and thus is an equivalence by assumption. We conclude noticing that the upper horizontal map of the above square is a retraction of the lower horizontal map via the unit morphism for the $\tau_0(E)$ -module $\tau_n(E)$. ■

PROPOSITION 4.3.2. *Let E be an object of \mathcal{M} satisfying points (1)–(3) of Assumption 4.2.1. Then*

$$\langle \tau_0 E \rangle \leq \langle E \rangle \leq \langle M \tau_0 E \rangle.$$

PROOF. Since $E \in \mathcal{M}_{\geq 0}$ by Assumption 4.2.1, the projection to the Postnikov truncation induces a map $p : E \rightarrow \tau_0 E$. As it follows from the discussion in Section 2.1.5, $\tau_0 E$ has a natural homotopy algebra structure for which p is a ring map. As a consequence $\tau_0(E)$ is a retract of $E \wedge \tau_0(E)$, and it follows that $\langle \tau_0(E) \rangle = \langle E \wedge \tau_0(E) \rangle \leq \langle E \rangle$.

It remains to show that $\langle E \rangle = \langle E \wedge M \tau_0 E \rangle$, which directly implies our claim that $\langle E \rangle \leq \langle M \tau_0 E \rangle$. For this, recall that

$$\langle M \tau_0 E \rangle = \langle C(f_1) \wedge \cdots \wedge C(f_n) \wedge \mathbf{1}[g^{-1}] \rangle.$$

From Corollary 3.4.7, the Bousfield class $\langle M \tau_0 E \rangle$ has a complement $\langle M \tau_0 E \rangle^c = \langle M \rangle$, where

$$M := \left(\bigoplus_{i \in I} \mathbf{1}[f_i^{-1}] \right) \oplus \left(\bigoplus_{j \in J} C(g_j) \right).$$

In particular,

$$\langle E \rangle = \langle E \rangle \wedge \langle (M \tau_0 E \oplus M) \rangle = \langle E \wedge M \tau_0 E \rangle \oplus \langle E \wedge M \rangle;$$

since smashing commutes with small sums, it suffices to see that $E \wedge \mathbf{1}[f_i^{-1}] = 0 = E \wedge C(g_j)$ for every $i \in I$ and every $j \in J$.

Let $f : L \rightarrow \mathbf{1}$ be any of the f_i and let $E' := E \wedge \mathbf{1}[f^{-1}]$. We claim that $\tau_0(E') \simeq 0$. For proving it, we show that multiplying by f on $\tau_0(E')$ is an isomorphism that factors through 0. In detail, the left multiplication $l_f(E')$ by f on E' is an equivalence and so is $\tau_0(l_f(E'))$. Lemma 3.2.7 implies that $\tau_0(l_f(E')) = l_f(\tau_0(E'))$ and that $l_f(\tau_0(E'))$ coincides with the map induced on τ_0 by

$$l_f(\tau_0(E)) \wedge \tau_0(\mathbf{1}[f^{-1}]) : L \wedge \tau_0(E) \wedge \tau_0(\mathbf{1}[f^{-1}]) \rightarrow \tau_0(E) \wedge \tau_0(\mathbf{1}[f^{-1}]),$$

since $E' \in \mathcal{M}_{\geq 0}$ by Corollary 3.4.6. By assumption we have $l_f(\tau_0(E)) = 0$. Indeed $\tau_0(E)$ is a $\tau_0(\mathbf{1})/\mathcal{I}$ -module, thus $l_f(\tau_0(E))$ is a retraction of $l_f(\tau_0(\mathbf{1})/\mathcal{I}) \wedge \tau_0(E)$, and $l_f(\tau_0(\mathbf{1})/\mathcal{I}) = 0$ directly by its definition. It follows that $\tau_0(E') \simeq 0$, and in particular $E' \in \mathcal{M}_{\geq 1}$. Since E' has a homotopy unital multiplication, the equivalence $\mathbf{1} \wedge E' \simeq E'$ factors through $E' \wedge E' \in \mathcal{M}_{\geq 2}$, so that $\tau_1(E') \simeq 0$. Inductively this shows that $\tau_i(E') \simeq 0$ for all i . Since $E' \in \mathcal{M}_{\geq 0}$ and the t -structure is left complete, we conclude that $E' \simeq 0$. Similarly, let $g : L \rightarrow \mathbf{1}$ be any of the g_j . Multiplication by g on E is an equivalence by Lemma 4.3.1, since it is so on $\tau_0(E)$, and thus $E \wedge C(g) \simeq 0$. ■

LEMMA 4.3.3. *Let X be any k -connected object of \mathcal{M} for some integer k . Then X is $\tau_0(\mathbf{1})$ -local.*

PROOF. By the multiplicative properties of the Postnikov tower (see Section 2.1.3), the homotopy objects $\tau_p(X)$ are $\tau_0(\mathbf{1})$ -modules and hence $\tau_0(\mathbf{1})$ -local. By connectivity of X , every stage of the Postnikov tower $P^n(X)$ is a finite extension of suspensions of the $\tau_p(X)$, and hence it is $\tau_0(\mathbf{1})$ -local. Finally, $X \simeq \lim_n P^n(X)$ by left completeness of the t -structure. ■

LEMMA 4.3.4. *Let $\{\mathbf{A}, \otimes, \mathbf{1}\}$ be a symmetric monoidal category where \mathbf{A} is also abelian and such that \otimes is right exact. Let R be a commutative monoid in \mathbf{A} and denote by e_R and μ_R its unit and multiplication. Let $L \in \mathbf{A}$ and assume we have a map $f : L \rightarrow R$. Then:*

- (1) *There exists a unique R -linear map $\cdot f : R \otimes L \rightarrow R$ that, composed with*

$$L \simeq \mathbf{1} \otimes L \xrightarrow{e_R \otimes \text{id}_L} R \otimes L,$$

gives back f .

- (2) *The R -module $C := \text{coker}(\cdot f)$ has a unique structure of commutative R -algebra having the natural projection $p : R \rightarrow C$ as unit.*
- (3) *If the multiplication of R is an isomorphism, so is that of C .*
- (4) *If L is a \otimes -invertible object of \mathbf{A} , then $K := \ker(\cdot f)$ has a unique structure of C -module, making the inclusion $i : K \hookrightarrow R \otimes L$ into a map of R -modules.*
- (5) *If M is an R -module in \mathbf{A} with action map $\alpha : R \otimes M \rightarrow M$ and*

$$\cdot f_M = \alpha \circ (f \otimes \text{id}) : L \otimes M \rightarrow R \otimes M \rightarrow M$$

is the induced multiplication by f on M , then $\text{coker}(\cdot f_M)$ and $\ker(\cdot f_M)$ have a unique structure of C -module induced by α .

PROOF. Point (1) follows from the usual free-forget adjunction. Point (2) and (3) follow from elementary diagram chases. Regarding (4) one needs to show that the natural action of R on K factors through C . Let us denote by i the monomorphism $K \subseteq R \otimes L$. Consider the map

$$\phi : R \otimes L \otimes R \otimes L \rightarrow R \otimes L \quad \text{defined by} \quad \phi = \mu \otimes \text{id}_L \circ \cdot f \otimes \text{id}_R \otimes \text{id}_L,$$

so that, with an abuse of notation, $\phi(r_1 \otimes l_1 \otimes r_2 \otimes l_2) = r_1 f(l_1) r_2 \otimes l_2$. An easy diagram chase shows that the required factorization exists if and only if the map ϕ composed with $j := \text{id} \otimes \text{id} \otimes i : R \otimes L \otimes K \rightarrow R \otimes L \otimes R \otimes L$ is zero. What we know, however, is that $\phi \circ t_{(12),(34)} \circ j = 0$, where $t_{(12),(34)}$ is the switching of the first and second pair of terms on $R \otimes L \otimes R \otimes L$. We claim that ϕ is a multiple of $\phi \otimes t_{(12),(34)}$ by a unit $\varepsilon \in \text{End}_{\mathbf{A}}(\mathbf{1})$. For this note that the permutation $t_{(12),(34)} = t_{1,3} \circ t_{2,4}$

of the R terms gives no trouble, since μ is commutative, so $\phi = \phi \circ t_{1,3}$. Here $t_{1,3}$ (resp. $t_{2,4}$) denotes the switching of the first and third (resp. second and fourth) tensor factors of $R \otimes L \otimes R \otimes L$. Since L is \otimes -invertible, $t_{2,4}$ can be identified with left multiplication by a suitable unit $\varepsilon \in \text{End}_{\mathbf{A}}(\mathbf{1})$. In conclusion, $\phi \circ t_{(12),(34)} = \phi \circ l_{\varepsilon}(R \otimes L \otimes R \otimes L)$, and thus

$$\begin{aligned} \phi \circ j &= \phi \circ t_{(12),(34)} \circ l_{\varepsilon}(R \otimes L \otimes R \otimes L)^{-1} \circ j \\ &= \phi \circ t_{(12),(34)} \circ j \circ l_{\varepsilon}(R \otimes L \otimes R \otimes L)^{-1} = 0, \end{aligned}$$

since multiplication by a unit commutes with every map in \mathbf{A} by Section 3.2.3.

The proof of point (5) works similarly, and we omit it. \blacksquare

LEMMA 4.3.5. *Let \mathcal{R} be a commutative algebra in \mathcal{M}^{\heartsuit} and let $f : \tau_0(L) \rightarrow \mathcal{R}$ be a map in \mathcal{M}^{\heartsuit} where L is a tif object of \mathcal{M} . Let C denote the cofiber of the induced map $r_f(\mathcal{R}) = \cdot f : \mathcal{R} \wedge L \simeq \mathcal{R} \otimes^{\heartsuit} \tau_0(L) \rightarrow \mathcal{R}$ given by Lemma 4.3.4 in \mathcal{M}^{\heartsuit} . Then $\tau_0(C) \simeq \text{coker}(\cdot f)$ and $\langle C \rangle \leq \langle \tau_0(C) \rangle$.*

PROOF. From the exact sequence of homotopy objects

$$0 \rightarrow \mathcal{K} \rightarrow L \wedge \mathcal{R} \xrightarrow{\cdot f} \mathcal{R} \rightarrow \text{coker}(\cdot f) \rightarrow 0$$

we deduce

$$\tau_k(C) = \begin{cases} \text{coker}(\cdot f) & \text{if } k = 0, \\ \ker(\cdot f) & \text{if } k = 1, \\ 0 & \text{otherwise.} \end{cases}$$

In particular, it follows that we have a fiber sequence

$$\Sigma^1 \tau_1 C \rightarrow C \rightarrow \tau_0 C$$

relating C with its truncations. Observe that, thanks to Lemma 4.3.4, $\tau_0(C)$ is a commutative algebra in \mathcal{M}^{\heartsuit} and $\Sigma^1 \tau_1(C)$ is a module in \mathcal{M}^{\heartsuit} over $\tau_0(C)$. More generally, for every $X \in \mathcal{M}$, the object $\tau_1(C) \wedge X$ is also a $\tau_0(C)$ -module in \mathcal{M} . In particular, if X is $\tau_0(C)$ -acyclic, then it is also $\Sigma^1 \tau_1(C)$ -acyclic, and by the above fiber sequence X is C -acyclic too. \blacksquare

COROLLARY 4.3.6. *Assume that \mathcal{M} is a presentably symmetric monoidal stable ∞ -category endowed with a left-complete multiplicative t -structure. Assume that E is a homotopy commutative algebra in \mathcal{M} satisfying Assumption 4.2.1, and in case $J \neq \emptyset$ we assume in addition that the t -structure of \mathcal{M} is compatible with filtered colimits. Then $\langle \tau_0 \mathbf{1} \wedge M \tau_0(E) \rangle \leq \langle \tau_0 E \rangle$.*

PROOF. Readily Lemma 4.3.5 implies that $\langle \tau_0(\mathbf{1}) \wedge C(f_1) \rangle \leq \langle \tau_0(\mathbf{1})/(f_1) \rangle$. Recall that, by definition,

$$M\tau_0 E = C(f_1) \wedge \cdots \wedge C(f_r) \wedge \mathbf{1}[\mathcal{J}^{-1}],$$

so we can proceed in order by smashing with one $C(f_i)$ at the time. Indeed, by the previous case

$$\langle \tau_0(\mathbf{1}) \wedge C(f_1) \wedge C(f_2) \rangle \leq \langle \tau_0(\mathbf{1})/(f_1) \wedge C(f_2) \rangle$$

and finally, using Lemma 4.3.5 with $\mathcal{R} = \tau_0(\mathbf{1})/(f_1)$ and $f = f_2$, we get that

$$\langle \tau_0(\mathbf{1})/(f_1) \wedge C(f_2) \rangle \leq \langle (\tau_0(\mathbf{1})/(f_1))/(f_2) \rangle = \langle \tau_0(\mathbf{1})/(f_1, f_2) \rangle,$$

so we conclude that

$$\langle \tau_0(\mathbf{1}) \wedge C(f_1) \wedge C(f_2) \rangle \leq \langle \tau_0(\mathbf{1})/(f_1, f_2) \rangle.$$

Inductively we arrive at

$$\langle \tau_0(\mathbf{1}) \wedge M(\underline{f}) \rangle \leq \langle \tau_0(\mathbf{1})/(f_1, \dots, f_r) \rangle,$$

where $M(\underline{f}) = C(f_1) \wedge \cdots \wedge C(f_r)$. Finally, we observe that

$$\langle \tau_0(\mathbf{1}) \wedge M\tau_0 E \rangle = \langle \tau_0(\mathbf{1}) \wedge M(\underline{f}) \wedge \mathbf{1}[\mathcal{J}^{-1}] \rangle \leq \langle \tau_0(\mathbf{1})/(f_1, \dots, f_r) \wedge \mathbf{1}[\mathcal{J}^{-1}] \rangle,$$

and that

$$\langle \tau_0(\mathbf{1})/(f_1, \dots, f_r) \wedge \mathbf{1}[\mathcal{J}^{-1}] \rangle = \langle \tau_0 E \rangle,$$

since $-\wedge \mathbf{1}[\mathcal{J}^{-1}]$ is t -exact according to Corollary 3.4.6. ■

THEOREM 4.3.7. *Let E be a homotopy commutative algebra in \mathcal{M} satisfying Assumption 4.2.1 in the special case that $J = \emptyset$. Then for every integer k , and every $X \in \mathcal{M}_{\geq k}$ we have that*

$$X_{M\tau_0 E} \simeq X_E.$$

PROOF. Thanks to Proposition 4.3.2 we know that for any $X \in \mathcal{M}_{\geq k}$ the localization map $X \rightarrow X_{M\tau_0 E}$ is an E -equivalence, so we only have to check that $X_{M\tau_0 E}$ is E -local. Now consider that $X \rightarrow X_{\tau_0(\mathbf{1})}$ is an equivalence by Proposition 4.3.3 so that

$$X_{M\tau_0 E} \rightarrow (X_{\tau_0(\mathbf{1})})_{M\tau_0 E}$$

is an equivalence too. In particular, by combining this with the result of Corollary 4.1.2 we deduce that

$$X_{M\tau_0 E} \xrightarrow{\simeq} (X_{\tau_0(\mathbf{1})})_{M\tau_0 E} \simeq X_{\tau_0(\mathbf{1}) \wedge M\tau_0 E}.$$

Finally, we apply Proposition 4.3.2 and Corollary 4.3.6 to deduce that $X_{\tau_0(\mathbf{1}) \wedge M\tau_0 E}$ is E -local. This concludes the proof. ■

5. Examples and applications

We provide some samples of usage in the motivic setting of the results seen so far. In this section, S denotes a Noetherian scheme of finite Krull dimension.

5.1 – Algebraic cobordisms

Let MGL (resp. MSL) be the spectrum representing Voevodsky’s algebraic cobordism (resp. special linear cobordism) over S . In [27] (resp. [30]), Panin et al. construct MGL (resp. MSL) as a commutative monoid in a monoidal model category presenting $\mathcal{SH}(S)$. By [13, Theorem 3.8], MGL satisfies Assumption 4.2.1; we conclude that the MGL-localization map is canonically identified with the η -completion map $\chi_\eta(X) : X \rightarrow X_\eta^\wedge$ on bounded-below spectra. Similarly in [38], when S is the spectrum of a perfect field of characteristic not 2, Yakerson verifies that MSL satisfies Assumption 4.2.1 with $I = J = \emptyset$, and as a consequence $(-)_\text{MSL} \simeq \text{id}$ on bounded-below spectra.

5.2 – Motivic cohomologies

Let S be a finite-dimensional Noetherian scheme, which is essentially smooth over a field K . Let HZ be the spectrum representing Voevodsky’s motivic cohomology with integral coefficients. Recall that we have a category of motives $\mathcal{DM}(S, \mathbb{Z})$ which is related to $\mathcal{SH}(S)$ by an adjunction

$$(5.1) \quad \mathbb{Z}_\text{tr} : \mathcal{SH}(S) \rightleftarrows \mathcal{DM}(S, \mathbb{Z}) : u_\text{tr}.$$

Since u_tr respects algebras, as it follows from [13, Section 4], $\text{HZ} = u_\text{tr} \mathbf{1}$ is a commutative algebra in $\mathcal{SH}(S)$. Moreover, [13, Theorem 7.4] implies that Assumption 4.2.1 is verified, and that $\tau_0(\text{HZ}) \simeq \tau_0(\mathbb{S}/\eta)$.

We can thus apply Theorem 4.3.7, to deduce that the HZ -localization of a bounded-below spectrum X is identified with the η -completion map $X \rightarrow X_\eta^\wedge$. In a similar fashion, let $E = \text{HZ}/\ell$ be the spectrum representing motivic cohomology with modulo ℓ coefficients. The same considerations allow us to conclude that the HZ/ℓ -localization map $\lambda_{\text{HZ}/\ell}(X) : X \rightarrow X_{\text{HZ}/\ell}$ of a bounded-below spectrum X is identified with the formal completion map $\chi_{\ell, \eta}(X) : X \rightarrow X_{\ell, \eta}^\wedge$.

The formalism of motives we have just recalled has a quadratic analogue. Let now S be the spectrum of an infinite perfect field K of characteristic not 2. We have a category of Chow–Witt motives $\widetilde{\mathcal{DM}}(K, \mathbb{Z})$ with a pair of adjoint functors

$$(5.2) \quad \widetilde{\mathbb{Z}}_\text{tr} : \mathcal{SH}(K) \rightleftarrows \widetilde{\mathcal{DM}}(K) : \widetilde{u}_\text{tr}$$

which are the stabilizations of the functors which respectively add and forget generalized transfers. The discussion in [4, Chapter 6, Section 4.1] shows that $\tilde{\mathrm{HZ}}$ is a commutative algebra in $\mathcal{SH}(K)$. By combining [4, Chapter 3, Proposition 4.1.2] and [4, Chapter 3, Theorem 4.2.3], we see that Assumption 4.2.1 is satisfied. It thus follows by Theorem 4.3.7 that for every bounded-below spectrum X , the $\tilde{\mathrm{HZ}}$ -localization is an equivalence.

5.3 – Slice completion

We recall the following result.

THEOREM 5.3.1 ([33, Theorem 3.50]). *Let K be a field of exponential characteristic p . Suppose that X is a spectrum having a cell presentation of finite type in $\mathrm{Mod}_A(K)$ where $A = \mathbb{S}[\frac{1}{p}]$. Then we have a canonical commutative square in $\mathrm{Mod}_A(K)$:*

$$\begin{array}{ccc} X & \xrightarrow{\chi_\eta(X)} & X_\eta^\wedge \\ \downarrow \sigma(X) & & \downarrow \sigma(X)_\eta^\wedge \\ X_{\mathrm{sc}} & \xrightarrow{\chi_\eta(X_{\mathrm{sc}})} & (X_{\mathrm{sc}})_\eta^\wedge, \end{array}$$

where the maps $\chi_\eta(X_{\mathrm{sc}})$ and $\sigma(X)_\eta^\wedge$ are equivalences. In particular, there is a natural isomorphism $X_{\mathrm{sc}} \simeq X_\eta^\wedge$ in $\mathrm{Mod}_A(K)$ under which the slice completion map $\sigma(X)$ and the η -completion map $\chi_\eta(X)$ are identified.

By combining Section 5.2 with the previous result, we deduce that for a cell spectrum X of finite type, $\mathrm{HZ} \wedge X = 0$ if and only if $X_{\mathrm{sc}} = 0$. In other words, if $f : X \rightarrow Y \in \mathrm{Mod}_A(K)$ is a map between objects that have a cell presentation of finite type, f is an equivalence on slice completions if and only if f induces an equivalence on motivic homology.

5.4 – Motives of spectra

Let K be a field of exponential characteristic p and $A = \mathbb{S}[\frac{1}{p}]$ and $R = \mathbb{Z}[\frac{1}{p}]$. Recall that the adjunction (5.1) factors as

$$\begin{array}{ccc} \mathcal{SH}(K)[\frac{1}{p}] \simeq \mathrm{Mod}_A(K) & \begin{array}{c} \xleftarrow{R_{\mathrm{tr}}} \\ \xrightarrow{u_{\mathrm{tr}}} \end{array} & \mathcal{DM}(K, R) \\ \begin{array}{c} \swarrow \mathrm{HR} \wedge_A - \\ \searrow U_{\mathrm{HR}} \end{array} & & \begin{array}{c} \swarrow \Phi \\ \searrow \Psi \end{array} \\ & & \mathrm{Mod}_{\mathrm{HR}}(K), \end{array}$$

where $R_{\mathrm{tr}}(-) \simeq \Phi(\mathrm{HR} \wedge_A -)$ and $u_{\mathrm{tr}} \simeq U_{\mathrm{HR}}(\Psi(-))$.

Similarly, at least when $p \neq 2$, the adjunction (5.2) factors as

$$\begin{array}{ccc}
 \mathcal{SH}(K)[\frac{1}{p}] \simeq \mathcal{M}od_A(K) & \begin{array}{c} \xleftarrow{\tilde{R}_{\text{tr}}} \\ \xrightarrow{\tilde{u}_{\text{tr}}} \end{array} & \widetilde{\mathcal{DM}}(K, R) \\
 \begin{array}{c} \xleftarrow{\tilde{H}R \wedge_A -} \\ \xrightarrow{\tilde{U}_{\text{HR}}} \end{array} & & \begin{array}{c} \xrightarrow{\tilde{\Phi}} \\ \xleftarrow{\tilde{\Psi}} \end{array} \\
 & & \mathcal{M}od_{\tilde{H}R}(K).
 \end{array}$$

The previous observations prove the following statement.

COROLLARY 5.4.1. *Let K be a perfect field of exponential characteristic $p \neq 2$. Then:*

- (1) *If K is not formally real the functor R_{tr} is conservative on bounded-below $\mathbb{S}[\frac{1}{2}]$ -modules.*
- (2) *If $\text{cd}_2(K) < \infty$, then R_{tr} is conservative on strongly dualizable objects, and in particular on compact objects.*
- (3) *If K is infinite, then \tilde{R}_{tr} is conservative on bounded-below objects.*

PROOF. By [32, Theorem 1] (for $p = 0$) or [15, Theorem 5.8] (in general), we have that (Φ, Ψ) is a pair of adjoint equivalences. Now $X \simeq X[\frac{1}{2}]$, so let us assume that $\text{HR} \wedge_A X = 0$. Then by the assumption on X , Theorem 4.3.7, and Proposition 3.2.14, we have that

$$0 = X_{\text{HR}} = X_{\eta}^{\wedge}.$$

If K is not formally real, Lemma 3.3.1 implies that $\chi_{\eta}(X) : X \rightarrow X_{\eta}^{\wedge}$ is an equivalence in $\mathcal{M}od_A(K)$ and hence $X \simeq 0$. Let us now assume that K has finite 2-cohomological dimension. We achieve the second point by running the same argument, but using Lemma 3.3.2 instead of Lemma 3.3.1. We deduce that, if X is a dualizable object with $\text{HR} \wedge_A X = 0$, then $X \simeq 0$ in $\mathcal{M}od_A(K)$. Since p is inverted in the coefficients, strongly dualizable objects and compact objects of $\mathcal{SH}(K)[\frac{1}{p}]$ are the same, cf. [12, Lemma 2.3]. The third point works similarly to the first, but using only Theorem 4.3.7, combined with the fact that $(\tilde{\Phi}, \tilde{\Psi})$ are mutually inverse equivalences, which follows from [12, Theorem 5.2]. \blacksquare

REMARK 5.4.2. In [3, Theorem 1], Bachmann proves that over a perfect field K of exponential characteristic $p \neq 2$ and with $\text{cd}_2(K) < \infty$, the functor R_{tr} is conservative on effective and bounded-below spectra where p acts invertibly. In the case where 2 is inverted in the coefficients, Corollary 5.4.1 is only apparently more general. A direct inspection of Bachmann's argument shows that, upon inverting two, he does not need to assume that $\text{cd}_2(K) < \infty$.

However, if we do not wish to invert the prime 2 in the coefficients, Bachmann's argument needs an extra non-trivial input, namely a description of the slice filtration on homotopy groups of spectra, coming from [20]. In this case, our approach is genuinely different and not as powerful, but in any case it recovers a very non-trivial portion of the stated result of Bachmann.

Nevertheless, we have not been able to use Bachmann's results to recover our results on HZ -localizations. It would probably be interesting to employ his techniques, particularly those using the real étale topology, for the study of homology localizations.

5.5 – K -theories

Let S be a finite-dimensional Noetherian scheme, which is essentially smooth over a field K of exponential characteristic p (resp. let S be the spectrum of a field K of characteristic $p \neq 2$). Recall that we have spectra KGL (resp. KQ) in $\mathcal{SH}(S)[\frac{1}{p}]$ representing algebraic K -theory (resp. algebraic Hermitian K -theory). The tensor product on bundles induces natural homotopy commutative algebra structures on KGL and on KQ . This can be found respectively in [28, Theorem 2.2.1] and in [29, Theorem 1.5]: in both cases, the multiplicative structure is constructed over $\mathrm{Spec}(\mathbb{Z})$ and $\mathrm{Spec}(\mathbb{Z}[\frac{1}{2}])$ respectively, and then pulled back over more general bases. Thanks to the multiplicative properties of the slice tower (resp. the very effective slice tower), the effective cover $f_0(\mathrm{KGL})$ (resp. the very effective cover $\tilde{f}_0(\mathrm{KQ})$) has a structure of homotopy commutative algebra as well.

We first deal with KGL . Consider the fiber sequence

$$f_1 \mathrm{KGL} \rightarrow f_0 \mathrm{KGL} \rightarrow s_0 \mathrm{KGL}.$$

On the one hand we have an isomorphism of commutative algebras $s_0 \mathrm{KGL} \xrightarrow{\cong} \mathrm{HZ}[\frac{1}{p}]$ in $\mathcal{SH}(S)[\frac{1}{p}]$, see [19, Section 11] and [13, Theorem 8.5]. On the other hand we have $f_i \mathrm{KGL} \in \mathcal{SH}(S)[\frac{1}{p}]_{\geq i}$ thanks to [13, Lemma 8.11]: this can be applied since KGL is the spectrum representing the cohomology theory associated with the Landweber exact formal group law $X + Y - \beta XY$ on $\mathbb{Z}[\beta, \beta^{-1}]$ by [36]. In conclusion, $f_0 \mathrm{KGL}$ satisfies Assumption 4.2.1 with $\tau_0 f_0 \mathrm{KGL} \simeq \tau_0(S/\eta)$.

Now we deal with KQ . Consider the fiber sequence

$$\tilde{f}_1 \mathrm{KQ} \rightarrow \tilde{f}_0 \mathrm{KQ} \rightarrow \tilde{s}_0 \mathrm{KQ}.$$

This time we have an isomorphism of commutative algebras $\tau_0 \tilde{s}_0 \mathrm{KQ} \simeq \tilde{\mathrm{H}}\mathbb{Z}[\frac{1}{p}]$ by [2, Theorem 16], while $\tilde{f}_i \mathrm{KQ} \in \mathcal{SH}(K)[\frac{1}{p}]_{\geq i}$ by construction. As a consequence $\tilde{f}_0 \mathrm{KQ}$ satisfies Assumption 4.2.1 with $\tau_0 \tilde{f}_0 \mathrm{KQ} \simeq \tau_0(S)$.

6. The E -based Adams–Novikov spectral sequence

In this section, we briefly recall the construction of the Adams–Novikov spectral sequence based on a homotopy commutative algebra in \mathcal{M} . This section works in any symmetric monoidal stable ∞ -category \mathcal{M} .

6.1 – Construction of the spectral sequence

Let E be a homotopy commutative algebra in \mathcal{M} with multiplication $\mu : E \wedge E \rightarrow E$ and unit $e : \mathbf{1} \rightarrow E$. We start by considering the fiber sequence

$$(6.1) \quad \bar{E} \xrightarrow{\bar{e}} \mathbf{1} \xrightarrow{e} E$$

where $\bar{E} := \text{fib}(e : \mathbf{1} \rightarrow E)$. We set the notation $\bar{E}^1 = \bar{E}$ and $\bar{E}^0 = \mathbf{1}$. By induction, assuming we have already defined \bar{E}^n , we obtain a new fiber sequence by applying $-\wedge \bar{E}^n$ to the fiber sequence (6.1): we get the fiber sequence

$$(6.2) \quad \bar{E} \wedge \bar{E}^n \xrightarrow{\bar{e} \wedge \text{id}} \mathbf{1} \wedge \bar{E}^n \xrightarrow{e \wedge \text{id}} E \wedge \bar{E}^n.$$

We set $\bar{E}^{n+1} := \bar{E} \wedge \bar{E}^n$ and as well $W_n := E \wedge \bar{E}^n$. Furthermore, we name the maps $\bar{e}^{n+1} := \bar{e} \wedge \text{id}_{\bar{E}^n}$ and $e^{n+1} := e \wedge \text{id}_{\bar{E}^n}$. This way we have produced a tower $\{\bar{E}^n\}_{n \in \mathbb{N}}$ over $\mathbf{1}$ fitting in the diagram

$$(6.3) \quad \begin{array}{ccccccc} \mathbf{1} = \bar{E}^0 & \xleftarrow{\bar{e}^1} & \bar{E}^1 & \xleftarrow{\bar{e}^2} & \bar{E}^2 & \xleftarrow{\bar{e}^3} & \dots \\ \downarrow e & \nearrow \text{dashed} & \downarrow e^2 & \nearrow \text{dashed} & \downarrow e^3 & \nearrow \text{dashed} & \\ E = W_0 & & E \wedge \bar{E}^1 = W_1 & & E \wedge \bar{E}^2 = W_2 & & \dots \end{array}$$

where each dashed arrow is pictured to remind that the triangle it bounds is a fiber sequence. Given any object X we can smash every part of the previous construction with X and get a tower $\{X \wedge \bar{E}^n, X \wedge \bar{e}^n\}_{n \in \mathbb{N}}$ over X and actually a whole diagram similar to (6.3).

6.1.1. We could use an exact couple coming from (6.3) to construct an Adams spectral sequence, but for our purposes it is more helpful to consider the tower under $\mathbf{1}$ of cofibers induced by (6.3). With this aim in mind we proceed. The techniques of Section 3.1 allow us to upgrade (6.3) to a diagram $\mathbb{N}(\mathbb{N}^{\text{op}}) \rightarrow \mathcal{M}$, and thus to a fiber sequence of towers:

$$(6.4) \quad \bar{E}^\bullet \xrightarrow{\varepsilon^\bullet} \mathbf{1} \rightarrow \bar{E}_{\bullet-1}.$$

We visualize it as a commutative ladder of fiber sequences in \mathcal{M} :

$$\begin{array}{ccccc} \bar{E}^{n+1} & \xrightarrow{\varepsilon^{n+1}} & \mathbf{1} & \longrightarrow & \bar{E}_n \\ \bar{e}^{n+1} \downarrow & & \parallel & & \downarrow f_n \\ \bar{E}^n & \xrightarrow{\varepsilon^n} & \mathbf{1} & \longrightarrow & \bar{E}_{n-1}, \end{array}$$

where ε^n denotes an n -fold composition $\bar{e}^1 \circ \dots \circ \bar{e}^n$. Note that implicitly we have $\bar{E}_{-1} \simeq 0$ and $\bar{E}_0 \simeq E$. Moreover, we have equivalences $W_n = \text{cofib}(\bar{e}^{n+1}) \simeq \text{fib}(f_n)$ in \mathcal{M} , and in particular fiber sequences

$$(6.5) \quad W_n \xrightarrow{l_n} \bar{E}_n \xrightarrow{f_n} \bar{E}_{n-1} \xrightarrow{\partial_n} \Sigma^1 W_n.$$

We thus get a new diagram

$$(6.6) \quad \begin{array}{ccccccc} & & W_3 & & W_2 & & W_1 & & W_0 & & \\ & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \\ \dots & & \bar{E}_3 & \xrightarrow{f_3} & \bar{E}_2 & \xrightarrow{f_2} & \bar{E}_1 & \xrightarrow{f_1} & \bar{E}_0 & \xrightarrow{f_0} & 0 \end{array}$$

where again the dashed arrows are pictured to remind us that the triangles they bound are fiber sequences. Note that now the maps f_n form a tower under $\mathbf{1}$.

As we did above, given any object X we can build similar diagrams by applying $X \wedge -$ to (6.6). We obtain the following:

$$(6.7) \quad \begin{array}{ccccccc} & & X \wedge W_3 & & X \wedge W_2 & & X \wedge W_1 & & X \wedge W_0 & & \\ & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \\ \dots & & X \wedge \bar{E}_3 & \xrightarrow{f_3} & X \wedge \bar{E}_2 & \xrightarrow{f_2} & X \wedge \bar{E}_1 & \xrightarrow{f_1} & X \wedge \bar{E}_0 & \xrightarrow{f_0} & 0 \end{array}$$

Here we abuse the notation and keep denoting the maps involved in (6.7) with the same names used above in (6.6).

DEFINITION 6.1.2. The tower under X

$$\dots \xrightarrow{f_{n+1}} X \wedge \bar{E}_n \xrightarrow{f_n} \dots \xrightarrow{f_2} X \wedge \bar{E}_1 \xrightarrow{f_1} X \wedge \bar{E}_0 \rightarrow 0$$

is called the *standard E-Adams tower*. The *E-nilpotent completion* of X is the object

$$X_E^\wedge := \lim_{N(\mathbb{N}^{\text{op}})} (X \wedge \bar{E}_n) \in \mathcal{M}.$$

The natural map $\alpha_E(X) : X \rightarrow X_E^\wedge$ is called the *E-nilpotent completion map* of X .

6.1.3. For every object Y we can apply the functor $[Y, -]$ to (6.7) and get an exact couple

$$(6.8) \quad \begin{array}{ccc} [\Sigma^\bullet Y, X \wedge \bar{E}_\bullet] & \xrightarrow{j} & [\Sigma^\bullet Y, X \wedge \bar{E}_\bullet] \\ & \swarrow i \quad \searrow k & \\ & [\Sigma^\bullet Y, X \wedge W_\bullet] & \end{array}$$

Here the map

$$j : [\Sigma^p Y, X \wedge \bar{E}_n] \rightarrow [\Sigma^p Y, X \wedge \bar{E}_{n-1}]$$

is the natural map induced by f_n and has bi-degree $(0, -1)$; the map

$$k : [\Sigma^p Y, X \wedge \bar{E}_n] \rightarrow [\Sigma^{p-1} Y, X \wedge W_{n+1}]$$

is the natural map induced by the dashed arrow ∂_{n+1} and has bi-degree $(-1, 1)$. Finally, the map

$$i : [\Sigma^p Y, X \wedge W_n] \rightarrow [\Sigma^{p-1} Y, X \wedge \bar{E}_n]$$

is the map induced by l_n and has bi-degree $(0, 0)$.

6.1.4. The spectral sequence obtained from the exact couple (6.8) is the E -based Adams–Novikov spectral sequence. Note that this is an example of the general procedure described in [9, Chapter IX.4] for associating the so-called *homotopy spectral sequence* to a tower of fibrations under a given space. In our specific example, the tower we used is $\{X \wedge \bar{E}_n, f_n\}$.

7. Nilpotent resolutions

In this section, we introduce E -nilpotent (resp. strongly \mathcal{R} -nilpotent) resolutions associated with a homotopy commutative algebra E in \mathcal{M} (resp. a commutative algebra \mathcal{R} in \mathcal{M}^\heartsuit). This takes place in Section 7.1 (resp. 7.2). We use these constructions to describe a universal property for the Adams tower after passing to pro-objects, obtaining thus a more ductile construction of E -nilpotent completions. As an application we obtain an explicit description (see Theorem 7.3.5) for the E -nilpotent completion of a k -connected object. Throughout this section \mathcal{M} is a presentably symmetric monoidal stable ∞ -category, endowed with a left-complete multiplicative t -structure.

7.1 – E -nilpotent resolutions

DEFINITION 7.1.1. Let E be a homotopy commutative algebra in \mathcal{M} . We define the ∞ -category of E -nilpotent objects as the smallest full sub- ∞ -category $\text{Nilp}(E) \subseteq \mathcal{M}$ satisfying the following properties:

- (1) $E \in \text{Nilp}(E)$.
- (2) Given any $X \in \mathcal{M}$ and any $F \in \text{Nilp}(E)$, we have $X \wedge F \in \text{Nilp}(E)$.
- (3) $\text{Nilp}(E)$ has the 2-out-of-3 property on fiber sequences, i.e., given a fiber sequence $X \rightarrow Y \rightarrow Z$ in \mathcal{M} where any two of the three objects X, Y, Z are in $\text{Nilp}(E)$, the third is in $\text{Nilp}(E)$ as well.
- (4) $\text{Nilp}(E)$ is closed under retracts.

REMARK 7.1.2. If R is a homotopy algebra and M is a homotopy R -module, then the action map $R \wedge M \rightarrow M$ is split by the unit. So if R is in $\text{Nilp}(E)$, then M is E -nilpotent too.

LEMMA 7.1.3. *If E is a homotopy commutative algebra and X is any E -nilpotent object, then X is E -local.*

PROOF. The proof goes exactly as in [8, Lemma 3.8]. We filter $\text{Nilp}(E)$ by inductively constructed subcategories C_i . C_0 is defined as the full sub- ∞ -category of \mathcal{M} whose objects are equivalent to $E \wedge X$ for some $X \in \mathcal{M}$. If $i \geq 1$, we set C_i to be the full subcategory of \mathcal{M} of those objects that are equivalent to a retract of an object in C_{i-1} or an extension of objects in C_{i-1} . It is formal to check that the union of the C_i coincides with $\text{Nilp}(E)$. Indeed, thanks to Remark 7.1.2 we have that $C_0 \subset \text{Nilp}(E)$, and since E -nilpotent objects are closed under retractions and extensions, we get by induction that each of the C_i is contained in $\text{Nilp}(E)$. Now the C_i form an increasing sequence of subcategories of $\text{Nilp}(E)$ and we need to check that their union, which we denote by C , is the whole $\text{Nilp}(E)$. However this is clear: by construction C satisfies all the four axioms of Definition 7.1.1 so we must have $C \supseteq \text{Nilp}(E)$, and so $\text{Nilp}(E) = C$. For proving the E -locality: E -modules are E -local, so $C_0 \subseteq \text{Loc}(E)$; since E -local objects are closed under extensions and retractions, we have $C_i \subseteq \text{Loc}(E)$, and hence $\text{Nilp}(E) = \bigcup_i C_i \subseteq \text{Loc}(E)$. \blacksquare

DEFINITION 7.1.4. Let E be a homotopy commutative algebra of \mathcal{M} . An object X is called E -pre-nilpotent if X_E is E -nilpotent.

PROPOSITION 7.1.5. *Let E be a homotopy commutative algebra of \mathcal{M} . Then the following are equivalent:*

- (P1) $\mathbf{1}$ is E -pre-nilpotent, i.e., $\mathbf{1}_E$ is E -nilpotent.
- (P2) For every object X , $\mathbf{1}_E \wedge X$ is E -nilpotent.
- (P3) Every object X is E -pre-nilpotent, i.e., X_E is E -nilpotent for every X .
- (P4) $\text{Nilp}(E) = \text{Loc}(E)$.

Moreover, the following are equivalent:

(S1) For every $X \in \mathcal{M}$, the map $\lambda_E(\mathbf{1}) \wedge \text{id}_X : X \rightarrow \mathbf{1}_E \wedge X$ is an E -localization of X .

(S2) The multiplication map of the E -local sphere $\mathbf{1}_E \wedge \mathbf{1}_E \rightarrow \mathbf{1}_E$ is an equivalence and the natural inequality $\langle E \rangle \leq \langle \mathbf{1}_E \rangle$ is an equality.

In addition, statement (P2) implies (S1). Furthermore, if E has a multiplication map $E \wedge E \rightarrow E$ which is an equivalence, then the unit $e : \mathbf{1} \rightarrow E$ is a localization map $\lambda_E(\mathbf{1})$, and condition (P1) holds for E .

PROOF. We start by observing that a localization map $\lambda_E(X)$ factors as

$$(7.1) \quad \begin{array}{ccc} X & \xrightarrow{\lambda_E(X)} & X_E \\ \lambda_E(\mathbf{1}) \wedge \text{id}_X \downarrow & \nearrow \tilde{\lambda}(X) & \\ \mathbf{1}_E \wedge X & & \end{array}$$

Since all the maps in the diagram are E -equivalences, $\tilde{\lambda}(X)$ is an equivalence if and only if $\mathbf{1}_E \wedge X$ is E -local.

E -nilpotent objects are closed under smashing with arbitrary objects, so that $\mathbf{1}_E$ is E -nilpotent if and only if for every $X \in \mathcal{M}$, $\mathbf{1}_E \wedge X$ is E -nilpotent too (i.e., (P1) is equivalent to (P2)). By Lemma 7.1.3, if $\mathbf{1}_E \wedge X$ is E -nilpotent, then it is E -local and hence $\tilde{\lambda}(X)$ is an equivalence in view of (7.1) (i.e., (P2) implies (P3)). Clearly (P3) implies (P1) and (P3) is equivalent to (P4).

Using Lemma 7.1.3, we immediately deduce that (P2) implies (S1).

We now prove that (S1) is equivalent to (S2). By applying (S1) to $X = \mathbf{1}_E$, in view of (7.1), we deduce that the multiplication map $\tilde{\lambda}(\mathbf{1}_E)$ of $\mathbf{1}_E$ is an equivalence. On the other hand, by smashing the fiber sequence

$$E\mathbf{1} \rightarrow \mathbf{1} \rightarrow \mathbf{1}_E$$

with an object X , we deduce that if X is $\mathbf{1}_E$ -acyclic, then it is also E -acyclic. This means that $\langle E \rangle \leq \langle \mathbf{1}_E \rangle$. The reverse inequality is immediate from (S1), so (S1) implies (S2). Assume now (S2). Since the multiplication of $\mathbf{1}_E$ is an equivalence, for every $X \in \mathcal{M}$ the map $\lambda_E(\mathbf{1}) \wedge \text{id}_X : X \rightarrow \mathbf{1}_E \wedge X$ is an $\mathbf{1}_E$ -localization of X ; however $\langle \mathbf{1}_E \rangle = \langle E \rangle$ so that (S2) implies (S1).

If E is a homotopy commutative algebra with the property that the multiplication $E \wedge E \rightarrow E$ is an equivalence, then the unit $e : \mathbf{1} \rightarrow E$ is an E -equivalence. Since E is E -nilpotent, and thus E -local, we conclude. \blacksquare

DEFINITION 7.1.6. We say that an object E induces a smashing localization if the map $\tilde{\lambda}(X) : \mathbf{1}_E \wedge X \rightarrow X_E$ of (7.1) is an equivalence in \mathcal{M} .

EXAMPLE 7.1.7. The objects $\mathbf{1}^{\mathcal{B}}$ appearing in Proposition 3.4.3 induce smashing localizations of \mathcal{M} .

Let S be the spectrum of a field (although this argument works over more general bases), $\mathcal{M} = \mathcal{SH}(S)$, and $E = \mathbb{H}\mathbb{Q}$ be the spectrum representing Voevodsky's motivic cohomology with rational coefficients. Combining [11, Proposition 14.1.6 with Corollary 16.1.7], we deduce that the multiplication map of $\mathbb{H}\mathbb{Q}$ is an equivalence. In particular, the localization at $\mathbb{H}\mathbb{Q}$ is smashing.

LEMMA 7.1.8. *Let E be an object inducing a smashing localization, and let $\lambda_E(\mathbf{1}) : \mathbf{1} \rightarrow \mathbf{1}_E$ be an E -localization of $\mathbf{1}$. Then the forgetful functor $U_{\mathbf{1}_E} : \mathcal{M}od_{\mathbf{1}_E}(\mathcal{M}) \rightarrow \mathcal{M}$ factors through an equivalence $U_{\mathbf{1}_E} : \mathcal{M}od_{\mathbf{1}_E}(\mathcal{M}) \rightarrow \text{Loc}_E$.*

PROOF. Since $\mathbf{1}_E$ -modules are $\mathbf{1}_E$ -local and $\text{Loc}_{\mathbf{1}_E} = \text{Loc}_E$ by (S2), we have the desired factorization of $U_{\mathbf{1}_E}$ through the inclusion $\text{Loc}_E \subseteq \mathcal{M}$. Note that every E -local object is in the underlying object of a $\mathbf{1}_E$ -module, since by definition $\lambda_E(X) : X \rightarrow X_E \simeq \mathbf{1}_E \wedge X$ is an equivalence. In order to show that U is fully-faithful, we just need to see that for every $\mathbf{1}_E$ -module X , the natural ‘‘action map’’ $\alpha : F_{\mathbf{1}_E} U_{\mathbf{1}_E}(X) \rightarrow X$ is an equivalence. Indeed, we have a commutative diagram of spaces

$$\begin{array}{ccc} \text{Map}_{\mathbf{1}_E}(X, Y) & \xrightarrow{\quad\quad\quad} & \text{Map}(U_{\mathbf{1}_E}(X), U_{\mathbf{1}_E}(Y)) \\ & \searrow \alpha^* & \nearrow \simeq \\ & \text{Map}_{\mathbf{1}_E}(F_{\mathbf{1}_E} U_{\mathbf{1}_E}(X), Y) & \end{array}$$

where $\text{Map}_{\mathbf{1}_E}(-, -)$ denotes the mapping space in $\mathbf{1}_E$ -modules. Checking that α is an equivalence can be done after forgetting to \mathcal{M} , where α has a right inverse induced by the unit $\lambda_E(\mathbf{1}) \wedge \text{id} : \mathbf{1} \wedge U_{\mathbf{1}_E}(X) \rightarrow \mathbf{1}_E \wedge U_{\mathbf{1}_E}(X)$. However $\lambda_E(\mathbf{1}) \wedge \text{id}$ is an equivalence, since E is smashing and $U_{\mathbf{1}_E}(X)$ is E -local, so α is an equivalence too. \blacksquare

7.1.9. Let E be a homotopy commutative algebra. We wish to point out how Definition 6.1.2 and Lemma 7.1.3 imply that, for every $X \in \mathcal{M}$, the E -nilpotent completion X_E^\wedge is E -local. Indeed,

$$X_E^\wedge = \lim(\cdots \xrightarrow{f_{n+1}} X \wedge \bar{E}_n \xrightarrow{f_n} \cdots \rightarrow X \wedge \bar{E}_0 \rightarrow 0),$$

and each of the maps in the tower sits in a fiber sequence

$$E \wedge \bar{E}^n \wedge X \rightarrow \bar{E}_n \wedge X \xrightarrow{f_n} \bar{E}_{n-1} \wedge X$$

that we have deduced from (6.5). As a consequence, by induction, each of the terms in the tower is E -nilpotent, hence E -local, and thus X_E^\wedge is E -local too. In particular, the

map $\alpha_E(X) : X \rightarrow X_E^\wedge$ factors as

$$(7.2) \quad \begin{array}{ccc} X & \xrightarrow{\alpha_E(X)} & X_E^\wedge \\ & \searrow \lambda_E(X) & \nearrow \beta_E(X) \\ & X_E & \end{array}$$

It follows that $\alpha_E(X)$ is an E -equivalence if and only if the induced map $\beta_E(X)$ is an equivalence.

7.1.10. We wish to point out another fact. On the one hand, if $X \rightarrow Y$ is an E -equivalence, then it induces an equivalence of the standard E -Adams towers (Definition 6.1.2) associated to X and Y , so that the natural map induced on homotopy inverse limits $X_E^\wedge \rightarrow Y_E^\wedge$ is an equivalence. On the other hand, the composition of $\alpha_E(X)$ with the projection to the 0-th term of the tower

$$X \rightarrow X_E^\wedge \rightarrow X \wedge \bar{E}_0 = X \wedge E$$

is identified with $\text{id}_X \wedge e$, where $e : \mathbf{1} \rightarrow E$ is the unit of the algebra E . Thus, the map $\alpha_E(X) \wedge E : X \wedge E \rightarrow X_E^\wedge \wedge E$ has a retraction which is functorial in X . So if $f : X \rightarrow Y$ is a map inducing an equivalence on E -nilpotent completions $X_E^\wedge \rightarrow Y_E^\wedge$, then f is an E -equivalence. We conclude that $\alpha_E(X)$ is an E -equivalence if and only if the induced map

$$\alpha_E(X)_E^\wedge : X_E^\wedge \rightarrow (X_E^\wedge)_E^\wedge$$

is an equivalence.

DEFINITION 7.1.11. For an object $X \in \mathcal{M}$, an E -nilpotent resolution of X is a tower of objects under X ,

$$X \rightarrow \cdots \rightarrow X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_0,$$

satisfying the following two properties:

- (1) $X_n \in \text{Nilp}(E)$ for every $n \in \mathbb{N}$.
- (2) For any $Y \in \text{Nilp}(E)$ the map of pro-objects $\{X\} \rightarrow \{X_\bullet\}$ defined by the tower induces an equivalence

$$\text{Map}_{\text{Pro}(\mathcal{M})}(\{X_\bullet\}, \{Y\}) \rightarrow \text{Map}_{\text{Pro}(\mathcal{M})}(\{X\}, \{Y\}),$$

where X and Y are considered as constant pro-objects.

7.1.12. Recall that for pro-objects X_\bullet, Y_\bullet in an ∞ -category \mathcal{C} ,

$$\mathrm{Map}_{\mathrm{Pro}(\mathcal{M})}(X_\bullet, Y_\bullet) \simeq \lim_m \mathrm{colim}_n \mathrm{Map}_{\mathcal{M}}(X_n, Y_m)$$

(cf. Lemma A.1.2). In our situation, by applying homotopy groups (of the geometric realization) to the previous formula, we get that a tower $X \rightarrow X_\bullet$ of E -nilpotent objects under a given $X \in \mathcal{M}$ is an E -nilpotent resolution if and only if for every E -nilpotent object $Y \in \mathcal{M}$ the induced map

$$\pi_i \mathrm{colim}_n \mathrm{Map}_{\mathcal{M}}(X_n, Y) \rightarrow \pi_i \mathrm{Map}_{\mathcal{M}}(X, Y)$$

is an isomorphism for all $i \in \mathbb{N}$. Since taking homotopy groups commutes with filtered colimits, and since E -nilpotent objects are closed under shifts, this is equivalent to asking that for every E -nilpotent object $Y \in \mathcal{M}$ the natural map

$$\mathrm{colim}_n [X_n, Y] \rightarrow [X, Y]$$

is an isomorphism. The definition we have given is thus compatible with the classical definition for the stable homotopy category.

PROPOSITION 7.1.13. *Let X be any object of \mathcal{M} . Then:*

- (1) *The standard Adams tower $\bar{E}_\bullet \wedge X$ is an E -nilpotent resolution of X .*
- (2) *The pro-object under X associated with an E -nilpotent resolution of X is unique up to a contractible space of choices.*
- (3) *If $X \rightarrow X_\bullet$ is an E -nilpotent resolution of X , then there is an equivalence $\lim_{\mathbb{N}(\mathbb{N}^{\mathrm{op}})} X_\bullet \simeq X_E^\wedge$ in \mathcal{M}_X .*

PROOF. We start with (1). As we already observed in Section 7.1.9, the terms $\bar{E}_n \wedge X$ of the tower are E -nilpotent. Let Y be any object of \mathcal{M} . By smashing the fiber sequence of towers (6.4) with X , we get a commutative ladder of long exact sequences

$$\begin{array}{ccccccc} \cdots & \longrightarrow & [\bar{E}_n \wedge X, Y] & \longrightarrow & [X, Y] & \longrightarrow & [\bar{E}^{n+1} \wedge X, Y] \longrightarrow \cdots \\ & & \uparrow (f_n \wedge \mathrm{id}_X)^* & & \parallel & & \uparrow (\bar{e}^{n+1} \wedge \mathrm{id}_X)^* \\ \cdots & \longrightarrow & [\bar{E}_{n-1} \wedge X, Y] & \longrightarrow & [X, Y] & \longrightarrow & [\bar{E}^n \wedge X, Y] \longrightarrow \cdots \end{array}$$

We deduce that we only need to show that $\varinjlim_n [\bar{E}^n \wedge X, Y] = 0$ for every E -nilpotent object Y . We will proceed by induction on the family of subcategories C_i that we used in the proof of Lemma 7.1.3. Assume thus that $Y \in C_0$, i.e., that $Y \simeq E \wedge Z$ for some $Z \in \mathcal{M}$. We will show that the transition maps in the colimit vanish, hence the colimit

vanishes too. For this we look at the fiber sequence (6.2): it gives a long exact sequence

$$\begin{aligned} \cdots [E \wedge \bar{E}^n \wedge X, E \wedge Z] &\xrightarrow{(e \wedge \text{id})^*} [\bar{E}^n \wedge X, E \wedge Z] \\ &\xrightarrow{(\bar{e}^{n+1} \wedge \text{id}_X)^*} [\bar{E}^{n+1} \wedge X, E \wedge Z] \cdots, \end{aligned}$$

where the maps $(e \wedge \text{id})^*$ are surjective since $E \wedge Z$ is a homotopy E -module. The transition maps in the direct limit are thus 0. Now observe that the property $\varinjlim_n [\bar{E}^n \wedge X, Y] = 0$ is stable in the Y variable under retracts and extensions. This implies that if every $Y \in C_{i-1}$ satisfies this property, then every $Y \in C_i$ does as well. Since the union of the C_i exhausts $\text{Nilp}(E)$, the first point is done.

For (2): existence is (1), and for uniqueness we argue as follows. Let $X_\bullet \xleftarrow{p} X \xrightarrow{q} Y_\bullet$ be E -nilpotent resolutions. Then we have a natural commutative square

$$\begin{array}{ccc} \text{Map}_{\text{Pro}(\mathcal{M})}(\{X_\bullet\}, \{Y_\bullet\}) & \xrightarrow{\cong} & \lim_n \text{Map}_{\text{Pro}(\mathcal{M})}(\{X_\bullet\}, \{Y_n\}) \\ \downarrow p^* & & \downarrow p^* \\ \text{Map}_{\text{Pro}(\mathcal{M})}(\{X\}, \{Y_\bullet\}) & \xrightarrow{\cong} & \lim_n \text{Map}_{\text{Pro}(\mathcal{M})}(\{X\}, \{Y_n\}), \end{array}$$

where the right vertical map is an equivalence, since each Y_n is E -nilpotent. This is enough to conclude that the diagram of pro-objects $\{X_\bullet\} \xleftarrow{p} \{X\} \xrightarrow{q} \{Y_\bullet\}$ can be filled essentially in a unique way to a 2-simplex

$$\begin{array}{ccc} & \{X\} & \\ p \swarrow & & \searrow q \\ \{X_\bullet\} & \xrightarrow{f} & \{Y_\bullet\} \end{array}$$

as the next lemma shows. The same argument with X_\bullet and Y_\bullet interchanged implies that any choice for f must be an equivalence.

Point (3) follows by combining (1) and (2) with the observation that the operation of taking inverse limits factors through pro-objects. \blacksquare

LEMMA 7.1.14. *Let $X \xleftarrow{p} Z \xrightarrow{q} Y$ be a diagram in an ∞ -category \mathcal{C} , and assume that composition with p induces an equivalence $\text{Map}(X, Y) \rightarrow \text{Map}(Z, Y)$. Then there exists a unique 2-simplex of \mathcal{C} extending the horn (p, q) up to a contractible space of choices.*

PROOF. Consider the over-category $\mathcal{C}_{p/}$. The restriction along the source and target of p respectively induces functors

$$\mathcal{C}_{Z/} \xleftarrow{\sigma} \mathcal{C}_{p/} \xrightarrow[\cong]{\tau} \mathcal{C}_{X/},$$

where σ is a left fibration [21, Proposition 2.1.2.1], and τ is a trivial Kan fibration [21, Proposition 2.1.2.5]. Moreover, the three categories are themselves total spaces of left fibrations over \mathcal{C} , and both σ and τ commute with the projection to \mathcal{C} . When we take fibers over $Y \in \mathcal{C}$, we get

$$\mathrm{Map}^L(Z, Y) \xleftarrow{\sigma_Y} (\mathcal{C}_{p/})_Y \xrightarrow[\cong]{\tau_Y} \mathrm{Map}^L(X, Y),$$

where $\mathrm{Map}^L(-, -)$ denotes the left mapping space. Here σ_Y is a left fibration over a Kan complex, and thus it is a Kan fibration by [21, Lemma 2.1.3.3]. Similarly, τ_Y is again a trivial Kan fibration. Our assumption then implies that σ_Y is a trivial Kan fibration. By definition of the left mapping space, the zero simplices of $\mathrm{Map}^L(Z, Y)$ are exactly the arrows $Z \rightarrow Y$ of \mathcal{C} , while the zero simplices of $(\mathcal{C}_{p/})_Y$ are commutative triangles

$$\begin{array}{ccc} & Z & \\ & \swarrow & \searrow \\ X & \xrightarrow{p} & Y. \end{array}$$

In particular, the fiber of σ_Y over q is contractible, which is what we wanted to prove. ■

7.2 – Strongly \mathcal{R} -nilpotent resolutions

LEMMA 7.2.1. *Let \mathcal{R} be a commutative algebra in \mathcal{M}^\heartsuit with the property that the multiplication map of $\mu_{\mathcal{R}} : \mathcal{R} \otimes^\heartsuit \mathcal{R} \rightarrow \mathcal{R}$ of \mathcal{R} is an isomorphism. Then:*

- (1) *For every \mathcal{R} -module \mathcal{M} in \mathcal{M}^\heartsuit , the action map $\mathcal{R} \otimes^\heartsuit \mathcal{M} \rightarrow \mathcal{M}$ is an isomorphism. In particular, an object \mathcal{M} of \mathcal{M}^\heartsuit has at most one \mathcal{R} -module structure.*
- (2) *Every map $\phi : \mathcal{M} \rightarrow \mathcal{N}$ in \mathcal{M}^\heartsuit where \mathcal{M} and \mathcal{N} are \mathcal{R} -modules is \mathcal{R} -linear. In particular, the category of \mathcal{R} -modules is a full subcategory of \mathcal{M}^\heartsuit .*

PROOF. For (1) observe that, given an \mathcal{R} -module \mathcal{M} in \mathcal{M}^\heartsuit , we have a co-equalizer diagram defining the monoidal product on the category of \mathcal{R} -modules:

$$\mathcal{R} \otimes^\heartsuit \mathcal{R} \otimes^\heartsuit \mathcal{M} \begin{array}{c} \xrightarrow{1 \otimes a} \\ \xrightarrow[\mu_{\mathcal{R}} \otimes 1]{} \end{array} \mathcal{R} \otimes^\heartsuit \mathcal{M} \xrightarrow{q} \mathcal{R} \otimes_{\mathcal{R}}^\heartsuit \mathcal{M}.$$

Moreover, the action map $a : \mathcal{R} \otimes^\heartsuit \mathcal{M} \rightarrow \mathcal{M}$ induces an isomorphism $\bar{a} : \mathcal{R} \otimes_{\mathcal{R}}^\heartsuit \mathcal{M} \rightarrow \mathcal{M}$. The map $\mathcal{R} \otimes^\heartsuit \mathcal{M} \rightarrow \mathcal{R} \otimes^\heartsuit \mathcal{R} \otimes^\heartsuit \mathcal{M}$ defined by $r \otimes m \mapsto r \otimes 1 \otimes m$ is an inverse of both $\mu_{\mathcal{R}}$ and a , so that $1 \otimes a = \mu_{\mathcal{R}} \otimes 1$, thus q is an isomorphism. In particular, every \mathcal{R} -module \mathcal{M} is isomorphic to the free \mathcal{R} -module on the object \mathcal{M} .

Point (2) follows by combining the free-forget adjunction and point (1). ■

LEMMA 7.2.2. *Assume E is a homotopy commutative algebra satisfying Assumption 4.2.1. Then $\mathcal{R} = \tau_0(E) \in \mathcal{M}^\heartsuit$ and its multiplication map $\mathcal{R} \otimes^\heartsuit \mathcal{R} \rightarrow \mathcal{R}$ is an isomorphism.*

PROOF. Surely $\tau_0(E)$ is a commutative algebra in \mathcal{M}^\heartsuit by the multiplicative properties of the Postnikov tower (see Section 2.1.5). Let us start from the special case $J = \emptyset$, so that $\mathcal{R} = \tau_0 E \simeq \tau_0(\mathbf{1})/(f_1, \dots, f_r)$ for some $f_i : L_i \rightarrow \mathbf{1}$. In this case, the map

$$\sum_{i=1}^r f_i : \bigoplus_i L_i \rightarrow \mathbf{1}$$

induces

$$\phi := \sum_{i=1}^r \ell_{f_i}(\tau_0 \mathbf{1}) : \bigoplus_i L_i \wedge \tau_0(\mathbf{1}) \rightarrow \tau_0(\mathbf{1})$$

in \mathcal{M}^\heartsuit . Since $\tau_0 E \simeq \text{coker}(\phi)$, Lemma 4.3.4 implies that the multiplication of \mathcal{R} is an isomorphism.

Let us now consider the case $J \neq \emptyset$. Denote by \mathcal{C} the object $\tau_0(\mathbf{1})/\text{Im}(\phi)$ so that we already know that $m : \mathcal{C} \otimes^\heartsuit \mathcal{C} \rightarrow \mathcal{C}$ is an isomorphism by the above argument, and that $\tau_0(E) \simeq \tau_0(\mathcal{C}[\mathcal{J}^{-1}])$ where $\mathcal{J} = \{g_j\}_{j \in J}$. Now by Corollary 3.4.6 the functor $[\mathcal{J}^{-1}]$ is right t -exact, so we have a natural equivalence

$$\tau_0(\mathcal{C}[\mathcal{J}^{-1}]) \otimes^\heartsuit \tau_0(\mathcal{C}[\mathcal{J}^{-1}]) \simeq \tau_0((\mathcal{C} \otimes^\heartsuit \mathcal{C})[\mathcal{J}^{-1}])$$

under which the multiplication of $\tau_0(\mathcal{C}[\mathcal{J}^{-1}])$ is identified with the map $\tau_0(m[\mathcal{J}^{-1}])$, which is an equivalence by the previous part. ■

7.2.3. In the rest of the section we fix a commutative algebra \mathcal{R} in \mathcal{M}^\heartsuit with the property that its multiplication is an isomorphism.

DEFINITION 7.2.4. We say that an object \mathcal{M} of \mathcal{M}^\heartsuit is *strongly \mathcal{R} -nilpotent* if it has a finite filtration $\mathcal{M} = \mathcal{M}_0 \supseteq \mathcal{M}_1 \supseteq \dots \supseteq \mathcal{M}_r$ in \mathcal{M}^\heartsuit such that $\mathcal{M}_i/\mathcal{M}_{i+1}$ has an \mathcal{R} -module structure for every i . An object X of \mathcal{M} is called *strongly \mathcal{R} -nilpotent* if for each $k \in \mathbb{Z}$ the homotopy object $\tau_k(X)$ is strongly \mathcal{R} -nilpotent and there exist integers s and t such that $X \simeq P_s(X) \simeq P^t(X)$. We denote the full subcategory of strongly \mathcal{R} -nilpotent objects of \mathcal{M} by $\text{SNilp}(\mathcal{R})$.

7.2.5. Observe that if $\mathcal{M} \in \mathcal{M}^\heartsuit$ is strongly \mathcal{R} -nilpotent, then it is strongly \mathcal{R} -nilpotent as an object of \mathcal{M} ; conversely any strongly \mathcal{R} -nilpotent object of \mathcal{M} which is concentrated in degree 0 for the t -structure is a strongly \mathcal{R} -nilpotent object of \mathcal{M}^\heartsuit . If an object X in \mathcal{M} is strongly \mathcal{R} -nilpotent, then it is \mathcal{R} -nilpotent in the sense of Definition 7.1.1. Indeed, since X is bounded in the t -structure of \mathcal{M} , X is an iterated extension of finitely

many of its homotopy objects $\Sigma^i \tau_i(X)$ which are, on their turn, iterated extensions of \mathcal{R} -modules. In particular, if E is a homotopy commutative algebra in $\mathcal{M}_{\geq 0}$ and $\tau_0(E) \simeq \mathcal{R}$, then any strongly \mathcal{R} -nilpotent object X is also E -nilpotent.

DEFINITION 7.2.6. Let X be an object in \mathcal{M} . A *strongly \mathcal{R} -nilpotent resolution* of X is a tower of objects under X ,

$$X \rightarrow \cdots \rightarrow X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_0,$$

satisfying the following two properties:

- (1) $X_n \in \text{SNilp}(\mathcal{R})$ for every $n \in \mathbb{N}$.
- (2) For any $Y \in \text{SNilp}(\mathcal{R})$ the induced map of pro-objects $\{X\} \rightarrow \{X_\bullet\}$ defined by the tower induces an equivalence

$$\text{Map}_{\text{Pro}(\mathcal{M})}(\{X_\bullet\}, \{Y\}) \rightarrow \text{Map}_{\text{Pro}(\mathcal{M})}(\{X\}, \{Y\}),$$

where X and Y are considered as constant pro-objects.

We now prepare for an analogue of Proposition 7.1.13 for strongly \mathcal{R} -nilpotent resolutions.

LEMMA 7.2.7. *If $\mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{O} \rightarrow \mathcal{P} \rightarrow \mathcal{Q}$ is an exact sequence in \mathcal{M}^\heartsuit where $\mathcal{M}, \mathcal{N}, \mathcal{P}, \mathcal{Q}$ are strongly \mathcal{R} -nilpotent, then \mathcal{O} is strongly \mathcal{R} -nilpotent too.*

PROOF. By breaking up the exact sequence in shorter pieces, the statement follows by combining Lemmas 7.2.8 and 7.2.9. ■

LEMMA 7.2.8. *If $0 \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{O} \rightarrow 0$ is a short exact sequence in \mathcal{M}^\heartsuit and both \mathcal{M} and \mathcal{O} are strongly \mathcal{R} -nilpotent, then \mathcal{N} is strongly \mathcal{R} -nilpotent too.*

PROOF. A suitable filtration on \mathcal{N} can be obtained by combining the filtration on \mathcal{M} and the pre-image in \mathcal{N} of the filtration on \mathcal{O} . ■

LEMMA 7.2.9. *If $\phi : \mathcal{C}^0 \rightarrow \mathcal{C}^1$ is a map in \mathcal{M}^\heartsuit and both \mathcal{C}^0 and \mathcal{C}^1 are strongly \mathcal{R} -nilpotent, then both $\mathcal{H}^0 = \text{Ker}\phi$ and $\mathcal{H}^1 = \text{Coker}\phi$ are strongly \mathcal{R} -nilpotent too.*

PROOF. Up to increasing the length of the filtrations \mathcal{C}_i^0 and \mathcal{C}_i^1 we can assume that ϕ respects the filtrations. As a consequence $(\mathcal{C}^\bullet, \phi)$ is a filtered complex in \mathcal{M}^\heartsuit . With respect to this filtration we use the spectral sequence for filtered complexes. By assumption, the terms in the E_2 -page are \mathcal{R} -modules and the differentials are \mathcal{R} -linear by Lemma 7.2.1. Hence we gain a finite filtration on the cohomology of \mathcal{C}^\bullet with associated graded \mathcal{R} -module quotients. ■

LEMMA 7.2.10. *Let $f : L \rightarrow \tau_0(\mathbf{1})$ be a morphism in \mathcal{M} , where L is a tif object. Denote by $\cdot f$ the induced map $\mathcal{R} \wedge L \rightarrow \mathcal{R}$ in \mathcal{M}^\heartsuit given by Lemma 4.3.4, and let $\mathcal{S} := \text{coker}(\cdot f)$. Then \mathcal{S} is a commutative \mathcal{R} -algebra in \mathcal{M}^\heartsuit and its multiplication map $\mu_{\mathcal{S}} : \mathcal{S} \otimes^\heartsuit \mathcal{S} \rightarrow \mathcal{S}$ is an isomorphism. Furthermore, for every strongly \mathcal{R} -nilpotent object \mathcal{M} of \mathcal{M}^\heartsuit both kernel and cokernel of the multiplication by f on \mathcal{M} are strongly \mathcal{S} -nilpotent.*

PROOF. First of all, since $L \in \mathcal{M}_{\geq 0}$, f induces a map $\tau_0(L) \rightarrow \tau_0(\mathbf{1}) \rightarrow \mathcal{R}$, and $\tau_0(L)$ is a strongly dualizable object of \mathcal{M}^\heartsuit , so Lemma 4.3.4 fully applies. We deduce that f extends uniquely to the map $\cdot f : \mathcal{R} \wedge L \rightarrow \mathcal{R}$, the commutative algebra structure of \mathcal{R} in \mathcal{M}^\heartsuit descends uniquely to a commutative algebra structure on \mathcal{S} , such that the projection $\mathcal{R} \rightarrow \mathcal{S}$ is an algebra map. Furthermore, the multiplication induced on \mathcal{S} is an isomorphism.

Let $0 = \mathcal{M}_n \subseteq \mathcal{M}_{n-1} \subseteq \dots \subseteq \mathcal{M}_0 = \mathcal{M}$ be a filtration of \mathcal{M} by objects \mathcal{M}_i of \mathcal{M}^\heartsuit whose associated graded pieces are \mathcal{R} -modules, and denote by $r_f(\mathcal{M}) : \mathcal{M} \wedge L \rightarrow \mathcal{M}$ the induced right multiplication by f on \mathcal{M} . If $n = 1$, then \mathcal{M} is an \mathcal{R} -module, $r_f(\mathcal{M})$ is automatically \mathcal{R} -linear by Lemma 7.2.1, and thus kernel and co-kernel of $r_f(\mathcal{M})$ are \mathcal{S} -modules by Lemma 4.3.4. If $n > 1$, one can proceed by induction. Indeed, since the multiplication map $r_f(\mathcal{M}) : \mathcal{M} \wedge L \rightarrow \mathcal{M}$ commutes with any map in \mathcal{M} by Section 3.2.3, it respects the filtration. As a consequence, for every integer $k \geq 1$, $r_f(\mathcal{M})$ induces a map of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{M}_{k-1} & \longrightarrow & \mathcal{M}_k & \longrightarrow & \mathcal{M}_k / \mathcal{M}_{k-1} \longrightarrow 0 \\ & & \downarrow r_f(\mathcal{M}) & & \downarrow r_f(\mathcal{M}) & & \downarrow r_f(\mathcal{M}) \\ 0 & \longrightarrow & \mathcal{M}_{k-1} & \longrightarrow & \mathcal{M}_k & \longrightarrow & \mathcal{M}_k / \mathcal{M}_{k-1} \longrightarrow 0. \end{array}$$

Thus, by combining the snake lemma together with Lemma 7.2.7 and the inductive assumption, we conclude. \blacksquare

7.3 – Relation between localizations and nilpotent completions

NOTATION 7.3.1. For the rest of the section we fix a homotopy commutative algebra E in $\mathcal{M}_{\geq 0}$, and we assume that the induced multiplication on $\tau_0(E)$ is an isomorphism. We also denote by \mathcal{R} the homotopy object $\tau_0 E$.

PROPOSITION 7.3.2. *Let X be an object of $\mathcal{M}_{\geq k}$ for some integer k . Then the tower $X \rightarrow P^\bullet(\bar{E}_\bullet \wedge X)$ is a strongly \mathcal{R} -nilpotent resolution of X .*

PROOF. We first need to check that $P^n(\bar{E}_n \wedge X) \in \text{SNilp}(\mathcal{R})$ for every $n \in \mathbb{Z}$. By the connectivity of X , each of the $P^n(\bar{E}_n \wedge X)$ is bounded in the t -structure, so we

only need to check that $\tau_k(\bar{E}_n \wedge X)$ is strongly \mathcal{R} -nilpotent for every pair of integers k, n . Recall that for every n in \mathbb{N} we have a fiber sequence of the form

$$(7.3) \quad X \wedge E \wedge \bar{E}^n \rightarrow X \wedge \bar{E}_n \xrightarrow{f_n} X \wedge \bar{E}_{n-1}$$

obtained from (6.5) by smashing with X . By construction, for $n = 0$, such a fiber sequence is the cone sequence of the identity of $X \wedge E$. In particular, for every $k \in \mathbb{Z}$, $\tau_k(X \wedge \bar{E}_0)$ is a $\tau_0 E$ -module and thus it is strongly \mathcal{R} -nilpotent ($\mathcal{R} = \tau_0 E$). We can now proceed by induction on n . Note that $\tau_k(X \wedge \bar{E}^n \wedge E)$ is an \mathcal{R} -module too and thus is strongly \mathcal{R} -nilpotent. This observation, once combined with the inductive assumption, allows to apply Lemma 7.2.7 to the long exact sequence of homotopy objects associated to the fiber sequence (7.3).

As a second step we need to prove that for any $Y \in \text{SNilp}(\mathcal{R})$ the projection $p : X \rightarrow P^{\leq \bullet}(\bar{E}_\bullet \wedge X)$ induces an equivalence

$$\text{Map}_{\text{Pro}(\mathcal{M})}(\{P^{\leq \bullet}(\bar{E}_\bullet \wedge X)\}, \{Y\}) \rightarrow \text{Map}_{\text{Pro}(\mathcal{M})}(\{X\}, \{Y\}).$$

However the factorization through the Adams tower $X \rightarrow \bar{E}_\bullet \wedge X$ induces a commutative diagram

$$\begin{array}{ccc} \text{colim}_n \text{Map}(P^n(\bar{E}_n \wedge X), Y) & \xleftarrow{\simeq} & \text{Map}_{\text{Pro}(\mathcal{M})}(\{P^{\leq \bullet}(\bar{E}_\bullet \wedge X)\}, \{Y\}) \\ \downarrow \pi_n & & \downarrow \pi \\ \text{colim}_n \text{Map}(\bar{E}_n \wedge X, Y) & \xleftarrow{\simeq} & \text{Map}_{\text{Pro}(\mathcal{M})}(\{\bar{E}_\bullet \wedge X\}, \{Y\}) \longrightarrow \text{Map}(X, Y), \end{array}$$

where the map π and the maps π_n are induced by the projection maps to the Postnikov truncations. The facts that Y is E -nilpotent (Section 7.2.5) and that the standard Adams tower $X \rightarrow \bar{E}_\bullet \wedge X$ is an E -nilpotent resolution (Proposition 7.1.13) imply that the rightmost horizontal map is an equivalence. We are left to show that π is an equivalence too. This follows from the fact that Y is bounded in the homotopy t -structure on \mathcal{M} , so π_n is an equivalence for large n . \blacksquare

PROPOSITION 7.3.3. *For every object $X \in \mathcal{M}_{\geq k}$, where $k \in \mathbb{Z}$, the following holds.*

- (1) *The Postnikov truncation of the standard Adams tower $X \rightarrow P^\bullet(\bar{E}_\bullet \wedge X)$ is a strongly \mathcal{R} -nilpotent resolution of X in \mathcal{M} .*
- (2) *The pro-object under X associated to a strongly \mathcal{R} -nilpotent resolution of X is unique up to a contractible space of choices.*
- (3) *If $X \rightarrow X_\bullet$ is a strongly \mathcal{R} -nilpotent resolution of X , there is an equivalence $\lim_{\mathbb{N}(\mathbb{N}^{\text{op}})} X_\bullet \simeq X_E^\wedge$ in \mathcal{M}_X .*

PROOF. Part (1) is Proposition 7.3.2. Part (2) works similarly to (2) in Proposition 7.1.13. For property (3) we combine the equivalence $\{X_\bullet\} \simeq \{P^\bullet(\bar{E}_\bullet \wedge X)\}$ of pro-objects under X coming from the previous points with the τ -equivalence (cf. Definition A.1.3) $\{P^\bullet(\bar{E}_\bullet \wedge X)\} \leftarrow \{\bar{E}_\bullet \wedge X\}$ of pro-objects under X induced by the projection to the tower of truncations. The inverse limit of a composition of these two maps induces an equivalence under X as required in the statement (cf. Section A.1.4). \blacksquare

LEMMA 7.3.4. *Consider a collection of tif objects L_1, \dots, L_r of \mathcal{M} . For every $i = 1, \dots, r$ let $f_i : L_i \rightarrow \tau_0(\mathbf{1})$ be a map in \mathcal{M} , and let \mathcal{R} be the commutative algebra in \mathcal{M}^\heartsuit defined by $\mathcal{R} = \tau_0(\mathbf{1})/(f_1, \dots, f_r)$. Then for every bounded-below object $X \in \mathcal{M}$, the tower $X \rightarrow P^\bullet(C(f_1^\bullet) \wedge \dots \wedge C(f_r^\bullet) \wedge X)$ is a strongly \mathcal{R} -nilpotent resolution of X .*

PROOF. We need to check that for every pair of integers k, n the homotopy objects $\tau_k(C(f_1^n) \wedge \dots \wedge C(f_r^n) \wedge X)$ are strongly \mathcal{R} -nilpotent. We accomplish this by induction on n , the base case being that of $n = 1$. Assume thus that $n = 1$. We proceed by induction on r . For $r = 0$ we have $\mathcal{R} = \tau_0(\mathbf{1})$, and thus the homotopy objects $\tau_k(X)$ are strongly $\tau_0(\mathbf{1})$ -nilpotent. When $r \geq 1$, we set $\mathcal{R}_a := \tau_0(\mathbf{1})/(f_1, \dots, f_a)$. We can apply Lemma 7.2.10 to the homotopy objects $\mathcal{M} := \tau_k(C(f_1) \wedge \dots \wedge C(f_a) \wedge X)$ and $\mathcal{M}' := \tau_{k-1}(C(f_1) \wedge \dots \wedge C(f_a) \wedge X)$, which are strongly \mathcal{R}_a -nilpotent by the inductive assumption. It follows that the external homotopy objects of the exact sequences (for varying k)

$$\begin{aligned} 0 \rightarrow \operatorname{coker}(r_{f_{a+1}}(\mathcal{M})) &\rightarrow \tau_k(C(f_1) \wedge \dots \wedge C(f_{a+1}) \wedge X) \\ &\rightarrow \ker(r_{f_{a+1}}(\mathcal{M}')) \rightarrow 0 \end{aligned}$$

are strongly $\mathcal{R}_a/(f_{a+1})$ -nilpotent. In particular, they are strongly \mathcal{R}_{a+1} -nilpotent, since $\mathcal{R}_a/(f_{a+1}) \simeq \mathcal{R}_{a+1}$, and by Lemma 7.2.8 we conclude that the central homotopy objects are strongly \mathcal{R}_{a+1} -nilpotent too. Given now any r -tuple of positive integers (n_1, \dots, n_r) , we can show that $\tau_k(C(f_1^{n_1}) \wedge \dots \wedge C(f_r^{n_r}) \wedge X)$ is strongly \mathcal{R} -nilpotent by induction, using the fiber sequences

$$L_i^{\wedge n_i - 1} \wedge C(f_i) \rightarrow C(f_i^{n_i}) \rightarrow C(f_i^{n_i - 1})$$

for $i = 1, \dots, r$ and Lemma 7.2.7 to reduce to the above base case.

Let us assume now that Y is a strongly \mathcal{R} -nilpotent object of \mathcal{M} . Let us denote by C_n the object $C(f_1^n) \wedge \dots \wedge C(f_r^n) \in \mathcal{M}$. We thus need to check that the map

$$\operatorname{colim}_n \operatorname{Map}(P^n(C_n \wedge X), Y) \rightarrow \operatorname{Map}(X, Y),$$

induced by the projection to the tower, is an equivalence. Since $\tau_k(Y)$ is strongly \mathcal{R} -nilpotent, each f_i acts on such a homotopy object nilpotently. It follows that there

is an integer N for which each f_i^N acts by 0 on each homotopy object of Y . Moreover, since Y is bounded in the t -structure, up to enlarging N we may actually assume that each f_i^k acts by 0 on Y whenever $k \geq N$. Since Y is bounded in the t -structure, we only need to check that the map

$$(7.4) \quad \operatorname{colim}_n \operatorname{Map}(C_n \wedge X, Y) \rightarrow \operatorname{Map}(X, Y)$$

induced by the projection to the tower $X \rightarrow C_\bullet \wedge X$ is an equivalence. If $r = 1$, then the fiber of the map of towers $X \rightarrow C_\bullet \wedge X$ is the tower $L^{\wedge \bullet} \wedge X$ with maps induced by $r_f(L^{\wedge n-1}) : L^{\wedge n} \rightarrow L^{\wedge n-1}$. From the above argument we readily deduce that

$$\operatorname{colim}_n \pi_k \operatorname{Map}(L^{\wedge n} \wedge X, Y) \simeq 0,$$

guaranteeing that the map (7.4) is an equivalence. In the case $r \geq 2$, one reduces to the case just proved using that by cofinality

$$\operatorname{colim}_{n_1, \dots, n_r} \operatorname{Map}(C(f_1^{n_1}) \wedge \dots \wedge C(f_r^{n_r}) \wedge X, Y) \simeq \operatorname{colim}_n \operatorname{Map}(C_n \wedge X, Y). \quad \blacksquare$$

THEOREM 7.3.5. *Let E be a homotopy commutative algebra of \mathcal{M} satisfying Assumption 4.2.1 in the special case that $J = \emptyset$. Then for every k -connected object of \mathcal{M} the natural map $\alpha_E(X) : X \rightarrow X_E^\wedge$ is an E -equivalence. In particular, the map $\beta_E(X) : X_E \rightarrow X_E^\wedge$ of (7.2) is an equivalence in \mathcal{M} .*

PROOF. Let $C_n := C(f_1^n) \wedge \dots \wedge C(f_r^n)$. Proposition 7.3.2 and Lemma 7.3.4 imply that both the tower $P^\bullet(\bar{E}_\bullet \wedge X)$ and the tower $P^\bullet(C_\bullet \wedge X)$ are strongly \mathcal{R} -nilpotent resolutions of X . Thanks to Proposition 7.3.3 there is an equivalence ϕ of their associated pro-objects under X . As a consequence we have an induced equivalence under X between their limits:

$$\psi : \lim_{\mathbf{N}(\mathbf{N}^{\text{op}})} P^\bullet(\bar{E}_\bullet \wedge X) \xrightarrow{\cong} \lim_{\mathbf{N}(\mathbf{N}^{\text{op}})} P^\bullet(C_\bullet \wedge X).$$

Since these limits are naturally identified with X_E^\wedge and $X_{f_1, \dots, f_r}^\wedge$ respectively, we get a commutative square of pro-objects under X :

$$(7.5) \quad \begin{array}{ccc} \{X_E^\wedge\} & \xrightarrow[\psi]{\cong} & \{X_{f_1, \dots, f_r}^\wedge\} \\ \downarrow & & \downarrow \\ \{P^{\leq \bullet}(\bar{E}_\bullet \wedge X)\} & \xrightarrow{\phi} & \{P^\bullet(C_\bullet \wedge X)\}, \end{array}$$

where the vertical maps are the natural projections.

After smashing the towers in (7.5) with E , we obtain a new commutative square of pro-objects:

$$(7.6) \quad \begin{array}{ccc} \{E \wedge X_E^\wedge\} & \longrightarrow & \{E \wedge X_{f_1, \dots, f_r}^\wedge\} \\ \downarrow & & \downarrow \\ \{E \wedge P^\bullet(\bar{E} \wedge X)\} & \longrightarrow & \{E \wedge P^\bullet(C \wedge X)\}. \end{array}$$

Here the lower horizontal map remains an equivalence of pro-objects. We claim that the right vertical map of (7.6) is a τ -equivalence of pro-objects.

For showing this claim, consider that the vertical map $X_{f_1, \dots, f_r}^\wedge \rightarrow \{P^\bullet(C \wedge X)\}$ on the right-hand side of (7.5) factors as the composition of two maps: the projection

$$(7.7) \quad \{X_{f_1, \dots, f_r}^\wedge\} \rightarrow \{C \wedge X\}$$

and the projection to the Postnikov tower

$$(7.8) \quad \{C \wedge X\} \rightarrow \{P^\bullet(C \wedge X)\}.$$

The map (7.8) is a τ -equivalence by Section A.1.4 and stays a τ -equivalence after smashing with E by Lemma A.1.7. Concerning (7.7), we observe that by construction we have a commutative diagram of pro-objects:

$$(7.9) \quad \begin{array}{ccc} X_{f_1, \dots, f_r}^\wedge & \longrightarrow & \{C_n \wedge X\} \\ & \nwarrow \chi & \uparrow \\ & & X. \end{array}$$

Since χ is an E -equivalence by Propositions 4.3.2 and 3.2.14, in order to finish the proof of the above claim we are left to show that the vertical map of (7.9) induces, after smashing with E , a τ -equivalence of pro-objects in \mathcal{M} .

To accomplish this task we consider the tower $F_\bullet^{(r)} = \text{fib}(X \rightarrow C \wedge X)$, and show that the pro-object $\{E \wedge F_\bullet^{(r)}\}$ is τ -equivalent to 0. For this, remember that in the tower C the transition maps

$$\psi_n : C_n = C(f_1^n) \wedge \cdots \wedge C(f_r^n) \rightarrow C(f_1^{n-1}) \wedge \cdots \wedge C(f_r^{n-1}) = C_{n-1}$$

are defined as $\psi_n = p_n(f_r) \circ \cdots \circ p_n(f_1)$: here for every integer n the maps $p_n(f_i)$ are induced by the maps p_n defined in Section 3.2.9 and displayed in (3.1). If $r = 1$, we have $F_\bullet^{(1)} = L^\wedge \wedge X$ and its transition maps are $r_{f_1}(L^{\wedge n-1}) \wedge X$, as in (3.1). Since r_{f_1} acts trivially on the towers of homotopy objects $\tau_k(E \wedge F_\bullet^{(1)})$, we deduce that $\{E \wedge F_\bullet^{(1)}\}$ is τ -equivalent to 0. If $r > 1$, one can argue by induction. Indeed, using

the fiber sequence $\text{fib}(\alpha) \rightarrow \text{fib}(\beta \circ \alpha) \rightarrow \text{fib}(\beta)$ associated to the composable arrows $a \xrightarrow{\alpha} b \xrightarrow{\beta} c$ of a stable ∞ -category, we can find a fiber sequence of towers

$$F_{\bullet}^{(s-1)} \rightarrow F_{\bullet}^{(s)} \rightarrow G_{\bullet}^{(s)}$$

having the following properties: the pro-object $\{E \wedge F_{\bullet}^{(s-1)}\}$ is τ -equivalent to 0 by the inductive assumption, and $\{E \wedge G_{\bullet}^{(s)}\}$ is τ -equivalent to 0 by the case $r = 1$ treated above. Hence, thanks to Corollary A.1.6, we conclude that (7.7) becomes a τ -equivalence after smashing with E .

Let us consider now the commutative square of towers

$$(7.10) \quad \begin{array}{ccc} X & \longrightarrow & X_E^\wedge \\ \downarrow & & \downarrow \\ \bar{E}_{\bullet} \wedge X & \longrightarrow & P^*(\bar{E}_{\bullet} \wedge X) \end{array}$$

and note that, in order to conclude, we only need to show that the map $E \wedge \alpha_E(X) : E \wedge X \rightarrow E \wedge X_E^\wedge$ is an equivalence. This map is the upper horizontal arrow of the diagram

$$(7.11) \quad \begin{array}{ccc} E \wedge X & \longrightarrow & E \wedge X_E^\wedge \\ \downarrow & & \downarrow \\ E \wedge (\bar{E}_{\bullet} \wedge X) & \longrightarrow & E \wedge P^*(\bar{E}_{\bullet} \wedge X) \end{array}$$

which is obtained from (7.10) by smashing with E . In (7.11), the right vertical map induces a τ -equivalence on pro-objects as it follows from the previous part of the argument. The lower horizontal map also induces a τ -equivalence of the associated pro-objects by Lemma A.1.7. Finally, the left vertical map of (7.11) induces a τ -equivalence too, as we now explain. We have a fiber sequence of towers

$$(7.12) \quad \bar{E}_{\bullet} \wedge X \rightarrow X \rightarrow \bar{E}_{\bullet-1} \wedge X$$

which we obtain from diagram (6.4) upon smashing with X . The left vertical map in the square (7.10) is the map induced by the right-hand side maps of (7.12). We claim that, after smashing (7.12) with E , the tower $\bar{E}_{\bullet} \wedge X$ on the left-hand side of (7.12) becomes τ -equivalent to zero. Indeed, by the very inductive definition of \bar{E}^n we have fiber sequences deduced from (6.2):

$$\bar{E}^{n+1} \wedge X \xrightarrow{\bar{e} \wedge \text{id}} \bar{E}^n \wedge X \xrightarrow{e_E \wedge \text{id}} E \wedge \bar{E}^n \wedge X.$$

Here $e_E : \mathbf{1} \rightarrow E$ is the unit of the algebra E , while $\bar{e} \wedge \text{id}$ (see (6.2)) appears as the transition map in the tower $E \wedge \bar{E}_{\bullet} \wedge X$. After smashing with E we thus have an

induced long exact sequences of homotopy objects

$$\cdots \rightarrow \tau_k(E \wedge \bar{E}^{n+1} \wedge X) \rightarrow \tau_k(E \wedge \bar{E}^n \wedge X) \rightarrow \tau_k(E \wedge E \wedge \bar{E}^n \wedge X) \rightarrow \cdots$$

and the maps $\tau_k(E \wedge \bar{E}^n \wedge X) \rightarrow \tau_k(E \wedge E \wedge \bar{E}^n \wedge X)$ are split by the multiplication of E . In this case, the previous map in the long exact sequence, which is the same as the transition map in the tower $\tau_k(E \wedge \bar{E}^\bullet \wedge X)$ is zero, and hence the associated pro-object is equivalent to 0 for every k . Using Corollary A.1.6, we deduce that the upper horizontal map of (7.11) is a τ -equivalence. Both source and target of this map are constant towers, so the map is actually an equivalence in \mathcal{M} , and this concludes the proof. \blacksquare

7.3.6. The combination of Theorem 7.3.5, Theorem 4.3.7 and Proposition 3.2.15 recovers and generalizes the results of [16] using different techniques. Similar results have been published by Bachmann and Østvær in [5] using techniques similar to ours. They have actually found a more efficient strategy for reaching the conclusions of Theorem 7.3.5.

7.3.7. The content of Theorem 7.3.5 is sometimes summed up by saying the E -based Adams–Novikov spectral sequence conditionally converges for k -connected objects. Of course one could ask for stronger notions of convergence. In the topological setting, work of Bousfield [8, Theorem 6.12] shows how better convergence properties may be related to structural properties of E . In the motivic setting, little is known in this direction. In a different direction, however, work of Kylling and Wilson [18] investigates strong convergence properties for the Adams spectral sequence for the sphere spectrum over fields.

LEMMA 7.3.8. *Let E be a homotopy commutative algebra in \mathcal{M} satisfying Assumption 4.2.1 in the special case that $I = \emptyset$. Then for every k -connected object X , the tower $\{P^n(\mathbf{1}[\mathcal{J}^{-1}] \wedge X)\}_n$ is a strongly $\tau_0 E$ -nilpotent resolution of X .*

PROOF. Since the unit $\mathbf{1} \rightarrow E$ induces an equivalence $\tau_0(\mathbf{1}[\mathcal{J}^{-1}]) \xrightarrow{\cong} \mathcal{R}$, we immediately conclude that the homotopy objects $\tau_k(X \wedge \mathbf{1}[\mathcal{J}^{-1}])$ are all \mathcal{R} -modules and hence they are strongly \mathcal{R} -nilpotent. The mapping property of strongly \mathcal{R} -nilpotent resolutions is an immediate consequence of the universal property of the Postnikov truncations. \blacksquare

THEOREM 7.3.9. *Let E be a homotopy commutative algebra in \mathcal{M} satisfying Assumption 4.2.1 in the special case that $I = \emptyset$. Then for every k -connected object X , the natural map $\lambda_E(X) : X \rightarrow X_E^\wedge$ is an E -equivalence. In particular, the map $\beta_E(X) : X_E \rightarrow X_E^\wedge$ of (7.2) is an equivalence in \mathcal{M} .*

PROOF. The proof proceeds along the same lines as the proof of Theorem 7.3.5. More precisely we start by observing that the towers $P^\bullet(\bar{E} \wedge X)$ and $P^\bullet(X \wedge \mathbf{1}[\mathcal{J}^{-1}])$ are both strongly \mathcal{R} -nilpotent resolutions of X by Proposition 7.3.2 and Lemma 7.3.8 respectively. We deduce, as in the proof of Theorem 7.3.5, that there is an equivalence $\psi : X \hat{E} \rightarrow X \wedge \mathbf{1}[\mathcal{J}^{-1}]$ under X making the following square of pro-objects under X commutative:

$$(7.13) \quad \begin{array}{ccc} \{X \hat{E}\} & \xrightarrow{u} & \{X \wedge \mathbf{1}[\mathcal{J}^{-1}]\} \\ \downarrow & & \downarrow \\ \{P^\bullet(\bar{E} \wedge X)\} & \xrightarrow[\phi]{\cong} & \{P^\bullet(X \wedge \mathbf{1}[\mathcal{J}^{-1}])\}. \end{array}$$

After smashing (7.13) with E , the lower horizontal map remains an equivalence of pro-objects. The vertical map on the right-hand side of (7.13) is a τ -equivalence by Section A.1.4, and stays a τ -equivalence after smashing with E by Lemma A.1.7. These two observations show that, after smashing with E , also the left vertical map of (7.13) is a τ -equivalence. The remaining part of the proof follows step by step the proof of Theorem 7.3.5. \blacksquare

A. Appendices

A.1 – Pro-objects

We gather here some well-known statements on pro-objects in an ∞ -category \mathcal{C} that we have freely used in the previous sections. We will also recall some properties of pro-objects in a symmetric monoidal stable ∞ -category \mathcal{M} endowed with a left-complete multiplicative t -structure.

A.1.1. Recall that the ∞ -category of pro-objects of \mathcal{C} , denoted by $\text{Pro}(\mathcal{C})$, is defined as $\text{Ind}(\mathcal{C}^{\text{op}})^{\text{op}}$ (cf. [21, Section 5.3.5]). Recall that every diagram $p : I \rightarrow \mathcal{C}$ indexed on a cofiltered simplicial set I gives a pro-object of \mathcal{C} : in \mathcal{C}^{op} we have a filtered diagram $I^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$, and the colimit of its composition with the Yoneda embedding

$$I^{\text{op}} \xrightarrow{p} \mathcal{C}^{\text{op}} \xrightarrow{j} \mathcal{P}(\mathcal{C}^{\text{op}})$$

is the desired pro-object. Starting with Section 3 we have considered towers in \mathcal{C} , i.e., diagrams indexed on $\mathbb{N}(\mathbb{N}^{\text{op}})$; these are clearly cofiltered simplicial sets, and thus every diagram in \mathcal{C} indexed on them gives rise to a pro-object of \mathcal{C} . Along the text we have used the symbol $\{X_\bullet\}$ to denote the pro-object associated with a diagram X_\bullet . We denote by $\text{Tow}(\mathcal{C})$ the full sub- ∞ -category of $\text{Pro}(\mathcal{C})$ spanned by pro-objects

associated with diagrams indexed on $N(\mathbb{N}^{\text{op}})$. We have used the term ‘tower’ to denote diagrams indexed on $N(\mathbb{N}^{\text{op}})$ as well as their associated pro-object.

LEMMA A.1.2. *Let $X_\bullet, Y_\bullet : I \rightarrow \mathcal{C}$ be cofiltered diagrams in \mathcal{C} . Then for the associated pro-objects, we have*

$$\text{Map}_{\text{Pro}(\mathcal{C})}(\{X_\bullet\}, \{Y_\bullet\}) \simeq \lim_n \text{colim}_k \text{Map}_{\mathcal{C}}(X_k, Y_n).$$

PROOF. The lemma follows from the facts that $\mathcal{P}(\mathcal{C}^{\text{op}})$ admits small colimits [21, Corollary 5.1.2.3] and constant ind-objects are compact [21, Proposition 5.3.5.5]. ■

DEFINITION A.1.3. Assume that \mathcal{M} is a stable ∞ -category endowed with a t -structure. A map of pro-objects $f : \{X_\bullet\} \rightarrow \{Y_\bullet\}$ is a τ -equivalence if, for every integer p , the induced map $\{\tau_p(X_\bullet)\} \rightarrow \{\tau_p(Y_\bullet)\}$ is an equivalence in $\text{Pro}(\mathcal{C}^\heartsuit)$. A pro-object $\{X_\bullet\}$ is τ -equivalent to 0 if, for every integer p , the homotopy objects $\{\tau_p(X_\bullet)\}$ are equivalent to 0 as objects of $\text{Pro}(\mathcal{C}^\heartsuit)$.

A.1.4. Of course every equivalence of pro-objects is a τ -equivalence. Moreover, given any object $\{X_\bullet\}$ of $\text{Tow}(\mathcal{M})$, the projection to the Postnikov tower

$$\pi_k : X_k \rightarrow P^k(X_k)$$

induces a τ -equivalence $\{\pi_\bullet\} : \{X_\bullet\} \rightarrow \{P^\bullet(X_\bullet)\}$. Note that in general the projection map $\{\pi_\bullet\}$ does not need to be a pro-equivalence. By cofinality, the map $\{X_\bullet\} \rightarrow \{P^\bullet(X_\bullet)\}$ induces an equivalence between the respective inverse limits.

A.1.5. Since \mathcal{M}^\heartsuit is an abelian category, $\text{Pro}(\mathcal{M}^\heartsuit)$ is an abelian category by [1, Appendix, Proposition 4.5]. Moreover, the full subcategory $\text{Tow}(\mathcal{M}^\heartsuit) \subseteq \text{Pro}(\mathcal{M}^\heartsuit)$ is closed under finite limits and colimits: this can be proved following the strategy of the proof of [6, Proposition 2.7]. In particular, $\text{Tow}(\mathcal{M}^\heartsuit)$ is an abelian category. It follows that a map $\{f_\bullet\} : \{M_\bullet\} \rightarrow \{N_\bullet\}$ of $\text{Tow}(\mathcal{M}^\heartsuit)$ is an equivalence if and only if both $\ker(\{f_\bullet\})$ and $\text{coker}(\{f_\bullet\})$ are pro-objects equivalent to 0. In particular, we conclude the following.

COROLLARY A.1.6. *Let $X_\bullet, Y_\bullet, Z_\bullet$ be towers in \mathcal{M} . Assume we have a fiber sequence of towers*

$$X_\bullet \xrightarrow{f_\bullet} Y_\bullet \xrightarrow{g_\bullet} Z_\bullet.$$

Then the induced map of pro-objects $\{g_\bullet\} : \{Y_\bullet\} \rightarrow \{Z_\bullet\}$ is a τ -equivalence if and only if the pro-object $\{X_\bullet\}$ is τ -equivalent to 0.

LEMMA A.1.7. *Assume that $\{X_\bullet\}$ is an object of $\text{Tow}(\mathcal{M})$ and that the t -structure on \mathcal{M} is multiplicative. Then for every object $E \in \mathcal{M}_{\geq k}$, the projection to the Postnikov tower induces a τ -equivalence*

$$\{E \wedge X_\bullet\} \rightarrow \{E \wedge P^\bullet(X_\bullet)\}.$$

PROOF. Consider the fundamental fiber sequence

$$E \wedge P_{n+1}(X_n) \rightarrow E \wedge X_n \rightarrow E \wedge P^n(X_n)$$

induced from (2.1). If $E \in \mathcal{M}_{\geq k}$, then $E \wedge P_{n+1}(X_n) \in \mathcal{M}_{\geq k+n+1}$, and in particular $\{\tau_p(E \wedge P_{\bullet+1}(X_\bullet))\}$ is equivalent to zero. The statement then readily follows from Corollary A.1.6. \blacksquare

A.2 – Categorical recollection

We quickly gather some known general properties of the ∞ -category $\mathcal{SH}(S)$, where S is a Noetherian scheme of finite Krull dimension. Using the formalism of [22], we deduce analogous properties for categories of modules over commutative algebras in $\mathcal{SH}(S)$. Our arguments are by no mean original; they are rather a reader's guide to the navigation of the relevant statements of [22].

A.2.1. In [17], Jardine introduces the motivic model structure on symmetric T -spectra $\mathbf{Spt}_T^\Sigma(S)$, which is the base for our constructions. We can combine [17, Section 4, in particular Theorems 4.15 and 4.31] together with [13, Lemma 4.2] to get the following result.

THEOREM A.2.2. *The stable motivic model structure on $\mathbf{Spt}_T^\Sigma(S)$ is simplicial, stable, proper, cofibrantly generated and combinatorial. The smash product of motivic symmetric T -spectra can be completed to the datum of a closed symmetric monoidal structure on $\mathbf{Spt}_T^\Sigma(S)$, making it a simplicial symmetric monoidal model category.*

DEFINITION A.2.3. A symmetric monoidal ∞ -category is an infinity category \mathcal{C}^\otimes with a coCartesian fibration $p : \mathcal{C}^\otimes \rightarrow \mathbf{N}(\text{Fin}_*)$ such that the functors $\rho_!^i : \mathcal{C}_{(n)}^\otimes \rightarrow \mathcal{C}_{(1)}^\otimes$ induce an equivalence of ∞ -categories

$$\prod_i \rho_!^i : \mathcal{C}_{(n)}^\otimes \simeq (\mathcal{C}_{(1)}^\otimes)^n$$

(see [22, Definition 2.0.0.7]). The ∞ -category $\mathcal{C}_{(1)}^\otimes$ is called the underlying ∞ -category associated with \mathcal{C}^\otimes , and is usually abusively called a symmetric monoidal ∞ -category, hiding the reference to the map p .

A.2.4. As in ordinary category theory, colored operads generalize the notion of symmetric monoidal category, similarly in higher category theory ∞ -operads generalize the notion of symmetric monoidal ∞ -categories. We do not recall the definition of an ∞ -operad which can be found in [22, Definition 2.1.1.10]. We however record that, as a consequence of [22, Proposition 2.1.2.12], a coCartesian fibration $p : \mathcal{C}^\otimes \rightarrow \mathbf{N}(\mathbf{Fin}_*)$ between ∞ -categories is a ∞ -operad if and only if it is a symmetric monoidal ∞ -category.

A.2.5. $\mathcal{SH}(S)$ is symmetric monoidal, presentable and stable. In the case of interest for us, the work of Lurie on ∞ -categories gives us a streamlined way of producing a symmetric monoidal ∞ -category $\mathcal{SH}(S)$ associated with a base scheme S . One possible construction is to set

$$\mathcal{SH}(S) := \mathbf{N}_\Delta(\mathbf{Spt}_T^\Sigma(S)^\circ),$$

where \mathbf{N}_Δ denotes the simplicial nerve construction, and $\mathbf{Spt}_T^\Sigma(S)^\circ \subseteq \mathbf{Spt}_T^\Sigma(S)$ denotes the full subcategory spanned by cofibrant-fibrant objects.

With this definition, using [23] allows to conclude that we have equivalences of homotopy categories

$$h\mathbf{N}_\Delta(\mathbf{Spt}_T^\Sigma(S)^\circ) \simeq \mathbf{Ho}(\mathbf{Spt}_T^\Sigma(S)^\circ) \simeq \mathbf{Ho}(\mathbf{Spt}_T^\Sigma(S)).$$

By design, $\mathcal{SH}(S)$ is the fiber over $\langle 1 \rangle$ of the operadic nerve construction

$$p : \mathbf{N}^\otimes(\mathbf{Spt}_T^\Sigma(S)^\circ) \rightarrow \mathbf{N}(\mathbf{Fin}_*),$$

see [22, Notation 2.1.1.22, Definition 2.1.1.23]. Combining Theorem A.2.2 with [22, Proposition 4.1.7.10], we conclude that $\mathcal{SH}(S)$ is the underlying ∞ -category of a symmetric monoidal ∞ -category. The monoidal product, which is induced by the smash product of spectra is denoted by $- \wedge -$ and the unit is denoted by \mathbb{S} .

Recall that $\mathbf{Spt}_T^\Sigma(S)$ is a combinatorial simplicial model category. In this situation, Proposition A.3.7.6 of [21] implies that $\mathcal{SH}(S)$ is a presentable ∞ -category. In particular, $\mathcal{SH}(S)$ admits all small limits and colimits [21, Proposition 4.2.4.8].

In addition, $\mathcal{SH}(S)$ is a stable ∞ -category in the sense of [22, Definition 1.1.1.9]. Indeed, in view of [22, Corollary 1.4.2.27], we only need to check that $\mathcal{SH}(S)$ has finite colimits and the suspension functor $\Sigma : \mathcal{SH}(S) \rightarrow \mathcal{SH}(S)$ is an equivalence. Now the first condition is satisfied since $\mathcal{SH}(S)$ is presentable. The second condition is implied by the fact that the suspension functor Σ is induced from the Quillen equivalence $\Sigma_{\mathbb{S}^1} : \mathbf{Spt}_T^\Sigma(S) \rightleftarrows \mathbf{Spt}_T^\Sigma(S) : \Omega_{\mathbb{S}^1}$. Indeed, from [21, Theorem 4.2.4.1] it follows that \mathbf{N}_Δ carries homotopy (co)limit diagrams in $\mathbf{Spt}_T^\Sigma(S)^\circ$ to (co)limit diagrams in $\mathcal{SH}(S)$.

A more informative construction of $\mathcal{SH}(S)$ was given by Robalo [31, Theorem 2.26] who proves that $\mathcal{SH}(S)$ is the stable presentable symmetric monoidal ∞ -category initial with respect to the property that $\mathbb{P}^1 \wedge -$ is an invertible endofunctor.

A.2.6. $\mathcal{SH}(S)$ is presentably symmetric monoidal. The properties of the monoidal structure on $\mathcal{SH}(S)$ are actually a bit stronger than what already stated, making it a presentably symmetric monoidal ∞ -category according to [22, Definition 3.4.4.1]. The essential point is that $\mathcal{SH}(S)$ is the underlying ∞ -category of a combinatorial simplicial symmetric monoidal model category. Indeed:

- (1) p is a coCartesian fibration of ∞ -operads, as recorded in Section A.2.5.
- (2) p is compatible with small colimits according to [22, Definition 3.1.1.18]. Indeed, $\mathbf{Spt}_T^{\Sigma}(S)$ is co-complete and the smash product of spectra commutes with all colimits, being a left adjoint.
- (3) For each object $\langle n \rangle \in N(\mathbf{Fin}_*)$, the fiber $\mathcal{SH}(S)_{\langle n \rangle}^{\otimes}$ is presentable. Indeed,

$$\mathcal{SH}(S)_{\langle n \rangle}^{\otimes} \simeq \prod_{i=1}^n \mathcal{SH}(S)_{\langle 1 \rangle}^{\otimes} \simeq \prod_{i=1}^n \mathcal{SH}(S),$$

whose presentability was addressed in Section A.2.5.

DEFINITION A.2.7 ([22, Definition 2.1.3.1]). A commutative algebra of $\mathcal{SH}(S)$ is a map of ∞ -operads $s : N(\mathbf{Fin}_*) \rightarrow \mathcal{SH}(S)^{\otimes}$ which is a section of the natural projection $p : \mathcal{SH}(S)^{\otimes} \rightarrow N(\mathbf{Fin}_*)$. The object $s(\langle 1 \rangle) : \Delta \rightarrow \mathcal{SH}(S)$ is called the commutative algebra underlying s . Along this section we will often refer to $s(\langle 1 \rangle)$ as commutative algebra in $\mathcal{SH}(S)$, leaving the section s implicit.

A.2.8. Modules over a commutative algebra. Let us denote $\mathcal{SH}(S)$ simply by \mathcal{C} in order to make references to [22] more directly traceable. Let A be a commutative algebra in \mathcal{C} . Lurie constructs in [22, Section 3.3.3] a fibration of ∞ -operads $p_A : \mathbf{Mod}_A(\mathcal{C})^{\otimes} \rightarrow N(\mathbf{Fin}_*)$. What we denoted in Section 2.2.2 by $\mathcal{M}od_A(S)$ is the ∞ -category $\mathbf{Mod}_A(\mathcal{C}) := \mathbf{Mod}_A(\mathcal{C})_{\langle 1 \rangle}^{\otimes}$. We omit the relevant ∞ -operad from the notation, since we are only dealing with $N(\mathbf{Fin}_*)$. The construction of $\mathbf{Mod}_A(\mathcal{C})^{\otimes}$ also produces a forgetful functor $\phi : \mathbf{Mod}_A(\mathcal{C})^{\otimes} \rightarrow \mathcal{C}^{\otimes}$. In Section 2.2.2, we denoted by $U_A = \phi_{\langle 1 \rangle}$ the functor induced by ϕ on the underlying ∞ -categories.

A.2.9. U_A commutes with small limits and colimits. We can use [22, Theorem 3.4.4.2] and [22, Corollary 3.4.4.6] to deduce that $p_A : \mathbf{Mod}_A(\mathcal{C})^{\otimes} \rightarrow N(\mathbf{Fin}_*)$ is actually a presentably symmetric monoidal ∞ -category and that the forgetful functor

$$\phi : \mathbf{Mod}_A(\mathcal{C})^{\otimes} \rightarrow \mathcal{C}^{\otimes}$$

detects and commutes with all small colimits. Similarly, using [22, Corollary 3.4.3.6], we deduce that ϕ detects and commutes with all small limits. In particular, the underlying functor $U_A = \phi_{(1)} : \text{Mod}_A(\mathcal{C}) \rightarrow \mathcal{C}$ detects and commutes with all small limits and colimits.

A.2.10. U_A is conservative. This is not spelled out explicitly in [22], but it can be easily deduced from the combination of some of the main statements. In first place, we use that forgetting the commutativity of A induces a canonical equivalence between A -modules and left modules over the associative algebra A

$$\text{Mod}_A(\mathcal{C}) \xrightarrow{\cong} \text{LMod}_A(\mathcal{C}),$$

as proven in [22, Corollary 4.5.1.6]. In second place, we use that $\text{Mod}_A(\mathcal{C})$ is naturally identified as the fiber over A of a cartesian fibration $\vartheta : \text{Mod}(\mathcal{C}) \rightarrow \text{Alg}(\mathcal{C})$. More precisely we have a diagram of ∞ -categories:

$$\begin{array}{ccc} \text{LMod}(\mathcal{C}) & \xrightarrow{U} & \mathcal{C} \\ \downarrow \vartheta & & \\ \text{Alg}(\mathcal{C}) & & \end{array}$$

The behavior of ϑ on objects is mapping $(R, M) \mapsto R$, where R is an associative algebra and M a left R -module. The functor U instead operates on objects as $(R, M) \mapsto M$. The functor ϑ is a cartesian fibration of ∞ -categories and an arrow of $\text{LMod}(\mathcal{C})$ is ϑ -cartesian if and only if its image in \mathcal{C} is an equivalence (see [22, Corollary 4.2.3.2]). However, an A -module map $a : \Delta^1 \rightarrow \text{LMod}_A(\mathcal{C})$ is a map of $\text{LMod}(\mathcal{C})$ covering $\text{id}_A : \Delta^1 \rightarrow \text{Alg}(\mathcal{C})$, and thus a is ϑ -cartesian if and only if a is an equivalence in $\text{LMod}(\mathcal{C})$ (see for instance [21, Proposition 2.4.4.3]) if and only if a is an equivalence in $\text{LMod}_A(\mathcal{C})$. Hence ϕ is conservative.

A.2.11. The left adjoint of U_A . We keep using the identification mentioned above: $\text{Mod}_A(\mathcal{C}) \xrightarrow{\cong} \text{LMod}_A(\mathcal{C})$. The functor $U_A : \text{LMod}_A(\mathcal{C}) \rightarrow \mathcal{C}$ has a left adjoint F_A , that on objects acts by mapping a spectrum X to the free A -module generated by X . The composition $U_A \circ F_A$ is equivalent to $A \wedge -$. Moreover, if $\lambda : F_A(X) \xrightarrow{\cong} M$ is a free A -module, then λ induces an equivalence of spaces

$$\text{Map}_A(M, N) \rightarrow \text{Map}_{\mathbb{S}}(X, U_A(N)).$$

All these claims can be found in [22, Corollaries 4.2.4.6 and 4.2.4.8].

The functor $\phi : \text{Mod}_A(\mathcal{C})^{\otimes} \rightarrow \mathcal{C}^{\otimes}$ has a symmetric monoidal left adjoint $\psi : \mathcal{C}^{\otimes} \rightarrow \text{Mod}_A(\mathcal{C})^{\otimes}$. This follows from [22, Theorem 4.5.3.1, Remark 4.5.3.2], whose

assumptions are automatically satisfied, since p_A is a presentably symmetric monoidal ∞ -category. In particular, F_A is equivalent to $\psi_{(1)}$.

ACKNOWLEDGMENTS – I wish to thank Marc Levine, Joseph Ayoub and Andrew Kresch for their encouragement in editing the arguments of [24]. A special thank goes to Tom Bachmann for the interest in a previous version of this paper. Another special thanks goes to Jones Kylling. His enthusiasm and interest on this project have helped me focusing on this work in several occasions. Furthermore, he found a mistake in a previous version of [24], and pointed out a number of typos. I wish to thank John Rognes for helpful comments on an earlier version of this paper, and for spotting a mistake. I wish to thank Georg Tamme for his helpful comments and for suggesting a reference. I wish to thank the anonymous referee for her/his careful reading, for suggesting further references, and for tangibly improving the overall quality of the exposition. I also wish to thank Mauro Porta who helped me navigating the higher categorical language. Finally, I wish to thank the algebraic geometry group of the University of Zürich for providing a comfortable working environment.

FUNDING – This work was supported by the *Swiss National Science Foundation* (SNF), project 200020_178729.

REFERENCES

- [1] M. ARTIN – B. MAZUR, *Etale homotopy*. Lecture Notes in Math. 100, Springer, Berlin, 1969. Zbl [0182.26001](#) MR [0245577](#)
- [2] T. BACHMANN, [The generalized slices of Hermitian \$K\$ -theory](#). *J. Topol.* **10** (2017), no. 4, 1124–1144. Zbl [1453.14065](#) MR [3743071](#)
- [3] T. BACHMANN, [On the conservativity of the functor assigning to a motivic spectrum its motive](#). *Duke Math. J.* **167** (2018), no. 8, 1525–1571. Zbl [1390.14064](#) MR [3807316](#)
- [4] T. BACHMANN – B. CALMÈS – F. DÉGLISE – J. FASEL – P. A. ØSTVÆR, *Milnor–Witt motives*. 2022, arXiv:[2004.06634v2](#).
- [5] T. BACHMANN – P. A. ØSTVÆR, [Topological models for stable motivic invariants of regular number rings](#). *Forum Math. Sigma* **10** (2022), Paper No. e1. Zbl [1490.14033](#) MR [4365078](#)
- [6] D. BLANC, *Colimits for the pro-category of towers of simplicial sets*. *Cahiers Topologie Géom. Différentielle Catég.* **37** (1996), no. 4, 258–278. Zbl [0867.18001](#) MR [1428551](#)
- [7] A. K. BOUSFIELD, [The Boolean algebra of spectra](#). *Comment. Math. Helv.* **54** (1979), no. 3, 368–377. Zbl [0421.55002](#) MR [543337](#)
- [8] A. K. BOUSFIELD, [The localization of spectra with respect to homology](#). *Topology* **18** (1979), no. 4, 257–281. Zbl [0417.55007](#) MR [551009](#)

- [9] A. K. BOUSFIELD – D. M. KAN, *Homotopy limits, completions and localizations*. Lecture Notes in Math. 304, Springer, Berlin, 1972. Zbl [0259.55004](#) MR [0365573](#)
- [10] U. BUNKE – T. NIKOLAUS – G. TAMME, *The Beilinson regulator is a map of ring spectra*. *Adv. Math.* **333** (2018), 41–86. Zbl [1400.14063](#) MR [3818072](#)
- [11] D.-C. CISINSKI – F. DÉGLISE, *Triangulated categories of mixed motives*. Springer Monogr. Math., Springer, Cham, 2019. Zbl [07138952](#) MR [3971240](#)
- [12] E. ELMANTO – H. KOLDERUP, *On modules over motivic ring spectra*. *Ann. K-Theory* **5** (2020), no. 2, 327–355. Zbl [1440.14120](#) MR [4113773](#)
- [13] M. HOYOIS, *From algebraic cobordism to motivic cohomology*. *J. Reine Angew. Math.* **702** (2015), 173–226. Zbl [1382.14006](#) MR [3341470](#)
- [14] M. HOYOIS, *Cdh descent in equivariant homotopy K-theory*. *Doc. Math.* **25** (2020), 457–482. Zbl [1453.14068](#) MR [4124487](#)
- [15] M. HOYOIS – S. KELLY – P. A. ØSTVÆR, *The motivic Steenrod algebra in positive characteristic*. *J. Eur. Math. Soc. (JEMS)* **19** (2017), no. 12, 3813–3849. Zbl [1386.14087](#) MR [3730515](#)
- [16] P. HU – I. KRIZ – K. ORMSBY, *Convergence of the motivic Adams spectral sequence*. *J. K-Theory* **7** (2011), no. 3, 573–596. Zbl [1309.14018](#) MR [2811716](#)
- [17] J. F. JARDINE, *Motivic symmetric spectra*. *Doc. Math.* **5** (2000), 445–552. Zbl [0969.19004](#) MR [1787949](#)
- [18] J. I. KYLLING – G. M. WILSON, *Strong convergence in the motivic Adams spectral sequence*. 2019, arXiv:[1901.03399](#).
- [19] M. LEVINE, *The homotopy coniveau tower*. *J. Topol.* **1** (2008), no. 1, 217–267. Zbl [1154.14005](#) MR [2365658](#)
- [20] M. LEVINE, *The slice filtration and Grothendieck-Witt groups*. *Pure Appl. Math. Q.* **7** (2011), no. 4, 1543–1584. Zbl [1316.14043](#) MR [2918174](#)
- [21] J. LURIE, *Higher topos theory*. Ann. of Math. Stud. 170, Princeton University Press, Princeton, NJ, 2009. Zbl [1175.18001](#) MR [2522659](#)
- [22] J. LURIE, *Higher algebra*. <https://www.math.ias.edu/~lurie/papers/HA.pdf>, 2017.
- [23] J. LURIE, *Kerodon*. <https://kerodon.net>, 2018.
- [24] L. MANTOVANI, *Localizations and completions in motivic homotopy theory*. 2018, arXiv:[1810.04134](#).
- [25] F. MOREL, *On the motivic π_0 of the sphere spectrum*. In *Axiomatic, enriched and motivic homotopy theory*, pp. 219–260, NATO Sci. Ser. II Math. Phys. Chem. 131, Kluwer, Dordrecht, 2004. Zbl [1130.14019](#) MR [2061856](#)
- [26] F. MOREL, *A^1 -algebraic topology over a field*. Lecture Notes in Math. 2052, Springer, Heidelberg, 2012. Zbl [1263.14003](#) MR [2934577](#)

- [27] I. PANIN – K. PIMENOV – O. RÖNDIGS, [A universality theorem for Voevodsky’s algebraic cobordism spectrum](#). *Homology Homotopy Appl.* **10** (2008), no. 2, 211–226. Zbl [1162.14013](#) MR [2475610](#)
- [28] I. PANIN – K. PIMENOV – O. RÖNDIGS, [On Voevodsky’s algebraic \$K\$ -theory spectrum](#). In *Algebraic topology*, pp. 279–330, Abel Symp. 4, Springer, Berlin, 2009. Zbl [1179.14022](#) MR [2597741](#)
- [29] I. PANIN – C. WALTER, [On the motivic commutative ring spectrum \$\mathbf{BO}\$](#) . *St. Petersburg Math. J.* **30** (2019), no. 6, 933–972. Zbl [1428.14011](#) MR [3882540](#)
- [30] I. PANIN – C. WALTER, [On the algebraic cobordism spectra \$\mathbf{MSL}\$ and \$\mathbf{MSp}\$](#) . *St. Petersburg Math. J.* **34** (2023), no. 1, 109–141.
- [31] M. ROBALO, [\$K\$ -theory and the bridge from motives to noncommutative motives](#). *Adv. Math.* **269** (2015), 399–550. Zbl [1315.14030](#) MR [3281141](#)
- [32] O. RÖNDIGS – P. A. ØSTVÆR, [Modules over motivic cohomology](#). *Adv. Math.* **219** (2008), no. 2, 689–727. Zbl [1180.14015](#) MR [2435654](#)
- [33] O. RÖNDIGS – M. SPITZWECK – P. A. ØSTVÆR, [The first stable homotopy groups of motivic spheres](#). *Ann. of Math. (2)* **189** (2019), no. 1, 1–74. Zbl [1406.14018](#) MR [3898173](#)
- [34] W. SCHARLAU, [Quadratic and Hermitian forms](#). Grundlehren Math. Wiss. 270, Springer, Berlin, 1985. Zbl [0584.10010](#) MR [770063](#)
- [35] J. SCHMIDT – F. STRUNK, [Stable \$\mathbb{A}^1\$ -connectivity over Dedekind schemes](#). *Ann. K-Theory* **3** (2018), no. 2, 331–367. Zbl [1423.14158](#) MR [3781430](#)
- [36] M. SPITZWECK – P. A. ØSTVÆR, [The Bott inverted infinite projective space is homotopy algebraic \$K\$ -theory](#). *Bull. Lond. Math. Soc.* **41** (2009), no. 2, 281–292. Zbl [1213.55006](#) MR [2496504](#)
- [37] R. THORNTON, [The homogeneous spectrum of Milnor-Witt \$K\$ -theory](#). *J. Algebra* **459** (2016), 376–388. Zbl [1348.14064](#) MR [3503978](#)
- [38] M. YAKERSON, [The unit map of the algebraic special linear cobordism spectrum](#). *J. Inst. Math. Jussieu* **20** (2021), no. 6, 1905–1930. Zbl [1481.14042](#) MR [4332780](#)

Manoscritto pervenuto in redazione il 12 maggio 2021.