

Holomorphic Extension from Weakly Pseudoconcave CR Manifolds

ANDREA ALTOMANI (*) - C. DENSON HILL (**) - MAURO NACINOVICH (***)
EGMONT PORTEN (* **)

ABSTRACT - Let M be a smooth locally embeddable CR manifold, having some CR dimension m and some CR codimension d . We find an improved local geometric condition on M which guarantees, at a point p on M , that germs of CR distributions are smooth functions, and have extensions to germs of holomorphic functions on a full ambient neighborhood of p . Our condition is a form of weak pseudoconcavity, closely related to essential pseudoconcavity as introduced in [HN1]. Applications are made to CR meromorphic functions and mappings. Explicit examples are given which satisfy our new condition, but which are not pseudoconcave in the strong sense. These results demonstrate that for codimension $d > 1$ there are additional phenomena, which are invisible when $d = 1$.

1. Introduction.

The goal of the present article is to give improved geometric conditions on a generic CR manifold $M \subset \mathbb{C}^n$ which guarantee that all local CR functions extend holomorphically to a full neighborhood of a given point. This is well known to be true for strictly pseudoconcave CR manifolds,

(*) Indirizzo dell'A.: Research Unity in Mathematics, University of Luxembourg, 6, rue Coudenhove-Kalergi, L-1359 Luxembourg.

E-mail address: andrea.altomani@uni.lu

(**) Indirizzo dell'A.: Department of Mathematics, Stony Brook University, Stony Brook NY 11794 (USA).

E-mail address: dhill@math.sunysb.edu

(***) Indirizzo dell'A.: Dipartimento di Matematica, II Università di Roma "Tor Vergata", Via della Ricerca Scientifica, 00133 Roma (Italy).

E-mail address: nacinovi@mat.uniroma2.it

(* **) Indirizzo dell'A.: Department of Mathematics, Mid Sweden University, 85170 Sundsvall (Sweden)

E-mail address: Egmont.Porten@miun.se

i.e. in the case where the Levi form has one negative eigenvalue in each characteristic conormal direction. For hypersurfaces this is a classical result of H. Kneser and H. Lewy; for M of higher codimension it was proved independently by a number of authors (see [BP], [NV] for CR distributions, and [HN4] for two different proofs, one very short). Despite numerous efforts, the general problem to characterize those weakly pseudoconcave manifolds for which one has extension to full neighborhoods is still far from being completely understood, even for real analytic hypersurfaces.

Subtle sufficient conditions (sector and ray property) are known for weakly pseudoconcave *hypersurfaces* of finite type (see [BT2], [FR] for results and references). In *higher codimension* there are several options to approach the weakly pseudoconcave case. Manifolds which are Levi flat at the reference point to a certain order, and have all relevant concavity in the generalized Levi form determined by the next-order terms are studied in [Bo]. Here we aim at the opposite case where effects of different orders (counted with respect to bracket length) are combined. Actually our main motivation stems from homogeneous CR manifolds which bi-holomorphically look the same near every point. These higher codimensional homogeneous CR manifolds are abundant, occurring naturally in mathematics, and they have a strong tendency to be weakly pseudoconcave (see [MN1], [MN2], [MN3], [MN4], [AMN]). The main result of the present article reveals that there are additional phenomena which are invisible in codimension one, and indicates that finite type together with a suitable notion of weak pseudoconcavity should imply extension to a full neighborhood. To avoid confusion, we stress that the problem under consideration is different in nature from the problem of holomorphic wedge extension, for which a definitive answer is known ([T1], [Tu1], [J], [M], see also [MP2]). In fact, this definitive answer was obtained without having explicit control on the directions of extension, which is crucial for the problem at hand.

Let $M \subset \mathbb{C}^n$ be a smooth CR manifold. We denote by J the complex structure tensor on $T\mathbb{C}^n$, by $HM = TM \cap JTM$ the holomorphic tangent bundle of M , which is the real subbundle of TM invariant under J , and by $H^0M \subset T^*M$ the characteristic bundle, defined fiberwise as the annihilator of HM . We define the *vector valued* Levi form

$$\mathcal{L}_{M,p} = \mathcal{L}_p : H_pM \times H_pM \rightarrow \mathbb{C} \otimes (T_pM/H_pM)$$

by

$$\mathcal{L}_{M,p}(X, Y) = -[\tilde{X}, J\tilde{Y}](p) + i[\tilde{X}, \tilde{Y}](p) \bmod \mathbb{C} \otimes H_pM,$$

where $\tilde{X}, \tilde{Y} \in \Gamma(M, HM)$ are smooth extensions of X, Y , respectively. The usual Levi form $\mathcal{L}_{M, \xi} = \mathcal{L}_\xi$ is parameterized by the characteristic codirections $\xi \in H_p^0 M$. It is defined by $\mathcal{L}_{M, \xi}(X, Y) = \xi(\mathcal{L}_p(X, Y))$, for $\xi \in H_p^0 M$, $X, Y \in H_p M$ (where we read ξ as a form on $\mathbb{C} \otimes (T_p M / H_p^0 M)$ in the canonical way). Most often we shall work with the corresponding real hermitian forms $\mathcal{L}_p(X) = \mathcal{L}_p(X, X) \in T_p M / H_p M$ and $\mathcal{L}_\xi(X) = \mathcal{L}_\xi(X, X)$. The reader should take note of the subtle difference in notation between $\mathcal{L}_p(X)$ and $\mathcal{L}_\xi(X)$; the former is vector valued, and the latter is scalar valued.

A CR manifold M is *strictly (weakly) pseudoconcave* at a point $p \in M$ if for every $\xi \in H_p^0 M$, $\xi \neq 0$, \mathcal{L}_ξ has a negative (nonpositive) eigenvalue. Replacing ξ by $-\xi$, we see that \mathcal{L}_ξ has actually eigenvalues of both signs (in the strictly pseudoconcave case). Following [HN1], we call M *trace pseudoconcave* at $p \in M$ if for every $\xi \in H_p^0 M$, \mathcal{L}_ξ is either zero or has eigenvalues of both signs. Trace pseudoconcavity isolates one of the properties of essential pseudoconcavity introduced in [HN1]. We refer to that article for background information.

Let \mathcal{G}_1 be the sheaf of germs of smooth (real) CR vector fields on M (i.e. sections of HM). For every positive integer k we define inductively \mathcal{G}_{k+1} as the sheaf generated by \mathcal{G}_k and $[\mathcal{G}_1, \mathcal{G}_k]$. Let $G_{k,p} \subset T_p M$ be the vector space generated by pointwise evaluations of germs in \mathcal{G}_k at a point $p \in M$. We say that M is of *kind k* at p if $G_{k,p} = T_p M$ but $G_{j,p} \subsetneq T_p M$ for $j < k$. We say that M satisfies the *constant rank condition* if the spaces $G_{k,p}$ have dimension independent of p , i.e. if they form vector bundles $G_k = \bigcup_{p \in M} G_{k,p}$.

Now we can formulate our main result.

THEOREM 1.1. *Let M be a smooth generic CR manifold in \mathbb{C}^n and $p_0 \in M$. Assume that in a neighborhood of p_0 , M is trace pseudoconcave, satisfies the constant rank condition and is of kind less or equal to 3. Then for every open neighborhood U of p_0 in M , there is an open neighborhood V of p_0 in \mathbb{C}^n such that every CR distribution on U is smooth on $M \cap V$ and has a unique holomorphic extension to V .*

We emphasize that Theorem 1.1 reveals a phenomenon which remains invisible in codimension 1. In fact, in the hypersurface case its assumptions imply that M is of kind 2, hence strictly pseudoconcave. We expect the result to extend to arbitrary finite kind. In [HN1], the weak identity principle for CR functions (coincidence on open sets implies coincidence everywhere) was shown for essentially pseudoconcave CR manifolds. For those CR manifolds covered by the assumptions of Theorem 1.1 our result immediately yields the strong identity principle

(coincidence of Taylor coefficients at some point yields coincidence in a neighborhood).

Theorem 1.1 is proved in Sections 2-4. The rest of the article is devoted to applications, extensions and examples. In Section 5 we observe that CR manifolds as in Theorem 1.1 enjoy the local extension property E introduced in [HN2]. The results in [HN2], [HN3] yield far reaching global consequences for fields of CR meromorphic functions on such manifolds. Here CR meromorphic functions are functions which are locally representable as fractions of CR functions. An alternative approach to CR meromorphic mappings originates from work of Harvey and Lawson [HL]. The idea is to require the graph to look like a CR manifold with appropriate singularities. In general, extension of such mappings is complicated. Based on [MP2], we prove in Section 6 that such CR meromorphic functions extend meromorphically from manifolds with property E to full ambient neighborhoods and are in particular representable as local quotients. In Section 7 we present several classes of homogeneous CR manifolds to which all the local and global results indicated above apply. These were discovered in a much broader context (see [AMN]). For the reader's comfort, we give a reasonably self-contained presentation.

2. Preliminaries.

We will use some standard facts about the bundles $G_1 \subset G_2 \subset \dots$: If $G_k = G_{k+1}$ then all $G_j, j \geq k$, are equal (the proof is an application of the Jacobi identity). This means in particular that G_k is integrable in the sense of Frobenius. Moreover the map associating to smooth sections $X \in \Gamma(U, G_1), Y \in \Gamma(U, G_k)$, the section $[X, Y] \bmod G_k \in \Gamma(U, G_{k+1}/G_k)$ is tensorial, i.e. $[X, Y](p) \bmod G_{k,p}$ depends only on $X(p)$ and $Y(p)$.

Let us now have a closer look at G_2 . First we note that independently of concavity $G_{2,p}/H_pM$ is spanned as a real vector space by the image $C_p = \{\mathcal{L}_p(X) : X \in H_pM\}$ of the vector valued Levi form. Indeed, $L = \text{span}_{\mathbb{R}} C_p$ is contained in $G_{2,p}/H_pM$ by definition. On the other hand, polarization shows that $L \otimes \mathbb{C} = \text{span}_{\mathbb{C}} \{\mathcal{L}_p(X, Y) : X, Y \in H_pM\}$. Since the imaginary part of $\mathcal{L}_p(X, Y)$ is essentially $[X, Y]$, we obtain $G_{2,p}/H_pM \subset L$. The above is equivalent to the fact that $G_{2,p}$ is spanned by the preimage of C_p under the canonical projection $T_pM \rightarrow T_pM/H_pM$.

A simple but crucial observation is that trace pseudoconcavity allows us to replace linear spans by convex hulls.

LEMMA 2.1. *Assume that M is trace pseudoconcave at $p \in M$. Then $G_{2,p}/H_pM$ is the convex hull of $C_p = \{\mathcal{L}_p(X) : X \in H_pM\}$.*

PROOF. If the lemma fails, there is a nonzero linear functional ξ on $G_{2,p}/H_pM$ such that $C_p \subset \{\xi \geq 0\}$. We may extend ξ to an element of H_p^0M . Since $G_{2,p}/H_pM$ is the linear span of C_p , there is an $X \in C_p$ with $\xi(X) > 0$. Any $Y \in H_pM$ with $\mathcal{L}_p(Y) = X$ satisfies $\mathcal{L}_\xi(Y) > 0$. But this implies that \mathcal{L}_ξ has also some negative eigenvalue, in contradiction to $C_p \subset \{\xi \geq 0\}$ \square

To keep track of directions of extension, we will use an analogue of the analytic wave front set, denoted by WF_u . It is defined for CR distributions u via the FBI transform in [S], see also [T2]. For U open in M , let $CR(U)$ denote the space of continuous CR functions defined on U . We do not even have to recall the definition of WF_u , since the following basic properties will suffice for our purposes:

- (a) Let u be a CR distribution defined on $U \subset M$. Then WF_u is a cone, closed in the pointed characteristic bundle $H^0U \setminus o$ (o denoting the zero section).
- (b) $WF_u \cap H_p^0M = \emptyset$ holds if and only if the CR distribution u extends holomorphically to an ambient neighborhood of p .
- (c) Let $u \in CR(U)$. If CR extension from U holds at (p, X) , $p \in U$, $X \in T_pM \setminus H_pM$, then for any $\xi \in WF_u \cap H_p^0M$ we have $\xi(X) \geq 0$.

In (c) we use the following terminology: We say that *CR extension from U* holds at (p, X) , $p \in U$, $X \in T_pM \setminus H_pM$, if there is a C^2 -smooth ($\dim M + 1$)-dimensional CR manifold \tilde{M} attached to U along some U -neighborhood U' of p such that (i) for a representative of X , JX points into \tilde{M} and (ii) every $u \in CR(U)$ has a continuous extensions to $\tilde{M} \cup U'$ which is CR on \tilde{M} . Neglecting the dependence on U , we will sometimes call (p, X) or just X , a direction of CR extension. For (a), (b), see [S], whereas (c) is observed in [T2].

Theorem 1.1 is a consequence of the following more precise result which does not require kind 3.

THEOREM 2.2. *Let M be a smooth generic CR manifold in \mathbb{C}^n . Assume that on an open set $U \subset M$, M is trace pseudoconcave and that G_2, G_3 are bundles. Then for every continuous CR function u defined on U , we have $WF_u \subset G_3^\perp$.*

Theorem 2.2 will be proved in the next two sections. It implies Theorem 1.1 in the following way: Kind 3 means that G_3^\perp is the zero bundle near

p_0 . By **(b)** a continuous CR function u extends holomorphically to an ambient neighborhood of every p contained in some neighborhood U' of p_0 in M . By a standard gluing argument one obtains extension to an ambient neighborhood V' of U , which may a priori depend on u . Note that holomorphic extension in particular shows that $u \in C^\infty(U')$. Now a Baire category argument as in [HN2] yields extension to a neighborhood of V whose size only depends on U . This proves the theorem for continuous CR functions.

If u is a CR distribution, we may use a method from [BT1], [T], to represent it near p_0 as $u = \Delta_M^k f$, where f is a continuous CR function. Here k is a sufficiently large integer, and Δ_M is a variant of the Laplace operator which is defined in an ambient neighborhood of p_0 and restricts nicely to M . If \tilde{f} is a holomorphic extension of f , then the various properties of Δ_M imply that $\Delta_M^k \tilde{f}$ is the desired extension of u . Hence Theorem 1.1 follows from Theorem 2.2.

3. Proof plan for Theorem 2.2.

Here we will prove Theorem 2.2 modulo some more technical results on CR extension which are postponed to the following section. Pick some $u \in CR(U)$.

STEP 1. – $WF_u \subset G_2^\perp$. This will follow from trace pseudoconcavity. It is a consequence of the following lemma which holds without constant rank assumptions.

LEMMA 3.1. *Assume that $M \subset \mathbb{C}^n$ is trace pseudoconcave at $p \in M$. Then for every continuous CR function u defined near p we have $WF_u \cap H_p^0 M \subset G_{2,p}^\perp$.*

PROOF. Let $\zeta \in WF_u \cap H_p^0 M$. We will consider ζ both as a functional acting on $T_p M$ and $T_p M / H_p M$. By [Tu2], every element $X \in C_p$ can be approximated by directions of CR extensions $X_j \in T_p M / H_p M$. From property **(c)** and continuity we get $\zeta(X) \geq 0$. It follows that ζ is nonnegative on the convex hull of C_p . Since this convex hull is the vector space $G_{2,p} / H_p M$, ζ vanishes on $G_{2,p}$. \square

Notice that the lemma together with property **(b)** already imply extension to a full neighborhood for strictly pseudoconcave CR manifolds.

STEP 2. – $WF_u \subset G_3^\perp$. By Step 1 it suffices to show that a covector ξ_0 that annihilates G_2 but not G_3 is not contained in WF_u . Let $p_0 \in U$ denote the base point to which ξ_0 projects. It is our aim to show $\xi_0 \notin WF_u$ by constructing an appropriate CR extension and applying property (c).

Since the image C_{p_0} of the vector valued Levi form spans $G_{2,p_0}/H_{p_0}M$ (see the remarks before Lemma 2.1), we may select vectors $X_1, \dots, X_k \in H_{p_0}M$ such that the vectors $\tilde{Y}_j = \mathcal{L}_{M,p_0}(X_j), j = 1, \dots, k$, form a basis of $G_{2,p_0}/H_{p_0}M$. Extending the X_j smoothly to CR vector fields defined near p_0 , we obtain a local basis $\tilde{Y}_j(p) = \mathcal{L}_{M,p}(X_j)$ of G_2/HM . Set $Y_j = [JX_j, X_j]$, and choose a local basis Z_1, \dots, Z_{2m} of HM . Then the Z_i form together with the Y_j a local basis of G_2 .

First we claim that G_3 is spanned in some neighborhood of p_0 by the Z_i, Y_j , together with the brackets $[Z_i, Y_j]$. Indeed, by definition $G_{3,p}$ is spanned by $G_{2,p}$ and vectors of the form $[Z, Y](p)$ where $Z \in \mathcal{G}_{1,p}, Y \in \mathcal{G}_{2,p}$ for p near p_0 . Around p we may write $Z = \sum z_i Z_i, Y = \sum \tilde{z}_i \tilde{Y}_i + \sum y_j Y_j$, with smooth coefficients z_i, \tilde{z}_i, y_j . This yields

$$[Z, Y] = \sum z_i y_j [Z_i, Y_j] + R,$$

where R is a germ in $\mathcal{G}_{2,p}$. This proves the claim.

In the sequel, we will only need the following consequence: Since ξ_0 does not annihilate G_{3,p_0} , there are i_0, j_0 such that $\xi_0([Z_{i_0}, Y_{j_0}](p_0)) \neq 0$. For notational convenience we will write from now on $Y = Y_{j_0}, Z = Z_{i_0}$.

The following proposition, which will be proved in Section 4, yields CR extension at $(p_0, Y(p_0))$ realized by a CR manifold to which Y is complex tangent in a neighborhood of p_0 in M .

PROPOSITION 3.2. *Let $M \subset \mathbb{C}^n$ be a smooth generic CR manifold of CR dimension m and codimension d . Let $p_0 \in M$ and let U be an open neighborhood of p_0 in M . Let X be a smooth CR vector field on U with $\mathcal{L}_{M,p_0}(X) \neq 0$. Then there is a local C^4 -smooth generic CR manifold \tilde{M} of dimension $\dim M + 1$ with the following properties:*

- (a) $M \cap \tilde{M}$ is a neighborhood of p_0 in U and $\tilde{M} \setminus M$ has two connected components \tilde{M}^\pm .
- (b) The distribution on $M \cap \tilde{M}$ spanned by HM and $[JX, X]$ coincides with $(H\tilde{M} \cap T\tilde{M})|_{M \cap \tilde{M}}$.
- (c) Denoting by \tilde{M}^+ the side into which $J[JX, X](p_0)$ points, we have CR extension from U to \tilde{M}^+ .

Since ξ_0 annihilates $H_{p_0}\tilde{M} \cap T_{p_0}M$, there is a unique extension $\tilde{\xi}_0 \in H_{p_0}^0\tilde{M}$.

We claim that $\mathcal{L}_{\tilde{M}, \tilde{\zeta}_0}$ has eigenvalues of both signs. Indeed, both Z and Y may be extended to \mathcal{C}^3 -smooth CR vector fields on \tilde{M} which we denote by the same symbols. For Y we use here that Y is *complex* tangent to \tilde{M} along M by Proposition 3.2. The fact that $\tilde{\zeta}_0$ annihilates the image of the Levi form of M implies $\mathcal{L}_{\tilde{M}, \tilde{\zeta}_0}(Z(p_0)) = \mathcal{L}_{M, \zeta_0}(Z(p_0)) = 0$. Since $\tilde{\zeta}_0([Z, Y](p_0)) \neq 0$ is essentially the imaginary part of the sesquilinear Levi form, Z does not lie in the kernel of $\mathcal{L}_{\tilde{M}, \tilde{\zeta}_0}$. Hence $\mathcal{L}_{\tilde{M}, \tilde{\zeta}_0}(\cdot)$ takes both positive and negative values in every neighborhood of $Z(p_0)$ in $H_{p_0}\tilde{M}$. The claim follows.

Let X^+, X^- be positive and negative eigenvectors of $\mathcal{L}_{\tilde{M}, \tilde{\zeta}_0}$, respectively. Since all vectors in $H_{p_0}M$ are null vectors of $\mathcal{L}_{\tilde{M}, \tilde{\zeta}_0}$, we may assume, after replacing X^\pm by an appropriate complex multiple if necessary, that X^\pm both point into \tilde{M}^+ . But now the following proposition, whose proof is postponed to Section 4, yields for M two directions of CR extension $Y^\pm \in T_{p_0}M/H_{p_0}M$ with $\zeta_0(Y^+) > 0$ and $\zeta_0(Y^-) < 0$. This implies $\zeta_0 \neq WF_u$ by property (c). Hence the proof of Theorem 2.2, and thus also of Theorem 1.1, will be complete as soon as we have shown the following.

PROPOSITION 3.3. *Let $M, \tilde{M}, \tilde{M}^\pm$ and $p_0 \in M$ be as above. Assume that there is $\tilde{\zeta} \in H_{p_0}^0\tilde{M}$ and $X \in H_{p_0}\tilde{M} \setminus T_{p_0}M$ pointing into \tilde{M}^+ such that $\mathcal{L}_{\tilde{M}, \tilde{\zeta}}(X) > 0$. Then there is a direction $Z \in T_{p_0}M/H_{p_0}M$ of CR extension from U satisfying $\tilde{\zeta}(Z) > 0$.*

4. CR extension.

In this section we will prove Propositions 3.2 and 3.3. First we recall some basic material on Bishop discs introduced in the seminal paper [B]. Consider a generic CR manifold $M \subset \mathbb{C}^n$ of CR dimension m and codimension d . A \mathcal{C}^* -smooth analytic disc is a mapping $A(\zeta) = (Z(\zeta), W(\zeta)) \in \mathcal{C}^*(\bar{\Delta}, \mathbb{C}^n) \cap \mathcal{O}(\Delta, \mathbb{C}^n)$ where $\Delta = \{\zeta \in \mathbb{C} : |\zeta| < 1\}$. We say that A is attached to M if $A(S^1) \subset M$.

For $|\mu| \leq 1$ let \mathcal{T}_μ denote the Hilbert transform of a function $U : S^1 \rightarrow \mathbb{R}$ to its harmonic conjugate $\mathcal{T}_\mu U$, normalized by the condition that the harmonic extension of $\mathcal{T}_\mu U$ vanishes at $\zeta = \mu$. It is known that \mathcal{T}_μ is a continuous linear operator on the Hölder spaces $\mathcal{C}^{k,\alpha}(S^1, \mathbb{R})$ if $k \in \mathbb{N}_0$, $0 < \alpha < 1$. We shall use the same notation for the Hilbert transform applied componentwise to vector valued functions of $\mathcal{C}^{k,\alpha}(S^1, \mathbb{R}^d)$.

We work in coordinates

$$(1) \quad z_1 = x_1 + iy_1, \dots, z_m = x_m + iy_m, w_1 = u_1 + iw_1, \dots, w_d = u_d + iw_d$$

centered at the origin in which M is locally given as a graph $v = h(z, u)$ with $h(0) = 0$, $dh(0) = 0$. The Bishop equation is the nonlinear functional equation

$$(2) \quad U = -\mathcal{T}_\mu(h(Z, U)) + u.$$

Here $U = U(\zeta)$ is the unknown function mapping the unit circle $S^1 = \{|\zeta| = 1\} \subset \mathbb{C}$ to \mathbb{R}^d , whereas $Z = Z(\zeta)$ is the boundary value of a given holomorphic function from Δ to \mathbb{C}^m and $u \in \mathbb{R}^d$ is a prescribed vector. It is known that the Bishop equation can be solved in $C^{k,\alpha}(S^1, \mathbb{R}^d)$ provided the data $Z(\zeta)$ are $C^{k,\alpha}$ -small and $|u|$ is small (see [MP4] for detailed information). The solution U corresponds to a unique analytic disc $A : \bar{\Delta} \rightarrow \mathbb{C}^n$ whose restriction to S^1 is $(Z(\zeta), U + ih(Z(\zeta), U(\zeta)))$. More precisely, A is holomorphic on Δ and $C^{k,\alpha}$ -smooth up to S^1 . By construction, A is attached to M .

PROOF OF PROPOSITION 3.2. – The construction of \tilde{M}^+ will be a refinement of [HT, proof of Theorem 9.1]. We may assume $p_0 = 0$ and choose coordinates (z, w) as in (1). After appropriate rotations and dilations in z and w , we may furthermore assume $\mathcal{L}_{M,0}\left(\frac{\partial}{\partial x_1}\right) = \frac{\partial}{\partial u_1} \bmod H_0M$. We will construct \tilde{M}^+ as the union of real curves $\gamma_p(s)$, $s \in [0, \varepsilon_1)$, having their initial points at $\gamma_p(0) = p \in M$ and such that the segments $\gamma_p((0, \varepsilon_1))$ foliate \tilde{M}^+ .

Let us start by the construction of γ_0 . Following [HT], we first simplify the defining equations by removing some of the pure terms. After a change of the w_1 -variable

$$w_1 \mapsto w_1 - i \left(\sum_{j,k=1}^m \frac{\partial^2 h_1}{\partial z_j \partial \bar{z}_k}(0) z_j \bar{z}_k + \sum_{j=1}^d \frac{\partial^2 h_1}{\partial z_1 \partial u_j}(0) z_1 w_j + \sum_{j,k=1}^d \frac{\partial^2 h_1}{\partial u_j \partial u_k}(0) w_j w_k \right),$$

we have

$$h_1(z, u) = \sum_{j,k=1}^m a_{j,k} z_j \bar{z}_k + O_3(z, u),$$

with $a_{1,1} = 1$. In particular, we still have $\mathcal{L}_{M,0}\left(\frac{\partial}{\partial x_1}\right) = \frac{\partial}{\partial u_1}$.

For a small $\varepsilon_1 > 0$, to be specified later, and a parameter $0 \leq t < \sqrt{\varepsilon_1}$, we let U_t be the solutions of the parameter-dependent Bishop equation

$$(3) \quad U_t = -\mathcal{T}_0(h(W_t, U_t)), \text{ where } W_t = (t\zeta, 0, \dots, 0).$$

Since the data are smooth, we can solve this in the Hölder spaces $C^{k,\alpha}(S^1, \mathbb{R}^d)$ ($k \geq 1$, $0 < \alpha < 1$) for ε_1 sufficiently small. Let the $A_t(\zeta)$ be the corresponding holomorphic disc and set $\gamma_0(s) = A_{\sqrt{s}}(0)$. One reads off from

(3) that the curve γ_0 starts at the origin and runs in the space $i\mathbb{R}_v^d = \{z = u_1 = \dots = u_n = 0\}$. It is shown in [HT] that (i) $\frac{d\gamma_0}{ds}(0)$ is a positive multiple of $\frac{\partial}{\partial v_1}$ and that (ii) γ_0 is on $[0, \varepsilon_1)$ as smooth as we please if ε_1 is sufficiently small.

The curves γ_p will be obtained by varying the initial point in this construction. To this end we first produce a family of coordinates (z_p, w_p) centered at p such that $T_{p=0}M = \{v_p = 0\}$. Clearly this can be achieved by an affine linear change of coordinates Φ_p which depends smoothly on p , where p ranges in some M -neighborhood of the origin. Thus (z_p, w_p) are holomorphic for p fixed, but are only smooth in p . Next we rotate and dilate in such a way that $d\Phi_p(X)$ transforms to $\frac{\partial}{\partial x_1}$ at the origin and such that $\mathcal{L}_{M,0}\left(\frac{\partial}{\partial x_1}\right) = \frac{\partial}{\partial u_1}$. Then we modify (z_p, w_p) again as before in order to simplify second-order terms. Clearly all this can be done by a family of local biholomorphisms depending smoothly on the parameter p . In these last coordinates, which we still denote by (z_p, w_p) , we construct a curve $\tilde{\gamma}_p(s)$, $0 \leq s < \varepsilon_1$, verbatim by the same construction as above. Thus $\tilde{\gamma}_p(s)$ starts at the origin in (z_p, w_p) -space, which corresponds to the point p , and its time derivative at $s = 0$ is a positive multiple of $\frac{\partial}{\partial v_1}$. The desired curve γ_p with $\gamma_p(0) = p$ is then obtained by reversing the coordinate transformations. The nature of the process implies that $\frac{d\gamma_p}{ds}(0)$ is a positive multiple of $X(p)$.

We claim that $\tilde{M}^+ \cup W = \bigcup_{p \in W, 0 \leq s < \varepsilon_1} \gamma_p(s)$ is a manifold with boundary W attached to M along W , provided W is a sufficiently small open neighborhood of the origin in M , and ε_1 is sufficiently small. Actually, optimal regularity results for the Bishop equation yield that the solution, which depends on a finite dimensional set of parameters, has arbitrarily small loss of smoothness with respect to the parameters ([Tu2], see also [MP4]). Thus we have that data depending $C^{k,\alpha}$ -smoothly on all variables and parameters lead to $C^{k,\alpha-0}$ -smooth solutions. As the smoothness in s is as good as needed and we can assume k as large as we please, the mapping $(p, s) \mapsto \gamma_p(s)$ is as smooth as we please. First, the inverse function theorem implies the claim about $\tilde{M}^+ \cup W$. Second, we may extend $(p, s) \mapsto \gamma_p(s)$ to $W \times (-\eta, \varepsilon_1)$ for some $0 < \eta \ll 1$, and obtain an extended manifold $\tilde{M} = \bigcup_{p \in W, -\eta < s < \varepsilon_1} \gamma_p(s)$ as required in (a). The before mentioned fact that $\frac{d\gamma_p}{ds}(0)$ is a positive multiple of $X(p)$ completes the proof of (b).

The proof of (c) is standard. One chooses an M -neighborhood $W' \subset \subset U$ of p_0 so small that every $u \in CR(U)$ can be uniformly approximated on $\overline{W'}$ by holomorphic polynomials P_j , using the Baouendi-Treves approximation theorem [BT1]. Restricting domains again, one restricts the above construction using only discs attached to W' . Then the maximum modulus principle applied to the discs implies that the P_j converge uniformly on $\tilde{M}^+ \cup W$ to a continuous function u_+ , which is CR on \tilde{M}^+ . The proof of Proposition 3.2 is complete \square

PROOF OF PROPOSITION 3.3. – Let us first sketch the geometrical idea. Since $Y = \mathcal{L}_{\tilde{M}, p_0}(X)$ is not complex tangent to \tilde{M} , it is classical (see [HT]) that appropriate $C^{2,\alpha}$ -small discs attached near p_0 to \tilde{M} whose z -coordinates are parallel to $\mathbb{C}X$ are nontangent to \tilde{M} along their boundaries, sticking out along directions which are approximately Y modulo $H_{p_0}\tilde{M}$. This would be enough in order to extend CR functions defined on all of \tilde{M} . In the case at hand, we are only allowed to use discs attached to $\tilde{M}^+ \cup W$. We will construct a family of discs whose boundaries touch M quadratically in exactly one point and obtain CR extension to a $(\dim M + 1)$ -dimensional manifold (distinct from \tilde{M}^+) contained in the union of the discs. Since essentially the same construction is explained in great detail in [MP3, Section 5], it will suffice to give a concise review of what has to be done.

We will first construct a single disc attached to $\tilde{M}^+ \cup W$ whose boundary touches M at p_0 . Choose local coordinates

$$z_1, \dots, z_{m+1}, w_1 = u_1 + iv_1, \dots, w_{d-1} = u_{d-1} + iv_{d-1}$$

centered at p_0 such that \tilde{M} is locally given as a graph $v = \tilde{h}(z, u)$, with $\tilde{h}(0) = 0$, $d\tilde{h}(0) = 0$. After convenient rotations, we may assume that $X = \frac{\partial}{\partial x_1}$, $\mathcal{L}_{\tilde{M}, 0}X = \frac{\partial}{\partial u_1} \bmod H_0\tilde{M}$ and $T_0M = i\mathbb{R}_{y_1} \oplus \mathbb{C}_{z_2, \dots, z_{m+1}}^m \oplus \mathbb{R}_{u_1, \dots, u_{d-1}}^{d-1}$ (by multiplying by some $\zeta \in S^1$, we have rotated X so that $JX \in T_0M$). For technical reasons, we also arrange that

$$(4) \quad \tilde{h}(z_1, 0, \dots, 0, u_1, \dots, u_{d-1}) = c|z_1|^2 + O(|(z, u)|^3), \quad c > 0,$$

by eliminating pure terms of second order.

For $Z_r(\zeta) = (r(1 - \zeta), 0, \dots, 0)$, $0 < r \ll 1$, let

$$A_r(\zeta) = (Z_r(\zeta), U_r(\zeta) + iV_r(\zeta)) = (Z_r(\zeta), U_r(\zeta) + ih(Z_r(\zeta), U_r(\zeta)))$$

be the analytic disc obtained by solving the Bishop equation

$$(5) \quad U_r = -\mathcal{T}_1(h(Z_r, U_r)).$$

Note that $A_r(1) = 0$. The crucial point is that $r(1 - \zeta)$ lies in the right z_1 -halfplane and touches the imaginary axis quadratically at the origin, and that the curvature of its boundary at the origin becomes large for $r \rightarrow 0$. First, it is proved in [MP3, 5.5] that for $r > 0$ small, we have $A_r(\bar{A} \setminus \{1\}) \subset \tilde{M}^+$ and that $A_r(S^1)$ touches M quadratically at 0. There is a geometric estimate for admissible r which is stable under $C^{2,\alpha}$ -small deformations of \tilde{h} . Second, the usual classical argument (see [HT]) based on (4) shows that

$$(6) \quad -\left. \frac{\partial V_r}{\partial \lambda} \right|_{\zeta=1} = r(c', 0, \dots, 0) + o(r),$$

with some $c' > 0$ independent of r (here $\zeta = \lambda + i\eta$).

Let us take for granted, for the moment, that we can construct for a small fixed r a $(\dim M + 1)$ -dimensional manifold \hat{M}^+ attached to M , and containing the image $A_r((1 - \varepsilon, 1))$ of the segment $(1 - \varepsilon, 1)$, such that CR functions extend from M to \hat{M}^+ . Then (6) yields for the normalized outgoing direction at the origin

$$-\left. \frac{\partial v_r}{\partial \lambda}(1) \right| \Big/ \left. \frac{\partial v_r}{\partial \lambda}(1) \right| \rightarrow (c', 0, \dots, 0), \text{ as } r \rightarrow 0.$$

But this implies that we can approximate the element defined by $\frac{\partial}{\partial v_1}$ in T_0M/H_0M by directions of CR extensions. Since $\tilde{\zeta}\left(\frac{\partial}{\partial v_1}\right) > 0$ we have found a direction of extension Z as desired. Now since a sufficiently small $r > 0$ has been fixed, we drop it from the notation and write A .

It remains (i) to construct the manifold \hat{M}^+ and (ii) to establish CR extension from M to \hat{M}^+ . The method for getting (i) is very similar to that of the proof of Proposition 3.2. We construct a family of local holomorphic coordinates

$$z_{1,p}, \dots, z_{m+1,p}, w_{1,p}, \dots, w_{d-1,p}$$

coinciding with the above coordinates for $p = 0$ and satisfying the following properties:

- a) (z_p, w_p) depend C^3 -smoothly on the parameter p , which ranges in a small M -neighborhood W'' of the origin.
- b) For p fixed, (z_p, w_p) are holomorphic and centered at p , and we have $T_0M = i\mathbb{R}_{y_{1,p}} \oplus \mathbb{C}_{z_{2,p}, \dots, z_{m+1,p}}^m \oplus \mathbb{R}_{u_{1,p}, \dots, u_{d-1,p}}^{d-1}$.

We apply the above construction with dependence on the parameter p . This yields discs A_p attached to $\tilde{M}^+ \cup W'$ and touching M quadratically at $A_p(1) = p$. Using regularity results for the Bishop equation ([Tu2], also

[MP4]), we see that $A_p(\zeta)$ depends $C^{2,1/2}$ -smoothly on p and $\zeta \in \bar{\Delta}$. After a further shrinking of W'' and with a smaller ε , we obtain a manifold $\tilde{M}^+ \cup W'' = \bigcup_{p \in W''} A_p((1 - \varepsilon, 1])$ as required in (i).

To show (ii) we just have to remember from the construction of \tilde{M}^+ that every $u \in CR(U)$ can be approximated by holomorphic polynomials P_j which converge uniformly on $\overline{W'} \cup \tilde{M}^+$. Hence the P_j also converge on $\hat{M}^+ \cup W''$ to a continuous CR function extending u . This yields (ii) and completes the proofs of Proposition 3.3, and also of Theorems 1.1 and 2.2 \square

5. Applications to CR meromorphic functions.

One of our motivations is to find concrete applications for the Siegel-type theorems proved in [HN2], [HN3]. In these papers, far reaching global consequences for the field of CR meromorphic functions are proved for CR manifolds satisfying a local extension property E. For a C^∞ -smooth generic CR submanifold M of a complex manifold N , property E means that the canonical restriction mapping $\mathcal{O}_{N,p} \rightarrow \mathcal{CR}_{M,p}$ is surjective for every $p \in M$. Here \mathcal{O}_N and \mathcal{CR}_M denote the sheaf of germs of holomorphic functions on N and the sheaf of germs of C^∞ -smooth CR functions on M , respectively, and $\mathcal{O}_{N,p}$, $\mathcal{CR}_{M,p}$ are their stalks at p . We obtain immediately that a manifold satisfying the assumptions of Theorem 1.1 in a coordinate neighborhood of each of its points has property E. Hence we obtain all results proved in [HN2], [HN3] by carrying the local situation studied here to general manifolds.

Actually the main results in [HN2], [HN3] concern CR meromorphic functions rather than CR functions. Similarly as ordinary meromorphic functions, we define *CR meromorphic functions in the usual sense* on $U \subset M$ as functions which are defined on a dense open subset of U and can be represented near every point $p \in U$ as the quotient p/q of C^∞ -smooth CR functions p, q , where q does not vanish identically on any nonempty open subset. If M has property E, every CR meromorphic function on U is the restriction of a meromorphic function defined on some ambient neighborhood of U in N .

Let M be a smooth compact locally embeddable CR manifold of CR dimension m and CR codimension d , which at each point satisfies the hypotheses of Theorem 1.1. Then the field $\mathcal{K}(M)$ of CR meromorphic functions on M has transcendence degree $k \leq m + d$. If f_1, \dots, f_k is a maximal set of algebraically independent CR meromorphic functions on M , then

$\mathcal{K}(M)$ is a simple finite algebraic extension of the field $\mathbb{C}(f_1, \dots, f_k)$ of rational functions of the f_1, f_2, \dots, f_k . Assuming that M is connected, there is also an equivalence between the algebraic dependence over \mathbb{C} , and the analytic dependence, of a finite set of CR meromorphic functions in $\mathcal{K}(M)$. When M has a projective embedding there is an analogue of Chow's theorem, and $\mathcal{K}(M)$ is isomorphic to the field $\mathcal{R}(Y)$ of rational functions on an irreducible projective algebraic variety Y , and M has a CR embedding in $\text{reg } Y$. For details, and further applications and remarks, see [HN2], [HN3].

6. CR meromorphic mappings according to Harvey and Lawson

An alternative notion of CR meromorphic functions and mappings was suggested by Harvey and Lawson in the context of the complex Plateau problem and studied in [HL], [DH], [DS], [MP1], [MP2]. The following definitions appear in [HL] for hypersurfaces and in [DS] for CR manifolds of arbitrary codimension. Let M be a smooth generic CR submanifold of a complex manifold N of CR dimension m and codimension d , and let X be an arbitrary complex manifold. Then a *CR meromorphic mapping* F in the sense of Harvey-Lawson of an open $U \subset M$ with values in X is given by a triple $(F, \mathcal{D}_F, \Gamma_F)$ with the following properties:

- (a) \mathcal{D}_F is an open dense subset of U ,
- (b) $F : \mathcal{D}_F \rightarrow X$ is a \mathcal{C}^1 -smooth CR mapping,
- (c) the closure of the graph of F in $U \times X$ equals Γ_F and is a local scarred CR cycle of CR dimension m and dimension $2m + d$ in $N \times X$.

In (c) we mean that Γ_F is of locally finite $(\dim M)$ -dimensional Hausdorff measure and contains a closed subset σ (the scar set) of $(\dim M)$ -dimensional Hausdorff measure zero such that (i) $\Gamma_F \setminus \sigma$ is a \mathcal{C}^1 -smooth CR manifold of same dimension and CR dimension as M and (ii) in a neighborhood of every $(p, x) \in \Gamma_F$, integration over $\Gamma_F \setminus \sigma$ yields a *closed current* (see [HL], [DS], [MP1] for full details). If X equals \mathbb{P}^1 , the complex projective line, we also speak of CR meromorphic functions in the sense of Harvey-Lawson.

Meromorphic extension of these CR meromorphic mappings is technically complicated because of the presence of the scar set. Actually it requires a certain machinery to derive the counterpart of Theorem 1.1 for

CR meromorphic functions in the sense of Harvey-Lawson. We can prove this for all manifolds with property E.

THEOREM 6.1. *Let M be a smooth generic CR submanifold of a complex manifold N with property E. Then every CR meromorphic function F in the sense of Harvey-Lawson, defined on an open set $U \subset M$ and with values in \mathbb{P}^1 has a meromorphic extension \tilde{F} to a neighborhood V of U in N . More precisely, the graph $\Gamma_{\tilde{F}} \subset V \times \mathbb{P}^1$ of \tilde{F} satisfies $\Gamma_{\tilde{F}} \cap (U \times \mathbb{P}^1) = \Gamma_F$. In particular, F is a CR meromorphic function in the usual sense.*

A CR manifold M is called *minimal at a point $p \in M$* (in the sense of Tumanov) if there is no germ of a CR manifold $N \subset M$ of the same CR dimension as M and of lower dimension than M containing p .

LEMMA 6.2. *Let M be a smooth generic CR manifold in \mathbb{C}^n with property E. Then M is minimal at every point $p \in M$.*

PROOF. Let us assume that M is not minimal at p_0 . By [BR] there is a smooth CR function u defined on an open neighborhood $U' \subset U$ of p_0 which does not extend holomorphically to any open wedge attached to M near p_0 , in contradiction to property E. □

PROOF OF THEOREM 6.1. – Let F be a \mathbb{P}^1 -valued CR meromorphic function in the sense of Harvey-Lawson defined on $U \subset M$. First we construct a local extension of F to an ambient neighborhood of a given $p_0 \in U$. Because of Lemma 6.2, [MP2, Theorem 1.2] gives meromorphic extension to an open wedge \mathcal{W} attached to a neighborhood U' of p_0 in M . More precisely, there is an open truncated cone $C \subset \mathbb{C}^n$ with vertex at the origin and a meromorphic function \tilde{F} on $\mathcal{W} = U' + C$ which attains F as continuous boundary value on \mathcal{D}_F .

From property E and a Baire category argument (see [HN2]), it follows that smooth CR functions on U' extend holomorphically to a *uniform* ambient neighborhood V' of U' . Since the envelope of meromorphy coincides with the envelope of holomorphy for domains in \mathbb{C}^n , functions which are meromorphic on an arbitrarily thin neighborhood of U' extend meromorphically to V' . For fixed $c \in C$, the rigid translates $U'_\varepsilon = U' + \{\varepsilon c\}$ approach U' for $\varepsilon \downarrow 0$. To obtain the desired extension to a neighborhood of p_0 , it suffices to choose $\varepsilon \in (0, 1)$ so small that $p_0 \in V' + \{\varepsilon c\}$ and to extend \tilde{F} to $V' + \{\varepsilon c\}$.

Now a standard gluing argument yields a meromorphic function \tilde{F} which is defined on a neighborhood V of U in \mathbb{C}^n and coincides with F on \mathcal{D}_F . It remains to prove that $\Gamma_{\tilde{F}} \cap (U \times \mathbb{P}^1) = \Gamma_F$. Near points in \mathcal{D}_F this is obvious. From **(b)** in the definition of CR meromorphic mappings and the corresponding (well known) property of meromorphic functions, we deduce $\Gamma_F \subset \Gamma_{\tilde{F}} \cap (U \times \mathbb{P}^1)$. Assume that there is $(p_0, \zeta_0) \in \Gamma_{\tilde{F}} \cap (U \times \mathbb{P}^1) \setminus \Gamma_F$. This is obviously impossible if p_0 is a point near which \tilde{F} is a smooth mapping. Hence it remains to consider the case in which p_0 lies in the indeterminacy set $\Sigma_{\tilde{F}} = \{p : \{p\} \times \mathbb{P}^1 \subset \Gamma_{\tilde{F}}\}$.

Following [DS], we also consider the indeterminacy set Σ_F of F defined by

$$\Sigma_F = \{p \in U : \{p\} \times \mathbb{P}^1 \subset \Gamma_F\}.$$

Our assumption on p_0 means that $p_0 \notin \Sigma_F$. Let α be a biholomorphism of \mathbb{P}^1 mapping ζ_0 to ∞ . It is observed in [DS] that the set-valued function $\alpha \circ F$ naturally induces a CR distribution g of order one on a neighborhood U_{p_0} of p_0 in M . As M is minimal in p_0 , Tumanov's theorem [Tu1] and the usual extension techniques for CR distributions yield holomorphic extension to an open wedge attached to M at p_0 (assuming g as boundary value in the weak sense). Then an argument with approach manifolds as above yields a holomorphic extension \tilde{g} to a full neighborhood of p_0 . Observing that $\alpha \circ F$ and \tilde{g} are smooth and coincide at points of \mathcal{D}_F , we obtain that the equality $\tilde{F} = \alpha^{-1} \circ \tilde{g}$ holds near p_0 . In particular, \tilde{F} is smooth near p_0 , in contradiction to $p_0 \in \Sigma_{\tilde{F}}$. The proof of Theorem 6.1 is complete \square

REMARK 6.3. It requires only little extra work to derive a corresponding result for CR meromorphic mappings with values in a projective manifold X . Note that in the general case we can only expect $\Gamma_F \subset \Gamma_{\tilde{F}} \cap (U \times X)$.

7. Homogeneous examples

At first glance, it may seem hard to find examples of CR manifolds of kind 3 satisfying the conditions of Theorem 1.1. However, the theory of homogeneous CR manifolds provides many of them in a very natural way (see [MN1], [MN2], [MN3], [MN4], [AMN]). We give a concise description of a class of such homogeneous CR manifolds, and refer to [AMN] for more details.

Let $G^{\mathbb{C}}$ be a complex connected semisimple Lie group, with Lie algebra $\mathfrak{g}^{\mathbb{C}}$, and G a connected real form of $G^{\mathbb{C}}$, with Lie algebra \mathfrak{g} . Fix a Cartan

subgroup H of G , which is maximally noncompact, that is a Cartan subgroup such that a maximal compact torus in H has minimal dimension, and denote by \mathfrak{h} and $\mathfrak{h}^{\mathbb{C}}$ the Lie algebras of H and of its complexification $H^{\mathbb{C}}$. In the set of roots $\mathcal{R} = \mathcal{R}(\mathfrak{g}^{\mathbb{C}}, \mathfrak{h}^{\mathbb{C}})$ choose a subset \mathcal{R}^+ of positive roots, adapted to \mathfrak{g} (cf. [AMN], Proposition 6.1), and let \mathcal{B} be the corresponding set of positive simple roots. To any subset $\Phi \subset \mathcal{B}$ we associate the parabolic subalgebra and subgroup

$$\mathfrak{q}_{\Phi} = \mathfrak{h}^{\mathbb{C}} + \sum_{\alpha \in \mathcal{R}^+} \mathfrak{g}_{\alpha}^{\mathbb{C}} + \sum_{\substack{\alpha \in -\mathcal{R}^+ \\ (\text{supp } \alpha) \cap \Phi = \emptyset}} \mathfrak{g}_{\alpha}^{\mathbb{C}},$$

$$Q_{\Phi} = \text{Norm}_{G^{\mathbb{C}}}(\mathfrak{q}_{\Phi}).$$

Here $\mathfrak{g}_{\alpha}^{\mathbb{C}}$ is the eigenspace in $\mathfrak{g}^{\mathbb{C}}$ of a root α , and $\text{supp } \alpha$ is the support of a root α in \mathcal{B} . The group Q_{Φ} has Lie algebra \mathfrak{q}_{Φ} .

The group $G^{\mathbb{C}}$ acts via the adjoint representation on $\mathfrak{g}^{\mathbb{C}}$ and on linear subspaces of $\mathfrak{g}^{\mathbb{C}}$. Fix a subset $\Phi \subset \mathcal{B}$ and let d be the dimension of \mathfrak{q}_{Φ} . The orbit

$$Y = \{\text{Ad}_{g^{\mathbb{C}}}(g)(\mathfrak{q}_{\Phi}) \mid g \in G^{\mathbb{C}}\} \subset \text{Gr}_d(\mathfrak{g}^{\mathbb{C}})$$

through \mathfrak{q}_{Φ} in the Grassmannian of d -planes in $\mathfrak{g}^{\mathbb{C}}$ is the flag manifold of parabolic subalgebras of $\mathfrak{g}^{\mathbb{C}}$ conjugate by an inner automorphism to \mathfrak{q}_{Φ} . It is a smooth irreducible projective subvariety of $\text{Gr}_d(\mathfrak{g}^{\mathbb{C}})$, isomorphic to $G^{\mathbb{C}}/Q_{\Phi}$. The orbit

$$M = \{\text{Ad}_{g^{\mathbb{C}}}(g)(\mathfrak{q}_{\Phi}) \mid g \in G\} \subset \text{Gr}_d(\mathfrak{g}^{\mathbb{C}})$$

of G through \mathfrak{q}_{Φ} is a smooth generic CR submanifold of Y which, by our choice of the Cartan subgroup and of the system of positive roots, is compact, and is called the minimal orbit of G in Y .

If all local CR functions near a point p of M extend to a full neighborhood of p in Y , then the pair (M, Y) has property E of [HN2], hence the field $\mathcal{K}(M)$ of CR meromorphic functions is isomorphic to the field $\mathcal{R}(Y)$ of rational functions on Y , because Y is the smallest projective variety containing M (see [HN2]).

EXAMPLE 7.1. Identify \mathbb{C}^6 , with the standard basis $\{e_j\}_{1 \leq j \leq 6}$, with the quaternionic vector space \mathbb{H}^3 by setting, for $\lambda \in \mathbb{C}$,

$$\mathbf{j}\lambda e_{2j-1} = \bar{\lambda} e_{2j}, \quad \mathbf{j}\lambda e_{2j} = -\bar{\lambda} e_{2j-1}.$$

Consider the complex flag manifold

$$Y = \{\ell_1 \subset \ell_3 \subset \ell_5 \subset \mathbb{C}^6 \mid \dim \ell_{2j-1} = 2j - 1, 1 \leq j \leq 3\}.$$

Then Y is a compact 13-dimensional complex manifold, homogeneous for the action of $G^{\mathbb{C}} = \mathrm{SL}(6, \mathbb{C})$. Near the point

$$o = (\langle e_1 \rangle_{\mathbb{C}}, \langle e_1, e_2, e_3 \rangle_{\mathbb{C}}, \langle e_1, e_2, e_3, e_4, e_5 \rangle_{\mathbb{C}})$$

it admits a holomorphic chart given by the nonconstant entries of the matrix

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ z_1 & 1 & 0 & 0 & 0 \\ z_2 & 0 & 1 & 0 & 0 \\ z_3 & z_6 & z_9 & 1 & 0 \\ z_4 & z_7 & z_{10} & 0 & 1 \\ z_5 & z_8 & z_{11} & z_{12} & z_{13} \end{pmatrix}.$$

Let M be the real submanifold of Y given by

$$M = \{(\ell_1, \ell_3, \ell_5) \in Y \mid \mathrm{H}\ell_1 \subset \ell_3, \mathrm{H}\ell_3 \subset \ell_5\}.$$

Then M is homogeneous for the action of the real form $G = \mathrm{SL}(3, \mathbb{H})$ of G , and it is a compact real-analytic homogeneous generic CR submanifold of Y . Denoting by A_j the j -th column of A , near the point o the manifold M is defined by the system of equations

$$\begin{cases} \mathrm{rk}(A_1, A_2, A_3, \mathbf{j}A_1) = 3, \\ \mathrm{rk}(A_1, A_2, A_3, A_4, A_5, \mathbf{j}A_1, \mathbf{j}A_2, \mathbf{j}A_3) = 5, \end{cases}$$

which, in the coordinates $\{z_j\}_{1 \leq j \leq 13}$, are

$$\begin{cases} z_2 - \bar{z}_6 + z_1\bar{z}_3 + z_3\bar{z}_9 - z_1\bar{z}_1\bar{z}_6 - z_1\bar{z}_2\bar{z}_9 = 0, \\ z_4 - \bar{z}_8 + z_1\bar{z}_5 + z_3\bar{z}_{11} - z_1\bar{z}_1\bar{z}_8 - z_1\bar{z}_2\bar{z}_{10} = 0, \\ z_{10} - \bar{z}_{12} + z_9\bar{z}_{11} + z_{11}\bar{z}_{13} - z_9\bar{z}_9\bar{z}_{12} - z_9\bar{z}_{10}\bar{z}_{13} = 0, \\ z_5 + \bar{z}_7 - z_1\bar{z}_4 - z_3\bar{z}_{10} + z_1\bar{z}_1\bar{z}_7 + z_1\bar{z}_2\bar{z}_{10} = 0, \end{cases}$$

thus M has CR dimension 5 and codimension 8. The holomorphic tangent space $T_o^{1,0}M$ admits the basis $\{\partial/\partial z_j\}_{j=1,3,9,11,13}$. The space of Levi forms has real dimension 6, and in the basis $\{\partial/\partial z_j\}_{j=1,3,9,11,13}$ it is the space of Hermitian symmetric matrices of the form

$$\begin{pmatrix} 0 & \alpha & 0 & 0 & 0 \\ \bar{\alpha} & 0 & \alpha & \beta & 0 \\ 0 & \bar{\alpha} & 0 & \gamma & 0 \\ 0 & \bar{\beta} & \bar{\gamma} & 0 & \gamma \\ 0 & 0 & 0 & \bar{\gamma} & 0 \end{pmatrix}, \quad \alpha, \beta, \gamma \in \mathbb{C},$$

as the fourth equation does not contribute to the Levi form. In some

characteristic codirections the Levi forms are zero, but in the remaining characteristic codirections, the Levi forms all have signature $(1, 1)$ or signature $(2, 2)$, and M is trace pseudoconcave. Moreover M has kind 3, indeed G_2 has real dimension 16 and G_3 is the whole tangent space of M . By Theorem 1.1 germs of CR distributions on M are real-analytic and extend to germs of holomorphic functions on Y .

With the notation of [AMN], M is the minimal orbit of the simple Lie group of type AII_5 associated to the parabolic subalgebra $\mathfrak{q} = \mathfrak{q}_{\{\alpha_1, \alpha_3, \alpha_5\}}$.

EXAMPLE 7.2. The example above can be generalized to the following pairs of complex flag manifolds, homogeneous for the action of $\text{SL}(2n, \mathbb{C})$, and compact generic CR submanifolds, homogeneous for the action of $\text{SL}(n, \mathbb{H})$:

$$\left\{ \begin{array}{l} Y = \{ \ell_{2j_1-1} \subset \ell_{2j_2-1} \subset \ell_{2j_3-1} \subset \mathbb{C}^{2n} \mid \dim \ell_{2j_k-1} = 2j_k - 1, 1 \leq k \leq 3 \}, \\ M = \left\{ (\ell_{2j_1-1}, \ell_{2j_2-1}, \ell_{2j_3-1}) \in Y \mid \begin{array}{l} \dim(\ell_{2j_k-1} \cap \mathbb{H}\ell_{2j_k-1}) = 2j_k - 2, \quad 1 \leq k \leq 3 \\ \mathbb{H}\ell_{2j_k-1} \subset \ell_{2j_{k+1}-1}, \quad 1 \leq k \leq 2 \end{array} \right\}, \\ \text{for } 1 = j_1 < j_2 < j_3 \leq n. \end{array} \right.$$

In this case the complex dimension of Y , the CR dimension of M , and the CR codimension of M are:

$$\begin{aligned} \dim_{\mathbb{C}} Y &= (2n - 1) + 2(j_2 - 1)(2n - 2j_2 + 1) + 2(j_3 - j_2)(2n - 2j_3 + 1), \\ \dim_{\text{CR}} M &= 2n + 2j_3 - 7, \\ \text{codim}_{\text{CR}} M &= 4((n - j_3)(j_3 - 1) + (j_3 - j_2)(j_2 - 1) + 1). \end{aligned}$$

The space of Levi forms of M has dimension

$$\begin{aligned} (2j_2 - 3)(2j_3 - 2j_2 + 1) + (2j_3 - 2j_2 - 1)(2n - 2j_3 + 2j_2 - 1) + \\ + (2n - 2j_3)(2j_3 - 2j_2), \end{aligned}$$

and all of them, in a suitable basis (as described in [AMN]), have all diagonal entries equal to zero. Hence M is trace pseudoconcave, and has not kind 2. It can be checked that M has kind 3, thus M satisfies the hypotheses of Theorem 1.1 and germs of CR distributions extend holomorphically to a full neighborhood.

With the notation of [AMN] M is the minimal orbit of the simple Lie group of type AII_{2n-1} associated to the parabolic subalgebra $\mathfrak{q} = \mathfrak{q}_{\{\alpha_1, \alpha_{2j_2-1}, \alpha_{2j_3-1}\}}$. By duality, a completely analogous statement holds for the case $1 \leq j_1 < j_2 < j_3 = n$.

EXAMPLE 7.3. Similar to the previous one is the case of pairs of complex flag manifolds, homogeneous for the action of $\mathrm{SL}(2n, \mathbb{C})$, and compact generic CR submanifolds, homogeneous for the action of $\mathrm{SL}(n, \mathbb{H})$:

$$\left\{ \begin{array}{l} Y = \{ \ell_{2j_1-1} \subset \ell_{2j_2-1} \subset \mathbb{C}^{2n} \mid \dim \ell_{2j_k-1} = 2j_k - 1, k = 1, 2 \}, \\ M = \left\{ (\ell_{2j_1-1}, \ell_{2j_2-1}) \in Y \mid \begin{array}{l} \dim(\ell_{2j_k-1} \cap \mathbb{H}\ell_{2j_k-1}) = 2j_k - 2, \quad k = 1, 2 \\ \mathbb{H}\ell_{2j_1-1} \subset \ell_{2j_2-1} \end{array} \right\}, \\ \text{for } 1 < j_1 < j_2 < n. \end{array} \right.$$

Also in this case, M has kind 3 and is trace pseudoconcave.

EXAMPLE 7.4. Let $G^{\mathbb{C}}$ be the connected and simply connected simple complex Lie group of type F_4 , and let $\mathfrak{g}^{\mathbb{C}}$ its Lie algebra. Fix a Cartan subalgebra $\mathfrak{h}^{\mathbb{C}}$ of $\mathfrak{g}^{\mathbb{C}}$ and a system of simple roots $\mathcal{B} = \{\alpha_j\}_{1 \leq j \leq 4}$ of the root system $\mathcal{R} = \mathcal{R}(\mathfrak{g}^{\mathbb{C}}, \mathfrak{h}^{\mathbb{C}})$ (we use the root numbering scheme of Bourbaki, see [AMN], Appendix). Let $\{\omega_j\}_{1 \leq j \leq 4}$ be the set of fundamental weights dual to \mathcal{B} and let $V = V_{\omega_2}$ be the fundamental representation of $G^{\mathbb{C}}$ with highest weight ω_2 . Let Y be the $G^{\mathbb{C}}$ -orbit, in $\mathbb{P}(V)$, of the highest weight root space. With the notation of [AMN], Y is the complex flag manifold of $G^{\mathbb{C}}$ consisting of parabolic subalgebras of $\mathfrak{g}^{\mathbb{C}}$ that are conjugate to $\mathfrak{q}_{\{\alpha_2\}}$.

Inside Y consider the minimal orbit M of the real form of $G^{\mathbb{C}}$ of type F II (also denoted by $F_{4(-20)}$). It is a CR manifold of CR dimension 9 and CR codimension 11. By [AMN], Theorem 9.1, it is of finite type, and by [AMN], Theorem 13.5, it is trace pseudoconcave. Direct computation shows that G_2 has codimension 4 and G_3 is the whole tangent space of M , thus M has kind 3. Hence also in this case we have holomorphic extension for germs of CR distributions.

Acknowledgments. The second and the fourth author would like to thank the Mathematisches Institut der Humboldt-Universität zu Berlin, and Professor Jürgen Leiterer in particular, for their kind hospitality. Furthermore we thank Anna Siano for information about the literature.

REFERENCES

- [AMN] A. ALTOMANI - C. MEDORI - M. NACINOVICH, *The CR structure of minimal orbits in complex flag manifolds*, J. Lie Theory, **16** (2006), pp. 483–530.
 [BR] M. S. BAOUENDI - L. P. ROTHSCHILD, *Cauchy Riemann functions on manifolds of higher codimension in complex space*, Invent. Math., **101** (1990), pp. 45–56.

- [BT1] M. S. BAOUENDI - F. TREVES, *A property of the functions and distributions annihilated by a locally integrable system of complex vector fields*, Ann. Math., **113** (1981), pp. 387–421.
- [BT2] M. S. BAOUENDI - F. TREVES, *About the holomorphic extension of CR functions on real hypersurfaces in complex space*, Duke Math. J., **51**, no. 1 (1984), pp. 77–107.
- [B] E. BISHOP, *Differentiable manifolds in complex Euclidean space*, Duke Math. J., **32** (1965), pp. 1–21.
- [Bo] A. BOGGESE, *CR extendability near a point where the first Levi form vanishes*, Duke Math. J., **48** (1981), pp. 665–684.
- [BP] A. BOGGESE - J. POLKING, *Holomorphic extensions of CR functions*, Duke Math. J., **49** (1982), pp. 757–784.
- [DS] T. C. DINH - F. SARKIS, *Wedge removability of metrically thin sets and application to the CR meromorphic extension*, Math. Z., **238** (2001), pp. 639–653.
- [DH] P. DOLBEAULT - G. HENKIN, *Chaînes holomorphes de bord donné dans un ouvert q -concave de $\mathbb{C}P^n$* , Bull. Soc. Math. France, **125** (1997), pp. 383–445.
- [FR] J. E. FORNAESS - C. REA, *Local holomorphic extendability and nonextendability of CR-functions on smooth boundaries*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), **12**, no. 3 (1985), pp. 491–502.
- [HL] R. HARVEY - B. LAWSON, *On boundaries of complex analytic varieties, Part II*, Ann. Math., **106** (1977), pp. 213–238.
- [HN1] C. D. HILL - M. NACINOVICH, *A weak pseudoconcavity condition for abstract almost CR manifolds*, Invent. Math., **142** (2000), pp. 251–283.
- [HN2] C. D. HILL - M. NACINOVICH, *Fields of CR meromorphic functions*, Rend. Sem. Mat. Univ. Padova, **111** (2004), pp. 179–204.
- [HN3] C. D. HILL - M. NACINOVICH, *Elementary pseudoconcavity and fields of CR meromorphic functions*, Rend. Sem. Mat. Univ. Padova, **113** (2005), pp. 99–115.
- [HN4] C. D. HILL - M. NACINOVICH, *Conormal suspensions of differential complexes*, J. Geom. Anal., **10**, 3 (2000), pp. 496–537.
- [HT] C. D. HILL - G. TAIANI, *Families of analytic discs in \mathbb{C}^n with boundaries on a prescribed CR submanifold*, Ann. Sc. Norm. Pisa, **5** (1978), pp. 327–380.
- [J] B. JÖRNICKE, *Deformation of CR-manifolds, minimal points and CR-manifolds with the microlocal analytic extension property*, J. Geom. Anal., **6** (1996), pp. 555–611.
- [MN1] C. MEDORI - M. NACINOVICH, *Levi-Tanaka algebras and homogeneous CR manifolds*, Compositio Mathematica, **109** (1997), pp. 195–250.
- [MN2] C. MEDORI - M. NACINOVICH, *Classification of semisimple Levi-Tanaka algebras*, Ann. Mat. Pura Appl., **CLXXIV** (1998), pp. 285–349.
- [MN3] C. MEDORI - M. NACINOVICH, *Complete nondegenerate locally standard CR manifolds*, Math. Ann., **317** (2000), pp. 509–526.
- [MN4] C. MEDORI - M. NACINOVICH, *Algebras of infinitesimal CR automorphisms*, J. Algebra, **287** (2005), pp. 234–274.
- [M] J. MERKER, *Global minimality of generic manifolds and holomorphic extendibility of CR functions*, IMRN, **8** (1994), pp. 329–342.
- [MP1] J. MERKER - E. PORTEN, *On the local meromorphic extension of CR meromorphic mappings*, Ann. Polon. Math., **70** (1998), pp. 163–193.

- [MP2] J. MERKER - E. PORTEN, *Metrically thin singularities of integrable CR functions*, Internat. J. Math., **11** (2000), pp. 857–872.
- [MP3] J. MERKER - E. PORTEN, *Characteristic foliations on maximally real submanifolds of \mathbb{C}^n and removable singularities for CR functions*, IMRP, Volume **2006** (2006), Article ID 72069, pp.1–131.
- [MP4] J. MERKER - E. PORTEN, *Holomorphic Extension of CR Functions, Envelopes of Holomorphy, and Removable Singularities*, IMRS, Volume **2006** (2006), Article ID 28925, pp.1–286.
- [NV] M. NACINOVICH - G. VALLI, *Tangential Cauchy-Riemann complexes on distributions*, Ann. Mat. Pura Appl., **146** (1987), pp. 123–160.
- [S] J. SJÖSTRAND, *The FBI transform for CR submanifolds of \mathbb{C}^n* , Prépublications Mathématiques Orsay 1982.
- [T1] J.-M. TRÉPREAU, *Sur le prolongement holomorphe des fonctions CR définies sur une hypersurface réelle de classe C^2* , Invent. Math., **83** (1986), pp. 583–592.
- [T2] J.-M. TRÉPREAU, *Sur la propagation des singularités dans les variétés CR*, Bull. Soc. Math. Fr., **118** (1990), pp. 403–450.
- [T] F. TREVES, *Hypoanalytic Structures: Local Theory*, Princeton Univ. Press 1992.
- [Tu1] A. E. TUMANOV, *Extension of CR-functions into a wedge from a manifold of finite type*, Math. USSR Sb., **64** (1989), pp. 129–140.
- [Tu2] A. E. TUMANOV, *On the propagation of extendability of CR functions*, Complex Analysis and Geometry, Proc. of the Conference in Trento 1995. Lecture Notes in Pure and Appl. Math., **173** (Dekker 1996), pp. 479–498.

Manoscritto pervenuto in redazione il 17 marzo 2009.