

Lasry-Lions Regularization and a Lemma of Ilmanen

PATRICK BERNARD (*)

Let H be a Hilbert space. We define the following inf (sup) convolution operators acting on bounded functions $u : H \rightarrow \mathbb{R}$:

$$T_t u(x) := \inf_y \left(u(y) + \frac{1}{t} \|y - x\|^2 \right)$$

and

$$\tilde{T}_t u(x) := \sup_y \left(u(y) - \frac{1}{t} \|y - x\|^2 \right).$$

We have the relation

$$T_t(-u) = -\tilde{T}_t(u).$$

Recall that these operators form semi-groups, in the sense that

$$T_t \circ T_s = T_{t+s} \quad \text{and} \quad \tilde{T}_t \circ \tilde{T}_s = \tilde{T}_{t+s}$$

for all $t \geq 0$ and $s \geq 0$, as can be checked by direct calculation. Note also that

$$\inf u \leq T_t u(x) \leq u(x) \leq \tilde{T}_t u(x) \leq \sup u$$

for each $t \geq 0$ and each $x \in H$. A function $u : H \rightarrow \mathbb{R}$ is called k -semi-concave, $k > 0$, if the function $x \rightarrow u(x) - \|x\|^2/k$ is concave. We will occasionally consider semi-concave functions which take values in $[-\infty, +\infty)$. The function u is called k -semi-convex if $-u$ is k -semi-concave. A function u is t -semi-concave and upper semi-continuous if and only if it belongs to the image of the operator T_t , this follows from Lemma 1 and Lemma 3 below. A function is called semi-concave if it is k -semi-concave for some $k > 0$. A function u is said $C^{1,1}$ if it is Frechet differentiable and if the gradient of u is Lipschitz. Note that a continuous function $u : H \rightarrow \mathbb{R}$ is $C^{1,1}$

(*) Indirizzo dell'A.: CEREMADE, Université Paris-Dauphine, Place du Maréchal de Lattre de Tassigny, 75775, Paris Cedex 16, France.

E-mail: patrick.bernard@ceremade.dauphine.fr

if and only if it is semi-concave and semi-convex, see Lemma 5. Let us recall two important results in that language:

THEOREM 1 (Lasry-Lions, [6]). *Let u be a bounded function. For $0 < s < t$, the function $\tilde{T}_s \circ T_t u$ is $C^{1,1}$ and, if u is uniformly continuous, then it converges uniformly to u when $t \rightarrow 0$.*

THEOREM 2 (Ilmanen, [5]). *Let $u \geq v$ be two bounded functions on H such that u and $-v$ are semi-concave. Then there exists a $C^{1,1}$ function w such that $u \geq w \geq v$.*

Our goal in the present paper is to “generalize” simultaneously both of these results as follows:

THEOREM 3. *The operator $R_t := \tilde{T}_t \circ T_{2t} \circ \tilde{T}_t$ has the following properties:*

- *Regularization : For each function $f : H \rightarrow \mathbb{R}$ and each $t > 0$, the function $R_t(f)$ is $C^{1,1}$ provided it is locally bounded. This holds for all $t > 0$ if f is bounded.*
- *Approximation : If $f : H \rightarrow \mathbb{R}$ is uniformly continuous, then $R_t(f)$ is $C^{1,1}$ and converges uniformly to f as $t \rightarrow 0$.*
- *Pinching: If there exists a k -semi-concave continuous function u and a k -semi-convex continuous function v such that $v \leq f \leq u$, then, for all $t \in]0, k]$, we have $u \geq R_t(f) \geq v$, and $R_t(f)$ is $C^{1,1}$.*

Theorem 3 does not, properly speaking, generalize Theorem 5. However, it offers a new (although similar) answer to the same problem: approximating uniformly continuous functions on Hilbert spaces by $C^{1,1}$ functions with a simple explicit formula.

Because of its symmetric form, the regularizing operator R_t enjoys some nicer properties than the Lasry-Lions operators. For example, if f is $C^{1,1}$, then it follows from the pinching property that $R_t f = f$ for t small enough.

Theorem 2, can be proved using Theorem 3 by taking $w = R_t u$, for t small enough. Note, in view of Lemma 3 bellow, that $R_t u = \tilde{T}_t \circ T_t u$ when t is small enough.

Theorem 3 can be somehow extended to the case of finite dimensional open sets or manifolds via partition of unity, at the price of loosing the simplicity of explicit expressions. Let M be a paracompact manifold of dimension n , equipped once and for all with an atlas $(\phi_i, i \in \mathfrak{S})$ composed of

charts $\phi_i : B^n \rightarrow M$, where B^n is the open unit ball of radius one centered at the origin in \mathbb{R}^n . We assume in addition that the image $\phi_i(B^n)$ is a relatively compact open set, and that the sets $\phi_i(B^n)$, $i \in \mathfrak{S}$ form a locally finite open covering of the manifold M . Let us fix, once and for all, a partition of the unity g_i subordinated to the open covering $(\phi_i(B^n), i \in \mathfrak{S})$. It means that the function g_i is non-negative with support inside $\phi_i(B^n)$, and that $\sum_i g_i = 1$ (note that this sum is finite at each point). Let us define the operator

$$G_t(u) := \sum_i [R_{ta_i}((g_i u) \circ \phi_i)] \circ \phi_i^{-1},$$

where $a_i, i \in \mathfrak{S}$ are positive real numbers. In this expression, we consider each of the terms $[R_{ta_i}((g_i u) \circ \phi_i)] \circ \phi_i^{-1}$ as defined on the whole manifold M with the value 0 outside of $\phi_i(B^n)$. The sum is then locally finite hence well-defined. We say that a function $u : M \rightarrow \mathbb{R}$ is locally semi-concave if, for each $i \in \mathfrak{S}$, there exists a constant b_i such that the function $u \circ \phi_i - \|\cdot\|^2/b_i$ is concave on B^n .

THEOREM 4. *Let $u \geq v$ be two continuous functions on M such that u and $-v$ are locally semi-concave. Then, the real numbers a_i can be chosen such that, for each $t \in]0, 1]$ and each function f satisfying $u \geq f \geq v$, we have:*

- *The function $G_t(f)$ is locally $C^{1,1}$.*
- *If f is continuous, then $G_t(f)$ converges locally uniformly to f as $t \rightarrow 0$.*
- *$u \geq G_t(f) \geq v$.*

We will give some properties, most of which are well-known, of the operators T_t and \tilde{T}_t in Section 1, and derive the proof of the main results in Section 2.

Notes and Acknowledgements. Theorem 2 appears in Ilmanen's paper [5] as Lemma 4G. Several proofs are sketch there but none is detailed. The proof we detail here follows lines similar to one of the sketches of Ilmanen. This statement also has a more geometric counterpart, Lemma 4E in [5]. A detailed proof of this geometric version is given in [2], Appendix. My attention was attracted to these statements and their relations with recent progresses on sub-solutions of the Hamilton-Jacobi equation (see [4, 1, 7]) by Pierre Cardialaguet, Albert Fathi and Maxime Zavidovique. Albert Fathi and Maxime Zavidovique also recently wrote a detailed proof of

Theorem 1, see [3]. This paper also proves how the geometric version follows from Theorem 2. There are many similarities between the tools used in the present paper and those used in [1]. Moreover, Maxime Zavidovique observed in [7] that the existence of $C^{1,1}$ subsolutions of the Hamilton-Jacobi equation in the discrete case can be deduced from Theorem 2. However, it seems that the main result of [1] (the existence of $C^{1,1}$ subsolutions in the continuous case) can't be deduced easily from Theorem 2. Neither can Theorem 2 be deduced from it.

1. The operators T_t and \tilde{T}_t on Hilbert spaces.

The proofs of the theorems follow from standard properties of the operators T_t and \tilde{T}_t that we now recall in details.

LEMMA 1. *For each function $u : H \rightarrow \mathbb{R}$, the function $T_t u$ (which takes values in $[-\infty, +\infty)$), is t -semi-concave and upper semi-continuous. The function $\tilde{T}_t u$ (which takes values in $(-\infty, +\infty]$), is t -semi-convex and lower semi-continuous. Moreover, if u is k -semi-concave, then for each $t < k$ the function $\tilde{T}_t u$ is $(k - t)$ -semi-concave. Similarly, if u is k -semi-convex, then for each $t < k$ the function $T_t u$ is $(k - t)$ -semi-convex.*

PROOF. We shall prove the statements concerning T_t . We have

$$\begin{aligned} T_t u(x) - \|x\|^2/t &= \inf_y (u(y) + \|y - x\|^2/t - \|x\|^2/t) = \\ &= \inf_y (u(y) + \|y\|^2/t - 2x \cdot y/t), \end{aligned}$$

this function is concave and upper semi-continuous as an infimum of continuous linear functions. On the other hand, we have

$$T_t u(x) + \|x\|^2/l = \inf_y (u(y) + \|y - x\|^2/t + \|x\|^2/l).$$

Setting $f(x, y) := u(y) + \|y - x\|^2/t + \|x\|^2/l$, the function $\inf_y f(x, y)$ is a convex function of x if f is a convex function of (x, y) . This is true if u is k -semi-convex, $t < k$, and $l = k - t$ because we have the expression

$$f(x, y) = u(y) + \|y - x\|^2/t + \|x\|^2/l = (u(y) + \|y\|^2/k) + \left\| \sqrt{\frac{l}{kt}}y - \sqrt{\frac{k}{lt}}x \right\|^2.$$

□

Given a uniformly continuous function $u : H \rightarrow \mathbb{R}$, we define its modulus of continuity $\rho(r) : [0, \infty) \rightarrow [0, \infty)$ by the expression $\rho(r) = \sup_{x,e} u(x + re) - u(x)$, where the supremum is taken on all $x \in H$ and all e in the unit ball of H . The function ρ is non-decreasing, it satisfies $\rho(r + r') \leq \rho(r) + \rho(r')$, and it converges to zero in zero (this last fact is equivalent to the uniform continuity of u). We say that a function $\rho : [0, \infty) \rightarrow [0, \infty)$ is a modulus of continuity if it satisfies these properties. Given a modulus of continuity $\rho(r)$, we say that a function u is ρ -continuous if $|u(y) - u(x)| \leq \rho(\|y - x\|)$ for all x and y in H .

LEMMA 2. *If $u : H \rightarrow \mathbb{R}$ is uniformly continuous, then the functions $T_t u$ and $\tilde{T}_t u$ converge uniformly to u when $t \rightarrow 0$. Moreover, given a modulus of continuity ρ , there exists a non-decreasing function $\varepsilon(t) : [0, \infty) \rightarrow [0, \infty)$ satisfying $\lim_{t \rightarrow 0} \varepsilon(t) = 0$ and such that, for each ρ -continuous bounded function u , we have:*

- $T_t u$ and $\tilde{T}_t u$ are ρ -continuous for each $t \geq 0$.
- $u - \varepsilon(t) \leq T_t u(x) \leq u$ and $u \leq \tilde{T}_t u \leq u + \varepsilon(t)$ for each $t \geq 0$.

PROOF. Let us fix $y \in H$, and set $v(x) = u(x + y)$. We have $u(x) - \rho(\|y\|) \leq v(x) \leq u(x) + \rho(\|y\|)$. Applying the operator T_t gives $T_t u(x) - \rho(y) \leq T_t v(x) \leq T_t u(x) + \rho(y)$. On the other hand, we have

$$T_t v(x) = \inf_z (u(z + y) + \|z - x\|^2/t) = \inf_z (u(z) + \|z - (x + y)\|^2/t) = T_t u(x + y),$$

so that

$$T_t u(x) - \rho(\|y\|) \leq T_t u(x + y) \leq T_t u(x) + \rho(\|y\|).$$

We have proved that $T_t u$ is ρ continuous if u is, the proof for $\tilde{T}_t u$ is the same.

In order to study the convergence, let us set $\varepsilon(t) = \sup_{r>0} (\rho(r) - r^2/t)$. We have

$$\varepsilon(t) = \sup_{r>0} (\rho(r\sqrt{t}) - r^2) \leq \sup_{r>0} ((r + 1)\rho(\sqrt{t}) - r^2) = \rho(\sqrt{t}) + \rho^2(\sqrt{t})/4.$$

We conclude that $\lim_{t \rightarrow 0} \varepsilon(t) = 0$. We now come back to the operator T_t , and observe that

$$u(y) - \|y - x\|^2/t \geq u(x) - \rho(\|y - x\|) + \|y - x\|^2/t \geq u(x) - \varepsilon(t)$$

for each x and y , so that

$$u - \varepsilon(t) \leq T_t u \leq u.$$

□

LEMMA 3. For each function $u : H \rightarrow (-\infty, +\infty]$, we have $\check{T}_t \circ T_t(u) \leq u$ and the equality $\check{T}_t \circ T_t(u) = u$ holds if and only if u is t -semi-convex and lower semi-continuous. Similarly, given a function $v : H \rightarrow [-\infty, +\infty)$, we have $T_t \circ \check{T}_t(v) \geq v$, with equality if and only if v is t -semi-concave and upper semi-continuous.

PROOF. Let us write explicitly

$$\check{T}_t \circ T_t u(x) = \sup_y \inf_z (u(z) + \|z - y\|^2/t - \|y - x\|^2/t).$$

Taking $z = x$, we obtain the estimate $\check{T}_t \circ T_t u(x) \leq \sup_y u(x) = u(x)$. Let us now write

$$\check{T}_t \circ T_t u(x) + \|x\|^2/t = \sup_y \inf_z (u(z) + \|z\|^2/t + (2y/t) \cdot (x - z))$$

which by an obvious change of variable leads to

$$\check{T}_t \circ T_t u(x) + \|x\|^2/t = \sup_y \inf_z (u(z) + \|z\|^2/t + y \cdot (x - z)).$$

We recognize here that the function $\check{T}_t \circ T_t u(x) + \|x\|^2/t$ is the Legendre bidual of the function $u(x) + \|x\|^2/t$. It is well-known that a function is equal to its Legendre bidual if and only if it is convex and lower semi-continuous. \square

2. Proof of the main results.

PROOF OF THEOREM 3. For each function f and each $t > 0$, the function $\check{T}_t \circ T_{2t} \circ \check{T}_t f$ is both t -semi-concave and t -semi-convex. It is t -semi-convex by Lemma 1, and it is semi-concave because $T_{2t}(\check{T}_t f)$ is $2t$ -semi-concave by Lemma 1, which implies, still by Lemma 1, that $\check{T}_t \circ T_{2t} \circ \check{T}_t f$ is t -semi-concave. As a consequence, Lemma 5 below implies that the function $R_t f$ is $C^{1,1}$ provided it is locally bounded. The function $R_t(f)$ is bounded if f is bounded, hence it is $C^{1,1}$ in this case.

In the case where f is uniformly continuous, Lemma 2 implies that

$$f - \varepsilon(2t) \leq R_t(f) \leq f + 2\varepsilon(t).$$

As a consequence, $R_t(f)$ is converging uniformly to f , and it is locally bounded hence $C^{1,1}$.

We now consider two continuous functions u and v such that u and $-v$ are k semi-concave, and such that $v \leq u$. We claim that

$$u \geq f \geq v \implies u \geq T_t \circ \check{T}_t f \geq v \text{ and } u \geq \check{T}_t \circ T_t f \geq v$$

for $t \leq k$. This claim implies that $u \geq \check{T}_t \circ T_{2t} \circ \check{T}_t f \geq v$ when $u \geq f \geq v$ and $t \leq k$. Let us now prove the claim concerning $\check{T}_t \circ T_t$, the other part being similar. Since v is k -semi-convex and continuous, we have $\check{T}_t \circ T_t v = v$ for $t \leq k$, by Lemma 3. Then,

$$u \geq f \geq \check{T}_t \circ T_t f \geq \check{T}_t \circ T_t v = v$$

where the second inequality follows from Lemma 3, and the third from the obvious fact that the operators T_t and \check{T}_t are order-preserving.

We have proved that $v \leq R_t(f) \leq u$ if $v \leq f \leq u$ and $t \leq k$. For $t \in]0, k]$, the function $R_t(f)$ is thus locally bounded hence $C^{1,1}$. \square

PROOF OF THEOREM 4. Let a_i be chosen such that the functions $(g_i u) \circ \phi_i$ and $-(g_i v) \circ \phi_i$ are a_i -semi-concave on \mathbb{R}^n (when extended by 0 outside of B^n). The existence of real numbers a_i with this property follows from Lemma 4 below. Given $u \geq f \geq v$, we can apply Theorem 3 for each i to the functions

$$(g_i u) \circ \phi_i \geq (g_i f) \circ \phi_i \geq (g_i v) \circ \phi_i$$

extended by zero outside of B^n . We conclude that, for $t \in]0, 1]$, the function $R_{ta_i}((g_i f) \circ \phi_i)$ is $C^{1,1}$ and satisfies

$$(g_i u) \circ \phi_i \geq R_{ta_i}((g_i f) \circ \phi_i) \geq (g_i v) \circ \phi_i.$$

As a consequence, the function

$$[R_{ta_i}((g_i f) \circ \phi_i)] \circ \phi_i^{-1},$$

extended as a function on M equal to 0 outside of $\phi_i(B^n)$, is $C^{1,1}$. The function $G_t(f)$ is thus locally a finite sum of $C^{1,1}$ functions hence it is locally $C^{1,1}$. Moreover, we have

$$u = \sum_i g_i u \geq G_t(f) \geq \sum_i g_i v = v.$$

\square

We have used:

LEMMA 4. *Let $u : B^n \rightarrow \mathbb{R}$ be a bounded function such that $u - \|\cdot\|^2/a$ is concave, for some $a > 0$. For each compactly supported non-negative C^2 function $g : B^n \rightarrow \mathbb{R}$, the product gu (extended by zero outside of B^n) is semi-concave on \mathbb{R}^n .*

PROOF. Since u is bounded, we can assume that $u \geq 0$ on B^n . Let $K \subset B^n$ be a compact subset of the open ball B^n which contains the support

of g in its interior. Since the function $u - \|\cdot\|^2/a$ is concave on B_1 it admits super-differentials at each point. As a consequence, for each $x \in B^n$, there exists a linear form l_x such that

$$0 \leq u(y) \leq u(x) + l_x \cdot (y - x) + \|y - x\|^2/a$$

for each $y \in B^1$. Moreover, the linear form l_x is bounded independently of $x \in K$. We also have

$$0 \leq g(y) \leq g(x) + dg_x \cdot (y - x) + C\|y - x\|^2$$

for some $C > 0$, for all x, y in \mathbb{R}^n . Taking the product, we get, for $x \in K$ and $y \in B^n$,

$$u(y)g(y) \leq u(x)g(x) + (g(x)l_x + u(x)dg_x) \cdot (y - x) + C\|y - x\|^2 + C\|y - x\|^3 + C\|y - x\|^4$$

where $C > 0$ is a constant independent of $x \in K$ and $y \in B^n$, which may change from line to line. As a consequence, setting $L_x = g(x)l_x + u(x)dg_x$, we obtain the inequality

$$(L) \quad (gu)(y) \leq (gu)(x) + L_x \cdot (y - x) + C\|y - x\|^2$$

for each $x \in K$ and $y \in B^n$. If we set $L_x = 0$ for $x \in \mathbb{R}^n - K$, the relation (L) holds for each $x \in \mathbb{R}^n$ and $y \in \mathbb{R}^n$. For $x \in K$ and $y \in B^n$, we have already proved it. Since the linear forms L_x , $x \in K$ are uniformly bounded, we can assume that $L_x \cdot (y - x) + C\|y - x\|^2 \geq 0$ for all $x \in K$ and $y \in \mathbb{R}^n - B^n$ by taking C large enough. Then, (L) holds for all $x \in K$ and $y \in \mathbb{R}^n$. For $x \in \mathbb{R}^n - K$ and y outside of the support g , the relation (L) holds in an obvious way, because $gu(x) = gu(y) = 0$, and $L_x = 0$. For $x \in \mathbb{R}^n - K$ and y in the support of g , the relation holds provided that $C \geq \max(gu)/d^2$, where d is the distance between the complement of K and the support of g . This is a positive number since K is a compact set containing the support of g in its interior. We conclude that the function (gu) is semi-concave on \mathbb{R}^n . \square

For completeness, we also prove, following Fathi:

LEMMA 5. *Let $u : H \rightarrow \mathbb{R}$ be a locally bounded function which is both k -semi-concave and k -semi-convex. Then the function u is $C^{1,1}$, and $6/k$ is a Lipschitz constant for the gradient of u .*

PROOF. It is well known that a locally bounded convex function is continuous. We conclude that u is continuous. Let u be a continuous function which is both k -semi-concave and k -semi-convex. Then, for each $x \in H$,

there exists a unique $l_x \in H$ such that

$$|u(x+y) - u(x) - l_x \cdot y| \leq \|y\|^2/k.$$

We conclude that l_x is the gradient of u at x , and we have to prove that the map $x \mapsto l_x$ is Lipschitz. We have, for each x, y and z in H :

$$\begin{aligned} l_x \cdot (y+z) - \|y+z\|^2/k &\leq u(x+y+z) - u(x) \leq l_x \cdot (y+z) + \|y+z\|^2/k \\ l_{(x+y)} \cdot (-y) - \|y\|^2/k &\leq u(x) - u(x+y) \leq l_{(x+y)} \cdot (-y) + \|y\|^2/k \\ l_{(x+y)} \cdot (-z) - \|z\|^2/k &\leq u(x+y) - u(x+y+z) \leq l_{(x+y)} \cdot (-z) + \|z\|^2/k. \end{aligned}$$

Taking the sum, we obtain

$$|(l_{x+y} - l_x) \cdot (y+z)| \leq \|y+z\|^2/k + \|y\|^2/k + \|z\|^2/k.$$

By a change of variables, we get

$$|(l_{x+y} - l_x) \cdot (z)| \leq \|z\|^2/k + \|y\|^2/k + \|z-y\|^2/k.$$

Taking $\|z\| = \|y\|$, we obtain

$$|(l_{x+y} - l_x) \cdot (z)| \leq 6\|z\|\|y\|/k$$

for each z such that $\|z\| = \|y\|$, we conclude that

$$\|l_{x+y} - l_x\| \leq 6\|y\|/k.$$

□

REFERENCES

- [1] P. BERNARD, Existence of $C^{1,1}$ critical sub-solutions of the Hamilton-Jacobi equation on compact manifolds, *Ann. Sci. E. N. S.* (4), **40**, no. 3 (2007), pp. 445–452.
- [2] P. CARDALIAGUET, *Front Propagation Problems with Nonlocal Terms 2*, *J. Math. An. Appl.*, **260** (2001), pp. 572–601.
- [3] A. FATHI - M. ZAVIDOVIQUE, *Insertion of $C^{1,1}$ functions and Ilmanen's Lemma*.
- [4] A. FATHI - A. SICONOLFI, *Existence of C^1 critical subsolutions of the Hamilton-Jacobi equation*. *Invent. Math.*, **155**, no. 2 (2004), pp. 363–388.
- [5] T. ILMANEN, *The Level-Set Flow on a Manifold*, *Proc. Symposia Pure Math.*, **54**, no. 1 (1993), pp. 193–204.
- [6] J. M. LASRY - P. L. LIONS, *A Remark on Regularization in Hilbert Spaces*, *Israel Math. J.*, **55**, no. 3 (1996), pp. 257–266.
- [7] M. ZAVIDOVIQUE, *Existence of $C^{1,1}$ critical subsolutions in discrete weak KAM theory*. To appear in *Journal of Modern Dynamics*.

Manoscritto pervenuto in redazione il 6 luglio 2010.