

# The Schur Multiplier of a Generalized Baumslag-Solitar Group

DEREK J. S. ROBINSON

ABSTRACT - The structure of the Schur multiplier of an arbitrary generalized Baumslag-Solitar group is determined and applications to central extensions are described.

## 1. Introduction and Results.

A *generalized Baumslag-Solitar group*, or *GBS-group*, is the fundamental group of a finite connected graph of groups with infinite cyclic vertex and edge groups. In detail let  $\Gamma$  be a finite connected graph – multiple edges and loops are allowed – with vertex set  $V(\Gamma)$  and edge set  $E(\Gamma)$ . For each edge  $e$  we choose endpoints  $e^+$  and  $e^-$ , and hence a direction for the edge. Infinite cyclic groups  $\langle g_x \rangle$  and  $\langle u_e \rangle$  are assigned to each vertex  $x$  and edge  $e$ . Injective homomorphisms  $\langle u_e \rangle \rightarrow \langle g_{e^+} \rangle$  and  $\langle u_e \rangle \rightarrow \langle g_{e^-} \rangle$  are defined by the assignments

$$u_e \mapsto g_{e^+}^{\omega^+(e)} \text{ and } u_e \mapsto g_{e^-}^{\omega^-(e)},$$

where  $\omega^+(e), \omega^-(e) \in \mathbb{Z}^* = \mathbb{Z} \setminus \{0\}$ . Thus we have a weight function

$$\omega : E(\Gamma) \rightarrow \mathbb{Z}^* \times \mathbb{Z}^*$$

where  $\omega(e) = (\omega^-(e), \omega^+(e))$ . The weighted graph  $(\Gamma, \omega)$  is called a *GBS-graph*.

The GBS-group determined by the weighted graph  $(\Gamma, \omega)$  is the fundamental group

$$G = \pi_1(\Gamma, \omega).$$

(\*) Indirizzo dell'A.: Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA.

E-mail: dsrobins@illinois.edu

To obtain a presentation of  $G$  choose a maximal subtree  $T$  of  $\Gamma$ . Then  $G$  has generators

$$g_x, (x \in V(\Gamma)), \text{ and } t_e, (e \in E(\Gamma) \setminus E(T)),$$

with defining relations

$$\begin{cases} g_{e^+}^{\omega^+(e)} &= g_{e^-}^{\omega^-(e)}, \text{ for } e \in E(T), \\ (g_{e^+}^{\omega^+(e)})^{t_e} &= g_{e^-}^{\omega^-(e)}, \text{ for } e \in E(\Gamma) \setminus E(T). \end{cases}$$

It is known that up to isomorphism  $G$  is independent of the choice of  $T$ : for this and other basic properties of graphs of groups see [1], [3], [10]. If  $\Gamma$  consists of a single loop with weight  $(n, m)$ , then  $\pi_1(\Gamma, \omega)$  is a Baumslag-Solitar group

$$BS(m, n) = \langle t, g \mid (g^m)^t = g^n \rangle.$$

It is easy to see that GBS-groups are torsion-free. They are obviously finitely presented, and in fact every finitely generated subgroup of a GBS-group is either free or a GBS-group ([6], 2.7: see also [4], 1.2), so such groups are coherent, i.e., all finitely generated subgroups are finitely presented. By an important result of Kropholler ([7]) the non-cyclic GBS-groups are exactly the finitely generated groups of cohomological dimension 2 which have an infinite cyclic subgroup commensurable with its conjugates. It is therefore natural to enquire about homology and cohomology of GBS-groups in dimensions 1 and 2.

Here we are concerned with integral homology: of course  $H_1(G) \simeq G_{ab}$ , the abelianization, while  $H_2(G) = M(G)$  is the Schur multiplier of  $G$ . Our principal result describes the structure of the Schur multiplier of an arbitrary GBS-group.

**THEOREM 1.** *Let  $G$  be a generalized Baumslag-Solitar group. Then  $M(G)$  is free abelian of rank  $r_0(G) - 1$  where  $r_0(G)$  is the torsion-free rank of  $G_{ab}$ .*

**COROLLARY 1.** *The Euler characteristic of a GBS-group is 0.*

This follows since the homology groups of a GBS-group  $G$  in dimensions 0, 1, 2 have torsion-free ranks 1,  $r_0(G)$ ,  $r_0(G) - 1$  respectively and the alternating sum of these is zero.

We remark that associated with any GBS-group there is a complex  $K(\Gamma, \omega)$  defined in [4]. It can be shown that the Euler characteristic of this complex is zero and this observation is the basis for a topological – but not necessarily shorter – treatment of Theorem 1. Details will appear elsewhere ([5]).

**T-dependence.**

The structure of  $G_{ab}$ , and hence  $r_0(G)$ , can be found from the abelian presentation of  $G_{ab}$  arising from the standard presentation of the GBS-group  $G$  by the usual method of Smith normal form. However, this is a lengthy process and, as only  $r_0(G)$  is required in order to compute  $M(G)$ , it is worthwhile to give a simpler method.

Let  $G = \pi_1(\Gamma, \omega)$  be a GBS-group and let  $T$  be the chosen maximal subtree of  $\Gamma$ . Suppose that  $e = \langle x, y \rangle \in E(\Gamma) \setminus E(T)$  where  $x \neq y$ . Now there is a unique path in the tree  $T$  from  $x$  to  $y$ , say  $x = x_0, x_1, \dots, x_n = y$ . By reading along this path, we obtain a relation  $g_x^{p_1(e)} = g_y^{p_2(e)}$  where  $p_1(e)$  and  $p_2(e)$  are the respective products of the left and right weight values of the edges in the path from  $x$  to  $y$ . If the vector  $(\omega^-(e), \omega^+(e))$  is a rational multiple of  $(p_1(e), p_2(e))$ , then  $e$  is said to be *T-dependent*, and otherwise  $e$  is *T-independent*. If  $e$  is a loop, then by convention  $p_1(e) = 1 = p_2(e)$  and  $e$  is *T-dependent* precisely when  $\omega^-(e) = \omega^+(e)$ .

The definition of *T-dependence* may be restated as follows.

LEMMA 1. *With the above notation, a non-tree edge  $e = \langle x, y \rangle$  of a GBS-graph is T-dependent if and only if*

$$\frac{\omega^-(e)}{\omega^+(e)} = \frac{p_1(e)}{p_2(e)}.$$

If every non-tree edge of a GBS-graph is *T-dependent*, the GBS-graph is said to be *tree-dependent*. The torsion-free rank of the abelianization of a GBS-group can be computed from the following result.

THEOREM 2. *Let  $G = \pi_1(\Gamma, \omega)$  be a generalized Baumslag-Solitar group. Then*

$$r_0(G) = |E(\Gamma)| - |V(\Gamma)| + 1 + \varepsilon(\Gamma, \omega)$$

where  $\varepsilon(\Gamma, \omega) = 1$  if  $(\Gamma, \omega)$  is *tree-dependent* and otherwise equals 0.

(A variant of this result with a different proof appears in [8], Theorem 1.1). We note that, as a consequence of Theorem 2,  $r_0(G)$  can be found by simply inspecting the graph of the GBS-group  $G$ . Notice also that  $\varepsilon(\Gamma, \omega)$  depends only on the GBS-graph  $(\Gamma, \omega)$ , not on the choice of maximal subtree. Thus the property of tree-dependence is independent of the maximal subtree selected.

We remark that the invariant  $\varepsilon$  is closely related to the centre of a GBS-group and is an important tool in the theory of GBS-groups: it is the subject of an ongoing investigation.

As is well known, knowledge of the structure of the Schur multiplier of a group allows one to draw conclusions about central extensions by the group. As a consequence of Theorem 1 one can determine when all central extensions by a GBS-group  $G$  split, i.e., they are direct products. It is shown in Corollary 4 below that *every central extension by a generalized Baumslag-Solitar group  $G$  splits if and only if  $G_{ab}$  is infinite cyclic.*

## 2. Proof of Theorem 2.

Let  $G = \pi_1(\Gamma, \omega)$  be a GBS-group with  $T$  a maximal subtree of  $\Gamma$ . Then  $G$  has an abelian presentation with generators  $g_x, t_e$ , where  $x \in V(\Gamma)$ ,  $e \in E(\Gamma) \setminus E(T)$ , subject to the defining relations  $g_e^{\omega^-(e)} = g_e^{\omega^+(e)}$ , ( $e \in E(\Gamma)$ ). Put  $G_0 = \langle g_x \mid x \in V(\Gamma) \rangle$ ; then  $G_0 \simeq \pi_1(T, \omega)$  and  $r_0(G_0) \leq 1$  since each pair of generators of  $G_0$  is linearly dependent. Since  $G_0$  has fewer relations than generators, it is infinite and  $r_0(G_0) = 1$ . Of course, the stable elements  $t_e$ , are linearly independent modulo the torsion subgroup of  $G_{ab}$ . Therefore

$$r_0(G) = |E(\Gamma) \setminus E(T)| + \varepsilon,$$

where  $\varepsilon = 1$  if each vertex generator has infinite order modulo  $G'$  and otherwise  $\varepsilon = 0$ . If some non-tree edge  $e$  is  $T$ -independent, then, in the notation of Lemma 1, the relations  $g_e^{\omega^-(e)} = g_e^{\omega^+(e)}$  and  $g_e^{p_1(e)} = g_e^{p_2(e)}$  are independent, which forces each vertex generator to have finite order modulo  $G'$ ; hence  $\varepsilon = 0$ . On the other hand, if all such edges are  $T$ -dependent, i.e.,  $(\Gamma, \omega)$  is tree-dependent, then all vertex generators have infinite order and  $\varepsilon = 1$ . Since

$$|E(\Gamma) \setminus E(T)| = |E(\Gamma)| - (|V(\Gamma)| - 1) = |E(\Gamma)| - |V(\Gamma)| + 1,$$

the result follows on setting  $\varepsilon(\Gamma, \omega) = \varepsilon$ . □

### 3. Proof of Theorem 1.

Let  $G = \pi_1(\Gamma, \omega)$  be a GBS-group with  $T$  a maximal subtree of  $\Gamma$ . We recall the following inequality, which is valid for any finitely presented group  $H$  with  $n$  generators and  $r$  relators:

$$n - r \leq r_0(H) - d(M(H)),$$

where  $d(X)$  is the minimal number of generators of a group  $X$ , (see, for example, [9], p.550). In the present situation we have  $n = |V(\Gamma)| + |E(\Gamma) \setminus E(T)|$  and  $r = |E(\Gamma)|$ , so  $n - r = 1$ . Thus  $d(M(G)) \leq r_0(G) - 1$  and it suffices to prove that  $r_0(M(G)) \geq r_0(G) - 1$ . The proof is by induction on  $|E(\Gamma)|$ , which may be assumed positive.

(i) *We can assume that  $r_0(G) > 1$ , so that  $\Gamma$  is not a tree.*

For if  $r_0(G) = 1$ , then  $d(M(G)) = 0$ . Note that if  $\Gamma$  is a tree, then  $r_0(G) = 1$  since each pair of vertex generators is linearly independent.

(ii) *Case:  $\Gamma$  has a single non-tree edge.*

Let  $e = \langle x, y \rangle$  be the edge which is not in  $T$ . Now  $r_0(G) \leq 2$  by Theorem 2, so  $r_0(G) = 2$  and  $\varepsilon(\Gamma, \omega) = 1$ ; thus  $e$  must be  $T$ -dependent. Apply the five-term homology sequence for the exact sequence  $G' \twoheadrightarrow G \twoheadrightarrow G_{ab}$  to get

$$M(G) \rightarrow M(G_{ab}) \rightarrow G'/[G', G] \rightarrow G_{ab} \rightarrow G_{ab} \rightarrow 1.$$

Note that  $r_0(M(G_{ab})) = 1$  since  $M(G_{ab}) \simeq G_{ab} \wedge G_{ab}$ .

We claim that  $G'/[G', G]$  is finite. To see this write  $t = t_e$  and let  $\omega(e) = (h, k)$ , so that  $(g_y^k)^t = g_x^h$ . Also  $\langle g_x \rangle \cap \langle g_y \rangle = \langle g_x^m = g_y^n \rangle$  where  $m, n \in \mathbb{Z}^*$ . By  $T$ -dependence  $(h, k)$  is a rational multiple of  $(m, n)$ , say  $ih = jm$  and  $ik = jn$ , with  $i, j \in \mathbb{Z}^*$ . Then

$$[g_y, t]^{ik} \equiv [g_y^{ik}, t] \equiv g_y^{-ik} g_x^{ih} \equiv g_y^{-jn} g_x^{jm} \equiv 1 \pmod{[G', G]}.$$

Next for any vertex generator  $g_z$  we have  $g_z^r = g_y^s$  for some  $r, s \in \mathbb{Z}^*$ , and hence

$$[g_z, t]^r \equiv [g_y^s, t] \equiv [g_y, t]^s \pmod{[G', G]}.$$

Finally,  $[g_u, g_v][G', G]$  has finite order for any vertex generators  $g_u, g_v$ . It follows that  $G'/[G', G]$  is periodic, so it is finite.

Returning to the exact homology sequence above, we conclude that  $M(G)$  must be infinite, so that  $r_0(M(G)) \geq 1 = r_0(G) - 1$ , as required.

(iii) *From now on we assume that  $\Gamma$  has at least two non-tree edges.*

Let  $e = \langle x, y \rangle$  be one of the non-tree edges and let the unique path in  $T$  from  $x$  to  $y$  be

$$\langle x_1, x_2 \rangle, \langle x_2, x_3 \rangle, \dots, \langle x_{k-1}, x_k \rangle,$$

where  $x = x_1$  and  $y = x_k$ . Define subgroups  $G_1 = \langle t_e, g_{x_1}, \dots, g_{x_k} \rangle$  and

$$G_2 = \langle t_f, g_x \mid x \in V(\Gamma), f \in E(\Gamma) \setminus E(T), f \neq e \rangle.$$

Then  $G_i = \pi_1(\Gamma_i, \omega)$ ,  $i = 1, 2$ , where  $\Gamma_1, \Gamma_2$  are subgraphs of  $\Gamma$  with  $V(\Gamma_1) = \{x_1, \dots, x_k\}$ ,  $V(\Gamma_2) = V(\Gamma)$  and respective edge sets  $\{e, \langle x_j, x_{j+1} \rangle \mid j = 1, 2, \dots, k-1\}$  and  $E(\Gamma) \setminus \{e\}$ , with restricted weight functions. Furthermore

$$G = G_1 *_U G_2$$

where  $U = \langle g_{x_1}, g_{x_2}, \dots, g_{x_k} \rangle$ . Since  $U \simeq \pi_1(T_0, \omega)$ , with  $T_0$  the path  $\langle x_1, x_2 \rangle, \langle x_2, x_3 \rangle, \dots, \langle x_{k-1}, x_k \rangle$ , we have  $r_0(U) = 1$  and  $M(U) = 0$  by (i).

Next we form the Mayer-Vietoris sequence for the generalized free product  $G = G_1 *_U G_2$ , ([2], p. 51),

$$0 = M(U) \rightarrow M(G_1) \oplus M(G_2) \rightarrow M(G) \rightarrow U_{ab} \rightarrow$$

$$(*) \quad (G_1)_{ab} \oplus (G_2)_{ab} \rightarrow G_{ab} \rightarrow 1.$$

At this point we must distinguish two cases.

(iv) *Case: the graph  $\Gamma$  has a non-tree edge  $e$  which is  $T$ -dependent.*

Apply the Mayer-Vietoris sequence above for the edge  $e$ . Since  $\Gamma_1$  has just one non-tree edge  $e$  and it is  $T$ -dependent in  $\Gamma_1$ , we conclude that  $r_0(G_1) = 2$  and  $M(G_1) \simeq \mathbb{Z}$  by (ii). Also  $UG'_1/G'_1$  is infinite, so the image of  $(G_1)_{ab}$  in the exact sequence (\*) has infinite projection into  $(G)_{ab}$ . Therefore

$$r_0(G) \leq r_0(G_1) + r_0(G_2) - 1 = r_0(G_2) + 1$$

and  $r_0(G_2) \geq r_0(G) - 1$ . By induction on  $|E(\Gamma)|$  the result is true for  $G_2$ , so we have

$$r_0(M(G)) \geq r_0(M(G_1) \oplus M(G_2)) \geq 1 + (r_0(G) - 2) = r_0(G) - 1,$$

as required.

We are now left with the situation:

(v) *Case: all non-tree edges in  $\Gamma$  are  $T$ -independent.*

Choose any non-tree edge  $e$  and apply the sequence (\*) in (iii) for this edge. Since  $e$  is  $T$ -independent,  $r_0(G_1) = 1$  and  $M(G_1) = 0$ . Also  $UG'_1/G'_1$  is finite because  $e$  is  $T$ -independent. By (iii) there is non-tree edge  $f \neq e$  and

$\bar{t} = t_f \in G_2$ . Since  $f$  is  $T$ -independent,  $r_0(\langle \bar{t}, U \rangle) = 1$  and also  $UG'_2/G'_2$  is finite. Consequently the image of  $U_{ab}$  in  $(G_1)_{ab} \oplus (G_1)_{ab}$  is finite. Since  $U_{ab}$  is infinite, it follows from the sequence (\*) that the cokernel of the map  $M(G_1) \oplus M(G_2) \rightarrow M(G)$  is infinite.

By induction hypothesis the result holds for  $G_2$ , so we conclude that

$$r_0(M(G)) \geq r_0(M(G_1)) + r_0(M(G_2)) + 1 = (r_0(G_2) - 1) + 1 = r_0(G_2),$$

since  $M(G_1) = 0$ . Finally, the image of  $U_{ab}$  in the sequence (\*) being finite, we obtain

$$r_0(G) = r_0((G_1)_{ab} \oplus (G_2)_{ab}) = r_0(G_1) + r_0(G_2) = 1 + r_0(G_2).$$

Hence  $r_0(G_2) = r_0(G) - 1$  and  $r_0(M(G)) \geq r_0(G) - 1$ , as required. □


**COROLLARY 2.** *The GBS-group  $\pi_1(\Gamma, \omega)$  has trivial Schur multiplier if and only if  $\Gamma$  is either a tree or else a tree with one further edge and  $\Gamma$  is not tree-dependent.*

**PROOF.** By the theorem  $M(G) = 0$  if and only if  $r_0(G) = 1$ . This condition requires there to be at most one non-tree edge and by Theorem 2 it must be  $T$ -independent. □

**COROLLARY 3.** *Every GBS-group has deficiency 1.*

**PROOF.** Recall that the deficiency  $\text{def}(G)$  of group  $G$  is equal to  $\sup\{n - r\}$  where  $n$  and  $r$  are the respective numbers of generators and relations in an arbitrary finite presentation. If  $G$  is a GBS-group, then  $1 \leq \text{def}(G) \leq r_0(G) - d(M(G)) = 1$ . □

**EXAMPLE.** Consider the GBS-group  $G$  arising from the following GBS-graph,



where the maximal subtree chosen is the path  $x, y, z, u$  and the stable elements are  $r, s, t$  as indicated. Then  $G$  has a presentation with generators

$$r, s, t, g_x, g_y, g_z, g_u$$

and relations

$$(g_x^2)^r = g_x^2, g_x^2 = g_y^{-3}, g_y^4 = g_z^4, g_z^5 = g_u^3, (g_u^{12})^t = g_y^{20}, (g_x^4)^s = g_y^{-1}.$$

All non-tree edges with the exception of  $\langle y, x \rangle$  are  $T$ -dependent. Therefore  $(\Gamma, \omega)$  is not tree-dependent,  $\varepsilon(\Gamma, \omega) = 0$  and  $r_0(G) = |E(\Gamma)| - |V(\Gamma)| + 1 = 3$ . Thus  $M(G) \simeq \mathbb{Z} \oplus \mathbb{Z}$ .

#### 4. Applications to Central Extensions.

We will now apply our results to yield information about central extensions by GBS-groups. Let  $G$  be a GBS-group and  $C$  an abelian group regarded as a trivial  $G$ -module. Denote by  $F$  the periodic subgroup of  $G_{ab}$ ; thus  $G_{ab} \simeq F \oplus \mathbb{Z}^{r_0(G)}$  where  $F$  is finite. By the Universal Coefficients Theorem

$$H^2(G, C) \simeq \text{Ext}(G_{ab}, C) \oplus \text{Hom}(M(G), C) \simeq \text{Ext}(F, C) \oplus \text{Dr } C^{r_0(G)-1}.$$

First we determine when all central extensions of  $C$  by  $G$  are direct products, i.e., when  $H^2(G, C) = 0$ .

**THEOREM 3.** *Let  $G$  be a generalized Baumslag-Solitar group and let  $C \neq 1$  be an abelian group regarded as a trivial  $G$ -module. Then  $H^2(G, C) = 0$  if and only if  $r_0(G) = 1$  and  $C$  is divisible by all primes  $p \in \pi(G_{ab})$ .*

**PROOF.** With the notation used above,  $H^2(G, C) = 0$  if and only if  $\text{Ext}(F, C) = 0$  and  $r_0(G) = 1$ . Since  $F$  is finite and  $\text{Ext}(\mathbb{Z}_n, C) \simeq C/C^n$ , it follows that  $\text{Ext}(F, C) = 0$  if and only if  $C = C^p$  for all  $p \in \pi(G_{ab})$ . (For the elementary properties of  $\text{Ext}$  used here see [9], 7.2).  $\square$

**COROLLARY 4.** *The following conditions on a generalized Baumslag-Solitar group  $G$  are equivalent*

- (i)  $H^2(G, \mathbb{Z}) = 0$ ;
- (ii)  $G_{ab} \simeq \mathbb{Z}$ ;
- (iii)  $H^2(G, C) = 0$  for all abelian groups  $C$ .

**PROOF.** Clearly condition (i) implies that  $\pi(G_{ab})$  is empty and so (ii) holds. Also (ii) implies (iii), while trivially (iii) implies (i).  $\square$



For example, if  $G = BS(m, n)$ , then  $G_{ab} \simeq \mathbb{Z} \oplus \mathbb{Z}_{|m-n|}$ , so that  $G$  has the property of Corollary 4 if and only if  $|m - n| = 1$ .

There are corresponding results for homology, which can be proved in an analogous way by using the Universal Coefficients Theorem for homology,

$$H_2(G, C) \simeq \text{Tor}(G_{ab}, C) \oplus (M(G) \otimes C),$$

and elementary properties of Tor, (see [9], 7.1).

**THEOREM 4.** *Let  $G$  be a generalized Baumslag-Solitar group and let  $C \neq 1$  an abelian group regarded as a trivial  $G$ -module. Then  $H_2(G, C) = 0$  if and only if  $r_0(G) = 1$  and  $C_p = 1$  for all primes  $p \in \pi(G_{ab})$ .*

**COROLLARY 5.** *The following conditions on a generalized Baumslag-Solitar group  $G$  are equivalent:*

- (i)  $H_2(G, \mathbb{Q}/\mathbb{Z}) = 0$ ;
- (ii)  $G_{ab} \simeq \mathbb{Z}$ ;
- (iii)  $H_2(G, C) = 0$  for all abelian groups  $C$ .

#### REFERENCES

- [1] O. BOGOPOLSKI, *Introduction to Group Theory*, European Math. Soc., Zürich, 2008.
- [2] K. S. BROWN, *Cohomology of Groups*, Graduate Texts, vol. 87, Springer, New York, 1982.
- [3] D. E. COHEN, *Combinatorial Group Theory: a Topological Approach*, London Math. Soc., Student Texts, vol. 14, Cambridge, 1989.
- [4] A. L. DELGADO - D. J. S. ROBINSON - M. TIMM, *Generalized Baumslag-Solitar groups and geometric homomorphisms*, J. Pure Appl. Algebra, **215** (2011), pp. 398–410.
- [5] A. L. DELGADO - D. J. S. ROBINSON - M. TIMM, *The topology of generalized Baumslag-Solitar groups*, preprint.
- [6] M. FORESTER, *On uniqueness of JSG decompositions of finitely generated groups*, Comment. Math. Helv., **78** (2003), pp. 740–741.
- [7] P. H. KROPHOLLER, *Baumslag-Solitar groups and some other groups of cohomological dimension two*, Comment. Math. Helv., **65** (4) (1990), pp. 547–558.
- [8] G. LEVITT, *On the automorphism group of generalized Baumslag-Solitar groups*, Geom. Topol., **11** (2007), pp. 473–515.
- [9] J. J. ROTMAN, *An Introduction to Homological Algebra*, 2nd ed., Springer, New York, 2008.
- [10] J.-P. SERRE, *Trees, translated from the French by J. Stillwell*, Springer, Berlin, 1960.

