

Classification of p -groups via their 2-nilpotent multipliers

PEYMAM NIROOMAND (*) – MOHSEN PARVIZI (**)

ABSTRACT – For a p -group of order p^n , it is known that the order of 2-nilpotent multiplier is equal to $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{2}n(n-1)(n-2)+3-s_2(G)}$, for an integer $s_2(G)$. In this article, we characterize all non-abelian p -groups satisfying $s_2(G) \in \{1, 2, 3\}$.

MATHEMATICS SUBJECT CLASSIFICATION (2020) – Primary 20C25; Secondary 20D15.

KEYWORDS – Nilpotent multiplier, Schur multiplier, non-abelian p -groups, 2-capable groups, capable groups, extra-special groups.

1. Preliminaries

The 2-nilpotent multiplier of a group is a generalization of the well-known notion of Schur multiplier. The latter was introduced by J. Schur in his works on projective representations in [15] and plays a considerable role in classifying groups. In fact, 2-nilpotent multiplier is a special case of the more general notion of Baer invariant.

For a group G with a free presentation $G \cong F/R$, the c -nilpotent multiplier of G , $\mathcal{M}^{(c)}(G)$, is defined as

$$\frac{R \cap \gamma_{c+1}(F)}{[R, {}_c F]},$$

in which $\gamma_{c+1}(F)$ is the c -th term of the lower central series of F , and $[R, {}_c F] = [[R, {}_{c-1} F], F]$ (see [4]).

The motivation of studying the 2-nilpotent multiplier comes from [4]. It is the connection to isologism of groups which is an important tool in classifying groups.

(*) *Indirizzo dell'A.*: School of Mathematics and Computer Science, Damghan University, Damghan, Iran; niroomand@du.ac.ir, p_niroomand@yahoo.com

(**) *Indirizzo dell'A.*: Department of Pure Mathematics, Ferdowsi University of Mashhad, Mashhad, Iran; parvizi@math.um.ac.ir

Recall from [6] that a group G which is isomorphic to $H/Z_2(H)$, for some group H , is called 2-capable. Choose a free presentation $G \cong F/R$, and consider the natural epimorphism $\alpha: F/[R, F, f] \rightarrow G$. We may define $Z_2^*(G) = \alpha(Z_2(F/[R, F, F]))$. Proposition 1.2 in [4] allows us to decide when a group G is 2-capable. More precisely, G is 2-capable if and only if $Z_2^*(G) = 1$. There is a somehow different way for detecting 2-capable groups using the notion of 2-nilpotent multiplier. In more detail, for a group G , the natural epimorphism $\mathcal{M}^{(2)}(G) \rightarrow \mathcal{M}^{(2)}(G/N)$ is a monomorphism if and only if N is a subgroup of $Z_2^*(G)$ (see [4, Lemma 2.1]).

Now, we restrict our study to finite p -groups. A famous result of Green shows that for a given finite 2-group G of order p^n , $|\mathcal{M}(G)| = p^{\frac{1}{2}n(n-1)-t(G)}$ for some integer $t(G) \geq 0$. Several authors worked on classifying the structure of G in term of $t(G)$ when $0 \leq t(G) \leq 5$ (see [1, 12–14, 16]). In [10], considering only non-abelian finite p -groups, a Green-type inequality was obtained. The first-named author showed that $|\mathcal{M}(G)| \leq p^{\frac{1}{2}(n-1)(n-2)+1}$, where G is a finite p -group of order p^n , and hence there is an integer $s(G)$ such that $|\mathcal{M}(G)| = p^{\frac{1}{2}(n-1)(n-2)+1-s(G)}$. A similar result for the 2-nilpotent multiplier of finite p -groups appeared in [14]. The authors proved for a non-abelian p -group of order p^n that there exists an integer $s_2(G)$ such that $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{2}n(n-1)(n-2)+3-s_2(G)}$, and the structure of all p -groups are classified when $s_2(G) = 0$. In the present paper, by the same motivation as in [1, 13, 14, 16], we are interested in characterizing p -groups up to isomorphisms when $s_2(G) \in \{1, 2, 3\}$.

Let us start by stating some lemmas which are needed for the present work. In the following lemma, $G_1 \otimes G_2$ denotes the non-abelian tensor product of two arbitrary groups G_1 and G_2 , and $G_1 \wedge G_2$ denotes the non-abelian exterior product. For more information on these two concepts one may see [2]. It is worth noting that if G_1 and G_2 are two groups acting trivially on each other, then $G_1 \otimes G_2$ coincides with the usual tensor product $G_1/G'_1 \otimes G_2/G'_2$ of abelian groups, by [3, Proposition 2.4].

LEMMA 1.1 ([5, Proposition 2], [7, 9]). *Let G be a finite group and $B \trianglelefteq G$. Set $A = G/B$.*

(i) (a) *If $B \subseteq Z_2(G)$, then*

$$|\mathcal{M}^{(2)}(G)| |B \cap \gamma_3(G)| \text{ divides } |\mathcal{M}^{(2)}(A)| \left| \left(B \otimes \frac{G}{\gamma_3(G)} \right) \otimes \frac{G}{\gamma_3(G)} \right|.$$

(b) *The sequence*

$$(B \wedge G) \wedge G \rightarrow \mathcal{M}^{(2)}(G) \rightarrow \mathcal{M}^{(2)}(G/B) \rightarrow B \cap \gamma_3(G) \rightarrow 1$$

is exact.

(ii) $|\mathcal{M}^{(2)}(A)|$ *divides* $|\mathcal{M}^{(2)}(G)| |B \cap \gamma_3(G)| / |[B, G], G]|$.

The following result plays an essential role in the rest of the paper.

LEMMA 1.2 ([8]). *Let G be a finite group. Put $G^{ab} = G/G'$. Then there is a natural isomorphism*

$$\begin{aligned} \mathcal{M}^{(2)}(G \times H) &\cong \mathcal{M}^{(2)}(G) \times \mathcal{M}^{(2)}(H) \\ &\quad \times (G^{ab} \otimes G^{ab}) \otimes H^{ab} \times (H^{ab} \otimes H^{ab}) \otimes G^{ab}. \end{aligned}$$

The following two lemmas are from [14].

LEMMA 1.3. *Let G be an extra-special p -group of order p^{2n+1} .*

- (i) *If $n > 1$, then $\mathcal{M}^{(2)}(G)$ is an elementary abelian p -group of order $p^{\frac{1}{3}(8n^3-2n)}$.*
- (ii) *Suppose that $|G| = p^3$ and p is odd. Then $\mathcal{M}^{(2)}(G) = \mathbb{Z}_p^{(5)}$ if G is of exponent p and $\mathcal{M}^{(2)}(G) = \mathbb{Z}_p \times \mathbb{Z}_p$ if G is of exponent p^2 .*
- (iii) *The quaternion group of order 8 has Klein four-group as the 2-nilpotent multiplier, whereas the 2-nilpotent multiplier of the dihedral group of order 8 is $\mathbb{Z}_2 \oplus \mathbb{Z}_4$.*

LEMMA 1.4. *Let $G = \mathbb{Z}_{p^{m_1}} \oplus \mathbb{Z}_{p^{m_2}} \oplus \dots \oplus \mathbb{Z}_{p^{m_k}}$, where $m_1 \geq m_2 \geq \dots \geq m_k$ and $\sum_{i=1}^k m_i = n$. Then*

- (i) *$|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n+1)}$ if and only if $m_i = 1$ for all i ;*
- (ii) *$|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}n(n-1)(n-2)}$ if and only if $m_1 \geq 2$.*

2. Main results

As mentioned above, we know that the order of the 2-nilpotent multiplier of a finite non-abelian p -group of order p^n is bounded by $p^{\frac{1}{3}n(n-1)(n-2)+3}$, therefore for any group G there exists a non-negative integer $s_2(G)$ for which

$$|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)+3-s_2(G)}.$$

In this paper, we characterize the explicit structures of finite non-abelian p -groups when $s_2(G) \in \{1, 2, 3\}$.

First, we state the following theorem from [14] to prove that the only groups which may have the desired property are those with small derived subgroups.

THEOREM 2.1. *Let G be a p -group of order p^n with $|G'| = p^m$ ($m \geq 1$). Then*

$$|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}(n-m)((n+2m-2)(n-m-1)+3(m-1))+3}$$

and the equality holds if and only if $G \cong E_1 \times \mathbb{Z}_p^{(n-3)}$.

LEMMA 2.2. *Let G be a non-abelian p -group of order p^n with $|G'| \geq p^3$. Then $|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}n(n-1)(n-2)-2}$.*

PROOF. Just use Theorem 2.1 and the fact that n is at least 5. ■

The following lemma has a completely similar proof to that of Lemma 2.2.

LEMMA 2.3. *Let G be a non-abelian p -group of order p^n with $|G'| = p^2$. Then $|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}n(n-1)(n-2)+1}$.*

The following theorem gives an upper bound for the order of the 2-nilpotent multiplier of a finite group G . Since B and G/B act trivially on each other, $B \otimes G/B$ is isomorphic to the usual tensor product $B \otimes (G/G'B)$, by [3, Proposition 2.4].

THEOREM 2.4. *Let G be a p -group and B be a cyclic central subgroup of G . Then*

$$|\mathcal{M}^{(2)}(G)| \leq |\mathcal{M}^{(2)}(G/B)| |(B \otimes G/G'B) \otimes G/G'|.$$

PROOF. Let $G = F/R$ and $B = S/R$ be free presentations for G and B , respectively. Since B is central, we have $[S, F] \subseteq R$, and also $R \cap S' = [R, S]$ because B is cyclic. Now $S' \subseteq R$, and so $S' = [R, S]$.

By definition, we have

$$\mathcal{M}^{(2)}(G) \cong \frac{R \cap \gamma_3(F)}{[R, F, F]} \quad \text{and} \quad \mathcal{M}^{(2)}(G/B) \cong \frac{S \cap \gamma_3(F)}{[S, F, F]},$$

and so

$$|\mathcal{M}^{(2)}(G)| \leq |\mathcal{M}^{(2)}(G/B)| \left| \frac{[S, F, F]}{[R, F, F]} \right|.$$

The proof is completed if there exists a well-defined epimorphism

$$\bar{\psi}: S/R \otimes F/SF' \otimes F/RF' \longrightarrow \frac{[S, F, F]}{[R, F, F]}.$$

To get this, considering the universal property of the usual tensor product of abelian groups, it is enough to produce a well-defined multi-linear map ψ by the rule

$$\psi(sR, f_1SF', f_2RF') = [s, f_1, f_2][R, F, F].$$

First we show that

$$[sr, f_1s'\gamma', f_2r'\gamma] \equiv [s, f_1, f_2] \pmod{[R, F, F]}$$

where $r, r' \in R, s, s' \in S$ and $\gamma, \gamma' \in F'$.

Expanding the commutator on the left hand side we have $[sr, f_1s'\gamma', f_2r'\gamma] = [sr, f_1s'\gamma', r'\gamma][sr, f_1s'\gamma', f_2][sr, f_1s'\gamma', f_2, r'\gamma]$. Trivially, $[sr, f_1s'\gamma', f_2, r'\gamma] \in [S, F, F, F]$, but $[S, F] \subseteq R$, hence $[S, F, F, F] \subseteq [R, F, F]$. On the other hand, $[sr, f_1s'\gamma', r'\gamma] = [sr, f_1s'\gamma', \gamma][sr, f_1s'\gamma', r']$, which is contained in $[S, F, F][S, F, R]$. A simple use of the three subgroup lemma shows that the latter is contained in $[R, F, F]$. We claim that $[sr, f_1s'\gamma', f_2r'\gamma] \equiv [sr, f_1s'\gamma', f_2] \pmod{[R, F, F]}$. Using commutator calculus again, we get

$$[sr, f_1s'\gamma', f_2] = [sr, s'\gamma', f_2][sr, s'\gamma', f_2, [sr, f_1]^{s'\gamma'}][[sr, f_1]^{s'\gamma'}, f_2].$$

It is easy to see that

$$[sr, s'\gamma', f_2][sr, s'\gamma', f_2, [sr, f_1]^{s'\gamma'}] \in [S, SF', F] = [S, S, F][S, F', F]$$

but we have

$$[S, S, F] = [S', F] = [R, S, F] \subseteq [R, F, F]$$

and

$$[S, F', F] \subseteq [S, F, F, F] = [R, F, F].$$

Finally, $[[sr, f_1]^{s'\gamma'}, f_2] = [sr, f_1, f_2][sr, f_1, f_2, [sr, f_1, s'\gamma']][sr, f_1, s'\gamma', f_2]$, and for the last two we have $[sr, f_1, f_2, [sr, f_1, s'\gamma']][sr, f_1, s'\gamma', f_2] \in [S, F, F, F] \subseteq [R, F, F]$. The first one can be decomposed as

$$\begin{aligned} [sr, f_1, f_2] &= [s, f_1, f_2][s, f_1, f_2, [s, f_1, r]] \\ &\quad \cdot [s, f_1, r, f_2][[s, f_1]^r, f_2, [r, f_1]][r, f_1, f_2], \end{aligned}$$

and we have

$$\begin{aligned} [s, f_1, f_2, [s, f_1, r]] &[s, f_1, r, f_2] \cdot [[s, f_1]^r, f_2, [r, f_1]][r, f_1, f_2] \\ &\in [S, F, F, F][R, F, F] \subseteq [R, F, F]. \end{aligned}$$

The multi-linearity of this mapping follows by a straightforward application of commutator calculus. ■

Considering Lemmas 2.2 and 2.3, in order to characterize all p -groups with $s_2(G) \in \{1, 2, 3\}$, it is enough to work with p -groups with $|G'| \leq p^2$. First we deal with those groups having commutator subgroup of order p . If G/G' is not elementary abelian, we have:

LEMMA 2.5. *Let G be a p -group of order p^n with G' of order p . If G/G' is not elementary abelian, then*

$$|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}n(n-1)(n-2)-2}.$$

PROOF. We use Theorem 2.4 with $B = G'$, to get

$$|\mathcal{M}^{(2)}(G)| \leq |\mathcal{M}^{(2)}(G/G')| |G' \otimes G/G' \otimes G/G'|.$$

Since G/G' is not elementary abelian, by using Lemma 1.4 we have

$$|\mathcal{M}^{(2)}(G/G')| \leq p^{\frac{1}{3}(n-1)(n-2)(n-3)}.$$

Since $|G' \otimes G/G' \otimes G/G'| \leq p^{(n-2)^2}$, we get the result. ■

Now we may assume that G/G' is elementary abelian. In [10, Lemma 2.1] p -groups with $G' = \phi(G)$ (the Frattini subgroup) of order p are classified as the central product of an extra-special p -group H by the center $Z(G)$ of G ; that is, $G = H \cdot Z(G)$. Now, depending on how G' embeds into $Z(G)$, we have the following lemma which has a straightforward proof.

LEMMA 2.6. *Let G be a p -group with $G' = \phi(G)$ of order p . Then:*

- (i) *If G' is a direct summand of $Z(G)$, then $G = H \times K$ for some finite abelian group K .*
- (ii) *If G' is not a direct summand of $Z(G)$, then $G = (H \cdot \mathbb{Z}_{p^2}) \times K$ where K is a finite abelian p -group.*

PROOF. As G is a p -group and $|G'| = p$, we have $G' \subseteq Z(G)$. Consider G/G' as a vector space over \mathbb{Z}_p and let H/G' be a complement to $Z(G)/G'$ in it. It is easy to see that $G = H \cdot Z(G)$ and $H \cap Z(G) = G'$. Now, if G' is a direct summand of $Z(G)$, then we have $Z(G) = G' \times K$ for some abelian subgroup K of $Z(G)$ and hence $G = H \times K$. If G' is not a direct summand of $Z(G)$, we have $\exp(Z(G)) = p^2$, because G/G' is an elementary abelian p -group and $G' \subseteq Z(G)$. Now it is easy to see that $Z(G) = \mathbb{Z}_{p^2} \times K$ and $G' \subseteq \mathbb{Z}_{p^2}$, so we can write $G = (H \cdot \mathbb{Z}_{p^2}) \times K$. ■

As we consider the groups for which G/G' is elementary abelian, we have only the following two cases:

- (1) $G = H \times T$,
- (2) $G = H \cdot \mathbb{Z}_{p^2} \times T$,

where T is an elementary abelian p -group. By Lemma 1.2, without loss of generality we can assume that $Z(G) = \mathbb{Z}_{p^2}$. For the groups of type (1) we have the following theorem.

THEOREM 2.7. *Let $G = H \times T$, where H is an extra-special p -group and T is an elementary abelian p -group. Then:*

- (i) *If $H = E_1$ then $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)+3}$.*
- (ii) *If $H = D_8$ then $|\mathcal{M}^{(2)}(G)| = 2^{\frac{1}{3}n(n-1)(n-2)+1}$.*
- (iii) *In all other cases, $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)}$.*

PROOF. It is just straightforward computations using Lemmas 1.2 and 1.3. ■

For the groups of type (2), first we compute the order of the 2-nilpotent multiplier of $H \cdot \mathbb{Z}_{p^2}$. It should be noted that, as mentioned before Theorem 2.7, we may assume that $Z(G) = \mathbb{Z}_{p^2}$.

THEOREM 2.8. *With the above notation and assumptions, let $G = H \cdot \mathbb{Z}_{p^2}$ be of order p^n . Then $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)}$.*

PROOF. Using Theorem 2.4 with $B = \mathbb{Z}_{p^2}$, we get

$$|\mathcal{M}^{(2)}(G)| \leq |\mathcal{M}^{(2)}(G/\mathbb{Z}_{p^2})| |\mathbb{Z}_{p^2} \otimes G/\mathbb{Z}_{p^2} \otimes G/G'|.$$

In order to compute $|\mathcal{M}^{(2)}(G/\mathbb{Z}_{p^2})|$, we have

$$\frac{G}{\mathbb{Z}_{p^2}} = \frac{H \cdot \mathbb{Z}_{p^2}}{\mathbb{Z}_{p^2}} \cong \frac{H}{H \cap \mathbb{Z}_{p^2}}.$$

But as we had in the proof of Lemma 2.6, $H \cap \mathbb{Z}_{p^2} = G'$. Therefore, $G/\mathbb{Z}_{p^2} \cong H/G'$. By assumption, $|H| = p^{2m+1}$, so H/G' is an elementary abelian p -group of order p^{2m} , hence using Lemma 1.2 and the multi-linearity of the tensor product of abelian groups, we have

$$|\mathcal{M}^{(2)}(G/\mathbb{Z}_{p^2})| = p^{\frac{1}{3}2m(2m+1)(2m-1)} \quad \text{and} \quad |\mathbb{Z}_{p^2} \otimes G/\mathbb{Z}_{p^2} \otimes G/G'| = p^{(2m+1)^2}.$$

After some computations, one gets $|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}n(n-1)(n-2)}$. Now, Lemma 1.1(a) with $B = G'$ shows that $|\mathcal{M}^{(2)}(G/G')| \leq |\mathcal{M}^{(2)}(G)|$. The result now follows by using Lemma 1.4. ■

Now the following theorem, whose proof is completely similar to the last two ones, completes the groups of type (2).

THEOREM 2.9. *Let $G = H \cdot \mathbb{Z}_{p^2} \times T$ be of order p^n , where T is an elementary abelian p -group and H is an extra-special p -groups. Then $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)}$.*

In the rest we concentrate on the groups with the derived subgroup of order p^2 .

LEMMA 2.10. *Let G be a p -group of order p^n with G' of order p^2 . If $Z(G)$ is not elementary abelian, then $|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}n(n-1)(n-2)-2}$.*

PROOF. Choose $B \subseteq Z(G)$ cyclic of order p^2 and use Theorem 2.4 to obtain

$$|\mathcal{M}^{(2)}(G)| \leq |\mathcal{M}^{(2)}(G/B)| |B \otimes G/B \otimes G/G'|.$$

Since

$$|\mathcal{M}^{(2)}(G/B)| \leq p^{\frac{1}{3}(n-1)(n-2)(n-3)} \quad \text{and} \quad |B \otimes G/B \otimes G/G'| \leq p^{(n-2)^2},$$

we have $|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}n(n-1)(n-2)-2}$, and the result follows. ■

In the class of groups with an elementary abelian center we must consider the following two lemmas.

LEMMA 2.11. *Let G be a p -group of order p^n with G' of order p^2 . Let $Z(G)$ be elementary abelian. If $|Z(G)| \geq p^3$ or $|Z(G)| = p^2$, and $G' \neq Z(G)$, then $|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}n(n-1)(n-2)-2}$.*

PROOF. Let K be a central subgroup of order p with $K \cap G' = 1$. By Lemma 1.1(a), we have $|\mathcal{M}^{(2)}(G)| \leq |\mathcal{M}^{(2)}(G/K)| |K \otimes G/\gamma_3(G) \otimes G/\gamma_3(G)|$. But G/K is a non-abelian p -group with $|(G/K)'| = p^2$, thus $|\mathcal{M}^{(2)}(G/K)| \leq p^{\frac{1}{3}(n-1)(n-2)(n-3)+1}$ by Lemma 2.9. Since $|K \otimes G/\gamma_3(G) \otimes G/\gamma_3(G)| \leq p^{(n-2)^2}$, the result follows. ■

LEMMA 2.12. *Let G be a p -group of order p^n with G' of order p^2 . If G/G' is not elementary, then $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)-2}$.*

PROOF. The result is obtained by a similar argument used in the proof of Lemma 2.5 and Theorems 2.7 and 2.8. ■

The next lemma shows that the same upper bound in Lemma 2.11 works when $Z(G)$ is of order p .

LEMMA 2.13. *Let G be a p -group of order p^n with G' of order p^2 . If $|Z(G)| = p$, then $|\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}n(n-1)(n-2)-2}$.*

PROOF. By using Lemma 1.1(a) when $B = Z(G)$, and Theorems 2.7 and 2.8, the result follows. ■

The last case is the one for which $G' = Z(G) \cong \mathbb{Z}_p \oplus \mathbb{Z}_p$.

THEOREM 2.14. *There is no finite p -group of order p^n with $G' = Z(G) \cong \mathbb{Z}_p \oplus \mathbb{Z}_p$ such that $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)}$.*

PROOF. By contradiction, assume that there is a finite p -group G of order p^n such that $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)}$ and $G' = Z(G) \cong \mathbb{Z}_p \oplus \mathbb{Z}_p$. Let K be a central subgroup of order p in G' ; by Lemma 1.1(a), we have $|\mathcal{M}^{(2)}(G)| \leq |\mathcal{M}^{(2)}(G/K)| |K \otimes G/G' \otimes G/G'|$. Now Theorems 2.7 and 2.8 show that

$$|\mathcal{M}^{(2)}(G/K)| \leq p^{\frac{1}{3}(n-1)(n-2)(n-3)+3},$$

whereas G/G' is elementary abelian by Lemma 2.12. Therefore, $p^{\frac{1}{3}n(n-1)(n-2)} = |\mathcal{M}^{(2)}(G)| \leq p^{\frac{1}{3}(n-1)(n-2)(n-3)+3} p^{(n-2)^2}$, whence $n \leq 5$. Since $n \neq 4$, we have $n = 5$. Now [11, page 345] shows that $G \cong \mathbb{Z}_p^{(4)} \rtimes \mathbb{Z}_p$. By a similar argument used in the proof of [14, Theorem 3.5], we have $|\mathcal{M}^{(2)}(G)| = p^{18}$, which is a contradiction. Hence, the assumption is false and the result follows. ■

We conclude summarizing the achieved results.

THEOREM 2.15. *Let G be a non-abelian p -group of order p^n . Then:*

- (i) *There is no group G with $|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)+2}$.*
- (ii) *$|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)+1}$ if and only if $p = 2$ and $G \cong D_8 \times \mathbb{Z}_2^{(n-3)}$.*
- (iii) *$|\mathcal{M}^{(2)}(G)| = p^{\frac{1}{3}n(n-1)(n-2)}$ if and only if $G \cong H_m \times \mathbb{Z}_p^{(n-2m-1)}$, where H_m is an extra-special p -group of order p^{2m+1} and $m \geq 2$ or $G \cong H_m \cdot \mathbb{Z}_{p^2} \times \mathbb{Z}_p^{(n-2m-2)}$.*

REFERENCES

- [1] Y. G. BERKOVICH, [On the order of the commutator subgroup and the Schur multiplier of a finite \$p\$ -group](#). *J. Algebra* **144** (1991), no. 2, 269–272. Zbl 0739.20005 MR 1140606
- [2] R. BROWN – D. L. JOHNSON – E. F. ROBERTSON, [Some computations of nonabelian tensor products of groups](#). *J. Algebra* **111** (1987), no. 1, 177–202. Zbl 0626.20038 MR 913203
- [3] R. BROWN – J.-L. LODAY, [Van Kampen theorems for diagrams of spaces](#). *Topology* **26** (1987), no. 3, 311–335. Zbl 0622.55009 MR 899052
- [4] J. BURNS – G. ELLIS, [On the nilpotent multipliers of a group](#). *Math. Z.* **226** (1997), no. 3, 405–428. Zbl 0892.20024 MR 1483540
- [5] J. BURNS – G. ELLIS, [Inequalities for Baer invariants of finite groups](#). *Canad. Math. Bull.* **41** (1998), no. 4, 385–391. Zbl 0943.20029 MR 1658215
- [6] M. HALL, JR. – J. K. SENIOR, *The groups of order 2^n ($n \leq 6$)*. The Macmillan Company, New York; Collier Macmillan Ltd., London, 1964. MR 0168631

- [7] A. S.-T. LUE, [The Ganea map for nilpotent groups](#). *J. London Math. Soc. (2)* **14** (1976), no. 2, 309–312. Zbl [0357.20030](#) MR [430103](#)
- [8] M. R. R. MOGHADDAM, [The Baer-invariant of a direct product](#). *Arch. Math. (Basel)* **33** (1979/80), no. 6, 504–511. Zbl [0413.20025](#) MR [570485](#)
- [9] M. R. R. MOGHADDAM, [Some inequalities for the Baer-invariant of a finite group](#). *Bull. Iranian Math. Soc.* **9** (1981/82), no. 1, 5–10. MR [660335](#)
- [10] P. NIROOMAND, [On the order of Schur multiplier of non-abelian \$p\$ -groups](#). *J. Algebra* **322** (2009), no. 12, 4479–4482. Zbl [1186.20013](#) MR [2558872](#)
- [11] P. NIROOMAND, [A note on the Schur multiplier of groups of prime power order](#). *Ric. Mat.* **61** (2012), no. 2, 341–346. Zbl [1305.20021](#) MR [3000665](#)
- [12] P. NIROOMAND, [Characterizing finite \$p\$ -groups by their Schur multipliers, \$t\(G\) = 5\$](#) . *Math. Rep. (Bucur.)* **17(67)** (2015), no. 2, 249–254. Zbl [1374.20017](#) MR [3375732](#)
- [13] P. NIROOMAND, [Classifying \$p\$ -groups by their Schur multipliers](#). *Math. Rep. (Bucur.)* **20(70)** (2018), no. 3, 279–284. Zbl [1424.20006](#) MR [3873102](#)
- [14] P. NIROOMAND – M. PARVIZI, [On the 2-nilpotent multiplier of finite \$p\$ -groups](#). *Glasg. Math. J.* **57** (2015), no. 1, 201–210. Zbl [1311.20010](#) MR [3292687](#)
- [15] J. SCHUR, [Untersuchungen über die Darstellung der endlichen Gruppen durch gebrochene lineare Substitutionen](#). *J. Reine Angew. Math.* **132** (1907), 85–137. Zbl [38.0174.02](#) MR [1580715](#)
- [16] X. M. ZHOU, [On the order of Schur multipliers of finite \$p\$ -groups](#). *Comm. Algebra* **22** (1994), no. 1, 1–8. Zbl [0832.20038](#) MR [1255666](#)

Manoscritto pervenuto in redazione il 3 ottobre 2020.