

Transposed Poisson structures on the Lie algebra of upper triangular matrices

Ivan Kaygorodov and Mykola Khrypchenko

Abstract. We describe transposed Poisson structures on the upper triangular matrix Lie algebra $T_n(F)$, $n > 1$, over a field F of characteristic zero. We prove that, for $n > 2$, any such structure is either of Poisson type or the orthogonal sum of a fixed non-Poisson structure with a structure of Poisson type, and for $n = 2$, there is one more class of transposed Poisson structures on $T_n(F)$. We also show that, up to isomorphism, the full matrix Lie algebra $M_n(F)$ admits only one non-trivial transposed Poisson structure, and it is of Poisson type.

Introduction

Since their origin in the 1970s in Poisson geometry, Poisson algebras have appeared in several areas of mathematics and physics, such as algebraic geometry, operads, quantization theory, quantum groups, and classical and quantum mechanics. One of the natural tasks in the theory of Poisson algebras is the description of all such algebras with fixed Lie or associative part [7, 9, 19].

Recently, Bai, Bai, Guo, and Wu [1] have introduced a dual notion of the Poisson algebra, called a *transposed Poisson algebra*, by exchanging the roles of the two multiplications in the Leibniz rule defining a Poisson algebra. A transposed Poisson algebra defined this way not only shares some properties of a Poisson algebra, such as the closedness under tensor products and the Koszul self-duality as an operad, but also admits a rich class of identities [1, 3, 4, 14, 15, 17]. It is important to note that a transposed Poisson algebra naturally arises from a Novikov–Poisson algebra by taking the commutator Lie algebra of its Novikov part.

Thanks to [3], any unital transposed Poisson algebra is a particular case of a “contact bracket” algebra and a quasi-Poisson algebra. In a recent paper by Ferreira,

Mathematics Subject Classification 2020: 17A30 (primary); 15B30, 17B40, 17B63 (secondary).

Keywords: transposed Poisson algebra, upper triangular matrix Lie algebra, full matrix Lie algebra, δ -derivation.

Kaygorodov, and Lopatkin a relation between $\frac{1}{2}$ -derivations of Lie algebras and transposed Poisson algebras has been established [5]. These ideas were used to describe all transposed Poisson structures on Witt and Virasoro algebras in [5]; on twisted Heisenberg–Virasoro, Schrödinger–Virasoro and extended Schrödinger–Virasoro algebras in [20]; on oscillator Lie algebras in [3]; on Schrödinger algebra in $(n + 1)$ -dimensional space-time in [18]; on Witt type Lie algebras in [12]; on generalized Witt algebras in [11] and Block Lie algebras in [10, 11]. Any complex finite-dimensional solvable Lie algebra was proved to admit a non-trivial transposed Poisson structure [13]. The algebraic and geometric classification of 3-dimensional transposed Poisson algebras was given in [2]. Also, see [8, Section 7.3], and the references therein for similar studies. For the list of actual open questions on transposed Poisson algebras, see [3].

In this paper, we describe transposed Poisson structures on the upper triangular matrix Lie algebra $T_n(F)$ over a field F of characteristic zero. To this end, we first characterize $\frac{1}{2}$ -derivations of $T_n(F)$ in Proposition 10. Then the case $n > 2$ is solved in Theorem 11, and the case $n = 2$ is treated separately in Theorem 12. Namely, we prove in Theorem 11 that any transposed Poisson structure on $T_n(F)$, $n > 2$, is either of Poisson type or the orthogonal sum of the fixed non-Poisson structure

$$e_{11} \cdot e_{11} = -e_{11} \cdot e_{nn} = e_{nn} \cdot e_{nn} = e_{1n}$$

with a structure of Poisson type. If $n = 2$, then there appears one more separate class of transposed Poisson structures on $T_n(F)$ consisting of the (non-orthogonal) sums of a structure of the family

$$\begin{aligned} e_{11} \cdot e_{11} &= ce_{11}, & e_{11} \cdot e_{12} &= -e_{12} \cdot e_{22} = ce_{12}, \\ e_{11} \cdot e_{22} &= -e_{22} \cdot e_{22} = ce_{22}, & c &\neq 0, \end{aligned}$$

with a structure of Poisson type. As a complementary result, we prove in Theorem 14 that the full matrix Lie algebra $M_n(F)$ admits only one non-trivial transposed Poisson structure $e_{ii} \cdot e_{jj} = \delta$, $1 \leq i, j \leq n$, and it is of Poisson type. In fact, it is isomorphic to the extension by zero of the product $\delta \cdot \delta = \delta$ as observed in Remark 15.

1. Definitions and preliminaries

All the algebras below will be over a field F of characteristic zero and all the linear maps will be F -linear, unless otherwise stated. The notation $\langle S \rangle$ means the F -subspace generated by S .

Definition 1. Let \mathfrak{L} be a vector space equipped with two non-zero bilinear operations \cdot and $[\cdot, \cdot]$. The triple $(\mathfrak{L}, \cdot, [\cdot, \cdot])$ is called a *transposed Poisson algebra* if (\mathfrak{L}, \cdot)

is a commutative associative algebra and $(\mathfrak{L}, [\cdot, \cdot])$ is a Lie algebra that satisfies the following compatibility condition:

$$2z \cdot [x, y] = [z \cdot x, y] + [x, z \cdot y]. \quad (1)$$

Transposed Poisson algebras were first introduced in a paper by Bai, Bai, Guo, and Wu [1].

Definition 2. Let $(\mathfrak{L}, [\cdot, \cdot])$ be a Lie algebra. A *transposed Poisson algebra structure* on $(\mathfrak{L}, [\cdot, \cdot])$ is a commutative associative multiplication \cdot on \mathfrak{L} which makes $(\mathfrak{L}, \cdot, [\cdot, \cdot])$ a transposed Poisson algebra.

Definition 3. Let $(\mathfrak{L}, [\cdot, \cdot])$ be an algebra and $\varphi : \mathfrak{L} \rightarrow \mathfrak{L}$ a linear map. Then φ is a $\frac{1}{2}$ -*derivation* if it satisfies

$$\varphi([x, y]) = \frac{1}{2}([\varphi(x), y] + [x, \varphi(y)]). \quad (2)$$

Observe that $\frac{1}{2}$ -derivations are a particular case of δ -derivations introduced by Filippov in [6]. The space of all $\frac{1}{2}$ -derivations of an algebra \mathfrak{L} will be denoted by $\Delta(\mathfrak{L})$. It is easy to see from (2) that $[\mathfrak{L}, \mathfrak{L}]$ and $\text{Ann}(\mathfrak{L})$ are invariant under any $\frac{1}{2}$ -derivation of \mathfrak{L} .

Definitions 1 and 3 immediately imply the following key Lemma.

Lemma 4. Let $(\mathfrak{L}, [\cdot, \cdot])$ be a Lie algebra and \cdot a new binary (bilinear) operation on \mathfrak{L} . Then $(\mathfrak{L}, \cdot, [\cdot, \cdot])$ is a transposed Poisson algebra if and only if \cdot is commutative and associative and for every $z \in \mathfrak{L}$ the multiplication by z in (\mathfrak{L}, \cdot) is a $\frac{1}{2}$ -derivation of $(\mathfrak{L}, [\cdot, \cdot])$.

The basic example of a $\frac{1}{2}$ -derivation is the multiplication by a field element. Such $\frac{1}{2}$ -derivations will be called *trivial*.

Theorem 5. Let \mathfrak{L} be a Lie algebra without non-trivial $\frac{1}{2}$ -derivations. Then all transposed Poisson algebra structures on \mathfrak{L} are trivial.

Another well-known class of $\frac{1}{2}$ -derivations of $(\mathfrak{L}, [\cdot, \cdot])$ is formed by linear maps $\mathfrak{L} \rightarrow \text{Ann}(\mathfrak{L})$ annihilating $[\mathfrak{L}, \mathfrak{L}]$. If $(\mathfrak{L}, [\cdot, \cdot])$ is a Lie algebra, such $\frac{1}{2}$ -derivations of \mathfrak{L} correspond to the following transposed Poisson structures on \mathfrak{L} . Denote by $Z(\mathfrak{L})$ the center of \mathfrak{L} and fix a complement V of $[\mathfrak{L}, \mathfrak{L}]$ in \mathfrak{L} . Then any commutative associative product $* : V \times V \rightarrow Z(\mathfrak{L})$ defines a transposed Poisson algebra structure \cdot on \mathfrak{L} by means of

$$(a_1 + a_2) \cdot (b_1 + b_2) = a_1 * b_1, \quad (3)$$

where $a_1, b_1 \in V$ and $a_2, b_2 \in [\mathfrak{L}, \mathfrak{L}]$. Indeed, the right-hand side of (1) is zero, because $z \cdot x, z \cdot y \in Z(\mathfrak{L})$, and the left-hand side of (1) is zero by (3), because $[x, y] \in [\mathfrak{L}, \mathfrak{L}]$.

We say that \cdot is *the extension by zero* of $*$. Observe that \cdot is, at the same time, a usual Poisson structure on $(\mathfrak{L}, [\cdot, \cdot])$. Thus, such transposed Poisson structures are said to be *of Poisson type*.

Given two transposed Poisson structures \cdot_1 and \cdot_2 on $(\mathfrak{L}, [\cdot, \cdot])$, their *sum* $*$, defined by

$$a * b = a \cdot_1 b + a \cdot_2 b,$$

is clearly commutative and satisfies (1). In general, $*$ may be non-associative, but it is associative, if

$$\mathfrak{L} \cdot_1 \mathfrak{L} \subseteq \text{Ann}(\mathfrak{L}, \cdot_2) \quad \text{and} \quad \mathfrak{L} \cdot_2 \mathfrak{L} \subseteq \text{Ann}(\mathfrak{L}, \cdot_1).$$

In this case we say that \cdot_1 and \cdot_2 are *orthogonal*, and $*$ is the *orthogonal sum* of \cdot_1 and \cdot_2 .

Let \cdot be a transposed Poisson algebra structure on a Lie algebra $(\mathfrak{L}, [\cdot, \cdot])$. Then any automorphism ϕ of $(\mathfrak{L}, [\cdot, \cdot])$ induces the transposed Poisson algebra structure $*$ on $(\mathfrak{L}, [\cdot, \cdot])$ given by

$$x * y = \phi(\phi^{-1}(x) \cdot \phi^{-1}(y)), \quad x, y \in \mathfrak{L}.$$

Clearly, ϕ is an isomorphism of transposed Poisson algebras $(\mathfrak{L}, \cdot, [\cdot, \cdot])$ and $(\mathfrak{L}, *, [\cdot, \cdot])$.

2. Transposed Poisson structures on the upper triangular matrix Lie algebra

2.1. Upper triangular matrix algebra

Let n be a positive integer. Denote by $T_n(F)$ the algebra of upper triangular $n \times n$ matrices over F . The usual matrix product on $T_n(F)$ will be denoted by the concatenation, and the commutator product by $[a, b] = ab - ba$. Following the terminology of incidence algebras, we denote the identity matrix by δ and, given $a \in T_n(F)$ and $1 \leq i, j \leq n$, we write $a(i, j)$ for the (i, j) -entry of a . The algebra $T_n(F)$ has the natural basis formed by the matrix units e_{ij} , $1 \leq i \leq j \leq n$, where $e_{ij}(k, l) = \delta(i, k)\delta(j, l)$. It is well known that

$$Z(T_n(F)) = \langle \delta \rangle \quad \text{and} \quad [T_n(F), T_n(F)] = \langle e_{ij} \mid 1 \leq i < j \leq n \rangle \quad \text{for all } n \geq 1.$$

Moreover,

$$Z([T_n(F), T_n(F)]) = \langle e_{1n} \rangle \quad \text{for all } n > 1. \quad (4)$$

Clearly, for all $a \in T_n(F)$ and $1 \leq i \leq j \leq n$ we have

$$e_{ij}a = \sum_{j \leq k} a(j, k)e_{ik}, \quad ae_{ij} = \sum_{l \leq i} a(l, i)e_{lj},$$

so

$$e_{ij}ae_{kl} = \begin{cases} a(j, k)e_{il}, & j \leq k, \\ 0, & j > k. \end{cases}$$

These equalities will be used numerous times throughout the text without any reference.

2.2. On $\frac{1}{2}$ -derivations of the upper triangular matrix Lie algebra

We denote by $\Delta(T_n(F))$ the space of $\frac{1}{2}$ -derivations of the Lie algebra $(T_n(F), [\cdot, \cdot])$.

Lemma 6. *Let $\varphi \in \Delta(T_n(F))$. Then for all $1 \leq i < j \leq n$ we have*

- (i) $\varphi(e_{ij})e_{ii} = 0$;
- (ii) $e_{ij}\varphi(e_{ii}) = \varphi(e_{ii})(j, j)e_{ij}$;
- (iii) $\varphi(e_{ii})(i, j) = -\varphi(e_{jj})(i, j)$;
- (iv) $\varphi(e_{ii})e_{ij} = \varphi(e_{ii})(i, i)e_{ij}$.

Proof. (i) Since $e_{ij} = [e_{ij}, e_{jj}]$, then

$$\begin{aligned} 2\varphi(e_{ij}) &= [\varphi(e_{ij}), e_{jj}] + [e_{ij}, \varphi(e_{jj})] \\ &= \varphi(e_{ij})e_{jj} - e_{jj}\varphi(e_{ij}) + e_{ij}\varphi(e_{jj}) - \varphi(e_{jj})e_{ij}. \end{aligned} \quad (5)$$

Multiplying (5) by e_{ii} on the right (under \cdot), we obtain the desired equality.

(ii) Apply φ to $[e_{ii}, e_{jj}] = 0$:

$$0 = \varphi([e_{ii}, e_{jj}]) = \varphi(e_{ii})e_{jj} - e_{jj}\varphi(e_{ii}) + e_{ii}\varphi(e_{jj}) - \varphi(e_{jj})e_{ii}. \quad (6)$$

If there exists $k > j$, then, multiplying (6) by e_{jj} on the left and by e_{kk} on the right, we obtain

$$\varphi(e_{ii})(j, k) = 0 \quad \text{for } j < k. \quad (7)$$

It follows that

$$e_{ij}\varphi(e_{ii}) = \sum_{j \leq k} \varphi(e_{ii})(j, k)e_{ik} = \varphi(e_{ii})(j, j)e_{ij},$$

as needed.

(iii) This follows by multiplying (6) by e_{ii} on the left and by e_{jj} on the right.

(iv) Applying φ to $e_{ij} = [e_{ii}, e_{ij}]$, we have

$$\begin{aligned} 2\varphi(e_{ij}) &= [\varphi(e_{ii}), e_{ij}] + [e_{ii}, \varphi(e_{ij})] \\ &= \varphi(e_{ii})e_{ij} - e_{ij}\varphi(e_{ii}) + e_{ii}\varphi(e_{ij}) - \varphi(e_{ij})e_{ii}. \end{aligned} \quad (8)$$

If there exists $l < i$, then the multiplication of (8) by e_{ll} on the left and by e_{jj} on the right gives

$$2\varphi(e_{ij})(l, j) = \varphi(e_{ii})(l, i). \quad (9)$$

On the other hand, $[e_{ll}, e_{ij}] = 0$, so

$$0 = \varphi(e_{ll})e_{ij} - e_{ij}\varphi(e_{ll}) + e_{ll}\varphi(e_{ij}) - \varphi(e_{ij})e_{ll}.$$

Multiplying this by e_{ll} on the left and by e_{jj} on the right, we come to

$$\varphi(e_{ij})(l, j) = -\varphi(e_{ll})(l, i). \quad (10)$$

The latter is $\varphi(e_{ii})(l, i)$ by item (iii). Thus, (9) and (10) result in

$$\varphi(e_{ii})(l, i) = 0 \quad \text{for } l < i. \quad (11)$$

Consequently,

$$\varphi(e_{ii})e_{ij} = \sum_{l \leq i} \varphi(e_{ii})(l, i)e_{ik} = \varphi(e_{ii})(i, i)e_{ij},$$

as desired. ■

Lemma 7. *Let $\varphi \in \Delta(T_n(F))$. Then for all $1 \leq i < j \leq n$ we have*

$$\varphi(e_{ij}) = (\varphi(e_{ii})(i, i) - \varphi(e_{ii})(j, j))e_{ij} = (\varphi(e_{jj})(j, j) - \varphi(e_{jj})(i, i))e_{ij}. \quad (12)$$

Proof. By (8) and (i), (ii) and (iv) of Lemma 6 we have

$$2\varphi(e_{ij}) = (\varphi(e_{ii})(i, i) - \varphi(e_{ii})(j, j))e_{ij} + e_{ii}\varphi(e_{ij}). \quad (13)$$

Multiplying (13) by e_{ii} on the left, we get

$$2e_{ii}\varphi(e_{ij}) = (\varphi(e_{ii})(i, i) - \varphi(e_{ii})(j, j))e_{ij} + e_{ii}\varphi(e_{ij}),$$

whence

$$e_{ii}\varphi(e_{ij}) = (\varphi(e_{ii})(i, i) - \varphi(e_{ii})(j, j))e_{ij}.$$

Substituting this into (13) and dividing by 2 (recall that $\text{char}(F) = 0$), we prove the first equality of (12). The second one is obtained from (5) using $\varphi(e_{ij}) \in \langle e_{ij} \rangle$ (which holds by the first equality of (12)). ■

Lemma 8. *Let $\varphi \in \Delta(T_n(F))$.*

- (i) *For all $1 \leq i \leq n$ and $1 \leq j \leq k \leq n$ with $i \notin \{j, k\}$ we have $\varphi(e_{ii})(j, j) = \varphi(e_{ii})(k, k)$.*
- (ii) *For all $1 \leq i < j \leq n$ we have $\varphi(e_{ij})(i, j) = \varphi(e_{1n})(1, n)$.*
- (iii) *For all $1 \leq i \leq n$ and $1 \leq j < k \leq n$ we have $\varphi(e_{ii})(j, k) = 0$, unless $i \in \{1, n\}$ and $(j, k) = (1, n)$.*

Proof. (i) Applying φ to $[e_{ii}, e_{jk}] = 0$, we get

$$0 = \varphi(e_{ii})e_{jk} - e_{jk}\varphi(e_{ii}) + e_{ii}\varphi(e_{jk}) - \varphi(e_{jk})e_{ii}.$$

Then the desired equality is obtained by the left multiplication by e_{jj} and right multiplication by e_{kk} .

(ii) Since $i < j$, then $i \neq n$, so $i \notin \{j, n\}$ and $n \notin \{1, i\}$. We use item (i) and (12),

$$\begin{aligned} \varphi(e_{ij})(i, j) &= \varphi(e_{ii})(i, i) - \varphi(e_{ii})(j, j) = \varphi(e_{ii})(i, i) - \varphi(e_{ii})(n, n) \\ &= \varphi(e_{nn})(n, n) - \varphi(e_{nn})(i, i) = \varphi(e_{nn})(n, n) - \varphi(e_{nn})(1, 1) \\ &= \varphi(e_{1n})(1, n). \end{aligned}$$

(iii) We have already seen in (7) that $\varphi(e_{ii})(j, k) = 0$ for $j < k$ and $i \notin \{j, k\}$ (although it is required that $i < j$ in the statement of Lemma 6, the proof of (7) uses only $j < k$ and $i \notin \{j, k\}$). Moreover, $\varphi(e_{ii})(l, i) = 0$ for all $l < i$ by (11) under the assumption that there exists $j > i$ (i.e., $i < n$). Similarly, (5) and Lemma 7 imply that $\varphi(e_{jj})(j, k) = 2\varphi(e_{ij})(j, k) = 0$ for all $i < j < k$, so $\varphi(e_{ii})(i, k) = 0$ for all $1 < i < k$. It remains to prove that

$$\varphi(e_{11})(1, k) = \varphi(e_{nn})(j, n) = 0 \quad \text{for } 1 < j, k < n.$$

But Lemma 6 (iii) shows that $\varphi(e_{11})(1, k) = -\varphi(e_{kk})(1, k)$, which is proved to be 0 for $1 < k < n$. Similarly, $\varphi(e_{nn})(j, n) = -\varphi(e_{jj})(j, n) = 0$ for $1 < j < n$. ■

Lemma 9. *Let $n > 1$. Then the linear map $\alpha : T_n(F) \rightarrow T_n(F)$ given by*

$$\alpha(e_{ij}) = \begin{cases} e_{1n}, & (i, j) = (1, 1), \\ -e_{1n}, & (i, j) = (n, n), \\ 0, & (i, j) \notin \{(1, 1), (n, n)\}, \end{cases}$$

is a $\frac{1}{2}$ -derivation of $T_n(F)$.

Proof. We are going to prove that $\varphi = \alpha$ satisfies (2) for $x = e_{ij}$ and $y = e_{kl}$. Since α annihilates the commutators of $T_n(F)$, the left-hand side of (2) is always zero. In view of the anti-commutativity of $[\cdot, \cdot]$, we have to deal only with the following 2 cases.

Case 1. $(i, j) = (1, 1)$. Then

$$[\alpha(e_{ij}), e_{kl}] + [e_{ij}, \alpha(e_{kl})] = [e_{1n}, e_{kl}] + [e_{11}, \alpha(e_{kl})]. \quad (14)$$

Both summands on the right-hand side of (14) are zero, unless $(k, l) \in \{(1, 1), (n, n)\}$. If $(k, l) = (1, 1)$, then the right-hand side of (14) is zero due to the anti-commutativity of $[\cdot, \cdot]$. And if $(k, l) = (n, n)$, then $[e_{1n}, e_{kl}] + [e_{11}, \alpha(e_{kl})] = e_{1n} - e_{1n} = 0$.

Case 2. $(i, j) = (n, n)$. Then

$$[\alpha(e_{ij}), e_{kl}] + [e_{ij}, \alpha(e_{kl})] = -[e_{1n}, e_{kl}] + [e_{nn}, \alpha(e_{kl})].$$

We again have only 2 non-trivial subcases $(k, l) = (1, 1)$ and $(k, l) = (n, n)$ that are similar to the corresponding subcases of Case 1. ■

We also introduce the linear maps $\beta_i : T_n(F) \rightarrow T_n(F)$, $1 \leq i \leq n$, by

$$\beta_i(e_{jk}) = \begin{cases} \delta, & (j, k) = (i, i), \\ 0, & (j, k) \neq (i, i). \end{cases}$$

Obviously, β_i , $1 \leq i \leq n$, constitute a basis of the space of linear maps $T_n(F) \rightarrow Z(T_n(F))$ annihilating $[T_n(F), T_n(F)]$. In particular, $\beta_i \in \Delta(T_n(F))$ for all $1 \leq i \leq n$.

Proposition 10. *Let $n > 1$. Then $\Delta(T_n(F)) = \langle \text{id}, \alpha \rangle \oplus \langle \beta_i \mid 1 \leq i \leq n \rangle$.*

Proof. In view of Lemma 9 we only need to prove that each $\varphi \in \Delta(T_n(F))$ is a linear combination of id , α and β_i , $1 \leq i \leq n$. Setting $a = \varphi(e_{1n})(1, n)$, we have

$$\varphi(e_{ij}) = ae_{ij} \quad \text{for all } 1 \leq i < j \leq n \quad (15)$$

by Lemma 7 and Lemma 8 (ii). Then

$$\varphi(e_{ii})(i, i) - \varphi(e_{ii})(j, j) = a = \varphi(e_{ii})(i, i) - \varphi(e_{ii})(k, k)$$

for all $k < i < j$ by (12). Hence, defining $b_i = \varphi(e_{ii})(j, j)$ for some $j \neq i$, we have

$$\varphi(e_{ii})(j, j) = \begin{cases} a + b_i, & j = i, \\ b_i, & j \neq i. \end{cases} \quad (16)$$

Finally, let $c = \varphi(e_{11})(1, n)$. By Lemma 6 (iii) and Lemma 8 (iii) for all $1 \leq i \leq n$ and $1 \leq j < k \leq n$ we have

$$\varphi(e_{ii})(j, k) = \begin{cases} c, & (i, j, k) = (1, 1, n), \\ -c, & (i, j, k) = (n, 1, n), \\ 0, & (i, j, k) \notin \{(1, 1, n), (n, 1, n)\}. \end{cases} \quad (17)$$

It follows from (15)–(17) that $\varphi = a \cdot \text{id} + c\alpha + \sum_{i=1}^n b_i \beta_i$. ■

Theorem 11. *Let $\text{char}(F) = 0$ and $n > 2$. Then any transposed Poisson algebra structure on $T_n(F)$ is of one of the following two non-isomorphic forms:*

- (i) *transposed Poisson algebra structure of Poisson type;*
- (ii) *the orthogonal sum of the transposed Poisson algebra structure*

$$e_{11} \cdot e_{11} = -e_{11} \cdot e_{nn} = e_{nn} \cdot e_{nn} = e_{1n} \quad (18)$$

with a transposed Poisson algebra structure of Poisson type.

Proof. Let \cdot be a transposed Poisson algebra structure on $T_n(F)$. By Proposition 10 and Lemma 4 for all $1 \leq i \leq j \leq n$ there exist $x_{ij}, y_{ij} \in F$ and $\{z_{ij}^k\}_{k=1}^n \subseteq F$, such that

$$e_{ij} \cdot e_{kl} = x_{ij}e_{kl} + y_{ij}\alpha(e_{kl}) + \sum_{s=1}^n z_{ij}^s \beta_s(e_{kl})$$

$$= \begin{cases} x_{ij}e_{kl}, & k < l, \\ x_{ij}e_{kk} + z_{ij}^k \delta, & k = l \notin \{1, n\}, \\ x_{ij}e_{11} + y_{ij}e_{1n} + z_{ij}^1 \delta, & k = l = 1, \\ x_{ij}e_{nn} - y_{ij}e_{1n} + z_{ij}^n \delta, & k = l = n. \end{cases}$$

Let $1 \leq i < j \leq n$ and $1 < k < n$. Then $e_{kk} \cdot e_{ij} = x_{kk}e_{ij}$, while $e_{ij} \cdot e_{kk} = x_{ij}e_{kk} + z_{ij}^k \delta$, whence

$$x_{kk} = 0 \quad \text{and} \quad x_{ij} = z_{ij}^k = 0 \quad \text{for } 1 \leq i < j \leq n \text{ and } 1 < k < n. \quad (19)$$

Now take $1 \leq i < j \leq n$ with $(i, j) \neq (1, n)$. Then $e_{11} \cdot e_{ij} = x_{11}e_{ij}$, while $e_{ij} \cdot e_{11} = y_{ij}e_{1n} + z_{ij}^1 \delta$. Similarly, $e_{nn} \cdot e_{ij} = x_{nn}e_{ij}$, while $e_{ij} \cdot e_{nn} = -y_{ij}e_{1n} + z_{ij}^n \delta$. Hence,

$$x_{11} = x_{nn} = 0 \quad \text{and} \quad y_{ij} = z_{ij}^1 = z_{ij}^n = 0 \quad (20)$$

for $1 \leq i < j \leq n$ with $(i, j) \neq (1, n)$.

Considering the products $e_{11} \cdot e_{1n} = 0$, $e_{1n} \cdot e_{11} = y_{1n}e_{1n} + z_{1n}^1 \delta$, $e_{nn} \cdot e_{1n} = 0$ and $e_{1n} \cdot e_{nn} = -y_{1n}e_{1n} + z_{1n}^n \delta$, we have

$$y_{1n} = z_{1n}^1 = z_{1n}^n = 0. \quad (21)$$

Now, for any $1 < i < n$ it follows from $e_{11} \cdot e_{ii} = z_{11}^i \delta$, $e_{ii} \cdot e_{11} = y_{ii}e_{1n} + z_{ii}^1 \delta$, $e_{nn} \cdot e_{ii} = z_{nn}^i \delta$ and $e_{ii} \cdot e_{nn} = -y_{ii}e_{1n} + z_{ii}^n \delta$ that

$$y_{ii} = 0, \quad z_{ii}^1 = z_{11}^i \quad \text{and} \quad z_{ii}^n = z_{nn}^i \quad \text{for } 1 < i < n. \quad (22)$$

Similarly, for $1 < i, j < n$, it follows from $e_{ii} \cdot e_{jj} = z_{ii}^j \delta$ and $e_{jj} \cdot e_{ii} = z_{jj}^i \delta$ that

$$z_{ii}^j = z_{jj}^i \quad \text{for } 1 < i, j < n. \quad (23)$$

Finally, $e_{11} \cdot e_{nn} = -y_{11}e_{1n} + z_{11}^n \delta$ and $e_{nn} \cdot e_{11} = y_{nn}e_{1n} + z_{nn}^1 \delta$ yield

$$y_{nn} = -y_{11} \quad \text{and} \quad z_{11}^n = z_{nn}^1. \quad (24)$$

Combining (19)–(24) and denoting $a_{ij} := z_{ii}^j$, $b = y_{11}$, we see that the only (possibly) non-zero products $e_{ij} \cdot e_{kl}$ are

$$e_{ii} \cdot e_{jj} = a_{ij} \delta, \quad 1 < i, j < n, \quad (25)$$

$$e_{11} \cdot e_{11} = be_{1n} + a_{11} \delta, \quad e_{nn} \cdot e_{nn} = be_{1n} + a_{nn} \delta, \quad (26)$$

$$e_{11} \cdot e_{nn} = e_{nn} \cdot e_{11} = -be_{1n} + a_{1n} \delta,$$

where $a_{ij} = a_{ji}$ for all $1 \leq i, j \leq n$.

If $b = 0$, then \cdot is of Poisson type, so we are in the case (i). Otherwise, choosing ϕ to be the conjugation by $(b^{-1} - 1)e_{11} + \delta$, we have

$$\phi(e_{ij}) = \begin{cases} e_{ij}, & 1 < i \leq j \leq n \text{ or } i = j = 1, \\ b^{-1}e_{ij}, & 1 = i < j \leq n. \end{cases}$$

It follows that, applying ϕ , we can replace $b \neq 0$ by $b = 1$ in (26), and we come to the form (ii).

To prove that the structures (i) and (ii) are not isomorphic, observe from [16] that for any automorphism ϕ of $(T_n(F), [\cdot, \cdot])$

$$\text{either } \phi(e_{ii}) \in e_{ii} + (\langle \delta \rangle \oplus \langle e_{ij} \mid i < j \rangle) \quad \text{for all } 1 \leq i \leq n, \quad (27)$$

$$\text{or } \phi(e_{ii}) \in -e_{n-i+1, n-i+1} + (\langle \delta \rangle \oplus \langle e_{ij} \mid i < j \rangle) \quad \text{for all } 1 \leq i \leq n. \quad (28)$$

Consider the case (27). Since $e_{ii} \cdot \delta \in \langle \delta \rangle = Z(T_n(F), [\cdot, \cdot])$ and $e_{ii} \cdot e_{jk} = 0$ for all $1 \leq i \leq n$ and $1 \leq j < k \leq n$ by (25) and (26), any such ϕ leads to the product

$$e_{ii} * e_{jj} = \phi(\phi^{-1}(e_{ii}) \cdot \phi^{-1}(e_{jj})) \in \phi(e_{ii} \cdot e_{jj} + \langle \delta \rangle) = \phi(e_{ii} \cdot e_{jj}) + \langle \delta \rangle,$$

where $\phi(e_{ii} \cdot e_{jj})$ belongs either to $\langle \delta \rangle$ (whenever $e_{ii} \cdot e_{jj} \in \langle \delta \rangle$) or to $\pm b\phi(e_{1n}) + \langle \delta \rangle$ (whenever $e_{ii} \cdot e_{jj} \in \pm be_{1n} + \langle \delta \rangle$). But $\phi(e_{1n})$ is a non-zero multiple of e_{1n} by (4), so

$$e_{ii} * e_{jj} \in \langle \delta \rangle \Leftrightarrow e_{ii} \cdot e_{jj} \in \langle \delta \rangle \quad (29)$$

for all $1 \leq i, j \leq n$. This shows that any structure of type (i) can be isomorphic only to a structure of type (i). As to the case (28), we similarly have

$$e_{ii} * e_{jj} = \phi(\phi^{-1}(e_{ii}) \cdot \phi^{-1}(e_{jj})) \in \phi(e_{n-i+1, n-i+1} \cdot e_{n-j+1, n-j+1} + \langle \delta \rangle).$$

However,

$$\begin{aligned} e_{n-i+1, n-i+1} \cdot e_{n-j+1, n-j+1} \in \langle \delta \rangle &\Leftrightarrow (i, j) \notin \{(1, 1), (1, n), (n, 1), (n, n)\} \\ &\Leftrightarrow e_{ii} \cdot e_{jj} \in \langle \delta \rangle. \end{aligned}$$

Thus, we again conclude that (29) holds for all $1 \leq i, j \leq n$ showing that the structures (i) and (ii) cannot be isomorphic.

Conversely, it is directly verified that (18) is a transposed Poisson algebra structure on $T_n(F)$, and it is orthogonal to any transposed Poisson algebra structure of Poisson type on $T_n(F)$. ■

The following Theorem 12 describes transposed Poisson structures on $T_2(F)$, and its proof is similar to that of Theorem 11 with the only difference that as a result we will have 3 types of the structures, and inside each of the 3 types the structures can be fully classified up to isomorphism. On the other hand, $T_2(F)$ is isomorphic to the 3-dimensional Lie algebra \mathfrak{g}_2^0 from [2] whose transposed Poisson structures were fully described in [2, Proposition 14]. By this reason, we have chosen to omit the proof of Theorem 12 and indicate on the left of each of the found structures its isomorphic version from [2].

Theorem 12. *Let $\text{char}(F) = 0$. Then any transposed Poisson algebra structure on $T_2(F)$ is isomorphic to exactly one of the following structures:*

- (i) *transposed Poisson algebra structure of Poisson type:*
 - (a) $T_{09}^{0,0}$: *the trivial one;*
 - (b) T_{17}^0 : $e_{11} \cdot e_{11} = \delta$;
 - (c) T_{10}^0 : $e_{11} \cdot e_{11} = -e_{11} \cdot e_{22} = e_{22} \cdot e_{22} = \delta$;
- (ii) *the orthogonal sum of the transposed Poisson algebra structure*

$$e_{11} \cdot e_{11} = -e_{11} \cdot e_{22} = e_{22} \cdot e_{22} = e_{12}$$

with a transposed Poisson algebra structure of Poisson type, which results in the structures:

- (a) T_{16} : $e_{11} \cdot e_{11} = -e_{11} \cdot e_{22} = e_{22} \cdot e_{22} = e_{12}$;
- (b) T_{18} : $e_{11} \cdot e_{11} = e_{12} + \delta$, $-e_{11} \cdot e_{22} = e_{22} \cdot e_{22} = e_{12}$;
- (c) T_{11}^0 : $e_{11} \cdot e_{11} = -e_{11} \cdot e_{22} = e_{22} \cdot e_{22} = e_{12} + \delta$;
- (iii) *the (non-orthogonal) sum of a transposed Poisson algebra structure of the family*

$$\begin{aligned} e_{11} \cdot e_{11} &= ce_{11}, & e_{11} \cdot e_{12} &= -e_{12} \cdot e_{22} = ce_{12}, \\ e_{11} \cdot e_{22} &= -e_{22} \cdot e_{22} = ce_{22}, & c &\neq 0, \end{aligned}$$

with a transposed Poisson algebra structure of Poisson type, which results in the structures

- (a) $T_{17}^{-c}: e_{11} \cdot e_{11} = ce_{11}, e_{11} \cdot e_{12} = -e_{12} \cdot e_{22} = ce_{12}, e_{11} \cdot e_{22} = -e_{22} \cdot e_{22} = ce_{22};$
- (b) $T_{09}^{0,-c}: e_{11} \cdot e_{11} = ce_{11}, e_{11} \cdot e_{12} = -e_{12} \cdot e_{22} = ce_{12}, e_{11} \cdot e_{22} = ce_{22}, e_{22} \cdot e_{22} = -c(e_{22} + \delta);$
- (c) $T_{19}^{-c}: e_{11} \cdot e_{11} = ce_{11}, e_{11} \cdot e_{12} = -e_{12} \cdot e_{22} = ce_{12}, -e_{11} \cdot e_{22} = e_{22} \cdot e_{22} = ce_{11}.$

3. Transposed Poisson structures on the full matrix Lie algebra

In this short section we describe transposed Poisson structures on the full matrix algebra $M_n(F)$. As above, we denote by e_{ij} , $1 \leq i, j \leq n$, the matrix units and by δ the identity matrix. Recall that

$$\begin{aligned} Z(M_n(F)) &= \langle \delta \rangle, \\ [M_n(F), M_n(F)] &= \mathfrak{sl}_n(F) = \{a \in M_n(F) \mid \text{tr}(a) = 0\} \\ &= \langle e_{ij} \mid 1 \leq i \neq j \leq n \rangle \oplus \langle e_{11} - e_{ii} \mid 1 < i \leq n \rangle. \end{aligned}$$

Denote by $\Delta(M_n(F))$ the space of $\frac{1}{2}$ -derivations of the Lie algebra $(M_n(F), [\cdot, \cdot])$. The linear map $\gamma: M_n(F) \rightarrow M_n(F)$ given by

$$\gamma(e_{ij}) = \begin{cases} \delta, & i = j, \\ 0, & i \neq j, \end{cases}$$

belongs to $\Delta(M_n(F))$ as a linear map $M_n(F) \rightarrow Z(M_n(F))$ annihilating $[M_n(F), M_n(F)]$.

Proposition 13. *Let $n > 1$. Then $\Delta(M_n(F)) = \langle \text{id}, \gamma \rangle$.*

Proof. It is easy to see that, as a Lie algebra, $M_n(F)$ is the direct sum

$$[M_n(F), M_n(F)] \oplus Z(M_n(F)).$$

Namely, every $a \in M_n(F)$ decomposes uniquely as

$$a = \left(a - \frac{\text{tr}(a)}{n} \delta \right) + \frac{\text{tr}(a)}{n} \delta.$$

Since $\frac{1}{2}$ -derivations of a Lie algebra \mathfrak{L} leave $[\mathfrak{L}, \mathfrak{L}]$ and $Z(\mathfrak{L})$ invariant, we have

$$\Delta(M_n(F)) = \Delta([M_n(F), M_n(F)]) \oplus \Delta(Z(M_n(F))).$$

By [6, Corollary 3] any $\varphi \in \Delta([M_n(F), M_n(F)])$ is trivial. As $\dim(Z(M_n(F))) = 1$, any $\varphi \in \Delta(Z(M_n(F)))$ is trivial as well. Hence, given $\varphi \in \Delta(M_n(F))$, there exist $k_1, k_2 \in F$ such that $\varphi = k_1 \cdot \text{id}_{[M_n(F), M_n(F)]} + k_2 \cdot \text{id}_{Z(M_n(F))}$. Clearly,

$$\begin{aligned}\varphi(e_{ij}) &= k_1 e_{ij}, \quad \text{if } i \neq j, \\ \varphi(e_{ii}) &= k_1 \left(e_{ii} - \frac{1}{n} \delta \right) + \frac{k_2}{n} \delta = k_1 e_{ii} + \frac{k_2 - k_1}{n} \delta.\end{aligned}$$

It follows that $\varphi = k_1 \cdot \text{id}_{M_n(F)} + \frac{k_2 - k_1}{n} \cdot \gamma$. Thus, $\Delta(M_n(F)) \subseteq \langle \text{id}, \gamma \rangle$. The converse inclusion is trivial. ■

Theorem 14. *Let $\text{char}(F) = 0$ and $n > 1$. Then, up to isomorphism, there is only one non-trivial transposed Poisson algebra structure on $M_n(F)$. It is given by*

$$e_{ii} \cdot e_{jj} = \delta, \quad 1 \leq i, j \leq n, \quad (30)$$

and it is of Poisson type.

Proof. Let \cdot be a transposed Poisson algebra structure on $M_n(F)$. By Proposition 13 and Lemma 4 for all $1 \leq i, j \leq n$ there exist x_{ij} and $y_{ij} \in F$, such that

$$e_{ij} \cdot e_{kl} = x_{ij} e_{kl} + y_{ij} \gamma(e_{kl}) = \begin{cases} x_{ij} e_{kl}, & k \neq l, \\ x_{ij} e_{kk} + y_{ij} \delta, & k = l. \end{cases}$$

Let $i \neq j$ and $1 \leq k \leq n$. Then $e_{kk} \cdot e_{ij} = x_{kk} e_{ij}$, while $e_{ij} \cdot e_{kk} = x_{ij} e_{kk} + y_{ij} \delta$, whence

$$x_{kk} = 0 \quad \text{and} \quad x_{ij} = y_{ij} = 0 \quad \text{for } i \neq j \text{ and } 1 \leq k \leq n.$$

It follows from $e_{ii} \cdot e_{jj} = y_{ii} \delta$ and $e_{jj} \cdot e_{ii} = y_{jj} \delta$ that

$$y_{ii} = y_{jj} \quad \text{for } 1 \leq i, j \leq n.$$

Thus, denoting y_{ii} by c , we have the only (possibly) non-zero products

$$e_{ii} \cdot e_{jj} = c \delta, \quad 1 \leq i, j \leq n.$$

If $c = 0$, then \cdot is trivial. Otherwise, choosing the automorphism ϕ of $(M_n(F), [\cdot, \cdot])$ given by

$$\begin{aligned}\phi(e_{ij}) &= e_{ij}, \quad \text{if } i \neq j, \\ \phi(e_{ii}) &= e_{ii} - \frac{1}{n} \delta + \frac{c}{n} \delta = e_{ii} + \frac{c-1}{n} \delta,\end{aligned}$$

we obtain the isomorphic structure (30).

It remains to see that (30) is of Poisson type. For all $a, b \in M_n(F)$ we have

$$\begin{aligned} a \cdot b &= \sum_{i,j=1}^n a(i, i)a(j, j)e_{ii} \cdot e_{jj} = \sum_{i,j=1}^n a(i, i)a(j, j)\delta \\ &= \operatorname{tr}(a)\operatorname{tr}(b)\delta = \frac{\operatorname{tr}(a)}{n}\delta \cdot \frac{\operatorname{tr}(b)}{n}\delta. \end{aligned}$$

Hence, \cdot is the extension by zero of a product on the complement $Z(M_n(F))$ of $[M_n(F), M_n(F)]$ in $M_n(F)$ with values in $Z(M_n(F))$. ■

Remark 15. The structure (30) is isomorphic to the extension by zero of the product $\delta \cdot \delta = \delta$.

Funding. The work was supported by FCT UIDB/MAT/00212/2020, UIDP/MAT/00212/2020, 2022.02474.PTDC and by CMUP, member of LASI, which is financed by national funds through FCT – Fundação para a Ciência e a Tecnologia, I.P., under the project with reference UIDB/00144/2020.

References

- [1] C. Bai, R. Bai, L. Guo, and Y. Wu, [Transposed Poisson algebras, Novikov–Poisson algebras and 3-Lie algebras](#). *J. Algebra* **632** (2023), 535–566 Zbl 07710371 MR 4607568
- [2] P. D. Beites, A. Fernández Ouaridi, and I. Kaygorodov, [The algebraic and geometric classification of transposed Poisson algebras](#). *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM* **117** (2023), no. 2, article no. 55 Zbl 1507.17003 MR 4534525
- [3] P. D. Beites, B. L. M. Ferreira, and I. Kaygorodov, [Transposed Poisson structures](#). *Results Math.*, **79** (2024), article no. 93
- [4] A. Fernández Ouaridi, [On the simple transposed Poisson algebras and Jordan superalgebras](#). *J. Algebra* **641** (2024), 173–198 Zbl 07787769 MR 4673293
- [5] B. L. M. Ferreira, I. Kaygorodov, and V. Lopatkin, [\$\frac{1}{2}\$ -derivations of Lie algebras and transposed Poisson algebras](#). *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM* **115** (2021), no. 3, article no. 142 Zbl 1507.17037 MR 4272639
- [6] V. T. Filippov, [On \$\delta\$ -derivations of Lie algebras](#). *Sib. Math. J.* **39** (1998), no. 6, 1218–1230 Zbl 0936.17020 MR 1672673
- [7] A. Jaworska-Pastuszek and Z. Pogorzały, [Poisson structures for canonical algebras](#). *J. Geom. Phys.* **148** (2020), article no. 103564 Zbl 1487.17045 MR 4039470
- [8] I. Kaygorodov, [Non-associative algebraic structures: classification and structure](#). *Commun. Math.* **32** (2024), no. 3, 1–62 MR 4683663
- [9] I. Kaygorodov and M. Khrypchenko, [Poisson structures on finitary incidence algebras](#). *J. Algebra* **578** (2021), 402–420 Zbl 1481.17031 MR 4237454

- [10] I. Kaygorodov and M. Khrypchenko, [Transposed Poisson structures on Block Lie algebras and superalgebras](#). *Linear Algebra Appl.* **656** (2023), 167–197 Zbl 1516.17002 MR 4492612
- [11] I. Kaygorodov and M. Khrypchenko, [Transposed Poisson structures on generalized Witt algebras and Block Lie algebras](#). *Results Math.* **78** (2023), no. 5, article no. 186 Zbl 07720518 MR 4617175
- [12] I. Kaygorodov and M. Khrypchenko, [Transposed Poisson structures on Witt type algebras](#). *Linear Algebra Appl.* **665** (2023), 196–210 Zbl 07667597 MR 4550293
- [13] I. Kaygorodov, V. Lopatkin, and Z. Zhang, [Transposed Poisson structures on Galilean and solvable Lie algebras](#). *J. Geom. Phys.* **187** (2023), article no. 104781 Zbl 07673053 MR 4552380
- [14] G. Liu and C. Bai, [A bialgebra theory for transposed Poisson algebras via anti-pre-Lie bialgebras and anti-pre-Lie-Poisson bialgebras](#). *Commun. Contemp. Math.*, 2023, to appear
- [15] G. Liu and C. Bai, [New Splittings of Operations of Poisson Algebras and Transposed Poisson Algebras and Related Algebraic Structures](#). In *Algebra without Borders – Classical and Constructive Nonassociative Algebraic Structures*, pp. 49–96, STEAM-H: Sci. Technol. Eng. Agric. Math. Health, Springer, Cham, 2023 MR 4696690
- [16] L. W. Marcoux and A. R. Sourour, [Commutativity preserving linear maps and Lie automorphisms of triangular matrix algebras](#). *Linear Algebra Appl.* **288** (1999), no. 1-3, 89–104 Zbl 0933.15029 MR 1670535
- [17] B. Sartayev, [Some generalizations of the variety of transposed Poisson algebras](#). *Commun. Math.* **32** (2024), no. 2, 55–62 MR 4683661
- [18] Y. Yang, X. Tang, and A. Khudoyberdiyev, [Transposed Poisson structures on Schrodinger algebra in \$\(n + 1\)\$ -dimensional space-time](#). 2023, arXiv:2303.08180v1
- [19] Y. Yao, Y. Ye, and P. Zhang, [Quiver Poisson algebras](#). *J. Algebra* **312** (2007), no. 2, 570–589 Zbl 1180.17007 MR 2333173
- [20] L. Yuan and Q. Hua, [\$\frac{1}{2}\$ -\(bi\)derivations and transposed Poisson algebra structures on Lie algebras](#). *Linear Multilinear Algebra* **70** (2022), no. 22, 7672–7701 Zbl 07681820 MR 4571051

Received 10 May 2023.

Ivan Kaygorodov

CMA-UBI, Universidade da Beira Interior, Covilhã, Portugal; kaygorodov.ivan@gmail.com

Mykola Khrypchenko

Departamento de Matemática, Universidade Federal de Santa Catarina, Florianópolis, Brazil; and CMUP, Departamento de Matemática, Faculdade de Ciências, Universidade do Porto, Porto, Portugal; nskhrpchenko@gmail.com