

## Existence, uniqueness and decay rates for evolution equations on trees

Leandro M. Del Pezzo<sup>1</sup>, Carolina A. Mosquera<sup>2</sup> and Julio D. Rossi<sup>3</sup>

**Abstract.** We study evolution equations governed by an averaging operator on a directed tree, showing existence and uniqueness of solutions. In addition we find conditions of the initial condition that allows us to find the asymptotic decay rate of the solutions as  $t \rightarrow \infty$ . It turns out that this decay rate is not uniform, it strongly depends on how the initial condition goes to zero as one goes down in the tree.

**Mathematics Subject Classification (2010).** 35B40, 35K55, 91A22.

**Keywords.** Evolution equations, averaging operators, decay estimates.

### 1. Introduction

Let  $\mathbb{T}_m$  be a directed tree with  $m$ -branching, we denote by  $x$  the vertices of the tree. Given a function  $f : \mathbb{T}_m \rightarrow \mathbb{R}$ , in this work we study the following Cauchy problem

$$\begin{cases} u_t(x, t) - \Delta_F u(x, t) = 0 & \text{in } \mathbb{T}_m \times (0, +\infty), \\ u(x, 0) = f(x) & \text{in } \mathbb{T}_m, \end{cases} \quad (1.1)$$

where

$$\Delta_F u(x, t) = F(u((x, 0), t), \dots, u((x, m-1), t)) - u(x, t),$$

being  $F$  an averaging operator, see the precise definition in Section 2. The simplest linear example of an averaging operator is the usual average

$$F(x_1, \dots, x_m) = \frac{1}{m} \sum_{j=1}^m x_j$$

---

<sup>1</sup>Leandro M. Del Pezzo was partially supported by UBACyT 20020110300067 and CONICET PIP 5478/1438 (Argentina).

<sup>2</sup>Carolina A. Mosquera was partially supported by UBACyT 20020100100638, PICT 0436 and CONICET PIP 112 200201 00398 (Argentina).

<sup>3</sup>Julio D. Rossi was partially supported by MTM2011-27998 (Spain).

but we can include nonlinear functions as

$$F(x_1, \dots, x_m) = \frac{\alpha}{2} \left( \max_{1 \leq j \leq m} \{x_j\} + \min_{1 \leq j \leq m} \{x_j\} \right) + \frac{1-\alpha}{m} \sum_{j=1}^m x_j,$$

with  $0 < \alpha < 1$ .

We can see that  $u$  is a solution of (1.1) if and only if it is a solution of the integral equation

$$u(x, t) = K_f u(x, t), \tag{1.2}$$

where

$$K_f u(x, t) := \int_0^t e^{z-t} F(u((x, 0), z), \dots, u((x, m-1), z)) dz + e^{-t} f(x).$$

We first prove existence and uniqueness of a locally bounded global in time solution using a fixed point argument for  $K_f$ .

**Theorem 1.1.** *Let  $f \in L^\infty(\mathbb{T}_m, \mathbb{R})$ . Then there exists a unique solution  $u$  in*

$$L_{\text{loc}}^\infty([0, +\infty); L^\infty(\mathbb{T}_m, \mathbb{R})) := \{v \in L^\infty(\mathbb{T}_m \times [0, T], \mathbb{R}) \ \forall T > 0\}$$

of (1.1).

In addition, a comparison principle holds.

**Theorem 1.2.** *Let  $F$  be an averaging operator,  $f, g \in L^\infty(\mathbb{T}_m, \mathbb{R})$  such that  $f \leq g$  in  $\mathbb{T}_m$ , and  $u, v \in L_{\text{loc}}^\infty([0, +\infty); L^\infty(\mathbb{T}_m, \mathbb{R}))$  such that*

$$u(x, t) \leq K_f u(x, t) \quad \text{and} \quad v(x, t) \geq K_g v(x, t) \tag{1.3}$$

for all  $(x, t) \in \mathbb{T}_m \times [0, +\infty)$ . Then  $u \leq v$  in  $\mathbb{T}_m \times [0, +\infty)$ .

Once we have established existence and uniqueness of global in time solutions a natural question is to look for its asymptotic behaviour as  $t \rightarrow \infty$ . We find conditions on the initial condition  $f$  (that involve the speed at which they go to zero as one goes down in the tree) that guarantee that solutions go to zero as  $t \rightarrow \infty$ . Under these conditions we can find bounds for the decay rate. Surprisingly the decay rate for solutions to (1.1) is not uniform. It strongly depends on the decay of the initial condition  $f$ . For example, for initial conditions with finite support (only a finite number of vertices have non-zero values) we find a decay of the form  $t^\mu e^{-t}$  (here  $\mu$  depends on the size of the support of  $f$ ), while for data without finite

support we find a decay of the form  $e^{-\lambda t}$  (with  $0 < \lambda < 1$  depending on the decay of  $f$ ). This is the content of our next results whose proof rely mostly on comparison arguments. For the statements we need to introduce the following notations.

Let  $f \in L^\infty(\mathbb{T}_m, \mathbb{R})$ . We will say that  $f$  has finite support if there exists  $n \in \mathbb{N}_0$  such that  $f(x) = 0$  for all  $x \in \mathbb{T}_m$  with  $l(x) \geq n$ , where  $l(x)$  denotes the level of  $x$ . We also define

$$a(f) := \min\{j \in \mathbb{N}_0 : f(x) = 0, \forall x \in \mathbb{T}_m \text{ with } l(x) \geq j\},$$

and

$$\mu(f) := \begin{cases} a(f) - 1 & \text{if } a(f) > 0, \\ 0 & \text{if } a(f) = 0. \end{cases}$$

**Theorem 1.3.** *Let  $F$  be an averaging operator and  $f \in L^\infty(\mathbb{T}_m, \mathbb{R})$  with finite support. If  $u \in L^\infty_{\text{loc}}([0, +\infty); L^\infty(\mathbb{T}_m, \mathbb{R}))$  is the solution of (1.1) with initial condition  $f$ , then*

$$\max_{x \in \mathbb{T}_m} |u(x, t)| \leq \frac{t^{\mu(f)} e^{-t}}{\mu(f)!} \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})}, \quad (1.4)$$

for  $t$  large enough.

The above bound is optimal, see Remark 4.1.

For  $f$  that are not finitely supported we have the following result.

**Theorem 1.4.** *Let  $F$  be an averaging operator and  $f \in L^\infty(\mathbb{T}_m, \mathbb{R})$  such that there exist  $\lambda \in (0, 1)$  and  $k \in \mathbb{R}_{>0}$  such that*

$$|f(x)| \leq k(1 - \lambda)^{l(x)} \quad \forall x \in \mathbb{T}_m.$$

If  $u \in L^\infty_{\text{loc}}([0, +\infty); L^\infty(\mathbb{T}_m, \mathbb{R}))$  is the solution of (1.1) with initial condition  $f$ , then

$$\max_{x \in \mathbb{T}_m} |u(x, t)| \leq k e^{-\lambda t} \quad \forall t \in \mathbb{R}.$$

Again this bound is optimal, see Proposition 4.2.

In the next result we show that we can construct a solution with quite different behaviours at  $\emptyset$ , the first node of our tree.

**Theorem 1.5.** *Let  $F$  be an averaging operator and  $a_0(t) \in C^\infty([0, \infty), \mathbb{R})$ , then there is a solution of  $u_t(x, t) - \Delta_F u(x, t) = 0$  in  $\mathbb{T}_m \times (0, +\infty)$ , such that*

$$u(\emptyset, t) = a_0(t) \quad \forall t \in \mathbb{R}.$$

Let us end the introduction with a brief comment on previous bibliography that concerns mostly the stationary problem. For nonlinear mean values on a finite graph we refer to [8] and references therein. For equations on trees like the ones considered here, see [1], [6], [7] and [9], [10], where for the stationary problem it is proved the existence and uniqueness of a solution using game theory. See also [2], [5] where the authors study the unique continuation and find some estimates for the harmonic measure on trees. Here we use ideas from these references.

The time dependent diffusion equations on simple, connected, undirected graphs, have been used to model diffusion processes, such as, modeling energy flows through a network or vibration of molecules, [3], [4].

In the case when  $F$  is the usual average, it is possible to construct a fundamental solution for (1.1) on infinite, locally finite, connected graphs. See [11], [12] and the references therein.

This paper is a natural extension of the previously mentioned references since here we deal with the evolution problem associated to an averaging operator on a tree that is a directed graph.

This paper is organized as follows: in Section 2 we collect some preliminaries; in Section 3 we deal with existence and uniqueness of solutions and prove Theorem 1.1 and Theorem 1.2; in Section 4 we prove our results concerning the decay of solutions as  $t \rightarrow \infty$  proving Theorem 1.3, Theorem 1.4 and Theorem 1.5.

## 2. Preliminaries

We begin with a review of the basic results that will be needed in subsequent sections. The known results are generally stated without proofs, but we provide references where the proofs can be found. Also, we introduce some of our notational conventions.

**2.1. Directed tree.** Let  $m \in \mathbb{N}_{>2}$ . In this work we consider a directed tree  $\mathbb{T}_m$  with regular  $m$ -branching, that is,  $\mathbb{T}_m$  consists of the empty set  $\emptyset$  and all finite sequences  $(a_1, a_2, \dots, a_k)$  with  $k \in \mathbb{N}$ , whose coordinates  $a_i$  are chosen from  $\{0, 1, \dots, m-1\}$ . The elements in  $\mathbb{T}_m$  are called vertices. Each vertex  $x$  has  $m$  successors, obtained by adding another coordinate. We will denote by  $\mathcal{S}(x)$  the set of successors of the vertex  $x$ . A vertex  $x \in \mathbb{T}_m$  is called an  $n$ -level vertex ( $n \in \mathbb{N}$ ) if  $x = (a_1, a_2, \dots, a_n)$ , and we will denote by  $l(x)$  the level of vertex  $x$ . The set of all  $n$ -level vertices is denoted by  $\mathbb{T}_m^n$ .

A branch of  $\mathbb{T}_m$  is an infinite sequence of vertices, each followed by its immediate successor. The collection of all branches forms the boundary of  $\mathbb{T}_m$ , denoted by  $\partial\mathbb{T}_m$ .

We now define a metric on  $\mathbb{T}_m \cup \partial\mathbb{T}_m$ . The distance between two sequences (finite or infinite)  $\pi = (a_1, \dots, a_k, \dots)$  and  $\pi' = (a'_1, \dots, a'_k, \dots)$  is  $m^{-K+1}$  when  $K$  is the first index  $k$  such that  $a_k \neq a'_k$ ; but when  $\pi = (a_1, \dots, a_K)$  and  $\pi' = (a_1, \dots, a_K, a'_{K+1}, \dots)$ , the distance is  $m^{-K}$ . Hausdorff measure and Hausdorff dimension are defined using this metric. We have that  $\mathbb{T}_m$  and  $\partial\mathbb{T}_m$  have diameter one and  $\partial\mathbb{T}_m$  has Hausdorff dimension one. Now, we observe that the mapping  $\psi : \partial\mathbb{T}_m \rightarrow [0, 1]$  defined as

$$\psi(\pi) := \sum_{k=1}^{+\infty} \frac{a_k}{m^k}$$

is surjective, where  $\pi = (a_1, \dots, a_k, \dots) \in \partial\mathbb{T}_m$  and  $a_k \in \{0, 1, \dots, m-1\}$  for all  $k \in \mathbb{N}$ . Whenever  $x = (a_1, \dots, a_k)$  is a vertex, we set

$$\psi(x) := \psi(a_1, \dots, a_k, 0, \dots, 0, \dots).$$

We can also associate to a vertex  $x$  an interval  $I_x$  of length  $\frac{1}{m^k}$  as follows

$$I_x := \left[ \psi(x), \psi(x) + \frac{1}{m^k} \right].$$

Observe that for all  $x \in \mathbb{T}_m$ ,  $I_x \cap \partial\mathbb{T}_m$  is the subset of  $\partial\mathbb{T}_m$  consisting of all branches that start at  $x$ . With an abuse of notation, we will write  $\pi = (x_1, \dots, x_k, \dots)$  instead of  $\pi = (a_1, \dots, a_k, \dots)$  where  $x_1 = a_1$  and  $x_k = (a_1, \dots, a_k) \in \mathcal{S}(x_{k-1})$  for all  $k \in \mathbb{N}_{\geq 2}$ .

Finally we will denote by  $\mathbb{T}_m^x$  the set of the vertices  $y \in \mathbb{T}_m$  such that  $I_y \subset I_x$ .

**Example 2.1.** Then Let  $\kappa \in \mathbb{N}$  be at least 3. A  $1/\kappa$ -Cantor set, which we denote by  $C_{1/\kappa}$ , is the set of all  $x \in [0, 1]$  that have a base  $\kappa$  expansion without the digit 1, that is,  $x = \sum a_j \kappa^{-j}$  with  $a_j \in \{0, 1, \dots, \kappa-1\}$  with  $a_j \neq 1$ . Thus  $C_{1/\kappa}$  is obtained from  $[0, 1]$  by removing the second  $\kappa$ -th part of the line segment  $[0, 1]$ , and then removing the second interval of length  $1/\kappa$  from the remaining intervals, and so on. This set can be thought of as a directed tree with regular  $m$ -branching with  $m = \kappa - 1$ .

For example, if  $\kappa = 3$ , we identify  $[0, 1]$  with  $\emptyset$ , the sequence  $(\emptyset, 0)$  with the first interval right  $[0, 1/3]$ , the sequence  $(\emptyset, 1)$  with the central interval  $[1/3, 2/3]$  (that is removed), the sequence  $(\emptyset, 2)$  with the left interval  $[2/3, 1]$ , the sequence  $(\emptyset, 0, 0)$  with the interval  $[0, 1/9]$  and so on.

**2.2. Averaging operator.** The following definition is taken from [1]. Let

$$F : \mathbb{R}^m \rightarrow \mathbb{R}$$

be a continuous function. We call  $F$  an averaging operator if it satisfies the following set of conditions:

- (i)  $F(0, \dots, 0) = 0$  and  $F(1, \dots, 1) = 1$ ;
- (ii)  $F(tx_1, \dots, tx_m) = tF(x_1, \dots, x_m)$  for all  $t \in \mathbb{R}$ ;
- (iii)  $F(t + x_1, \dots, t + x_m) = t + F(x_1, \dots, x_m)$  for all  $t \in \mathbb{R}$ ;
- (iv)  $F(x_1, \dots, x_m) < \max\{x_1, \dots, x_m\}$  if not all  $x_j$ 's are equal;
- (v)  $F$  is nondecreasing with respect to each variable.

**Remark 2.2.** It holds that, if  $(x_1, \dots, x_m), (y_1, \dots, y_m) \in \mathbb{R}^m$ , then

$$x_j \leq y_j + \max_{1 \leq j \leq m} \{x_j - y_j\}$$

for all  $j \in \{1, \dots, m\}$ . Let  $F$  be an averaging operator. Then, by (iii) and (v),

$$F(x_1, \dots, x_m) \leq F(y_1, \dots, y_m) + \max_{1 \leq j \leq m} \{x_j - y_j\}.$$

Therefore

$$F(x_1, \dots, x_m) - F(y_1, \dots, y_m) \leq \max_{1 \leq j \leq m} \{x_j - y_j\},$$

and moreover

$$|F(x_1, \dots, x_m) - F(y_1, \dots, y_m)| \leq \max_{1 \leq j \leq m} \{|x_j - y_j|\}.$$

Now we give some examples.

**Example 2.3.** This example is taken from [6]. For  $1 < p < +\infty$ , the operator

$$F^p(x_1, \dots, x_m) = t$$

from  $\mathbb{R}^m$  to  $\mathbb{R}$  defined implicitly by

$$\sum_{j=1}^m (x_j - t)|x_j - t|^{p-2} = 0$$

is a permutation invariant averaging operator.

**Example 2.4.** For  $0 \leq \alpha, \beta \leq 1$  with  $\alpha + \beta = 1$ , let us consider

$$F_1(x_1, \dots, x_m) = \alpha \operatorname{median}_{1 \leq j \leq m} \{x_j\} + \frac{\beta}{m} \sum_{j=1}^m x_j,$$

$$F_2(x_1, \dots, x_m) = \alpha \operatorname{median}_{1 \leq j \leq m} \{x_j\} + \frac{\beta}{2} \left( \max_{1 \leq j \leq m} \{x_j\} + \min_{1 \leq j \leq m} \{x_j\} \right),$$

where

$$\operatorname{median}_{1 \leq j \leq m} \{x_j\} := \begin{cases} y_{(m+1)/2} & \text{if } m \text{ is even,} \\ \frac{y_{m/2} + y_{(m/2+1)}}{2} & \text{if } m \text{ is odd,} \end{cases}$$

with  $\{y_1, \dots, y_m\}$  a nondecreasing rearrangement of  $\{x_1, \dots, x_m\}$ .

It holds that  $F_1$  and  $F_2$  are permutation invariant averaging operators.

### 3. Existence and uniqueness

First we show that there exists a unique solution of problem (1.1) in the space  $L_{\text{loc}}^\infty([0, +\infty); L^\infty(\mathbb{T}_m, \mathbb{R}))$ .

*Proof of Theorem 1.1.* Existence: Let  $T > 0$  and

$$C_T := \{u \in L^\infty(\mathbb{T}_m \times [0, T], \mathbb{R}) : u(x, t) \text{ is continuous in } t\}.$$

Observe that  $C_T$  is a Banach space with the  $L^\infty$ -norm.

We can see that  $K_f$  is a contraction on  $C_T$ . In fact, using Remark 2.2, we have that

$$\|K_f u_1 - K_f u_2\|_\infty \leq \int_0^t e^{z-t} dz \|u_1 - u_2\|_\infty \leq (1 - e^{-T}) \|u_1 - u_2\|_\infty,$$

for all  $u_1, u_2 \in C_T$ . Therefore, by the Brouwer fixed-point theorem,  $K_f$  has a unique fixed point  $u \in C_T$ .

Since  $T > 0$  is arbitrary, we can obtain a globally defined solution of (1.2),  $u$ .

Uniqueness: Let  $u, v$  be two solutions of (1.1) such that

$$u, v \in L_{\text{loc}}^\infty([0, +\infty); L^\infty(\mathbb{T}_m, \mathbb{R})).$$

Then  $u, v$  are solutions of (1.2) and therefore they are fixed points of  $K_f$ . Thus  $u \equiv v$  in  $\mathbb{T}_m \times [0, T]$  for all  $T > 0$  due to  $K_f$  is a contraction operator. Therefore  $u \equiv v$  in  $\mathbb{T}_m \times [0, +\infty)$ .  $\square$

**Remark 3.1.** We note that there is no need of a “boundary condition”. This problem can be regarded as the analogous for the tree to the Cauchy problem for a PDE, as  $u_t = \Delta u$  in  $\mathbb{R}^n \times (0, \infty)$  with  $u(x, 0) = f(x)$  in  $\mathbb{R}^n$ . Here we consider  $f \in L^\infty$ , but the result can be slightly improved to allow for an unbounded initial condition; see Remark 3.3 below.

Next we show a comparison principle.

*Proof of Theorem 1.2.* Let  $T > 0$ . We consider

$$M_T := \sup_{\mathbb{T}_m \times [0, T]} \{u - v\}.$$

Then, given  $\varepsilon > 0$ , there exists  $(\tilde{x}, \tilde{t}) \in \mathbb{T}_m \times [0, T]$  such that

$$M_T - \varepsilon \leq u(\tilde{x}, \tilde{t}) - v(\tilde{x}, \tilde{t}).$$

Now, by (1.3), we obtain that

$$\begin{aligned} M_T - \varepsilon &\leq u(\tilde{x}, \tilde{t}) - v(\tilde{x}, \tilde{t}) \\ &\leq \int_0^{\tilde{t}} e^{z-\tilde{t}} (F(u((\tilde{x}, 0), z), \dots, u((\tilde{x}, m-1), z)) \\ &\quad - F(v((\tilde{x}, 0), z), \dots, v((\tilde{x}, m-1), z))) dz + e^{-\tilde{t}} (f(\tilde{x}) - g(\tilde{x})). \end{aligned}$$

Thus, using that  $f \leq g$  in  $\mathbb{T}_m$  and Remark 2.2, we have

$$M_T - \varepsilon \leq M_T(1 - e^{-T}),$$

and therefore  $e^{-T} M_T \leq \varepsilon$  for all  $\varepsilon > 0$ . Then, using that  $e^{-T} > 0$ , we obtain that  $M_T \leq 0$  and this implies that  $u(x, t) \leq v(x, t)$  for all  $(x, t) \in \mathbb{T}_m \times [0, T]$ .

Since  $T > 0$  is arbitrary, we can conclude that  $u \leq v$  in  $\mathbb{T}_m \times [0, +\infty)$ .  $\square$

**Corollary 3.2.** *Let  $F$  be an averaging operator and  $f \in L^\infty(\mathbb{T}_m, \mathbb{R})$ . Then any bounded solution  $u$  of (1.1) with initial condition  $f$  satisfies the inequality*

$$|u(x, t)| \leq \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})}$$

for all  $(x, t) \in \mathbb{T}_m \times [0, +\infty)$ .

*Proof.* We just observe that  $w(x, t) = M = \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})}$  is the solution of (1.1) with initial condition  $M$ . Since  $f(x) \leq M$ , from Theorem 1.2, we obtain that

$$u(x, t) \leq M \quad \text{for all } (x, t) \in \mathbb{T}_m \times [0, +\infty).$$



In a similar way, we can prove that  $u(x, t) \geq -M$  for all  $(x, t) \in \mathbb{T}_m \times [0, +\infty)$ . Therefore,

$$|u(x, t)| \leq \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} \quad \forall (x, t) \in \mathbb{T}_m \times [0, +\infty).$$

This completes the proof.  $\square$

**Remark 3.3.** We remark that we can have existence of a solution even if the initial condition  $f$  is not bounded. In fact, we just observe that

$$u(x, t) = Ce^{(\lambda-1)t} \lambda^{l(x)},$$

with  $\lambda > 0$  is a solution of (1.1) with initial condition  $f(x) = C\lambda^{l(x)}$ .

Then there is a solution of (1.1) for any initial condition such that

$$0 \leq f(x) \leq C\lambda^{l(x)}.$$

To obtain such a solution we generate a sequence of approximating solutions using truncations of the initial condition. In fact, let

$$f_n(x) = \min\{f(x), n\}, \quad u_n(x, t) = \min\{u(x, t), n\},$$

and take  $w_n(x, t) \in L^\infty_{\text{loc}}([0, +\infty); L^\infty(\mathbb{T}_m, \mathbb{R}))$  the unique solution of (1.1) with initial condition  $f_n$  (Theorem 1.1).

We can see that,  $u_n \rightarrow u$  as  $n \rightarrow +\infty$ ,  $K_{f_n} u_n \leq u_n$ , and, by the comparison principle,  $w_n$  is increasing with  $n$  and  $w_n \leq u_n$ .

Finally, taking the limit as  $n \rightarrow \infty$  in the form of the equation given by (1.2), we obtain that  $w(x, t) := \lim_{n \rightarrow +\infty} w_n(x, t)$  is a solution of (1.1) with initial condition  $w(x, 0) = f(x)$ .

#### 4. Decay estimates

First, we prove Theorem 1.3.

*Proof of Theorem 1.3.* We begin by observing that if  $f \equiv 0$  on  $\mathbb{T}_m$  then  $u \equiv 0$  on  $\mathbb{T}_m \times [0, +\infty)$ . Therefore, (1.4) holds trivially in this case.

Now we consider the case  $f \not\equiv 0$ . Then  $a(f) \neq 0$ ,  $f(x) \neq 0$  for some  $x \in \mathbb{T}_m^{\mu(f)}$  and  $f(x) = 0$  for all  $x$  such that  $l(x) > \mu(f)$ . Thus, by Theorem 1.1,  $u(x, t) = 0$  for all  $x$  such that  $l(x) > \mu(f)$ . Therefore, if  $x \in \mathbb{T}_m^{\mu(f)}$ , we have that

$$\begin{aligned} u_t(x, t) &= F(u((x, 0), t), \dots, u((x, m-1), t)) - u(x, t) \\ &= F(0, \dots, 0) - u(x, t) = -u(x, t). \end{aligned}$$

Then

$$\frac{d}{dt}(e^t u(x, t)) = 0.$$

Since  $u(x, 0) = f(x)$  for all  $x \in \mathbb{T}_m$ , we get

$$u(x, t) = f(x)e^{-t} \quad \forall x \in \mathbb{T}_m^{\mu(f)}.$$

Thus, for any  $x \in \mathbb{T}_m^{\mu(f)-1}$  we have that

$$\begin{aligned} u_t(x, t) &= F(u((x, 0), t), \dots, u((x, m-1), t)) - u(x, t) \\ &= F(f(x, 0)e^{-t}, \dots, f(x, m-1)e^{-t}) - u(x, t) \\ &= F(f(x, 0), \dots, f(x, m-1))e^{-t} - u(x, t). \end{aligned}$$

Then

$$\frac{d}{dt}(e^t u(x, t)) = \mathcal{A}_x^1,$$

where  $\mathcal{A}_x^1 = F(f(x, 0), \dots, f(x, m-1))$ . Therefore,

$$u(x, t) = (\mathcal{A}_x^1 t + f(x))e^{-t} \quad \forall x \in \mathbb{T}_m^{\mu(f)-1}. \quad (4.5)$$

Observe that

$$|\mathcal{A}_x^1| \leq \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} \quad \forall x \in \mathbb{T}_m^{\mu(f)-1}, \quad (4.6)$$

due to the fact that  $F$  is nondecreasing with respect to each variable.

Arguing as before, using (4.5), we obtain

$$\frac{d}{dt}(e^t u(x, t)) = F(\mathcal{A}_{(x,0)}^1 t + f(x, 0), \dots, \mathcal{A}_{(x,m-1)}^1 t + f(x, m-1)),$$

for every  $x \in \mathbb{T}_m^{\mu(f)-2}$ . Then, since  $F$  is nondecreasing with respect to each variable, we have

$$\mathcal{A}_x^2 t - \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} \leq \frac{d}{dt}(e^t u(x, t)) \leq \mathcal{A}_x^2 t + \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} \quad \forall x \in \mathbb{T}_m^{\mu(f)-2},$$

where

$$\mathcal{A}_x^2 = F(\mathcal{A}_{(x,0)}^1, \dots, \mathcal{A}_{(x,m-1)}^1).$$

Therefore,

$$e^{-t} \left( \mathcal{A}_x^2 \frac{t^2}{2} - \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} t + f(x) \right) \leq u(x, t) \leq e^{-t} \left( \mathcal{A}_x^2 \frac{t^2}{2} + \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} t + f(x) \right)$$

for all  $x \in \mathbb{T}_m^{\mu(f)-2}$ .

By (4.6), using again that  $F$  is nondecreasing with respect to each variable, we obtain

$$|\mathcal{A}_x^2| \leq \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} \quad \forall x \in \mathbb{T}_m^{\mu(f)-2}.$$

Continuing in the same manner, we can prove

$$e^{-t} p_1(t) \leq u(\emptyset, t) \leq e^{-t} p_2(t),$$

where

$$\begin{aligned} p_1(t) &= \mathcal{A}_\emptyset^{\mu(f)} \frac{t^{\mu(f)}}{\mu(f)!} - \sum_{j=1}^{\mu(f)-1} \frac{t^j}{j!} \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} + f(\emptyset), \\ p_2(t) &= \mathcal{A}_\emptyset^{\mu(f)} \frac{t^{\mu(f)}}{\mu(f)!} + \sum_{j=1}^{\mu(f)-1} \frac{t^j}{j!} \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} + f(\emptyset), \\ \mathcal{A}_\emptyset^{\mu(f)} &= F(\mathcal{A}_{(\emptyset, 0)}^{\mu(f)-1}, \dots, \mathcal{A}_{(\emptyset, m-1)}^{\mu(f)-1}). \end{aligned}$$

Arguing as before, we have

$$|\mathcal{A}_\emptyset^{\mu(f)}| \leq \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})}.$$

Thus,

$$\max_{x \in \mathbb{T}_m} |u(x, t)| \leq \frac{t^{\mu(f)} e^{-t}}{\mu(f)!} \|f\|_{L^\infty(\mathbb{T}_m, \mathbb{R})}$$

for  $t$  large enough. □

**Remark 4.1.** The bound that we obtained in Theorem 1.3 is optimal. In fact, let  $n \in \mathbb{N}$ ,  $F$  be an averaging operator and  $f_n \in L^\infty(\mathbb{T}_m, \mathbb{R})$  defined by

$$f_n(x) := \begin{cases} n! & \text{if } l(x) = n, \\ 0 & \text{if } l(x) \neq n. \end{cases}$$

Note that  $\|f_n\|_{L^\infty(\mathbb{T}_m, \mathbb{R})} = n!$  and  $\mu(f_n) = n$ . Let

$$z_n(x, t) := \begin{cases} \frac{n!}{(n-l(x))!} t^{(n-l(x))} & \text{if } 0 \leq l(x) \leq n, \\ 0 & \text{if } l(x) > n. \end{cases}$$

Then we observe that  $u_n(x, t) := e^{-t} z_n(x, t) \in L^\infty_{\text{loc}}([0, +\infty); L^\infty(\mathbb{T}_m, \mathbb{R}))$ ,  $u_n$  is the solution of (1.1) with initial condition  $f_n$ , and

$$\max_{x \in \mathbb{T}_m} |u_n(x, t)| = t^n e^{-t} = \frac{t^{\mu(f_n)} e^{-t}}{\mu(f_n)!} \|f_n\|_{L^\infty(\mathbb{T}_m, \mathbb{R})}.$$

**Proposition 4.2.** *Let  $F$  be an averaging operator and  $f(x) = (1 - \lambda)^{l(x)}$  for some  $\lambda \in (0, 1)$ . Then  $u(x, t) = e^{-\lambda t} f(x)$  is the solution to (1.1).*

*Proof.* We have  $u(x, 0) = f(x)$  for all  $x \in \mathbb{T}_m$  and

$$\begin{aligned} \Delta_F u(x, t) &= F(u((x, 0), t), \dots, u((x, m-1), t)) - u(x, t) \\ &= F(e^{-\lambda t} f(x, 0), \dots, e^{-\lambda t} f(x, m-1)) - e^{-\lambda t} f(x) \\ &= e^{-\lambda t} F(f(x, 0), \dots, f(x, m-1)) - e^{-\lambda t} f(x) \\ &= e^{-\lambda t} F((1 - \lambda)^{l(x)+1}, \dots, (1 - \lambda)^{l(x)+1}) - e^{-\lambda t} (1 - \lambda)^{l(x)} \\ &= e^{-\lambda t} (1 - \lambda)^{l(x)+1} - e^{-\lambda t} (1 - \lambda)^{l(x)} \\ &= e^{-\lambda t} (1 - \lambda)^{l(x)} (1 - \lambda + 1) \\ &= -\lambda e^{-\lambda t} (1 - \lambda)^{l(x)} \\ &= u_t(x, t) \end{aligned}$$

for all  $(x, t) \in \mathbb{T}_m \times (0, +\infty)$ . □

We observe that for this particular solution we have

$$\max_{x \in \mathbb{T}_m} u(x, t) = e^{-\lambda t} \max_{x \in \mathbb{T}_m} f(x) = e^{-\lambda t} = u(\emptyset, t).$$

Therefore, using the comparison principle stated in Theorem 1.2, we obtain Theorem 1.4 as an immediate consequence.

Proposition 4.2 shows that the bound is optimal.

Finally, let us prove that there are solutions with any prescribed behaviour of  $u(\emptyset, t)$ .

*Proof of Theorem 1.5.* We just consider  $u(x, t) \equiv a_{l(x)}(t)$  (that is, we take  $u$  to be constant at every level). Then the equation reduces to find  $a_1, a_2, \dots, a_n, \dots$

such that

$$a_i'(t) = a_{i+1}(t) - a_i(t),$$

that is,

$$a_{i+1}(t) = a_i'(t) + a_i(t).$$

Hence, given  $a_0$ , we can construct

$$\begin{aligned} a_1(t) &= a_0'(t) + a_0(t), \\ a_2(t) &= a_0''(t) + 2a_0'(t) + a_0(t) \end{aligned}$$

etc., that is, at level  $n$ , we have

$$a_n(t) = \sum_{j=0}^n \binom{n}{j} a_0^{(j)}(t).$$

Therefore

$$u(x, t) = \sum_{j=0}^{l(x)} \binom{l(x)}{j} a_0^{(j)}(t)$$

is a solution of the equation. □

Remark that depending on the behaviour of the derivatives of  $a_0$  it may hold that

$$u(\emptyset, t) = a_0(t) = \max_{x \in \mathbb{U}_m} u(x, t).$$

If we have

$$a_0(t) = (1 + t)^{-\alpha} \quad (\alpha > 0),$$

then we get

$$u(x, t) = \sum_{j=0}^{l(x)} \binom{l(x)}{j} a_0^{(j)}(t) = \sum_{j=0}^{l(x)} \binom{l(x)}{j} (-1)^j \left( \prod_{i=0}^{j-1} (\alpha + i) \right) (1 + t)^{-\alpha-j}.$$

Note that we have as initial condition for this particular solution

$$f(x) = \sum_{j=0}^{l(x)} \binom{l(x)}{j} (-1)^j \left( \prod_{i=0}^{j-1} (\alpha + i) \right).$$

Note that this initial condition can be unbounded. For example, for  $\alpha = 1$  we have

$$a_0(t) = (1 + t)^{-1}.$$

Then we get that

$$\begin{aligned} u(x, t) &= \sum_{j=0}^{l(x)} \binom{l(x)}{j} a_0^{(j)}(t) \\ &= \sum_{j=0}^{l(x)} (-1)^j \binom{l(x)}{j} j! (1 + t)^{-(j+1)} \\ &= \sum_{j=0}^{l(x)} (-1)^j \frac{l(x)!}{(l(x) - j)!} (1 + t)^{-(j+1)} \end{aligned}$$

is a solution of (1.1) with initial condition

$$f(x) = \sum_{j=0}^{l(x)} (-1)^j \frac{l(x)!}{(l(x) - j)!} = (-1)^{l(x)} l(x)!,$$

where  $!n$  denotes the subfactorial of  $n$ .

We observe that  $f(x)$  is an oscillating function with

$$|f(x)| \rightarrow +\infty \quad \text{as } l(x) \rightarrow +\infty.$$

## References

- [1] V. Alvarez, J. M. Rodríguez, and D. V. Yakubovich, Estimates for nonlinear harmonic “measures” on trees. *Michigan Math. J.* **49** (2001), 47–64. Zbl 1006.31006 MR 1827075
- [2] L. M. Del Pezzo, C. A. Mosquera and J. D. Rossi, The unique continuation property for a nonlinear equation on trees. *J. London Math. Soc.*, to appear. doi: 10.1112/jlms/jdt067
- [3] S.-Y. Chung and C. A. Berenstein,  $\omega$ -harmonic functions and inverse conductivity problems on networks. *SIAM J. Appl. Math.* **65** (2005), 1200–1226. Zbl 1068.05040 MR 2147325
- [4] E. B. Curtis and J. A. Morrow, Determining the resistors in a network. *SIAM J. Appl. Math.* **50** (1990), 918–930. Zbl 0717.35092 MR 1050922
- [5] L. M. Del Pezzo, C. A. Mosquera and J. D. Rossi, Estimates for nonlinear harmonic measures on trees. Preprint 2013. arXiv:1303.6521

- [6] R. Kaufman, J. G. Llorente, and J.-M. Wu, Nonlinear harmonic measures on trees. *Ann. Acad. Sci. Fenn. Math.* **28** (2003), 279–302. Zbl 1033.31004 MR 1996439
- [7] R. Kaufman and J.-M. Wu, Fatou theorem of  $p$ -harmonic functions on trees. *Ann. Probab.* **28** (2000), 1138–1148. Zbl 1038.31007 MR 1797306
- [8] J. J. Manfredi, A. Oberman and A. Sviridov. Nonlinear elliptic PDEs and  $p$ -harmonic functions on graphs. Preprint 2012. arXiv:1212.0834
- [9] A. P. Sviridov, Elliptic equations in graphs via stochastic games. Ph.D. thesis, University of Pittsburgh, Pittsburgh 2011. <http://d-scholarship.pitt.edu/9581/>
- [10] A. P. Sviridov,  $p$ -harmonious functions with drift on graphs via games. *Electron. J. Differential Equations* (2011), No. 114. Zbl 1229.35304 MR 2836795
- [11] A. Weber, Analysis of the physical Laplacian and the heat flow on a locally finite graph. *J. Math. Anal. Appl.* **370** (2010), 146–158. Zbl 1214.35075 MR 2651136
- [12] R. K. Wojciechowski, Heat kernel and essential spectrum of infinite graphs. *Indiana Univ. Math. J.* **58** (2009), 1419–1441. Zbl 1231.05186 MR 2542093

Received September 29, 2013; revised February 26, 2014

L. M. Del Pezzo, CONICET and Departamento de Matemática, FCEyN, Universidad de Buenos Aires, Pabellon I, Ciudad Universitaria (1428), Buenos Aires, Argentina  
E-mail: [ldpezzo@dm.uba.ar](mailto:ldpezzo@dm.uba.ar)

C. A. Mosquera, CONICET and Departamento de Matemática, FCEyN, Universidad de Buenos Aires, Pabellon I, Ciudad Universitaria (1428), Buenos Aires, Argentina  
E-mail: [mosquera@dm.uba.ar](mailto:mosquera@dm.uba.ar)

J. D. Rossi, Departamento de Análisis Matemático, Universidad de Alicante, Ap. correo 99, 03080, Alicante, Spain  
E-mail: [julio.rossi@ua.es](mailto:julio.rossi@ua.es)