

On multisetsemigroups

Ganna Kudryavtseva and Volodymyr Mazorchuk

(Communicated by Jorge Almeida)

Abstract. Motivated by recent appearance of multivalued structures in categorification, tropical geometry and other areas, we study basic properties of abstract multisetsemigroups. We give many new and old examples and general constructions for multisetsemigroups. Special attention is paid to simple and nilpotent multisetsemigroups. We also show that “almost all” randomly chosen multivalued binary operations define multisetsemigroups.

1. Introduction and motivation

A *multisetsemigroup* is a “semigroup in which multiplication is multivalued”. More precisely, a multisetsemigroup is a pair $(S, *)$, where S is a non-empty set and $*$ is a map (a so-called *multivalued operation* or *multioperation*) from $S \times S$ to 2^S , the power set of S , such that the following associativity axiom is satisfied: for every $a, b, c \in S$ we have

$$\bigcup_{t \in b * c} a * t = \bigcup_{s \in a * b} s * c. \quad (1.1)$$

Every semigroup is a multisetsemigroup in an obvious way, however, there are many natural examples of multisetsemigroups which are not semigroups.

Our motivation for the present paper comes from the fact observed in [MM2] that multisetsemigroups appear naturally in higher representation theory and categorification, see Subsection 3.8 of this paper and also [MM2] for more details.

Definition of multistructures in general goes back at least to the 1934 paper [Mar] by Marty. Various aspects and properties of multistructures, in particular, multisetsemigroups (usually under different names which will be reviewed in the next section), were studied by several authors, see for example [Ca], [DO], [Ea], [Ha], [Ko], [Wa]. We also refer the reader to the recent survey paper [Vi] by Viro, which mostly deals with multigroups and multirings, for more historical information. The paper [Vi] shows that some multistructures (mainly multifields) are relevant to the study of tropical geometry.

In the present paper we make an attempt to establish basic properties of abstract multisemigroups in analogy with those of semigroups. We give many examples and counterexamples illustrating the scopes of the theory we are developing. We will see that in many respects semigroups and multisemigroups are similar but in many other respects they are very different. Section 2 fixes notation and vocabulary for multivalued analogues of binary operations and corresponding structures, including multisemigroups, hypergroups and multigroups. In Section 3 we propose many different examples and constructions of multisemigroups. These include both concrete examples of multisemigroups of small cardinality (Subsections 3.1, 3.2 and 3.15), the multisemigroup of cosets of a group modulo a subgroup (Subsection 3.3) and various constructions associated to different kinds of ideals, variants, subwords etc. (Subsections 3.4, 3.5, 3.6, 3.11, 3.12 and 3.14). We also mention several multisemigroups related to our motivation: the multisemigroup associated to a fully additive bicategory (Subsection 3.8) and its disguise via a positive basis in an associative algebra (Subsection 3.7). We also show that the underlying set of every Weyl group can be equipped with two very different multisemigroups structures: the first one coming from the Kazhdan-Lusztig combinatorics (Subsection 3.9) and the other one coming from the standard basis of the corresponding Hecke algebra for the values of the parameter which guarantee that this standard basis has necessary positivity properties (Subsection 3.10).

In Section 4 we collect basic notions and properties: Green's relations, various types of elements, ideals, homomorphisms, congruences, quotients, representations by binary relations and connections to dioids, quantales and Boolean algebras. In Section 5 we study simple multisemigroups (that is multisemigroups with a unique two-sided ideal) and a special class of simple multisemigroups which we call *strongly simple*. A simple multisemigroup is strongly simple provided that it is a union of its minimal left ideals and it is a union of its minimal right ideals. Strongly simple multisemigroups can be viewed as analogues of completely 0-simple or 0-bisimple semigroups. For strongly simple multisemigroups we establish a structure theory similar to the classical structure theory of 0-bisimple semigroups. However, there are significant differences: The role of idempotent \mathcal{H} -classes is now played by hypergroups (and the latter do not need to have any idempotents or quasi-idempotents, see an example in Subsection 3.15). There is only a partial analogue of Green's lemma, namely minimal right (left) ideals are no longer connected by bijective translations but by surjective multivalued translations. We give explicit examples showing that the general case behaves much more complicated than the case of classical semigroups (see Subsections 5.3, 5.4, 5.5, 5.6 and 5.7).

In Section 6 we establish another essential difference between semigroups and multisemigroups. Given a set with n elements, one can randomly choose a binary operation on it and ask what is the probability that it is associative

(hence defining a semigroup). The answer is that this probability is “small” in the sense that it tends to 0 when n tends to infinity. For multisetsemigroups the picture turns out to be exactly the opposite: when n tends to infinity, then the probability that a randomly chosen multivalued operation on an n -element set defines a multisetsemigroup approaches 1.

Finally, in Section 7 we look closer at nilpotent multisetsemigroups, characterize them in terms of the action graphs, extend to finite multisetsemigroups two classical characterizations of finite nilpotent semigroups, define and characterize the radical of a multisetsemigroup and study maximal (with respect to inclusions) nilpotent submultisetsemigroups of strongly simple multisetsemigroups.

Acknowledgment. This work was done during the visit of the first author to Uppsala University, which was supported by the Department of Mathematics, Uppsala University and the Swedish Research Council. Hospitality and support of Uppsala University is gratefully acknowledged. The research of the first author was also partially supported by the ARRS grant P1-0288. For the second author the research was partially supported by the Royal Swedish Academy of Sciences and the Swedish Research Council. We thank Denis Saveliev for useful comments and discussions. We are really thankful to the referees for very careful reading of the paper, many very helpful suggestions and for pointing out several subtle inaccuracies in the original version of the paper.

2. Multistructures and their aliases

2.1. Multisetsemigroups. Multisetsemigroups, as defined above, have appeared in the literature under many different names. In particular, we have seen the following names: hypersemigroups, polysemigroups, semihypergroups and associative multiplicative systems (the list disregards different hyphenations of the names). Following [MM2], we use the name “multisetsemigroups” as we think that, compared to all other aliases, it describes the essence of the structure best. Note that the operation $*$ of a multisetsemigroup $(S, *)$ can also be understood as a binary relation from $S \times S$ to S (satisfying the usual associativity axiom).

An element 1 of a multisetsemigroup $(S, *)$ is called an *identity* or *unity* element provided that $1 * a = a * 1 = a$ for all $a \in S$. A standard argument shows that the identity element is unique, if exists. For any multisetsemigroup $(S, *)$ and any element $e \notin S$, the set $S \cup \{e\}$ has the natural structure $(S \cup \{e\}, \circ)$ of a multisetsemigroup defined, for $a, b \in S \cup \{e\}$, as follows:

$$a \circ b := \begin{cases} a * b, & a, b \in S; \\ a, & b = e; \\ b, & a = e. \end{cases}$$

The element e is the identity of $(S \cup \{e\}, \circ)$. As usual, we denote by S^1 the multisemigroup S if the latter has an identity and $S \cup \{e\}$ defined as above if S does not have any identity (and we denote this identity by 1 and the operation on S^1 by $*$).

An element e of a multisemigroup $(S, *)$ is called an *idempotent* provided that $e * e = e$ and a *quasi-idempotent* provided that $e \in e * e$.

Multisemigroups are closely connected to some other *different* algebraic structures. Here we describe two of such connections. A map $*$: $S \times S \rightarrow 2^S$, $(a, b) \mapsto a * b$, can be extended to a binary operation $*$: $2^S \times 2^S \rightarrow 2^S$ by setting, for $A, B \in 2^S$,

$$A * B := \bigcup_{a \in A, b \in B} a * b. \quad (2.1)$$

The equality (1.1) can now be written as $a * (b * c) = (a * b) * c$. In this way, for a multisemigroup $(S, *)$, the power set 2^S inherits a natural structure of a semigroup, where the associativity follows directly from (1.1). Thus, a multisemigroup can be seen as a non-empty set S along with an associative binary operation $*$ on 2^S satisfying (2.1) for any $A, B \subset S$. Notice that to show that such a binary operation $*$ on 2^S defines a multisemigroup on S it is enough to show that $a * (b * c) = (a * b) * c$ for all $a, b, c \in S$. From (2.1) it follows that for any $A, B_i \in 2^S$, $i \in I$, we have the following property:

$$A * \left(\bigcup_i B_i \right) = \bigcup_i (A * B_i) \quad \text{and} \quad \left(\bigcup_i B_i \right) * A = \bigcup_i (B_i * A). \quad (2.2)$$

Recall, see e.g. [Gu], 2.2, that a *semiring* is a tuple $(R, +, *, 0, 1)$ where R is a non-empty set, $+$ and $*$ are binary operations on R and 0 and 1 are two distinguished elements of R such that the following conditions are satisfied:

- $(R, +, 0)$ is a commutative monoid with identity 0;
- $(R, *, 1)$ is a monoid with identity 1;
- $r * 0 = 0 * r = 0$ for all $r \in R$ (i.e. 0 is *absorbing* with respect to $*$);
- $r * (s + t) = r * s + r * t$ and $(s + t) * r = s * r + t * r$ for all $r, s, t \in R$.

A semiring $(R, +, *, 0, 1)$ for which $r + r = r$ for all $r \in R$ is called an *idempotent semiring* or *dioid*. The previous paragraph implies that any multisemigroup $(S, *)$ induces a natural dioid structure $(2^{S^1}, \cup, *, \emptyset, \{1\})$ on 2^{S^1} . If Q is finite, then every dioid structure $(2^Q, \cup, *, \emptyset, 1)$, where 1 is a singleton, gives, by restriction to elements of Q , a multisemigroup $(Q, *)$ possessing a unit element.

A notable difference between multisemigroups and dioids is revealed comparing the arbitrary distributivity property for multisemigroup given by (2.2) with the finite distributivity property for dioids. This discrepancy motivates connection of multisemigroups with quantales.

Recall, see [Re], that a *sup-lattice* is a partially ordered set in which every subset has a *supremum*, or a *join*. A *quantale* is a sup-lattice equipped with an associative product $(a, b) \mapsto a * b$ such that the multiplication distributes over arbitrary suprema, that is for any $X \subset Q$ and $a \in Q$ we have

$$a * \left(\sup_{x \in X} x \right) = \sup_{x \in X} (a * x) \quad \text{and} \quad \left(\sup_{x \in X} x \right) * a = \sup_{x \in X} (x * a).$$

A *unital quantale* is a quantale possessing a unit element with respect to the product operation. Unital quantales form a special class of dioids, the so-called *complete dioids*.

The discussion above shows that any multisetsemigroup $(S, *)$ induces a natural quantale structure on 2^S . Being defined on a power set, this quantale is a *complete atomic Boolean algebra* where atomic means that every nonzero element lies above an atom, a minimal nonzero element. Conversely, every quantale structure on a complete atomic Boolean algebra Q induces a natural structure of a multisetsemigroup on the set $S = S(Q)$ of atoms of Q . This can be used to identify multisetsemigroups and quantale structures on complete atomic Boolean algebras.

2.2. Multigroups and hypergroups. Unlike multisetsemigroups, whose definition is more or less uncontroversial despite of many different aliases, there exist many *different* multistructures considered in the literature as multivalued analogues of groups. The present paper is not about these structures, so to fix terminology for the present paper we will just define those of them which we will use later. We refer the reader to [DO], [Vi], [Wa] and references therein for further details.

Following [Vo], a multisetsemigroup $(S, *)$ will be called a *hypergroup* provided that it satisfies the following *reproduction axiom*: $S * a = a * S = S$ for any $a \in S$.

Following [Vi], a multisetsemigroup $(S, *)$ with identity 1 will be called a *multigroup* provided that the following two conditions are satisfied:

- for each $a \in S$ there are unique $b, c \in S$ such that $1 \in a * b$ and $1 \in c * a$, moreover, $b = c$ (this unique element will be denoted by a^{-1});
- for any $a, b, c \in S$ we have $c \in a * b$ if and only if $c^{-1} \in b^{-1} * a^{-1}$.

It is clear that $1^{-1} = 1$ and $(a^{-1})^{-1} = a$ for any element a of a multigroup S (cf. [Vi], 3.2).

2.3. Involution. If $(S, *)$ is a multisetsemigroup, then an *involution* on S is a transformation $\star : S \rightarrow S$, written $s \mapsto s^*$, which is involutive, that is $(s^*)^* = s$ for all $s \in S$, and satisfies, for all $a, b \in S$, the following equality:

$$b^* * a^* = \{s^* \mid s \in a * b\}.$$

For example, if $(S, *)$ is a multigroup, then $a \mapsto a^{-1}$ is an involution.

3. Some examples of multisemigroups

In this section we collect many examples of multisemigroups, for further examples of multigroups we refer the reader to [Vi] and references therein.

3.1. The trivial multisemigroups. For any non-empty set S and a subset $X \subset S$ setting $s *_X t := X$ for all $s, t \in S$ defines on S the structure of a multisemigroup. In particular, we have two *trivial* multisemigroup structures on S , namely, $\diamond := *_\emptyset$ and $\bullet := *_S$. We note that (S, \bullet) is always a hypergroup, but it is not a multigroup if $|S| > 1$.

3.2. A two-element multisemigroup. Define the operation $*$ on $S := \{a, b\}$ using the following Cayley table:

| | | |
|-----|-----|------------|
| $*$ | a | b |
| a | a | $\{a, b\}$ |
| b | a | b |

It is straightforward to verify that this is a multisemigroup. We note that $x * S = S$ for any $x \in S$ while $S * a = a \neq S$ (in particular, this is a *right* hypergroup but not a hypergroup).

3.3. The coset multisemigroup. Let (G, \cdot) be a group and H a (not necessarily normal) subgroup of G . Define a multivalued operation $*$ on G as follows: for every $a, b \in G$ we set $a * b := HaHb$. It is straightforward to verify that $(a * b) * c = a * (b * c) = HaHbHc$ for all $a, b, c \in G$ and hence $(G, *)$ is a multisemigroup.

We can also consider the set $H \setminus G$ of left H -cosets in G . Then for any $a, b \in G$ the set $HaHb$ is a union of cosets and hence we may define

$$Ha * Hb := \{Hc \mid c \in G \text{ and } Hc \subset HaHb\}.$$

This turns $(H \setminus G, *)$ into a multisemigroup and even a hypergroup (but not a multigroup if H is not normal since an identity would be necessarily H , and H is normal if and only if, for all $a, b \in G$, $H \subset HaHb$ implies $Hb = Ha^{-1}$). If H is a normal subgroup of G , then the operation $*$ on $H \setminus G$ is, in fact, single-valued and hence $(H \setminus G, *)$ is a group.

3.4. Inflations of multisemigroups. Let $(S, *)$ be a multisemigroup, X an arbitrary set, and $f : X \rightarrow S$ a surjective map. For $x, y \in X$ define

$$x *_f y := \{z \in X \mid f(z) \in f(x) * f(y)\}.$$

Then it is straightforward to verify that $(X, *_f)$ is a multisetsemigroup called the *inflation* of S with respect to f . Note that the trivial multisetsemigroup (S, \bullet) defined in Subsection 3.1 can be viewed as an inflation of a singleton group.

3.5. Multisetsemigroups of ideals. Let (S, \cdot) be a semigroup. Define multioperations $*_L$ and $\hat{*}_L$ on S as follows: for $a, b \in S$ set

$$a *_L b := S^1 a S^1 b \quad \text{and} \quad a \hat{*}_L b := S^1 a \cap S^1 b.$$

Then it is straightforward to verify that both $(S, *_L)$ and $(S, \hat{*}_L)$ are multisetsemigroups. Similarly one defines multisetsemigroups $(S, *_R)$ and $(S, \hat{*}_R)$ using right ideals and multisetsemigroups $(S, *_J)$ and $(S, \hat{*}_J)$ using two-sided ideals. The multisetsemigroups $(S, \hat{*}_L)$, $(S, \hat{*}_R)$ and $(S, \hat{*}_J)$ are commutative.

3.6. Monogenic associated multisetsemigroups. Let (S, \cdot) be a semigroup. For $a \in S$ let $\langle a \rangle$ denote the subsemigroup of S consisting of all elements of the form a^i , $i > 0$ (the so-called “monogenic subsemigroup” generated by a). Define the multioperation $*$ on S as follows: for $a, b \in S$ set

$$a * b := \langle a \rangle \cap \langle b \rangle.$$

Then it is straightforward to verify that $(S, *)$ is a commutative multisetsemigroup.

3.7. Multisetsemigroups of positive bases in associative algebras. Let A be an associative algebra over some subring \mathbb{k} of real numbers. Assume that A has a basis $\mathbf{a} := \{a_i \mid i \in S\}$ with non-negative structure constants, that is

$$a_i a_j = \sum_{k \in S} c_{i,j}^k a_k \quad \text{and} \quad c_{i,j}^k \geq 0 \quad \text{for all } i, j, k \in S.$$

Define the multioperation $*$ on S as follows: for $i, j \in S$ set

$$i * j := \{k \mid c_{i,j}^k > 0\}.$$

Then the associativity of A implies that $(S, *)$ is a multisetsemigroup.

A similar construction works if instead of a subring of real numbers one considers, for example, the Boolean algebra $\mathbb{B} := \{0, 1\}$ (with respect to the usual meet and join operations).

3.8. Multisetsemigroups of fully additive bicategories. This example is taken from [MM2], Subsection 3.3. Let \mathcal{C} be a small additive bicategory with skeletally small, fully additive and Krull-Schmidt categories of morphisms. Let $S[\mathcal{C}]$ be the set of isomorphism classes of indecomposable 1-morphisms in \mathcal{C} . For

an indecomposable 1-morphism F we denote by $[F]$ its class in $S[\mathcal{C}]$. For $[F], [G] \in S[\mathcal{C}]$ set

$$[F] * [G] := \{[H] \in S[\mathcal{C}] : H \text{ is isomorphic to a direct summand of } F \circ G\}.$$

Then the associativity axiom for \mathcal{C} implies that $(S[\mathcal{C}], *)$ is a multisemigroup. Via decategorification (i.e. taking the split Grothendieck group of \mathcal{C}) this example can be considered as a special case of the previous example.

As a more concrete example of this construction, consider \mathcal{C} to be a bicategory with one object \bullet and such that $\mathcal{C}(\bullet, \bullet)$ is the category of all finite dimensional representations of a semi-simple complex finite dimensional Lie algebra \mathfrak{g} , with horizontal composition given by the usual tensor product of \mathfrak{g} -modules. It is easy to see that in this case the obtained multisemigroup is, in fact, a multigroup. In the case of the algebra \mathfrak{sl}_2 , isomorphism classes of simple finite dimensional modules are in a natural bijection with the set \mathbb{N}_0 of non-negative integers (this bijection is given by taking the highest weight of a module, see [Maz2], Theorem 1.22). From the classical Clebsch-Gordan rule (see e.g. [Maz2], Theorem 1.39) it follows that for $k, l \in \mathbb{N}_0$ the multisemigroup operation is given by the following:

$$k * l = \{m : |k - l| \leq m \leq k + l, m \equiv k + l \pmod{2}\}.$$

Remark 1. The concrete example above can be generalized. For \mathfrak{g} as above consider the BGG category \mathcal{O} and let $\bar{\mathcal{O}}$ be the tensor category which \mathcal{O} generates (see e.g. [Kaa]). The category $\bar{\mathcal{O}}$ is no longer semi-simple but all objects in $\bar{\mathcal{O}}$ have well-defined composition multiplicities (see e.g. [Kaa]). There is a natural multisemigroup structure on the set of isomorphism classes of simple objects in $\bar{\mathcal{O}}$ defined as follows: For two simple objects L and L' define $[L] * [L']$ to be the set of isomorphism classes of all simple subquotients which appear in the tensor product of L and L' .

3.9. The Kazhdan-Lusztig multisemigroup of a Weyl group. Let Δ be a finite root system and W the corresponding Weyl group. Let $\{\underline{H}_w : w \in W\}$ be the Kazhdan-Lusztig basis of $\mathbb{Z}[W]$, see [KL] or [Maz1], Section 7. By [KL], this basis has positive structure constants and hence the construction of Subsection 3.7 gives a multisemigroup structure $(W, *)$. Remark that this multisemigroup can be also obtained by the construction of Subsection 3.8 considering the bicategory of Soergel bimodules acting on the principal block of category \mathcal{O} for the Lie algebra \mathfrak{g} associated with Δ , see [Maz1], Section 8 and [MM1], [MM2] for details. Mapping $w \mapsto w^{-1}$ defines an involution on this multisemigroup (it corresponds to “taking the adjoint functor” in the categorical picture of Soergel bimodules).

To give an explicit example, let Δ be of type A_2 , that is $W \simeq S_3 = \{e, s, t, st, ts, sts\}$, where $s^2 = t^2 = e$ and $sts = tst$. Then we have

$$\begin{aligned} \underline{H}_e &= e, & \underline{H}_s &= e + s, & \underline{H}_t &= e + t, & \underline{H}_{st} &= e + s + t + st, \\ \underline{H}_{ts} &= e + s + t + ts, & \underline{H}_{sts} &= e + s + t + st + ts + sts; \end{aligned}$$

and one easily obtains the following Cayley table for $(S_3, *)$:

| * | e | s | t | st | ts | sts |
|-------|-------|--------------|--------------|---------------|---------------|-------|
| e | e | s | t | st | ts | sts |
| s | s | s | st | st | $\{sts, s\}$ | sts |
| t | t | ts | t | $\{sts, t\}$ | ts | sts |
| st | st | $\{sts, s\}$ | st | $\{sts, st\}$ | $\{sts, s\}$ | sts |
| ts | ts | ts | $\{sts, t\}$ | $\{sts, t\}$ | $\{sts, ts\}$ | sts |
| sts | sts | sts | sts | sts | sts | sts |

3.10. The Boolean Hecke multigroup of a Weyl group. This example is taken from [Tr] where it is given in different terms. Let Δ be a finite root system and W the corresponding Weyl group. Choose some basis π in Δ and let $S \subset W$ be the corresponding system of simple reflections. Then (W, S) is a Coxeter system. Let $l : W \rightarrow \mathbb{N}_0$ be the corresponding length function. Fix some $q \in \mathbb{R}$ such that $q > 1$ and let \mathbb{H}_q be the corresponding Hecke algebra (over \mathbb{R}), that is the associative algebra with generators $H_s, s \in S$, satisfying the braid relations for (W, S) together with the relations

$$H_s^2 = (q - 1)H_s + qH_e, \quad s \in S. \tag{3.1}$$

Note that under our choice of q the latter relation has positive coefficients.

For each $w \in W$ fix some reduced expression $w = s_1 s_2 \dots s_k$ and set $H_w := H_{s_1} H_{s_2} \dots H_{s_k}$. Since the H_s 's satisfy braid relations, the element H_w does not depend on the choice of a reduced expression for w . Then $\{H_w \mid w \in W\}$ is the standard basis of \mathbb{H}_q .

Lemma 2. *All structure constants for the standard basis are non-negative.*

Proof. As $H_w := H_{s_1} H_{s_2} \dots H_{s_k}$, it is enough to show that for any $s \in S$ and $x \in W$ the element $H_s H_x$ is a linear combination of basis elements with non-negative coefficients. If $l(sx) > l(x)$, then we have $H_s H_x = H_{sx}$. In the other case we have $x = sy$ for some $y \in W$ such that $l(y) < l(x)$. Then, using (3.1), we have

$$H_s H_x = H_s H_s H_y = ((q - 1)H_s + qH_e)H_y = (q - 1)H_x + qH_y$$

and the claim follows. □

From Lemma 2 it follows that the construction of Subsection 3.7 gives a multi-semigroup structure $(W, *)$. This multi-semigroup is called the *Boolean Hecke hypermonoid* in [Tr]. If Δ is of type A_2 , then for $W \simeq S_3 = \{e, s, t, st, ts, sts\}$ we get the following Cayley table:

| $*$ | e | s | t | st | ts | sts |
|-------|-------|---------------|---------------|----------------------|----------------------|----------------------------|
| e | e | s | t | st | ts | sts |
| s | s | $\{e, s\}$ | st | $\{t, st\}$ | sts | $\{ts, sts\}$ |
| t | t | ts | $\{e, t\}$ | sts | $\{s, ts\}$ | $\{st, sts\}$ |
| st | st | sts | $\{s, st\}$ | $\{ts, sts\}$ | $\{e, s, sts\}$ | $\{t, st, ts, sts\}$ |
| ts | ts | $\{t, ts\}$ | sts | $\{e, t, sts\}$ | $\{st, sts\}$ | $\{s, st, ts, sts\}$ |
| sts | sts | $\{st, sts\}$ | $\{ts, sts\}$ | $\{s, st, ts, sts\}$ | $\{t, st, ts, sts\}$ | $\{e, s, t, st, ts, sts\}$ |

Proposition 3. *The multi-semigroup $(W, *)$ is, in fact, a multigroup.*

Proof. We have to check that both additional conditions from Subsection 2.2 are satisfied. Let $x \in W$. First we show, by induction on $l(x)$, that $e \in x * x^{-1}$. This is clear if $x = e$, so to prove the induction step assume that the claim is true for some x and that $s \in S$ is such that $l(sx) > l(x)$. Then, using associativity and definitions, we have $(sx) * (sx)^{-1} = s * x * x^{-1} * s$. The latter set contains $s * e * s = \{e, s\}$ as $e \in x * x^{-1}$ by the inductive assumption.

Now let $x, y \in W$ be such that $x \neq y^{-1}$. Consider the case $l(x) \leq l(y)$, in particular, $y \neq e$ (the other case is dealt with by similar arguments). Let us prove, by induction on $l(x)$, the following three claims:

- (i) any $w \in x * y$ satisfies $l(w) \geq l(y) - l(x)$;
- (ii) $x * y$ contains some w satisfying $l(w) = l(y) - l(x)$ if and only if $y = x^{-1}w$;
- (iii) $e \notin x * y$.

Note that the last claim obviously follows from the first two. The basis $x = e$ of the induction is obvious. To prove the induction step let $s \in S$ be a simple reflection such that $l(x) < l(sx) \leq l(y)$. Then we have $sx = s * x$ and hence $(sx) * y = s * (x * y)$ by associativity. By the inductive assumption, any $w \in x * y$ satisfies $l(w) \geq l(y) - l(x)$. We have $s * w = sw$ if $l(sw) > l(w)$ and $s * w = \{w, sw\}$ otherwise. In the second case $l(sw) \geq l(y) - l(x) - 1 = l(y) - l(sx)$ which proves claim (i). This argument also shows that if k is maximal such that $x * y$ contains some w of length $l(y) - k$, then every element in $s * (x * y)$ has length at least $l(y) - k - 1$. Therefore, for $s * (x * y)$ to contain an element of length $l(y) - l(sx)$, the set $x * y$ must contain an element of length $l(y) - l(x)$. By (ii) of

the inductive assumption, the only element of $x * y$ with this property is xy and this is possible if and only if $l(xy) = l(y) - l(x)$. In the latter case $s * xy$ contains an element of length $l(y) - l(x) - 1$ if and only if $l(sxy) = l(y) - l(x) - 1$ and this is the case if and only if $l(sxy) = l(y) - l(sx)$ which implies claim (ii). This proves the first condition from Subsection 2.2.

Mapping $H_s \mapsto H_s$ extends to an anti-involution on \mathbb{H}_q (as the ideal generated by the defining relations is invariant under this map). This anti-involution maps H_w to $H_{w^{-1}}$ for any $w \in W$ and hence mapping $w \mapsto w^{-1}$ is an involution on $(W, *)$. This proves the second condition from Subsection 2.2. \square

3.11. Double variants of multisetsemigroups. Let (S, \cdot) and (S, \circ) be two multisetsemigroups with the same underlying set S . Assume further that for any $a, b, c \in S$ we have the following equalities in 2^S :

$$(a \cdot b) \circ c = a \cdot (b \circ c) \quad \text{and} \quad (a \circ b) \cdot c = a \circ (b \cdot c). \quad (3.2)$$

For $a, b \in S$ set $a * b := (a \cdot b) \cup (a \circ b)$.

Proposition 4. $(S, *)$ is a multisetsemigroup.

Proof. For $a, b, c \in S$ we have the following:

$$\begin{aligned} (a * b) * c &= ((a \cdot b) \cup (a \circ b)) * c \\ &= ((a \cdot b) * c) \cup ((a \circ b) * c) \\ &= ((a \cdot b) \cdot c) \cup ((a \cdot b) \circ c) \cup ((a \circ b) \cdot c) \cup ((a \circ b) \circ c) \\ &\stackrel{(3.2)}{=} (a \cdot (b \cdot c)) \cup (a \cdot (b \circ c)) \cup (a \circ (b \cdot c)) \cup (a \circ (b \circ c)) \\ &= (a \cdot (b \cdot c)) \cup (a \circ (b \cdot c)) \cup (a \cdot (b \circ c)) \cup (a \circ (b \circ c)) \\ &= (a * (b \cdot c)) \cup (a * (b \circ c)) \\ &= a * ((b \cdot c) \cup (b \circ c)) \\ &= a * (b * c). \end{aligned}$$

The claim follows. \square

A typical situation in which this construction applies is the following multisetsemigroup version of the *variant* (or *sandwich*) construction for semigroups: Let (S, \bowtie) be a multisetsemigroup and $X, Y \subset S$. Then for $a, b \in S$ set

$$a \cdot b := a \bowtie X \bowtie b, \quad \text{and} \quad a \circ b := a \bowtie Y \bowtie b.$$

Then both (S, \cdot) and (S, \circ) are *variant* multisetsemigroups of (S, \bowtie) , moreover, condition (3.2) is obviously satisfied. Thus $(S, *)$ is a new multisetsemigroup which

is natural to call a *double variant* of (S, \bowtie) . It is easy to see that $a * b = a \bowtie (X \cup Y) \bowtie b$. In the case when (S, \bowtie) is a semigroup and $|X| = |Y| = 1$ we have that $(S, *)$ is a multisemigroup such that $1 \leq |a * b| \leq 2$ for all $a, b \in S$.

3.12. Multisemigroup of subwords. Let A be an alphabet and A^* the monoid of all finite words over A . For $u, v \in A^*$ define $u \circledast v$ to be the set of all scattered (that is, not necessarily connected) subwords of uv . It is straightforward to verify that for any $u, v, w \in A^*$ both $(u \circledast v) \circledast w$ and $u \circledast (v \circledast w)$ coincide with the set of all scattered subwords of uvw and hence (A^*, \circledast) is a multisemigroup.

3.13. Disconnected unions of multisemigroups. Let (S, \cdot) and (T, \bullet) be multisemigroups and assume that $S \cap T = \emptyset$. Define a multivalued operation $*$ on $S \cup T$ as follows: for $a, b \in S \cup T$ set

$$a * b := \begin{cases} a \cdot b, & a, b \in S; \\ a \bullet b, & a, b \in T; \\ \emptyset, & \text{otherwise.} \end{cases}$$

It is straightforward to verify that this turns $(S \cup T, *)$ into a multisemigroup, which we call the *disconnected union* of S and T .

3.14. Reproductive construction. The following general approach to construction of multisemigroups is inspired by [Vo]. Let (S, \cdot) be a semigroup and $f : S \rightarrow 2^S$ a map. For $A \subset S$ set $f(A) := \bigcup_{a \in A} f(a)$. For $a, b \in S$ define $a * b := f(a)f(b)$.

Lemma 5. *Assume that for any $a, b \in S$ we have $f(f(a)f(b)) = f(a)f(b)$. Then $(S, *)$ is a multisemigroup.*

The condition $f(f(a)f(b)) = f(a)f(b)$ resembles the *reproductive condition* in [Vo].

Proof. Using our assumption, for $a, b, c \in S$ we have:

$$\begin{aligned} (a * b) * c &= \bigcup_{s \in f(a)f(b)} (f(s)f(c)) = \left(\bigcup_{s \in f(a)f(b)} f(s) \right) f(c) \\ &= f(f(a)f(b))f(c) = f(a)f(b)f(c). \end{aligned}$$

Similarly one checks that $a * (b * c) = f(a)f(b)f(c)$. □

Some of the previous examples can be obtained using the reproductive construction. For instance, the example in Subsection 3.12 is obtained if we define f to be the map which sends a word w to the set of all scattered subwords of w ; the

first example in Subsection 3.3 is obtained if we define f to be the map which sends a to Ha ; and the first example in Subsection 3.5 is obtained if we define f to be the map which sends a to S^1a .

3.15. A hypergroup without quasi-idempotents. Let S be a set satisfying $|S| \geq 3$. For $a, b \in S$ we define

$$a * b := \begin{cases} S, & a \neq b; \\ S \setminus \{a\}, & a = b. \end{cases}$$

Then $|a * b| > 1$ for any $a, b \in S$, which implies $(a * b) * c = a * (b * c) = S$ for all $a, b, c \in S$. Thus $(S, *)$ is a multisetsemigroup. Obviously, S is a hypergroup and it does not contain any quasi-idempotent.

4. Elementary properties of multisetsemigroups

In this section we provide multisetsemigroup analogues of some basic notions from semigroup theory as well as record some basic properties of multisetsemigroups.

4.1. Ideals and Green's relations. Let $(S, *)$ be a multisetsemigroup. A subset $I \subset S$ is called a *left ideal* (resp. *right ideal*, *two-sided ideal*) provided that for any $a \in I$ and $s \in S$ we have $s * a \subset I$ (resp. $a * s \subset I$; $a * s, s * a \subset I$). For example, for every $a \in S$ the set $S^1 * a$ is the smallest left ideal containing a , called the *principal left ideal* generated by a . Similarly one has the *principal right ideal* $a * S^1$ and the *principal two-sided ideal* $S^1 * a * S^1$. We define the left pre-order \leq_L , the right pre-order \leq_R and the two-sided pre-order \leq_J on S as follows: for $a, b \in S$ set $b \leq_L a$ if and only if $S^1 * b \subset S^1 * a$, $b \leq_R a$ if and only if $b * S^1 \subset a * S^1$, and $b \leq_J a$ if and only if $S^1 * b * S^1 \subset S^1 * a * S^1$.

Following [Gr], we define an equivalence relation \mathcal{L} on S as the equivalence relation induced by \leq_L , i.e. $a \mathcal{L} b$ if and only if $a \leq_L b$ and $b \leq_L a$. Similarly we define relations \mathcal{R} and \mathcal{J} (see also [Ha], [MM2]). We set $\mathcal{H} := \mathcal{L} \cap \mathcal{R}$ and denote by \mathcal{D} the minimal equivalence relation containing both \mathcal{L} and \mathcal{R} . The relations \mathcal{L} , \mathcal{R} , \mathcal{J} , \mathcal{H} and \mathcal{D} are called *Green's relations*. Obviously, $\mathcal{D} \subset \mathcal{J}$. Note that the equality $\mathcal{L} \circ \mathcal{R} = \mathcal{R} \circ \mathcal{L}$ (where \circ denotes the usual product of binary relations) which holds for any semigroup fails for multisetsemigroups in general, see Subsection 5.5. For an element $a \in S$ we denote by \mathcal{L}_a the \mathcal{L} -class of S containing a . We define $\mathcal{R}_a, \mathcal{H}_a, \mathcal{J}_a$ and \mathcal{D}_a similarly.

Example 6. Let (S, \cdot) be a semigroup and $(S, *_L)$ be the corresponding multisetsemigroup of left ideals defined in Subsection 3.5. For $a \in S$ we have $S^1a = S^1 *_L a$ and hence the \mathcal{L} -classes in (S, \cdot) and in $(S, *_L)$ coincide. Any two-sided

ideal of (S, \cdot) is, in particular, a left ideal, which implies that the \mathcal{J} -classes in (S, \cdot) and in $(S, *_L)$ coincide as well. On the other hand, for any $a \in S$ we have $a *_L S^1 = S^1 (aS^1) = S^1 *_L a *_L S^1$, which implies that the relations \mathcal{R} and \mathcal{J} for $(S, *_L)$ coincide with \mathcal{J} in (S, \cdot) . Hence we also have $\mathcal{D} = \mathcal{J}$ and $\mathcal{L} = \mathcal{H}$ for the multisetimigroup $(S, *_L)$.

4.2. Rees quotients. Let $(S, *)$ be a multisetimigroup and $I \subset S$ a two-sided ideal different from S (possibly empty). Consider the set $T := S \setminus I$ and for $a, b \in T$ set $a \bullet b := (a * b) \setminus I$. It is straightforward to verify that this turns (T, \bullet) into a multisetimigroup. Making a parallel with the classical semigroup theory, we will call the multisetimigroup (T, \bullet) the *Rees quotient* of S modulo the ideal I .

4.3. Zero elements. Let $(S, *)$ be a multisetimigroup. An element $z \in S$ is called a *zero element* provided that for every $a \in S$ we have $a * z = z * a = z$. A zero element is necessarily unique, if exists, and therefore it is natural to denote this unique zero element by 0 .

Let $(S, *)$ be a multisetimigroup with the zero 0 and suppose that $S \neq \{0\}$. Then we claim that for any $a, b \in S$ we have $a * b \neq \emptyset$. Indeed, assume that $a * b = \emptyset$, then, on the one hand, $(a * b) * 0 = \emptyset$, but, on the other hand, $a * (b * 0) = a * 0 = 0$, a contradiction. Consider the set $T := S \setminus \{0\}$ and for $a, b \in T$ set $a \bullet b := (a * b) \setminus \{0\}$. It is straightforward to verify that (T, \bullet) is a multisetimigroup (see also Subsection 4.2).

Conversely, let $(S, *)$ be a multisetimigroup without a zero element. Consider the set $S^0 := S \cup \{0\}$, where we assume $0 \notin S$, and for $a, b \in S^0$ define

$$a \bullet b := \begin{cases} (a * b) \cup \{0\}, & a, b \in S; \\ \{0\}, & \text{otherwise.} \end{cases}$$

Isomorphism of multisetimigroups is defined in the obvious way, that is, two multisetimigroups (X_1, \circ_1) and (X_2, \circ_2) are called *isomorphic*, denoted $X_1 \cong X_2$, provided that there is a bijection $\varphi : X_1 \rightarrow X_2$ such that for any $x, y \in X_1$ we have

$$\{\varphi(s) : s \in x \circ_1 y\} = \varphi(x) \circ_2 \varphi(y).$$

Lemma 7. (a) *The construct (S^0, \bullet) is a multisetimigroup with the zero 0 .*

(b) *The set $\{0\}$ is an ideal of (S^0, \bullet) .*

(c) *We have $(S^0, \bullet) \setminus \{0\} \cong (S, *)$.*

Proof. Let $a, b, c \in S^0$. If one of these elements equals 0 , then both sides of (1.1), with \bullet in place of $*$, are equal to 0 . If $a, b, c \in S$, then both sides of (1.1), with \bullet in place of $*$, are equal to $(a * b * c) \cup \{0\}$. This proves claim (a). Claims (b) and (c) follow from the definition of \bullet . \square

By the above one can consider multisetsemigroups *without* zero elements and understand that the role of the zero element is played by the “undefined” multiplication, that is the case $a * b = \emptyset$. This, in particular, unifies the notions of “simple” and “0-simple” multisetsemigroups and semigroups.

A multisetsemigroup $(S, *)$ will be called a *quasi-semigroup* provided that for any $a, b \in S$ the product $a * b$ is either empty or an element of S . Quasi-semigroups can be identified with semigroups with zero elements: Given a semigroup with a zero element we can take this zero element away and redefine the product to be empty whenever it was zero to obtain a quasi-semigroup. Conversely, let $(S, *)$ be a quasi-semigroup. Consider the set $S^0 := S \cup \{0\}$, where we assume $0 \notin S$, and for $a, b \in S^0$ define

$$a \odot b := \begin{cases} a * b; & a, b \in S, a * b \neq \emptyset; \\ \{0\}, & \text{otherwise.} \end{cases}$$

Then (S^0, \odot) becomes a semigroup with a zero element. Note that the previous construction (S^0, \bullet) produces in this case a multisetsemigroup, not a semigroup.

4.4. Homomorphisms. Let $(S, *)$ and (T, \bullet) be multisetsemigroups. A *strong homomorphism* from S to T is a map $\varphi : S \rightarrow T$ such that for any $a, b \in S$ we have

$$\{\varphi(s) : s \in a * b\} = \varphi(a) \bullet \varphi(b).$$

Comparing this with the notion of isomorphism defined in the previous subsection, we see that an isomorphism is just a bijective strong homomorphism. If $\varphi : S \rightarrow T$ is a strong homomorphism, it extends uniquely to a quantale homomorphism $\bar{\varphi} : 2^S \rightarrow 2^T$ by setting $\bar{\varphi}(A) := \bigcup_{a \in A} \{\varphi(a)\}$ for $A \subset S$. By definition, $\bar{\varphi}$ maps atoms of 2^S to atoms of 2^T . A homomorphism of atomic quantales which maps atoms to atoms is called *atomic*. Conversely, any atomic homomorphism from 2^S to 2^T gives, via restriction to atoms, a strong homomorphism from S to T .

Let **MSemi** denote the category of multisetsemigroups with strong homomorphisms. By the above, this category is equivalent to the category **QCABA'** of quantale structures on complete atomic Boolean algebras with atomic homomorphisms.

As multisetsemigroups are defined in a multi-setting anyway, it is natural to extend the above as follows: Let $(S, *)$ and (T, \bullet) be multisetsemigroups. A *weak homomorphism* from S to T is a map $\varphi : S \rightarrow 2^T$ such that for any $a, b \in S$ we have

$$\bigcup_{s \in a * b} \varphi(s) = \varphi(a) \bullet \varphi(b).$$

If $\varphi : S \rightarrow 2^T$ is a weak homomorphism, it extends uniquely to a quantale homomorphism $\bar{\varphi} : 2^S \rightarrow 2^T$ by setting $\bar{\varphi}(A) := \bigcup_{a \in A} \varphi(a)$ for $A \subset S$. Conversely, any quantale homomorphism from 2^S to 2^T gives, via restriction to atoms of S , a weak homomorphism from S to T .

Let $\overline{\mathbf{MSemi}}$ denote the category multisemigroups with weak homomorphisms. By the above, this category is equivalent to the category \mathbf{QCABA} of quantale structures on complete atomic Boolean algebras with usual homomorphisms.

4.5. Submultisemigroups. A non-empty subset T of a multisemigroup $(S, *)$ is called a *submultisemigroup* provided that $a * b \subset T$ for all $a, b \in T$. Any submultisemigroup T is a multisemigroup with respect to the restriction of $*$ to T .

Let $(S, *)$ be a multisemigroup and $X \subset S$ a non-empty subset. Then the *submultisemigroup* $\langle X \rangle$ of S generated by X is the minimal (with respect to inclusion) submultisemigroup of S containing X . Alternatively, $\langle X \rangle$ is the intersection of all submultisemigroups of S containing X .

Let $(S, *)$ be a multisemigroup and J a \mathcal{J} -class of S . Set $T = \langle J \rangle$. As J is a \mathcal{J} -class, $T \setminus J$ is an ideal of T and hence we can consider the corresponding Rees quotient $T \setminus (T \setminus J) = J$. Note that J might not be a semigroup even if S is a semigroup. This construction is a natural multisemigroup analogue of the standard construction of the “semigroup associated with a \mathcal{J} -class”.

4.6. Congruences and quotients. Let $(S, *)$ be a multisemigroup. An equivalence relation \sim on S is called a *left congruence* provided that the following condition is satisfied: for any $a, b, c \in S$ such that $a \sim b$ and for any $s \in c * a$ and $t \in c * b$ there are $s' \in c * b$ and $t' \in c * a$ such that $s \sim s'$ and $t \sim t'$. A *right congruence* is defined similarly (using $a * c$ and $b * c$ at appropriate places) and a *congruence* is an equivalence relation which is both a left and a right congruence (confer [Da]).

Let \sim be a congruence on $(S, *)$. Then the usual argument shows that the quotient set S/\sim has the natural structure of a multisemigroup with respect to the induced multioperation \circ defined for $A, B \in S/\sim$ as follows:

$$A \circ B = \{C \in S/\sim \mid \text{there exist } a \in A, b \in B, c \in C \text{ such that } c \in a * b\}.$$

Mapping $s \in S$ to its \sim -class \bar{s} is obviously a strong homomorphism from the multisemigroup $(S, *)$ to the multisemigroup $(S/\sim, \circ)$. We have $A \circ B = \{\bar{s} \mid s \in A * B\}$ for $A, B \in S/\sim$, furthermore, for any $a, b \in S$ we have $\bar{a} \circ \bar{b} = \{\bar{s} \mid s \in a * b\}$.

Let (T, \cdot) be another multisemigroup and $\varphi : S \rightarrow T$ a strong homomorphism. Then the equivalence relation $\text{Ker}(\varphi)$ on S is easily seen to be a congruence. Hence, as usual, congruences on multisemigroups are exactly kernels of strong homomorphisms.

Let \sim be a congruence on a multisetsemigroup $(S, *)$. Then the map j which sends $a \in S$ to its \sim -class extends to the map $j : 2^S \rightarrow 2^S$, by $j(A) = \bigcup_{a \in A} j(a)$. It is straightforward to verify that this map is monotone, extensive, idempotent and satisfies $j(A) * j(B) \leq j(A * B)$. It follows that j is a quantic nucleus, see [Ro], Chapter 3. The *quotient quantale defined by j* is a complete atomic Boolean algebra with the atoms $j(a)$, $a \in S$. Setting $T = S/\sim$, it is easy to see that this quotient quantale is just the quantale 2^T associated to the multisetsemigroup T . Conversely, assume that we are given a complete atomic Boolean algebra 2^X with a quantale structure on it and assume that $j : 2^X \rightarrow 2^X$ is a nucleus such that the quotient quantale is a complete atomic Boolean algebra 2^Y . On the multisetsemigroup X we define a relation \sim as follows: $x \sim y$ if and only if $j(\{x\}) = j(\{y\})$. This relation is a congruence and the quotient multisetsemigroup is Y , the multisetsemigroup of atoms of 2^Y . It follows that congruences on multisetsemigroups are in a bijective correspondence with congruences on quantale structures on complete atomic Boolean algebras.

4.7. Representations of multisetsemigroups by binary relations. Regular representations of multisetsemigroups by binary relations were constructed in [Ea] in the following way: Let $(S, *)$ be a multisetsemigroup. Then $*$ can be viewed as a binary relation $S \times S \rightarrow S$, in particular, every $a \in S$ defines a binary relation τ_a on S via $y \tau_a x$ if and only if $y \in a * x$ (this corresponds to the usual convention that maps operate from the right to the left). Our convention for multiplication \circ of binary relations is as follows: for two binary relations τ and σ we have $y(\tau \circ \sigma)x$ if and only if there is z such that $y \tau z$ and $z \sigma x$. This allows us to identify binary relations on S with square matrices over \mathbb{B} , whose rows and columns are indexed by elements of S (i.e. $a \tau b$ if and only if the intersection of row a and column b of the matrix of τ contains 1). Then for the usual sum $+$ and product \circ of binary relations (i.e. sum and product of Boolean matrices) we have

$$\tau_a \circ \tau_b = \sum_{s \in a * b} \tau_s.$$

This is the left regular representation of $(S, *)$. The right regular representation is defined similarly.

This representation extends to a homomorphism from the quantale 2^S to the quantale $\mathbf{B}(S)$ of binary relations on S . Therefore it is natural to generalize this and call a *representation* of $(S, *)$ (by binary relations) any quantale homomorphism from 2^S to any quantale of binary relations.

Similarly, given an \mathcal{L} -class L in S , for $a \in S$ we define the binary relation λ_a on L via $y \lambda_a x$ if and only if $y \in a * x$. This extends, using addition of binary relations, to a homomorphism from the quantale 2^S to the quantale $\mathbf{B}(L)$ of

binary relations on L . This is the representation of S associated to L (confer [GM2], Section 10).

5. Strongly simple multisemigroups

5.1. Simple and strongly simple multisemigroups. A multisemigroup $(S, *)$ is called *simple* if for any $a \in S$ we have $S^1 * a * S^1 = S$, that is S has a unique \mathcal{J} -class (confer [JSM]). From now on, unless stated otherwise, we assume that S does not contain a zero element (see Subsection 4.3). This excludes one special case: when $(S, *)$ is a singleton group. In the latter case $(S, *)$ contains both the identity and the zero elements and they coincide. This case is usually excluded in the classical ring theory as well.

Recall the example of a two-element multisemigroup, $(S, *)$, from Subsection 3.2. This multisemigroup is finite simple with a unique \mathcal{R} -class but two different \mathcal{L} -classes which are, moreover, comparable with respect to \leq_L . This shows a significant difference between multisemigroups and semigroups as no analogous situation is possible for semigroups.

From now on, if the converse is not explicitly stated, by a *minimal* left ideal of S we mean a minimal non-empty left ideal, that is a non-empty left ideal I of S such that for any left ideal J of S the inclusion $J \subseteq I$ implies that $J = I$ or $J = \emptyset$. In a similar way we will also use the notions of a minimal right ideal and a minimal (two-sided) ideal.

If $(S, *)$ is a multisemigroup, an element $s \in S$ will be called a *quark* provided that $S^1 * s$ is a minimal left ideal and $s * S^1$ is a minimal right ideal. For any quark s we thus have $\mathcal{L}_s = S^1 * s$ and $\mathcal{R}_s = s * S^1$. We denote by $Q(S)$ the set of all quarks in $(S, *)$. The set $Q(S)$ will be called the *support* of S . A simple multisemigroup $(S, *)$ will be called *strongly simple* if $S = Q(S)$. For instance, any completely simple semigroup is a strongly simple multisemigroup. Furthermore, the quasi-semigroup associated to a completely 0-simple semigroup as described at the end of Subsection 4.3 is a strongly simple multisemigroup. A special case of the last example is the singleton multisemigroup $\mathbf{0} := (\{0\}, *)$, where $0 * 0 = \emptyset$. Moreover, the multisemigroup $\mathbf{0}$ is very special because of the following:

Proposition 8. *Let $(S, *)$ be a multisemigroup. If $(S, *)$ contains only one \mathcal{H} -class, then either $S \cong \mathbf{0}$ or S is a hypergroup.*

Proof. Assume that $S = \mathcal{H}_a$ for every $a \in S$. If $|S| = 1$ then $S \cong \mathbf{0}$ (note that S cannot be isomorphic to the trivial semigroup as the trivial semigroup contains a zero element) and hence is a hypergroup. Suppose that $|S| > 1$. Let $a \in S$. We have $a * S^1 = b * S^1 \ni b$, for any $b \in S$, which implies that $a * S \supset S \setminus \{a\}$.

Assume that $a * S = S \setminus \{a\}$. Then $S \setminus \{a\}$ is a proper right ideal of S and hence for any $b \in S \setminus \{a\}$ we have $b * S^1 \subset S \setminus \{a\}$, a contradiction. Hence $a * S = S$. Similarly one shows that $S * a = S$ and the claim follows. \square

For $Q(S)$ we have the following elementary property:

Proposition 9. *Let $(S, *)$ be a multisetsemigroup with non-empty support. Then we have:*

- (a) $Q(S)$ and every non-empty intersection of $Q(S)$ with a \mathcal{J} -class of S is a submultisetsemigroup.
- (b) The multisetsemigroup $Q(S)$ is a disconnected union (as defined in Subsection 3.13) of its non-empty intersections with \mathcal{J} -classes of S .

Proof. Let $a, b \in Q(S)$. Every element $s \in a * b$ belongs to $S^1 * b \cap a * S^1$ and hence to $Q(S)$. This shows that $Q(S)$ is a submultisetsemigroup and claim (a) follows.

Let $a, b \in Q(S)$ be such that a and b belong to different \mathcal{J} -classes of S . If $s \in a * b$, then $s \in S^1 * b$ and hence $s \mathcal{L} b$ by the minimality of $S^1 * b$. At the same time $s \in a * S^1$ and hence $s \mathcal{R} a$ by the minimality of $a * S^1$. This implies $s \mathcal{J} b$ and $s \mathcal{J} a$ which means that $a \mathcal{J} b$ contradicting our assumptions. This implies that $a * b = \emptyset$ and hence $Q(S)$ is the disconnected union of its intersections with \mathcal{J} -classes of S . \square

Lemma 10. *Let $(S, *)$ be a simple multisetsemigroup, I a non-empty left ideal of S and J a non-empty right ideal of S . Then $I \cap J \neq \emptyset$ and $J * I \neq \emptyset$.*

Proof. The claim is obvious in the case $|S| = 1$, so we assume that $|S| > 1$. Since $I \cap J \supset J * I$, it is enough to show that $J * I \neq \emptyset$. Assume that this is not the case, that is $J * I = \emptyset$. Since $S^1 * J$ is a non-empty ideal of S , we have $S = S^1 * J$ since S is simple. Similarly $S = I * S^1$. But then

$$S * S = S^1 * J * I * S^1 = \emptyset.$$

This is, however, not possible if S is simple and $|S| > 1$. This completes the proof. \square

We say that a multisetsemigroup $(S, *)$ is of *finite type* provided that every (non-empty) left ideal of S contains a minimal left ideal and every (non-empty) right ideal of S contains a minimal right ideal. Clearly, every finite multisetsemigroup is of finite type.

Corollary 11. *Let $(S, *)$ be a simple multisetsemigroup of finite type. Then $Q(S) \neq \emptyset$.*

Lemma 12. *Let $(S, *)$ be a multisemigroup and $a, b \in S$. Then the following statements are equivalent:*

- (a) $a * b = \emptyset$.
- (b) $(S^1 * a) * (b * S^1) = \emptyset$.
- (c) $\mathcal{L}_a * \mathcal{R}_b = \emptyset$.
- (d) *There is $s \in \mathcal{L}_a$ and $t \in \mathcal{R}_b$ such that $s * t = \emptyset$.*

Proof. That (b) implies (c) is obvious. That (c) implies (d) is obvious. That (a) implies (b) follows from the following computation, which uses associativity: $(S^1 * a) * (b * S^1) = S^1 * (a * b) * S^1 = \emptyset$. The same argument shows that (d) implies $(S^1 * s) * (t * S^1) = \emptyset$ and, as $a \in S^1 * s$ and $b \in t * S^1$, we get that (d) implies (a). \square

Corollary 13. *Let $(S, *)$ be a multisemigroup and H an \mathcal{H} -class in S . Then the following conditions are equivalent:*

- (i) $H * H \neq \emptyset$.
- (ii) *There exist $s, t \in H$ such that $s * t \neq \emptyset$.*
- (iii) *For all $s, t \in H$ we have $s * t \neq \emptyset$.*

Proof. Obviously, condition (iii) implies condition (i) and condition (i) implies condition (ii). For $a \in H$ we have $H = \mathcal{L}_a \cap \mathcal{R}_a$ and the fact that condition (ii) implies condition (iii) follows from Lemma 12. \square

Proposition 14. *Let $(S, *)$ be a multisemigroup and $a, b \in Q(S)$. Then $a * b \subset \mathcal{L}_b \cap \mathcal{R}_a$. Moreover, if S is simple and $H := \mathcal{L}_a \cap \mathcal{R}_b$, then $a * b \neq \emptyset$ if and only if $H * H \neq \emptyset$.*

Proof. The first claim is obvious, so we prove the second one. Suppose $a * b \neq \emptyset$. Since $a, b \in Q(S)$ it follows that $\mathcal{L}_a, \mathcal{L}_b$ are minimal left ideals and $\mathcal{R}_a, \mathcal{R}_b$ are minimal right ideals. By Lemma 10 we have $H \neq \emptyset$ and hence there exists $y \in H$. We have $y * y \neq \emptyset$ by Lemma 12 and thus $H * H \neq \emptyset$. Conversely, if $H * H \neq \emptyset$, then $a * b \neq \emptyset$ by Lemma 12. \square

Proposition 15. *Let $(S, *)$ be a multisemigroup such that $Q(S) \neq \emptyset$. Then $Q(S)$ is a union of \mathcal{H} -classes of S , the restriction \sim of the relation \mathcal{H} to $Q(S)$ is a congruence on $Q(S)$ and $Q(S)/\sim$ is a quasi-semigroup with singleton \mathcal{H} -classes.*

Proof. If $a \in Q(S)$ and $b \mathcal{H} a$, then clearly $b \in Q(S)$, so $Q(S)$ is a union of \mathcal{H} -classes. Let $a, b \in Q(S)$ be such that $a \sim b$. Then there exist a minimal left ideal I and a minimal right ideal J of S such that $a, b \in I \cap J$. Let further $c \in Q(S)$.

Then there exist a minimal left ideal I' and a minimal right ideal J' of S such that $c \in I' \cap J'$. But then we have both $a * c \in I' \cap J$ and $b * c \in I' \cap J$, which implies that \sim is a right congruence on $Q(S)$. That \sim is a left congruence is proved similarly. Hence \sim is a congruence on $Q(S)$. The elements of $Q(S)/\sim$ can be identified with pairs (I, J) , where I is a minimal left ideal in S and J is a minimal right ideal in S such that $I \cap J \neq \emptyset$ (the pair (I, J) corresponds to $I \cap J$). The above also shows that the multiplication in $Q(S)/\sim$ is given by

$$(I, J) \cdot (I', J') = \begin{cases} (I', J), & (I \cap J) * (I' \cap J') \neq \emptyset; \\ \emptyset, & \text{otherwise.} \end{cases}$$

This is obviously a quasi-semigroup with singleton \mathcal{H} -classes. □

5.2. Structure of strongly simple multisetsemigroups. Let $(S, *)$ be a strongly simple multisetsemigroup. Then S is both, the (disjoint) union of its minimal left ideals and the (disjoint) union of its minimal right ideals. At the same time, if S is simple and every element of S generates a minimal left and a minimal right ideal, then S is strongly simple. Furthermore, $S^1 * a = \mathcal{L}_a$ and $a * S^1 = \mathcal{R}_a$ for any $a \in S$. In this subsection we show how the results of the previous subsection can be strengthened in the case of strongly simple multisetsemigroups. It turns out that hypergroups arise naturally and play a very important role in this case (similar to the role of group \mathcal{H} -classes for completely 0-simple semigroups).

Theorem 16 (Structure of strongly simple multisetsemigroups). *Let $(S, *)$ be a strongly simple multisetsemigroup.*

- (a) *For any $a, b \in S$ we have $\mathcal{L}_a \cap \mathcal{R}_b \neq \emptyset$.*
- (b) *If H is an \mathcal{H} -class in S , then either $H * H = \emptyset$ or H is a hypergroup.*
- (c) *For $a, b \in S$ we have $a * b \neq \emptyset$ if and only if $\mathcal{L}_a \cap \mathcal{R}_b$ is a hypergroup.*
- (d) *Assume $S \not\cong \mathbf{0}$. Then every \mathcal{L} -class and every \mathcal{R} -class in S contains at least one hypergroup \mathcal{H} -class.*
- (e) *Let $a, b \in S$ be such that $a \mathcal{R} b$ and let $s \in S^1$ be such that $b \in a * s$. Then the multivalued map $x \mapsto x * s$ is surjective from \mathcal{L}_a to \mathcal{L}_b and preserves both \mathcal{R} - and \mathcal{H} -classes, that is, for any \mathcal{H} -class $H \subset \mathcal{L}_a$ we have $H * s$ is an \mathcal{H} -class in \mathcal{L}_b and the two \mathcal{H} -classes H and $H * s$ are contained in the same \mathcal{R} -class.*
- (f) *Assume $S \not\cong \mathbf{0}$. Let I be a minimal left ideal of S and J a minimal right ideal of S . Then $I \cap J = J * I$.*
- (g) *\mathcal{H} is a congruence on S and the quasi-semigroup S/\mathcal{H} is bisimple (i.e. contains a unique \mathcal{D} -class) with singleton \mathcal{H} -classes.*

Proof. Claim (a) is a special case of Lemma 10.

Let us prove claim (b). Let H be an \mathcal{H} -class in S and assume that $H * H \neq \emptyset$. Let L and R be the \mathcal{L} -class and the \mathcal{R} -class whose intersection is H . We have that $L * R$ is an ideal of S and is non-empty (since $H * H \neq \emptyset$). Hence $L * R = S$. We first show that $H * H = H$. Let $a \in H$. Since $L * R = S \supseteq H$, it follows that $a \in b * c$ for some $b \in L$ and $c \in R$. But $b * c \subseteq b * S^1 \cap S^1 * c$. Since the latter is an \mathcal{H} -class and $b * c \cap H \neq \emptyset$ then $H = b * S^1 \cap S^1 * c$. Since also $H = S^1 * b \cap c * S^1$ it follows that $b \in H$ and $c \in H$. Therefore $H = H * H$, as required. Let $a \in H$. Show that $H * a = H$. Let $c \in H$. As $H = H * H$, there are some $x, y \in H$ such that $c \in x * y$. Now $a \mathcal{R} x$ implies $x \in a * t$ for some $t \in S^1$. We then have $c \in x * y \subseteq a * t * y$. So $c \in a * L$ (because $t * y \subseteq L$). We have shown that $H \subseteq a * L$. So $H = a * L$ since clearly $a * L \subseteq H$. Observe that $L * a$ is a non-empty left ideal and so $L * a = L$. Hence we have

$$H * a = (a * L) * a = a * (L * a) = a * L = H.$$

Similarly we verify that $a * H = H$ and claim (b) follows.

Claim (c) follows from claim (b) and Proposition 14.

We prove claim (d) for \mathcal{L} -classes (for \mathcal{R} -classes the proof is similar). The case $|S| = 1$ cannot occur by our assumptions that S has no zero element and $S \not\cong \mathbf{0}$. Consider the case $|S| > 1$. Then for any different $a, b \in S$ we have $S^1 * a * S^1 = S^1 * b * S^1 \ni b$, which implies $S * S \neq \emptyset$, in particular, there exist $s, t \in S$ such that $s * t \neq \emptyset$. Set $H := \mathcal{L}_s \cap \mathcal{R}_t$. Claim (c) implies that H is a hypergroup \mathcal{H} -class in \mathcal{L}_s . Let L be an \mathcal{L} -class different from \mathcal{L}_s , $x \in L \cap \mathcal{R}_t$ and $y \in H$. Then $y \neq x$ and $y \mathcal{R} x$ and hence there is $u \in S$ such that $y \in x * u$. Consider $H' = S^1 * x \cap u * S^1$, which is an \mathcal{H} -class by Lemma 10. Proposition 14 along with claim (b) imply that H' is a hypergroup. Claim (d) follows.

In the setup of claim (e) we have $\mathcal{L}_a * s \neq \emptyset$ and hence $\mathcal{L}_a * s$ is a left ideal of S contained in $\mathcal{L}_s = \mathcal{L}_b$. Then $\mathcal{L}_a * s = \mathcal{L}_b$ by minimality of \mathcal{L}_b . If $c \in \mathcal{L}_b$ is such that $c \in x * s$, then $c \mathcal{R} x$ and hence the map preserves \mathcal{R} -classes and, consequently, \mathcal{H} -classes. This proves claim (e).

To prove claim (f) we note that $J * I \neq \emptyset$ by Lemma 10 and $J * I \subset I \cap J$. Let $a \in J$ and $b \in I$ be such that $a * b \neq \emptyset$. Then claim (e) implies that $\mathcal{H}_a * b = I \cap J$. This proves claim (f).

Finally, claim (g) follows from Proposition 15 and claims (c) and (d). \square

5.3. Example: a finite simple multisemigroup with nontrivial inclusions of principal one-sided ideals. Let (X, \leq) be a partially ordered set possessing a unique minimal element y . Consider the rectangular band $T = X \times X$. For $a \in X$ let $a^\downarrow = \{x \in X : x \leq a\}$ be the *downward closure* of a . We define the function $f : T \rightarrow 2^T$ by setting

$$f(a, b) = a^\downarrow \times b^\downarrow. \quad (5.1)$$

It is easy to check that the function f satisfies the equality $f(f(x)f(y)) = f(x)f(y)$ and therefore $S = (T, *)$ with $(a, b) * (c, d) = f(a, b)f(c, d)$ is a multisetsemigroup (see Subsection 3.14). Moreover, one checks that S is a simple multisetsemigroup and $Q(S) = \{(y, y)\}$. Clearly $Q(S)$ is a strongly simple multisetsemigroup. Note that $(a, b) * S^1$ is a minimal right ideal if and only if $a = y$; $S^1 * (a, b)$ is a minimal left ideal if and only if $b = y$. Let $a_1 < \dots < a_m$ be a chain in X and $c \in X$ be any element. Then we have a chain of principal left ideals

$$S^1 * (c, a_1) \subsetneq S^1 * (c, a_2) \subsetneq \dots \subsetneq S^1 * (c, a_m). \quad (5.2)$$

The above example can be generalized as follows. Let I be a set and let (X_i, \leq_i) , $i \in I$, be partially ordered sets each possessing a unique minimal element y_i , $i \in I$. We assume that the sets X_i are pairwise disjoint. Let $Y = \{y_i : i \in I\}$. Let $X = \bigcup_{i \in I} X_i$. We define on X the partial order \leq by setting $x \leq y$ provided that there is i such that $x, y \in X_i$ and $x \leq_i y$. Consider the rectangular band $T = X \times X$ and define $f : T \rightarrow 2^T$ as given in (5.1).

The map f gives a simple multisetsemigroup $S = (T, *)$ according to Subsection 3.14. The submultisetsemigroup $Q(S)$ of S is equal to $Y \times Y$ and is a rectangular band, hence is a strongly simple multisetsemigroup (in fact, a semigroup); $(a, b) * S^1$ is a minimal right ideal if and only if $a \in Y$; $S^1 * (a, b)$ is a minimal left ideal if and only if $b \in Y$. Let $a_1 < \dots < a_m$ be a chain in X . Let $c \in Z$ be any element. Then (5.2) gives a chain of left ideals.

5.4. Example: a finite simple multisetsemigroup whose support is not simple. Let $X = \{1, 2\}$ with the order given by $1 < 2$. Let $S = X \times X$. Let $(S, *) = S(X)$ be the multisetsemigroup constructed in Subsection 5.3. That is, the multiplication $*$ on S is given by the map $* : S \times S \rightarrow 2^S$:

$$(i, j) * (k, l) = i^\downarrow \times l^\downarrow.$$

Let $T = S \cup (1', 1')$ (here $1'$ is an element different from both 1 and 2). Define the map $\pi : T \rightarrow S$ by setting $\pi(a) = a$, if $a \in S$, and $\pi((1', 1')) = (1, 1)$. Let $\circ : T \times T \rightarrow 2^T$ be defined as follows:

$$x \circ y = \begin{cases} \pi(x) * \pi(y), & x \in \{(1, 1), (1', 1'), (1, 2)\} \text{ and} \\ & y \in \{(1, 1), (1', 1'), (2, 1)\}; \\ (\pi(x) * \pi(y)) \cup (1', 1'), & \text{otherwise.} \end{cases}$$

Lemma 17. (T, \circ) is a multisetsemigroup.

Proof. Observe that we have $(1, 1) \circ x = (1', 1') \circ x$ and $x \circ (1, 1) = x \circ (1', 1')$ for any $x \in T$.

Let $x, y, z \in T$. We have to verify that $(x \circ y) \circ z = x \circ (y \circ z)$. Note that $(1, 1) \in s \circ t$ for any $s, t \in T$. It follows that $(x \circ y) \circ z = (\pi(x) * \pi(y)) \circ z$ and, similarly, $x \circ (y \circ z) = x \circ (\pi(y) * \pi(z))$.

By the definition, $(\pi(x) * \pi(y)) \circ z$ equals either $(\pi(x) * \pi(y)) * \pi(z)$ or $((\pi(x) * \pi(y)) * \pi(z)) \cup \{(1', 1')\}$. Let us establish when it equals $(\pi(x) * \pi(y)) * \pi(z)$. From the definition we have that this happens if and only if $z \in \{(1, 1), (1', 1'), (2, 1)\}$ and $\pi(x) * \pi(y) \subseteq \{(1, 1), (1, 2)\}$. The latter inclusion holds if and only if $x \in \{(1, 1), (1', 1'), (1, 2)\}$.

Similarly, $x \circ (\pi(y) * \pi(z))$ equals either $\pi(x) * (\pi(y) * \pi(z))$ or $(\pi(x) * (\pi(y) * \pi(z))) \cup \{(1', 1')\}$ and one shows that $x \circ (\pi(y) * \pi(z)) = \pi(x) * (\pi(y) * \pi(z))$ if and only if $x \in \{(1, 1), (1', 1'), (1, 2)\}$ and $z \in \{(1, 1), (1', 1'), (2, 1)\}$. The equality $(x \circ y) \circ z = x \circ (y \circ z)$ follows. \square

It is easy to check that T is a simple multisetsemigroup. For $x \in T$ we have $x \circ T^1 = (\pi(x) * S^1) \cup \{(1', 1')\}$, $T^1 \circ x = (S^1 * \pi(x)) \cup \{(1', 1')\}$. Furthermore, $Q(T) = \{(1, 1), (1', 1')\}$, $(1, 1) \circ T^1 \cap T^1 \circ (1, 1) = \{(1, 1), (1', 1')\}$. Both $(1, 1)$ and $(1', 1')$ belong to the same \mathcal{H} -class of T . At the same, $Q(T) \circ Q(T) = \{(1, 1)\}$ and hence $Q(T)$ is not a hypergroup. Clearly, $(1, 1)$ and $(1', 1')$ are not in the same \mathcal{J} -class of $Q(T)$ and so $Q(T)$ is not simple.

5.5. Example: a finite simple multisetsemigroup for which $\mathcal{R} \circ \mathcal{L} \neq \mathcal{L} \circ \mathcal{R}$. Let $X = \{1, 2\}$ and $S = X \times X$. Let $(S, *) = S(X)$ be the multisetsemigroup from Subsection 5.4. We have $(i, j) * (k, l) * (p, q) = i^\perp \times q^\perp$.

Let $T = S \setminus \{(2, 2)\}$. We define a map $T \times T \rightarrow 2^T$, $(x, y) \mapsto x \bullet y$, by

$$(i, j) \bullet (k, l) = ((i, j) * (k, l)) \setminus \{(2, 2)\}.$$

Lemma 18. (T, \bullet) is a multisetsemigroup.

Proof. Let $(i, j), (k, l), (p, q) \in T$. Observe that if $(i, j) * (k, l) \ni (2, 2)$, then $(i, j) * (k, l) \ni (2, 1)$ and hence $((i, j) * (k, l)) * (p, q) = (((i, j) * (k, l)) \setminus \{(2, 2)\}) * (p, q)$. Using this we calculate

$$\begin{aligned} ((i, j) \bullet (k, l)) \bullet (p, q) &= (((i, j) * (k, l)) \setminus \{(2, 2)\}) * (p, q) \setminus \{(2, 2)\} \\ &= (((i, j) * (k, l)) * (p, q)) \setminus \{(2, 2)\}. \end{aligned}$$

Similarly we show that

$$(i, j) \bullet ((k, l) \bullet (p, q)) = ((i, j) * ((k, l) * (p, q))) \setminus \{(2, 2)\}.$$

Associativity of \bullet follows. \square

It is easy to see that T is simple and $Q(T) = \{(1, 1)\}$. Further, it is easy to calculate Green's relations on T . We have $\mathcal{R}_{(1,1)} = \{(1, 1), (1, 2)\}$ and $\mathcal{R}_{(2,1)} = \{(2, 1)\}$. Also $\mathcal{L}_{(1,1)} = \{(1, 1), (2, 1)\}$ and $\mathcal{L}_{(1,2)} = \{(1, 2)\}$. We see that $(1, 2) \mathcal{R} (1, 1) \mathcal{L} (2, 1)$ so that $((1, 2), (2, 1)) \in \mathcal{R} \circ \mathcal{L}$. But since $\mathcal{L}_{(1,2)} \cap \mathcal{R}_{(2,1)} = \emptyset$, we have $((1, 2), (2, 1)) \notin \mathcal{L} \circ \mathcal{R}$. In particular, $\mathcal{L} \circ \mathcal{R} \neq \mathcal{R} \circ \mathcal{L}$. Note that we always have $\mathcal{L} \circ \mathcal{R} = \mathcal{R} \circ \mathcal{L}$ for semigroups.

5.6. Simple multisetsemigroups with identity. A simple finite semigroup with identity is a group. For multisetsemigroups the situation is much more complicated.

Lemma 19. *Let $(S, *)$ be a strongly simple multisetsemigroup with identity. Then S is a hypergroup.*

Proof. Since S is strongly simple, $S = S * 1$ is a minimal left ideal and $S = 1 * S$ is a minimal right ideal. Hence $S * a = S$ and $a * S = S$ for any $a \in S$ and thus S is a hypergroup. □

One could expect that even a simple finite multisetsemigroup with identity should be a hypergroup. Unfortunately, this is not the case. Indeed, consider $S = \{1, a, b, t\}$ with the multioperation $*$ defined by the following Cayley table:

| | | | | | |
|-----|--|--|--|--|--|
| $*$ | | | | | |
| $*$ | | | | | |
| 1 | | | | | |
| a | | | | | |
| b | | | | | |
| t | | | | | |

It is straightforward to verify that $(S, *)$ is a multisetsemigroup. This multisetsemigroup is simple; it contains an identity element; it consists of idempotents; it has a unique minimal left ideal, namely $\{a, t\}$; it has a unique minimal right ideal, namely $\{b, t\}$; we have $Q(S) = \{t\}$ and S is not a hypergroup.

Below we collect some properties of simple multisetsemigroups of finite type with identity:

Proposition 20. *Let $(S, *)$ be a simple multisetsemigroup of finite type with identity.*

- (a) *If I is a non-empty left ideal of S and J is a non-empty right ideal of S , then $I \cap \mathcal{R}_1 \neq \emptyset$ and $J \cap \mathcal{L}_1 \neq \emptyset$.*
- (b) *If I and I' are two non-empty left ideals of S , then $I * I' \neq \emptyset$.*
- (c) *If J and J' are two non-empty right ideals of S , then $J * J' \neq \emptyset$.*

Proof. Since S contains the identity, it contains a unique (possibly empty) maximal proper right ideal (the union of all right ideals of S which do not contain the identity), call it \mathcal{K} . Clearly, $\mathcal{R}_1 = S \setminus \mathcal{K}$.

Let I be a non-empty left ideal of S . Then $X := I * S$ is a non-empty two-sided ideal of S since S contains the identity and hence $X = S$ since S is simple. If $I \subset \mathcal{K}$, then $X \subset \mathcal{K}$ and hence X cannot be equal to S , a contradiction. Therefore I intersects $S \setminus \mathcal{K} = \mathcal{R}_1$ non-trivially. Similarly one shows that every non-empty right ideal intersects \mathcal{L}_1 non-trivially, which proves claim (a).

Let I and I' be two non-empty left ideals in of S . Then from the previous paragraph we have: $(I * I') * S = I * (I' * S) = I * S = S$, thus $I * I' \neq \emptyset$. This proves claim (b). Claim (c) is proved similarly to claim (b). \square

5.7. A Kazhdan-Lusztig example in type B. In Subsection 5.2 we established many elementary properties of strongly simple multisemigroups that are similar to properties of 0-bisimple semigroups. One could observe that the multisemigroup version of Green's lemma does not assert that $c \mapsto c * x$ is a *bijection* from H to $H * x$. This turns out to be false for multisemigroups in general. Here we give an explicit example.

Let $W = \{e, s, t, st, ts, sts, tst, stst\}$ be a Weyl group of type B_2 (the generators s and t satisfy $s^2 = t^2 = e$ and $stst = tsts$) and $(W, *)$ be the corresponding Kazhdan-Lusztig multisemigroup as explained in Subsection 3.9. Denote by T the Rees quotient of the submultisemigroup $W \setminus \{e\}$ by the zero element $\{stst\}$. Then a direct computation shows that T is a strongly simple multisemigroup with the following Cayley table:

| * | s | t | st | ts | sts | tst |
|-------|--------------|--------------|--------------|--------------|--------------|--------------|
| s | s | st | st | $\{s, sts\}$ | sts | st |
| t | ts | t | $\{t, tst\}$ | ts | ts | tst |
| st | $\{s, sts\}$ | st | st | $\{s, sts\}$ | $\{s, sts\}$ | st |
| ts | ts | $\{t, tst\}$ | $\{t, tst\}$ | ts | ts | $\{t, tst\}$ |
| sts | sts | st | st | $\{s, sts\}$ | s | st |
| tst | ts | tst | $\{t, tst\}$ | ts | ts | t |

It follows that T has the following egg-box diagram (in which all \mathcal{H} -classes are hypergroups):

| | \mathcal{L}_s | \mathcal{L}_t |
|-----------------|-----------------|-----------------|
| \mathcal{R}_s | $\{s, sts\}$ | $\{st\}$ |
| \mathcal{R}_t | $\{ts\}$ | $\{t, tst\}$. |

6. The number of multisetsemigroups

In this section we compare asymptotic properties of associativity for ordinary binary operations and for multivalued binary operations.

6.1. The number of semigroups. Let n be a positive integer and $N_n := \{1, 2, \dots, n\}$. A binary operation on N_n corresponds to the choice of a square $n \times n$ matrix with coefficients in N_n . Let $\text{Mat}_{n \times n}(N_n)$ be the set of all such matrices. Clearly, $|\text{Mat}_{n \times n}(N_n)| = n^{n^2}$. Let X denote the subset of $\text{Mat}_{n \times n}(N_n)$ which consists of all matrices corresponding to *associative* binary operation. By [KRS], Equation (3.6), we have

$$|X| = \left(\frac{n}{(2e + o(1)) \ln n} \right)^{n^2}$$

and hence, dividing by n^{n^2} , we obtain the following corollary, which says that “almost all” binary operations are not associative:

Corollary 21. *We have $\frac{|X|}{n^{n^2}} \rightarrow 0$ when $n \rightarrow \infty$.*

6.2. The number of multisetsemigroups. A multivalued binary operation on N_n corresponds to the choice of a square $n \times n$ matrix with coefficients in 2^{N_n} . Let $\text{Mat}_{n \times n}(2^{N_n})$ be the set of all such matrices. Clearly, $|\text{Mat}_{n \times n}(2^{N_n})| = 2^{n^3}$. Let Y denote the subset of $\text{Mat}_{n \times n}(2^{N_n})$ which consists of all matrices corresponding to *associative* multivalued binary operation (i.e. those satisfying (1.1)). Let Y' denote the subset of $\text{Mat}_{n \times n}(2^{N_n})$ which consists of all matrices corresponding to multivalued binary operation defining a hypergroup. The following claim, which says that “almost all” multivalued binary operations are associative, is in striking contrast with Corollary 21:

Theorem 22. *We have both $\frac{|Y|}{2^{n^3}} \rightarrow 1$ and $\frac{|Y'|}{2^{n^3}} \rightarrow 1$ when $n \rightarrow \infty$.*

Proof. Denote by Z the subset of $\text{Mat}_{n \times n}(2^{N_n})$ which consists of all matrices such that the corresponding multivalued binary operation $*$ for every $a, b, c \in N_n$ satisfies

$$\bigcup_{s \in a * b} s * c = \bigcup_{t \in b * c} a * t = N_n.$$

Then $Z \subset Y' \subset Y$ and hence it is enough to show that $\frac{|Z|}{2^{n^3}} \rightarrow 1$ when $n \rightarrow \infty$.

It is enough to show that for the subset U of $\text{Mat}_{n \times n}(2^{N_n})$ which consists of all matrices such that the corresponding multivalued binary operation $*$ for every $a, b, c \in N_n$ satisfies

$$\bigcup_{t \in b * c} a * t = N_n, \tag{6.1}$$

we have $\frac{|U|}{2^{n^3}} \rightarrow 1$ when $n \rightarrow \infty$. Indeed, the latter, by symmetry, also implies that $\frac{|U'|}{2^{n^3}} \rightarrow 1$ for $n \rightarrow \infty$, where U' denotes the subset of $\text{Mat}_{n \times n}(2^{N_n})$ corresponding to all operations $*$ satisfying

$$\bigcup_{s \in a * b} s * c = N_n$$

for all $a, b, c \in N_n$, so that we have $\frac{|U \cap U'|}{2^{n^3}} \rightarrow 1$ for $n \rightarrow \infty$.

Given a multivalued binary operation $*$ on N_n , with each $a \in N_n$ we can associate, similarly to Subsection 4.7, a binary relation τ_a on N_n defined as follows: $i \tau_a j$ if and only if $i \in a * j$. Let \mathbf{f} denote the full relation on N_n . Then (6.1) is equivalent to the fact that $\tau_a \tau_b = \mathbf{f}$ for all $a, b \in N_n$.

To prove the latter claim we can adopt the classical argument from [KR], Theorem 4. Our choice of an element $M = (M_{i,j})$ of $\text{Mat}_{n \times n}(2^{N_n})$ corresponds to a random choice of n binary relations σ_a , $a \in N_n$, on N_n (here $i \sigma_a j$ if and only if $i \in M_{a,j}$). We claim that the probability of the random event that the product of any two of these n relations equals the full relation tends to 1 when n tends to infinity ([KR], Theorem 4 claims this just for two instead of n random binary relations). Indeed, the probability that the product of two random elements of the two-element Boolean algebra is zero equals $\frac{3}{4}$. This implies that the probability that the element in a fixed entry in a product of two random Boolean $n \times n$ matrices is zero is at most $(\frac{3}{4})^n$. We have n random elements σ_a , $a \in N_n$. We can form n^2 pairs (σ_a, σ_b) , $a, b \in N_n$, from these elements. The Boolean $n \times n$ matrix corresponding to the product $\sigma_a \sigma_b$ has n^2 entries, each of which is zero with probability at most $(\frac{3}{4})^n$. Hence the probability that at least one of these entries in at least one of the products of the form $\sigma_a \sigma_b$ is zero is at most $n^4 (\frac{3}{4})^n$. Since $n^4 (\frac{3}{4})^n \rightarrow 0$ for $n \rightarrow \infty$, the claim follows. \square

7. Nilpotent multisemigroups

In this section we establish some basic fact about nilpotent multisemigroups (see e.g. [GM1], [GM2] for more advanced semigroup analogues).

7.1. Nilpotent multisemigroups and their characterization. Let $(S, *)$ be a multisemigroup. As usual, for an element $s \in S$ and $k \in \mathbb{N}$ we write s^k for the product $s * s * \dots * s$ of length k . Similarly, for $X \subset S$ we write X^k for the union of all $x_1 * x_2 * \dots * x_k$, where $x_i \in X$ for all i . An element $s \in S$ is said to be *nilpotent* provided that $s^k = \emptyset$ for some k . The multisemigroup S is called *nilpotent* provided that $S^k = \emptyset$ for some k . The minimal such k is called the *nilpotency degree* of S . Similarly, a subset $X \subset S$ is called *nilpotent* provided that $X^k = \emptyset$ for some k (in particular, the empty set is nilpotent). The notion of nilpotent

multisetsemigroups generalizes that of nilpotent semigroups. Nilpotent semigroups correspond, via the construction given at the end of Subsection 4.3, to nilpotent quasi-semigroups.

For a multisetsemigroup $(S, *)$ define the *action digraph* $\Gamma = \Gamma_S$ as follows: the set of vertices of Γ is S and for $s, t \in S$ (not necessarily different) we have an oriented edge $s \rightarrow t$ if and only if there exists $a \in S$ such that $t \in a * s$.

Proposition 23. *Let $(S, *)$ be a multisetsemigroup.*

- (a) *The multisetsemigroup $(S, *)$ is nilpotent if and only if there is $m \in \mathbb{N}$ such that the length of any directed path in Γ is smaller than m .*
- (b) *If $(S, *)$ is nilpotent, then the nilpotency degree of S is exactly the length of the longest directed path in Γ minus two.*

Proof. Let $(S, *)$ be nilpotent of nilpotency degree k and $s_0 \rightarrow s_1 \rightarrow \dots \rightarrow s_m$ be an oriented path in Γ of length m . Then there exist a_i such that $s_i \in a_i * s_{i-1}$, $i = 1, 2, \dots, m$. This means that $a_m * a_{m-1} * \dots * a_1 * s_0 \neq \emptyset$ and hence $m + 1 < k$.

On the other hand, if the length of any oriented path in Γ is at most m , then for any $m + 2$ elements $s_1, \dots, s_{m+2} \in S$ we have $s_{m+2} * s_{m+1} * \dots * s_1 = \emptyset$ and hence S is nilpotent of nilpotency degree at most $m + 2$. This implies both claims of the proposition. \square

Corollary 24. *Let $(S, *)$ be a nilpotent multisetsemigroup. Then all Green's relations on S coincide with the equality relation.*

Proof. This follows directly from the fact that Γ does not have any oriented cycles. The latter is a direct consequence of Proposition 23. \square

7.2. Finite nilpotent multisetsemigroups. Similarly to the case of finite nilpotent semigroups (see [Ar], Chapter 7, Fact 2.30), for finite nilpotent multisetsemigroups we have:

Proposition 25. *A finite multisetsemigroup $(S, *)$ is nilpotent if and only if every element of S is nilpotent.*

Proof. The “only if” part is obvious, so we prove the “if” part. Assume that every element of S is nilpotent. We claim that the digraph Γ does not have oriented cycles (in particular, loops). Indeed, assume that this is not the case and let $s_1 \rightarrow s_2 \rightarrow \dots \rightarrow s_k \rightarrow s_{k+1} = s_1$ be an oriented cycle in Γ . For $i = 1, 2, \dots, k$ let $a_i \in S$ be such that $s_{i+1} \in a_i * s_i$. Then there exists $t \in a_k * a_{k-1} * \dots * a_1$ such that $s_1 \in t * s_1$. This implies that $s_1 \in t^n * s_1$ for any $n \in \mathbb{N}$ and hence t is not nilpotent, a contradiction.

As S is finite and Γ does not have oriented cycles, then the length of any oriented path in Γ is strictly smaller than $|S|$. This means that S is nilpotent by Proposition 23. \square

Finite nilpotent semigroups can be characterized as finite semigroups with unique idempotent which, moreover, is the zero element. Equivalently, a finite semigroup (S, \cdot) is nilpotent if and only if for any non-singleton subsemigroup T of S we have $|T \cdot T| < |T|$. The following gives an analogue of the latter characterization of nilpotency for multisemigroups.

Theorem 26. *A finite multisemigroup $(S, *)$ is nilpotent if and only if for any (non-empty) submultisemigroup T of S we have $|T * T| < |T|$ (or, equivalently, $T * T \neq T$).*

Proof. If S is nilpotent and T is a submultisemigroup of S , then T is nilpotent as well. However, $T * T = T$ implies $T^k = T$ for all k which contradicts nilpotency of T . Therefore $T * T \neq T$.

If S is not nilpotent, we can use the fact that S is finite and choose some non-nilpotent submultisemigroup T of S which is minimal with respect to inclusion. We claim that $T * T = T$. If this were not the case, then $U := T * T \neq \emptyset$ would be a submultisemigroup of S properly contained in T . Hence U must be nilpotent by minimality of T . Therefore $U^k = \emptyset$ for some k , which implies that $(T * T)^k = T^{2k} = \emptyset$, a contradiction. \square

7.3. The radical. Let $(S, *)$ be a multisemigroup. Following [GM1], by the radical $R(S)$ of S we will mean the set

$$R(S) := \{s \in S \mid S^1 * s * S^1 \text{ is nilpotent}\}.$$

Then $R(S)$, which might be empty, is a union of two-sided ideals of S and hence is a two-sided ideal of S .

Lemma 27. *Let S be a finite multisemigroup.*

- (a) *The set $R(S)$ is the maximal (with respect to inclusion) nilpotent two-sided ideal of S .*
- (b) *The multisemigroup S is nilpotent if and only if $S = R(S)$.*

Proof. For $s \in R(S)$ we have that $S^1 * s * S^1$ is nilpotent, in particular, s is nilpotent. Hence every element of $R(S)$ is nilpotent and thus the fact that $R(S)$ is nilpotent follows from Proposition 25 and the fact that $R(S)$ is finite (since S is finite).

On the other hand, if I is a nilpotent two-sided ideal of S and $s \in I$, then $S^1 * s * S^1 \subset I$ is nilpotent and hence $s \in R(S)$. Claim (a) follows. Claim (b) follows from claim (a). \square

Our next observation is the following:

Proposition 28. *Let S be a finite multisetsemigroup and T a nilpotent submultisetsemigroup of S .*

- (a) *The set $T \cup R(S)$ is a nilpotent submultisetsemigroup of S .*
- (b) *If T is maximal with respect to inclusion, then $R(S) \subset T$.*

Proof. That $T \cup R(S)$ is submultisetsemigroup follows directly from the facts that T is a submultisetsemigroup and $R(S)$ is a two-sided ideal of S . That $T \cup R(S)$ is nilpotent follows from Proposition 25. This proves claim (a). Claim (b) follows from claim (a). \square

Corollary 29. *Let S be a finite multisetsemigroup.*

- (a) *If $R(S)$ is a maximal nilpotent submultisetsemigroup of S , in particular, if $S = R(S)$, then $R(S)$ is the unique maximal nilpotent submultisetsemigroup of S .*
- (b) *If $R(S)$ is not a maximal nilpotent submultisetsemigroup of S , then the map $X \mapsto X \setminus R(S)$ is a bijection from the set of all maximal nilpotent submultisetsemigroups of S to the set of all maximal nilpotent submultisetsemigroups of the Rees quotient $S \setminus R(S)$.*

Proof. Claim (a) follows directly from Proposition 28(b).

To prove claim (b), let X be a maximal nilpotent submultisetsemigroup of S . Then $X \neq R(S)$ and $X \setminus R(S)$ is a nilpotent submultisetsemigroup of $S \setminus R(S)$. Assume Y is a nilpotent submultisetsemigroup of $S \setminus R(S)$ containing $X \setminus R(S)$. Then from Lemma 27 it follows that $Y \cup R(S)$ is a nilpotent submultisetsemigroup of S . Therefore $Y = X \setminus R(S)$ by maximality of X , that is $X \setminus R(S)$ is maximal.

Conversely, let Y be a maximal nilpotent submultisetsemigroup of $S \setminus R(S)$. Then, similarly to the above, $Y \cup R(S)$ is a nilpotent submultisetsemigroup of S . If X is a nilpotent submultisetsemigroup of S containing $Y \cup R(S)$, then $X \setminus R(S)$ is a nilpotent submultisetsemigroup of $S \setminus R(S)$ containing Y . Hence $X = Y \cup R(S)$ by maximality of Y . This proves claim (b). \square

7.4. Maximal nilpotent submultisetsemigroups of strongly simple multisetsemigroups. Let $(S, *)$ be a strongly simple multisetsemigroup which is not isomorphic to the singleton multisetsemigroup $\mathbf{0}$. Recall that \mathcal{H} is a congruence on S and that S/\mathcal{H} is a quasi-semigroup.

Proposition 30. *The canonical strong surjective homomorphism $\varphi : S \rightarrow S/\mathcal{H}$ induces a bijection between the set of maximal (with respect to inclusion) nilpotent submultisemigroups of S and maximal nilpotent sub-quasi-semigroups of S/\mathcal{H} . This bijection preserves nilpotency degree.*

Proof. From Lemma 12 it follows that for $a, b \in S$ we have $a * b \neq \emptyset$ if and only if $\mathcal{H}_a * \mathcal{H}_b \neq \emptyset$. Furthermore, from Proposition 14 it also follows that $\mathcal{H}_a * \mathcal{H}_b$, when non-empty, is contained in a single \mathcal{H} -class of S , namely $\mathcal{R}_a \cap \mathcal{L}_b$. Hence, by induction, for $a_1, a_2, \dots, a_k \in S$ we have $a_1 * a_2 * \dots * a_k \neq \emptyset$ if and only if $\mathcal{H}_{a_1} * \mathcal{H}_{a_2} * \dots * \mathcal{H}_{a_k} \neq \emptyset$, moreover, $\mathcal{H}_{a_1} * \mathcal{H}_{a_2} * \dots * \mathcal{H}_{a_k}$, when non-empty, is contained in a single \mathcal{H} -class of S .

Consequently, if T is a nilpotent submultisemigroup of S , then $T' := \bigcup_{I \in T} \mathcal{H}_I$ is a nilpotent submultisemigroup of S of the same nilpotency degree, moreover, $T \subset T'$. Hence any maximal nilpotent submultisemigroup of S is a union of \mathcal{H} -classes of S .

If T is a nilpotent submultisemigroup of S , then $\varphi(T)$ is a nilpotent sub-quasi-semigroup of S/\mathcal{H} (since φ is a homomorphism), moreover, the above implies that $\varphi(T)$ has the same nilpotency degree as T . Conversely, if K is a nilpotent sub-quasi-semigroup of S/\mathcal{H} , then the above observations imply that $\varphi^{-1}(K)$ is a nilpotent submultisemigroup of S of the same nilpotency degree as K . Note also that $\varphi^{-1}(K)$ is a union of \mathcal{H} -classes of S by construction. We have $\varphi^{-1}(\varphi(T)) = T'$ in the above notation and also we have $\varphi(\varphi^{-1}(K)) = K$, in particular, φ^{-1} maps maximal nilpotent sub-quasi-semigroups of S/\mathcal{H} to maximal nilpotent submultisemigroups of S . The above also implies that if T is a maximal nilpotent submultisemigroup of S , then $\varphi(T)$ is a maximal nilpotent sub-quasi-semigroup of S/\mathcal{H} . Thus the first claim of the proposition follows. The second claim follows from the above observations that both φ and φ^{-1} preserve nilpotency degree. \square

Let $(S, *)$ be a finite strongly simple multisemigroup, I_1, \dots, I_m —the list of all \mathcal{L} -classes, and J_1, \dots, J_n —the list of all \mathcal{R} -classes in S . Let $\mathbf{I}(S)$ denote the Boolean matrix (a_{ij}) , $i = 1, \dots, m$, $j = 1, \dots, n$, defined as follows:

$$a_{ij} = \begin{cases} 1, & I_i \cap J_j \text{ is a hypergroup;} \\ 0, & \text{otherwise.} \end{cases}$$

The matrix $\mathbf{I}(S)$ is called the *incidence matrix* of S . Recall that $S \not\cong \mathbf{0}$. Then, by Theorem 16, all rows and columns of S are non-zero. Assume that $\mathbf{I}(S)$ has the form

$$\left(\begin{array}{c|c} E_k & B \\ \hline A & A \cdot B \end{array} \right)$$

for some positive integer k , where E_k is the identity matrix and A and B are Boolean matrices such that each row of A is not zero and each column of B is not zero, and $A \cdot B$ is the Boolean matrix which is the product of A and B . In this case a classification of maximal nilpotent sub-quasi-semigroups of S/\mathcal{H} can be found in [GM1], Subsection 6.10. In particular, there are exactly $k!$ such sub-quasi-semigroups and each of them has nilpotency degree k .

References

- [Ar] M. Arbib (Ed.). Algebraic Theory of Machines, Languages, and Semigroups. With a Major Contribution by Kenneth Krohn and John L. Rhodes, Academic Press, New York/London, 1968.
- [Ca] H. Campaigne. Partition hypergroups. *Amer. J. Math.* **62** (1940), no. 1, 599–612.
- [Da] B. Davvaz. Some results on congruences on semihypergroups. *Bull. Malays. Math. Sci. Soc. (2)* **23** (2000), no. 1, 53–58.
- [Di] J. Dixmier. Enveloping algebras. Graduate Studies in Mathematics, **11**. American Mathematical Society, Providence, RI, 1996.
- [DO] M. Dresher, O. Ore. Theory of Multigroups. *Amer. J. Math.* **60** (1938), no. 3, 705–733.
- [Ea] J. E. Eaton. Associative multiplicative systems. *Amer. J. Math.* **62** (1940), no. 1, 222–232.
- [GM1] O. Ganyushkin, V. Mazorchuk. On classification of maximal nilpotent subsemigroups. *J. Algebra* **320** (2008), no. 8, 3081–3103.
- [GM2] O. Ganyushkin, V. Mazorchuk. Classical finite transformation semigroups, An introduction. Algebra and Applications, Vol. **9**, Springer-Verlag London, Ltd., London, 2009.
- [Gr] J. Green. On the structure of semigroups. *Ann. of Math. (2)* **54** (1951), 163–172.
- [Gu] J. Gunawardena. An introduction to idempotency. *Idempotency (Bristol, 1994)*, 1–49, Publ. Newton Inst., **11**, Cambridge Univ. Press, Cambridge, 1998.
- [Ha] A. Hasankhani. Ideals in a semihypergroup and Green’s relations. *Hyperstructures and their applications in criptography, geometry and uncertainty treatment (Pesara, 1995)*. *Ratio Math. No.* **13** (1999), 29–36.
- [JSM] H. Jafarabadi, N. Sarmin, M. Molaei. Completely simple and regular semi hypergroups; *Bull. Malays. Math. Sci. Soc. (2)* **35** (2012), no. 2, 335–343.
- [KL] D. Kazhdan, G. Lusztig. Representations of Coxeter groups and Hecke algebras. *Invent. Math.* **53** (1979), no. 2, 165–184.
- [KR] K. Kim, F. Roush. Two-generator semigroups of binary relations. *J. Math. Psych.* **17** (1978), no. 3, 236–246.
- [KRS] D. Kleitman, B. Rothschild, J. Spencer. The number of semigroups of order n . *Proc. Amer. Math. Soc.* **55** (1976), no. 1, 227–232.

- [Ko] M. Koskas. Structures algébriques multivoques. Applications. Bull. Soc. Math. France Suppl. Mém. **23** (1970), 84 pp.
- [Kaa] J. Kåhrström. Tensoring with infinite-dimensional modules in \mathcal{O} . Algebr. Represent. Theory **13** (2010), no. 5, 561–587.
- [Mar] F. Marty. Sur une généralisation de la notion de groupe. Actes du 8^{me} Congrès des Mathématiciens Scandinaves, Stockholm, 1934, 45–49.
- [Maz1] V. Mazorchuk. Lectures on algebraic categorification. QGM Master Class Series. European Mathematical Society (EMS), Zürich, 2012.
- [Maz2] V. Mazorchuk. Lectures on $\mathfrak{sl}_2(\mathbb{C})$ -modules. Imperial College Press, London, 2010.
- [MM1] V. Mazorchuk, V. Miemietz. Cell 2-representations of finitary 2-categories. Compos. Math. **147** (2011), no. 5, 1519–1545.
- [MM2] V. Mazorchuk, V. Miemietz. Additive versus abelian 2-representations of fiat 2-categories. Mosc. Math. J. **14** (2014), no. 3, 595–615.
- [Re] P. Resende. Lectures on étale groupoids, inverse semigroups and quantales. Preprint available from: www.math.ist.utl.pt
- [Ro] K. Rosenthal. Quantales and their applications. Pitman Res. Notes Math. Ser., vol. **234**, Longman Scientific & Technical, 1990.
- [Tr] T. Trimble. Hypermonoid. <http://ncatlab.org/nlab/show/hypermonoid>
- [Vi] O. Viro. Hyperfields for Tropical Geometry I. Hyperfields and dequantization. Preprint arXiv:1006.3034.
- [Vo] T. Vougiouklis. Representations of hypergroups by generalized permutations. Algebra Universalis **29** (1992), no. 2, 172–183.
- [Wa] H. S. Wall. Hypergroups. Amer. J. Math. **59** (1937), no. 1, 77–98.

Received August 21, 2012; revised July 3, 2014

G. Kudryavtseva, Institute Jožef Stefan, Jamova cesta 39, 1000 Ljubljana, Slovenia

E-mail: ganna.kudryavtseva@ijs.si

V. Mazorchuk, Department of Mathematics, Uppsala University, SE 471 06, Uppsala, Sweden

E-mail: mazor@math.uu.se