

Failure of the local chain rule for the fractional variation

Giovanni E. Comi and Giorgio Stefani

Abstract. We prove that the local version of the chain rule cannot hold for the fractional variation defined in our previous article (2019). In the case $n = 1$, we prove a stronger result, exhibiting a function $f \in BV^\alpha(\mathbb{R})$ such that $|f| \notin BV^\alpha(\mathbb{R})$. The failure of the local chain rule is a consequence of some surprising rigidity properties for non-negative functions with bounded fractional variation which, in turn, are derived from a fractional Hardy inequality localized to half-spaces. Our approach exploits the distributional techniques developed in our previous works (2019–2022). As a byproduct, we refine the fractional Hardy inequality obtained in works of Shieh and Spector (2018) and Spector (2020) and we prove a fractional version of the closely related Meyers–Ziemer trace inequality.

1. Introduction

1.1. The fractional variation

Let $\alpha \in (0, 1)$. The *fractional α -gradient* of a function $f \in \text{Lip}_c(\mathbb{R}^n)$ is defined as

$$\nabla^\alpha f(x) = \mu_{n,\alpha} \int_{\mathbb{R}^n} \frac{(y-x)(f(y) - f(x))}{|y-x|^{n+\alpha+1}} dy, \quad x \in \mathbb{R}^n, \quad (1.1)$$

where

$$\mu_{n,\alpha} = 2^\alpha \pi^{-\frac{n}{2}} \frac{\Gamma(\frac{n+\alpha+1}{2})}{\Gamma(\frac{1-\alpha}{2})}$$

is a renormalizing constant controlling the behavior of ∇^α as $\alpha \rightarrow 1^-$. A simple computation (see [7, Proposition 2.2] for instance) shows that one can equivalently write $\nabla^\alpha f = \nabla I_{1-\alpha} f$ whenever $f \in C_c^\infty(\mathbb{R}^n)$ (and even for less regular functions, see [7, Lemma 3.28(i)] for a more precise statement), where

$$I_s f(x) = 2^{-s} \pi^{-\frac{n}{2}} \frac{\Gamma(\frac{n-s}{2})}{\Gamma(\frac{s}{2})} \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-s}} dy, \quad x \in \mathbb{R}^n,$$

is the *Riesz potential* of order $s \in (0, n)$.

2020 *Mathematics Subject Classification.* Primary 46E35; Secondary 28A12.

Keywords. Fractional gradient, fractional divergence, fractional variation, fractional Hardy inequality, chain rule.

The literature around the operator ∇^α has been quickly growing in the recent years in various research directions. On the one side, we refer the reader to [15,22–25,27,28] for well-posedness results concerning solutions of PDEs and minimizers of functionals involving this fractional operator, and to [3,4,13] for the study of polyconvexity and quasiconvexity in connection with the present fractional setting. On the other side, the properties of ∇^α led to the discovery of new (optimal) embedding inequalities [26,30,31] and the development of a distributional and asymptotic analysis in this fractional framework [5–9,29]. For a general panoramic on the fractional framework, the reader may consult the survey [32] and the monograph [19].

At least for sufficiently smooth functions, the operator ∇^α obeys the following natural fractional integration-by-parts formula

$$\int_{\mathbb{R}^n} f \operatorname{div}^\alpha \varphi \, dx = - \int_{\mathbb{R}^n} \varphi \cdot \nabla^\alpha f \, dx, \quad (1.2)$$

where

$$\operatorname{div}^\alpha \varphi(x) = \mu_{n,\alpha} \int_{\mathbb{R}^n} \frac{(y-x) \cdot (\varphi(y) - \varphi(x))}{|y-x|^{n+\alpha+1}} \, dy, \quad x \in \mathbb{R}^n,$$

is the *fractional α -divergence* of the vector field $\varphi \in \operatorname{Lip}_c(\mathbb{R}^n; \mathbb{R}^n)$.

Equality (1.2) is the fundamental basis of the distributional theory in the present fractional setting developed in the previous papers [5–9]. In more precise terms, by imitating the classical definition of BV functions, for a given exponent $p \in [1, +\infty]$, we define the (total) fractional variation of a function $f \in L^p(\mathbb{R}^n)$ as

$$|D^\alpha f|(\mathbb{R}^n) = \sup \left\{ \int_{\mathbb{R}^n} f \operatorname{div}^\alpha \varphi \, dx : \varphi \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^n), \|\varphi\|_{L^\infty(\mathbb{R}^n; \mathbb{R}^n)} \leq 1 \right\}. \quad (1.3)$$

The above definition naturally gives rise to the linear space of L^p functions with bounded *fractional α -variation*

$$BV^{\alpha,p}(\mathbb{R}^n) = \{f \in L^p(\mathbb{R}^n) : |D^\alpha f|(\mathbb{R}^n) < +\infty\}$$

that can be endowed with the norm

$$\|f\|_{BV^{\alpha,p}(\mathbb{R}^n)} = \|f\|_{L^p(\mathbb{R}^n)} + |D^\alpha f|(\mathbb{R}^n), \quad f \in BV^{\alpha,p}(\mathbb{R}^n).$$

The resulting normed space is Banach and, moreover, one easily checks that $f \in L^p(\mathbb{R}^n)$ belongs to $BV^{\alpha,p}(\mathbb{R}^n)$ if and only if there exists a finite vector-valued Radon measure $D^\alpha f \in \mathcal{M}(\mathbb{R}^n; \mathbb{R}^n)$, the *fractional α -variation measure* of f , such that

$$\int_{\mathbb{R}^n} f \operatorname{div}^\alpha \varphi \, dx = - \int_{\mathbb{R}^n} \varphi \cdot dD^\alpha f$$

for all $\varphi \in \operatorname{Lip}_c(\mathbb{R}^n; \mathbb{R}^n)$, see [6, Theorem 3].

In a very similar way, one can define the *distributional fractional Sobolev space*

$$S^{\alpha,p}(\mathbb{R}^n) = \{f \in L^p(\mathbb{R}^n) : \nabla^\alpha f \in L^p(\mathbb{R}^n; \mathbb{R}^n)\}$$

where $\nabla^\alpha f$ stands for the *distributional fractional α -gradient*, see [7, Definition 3.9]. As proved in [5, Corollary 1] and in [13, Theorem 2.7], $S^{\alpha,p}(\mathbb{R}^n) = L^{\alpha,p}(\mathbb{R}^n)$ whenever $p \in (1, +\infty)$, where $L^{\alpha,p}(\mathbb{R}^n)$ stands for the *Bessel potential space*. We refer the reader to [5, Section 2.1] and to the references therein for an agile account on Bessel potential spaces, and to the discussion in [7, Section 3.9] for the relations between $L^{\alpha,p}(\mathbb{R}^n)$ and the *Gagliardo–Sobolev–Slobodeckij fractional space* $W^{\alpha,p}(\mathbb{R}^n)$.

The study of the space $BV^\alpha(\mathbb{R}^n) = BV^{\alpha,1}(\mathbb{R}^n)$ in the *geometric regime* $p = 1$ was initiated in [7], also in connection with the naturally associated notion of *fractional Caccioppoli perimeter* (see [7, Definition 4.1]), and then further investigated in the subsequent works [5, 8]. The fractional variation of an L^p function for an arbitrary exponent $p \in [1, +\infty]$ has been explored in [6, 8, 9].

Throughout this paper, with a slight abuse of notation (that, however, can be rigorously justified thanks to the analysis done in the previous works [5–9]), in the integer case $\alpha = 1$ we let

$$BV^{1,p}(\mathbb{R}^n) = \{f \in L^p(\mathbb{R}^n) : Df \in \mathcal{M}(\mathbb{R}^n; \mathbb{R}^n)\}$$

be the space of L^p functions, $p \in [1, +\infty]$, with bounded variation.

1.2. Hardy inequality and chain rule

Due to the central role played by the classical Hardy inequality in the theory of integer as well as of fractional Sobolev spaces, see [18] for an account, in [28], Shieh and Spector investigated the validity of the natural analogue of the Hardy inequality in the present fractional setting. In [28, Theorem 1.2], they proved the validity of the following inequality:

$$c_{n,\alpha} \int_{\mathbb{R}^n} \frac{|f(x)|}{|x|^\alpha} dx \leq \int_{\mathbb{R}^n} |\nabla^\alpha |f|| dx \quad (1.4)$$

for all measurable functions f such that $\nabla^\alpha |f| = \nabla I_{1-\alpha} |f| \in L^1(\mathbb{R}^n; \mathbb{R}^n)$, where $c_{n,\alpha} > 0$ is a constant depending on $\alpha \in (0, 1)$ and $n \geq 2$ only. Actually, the validity of (1.4) for $n = 1$ is not explicitly shown in [28], but one can still recover it via an *ad hoc* modification of their argument.

Motivated by (1.4), the authors in [28] asked if it is possible to remove the modulus in the right-hand side of (1.4), that is, more generally, if the following chain rule for the fractional gradient

$$\int_{\mathbb{R}^n} |\nabla^\alpha |f|| dx \leq c_{n,\alpha} \int_{\mathbb{R}^n} |\nabla^\alpha f| dx \quad (1.5)$$

holds whenever f is measurable with $\nabla I_{1-\alpha} f \in L^1(\mathbb{R}^n; \mathbb{R}^n)$, where $c_{n,\alpha} > 0$ is a constant depending on α and n only, see [28, Open Problem 1.4].

Later, Spector proved the validity of the fractional Hardy inequality

$$c_{n,\alpha} \int_{\mathbb{R}^n} \frac{|f(x)|}{|x|^\alpha} dx \leq \int_{\mathbb{R}^n} |\nabla^\alpha f| dx \quad (1.6)$$

for $n \geq 2$, whenever $f \in L^p(\mathbb{R}^n)$ with $p \in [1, \frac{n}{1-\alpha})$ and $\nabla^\alpha f = \nabla I_{1-\alpha} f \in L^1(\mathbb{R}^n; \mathbb{R}^n)$, see [31, Theorem 1.4]. The approach used in [31] completely bypasses the validity of (1.5) and instead relies on an optimal embedding in Lorentz spaces for the Riesz potential, see [31, Theorem 1.1].

The relation between the Hardy inequality in (1.4), as well as the one in (1.6), with the one valid in the usual fractional Sobolev space $W^{\alpha,1}(\mathbb{R}^n)$ easily follows from the elementary inequality

$$\|\nabla^\alpha f\|_{L^1(\mathbb{R}^n; \mathbb{R}^n)} \leq \mu_{n,\alpha} [f]_{W^{\alpha,1}(\mathbb{R}^n)}$$

naturally available for all functions $f \in W^{\alpha,1}(\mathbb{R}^n)$, see [7, Section 1.1]. Similar considerations can be done for the Hardy inequalities in the integrability regime $p \in (1, +\infty)$, see the introductions of [27, 28].

Up to our knowledge, the validity of a chain rule for the fractional gradient ∇^α like (1.5) is still an open problem. Somehow complementing the validity of (1.6) for $n \geq 2$, in the present work we disprove the validity of (1.5) in the case $n = 1$. More precisely, we prove the following result.

Theorem 1.1 (Failure of the chain rule for $n = 1$). *Let $\alpha \in (0, 1)$. The function*

$$f_\alpha(x) = \mu_{1,-\alpha} (|x|^{\alpha-1} \operatorname{sgn} x - |x-1|^{\alpha-1} \operatorname{sgn}(x-1)), \quad x \in \mathbb{R} \setminus \{0, 1\},$$

is such that $f_\alpha \in BV^\alpha(\mathbb{R})$ but $|f_\alpha| \notin BV^\alpha(\mathbb{R})$.

The proof of Theorem 1.1 works by contradiction. Precisely, if $D^\alpha |f_\alpha| \in \mathcal{M}(\mathbb{R})$, then a generalized version of inequality (1.4) for $n = 1$ would hold (see Theorem 1.2 below). Thus we would get $f_\alpha \in L^1(\mathbb{R}; |x|^{-\alpha} \mathcal{L}^1)$, which is clearly false. Actually, inequality (1.4) cannot be directly applied to the function f_α in Theorem 1.1, since $D^\alpha f_\alpha = \delta_0 - \delta_1 \notin L^1(\mathbb{R})$, see [7, Theorem 3.26]. However, one can exploit the regularization properties of $BV^{\alpha,p}$ functions [6, Theorem 4] to suitably extend (1.4), as well as (1.6), to this more general framework.

Theorem 1.2 (Hardy inequality in $BV^{\alpha,p}(\mathbb{R}^n)$ for $p \in [1, \frac{n}{1-\alpha})$). *Let $\alpha \in (0, 1)$ and $p \in [1, \frac{n}{1-\alpha})$. If $f \in BV^{\alpha,p}(\mathbb{R}^n)$, with $f \geq 0$ if $n = 1$, then*

$$c_{n,\alpha} \int_{\mathbb{R}^n} \frac{|f(x)|}{|x-x_0|^\alpha} dx \leq |D^\alpha f|(\mathbb{R}^n)$$

for all $x_0 \in \mathbb{R}^n$, where $c_{n,\alpha} > 0$ is a constant depending on n and α only. In particular, if $n = 1$, the optimal constant is $c_{1,\alpha} = \frac{2\mu_{1,\alpha}}{\alpha}$.

1.3. Local chain rule

We do not know if a counterexample to the chain rule (1.5) like the one in Theorem 1.1 can be provided also for $n \geq 2$.

The current lack of a counterexample to (1.5) may suggest that a stronger version of the chain rule could be valid for the fractional variation for $n \geq 2$, in analogy with the chain rule available for BV functions. More precisely, for a given $\Phi \in \text{Lip}(\mathbb{R})$ such that $\Phi(0) = 0$, one may wonder if the *local* chain rule

$$|D^\alpha \Phi(f)| \leq C(\Phi) |D^\alpha f| \quad \text{in } \mathcal{M}(\mathbb{R}^n) \quad (1.7)$$

holds for all $f \in BV^{\alpha,p}(\mathbb{R}^n)$ with $n \geq 2$, where $C(\Phi) > 0$ is a constant depending on the chosen function Φ only. In the present work, we disprove the validity of (1.7) for all $n \geq 2$ and, actually, we prove the following stronger result.

Theorem 1.3 (Failure of the local chain rule). *Let $\alpha \in (0, 1)$ and $p \in [1, \frac{n}{n-\alpha})$. Let $\Phi \in \text{Lip}(\mathbb{R})$ be such that $\Phi(0) = 0$ and $\Phi \geq 0$. If $\Phi(f) \in BV^{\alpha,p}(\mathbb{R}^n)$ with*

$$\text{supp } |D^\alpha \Phi(f)| \subset \text{supp } |D^\alpha f|$$

for all $f \in BV^{\alpha,p}(\mathbb{R}^n)$, then $\Phi \equiv 0$.

In particular, if we consider $\Phi(t) = |t|$ for $t \in \mathbb{R}$, an immediate consequence of Theorem 1.3 is that, for all $\alpha \in (0, 1)$ and $p \in [1, \frac{n}{n-\alpha})$, there exists a function $f \in BV^{\alpha,p}(\mathbb{R}^n)$ such that $\text{supp } |D^\alpha |f||$ is not contained in $\text{supp } |D^\alpha f|$.

The validity of Theorem 1.3 is a simple consequence again of the analysis made in [7] and of a new surprising rigidity property of non-negative $BV^{\alpha,p}$ functions with $p \in [1, \frac{n}{n-\alpha})$, see Theorem 1.4 below. Here and in the rest of the paper, given $v \in \mathbb{S}^{n-1}$ and $x_0 \in \mathbb{R}^n$, we let

$$H_v^+(x_0) = \{y \in \mathbb{R}^n : (y - x_0) \cdot v > 0\}$$

and

$$H_v(x_0) = \{y \in \mathbb{R}^n : (y - x_0) \cdot v = 0\}.$$

In the case $x_0 = 0$, we simply write $H_v^+ = H_v^+(0)$ and $H_v = H_v(0)$. Moreover, for $\alpha \in (0, 1)$ and $p \in [1, +\infty]$, we let

$$BV_+^{\alpha,p}(\mathbb{R}^n) = \{f \in BV^{\alpha,p}(\mathbb{R}^n) : f \geq 0\}.$$

Theorem 1.4 (Rigidity property in $BV_+^{\alpha,p}(\mathbb{R}^n)$ for $p \in [1, \frac{n}{n-\alpha})$). *Let $\alpha \in (0, 1)$, $p \in [1, \frac{n}{n-\alpha})$ and $f \in BV_+^{\alpha,p}(\mathbb{R}^n)$. If either*

$$\text{supp } |D^\alpha f| \text{ is bounded,} \quad (1.8)$$

or

$$|D^\alpha f|(\overline{H_\nu^+(x_0)}) = 0 \quad \text{for some } x_0 \in \mathbb{R}^n, \nu \in \mathbb{S}^{n-1}, \quad (1.9)$$

or

$$f \in L^\infty(\mathbb{R}^n) \text{ and } D^\alpha f(H_\nu^+(x_0)) = 0 \quad \text{for some } x_0 \in \mathbb{R}^n, \nu \in \mathbb{S}^{n-1}, \quad (1.10)$$

then $f \equiv 0$.

The rigidity property given by Theorem 1.4 strongly underlines the difference between the non-local operator ∇^α and its local integer counterpart ∇ . Indeed, it is easily seen that $BV^{1,p}$ functions do not possess such a rigidity property for any given $p \in [1, +\infty]$, due to the locality of the classical variation measure (for instance, one may consider the characteristic function of the unit ball).

In addition, we recall that, despite of the non-local nature of the fractional gradient, there exist functions $f \in BV^{\alpha,p}(\mathbb{R}^n)$, for $p \in [1, \frac{n}{n-\alpha})$, such that $|D^\alpha f|$ is a finite Radon measure with compact support, see the function defined in Theorem 1.1 for $n = 1$, and [7, Lemma 3.28] as well as [6, Proposition 4] for the general case. Hence, Theorem 1.4 immediately tells us that such functions cannot have constant sign. Conversely, as observed in [13, Section 2.2] in the case $n = 1$, given any non-zero function $f \in C_c^\infty(\mathbb{R}^n)$ with $f \geq 0$ and $\text{supp } f \subset (-L, L)^n$ for some $L > 0$, for each $j \in \{1, \dots, n\}$ we have

$$\nabla_j^\alpha f(x) = e_j \cdot \nabla^\alpha f(x) \neq 0 \text{ at each } x \in \mathbb{R}^n \text{ with } |x_j| \geq L,$$

where e_j is the j -th vector of the standard coordinate basis of \mathbb{R}^n .

We end this section by stating a simple consequence of Theorem 1.2. To this purpose, we define

$$\text{LSC}_b(\mathbb{R}^n) = \{f : \mathbb{R}^n \rightarrow \mathbb{R} : f \text{ lower semicontinuous and bounded}\}.$$

Corollary 1.5. *Let $\alpha \in (0, 1)$ and $p \in [1, \frac{n}{1-\alpha})$. The operator*

$$I_{n-\alpha} : BV_+^{\alpha,p}(\mathbb{R}^n) \rightarrow \text{LSC}_b(\mathbb{R}^n)$$

is continuous. In addition, if $n \geq 2$, then $I_{n-\alpha} : BV^{\alpha,p}(\mathbb{R}^n) \rightarrow L^\infty(\mathbb{R}^n)$ is continuous.

1.4. Integration-by-parts formulas

The rigidity property of non-negative $BV^{\alpha,p}$ functions stated in Theorem 1.4 is, in turn, a consequence of a fractional Gauss–Green formula on half-spaces, see Theorem 1.6 below, which can be regarded as a ‘vectorial’ Hardy-type equality for the fractional variation. Here and in the rest of the paper, for $\alpha \in (0, 1)$ and $p, q \in [1, +\infty]$, we let

$$B_{p,q}^{\alpha}(\mathbb{R}^n) = \{u \in L^p(\mathbb{R}^n) : [u]_{B_{p,q}^{\alpha}(\mathbb{R}^n)} < +\infty\}$$

be the space of *Besov functions* on \mathbb{R}^n , see [14, Chapter 17] for its precise definition and main properties, where

$$[u]_{B_{p,q}^{\alpha}(\mathbb{R}^n)} = \begin{cases} \left(\int_{\mathbb{R}^n} \frac{\|u(\cdot+h)-u\|_{L^p(\mathbb{R}^n)}^q}{|h|^{n+q\alpha}} dh \right)^{\frac{1}{q}} & \text{if } q \in [1, +\infty), \\ \sup_{h \in \mathbb{R}^n \setminus \{0\}} \frac{\|u(\cdot+h)-u\|_{L^p(\mathbb{R}^n)}}{|h|^{\alpha}} & \text{if } q = +\infty. \end{cases}$$

Theorem 1.6 (Fractional Gauss–Green formula on half-spaces). *Let $\alpha \in (0, 1)$, $p \in [1, \frac{n}{n-\alpha})$ and $q \in (\frac{n}{\alpha}, +\infty]$ be such that $\frac{1}{p} + \frac{1}{q} = 1$. If $f \in BV^{\alpha,p}(\mathbb{R}^n) \cap L^{\infty}(\mathbb{R}^n)$, then*

$$\frac{\mu_{1,\alpha}}{\alpha} \lim_{R \rightarrow +\infty} \int_{\mathbb{R}^n} \eta_R(x) \frac{f(x)v}{|(x-x_0) \cdot v|^{\alpha}} dx = -\eta(0) D^{\alpha} f(H_v^+(x_0)) \quad (1.11)$$

whenever $v \in S^{n-1}$ and $x_0 \in \mathbb{R}^n$, where $\eta_R(x) = \eta(\frac{x}{R})$ for $x \in \mathbb{R}^n$ and $R > 0$, for some fixed $\eta \in B_{q,1}^{\alpha}(\mathbb{R}^n)$ with compact support. In particular, if either $\text{supp } f$ is bounded or f has constant sign, then

$$\frac{\mu_{1,\alpha}}{\alpha} \int_{\mathbb{R}^n} \frac{f(x)v}{|(x-x_0) \cdot v|^{\alpha}} dx = -D^{\alpha} f(H_v^+(x_0)). \quad (1.12)$$

We let the reader note that the requirement that $\eta \in B_{q,1}^{\alpha}(\mathbb{R}^n)$ naturally comes from the general integration-by-parts formula obtained in [9, Theorem 1.1], see (2.1) below for a more detailed account.

Actually, Theorem 1.6 is a particular case of the following result, which can be seen as an extension of the integration-by-parts formula (1.2) in the spirit of the fractional Gauss–Green formulas established in [9, Section 3.3]. Here and in the following, we let

$$f^*(x) = \begin{cases} \lim_{r \rightarrow 0^+} \int_{B_r(x)} f(y) dy & \text{if the limit exists,} \\ 0 & \text{otherwise,} \end{cases} \quad (1.13)$$

be the *precise representative* of $f \in L^1_{\text{loc}}(\mathbb{R}^n)$.

Theorem 1.7 (Limit integration-by-parts formula). *Let $\alpha \in (0, 1)$ and let $p \in [1, \frac{n}{n-\alpha})$ and $q \in (\frac{n}{\alpha}, +\infty]$ be such that $\frac{1}{p} + \frac{1}{q} = 1$. If $f \in BV^{\alpha,p}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ and $g \in W_{\text{loc}}^{\alpha,1}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ with $\nabla^\alpha g \in L^1_{\text{loc}}(\mathbb{R}^n; \mathbb{R}^n)$, then*

$$\lim_{R \rightarrow +\infty} \int_{\mathbb{R}^n} \eta_R f \nabla^\alpha g \, dx = -\eta(0) \int_{\mathbb{R}^n} g^* \, dD^\alpha f,$$

where η_R is as in Theorem 1.6 and the limit in (1.13) defining $g^*(x)$ exists for $|D^\alpha f|$ -a.e. $x \in \mathbb{R}^n$.

In order to apply Theorem 1.7 to get Theorem 1.6, one then just needs to explicitly compute the fractional gradient of the characteristic function of a half-space.

Proposition 1.8 (∇^α of a half-space). *Let $\alpha \in (0, 1)$, $v \in \mathbb{S}^{n-1}$ and $x_0 \in \mathbb{R}^n$. We have*

$$\nabla^\alpha \chi_{H_v^+(x_0)}(x) = \frac{\mu_{1,\alpha}}{\alpha} \frac{v}{|(x - x_0) \cdot v|^\alpha} \quad (1.14)$$

for $x \in \mathbb{R}^n \setminus H_v(x_0)$.

It is worth noticing that Theorem 1.6 immediately implies the following version of the fractional Hardy inequality for non-negative $BV^{\alpha,p}$ functions in the regime $p \in [1, \frac{n}{n-\alpha})$, where the right-hand side does not involve the knowledge of the fractional variation on the whole space, but just on a specific half-space.

Corollary 1.9 (Fractional Hardy inequality in $BV_+^{\alpha,p}(\mathbb{R}^n)$ for $p \in [1, \frac{n}{n-\alpha})$). *Let $\alpha \in (0, 1)$ and $p \in [1, \frac{n}{n-\alpha})$. If $f \in BV_+^{\alpha,p}(\mathbb{R}^n)$, then*

$$\frac{\mu_{1,\alpha}}{\alpha} \int_{\mathbb{R}^n} \frac{f(x)}{|(x - x_0) \cdot v|^\alpha} \, dx \leq |D^\alpha f|(\overline{H_v^+(x_0)}) \quad (1.15)$$

for all $x_0 \in \mathbb{R}^n$ and $v \in \mathbb{S}^{n-1}$.

As the reader may notice, Theorem 1.6 allows to prove Theorem 1.4 under the assumption (1.10). To deal with the assumption (1.8), one needs to perform a further integration with respect to the direction $v \in \mathbb{S}^{n-1}$ and obtains the following fractional weighted inequality of Hardy-type. Again, we underline that the fractional variation appearing in the right-hand side is not computed on the whole space, but just on the complement of a particular ball.

Corollary 1.10 (Weighted fractional Hardy-type inequality). *Let $\alpha \in (0, 1)$ and $p \in [1, \frac{n}{n-\alpha})$. If $f \in BV_+^{\alpha,p}(\mathbb{R}^n)$, then*

$$\int_{\mathbb{R}^n} f(x) w_{n,\alpha}(|x - x_0|, r) \, dx \leq |D^\alpha f|(\mathbb{R}^n \setminus B_r(x_0))$$

for all $x_0 \in \mathbb{R}^n$ and $r > 0$, where

$$w_{n,\alpha}(t, r) = \begin{cases} \frac{(n-1)\omega_{n-1}}{n\omega_n} \frac{\mu_{1,\alpha}}{\alpha} \int_{-1}^1 \frac{(1-s^2)^{\frac{n-3}{2}}}{|st-r|^\alpha} ds & \text{for } n \geq 2, \\ \frac{\mu_{1,\alpha}}{2\alpha} \left(\frac{1}{|t-r|^\alpha} + \frac{1}{|t+r|^\alpha} \right) & \text{for } n = 1. \end{cases}$$

In the particular geometric case $f = \chi_E$ for some measurable set $E \subset \mathbb{R}^n$, the above results read as follows (recall that, by [7, Corollary 5.4], if $\chi_E \in BV^\alpha(\mathbb{R}^n)$, then we have $|D^\alpha \chi_E| \ll \mathcal{H}^{n-\alpha}$). Here and in the following, $\mathcal{F}^\alpha E$ denotes the *fractional reduced boundary* in the sense of De Giorgi, see [7, Definition 4.7].

Corollary 1.11 (Geometric case). *Let $\alpha \in (0, 1)$. If $\chi_E \in BV^\alpha(\mathbb{R}^n)$, then*

$$\begin{aligned} \frac{\mu_{1,\alpha}}{\alpha} \int_E \frac{v}{|(x-x_0) \cdot v|^\alpha} dx &= -D^\alpha \chi_E(H_v^+(x_0)), \\ \int_E |\nabla^\alpha \chi_{H_v^+(x_0)}| dx &\leq |D^\alpha \chi_E|(H_v^+(x_0)), \\ \int_E w_{n,\alpha}(|x-x_0|, r) dx &\leq |D^\alpha \chi_E|(\mathbb{R}^n \setminus B_r(x_0)), \end{aligned}$$

for $x_0 \in \mathbb{R}^n$, $v \in \mathbb{S}^{n-1}$ and $r > 0$, where $w_{n,\alpha}$ is as in Corollary 1.10. Moreover, if either $\text{supp } |D^\alpha \chi_E|$ is bounded or $D^\alpha \chi_E(H_v^+(x_0)) = 0$ for some $x_0 \in \mathbb{R}^n$, $v \in \mathbb{S}^{n-1}$, then $|E| = 0$. In particular, if $|E| > 0$, then $\mathcal{F}^\alpha E$ must be unbounded and must intersect all half-spaces.

1.5. Fractional Meyers–Ziemer trace inequalities

As discussed in [32], the Hardy inequality in (1.6) can be also seen as a particular consequence of known interpolation inequalities in Lorentz spaces. Precisely, one recognizes that

$$\frac{1}{|\cdot|^\alpha} \in L^{\frac{n}{\alpha}, \infty}(\mathbb{R}^n),$$

so that (1.6) follows by combining the Hölder inequality

$$\int_{\mathbb{R}^n} \frac{|f(x)|}{|x|^\alpha} dx \leq \|f\|_{L^{\frac{n}{n-\alpha}, 1}(\mathbb{R}^n)} \left\| \frac{1}{|\cdot|^\alpha} \right\|_{L^{\frac{n}{\alpha}, \infty}(\mathbb{R}^n)}$$

with the bound

$$\|f\|_{L^{\frac{n}{n-\alpha}, 1}(\mathbb{R}^n)} \leq c_{n,\alpha} |D^\alpha f|(\mathbb{R}^n),$$

valid for $n \geq 2$, which, in turn, is a consequence of [31, Theorem 1.1].

In the classical integer case, an even more general approach is possible. Indeed, if $f \in BV(\mathbb{R}^n)$ and $\mu \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^n)$ is a non-negative locally finite measure, then

$$\int_{\mathbb{R}^n} |f^*| d\mu \leq c_n \|\mu\|_{n-1} |Df|(\mathbb{R}^n), \quad (1.16)$$

for a dimensional constant $c_n > 0$, where f^* is as in (1.13) and

$$\|\mu\|_s = \sup_{x \in \mathbb{R}^n, r > 0} \frac{\mu(B_r(x))}{r^s}$$

whenever $s \in [0, n]$. The inequality in (1.16) can be found in [17, Theorem 4.7] and is nowadays called the *Meyers–Ziemer trace inequality*. We also refer the reader to the recent work [21] for an interesting historical panoramic around the inequality (1.16). In particular, the authors of [21] note that V. G. Maz'ya proved such an inequality in [16], a few years before the aforementioned [17].

Inequality (1.16) plays a central role in the classical BV framework, since it can be considered as the mother inequality of several embedding inequalities, like the Hardy inequality

$$\int_{\mathbb{R}^n} \frac{|f(x)|}{|x|} dx \leq c_n |Df|(\mathbb{R}^n), \quad (1.17)$$

the Gagliardo–Nirenberg–Sobolev inequality

$$\|f\|_{L^{\frac{n}{n-1}}(\mathbb{R}^n)} \leq c_n |Df|(\mathbb{R}^n), \quad (1.18)$$

and its refinement, the Alvino inequality

$$\|f\|_{L^{\frac{n}{n-1},1}(\mathbb{R}^n)} \leq c_n |Df|(\mathbb{R}^n). \quad (1.19)$$

For a more detailed discussion, we refer the reader to [30, Section 1] and [32, Section 6]. Indeed, as soon as $g \in L^{n,\infty}(\mathbb{R}^n)$, one immediately recognizes that the measure

$$\mu(A) = \int_A g(x) dx, \quad A \subset \mathbb{R}^n,$$

satisfies

$$\|\mu\|_{n-1} \leq c_n \|g\|_{L^{n,\infty}(\mathbb{R}^n)}$$

for some dimensional constant $c_n > 0$ (for instance, see [32, Section 6]), so that one can recover the above inequalities (1.17), (1.18) and (1.19) from (1.16) via known interpolation inequalities in Lorentz spaces.

Motivated by the analogy between BV and BV^α functions, one would be tempted to say that, at least for $n \geq 2$, an inequality of the form

$$\int_{\mathbb{R}^n} |f| d\mu \leq c_{n,\alpha} \|\mu\|_{n-\alpha} \int_{\mathbb{R}^n} |\nabla^\alpha f| dx, \quad (1.20)$$

that is, equivalently,

$$\int_{\mathbb{R}^n} |I_\alpha f| d\mu \leq c_{n,\alpha} \|\mu\|_{n-\alpha} \int_{\mathbb{R}^n} |Rf| dx, \quad (1.21)$$

may hold for all sufficiently regular functions f , see [32, Question 7.1], where

$$Rf(x) = \pi^{-\frac{n+1}{2}} \Gamma\left(\frac{n+1}{2}\right) \lim_{\varepsilon \rightarrow 0^+} \int_{\{|y|>\varepsilon\}} \frac{yf(x+y)}{|y|^{n+1}} dy, \quad x \in \mathbb{R}^n,$$

is the (vector-valued) *Riesz transform* of f . Unfortunately, in [30, Theorem 1.3], Spector ruled out the validity of (1.20), as well as of (1.21), whenever $\alpha \in (0, 1)$.

Nonetheless, recalling that $\nabla^\alpha f = \nabla I_{1-\alpha} f$, one may apply the Meyers–Ziemer trace inequality (1.16) to the function $I_{1-\alpha} f$ to get

$$\int_{\mathbb{R}^n} |(I_{1-\alpha} f)^*| d\mu \leq c_n \|\mu\|_{n-1} \int_{\mathbb{R}^n} |\nabla^\alpha f| dx. \quad (1.22)$$

Interestingly, inequality (1.22) turns out to behave as the mother inequality for the Meyers–Ziemer trace inequality (1.16) as well as for the fractional Hardy inequality (1.4). Indeed, on the one side, taking the limit as $\alpha \rightarrow 1^-$ in (1.22), then one gets inequality (1.16) back. On the other side, if one takes $f \geq 0$ and $\mu = \frac{1}{|\cdot|} \mathcal{L}^n$, then one easily recognizes that

$$\int_{\mathbb{R}^n} I_{1-\alpha} f \frac{dx}{|x|} = c_n \int_{\mathbb{R}^n} I_{1-\alpha} f I_{n-1} dx = c_n \int_{\mathbb{R}^n} I_{n-\alpha} f dx = c_{n,\alpha} \int_{\mathbb{R}^n} \frac{f(x)}{|x|^\alpha} dx,$$

recovering (1.4).

Having the above observations in mind, our last main result is the following rigorous statement of the inequality (1.22).

Theorem 1.12 (Fractional Meyers–Ziemer trace inequality). *Let $\alpha \in (0, 1)$ and $p \in [1, \frac{n}{1-\alpha})$. There exists a dimensional constant $c_n > 0$ such that*

$$\int_{\mathbb{R}^n} |(I_{1-\alpha} f)^*| d\mu \leq c_n \|\mu\|_{n-1} |D^\alpha f|(\mathbb{R}^n) \quad (1.23)$$

for all $f \in BV^{\alpha,p}(\mathbb{R}^n)$ and all $\mu \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^n)$.

As formally observed above, besides providing an alternative route for the proof of Theorem 1.2, Theorem 1.12 leads to the following consequences. Here and in the rest of the paper, we let

$$\mathcal{H}^1(\mathbb{R}^n) = \{f \in L^1(\mathbb{R}^n) : Rf \in L^1(\mathbb{R}^n; \mathbb{R}^n)\}$$

be the (real) *Hardy space*, see [12, 33] for a detailed exposition.

Corollary 1.13 (Meyers–Ziemer trace inequalities). *There exists a dimensional constant $c_n > 0$ with the following properties:*

- (i) *If $f \in BV^{1,p}(\mathbb{R}^n)$ for some $p \in [1, +\infty)$, with $p \leq \frac{n}{n-1}$ if $n \geq 2$, and $\mu \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^n)$, then*

$$\int_{\mathbb{R}^n} |f^*| d\mu \leq c_n \|\mu\|_{n-1} |Df|(\mathbb{R}^n). \quad (1.24)$$

- (ii) *If $f \in \mathcal{H}^1(\mathbb{R}^n)$ and $\mu \in \mathcal{M}_{\text{loc}}^+(\mathbb{R}^n)$, then*

$$\int_{\mathbb{R}^n} |(I_1 f)^*| d\mu \leq c_n \|\mu\|_{n-1} \|Rf\|_{L^1(\mathbb{R}^n; \mathbb{R}^n)}. \quad (1.25)$$

We notice that Corollary 1.13 (ii) positively answers [32, Question 7.1] in the (solely possible) case $\alpha = 1$ and, as well-known, it implies the following stronger version of the Stein–Weiss inequality:

$$\|I_1 f\|_{L^{\frac{n}{n-1}}(\mathbb{R}^n)} \leq c_n \|Rf\|_{L^1(\mathbb{R}^n; \mathbb{R}^n)} \quad (1.26)$$

for all $f \in \mathcal{H}^1(\mathbb{R}^n)$, see [32, Section 1] for a more detailed discussion. Consequently, once again choosing the measure $\mu = \frac{1}{|\cdot|} \mathcal{L}^n$, inequality (1.25) implies the Hardy-type inequality

$$\int_{\mathbb{R}^n} \frac{|I_1 f(x)|}{|x|} dx \leq c_n \|Rf\|_{L^1(\mathbb{R}^n; \mathbb{R}^n)} \quad (1.27)$$

whenever $f \in \mathcal{H}^1(\mathbb{R}^n)$. Inequality (1.27), in turn, can be also inferred from the Hardy inequality (1.17), thanks to the continuity of the map $I_1: \mathcal{H}^1(\mathbb{R}^n) \rightarrow BV^{1, \frac{n}{n-1}}(\mathbb{R}^n)$ provided by (1.26) (see [5, Proposition 3.4 (i)] for the fractional case $\alpha \in (0, 1)$).

1.6. Organization of the paper

The paper is organized as follows. Section 2 is dedicated to the proof of Theorem 1.7. In Section 3, we apply it first to prove Theorem 1.6 and then, in turn, its consequences Corollary 1.10, Theorem 1.4, Theorem 1.1. Finally, in Section 4, we prove Theorem 1.12 and its consequences in Corollary 1.13.

2. Proof of Theorem 1.7

In the proof Theorem 1.7, we take advantage of the following non-local Leibniz rule for $BV^{\alpha,p}$ functions, see [9, Theorem 1.1 and Corollary 2.7]. For $p \in [1, \frac{n}{n-\alpha})$ and

$q \in (\frac{n}{\alpha}, +\infty]$ such that $\frac{1}{p} + \frac{1}{q} = 1$, if $f \in BV^{\alpha,p}(\mathbb{R}^n)$ and $g \in B_{q,1}^{\alpha}(\mathbb{R}^n)$, then $fg \in BV^{\alpha,r}(\mathbb{R}^n)$ for all $r \in [1, p]$, with $\nabla_{\text{NL}}^{\alpha}(f, g) \in L^1(\mathbb{R}^n; \mathbb{R}^n)$ and

$$D^{\alpha}(fg) = g^* D^{\alpha} f + f \nabla^{\alpha} g \mathcal{L}^n + \nabla_{\text{NL}}^{\alpha}(f, g) \mathcal{L}^n \quad \text{in } \mathcal{M}(\mathbb{R}^n; \mathbb{R}^n). \quad (2.1)$$

Here and in the rest of the paper, we let

$$\nabla_{\text{NL}}^{\alpha}(f, g)(x) = \mu_{n,\alpha} \int_{\mathbb{R}^n} \frac{(y-x)(f(y)-f(x))(g(y)-g(x))}{|y-x|^{n+\alpha+1}} dy,$$

for a.e. $x \in \mathbb{R}^n$, be the *non-local fractional α -gradient* of the couple (f, g) .

Proof of Theorem 1.7. Let $R > 0$ be fixed. Since $\eta_R \in B_{q,1}^{\alpha}(\mathbb{R}^n)$ with $q \in (\frac{n}{\alpha}, +\infty]$, by the Sobolev Embedding Theorem (see [1, Theorem 7.34(c)] and [14, Theorem 17.52] for instance) we know that $\eta_R \in C_b(\mathbb{R}^n)$. Now let $(\varrho_{\varepsilon})_{\varepsilon>0}$ be a family of standard mollifiers (see [7, Section 3.3] for example) and let $g_{\varepsilon} = \varrho_{\varepsilon} * g$ for all $\varepsilon > 0$. We note that $g_{\varepsilon} \in \text{Lip}_b(\mathbb{R}^n)$ and $\nabla^{\alpha} g_{\varepsilon} = \varrho_{\varepsilon} * \nabla^{\alpha} g$ for all $\varepsilon > 0$, so that

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^n} \eta_R f \nabla^{\alpha} g_{\varepsilon} dx = \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^n} \varrho_{\varepsilon} * (f \eta_R) \nabla^{\alpha} g dx = \int_{\mathbb{R}^n} \eta_R f \nabla^{\alpha} g dx$$

by the Dominated Convergence Theorem, since

$$|\varrho_{\varepsilon} * (f \eta_R)| \leq \|f\|_{L^{\infty}(\mathbb{R}^n)} \|\eta_R\|_{L^{\infty}(\mathbb{R}^n)} \chi_{A_R}$$

for all $\varepsilon > 0$ sufficiently small, where $A_R \subset \mathbb{R}^n$ is a bounded set such that $A_R \supset \text{supp } \eta_R$. Now let $\varepsilon > 0$ be fixed. By (2.1), we have that $f \eta_R \in BV^{\alpha}(\mathbb{R}^n)$, with

$$D^{\alpha}(f \eta_R) = \eta_R D^{\alpha} f + f \nabla^{\alpha} \eta_R \mathcal{L}^n + \nabla_{\text{NL}}^{\alpha}(f, \eta_R) \mathcal{L}^n \quad \text{in } \mathcal{M}(\mathbb{R}^n; \mathbb{R}^n).$$

Consequently, by [8, Proposition 2.7], we can compute

$$\begin{aligned} \int_{\mathbb{R}^n} \eta_R f \nabla^{\alpha} g_{\varepsilon} dx &= - \int_{\mathbb{R}^n} g_{\varepsilon} dD^{\alpha}(f \eta_R) \\ &= - \int_{\mathbb{R}^n} \eta_R g_{\varepsilon} dD^{\alpha} f - \int_{\mathbb{R}^n} f g_{\varepsilon} \nabla^{\alpha} \eta_R dx \\ &\quad - \int_{\mathbb{R}^n} g_{\varepsilon} \nabla_{\text{NL}}^{\alpha}(f, \eta_R) dx. \end{aligned}$$

On the one side, we can estimate

$$\begin{aligned} \left| \int_{\mathbb{R}^n} f g_{\varepsilon} \nabla^{\alpha} \eta_R dx \right| &\leq \|g_{\varepsilon}\|_{L^{\infty}(\mathbb{R}^n)} \int_{\mathbb{R}^n} |f| |\nabla^{\alpha} \eta_R| dx \\ &\leq \|g\|_{L^{\infty}(\mathbb{R}^n)} \|f\|_{L^p(\mathbb{R}^n)} \|\nabla^{\alpha} \eta_R\|_{L^q(\mathbb{R}^n; \mathbb{R}^n)} \\ &\leq \mu_{n,\alpha} R^{\frac{n}{q}-\alpha} \|f\|_{L^p(\mathbb{R}^n)} \|g\|_{L^{\infty}(\mathbb{R}^n)} [\eta]_{B_{q,1}^{\alpha}(\mathbb{R}^n)}, \end{aligned}$$

thanks to [9, Corollary 2.3]. On the other side, in a similar way, we can bound

$$\begin{aligned} \left| \int_{\mathbb{R}^n} f \nabla_{\text{NL}}^\alpha(g_\varepsilon, \eta_R) dx \right| &\leq \int_{\mathbb{R}^n} |f| |\nabla_{\text{NL}}^\alpha(g_\varepsilon, \eta_R)| dx \\ &\leq \|f\|_{L^p(\mathbb{R}^n)} \|\nabla_{\text{NL}}^\alpha(g_\varepsilon, \eta_R)\|_{L^q(\mathbb{R}^n; \mathbb{R}^n)} \\ &\leq 2\mu_{n,\alpha} \|f\|_{L^p(\mathbb{R}^n)} \|g_\varepsilon\|_{L^\infty(\mathbb{R}^n)} [\eta_R]_{B_{q,1}^\alpha(\mathbb{R}^n)} \\ &\leq 2\mu_{n,\alpha} R^{\frac{n}{q}-\alpha} \|f\|_{L^p(\mathbb{R}^n)} \|g\|_{L^\infty(\mathbb{R}^n)} [\eta]_{B_{q,1}^\alpha(\mathbb{R}^n)} \end{aligned}$$

thanks to [9, Corollary 2.7]. Therefore, thanks to these estimate (which are uniform in ε), we get the limit

$$\lim_{R \rightarrow +\infty} \sup_{\varepsilon > 0} \left| \int_{\mathbb{R}^n} f g_\varepsilon \nabla^\alpha \eta_R dx \right| + \left| \int_{\mathbb{R}^n} f \nabla_{\text{NL}}^\alpha(g_\varepsilon, \eta_R) dx \right| = 0.$$

Now we need to show that

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^n} \eta_R g_\varepsilon dD^\alpha f = \int_{\mathbb{R}^n} \eta_R g^* dD^\alpha f. \quad (2.2)$$

Indeed, since $f \in BV^{\alpha,\infty}(\mathbb{R}^n)$, by [6, Theorem 1] we have that $|D^\alpha f| \ll \mathcal{H}^{n-\alpha}$. Moreover, being $g \in W_{\text{loc}}^{\alpha,1}(\mathbb{R}^n)$, by [20, Proposition 3.1] we can infer that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0^+} g_\varepsilon(x) &= \lim_{\varepsilon \rightarrow 0^+} \varrho_\varepsilon * g(x) \\ &= \lim_{r \rightarrow 0^+} \int_{B_r(x)} g(y) dy = g^*(x) \quad \text{for } \mathcal{H}^{n-\alpha}\text{-a.e. } x \in \mathbb{R}^n, \end{aligned}$$

so that (2.2) immediately follows by the Dominated Convergence Theorem (with respect to the finite measure $|D^\alpha f|$). Finally, since

$$\lim_{R \rightarrow +\infty} \eta_R(x) = \lim_{R \rightarrow +\infty} \eta\left(\frac{x}{R}\right) = \eta(0)$$

for all $x \in \mathbb{R}^n$, by the Dominated Convergence Theorem (with respect to the finite measure $|g^*| |D^\alpha f|$) we conclude that

$$\lim_{R \rightarrow +\infty} \int_{\mathbb{R}^n} \eta_R g^* dD^\alpha f = \eta(0) \int_{\mathbb{R}^n} g^* dD^\alpha f$$

and the proof is complete. \blacksquare

3. Hardy inequalities and failure of the chain rule

3.1. Integration by parts on half-spaces

We begin with the proof of the formula for the fractional gradient of the characteristic function of a half-space.

Proof of Proposition 1.8. By the translation invariance of the fractional gradient (recall [29, Theorem 2.2]), we have

$$\nabla^\alpha \chi_{H_v^+(x_0)}(x) = \nabla^\alpha \chi_{H_v^+}(x - x_0)$$

for all $x \in \mathbb{R}^n$ and so we can assume $x_0 = 0$ without loss of generality. Since $\chi_{H_v^+} \in BV_{\text{loc}}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ and clearly

$$|D\chi_{H_v^+}|(\partial B_R) = \mathcal{H}^{n-1}(H_v \cap \partial B_R) = 0$$

for all $R > 0$, by [8, Proposition 3.5] we get $\nabla^\alpha \chi_{H_v^+} \in L^1_{\text{loc}}(\mathbb{R}^n; \mathbb{R}^n)$ and

$$\begin{aligned} \int_{\mathbb{R}^n} \varphi \cdot \nabla^\alpha \chi_{H_v^+} dx &= \lim_{R \rightarrow +\infty} \int_{\mathbb{R}^n} \varphi \cdot I_{1-\alpha}(\chi_{B_R} D\chi_{H_v^+}) dx \\ &= \lim_{R \rightarrow +\infty} \nu \cdot \int_{\mathbb{R}^n} \varphi I_{1-\alpha}(\chi_{B_R} \mathcal{H}^{n-1} \llcorner H_v) dx \end{aligned} \quad (3.1)$$

for all $\varphi \in \text{Lip}_c(\mathbb{R}^n; \mathbb{R}^n)$. By the Monotone Convergence Theorem, we get

$$\lim_{R \rightarrow +\infty} I_{1-\alpha}(\chi_{B_R} \mathcal{H}^{n-1} \llcorner H_v)(x) = I_{1-\alpha}(\mathcal{H}^{n-1} \llcorner H_v)(x)$$

for \mathcal{L}^n -a.e. $x \in \mathbb{R}^n$. We now claim that

$$I_{1-\alpha}(\mathcal{H}^{n-1} \llcorner H_v)(x) = \frac{\mu_{1,\alpha}}{\alpha} \frac{1}{|x \cdot \nu|^\alpha} \quad \text{for all } x \notin H_v, \quad (3.2)$$

which defines a function in $L^1_{\text{loc}}(\mathbb{R}^n)$. The case $n = 1$ is easy. For $n \geq 2$, we argue as follows. Let $\mathcal{R} \in \text{SO}(n)$ be such that $\mathcal{R}\nu = e_1$, so that

$$(\mathcal{R}x)_1 = (\mathcal{R}x) \cdot e_1 = x \cdot {}^t\mathcal{R}e_1 = x \cdot \nu.$$

By simple changes of variables, we get

$$\begin{aligned} \int_{H_v} \frac{d\mathcal{H}^{n-1}(y)}{|y-x|^{n+\alpha-1}} &= \int_{H_{e_1}} \frac{d\mathcal{H}^{n-1}(y)}{|y-\mathcal{R}x|^{n+\alpha-1}} \\ &= \int_{\mathbb{R}^{n-1}} \frac{dy_2 \cdots dy_n}{\left((\mathcal{R}x)_1^2 + \sum_{j=2}^n (y_j - (\mathcal{R}x)_j)^2 \right)^{\frac{n+\alpha-1}{2}}} \\ &= \int_{\mathbb{R}^{n-1}} \frac{1}{|(\mathcal{R}x)_1|^\alpha} \frac{dy_2 \cdots dy_n}{(1 + |(y_2, \dots, y_n)|^2)^{\frac{n+\alpha-1}{2}}} \\ &= \frac{(n-1)\omega_{n-1}}{|x \cdot \nu|^\alpha} \int_0^{+\infty} \frac{\varrho^{n-2}}{(1 + \varrho^2)^{\frac{n+\alpha-1}{2}}} d\varrho \end{aligned}$$

whenever $x \notin H_v$. By known properties of the Gamma function, it is not difficult to recognize that

$$\int_0^{+\infty} \frac{\varrho^{n-2}}{(1+\varrho^2)^{\frac{n+\alpha-1}{2}}} d\varrho = \frac{\Gamma(\frac{\alpha}{2})\Gamma(\frac{n-1}{2})}{2\Gamma(\frac{n+\alpha-1}{2})},$$

so that

$$I_{1-\alpha}(\mathcal{H}^{n-1} \llcorner H_\nu)(x) = \frac{\mu_{n,\alpha}}{(n+\alpha-1)} \frac{\Gamma(\frac{\alpha}{2})\Gamma(\frac{n-1}{2})}{2\Gamma(\frac{n+\alpha-1}{2})} \frac{(n-1)\omega_{n-1}}{|x \cdot \nu|^\alpha} = \frac{\mu_{1,\alpha}}{\alpha} \frac{1}{|x \cdot \nu|^\alpha}$$

whenever $x \notin H_\nu$, proving (3.2). Therefore, we can apply the Dominated Convergence Theorem in (3.1) to obtain

$$\int_{\mathbb{R}^n} \varphi \cdot \nabla^\alpha \chi_{H_\nu^+} dx = \nu \cdot \int_{\mathbb{R}^n} \varphi I_{1-\alpha}(\mathcal{H}^{n-1} \llcorner H_\nu) dx = \frac{\mu_{1,\alpha}}{\alpha} \nu \cdot \int_{\mathbb{R}^n} \frac{\varphi(x)}{|x \cdot \nu|^\alpha} dx$$

and the conclusion immediately follows. \blacksquare

Having Proposition 1.8 at disposal, we can easily deduce the limit Gauss–Green formula on half-spaces.

Proof of Theorem 1.6. The validity of (1.11) is an immediate consequence of Theorem 1.7, since $\chi_{H_\nu^+(x_0)} \in BV_{\text{loc}}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ with $\nabla^\alpha \chi_{H_\nu^+(x_0)} \in L^1_{\text{loc}}(\mathbb{R}^n; \mathbb{R}^n)$ thanks to Proposition 1.8. For the proof of (1.12), we can simply choose $\eta \in C_c^\infty(\mathbb{R}^n)$ such that $0 \leq \eta \leq 1$ and $\eta(x) = 1$ for $x \in B_1$, so that, arguing component-wise,

$$\lim_{R \rightarrow +\infty} \int_{\mathbb{R}^n} \eta_R(x) \frac{f(x)\nu}{|(x-x_0) \cdot \nu|^\alpha} dx = \int_{\mathbb{R}^n} \frac{f(x)\nu}{|(x-x_0) \cdot \nu|^\alpha} dx$$

either trivially if $\text{supp } f$ is bounded, or by the Monotone Convergence Theorem if f has constant sign. Thus, the proof is complete. \blacksquare

3.2. Fractional Hardy inequalities

We can now deal with the proofs of the fractional Hardy inequalities in Corollary 1.9, Corollary 1.10 and Theorem 1.2.

Proof of Corollary 1.9. Let $(\varrho_\varepsilon)_{\varepsilon>0}$ be a family of standard mollifiers and set $f_\varepsilon = \varrho_\varepsilon * f$ for all $\varepsilon > 0$. Clearly, $f_\varepsilon \in BV_+^{\alpha,p}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$, so that (1.12) implies

$$\frac{\mu_{1,\alpha}}{\alpha} \int_{\mathbb{R}^n} \frac{f_\varepsilon(x)}{|(x-x_0) \cdot \nu|^\alpha} dx \leq |D^\alpha f_\varepsilon|(H_\nu^+(x_0))$$

for all $x_0 \in \mathbb{R}^n$ and $\nu \in \mathbb{S}^{n-1}$. On the left-hand side, we employ Fatou's Lemma to obtain

$$\liminf_{\varepsilon \rightarrow 0^+} \frac{\mu_{1,\alpha}}{\alpha} \int_{\mathbb{R}^n} \frac{f_\varepsilon(x)}{|(x-x_0) \cdot \nu|^\alpha} dx \geq \frac{\mu_{1,\alpha}}{\alpha} \int_{\mathbb{R}^n} \frac{f(x)}{|(x-x_0) \cdot \nu|^\alpha} dx.$$

As for the right-hand side, thanks to [2, Theorem 2.2 (b)] we notice that

$$\limsup_{\varepsilon \rightarrow 0^+} |D^\alpha f_\varepsilon|(H_v^+(x_0)) \leq \limsup_{\varepsilon \rightarrow 0^+} |D^\alpha f|(H_v^+(x_0) + B_\varepsilon) \leq |D^\alpha f|(\overline{H_v^+(x_0)})$$

and this proves (1.15). \blacksquare

Proof of Corollary 1.10. At first, let us also assume that $f \in BV_+^{\alpha,p}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$. Let $r > 0$ be fixed. Choosing $x_0 + r\nu$ in place of x_0 in (1.12) and taking the integral average on \mathbb{S}^{n-1} , we get

$$\begin{aligned} & \frac{\mu_{1,\alpha}}{\alpha} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^n} \frac{f(x)}{|(x-x_0) \cdot \nu - r|^\alpha} dx d\mathcal{H}^{n-1}(\nu) \\ &= - \int_{\mathbb{S}^{n-1}} \nu \cdot D^\alpha f(H_v^+(x_0 + r\nu)) d\mathcal{H}^{n-1}(\nu) \\ &\leq |D^\alpha f|(\mathbb{R}^n \setminus B_r(x_0)). \end{aligned}$$

By Tonelli's Theorem, we can compute

$$\begin{aligned} & \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}^n} \frac{f(x)}{|(x-x_0) \cdot \nu - r|^\alpha} dx d\mathcal{H}^{n-1}(\nu) \\ &= \int_{\mathbb{R}^n} f(x) \int_{\mathbb{S}^{n-1}} \frac{d\mathcal{H}^{n-1}(\nu)}{|(x-x_0) \cdot \nu - r|^\alpha} dx \\ &= \int_{\mathbb{R}^n} f(x) w_{n,\alpha}(|x-x_0|, r) dx, \end{aligned}$$

where in the last inequality we exploited the formula proved in [11, Section D.3] for $n \geq 2$ (the case $n = 1$ being trivial).

Now let $f \in BV_+^{\alpha,p}(\mathbb{R}^n)$ be possibly unbounded. Let $(\varrho_\varepsilon)_{\varepsilon>0}$ be a family of standard mollifiers and set $f_\varepsilon = \varrho_\varepsilon * f$ for all $\varepsilon > 0$. Clearly, $f_\varepsilon \in BV_+^{\alpha,p}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$, so that

$$\int_{\mathbb{R}^n} f_\varepsilon(x) w_{n,\alpha}(|x-x_0|, r) dx \leq |D^\alpha f_\varepsilon|(\mathbb{R}^n \setminus B_r(x_0))$$

for all $\varepsilon > 0$. On the one side, we have

$$\liminf_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^n} f_\varepsilon(x) w_{n,\alpha}(|x-x_0|, r) dx \geq \int_{\mathbb{R}^n} f(x) w_{n,\alpha}(|x-x_0|, r) dx$$

by Fatou's Lemma. On the other side, thanks to [6, Theorem 4] as well as [2, Theorem 2.2(b)], we can estimate

$$|D^\alpha f_\varepsilon|(\mathbb{R}^n \setminus B_r(x_0)) \leq (\varrho_\varepsilon * |D^\alpha f|)(\mathbb{R}^n \setminus B_r(x_0)) \leq |D^\alpha f|(\mathbb{R}^n \setminus B_{r-\varepsilon}(x_0))$$

for all $\varepsilon \in (0, r)$. Consequently, we get

$$\begin{aligned} \int_{\mathbb{R}^n} f(x) w_{n,\alpha}(|x - x_0|, r) dx &\leq \lim_{\varepsilon \rightarrow 0^+} |D^\alpha f|(\mathbb{R}^n \setminus B_{r-\varepsilon}(x_0)) \\ &= |D^\alpha f|(\mathbb{R}^n \setminus B_r(x_0)) \end{aligned}$$

by monotonicity and the proof is complete. \blacksquare

Proof of Theorem 1.2. At first, let $n \geq 2$ and $f \in BV^{\alpha,p}(\mathbb{R}^n)$ with $p \in [1, \frac{n}{1-\alpha})$. Up to a translation, we can assume $x_0 = 0$. Let $(\varrho_\varepsilon)_{\varepsilon>0}$ be a family of standard mollifiers and let $f_\varepsilon = \varrho_\varepsilon * f$ for all $\varepsilon > 0$. By [6, Theorem 4], we know that $f_\varepsilon \in BV^{\alpha,p}(\mathbb{R}^n)$ and $\nabla^\alpha f_\varepsilon = \varrho_\varepsilon * D^\alpha f$ for all $\varepsilon > 0$. Moreover, thanks to [7, Lemma 3.28 (i)] and [6, Proposition 4 (i)], we have $\nabla^\alpha f_\varepsilon = \nabla I_{1-\alpha} f_\varepsilon \in L^1(\mathbb{R}^n; \mathbb{R}^n)$. Therefore, the conclusion follows by applying (1.6) to f_ε and then passing to the limit as $\varepsilon \rightarrow 0^+$ via Fatou's Lemma and [6, Theorem 4]. Let now $n = 1$, $f \in BV_+^{\alpha,p}(\mathbb{R})$ with $p \in [1, \frac{1}{1-\alpha})$, and $f_\varepsilon = \varrho_\varepsilon * f$ for all $\varepsilon > 0$. Clearly, $f_\varepsilon \geq 0$, so that we may employ Corollary 1.9 to get

$$\frac{\mu_{1,\alpha}}{\alpha} \int_{\mathbb{R}} \frac{f_\varepsilon(x)}{|x - x_0|^\alpha} dx = \frac{\mu_{1,\alpha}}{\alpha} \int_{\mathbb{R}} \frac{f_\varepsilon(x)}{|(x - x_0) \cdot \nu|^\alpha} dx \leq |D^\alpha f_\varepsilon|(H_\nu^+(x_0)) \quad (3.3)$$

for all $x_0 \in \mathbb{R}$ and $\nu \in \{\pm 1\}$, since $f_\varepsilon \in BV^{\alpha,\infty}(\mathbb{R})$, and so $|D^\alpha f_\varepsilon| \ll \mathcal{H}^{1-\alpha}$ by [6, Theorem 1]. Hence, if we substitute ν with $-\nu$ in (3.3) and then add the two inequalities, we get

$$\frac{2\mu_{1,\alpha}}{\alpha} \int_{\mathbb{R}} \frac{f_\varepsilon(x)}{|x - x_0|^\alpha} dx \leq |D^\alpha f_\varepsilon|(\mathbb{R}) \leq |D^\alpha f|(\mathbb{R}).$$

Thus, we can pass to the limit as $\varepsilon \rightarrow 0^+$ exploiting again Fatou's Lemma. In order to prove the optimality of the constant $c_{1,\alpha} = \frac{2\mu_{1,\alpha}}{\alpha}$, we choose $f = \chi_{(x_0-1, x_0+1)}$, so that

$$\int_{\mathbb{R}} \frac{\chi_{(x_0-1, x_0+1)}(x)}{|x - x_0|^\alpha} dx = \frac{2}{1-\alpha}.$$

Since

$$|D^\alpha \chi_{(x_0-1, x_0+1)}|(\mathbb{R}) = \frac{4\mu_{1,\alpha}}{\alpha(1-\alpha)},$$

thanks to [7, Example 4.11], we get the optimality of $c_{1,\alpha}$ and the proof is complete. \blacksquare

Remark 3.1. Let $\alpha \in (0, 1)$ and $p \in [1, \frac{n}{n-\alpha})$. Arguing as in the second part of the proof of Theorem 1.2, it is possible to show that

$$\frac{2\mu_{1,\alpha}}{\alpha} \int_{\mathbb{R}^n} \frac{f(x)}{|x - x_0|^\alpha} dx \leq |D^\alpha f|(\mathbb{R}^n)$$

for all $f \in BV_+^{\alpha,p}(\mathbb{R}^n)$. Combining this with [28, Theorem 1.2], we deduce that

$$c_{n,\alpha} \int_{\mathbb{R}^n} \frac{f(x)}{|x - x_0|^\alpha} dx \leq |D^\alpha f|(\mathbb{R}^n) \quad \text{for all } f \in BV_+^{\alpha,p}(\mathbb{R}^n),$$

where

$$c_{n,\alpha} = \max \left\{ \frac{2\mu_{1,\alpha}}{\alpha}, \gamma_{n,\alpha} \right\}$$

and

$$\gamma_{n,\alpha} = \frac{2^\alpha \Gamma(\frac{\alpha}{2}) \Gamma(\frac{n+1}{2})}{\pi^{1-\frac{\alpha}{2}} \Gamma(\frac{n-\alpha}{2})}.$$

However, for $n \geq 3$, one can see that $\gamma_{n,\alpha} > \frac{2\mu_{1,\alpha}}{\alpha}$ for all $\alpha \in (0, 1)$. To the best of our knowledge, it is not known whether $c_{n,\alpha}$ is the optimal constant for some $n \geq 2$ and $\alpha \in (0, 1)$.

3.3. Failure of the fractional chain rule

We begin with the proof of the rigidity property contained in Theorem 1.4.

Proof of Theorem 1.4. If $\text{supp}|D^\alpha f|$ is bounded, then $|D^\alpha f|(\mathbb{R}^n \setminus B_r) = 0$ for some $r > 0$. Hence, by Corollary 1.10, we must have $f = 0$ \mathcal{L}^n -a.e. in \mathbb{R}^n , being $w_{n,\alpha} > 0$. If, instead, $|D^\alpha f|(H_v^+(x_0)) = 0$ or $f \in L^\infty(\mathbb{R}^n)$ and $D^\alpha f(H_v^+(x_0)) = 0$ for some $x_0 \in \mathbb{R}^n$ and $v \in \mathbb{S}^{n-1}$, then we similarly conclude by (1.12) in Theorem 1.6 and Corollary 1.9. ■

We can now end this section by showing the failure of the fractional chain rule. Here and in the following, we let

$$(-\Delta)^{\frac{\beta}{2}} f(x) = v_{n,\beta} \int_{\mathbb{R}^n} \frac{f(x+y) - f(x)}{|y|^{n+\beta}} dy, \quad \text{for a.e. } x \in \mathbb{R}^n,$$

be the *fractional Laplacian* of order $\beta \in (0, 1)$ of the function $f \in W^{\beta,1}(\mathbb{R}^n)$, where

$$v_{n,\beta} = 2^\beta \pi^{-\frac{n}{2}} \frac{\Gamma(\frac{n+\beta}{2})}{\Gamma(-\frac{\beta}{2})}.$$

Note that $(-\Delta)^{\frac{\beta}{2}}: W^{\beta,1}(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$ is continuous (see [7, Section 3.10] for a more detailed discussion). In particular, this operator is well posed on BV functions.

Proof of Theorem 1.3. Let $Q_1 = (-1, 1)^n$. We consider the function

$$f = (-\Delta)^{\frac{1-\alpha}{2}} \chi_{Q_1},$$

that is,

$$\begin{aligned} f(x) = v_{n,1-\alpha} & \left(-\chi_{Q_1}(x) \int_{\mathbb{R}^n \setminus Q_1} \frac{1}{|y-x|^{n+1-\alpha}} dy \right. \\ & \left. + \chi_{\mathbb{R}^n \setminus Q_1}(x) \int_{Q_1} \frac{1}{|y-x|^{n+1-\alpha}} dy \right), \end{aligned} \quad (3.4)$$

for $x \in \mathbb{R}^n \setminus \partial Q_1$. Thanks to [7, Lemma 3.28 (ii)], we know that $f \in BV^\alpha(\mathbb{R}^n)$ with $D^\alpha f = D\chi_{Q_1}$. By [6, Theorem 6], we also have that $f \in BV^{\alpha,p}(\mathbb{R}^n)$ for all $p \in [1, \frac{n}{n-\alpha})$. Now let $\Phi \in \text{Lip}_b(\mathbb{R})$ be such that $\Phi(0) = 0$ and $\Phi \geq 0$ and assume that $\Phi(f) \in BV^\alpha(\mathbb{R}^n)$ with

$$\text{supp } |D^\alpha \Phi(f)| \subset \text{supp } |D^\alpha f| = \text{supp } |D\chi_{Q_1}| = \partial Q_1.$$

Note that, again by [6, Theorem 6], $\Phi(f) \in BV_+^{\alpha,p}(\mathbb{R}^n)$ for all $p \in [1, \frac{n}{n-\alpha})$. Consequently, $\text{supp } |D^\alpha \Phi(f)|$ is compact, so that $\Phi(f) \equiv 0$ thanks to Theorem 1.4. Since $v_{n,1-\alpha} < 0$, we observe that $f(x) \rightarrow 0^-$ as $|x| \rightarrow +\infty$ and, moreover,

$$\begin{aligned} \liminf_{t \rightarrow 1^+} \int_{Q_1} \frac{1}{|y-te_1|^{n+1-\alpha}} dy & \geq \int_{Q_1} \frac{dy}{|y-e_1|^{n+\alpha-1}} \\ & \geq \sup_{\varepsilon \in (0,1)} \int_{(-1,1-\varepsilon) \times (-\varepsilon,\varepsilon)^{n-1}} \frac{dy}{|y-e_1|^{n+\alpha-1}} \\ & \geq c_{n,\alpha} \sup_{\varepsilon \in (0,1)} \int_{(-1,1-\varepsilon) \times (-\varepsilon,\varepsilon)^{n-1}} \frac{dy}{|y_1-1|^{n+\alpha-1}} \\ & = c_{n,\alpha} \sup_{\varepsilon \in (0,1)} \varepsilon^{n-1} (\varepsilon^{\alpha-n} - 2^{\alpha-n}) = +\infty, \end{aligned}$$

thanks to Fatou's Lemma. As a consequence, $f(\mathbb{R}^n) \supset (-\infty, 0)$ and thus $\Phi(t) = 0$ for all $t \in (-\infty, 0)$. Replacing f with $-f$, we also get that $\Phi(t) = 0$ for all $t \in (0, +\infty)$, proving that $\Phi \equiv 0$ and the proof is complete. ■

Proof of Theorem 1.1. By [7, Theorem 3.26], we know that $f_\alpha \in BV^\alpha(\mathbb{R})$. We claim that $|f_\alpha| \notin BV^\alpha(\mathbb{R})$. By contradiction, if $|f_\alpha| \in BV^\alpha(\mathbb{R})$, then Theorem 1.2 implies that

$$c_\alpha \int_{\mathbb{R}} \frac{|f_\alpha(x)|}{|x|^\alpha} dx \leq |D^\alpha |f_\alpha||(\mathbb{R}) < +\infty. \quad (3.5)$$

However, for $x \in (0, 1)$, we have

$$\frac{|f_\alpha(x)|}{|x|^\alpha} = |\mu_{1,-\alpha}| \left(\frac{1}{x} + \frac{(1-x)^{\alpha-1}}{x^\alpha} \right),$$

contradicting (3.5) and the proof is complete. ■

4. Fractional Meyers–Ziemer trace inequality

We begin by noticing that, somehow formulating in a more rigorous way the ideas sketched in the introduction, one can prove Theorem 1.12 by directly applying the standard Meyers–Ziemer trace inequality (1.16) to the function $u = I_{1-\alpha}f$ whenever $f \in BV^{\alpha,p}(\mathbb{R}^n)$ with $p \in (1, \frac{n}{1-\alpha})$, since

$$I_{1-\alpha}: BV^{\alpha,p}(\mathbb{R}^n) \rightarrow BV^{1,\frac{n}{n-1}}(\mathbb{R}^n)$$

with $Du = D^\alpha f$ in $\mathcal{M}(\mathbb{R}^n; \mathbb{R}^n)$, thanks to [6, Proposition 4 (i)]. In the case $p = 1$, we only have

$$I_{1-\alpha}: BV^\alpha(\mathbb{R}^n) \rightarrow BV^{1,q}(\mathbb{R}^n)$$

for all $q \in (\frac{n}{n-1+\alpha}, \frac{n}{n-1})$ with $Du = D^\alpha f$ in $\mathcal{M}(\mathbb{R}^n; \mathbb{R}^n)$, in virtue of [7, Remark 3.29] and [6, Theorem 6], but this is still enough in order to directly exploit (1.16).

Below, we instead outline a direct argument showing that the very same line of reasoning used in [17] (see [32, Section 7] for a more detailed explanation) to prove (1.16) works as well for proving Theorem 1.12.

Proof of Theorem 1.12. Let $f \in BV^{\alpha,p}(\mathbb{R}^n)$ for some $p \in [1, \frac{n}{1-\alpha})$. Let $(\varrho_\varepsilon)_{\varepsilon>0}$ be a family of standard mollifiers and let $f_\varepsilon = \varrho_\varepsilon * f$ for all $\varepsilon > 0$. By [6, Theorem 4], we know that $f_\varepsilon \in BV^{\alpha,p}(\mathbb{R}^n)$ with $\nabla^\alpha f_\varepsilon = \varrho_\varepsilon * D^\alpha f$ for all $\varepsilon > 0$. Now let $u_\varepsilon = I_{1-\alpha}f_\varepsilon$ for all $\varepsilon > 0$. By what we have just observed above, it is not difficult to see that $u_\varepsilon \in BV^{1,q}(\mathbb{R}^n) \cap C^\infty(\mathbb{R}^n)$ for some $q \in (\frac{n}{n-1+\alpha}, \frac{n}{n-1}]$ with

$$|\nabla u_\varepsilon| = |\nabla^\alpha f_\varepsilon| \leq \varrho_\varepsilon * |D^\alpha f| \quad \text{in } L^1(\mathbb{R}^n).$$

Therefore, we can estimate

$$\int_{\mathbb{R}^n} |\nabla u_\varepsilon| dx = \int_{\mathbb{R}^n} |\nabla^\alpha f_\varepsilon| dx \leq |D^\alpha f|(\mathbb{R}^n) < +\infty$$

and, moreover, the set

$$E_t^\varepsilon = \{x \in \mathbb{R}^n : |u_\varepsilon(x)| > t\}$$

is open with finite perimeter for a.e. $t > 0$. Since

$$\frac{|E_t^\varepsilon \cap B_r(x)|}{|B_r(x)|} \leq \frac{\min\{|E_t^\varepsilon|, |B_r(x)|\}}{|B_r(x)|}$$

and

$$|E_t^\varepsilon| = |\{x \in \mathbb{R}^n : |I_{1-\alpha}f_\varepsilon(x)| > t\}| \leq c_{n,\alpha,p} \left(\frac{\|f\|_{L^p(\mathbb{R}^n)}}{t} \right)^{\frac{np}{n-(1-\alpha)p}} < +\infty$$

by the Hardy–Littlewood–Sobolev inequality (see [12, Theorem 1.2.3] for instance), for each given $x \in E_t^\varepsilon$ the function

$$r \mapsto \frac{|E_t^\varepsilon \cap B_r(x)|}{|B_r(x)|}$$

is continuous, equals 1 for small $r > 0$ (since E_t^ε is open) and tends to zero as $r \rightarrow +\infty$. Thus, reasoning exactly as in [30, Section 6], via a routine Vitali covering argument we can estimate

$$\mu(E_t^\varepsilon) \leq c_n \|\mu\|_{n-1} |D\chi_{E_t^\varepsilon}|(\mathbb{R}^n)$$

for a.e. $t > 0$, where $c_n > 0$ is a dimensional constant. Therefore, by the coarea formula and the chain rule for functions with bounded variation, we can estimate

$$\begin{aligned} \int_{\mathbb{R}^n} |u_\varepsilon| d\mu &= \int_{\mathbb{R}} \mu(E_t^\varepsilon) dt \\ &\leq c_n \|\mu\|_{n-1} \int_{\mathbb{R}} |D\chi_{E_t^\varepsilon}|(\mathbb{R}^n) dt \\ &= c_n \|\mu\|_{n-1} \int_{\mathbb{R}^n} |\nabla u_\varepsilon| dx \\ &\leq c_n \|\mu\|_{n-1} |D^\alpha f|(\mathbb{R}^n) \end{aligned}$$

for all $\varepsilon > 0$. Now, assuming $\|\mu\|_{n-1} < +\infty$ without loss of generality, it is standard to see that $\mu \ll \mathcal{H}^{n-1}$, see [20] and the references therein for a more detailed discussion. Therefore, since

$$\lim_{\varepsilon \rightarrow 0^+} u_\varepsilon(x) = \lim_{\varepsilon \rightarrow 0^+} \varrho_\varepsilon * u(x) = u^*(x) \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in \mathbb{R}^n$$

(see [10, Section 5.9] for instance), by the Fatou's Lemma we conclude that

$$\int_{\mathbb{R}^n} |(I_{1-\alpha} f)^*| d\mu \leq \liminf_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^n} |u_\varepsilon| d\mu \leq c_n \|\mu\|_{n-1} |D^\alpha f|(\mathbb{R}^n)$$

and the proof is complete. \blacksquare

We now conclude our paper with the proof of Corollary 1.13.

Proof of Corollary 1.13. The validity of (i) for any $f \in C_c^\infty(\mathbb{R}^n)$ follows directly from Theorem 1.12 combined with the asymptotic analysis obtained in [8]. For a general $f \in BV^{1, \frac{n}{n-1}}(\mathbb{R}^n)$, one just needs to perform a routine approximation argument thanks to [6, Proposition 1]. The validity of (ii) follows in a similar way, this time relying on the asymptotic analysis carried out in [5]. \blacksquare

Acknowledgments. The authors thank Daniel Spector for many valuable comments on a preliminary version of the present work and for pointing out the references [16, 21]. Part of this work was undertaken while the authors were visiting each other at the University of Pisa and the Scuola Internazionale Superiore di Studi Avanzati (SISSA) in Trieste. They would like to thank these institutions for the support and warm hospitality during the visits.

Funding. The authors are members of the Istituto Nazionale di Alta Matematica (INdAM), Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA). The first author is partially supported by the INdAM–GNAMPA 2022 Project *Alcuni problemi associati a funzionali integrali: riscoperta strumenti classici e nuovi sviluppi*, codice CUP_E55F22000270001, and has received funding from the MIUR PRIN 2017 Project “Gradient Flows, Optimal Transport and Metric Measure Structures”. The second author is partially supported by the INdAM–GNAMPA 2022 Project *Analisi geometrica in strutture subriemanniane*, codice CUP_E55F22000270001, and has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement No. 945655).

References

- [1] R. A. Adams and J. J. F. Fournier, *Sobolev spaces*. Second edn., Pure Appl. Math., Amst. 140, Elsevier/Academic Press, Amsterdam, 2003 Zbl [1098.46001](#) MR [2424078](#)
- [2] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of bounded variation and free discontinuity problems*. Oxford Math. Monogr., The Clarendon Press, Oxford University Press, New York, 2000 Zbl [0957.49001](#) MR [1857292](#)
- [3] J. C. Bellido, J. Cueto, and C. Mora-Corral, Fractional Piola identity and polyconvexity in fractional spaces. *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **37** (2020), no. 4, 955–981 Zbl [1442.35445](#) MR [4104831](#)
- [4] J. C. Bellido, J. Cueto, and C. Mora-Corral, Γ -convergence of polyconvex functionals involving s -fractional gradients to their local counterparts. *Calc. Var. Partial Differential Equations* **60** (2021), no. 1, Paper No. 7 Zbl [1455.49008](#) MR [4179861](#)
- [5] E. Bruè, M. Calzi, G. E. Comi, and G. Stefani, A distributional approach to fractional Sobolev spaces and fractional variation: asymptotics II. *C. R. Math. Acad. Sci. Paris* **360** (2022), 589–626 Zbl [07547261](#) MR [4449863](#)
- [6] G. E. Comi, D. Spector, and G. Stefani, The fractional variation and the precise representative of $BV^{\alpha,p}$ functions. *Fract. Calc. Appl. Anal.* **25** (2022), no. 2, 520–558 Zbl [07636540](#) MR [4437291](#)
- [7] G. E. Comi and G. Stefani, A distributional approach to fractional Sobolev spaces and fractional variation: existence of blow-up. *J. Funct. Anal.* **277** (2019), no. 10, 3373–3435 Zbl [1437.46039](#) MR [4001075](#)

- [8] G. E. Comi and G. Stefani, A distributional approach to fractional Sobolev spaces and fractional variation: asymptotics I. *Rev. Mat. Complut.* (2022)
DOI [10.1007/s13163-022-00429-y](https://doi.org/10.1007/s13163-022-00429-y)
- [9] G. E. Comi and G. Stefani, Leibniz rules and Gauss–Green formulas in distributional fractional spaces. *J. Math. Anal. Appl.* **514** (2022), no. 2, Paper No. 126312
Zbl [07545051](https://zbmath.org/?q=ser/07545051) MR [4422400](https://www.ams.org/mathscinet-getitem?mr=4422400)
- [10] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*. Revised edn., Textb. Math., CRC Press, Boca Raton, FL, 2015 Zbl [1310.28001](https://zbmath.org/?q=ser/1310.28001) MR [3409135](https://www.ams.org/mathscinet-getitem?mr=3409135)
- [11] L. Grafakos, *Classical Fourier analysis*. Third edn., Grad. Texts Math. 249, Springer, New York, 2014 Zbl [1304.42001](https://zbmath.org/?q=ser/1304.42001) MR [3243734](https://www.ams.org/mathscinet-getitem?mr=3243734)
- [12] L. Grafakos, *Modern Fourier analysis*. Third edn., Grad. Texts Math. 250, Springer, New York, 2014 Zbl [1304.42002](https://zbmath.org/?q=ser/1304.42002) MR [3243741](https://www.ams.org/mathscinet-getitem?mr=3243741)
- [13] C. Kreisbeck and H. Schönberger, Quasiconvexity in the fractional calculus of variations: characterization of lower semicontinuity and relaxation. *Nonlinear Anal.* **215** (2022), Paper No. 112625 Zbl [1478.49011](https://zbmath.org/?q=ser/1478.49011) MR [4330183](https://www.ams.org/mathscinet-getitem?mr=4330183)
- [14] G. Leoni, *A first course in Sobolev spaces*. Second edn., Grad. Stud. Math. 181, American Mathematical Society, Providence, RI, 2017 Zbl [1382.46001](https://zbmath.org/?q=ser/1382.46001) MR [3726909](https://www.ams.org/mathscinet-getitem?mr=3726909)
- [15] C. W. K. Lo and J. F. Rodrigues, On a class of fractional obstacle type problems related to the distributional Riesz derivative. 2021 arXiv:[2101.06863](https://arxiv.org/abs/2101.06863)
- [16] V. G. Maz'ja, *The summability of functions belonging to Sobolev spaces*. (in Russian) Problems of mathematical analysis, No. 5: Linear and nonlinear differential equations, Differential operators (Russian), pp. 66–98. Izdat. Leningrad. Univ., Leningrad, 1975
MR [0511931](https://www.ams.org/mathscinet-getitem?mr=0511931)
- [17] N. G. Meyers and W. P. Ziemer, Integral inequalities of Poincaré and Wirtinger type for BV functions. *Amer. J. Math.* **99** (1977), no. 6, 1345–1360 Zbl [0416.46025](https://zbmath.org/?q=ser/0416.46025)
MR [507433](https://www.ams.org/mathscinet-getitem?mr=507433)
- [18] P. Mironescu, The role of the Hardy type inequalities in the theory of function spaces. *Rev. Roumaine Math. Pures Appl.* **63** (2018), no. 4, 447–525 Zbl [1424.46050](https://zbmath.org/?q=ser/1424.46050) MR [3892576](https://www.ams.org/mathscinet-getitem?mr=3892576)
- [19] A. C. Ponce, *Elliptic PDEs, measures and capacities*. EMS Tracts Math. 23, European Mathematical Society (EMS), Zürich, 2016 Zbl [1357.35003](https://zbmath.org/?q=ser/1357.35003) MR [3675703](https://www.ams.org/mathscinet-getitem?mr=3675703)
- [20] A. C. Ponce and D. Spector, A boxing inequality for the fractional perimeter. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **20** (2020), no. 1, 107–141 Zbl [1466.46029](https://zbmath.org/?q=ser/1466.46029) MR [4088737](https://www.ams.org/mathscinet-getitem?mr=4088737)
- [21] B. Raiță, D. Spector, and D. Stolyarov, A trace inequality for solenoidal charges. *Potential Anal.* (2022) DOI [10.1007/s11118-022-10008-x](https://doi.org/10.1007/s11118-022-10008-x)
- [22] J. F. Rodrigues and L. Santos, On nonlocal variational and quasi-variational inequalities with fractional gradient. *Appl. Math. Optim.* **80** (2019), no. 3, 835–852 Zbl [1429.49011](https://zbmath.org/?q=ser/1429.49011)
MR [4026601](https://www.ams.org/mathscinet-getitem?mr=4026601)
- [23] J. F. Rodrigues and L. Santos, Correction to: On nonlocal variational and quasi-variational inequalities with fractional gradient. *Appl. Math. Optim.* **84** (2021), no. 3, 3565–3567
Zbl [1472.49021](https://zbmath.org/?q=ser/1472.49021) MR [4308239](https://www.ams.org/mathscinet-getitem?mr=4308239)
- [24] A. Schikorra, T.-T. Shieh, and D. Spector, L^p theory for fractional gradient PDE with VMO coefficients. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **26** (2015), no. 4, 433–443 Zbl [1332.35384](https://zbmath.org/?q=ser/1332.35384) MR [3420498](https://www.ams.org/mathscinet-getitem?mr=3420498)

- [25] A. Schikorra, T.-T. Shieh, and D. E. Spector, Regularity for a fractional p -Laplace equation. *Commun. Contemp. Math.* **20** (2018), no. 1, Paper No. 1750003 Zbl [1386.35455](#) MR [3714833](#)
- [26] A. Schikorra, D. Spector, and J. Van Schaftingen, An L^1 -type estimate for Riesz potentials. *Rev. Mat. Iberoam.* **33** (2017), no. 1, 291–303 Zbl [1375.47039](#) MR [3615452](#)
- [27] T.-T. Shieh and D. E. Spector, On a new class of fractional partial differential equations. *Adv. Calc. Var.* **8** (2015), no. 4, 321–336 Zbl [1330.35510](#) MR [3403430](#)
- [28] T.-T. Shieh and D. E. Spector, On a new class of fractional partial differential equations II. *Adv. Calc. Var.* **11** (2018), no. 3, 289–307 Zbl [1451.35257](#) MR [3819528](#)
- [29] M. Šilhavý, Fractional vector analysis based on invariance requirements (critique of coordinate approaches). *Contin. Mech. Thermodyn.* **32** (2020), no. 1, 207–228 Zbl [1443.26004](#) MR [4048032](#)
- [30] D. Spector, A noninequality for the fractional gradient. *Port. Math.* **76** (2019), no. 2, 153–168 Zbl [1453.46035](#) MR [4065096](#)
- [31] D. Spector, An optimal Sobolev embedding for L^1 . *J. Funct. Anal.* **279** (2020), no. 3, Paper No. 108559 Zbl [1455.46039](#) MR [4093790](#)
- [32] D. Spector, New directions in harmonic analysis on L^1 . *Nonlinear Anal.* **192** (2020), Paper No. 111685 Zbl [1437.42037](#) MR [4034690](#)
- [33] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. Princeton Math. Ser. 43, Princeton University Press, Princeton, NJ, 1993 Zbl [0821.42001](#) MR [1232192](#)

Received 7 June 2022; revised 2 December 2022.

Giovanni E. Comi

Centro di Ricerca Matematica “Ennio De Giorgi”, Scuola Normale Superiore, Piazza dei Cavalieri 3, 56126 Pisa, Italy; giovanni.comi@sns.it

Giorgio Stefani

Scuola Internazionale Superiore di Studi Avanzati (SISSA), Via Bonomea 265, 34136 Trieste, Italy; giorgio.stefani.math@gmail.com