

---

---

## *Short note*     **A new refinement of the Garfunkel–Bankoff inequality**

---

---

Quang Hung Tran

**Abstract.** This note introduces an innovative refinement of the Garfunkel–Bankoff inequality, further improving upon the Finsler–Hadwiger inequality and the Lukarevski–Marinescu inequality within a triangle.

### **1 Introduction**

In the triangle  $ABC$ , we use the usual symbols, where  $A$ ,  $B$ , and  $C$  denote the measures of the three angles, and  $a$ ,  $b$ , and  $c$  represent the lengths of the sides opposite angles  $A$ ,  $B$ , and  $C$ , respectively. Additionally,  $R$ ,  $r$ , and  $s$  stand for the circumradius, inradius, and semiperimeter of triangle  $ABC$ .

In [3], J. Garfunkel proposed the following inequality:

$$\tan^2 \frac{A}{2} + \tan^2 \frac{B}{2} + \tan^2 \frac{C}{2} \geq 2 - 8 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}. \quad (1)$$

The first proof of inequality (1) was provided by L. Bankoff in [1], so inequality (1) is also known as the Garfunkel–Bankoff inequality. It is worth noting that inequality (1) also represents an improvement of the Finsler–Hadwiger inequality (refer to [8, 9])

$$a^2 + b^2 + c^2 \geq 4\sqrt{3}S + (b - c)^2 + (c - a)^2 + (a - b)^2,$$

and at the same time, it serves as an equivalent form of the Kooi inequality (see [6])

$$s^2 \leq \frac{R(4R + r)^2}{2(2R - r)}.$$

In [10], Wei-Dong Jiang proposed the following inequality:

$$\tan^2 \frac{A}{2} + \tan^2 \frac{B}{2} + \tan^2 \frac{C}{2} \geq 2 - 8 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} + \frac{r^2(R - 2r)}{4R^2(R - r)}. \quad (2)$$

This represents a strengthened version of inequality (1), and therefore, inequality (2) is an improvement upon the Finsler–Hadwiger and Kooi inequalities. It is worth noting that M. Lukarevski and D. S. Marinescu also provided a refinement of Kooi’s inequality

(see [4–6]),

$$s^2 \leq \frac{R(4R+r)^2}{2(2R-r)} - \frac{r^2(R-2r)}{4R}. \quad (3)$$

However, inequality (2) remains stronger than inequality (3) (see [10]).

In this article, we propose a new inequality that is an improvement of inequality (2) as follows:

$$\tan^2 \frac{A}{2} + \tan^2 \frac{B}{2} + \tan^2 \frac{C}{2} \geq 2 - 8 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} + \frac{r^2(R-2r)}{2(2R^2-r^2)(R-r)}. \quad (4)$$

Clearly, inequality (4) directly extends the Garfunkel–Bankoff inequality (inequality (1)). From inequality (4), we also have new improvements for the Finsler–Hadwiger and Lukarevski–Marinescu inequalities as follows:

- a new improvement of the Finsler–Hadwiger inequality,

$$a^2 + b^2 + c^2 \geq 4 \sqrt{4 - \frac{2r}{R} + \frac{r^2(R-2r)}{2(2R^2-r^2)(R-r)}} S + (b-c)^2 + (c-a)^2 + (a-b)^2,$$

- a new improvement of the Lukarevski–Marinescu inequality (including the Kooi inequality),

$$s^2 \leq \frac{R(4R+r)^2}{2(2R-r)} - \frac{r^2(R-2r)}{2(2R^2-r^2)(R-r)}.$$

These extensions are all stronger than the extensions by Wei-Dong Jiang because inequality (4) is stronger than inequality (2), as it is evident from the fact that

$$\frac{r^2(R-2r)}{2(2R^2-r^2)(R-r)} \geq \frac{r^2(R-2r)}{4R^2(R-r)} \quad \text{or} \quad 4R^2 > 4R^2 - 2r^2.$$

## 2 Proof of inequality (4)

In this proof, we employ the following fundamental inequality within a triangle:

$$s^2 \leq 2R^2 + 10Rr - r^2 + 2(R-2r)\sqrt{R(R-2r)}.$$

Historical references to the method of usage, as well as the nomenclature of this inequality, can be found in [7].

*Proof.* Using the well-known identities

$$\sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} = \frac{r}{4R}$$

and

$$\tan^2 \frac{A}{2} + \tan^2 \frac{B}{2} + \tan^2 \frac{C}{2} = \frac{(4R+r)^2}{s^2} - 2,$$

inequality (4) is equivalent to

$$s^2 \leq \frac{2R(32R^5 + r^5 + 7Rr^4 + 6R^2r^3 - 30R^3r^2 - 16R^4r)}{16R^4 - 4r^4 + 10Rr^3 + R^2r^2 - 24R^3r}.$$

Using the fundamental inequality of a triangle (see [2, 7])

$$s^2 \leq 2R^2 + 10Rr - r^2 + 2(R - 2r)\sqrt{R(R - 2r)},$$

we must prove that

$$\begin{aligned} & 2R^2 + 10Rr - r^2 + 2(R - 2r)\sqrt{R(R - 2r)} \\ & \leq \frac{2R(32R^5 + r^5 + 7Rr^4 + 6R^2r^3 - 30R^3r^2 - 16R^4r)}{16R^4 - 4r^4 + 10Rr^3 + R^2r^2 - 24R^3r}. \end{aligned} \quad (5)$$

Putting  $t = \frac{r}{R}$ , we have  $0 < t \leq \frac{1}{2}$ . Inequality (5) is equivalent to

$$2 + 10t - t^2 + 2(1 - 2t)\sqrt{1 - 2t} \leq \frac{2(t^5 + 7t^4 + 6t^3 - 30t^2 - 16t + 32)}{-4t^4 + 10t^3 + t^2 - 24t + 16}.$$

This is true since

$$\begin{aligned} & \left[ \frac{2(t^5 + 7t^4 + 6t^3 - 30t^2 - 16t + 32)}{-4t^4 + 10t^3 + t^2 - 24t + 16} - (2 + 10t - t^2) \right]^2 - [2(1 - 2t)\sqrt{1 - 2t}]^2 \\ & = \frac{t^5(16t^7 + 96t^6 - 8t^5 - 504t^4 + 393t^3 + 724t^2 - 784t + 192)}{(-4t^4 + 10t^3 + t^2 - 24t + 16)^2} \\ & = \frac{t^5(t + 4)^2(2t - 1)^2(2t + 3)(2t^2 - 5t + 4)}{(-4t^4 + 10t^3 + t^2 - 24t + 16)^2} \geq 0 \end{aligned}$$

for  $0 < t \leq \frac{1}{2}$ . This completes our proof. ■

**Acknowledgments.** The author thanks the editor for his enthusiasm throughout the process of this manuscript. The author also thanks the referee, who has made important comments, helping the author to complete the article.

## References

- [1] L. Bankoff, Solution of Problem 825, *Crux Math.* **10** (1984), no. 5, 168.
- [2] O. Bottema, R. Z. Djordjevic, R. R. Janic, D. S. Mitrinovic, and P. M. Vasic, *Geometric inequalities*, Wolters–Noordhoff Publishing, Groningen, 1969.
- [3] J. Garfunkel, Problem 825, *Crux Math.* **9** (1983), no. 3, 79.
- [4] M. Lukarevski, Problem 11938, *Amer. Math. Monthly* **123** (2016), no. 9, 941.
- [5] M. Lukarevski, A simple proof of Kooi’s inequality, *Math. Mag.* **93** (2020), 225.
- [6] M. Lukarevski and D. S. Marinescu, A refinement of the Kooi’s inequality, Mittenpunkt and applications, *J. Math. Inequal.* **13** (2019), no. 3, 827–832.

- 
- [7] M. Lukarevski, A new look at the fundamental triangle inequality, *Math. Mag.* **96** (2023), 141–149.
- [8] P. Von Finsler and H. Hadwiger, Einige Relationen im Dreieck, *Comment. Math. Helv.* **10** (1937), no. 1, 316–326.
- [9] S. Wu and L. Debnath, Parametrized Garfunkel–Bankoff inequality and improved Finsler–Hadwiger inequality, *Appl. Math. Lett.* **23** (2010), 331–336.
- [10] W. D. Jiang, The Garfunkel–Bankoff inequality and the Finsler–Hadwiger inequality, *Elem. Math.* **78** (2023), 82–84.

Quang Hung Tran  
High School for Gifted Students, Hanoi University of Science  
Vietnam National University, Hanoi, Vietnam  
[tranquanghung@hus.edu.vn](mailto:tranquanghung@hus.edu.vn)