

On split extensions of preordered groups

Maria Manuel Clementino and Carla Ruivo

Abstract. We investigate the behaviour of split extensions in the category OrdGrp of (pre)-ordered groups. Namely we show that the lexicographic order plays a key role on the existence of compatible orders for semidirect products, establishing necessary and sufficient conditions for such existence; we prove that the Split Short Five Lemma holds for (stably) strong split extensions, and identify classes of split extensions which admit a classifier.

1. Introduction

In [10], the authors studied the behaviour of the category OrdGrp of preordered groups and monotone group homomorphisms. They show in particular that, unlike the categories Grp of groups and TopGrp of topological groups, OrdGrp is not protomodular, and, consequently, the Split Short Five Lemma does not hold. This relies essentially on the study of possible orders (as very common in the literature, throughout by *order* we mean preorder) in a semidirect product $X \rtimes_{\varphi} B$ in Grp of two ordered groups X and B so that

$$X \xrightarrow{\langle 1,0 \rangle} X \rtimes_{\varphi} B \xleftarrow[\pi_2]{\langle 0,1 \rangle} B$$

is a split extension in OrdGrp . Calling these orders *compatible*, it is shown in [10] that compatible orders must contain the product order and be contained in the (reverse) *lexicographic order*, and that they may not exist, or there may be plenty of them.

This note complements the study of split extensions presented in [10]. Indeed, we establish necessary and sufficient conditions in order to a compatible order in $X \rtimes_{\varphi} B$ exist, and show that the compatibility of the lexicographic order is essential. In case it exists, we identify both the maximal order, which is the lexicographic order, and the minimal one, which in general does not coincide with the product order.

Strong split extensions, or strong points, are exactly those with minimal order. It is shown that strong points need not to be stable under pullback, and that the Split

Short Five Lemma holds for (stably) strong points. This way we identify a sort of relative protomodularity which is possibly weaker than the notion introduced in [7], but which still guarantees the validity of the Split Short Five Lemma, and, consequently, the reflection of isomorphisms by the corresponding change of base functors, as for protomodularity [1,4].

Finally, we investigate the existence of split extension classifier, showing that this is possible only for special classes of split extensions, as for instance for those that may be identified as ralis (= right adjoints and left inverses) for the Ord-enrichment of OrdGrp considered in [11].

2. Preliminaries

Let OrdGrp be the category of ordered groups and monotone group homomorphisms. By an ordered group we mean a (non-necessarily abelian) group X equipped with an *order* (i.e., a reflexive and transitive) relation \leq such that the group operation (here denoted by $+$) is monotone. We point out that in general the group inversion is not monotone; it is in fact necessarily anti-monotone. The order is completely determined by its *positive cone* $P = \{x \in X; x \geq 0\}$, which is a submonoid of X closed under conjugation. Moreover, for any group X , any submonoid closed under conjugation defines an order on X .

Remark 2.1. Given a subset A of a group X , the least order on X whose positive cone contains A is obtained in two steps: first we consider the closure \hat{A} of A under conjugation and then the closure of \hat{A} under addition, which we denote by $\langle A \rangle$.

The category OrdGrp has both an algebraic and a topological flavour; indeed, it was shown in [10] that the forgetful functors $\text{OrdGrp} \rightarrow \text{Grp}$ and $\text{OrdGrp} \rightarrow \text{Ord}$ are, respectively, topological and monadic. These functors allow us to construct limits and colimits easily, and are the basis for the categorical study of OrdGrp. Therefore, in order to study the behaviour of split extensions in OrdGrp we start by recalling briefly the behaviour of split extensions in Grp. By *split extension* we mean a short exact sequence

$$(X \xrightarrow{k} A \xrightarrow{f} B),$$

with $k = \ker f$ and $f = \text{coker } k$, where f is a split epimorphism and a splitting of f is given,

$$X \xrightarrow{k} A \xrightleftharpoons[f]{s} B. \tag{2.i}$$

A morphism between split extensions is a triple (a, b, c) making the following diagram commutative:

$$\begin{array}{ccccc}
 X & \xrightarrow{k} & A & \xleftarrow{s} & B \\
 a \downarrow & & \downarrow b & \xleftarrow{f} & \downarrow c \\
 X' & \xrightarrow{k'} & A' & \xleftarrow{s'} & B' \\
 & & & \xleftarrow{f'} &
 \end{array} \tag{2.ii}$$

so that $k' \cdot a = b \cdot k$, $f' \cdot b = c \cdot f$, and $s' \cdot c = b \cdot s$.

It is well known that every split extension in Grp is isomorphic to one given by a *semidirect product*, i.e., A is necessarily isomorphic to the group $X \rtimes_{\varphi} B$ having as underlying set the cartesian product $X \times B$ and, for $(x, b), (x', b')$ in $X \times B$,

$$(x, b) + (x', b') = (x + \varphi(b, x'), b + b'),$$

where $\varphi: B \times X \rightarrow X$ is an *action* of B on X (so that $\varphi(0, x) = x$, $\varphi(b, x + x') = \varphi(b, x) + \varphi(b, x')$, $\varphi(b', \varphi(b, x)) = \varphi(b' + b, x)$). This induces an isomorphism of split extensions

$$\begin{array}{ccccc}
 X & \xrightarrow{k} & A & \xleftarrow{s} & B \\
 \parallel & & \downarrow \theta & \xleftarrow{f} & \parallel \\
 X & \xrightarrow{\langle 1, 0 \rangle} & X \rtimes_{\varphi} B & \xleftarrow{\pi_2} & B,
 \end{array}$$

with $\theta(a) = (k^{-1}(a - sf(a)), f(a))$ for every $a \in A$, and $\varphi_b(x) = \varphi(b, x) = s(b) + k(x) - s(b)$ (see for instance [9, Section 4.1] for details).

In Grp, split extensions with given kernel have a *classifier*, in the sense that the category of split extensions with kernel X has a terminal object, i.e., there exists a split extension with kernel X

$$X \longrightarrow X \rtimes \text{AUT}(X) \xleftarrow{\quad} \text{AUT}(X) \tag{2.iii}$$

such that, for each split extension (2.i) there exists exactly one morphism (a, b, c) from (2.i) to (2.iii) with $a = \text{id}_X$; here $\text{AUT}(X)$ is the group

$$\{\alpha: X \rightarrow X; \alpha \text{ is an automorphism}\},$$

with the operation given by composition, and the addition on the semidirect product $X \rtimes \text{AUT}(X)$ given by

$$(x, \alpha) + (x', \alpha') = (x + \alpha(x'), \alpha \cdot \alpha').$$

The claimed morphism of split extensions is then given by

$$\begin{array}{ccccc}
 X & \xrightarrow{k} & A & \xleftarrow{s} & B \\
 \parallel & & \downarrow \theta & \xrightarrow{f} & \parallel \\
 X & \xrightarrow{\langle 1,0 \rangle} & X \rtimes_{\varphi} B & \xleftarrow{\pi_2} & B \\
 \parallel & & \downarrow 1 \times \bar{\varphi} & & \downarrow \bar{\varphi} \\
 X & \xrightarrow{\langle 1,0 \rangle} & X \rtimes \text{AUT}(X) & \xleftarrow{\pi_2} & \text{AUT}(X)
 \end{array} \tag{2.iv}$$

where $\bar{\varphi}(b) = \varphi_b$.

Alternatively one also says that in Grp *actions are representable*, since this property can be stated as *representability of a functor* into Set (see [2, 3] for details).

3. Compatible orders

Throughout this section

$$(X, P_X) \xrightarrow{\langle 1,0 \rangle} X \rtimes_{\varphi} B \xleftarrow[\pi_B]{\langle 0,1 \rangle} (B, P_B) \tag{3.i}$$

is a split extension in Grp, and X and B are ordered groups, with positive cones P_X and P_B respectively. As shown in [10] there may be no – or there may be plenty of – orders in $X \rtimes_{\varphi} B$ making (3.i) a split extension in OrdGrp. Here we will identify exactly those split extensions (3.i) for which there exist compatible orders. For that we make use of

- the *product order*, with positive cone $P_{\text{prod}} = P_X \times P_B$, and
- the *lexicographic order*, with positive cone

$$P_{\text{lex}} = \{(x, b) \in X \times B; b > 0 \text{ or } (b \sim 0 \text{ and } x \geq 0)\}.$$

We point out that these orders need not to be compatible in (3.i), and that the definition of lexicographic order above is not the one given in [10], where the authors neglected in the definition of P_{lex} the possible existence of elements $b \sim 0$ in B . In fact, the right formulation of P_{lex} led to improvements of the results, on the lexicographic order, of [10, Section 5] we present in the sequel.

Proposition 3.1. *For a positive cone P in $X \rtimes_{\varphi} B$, the following conditions are equivalent:*

- (i) P is compatible in (3.i);
- (ii) $P_{\text{prod}} \subseteq P \subseteq P_{\text{lex}}$.

Proof. (i) \Rightarrow (ii): Monotonicity of $X \xrightarrow{\langle 1,0 \rangle} X \rtimes_{\varphi} B \xleftarrow{\langle 0,1 \rangle} B$ implies $(x, 0) \in P$ and $(0, b) \in P$ when $x \in P_X$ and $b \in P_B$, and therefore $(x, b) = (x, 0) + (0, b) \in P$, that is $P_{\text{prod}} \subseteq P$.

If $(x, b) \in P$ then necessarily $b \geq 0$ because π_B is monotone. When $b \sim 0$ and $(x, b) \in P$, then $(0, b) \sim (0, 0)$ and so $(x, 0) = (x, b) - (0, b) \in P$, which implies $x \geq 0$ because $\langle 1, 0 \rangle: X \rightarrow X \rtimes_{\varphi} B$ is a kernel in OrdGrp .

(ii) \Rightarrow (i): For every $b \in P_B$, $x \in X$, $(x, b) \geq 0$ implies $b \geq 0$, hence π_B is monotone. For every $x \in X$, $(x, 0) \in P$ if and only if $x \in P_X$, and therefore $\langle 1, 0 \rangle: (X, P_X) \rightarrow (X \rtimes_{\varphi} B, P)$ is the kernel of π_B . Finally, if $b \geq 0$ then $(0, b) \in P_{\text{prod}} \subseteq P$, and so $\langle 0, 1 \rangle: (B, P_B) \rightarrow (X \rtimes_{\varphi} B, P)$ is also monotone. ■

The lexicographic order plays an essential role here, since it is compatible as soon as there is a compatible order, as we show next.

Theorem 3.2. *Given (3.i), the following conditions are equivalent:*

- (i) *There is a compatible order in (3.i).*
- (ii) *For every $b \in B$, φ_b is monotone, and, if $b \sim 0$ then $\varphi_b \sim \text{id}$ (pointwise).*
- (iii) *The lexicographic order is compatible in (3.i).*

Proof. (i) \Rightarrow (ii): Let P be a compatible positive cone. If $x \geq 0$, then, for any $b \in B$, $(\varphi_b(x), 0) = (0, b) + (x, 0) - (0, b) \in P$, that is $\varphi_b(x) \geq 0$. Now let $b \sim 0$ in B . Then, for every $x \in X$, $(x - \varphi_b(x), b) = (x, 0) + (0, b) - (x, 0) \in P$ and so $x \geq \varphi_b(x)$; this, together with $-x \geq \varphi_b(-x) = -\varphi_b(x)$ gives $x \sim \varphi_b(x)$.

(ii) \Rightarrow (iii): We need to prove that P_{lex} is a positive cone, that is, it is closed under addition and conjugation. If $(x, b), (x', b') \in P$ and $b > 0$ or $b' > 0$, then, obviously, $(x, b) + (x', b') \in P$; if both $b \sim 0$ and $b' \sim 0$, then in

$$(x, b) + (x', b') = (x + \varphi_b(x'), b + b')$$

we have $b + b' \sim 0$ and $x + \varphi_b(x') \geq 0$ because both x and $\varphi_b(x')$ are positive. Now let $(x, b) \in P$ and $(y, a) \in X \rtimes_{\varphi} B$. Then

$$\begin{aligned} (y, a) + (x, b) - (y, a) &= (y, a) + (x, b) - (0, a) - (y, 0) \\ &= (y + \varphi_a(x) - \varphi_{a+b-a}(y), a + b - a); \end{aligned}$$

if $b > 0$, then $a + b - a > 0$ and so the pair above belongs to P_{lex} ; if $b \sim 0$, then $x \geq 0$ and $a + b - a \sim 0$, and so $y + \varphi_a(x) - \varphi_{a+b-a}(y) \sim y + \varphi_a(x) - y \geq 0$.

(iii) \Rightarrow (i) is trivial. ■

Corollary 3.3. *If the order in B is antisymmetric, then there is a compatible order in (3.i) if and only if φ_b is monotone for every $b \in B$.*

We may consider now the set \mathcal{P} of compatible positive cones for (3.i). We have just shown that either \mathcal{P} is empty or it has a top element, P_{lex} , when ordered by inclusion. A split extension (3.i) where $X \rtimes_{\varphi} B$ has the lexicographic order will be called *maximal*.

Proposition 3.4. *Either $\mathcal{P} = \emptyset$ or \mathcal{P} is a complete lattice.*

Proof. It is easily checked that the meet of compatible orders is a compatible order. ■

Hence, if $\mathcal{P} \neq \emptyset$, there is a least compatible order, which we call *minimal* and describe next.

Proposition 3.5. *Let $\mathcal{P} \neq \emptyset$.*

- (1) *The positive cone of the least compatible order for (3.i) is $\langle P_{\text{prod}} \rangle$, that is, the one generated by $P_X \times P_B$.*
- (2) *Moreover, it coincides with P_{prod} if and only if $\varphi_b \sim \text{id}$ for every positive element b of B .*

Proof. (1) is obvious. To show (2) we use [10, Proposition 5.2], which assures that P_{prod} is compatible if and only if $\varphi_b(x) \geq x$ for all $b \in P_B$ and $x \in X$. But this, together with $\varphi_b(-x) \geq -x$ gives $\varphi_b \sim \text{id}$ as claimed. ■

Remark 3.6. On one hand there are examples of (3.i) with no compatible order, like for instance

$$(\mathbb{Z}, \mathbb{N}) \xrightarrow{\langle 1,0 \rangle} \mathbb{Z} \rtimes \mathbb{Z} \xleftarrow[\pi_2]{\langle 0,1 \rangle} (\mathbb{Z}, \mathbb{Z})$$

with $\varphi_b(x) = (-1)^b x$, since φ_b is not monotone. On the other hand, [10, Example 5.8] shows that there may be plenty of possible positive cones, even in the case when both X and B are abelian and have antisymmetric orders: it is shown there that, for the split extension

$$(\mathbb{Z}, \mathbb{N}) \xrightarrow{\langle 1,0 \rangle} \mathbb{Z} \times \mathbb{Z} \xleftarrow[\pi_2]{\langle 0,1 \rangle} (\mathbb{Z}, \mathbb{N}),$$

\mathcal{P} is an uncountable set. Indeed, a positive cone P on $\mathbb{Z} \times \mathbb{Z}$ can be determined by a family of sets $(X_j = \{n \in \mathbb{Z}; (n, j) \in P\})_{j \in \mathbb{Z}}$, with $X_j = \emptyset$ if $j < 0$ and $X_j = \uparrow -x_n \subseteq \mathbb{Z}$ for $j \geq 0$ so that $x_0 = 0$ and $(x_n)_{n \in \mathbb{N}}$ is a sequence in $\mathbb{N} \cup \{\infty\}$ such that $x_{n+m} \geq x_n + x_m$.

Next we present a characterization of compatible orders inspired by the latter example. We say that family of subsets $(X_b)_{b \in B}$ of X is compatible if $P = \{(x, b); b \in B, x \in X_b\}$ is a compatible positive cone.

Proposition 3.7. *A family $(X_b)_{b \in P_B}$ of subsets of X is compatible if and only if it satisfies the following conditions:*

- (1) $X_b \neq \emptyset \Leftrightarrow b \in P_B \Leftrightarrow 0 \in X_b$;
- (2) $X_0 = P_X$;
- (3) $(\forall b, b' \in P_B) X_b + \varphi_b(X_{b'}) \subseteq X_{b+b'}$;
- (4) $(\forall a \in B) (\forall b \in P_B) (\forall x \in X) x + \varphi_a(X_b) \subseteq X_{a+b-a} + \varphi_{a+b-a}(x)$.

Proof. (1) and (2) guarantee that π_B is monotone and the order of X is inherited from the order of $X \rtimes_{\varphi} B$, while (3) and (4) are equivalent to the closure of P under addition and conjugation, respectively. ■

Remark 3.8. Although the previous result is just a way of formulating the closure under addition and conjugation of P , it gives interesting information on the behaviour of X_b under the action φ_a . Indeed, one concludes that, if $(X_b)_{b \in B}$ is compatible, then

$$(\forall a \in B) (\forall b \in P_B) \varphi_a(X_b) = X_{a+b-a}.$$

This gives that, in particular, for all $b \in P_B$, $\varphi_b(X_b) = X_b$, and that, for conjugate positive elements b, b' of B , X_b is isomorphic to $X_{b'}$, via the action φ .

4. On \mathcal{S} -protomodularity

The existence of different compatible orders shows that the Split Short Five Lemma fails in OrdGrp , and so this category is not protomodular. One may then ask whether OrdGrp is \mathcal{S} -protomodular (cf. [7, Definition 3.1]), for a suitable class \mathcal{S} of split extensions, which in this context are usually called *points*, due to the fact that a split epimorphism $f: A \rightarrow B$ in a category \mathcal{C} , together with its splitting $s: B \rightarrow A$, is nothing but a morphism from the terminal object $\text{id}: B \rightarrow B$ into $f: A \rightarrow B$ in the slice category \mathcal{C}/B over B , which we will refer to as a *point over B* . We will denote by $\text{Pt}(B)$ the category of points $(f: A \rightarrow B, s: B \rightarrow A)$ over B where a morphism $h: (f, s) \rightarrow (f', s')$ is a morphism $h: A \rightarrow A'$ in \mathcal{C} such that $f' \cdot h = f$ and $h \cdot s = s'$; $\text{Pt}_{\mathcal{S}}(B)$ is its full subcategory of points in \mathcal{S} .

We recall that a split extension

$$X \xrightarrow{k} A \begin{array}{c} \xleftarrow{s} \\ \xrightarrow{f} \end{array} B \tag{4.i}$$

or, equivalently, a point (f, s) with kernel k , is *strong* if k and s are jointly strongly epimorphic. It is *stably strong* if every pullback of it along any morphism $g: C \rightarrow B$ is a strong point.

As in every protomodular category, in Grp every point is strong, hence also stably strong, but that is not the case in OrdGrp. Below we identify the strong points in OrdGrp. Also, one may wonder whether they are related to the points (f, s) such that (f, s) is a *rali* (f is right adjoint and left inverse to s) with respect to the Ord-enrichment of OrdGrp studied in [11]: given two morphisms $g, h: X \rightarrow Y$ in OrdGrp, $g \leq h$ if $g(x) \leq h(x)$ for every *positive* element x of X .

Lemma 4.1. *Given a point (4.i) in OrdGrp, where we identify A with $X \rtimes_{\varphi} B$ as usual, and P is its positive cone,*

- (1) (f, s) is a *rali* if and only if $P = P_{\text{prod}}$;
- (2) (f, s) is strong if and only if P is minimal (i.e., P is generated by P_{prod}).

Proof. (1) Assume that (f, s) is a *rali*, i.e., $f \cdot s = \text{id}_B$ and $s \cdot f \leq \text{id}_A$. Then, for every $(x, b) \in P$, $(0, b) = s(f(x, b)) \leq (x, b)$, and therefore $(0, 0) \leq (x, 0)$, which is equivalent to $x \geq 0$ since k is an extremal monomorphism.

(2) Since (f, s) is always strong as a point in Grp, we only have to show that, if k and s factor through a bijective morphism, then it is an isomorphism, and this is easily seen to be the case exactly when P is minimal. ■

Remark 4.2. As a side remark we mention that a point (4.i) is a *rali* exactly when the split extension

$$P_X \xrightarrow{\langle 1, 0 \rangle} P \xleftarrow[\pi_2]{\langle 0, 1 \rangle} P_B$$

is a Schreier point (see [6]) in the category of monoids.

In [10, Proposition 6.2], it is shown that OrdGrp is \mathcal{S} -protomodular when \mathcal{S} is the class of split extensions with the product order, i.e., *rali* points. Here we analyse whether there is a larger class \mathcal{S} of points that makes OrdGrp an \mathcal{S} -protomodular category. Denoting the classes of *rali*, strong and stably strong points by *Rali*, *Strong* and *Strong** respectively, we know that

$$\text{Rali} \subseteq \text{Strong}^* \subseteq \text{Strong} \quad \text{and that we must have} \quad \text{Rali} \subseteq \mathcal{S} \subseteq \text{Strong}^* .$$

First we show that the two inclusions on the left are strict.

Proposition 4.3. (1) *Strong points are not stable under pullback.*

(2) *There is a stably strong point which is not a rali.*

Proof. (1) With \mathbb{Z}_0 and \mathbb{Z} the group of integers, respectively with $P = \{0\}$ and with the usual order, and $\varphi: \mathbb{Z}_0 \times \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $\varphi_1(x) = -x$, consider the strong point

$$\mathbb{Z}_0 \xrightarrow{\langle 1, 0 \rangle} \mathbb{Z}_0 \rtimes_{\varphi} \mathbb{Z} \xleftarrow[\pi_2]{\langle 0, 1 \rangle} \mathbb{Z};$$

that is, $\mathbb{Z}_0 \rtimes_{\varphi} \mathbb{Z}$ is equipped with the minimal order. This point is not stably strong, as we show next.

Indeed, for any strong point (4.i) such that there exists $b \geq 0$ with $\varphi_b \not\geq \text{id}$ but $\varphi_b^m \geq \text{id}$ for some natural number $m \neq 0$ (here by φ_b^m we mean φ_b computed m times), we may consider its pullback along $f: \mathbb{Z} \rightarrow B$ with $f(n) = nmb$,

$$\begin{array}{ccc} X \rtimes \mathbb{Z} & \longrightarrow & X \rtimes_{\varphi} B \\ \langle 0,1 \rangle \updownarrow \pi_2 & & \langle 0,1 \rangle \updownarrow \pi_2 \\ \mathbb{Z} & \xrightarrow{f} & B. \end{array}$$

Then the action of the point on the left is given by φ_b^m , hence the product order is the minimal compatible order in $X \rtimes \mathbb{Z}$. However, this point is not ordered by the product order: for $x \in X$ such that $\varphi_b(x) \not\geq x$ one has, since $b \geq 0$,

$$(-x, 0) + (0, b) + (x, 0) + (0, (m - 1)b) = (-x + \varphi_b(x), mb) \geq 0;$$

hence $(-x + \varphi_b(x), 1)$ is positive in $X \rtimes \mathbb{Z}$ although $-x + \varphi_b(x)$ is not positive, by assumption.

(2) Consider now the point

$$\mathbb{Q} \xrightarrow{\langle 1,0 \rangle} \mathbb{Q} \rtimes_{\varphi} \mathbb{Z} \xleftarrow[\pi_2]{\langle 0,1 \rangle} \mathbb{Z} \tag{4.ii}$$

with $\varphi_n(x) = 2^n x$ (hence monotone for every $n \in \mathbb{Z}$), equipped with the minimal order, and let $g: A \rightarrow \mathbb{Z}$ be any morphism in OrdGrp. In its pullback $\mathbb{Q} \rtimes_{\psi} A$ as in the diagram

$$\begin{array}{ccc} \mathbb{Q} \rtimes_{\psi} A & \longrightarrow & \mathbb{Q} \rtimes_{\varphi} \mathbb{Z} \\ \downarrow & & \downarrow \\ A & \xrightarrow{g} & \mathbb{Z}, \end{array}$$

the positive cone is given by $P = \{(x, a); a \geq 0 \text{ and } (x, g(a)) \geq 0\}$, and the action ψ is given by $\psi_a(x) = \varphi_{g(a)}(x)$. Let us check that $\mathbb{Q} \rtimes_{\psi} A$ has the minimal order: if $(x, a) \geq 0$ and $g(a) = 0$, then both a and x are positive; if $(x, a) \geq 0$ and $g(a) \neq 0$, then, for $r = \frac{x}{1-2^{g(a)}}$ one gets, by closure of P under conjugation,

$$(r, 0) + (0, a) - (r, 0) = (r - \psi_a(r), a) = (r - \varphi_{g(a)}(r), a) = (x, a) \in P.$$

Therefore, every pullback of the point (4.ii) is a strong point, as claimed. ■

Lemma 4.4. *The class Strong* of stably strong points in OrdGrp is closed under finite products in the category of points.*

Proof. Strong* contains the terminal object and is closed under binary products because they commute with pullbacks. ■

Theorem 4.5. *Let $\mathcal{S} = \text{Strong}^*$ be the class of stably strong points in OrdGrp. Then:*

- (1) *The Split Short Five Lemma holds with respect to \mathcal{S} ; that is, for any commutative diagram*

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & X & \xrightarrow{k} & A & \xrightleftharpoons[s]{s} & B & \longrightarrow & 0 \\
 & & \downarrow a & & \downarrow b & & \downarrow c & & \\
 0 & \longrightarrow & X' & \xrightarrow{k'} & A' & \xrightleftharpoons[f']{s'} & B' & \longrightarrow & 0
 \end{array} \tag{4.iii}$$

in the sense that $b \cdot k = k' \cdot a$, $c \cdot f = f' \cdot b$ and $b \cdot s = s' \cdot c$, if the rows are split extensions belonging to \mathcal{S} and a and c are isomorphisms, then b is an isomorphism as well.

- (2) *For any morphism $h: Y \rightarrow B$, the change of base functor $h^*: \text{Pt}_{\mathcal{S}}(B) \rightarrow \text{Pt}_{\mathcal{S}}(Y)$ is conservative.*

Proof. (1) Given diagram (4.iii), we know that b is an isomorphism of groups, because the Split Short Five Lemma holds in Grp. Since both orders are minimal, b is in fact an isomorphism in OrdGrp.

(2) Adapting the classical proof that the change of base functor between points is conservative provided that the Split Short Five Lemma holds, it is enough to observe that the change of base functor between points restricts to $\text{Pt}_{\mathcal{S}}$ since \mathcal{S} is pullback stable. ■

Remark 4.6. We point out that our argument for the validity of the Split Short Five Lemma can also be applied to the (larger) class Strong of strong points.

It is an open problem to know whether Strong* is stable under equalizers in the category of points, and consequently whether OrdGrp is Strong*-protomodular in the sense of [6, Definition 8.1.1], [7, Definition 3.1], although the previous theorem shows that OrdGrp has the desired properties for relative protomodularity with respect to Strong*. Moreover, OrdGrp is Strong*-protomodular in the sense of [5, Definition 8.5], where the author only imposes that \mathcal{S} is a stable class of strong points, and therefore this weaker notion does not assure that the change of base functor, restricted to \mathcal{S} , is conservative (hence neither the Split Short Five Lemma). To assure that the change of base functor, restricted to $\text{Pt}_{\mathcal{S}}$, is conservative, the authors of [7] impose that \mathcal{S} is stable under equalizers in the category of points. However, this property seems quite complicated to check, as it is the case of our example and, for instance, of the class of points studied in [12, Section 7.14].

Henceforth, we propose the following definition, which is in fact a translation of the notion of (absolute) protomodularity [1, Definition 3.1.3], not making use of equalizers but instead focussing on the key property of protomodularity.

Definition 4.7. If \mathcal{S} is a class of strong points of the category \mathcal{C} , \mathcal{C} is said to be *protomodular with respect to \mathcal{S}* if

- (1) \mathcal{C} has pullbacks of points in \mathcal{S} along any morphism, which belong also to \mathcal{S} .
- (2) For any morphism $h: Y \rightarrow B$, the change of base functor $h^*: \text{Pt}_{\mathcal{S}}(B) \rightarrow \text{Pt}_{\mathcal{S}}(Y)$ reflects isomorphisms.

Then we can compare these notions using [7, Proposition 3.2]:

Theorem 4.8. *If \mathcal{C} is \mathcal{S} -protomodular in the sense of [7], then the change of base functor is conservative when restricted to $\text{Pt}_{\mathcal{S}}$, and so \mathcal{C} is protomodular with respect to \mathcal{S} .*

5. On the existence of \mathcal{S} -classifiers

Unlike the category TopGrp of topological groups (see [8]), OrdGrp has no split extension classifiers, as we show in Theorem 5.7. Still, it is interesting to analyse the existence, in OrdGrp , of split extension classifiers for *special classes of points*, as we discuss in this last section.

Given a class \mathcal{S} of split extensions in OrdGrp , we denote by \mathcal{S}_X the category of split extensions in \mathcal{S} with kernel X with morphisms triples (a, b, c) as in (2.ii) with $a = \text{id}$.

Definition 5.1. If \mathcal{S} is a class of split extensions in OrdGrp , we say that OrdGrp has *\mathcal{S} -classifiers* if the category \mathcal{S}_X has a terminal object; that is, for every ordered group X there exists a split extension with kernel X

$$X \xrightarrow{\langle 1,0 \rangle} X \rtimes A(X) \xleftarrow[\pi_2]{\langle 0,1 \rangle} A(X) \tag{5.i}$$

in \mathcal{S} such that, for each split extension in \mathcal{S} with kernel X , there exists exactly one morphism in \mathcal{S}_X from it into (5.i).

Given an ordered group X , let $\text{Aut}_P(X)$ be the group

$$\text{Aut}(X) = \{\alpha: X \rightarrow X; \alpha \text{ is a monotone automorphism}\}$$

equipped with an order with positive cone P . Then, by Theorem 3.2, there is an order in $X \rtimes \text{Aut}(X)$ making

$$X \xrightarrow{\langle 1,0 \rangle} X \rtimes \text{Aut}_P(X) \xleftarrow[\pi_2]{\langle 0,1 \rangle} \text{Aut}_P(X) \tag{5.ii}$$

a split extension if and only if when $\alpha \sim \text{id}$ in $\text{Aut}_P(X)$ also $\alpha \sim \text{id}$ pointwise in X . We call such orders in $\text{Aut}(X)$ *admissible*.

Proposition 5.2. *Let \mathcal{S} be a class of split extensions, X an ordered group and P an admissible positive cone in $\text{Aut}(X)$ such that (5.ii) belongs to \mathcal{S} . If*

$$X \xrightarrow{\langle 1,0 \rangle} X \rtimes A(X) \xleftarrow[\pi_2]{\langle 0,1 \rangle} A(X)$$

is a classifier for \mathcal{S}_X , then $A(X)$ is isomorphic, as a group, to $\text{Aut}(X)$, and its positive cone P_A contains P .

Proof. By definition of classifiers, there exists a unique morphism γ making the following diagram commute:

$$\begin{CD} X @>\langle 1,0 \rangle>> X \rtimes \text{Aut}_P(X) @<<[\pi_2]{\langle 0,1 \rangle}<< \text{Aut}_P(X) \\ @| @VVV @VV\gamma V \\ X @>\langle 1,0 \rangle>> X \rtimes A(X) @<<[\pi_2]{\langle 0,1 \rangle}<< A(X). \end{CD}$$

On the other hand, being a split extension in Grp, there is a group homomorphism $\psi: A(X) \rightarrow \text{Aut}(X)$ as in diagram (2.iv). Since we are in OrdGrp, the image of ψ has only monotone automorphisms, i.e., ψ factors through $\text{Aut}(X)$ as $\bar{\psi}: A(X) \rightarrow \text{Aut}(X)$. Clearly $\bar{\psi} \cdot \gamma = \text{id}$ and $\gamma \cdot \bar{\psi} = \text{id}$, hence we may assume that $A(X)$, as a group, is $\text{Aut}(X)$; moreover, from the monotonicity of γ it follows that its positive cone P_A contains the positive cone P of $\text{Aut}_P(X)$. ■

Theorem 5.3. *For every ordered group X , let $\tilde{P} = \{\alpha; \alpha(x) \sim x \text{ for all } x \in X\}$. The rali point*

$$X \xrightarrow{\langle 1,0 \rangle} X \rtimes \text{Aut}_{\tilde{P}}(X) \xleftarrow[\pi_2]{\langle 0,1 \rangle} \text{Aut}_{\tilde{P}}(X) \tag{5.iii}$$

is a terminal object of Rali_X .

Proof. First of all it is easy to check that the product order in $X \rtimes \text{Aut}_{\tilde{P}}(X)$, which in fact coincides with the lexicographic order, is compatible in (5.iii). So (5.iii) is a rali point.

Given a rali point

$$X \rtimes_{\varphi} B \xleftarrow[\pi_2]{\langle 0,1 \rangle} B$$

with kernel X , we know that there is a unique group homomorphism $\bar{\varphi}: B \rightarrow \text{Aut}(X)$, which factors through $\text{Aut}(X)$, making diagram (2.iv) commute. Moreover, its co-restriction $\tilde{\varphi}: B \rightarrow \text{Aut}_{\tilde{P}}(X)$ is monotone – since $X \rtimes_{\varphi} B$ has the product order, by Proposition 3.5 (2), $\varphi_b \sim \text{id}$ for every $b \geq 0$ – and $1 \times \tilde{\varphi}: X \rtimes_{\varphi} B \rightarrow X \rtimes \text{Aut}_{\tilde{P}}(X)$ is also clearly monotone. Uniqueness of $\bar{\varphi}$ guarantees uniqueness of $\tilde{\varphi}$. ■

Corollary 5.4. *OrdGrp has Rali-classifiers.* ■

This result can be naturally extended in the following way.

Proposition 5.5. *Let X be an ordered group and P an admissible positive cone in $\text{Aut}(X)$. Then the maximal point*

$$X \xrightarrow{\langle 1,0 \rangle} X \rtimes \text{Aut}_P(X) \xleftarrow[\pi_2]{\langle 0,1 \rangle} \text{Aut}_P(X)$$

classifies the class \mathcal{S}_X of split extensions $X \rtimes_{\varphi} B \xleftarrow[\pi_2]{\langle 0,1 \rangle} B$ with kernel X such that

- (1) *for all $b \in P_B$, $\varphi_b \in P$;*
- (2) *for all $x \in X$, $x \geq 0$ provided that there exists $b \in P_B$ with $(x, b) \geq 0$ in $X \rtimes_{\varphi} B$ and $\varphi_b \sim \text{id}$ in $\text{Aut}_P(X)$.*

Proof. Given a point in \mathcal{S}_X , we know that there is a unique group homomorphism $\bar{\varphi}: B \rightarrow \text{Aut}(X)$ making the diagram

$$\begin{array}{ccccc} X & \xrightarrow{\langle 1,0 \rangle} & X \rtimes_{\varphi} B & \xleftarrow[\pi_2]{\langle 0,1 \rangle} & B \\ \parallel & & \downarrow 1 \times \bar{\varphi} & & \downarrow \bar{\varphi} \\ X & \xrightarrow{\langle 1,0 \rangle} & X \rtimes \text{Aut}_P(X) & \xleftarrow[\pi_2]{\langle 0,1 \rangle} & \text{Aut}_P(X) \end{array}$$

commute. Then condition (1) guarantees that $\bar{\varphi}$ is monotone, and (2) gives monotonicity of $1 \times \bar{\varphi}$. ■

Remark 5.6. Given an ordered group X such that, for every $x \in X$, either $x \geq 0$ or $-x \geq 0$, consider in $\text{Aut}(X)$ the admissible order defined by

$$P^+ = \{\alpha; (\forall x \in P_X) \alpha(x) \geq x\}.$$

We denote by $\text{Aut}^+(X)$ this ordered group and by \mathcal{S}^+ the class of split extensions defined as in the proposition above, that is, \mathcal{S}^+ is the union of \mathcal{S}_X^+ for all such ordered groups X . The split extensions classified by $\text{Aut}^+(X)$ depend very much on the action φ . For instance, $\text{Aut}^+(\mathbb{Q})$ classifies a split extension

$$\mathbb{Q} \xrightarrow{\langle 1,0 \rangle} \mathbb{Q} \rtimes_{\varphi} \mathbb{Z} \xleftarrow[\pi_2]{\langle 0,1 \rangle} \mathbb{Z}$$

when $\varphi = \text{id}$ only if it is a rali, while when $\varphi_n(x) = 2^n x$ it classifies any such split extension.

We could use analogously, for example, the order in $\text{Aut}(X)$ defined by $P^- = \{\alpha; (\forall x \leq 0) \alpha(x) \geq x\}$.

Theorem 5.7. *OrdGrp has no split extensions classifiers.*

Proof. For each ordered group X , assume that $X \rtimes \text{Aut}_P(X) \xrightleftharpoons[\pi_2]{(0,1)} \text{Aut}_P(X)$ classifies the split extensions with kernel X . As we remarked above, when X is an ordered group such that each $x \in X$ is either positive or negative, both P^+ and P^- are admissible positive cones in $\text{Aut}(X)$. Hence, both P^+ and P^- are contained in P . Since $\alpha \in P^+$ if and only if its inverse $\alpha^{-1} \in P^-$, this shows that if $\alpha \in P^+$ then $\alpha \sim \text{id}$ in P , and so $\alpha \sim \text{id}$ pointwise in X because P must be admissible. But this is not true in general: for instance, for $X = \mathbb{Q}$ and n a natural number larger than 1, $\alpha(x) = nx \in P^+$ but pointwise $\alpha \not\sim \text{id}$. ■

Acknowledgements. The results of this paper were obtained while the first author was a tutor of the second author under the programme *Novos Talentos em Matemática* from the Gulbenkian Foundation.

We thank Graham Manuell for valuable discussions on the subject of this paper. We also thank Diana Rodelo, Andrea Montoli and Manuela Sobral for useful discussions on the concept of \mathcal{S} -protomodularity.

Funding. This work was partially supported by the Centre for Mathematics of the University of Coimbra, UIDB/00324/2020, funded by the Portuguese Government through FCT/MCTES, and a grant from the Gulbenkian Foundation programme *Novos Talentos em Matemática*

References

- [1] F. Borceux and D. Bourn, *Mal'cev, protomodular, homological and semi-abelian categories*. Math. Appl. 566, Kluwer Academic Publishers, Dordrecht, 2004 Zbl 1061.18001 MR 2044291
- [2] F. Borceux, G. Janelidze, and G. M. Kelly, Internal object actions. *Comment. Math. Univ. Carolin.* **46** (2005), no. 2, 235–255 Zbl 1121.18004 MR 2176890
- [3] F. Borceux, G. Janelidze, and G. M. Kelly, On the representability of actions in a semi-abelian category. *Theory Appl. Categ.* **14** (2005), No. 11, 244–286 Zbl 1103.18006 MR 2182676
- [4] D. Bourn, *Normalization equivalence, kernel equivalence and affine categories*. In *Category theory (Como, 1990)*, pp. 43–62, Lecture Notes in Math. 1488, Springer, Berlin, 1991 Zbl 0756.18007 MR 1173004
- [5] D. Bourn, Partial Mal'tsevness and partial protomodularity. 2015, arXiv:1507.02886
- [6] D. Bourn, N. Martins-Ferreira, A. Montoli, and M. Sobral, *Schreier split epimorphisms in monoids and in semirings*. Textos Mat. Sér. B 45, Universidade de Coimbra, Departamento de Matemática, Coimbra, 2013 Zbl 1294.18001 MR 3157484

- [7] D. Bourn, N. Martins-Ferreira, A. Montoli, and M. Sobral, [Monoids and pointed \$S\$ -protomodular categories](#). *Homology Homotopy Appl.* **18** (2016), no. 1, 151–172
Zbl [1350.18019](#) MR [3485342](#)
- [8] F. Cagliari and M. M. Clementino, [Topological groups have representable actions](#). *Bull. Belg. Math. Soc. Simon Stevin* **26** (2019), no. 4, 519–526 Zbl [1432.54040](#) MR [4042398](#)
- [9] M. M. Clementino, [An invitation to topological semi-abelian algebras](#). In *New perspectives in algebra, topology and categories*, pp. 27–66, Coimbra Math. Texts 1, Springer, Cham, 2021 Zbl [1480.18008](#) MR [4367549](#)
- [10] M. M. Clementino, N. Martins-Ferreira, and A. Montoli, [On the categorical behaviour of preordered groups](#). *J. Pure Appl. Algebra* **223** (2019), no. 10, 4226–4245
Zbl [1468.18012](#) MR [3958089](#)
- [11] M. M. Clementino, A. Montoli, and D. Rodelo, [On lax protomodularity for Ord-enriched categories](#). *J. Pure Appl. Algebra* **227** (2023), no. 8, article no. 107348 MR [4549004](#)
- [12] A. Montoli, D. Rodelo, and T. Van der Linden, [Two characterisations of groups amongst monoids](#). *J. Pure Appl. Algebra* **222** (2018), no. 4, 747–777 Zbl [1386.18033](#)
MR [3720851](#)

Received 13 December 2022; revised 2 April 2023.

Maria Manuel Clementino

CMUC, Department of Mathematics, University of Coimbra, 3000-143 Coimbra, Portugal;
mmc@mat.uc.pt

Carla Ruivo

Department of Mathematics, University of Coimbra, 3000-143 Coimbra, Portugal;
cruivo@proton.me