

MATHEMATISCHES FORSCHUNGSINSTITUT OBERWOLFACH

Report No. 54/2005

## Heat Kernels, Stochastic Processes and Functional Inequalities

Organised by  
Thierry Coulhon (Cergy)  
Bruno Franchi (Bologna)  
Takashi Kumagai (RIMS, Kyoto)  
Karl-Theodor Sturm (Bonn)

November 27th – December 3rd, 2005

ABSTRACT. The conference brought together mathematicians belonging to several fields, essentially analysis, probability and geometry. One of the main unifying topics was certainly the study of heat kernels in various contexts: fractals, manifolds, domains of the Euclidean space, percolation clusters, infinite dimensional spaces, metric measure spaces.

*Mathematics Subject Classification (2000):* 31, 32W, 35J, 46T, 47D, 58J, 60H, 60J.

### Introduction by the Organisers

The workshop *Heat kernels, stochastic processes and functional inequalities* was organized by Thierry Coulhon (Cergy), Bruno Franchi (Bologna), Takashi Kumagai (Kyoto) and Karl-Theodor Sturm (Bonn). It was held from November 27th to December 3rd. The meeting was attended by 56 participants from Australia, Austria, Canada, Finland, France, Germany, Italy, Japan, Poland, Switzerland, United Kingdom, and USA. This workshop was sponsored by the European Union, which allowed the invitation of 18 young people, who contributed positively to the atmosphere of the meeting.

The conference brought together mathematicians belonging to several fields, essentially analysis, probability and geometry. One of the main unifying topics was certainly the study of heat kernels in various contexts: fractals, manifolds, domains of the Euclidean space, percolation clusters, infinite dimensional spaces, metric measure spaces. Some related aspects of geometric analysis were also considered such as  $L^p$ -cohomology and mass transportation. There was a stimulating exchange between probabilistic and analytic points of view, together with a geometric emphasis in most of the problems. We had 5 one hour survey lectures and

21 thirty-five minutes talks. A lot of time was devoted to discussions and exchange of ideas.

Among the highlights were relations between mass transportation, generalized Ricci bounds and contraction properties, connections between heat kernel estimates and percolation clusters, non-linear aspects of diffusions, functional analytic approach to parabolic regularity, geometric and functional analytic aspects of infinite dimensional analysis.

This diversity of topics and mix of participants stimulated many extensive and fruitful discussions. It also helped initiate new collaborations, in particular for the younger researchers.

## Heat Kernels, Stochastic Processes and Functional Inequalities

### Table of Contents

Pascal Auscher	
<i>The Kato square root problem: a review and new developments</i> . . . . .	3065
Martin Barlow	
<i>Heat kernels on percolation clusters</i> . . . . .	3069
Richard Bass (joint with Takashi Kumagai)	
<i>Symmetric Markov chains on <math>Z^d</math> with unbounded range</i> . . . . .	3072
Alexander Bendikov (joint with Th. Coulhon, L. Saloff-Coste)	
<i>Ultracontractivity and embedding into <math>L^\infty</math></i> . . . . .	3073
Krzysztof Burdzy (joint with R. Bass, Z.-Q. Chen)	
<i>On the Robin problem in fractal domains</i> . . . . .	3075
Eric Carlen	
<i>Correlation inequalities of Brascamp-Lieb inequalities and the heat kernel</i>	3078
Gilles Carron (joint with Th. Coulhon, A. Hassell)	
<i><math>L^p</math> cohomology and the boundedness of the Riesz transform</i> . . . . .	3081
Zhen-Qing Chen (joint with Masatoshi Fukushima, Jiengang Ying)	
<i>Boundary Processes of Markov Processes</i> . . . . .	3084
Bruce Driver	
<i>Limit theorems for geometric approximations to Wiener measure</i> . . . . .	3086
David Elworthy (joint with Y. LeJan, X.-M. Li-Hairer)	
<i>Intertwined diffusions</i> . . . . .	3088
Masatoshi Fukushima	
<i>One point extensions of general Markov processes</i> . . . . .	3091
Elton Hsu	
<i>Quasi-invariance of the Wiener Measure on the Loop Space of a</i>	
<i>Riemannian Manifold</i> . . . . .	3092
Moritz Kassmann (joint with R. Hussein and M. Barlow/R. Bass/Z.-Q. Chen)	
<i>Jump processes, <math>\mathcal{L}</math>-harmonic functions, continuity estimates</i> . . . . .	3095
Jun Kigami	
<i>Resistance forms and heat kernel estimates</i> . . . . .	3096
M. Ledoux (joint with D. Bakry)	
<i>A logarithmic Sobolev form of the Li-Yau inequality</i> . . . . .	3097

Pierre Mathieu	
<i>On the range of random walks on percolation clusters</i> . . . . .	3098
Robert J. McCann	
<i>Sharp decay rates for the fastest conservative diffusions</i> . . . . .	3099
Hervé Pajot	
<i>Besov Spaces on Fractals and Conformal Geometry</i> . . . . .	3100
M. Röckner (joint with J.-G. Ren, F.-Y. Wang)	
<i>Stochastic Generalized Porous Media Equations</i> . . . . .	3102
Takashi Shioya (joint with K. Kuwae)	
<i>Variational convergence over metric spaces</i> . . . . .	3104
Masayoshi Takeda	
<i>The principal eigenvalue for time-changed processes and applications</i> . . . .	3105
Anton Thalmaier (joint with Hélène Airault and Paul Malliavin)	
<i>Brownian motion on Jordan curves and stochastic calculus of variations</i> <i>on the diffeomorphism group of the circle</i> . . . . .	3108
Cedric Villani	
<i>Optimal transport, Ricci curvature and synthetic geometry</i> . . . . .	3111
Max-K von Renesse	
<i>Local Poincare Inequality by Transportation</i> . . . . .	3112
Wolfgang Woess (joint with Donald I. Cartwright)	
<i>The spectrum of the averaging operator on a network</i> . . . . .	3113

## Abstracts

### The Kato square root problem: a review and new developments

PASCAL AUSCHER

The following theorem has been proved in [2] in full generality.

Let  $B = B(x)$  be an  $n \times n$  ( $n \geq 1$  is all this paper) matrix of complex,  $L^\infty$  coefficients, defined on  $\mathbb{R}^n$ , and satisfying the ellipticity (or “accretivity”) condition

$$(1) \quad \lambda|\xi|^2 \leq \Re B\xi \cdot \xi^* \text{ and } |B\xi \cdot \zeta^*| \leq \Lambda|\xi||\zeta|,$$

for  $\xi, \zeta \in \mathbb{C}^n$  and for some  $\lambda, \Lambda$  such that  $0 < \lambda \leq \Lambda < \infty$ . Here,  $u \cdot v = u_1v_1 + \dots + u_nv_n$  and  $u^*$  is the complex conjugate of  $u$  so that  $u \cdot v^*$  is the usual inner product in  $\mathbb{C}^n$  and, therefore,  $B\xi \cdot \zeta^* \equiv \sum_{j,k} b_{j,k}(x)\xi_k \bar{\zeta}_j$ . We define a second order divergence form operator

$$(2) \quad Jf \equiv -\operatorname{div}(B\nabla f) = -\sum_{i,j=1}^n \frac{\partial}{\partial x_i} \left( b_{i,j} \frac{\partial}{\partial x_j} \right),$$

which we interpret in the usual weak sense via a sesquilinear form, where  $\nabla$  is the gradient operator and  $\operatorname{div}$  the divergence operator, its negative adjoint.

The accretivity condition (1) enables one to define a square root  $J^{1/2} \equiv \sqrt{J}$ , and a fundamental question is to determine whether one can solve the “Kato square root problem”, i.e. establish the estimate

$$(3) \quad \|\sqrt{J}f\|_2 \sim \|\nabla f\|_2,$$

where  $\sim$  is the equivalence in the sense of norms, with constants  $C$  depending only on  $n$ ,  $\lambda$  and  $\Lambda$ , and  $\|f\|_2 = (\int_{\mathbb{R}^n} |f(x)|_H^2 dx)^{1/2}$  denotes the usual norm for functions on  $\mathbb{R}^n$  valued in a Hilbert space  $H$ .

**Theorem 1.** *For any operator as above the domain of  $\sqrt{J}$  coincides with the Sobolev space  $W^{1,2}(\mathbb{R}^n)$  and  $\|\sqrt{J}f\|_2 \sim \|\nabla f\|_2$ .*

We first mention that an abstract Hilbert space formulation (where  $B$ ,  $\nabla$  are replaced by abstract operators) was disproved by McIntosh, hence this is really a theorem about elliptic differential operators. In fact, harmonic analysis methods involving square function estimates, Carleson measures, maximal functions, variants of the  $T(b)$  theorem and stopping-time arguments are at the heart of the proof. The case  $n = 1$  was proved back to 1982 in a celebrated paper of Coifman, McIntosh and Meyer.

The estimate contained in the statement applies to time-evolution parabolic, hyperbolic or elliptic partial differential equations involving  $J$  as the main elliptic part and where the matrix  $B$  could also depend smoothly on time. We review such applications in this lecture.

Let us concentrate here on one aspect, namely elliptic problems. This estimate also contains the  $L^2$  boundedness of a boundary Neumann to Dirichlet map for the following elliptic problem. Consider the elliptic problem

$$(4) \quad \begin{cases} \partial_t^2 u(x, t) - Ju(x, t) = 0 & \text{in } \mathbb{R}_+^{n+1} = \{(x, t) \in \mathbb{R}^n \times (0, \infty)\} \\ u(x, 0) = f(x) & \text{on } \mathbb{R}^n. \end{cases}$$

Then, a solution is given by the semigroup equation  $u(x, t) = e^{-t\sqrt{J}}f(x)$  and the Neumann data equals  $-\partial_t u(x, 0) = \sqrt{J}f(x)$  so that the Neumann to Dirichlet map alluded to is  $\nabla\sqrt{J}^{-1}$ . Hence, one can solve the Neumann problem with data in  $L^2$  with an estimate on  $\nabla u$  and  $\partial_t u$ . Also this map is invertible, with inverse  $-\sqrt{J}^{-1}\text{div}B$  acting on gradient fields, and one can solve the Dirichlet problem with Dirichlet data  $f$  having gradients in  $L^2$  with an estimate on  $\nabla u$  and  $\partial_t u$  (this is called the regularity problem). This holds for this general class of equations with complex coefficients. Furthermore, one can impose the solution to satisfy some control such as quadratic estimates and then uniqueness can be discussed.

This elliptic problem can be embedded in a larger class of problems where we allow mixed second order derivatives

$$(5) \quad \begin{cases} Lu(x, t) = 0 & \text{in } \mathbb{R}_+^{n+1} \\ u(x, 0) = f(x) & \text{on } \mathbb{R}^n. \end{cases}$$

where, now,

$$L = -\text{div}A\nabla \equiv -\sum_{i,j=1}^{n+1} \frac{\partial}{\partial x_i} \left( a_{i,j}(x) \frac{\partial}{\partial x_j} \right)$$

(we use the notational convention that  $t = x_{n+1}$  and  $\partial/\partial t = \partial/\partial x_{n+1}$  and  $\nabla$ ,  $\text{div}$  now denote the full gradient and divergence operators on  $\mathbb{R}^{n+1}$ ) where  $A = A(x)$  is an  $(n+1) \times (n+1)$  matrix of complex-valued  $L^\infty$  coefficients, defined on  $\mathbb{R}^n$  (i.e., independent of the  $t$  variable), and satisfying the uniform ellipticity condition

$$(6) \quad \lambda|\xi|^2 \leq \Re e A(x)\xi \cdot \xi^*, \quad \|A\|_{L^\infty(\mathbb{R}^n)} \leq \Lambda,$$

for some  $\lambda > 0$ ,  $\Lambda < \infty$ , and for all  $\xi \in \mathbb{C}^{n+1}$ ,  $x \in \mathbb{R}^n$ . The preceding case, we say that  $A$  is of Kato type, corresponds to  $a_{n+1,n+1} = 1$  and the non-diagonal coefficients  $a_{i,n+1} = a_{n+1,i} = 0$  if  $i = 1, \dots, n$ . We refer to the forthcoming articles [1] for references on what follows as well for the statements which are new.

In this situation, there no longer is semigroup structure for solving. The construction of the Neumann to Dirichlet map can be made formally using limits of the layer potentials at the boundary. However, the study of boundedness and invertibility of the Neumann to Dirichlet map becomes much harder in general.

A trivial case is when  $A$  is constant, as this map and its inverse are bounded (on  $L^2(\mathbb{R}^n)$ ) singular integrals of convolution type, that is Calderón-Zygmund operators. For example, if  $A$  is the identity, they are associated to the standard Riesz transforms on  $\mathbb{R}^n$ .

Another situation where things are known is the case where  $A$  has *real and symmetric*  $L^\infty$  coefficients. Then, Jerison and Kenig solved (I will not be precise

in saying in which class) the Dirichlet problem with  $L^2$  data by the method of harmonic measure. Later Kenig and Pipher solved (again, in some class) both the Neumann and regularity problems with datum in  $L^2(\mathbb{R}^n)$  using the previous result of Jerison and Kenig, and Rellich identities. These identities furnish the *a priori* comparison between tangential and normal components of the gradient of any solution at the boundary, namely

$$(7) \quad \int_{\mathbb{R}^n} |\partial_\nu u(x, 0)|^2 dx \sim \int_{\mathbb{R}^n} |\nabla_T u(x, 0)|^2 dx,$$

where  $\nabla_T$  is the tangential gradient, that is the gradient with respect to  $x$  and  $\partial_\nu$  is the exterior conormal derivative, that is  $-e_{n+1} \cdot A \nabla$  with  $e_{n+1}$  the unit upward vector along the  $x_{n+1}$  co-ordinate. Hence, one controls the full gradient of a solution at the boundary provided one controls either its conormal component (Neumann datum) or its tangential component (regularity datum), whence solvability. This method does not rely on boundedness of the Neumann to Dirichlet map. However, if this boundedness could be proved, then invertibility follows directly from the Rellich estimate. Such a boundedness is far from trivial. To indicate why, let us mention that this class of  $L$ 's includes pull-backs of the Laplace operator under a Lipschitz change of coordinate, and thus the class of Neumann to Dirichlet maps includes double layer potentials on Lipschitz graphs from potential theory, whose boundedness is due to Coifman, McIntosh and Meyer in the 1982 paper mentioned before. Then, Verchota understood in 1984 how to invert the Neumann to Dirichlet map incorporating the Rellich identities, so that he could use the method of layer potential to solve both the Neumann and regularity problems.

In the last 20 years some new technology has been developed to reprove the boundedness of the double layer potential, one of them is called the  $T(b)$  theorem, a criterion to check  $L^2$  boundedness for singular integrals (no longer of convolution type). For the purpose of solving the Kato problem and also some other geometric problems in complex function theory (the Painlevé problem), the  $T(b)$  theorem has been adapted to various settings and more powerful and flexible versions are available nowadays. One of them can be proved and applied to obtain the

**Theorem 2.** *Let  $L$  be real and symmetric, then the Neumann to Dirichlet map is bounded and invertible.*

As Verchota did in his situation, we can reprove the results of Kenig and Pipher mentioned above by the method of layer potentials. But that is not all. We can also develop a perturbation theory by allowing the coefficients of  $A$  to vary in  $L^\infty$  norm. Another type of perturbation theory allowing the coefficients to depend in a minimally smooth way on the transverse variable  $t = x_{n+1}$  was developed by R. Fefferman, Kenig, Pipher and then Kenig, Pipher.

**Theorem 3.** *Assume that  $A(x)$  is either constant and complex, or real and symmetric with  $L^\infty$  coefficients or real of Kato type (see above) with  $L^\infty$  coefficients. Then there exists  $\varepsilon > 0$  depending only on the ellipticity constants of  $A$  such that if  $A'(x)$  is another complex coefficients matrix with*

$$(8) \quad \|A' - A\|_\infty \leq \varepsilon$$

then the Neumann to Dirichlet map corresponding to  $L' = -\operatorname{div}A'\nabla$  is bounded and invertible. Hence, the Neumann and Regularity problems for  $L'$  with  $L^2$  data are solvable.

In fact, this theorem is a consequence of a more general statement that would be too complicated to explain here. The proof of boundedness is done by a perturbation method comparing the layer potentials associated to  $A$  and  $A'$ . A remarkable point of this proof is that the Kato part of  $A'$ , that is the matrix obtained by setting the non-diagonal term to 0 and replacing  $a'_{n+1,n+1}$  by 1, plays an important role and in fact many of the estimates proved to establish Theorem 1, and some similar ones, are instrumental in the argument. The proof of invertibility for the perturbed Neumann to Dirichlet map is done via the method of continuity. This is here that we need to have handled complex coefficients so as to use tools of analytic functions such as the Cauchy formula.

To finish, we mention that in [1], we also develop another approach to obtain solvability of Neumann and Regularity problems (and more general transmission problems with Lipschitz interface). More precisely we imbed the elliptic problem in a first order systems (generalized Cauchy-Riemann equations) in some exterior algebra  $\Lambda$ . This first order system takes the form

$$(9) \quad \partial_t F + TF = 0 \quad \text{in } \mathbb{R}_+^{n+1}$$

where  $F$  is  $\Lambda$ -valued and  $T$  is some bi-sectorial Dirac-type operator on  $L^2(\mathbb{R}^n, \Lambda)$  that is independent of  $t$ . In this setting, there is no layer potential but there is an essential operator,  $\operatorname{sgn}(T) = T\sqrt{T^2}^{-1}$ , and the goal is to prove that it is a bounded operator on  $L^2(\mathbb{R}^n, \Lambda)$ . The solvability of the BVP is equivalent to having specific numbers in the resolvent set of another related operator whose boundedness follows from that of  $\operatorname{sgn}(T)$ . And openness of resolvent sets allows perturbation. This allows somehow more general perturbation results than the ones stated above.

#### REFERENCES

- [1] M. Alfonseca, P. Auscher, A. Axelsson, S. Hofmann and S. Kim. Two papers: Stability of layer potentials and  $L^2$  solvability of boundary value problems for divergence form elliptic equations with complex  $L^\infty$  coefficients and Perturbations of transmission problems for Dirac operators with complex coefficients with applications to regularity and Neumann problems. preprints.
- [2] P. Auscher, S. Hofmann, M. Lacey, A. McIntosh, and P. Tchamitchian. The solution of the Kato Square Root Problem for Second Order Elliptic operators on  $\mathbb{R}^n$ , *Annals of Math.* **156** (2002), 633–654.



## Heat kernels on percolation clusters

MARTIN BARLOW

Let  $G$  be an infinite (connected) graph. For  $x, y \in G$  let  $\mu_{xy} = 1$  if  $\{x, y\}$  is edge, and  $\mu_{xy} = 0$  otherwise. Define the *vertex degree*  $\mu(x) = \sum_y \mu_{xy}$ . Assume  $\mu(x) < \infty$  for all  $x \in G$ , and extend  $\mu$  to a measure on  $G$ . Set  $V(x, r) = \mu(B(x, r))$ . We define a Dirichlet form by:

$$\mathcal{E}(f, f) = \frac{1}{2} \sum_{x, y} (f(x) - f(y))^2 \mu_{xy}.$$

The continuous time simple random walk on  $G$  (CTSRW) on  $G$  is the process  $Y = (Y_t, t \in [0, \infty), P^x, x \in G)$  associated with  $(\mathcal{E}, L^2(G, \mu))$ . This process waits at a point  $x$  for an exponential time with mean 1, then moves to  $y \sim x$  with probability  $\mu_{xy}/\mu_x$ . The *heat kernel* on  $G$  is

$$q_t(x, y) = P^x(Y_t = y) \mu_y^{-1}.$$

Now consider bond percolation on  $\mathbb{Z}^d$ ; write  $\mathcal{C}(x)$  for the open cluster containing  $x$ , and let  $\theta(p) = \mathbb{P}_p(|\mathcal{C}(x)| = \infty)$ . Recall that in the supercritical regime ( $p > p_c$ ) there exists a unique infinite cluster, which we denote by  $\mathcal{C}_\infty = \mathcal{C}_\infty(\omega)$ . The CTSRW on the infinite percolation cluster  $\mathcal{C}_\infty$  is just the CTSRW on the graph  $(\mathcal{C}_\infty(\omega), E(\omega)|_{\mathcal{C}_\infty(\omega)})$ ; denote its heat kernel by  $q_t^\omega(x, y)$ .

In this talk I considered the heat kernel aspects of the following theorem

**Theorem 4.** [1] *Let  $p > p_c$ . For each  $x \in \mathbb{Z}^d$  there exist r.v.  $N_x(\omega)$  with  $\mathbb{P}_p(N_x \geq n) \leq c \exp(-n^{\varepsilon_d})$  and (non-random) constants  $c_i = c_i(d, p)$  such that the transition density of  $X$  satisfies, for  $x, y \in \mathcal{C}_\infty(\omega)$ :*

$$\frac{c_1}{t^{d/2}} e^{-c_2|x-y|^2/t} \leq q_t^\omega(x, y) \leq \frac{c_3}{t^{d/2}} e^{-c_4|x-y|^2/t}, \quad (GE)$$

provided

$$t \geq N_x(\omega), \text{ and } t \geq |x - y|.$$

Let  $Q$  be a (large) box in  $\mathbb{Z}^d$  with side  $r$ ,  $\mathcal{C}(Q)$  be the largest connected cluster in  $Q$ , and  $Q'$  be a box with side  $(9/10)r$  and the same centre as  $Q$ . We use four properties of  $\mathcal{C}(Q)$ ; these all hold with probability at least  $1 - e^{-cr^\delta}$ .

1. ‘Good volume’:

$$c_1 r^d \leq \mu(\mathcal{C}(Q)) \leq c_2 r^d. \quad (V_d)$$

2. ‘Good distances’:  $|x - y|_1 \leq d_\omega(x, y) \leq c_2 |x - y|_1$  for  $x, y \in Q' \cap \mathcal{C}_\infty$ .

3. ‘Good isoperimetric inequality’. If  $A \subset Q \cap \mathcal{C}_\infty$  with  $\mu(A) \geq \frac{1}{2} \mu(\mathcal{C}(Q))$  then

$$\frac{\mu(A, \mathcal{C}(Q) - A)}{\mu(A)} \geq \frac{C}{r}. \quad (1)$$

4. ‘Good surface effects’:  $\mathcal{C}(Q)$  and  $\mathcal{C}_\infty \cap Q$  do not differ by much. In particular, one has

$$Q' \cap \mathcal{C}(Q) = Q' \cap \mathcal{C}_\infty.$$

(1) implies a Poincaré inequality for  $B = \mathcal{C}(Q)$ : if  $f : \mathcal{C}(Q) \rightarrow \mathbb{R}$  then

$$\int_{\mathcal{C}(Q)} (f - \bar{f})^2 d\mu \leq cr^2 \sum_{x,y \in \mathcal{C}(Q)} (f(x) - f(y))^2 \mu_{xy}. \quad (PI)$$

(As usual  $\bar{f}$  is the real number which minimises the left hand side.)

A guide to the expected behaviour of  $Y$  on  $\mathcal{C}_\infty$  is given by the following theorem of Delmotte, which translates into the graph setting work of Grigoryan [4] and Saloff-Coste [7] on manifolds.

**Theorem 5.** [3] *Let  $\Gamma$  be an infinite connected graph, with heat kernel  $q_t(x, y)$ . The following are equivalent:*

- (a)  $\Gamma$  satisfies (VD) (volume doubling) and (PI) (Poincaré inequality).  
 (b) For  $x, y \in G$ ,  $t \geq d(x, y)$ ,

$$\frac{c_1}{t^{d/2}} e^{-c_2|x-y|^2/t} \leq q_t(x, y) \leq \frac{c_3}{t^{d/2}} e^{-c_4|x-y|^2/t}. \quad (GE)$$

- (c)  $\Gamma$  satisfies (PHI).

*Proof.* Of course the hardest part of the argument is to obtain (b) or (c) from (a). Delmotte proved that (a)  $\Rightarrow$  (c) using Moser's argument.

Since  $\mathcal{C}_\infty$  is random one can consider 'annealed' bounds (i.e. on  $\mathbb{E}_p(q_t^\omega(x, y) | x, y \in \mathcal{C}_\infty)$ ) or 'quenched' bounds, on  $q_t^\omega(x, y)$ . The first quenched bounds were obtained in by Mathieu and Remy in [6]:

$$q_t^\omega(x, y) \leq c_1 t^{-d/2}, \quad t \geq N_x(\omega), \quad (UB1)$$

where  $\mathbb{P}_p(N_x < \infty | x \in \mathcal{C}_\infty) = 1$ .

Before considering how one can prove (UB1), it may be useful to see how not to do so. It is well known that the 'Nash inequality'

$$\mathcal{E}(f, f) \geq c \|f\|_2^{2+4/d} \|f\|_1^{-4/d}. \quad (N)$$

is equivalent to:

$$q_t(x, x) \leq Ct^{-d/2} \text{ for all } x, y \in G, t \geq 1. \quad (GUB)$$

However (GUB) is false for  $\mathcal{C}_\infty$ , since  $\mathcal{C}_\infty$  contains arbitrarily large 'bad' regions – for example one-sided strings of length  $n$  connected at one end to the rest of  $\mathcal{C}_\infty$ . Thus (N) (and related global Sobolev inequalities) must fail for  $\mathcal{C}_\infty$ , and one needs 'local' methods.

The basic idea is that if (VD) and (PI) hold for all large balls then one should also get (GE) for all large times  $t$ . To make this more precise we make the following definitions.

**Definition.**

1. A ball  $B = ga(x, r)$  is good if (VD) and (PI) hold for  $B$ .

2. Let  $\alpha = 1/(11(2+d))$ . The ball  $B = B_\omega(x, R)$  is *very good* (VG) if  $B_\omega(y, r)$  is good for all

$$B_\omega(y, r) \subset B \text{ and } R^\alpha \leq r \leq R.$$

3.  $B = B_\omega(x, R)$  is *exceedingly good* (EG) if  $B$  is very good and satisfies a further, rather complicated, condition.

**Theorem 6.** *Let  $\Gamma$  be an infinite connected graph such that  $V(x, r) \leq C_0 r^d$  for all  $x \in G$ ,  $r \geq 1$ .*

(a) *Suppose for each  $x \in G$  there exists  $r_x < \infty$  such that  $B(x, R)$  is VG for all  $R \geq r_x$ . Let  $N_x = (r_x)^c$ . Then*

$$q_t(x, y) \leq \frac{c_3}{t^{d/2}} e^{-c_4|x-y|^2/t}, \quad t \geq N_x \vee d(x, y),$$

and

$$q_t^\omega(x, y) \geq \frac{c_1}{t^{d/2}} e^{-c_2|x-y|^2/t}, \quad t \geq N_x \vee d(x, y)^3.$$

(b) *If  $B(x, R)$  is EG for all  $R \geq r_x$  then the lower bound holds for  $t \geq d(x, y) \vee N_x$ .*

One can prove that the hypotheses of Theorem 3(b) hold for  $\mathcal{C}_\infty$  a.s., and so one derives Theorem 1.

*Proof of Theorem 3.* The basic idea is to work on an infinite connected graph  $\Gamma$ , and prove that if a ball  $B(x_0, R)$  is good/VG/EG then we have the right upper and lower bounds on  $q_t(x, y)$  for  $x, y \in B(x_0, \frac{1}{2}R)$  and for suitable  $t$ . Ideally we would have  $t \approx R^2$ , but in fact the arguments needed  $t \leq R^2/\log R$ .

Then, since for Theorem 3(a) we suppose that  $B(x_0, R)$  is VG for all  $R \geq R_x$  we obtain bounds on  $q_t(x, y)$  for all large enough  $t$ .

*Outline (for experts).*

1. One can prove on-diagonal upper bounds on  $q_t(x, x)$  directly from the Poincaré inequalities, using an argument of Kusuoka-Zhou – see [5].
2. The hardest part is usually to obtain off-diagonal upper bounds. Of the approaches available one cannot easily use Davies' method, as it relies on (global) Sobolev inequalities. An alternative might be to prove a Harnack inequality first, as in [3], but in [1] I used a method due to Bass and Nash – see [2].
3. For near diagonal lower bounds – i.e.

$$q_t(x, y) \geq ct^{-d/2} \quad \text{if } d(x, y) \leq t^{1/2} \tag{2}$$

one can use the Fabes-Stroock method based on a weighted Poincaré inequality, which can be derived from (PI) by the method of [8].

4. Full lower bounds (i.e. if  $t \geq d(x, y) \gg t^{1/2}$ ) require a chaining argument which needs (2) in small balls. So 'very good' isn't good enough, and we need 'exceedingly good'.

## REFERENCES

- [1] M.T. Barlow. Random walks on supercritical percolation clusters. *Ann. Probab.* **32** (2004), 3024–3084.
- [2] R.F. Bass. On Aronsen’s upper bounds for heat kernels. *Bull. London Math. Soc.* **34** (2002), 415–419.
- [3] T. Delmotte. Parabolic Harnack inequality and estimates of Markov chains on graphs. *Rev. Math. Iberoamericana* **15** (1999), 181–232.
- [4] A.A. Grigor’yan. The heat equation on noncompact Riemannian manifolds. *Math. USSR Sbormik* **72** (1992), 47–77.
- [5] S. Kusuoka, X.Y. Zhou. Dirichlet form on fractals: Poincaré constant and resistance. *Probab. Theory Related Fields* **93** (1992) 169–196.
- [6] P. Mathieu, E. Remy. Isoperimetry and heat kernel decay on percolation clusters. *Ann. Probab.* **32** (2004), 100–128.
- [7] L. Saloff-Coste. A note on Poincaré, Sobolev, and Harnack inequalities. *Inter. Math. Res. Notices* **2** (1992) 27–38.
- [8] L. Saloff-Coste, D.W. Stroock. Opérateurs uniformément sous-elliptiques sur les groupes de Lie. *J. Funct. Anal.* **98** (1991) 97–121.

**Symmetric Markov chains on  $Z^d$  with unbounded range**

RICHARD BASS

(joint work with Takashi Kumagai)

Let  $X_n$  be a symmetric Markov chain on  $Z^d$ . We say that  $X_n$  has *bounded* range if there exists  $K > 0$  such that  $P(X_{n+1} = y \mid X_n = x) = 0$  whenever  $|y - x| \geq K$ . The range is *unbounded* if for every  $K$  there exists  $x$  and  $y$  (depending on  $K$ ) with  $|x - y| > K$  such that  $P(X_{n+1} = y \mid X_n = x) > 0$ . There is a great deal known about Markov chains on graphs when the chains have bounded range. The purpose of this talk is to discuss results for Markov chains on  $Z^d$  that have unbounded range.

Suppose  $C_{xy}$  is the conductance between  $x$  and  $y$ . We impose a condition on  $C_{xy}$  which essentially says that the  $C_{xy}$  satisfy a uniform second moment condition. Let  $Y_t$  be the continuous time Markov chain on  $Z^d$  determined by the  $C_{xy}$ , while  $X_n$  is the discrete time Markov chain determined by these conductances. The transition probabilities for the Markov chain  $X$  are defined by

$$P^x(X_1 = y) = \frac{C_{xy}}{\sum_z C_{xz}},$$

while the process  $Y_t$  is the Markov chain that has the same jumps as  $X$  but where the times between jumps are independent exponential random variables with parameter 1. When the second moment condition holds, together with two very mild regularity conditions, we obtain upper bounds on the transition probabilities of the form

$$P(Y_t = y \mid Y_0 = x) \leq ct^{-d/2}$$

and some corresponding lower bounds when  $x$  and  $y$  are not too far apart. Unlike the case of bounded range, reasonable universal bounds of Gaussian type need not

hold when the range is unbounded. We also obtain bounds on the exit probabilities  $P(\sup_{s \leq t} |Y_s - x| > \lambda t^{1/2})$ .

We say a uniform Harnack inequality holds for  $X$  if whenever  $h$  is nonnegative and harmonic for the Markov chain  $X$  in the ball  $B(x_0, R)$  of radius  $R > 1$  about a point  $x_0$ , then

$$h(x) \leq Ch(y), \quad |x - x_0|, |y - x_0| < R/2,$$

where  $C$  is independent of  $R$ . Even when  $X_n$  is a random walk, i.e., the increments  $X_n - X_{n-1}$  form an independent identically distributed sequence, a uniform Harnack inequality need not hold. However, if we impose an additional strong assumption on the conductances, then we can prove such a Harnack inequality.

We prove that if we have Markov chains  $X^{(n)}$  on  $Z^d$  satisfying the second moment condition uniformly in  $n$ , then the sequence of processes  $X_t^{(n)} = X_{[nt]}/\sqrt{n}$  is tight in the space  $D[0, \infty)$  of right continuous, left limit functions, and all subsequential limit points are continuous processes. Under an additional condition on the conductances (different than the one needed for the Harnack inequality), we then show that the  $X^{(n)}$  converge weakly as processes to the law of the diffusion corresponding to an elliptic operator

$$\mathcal{L}f(x) = \sum_{i,j=1}^d \frac{\partial}{\partial x_i} \left( a_{ij}(\cdot) \frac{\partial f}{\partial x_j}(\cdot) \right) (x)$$

in divergence form.

### Ultracontractivity and embedding into $L^\infty$

ALEXANDER BENDIKOV

(joint work with Th. Coulhon, L. Saloff-Coste)

One of the classical uses of Sobolev embedding theorem is to show that an  $L^2$  function on  $\mathbb{R}^d$  having  $k$  derivatives in  $L^2$  with  $k > d/2$  is a bounded function. This has been generalized as follows. Let  $e^{-tA}$  be a semigroup of self-adjoint operators on  $L^2(X, \mu)$ . Assume that, for all  $t \in (0, 1)$ ,

$$\|e^{-tA}\|_{2 \rightarrow \infty} = \sup_{\|g\|_2 \leq 1} \|e^{-tA}g\|_\infty \leq Ct^{-\nu/4}.$$

Then any function  $f \in L^2(X, \mu)$  such that  $A^k f \in L^2(X, \mu)$  for some  $k > \nu/4$  (roughly speaking, this corresponds to  $2k$  derivatives in  $L^2$ ) must be a bounded function. See, e.g., [2, Théorème 1] and the references therein.

The aim of the present paper is to obtain results in this spirit when the semigroup  $e^{-tA}$  satisfies an ultracontractivity bound of the type

$$(1) \quad \|e^{-tA}\|_{2 \rightarrow \infty} = \sup_{\|g\|_2 \leq 1} \|e^{-tA}g\|_\infty \leq e^{m(t)}, \quad t > 0$$

with a function  $m$  which tends to infinity at least as fast as  $\log 1/t$  as  $t$  tends to 0. We call such a function  $m$  an ultracontractivity function for  $e^{-tA}$ .

More precisely, we would like to obtain equivalences between (1) and properties such as

$$(2) \quad g \in \bigcap_0^\infty \text{Dom}(A^n) \text{ and } \limsup_{n \rightarrow \infty} \frac{\|A^n g\|_2^{1/n}}{\phi(n)} \leq 1 \implies g \in L^\infty(X, \mu)$$

where the function  $m$  in (1) and the function  $\phi$  in (2) are related in some explicit way. We call a function  $\phi$  such that (2) holds an embedding function for  $A$ .

We also relate these properties to Nash type inequalities and characterize those functions  $f$  on the real line such that  $\|e^{-f(A)}\|_{2 \rightarrow \infty} < \infty$ . Similar questions were discussed in [1] which focussed on problems related to the long time behavior of the semigroup. In this paper, the focus is on the short time behavior.

In fact, the connection between Nash inequalities and ultracontractivity bounds on the one hand, embedding properties of the form (2) (or, to follow the terminology of [2], generalized Gagliardo-Nirenberg inequalities) and ultracontractivity bounds on the other hand have a different range of validity, as far as the behaviour of the function  $m$  is concerned. Nash inequalities are relevant when the explosion of  $m$  at 0 is not too fast, embedding properties are relevant when the explosion of  $m$  at 0 is fast enough. There is a common zone where both operate (see Theorem 7), and two exclusive zones where only one of them operates.

We also consider the case where  $X$  has finite measure and  $A$  has discrete spectrum together with  $L^\infty$  bounds on the eigenfunctions. In that case, one can obtain the connection with generalized Gagliardo-Nirenberg inequalities even when  $m$  does not belong to the favorable zone. This applies in particular to left-invariant Markov generators acting on locally compact metric groups.

Finally, we exhibit families of concrete examples, namely invariant diffusions on infinite dimensional tori and symmetric Lévy semigroups on the real line, which display the whole variety of behaviours we have been considering.

**Definition 1.** Let  $M$  be a non-negative non-increasing function defined on  $(0, +\infty)$  and such that  $M(0_+) = \infty$ . For non-negative  $x$ , set

$$(3) \quad F(x) = F_M(x) = \inf_{t>0} \{tx + M(t)\}$$

and

$$(4) \quad \Phi(x) = \Phi_M(x) = \sup_{t>0} \left\{ \frac{x}{t} e^{-M(t)/x} \right\}.$$

**Definition 2.** Let  $M$  be a non-increasing non-negative function defined on  $(0, +\infty)$  and such that  $M(0_+) = \infty$ . For any real  $x$ , set

$$(5) \quad N(x) = \sup_{t>0} \{xt - tM(1/t)\}.$$

When  $M \in \mathcal{C}^1$ , set also

$$(6) \quad Q(x) = \begin{cases} -M' \circ M^{-1}(x) & \text{if } x \geq M(\infty) \\ 0 & \text{otherwise.} \end{cases}$$

We can now state our main result.

**Theorem 7.** *Let  $M, F, \Phi$  be as in the above definition. Assume that  $M$  is  $C^1$ , convex, and that there are constants  $a, b \in (0, \infty)$  such that  $aM(t) \leq -tM'(t) \leq bM(t)$  for all  $t$  small enough. Let  $-A$  be the infinitesimal generator of a sub-Markovian semigroup on  $L^2(X, \mu)$ . Then the following properties are equivalent:*

- (1) *There exists  $c_1 \in (0, \infty)$  such that, for all  $t > 0$ ,  $\|e^{-tA}\|_{2 \rightarrow \infty} \leq e^{c_1 M(t)}$ .*
- (2) *There exists  $t_0 > 0$  such that, for all  $t > t_0$ ,  $\|e^{-tF(A)}\|_{2 \rightarrow \infty} < \infty$ .*
- (3) *There exists  $C_1 \in (0, \infty)$  such that for any function  $f \in \bigcap_0^\infty \text{Dom}(A^n)$  we have*

$$\limsup_{n \rightarrow \infty} \left\{ \frac{\|A^n f\|_2^{1/n}}{\Phi(n)} \right\} \leq C_1 \implies f \in L^\infty(X, \mu).$$

- (4) *There exists  $C_2 \in (0, \infty)$  such that the Nash inequality*

$$\forall f \in \text{Dom}(A) \text{ with } \|f\|_1 \leq 1, \quad \|f\|_2^2 \Phi(\log \|f\|_2) \leq C_2 (\langle Af, f \rangle + \|f\|_2^2)$$

*is satisfied.*

#### REFERENCES

- [1] Coulhon T. and Meda S. *Subexponential ultracontractivity and  $L^p$ - $L^q$  functional calculus.* Math. Z. 244, 2003, 291-308.
- [2] Coulhon T. *Inégalités de Gagliardo-Nirenberg pour les semi-groupes d'opérateurs et applications.* Potential Anal. 1, 1992, 343-353.

### On the Robin problem in fractal domains

KRZYSZTOF BURDZY

(joint work with R. Bass, Z.-Q. Chen)

The Robin problem (also known as the “third” boundary problem) for a Euclidean domain  $D \subset \mathbf{R}^d$  is to find a function  $u$  such that

$$(1) \quad \Delta u(x) = 0, \quad x \in D,$$

$$(2) \quad \frac{\partial u}{\partial \mathbf{n}} = cu, \quad x \in \partial D,$$

with one or more side conditions, where  $\mathbf{n}$  is the unit inward normal vector field on  $\partial D$ ,  $\partial u / \partial \mathbf{n}$  is the normal derivative of  $u$  in the distributional sense and  $c > 0$  is a constant. See Gustafson and Abe [8] for the history of this problem.

Our interest in the Robin problem stems from some recent applications in physics, electrochemistry, heterogeneous catalysis and physiology; see [4], [5], [7], [11] and the references therein. Consider the mixed Dirichlet-Robin problem

$$(3) \quad \Delta u(x) = 0, \quad x \in D \setminus B_*,$$

$$(4) \quad \frac{\partial u}{\partial \mathbf{n}} = cu, \quad x \in \partial D,$$

together with the side condition

$$(5) \quad u(x) = 1, \quad x \in \partial B_*,$$

where  $B_* \subset D$  is a fixed closed ball with non-zero radius. The solution to (3)-(5) represents the steady state of a system in which some particles move randomly in  $D \setminus B_*$  and cross a semi-permeable membrane  $\partial D$ . The other part of the boundary,  $\partial B_*$ , is a source of particles and can be controlled so that we can assume a condition of type (5). The constant  $c$  in (4) is a physical characteristic of the membrane  $\partial D$ . The constant  $c$  will play no role in our theorems so we take  $c = 1$ .

In some applied situations, it is desirable to have as much flux through the boundary as possible. The points of a man-made or natural membrane  $\partial D$  where there is no flux can be considered an inefficient use of material. Hence, it is interesting to know when the flux is non-negligible through all points of the membrane. In other words, we would like to know whether  $\inf_{x \in \partial D} \partial u / \partial \mathbf{n}(x) > 0$ . In view of the relation (4) between the flux  $\partial u / \partial \mathbf{n}$  and the density  $u$  of particles and the maximum principle for the harmonic function  $u$ , this condition is equivalent to  $\inf_{x \in D \setminus B_*} u(x) > 0$ .

**Definition 1.1.** We say that the whole surface of  $D$  is active if

$$(6) \quad \inf_{x \in D \setminus B_*} u(x) > 0.$$

If it is not the case that the whole surface is active, we say part of the surface is nearly inactive.

In this paper we investigate the following problem.

**Problem 1.2.** Give necessary and sufficient conditions of a geometric nature for the whole surface of  $D$  to be active.

It is not difficult to show that the whole surface of a bounded Lipschitz domain is always active. We have posed Problem 1.2 in terms of  $u$  rather than  $\partial u / \partial \mathbf{n}$  because we are interested in non-Lipschitz domains  $D$ ; so there are some boundary points where  $\mathbf{n}$  is not well-defined while the solution  $u$  is always well-defined, and, in fact, is smooth in  $D \setminus B_*$ . We do not have a complete solution to Problem 1.2, but we give a fairly explicit answer for some natural families of domains with fractal boundary.

We will approach Problem 1.2 using probabilistic methods. This agrees well with the motivating physical models. Suppose that  $X$  is reflecting Brownian motion in  $D$ ,  $L$  is its local time on  $\partial D$ , and  $T_{B_*}$  is the hitting time of  $B_*$  by  $X$ . When  $D$  is a bounded  $C^3$ -smooth domain, it is known that (see [9] and [10])

$$(7) \quad u(x) = E_x \left[ \exp \left( -\frac{1}{2} L_{T_{B_*}} \right) \right].$$

This formula indicates that the third boundary problem (4) is more difficult to study from the probabilistic point of view than the corresponding Dirichlet and Neumann problems. This is because the Dirichlet problem corresponds to killed Brownian motion and killing on the boundary presents no technical problems. The Neumann boundary problem corresponds to reflecting Brownian motion. The construction of reflecting Brownian motion in an arbitrary domain  $D$  is a major



technical challenge. Although this feat has been accomplished long time ago by Fukushima [6] on an abstract compactification, called the Martin-Kumarochi compactification, of  $D$ , many questions about the construction of reflecting Brownian motion on the Euclidean closure of a domain remain open (see [1]). Formula (7) shows that the Robin boundary problem (3)-(5) requires the construction and understanding of the local time. This is harder than constructing reflecting Brownian motion itself, because it is known that reflecting Brownian motion does not have a semimartingale decomposition in some domains. For some results in this area, see, e.g., DeBlassie and Toby [3]. For information on the eigenvalue problem for the Laplacian with Robin boundary conditions, see Smits [12], [13].

The following are some results proved in this paper.

- (i) The solution of Problem 1.2 for a class of domains with fractal boundaries.
- (ii) A semimartingale decomposition of reflecting Brownian motion in a class of fractal domains.
- (iii) A sharp estimate for the Green function with Neumann boundary conditions in long and thin domains.
- (iv) A new version of the Neumann boundary Harnack principle, stronger than the one in [2].

A simple example illustrating our main theorems is a cusp domain, defined for a fixed  $\alpha > 1$  by

$$D = \left\{ x = (x_1, x_2, \dots, x_d) : 0 < x_1 < 1 \text{ and } x_1^\alpha > (x_2^2 + \dots + x_d^2)^{1/2} \right\}.$$

Applying the main results of this paper, we can show that the whole boundary of  $D$  is active if  $\alpha \in (1, 2)$ , and part of  $\partial D$  is nearly inactive if  $\alpha \geq 2$ .

#### REFERENCES

- [1] R. Bass, K. Burdzy and Z.-Q. Chen (2005), Uniqueness for reflecting Brownian motion in lip domains *Ann. I. H. Poincaré* **41**, 197–235.
- [2] R. Bass and P. Hsu (1991), Some potential theory for reflecting Brownian motion in Hölder and Lipschitz domains. *Ann. Probab.* **19**, 486–508.
- [3] D. DeBlassie and E. Toby (1993), On the semimartingale representation of reflecting Brownian motion in a cusp. *Probab. Theory Rel. Fields* **94**, 505–524.
- [4] M. Felici, B. Sapoval, and M. Filoche (2003) Renormalized random walk study of oxygen absorption in the human lung, *Phys. Rev. Lett.* **92**, 068101-1—068101-4.
- [5] M. Filoche and B. Sapoval (1999) Can one hear the shape of an electrode? The active zone in Laplacian transfer, II. Theory. *European Journal of Physics, B*, **9**, 754–763.
- [6] M. Fukushima (1967), A construction of reflecting barrier Brownian motions for bounded domains. *Osaka J. Math.*, **4**, 183-215.
- [7] D.S. Grebenkov, M. Filoche and B. Sapoval (2003) Spectral properties of the Brownian self-transport operator *Europhys. Journ. B.* **36**, 221–231.
- [8] K. Gustafson and T. Abe (1998), The third boundary condition—was it Robin’s? *Math. Intelligencer* **20**, 63–71.

- [9] Z.-M. Ma and R. Song (1990), Probabilistic methods in Schrödinger equations. *Seminar on Stochastic Processes, 1989*. 135–164, *Progr. Probab.*, **18**, Birkhäuser, Boston.
- [10] V.G. Papanicolaou (1990), The probabilistic solution of the third boundary value problem for second order elliptic equations. *Probab. Theory Rel. Fields* **87**, 27–77.
- [11] B. Sapoval (1996), Transport across irregular interfaces: fractal electrodes, membranes and catalysts, in *Fractals and disordered systems, 2nd ed.*, A. Bunde and S. Havlin, Eds. (Springer-Verlag) 232–261.
- [12] R. Smits (2005), Eigenvalue estimates and critical temperature in zero fields for enhanced surface superconductivity, preprint.
- [13] R. Smits (2005), Upper and lower bounds for the principal eigenvalue of the generalized Robin problem, preprint.

### Correlation inequalities of Brascamp-Lieb inequalities and the heat kernel

ERIC CARLEN

*We discuss recent work with Elliott Lieb and Michael Loss, showing that a simple heat kernel argument can be used to prove correlation inequalities of Brascamp-Lieb type, and explain some new examples obtained this way.*

#### 1. INTRODUCTION

Consider an integral of a product of functions, such as

$$(1) \quad \int_X \prod_{j=1}^N f_j(\pi_j(x)) d\mu$$

where the  $\pi_j$  are certain “projections” from the measure space  $X$  to  $\mathbb{R}$ . For example when  $X$  is the unit sphere in  $\mathbb{R}^N$ , one might have  $\pi_j(x) = \vec{e}_j \cdot x$ , where  $\vec{e}_j$  is the  $j$ th standard basis vector of  $\mathbb{R}^N$ .

When  $\mu$  is a probability measure, the  $\pi_j$ , and hence the  $f_j \circ \pi_j$  are random variables. In general, they will not be independent. This is the case in the example mentioned above when  $\mu$  is the normalized probability measure on the unit sphere. Then of course, the integral of the product will not equal the product of the integrals. Indeed, it can even be the case that the integral of the product is infinite, due to correlations, while each individual integral is finite.

A Brascamp-Lieb inequality is one that relates an integral of a product, such as the one in (1), to a product of the  $L^p$  norms of the  $f_j \circ \pi_j$ , as in

$$(2) \quad \prod_{j=1}^N \left( \int_X |f_j(\pi_j(x))|^{p_j} d\mu \right)^{1/p_j} .$$

The classical theorem of Brascamp and Lieb [2],[6] concerns the case in which  $X$  is  $\mathbb{R}^M$ , with  $M < N$ ,  $d\mu$  is Lebesgue measure, and  $\pi_j(x) = \vec{a}_j \cdot x$  where the  $\{\vec{a}_1, \dots, \vec{a}_N\}$  span  $\mathbb{R}^M$ . Given  $L^p$  indices  $p_j$  with  $1 \leq p_j \leq \infty$  for  $j = 1, 2, \dots, N$ , form the vector

$$(3) \quad \vec{p} = (1/p_1, 1/p_2, \dots, 1/p_N) ,$$

and define

$$(4) \quad D(\vec{p}) = \sup \left\{ \frac{\int_{\mathbb{R}^M} \prod_{j=1}^N f_j(\vec{a}_j \cdot x) d^N x}{\prod_{j=1}^N \|f_j\|_{p_j}} : f_j \in L^{p_j}(\mathbb{R}) \quad j = 1, 2, \dots, N \right\} .$$

Next, let  $\mathcal{G}$  denote the set of all centered Gaussian function functions on  $\mathbb{R}$ ; i.e., those of the form  $g(x) = e^{-(sx)^2/2}$  for some  $s > 0$ . Define  $D_{\mathcal{G}}(\vec{p})$  by

$$(5) \quad D_{\mathcal{G}}(\vec{p}) = \sup \left\{ \frac{\int_{\mathbb{R}^M} \prod_{j=1}^N g_j(\vec{a}_j \cdot x) d^N x}{\prod_{j=1}^N \|g_j\|_{p_j}} : g_j \in \mathcal{G} \quad j = 1, 2, \dots, N \right\} .$$

The classical theorem of Brascamp and Lieb states that  $D(\vec{p}) = D_{\mathcal{G}}(\vec{p})$ , and hence

$$(6) \quad \int_{\mathbb{R}^M} \prod_{j=1}^N f_j(\vec{a}_j \cdot x) d^N x \leq D_{\mathcal{G}}(\vec{p}) \prod_{j=1}^N \|f_j\|_{p_j}$$

for all non negative  $f_1, \dots, f_N$ .

An important special case arises when  $N = 3$ ,  $M = 2$  and

$$\vec{a}_1 = \vec{e}_1 \quad \vec{a}_2 = \vec{e}_1 - \vec{e}_2 \quad \text{and} \quad \vec{a}_3 = \vec{e}_2 .$$

Then, denoting  $f_1, f_2, f_3$  by  $f, g, h$  respectively, (1) becomes

$$\int_{\mathbb{R}^2} f(x)g(x-y)h(y)dx dy .$$

In this case, it is quite easy to compute  $D_{\mathcal{G}}(\vec{p})$ , and thus one obtain the sharp constant in Young's inequality. (This important special case was also treated independently and simultaneously by Beckner.)

Several new examples of Brascamp–Lieb type correlation inequalities have recently been proven by Lieb, Loss and the author [3],[4]. The first of these holds when, as discussed in the introduction, the measure space  $X$  in (1) is the unit sphere  $S^{N-1}$  in  $\mathbb{R}^N$  equipped with the uniform probability measure  $\mu$ :

**Theorem 8.** *For all  $N \geq 2$ , given non-negative measurable functions  $f_1, \dots, f_N$ , on  $[-1, 1]$ ,*

$$(7) \quad \int_{\mathbb{S}^{N-1}} \left( \prod_{j=1}^N f_j \circ \pi_j \right) d\mu \leq \prod_{j=1}^N \|f_j \circ \pi_j\|_{L^p(\mathbb{S}^{N-1})} .$$

*for all  $p \geq 2$ . Moreover, the  $L^2$  norm is optimal in that for each  $p < 2$ , there exist functions  $f_j$  so that  $\|f_j \circ \pi_j\|_{L^p(\mathbb{S}^{N-1})} < \infty$  for each  $j$ , while the integral on the left side of (7) diverges. Finally, for every  $p \geq 2$  and  $N \geq 3$ , there is equality in (7) if and only if some function  $f_j$  vanishes identically, or else each  $f_j$  is constant.*

In the second of these,  $X$  is  $\mathcal{S}^n$  denote the symmetric group on  $N$  letters; i.e., the group of all permutations  $\sigma$  of  $\{1, \dots, N\}$ . Let the (composition) product in  $\mathcal{S}^n$  be denoted by juxtaposition, and for each  $1 \leq i, j \leq N$  with  $i \neq j$ , let  $\sigma_{i,j}$  be

the pair permutation with  $\sigma_{i,j}(i) = j$ ,  $\sigma_{i,j}(j) = i$ ,  $\sigma_{i,j}(k) = k$  for  $k \neq i, j$ . Let  $\mu$  denote the uniform probability measure on  $\mathcal{S}^n$  so that if  $g$  is any function on  $\mathcal{S}^n$ ,

$$(8) \quad \int_{\mathcal{S}} g(\sigma) d\mu = \frac{1}{N!} \sum_{\sigma \in \mathcal{S}^n} g(\sigma) .$$

**Theorem 9.** For any  $N \geq 2$  given functions  $f_1, \dots, f_N$ , on  $\{1, \dots, N\}$ ,

$$(9) \quad \int_{\mathcal{S}^n} \left( \prod_{j=1}^N f_j \circ \pi_j \right) d\mu \leq \prod_{j=1}^N \|f_j \circ \pi_j\|_{L^p(\mathcal{S}^n)} .$$

there is equality in (9) if and only if at least one of the functions  $f_j$  is zero, or else each  $f_j(k) = C_j e^{i(\alpha_j - \beta_k)}$  for some  $\alpha_1, \dots, \alpha_N$ ,  $\beta_1, \dots, \beta_N$ , and  $C_1, \dots, C_N$

This can be reformulated as a theorem about permanents, if we identify functions  $f$  on  $\{1, \dots, N\}$  in the obvious way: Let  $F$  be an  $N \times N$  complex matrix whose  $j$ th column is the vector  $\vec{f}_j$  in  $\mathbb{C}^N$ . Let  $|f_j|^2$  denote the sum of the absolute squares of the entries of  $\vec{f}_j$ . Hadamard's inequality for determinants [5] states that  $|\det(F)| \leq \prod_{j=1}^N |\vec{f}_j|$ . Then it is easy to reformulate the previous theorem as follows:

**Theorem 10.** For any vectors  $\vec{f}_1, \dots, \vec{f}_N$  in  $\mathbb{C}^N$  we have the inequality

$$(10) \quad |\text{perm}(F)| \leq \frac{N!}{N^{N/2}} \prod_{j=1}^N |\vec{f}_j| .$$

For  $N > 2$ , there is equality in (10) if and only if at least one of the vectors  $\vec{f}_j$  is zero, or else  $F$  is a rank one matrix and, moreover, each of the vectors  $\vec{f}_j$  is a constant modulus vector; i.e., its entries all have the same absolute value.

All of the theorems mentioned above, including the original paradigm, can be proved using a simple heat kernel interpolation. Barthe [1] had earlier given a proof of the original Brascamp–Lieb Theorem using an interpolation constructed by means of optimal mass transportation. This works very well on  $\mathbb{R}^N$ , but it is not clear how it could be adapted to work on  $S^{N-1}$  or on  $\mathcal{S}^n$ . On the other hand, the heat kernel proof is very simple, and works in essentially the same way in each of the cases mentioned above. We now briefly describe how this works.

For any non negative function  $f$  on  $\{1, \dots, N\}$  and any  $1 \leq j \leq N$ , and any  $1 \leq p < \infty$ , consider the function defined by  $(e^{t\Delta}(f \circ \pi_j))^p$ . It is easy to see that the result is also of the form  $g \circ \pi_j$ . Moreover, the  $L^p$  norm is constant during the evolution. All one has to do is a calculation to check that the integral of the product increases with  $t$ . Finally, on  $S^{N-1}$  or on  $\mathcal{S}^n$ ,  $(e^{t\Delta}(f \circ \pi_j))^p$  tends to a constant, while on  $\mathbb{R}^M$ , it gets “more and more Gaussian”. For the details, see the cited papers.

## REFERENCES

- [1] F. Barthe: *Optimal Young's inequality and its converse, a simple proof* Geom. Func. Analysis., **80** 234–242 (1998)
- [2] H. J. Brascamp and E. H. Lieb: *Best constants in Young's inequality, its converse, and its generalization to more than three functions* Advances in Math., **20** 151-173 (1976)
- [3] E.A. Carlen, E.H. Lieb and M. Loss, *A Sharp analog of Young's Inequality on  $\mathbf{S}^N$  and Related Entropy Inequalities*, Jour. Geom. Anal. **14** , 487-520 (2004). arXiv math.FA/0408030.
- [4] E.A. Carlen, E.H. Lieb and M. Loss, *An inequality of Hadamard type for permanents*, preprint arXiv math.CA/0508096
- [5] J. Hadamard, *Résolution d'une question relatif aux déterminants*, Bull. Sci. Math. **28**, 240-246 (1893).
- [6] E. H. Lieb, *Gaussian kernels have only Gaussian maximizers* Invent Math **102**, 179–208 (1990).

 **$L^p$  cohomology and the boundness of the Riesz transform**

GILLES CARRON

(joint work with Th. Coulhon, A. Hassell)

Let  $M$  be a complete Riemannian manifold with infinite measure. The Riesz transform  $T$  on  $M$  is the operator

$$f \rightarrow d\Delta^{-1/2}f,$$

where  $\Delta$  is the positive Laplace operator on  $M$ . Thanks to the Green formula we have :

$$\forall f \in C_0^\infty(M), \|\Delta^{1/2}f\|_{L^2}^2 = \langle \Delta f, f \rangle = \|df\|_{L^2}^2$$

hence the Riesz transform is always a bounded map from  $L^2(M)$  to  $L^2(M; T^*M)$ . It is of interest to figure out the range of  $p$  for which  $T$  extends to a bounded map  $L^p(M) \rightarrow L^p(M; T^*M)$ . Equivalently, we can ask whether

$$\|df\|_p \leq \|\Delta^{1/2}f\|_p \text{ for all } f \in C_c^\infty(M).$$

There is a lot of result in this direction. I will only mention few of such results :

- i) On  $\mathbb{R}^n$ , for all  $p > 1$ , the Riesz transform is bounded on  $L^p$ .
- ii) On manifold with non negative Ricci curvature, the Riesz transform is bounded on  $L^p$  for all  $p > 1$  ([3]).
- iii) On certain Cartan-Hadamard manifolds, the Riesz transform is bounded on  $L^p$  for all  $p > 1$  ([7]).
- iv) On simply connected nilpotent Lie groups endowed with a left invariant metric, the Riesz transform is bounded on  $L^p$  for all  $p > 1$  ([1]).
- v) In [5], T. Coulhon and X.T. Duong have shown that if for some  $C > 0$  and  $\nu > 0$ ,  $(M, g)$  satisfies the the relative Faber-Krahn inequality :

$$\forall x \in M, R > 0 \text{ and } \Omega \subset B(x, R) \lambda_1(\Omega) \geq \frac{C}{R^2} \left( \frac{\text{vol}\Omega}{\text{vol}B(x, R)} \right)^{-2/\nu},$$

then for all  $p \in ]1, 2]$  the Riesz transform is bounded on  $L^p$ .

They also indicated that for  $p > n$ , the Riesz transform is not bounded on  $L^p$  on a connected sum of two Euclidean space.

Our result is the following :

**Theorem** *Let  $M$  be a complete Riemannian manifold of dimension  $n \geq 3$  which is the union of a compact part and a finite number of Euclidean ends. Then the Riesz transform is bounded from  $L^p(M)$  to  $L^p(M; T^*M)$  for  $1 < p < n$ , and is unbounded on  $L^p$  for all other values of  $p$  if the number of ends is at least two.*

Let's describe a key point in the proof in the case of a connected sum of two Euclidean space  $M = \mathbb{R}^n \# \mathbb{R}^n$ . Such manifold is topologically the product  $\mathbb{R} \times \mathbb{S}^{n-1}$  endowed with the metric

$$(dr)^2 + (1 + r^2)(d\theta)^2$$

where  $(d\theta)^2$  is the standard round metric on the sphere  $\mathbb{S}^{n-1}$  and  $r$  the radial variable. One of our argument is a precise description of the behavior for large  $x$  and  $y$  of the Schwarz kernel of the operator :

$$\Delta^{-1/2} = \frac{2}{\pi} \int_0^\infty (\Delta + k^2)^{-1} dk.$$

If  $P(x, y)$  is the kernel of this operator, we show that when  $x \in M$  and  $y = (r, \theta)$  then for  $r \rightarrow \pm\infty$

$$P(x, (r, \theta)) \simeq \frac{u_\pm(x)}{r^{n-1}}$$

where  $u_\pm$  is the harmonic function such that

$$\begin{aligned} \lim_{r \rightarrow \pm\infty} u_\pm(x) &= 1 \\ \lim_{r \rightarrow \mp\infty} u_\pm(x) &= 0 \end{aligned}$$

At infinity, we also get

$$|du_\pm(r, \omega)| = O(r^{1-n}).$$

With this estimate, we can show that the Riesz transform can be bounded only when  $p < n$ . To get the full result, we obtain sharper estimates by using the scattering calculus introduced by R. Melrose ([8]).

In ([2]), authors have remarked that if the Riesz transform is bounded on  $L^{p/(p-1)}$  and on  $L^p$  then the operator  $P = d\Delta^{-1}d^*$  extend to a bounded operator on  $L^p(T^*M)$ . On  $L^2(T^*M)$ , this operator  $P$  is the orthogonal projection on the closure of the space  $dC_0^\infty(M)$ . But on  $(M, g)$ , the space of  $L^2$  differential one-forms admits the Hodge decomposition

$$L^2(T^*M) = \mathcal{H}^1(M) \oplus \overline{dC_0^\infty(M)} \oplus \overline{d^*C_0^\infty(\Lambda^2 T^*M)},$$

where  $\mathcal{H}^1(M) = \{\alpha \in L^2(T^*M), d\alpha = 0 = d^*\alpha\}$  (see [6]). Let us recall now the definition of reduced  $L^p$ -cohomology: for  $p \geq 1$ , the first space of reduced  $L^p$  cohomology of  $(M, g)$  is

$$H_p^1(M) = \frac{\{\alpha \in L^p(T^*M), d\alpha = 0\}}{\overline{dC_0^\infty(M)}},$$

where we take the closure in  $L^p$ . On  $L^2$ , we have

$$\{\alpha \in L^2(T^*M), d\alpha = 0\} = \mathcal{H}^1(M) \oplus \overline{dC_0^\infty(M)},$$

hence the first space of reduced  $L^2$  cohomology can be identified with  $\mathcal{H}^1(M)$ .

We assume now that the manifold  $(M, g)$  satisfies the following conditions

$$\left( \int_M |f|^{\frac{2\nu}{\nu-2}} d\text{vol} \right)^{1-2/\nu} \leq C \int_M |df|^2 d\text{vol}, \quad \forall f \in C_0^\infty(M),$$

And that at one point  $x_0 \in M$ , we have a control on the growth of geodesic balls centered at  $x_0$  :

$$\text{vol} B(x_0, r) \leq Cr^\nu, \quad \forall r \geq 1.$$

**Proposition** *Under these two hypotheses, if the Riesz transform is bounded in  $L^p$  for some  $p > 2$ , then*

$$H_p^1(M) = \{\alpha \in L^p(M; T^*M) \mid d^*\alpha = d\alpha = 0\}.$$

*If moreover the Ricci curvature of  $M$  is bounded from below then there is a natural map*

$$H_2^1(M) \rightarrow H_p^1(M)$$

*which is injective.*

**Corollary** *If  $M$  has at least two ends, then the Riesz transform is not bounded on  $L^p$  for any  $p \geq \nu$ .*

Let's us describe two examples :

**First example:** The manifold  $M = \mathbb{R}^n \# \mathbb{R}^n = \mathbb{R} \times \mathbb{S}^{n-1}$  endowed with a metric  $(dr)^2 + (1+r^2)(d\theta)^2$ . Then it is easy to show by direct computation that

$$\{\alpha \in L^p(M; T^*M) \mid d^*\alpha = d\alpha = 0\} = \mathbb{R} \frac{dr}{(1+r^2)^{\frac{n-1}{2}}}.$$

Where as

$$H_p^1(M) = \begin{cases} \mathbb{R} & \text{if } p < n \\ \{0\} & \text{if } p \geq n \end{cases}$$

Hence the map  $H_2^1(M) \rightarrow H_p^1(M)$  is injective only for  $p < n$ . The proposition gives in this case the right range of  $p$ 's where the Riesz transform is bounded on  $L^p$ .

**Second example:** Let  $M$  be a connected sum of several (say  $l \geq 2$ ) copies of a simply connected nilpotent Lie group  $N$  endowed with a left invariant metric, let  $\nu$  be the homogeneous dimension of  $N$ ; that is

$$\nu = \lim_{R \rightarrow \infty} \frac{\log \text{vol} B(o, R)}{\log R},$$

Then  $M$  satisfies the Sobolev inequality, the upper bound on the volume growth of geodesic balls and more over it's clear that its Ricci curvature is bounded from below. We known that the Riesz transform is bounded on  $L^p$  for  $p \in ]1, 2]$  and not

bounded on  $L^p$  for any  $p \geq \nu$ . Moreover we can compute the  $L^p$  cohomology of  $M$ :

$$H_p^1(M) = \begin{cases} H_c^1(M) \simeq \mathbb{R}^{l-1} & \text{if } p < \nu \\ \{0\} & \text{if } p > \nu \end{cases}$$

Hence the map

$$H_2^1(M) \rightarrow H_p^1(M)$$

is injective when  $p < \nu$ . It is then tempting to propose the following conjecture :

*On such a manifold  $M$ , is the Riesz transform bounded on any  $p \in ]1, \nu[$  ?*

#### REFERENCES

- [1] G. Alexopoulos, An application of homogenization theory to harmonic analysis: Harnack inequalities and Riesz transforms on Lie groups of polynomial growth, *Canad. J. Math.* 44 (1992), 691–727.
- [2] P. Auscher, T. Coulhon, X.T. Duong and S. Hofmann, Riesz transform on manifolds and heat kernel regularity, *Ann. Sc. E.N.S.* 37 (2004), 911–957.
- [3] D. Bakry, Etude des transformations de Riesz dans les variétés riemanniennes à courbure de Ricci minorée, *Séminaire de Probabilités*, XXI, 137–172, Lecture Notes in Math., 1247, Springer, Berlin, 1987.
- [4] T. Coulhon, G. Carron and A. Hassell, Riesz transform and  $L^p$  cohomology for manifold with Euclidean ends, to appear in *Duke Math. Journal*. ArXiv math.AP/0411648.
- [5] T. Coulhon and X.T. Duong, Riesz transforms for  $1 \leq p \leq 2$ , *Trans. Amer. Math. Soc.* 351 (1999), 1151–1169.
- [6] G. De Rham, *Variétés différentiables, formes, courants, formes harmoniques*, Hermann, 3rd edition, 1973.
- [7] N. Lohoué, Comparaison des champs de vecteurs et des puissances du laplacien sur une variété riemannienne à courbure non positive, *J. Funct. Anal.* 61 (1985), no. 2, 164–201.
- [8] R.B. Melrose, Spectral and scattering theory for the Laplacian on asymptotically Euclidian spaces, in *Spectral and Scattering Theory*, M. Ikawa, ed., Marcel Dekker, 1994.

### Boundary Processes of Markov Processes

ZHEN-QING CHEN

(joint work with Masatoshi Fukushima, Jiangang Ying)

Time change is one of the most basic and very useful transformations for Markov processes, which has been studied by many authors. However a precise characterization of the time-changed process of a Markov process  $X$  on a state space  $E$  by a Revuz measure whose quasi-support  $F$  is a proper subset of  $E$  has only been started very recently. The time-changed process has  $F$  as its state space so it can be regarded as the trace process of  $X$  on  $F$ .

The following is a prototype of the problem we discussed in the talk. Suppose  $X$  is a Lévy process in  $\mathbf{R}^n$  that is the sum of a Brownian motion in  $\mathbf{R}^n$  and an independent spherically symmetric  $\alpha$ -symmetric stable process in  $\mathbf{R}^n$ , where  $n \geq 1$  and  $\alpha \in (0, 2)$ . Denote by  $B(x, r)$  the open ball in  $\mathbf{R}^n$  centered at  $x \in \mathbf{R}^n$  with radius  $r$ . Its Euclidean closure is denoted by  $\overline{B(x, r)}$ . Let  $F = \overline{B(0, 1)} \cup \partial B(x_0, 1)$ , where  $x_0 \in \mathbf{R}^n$  with  $|x_0| = 3$ . What is the trace process of  $X$  on the closed set



$F$ ? More precisely, let  $\mu(dx) := 1_{\overline{B(0,1)}}(x) dx + \sigma_{\partial B(x_0,1)}$ , where  $\sigma_{\partial B(x_0,1)}$  denotes the Lebesgue surface measure of  $\partial B(x_0,1)$ . It is easy to see that  $\mu$  is a smooth measure of  $X$  and it uniquely determines a positive continuous additive functional  $A^\mu = \{A_t^\mu, t \geq 0\}$  of  $X$  having  $\mu$  as its Revuz measure. Define its inverse

$$\tau_t := \inf\{s > 0 : A_s^\mu > t\} \quad \text{for } t \geq 0.$$

Then the time changed process  $Y_t := X_{\tau_t}$  is a symmetric Markov process on  $F$ , which can be regarded as the trace process of  $X$  on  $F$ . So the more precise question is

*Question:* Can we characterize the time changed process  $Y$ ?

In fact, one can ask such questions for a general irreducible Markov process  $X$  that has a weak dual process on a general state space  $E$  which not only can have discontinuous sample paths but also can have killings inside  $E$  or have finite lifetime, and for any *quasi-closed* subset  $F$  of  $E$ .

When  $X$  is a general  $m$ -symmetric Markov process, we have obtained a complete characterization of the time changed process  $Y$  in terms of its Burling-Deny decomposition in [1]. In particular, the characterization of Lévy system of  $Y$  is obtained in terms of Feller measure and supplement Feller measure. These measures are intrinsic quantities for the part process of  $X$  killed upon leaving  $E \setminus F$ .

We also studied time changes for non-symmetric Markov processes. Let  $(X, \widehat{X})$  be a pair of Borel standard processes on a Lusin space  $E$  that are in weak duality with respect to some  $\sigma$ -finite measure  $m$  that has full support on  $E$ . Let  $F$  be a finely closed subset of  $E$ . In [2], we have obtained the characterization of a Lévy system of the time changed process of  $X$  by a positive continuous additive functional (PCAF in abbreviation) of  $X$  having support  $F$ , under the assumption that every  $m$ -semipolar set of  $X$  is  $m$ -polar for  $X$ . The characterization of the Lévy system is again in terms of Feller measures, which are intrinsic quantities for the part process of  $X$  killed upon leaving  $E \setminus F$ . Along the way, various relations between the entrance law, exit system, Feller measures and the distribution of the starting and ending point of excursions of  $X$  away from  $F$  are studied. We also show that the time changed process of  $X$  is a special standard process having a weak dual and that the  $\mu$ -semipolar set of  $Y$  is  $\mu$ -polar for  $Y$ , where  $\mu$  is the Revuz measure for the PCAF used in the time change.

The research of this author is supported in part by NSF Grant DMS-0303310.

These work extend the previous results obtained by LeJan in [4]-[5] and by Fukushima, He and Ying in [3].

This talk is based on joint work with Masatoshi Fukushima and Jiangang Ying [1] and [2].

#### REFERENCES

- [1] Z.-Q. Chen, M. Fukushima and J. Ying, Traces of symmetric Markov processes and their characterizations. To appear in *Ann. Probab.*
- [2] Z.-Q. Chen, M. Fukushima and J. Ying, Entrance law, exit system and Lévy system of time changed processes. To appear in Doob's memorial issue of *Illinois J. Math.*

- [3] M. Fukushima, P. He and J. Ying, Time changes of symmetric diffusions and Feller measures. *Ann. Probab.* **32** (2004), 3138-3166
- [4] Y. LeJan, Balayage et formes de Dirichlet. *Z. Wahrsch. verw. Gebiete* **37** (1977), 297-319
- [5] Y. LeJan, Measure associées a une forme de Dirichlet. applications. *Bull.Soc. Math. France* **106** (1978), 61-112

### Limit theorems for geometric approximations to Wiener measure

BRUCE DRIVER

A typical path integral is an **informal** expression of the form

$$(1) \quad \frac{1}{Z} \int_{\mathcal{F}} f(\sigma) e^{-E(\sigma)} \mathcal{D}\sigma$$

where  $\mathcal{F}$  is a space of maps from one manifold to another,  $f$  is a real valued function on  $\mathcal{F}$ ,  $E(\sigma)$  is the energy of the map  $\sigma$ ,  $\mathcal{D}\sigma$  is “Lebesgue measure” and  $Z$  is a normalization constant. The use of path integrals for “quantizing” classical mechanical systems (whose classical energy is  $E$ ) started with Feynman in [3] with very early beginnings being traced back to Dirac [2]. We present three rigorous interpretations to the path integral in (1) in the simplest non-trivial case, namely when  $\mathcal{F} = \{\sigma \in C([0, 1] \rightarrow M) : \sigma(0) = o\}$  where  $M$  is a compact,  $d$ -dimensional Riemannian manifold and  $o$  is a fixed point in  $M$ . This is done by first replacing  $\mathcal{F}$  by an increasing sequence  $\{\mathcal{F}_n\}$  of approximating finite dimensional “submanifolds” consisting of piecewise geodesic paths. The formal measure  $\mathcal{D}\sigma$  on  $\mathcal{F}$  is then replaced by the Riemannian volume measure on  $\mathcal{F}_n$  relative to three different natural Riemannian metrics. The limit of these three approximating measures is shown to exist as a measure on  $\mathcal{F}$ . Each of the limiting measure are absolutely continuous relative to the Wiener measure ( $\nu$ ) on  $\mathcal{F}$  and their (distinct) Radon-Nikodym derivatives are explicitly computed.

In more detail, let  $\mathcal{P}_n := \{s_i := \frac{i}{n} : i = 0, 1, 2, \dots, n\}$  be a uniform partition of  $[0, 1]$ ,

$$\mathcal{F}_n = \{\sigma \in \mathcal{F} : \nabla \sigma'(s)/ds = 0 \text{ off } \mathcal{P}_n\}.$$

For  $\sigma \in \mathcal{F}_n$ , and  $X, Y \in T_\sigma \mathcal{F}_n$ , define:

(1) the  $H^1$ -Metric on  $\mathcal{F}_n$  by

$$G^1(X, Y) := \int_0^1 \left\langle \frac{\nabla X(s)}{ds}, \frac{\nabla Y(s)}{ds} \right\rangle ds,$$

(2) the **Riemann sum approximation** to the  $H^1$ -Metric on  $\mathcal{F}_n$  by

$$G_n^1(X, Y) := \sum_{i=1}^n \left\langle \frac{\nabla X(s_{i-1+})}{ds}, \frac{\nabla Y(s_{i-1+})}{ds} \right\rangle \frac{1}{n},$$

and

(3) the **Riemann sum approximation** to the  $L^2$ -Metric on  $\mathcal{F}$  by

$$G_n^0(X, Y) := \sum_{i=1}^n \langle X(s_i), Y(s_i) \rangle \frac{1}{n}.$$

Further define three measure  $\nu_n^0$ ,  $\nu_n^1$ , and  $\nu_n$  on  $\mathcal{F}_n$  as:

$$d\nu_n^0 := \frac{1}{Z_n^0} e^{-\frac{1}{2}E_M} \cdot d\text{Vol}_{G_n^0},$$

$$d\nu_n^1 := \frac{1}{Z_n^1} e^{-\frac{1}{2}E_M} \cdot d\text{Vol}_{G_n^1},$$

and

$$d\nu_n := \frac{1}{Z_n^1} e^{-\frac{1}{2}E_M} \cdot d\text{Vol}_{G^1|_{T\mathcal{F}_n}}$$

where

$$E_M(\sigma) := \int_0^1 |\sigma'(s)|_g^2 ds,$$

is the energy functional, and  $Z_n^0$  and  $Z_n^1$  are normalization constants given by

$$Z_n^0 := (2\pi/n^2)^{nd/2} \quad \text{and} \quad Z_n^1 := (2\pi)^{dn/2}.$$

In each of the three theorems below  $f$  is assumed to be a bounded and continuous (in the sup-norm topology) real valued function on  $\mathcal{F}$ .

**Theorem 11** (Andersson and Driver [1]). *The sequence of measures  $\{\nu_n^1\}_{n=1}^\infty$  satisfy,*

$$\lim_{n \rightarrow \infty} \int_{\mathcal{F}_n} f(\sigma) d\nu_n^1(\sigma) = \int_{\mathcal{F}} f(\sigma) d\nu(\sigma)$$

**Theorem 12** (Andersson and Driver [1]). *Let  $\text{Scal}(m)$  be the scalar curvature of  $M$  at a point  $m \in M$ . Then*

$$\lim_{n \rightarrow \infty} \int_{\mathcal{F}_n} f(\sigma) d\nu_n^0(\sigma) = \int_{\mathcal{F}} f(\sigma) e^{-\frac{1}{6} \int_0^1 \text{Scal}(\sigma(s)) ds} d\nu(\sigma).$$

For each  $m \in M$ , let  $R_m$  be the curvature tensor at  $m \in M$  and define  $\Gamma_m \in \text{End}(T_m M)$  by

$$\Gamma_m = \sum_{i,j=1}^d \left( R_m(e_i, R_m(e_i, \cdot)e_j) e_j \right. \\ \left. + R_m(e_i, R_m(e_j, \cdot)e_i) e_j + R_m(e_i, R_m(e_j, \cdot)e_j) e_i \right)$$

where  $\{e_i\}_{i=1,2,\dots,d}$  is any orthonormal basis for  $T_m(M)$ . For  $\sigma \in \mathcal{F}$ , let  $u(s) := //_s(\sigma)$  denote stochastic parallel translation along  $\sigma$  and define a linear operator on  $L^2([0, 1]; T_o M)$  by

$$(K_\sigma f)(t) := \int_0^1 t \wedge s \left[ u(s)^{-1} \Gamma_{\sigma(s)} u(s) \right] f(s) ds.$$

**Theorem 13** (Adrian Lim [4]). *Suppose the sectional curvatures of  $M$  are bounded below by 0 and above by  $1/(2 \cdot \dim(M))$ , then*

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathcal{F}_n} f(\sigma) d\nu_n(\sigma) \\ = \int_{\mathcal{F}} f(\sigma) e^{-\frac{1}{6} \int_0^1 \text{Scal}(\sigma(s)) ds} \cdot \det \left( I + \frac{1}{12} K_\sigma \right) d\nu(\sigma). \end{aligned}$$

We refer the reader to the references in [1] and [4] for more background and related references to these three theorems.

#### REFERENCES

- [1] Lars Andersson and Bruce K. Driver, *Finite-dimensional approximations to Wiener measure and path integral formulas on manifolds*, J. Funct. Anal. **165** (1999), no. 2, 430–498.
- [2] P. A. M. Dirac, *Physikalische Zeitschrift der Sowjetunion* **3** (1933), 64.
- [3] R. P. Feynman, *Space-time approach to non-relativistic quantum mechanics*, Rev. Modern Physics **20** (1948), 367–387.
- [4] Adrian Lim, *Finite Dimensional Approximations to Wiener Measure*, Ph.D. thesis in preparation, 2006.

#### Intertwined diffusions

DAVID ELWORTHY

(joint work with Y. LeJan, X.-M. Li-Hairer)

**A) Intertwined diffusion generators.** Consider a smooth surjective map  $p : N \rightarrow M$  between manifolds  $N$  and  $M$  with smooth diffusion generators  $\mathcal{A}$  and  $\mathcal{B}$  on the manifolds  $M$  and  $N$  respectively, intertwined by  $p$ . This means that for any smooth  $f : M \rightarrow \mathbb{R}$  we have

$$\mathcal{B}(f \circ p) = \mathcal{A}(f) \circ p.$$

They have symbols  $\sigma^{\mathcal{A}}$  and  $\sigma^{\mathcal{B}}$  related by the commutative diagram

$$\begin{array}{ccc} T_u^* N & \xrightarrow{\sigma_u^{\mathcal{B}}} & T_u N \\ (T_u p)^* \uparrow & & \downarrow T_u p \\ T_{p(u)}^* M & \xrightarrow{\sigma_{p(u)}^{\mathcal{A}}} & T_{p(u)} M \end{array}$$

where  $T_u p$  denotes the derivative map of  $p$  at the point  $u$ , a linear map between the relevant tangent spaces, and  $(T_u p)^*$  is its adjoint acting on cotangent spaces. Recall that the (principal) *symbol* of a second order differential operator such as  $\mathcal{A}$  is defined by

$$df_x(\sigma_x^{\mathcal{A}}(dg_x)) = \frac{1}{2} (\mathcal{A}(fg)(x) - f(x)\mathcal{A}(g)(x) - g(x)\mathcal{A}(f)(x))$$

for  $x \in M$  and smooth functions  $f$  and  $g$  on  $M$ . The symbols of our diffusion generators determine symmetric, positive semi-definite, bilinear forms on the cotangent spaces. Let  $E_x$  be the image of  $\sigma_x^{\mathcal{A}}$  and let  $\langle -, - \rangle$  be the inner product induced on it by  $\sigma_x^{\mathcal{A}}$ . In the elliptic case  $E_x$  the symbol is positive definite and  $E_x$  is the whole of the tangent space  $T_x M$ , furnishing  $M$  with a Riemannian metric.

There are obvious questions which arise from this set up. One is on the relationships between the operators  $\mathcal{A}$  and  $\mathcal{B}$ , for example is there a nice decomposition of  $\mathcal{B}$ ? are there spectral relationships? Others concern the diffusion processes generated by the operators, for example the existence of a skew product decomposition of  $\mathcal{B}$ -diffusions, or the filtering problem of finding the conditional law of the  $\mathcal{B}$ -diffusion given its projection on  $M$  by  $p$ . We have not considered relations between the spectra of  $\mathcal{A}$  and  $\mathcal{B}$ , for this in the special cases when  $M$  and  $N$  are Riemannian manifolds and the map  $p$  is a Riemannian submersion with  $\mathcal{A}$  and  $\mathcal{B}$  the corresponding Laplace -Beltrami operators, see [1]. However the other questions are discussed in [4], with an earlier version for the case when  $p : N \rightarrow M$  is a principal bundle, and so in particular  $M$  is the quotient of  $N$  by a group action in [3]. Most of our results require a "cohesiveness" condition on  $\mathcal{A}$ , described below.

Earlier results have mainly concerned the cases where  $\mathcal{B}$  is the Laplacian : see [2] for a skew product decomposition when  $p$  is Riemannian with totally geodesic fibres, with an application to a factorisation theorem for harmonic maps; there is an extension of this by Liao in [7] to more general Riemannian submersions; for a discussion of skew -product decompositions of Brownian motions, with many examples, see Pauwels& Rogers [9], or [10].

**B) Semi-Connections.** The key step in our discussions is the construction of a *semi-connection* associated to our intertwining. For this we need the symbol of  $\mathcal{A}$  to have constant rank,  $q$  say. this means that if  $E$  denotes the union of all the subspaces  $E_x, x \in M$  then  $E$  is a subbundle of the tangent bundle  $TM$ .

By a (non-linear) *semi-connection on  $p : N \rightarrow M$  over  $E$*  we will mean a smooth horizontal lift map  $\mathcal{H}$  giving for each  $u \in N$  a linear mapping  $\mathcal{H}_u : E_{p(u)} \rightarrow T_u N$  which is a right inverse to the derivative  $T_u p : T_u N \rightarrow T_{p(u)} M$  of  $p$  at  $u$ . For such a semi-connection let  $H_u$  denote the image of  $\mathcal{H}_u$ ; this is the *horizontal subspace* at  $u$ . Let  $F_u$  be the sum of  $H_u$  with the *vertical subspace*  $\text{Ker} T_u p$  and  $\Pi_u : F_u \rightarrow H_u$  the projection. When  $E = TM$  we have a (*non-linear*) *connection* [8].

If we assume that  $\sigma^{\mathcal{A}}$  has constant rank and so has image a subbundle  $E$  we obtain a semi-connection over  $E$  characterised by the requirement

$$\mathcal{H}_u \circ \sigma_{p(u)}^{\mathcal{A}} = \sigma^{\mathcal{B}}(T_u p)^*.$$

A semi-connection  $\mathcal{H}$  over  $E$  determines a covariant differentiation  $\nabla^{\mathcal{H}}$  in the  $E$ -directions acting on smooth sections  $f : M \rightarrow N$  of  $p$ . For this note that the derivative  $T_x f$  at a point  $x = p(u)$  of such a section maps  $E_x$  to  $F_{f(x)}$ . Then, by definition,  $\nabla_v^{\mathcal{H}} := T_x f(v) - \Pi_{f(x)} T_x f(v) \in \text{Ker} T_{f(x)} p$ , for all  $v \in E_x$ . Also any curve  $\sigma$  in  $M$  with  $\dot{\sigma}(t) \in E_{\sigma(t)}$  for all  $t$  has a unique maximal horizontal lift  $\tilde{\sigma}$  with  $\sigma(0)$  any given point above  $\sigma(0)$  and  $\dot{\tilde{\sigma}}(t) \in H_{\tilde{\sigma}(t)}$  for all  $t$  for which it is defined.

Using Stratonovich differentials there is the corresponding result for continuous semi-martingales in  $M$ .

To lift an  $\mathcal{A}$ -diffusion by this "corresponding result" we need to assume that  $\mathcal{A}$  is *cohesive* or *along*  $E$ . There are various possible definitions of this; one is that  $\mathcal{A}$  has a smooth Hormander form representation

$$\mathcal{A} = 1/2 \sum_j \mathcal{L}_{X^j} \mathcal{L}_{X^j} + \mathcal{L}_A$$

where  $\mathcal{L}_A$  denotes Lie differentiation along the vector field  $A$ , and  $X^j, j = 1, \dots, m$  and  $A$  are vector fields taking values in  $E$ .

Given this cohesiveness there is a canonical decomposition of  $\mathcal{B}$  into the sum of a "horizontal" and a "vertical" diffusion generator. If  $p : N \rightarrow M$  is a principal bundle and  $\mathcal{B}$  is equivariant then so is this decomposition and there is a skew product representation of  $\mathcal{B}$ -diffusions. In this case the semi-connection is equivariant, and is sometimes called a "partial connection" or "connection over  $E$ ".

C. Stochastic flows and diffusion of tensors. This theory is applied to stochastic flows to obtain a skew product decomposition of the flow (given a cohesive generator), and a representation of the conditioned flow given the one point motion from a chosen point of  $M$ . Also a stochastic flow for  $\mathcal{A}$  determines a diffusion of sections of tensor bundles giving a semi-group on tensor fields generated by the right hand side of our Hormander form for  $\mathcal{A}$  using the usual extension of lie differentiation to tensor fields. Our techniques give an expression for the 'filtering out of the redundant noise' from these diffusions, generalising results in [6] and [5].

#### REFERENCES

- [1] L. Bérard-Bergery and J.-P. Bourguignon. Laplacians and Riemannian submersions with totally geodesic fibres. *Illinois J. Math.*, 26(2):181–200, 1982.
- [2] K. D. Elworthy and W. S. Kendall. Factorization of harmonic maps and Brownian motions. In *From local times to global geometry, control and physics (Coventry, 1984/85)*, Pitman Res. Notes Math. Ser., 150, pages 75–83. Longman Sci. Tech., Harlow, 1986.
- [3] K. D. Elworthy, Y. Le Jan, and Xue-Mei Li. Equivariant diffusions on principal bundles. In *Stochastic analysis and related topics in Kyoto*, volume 41 of *Adv. Stud. Pure Math.*, pages 31–47. Math. Soc. Japan, Tokyo, 2004.
- [4] K. D. Elworthy, Yves Le Jan, and Xue-Mei Li. Intertwined diffusions (tentative title). In preparation.
- [5] K. D. Elworthy, Y. LeJan, and X.-M. Li. *On the geometry of diffusion operators and stochastic flows*, *Lecture Notes in Mathematics 1720*. Springer, 1999.
- [6] K. D. Elworthy and M. Yor. Conditional expectations for derivatives of certain stochastic flows. In J. Azéma, P.A. Meyer, and M. Yor, editors, *Sem. de Prob. XXVII. Lecture Notes in Maths. 1557*, pages 159–172. Springer-Verlag, 1993.
- [7] M. Liao. Factorization of diffusions on fibre bundles. *Trans. Amer. Math. Soc.*, 311(2):813–827, (1989).
- [8] Peter W. Michor. *Gauge theory for fiber bundles*, volume 19 of *Monographs and Textbooks in Physical Science. Lecture Notes*. Bibliopolis, Naples, 1991.
- [9] E. J. Pauwels and L. C. G. Rogers. Skew-product decompositions of Brownian motions. In *Geometry of random motion (Ithaca, N.Y., 1987)*, volume 73 of *Contemp. Math.*, pages 237–262. Amer. Math. Soc., 1988.

- [10] L. C. G. Rogers and D. Williams. *Diffusions, Markov processes, and martingales. Vol. 2.* Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2000.

### One point extensions of general Markov processes

MASATOSHI FUKUSHIMA

The 'boundary problem of Markov processes' was one of the popular subjects in probability theory in the 1960's ([1],[2],[3],[4], [5], [6]). Later developments in the theory of Markov processes, especially the excursion theory based on exit systems [8] and the time change theory based on Dirichlet forms [9] were rooted in the study of the boundary problem. In a series of joint works with H.Tanaka, Z.-Q.Chen and J.Ying ([11],[13],[14], [15]), I have been reconsidering those issues left open in the boundary problem by using the well developed current theory of Markov processes.

In [11] and [15], we are concerned with the one point extensions of general Markov processes originated in the seminal work of Itô [6] supplemented by Meyer [7]. Itô only gave an analysis and left open the synthesis, namely, construction problem starting with the minimal process. A motivation of Itô's work was in his study with McKean [1] on the Brownian motion on the half line, while FT was motivated by the extension of the absorbing Brownian motion on the multi-dimensional bounded domain to its one point compactification presented in [4] in terms of the associated Sobolev space.

I shall talk about [15] which extends [11] from the symmetric diffusion process to the pair of standard processes in weak duality.

More specifically, Let  $a$  be a non-isolated point of a topological space  $E$ . Suppose we are given standard processes  $X^0$  and  $\widehat{X}^0$  on  $E_0 = E \setminus \{a\}$  in weak duality with respect to a  $\sigma$ -finite measure  $m$  on  $E_0$  which are of no killing inside  $E_0$  but approachable to  $a$ . We first show that their right process extensions  $X$  and  $\widehat{X}$  to  $E$  admitting no sojourn at  $a$  and keeping the weak duality are uniquely determined by the approaching probabilities of  $X^0$ ,  $\widehat{X}^0$  and  $m$  up to a non-negative constant  $\delta_0$  representing the killing rate of  $X$  at  $a$ . We then construct, starting from  $X^0$ , such  $X$  by piecing together returning excursions around  $a$  and a possible non-returning excursion including the instant killing. This extends [11] which treats the case where  $X^0$ ,  $X$  are  $m$ -symmetric diffusions and  $X$  admits no sojourn nor killing at  $a$ . While the Dirichlet form theory plays a role in [11], the recent studies of exit systems in [12] and [14] replace the role. As typical examples, the cases where  $X^0$  is the censored symmetric stable process studied in [10] and a non-symmetric diffusion process on an Euclidean domain are presented.

#### REFERENCES

- [1] K. Itô and H.P.McKean.Jr., Brownian motions on a half line, Illinois J. Math. **7**(1963), 181-231

- [2] M. Motoo, Application of additive functionals to the boundary problem of Markov processes (Lévy system of U-processes), in Proc. 5th Berkeley Symp. Math. Statist. Probab. **III**, 75-110, Univ. of California Press 1965
- [3] M. Fukushima, A construction of reflecting barrier Brownian motions for bounded domains, Osaka J. Math. **4**(1967), 183-215
- [4] M. Fukushima, On boundary conditions for multi-dimensional Brownian motions with symmetric resolvent densities, J. Math. Soc. Japan **21**(1969), 58-93
- [5] H. Kunita, General boundary conditions for multi-dimensional diffusion processes, J. Math. Kyoto Univ. **10**(1970), 273-335
- [6] K. Itô, Poisson point processes attached to Markov processes, in: Proc. Sixth Berkeley Symp. Math. Stat. Probab. **III**, 1970, pp225-239
- [7] P.A. Meyer, Processus de Poisson ponctuels, d'après K. Itô, Séminaire de Probab. V, in: Lecture Notes in Math., Vol.191, Springer,1971, pp.177-190
- [8] B. Maisonneuve, Exit systems, Ann. Probab. **3**(1975), 399-411
- [9] M. Fukushima, Dirichlet forms and Markov processes, North Holland/Kodansha, 1980
- [10] K. Bogdan, K. Burdzy, Z.Q. Chen, Censored stable processes, Probab. Theory Relat. Fields **127**(2003), 89-152
- [11] M. Fukushima and H. Tanaka, Poisson point processes attached to symmetric diffusions, Ann. Inst. H. Poincaré Probab.Statist. **41**(2005), 419-459
- [12] P.J. Fitzsimmons and R.K. Gettoor, Excursion theory revisited, to appear in Illinois J. Math.
- [13] Z.Q. Chen, M. Fukushima and J. Ying, Traces of symmetric Markov processes and their characterizations, to appear in Ann. Probab.
- [14] Z.Q. Chen, M. Fukushima and J. Ying, Entrance law, exit system and Lévy system of time changed processes, to appear in Illinois J. Math.
- [15] Z.Q. Chen, M. Fukushima and J. Ying, Extending Markov processes in weak duality by Poisson point processes of excursions, to appear in Proceedings of the Abel symposium 2005 Stochastic Analysis and Applications-A Symposium in honor of Kiyosi Itô, Springer

### Quasi-invariance of the Wiener Measure on the Loop Space of a Riemannian Manifold

ELTON HSU

Let  $M$  be a compact Riemannian manifold and  $P_x(M)$  the space of continuous functions  $\gamma : [0, 1] \rightarrow M$  such that  $\gamma(0) = x$ . The standard filtration of Borel  $\sigma$ -fields is denoted by  $\{\mathcal{B}_s, 0 \leq s \leq 1\}$ . Let  $\mathbb{P}_x$  be the law of Brownian motion on  $M$  from  $x \in M$  and  $\mathbb{P}_{xy}$  be the law of Brownian bridge on  $M$  from  $x$  to  $y$  with time length  $[0, 1]$ . We know that these two measures are mutually equivalent on  $\mathcal{B}_s$  for all  $s < 1$ . The measure  $\mathbb{P}_{xy}$  is concentrated on  $P_{xy}(M) = \{\gamma \in P_x(M) : \gamma(1) = y\}$ .

Introduce the Cameron-Martin norm

$$|h|_{\mathcal{H}} = \sqrt{\int_0^1 |\dot{h}_s|^2 ds}$$

and the Cameron-Martin space

$$\mathcal{H}_0 = \{h \in P(\mathbb{R}^n) : h(0) = h(1) = 0 \text{ and } |h|_{\mathcal{H}} < \infty\}.$$

Consider the vector field  $D_h$  defined on  $P(M)$ :

$$D_h(\gamma)_s = U(\gamma)_s h_s,$$



where  $U(\gamma)$  is the horizontal lift of  $\gamma$  starting from a fixed frame  $u_x$  at  $x$ . It has been shown ([1] and [2]) that the vector field  $D_h$  generates a flow  $\{\zeta^t, t \in \mathbb{R}\}$  on  $P_{xy}(M)$  and the Wiener measure  $\mathbb{P}_{xy}$  is quasi-invariant under the flow. More precisely, the shifted Wiener measure  $P_{xy}^t = P_{xy} \circ \zeta^{-t}$  is mutually absolutely continuous with  $\mathbb{P}_{xy}$ . In this note, we outline a proof of this result which is simpler than the previous approaches. Details will be found in the forthcoming paper [3] coauthored with Fuzhou Gong.

For  $s < 1$ , the two measures  $\mathbb{P}_{xy}^t$  and  $\mathbb{P}_{xy}$  are equivalent on  $\mathcal{B}_s$ . The Radon-Nikodym derivative

$$l_s = \left. \frac{d\mathbb{P}_{xy}^{-t}}{d\mathbb{P}_{xy}} \right|_{\mathcal{B}_s}, \quad 0 \leq s < 1$$

is an exponential martingale. Hence, it must have the form  $l_s = e^{z_s}$ , where

$$z_s = \int_0^s \langle \beta_\tau, db_\tau \rangle - \frac{1}{2} \int_0^s |\beta_\tau|^2 d\tau.$$

Here  $b_s$  is a euclidean Brownian motion (the martingale part of the stochastic development  $w_s$  of  $\gamma_s$ ). The process  $\beta_s$  can be expressed as

$$\beta_s = (O_s^t)^* c(1-s, u_s^t) - c(1-s, u_s) + (O_s^t)^* A_s^t,$$

where

$$c(1-s, u_s) = u_s^{-1} \nabla \ln p_M(1-s, \gamma_s, y)$$

with  $p_M(\cdot, \cdot, \cdot)$  being the heat kernel of  $M$  and  $u_s^t$  being the horizontal lift of  $\gamma_s^t = \zeta(\gamma)_s^t$  starting from a fixed frame  $u_0$ . The processes  $\{O_s^t, A_s^t\}$  are given in terms of the flow  $\zeta^t$  as follows. Let  $J : P_0(\mathbb{R}^n) \rightarrow P_x(M)$  be the Itô map (stochastic development) and  $w^t = J^{-1}\gamma^t : P_x(M) \rightarrow P_0(\mathbb{R}^n)$  the image of the flow in the flat path space  $P_0(\mathbb{R}^n)$ . It can be uniquely written in the form

$$dw_s^t = O_s^t dw_s + A_s^t ds,$$

where  $O_s^t \in O(n)$ , the orthogonal group. The flow equation for the vector field  $D_h$  is equivalent to a flow equation for  $w^t$  (or equivalently, for  $\{O^t, A^t\}$ ):

$$O^t = I - \int_0^t K(w^\lambda) O^\lambda d\lambda, \quad A^t = O^t \int_0^t (O^\lambda)^{-1} \left\{ \dot{h} - \frac{1}{2} \text{Ric}_{u^\lambda} h \right\} d\lambda.$$

Here  $\text{Ric}_u$  is the scalarized Ricci transform at a frame  $u \in \mathcal{O}(M)$  and  $K_h(w^t)_s \in o(n)$  (antisymmetric matrices). We do not need to  $K(w^t)_s$  explicitly save for the fact that  $K(w^t)_s$  is the vertical component of  $d(u_s^t)/dt$ .

The mutual absolute continuity of  $\mathbb{P}_{xy}^{-t}$  and  $\mathbb{P}_{xy}$  on  $\mathcal{B}_1$  is equivalent to the uniform integrability of the exponential martingale  $l_s$ . Writing  $l_1 = e^{z_1} e^{z_1 - z_s}$  and using Jensen's inequality we have

$$l_s^{-1} \mathbb{E}_{xy} [l_1 | \mathcal{B}_s] = \mathbb{E}_{xy} [e^{z_1 - z_s} | \mathcal{B}_s] \geq e^{\mathbb{E}_{xy}[z_1 - z_s | \mathcal{B}_s]}.$$

Suppose that we can find a constant  $C$  independent of  $0 \leq s < 1$  such that

$$(1) \quad \mathbb{E}_{xy} [z_1 - z_s | \mathcal{B}_s] \geq -C.$$

Then  $0 \leq l_s \leq e^C \mathbb{E}_{xy} \{l_1 | \mathcal{B}_s\}$ , which immediately implies the uniform integrability of  $\{l_s, 0 \leq s < 1\}$ . Therefore it is enough to prove (1), or

$$(2) \quad \mathbb{E}_{xy} \left[ \int_s^1 |\beta_\tau^t|^2 d\tau \middle| \mathcal{B}_s \right] \leq C.$$

the process  $\beta_s$  has three terms. The last term is easy to deal with because

$$|O^{-t*} A_s^{-t}| \leq C |t| \{ |h_s| + |\dot{h}_s| \}.$$

The difference which forms the first two terms is the integral from 0 to  $t$  of the derivative of  $O_s^{t*} c(1 - s, u_s^t)$  (with respect to  $t$ ). From the flow equation we have

$$\frac{d}{dt} O_s^{t*} = O_s^{t*} K(w^t)_s.$$

Here we have used the antisymmetry of  $K(w^t)_s$ . As for the derivative  $du^t/dt$ , its horizontal component is  $u_s^t h_s$  by the definition of horizontal lift, and its vertical component is  $K(w^t)_s$ , as we have pointed out before. Therefore the derivative of  $c(1 - s, u_s^t)$  with respect to  $t$  is equal to

$$\left\langle (u_s^t)^{-1} \nabla^2 \ln p_M(1 - s, \gamma_s^t, y), h_s \right\rangle - K(w^t)_s c(1 - s, u_s^t),$$

the first term on the right side being just the matrix multiplication. Now, in the derivative of  $O_s^{t*} c(1 - s, u_s^t)$  with respect to  $t$ , the two terms involving  $K(w^t)_s$  cancel each other and we obtain simply

$$\frac{d}{dt} \{ O_s^{t*} c(1 - s, u_s^t) \} = O_s^{t*} \left\langle (u_s^t)^{-1} \nabla^2 \ln p_M(1 - s, \gamma_s^t, y), h_s \right\rangle.$$

This shows that the inequality (2) we wanted to prove is implied by

$$\mathbb{E}_{xy} \left\{ \int_s^1 |\nabla^2 \ln p_M(1 - s, \gamma_s^t, y)|^2 |h_\tau|^2 d\tau \right\} \leq C.$$

Using the estimate

$$|\nabla^k \ln p_M(s, z_1, z_2)| \leq C_k \left[ \frac{1}{\sqrt{s}} + \frac{d_M(z_1, z_2)}{s} \right]^k.$$

we see that the left side is bounded by

$$\mathbb{E}_{xy} \int_0^1 \left[ \frac{d_M(\gamma_s^t, y)^4}{(1 - s)^4} + \frac{1}{(1 - s)^2} \right] |h_s|^2 ds.$$

We have the bound

$$\mathbb{E}_{xy}, d_M(\gamma_s^t, y)^4 \leq C (1 + |h|_{\mathcal{H}}^4 |t|^4) (1 - s)^2.$$

It follows that there is a constant  $C$  (depending on  $t$  and  $|h|_{\mathcal{H}}$ ) such that

$$\mathbb{E}_{xy} \int_0^1 \left[ \frac{d_M(\gamma_s^t, y)^4}{(1 - s)^4} + \frac{1}{(1 - s)^2} \right] |h_s|^2 ds \leq C \int_0^1 \left| \frac{h_s}{1 - s} \right|^2 ds \leq 4C |h|_{\mathcal{H}}.$$

## REFERENCES

- [1] Driver, B., A Cameron-Martin type quasi-invariance theorem for pinned Brownian motion on a compact Riemannian manifold. *Trans. Amer. Math. Soc.*, **342**, no. 1, 375-395 (1994).  
 [2] Enchev, O. and Stroock, D., Integration by parts for pinned Brownian motion, *Math. Res. Lett.*, **2**, no. 2, 161-169 (1995).  
 [3] Gong, F. Z. and Hsu, Elton P., On the Quasi-invariance of Brownian Bridge on a Compact Riemannian Manifold, preprint (2006).

**Jump processes,  $\mathcal{L}$ -harmonic functions, continuity estimates**

MORITZ KASSMANN

(joint work with R. Husseini and M. Barlow/R. Bass/Z.-Q. Chen)

Given a family of Lévy measures  $\mu(x, dh)$  we study the following set of questions:

- When can one construct corresponding jump processes in  $\mathbb{R}^d$  ?
- What can one say about heat kernels ?
- Are harmonic functions with respect to these processes regular ?
- Does the generator extend to the generator of a Feller semi-group ?

This program is not new at all, it is rather natural to raise the questions above and there are several publications on this subject. One way to summarize existing results is: As long as  $x \mapsto \mu(x, dh)$  is smooth and  $A \mapsto \mu(x, A)$  is asymptotically rotational invariant and non-degenerate, one can construct corresponding Feller processes and harmonic functions are smooth.

In our presentation we concentrated on cases where these two assumptions may fail. First, we explained what can happen in general situations. Assume  $\mu(x, dh) = n(x, h) dh$  and

$$(1) \quad \frac{c_0}{|h|^{d+\alpha}} \leq n(x, h) \leq \frac{c_1}{|h|^{d+\beta}}, \quad n(x, h) = n(x+h, -h)$$

where  $d$  is the space dimension and  $\alpha, \beta \in (0, 2)$ ,  $\alpha < \beta$  are two real numbers. With the help of Dirichlet form techniques one can associate a strong Markov process to  $\mu(x, dh)$ . Quite astonishing is the following result which we presented:

**Theorem 14.** [2] *Under the above assumption (1) there are discontinuous  $\mathcal{L}$ -harmonic functions and the martingale problem for*

$$\mathcal{L}u(x) = \text{p. v.} \int (u(x+h) - u(x))n(x, h) dh$$

*is not well-posed.*

In [1] we concentrate on continuity a-priori estimates for harmonic functions. Hölder regularity of functions being harmonic w.r.t to diffusions or jump processes has been an important object of studies for many years. The standard method used to prove Hölder regularity in the general case goes back to Krylov and Safonov (1979). At the heart of these techniques are exit time estimates of the following type: There exists a  $c > 0$  such that, given any starting point  $x$ , a small ball  $B(x, r)$  and any measurable set  $A \subset B(x, r)$  not too small relative to the ball,

the probability of hitting  $A$  before leaving  $B(x, r)$  the first time is bounded from below by  $c$ .

Nonetheless often a much weaker uniform control on the modulus of continuity of harmonic functions is sufficient. In the talk we presented continuity estimates for a class of pure jump processes where the above exit time estimates do not necessarily hold. In fact the probability of hitting  $A$  before leaving  $B(x, r)$  may tend logarithmically to 0 for  $r \rightarrow 0$ , as can be illustrated by an example.

As an application a certain kind of uniqueness of the martingale problem is established in [1]. Combining the results of [1] and [2] the problem of well-posedness of the martingale problem for non-local operators is now much better understood than it was before.

Preprints with references to other papers can be found at the web page:

<http://www.iam.uni-bonn.de/~kassmann>

#### REFERENCES

- [1] R. Hussein, M. Kassmann. *Jump processes,  $\mathcal{L}$ -harmonic functions, continuity estimates and the Feller property*, preprint.
- [2] M. Barlow, R. Bass, Z.-Q. Chen, M. Kassmann. *Non-local Dirichlet forms and symmetric jump processes*, preprint.

### Resistance forms and heat kernel estimates

JUN KIGAMI

The notion of resistance form was introduced to characterize a space where every pair of points has a finite resistance. Let  $(\mathcal{E}, \mathcal{F})$  be a resistance form on  $X$ . Under mild assumptions, for any nonempty compact subset  $B$  of  $X$ , we have the  $B$ -Green function  $g_B : X \times X \rightarrow [0, \infty)$  which satisfies (GB1) and (GB2):

(GF1)  $g_B(x, y) = g_B(y, x)$  and  $g_B(x, y) = 0$  for any  $y \in B$ .

(GF2) Define  $g_B^x(y) = g_B(x, y)$ , then  $g_B^x \in \mathcal{F}$  and

$$\mathcal{E}(g_B^x, u) = u(x)$$

for any  $u \in \mathcal{F}$  with  $u|_B \equiv 0$ .

If  $(\mathcal{E}, \mathcal{F})$  is a local regular Dirichlet form on  $L^2(X, \mu)$  as well, we may associate a diffusion process and have

$$E_x(\tau_B) = \int_X g_B(x, y) \mu(dy),$$

where  $\tau_B = \inf\{t | X_t \in B\}$ .

Applying those facts, we will study an asymptotic behavior of the associated heat kernel. In particular, desired heat kernel estimates are the sub-Gaussian upper estimate (1) and the near diagonal lower sub-Gaussian estimate (2) below.

**The upper sub-Gaussian estimate:**

$$(1) \quad p(t, x, y) \leq \frac{c_1}{V(x, t^{1/\beta})} \exp\left(-c_2 \left(\frac{d(x, y)^\beta}{t}\right)^{\frac{1}{\beta-1}}\right),$$

where  $V(x, r) = \mu(\{y | d(x, y) < r\})$ .

**The near diagonal lower sub-Gaussian estimate:**

$$(2) \quad \frac{c}{V(x, t^{1/\beta})} \leq p(t, x, y)$$

if  $d(x, y) \leq \epsilon$ .

We will try to find a “good distance”  $d$  under which we have (1) and (2)

**Theorem** Let  $(\mathcal{E}, \mathcal{F})$  be a resistance form on  $X$ . Assume that  $(\mathcal{E}, \mathcal{F})$  is a local regular Dirichlet form on  $L^2(X, \mu)$ . Let  $p(t, x, y)$  be the heat kernel associated with  $(\mathcal{E}, \mathcal{F})$  and  $\mu$ . Then the following two conditions (A) and (B) are equivalent.

(A)  $\mu$  has the volume doubling property with respect to the resistance metric associated with the resistance form  $(\mathcal{E}, \mathcal{F})$ .

(B) There exists a distance  $d$  and  $\beta > 1$  such that  $d$  is quasisymmetric to the resistance metric and (1) and (2) hold.

### A logarithmic Sobolev form of the Li-Yau inequality

M. LEDOUX

(joint work with D. Bakry)

This is a summary of a paper to appear in *Revista Mat. Iberoamericana*.

We present a finite dimensional version of the logarithmic Sobolev inequality for heat kernel measures of non-negatively curved diffusion operators that contains and improves upon the Li-Yau parabolic inequality in Riemannian manifolds with non-negative Ricci curvature. This new inequality is of interest already in Euclidean space for the standard Gaussian measure, and provides a finite dimensional extension of the classical logarithmic Sobolev inequality. The result may also be seen as an extended version of the semigroup commutation properties under curvature conditions, and the functional inequality is actually equivalent to the curvature-dimension condition. The proof relies on the classical Bakry-Emery semigroup argument which is improved by the dimension hypothesis together with a suitable integration of the resulting differential inequality. Partial results under arbitrary curvature conditions are also discussed, although the optimal form seems difficult to obtain. The main result may be applied, as time goes to infinity, to reach, in this setting, optimal Euclidean logarithmic Sobolev inequalities for the invariant measure. Exponential Laplace differential inequalities through the Herbst argument furthermore yield diameter bounds and dimensional estimates on the heat kernel volume of balls.

## On the range of random walks on percolation clusters

PIERRE MATHIEU

Let

$$N(t) = \#\{X(s); 0 \leq s \leq t\}$$

denote the **range** up to time  $t$  of a nearest neighbor path in the lattice  $\mathbb{Z}^d$ , say  $(X(s), s \geq 0)$ .

When  $(X(t), t \geq 0)$  performs a **simple symmetric random walk** on  $\mathbb{Z}^d$ , the asymptotics of the Laplace transform of  $N(t)$  are given by the:

**Theorem 15.** (*M.D. Donsker, S.R.S. Varadhan, [3]*)

Let  $E$  denote the expectation with respect to the law of the simple symmetric random walk on  $\mathbb{Z}^d$ . Let  $\alpha \in ]0, 1[$ . As  $t$  tends to  $+\infty$ ,

$$t^{-d/(2+d)} \log E(\alpha^{N(t)})$$

converges to some limit:  $c_d(\alpha) \in ]-\infty, 0[$ .

This abstract reports on our attempt to obtain a similar bound for the symmetric random walk on a percolation cluster.

### Random walks on percolation clusters

Consider super critical Bernoulli bond percolation in  $\mathbb{Z}^d$ : let  $\mathbb{E}_d$  be the set of edges of the grid  $\mathbb{Z}^d$ . Let  $\Omega$  be the set of sub-graphs of  $(\mathbb{Z}^d, \mathbb{E}_d)$ . We identify  $\Omega$  with the product space  $\{0, 1\}^{\mathbb{E}_d}$  and define  $Q$  to be the product Bernoulli( $p$ ) measure on  $\Omega$ .

We assume that  $p > p_c$ , where  $p_c$  is the critical probability for the appearance of an infinite connected component in  $\omega$ , see [5]. Thus  $Q$ .a.s. the graph  $\omega$  has a unique infinite connected component, the so-called **infinite cluster**, that we will denote with  $\mathcal{C}(\omega)$ . Finally define

$$Q_0(\cdot) = Q(\cdot | 0 \in \mathcal{C}(\omega)).$$

For a given  $\omega \in \Omega$  such that  $0 \in \mathcal{C}(\omega)$ , let  $P^\omega$  be the law of the **simple symmetric random walk** on  $\mathcal{C}(\omega)$  started at point 0. We use the notation  $E^\omega$  for the expectation with respect to  $P^\omega$ .

**Theorem 16.** (*C. Rau, in preparation*)

Let  $\alpha \in ]0, 1[$ . There exist constants  $C_1 = C_1(\alpha, p, d)$  and  $C_2 = C_2(\alpha, p, d)$  such that  $Q_0$  almost surely, for large enough times  $t$  one has

$$e^{-C_1 t^{-d/(2+d)}} \leq E^\omega(\alpha^{N(t)}) \leq e^{-C_2 t^{-d/(2+d)}}.$$

Comparing with Theorem 15, one notices that the order of magnitude of the Laplace transform of  $N(t)$  is the same as for the random walk on the full grid  $\mathbb{Z}^d$ .

We do not know whether  $t^{-d/(2+d)} \log E^\omega(\alpha^{N(t)})$  actually has a limit or not.

**Some indication on the proof:** The lower bound is easy to prove. Below, we only discuss the upper bound when  $\alpha = 1/2$ .

Following [4], we use the interpretation of the Laplace transform of  $N(t)$  as a return probability for a Markov chain on the **wreath product**  $\mathcal{C}(\omega) \wr \{0, 1\}$ , say  $E^\omega[2^{-N(t)}1_{X(t)=0}] = P^\omega[Z(t) = 0]$ . To get an upper bound on  $P^\omega[Z(t) = 0]$ , we first obtain a sharp control on the isoperimetric profile of the cluster  $\mathcal{C}(\omega)$  in the spirit of [6], and then deduce isoperimetric bounds on the wreath product as in [4]. Using general results on the link between isoperimetric inequalities, Nash inequalities and return probabilities as in [2], we can conclude the proof.

#### REFERENCES

- [1] BARLOW M.T. Random walks on supercritical percolation clusters. *Ann Probability* **32**, 3024-3084 (2004).
- [2] COULHON T. Ultracontractivity and Nash type inequalities. *Journ Functional Anal.* **141**, 510-539 (1996).
- [3] DONSKER M.D. and VARADHAN S.R.S. On the number of distinct sites visited by a random walk. *Comm. Pure Applied Math* **32**, 721-747 (1979).
- [4] ERSCHLER A. On isoperimetric profiles of finitely generated groups. *Geometrica Dedicata* **100**, 157-171 (2003).
- [5] GRIMMETT, G. Percolation. Springer-Verlag, Berlin (1999).
- [6] MATHIEU P. and REMY E. Isoperimetry and heat kernel decay on percolation clusters. *Ann Probability* **32**, 100-128 (2004)

### Sharp decay rates for the fastest conservative diffusions

ROBERT J. MCCANN

In many diffusive settings, initial disturbances will gradually disappear and all but their crudest features — such as size and location — will eventually be forgotten. Quantifying the rate at which this information is lost is sometimes a question of central interest. Joint works with Yong Jung Kim (KAIST) and Dejan Slepcev (UCLA) address this issue for the conservative nonlinearities in a model problem known as the fast diffusion equation

$$u_t = \Delta(u^m), \quad (n-2)_+/n < m < 1 \quad u, t \geq 0, \quad x \in \mathbf{R}^n,$$

which governs the decay of any integrable, compactly supported initial density towards a characteristically spreading self-similar profile. For other values of the parameter  $m$ , this equation has been used to model heat transport, population spreading, fluid seepage, curvature flows, and avalanches in sandpiles.

For the fastest conservative nonlinearities  $m \leq n/(n+2)$ , we develop a potential theoretic comparison technique with Kim which establishes the sharp conjectured power law rate of decay  $1/t$  uniformly in relative error, and in weaker norms such as  $L^1(\mathbf{R}^n)$ . For nonlinearities  $m \geq (n-1)/n$  we attain nearly the same  $L^1(\mathbf{R}^n)$  rate as a second order asymptotic after centering the mass of the solution, using an entropy dissipation approach with Slepcev. This leaves a gap in dimensions  $n \geq 3$ .

## REFERENCES

- [1] Y.-J. Kim and R.J. McCann. Sharp decay rates for the fastest conservative diffusions. *C. R. Acad. Sci. Paris Ser. I Math.* **341** (2005) 157–162.
- [2] Y.-J. Kim and R.J. McCann. Potential theory and optimal convergence rates in fast nonlinear diffusion. To appear in *J. Math. Pures Appl.*
- [3] R.J. McCann and D. Slepcev. Second-order asymptotics for the fast-diffusion equation. Submitted.

**Besov Spaces on Fractals and Conformal Geometry**

HERVÉ PAJOT

The main goal of this talk is to explain how to define classical tools of analysis (like Besov spaces or Poincaré inequalities) in singular spaces and how to use them to study some problems in conformal geometry.

Let  $(X, d, \mu)$  be a metric measure space. Assume that  $\mu$  is  $Q$ -regular, that is there exists  $C > 0$  such that

$$C^{-1}R^Q \leq \mu(B(x, R)) \leq CR^Q$$

whenever  $x \in X$  and  $R \in (0, \text{diam}X)$ . Note that  $Q$  is the Hausdorff dimension of  $(X, d)$ . We will say that  $d$  is  $Q$ -regular if the Hausdorff measure of  $d$  is  $Q$ -regular.

For any  $p \geq 1$ , the Besov  $p$ -norm of the function  $u : X \rightarrow \mathbb{R}$  is given by

$$\|u\|_p = \left( \int_X \int_X \frac{|u(x) - u(y)|^p}{d(x, y)^{2Q}} d\mu(x) d\mu(y) \right)^{1/p}.$$

Given two functions  $u$  and  $v$ , we write  $u \sim v$  if  $u - v$  is constant almost everywhere (with respect to  $\mu$ ). For  $p \geq 1$ , we define the Besov space  $B_p(d)$  by

$$B_p(d) = \{u : X \rightarrow \mathbb{R} \text{ in } L^p(\mu); \|u\|_p < +\infty\} / \sim.$$

In the Euclidean case of  $\mathbb{R}^n$ ,  $B_p(d_{\text{eucl}})$  is the classical Besov space  $B_{p,p}^{n/p}(\mathbb{R}^n)$  (if  $n/p < 1$ ).

Recall that an homeomorphism  $f : (Z_1, d_1) \rightarrow (Z_2, d_2)$  is quasisymmetric (or shortly QS) if there exists an increasing homeomorphism  $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that

$$d_1(x, y) \leq td_1(x, z) \implies d_2(f(x), f(y)) \leq \phi(t)d_2(f(x), f(z))$$

whenever  $x, y, z \in Z_1$ ,  $t \in (0, +\infty)$ . The conformal gauge of the metric space  $(X, d)$  is defined by

$$\mathcal{J}(X, d) = \{\delta \text{ distance on } X; Id : (X, d) \rightarrow (X, \delta) \text{ is QS}\}.$$

The conformal dimension of  $(X, d)$  is

$$\text{Cdim}(X, d) = \inf\{\text{Hdim}(X, \delta); \delta \in \mathcal{J}(X, d)\}.$$



In [1], first groups of  $l_p$ -cohomology associated to the conformal gauge  $\mathcal{J}(X, d)$  are constructed (if  $(X, d)$  is also compact and uniformly perfect). They are denoted by  $l_p H^1(\mathcal{J}(x, d))$ . The critical exponent of  $l_p$  cohomology is given by

$$p(\mathcal{J}(X, d)) = \inf\{p \in [1, +\infty); l_p H^1(\mathcal{J}(x, d)) \neq \{0\}\}.$$

It turns out that  $l_p H^1(\mathcal{J}(x, d))$  and  $B_p(d)$  are isomorphic as Banach spaces. Hence,

$$p(\mathcal{J}(X, d)) = \inf\{p \in [1, +\infty), B_p(d) \neq \{0\}\}.$$

The conformal dimension and the critical exponent of  $l_p$  cohomology are invariant under QS homeomorphisms. Are they equal in general ?

**Theorem** [1]. *Let  $(X, d)$  be a compact, uniformly perfect metric space. Assume that there exists  $\delta \in \mathcal{J}(x, d)$  such that  $\delta$  is  $Q_0$ -regular and admits a  $(1, Q_0)$ -Poincaré inequality (see the definition below) for some  $Q_0 > 1$ . Then,*

$$Cdim(X, d) = p(\mathcal{J}(X, d)).$$

In fact, by a result of Tyson [3],  $Q_0 = Cdim(X, d)$ . Observe that Lipschitz functions are in  $B_p(d)$  for  $p > Q_0$  and hence  $p(\mathcal{J}(X, d)) \leq Q_0$ . To get the reverse inequality, we prove that any function in  $B_p(d)$  with  $p < Q_0$  is constant almost everywhere by using the Poincaré inequality.

Let  $p \geq 1$ . We say that  $(X, d, \mu)$  supports a  $(1, p)$  Poincaré inequality if there exists a constant  $C_p > 0$  such that

$$\frac{1}{\mu_X(B)} \int_B |u - u_B| d\mu_X \leq C_p \text{diam} B \left( \frac{1}{\mu_X(B)} \int_B \rho^p d\mu_X \right)^{\frac{1}{p}}$$

whenever

- $B$  is a open ball in  $X$ ,
- $u : X \rightarrow \mathbb{R}$  is continuous and bounded in  $B$ ,
- $\rho : X \rightarrow \mathbb{R}^+$  is an upper gradient of  $u$  in  $B$ ,

and where  $u_B$  denotes the mean value of  $u$  on  $B$ . Recall that  $\rho : X \rightarrow \mathbb{R}^+$  is an upper gradient of  $u$  in  $B$  if for any  $x \in X$ , any  $y \in X$ , any rectifiable curve  $\gamma$  in  $X$  connecting  $x$  to  $y$ , we have  $|u(x) - u(y)| \leq \int_{\gamma} \rho$ . We say that  $d$  admits a  $(1, p)$ -Poincaré if  $(X, d, \mu)$  (where  $\mu$  is the Hausdorff measure of  $d$ ) supports a  $(1, p)$ -Poincaré inequality in the previous sense.

What happens in fractals, for instance for the Sierpinski carpet (SC) and the Sierpinski gasket (SG) ?

$$\text{Hdim}(SC) = \log(3)/\log(2) \quad Cdim(SC) = 1 \quad p(SC) = ???,$$

$$\text{Hdim}(SG) = \log(8)/\log(3) \quad Cdim(SC) = ??? \quad p(SC) = ???.$$

In fact, the conformal dimension of other pcf sets in the sense of Kigami (like the Sierpinski gasket) is also one (see [4]. Note that, by standard arguments, we

get  $1 + \log(2)/\log(3) \leq \text{Hdim}(SC) < \log(8)/\log(3)$ . A good general reference concerning conformal dimension and related topics is the recent book of Heinonen [2].

#### REFERENCES

- [1] M. Bourdon, H. Pajot, *Cohomologie  $l_p$  et espaces de Besov*, J. reine angew. Math. **558** (2003), 85–108.
- [2] J. Heinonen, *Lectures on analysis on metric spaces*, Universitext, Springer (2001).
- [3] J. Tyson, *Quasiconformality and quasimetry in metric measure spaces* Ann. Acad. Sci. Fenn. (Math.) **23** (1998), 525–548.
- [4] J. Tyson, J. M. Wu, *Quasiconformal dimensions of self-similar sets*, to appear in Rev. Mat. Iberoamericana.

### Stochastic Generalized Porous Media Equations

M. RÖCKNER

(joint work with J.-G. Ren, F.-Y. Wang)

Recall that for  $\Delta$ , the Laplace operator on  $\mathbb{R}^d$ , the following differential equation

$$\frac{dX_t}{dt} = \Delta(|X_t|^r \text{sign } X_t)$$

is called the porous medium equation if  $r > 1$  and the fast diffusion equation if  $r \in (0, 1)$ . A special feature of this equation is that the solution decays algebraically fast in  $t$  when  $r > 1$  and decays to zero at finite time when  $r \in (0, 1)$ . For historical remarks and recent progress on this equation, we refer to [2, 1, 11] and the references therein.

In recent years, the stochastic version of the porous medium equation has been studied intensively, see [8] for the existence, uniqueness and long-time behavior of some stochastic generalized porous media equations with finite reference measures, [12] for the stochastic porous media equation on  $\mathbb{R}^d$  where the reference (Lebesgue) measure is infinite and [3, 7] for the analysis of the corresponding Kolmogorov equations. See also [19] for large deviation results for a class of generalized porous media equations.

The basic motivation of this work is to study, for example, the following stochastic porous media and fast diffusion equation:

$$(1) \quad dX_t = L\left(\sum_{i=1}^l \alpha_i |X_t|^{r_i} \text{sign } X_t\right) + B(t, X_t) dW_t,$$

where  $L$  is a negative definite self-adjoint operator on  $L^2(\mathbf{m})$  for some  $\sigma$ -finite measure  $\mathbf{m}$ ,  $W_t$  is the Brownian motion on a reference Hilbert space, and  $B$  a properly defined linear operator from this Hilbert space to the state space of  $X_t$ . Here,  $l \in \mathbb{N}$  and  $r_i, \alpha_i > 0$ ,  $i = 1, \dots, l$ , are fixed numbers. If  $r_i < 1$  and  $r_j > 1$  for some  $i, j \leq l$ , we call this equation the stochastic porous media and fast diffusion equation. As a consequence of our main result, this equation has a unique solution as soon as  $B$  satisfies a Lipschitz type condition.

To solve such an equation, we first establish a general result for “monotone” stochastic equations, which extends the corresponding one of Krylov and Rozovskii ([13, Theorems II.2.1 and II.2.2]). As a direct application, we are able to solve (1) with  $\sum_{i=1}^l \alpha_i |X_t|^{r_i} \text{sign}(X_t)$  replaced by  $\Psi(t, X_t)$  for a more general time-dependent function  $\Psi$  comparable with an  $N$ -function. The Orlicz space (cf. [16]) determined by  $N$  plays an essential role in our analysis.

Acknowledgement. This work has been supported in part by the DFG through the Forschergruppe “Spectral Analysis, Asymptotic Distributions and Stochastic Dynamics”, the BiBoS Research Centre, NNSFC(10121101) and RFDP.

## REFERENCES

- [1] D.G. Aronson and L.A. Peletier. Large time behaviour of solutions of the porous medium equation in bounded domains. *J. Differential Equations*, 39(3):378–412, 1981.
- [2] D.G. Aronson. The porous medium equation. In *Nonlinear diffusion problems (Montecatini Terme, 1985)*, volume 1224 of *Lecture Notes in Math.*, pages 1–46. Springer, Berlin, 1986.
- [3] V. Barbu, V.I. Bogachev, G. Da Prato, and M. Röckner. Weak solution to the stochastic porous medium equations: the degenerate case. BiBoS-Preprint 04-10-161, 22 pp., 2004.
- [4] V.I. Bogachev, G. Da Prato, and M. Röckner. Invariant measures of generalized stochastic equations of porous media. *Dokl. Akad. Nauk*, 396(1):7–11, 2004.
- [5] J.L. Doob. *Stochastic processes*. John Wiley & Sons Inc., New York, 1953.
- [6] G. Da Prato and M. Röckner. Invariant measures for a stochastic porous medium equation. In *Stochastic analysis and related topics in Kyoto*, volume 41 of *Adv. Stud. Pure Math.*, pages 13–29. Math. Soc. Japan, Tokyo, 2004.
- [7] G. Da Prato and M. Röckner. Weak solutions to stochastic porous media equations. *J. Evol. Equ.*, 4(2):249–271, 2004.
- [8] G. Da Prato, M. Röckner, B. Rozovski, and F.-Y. Wang. Strong solutions to stochastic generalized porous media equations: existence, uniqueness and ergodicity. BiBoS-Preprint 04-09-157, 15 pp., to appear in *Commun. PDE*, 2004.
- [9] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1992.
- [10] A. Dembo and O. Zeitouni. *Large deviations techniques and applications*, volume 38 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, second edition, 1998.
- [11] C. Ebmeyer. Regularity in Sobolev spaces for the fast diffusion and the porous medium equation. *J. Math. Anal. Appl.*, 307(1):134–152, 2005.
- [12] J.U. Kim. On the stochastic porous medium equation. Preprint, to appear in *J. Diff. Equ.*
- [13] N.V. Krylov and B.L. Rozovskii. Stochastic evolution equations. In *Current problems in mathematics, Vol. 14 (Russian)*, pages 71–147, 256. Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1979.
- [14] N.V. Krylov. On Kolmogorov’s equations for finite-dimensional diffusions. In *Stochastic PDE’s and Kolmogorov equations in infinite dimensions (Cetraro, 1998)*, volume 1715 of *Lecture Notes in Math.*, pages 1–63. Springer, Berlin, 1999.
- [15] B.L. Rozovskii. *Stochastic evolution systems*, volume 35 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1990. Linear theory and applications to nonlinear filtering, Translated from the Russian by A. Yarkho.
- [16] M.M. Rao and Z.D. Ren. *Applications of Orlicz spaces*, volume 250 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker Inc., New York, 2002.
- [17] J.-G. Ren, M. Röckner, and F.-Y. Wang. Stochastic generalized porous media and fast diffusion equations. Preprint, publication in preparation, 22 pp., 2005.
- [18] M. Röckner and F.-Y. Wang. Concentration of invariant measures for stochastic generalized porous media equations. Preprint, publication in preparation, 2005.

- [19] M. Röckner, F.-Y. Wang, and L. Wu. Large deviations for stochastic generalized porous media equations. Preprint, publication in preparation, 2004.

### Variational convergence over metric spaces

TAKASHI SHIOYA

(joint work with K. Kuwae)

The talk is on the joint work [7, 8] with K. Kuwae.

Let  $M_i \rightarrow M$  and  $Y_i \rightarrow Y$  ( $i = 1, 2, 3, \dots$ ) be two pointed Gromov-Hausdorff convergent sequences of proper metric spaces, where ‘*proper*’ means that any bounded subset is relatively compact, and let us give measures on  $M_i$  which converge to a measure on  $M$  with respect to the measured Gromov-Hausdorff topology. We are interested in the convergence and asymptotic behavior of maps  $u_i : M_i \rightarrow Y_i$  and also energy functionals  $E_i$  defined on a family of maps from  $M_i \rightarrow Y_i$ . Note that there are several attempts to define natural energy functionals on the mapping space from  $M$  to  $Y$  by the measured metric structure of  $M$  and the metric structure of  $Y$ . We introduce a natural definition of the  $L^p$ -convergence of  $u_i : M_i \rightarrow Y_i$  to  $u : M \rightarrow Y$ ,  $p \geq 1$ , and establish a general theory for energy functionals  $E_i$  by extending the theory of variational convergences, mainly studied by Mosco [9]. Mosco introduced the asymptotic compactness of energy functionals  $\{E_i\}$ , which is a generalization of the Rellich compactness. The asymptotic compactness is useful to obtain the convergence of energy minimizers, i.e., harmonic maps, and also to investigate spectral properties in the linear case. Under a uniform bound of Poincaré constants and some property of the metric on  $M$ , we prove the asymptotic compactness of  $\{E_i\}$ . We focus on a  $\Gamma$ -convergence with the asymptotic compactness, say *compact convergence*. If  $\{E_i\}$  is asymptotically compact, it has a compact convergent subsequence. We prove that the compact convergence is equivalent to the Gromov-Hausdorff convergence of the energy-sublevel sets, which is a geometric interpretation of compact convergence. This is connected with Gromov’s study of spectral concentration, Section 3 $\frac{1}{2}$ .57 of [1]. We also prove that the compact convergence of  $E_i$  is equivalent to a convergence of associated resolvents, provided that  $Y_i$  are all  $CAT(0)$ -spaces and  $E_i$  are lower semi-continuous convex functionals. Such the resolvent was defined by Jost [2] using the Moreau-Yosida approximation. As applications of our theory, we study the approximating energy functional and its spectral property. We also obtain the compactness of energy functionals if  $M_i$  are Riemannian manifolds with a lower bound of Ricci curvature. In the case where  $E_i$  are symmetric quadratic forms on  $L^2$ -spaces, there are some applications of our theory to a homogenization problem, [11], and convergence of Dirichlet forms, [3, 4, 10], including finite-dimensional approximation problems, [5, 6].

## REFERENCES

- [1] M. Gromov, *Metric structures for Riemannian and non-Riemannian spaces*, Birkhäuser Boston Inc., Boston, MA, 1999, Based on the 1981 French original, With appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates.
- [2] J. Jost, *Convex functionals and generalized harmonic maps into spaces of nonpositive curvature*, Comment. Math. Helv. **70** (1995), no. 4, 659–673.
- [3] A. Kasue, *Convergence of Riemannian manifold and Laplace operators. II*, preprint, 2003.
- [4] ———, *Convergence of metric graphs and energy forms*, preprint, 2004.
- [5] A. V. Kolesnikov, *Convergence of Dirichlet forms with changing speed measures on  $\mathbb{R}^d$* , Forum Math. **17** (2005), no. 2, 225–259.
- [6] ———, *Convergence of infinite-dimensional Dirichlet forms with changing speed measures*, preprint.
- [7] K. Kuwae and T. Shioya, *Convergence of spectral structures: a functional analytic theory and its applications to spectral geometry*, Comm. Anal. Geom. **11** (2003), no. 4, 599–673.
- [8] ———, *Variational convergence over metric spaces*, to appear in Trans. Amer. Math. Soc.
- [9] U. Mosco, *Composite media and asymptotic Dirichlet forms*, J. Funct. Anal. **123** (1994), no. 2, 368–421.
- [10] Y. Ogura, T. Shioya, and M. Tomisaki, *Collapsing of warped product spaces and one-dimensional diffusion processes*, in preparation.
- [11] C. Vernicos, *The macroscopic sound of tori*, Pacific J. Math. **213** (2004), no. 1, 121–156.

**The principal eigenvalue for time-changed processes and applications**

MASAYOSHI TAKEDA

## 1. GAUGEABILITY FOR FEYNMAN-KAC FUNCTIONALS

Chen [1] and Takeda [3] showed that the gaugeability, that is, the integrability of Feynman-Kac functionals, or the subcriticality of Schrödinger operators is equivalent to that the principal eigenvalue of time changed process is greater than one (Theorem 17 below). The objective of my talk is to give two applications of this result.

Let  $M$  be a non-compact, complete Riemannian manifold and  $m$  the volume. Let  $\mathbb{M} = (\mathbb{P}_x, B_t)$  be the Brownian motion on  $M$ . For a domain  $D \subset M$ , let  $\mathbb{M}^D$  be the absorbing Brownian motion on  $D$ . We assume that  $\mathbb{M}^D$  is transient. Chen [1] generalized the notion of *Green-tightness* for general transient Markov processes and introduced a new Kato class. We denote by  $\mathcal{S}_\infty^D$  this class associated with  $\mathbb{M}^D$ . For  $\mu \in \mathcal{S}_\infty^D$ , define

$$\lambda(\mu; D) = \inf \left\{ \frac{1}{2} \int_D (\nabla u, \nabla v) \, dm : u \in C_0^\infty(D), \int_D u^2(x) \mu(dx) = 1 \right\}.$$

Then  $\lambda(\mu; D)$  is the principal eigenvalue of the time-changed process of  $\mathbb{M}^D$  by  $A_t^\mu$ . Here  $A_t^\mu$  is the positive continuous additive functional in the Revus correspondence with  $\mu$ . Let  $p_t^{\mu, D}(x, y)$  be the integral kernel of *Feynman-Kac semigroup*:

$$p_t^{\mu, D} f(x) := \mathbb{E}_x[\exp(A_t^\mu) f(B_t); t < \tau_D] = \int_D p_t^{\mu, D}(x, y) f(y) dy$$

$(\tau_D = \inf\{t > 0 : B_t \not\subset D\})$  and  $G^{\mu,D}(x, y) := \int_0^\infty p_t^{\mu,D}(x, y) dt.$ @

**Theorem 17.** ([1],[3]) *Let  $\mu \in \mathcal{S}_\infty^D$ . Then the following statements are equivalent:*

- (1) (i) **(gaugeability)**  $\sup_{x \in D} \mathbb{E}_x[e^{A_{\tau_D}^\mu}] < \infty;$
- (ii) **(subcriticality)**  $G^{\mu,D}(x, y) < \infty$  for  $x, y \in D, x \neq y;$
- (iii)  $\lambda(\mu; D) > 1.$

2. APPLICATIONS OF THEOREM 17

(i) Let  $\bar{\mathbb{B}} = (\bar{\mathbb{P}}_x, \bar{B}_t)$  be a branching Brownian motion with **branching rate**  $k$  and **branching mechanism**  $\{p_n(x)\}_{n \geq 2}: \bar{\mathbb{P}}_x[T > t | \sigma(B)] = \exp(-A_t^k)$  ( $T$  is the **first splitting time**),  $\sum_{n=2}^\infty p_n(x) = 1$ . Set  $Q(x) = \sum_{n \geq 2} n p_n(x)$  and  $\mu(dx) = (Q(x) - 1)k(dx)$ . We assume that  $\sup_{x \in M} Q(x) < \infty$ .

**Theorem 18.** ([4]) *Let  $K$  be a closed set with  $\text{Cap}(K) > 0$  and let  $\mu \in \mathcal{S}_\infty^D$ ,  $D = M \setminus K$ . Then*

$$\lambda(\mu; D) > 1 \iff \sup_{x \in D} \bar{\mathbb{E}}_x[N_K] < \infty.$$

Here  $N_K$  is the number of branches hitting  $K$ .

R.Z. Khas'minskii gave a sufficient condition for gaugeability known as Khas'minskii lemma, and applied it for showing the finiteness of the expectation of  $N_K$ . To prove Theorem 18, we show that for  $\mu \in \mathcal{S}_\infty^D$

$$\sup_{x \in D} \mathbb{E}_x[\exp(A_{\tau_D}^\mu)] < \infty \iff \sup_{x \in D} \mathbb{E}_x[\exp(A_{\tau_D}^\mu); \tau_D < \infty] < \infty.$$

Then the identity

$$\bar{\mathbb{E}}_x[N_K] = \mathbb{E}_x[\exp(A_{\tau_D}^\mu); \tau_D < \infty]$$

leads us to Theorem 18.

(ii) Let  $d(x, y)$  be the distance. Let  $p(t, x, y)$  be the heat kernel. Suppose that it satisfies the Gaussian lower and upper bounds (**Li-Yau estimate**): for any  $x, y \in M$  and  $t > 0$ ,

$$\frac{C_1 \exp\left(-c_1 \frac{d^2(x,y)}{t}\right)}{m(B(x, \sqrt{t}))} \leq p(t, x, y) \leq \frac{C_2 \exp\left(-c_2 \frac{d^2(x,y)}{t}\right)}{m(B(x, \sqrt{t}))}$$

( $C_1, c_1, C_2, c_2$  are positive constants.  $B(x, r) = \{y \in M : d(x, y) < r\}$ ). Let  $p^\mu(t, x, y)$  be the heat kernel of the Schrödinger operator,  $\frac{1}{2}\Delta + \mu$ .

**Theorem 19.** ([5]) *Let  $\mu \in \mathcal{S}_\infty^M$ . Then  $p^\mu(t, x, y)$  satisfies the Li-Yau estimate if and only if  $\lambda(\mu; M) > 1$ .*

**Example:** Let  $M = \mathbb{R}^d$  ( $d \geq 3$ ) and  $\mu = \sigma_r$  the surface measure of  $\{|x| = r\}$ . Then

$$\lambda(\sigma_r; \mathbb{R}^d) = \frac{d-2}{2r}.$$

Hence  $p^{\sigma_r}(t, x, y)$  satisfies the Gaussian estimate, if and only if  $r < \frac{d-2}{2}$ .

**Remark 1.** Let  $d = 3$  and  $\lambda(\mu; \mathbb{R}^3) = 1$ , for example,  $\mu = \sigma_{1/2}$ . Then

$$p^\mu(t, x, y) \simeq \frac{C_1}{t^{3/2}} \left(1 + \frac{\sqrt{t}}{\langle x \rangle}\right) \left(1 + \frac{\sqrt{t}}{\langle y \rangle}\right) \exp\left(-c_1 \frac{|x-y|^2}{t}\right)$$

( $\langle x \rangle := 1 + |x|$ ) (cf. [2, Theorem 10.15]).

**Remark 2.** Let  $\mu = \mu^+ - \mu^-$  be a signed measures such that  $\mu^+ \in \mathcal{S}_\infty^M$  and  $\sup_{x \in M} G\mu^-(x) < \infty$ . Then the Li-Yau estimate of the heat kernel  $p^\mu(t, x, y)$  is equivalent to

$$\inf\{\mathcal{E}(u, u) + \int_M u^2 d\mu^- : u \in C_0^\infty(M), \int_M u^2 d\mu^+ = 1\} > 1.$$

For  $\mu = -\mu^-$ , the result above is shown in [2, Theorem 10.5].

**Remark 3.** Theorem 19 can be extended to the symmetric  $\alpha$ -stable process ( $d > \alpha$ ). Let  $\mu \in \mathcal{S}_\infty^{\mathbb{R}^d}$  with  $\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |x-y|^{\alpha-d} d\mu(x) d\mu(y) < \infty$ . Then the heat kernel of  $-\frac{1}{2}(-\Delta)^{\alpha/2} + \mu$  satisfies

$$C_1 \left( \frac{1}{t^{d/\alpha}} \wedge \frac{t}{|x-y|^{d+\alpha}} \right) \leq p^\mu(t, x, y) \leq C_2 \left( \frac{1}{t^{d/\alpha}} \wedge \frac{t}{|x-y|^{d+\alpha}} \right),$$

if and only if

$$\lambda(\mu) = \inf\{\mathcal{E}^\alpha(u, u) : u \in C_0^\infty(\mathbb{R}^d), \int_{\mathbb{R}^d} u^2 d\mu = 1\} > 1.$$

Here

$$\mathcal{E}^{(\alpha)}(u, u) = K(\alpha, d) \iint_{\mathbb{R}^d \times \mathbb{R}^d \setminus d} \frac{(u(x) - u(y))^2}{|x-y|^{d+\alpha}} dx dy.$$

#### REFERENCES

- [1] Z.-Q. Chen, Gaugeability and conditional gaugeability, *Trans. Amer. Math. Soc.* 354 (2002), 4639-4679.
- [2] A. Grigor'yan, Heat kernels on weighted manifolds and applications, to appear in *Cont. Math.*
- [3] M. Takeda, Subcriticality and conditional gaugeability of generalized Schrödinger operators, *J. Funct. Anal.* 191 (2002), 343-376.
- [4] M. Takeda, Gaugeability for Feynman-Kac functionals with applications to symmetric  $\alpha$ -stable processes, to appear in *Proc. Amer. Math. Soc.*
- [5] M. Takeda Gaussian bounds of heat kernels for Schrödinger operators on Riemannian manifolds, preprint.

**Brownian motion on Jordan curves and stochastic calculus of variations on the diffeomorphism group of the circle**

ANTON THALMAIER

(joint work with H el ene Airault and Paul Malliavin)

We report on recent progress of constructing unitarizing probability measures for the representation of the Virasoro algebra  $\mathcal{V}$ , using methods of Stochastic Analysis.

Denote by  $\text{Diff}(S^1)$  the group of orientation preserving diffeomorphisms of the circle  $S^1$  and by  $\text{diff}(S^1)$  its Lie algebra. With the identification  $\text{diff}(S^1) \cong C^\infty(S^1; \mathbb{R})$  the Lie bracket takes the form  $[u, v] = uv' - v'u$ . We are interested in central extensions of  $\text{Diff}(S^1)$  and  $\text{diff}(S^1)$ , the *Virasoro group*, resp., *Virasoro algebra*. In explicit terms the Virasoro algebra is given (modulo equivalence) by

$$\mathcal{V} \equiv \mathcal{V}_c = \mathbb{R} \oplus \text{diff}(S^1)$$

with the Lie bracket  $[s\kappa + u, t\kappa + v] = \frac{c}{12} \omega(u, v)\kappa + [u, v]$  ( $\kappa$  *central element*,  $c > 0$  *central charge*), where  $\omega$  an antisymmetric bilinear form satisfying the *cocycle condition*  $\omega([u, v], w) + \omega([v, w], u) + \omega([w, u], v) = 0$ . Such forms have been classified by Gelfand-Fuks and are modulo the choice of constants given by the fundamental cocycle (1.2.1) [2]. There are two well-known representations of  $\mathcal{V}$ :

- on the space  $\mathcal{U}^\infty$  of univalent functions on the disk
- on the Sato Grassmannian via an embedding of  $\text{Diff}(S^1)$  into  $\text{Sp}(\infty)$ .

For both the diffeomorphism group  $\text{Diff}(S^1)$  of the circle  $S^1$  is a key object [2].

**Brownian motion on  $\text{Diff}(S^1)$ .** Denote by  $\mathcal{H} \subset \text{Diff}(S^1)$  the restrictions to  $S^1$  of the Poincar e group of homographic transformations. The Lie algebra of  $\mathcal{H}$  is  $\mathfrak{su}(1, 1)$  and generated by the vector fields  $\cos \theta, \sin \theta, 1$ .

**Theorem 20** ([3]). *There exists a unique semi-Hilbertian metric on  $\text{diff}(S^1)$  which is invariant under the adjoint action of  $\mathfrak{su}(1, 1)$ .*

Let  $\text{Diff}(S^1)/S^1$  be the diffeomorphisms of  $S^1$  which fix a point on the circle and let  $\text{diff}_0(S^1) = \{u \in \text{diff}(S^1) : \int_{S^1} u(\theta) d\theta = 0\}$ . Developing  $u \in \text{diff}_0(S^1)$  in terms of a Fourier series  $u(\theta) = \sum_{k \geq 1} (a_k \cos k\theta + b_k \sin k\theta)$ , let  $\mathcal{J}u(\theta) = \sum_{k \geq 1} (-a_k \sin k\theta + b_k \cos k\theta)$ . Then  $\mathcal{J}^2 = -\text{id}$  and  $\mathcal{J}$  defines an *almost complex structure* on  $\text{diff}_0(S^1)$  which is integrable. The *K ahler metric* on  $\text{diff}_0(S^1)$  is

$$\|u\|^2 := \omega(u, \mathcal{J}u) = \sum_{k \geq 1} \alpha_k^2 (a_k^2 + b_k^2), \quad \alpha_k = (k^3 - k)^{1/2}.$$

The Lie subalgebra of  $\text{diff}(S^1)$  generated by  $1, \sin, \cos$  is  $\mathfrak{su}(1, 1)$ , and

$$\text{diff}(S^1)/\mathfrak{su}(1, 1) = \left\{ \theta \mapsto \sum_{k \geq 2} (a_k \cos k\theta + b_k \sin k\theta) \right\} \hat{=} H^{3/2}(S^1).$$

Brownian motion on the *universal Teichm uller space*  $\text{Diff}(S^1)/\text{SU}(1, 1)$ , with respect to the  $H^{3/2}$  metric, is constructed by exponentiating BM on  $\text{diff}(S^1)$ , see



[10], [5], [8]. Let  $x_k(t)$  be an independent sequence of scalar Brownian motions, and let

$$e_{2k+1} := \alpha_k^{-1} \sin k\theta, \quad e_{2k} := \alpha_k^{-1} \cos k\theta, \quad \alpha_k = \sqrt{k^3 - k}, \quad k > 1.$$

Brownian motion on  $\text{Diff}(S^1)$  is *formally* a solution of the Stratonovich SDE

$$(*) \quad dg_t = \sum_{k>1} \left( e_{2k}(g_t) \circ dx_{2k}(t) + e_{2k+1}(g_t) \circ dx_{2k+1}(t) \right).$$

Solving (\*) via *Abel regularization*, we fix  $\rho > 0$  and replace  $\alpha_k$  by  $\sqrt{k^3 - k}/\rho^k$ . For  $\rho < 1$ , the theory of stochastic flow of diffeomorphisms guarantees a unique solution  $g^\rho$  which takes values in the smooth diffeomorphisms of  $S^1$ .

**Theorem 21 (Limiting case  $\rho = 1$ ).** *As  $\rho \nearrow 1$ , the limit  $g_t(\theta)$  of  $g_t^\rho(\theta)$  exists uniformly in  $\theta$  and defines a solution of (\*), but  $(\theta \mapsto g_t^\rho(\theta)) \in \text{Homeo}(S^1)$  only.*

**Brownian motion on Jordan curves** [4]. Let  $\mathcal{J} = \{\Gamma \subset \mathbb{C} : \Gamma \text{ Jordan curve}\}$ . To  $\Gamma \in \mathcal{J}$  there is a continuous injective parametrization  $\varphi : S^1 \rightarrow \mathbb{C}$  such that  $\varphi(S^1) = \Gamma$ . *Holomorphic parametrizations* are constructed as follows. Each Jordan curve  $\Gamma$  splits the complex plane into two simply connected domains  $D_\Gamma^+$ ,  $D_\Gamma^-$ . Let  $D$  be the open unit disc in  $\mathbb{C}$ . By the Riemann mapping theorem, there exist biholomorphic mappings  $F^+ : D \rightarrow D_\Gamma^+$  and  $F^- : D \rightarrow D_\Gamma^-$  (unique modulo  $\text{SU}(1, 1)$ ) which by Caratheodory extend to homeomorphisms  $F^+ : \bar{D} \rightarrow \bar{D}_\Gamma^+$ ,  $F^- : \bar{D} \rightarrow \bar{D}_\Gamma^-$ . Then  $F^\pm|_{S^1}$  parametrize  $\Gamma$ , and  $g_\Gamma := (F^+)^{-1} \circ F^-|_{S^1}$  defines an orientation preserving homeomorphism of  $S^1$ . Now let  $\mathcal{J}^\infty := C^\infty$  Jordan curves

**Theorem 22** (Beurling-Ahlfors “conformal welding”). *The mapping  $\Gamma \mapsto g_\Gamma$  from  $\mathcal{J}^\infty$  to  $\text{Diff}(S^1)$  is surjective and induces a canonical isomorphism*

$$\mathcal{J}^\infty \cong \text{SU}(1, 1) \backslash \text{Diff}(S^1) / \text{SU}(1, 1).$$

To construct BM on Jordan curves from BM “on  $\text{Diff}(S^1)$ ” we have to factorize  $g_t$  as  $g_t = (F_t^+)^{-1} \circ F_t^-|_{S^1}$ . By conformal welding we first associate to the regularized process  $t \mapsto {}^\rho g_t$  on  $\text{Diff}(S^1)$  a process  $t \mapsto {}^\rho \Gamma_t$  with values in  $\mathcal{J}^\infty$ .

**Theorem 23** (Brownian motion on Jordan curves [4]). *As  $\rho \nearrow 1$ , the regularized process  $t \mapsto {}^\rho \Gamma_t$  converges to a diffusion on the space of continuous Jordan curves.*

Let  $\mathcal{U}^+$  denote the space of univalent functions on  $D$  which are continuous and injective on  $\bar{D}$ , resp.  $\mathcal{U}^-$  the equivalent space of functions, univalent on  $\{z \in \mathbb{C} : |z| > 1\}$ , continuous and injective on  $\{z \in \mathbb{C} : |z| \geq 1\}$ .

**Theorem 24** (Stochastic Sewing Theorem). *There exist processes  $f_t^+$ ,  $f_t^-$ , taking values in  $\mathcal{U}^+$ , resp. in  $\mathcal{U}^-$ , such that  $g_t = (f_t^+)^{-1} \circ f_t^-|_{S^1}$ .*

**Brownian motion on univalent functions.** Denote by  $\mathcal{U}^\infty$  the space of univalent functions  $f \in C^\infty(\bar{D}; \mathbb{C})$ , normalized by  $f(0) = 0$ ,  $f'(0) = 1$ . To  $f \in \mathcal{U}^\infty$  we associate the  $C^\infty$  Jordan curve  $\Gamma = f(\partial D)$  around the origin 0. The Riemann mapping theorem provides a biholomorphic map  $h_f : \{z \in \mathbb{C} : |z| > 1\} \xrightarrow{\sim} D_\Gamma^-$  with  $h_f(\infty) = \infty$ , which extends to a diffeomorphism  $h_f : \{z \in \mathbb{C} : |z| \geq 1\} \xrightarrow{\sim} \bar{D}_\Gamma^-$ .

By Kirillov (1982), the mapping  $\mathcal{U}^\infty \ni f \mapsto g_f := (f^{-1} \circ h_f)|_{S^1} \in \text{Diff}(S^1)$  induces an isomorphism  $\mathcal{U}^\infty \cong \text{Diff}(S^1)/S^1$ . In particular,  $\mathcal{U}^\infty$  is a homogeneous space under the left action of  $\text{Diff}(S^1)$ .

**Theorem 25** ([4]). *As  $\rho \nearrow 1$ , the regularized process with values in  $\mathcal{U}^\infty$  constructed from the motion  $t \mapsto \rho g_t$  on  $\text{Diff}(S^1)$  converges to a diffusion  $t \mapsto f_t$  taking values in the univalent functions on  $D$  which are continuous and injective on  $\bar{D}$ .*

**Unitarizing measures.** We read  $u \in \text{diff}(S^1)$  as right-invariant vector field  $\hat{u}$  on  $\text{Diff}(S^1)$ , and consider the Neretin differential form  $\Omega$  on  $\text{Diff}(S^1)$ , see (2.2.5) [2].

**Theorem 26.** *Let  $\mathcal{M} = \mathcal{U}^\infty = \text{Diff}(S^1)/S^1$  and denote by  $\mathcal{H}(\mathcal{M})$  the space of holomorphic functionals on  $\mathcal{M}$ . For any  $c > 1$ , we get a representation  $\rho$  of  $\mathcal{V}_c$  on  $\mathcal{H}(\mathcal{M})$  by defining  $\rho(u)\phi = \langle \hat{u}, d\phi \rangle + c \langle \hat{u}, \Omega \rangle$  and  $\rho(\kappa)\phi = i\phi c/12$ .*

There exists a function  $K$  (“Kähler potential”) on  $\text{Diff}(S^1)$ , invariant under  $\text{SU}(1, 1)$ , such that  $\partial\bar{\partial}K = \omega$  and  $3\bar{\partial}K = \Omega$ . Unitarizing measures should formally be of the type “ $\mu_\gamma = c_0 \exp(-cK) d\text{vol}$ ” where  $d\text{vol} \equiv$  “volume measure” of the infinite dimensional Kähler manifold  $\Delta_{\mathcal{M}_1} := \text{Diff}(S^1)/\Gamma$  with  $\Gamma = \text{SU}(1, 1)$

Consider the horizontal Laplacian  $\Delta^H = \frac{1}{2} \sum (\partial_{e_k}^r)^2$  on  $\text{Diff}(S^1)$  and the Laplace Beltrami operator  $\Delta_{\mathcal{M}_1}$  to the Kähler metric on  $\mathcal{M}_1$ . Let  $\pi$  be the projection from  $\text{Diff}(S^1)$  to  $\mathcal{M}_1$ . Then  $\Delta^H(\phi \circ \pi) = (\Delta_{\mathcal{M}_1}\phi) \circ \pi$ . By solving an appropriate SDE, a diffusion on  $\mathcal{M}_1$  is constructed to the operator  $\Delta_{\mathcal{M}_1} - c \nabla K \cdot \nabla$  where  $\nabla K \cdot \nabla = \sum_k (\partial_{e_k}^r K) \partial_{e_k}^r$ . The vector field  $\nabla K$  is called *unitarizing drift*.

**Theorem 27.** *The Ornstein-Uhlenbeck process associated to  $\Delta_{\mathcal{M}_1} - c \nabla K \cdot \nabla$  exists on the group of Hölder homeomorphisms.*

**Theorem 28** (Airault-Malliavin-Thalmaier [3]). *An unitarizing measure must be supported by  $\text{Homeo}(S^1)/\Gamma$  and will be an invariant measure for the Ornstein-Uhlenbeck diffusion to  $\Delta_{\mathcal{M}_1} - c \nabla K \cdot \nabla$ , where  $c$  is the central charge.*

**Theorem 29** (Bowick-Rajeev [6], Airault [1], Gordina-Lescot [9]).

$$\text{Ricci}(\text{Diff}(S^1)/\Gamma) = -13/6 \times \text{identity}.$$

**Theorem 30.** *For  $\gamma > 13/6$  an invariant measure exists and is unique.*

The proof uses the method of “confinement under curvature positivity” [7].

#### REFERENCES

- [1] H. Airault, *Riemannian connections and curvatures on the universal Teichmüller space*, C. R. Acad. Sci. Paris, Ser. I, **341** (2005), 253–258.
- [2] H. Airault, P. Malliavin, *Unitarizing probability measures for representations of Virasoro algebra*, J. Math. Pures Appl. **80** (2001), 627–667.
- [3] H. Airault, P. Malliavin and A. Thalmaier, *Support of Virasoro unitarizing measures*, C. R. Acad. Sci. Paris, Ser. I, **335** (2002), 621–626.
- [4] H. Airault, P. Malliavin and A. Thalmaier, *Canonical Brownian motion on the space of Jordan curves and resolution of Beltrami equations by a continuity method along stochastic flows*, J. Math. Pures Appl. **83** (2004), 955–1018.
- [5] H. Airault, J. Ren, *Modulus of continuity of the canonic Brownian motion “on the group of diffeomorphisms of the circle”*, J. Funct. Anal. **196** (2002) 395–426.

- [6] M.J. Bowik, A. Lahiri, *The Ricci curvature of  $\text{Diff } S^1/\text{SL}(2, \mathbb{R}^n)$* , J. Math. Phys. **29** (1988), no. 9, 1979–1981.
- [7] A.-B. Cruzeiro, P. Malliavin, *Nonperturbative construction of invariant measure through confinement by curvature*, J. Math. Pures Appl. (9) **77** (1998), no. 6, 527–537.
- [8] S. Fang, *Canonical Brownian motion on the diffeomorphism group of the circle*, J. Funct. Anal. **196** (2002) 162–179.
- [9] M. Gordina, P. Lescot, *Riemannian geometry of  $\text{Diff}(S^1)/S^1$  and representations of the Virasoro algebra*, J. Funct. Anal. (2006).
- [10] P. Malliavin, *The canonic diffusion above the diffeomorphism group of the circle*, C. R. Acad. Sci. Paris Sér. I Math. **329** (1999) 325–329.

## Optimal transport, Ricci curvature and synthetic geometry

CEDRIC VILLANI

In this survey talk, I reported on a recent direction of research mixing analysis, geometry and probability theory. I first illustrated the power of monotone changes of variables with an elementary proof of optimal Sobolev inequalities (joint works with Cordero-Erausquin, Nazaret and Maggi). On a Riemannian manifold, there is a “canonical” construction of monotone maps based on optimal transport theory with cost equal to the square of the geodesic distance. This leads to the interplay between optimal transport and Riemannian geometry, in particular Ricci curvature lower bounds (Otto and I; Cordero-Erausquin, McCann and Schmuckenschläger). These ideas were recently used by Lott and I on one hand, Sturm on the other hand, to start the development of a synthetic theory of Ricci curvature lower bounds on metric-measure length spaces: curvature-dimension bounds  $\text{CD}(K, N)$  are defined in terms of the behavior of certain nonlinear convex functions of the density along geodesics of optimal transport.

Much more information can be found in the following references:

### REFERENCES

- [1] CORDERO-ERAUSQUIN, D., NAZARET, B. AND VILLANI, C. A new approach to sharp Sobolev and Gagliardo-Nirenberg inequalities. *Adv. Math.* **182** (2004), 307–332.
- [2] CORDERO-ERAUSQUIN, D., MCCANN, R.J. AND SCHMUCKENSCHLÄGER, M. A Riemannian interpolation inequality à la Borell, Brascamp and Lieb, *Inv. Math.* **146** (2001), 219–257.
- [3] LOTT, J. AND VILLANI, C. Ricci curvature for metric-measure spaces via optimal transport. Preprint (2005), available online via <http://www.umpa.ens-lyon.fr/~cvillani>
- [4] MAGGI, F. AND VILLANI, C. Balls have the worst best Sobolev inequalities. *J. Geom. Anal.* **15**, 1 (2005), 83–121.
- [5] OTTO, F. AND VILLANI, C. Generalization of an inequality by Talagrand, and links with the logarithmic Sobolev inequality, *J. Funct. Anal.* **173** (2000), 361–400.
- [6] STURM, K.-T. On the geometry of metric measure spaces, I, II. Preprints (2004–2005), available online at <http://www-wt.iam.uni-bonn.de/~sturm/en/index.html>.
- [7] VILLANI, C. Topics in Optimal Transportation. *Graduate Studies in Mathematics* 58, American Mathematical Society, Providence (2003).
- [8] VILLANI, C. Optimal Transport, old and new. *Saint-Flour 2005 Lecture Notes*, preliminary version available online via <http://www.umpa.ens-lyon.fr/~cvillani>

## Local Poincare Inequality by Transportation

MAX-K VON RENESSE

The sharp geometric properties of optimal mass transportation [2], [8] can be used to develop a theory of singular metric measure spaces with generalized lower Ricci curvature bounds [7], [3].

The notion of dimension-curvature bounds obtained in this framework are a priori global in nature and do not allow for simple localization unless additional geodesic convexity of the respective domain is assumed.

In particular it is not at all obvious how to obtain local Poincare type inequalities which are known as a crucial ingredient for regularity and heat kernel estimates on metric measure spaces. For instance, the ad hoc technique of expanding the global logarithmic Sobolev inequality gives only a global Poincare estimate in the positive curvature case [3].

We show how to derive local Poincare inequalities from the related concept of the qualitative Measure Contraction Property (MCP) introduced in [5] and which is implied by the curvature-dimension bounds under weak additional assumptions. The MCP condition is a substitute for the Jacobian estimate for the exponential map in the smooth Riemannian case and it suffices to carry out the argument in [1] in order to obtain the 'segment inequality'. The latter implies a weak (1,1)-Poincare inequality in the sense of upper gradients.

The result is very similar to the work [6] which assumes a 'strong local doubling condition'. A different approach to local Poincare inequalities based on the notion of 'democratic coupling' in the non negative curvature case is taken in [4].

### REFERENCES

- [1] Jeff Cheeger and Tobias H. Colding. Lower bounds on Ricci curvature and the almost rigidity of warped products. *Ann. of Math. (2)*, 144(1):189–237, 1996.
- [2] Dario Cordero-Erausquin, Robert J. McCann, and Michael Schmuckenschläger. A Riemannian interpolation inequality à la Borell, Brascamp and Lieb. *Invent. Math.*, 146(2):219–257, 2001.
- [3] John Lott and Cedric Villani. Ricci curvature for metric measure spaces via optimal transportation. 2004. Preprint, <http://arxiv.org/abs/math.DG/0412127>.
- [4] John Lott and Cedric Villani. Weak curvature conditions and Poincaré inequalities 2005. Preprint
- [5] Shin-Ichi Ohta. On measure contraction property of metric measure spaces. 2005. Preprint
- [6] Alireza Ranjbar-Motlagh. On the Poincaré inequality for abstract spaces. 2002. Preprint, Sharif University of Technology, Teheran.
- [7] Karl-Theodor Sturm. Generalized Ricci bounds and convergence of metric measure spaces. *C. R. Math. Acad. Sci. Paris*, 340(3):235–238, 2005.
- [8] Max-K von Renesse and Karl-Theodor Sturm. Transport inequalities, gradient estimates, entropy and Ricci curvature. *Comm. Pure Appl. Math.*, 58(7):923–940, 2004.

### The spectrum of the averaging operator on a network

WOLFGANG WOESS

(joint work with Donald I. Cartwright)

Let  $X$  be a countable, connected graph with symmetric neighbourhood relation  $\sim$  and without loops and multiple edges. We view it as a one-complex, where each edge is a (homeomorphic) copy of the unit interval and edges are glued together at common endpoints (vertices). We write  $X^0$  for the vertex set and  $X^1$  for the one-skeleton of  $X$ . Every point in  $X^1$  is of the form  $(xy, t)$ , the point at distance  $t$  from  $x$  on the non-oriented edge  $[x, y] = [y, x]$ , where  $0 \leq t \leq 1$ , and  $x, y \in X^0$ ,  $x \sim y$ . Thus,  $(xy, 0) = x$  and  $(xy, t) = (yx, 1 - t)$ . The discrete graph metric  $d(\cdot, \cdot)$  on the vertex set has a natural extension to  $X^1$ . We equip each edge  $[x, y]$  with a positive *conductance*

$c(xy) = c(yx)$ . On  $X^0$ , we consider the discrete measure  $\mathfrak{m}^0$ , where  $\mathfrak{m}^0(x) = \sum_{y: y \sim x} c(xy)$ , and assume that

$\mathfrak{m}^0(x) < \infty$  for all  $x \in X^0$ . On  $X^1$ , we introduce the

continuous weighted ‘‘Lebesgue’’ measure  $\mathfrak{m}^1$  which at the point

$(xy, t)$  is given by  $c(xy) \cdot dt$ , if  $0 < t < 1$  (the vertex set has  $\mathfrak{m}^1$ -measure 0).

The pair  $(X, c)$ , together with these measures, is called a *network*. On a network, there are three natural operators.

The first is the *transition operator*  $P$  acting on functions  $g : X^0 \rightarrow \mathbb{C}$  by

$$Pg(x) = \frac{1}{\mathfrak{m}^0(x)} \sum_{y: y \sim x} c(xy) g(y).$$

The second is the *Laplace operator*  $\Delta$ . It can be defined via Dirichlet form theory, or by considering the space of all continuous functions  $F : X^1 \rightarrow \mathbb{C}$  which are twice differentiable in the interior of each edge and satisfy the Kirchhoff equations

$$\sum_{y: y \sim x} c(xy) F'(xy, 0+) = 0 \quad \text{for all } x \in X^0.$$

We then have

$$\Delta F(xy, t) = F''(xy, t),$$

the 2nd derivative with respect to  $t \in (0, 1)$ , and  $\Delta$  has to be closed suitably. See e.g. CATTANEO [1], SOLOMYAK [3] or EELLS AND FUGLEDE [2] for precise details.

The third operator, and main object of the talk, is the

*averaging operator*  $A$  over balls of radius 1. It acts on locally integrable functions  $F : X^1 \rightarrow \mathbb{C}$  by

$$\begin{aligned} AF(xy, t) &= \frac{1}{\mathfrak{m}^0(x)} \sum_{u \sim x} c(xu) \int_0^{1-t} F(xu, s) ds \\ &\quad + \frac{1}{\mathfrak{m}^0(y)} \sum_{v \sim y} c(yv) \int_0^t F(yv, s) ds. \end{aligned}$$

In the *regular* case, i.e., when  $\mathbf{m}^0(\cdot)$  is constant, this is just the  $\mathbf{m}^1$ -average of  $F$  over the ball with radius 1 centered at  $(xy, t)$ .

We are interested in the relation between the spectra of the operators  $A$  and  $P$ . CATTANEO [1] has given a complete description of the  $H^2$ -spectrum of  $\Delta$  in terms of the  $\ell^2$ -spectrum of  $P$ . Our plan is to describe the  $L^2$ -spectrum of  $A$  in terms of the  $\ell^2$ -spectrum of  $P$ .

This refers to the (complex) Hilbert spaces  $L^2(X^1, \mathbf{m}^1)$  and  $\ell^2(X^0, \mathbf{m}^0)$ . The inner product of the latter is given by

$$\langle g_1, g_2 \rangle = \sum_{x \in X^0} g_1(x) \overline{g_2(x)} \mathbf{m}^0(x),$$

and  $P$  is self-adjoint with  $\|P\| \leq 1$  on this space.

Analogously, the inner product on  $L^2(X^1, \mathbf{m}^1)$  is

$$\langle F_1, F_2 \rangle = \frac{1}{2} \sum_{x \in X^0} \sum_{y \in X^0: y \sim x} c(xy) \int_0^1 F_1(xy, t) \overline{F_2(xy, t)} dt.$$

Again,  $A$  is self-adjoint with norm bounded by 1 on  $L^2(X^1, \mathbf{m}^1)$ .

There is a large body of literature on the spectrum of transition (resp. adjacency and discrete Laplace) operators on finite graphs.

Since not much work has been done regarding the spectra of averaging operators on networks, it appears to be useful to have a method for translating the spectrum of  $P$  into the spectrum of  $A$ . Our main results are the following.

**Theorem 1.** *The spectrum of  $A$  is*

$$\text{spec}(A) = \{0\} \cup \left\{ \frac{\sin \omega}{\omega} : \omega \in \mathbb{R} \setminus \{0\}, \cos \omega \in \text{spec}(P) \right\} \cup \{1 : 1 \in \text{spec}(P)\}.$$

**Corollary.** *Let  $\rho = \rho(P)$  denote the spectral radius of  $P$ . Then the spectral radius of  $A$  is*

$$\rho(A) = \begin{cases} 1, & \text{if } \rho = 1, \\ \sqrt{1 - \rho^2} / \arccos(\rho), & \text{if } \rho < 1. \end{cases}$$

Let  $\text{spec}_p(P)$  denote the point spectrum of  $P$ , i.e., the set of  $\ell^2(X^0, \mathbf{m}^0)$ -eigenvalues of  $P$ .

**Theorem 2.** *We have*

$$\text{spec}_p(A) \setminus \{0\} = \{1 : 1 \in \text{spec}_p(P)\} \cup \left\{ \frac{\sin \omega}{\omega} : \omega \in \mathbb{R} \setminus \{0\}, \cos \omega \in \text{spec}_p(P) \right\}.$$

Moreover,  $0 \in \text{spec}_p(A)$  unless  $m^0(X^0) = \infty$  and  $X$  is a tree with the property that after removal of any edge, at least one of the two connected components is recurrent.

## REFERENCES

- [1] Cattaneo, Carla: *The spectrum of the continuous Laplacian on a graph*, Monatsh. Math. **124** (1997), 215–235.
- [2] Eells, J. and Fuglede, B.: *Harmonic maps between Riemannian polyhedra*. With a preface by M. Gromov. Cambridge Tracts in Mathematics, **142**. Cambridge University Press, Cambridge, 2001.
- [3] Solomyak, Michael: *On the spectrum of the Laplacian on regular metric trees*, Waves in Random Media **14** (2004) 155–171.

## Participants

**Prof. Dr. Pascal Auscher**  
pascal.auscher@math.u-psud.fr  
Mathematiques  
Universite Paris Sud (Paris XI)  
Centre d'Orsay, Batiment 425  
F-91405 Orsay Cedex

**Kathrin Bacher**  
kathrin.bacher@web.de  
Mechenstr. 15-17  
53129 Bonn

**Prof. Dr. Zoltan Balogh**  
zoltan.balogh@math-stat.unibe.ch  
Mathematisches Institut  
Universität Bern  
Sidlerstr. 5  
CH-3012 Bern

**Davide Barbieri**  
barbier@dm.unibo.it  
Dipartimento di Matematica  
Universita degli Studi di Bologna  
Piazza di Porta S. Donato, 5  
I-40126 Bologna

**Prof. Dr. Martin T. Barlow**  
barlow@math.ubc.ca  
Dept. of Mathematics  
University of British Columbia  
1984 Mathematics Road  
Vancouver, BC V6T 1Z2  
CANADA

**Prof. Dr. Richard F. Bass**  
bass@math.uconn.edu  
Department of Mathematics  
University of Connecticut  
196, Auditorium Road  
Storrs, CT 06269-3009  
USA

**Besma Ben Ali**  
besmath@yahoo.fr  
Mathematiques  
Universite Paris Sud (Paris XI)  
Centre d'Orsay, Batiment 425  
F-91405 Orsay Cedex

**Prof. Dr. Alexander Bendikov**  
bendikov@math.uni.wroc.pl  
Institute of Mathematics  
Wroclaw University  
pl. Grunwaldzki 2/4  
50-384 Wroclaw  
POLAND

**Fabrice Blache**  
blache@wiener.iam.uni-bonn.de  
Institut für Angewandte Mathematik  
Universität Bonn  
Wegelerstr. 6  
53115 Bonn

**Anca-Iuliana Bonciocat**  
anca@wiener.iam.uni-bonn.de  
Institut für Angewandte Mathematik  
Universität Bonn  
Wegelerstr. 6  
53115 Bonn

**Dr. Matteo Bonforte**  
bonforte@calvino.polito.it  
matteo\_bonforte@yahoo.it  
Dipartimento di Matematica  
Politecnico di Torino  
Corso Duca degli Abruzzi, 24  
I-10129 Torino



**Prof. Dr. Krzysztof Burdzy**

burdzy@math.washington.edu  
Department of Mathematics  
University of Washington  
Padelford Hall  
Box 354350  
Seattle, WA 98195-4350  
USA

**Prof. Dr. Eric Carlen**

carlen@math.gatech.edu  
School of Mathematics  
Georgia Institute of Technology  
Atlanta, GA 30332-0160  
USA

**Prof. Dr. Gilles Carron**

Gilles.Carron@math.univ-nantes.fr  
Dpt. Mathematiques  
Université de Nantes  
2 rue de la Houssinière  
F-44322 Nantes Cedex 03

**Prof. Dr. Zhenqing Chen**

zchen@math.washington.edu  
Department of Mathematics  
University of Washington  
Seattle, WA 98195  
USA

**Tom Christiansen**

tom@wiener.iam.uni-bonn.de  
Institut für Angewandte Mathematik  
Universität Bonn  
Wegelerstr. 6  
53115 Bonn

**Prof. Dr. Dario Cordero-Erausquin**

cordero@math.univ-mlv.fr  
Laboratoire d'Analyse et de  
Mathematiques Appliquees  
Universite de Marne la Vallee  
F-77454 Marne la Vallee Cedex 2

**Prof. Dr. Thierry Coulhon**

Thierry.Coulhon@math.u-cergy.fr  
Departement de Mathematiques  
Universite de Cergy-Pontoise  
Site Saint-Martin  
2, avenue Adolphe Chauvin  
F-95302 Cergy-Pontoise Cedex

**Prof. Dr. Thierry Delmotte**

delmotte@cict.fr  
Laboratoire de  
Statistiques et Probabilites  
Universite Paul Sabatier  
118 route de Narbonne  
F-31062 Toulouse Cedex

**Judith M. Dohmann**

dohmann@math.uni-bielefeld.de  
Fakultät für Mathematik  
Universität Bielefeld  
Universitätsstr. 25  
33615 Bielefeld

**Prof. Dr. Bruce K. Driver**

driver@math.ucsd.edu  
Dept. of Mathematics  
University of California, San Diego  
9500 Gilman Drive  
La Jolla, CA 92093-0112  
USA

**Prof. Dr. Andreas Eberle**

eberle@uni-bonn.de  
Mathematisches Institut  
Universität Bonn  
Beringstr. 6  
53115 Bonn

**Prof. Dr. Kenneth David Elworthy**

kde@maths.warwick.ac.uk  
Mathematics Institute  
University of Warwick  
Gibbet Hill Road  
GB-Coventry, CV4 7AL

**Dr. Uta Freiberg**

uta@mathematik.uni-jena.de  
Mathematisches Institut  
Universität Jena  
Ernst-Abbe-Platz 1-4  
07743 Jena

**Prof. Dr. Masatoshi Fukushima**

fuku@ipcku.kansai-u.ac.jp  
Dept. of Mathematics  
Faculty of Engineering  
Kansai University  
Suita  
Osaka 564-8680  
JAPAN

**Prof. Dr. Friedrich Götze**

goetze@math.uni-bielefeld.de  
Fakultät für Mathematik  
Universität Bielefeld  
Universitätsstr. 25  
33615 Bielefeld

**Dr. Masha Gordina**

gordina@math.uconn.edu  
Department of Mathematics  
University of Connecticut  
196, Auditorium Road  
Storrs, CT 06269-3009  
USA

**Prof. Dr. Ben Hambly**

hambly@maths.ox.ac.uk  
Mathematical Institute  
Oxford University  
24-29, St. Giles  
GB-Oxford OX1 3LB

**Prof. Dr. Elton P. Hsu**

elton@math.northwestern.edu  
Department of Mathematics  
Northwestern University  
2033 Sheridan Road  
Evanston, IL 60208-2730  
USA

**Nicolas Juillet**

Nicolas.Juillet@ujf-grenoble.fr  
Université de Grenoble I  
Institut Fourier, UMR 5582 du CNRS  
100, rue des Maths  
BP 74  
F-38402 St. Martin d'Herès

**Dr. Moritz Kassmann**

kassmann@iam.uni-bonn.de  
Institut für Angewandte Mathematik  
Universität Bonn  
Berlingstr. 6  
53115 Bonn

**Prof. Dr. Jun Kigami**

kigami@i.kyoto-u.ac.jp  
Graduate School of Informatics  
Kyoto University  
Kyoto 606-8501  
JAPAN

**Prof. Dr. Pekka Koskela**

pkoskela@math.jyu.fi  
Dept. of Mathematics  
University of Jyväskylä  
P.O.Box 35  
FIN-40351 Jyväskylä

**Prof. Dr. Takashi Kumagai**

kumagai@kurims.kyoto-u.ac.jp  
Research Institute for  
Mathematical Sciences  
Kyoto University  
Kitashirakawa, Sakyo-ku  
Kyoto 606-8502  
JAPAN

**Prof. Dr. Kazuhiro Kuwae**

kuwae@gpo.kumamoto-u.ac.jp  
Department of Mathematics  
Faculty of Education  
Kumamoto University  
Kumamoto 860-8555  
JAPAN

**Prof. Dr. Michel Ledoux**

ledoux@math.ups-tlse.fr  
Institut de Mathematiques  
Universite de Toulouse  
118, route de Narbonne  
F-31062 Toulouse Cedex 4

**Prof. Dr. Yves Le Jan**

yves.lejan@math.u-psud.fr  
Mathematiques  
Universite Paris Sud (Paris XI)  
Centre d'Orsay, Batiment 425  
F-91405 Orsay Cedex

**Prof. Dr. Pierre Mathieu**

pierre.mathieu@gyptis.univ-mrs.fr  
Centre de Mathematiques et  
d'Informatique  
Universite de Provence  
39, Rue Joliot-Curie  
F-13453 Marseille Cedex 13

**Prof. Dr. Robert J. McCann**

mccann@math.toronto.edu  
Department of Mathematics  
University of Toronto  
100 St. George Str.  
Toronto Ont. M5S 3G3  
CANADA

**Shin-ichi Ohta**

sohta@math.kyoto-u.ac.jp  
Dept. of Mathematics  
Faculty of Science  
Kyoto University  
Kitashirakawa, Sakyo-ku  
Kyoto 606-8502  
JAPAN

**Prof. Dr. Herve Pajot**

herve.pajot@ujf-grenoble.fr  
Institut Fourier  
Universite de Grenoble I  
BP 74  
F-38402 Saint-Martin d'Herès -Cedex

**Robert Philipowski**

philipowski@wiener.iam.uni-bonn.de  
Institut für Angewandte Mathematik  
Universität Bonn  
Wegelestr. 6  
53115 Bonn

**Prof. Dr. Jean Picard**

picard@math.univ-bpclermont.fr  
Jean.Picard@math.univ-bpclermont.fr  
Lab. de Mathematiques  
Universite Blaise Pascal  
Les Cezaux  
F-63177 Aubiere Cedex

**Cesar Poupaud**

Cesar.Poupaud@math.u-bordeaux.fr  
Universite Bordeaux 1  
Lab. Bordelais d'Analyse et  
Geometrie-CNRS  
351, Cours de la Liberation  
F-33405 Talence

**Dr. Max von Renesse**

mrenesse@math.tu-berlin.de  
Institut für Mathematik / FB 3  
Skr. MA 7-5  
Technische Universität Berlin  
Straße des 17. Juni 136  
10623 Berlin

**Prof. Dr. Michael Röckner**

roeckner@math.purdue.edu  
Department of Mathematics  
Purdue University  
150 N. University Street  
West Lafayette IN 47907-2067  
USA

**Ann-Kathrin Roewer**

roewer@wiener.iam.uni-bonn.de  
Mathematisches Institut  
Universität Bonn  
Berlingstr. 1  
53115 Bonn

**Prof. Dr. Ludger Rüschemdorf**  
ruschen@stochastik.uni-freiburg.de  
Institut für Mathematische  
Stochastik  
Universität Freiburg  
Eckerstr. 1  
79104 Freiburg

**Prof. Dr. Takashi Shioya**  
shioya@math.tohoku.ac.jp  
Mathematical Institute  
Tohoku University  
Aramaki, Aoba-Ku  
Sendai, 980-8578  
JAPAN

**Prof. Dr. Karl-Theodor Sturm**  
sturm@uni-bonn.de  
sturm@wiener.iam.uni-bonn.de  
Institut für Angewandte Mathematik  
Universität Bonn  
Wegelerstr. 6  
53115 Bonn

**Prof. Dr. Masayoshi Takeda**  
takeda@math.tohoku.ac.jp  
Mathematical Institute  
Tohoku University  
Aramaki, Aoba-Ku  
Sendai, 980-8578  
JAPAN

**Romain Tessera**  
Romain.Tessera@ens.fr  
tessera@clipper.ens.fr  
Universite de Cergy-Pontoise  
6, rue Vulpian  
F-75013 Paris

**Prof. Dr. Anton Thalmaier**  
thalmaier@math.univ-poitiers.fr  
Universite de Poitiers  
Departement de Mathematiques  
Teleport 2 - BP 30179  
Blvd. Marie et Pierre Curie  
F-86962 Futuroscope Chasseneuil Cedex

**Dr. Cedric Villani**  
cvillani@umpa.ens-lyon.fr  
Dept. de Mathematiques, U.M.P.A.  
Ecole Normale Superieure de Lyon  
46, Allee d'Italie  
F-69364 Lyon Cedex 07

**Prof. Dr. Wolfgang Woess**  
woess@weyl.math.tu-graz.ac.at  
Institut für Mathematik  
Technische Universität Graz  
Steyrergasse 30  
A-8010 Graz