

The sharp A_p constant for weights in a reverse-Hölder class

Martin Dindoš and Treven Wall

Abstract

Coifman and Fefferman established that the class of Muckenhoupt weights is equivalent to the class of weights satisfying the “reverse Hölder inequality”. In a recent paper V. Vasyunin [17] presented a proof of the reverse Hölder inequality with *sharp* constants for the weights satisfying the usual Muckenhoupt condition. In this paper we present the inverse, that is, we use the Bellman function technique to find the sharp A_p constants for weights in a reverse-Hölder class on an interval; we also find the sharp constants for the higher-integrability result of Gehring [7].

Additionally, we find sharp bounds for the A_p constants of reverse-Hölder-class weights defined on rectangles in \mathbb{R}^n , as well as bounds on the A_p constants for reverse-Hölder weights defined on cubes in \mathbb{R}^n , without claiming the sharpness.

1. Introduction

A weight w (a non-negative, measurable function) on an interval I is an $A_p(I)$ (or “Muckenhoupt”) weight ($1 < p < \infty$) if there is a constant $C < \infty$ such that the following inequality holds for every sub-interval $J \subset I$:

$$(1.1) \quad \langle w \rangle_J \langle w^{1-p'} \rangle_J^{p-1} \leq C.$$

Here $\langle w \rangle_I$ denotes $\frac{1}{|I|} \int_I w(t) dt$, the average of w over I , and p' is the conjugate exponent to p ($p' = \frac{p}{p-1}$).

2000 Mathematics Subject Classification: 42B20, 42B25.

Keywords: Reverse-Hölder class, Gehring class, A_p weight, Muckenhoupt weight, Bellman function.

The reverse-Hölder (or ‘‘Gehring’’) class¹ consists of all weights w for which there is a constant $C < \infty$ so that following inequality holds for every sub-interval $J \subset I$:

$$(1.2) \quad \frac{\langle w^p \rangle_J^{1/p}}{\langle w \rangle_J} \leq C.$$

If $w \in RH_p(I)$, then its RH_p constant, denoted by $RH_p(w)$, is defined to be the smallest constant C so that (1.2) holds for all $J \subset I$. We use $RH_p^\delta(I)$ to denote the class of all weights $w \in RH_p(I)$ such that $RH_p(w) \leq \delta$. Note that by Hölder’s inequality, the ratio (1.2) is never less than 1; hence we only consider $\delta \geq 1$.

We can also define $RH_\infty(I)$ by taking the limit as $p \rightarrow \infty$ of (1.2). Then, similarly, we say a weight w is in $RH_\infty^\delta(I)$ if for every sub-interval $J \subset I$,

$$(1.3) \quad \frac{\text{ess sup}_J w}{\langle w \rangle_J} \leq \delta.$$

It is worth noting that $RH_\infty(I)$ is strictly contained within $\bigcap_p RH_p(I)$. Among the RH_p classes, RH_∞ plays a role analogous to that of A_1 in the A_p classes. Several equivalent definitions of RH_∞ can be found in [4].

The class A_p was first described by Muckenhoupt [12], and its connection with the reverse-Hölder inequality was first explored by Coifman and C. Fefferman [3], who established that $\bigcup_p RH_p(I) = \bigcup_p A_p(I)$; this union is called $A_\infty(I)$. There is an alternative description of A_∞ weights as follows (see [8]); a weight w is in $A_\infty(I)$ if there is a constant C such that for all subintervals $J \subset I$, the following holds:

$$(1.4) \quad \langle w \rangle_J \exp(-\langle \log(w) \rangle_J) \leq C.$$

Our chief goal in this paper is to find the sharp constant \tilde{C} , depending only on p, q and δ such that any $w \in RH_p^\delta$ satisfies (1.1) (or (1.4)) with constant $C = \tilde{C}$. We will denote the class of weights satisfying (1.1) by $A_p^C(I)$ and those satisfying (1.4) by $A_\infty^C(I)$, respectively. This result is the reverse direction of Vasyunin’s work [17]. He found the sharp constant C such that any $w \in A_p^\delta$ belongs to the class RH_q^C . As a byproduct of our work on the above problem, we also are able to find the sharp constant C in the embedding of RH_p^δ into RH_t^C for $t > p$.

Our motivation to look at this problem arose when we attempted to establish a perturbation result for a certain class of nondivergence type elliptic operators. While studying this problem, we realized we needed to know to

¹There is no standard notation for this class. Some authors use the notation G_p , e.g., [1] and [11]; others denote it by B_p , e.g., [5] and [15]; our notation RH_p follows that of [4] and [14] and is, we feel, the most natural.

what A_q class a certain elliptic measure belongs, given that we know it satisfies the reverse Hölder inequality with a known constant.

We are very grateful to Sasha Volberg who provided valuable insight and brought our attention to his own results in the dyadic setting [13] as well as results of Vasyunin [17].

For our work on the perturbation problem for nondivergence elliptic operators we must establish these results not only on a real interval, but also on \mathbb{R}^n . This is the reason we have included the results in higher dimensions, despite the fact that in the cube case the constants are not sharp. The main point of Theorem 4 is the asymptotic as $\delta \rightarrow 1$. We prove that for fixed p and q , on a cube $Q \subset \mathbb{R}^n$,

$$RH_p^\delta(Q) \subset A_q^C(Q)$$

for some $C = C(\delta, p, q, n)$ and that $C \rightarrow 1$ as $\delta \rightarrow 1$.

2. Statement of Principal Results

To state our main results, we need to define the critical value of q . This value, $q^* = q^*(p, \delta)$, is the unique solution greater than one to

$$(2.1) \quad \frac{(q/\delta)^p - 1}{q - 1} = p.$$

It is fairly easy to see that:

- For every $1 < p < \infty$, $q^*(p, \delta) > \delta$.
- For a fixed $\delta \geq 1$, $\lim_{p \rightarrow \infty} q^*(p, \delta) = \delta$.
- For a fixed $1 < p < \infty$, $q^*(p, \delta) \sim \frac{(p\delta)^{p'}}{p}$ as $\delta \rightarrow \infty$.
- For a fixed $1 < p < \infty$, $q^*(p, \delta) \sim (\delta^{p'} - 1)^{1/p'} + 1$ as $\delta \rightarrow 1$.

Theorem 1. *For any weight $w \in RH_p^\delta(I)$, $1 < p < \infty$ we have that $w \in A_q^{C_q(p, \delta)}$, i.e.,*

$$(2.2) \quad \sup_{J \subset I} \langle w \rangle_J \langle w^{1-q'} \rangle_J^{q-1} \leq C_q(p, \delta)$$

holds, where

$$C_q(p, \delta) = \begin{cases} +\infty & 1 < q \leq q^*(p, \delta) \\ \frac{1}{q^*} \left(\frac{q-1}{q-q^*} \right)^{q-1} & q > q^*(p, \delta) \end{cases},$$

and $w \in A_\infty^{C_\infty(p, \delta)}$, i.e.,

$$(2.3) \quad \sup_{J \subset I} \langle w \rangle_J \exp(-\langle \log(w) \rangle_J) \leq C_\infty(p, \delta),$$

where

$$C_\infty(p, \delta) = \frac{1}{q^*} \exp(q^* - 1).$$

Here, as throughout the paper, $q^* = q^*(p, \delta)$ is the solution to (2.1), above. The constants $C_q(p, \delta)$ and $C_\infty(p, \delta)$ in this statement are the best possible.

Theorem 2. *If $w \in RH_\infty^\delta(I)$, then $w \in A_q^{C_q(\infty, \delta)}$, i.e.,*

$$(2.4) \quad \sup_{J \subset I} \langle w \rangle_J \langle w^{1-q'} \rangle_J^{q-1} \leq C_q(\infty, \delta),$$

where

$$C_q(\infty, \delta) = \begin{cases} +\infty & 1 < q \leq \delta \\ \frac{1}{\delta} \left(\frac{q-1}{q-\delta} \right)^{q-1} & q > \delta \end{cases},$$

and $w \in A_\infty^{C_\infty(\infty, \delta)}$, i.e.,

$$(2.5) \quad \sup_{J \subset I} \langle w \rangle_J \exp(-\langle \log(w) \rangle_J) \leq C_\infty(\infty, \delta),$$

where

$$C_\infty(\infty, \delta) = \frac{1}{\delta} \exp(\delta - 1).$$

Again, the constants $C_q(\infty, \delta)$ and $C_\infty(\infty, \delta)$ in this statement are the best possible.

Since, for a fixed $\delta \geq 1$, $\lim_{p \rightarrow \infty} q^*(p, \delta) = \delta$, Theorem 2 comes as no surprise considering Theorem 1. However, the proof of Theorem 1 must be adjusted to prove Theorem 2. We will primarily address the proof of Theorem 1, treating the proof of Theorem 2 as a special case where the need arises.

For the other endpoints $p = 1$ and $q = 1$, a few comments are in order. A moment's thought reveals that $RH_1(I)$ is not an interesting class to consider, as every positive L^1 function on I satisfies the condition. It is also evident from Theorem 1 that given any $\delta > 1$, and any $1 < p \leq \infty$ there is a weight $w \in RH_p^\delta(I)$ which is excluded from at least one $A_q(I)$ class; hence, since $A_1(I) \subset \bigcap_q A_q(I)$, there is no A_1 constant which can represent the entire class $RH_p^\delta(I)$.

Our method also allows us to find the sharp constants in Gehring's self-improvement result for the reverse-Hölder class [7]. We define the critical exponent $t^* = t^*(p, \delta)$ as the unique solution greater than p to $\left(\frac{\delta t^*}{t^* - 1}\right)^p \frac{t^* - p}{t^*} = 1$.

Theorem 3. For any weight $w \in RH_p^\delta(I)$, $1 < p < \infty$ we have that $w \in RH_t^{C_t(p,\delta)}$, i.e.,

$$(2.6) \quad \sup_{J \subset I} \frac{\langle w^t \rangle_J^{1/t}}{\langle w \rangle_J} \leq C_t(p, \delta)$$

holds, where

$$C_t(p, \delta) = \begin{cases} \frac{t^*-1}{t^*} \left(\frac{t^*}{t^*-t}\right)^{1/t} & p \leq t < t^*(p, \delta) \\ +\infty & t \geq t^*(p, \delta) \end{cases},$$

and the constant $C_t(p, \delta)$ is sharp.

In considering the n -dimensional analog of our results, we are no longer able to find the sharp constants. However, we find useful asymptotic information as $\delta \rightarrow 1$,

Theorem 4. Let $n > 1$ and let $Q \subset \mathbb{R}^n$ be a cube with sides parallel to the coordinate axes. Fix $p > 1$, $q > 1$ and $\eta > 1$. Then there is a $\delta > 1$ such that any weight $w \in RH_p^\delta(Q)$ is in $A_q^\eta(Q)$, that is,

$$(2.7) \quad \langle w \rangle_K \langle w^{1-q'} \rangle_K^{q-1} \leq \eta,$$

for every cube $K \subset Q$ with sides parallel to the coordinate axes.

The standard definition of RH_p in n -dimensions is based on cubes (or balls). However, if one strengthens the definition of $RH_p(I)$ to require the inequality

$$\frac{\langle w^p \rangle_J^{1/p}}{\langle w \rangle_J} \leq C$$

to hold for all bounded, open rectangles $J \subset I$, a new, smaller class of weights is formed (these classes are considered in, e.g., [2], [9]). We will call this class *strong* $RH_p(I)$ and denote it by $s-RH_p(I)$. Similarly, one can define *strong* A_q (denoted $s-A_q$). With these definitions in mind,

Theorem 5. Let $1 < p < \infty$, let $I \subset \mathbb{R}^n$ be a bounded, open rectangle, and assume $w \in s-RH_p^\delta(I)$. Then $w \in s-A_q^{C_q(p,\delta)}(I)$ and $w \in s-A_\infty^{C_\infty(p,\delta)}(I)$, where $C_q(p, \delta)$ and $C_\infty(p, \delta)$ are the constants in Theorem 1, independent of n . Also, $s-RH_\infty^\delta(I) \subset s-A_q^{C_q(\infty,\delta)}(I)$ and $s-RH_\infty^\delta(I) \subset s-A_\infty^{C_\infty(\infty,\delta)}(I)$, where $C_q(\infty, \delta)$ and $C_\infty(\infty, \delta)$ are the constants in Theorem 2. Finally, $s-RH_p^\delta(I) \subset s-RH_t^{C_t(p,\delta)}(I)$, where $C_t(p, \delta)$ is the constant in Theorem 3. In all cases, these constants are sharp.

The literature on A_p and RH_p weights is far too extensive to comprehensively cover here, but a small review is in order. The papers [12] and [3], mentioned earlier, contain foundational results on these weights. Both [6] and [16] are good references; they emphasize the connection to singular

integral operators. There are several factorization results relating RH_p and A_q (see, e.g., [4]), and C. J. Neugebauer, in [14], uses these to prove that if $w \in RH_\infty^\delta$, then $w \in A_q$ for all $q > \delta$. Additionally, he provides conditions for weights in RH_p^δ to be in A_q . However, the results there depend on specific factorizations of the weights and the A_q constants aren't provided. In [11], the one-dimensional embedding of RH_p into A_q with the best range of q is proven using rearrangements, but, again, this method doesn't find the A_q constants. In [1], the RH_∞ embedding result, Theorem 2, is found using rearrangements; we include it here because it follows with little extra work from the proof of our RH_p embedding result (Theorem 1). This same group of authors finds, in [2], the embedding of *strong* RH_∞ into *strong* A_p in n dimensions with the same constant. In improving upon Gehring's original result, [7], Korenovskii [10] found the sharp upper bound on t in the embedding $RH_p^\delta \subset RH_t$ in one dimension and Kinnunen [9] found the same upper bound for the *strong* RH_p classes in n dimensions; however, neither of these methods provide the RH_t constant of the embedded weight. Using the Bellman function technique, we are able to provide the sharp constants in one dimension and new results in n dimensions, including sharp constants for all *strong* RH_p embeddings (the technique is explained in [13], especially in the context of classical analysis problems). However, finding the sharp constants for the usual RH_p classes in n dimensions remains an open problem.

The paper will proceed as follows: first, we describe the setup for the Bellman function technique in section 3. We use $\mathbb{B}(x)$ to denote the Bellman function. Then, in section 4 we prove Theorems 1, 2 and 3 from an auxiliary theorem, Theorem 6. In section 5 we explain the heuristics behind our "guess" at the explicit formula for the Bellman function; we call this guess B . We then show that our guess is correct, proving Theorem 6, by verifying that $B(x) \leq \mathbb{B}(x)$ (Lemma 2) and $B(x) \geq \mathbb{B}(x)$ on Ω_δ (Lemma 4). Proving the former inequality requires finding a weight representing each x in Ω_δ (section 7), and proving the latter requires working with domains Ω_ϵ for $\epsilon > \delta$ (see section 9). Finally, we prove Theorems 4 and 5 in section 10. Throughout the paper, we alternate between heuristic calculations and rigorous proof to exhibit the philosophy of the Bellman function technique.

3. Bellman function ideas

Typically, when one uses the Bellman function technique, all one needs is to find an upper bound for the Bellman function which preserves concavity (or convexity, as needed). However, this approach doesn't allow for the calculation of sharp constants. Consequently, we find the actual formula for the Bellman function.

For all weights w , for $1 < p < \infty$, and for any interval J , $\langle w \rangle_J^p \leq \langle w^p \rangle_J$, by Hölder’s inequality. Hence, if $w \in RH_p^\delta(I)$, the point $x = (x_1, x_2) = (\langle w \rangle_I, \langle w^p \rangle_I)$ lies in the domain

$$\Omega_\delta(p) := \{(x_1, x_2) : x_1 > 0, x_1^p \leq x_2 \leq (\delta x_1)^p\}.$$

For our problem, the Bellman function for $1 < p < \infty$, $0 < q < \infty$ is

$$(3.1) \quad \mathbb{B}(x; p, q, \delta) := \sup\{\langle w^{1-q'} \rangle_I : x_1 = \langle w \rangle_I, x_2 = \langle w^p \rangle_I, \text{ and } w \in RH_p^\delta(I)\},$$

and for $q = \infty$,

$$(3.2) \quad \mathbb{B}(x; p, \infty, \delta) := \sup\{\exp(-\langle \log(w) \rangle_I) : x_1 = \langle w \rangle_I, x_2 = \langle w^p \rangle_I, w \in RH_p^\delta(I)\}.$$

Note that \mathbb{B} doesn’t depend on the interval I on which it is defined, since, given two intervals I_1 and I_2 , the affine mapping of one onto the other preserves the averages and puts $RH_p^\delta(I_1)$ in one-to-one correspondence with $RH_p^\delta(I_2)$. We are allowing for $0 < q < 1$ in order to prove Theorem 3; for q in this range, the exponent $1 - q'$ is greater than 1.

For $p = \infty$, we must adjust these coordinates. We set $x = (x_1, x_2) = (\langle w \rangle_I, \text{ess sup}_I w)$, whence

$$\begin{aligned} \Omega_\delta(\infty) &:= \{(x_1, x_2) : x_1 > 0, x_1 \leq x_2 \leq \delta x_1\}, \\ \mathbb{B}(x; \infty, q, \delta) &:= \\ &\sup \left\{ \langle w^{1-q'} \rangle_I : x_1 = \langle w \rangle_I, x_2 = \text{ess sup}_I w, w \in RH_\infty^\delta(I) \right\}, \\ \mathbb{B}(x; \infty, \infty, \delta) &:= \\ &\sup \left\{ \exp(-\langle \log(w) \rangle_I) : x_1 = \langle w \rangle_I, x_2 = \text{ess sup}_I w, w \in RH_\infty^\delta(I) \right\}. \end{aligned}$$

We consider \mathbb{B} as a function on Ω_δ , since each point $x \in \Omega_\delta$ can be represented by a weight $w \in RH_p^\delta$. We will demonstrate the existence of such weights in Lemma 2.

We will often split an interval J into the union of two disjoint subintervals which we will call J^- and J^+ , with $|J^\pm| = \alpha^\pm |J|$. Given a weight w defined on J , we split it into two weights w^\pm defined on their respective subintervals. As above, we relate these weights to points in \mathbb{R}^2 , letting the point x^0 correspond to the original weight w and the points x^\pm correspond to w^\pm . These points are co-linear: $x^0 = \alpha^- x^- + \alpha^+ x^+$. Also, if we start with $w \in RH_p^\delta(J)$, w^\pm are in $RH_p^\delta(J^\pm)$; consequently, x^0, x^- and x^+ are all points in Ω_δ .

We need some further notation. For $1 < p < \infty$, denote by u_p^\pm the functions inverse to

$$t \rightarrow \frac{(1 - pt)^{p-1}}{(1 - (p - 1)t)^p}$$

on the following domains: $u_p^+ : [0, 1] \rightarrow [0, \frac{1}{p}]$, $u_p^- : [0, 1] \rightarrow (-\infty, 0]$, i.e., the values $u_p^\pm(t)$ are the positive and negative solutions to the equation $\frac{(1-pu)^{p-1}}{(1-(p-1)u)^p} = t$ for $0 \leq t \leq 1$. Based on this, we define $s^\pm = s_p^\pm(\delta) := u_p^\pm(1/\delta^p)$ and $r^\pm = r_p^\pm(x, \delta) := u_p^\pm(x_2/(\delta x_1)^p)$. Finally, we set $\gamma := p + q' - 1$.

Theorems 1, 2 and 3 are consequences of the following

Theorem 6. *For $0 < q < \infty$ and $1 < p < \infty$, if $x_2 = x_1^p$ (or if $p = \infty$ and $x_2 = x_1$), we have*

$$\mathbb{B}(x; p, q, \delta) = x_1^{1-q'}$$

If $1 < p < \infty$ and $x_2 > x_1^p$, then

$$(3.3) \quad \mathbb{B}(x; p, q, \delta) = \begin{cases} x_1^{1-q'} \left(\frac{1-ps^+}{1-pr^+}\right)^{q'} \left(\frac{1-(p-1)r^+}{1-(p-1)s^+}\right)^{q'-1} \left(\frac{1-\gamma r^+}{1-\gamma s^+}\right) & \text{for } q > \frac{1-(p-1)s^+}{1-ps^+} \\ \infty & \text{for } \frac{1-(p-1)s^-}{1-ps^-} \leq q \leq \frac{1-(p-1)s^+}{1-ps^+} \\ x_1^{1-q'} \left(\frac{1-ps^-}{1-pr^-}\right)^{q'} \left(\frac{1-(p-1)r^-}{1-(p-1)s^-}\right)^{q'-1} \left(\frac{1-\gamma r^-}{1-\gamma s^-}\right) & \text{for } \frac{p-1}{p} < q < \frac{1-(p-1)s^-}{1-ps^-}. \end{cases}$$

If $p = \infty$ and $x_2 > x_1$, then

$$(3.4) \quad \mathbb{B}(x; \infty, q, \delta) = \begin{cases} x_2^{1-q'} \left(\frac{q-\frac{x_1}{x_2}\delta}{q-\delta}\right) & \text{for } q > \delta \\ \infty & \text{for } 1 < q \leq \delta. \end{cases}$$

4. Proof of Theorems 1, 2 and 3

One can easily check that the value $q^*(p, \delta)$ defined by (2.1) and used in Theorem 1 is the same as $\frac{1-(p-1)s^+}{1-ps^+}$ used in Theorem 6. Similarly, if we define $q_* = q_*(p, \delta)$ to be the unique solution to (2.1) between 0 and 1, $q_* = \frac{1-(p-1)s^-}{1-ps^-}$, the other bound in Theorem 6. The critical exponent in Theorem 3, t^* , satisfies $t^* = 1 - q'_* = \frac{1}{1-q_*}$.

Now, we assume Theorem 6. We consider $q > 1$ for Theorems 1 and 2 and $\frac{p-1}{p} < q < 1$ for Theorem 3 (in this range, $1 - q' > p$, which is all that we are interested in). Recall that $\mathbb{B}(x)$ represents the maximum of $\langle w^{1-q'} \rangle$ for all weights in RH_p^δ which are represented by x and that x_1 represents $\langle w \rangle$. For $q > 1$, the constant we desire is

$$\sup_{x \in \Omega_\delta} x_1 (\mathbb{B}(x; p, q, \delta))^{q-1},$$

and for $\frac{p-1}{p} < q < 1$, we seek

$$\sup_{x \in \Omega_\delta} \frac{1}{x_1} \mathbb{B}(x; p, q, \delta)^{1-q}.$$

With that in mind, we define $g := x_1^{q'-1} \mathbb{B}(x)$, that is,

$$g = \left(\frac{1 - ps^\pm}{1 - pr^\pm} \right)^{q'} \left(\frac{1 - (p-1)r^\pm}{1 - (p-1)s^\pm} \right)^{q'-1} \left(\frac{1 - \gamma r^\pm}{1 - \gamma s^\pm} \right).$$

$g \geq 0$ and

$$\frac{d}{dr^\pm} \log(g) = \frac{-q'(q'-1)r^\pm}{(1 - pr^\pm)(1 - (p-1)r^\pm)(1 - \gamma r^\pm)}.$$

For $q > 1$, we use r^+ and s^+ , so g' is negative. For $\frac{p-1}{p} < q < 1$, we use r^- and s^- , so g' is positive. Hence, the maximum of g in both cases is at $r^\pm = 0$. So, our best constant is

$$(g(0))^{q-1} = \frac{(1 - ps^+)^q}{(1 - (p-1)s^+)(1 - \gamma s^+)^{q-1}}, \text{ for } q > 1, \text{ and}$$

$$(g(0))^{1-q} = \frac{1 - (p-1)s^-}{(1 - ps^-)^q(1 - \gamma s^-)^{1-q}} \text{ for } \frac{p-1}{p} < q < 1.$$

Relating this first constant back to q^* , we find

$$\begin{aligned} \frac{(1 - ps^+)^q}{(1 - (p-1)s^+)(1 - \gamma s^+)^{q-1}} &= \frac{1}{q^*} \left(1 + \frac{s^+(1 - q')}{1 - ps^+} \right)^{1-q} \\ &= \frac{1}{q^*} \left(1 - \frac{q^* - 1}{q - 1} \right)^{1-q} = \frac{1}{q^*} \left(\frac{q - 1}{q - q^*} \right)^{q-1}, \end{aligned}$$

which is the constant in Theorem 1. For the second constant, $\frac{p-1}{p} < q < 1$ and we use $t = 1 - q'$ and $t^* = 1 - q'_*$ to see

$$\begin{aligned} \frac{(1 - (p-1)s^-)}{(1 - ps^-)^q(1 - \gamma s^-)^{1-q}} &= q_* \left(\frac{q - q_*}{q - 1} \right)^{q-1} \\ &= \frac{t^* - 1}{t^*} \left((1 - t) + t \left(\frac{t^* - 1}{t^*} \right) \right)^{-1/t} = \frac{t^* - 1}{t^*} \left(\frac{t^*}{t^* - t} \right)^{1/t}, \end{aligned}$$

which is the constant in Theorem 3.

To complete the proof for $p < \infty$, fix a point $x \in \Omega_\delta$ (fixing $r^\pm = r^\pm(x)$). Then, the weight

$$(4.1) \quad w_{c,a,\nu}(t) = \begin{cases} c \left(\frac{t}{a}\right)^\nu & \text{if } 0 \leq t \leq a \\ c & \text{if } a \leq t \leq 1, \end{cases}$$

with constants $\nu = \frac{s^\pm}{1-ps^\pm}$, $a = \frac{s^\pm-r^\pm}{s^\pm(1-pr^\pm)}$, $c = x_1 \frac{(1-pr^\pm)(1-(p-1)s^\pm)}{(1-(p-1)r^\pm)(1-ps^\pm)}$, is in $RH_p^\delta(I)$ and its A_q norm is infinite for any $q_* \leq q \leq q^*$. This is exhibited in the proof of Lemma 2.

For the case of $p = \infty$, the analysis is even easier. Given the definition of \mathbb{B} in (3.4), we see that for $q > \delta$,

$$x_1(\mathbb{B}(x; \infty, q, \delta))^{q-1} = \frac{x_1}{x_2} \left(\frac{q - \frac{x_1}{x_2}\delta}{q - \delta} \right)^{q-1}.$$

Letting $y = \frac{x_1}{x_2}$, we know that $\frac{1}{\delta} \leq y \leq 1$, and we see

$$\frac{d}{dy} \left[y \left(\frac{q - y\delta}{q - \delta} \right)^{q-1} \right] = q(1 - \delta y) \frac{(q - y\delta)^{q-2}}{(q - \delta)^{q-1}},$$

which is negative for $y > 1/\delta$. Thus, the maximum is at $y = 1/\delta$, which is exactly the constant in Theorem 2. For $q \leq \delta$, we again fix an $x \in \Omega_\delta$. Then, the weight in (4.1), with constants $\nu = \delta - 1$, $a = \frac{1-x_1/x_2}{1-1/\delta}$ and $c = x_2$, is in $RH_\infty^\delta(I)$ with infinite A_q norm for $1 \leq q \leq \delta$, which completes the proof. As before, this is contained in the proof of Lemma 2.

Finally, we address the case of $q = \infty$. Define, for $1 < p < \infty$,

$$(4.2) \quad \begin{aligned} B(x; p, \infty, \delta) &:= \lim_{q \rightarrow \infty} (\mathbb{B}(x; p, q, \delta))^{q-1} \\ &= \frac{1}{x_1} \frac{(1 - (p-1)r^+)(1 - ps^+)}{(1 - pr^+)(1 - (p-1)s^+)} \exp\left[\frac{s^+ - r^+}{(1 - ps^+)(1 - pr^+)}\right], \end{aligned}$$

and, for $p = \infty$,

$$(4.3) \quad B(x; \infty, \infty, \delta) := \lim_{q \rightarrow \infty} (\mathbb{B}(x; \infty, q, \delta))^{q-1} = \frac{1}{x_2} \exp[\delta(1 - \frac{x_1}{x_2})].$$

We want to establish that these functions satisfy

$$B(x; p, \infty, \delta) = \mathbb{B}(x; p, \infty, \delta) \quad \text{and} \quad B(x; \infty, \infty, \delta) = \mathbb{B}(x; \infty, \infty, \delta).$$

First, it is not difficult to check that the weights w_p and w_∞ defined by (4.1) with the respective constants for $p < \infty$, $p = \infty$ do not depend on q and satisfy

$$\exp(-\langle \log(w_p) \rangle) = B(x; p, \infty, \delta) \quad \text{and} \quad \exp(-\langle \log(w_\infty) \rangle) = B(x; \infty, \infty, \delta),$$

respectively. By the definition of \mathbb{B} , this gives the inequalities $B(x; p, \infty, \delta) \leq \mathbb{B}(x; p, \infty, \delta)$ and $B(x; \infty, \infty, \delta) \leq \mathbb{B}(x; \infty, \infty, \delta)$. The other inequality is a result of applying Jensen's inequality; namely,

$$\exp(-\langle \log(w) \rangle) \leq \langle w^{1-q'} \rangle^{q-1} \leq \mathbb{B}(x; p, q, \delta)^{q-1}.$$

So in (4.2) and (4.3), taking the limits establishes the desired equality of B and \mathbb{B} .

Given this, the A_∞ constants in Theorems 1 and 2 are easy to find. We simply calculate

$$\begin{aligned} \sup_{x \in \Omega_\delta} x_1 \mathbb{B}(x; p, \infty, \delta) &= \\ &= \sup_{x \in \Omega_\delta} \frac{(1 - (p - 1)r^+)(1 - ps^+)}{(1 - pr^+)(1 - (p - 1)s^+)} \exp \left[\frac{s^+ - r^+}{(1 - ps^+)(1 - pr^+)} \right]. \end{aligned}$$

Again, the maximum is at $r^+ = 0$, whence the constant is

$$\frac{(1 - ps^+)}{(1 - (p - 1)s^+)} \exp \left[\frac{s^+}{(1 - ps^+)} \right] = \frac{1}{q^*} \exp(q^* - 1).$$

And,

$$\sup_{x \in \Omega_\delta} x_1 \mathbb{B}(x; \infty, \infty, \delta) = \sup_{x \in \Omega_\delta} \frac{x_1}{x_2} \exp \left[\delta \left(1 - \frac{x_1}{x_2} \right) \right] = \frac{1}{\delta} \exp(\delta - 1).$$

This completes the proof of Theorems 1, 2 and 3 from Theorem 6.

5. Deriving the formula for \mathbb{B}

We start by examining the scaling properties of \mathbb{B} . Given $w \in RH_p^\delta$, and $\lambda > 0$ a constant, then $\tilde{w} := \lambda w$ is in RH_p^δ as well, and $\langle \tilde{w} \rangle_I = \lambda \langle w \rangle_I$, $\langle \tilde{w}^p \rangle_I = \lambda^p \langle w^p \rangle_I$. Consequently, $\mathbb{B}(\lambda x_1, \lambda^p x_2) = \lambda^{1-q'} \mathbb{B}(x_1, x_2)$. Letting $x_1 = \frac{1}{\lambda}$, we see $\mathbb{B}(x_1, x_2) = x_1^{1-q'} \mathbb{B}(1, \frac{x_2}{x_1^p})$. Thus, we define $g(y) := \mathbb{B}(1, y)$, and we see that $\mathbb{B}(x_1, x_2) = x_1^{1-q'} g(\frac{x_2}{x_1^p})$.

For $0 < q < \infty$, $\langle w^{1-q'} \rangle_I \geq \langle w \rangle_I^{1-q'}$, using Hölder's inequality, whence, $\mathbb{B}(x_1, x_2) \geq x_1^{1-q'}$. Therefore, $g(y) \geq 1$. Further, if $x_2 = x_1^p$, that is, if $\langle w^p \rangle_I = \langle w \rangle_I^p$, the weight w must be a constant. In that case, $\mathbb{B}(x) = x_1^{1-q'}$, and we see that $g(1) = 1$.

We expect \mathbb{B} to be a concave function, as the following illustrates. Given an interval J , split it into the disjoint union of subintervals J^- and J^+ . Assuming they exist, let w^\pm be two extremal weights (i.e., which satisfy $\langle (w^\pm)^{1-q'} \rangle_{J^\pm} = \mathbb{B}(x^\pm)$). Then, concatenate these two weights to form a new weight w on J . Thus, $\langle w^{1-q'} \rangle_J = \alpha^- \langle (w^-)^{1-q'} \rangle_{J^-} + \alpha^+ \langle (w^+)^{1-q'} \rangle_{J^+}$. The weight w corresponds to the point x^0 , and w^\pm to x^\pm . Then, $x^0 = \alpha^- x^- + \alpha^+ x^+$, and we have $\mathbb{B}(x^0) \geq \alpha^- \mathbb{B}(x^-) + \alpha^+ \mathbb{B}(x^+)$, which is the concavity condition (alternatively, using the terminology of [13], we expect a concave solution since the profit function is zero and there is no drift term). We ignore the substantive issues of whether extremal weights exist, and whether $x^0 \in \Omega_\delta$ due to the heuristic nature of our procedure. However,

later proofs lay these concerns to rest. We will also assume that the Hessian of \mathbb{B} is singular. This last assumption gives rise to an ODE that we can solve, which enables us to find \mathbb{B} explicitly. This assumption is frequently made in the application of the Bellman function technique; here it is reasonable because we expect extremal weights to exist.

To proceed further, we must calculate the Hessian of \mathbb{B} in terms of g (assuming, of course, that \mathbb{B} is sufficiently differentiable). Let $y = \frac{x_2}{x_1}$. Then,

$$\begin{aligned} \frac{\partial^2 \mathbb{B}}{\partial x_1^2} &= x_1^{-q'-1} [q'(q' - 1)g + (2q' + p - 1)pyg' + p^2y^2g''], \\ \frac{\partial^2 \mathbb{B}}{\partial x_1 \partial x_2} &= x_1^{-q'-p} [(1 - q' - p)g' - pyg''], \quad \text{and} \quad \frac{\partial^2 \mathbb{B}}{\partial x_2^2} = x_1^{1-q'} x_2^{-2} y^2 g''. \end{aligned}$$

Thus,

$$(5.1) \quad \text{Hess}(\mathbb{B}) = \begin{pmatrix} \frac{\partial^2 \mathbb{B}}{\partial x_1^2} & \frac{\partial^2 \mathbb{B}}{\partial x_1 \partial x_2} \\ \frac{\partial^2 \mathbb{B}}{\partial x_2 \partial x_1} & \frac{\partial^2 \mathbb{B}}{\partial x_2^2} \end{pmatrix} = x_1^{-q'-1} \begin{pmatrix} 1 & 1 \\ 0 & -\frac{x_1}{px_2} \end{pmatrix} R \begin{pmatrix} 1 & 0 \\ 1 & -\frac{x_1}{px_2} \end{pmatrix}$$

with

$$(5.2) \quad R = \begin{pmatrix} q'(q' - 1)g - p(p - 1)yg' & p\gamma yg' \\ p\gamma yg' & p^2y^2g'' \end{pmatrix},$$

where $\gamma = p + q' - 1$.

To force R to be singular, we require

$$(5.3) \quad (p^2y^2g'')(q'(q' - 1)g - p(p - 1)yg') = (p\gamma yg')^2.$$

Now, we make the substitution $h = \frac{yg'}{g}$. Then $g'' = \frac{gh' + hg' - g'}{y}$. Convert and divide both sides by $(pg)^2$ to get

$$(yh' - h + h^2)(q'(q' - 1) - p(p - 1)h) = \gamma^2 h^2,$$

which is separable. So,

$$y = C \left(\frac{h((p - 1)h + q')^{p-1}}{(ph + q' - 1)^p} \right).$$

We make one further change of variables,

$$h = \frac{q'(q' - 1)}{p(p - 1)(1 - \gamma r)} \quad \text{or} \quad r = \frac{1}{\gamma} - \frac{q'(q' - 1)}{\gamma p(p - 1)h},$$

which yields

$$(5.4) \quad y = C \frac{(1 - pr)^{p-1}}{(1 - (p - 1)r)^p}.$$

It turns out to be natural to choose $C = \delta^p$ in (5.4), and it is at this point that we see the origin of the function u_p , mentioned above. Recall that we have set u_p^\pm as the positive and negative inverses of the function

$$t \rightarrow \frac{(1-pt)^{p-1}}{(1-(p-1)t)^p}.$$

Also recall that $s^\pm = u_p^\pm(\frac{1}{\delta^p})$. We can see from (5.4), with $C = \delta^p$, that $r^\pm = u_p^\pm(\frac{x_2}{(\delta x_1)^p})$. Note also that if $r^\pm = s^\pm$, then $y = 1$.

We want to relate this all back to g , so we calculate

$$\frac{d}{dr} \log(y) = \frac{-p(p-1)}{1-pr} + \frac{p(p-1)}{1-(p-1)r}$$

and use this to see that

$$\begin{aligned} d(\log(g)) &= hd(\log(y)) = \frac{q'(q'-1)}{p(p-1)(1-\gamma r)} \left(\frac{-p(p-1)}{1-pr} + \frac{p(p-1)}{1-(p-1)r} \right) \\ &= \frac{pq'}{1-pr} - \frac{\gamma}{1-\gamma r} - \frac{(p-1)(q'-1)}{1-(p-1)r}. \end{aligned}$$

Since $g|_{y=1} = g|_{r=s} = 1$, we have

$$\begin{aligned} \log g &= \int_s^r \left(\frac{pq'}{1-pt} - \frac{\gamma}{1-\gamma t} - \frac{(p-1)(q'-1)}{1-(p-1)t} \right) dt \\ &= -q' \log\left(\frac{1-pr}{1-ps}\right) + (q'-1) \log\left(\frac{1-(p-1)r}{1-(p-1)s}\right) + \log\left(\frac{1-\gamma r}{1-\gamma s}\right), \end{aligned}$$

whence

$$(5.5) \quad g = \left(\frac{1-ps}{1-pr}\right)^{q'} \left(\frac{1-(p-1)r}{1-(p-1)s}\right)^{q'-1} \left(\frac{1-\gamma r}{1-\gamma s}\right).$$

The last thing is to discover whether we should use r^+ or r^- in the definition of g to ensure that R is negative semi-definite. Since R is singular and symmetric, it suffices to make the upper left-hand entry of R negative. That is, we must make sure that

$$p(p-1)yg' \geq q'(q'-1)g \quad \text{or} \quad h \geq \frac{q'(q'-1)}{p(p-1)}.$$

Note that

$$\frac{dh}{dr} = \frac{q'(q'-1)}{p(p-1)} \gamma (1-\gamma r)^2,$$

which is positive for $q > 1$ and negative for $\frac{p-1}{p} < q < 1$. Also, $h(0) = \frac{q'(q'-1)}{p(p-1)}$, so we need $h(r) \geq h(0)$ for $q > 1$, which is accomplished by choosing the positive solution r^+ . Accordingly, for $\frac{p-1}{p} < q < 1$, we use r^- .

Therefore, we have a candidate for \mathbb{B} : $B(x) = x_1^{1-q'} g(\frac{x_2}{x_1})$. It is occasionally helpful to have this expressed in two different ways, which we record here

$$(5.6) \quad B(x) = x_1^{1-q'} \left(\frac{1 - ps^\pm}{1 - pr^\pm} \right)^{q'} \left(\frac{1 - (p-1)r^\pm}{1 - (p-1)s^\pm} \right)^{q'-1} \left(\frac{1 - \gamma r^\pm}{1 - \gamma s^\pm} \right)$$

$$(5.7) \quad B(x) = x_1^{-\gamma} x_2 \left(\frac{1 - ps^\pm}{1 - pr^\pm} \right)^\gamma \left(\frac{1 - (p-1)r^\pm}{1 - (p-1)s^\pm} \right)^\gamma \left(\frac{1 - \gamma r^\pm}{1 - \gamma s^\pm} \right).$$

The second representation is obtained by using the definitions of r^\pm and s^\pm to see that

$$\frac{x_2}{x_1^p} = \left(\frac{1 - pr^\pm}{1 - ps^\pm} \right)^{p-1} \left(\frac{1 - (p-1)s^\pm}{1 - (p-1)r^\pm} \right)^p.$$

We note that since $s^- \leq r^- \leq 0 \leq r^+ \leq s^+ < 1/p$, the only concern with the denominator of g occurs when $(1 - \gamma s^\pm) = 0$. At this point, we have $q' = 1/s^\pm + 1 - p$, or $q = \frac{1-(p-1)s^\pm}{1-ps^\pm}$, which is q^* (or q_*), the critical values of q in Theorems 1 and 6.

For the case of $p = \infty$, we have two options. Either we can take limits, using the asymptotics as $p \rightarrow \infty$ of

$$(5.8) \quad s^+ \sim \frac{1}{p} - \frac{1}{(\delta - 1)p^2} \quad \text{and} \quad r^+ \sim \frac{1}{p} - \frac{1}{(\delta \tilde{x} - 1)p^2},$$

where $\tilde{x} := \frac{x_1}{x_2^{1/p}} \rightarrow \frac{x_1}{x_2}$. Or, we can carry out a similar analysis. We leave the former approach to the reader and illustrate the latter approach, as the ideas involved are useful for later. We start by recalling that for a weight w and an interval J , $x_2 = \text{ess sup}_J w$. This change of coordinates alters the effect of splitting; if we split an interval J into two parts, $J = J^- \cup J^+$, the point $x^0 = (\langle w \rangle_J, \text{ess sup}_J w)$ is no longer necessarily co-linear with the points $x^\pm = (\langle w \rangle_{J^\pm}, \text{ess sup}_{J^\pm} w)$. The first coordinate splits proportionally, $x_1^0 = \alpha^- x_1^- + \alpha^+ x_1^+$, but the second coordinate now satisfies $x_2^0 = \max\{x_2^-, x_2^+\}$. Nevertheless, we are still seeking a concavity condition as before, that is, we want to ensure that $\mathbb{B}(x^0) \geq \alpha^- \mathbb{B}(x^-) + \alpha^+ \mathbb{B}(x^+)$. However, this is not typical concavity, due to the behavior of the coordinates. To get our hands on an expression for the concavity, we look at the Taylor series for \mathbb{B} based at x^0 up to the second terms (assuming \mathbb{B} is sufficiently differentiable):

$$(5.9) \quad \mathbb{B}(x^\pm) \simeq \mathbb{B}(x^0) + \sum_{i=1}^2 \frac{\partial \mathbb{B}}{\partial x_i}(x^0)(x_i^\pm - x_i^0) + \frac{1}{2} \sum_{i,j=1}^2 \frac{\partial^2 \mathbb{B}}{\partial x_i \partial x_j}(x^0)(x_i^\pm - x_i^0)(x_j^\pm - x_j^0).$$

We know one of x_2^\pm is equal to x_2^0 , so we assume that $x_2^+ = x_2^0$; due to the symmetry of the situation, we lose no generality in this assumption. Recall that $x_1^0 = \alpha^- x_1^- + \alpha^+ x_1^+$, and define $\Delta_1 := x_1^+ - x_1^-$ and $\Delta_2 := x_2^0 - x_2^- \geq 0$. Then, for concavity, we want the following linear combination of these terms to be non-positive for small Δ_1 and Δ_2 :

$$(5.10) \quad \alpha^- \mathbb{B}(x^-) + \alpha^+ \mathbb{B}(x^+) - \mathbb{B}(x^0) \simeq \alpha^- \left(-\frac{\partial \mathbb{B}}{\partial x_2} \Delta_2 + \frac{1}{2} \alpha^+ \frac{\partial^2 \mathbb{B}}{\partial x_1^2} \Delta_1^2 + \alpha^+ \frac{\partial^2 \mathbb{B}}{\partial x_1 \partial x_2} \Delta_1 \Delta_2 + \frac{1}{2} \frac{\partial^2 \mathbb{B}}{\partial x_2^2} \Delta_2^2 \right).$$

Assume that $\text{Hess}(\mathbb{B})$ is singular, that $\frac{\partial \mathbb{B}}{\partial x_2} \geq 0$ and that $\frac{\partial^2 \mathbb{B}}{\partial x_1^2} \leq 0$; we will then demonstrate that (5.10) is non-positive. From the assumptions that $\text{Hess}(\mathbb{B})$ is singular and $\frac{\partial^2 \mathbb{B}}{\partial x_1^2} \leq 0$, we know that $\frac{\partial^2 \mathbb{B}}{\partial x_2^2} \leq 0$ and that the quadratic form of $\text{Hess}(\mathbb{B})$ is negative semi-definite, hence

$$\frac{\partial^2 \mathbb{B}}{\partial x_1 \partial x_2} \Delta_1 \Delta_2 \leq -\frac{1}{2} \left(\frac{\partial^2 \mathbb{B}}{\partial x_1^2} \Delta_1^2 + \frac{\partial^2 \mathbb{B}}{\partial x_2^2} \Delta_2^2 \right).$$

Therefore, the right-hand side of (5.10) is less than or equal to

$$\alpha^- \left(-\frac{\partial \mathbb{B}}{\partial x_2} \Delta_2 + \frac{1}{2} (1 - \alpha^+) \frac{\partial^2 \mathbb{B}}{\partial x_2^2} \Delta_2^2 \right),$$

which is non-positive. To get the differential equation which defines B , we supplement these conditions with yet another singularity assumption and arrive at two possibilities

$$(5.11) \quad \frac{\partial^2 \mathbb{B}}{\partial x_1^2} = 0 \quad \text{and} \quad \frac{\partial \mathbb{B}}{\partial x_2} \geq 0, \quad \text{or}$$

$$(5.12) \quad \frac{\partial \mathbb{B}}{\partial x_2} = 0 \quad \text{and} \quad \frac{\partial^2 \mathbb{B}}{\partial x_1^2} \leq 0.$$

We then turn to the scaling of \mathbb{B} . In this case, we see $\mathbb{B}(\lambda x_1, \lambda x_2) = \lambda^{1-q'} \mathbb{B}(x_1, x_2)$, so setting $\lambda = 1/x_1$ yields $\mathbb{B}(x_1, x_2) = x_1^{1-q'} \mathbb{B}(1, \frac{x_2}{x_1})$. We then define $g(y) = \mathbb{B}(1, y)$, so that $\mathbb{B}(x_1, x_2) = x_1^{1-q'} g(\frac{x_2}{x_1})$. Also, as before, we know that $\mathbb{B}(x) \geq x_1^{1-q'}$, whence $g(y) \geq 1$. Further, $\frac{x_2}{x_1} = 1$ only occurs when the weight is constant, in which case we have $\mathbb{B}(x_1, x_1; q, \delta) = x_1^{1-q'}$; that is, $g(1) = 1$.

If \mathbb{B} solves (5.11), then \mathbb{B} is linear in x_1 , which yields

$$\mathbb{B}(x_1, x_2) = a(x_2) + b(x_2)x_1 = x_1^{1-q'} g(x_2/x_1).$$

Since $g(1) = 1$, we set $x_1 = x_2$ and find

$$a(x_2) = x_2^{1-q'} - b(x_2)x_2.$$

Then substitute this in and multiply both sides by $x_2^{q'-1}$ to get

$$1 + x_2^{q'} b(x_2) \left(\frac{x_2}{x_1} - 1 \right) = \left(\frac{x_2}{x_1} \right)^{q'-1} g\left(\frac{x_2}{x_1}\right). \text{ So,}$$

$$x_2^{q'} b(x_2) = \frac{\left(\frac{x_2}{x_1}\right)^{q'-1} g\left(\frac{x_2}{x_1}\right) - 1}{\frac{x_1}{x_2} - 1} = c,$$

a constant. Then,

$$(5.13) \quad \mathbb{B}(x) = x_2^{1-q'} \left(1 - c + c \frac{x_1}{x_2} \right).$$

If we let $y = \frac{x_2}{x_1}$, then $y \in [1, \delta]$ and we see $g(y) = y^{-q'}[(1-c)y + c]$. Next,

$$0 \leq g(y) - 1 = y^{-q'}(y-1) \left(\frac{y-y^{q'}}{y-1} - c \right),$$

and since the function $y \rightarrow \frac{y-y^{q'}}{y-1}$ is monotone decreasing, we see that

$$(5.14) \quad c \leq \frac{\delta - \delta^{q'}}{\delta - 1} < 1 - q'.$$

Now, we calculate

$$\frac{\partial \mathbb{B}}{\partial x_2} = x_2^{1-q'} \left[(1-c)(1-q') - q'c \frac{x_1}{x_2} \right],$$

which needs to be non-negative to satisfy (5.11). At $y = 1$, the expression above is positive, given (5.14). If $q \leq \delta$, there is no possible value for c , because then $q' \geq \frac{\delta}{\delta-1}$, in which case $(1-c)(1-q') - \frac{q'c}{\delta} = c(q' - 1 - q'/\delta) + 1 - q' \geq 0$ contradicts (5.14). So, assuming $q > \delta$, we want to choose c in such a way that $\left[(1-c)(1-q') - q'c \frac{x_1}{x_2} \right]$ stays positive on the entire interval $[1, \delta]$. Solving for c , we get

$$c = \frac{\delta(q' - 1)}{\delta(q' - 1) - q'} = \frac{\delta}{\delta - q}.$$

Substituting this into (5.13) yields our candidate function for $\mathbb{B}(x; \infty, q, \delta)$,

$$B(x; \infty, q, \delta) = x_2^{1-q'} \left(\frac{q - \frac{x_1}{x_2} \delta}{q - \delta} \right),$$

which is what appears in (3.4).

If \mathbb{B} solves (5.12), then \mathbb{B} is constant in x_2 , in which case g must be constant, whence $g \equiv 1$ and $\mathbb{B} = x_1^{1-q'}$. But, then $\frac{\partial^2 \mathbb{B}}{\partial x_1^2} = (1-q')(-q')x_1^{-q'-1}$, which is positive. Therefore (5.12) has no solution for $q > 1$.

6. B is concave

We now proceed to verify, in several steps, that $B = \mathbb{B}$, proving Theorem 6. Our first lemma addresses the fact that B is, indeed, concave.

Lemma 1. Case 1, $p < \infty$: *Let x^\pm be two arbitrary points in Ω_δ . If the entire line segment joining these two points (denoted $[x^-, x^+]$) is contained within Ω_δ , then*

$$(6.1) \quad B(\alpha^- x^- + \alpha^+ x^+) \geq \alpha^- B(x^-) + \alpha^+ B(x^+)$$

holds for all non-negative numbers α^\pm with $\alpha^- + \alpha^+ = 1$.

Case 2, $p = \infty$: *Let x^\pm be two arbitrary points in Ω_δ and let α^\pm be a pair of non-negative numbers such that $\alpha^- + \alpha^+ = 1$. Define $x^0 = (x_1^0, x_2^0) := (\alpha^- x_1^- + \alpha^+ x_1^+, \max\{x_2^-, x_2^+\})$. If both of the points (x_1^\pm, x_2^0) are in Ω_δ , then*

$$(6.2) \quad B(x^0) \geq \alpha^- B(x^-) + \alpha^+ B(x^+).$$

Proof. For $p < \infty$, this is a direct calculation, since we simply need to check that the Hessian of B is negative (semi-)definite.

$$\begin{aligned} \frac{\partial^2 B}{\partial x_1^2}(x) &= \frac{-(1 - (p-1)r^\pm)^2 \gamma q'(q'-1)B(x)}{(1 - \gamma r^\pm)(p-1)^2 r^\pm x_1^2}, \\ \frac{\partial^2 B}{\partial x_1 \partial x_2}(x) &= \frac{(1 - (p-1)r^\pm) \gamma q'(q'-1)B(x)}{(1 - \gamma r^\pm)p(p-1)^2 r^\pm x_1 x_2}, \\ \frac{\partial^2 B}{\partial x_2^2}(x) &= \frac{-\gamma q'(q'-1)B(x)}{(1 - \gamma r^\pm)p^2(p-1)^2 r^\pm x_2^2}. \end{aligned}$$

The quadratic form given by the Hessian of B is

$$(6.3) \quad \sum_{i,j=1}^2 \frac{\partial^2 B}{\partial x_i \partial x_j}(x) \Delta_i \Delta_j = \frac{-(1 - (p-1)r^\pm)^2 \gamma q'(q'-1)B(x)}{(1 - \gamma r^\pm)(p-1)^2 r^\pm x_1^2} \left(\Delta_1 - \frac{x_1}{(1 - (p-1)r^\pm)p x_2} \Delta_2 \right)^2.$$

This is non-positive for $q > q^*$ because $(1 - \gamma r^+) > 0$; for $\frac{p-1}{p} < q < q^*$, $\gamma < 0$ so this is non-positive as well. Thus, $B(x; p, q, \delta)$ is concave.

For $p = \infty$, a slightly different approach is needed. First of all, we may assume that $x_2^0 = x_2^+$ due to the symmetry between x^- and x^+ . Also, B is linear in x_1 , so

$$\begin{aligned} B(x^0) - \alpha^- B(x^-) - \alpha^+ B(x^+) \\ &= \alpha^- B(x_1^-, x_2^+) + \alpha^+ B(x_1^+, x_2^+) - \alpha^- B(x^-) - \alpha^+ B(x^+) \\ &= \alpha^- (B(x_1^-, x_2^+) - B(x_1^-, x_2^-)). \end{aligned}$$

This leads us to investigate

$$\frac{\partial B(x_1^-, x_2)}{\partial x_2} = \frac{x_2^{-q'}}{q - \delta} \left[q' \left(\delta \frac{x_1^-}{x_2} - 1 \right) \right],$$

which is non-negative, since we have assumed that $q > \delta$ and that $\frac{x_1^-}{x_2} \geq \frac{1}{\delta}$ for any x_2 between x_2^- and x_2^+ . Hence, $B(x; \infty, q, \delta)$ is concave. \blacksquare

7. How to find extremal weights

We now want to show that $B(x) \leq \mathbb{B}(x)$ on Ω_δ . To do so, given a point $x \in \Omega_\delta$, we will find a weight $w \in RH_p^\delta$ which corresponds to x and which satisfies $B(x) = \langle w^{1-q'} \rangle$. We will (prematurely) call such a weight *extremal*, because once we show that $B = \mathbb{B}$, these weights achieve the supremum which defines \mathbb{B} . The heuristics for finding such weights follows.

Since we know that B is concave and that its Hessian has a kernel, we know that B is linear along certain lines in Ω_δ . We will show later that these lines actually cover Ω_δ . With that, the heuristics above for why B should be concave give us a pattern for how to find extremal weights. We start with $p < \infty$. Given an arbitrary point x^0 on the curve $\Gamma_\delta := \{(x_1, x_2) : x_2 = (\delta x_1)^p\}$, we find a maximal weight representing x^0 (this is far easier than doing so in general). Also, any weight represented by a point x on the graph $\Gamma_1 := \{(x_1, x_2) : x_2 = x_1^p\}$ is constant (and therefore maximal). So, given a point $\hat{x} \in \Omega_\delta$, we find the line along which B is linear which passes through \hat{x} . This line will intersect the graphs Γ_δ and Γ_1 at points which we call x^- and x^+ , respectively. We find the constants α^\pm such that $\hat{x} = \alpha^- x^- + \alpha^+ x^+$. Then, \hat{x} can be represented by the weight w which is the concatenation of the maximal weight for x^- on I^- and the maximal (constant) weight for x^+ on I^+ , re-scaling the intervals if necessary. Since B is linear along this line, we know that $B(\hat{x}) = \alpha^- B(x^-) + \alpha^+ B(x^+) = \alpha^- \langle w^{1-q'} \rangle_{I^-} + \alpha^+ \langle w^{1-q'} \rangle_{I^+} = \langle w^{1-q'} \rangle_I$, whence w is maximal.

We start by finding an extremal weight for a point on the curve Γ_δ . Given an arbitrary positive number x_1^0 , let $x_2^0 = (\delta x_1^0)^p$. Let $I = [0, 1]$. We seek a

weight $w \in RH_p^\delta(I)$ such that $\langle w \rangle_I = x_1^0$, $\langle w^p \rangle_I = x_2^0$ and such that $\langle w^{1-a'} \rangle_I$ is as large as possible. Therefore, for any $a \in [0, 1]$, we insist that

$$\left(\frac{1}{a} \int_0^a w^p(t) dt \right) \left(\frac{1}{a} \int_0^a w(t) dt \right)^{-p} = \delta^p.$$

Let $v(a) := \int_0^a w^p(t) dt$. Then, $w(t) = v'(t)^{1/p}$. Therefore,

$$\begin{aligned} \frac{v(a)}{a} &= \delta^p \left(\frac{1}{a} \int_0^a v'(t)^{1/p} dt \right)^p, \text{ hence} \\ \frac{a}{\delta} \left(\frac{v(a)}{a} \right)^{1/p} &= \int_0^a v'(t)^{1/p} dt. \end{aligned}$$

We take the derivative with respect to a and find

$$\begin{aligned} \delta^{-1} \left[\left(1 - \frac{1}{p}\right) t^{-1/p} v^{1/p} + \left(\frac{1}{p}\right) t^{1-1/p} v^{1/p-1} v' \right] &= (v')^{1/p}, \text{ so} \\ p\delta &= \left(p - 1 + \frac{tv'}{v} \right) \left(\frac{v}{tv'} \right)^{1/p}. \end{aligned}$$

From the definition of $s^\pm = s_p^\pm(\delta)$, we know that

$$p\delta = \left(p - 1 + \frac{1}{1 - ps^\pm} \right) (1 - ps^\pm)^{1/p},$$

so we must have $\frac{v}{tv'} = 1 - ps^\pm$. Consequently,

$$\begin{aligned} \frac{dt}{t} \frac{1}{1 - ps^\pm} &= \frac{dv}{v}, \text{ so} \\ v(t) &= Ct^{\frac{1}{1-ps^\pm}}, \text{ whence} \\ w(t) = v'(t)^{1/p} &= Ct^{\frac{s^\pm}{1-ps^\pm}}. \end{aligned}$$

We want $\langle w \rangle_I = x_1^0$, so we must set $C = x_1^0 \left(\frac{1-(p-1)s^\pm}{1-ps^\pm} \right)$. Putting it all together,

$$(7.1) \quad w(t) = x_1^0 \left(\frac{1 - (p-1)s^\pm}{1 - ps^\pm} \right) t^{\frac{s^\pm}{1-ps^\pm}}.$$

It is straightforward to check that with this constant, $\langle w^p \rangle_I = x_2^0 = (\delta x_1^0)^p$. So, we have found our candidate for an extremal weight representing a point on the curve Γ_δ .

Next, we must find the lines along which B is linear. By (6.3), we see that the vector field along which B is linear is

$$(7.2) \quad p(1 - (p - 1)r^\pm)dx_1 - \frac{x_1}{px_2}dx_2 = 0.$$

To find explicit formulae for the lines, we work with the definition of r^\pm . Recall

$$(7.3) \quad \frac{x_2}{x_1^p} = \delta^p \frac{(1 - pr^\pm)^{p-1}}{(1 - (p - 1)r^\pm)^p}.$$

Therefore,

$$\frac{dx_2}{x_2} - p\frac{dx_1}{x_1} = \frac{-rp(p - 1)dr^\pm}{(1 - pr^\pm)(1 - (p - 1)r^\pm)}.$$

Using (7.2), we see that

$$\frac{dx_1}{x_1} = dr^\pm \left[\frac{p}{1 - pr^\pm} - \frac{p - 1}{1 - (p - 1)r^\pm} \right] \quad \text{and} \quad \frac{dx_2}{x_2} = \frac{pdr^\pm}{1 - pr^\pm},$$

whence

$$(7.4) \quad x_1(r^\pm) = x_1(0) \frac{1 - (p - 1)r^\pm}{1 - pr^\pm} \quad \text{and} \quad x_2(r^\pm) = \frac{x_2(0)}{1 - pr^\pm}.$$

Taking $x_1(0) = b$ as a free parameter, (7.3) gives us $x_2(0) = (\delta b)^p$. And, eliminating r^\pm from (7.4) yields

$$(7.5) \quad \delta^p px_1 - b^{1-p}x_2 = \delta^p b(p - 1),$$

which is the equation of the line tangent to Γ_δ at the point $x = (b, (\delta b)^p)$. Notice that at $r^\pm = 0$, we have the point $(b, (\delta b)^p)$ on the line Γ_δ and at $r^\pm = s^\pm$, we have $x = \left(\frac{b(1 - (p - 1)s^\pm)}{1 - ps^\pm}, \frac{(\delta b)^p}{1 - ps^\pm} \right)$, which satisfies $x_2 = x_1^p$ and hence lies on Γ_1 . It is clear that by varying b , these segments cover Ω_δ .

We now check that B is, in fact, linear along these segments. We use (5.7) as an easier representation of B for this calculation:

$$\begin{aligned} B(x) &= x_1^{-\gamma} x_2 \left(\frac{1 - ps^\pm}{1 - pr^\pm} \right)^\gamma \left(\frac{1 - (p - 1)r^\pm}{1 - (p - 1)s^\pm} \right)^\gamma \left(\frac{1 - \gamma r^\pm}{1 - \gamma s^\pm} \right) \\ &= \left(\frac{b(1 - (p - 1)r^\pm)}{1 - pr^\pm} \right)^{-\gamma} x_2 \left(\frac{1 - ps^\pm}{1 - pr^\pm} \right)^\gamma \left(\frac{1 - (p - 1)r^\pm}{1 - (p - 1)s^\pm} \right)^\gamma \left(\frac{1 - \gamma r^\pm}{1 - \gamma s^\pm} \right) \\ &= \frac{1}{1 - \gamma s^\pm} \left(\frac{1 - ps^\pm}{b(1 - (p - 1)s^\pm)} \right)^\gamma x_2 (1 - \gamma r^\pm). \end{aligned}$$

However, $x_2 = \frac{(\delta b)^p}{1-pr^\pm}$, so $r^\pm = \frac{1}{p} - \frac{(\delta b)^p}{px_2}$ and therefore,

$$x_2(1 - \gamma r^\pm) = x_2\left(1 - \frac{\gamma}{p}\right) + \frac{\gamma(\delta b)^p}{p}.$$

So,

$$B(x) = \frac{1}{1 - \gamma s^\pm} \left(\frac{1 - ps^\pm}{b(1 - (p - 1)s^\pm)} \right)^\gamma \left(x_2\left(1 - \frac{\gamma}{p}\right) + \frac{\gamma(\delta b)^p}{p} \right),$$

and hence is linear.

We now work with an arbitrary point $x^0 \in \Omega_\delta$. Given such a point, one of the segments on which B is linear passes through x^0 ; finding it requires that we find the corresponding value of b . Further, given that x^0 corresponds to $I = [0, 1]$, we want to calculate where to split I so that x^- is on Γ_δ and x^+ is on Γ_1 . We determine b first. x^0 determines a value $r^0 = u_p^\pm \left(\frac{x_2^0}{(\delta x_1^0)^p} \right)$, from which we get

$$x^0 = \left(\frac{b(1 - (p - 1)r^0)}{1 - pr^0}, \frac{(\delta b)^p}{1 - pr^0} \right),$$

and we see

$$b = \frac{x_1^0(1 - pr^0)}{1 - (p - 1)r^0}.$$

Then, if we split I at a , we know that $x_2^0 - x_2^+ = a(x_2^- - x_2^+)$. So, we calculate

$$x_2^0 - x_2^+ = \delta^p b^p p \left(\frac{r^0 - s^\pm}{(1 - pr^0)(1 - ps^\pm)} \right) \quad \text{and} \quad x_2^- - x_2^+ = \frac{-(\delta b)^p ps^\pm}{1 - ps^\pm},$$

whence $a = \frac{s^\pm - r^0}{s^\pm(1 - pr^0)}$.

On I^+ , our weight should be constant, so

$$w(t)|_{I^+} \equiv x_1^+ = \frac{b(1 - (p - 1)s^\pm)}{1 - ps^\pm} = x_1^0 \frac{(1 - pr^0)(1 - (p - 1)s^\pm)}{(1 - (p - 1)r^0)(1 - ps^\pm)}.$$

On I^- , the weight should be maximal. So, we re-scale (7.1) and get

$$\begin{aligned} w(t)|_{I^-} &= b \left(\frac{1 - (p - 1)s^\pm}{1 - ps^\pm} \right) \left(\frac{t}{a} \right)^{\frac{s^\pm}{1 - ps^\pm}} \\ &= x_1^0 \frac{(1 - pr^0)(1 - (p - 1)s^\pm)}{(1 - (p - 1)r^0)(1 - ps^\pm)} \left(\frac{t}{a} \right)^{\frac{s^\pm}{1 - ps^\pm}}. \end{aligned}$$

Therefore, our (potential) extremal weight is

$$(7.6) \quad w(t) = \begin{cases} x_1^0 \frac{(1-pr^0)(1-(p-1)s^\pm)}{(1-(p-1)r^0)(1-ps^\pm)} \left(\frac{t}{a}\right)^{\frac{s^\pm}{1-ps^\pm}} & 0 \leq t \leq a \\ x_1^0 \frac{(1-pr^0)(1-(p-1)s^\pm)}{(1-(p-1)r^0)(1-ps^\pm)} & a \leq t \leq 1, \end{cases}$$

where $a = \frac{s^\pm - r^0}{s^\pm(1-pr^0)}$.

For the case of $p = \infty$, given the above work, finding the extremal weight is rather easy. If we look at the power of t in (7.6) and use the asymptotics in (5.8), we see the power of our extremal weight should be $\delta - 1$. Also, we want the weight to represent a given point $x^0 = (x_1^0, x_2^0)$; recall that the second coordinate when $p = \infty$ is just $x_2 = \text{ess sup}_I w$. Thus, the constant part of the weight must be equal to x_2^0 . This leaves us with the simple task of finding the appropriate splitting point a . We take

$$(7.7) \quad w(t) = \begin{cases} x_2^0 \left(\frac{t}{a}\right)^{\delta-1} & 0 \leq t \leq a \\ x_2^0 & a \leq t \leq 1, \end{cases}$$

and look for an a such that $\langle w \rangle_I = x_1^0$. But,

$$\langle w \rangle_I = x_2^0 \int_0^a \left(\frac{t}{a}\right)^{\delta-1} dt + (1-a)x_2^0 = x_2^0 - a \left(x_2^0 - \frac{x_2^0}{\delta}\right).$$

Therefore, $a = \frac{1-x_1^0/x_2^0}{1-1/\delta}$.

8. $B(x) \leq \mathbb{B}(x)$

Lemma 2. *For every $\delta \geq 1$, $1 < p \leq \infty$, $\frac{p-1}{p} < q < \infty$ and every $x \in \Omega_\delta$, $B(x; p, q, \delta) \leq \mathbb{B}(x; p, q, \delta)$.*

Proof. As usual, we address $p < \infty$ first. Since $x_2 = x_1^p$ if and only if $w(t) \equiv x_1$ is constant, we know that for such points $r_p^\pm = s_p^\pm$ and hence that $\mathbb{B}(x_1, x_1^p; p, q, \delta) = B(x_1, x_1^p; p, q, \delta) = x_1^{1-q}$ for all $x_1 > 0$, $1 < p < \infty$, and $0 < q < \infty$. So, we now consider $\delta > 1$ and points x with $x_2 x_1^{-p} > 1$.

Fix an arbitrary point $x^0 \in \Omega_\delta$ with $x_2^0(x_1^0)^{-p} > 1$. Fix $I = [0, 1]$ and let $a \in (0, 1]$. We define

$$w_{c,a,\nu}(t) = \begin{cases} c \left(\frac{t}{a}\right)^\nu & \text{if } 0 \leq t \leq a \\ c & \text{if } a \leq t \leq 1. \end{cases}$$

Then,

$$(8.1) \quad \langle w_{c,a,\nu}^\theta \rangle_I = \begin{cases} c^{\theta \frac{1+(1-a)\theta\nu}{1+\theta\nu}} & \text{if } \theta\nu > -1 \\ \infty & \text{if } \theta\nu \leq -1. \end{cases}$$

The weight $w_{c,a,\nu}$ has RH_p constant equal to $\frac{1+\nu}{(p\nu+1)^{1/p}}$, as is demonstrated in the appendix (Lemma 7).

As our earlier work suggests, to get an extremal function we use the values

$$(8.2) \quad \nu = \frac{s^\pm}{1 - ps^\pm}, \quad a = \frac{s^\pm - r^0}{s^\pm(1 - pr^0)}, \quad c = x_1^0 \frac{(1 - pr^0)(1 - (p - 1)s^\pm)}{(1 - (p - 1)r^0)(1 - ps^\pm)}.$$

Since the calculations for both cases are very similar, we only address the case of $q > 1$. With $\delta > 1$, we know $s^+ > 0$; further, $s^+ < 1/p$. Therefore, $\frac{s^+ - r^0}{s^+} < 1 - pr^0$, whence $0 < a < 1$. And, $\nu > 0$. Also, for this value of ν , the RH_p constant for $w_{c,a,\nu}$ is equal to δ , as can easily be checked. Using (8.1), we check that the weight $w_{c,a,\nu}$ does indeed represent the point x^0 . We see

$$\langle w_{c,a,\nu} \rangle_I = c \frac{1 + (1 - a)\nu}{1 + \nu}, \quad \text{but}$$

$$1 + \nu = \frac{1 - (p - 1)s^+}{1 - ps^+}, \quad 1 + (1 - a)\nu = \frac{1 - (p - 1)r^0}{1 - pr^0}, \quad \text{whence } \langle w_{c,a,\nu} \rangle_I = x_1^0.$$

Further,

$$\langle w_{c,a,\nu}^p \rangle_I = c^p \frac{1 + (1 - a)p\nu}{1 + p\nu},$$

$$c^p = (x_1^0)^p \frac{\delta^p}{1 - ps^+} \frac{x_2^0(1 - pr^0)}{\delta^p(x_1^0)^p} = x_2^0 \frac{1 - pr^0}{1 - ps^+},$$

$$1 + p\nu = \frac{1}{1 - ps^+}, \quad 1 + (1 - a)p\nu = \frac{1}{1 - pr^0}, \quad \text{thus } \langle w_{c,a,\nu}^p \rangle_I = x_2^0,$$

and $w_{c,a,\nu}$ does represent x^0 . Finally, we check that $w_{c,a,\nu}$ is maximal, assuming that $(1 - q')\nu > -1$:

$$\langle w_{c,a,\nu}^{1-q'} \rangle_I = c^{1-q'} \frac{1 + (1 - a)(1 - q')\nu}{1 + (1 - q')\nu},$$

$$c^{1-q'} = (x_1^0)^{1-q'} \left(\frac{1 - ps^+}{1 - pr^0} \right)^{q'-1} \left(\frac{1 - (p - 1)r^0}{1 - (p - 1)s^+} \right)^{q'-1},$$

$$1 + (1 - q')\nu = \frac{1 - \gamma s^+}{1 - ps^+}, \quad 1 + (1 - a)(1 - q')\nu = \frac{1 - \gamma r^0}{1 - pr^0},$$

hence $\langle w_{c,a,\nu}^{1-q'} \rangle_I = B(x^0)$.

Our assumption that $(1 - q')\nu > -1$ yields the restrictions on q in Theorems 1 and 6:

$$(1 - q')\nu = \frac{(1 - q')s^+}{1 - ps^+}, \quad \text{so}$$

$$(1 - q')\nu > -1 \Leftrightarrow q' < \frac{1 - (p - 1)s^+}{s^+} \Leftrightarrow q > \frac{1 - (p - 1)s^+}{1 - ps^+} = q^*(\delta, p);$$

similarly, for $\frac{p-1}{p} < q < 1$, $(1 - q')\nu > -1 \Leftrightarrow q < q_*$. Therefore, for $\frac{p-1}{p} < q < q_*$ and $q > q^*$, for all p and any $\delta \geq 1$, by the definition of $\mathbb{B}(x)$, we know that $B(x; p, q, \delta) \leq \mathbb{B}(x; p, q, \delta)$.

Also, for $q_* \leq q \leq q^*$, $\mathbb{B}(x)$ is infinite, since the average $\langle w_{c,a,\nu}^{1-q'} \rangle_I$ is infinite. Hence, for $q_* \leq q \leq q^*$, the inequality $B(x) \leq \mathbb{B}(x)$ is trivially true.

Now, for $p = \infty$, we follow the same path. As before, along the line $x_1 = x_2$, the weights are all constant, so $\mathbb{B}(x; \infty, q, \delta) = B(x; \infty, q, \delta)$. We thus consider only $x_2 > x_1$. Again we fix a point $x^0 \in \Omega_\delta$ and consider the weight $w_{c,a,\nu}$. We use our earlier work to inform our choices of $c = x_2^0$, $a = \frac{1-x_1^0/x_2^0}{1-1/\delta}$ and $\nu = \delta - 1$. We check that this weight represents x^0 , first by checking

$$\langle w_{c,a,\nu} \rangle_I = c \frac{1 + (1 - a)\nu}{1 + \nu} = \frac{x_2^0}{\delta} (\delta - \delta(1 - \frac{x_1^0}{x_2^0})) = x_1^0.$$

And, clearly, $\text{ess sup}_I w_{c,a,\nu} = x_2^0$. This weight has RH_∞ constant equal to δ , which, as before, we prove in the appendix (see Lemma 8). Finally, we check that this weight is maximal, assuming $(1 - q')\nu > -1$:

$$\begin{aligned} \langle w_{c,a,\nu}^{1-q'} \rangle_I &= c^{1-q'} \frac{1 + (1 - a)(1 - q')\nu}{1 + (1 - q')\nu} \\ &= (x_2^0)^{1-q'} \frac{1 + (1 - a)(q' - \delta(q' - 1) - 1)}{q' - \delta(q' - 1)} \\ &= (x_2^0)^{1-q'} \frac{q' - \delta(q' - 1) + \delta(q' - 1)(1 - \frac{x_1^0}{x_2^0})}{q' - \delta(q' - 1)} \\ &= (x_2^0)^{1-q'} \frac{q - \delta \frac{x_1^0}{x_2^0}}{q - \delta} = B(x^0). \end{aligned}$$

The restriction that $(1 - q')\nu > -1$ corresponds to $q > \delta$, as in Theorem 2, and the fact that the average $\langle w_{c,a,\nu}^{1-q'} \rangle$ is infinite for $q \leq \delta$ establishes the fact that $\mathbb{B}(x; \infty, q, \delta) \geq B(x; \infty, q, \delta)$ for all $q > 1$. ■

9. $B(x) \geq \mathbb{B}(x)$

We won't be able to prove this directly; instead, we will resort to an approximation procedure which will involve looking at domains Ω_ϵ for $\epsilon > \delta$. We would like to use Lemma 1, but there is a slight difficulty, in that the line joining two points in Ω_δ mentioned there might leave Ω_δ . Thus, our first task is to show that for a given δ , for every $\epsilon > \delta$, there is a way to split the interval in such a way that $[x^-, x^+]$ is contained inside Ω_ϵ .

Lemma 3. Case 1, $p < \infty$: Fix $\delta > 1$. Then for an arbitrary $\epsilon > \delta$ and an arbitrary weight $w \in RH_p^\delta(J)$, there is a splitting $J = J^- \cup J^+$, $|J^\pm| = \alpha^\pm |J|$ such that the entire interval with the endpoints $x^\pm = (\langle w \rangle_{J^\pm}, \langle w^p \rangle_{J^\pm})$ is in Ω_ϵ . Moreover, the splitting parameters α^\pm can be chosen bounded away from 0 and 1 uniformly with respect to w and, therefore, with respect to J as well.

Case 2, $p = \infty$: Fix $\delta > 1$. Then for an arbitrary $\epsilon > \delta$ and an arbitrary weight $w \in RH_\infty^\delta(J)$, there is a splitting $J = J^- \cup J^+$, $|J^\pm| = \alpha^\pm |J|$ such that $\frac{x_2^0}{x_1^0} \leq \epsilon$, where $x^\pm = (\langle w \rangle_{J^\pm}, \text{ess sup}_{J^\pm} w)$ and $x^0 = (\langle w \rangle_J, \text{ess sup}_J w)$. Moreover, the splitting parameters α^\pm can be chosen bounded away from 0 and 1 uniformly with respect to w and, therefore, with respect to J as well.

Proof. (ideas from [17]) We start with $p < \infty$. Picking a weight $w \in RH_p^\delta(J)$ fixes an interval J and a point x^0 in Ω_δ . Starting from this point, we first try $\alpha^\pm = \frac{1}{2}$. If the entire interval $[x^-, x^+] \subset \Omega_\epsilon$ for these parameters, then we fix this splitting and stop. Assuming that some point of $[x^-, x^+]$ is outside of Ω_ϵ for $\alpha^\pm = \frac{1}{2}$, we then consider the possible points of escape from Ω_ϵ . First of all, since $w \in RH_p^\delta(J)$, x^0, x^+ and x^- are all contained within Ω_δ . As they are co-linear, and as the boundary graph $\Gamma_\epsilon := \{(x_1, x_2) : x_2 = \epsilon^p x_1^p\}$ is convex, we know that the portion of $[x^-, x^+]$ which lies outside of Ω_ϵ must be either between x^- and x^0 or between x^0 and x^+ , but not both. Let $\xi = x^-$ in the first case and $\xi = x^+$ in the second case; that is, the only part of $[x^-, x^+]$ which lies outside of Ω_ϵ is contained within the segment $[x^0, \xi]$. We now proceed to change α^+ to bring $[x^-, x^+]$ entirely within Ω_ϵ . If $\xi = x^-$, we decrease α^+ ; if $\xi = x^+$, we increase α^+ . Let $\rho(\alpha^+)$ be the maximum value of $\frac{x_2^{1/p}}{x_1}$ along the segment $[x^0, \xi]$. We already know that $\rho(\frac{1}{2}) > \epsilon$, and, since $\epsilon > \delta$, for ξ sufficiently close to x^0 , $\rho(\alpha^+) < \epsilon$. Further, $\rho(\alpha^+)$ is continuous. Therefore, when changing α^+ from $\frac{1}{2}$, there is a value of α^+ such that $\rho(\alpha^+) = \epsilon$ for the first time; call this “stopping time” value ω^+ (with its corresponding ω^- such that $\omega^+ + \omega^- = 1$). We now check that ω^+ is bounded away from 0 and 1 uniformly with respect to w and I .

If $\xi = x^+$, then $\omega^+ > \frac{1}{2}$ and $\xi_1 - x_1^0 = x_1^+ - x_1^0 = \omega^-(x_1^+ - x_1^-)$. On the other hand, if $\xi = x^-$, then $\omega^- > \frac{1}{2}$ and $\xi_1 - x_1^0 = x_1^- - x_1^0 = \omega^+(x_1^- - x_1^+)$. Thus, $|\xi_1 - x_1^0| = \min\{\omega^\pm\} |x_1^- - x_1^+|$.

At ω^+ , the line passing through x^\pm and x^0 is tangent to Γ_ϵ and touches it at a point we will call τ . The equation for the line tangent to $x_2 = \epsilon^p x_1^p$ for any constant c at the point τ is

$$p\tau_2 x_1 - \tau_1 x_2 = (p - 1)\tau_1 \tau_2.$$

The equation for the points of intersection of this line with the graphs Γ_δ

and Γ_1 , reduces to

$$t\left(\frac{x_1}{\tau_1}\right)^p - p\frac{x_1}{\tau_1} = 1 - p,$$

for some fixed t which has the solutions $x_1 = \tau_1\left(1 + \frac{u_p^\pm(t)}{1 - pu_p^\pm(t)}\right)$. For Γ_δ , $t = \left(\frac{\delta}{\epsilon}\right)^p$ and for Γ_1 , $t = \frac{1}{\epsilon^p}$. Therefore, the line tangent to Γ_ϵ at τ intersects Γ_δ at the points

$$x(\delta)^\pm = \left(\tau_1 \left[1 + \frac{u_p^\pm\left(\left(\frac{\delta}{\epsilon}\right)^p\right)}{1 - pu_p^\pm\left(\left(\frac{\delta}{\epsilon}\right)^p\right)} \right], \tau_2 \left[\frac{1}{1 - pu_p^\pm\left(\left(\frac{\delta}{\epsilon}\right)^p\right)} \right] \right),$$

and it intersects Γ_1 at

$$x(1)^\pm = \left(\tau_1 \left[1 + \frac{u_p^\pm\left(\frac{1}{\epsilon^p}\right)}{1 - pu_p^\pm\left(\frac{1}{\epsilon^p}\right)} \right], \tau_2 \left[\frac{1}{1 - pu_p^\pm\left(\frac{1}{\epsilon^p}\right)} \right] \right).$$

This gives us the following string of inclusions, $[x(\delta)^-, x(\delta)^+] \subset [x^0, \xi] \subset [x^-, x^+] \subset [x(1)^-, x(1)^+]$. Therefore,

$$|x(\delta)_1^+ - x(\delta)_1^-| \leq |x_1^0 - \xi_1| = \min\{\omega^\pm\} |x_1^+ - x_1^-| \leq \min\{\omega^\pm\} |x(1)_1^+ - x(1)_1^-|,$$

and

$$\begin{aligned} \min\{\omega^\pm\} &\geq \frac{|x(\delta)_1^+ - x(\delta)_1^-|}{|x(1)_1^+ - x(1)_1^-|} \\ &= \frac{|(u_p^+\left(\left(\frac{\delta}{\epsilon}\right)^p\right) - u_p^-\left(\left(\frac{\delta}{\epsilon}\right)^p\right))(1 - pu_p^+\left(\frac{1}{\epsilon^p}\right))(1 - pu_p^-\left(\frac{1}{\epsilon^p}\right))|}{|(u_p^+\left(\frac{1}{\epsilon^p}\right) - u_p^-\left(\frac{1}{\epsilon^p}\right))(1 - pu_p^+\left(\left(\frac{\delta}{\epsilon}\right)^p\right))(1 - pu_p^-\left(\left(\frac{\delta}{\epsilon}\right)^p\right))|}, \end{aligned}$$

which is bounded away from zero and depends only on p , δ and ϵ , and neither w nor I .

Now, we turn to the case of $p = \infty$. Since $x_2^0 = \max\{x_2^\pm\}$ and both points x^\pm are in Ω_δ , at least one of the inequalities $\frac{x_2^0}{x_1^\pm} \leq \epsilon$ is always true. First we take $\alpha^- = \alpha^+ = 1/2$; if the required inequalities are both true, we fix this splitting. Otherwise, we start to change the splitting; namely, we increase α^+ if the point (x_1^+, x_2^0) is outside Ω_ϵ and reduce it in the opposite case. By symmetry, it suffices to examine one of the possible situations, say, the case where $\frac{x_2^0}{x_1^+} > \epsilon$ for $\alpha^+ = 1/2$. The points x^\pm do not, in general, depend continuously on α^+ , but the first coordinates x_1^\pm do. Since $x^0 \in \Omega_\delta$ and $\epsilon > \delta$, for all α^+ sufficiently close to 1, $\frac{x_2^0}{x_1^+} < \epsilon$, and, by our assumption, $\frac{x_2^0}{x_1^+} > \epsilon$ for $\alpha^+ = 1/2$. So, when we increase α^+ from $1/2$ (i.e., enlarge I^+), there is a value of α^+ such that $\frac{x_2^0}{x_1^+} = \epsilon$ for the first time. As before, we call

this “stopping time” ω^+ , with $\omega^- := 1 - \omega^+$. We want to check that ω^\pm are bounded away from 0 and 1, using the geometry of the situation. At the stopping time, we know $\epsilon x_1^+ = x_2^0$. Also, since $x_2^0 = x_2^-$, we know that x^0 and x^- both lie on the horizontal line through the point (x_1^+, x_2^0) . Moreover, since $x^- \in \Omega_\delta$, we know that $\frac{x_2^0}{\delta} \leq x_1^- \leq x_2^0$. Therefore, by examining the first coordinates of these points, we see

$$0 < \left(\frac{1}{\delta} - \frac{1}{\epsilon}\right) x_2^0 = \frac{x_2^0}{\delta} - x_1^+ \leq x_1^0 - x_1^+ = \omega^-(x_1^- - x_1^+) \leq \omega^- \left(1 - \frac{1}{\epsilon}\right) x_2^0,$$

from which we see that $\omega^- \geq \frac{\epsilon/\delta - 1}{\epsilon - 1} > 0$. Since $\omega^+ > 1/2$, this proves that ω^\pm are both bounded away from 0 and 1. Further, these bounds depend on δ and ϵ , and on neither w nor J . ■

We now prove the equality of \mathbb{B} and B by establishing the following inequality.

Lemma 4. *For every $\delta \geq 1$, every $x \in \Omega_\delta$, every $p \in (1, \infty]$, every $q \in (\frac{p-1}{p}, \infty)$ and every $\epsilon > \delta$,*

$$\mathbb{B}(x; p, q, \delta) \leq B(x; p, q, \epsilon)$$

With this established, we can pass to the limit as $\epsilon \rightarrow \delta$. For $p < \infty$ and $q > 1$, this is so because s^+ is a continuous, increasing function of δ . Thus, if $s^+(\delta) \geq \frac{1}{\gamma}$, then $s^+(\epsilon)$ is also, in which case $B(x; p, q, \epsilon)$ is infinite and the inequality is trivially true. On the other hand, if $s^+(\delta) < \frac{1}{\gamma}$, then there is a $\kappa > 0$ such that $s^+(\delta + \kappa) = \frac{1}{\gamma}$, and as long as $|\epsilon - \delta| < \kappa$, we know that $B(x; p, q, \epsilon)$ is finite and depends continuously upon ϵ . A similar argument applies to $\frac{p-1}{p} < q < 1$. For $p = \infty$, the continuity in δ is clear; if $\delta \geq q$, then $B(x; \infty, q, \epsilon)$ is infinite. If $\delta < q$, then as long as $|\epsilon - \delta| < (q - \delta)$, $B(x; \infty, q, \epsilon)$ is finite and depends continuously on ϵ . The following proof is due to Vasyunin [17].

Proof of Lemma 4. The statement of the lemma means that for an arbitrary weight $w \in RH_p^\delta(J)$ and for any $\epsilon > \delta$, we have

$$(9.1) \quad \langle w^{1-q'} \rangle_J \leq B(x; p, q, \epsilon).$$

Here $x = (x_1, x_2) = (\langle w \rangle_J, \langle w^p \rangle_J)$. It suffices to prove (9.1) for step functions w because an arbitrary weight w can be approximated by a sequence of step functions w_n so that all averages converge: $\langle w_n^{1-q'} \rangle_J \rightarrow \langle w^{1-q'} \rangle_J$, and $(\langle w_n \rangle_J, \langle w_n^p \rangle_J) \rightarrow (\langle w \rangle_J, \langle w^p \rangle_J)$, and we can pass to the limit in (9.1) because B is continuous in x if it is finite, and if it is infinite then there is nothing

to prove. So, we fix an interval J , a step function $w \in RH_p^\delta(J)$ and a number $\epsilon > \delta$. Starting with the interval $J^{0,0}$, we construct a chain of intervals $J^{n,m}$ in accordance with the rule in Lemma 3. For the intervals in the n th generation, we use the notation $J^{n,m}$, where $J^{n,2k} = (J^{n-1,k})^-$ and $J^{n,2k+1} = (J^{n-1,k})^+$. Consequently, the second index runs from 0 to $2^n - 1$. The corresponding mean values will be labeled by the same pair of indices and $\alpha^{n,m} = |J^{n,m}|/|J|$. By Lemma 1, we can write the following chain of inequalities:

$$\begin{aligned}
 B(x^{0,0}; p, q, \epsilon) &\geq \alpha^{1,0} B(x^{1,0}; p, q, \epsilon) + \alpha^{1,1} B(x^{1,1}; p, q, \epsilon) \\
 (9.2) \qquad \qquad \qquad &\geq \sum_{m=0}^{2^n-1} \alpha^{n,m} B(x^{n,m}; p, q, \epsilon).
 \end{aligned}$$

The latter sum tends to $\langle w^{1-q'} \rangle_J$ as $n \rightarrow \infty$. Indeed, for a fixed step function $w \in RH_p^\delta(I)$, the set $\{x = (\langle w \rangle_J, \langle w^p \rangle_J) : J \subset I\}$ is a compact subset of Ω_δ , and, therefore, the continuous function B is bounded on this subset, say $B \leq M$, i.e., $B(x^{n,m}; p, q, \epsilon) \leq M$ (excluding the case of $B = \infty$, when there's nothing to prove). So if N is the number of discontinuity points for the step function w , then the number of intervals $J^{n,m}$ where w is not a constant is at most N . On all other intervals w is a constant, in which case $B(x^{n,m}; p, q, \epsilon) = (x_1^{n,m})^{1-q'} = w^{1-q'}$, i.e., the corresponding summand is $\langle w^{1-q'} \rangle_{J^{n,m}}$ and the entire sum differs from $\langle w^{1-q'} \rangle_J$ by at most $NM \max_m \alpha^{m,n}$. This latter quantity tends to zero because by Lemma 3, the values α^\pm are bounded away from 0 and 1, and the maximum length of the n th generation intervals tends to 0 as $n \rightarrow \infty$. For $p = \infty$, the proof is nearly identical; simply use $x_2 = \text{ess sup}_J w$ instead of $x_2 = \langle w^p \rangle_J$. ■

10. n -dimensional results

We turn to the proof of Theorem 4. Lemma 6, below, and the maximization argument from section 4 yield an upper bound on the Bellman function. This gives us an upper bound, $C(\delta, p, q, n)$, on the A_q constant of the weights in question. We finish the proof by showing that for a fixed $n > 1$, $p > 1$ and $q > 1$, the limit of $C(\delta, p, q, n)$ as $\delta \rightarrow 1$ is 1.

We only consider n -cubes (equal side lengths) with sides parallel to the coordinate axes. Each time we divide a cube Q , we cut it into 2^n sub-cubes $\{Q_i\}$, each of size $|Q_i| = \frac{|Q|}{2^n}$. Given a cube $Q \subset \mathbb{R}^n$, a weight $w \in RH_p^\delta(Q)$ if

$$\frac{\langle w^p \rangle_K^{1/p}}{\langle w \rangle_K} \leq \delta$$

for all sub-cubes $K \subset Q$. Restricting ourselves to cubes doesn't allow us the flexibility we had before in choosing our splitting, and it is here that we lose the sharpness of the constant.

Given a cube Q and its 2^n sub-cubes $\{Q_i\}$, we first find bounds on the ratio $\frac{\langle w \rangle_{Q_i}}{\langle w \rangle_{Q_j}}$ for δ sufficiently small (Lemma 5). We can then use this bound to find a domain Ω_ϵ which contains the convex hull of the points $\{(\langle w \rangle_{Q_i}, \langle w^p \rangle_{Q_i})\}$ which arise from the splitting of Q . Finally, we use the concavity of our function to produce an upper bound for the Bellman function.

Lemma 5. *Fix $n > 1$, a cube $Q \subset \mathbb{R}^n$ and a weight $w \in RH_p^\delta(Q)$. Divide Q into 2^n equal sub-cubes Q_i in the manner described above. If $1 \leq \delta < (\frac{2^n}{2^n-1})^{1/p'}$, then*

$$\frac{1}{y} \leq \frac{\langle w \rangle_{Q_i}}{\langle w \rangle_{Q_j}} \leq y,$$

where y is given as the solution to

$$(10.1) \quad \left(2 + 2^n(\delta^{-p'} - 1)\right)^{p-1} = \frac{(1+y)^p}{1+y^p}.$$

Proof: First, an observation:

$$(10.2) \quad \frac{\sum_i \langle w \rangle_{Q_i}^p}{(\sum_i \langle w \rangle_{Q_i})^p} \leq \frac{\delta^p}{2^{n(p-1)}}.$$

This can be seen from the following, where we apply Hölder's inequality and then the assumed reverse-Hölder's inequality.

$$\frac{1}{2^n} \sum_{i=1}^{2^n} \langle w \rangle_{Q_i}^p \leq \frac{1}{2^n} \sum_{i=1}^{2^n} \langle w^p \rangle_{Q_i} = \langle w^p \rangle_Q \leq \delta^p \langle w \rangle_Q^p = \delta^p \left(\frac{1}{2^n} \sum_{i=1}^{2^n} \langle w \rangle_{Q_i}\right)^p.$$

The idea is that if δ is sufficiently small, (10.2) restricts the size of the quantity $\frac{\langle w \rangle_{Q_i}}{\langle w \rangle_{Q_j}}$.

Temporarily let $x_i := \langle w \rangle_{Q_i}$, $i = 1 \dots 2^n$. Then arrange the x_i in non-decreasing order and re-label so that $x_1 \leq x_2 \leq \dots \leq x_{2^n}$. Now, set $y_i := \frac{x_i}{x_1}$. Then $1 = y_1 \leq y_2 \leq \dots \leq y_{2^n}$. We are then after the solution to the following optimization problem: given the set

$$S := \left\{ (y_1, y_2, \dots, y_{2^n}) : 1 = y_1 \leq y_2 \leq \dots \leq y_{2^n} \text{ and } \frac{\sum_i y_i^p}{(\sum_i y_i)^p} \leq \frac{\delta^p}{2^{n(p-1)}} \right\},$$

what is

$$\sup \{y_{2^n} : \exists (y_1, y_2, \dots, y_{2^n}) \in S\}?$$

If this supremum is finite, then we have a bound on the ratio between the averages of our weight on different sub-cubes Q_i .

Define the function $k(y_1, y_2, \dots, y_{2^n}) := \frac{\sum_i y_i^p}{(\sum_i y_i)^p}$. A little calculus (and induction) allows one to see that, for any $n > 1$, given a value $y := y_{2^n}$, the minimum value of $k(1, x_2, \dots, x_{2^{n-1}}, y)$ on the region defined by $1 \leq x_2 \leq \dots \leq x_{2^{n-1}} \leq y$ is at the point $(1, a, a, \dots, a, y)$, where $a := \left(\frac{1+y^p}{1+y}\right)^{1/(p-1)}$.

Thus, if we choose our value y so that $k(1, a, \dots, a, y) = \frac{\delta^p}{2^{n(p-1)}}$, we have found our supremum. This value of y solves (10.1), which is only possible (for $1 \leq y < \infty$) if $1 \leq \delta < \left(\frac{2^n}{2^{n-1}}\right)^{1/p'}$. ■

For any weight $w \in RH_p^\delta(Q)$, the set of points

$$P := \{(\langle w \rangle_Q, \langle w^p \rangle_Q), (\langle w \rangle_{Q_1}, \langle w^p \rangle_{Q_1}), \dots, (\langle w \rangle_{Q_{2^n}}, \langle w^p \rangle_{Q_{2^n}})\}$$

lies in the domain $\Omega_\delta := \{(x_1, x_2) : x_1^p \leq x_2 \leq (\delta x_1)^p\}$. Our goal is to find an ϵ such that the convex hull of P lies within Ω_ϵ . Since the curve $\Gamma_\delta := \{(x_1, (\delta x_1)^p)\}$ is convex, if part of the convex hull of P lies outside of Ω_δ , only one part of one edge of the hull lies the furthest outside of Ω_δ . Thus, we can simply focus on pairs of points in P .

Since $\frac{1}{y}\langle w \rangle_{Q_i} \leq \langle w \rangle_{Q_j} \leq y\langle w \rangle_{Q_i}$, we know $\frac{1}{y}\langle w \rangle_Q \leq \langle w \rangle_{Q_j} \leq y\langle w \rangle_Q$. If we label $\langle w \rangle_Q = x_1^0$, as before, then we see that the worst that could happen is if $P_1 := (\frac{x_1^0}{y}, (\delta \frac{x_1^0}{y})^p)$ and $P_2 := (yx_1^0, (\delta yx_1^0)^p)$ are both in P . Thus, the smallest ϵ which guarantees that Ω_ϵ will contain the convex hull of P is such that the line between P_1 and P_2 is tangent to Γ_ϵ . Solving for ϵ yields

$$(10.3) \quad \epsilon = \delta \left[\frac{f_p(y)}{p} \cdot \left(\frac{f_p(y) - 1}{p - 1} \right)^{(1-p)/p} \right],$$

where $f_p(y) := \frac{y^2 - y^{2-2p}}{y^2 - 1}$. As long as $y \geq 1$, (10.3) gives that $\epsilon \geq \delta$ and is bounded.

Consequently, if $\delta < \left(\frac{2^n}{2^{n-1}}\right)^{1/p'}$, there is a value $y(\delta, n, p)$ which solves (10.1). Then, the ϵ which satisfies our conditions is given by (10.3), using $y(\delta, n, p)$, and this ϵ depends only upon δ, n and p .

Now, we proceed as before. The function $B(x; p, q, \delta)$ used in the one-dimensional case is still our “best guess” at the true Bellman function, as the scaling argument and calculations of section 5 are identical in the n -dimensional case. What changes is that we can only prove a restricted version of Lemma 4.

Lemma 6. *For every $n \geq 1$, every $p, q \in (1, \infty)$, every $\delta \in [1, (\frac{2^n}{2^n-1})^{1/p'}]$, every $x \in \Omega_\delta$, if ϵ solves (10.3), with the value $y(\delta, n, p)$ given by (10.1), then*

$$\mathbb{B}(x; p, q, \delta) \leq B(x; p, q, \epsilon)$$

The proof of this lemma, given the above work, is actually easier than the proof of Lemma 4. However, the argument is so similar that we won't repeat it here. The main change is that in the n -dimensional case, the splitting is determined and uniform.

Now, from Lemma 6, we can get an upper bound on the $A_q(Q)$ constant of a weight $w \in RH_p^\delta(Q)$.

$$(10.4) \quad \sup_{K \subset Q} \langle w \rangle_K \langle w^{1-q'} \rangle_K^{q-1} \leq C(\delta, p, q, n),$$

with

$$C(\delta, p, q, n) = \begin{cases} +\infty & 1 < q \leq q^*(p, \epsilon) \\ \frac{1}{q^*(p, \epsilon)} \left(\frac{q-1}{q-q^*(p, \epsilon)} \right)^{q-1} & q > q^*(p, \epsilon) \end{cases},$$

and where $q^*(p, \epsilon)$, is as defined in (2.1), but with δ replaced by the ϵ which solves (10.3), given the y which is a solution to (10.1).

What is important for the proof of Theorem 4 is the limit of this bound as δ approaches 1 for a fixed $p > 1$, $q > 1$, $n > 1$. It is not difficult to see, from (10.1), that for a fixed $n > 1$ and $p > 1$, $\lim_{\delta \rightarrow 1} y(\delta, n, p) = 1$. Also, for a fixed $p > 1$, the function $f_p(y) = \frac{y^2 - y^{2-2p}}{y^2 - 1}$ satisfies $\lim_{y \rightarrow 1} f_p(y) = p$. Consequently, the limit of the ϵ given by (10.3) as $\delta \rightarrow 1$ is 1. By our earlier work on q^* , we know that for a fixed $p > 1$, $\lim_{\epsilon \rightarrow 1} q^*(p, \epsilon) = 1$, whence $\lim_{\delta \rightarrow 1} C(\delta, p, q, n) = 1$. Therefore, given any $n > 1$, $p > 1$, $q > 1$, and $\eta > 1$, by taking δ close enough to 1, we can ensure that every weight $w \in RH_p^\delta(Q)$ satisfies

$$\sup_{K \subset Q} \langle w \rangle_K \langle w^{1-q'} \rangle_K^{q-1} \leq \eta,$$

whence $RH_p^\delta(Q) \subset A_q^\eta(Q)$. This proves Theorem 4.

Proof of Theorem 5. Our earlier work is nearly sufficient; as the Bellman function is dimension-blind, only the splitting of the rectangles and the extremal weights need to be addressed. At the start of the proof of Lemma 3, given the bounded rectangle $I \subset \mathbb{R}^n$ and a weight $w \in s-RH_p^\delta(I)$, re-scale I so that that the longest side(s) of I has length 1. Also, translate I so that (one of) the longest side(s) is the interval $(0, 1)$ in the direction which we will distinguish with the label x_1 (as before, translating and re-scaling I

doesn't affect the Bellman function). Then, split I by cutting this x_1 side a distance $0 < \alpha^- < 1$ from 0, producing two sub-rectangles I^\pm . As before, $|I^\pm| = \alpha^\pm |I|$ (where $\alpha^- + \alpha^+ = 1$), and we get two weights w^\pm defined on I^\pm . The remainder of the splitting is done by further sub-dividing I along the x_1 axis. Given this convention, the proof of Lemma 3 is exactly the same.

Moreover, the extremal weights we found earlier are sufficient here; for a point $x \in \Omega_\delta$, an extremal weight representing x on the cube $I := (0, 1)^n$ is simply

$$w(x_1, x_2, \dots, x_n) = w_{c,a,\nu}(x_1) = \begin{cases} c \left(\frac{x_1}{a}\right)^\nu & \text{if } 0 \leq x_1 \leq a \\ c & \text{if } a \leq x_1 \leq 1. \end{cases}$$

with c, a, ν as before. This is not difficult to check, and it completes the proof of Theorem 5. ■

11. Appendix: Finding the RH_p and RH_∞ constants

Lemma 7. *The RH_p constant for the weight*

$$w_{c,a,\nu}(t) = \begin{cases} c \left(\frac{t}{a}\right)^\nu & \text{if } 0 \leq t \leq a \\ c & \text{if } a \leq t \leq 1, \end{cases}$$

with $0 < a \leq 1$, $c \neq 0$ and $\nu > -\frac{1}{p}$ is $\frac{1+\nu}{(p\nu+1)^{1/p}}$.

Proof: We want to find the supremum of the expression

$$(11.1) \quad \frac{\langle w^p \rangle_J^{1/p}}{\langle w \rangle_J}$$

over all intervals $J \subset I = [0, 1]$. We first notice that the value of c is immaterial, as the ratio (11.1) is invariant if we multiply w by a constant. Therefore, we simplify the calculations and set $c = 1$. It suffices to restrict our attention to intervals $J = [\alpha, \beta]$, with $0 \leq \alpha < a \leq \beta \leq 1$. We will justify this restriction later. We now simply calculate

$$\begin{aligned} \langle w_{a,\nu} \rangle_{[\alpha,\beta]} &= \frac{\beta(\nu + 1) - a\nu - a^{-\nu}\alpha^{\nu+1}}{(\beta - \alpha)(\nu + 1)} \\ \langle w_{a,\nu}^p \rangle_{[\alpha,\beta]} &= \frac{\beta(p\nu + 1) - ap\nu - a^{-p\nu}\alpha^{p\nu+1}}{(\beta - \alpha)(p\nu + 1)}, \end{aligned}$$

so

$$(11.2) \quad \frac{\langle w_{a,\nu}^p \rangle_{[\alpha,\beta]}^{1/p}}{\langle w_{a,\nu} \rangle_{[\alpha,\beta]}} = \frac{(\nu + 1)(\beta - \alpha)^{1-1/p}(\beta(p\nu + 1) - a p\nu - a^{-p\nu} \alpha^{p\nu+1})^{1/p}}{(p\nu + 1)^{1/p}(\beta(\nu + 1) - a\nu - a^{-\nu} \alpha^{\nu+1})}.$$

If we define

$$\lambda := \frac{a^{-\nu} \alpha^{\nu+1}}{\beta(\nu + 1) - a\nu} \quad \text{and} \quad \mu := \frac{a^{-p\nu} \alpha^{p\nu+1}}{\beta(p\nu + 1) - a p\nu},$$

we see that $0 \leq \lambda < 1$ and $0 \leq \mu < 1$, by our restrictions on α , β and ν . Also, define

$$\theta := \frac{1 + \nu}{(p\nu + 1)^{1/p}}.$$

Using these substitutions, and pulling out β , the right-hand side of (11.2) becomes

$$\theta \frac{(1 - \mu)^{1/p}}{1 - \lambda} \left(1 - \frac{\alpha}{\beta}\right)^{1-1/p} \frac{(1 + p\nu(1 - \frac{\alpha}{\beta}))^{1/p}}{1 + \nu(1 - \frac{\alpha}{\beta})}.$$

We simplify further, using $\tau = \nu(1 - \frac{\alpha}{\beta})$, and $K = \frac{(1+p\tau)^{1/p}}{1+\tau}$. Note that for $\nu > -\frac{1}{p}$, we have $0 < K \leq 1$. Then we arrive at

$$(11.3) \quad \frac{\langle w_{a,\nu}^p \rangle_{[\alpha,\beta]}^{1/p}}{\langle w_{a,\nu} \rangle_{[\alpha,\beta]}} = \theta \frac{(1 - \mu)^{1/p}}{1 - \lambda} \left(1 - \frac{\alpha}{\beta}\right)^{1-1/p} K.$$

One further reduction is possible, as

$$(11.4) \quad \lambda^p \mu^{-1} = \beta^{1-p} \alpha^{p-1} K^p, \quad \text{whence} \quad \frac{\alpha}{\beta} = \left(\frac{\lambda}{K}\right)^{p'} \mu^{\frac{-1}{p-1}}$$

So, we arrive at the expression we want to maximize,

$$\theta \frac{(1 - \mu)^{1/p}}{1 - \lambda} \left(1 - \left(\frac{\lambda}{K}\right)^{p'} \mu^{\frac{-1}{p-1}}\right)^{1-1/p} K.$$

Let

$$\phi(\lambda, \mu) = \frac{(1 - \mu)^{1/p}}{1 - \lambda} \left(1 - \left(\frac{\lambda}{K}\right)^{p'} \mu^{\frac{-1}{p-1}}\right)^{1-1/p},$$

and the rest is straightforward calculus.

$$\frac{\partial \phi}{\partial \mu} = \frac{(1 - (\frac{\lambda}{K})^{p'} \mu^{\frac{-1}{p-1}})^{-1/p} (1 - \mu)^{-1/p'}}{p(1 - \lambda)} \left[\left(\frac{\lambda}{K\mu}\right)^{p'} - 1 \right],$$

which is zero at $\mu = \lambda/K$; further, this critical value is the location of a maximum. Consequently, we calculate

$$\phi\left(\lambda, \frac{\lambda}{K}\right) = \frac{1 - \frac{\lambda}{K}}{1 - \lambda},$$

and, we see from (11.4) that $\lambda/K = \mu^{\frac{1}{p}} \left(\frac{\alpha}{\beta}\right)^{\frac{1}{p'}} < 1$. Since $K \leq 1$, we know then that

$$0 \leq \phi\left(\lambda, \frac{\lambda}{K}\right) \leq 1.$$

Consequently,

$$(11.5) \quad \frac{\langle w_{a,\nu}^p \rangle_{[\alpha,\beta]}^{1/p}}{\langle w_{a,\nu} \rangle_{[\alpha,\beta]}} = \theta K \phi(\lambda, \mu) \leq \theta = \frac{1 + \nu}{(p\nu + 1)^{1/p}},$$

which is our desired result. This bound is achieved at $\alpha = 0, \beta = a$. We note here that this bound doesn't depend on a . Consequently, we don't need to treat the case of $\beta < a$ separately, since in that case a is simply a multiplicative constant which doesn't affect the norm. The case of $\beta = a$ (which we have considered) is sufficient to cover this. ■

Lemma 8. *The RH_∞ constant for the weight*

$$w_{c,a,\nu}(t) = \begin{cases} c \left(\frac{t}{a}\right)^\nu & \text{if } 0 \leq t \leq a \\ c & \text{if } a \leq t \leq 1, \end{cases}$$

with $0 < a \leq 1, c \neq 0$ and $\nu > 0$ is $\nu + 1$.

Proof: We seek the supremum of

$$(11.6) \quad \frac{\text{ess sup}_J w}{\langle w \rangle_J}$$

over all subintervals $J \subset I := [0, 1]$. We again notice that the value of c is immaterial, as (11.6) doesn't change when w is multiplied by a constant. Thus, we take $c = 1$. We work with intervals $J = [\alpha, \beta]$ and first prove that it is sufficient to consider $0 \leq \alpha < \beta \leq a$. Clearly, if $\alpha \geq a$, (11.6) is equal to one, which is not maximal; so we only consider $\alpha < a$. If $\beta > a$, then (11.6) is equal to

$$\frac{(\beta - \alpha)}{\int_\alpha^a (t/a)^\nu dt + (\beta - a)},$$

which is maximized when $\beta = a$.

Consequently, we consider $0 \leq \alpha < \beta \leq a$. With this, (11.6) becomes

$$\frac{(\beta/a)^\nu(\beta - \alpha)}{a^{-\nu} \int_\alpha^\beta t^\nu dt} = (\nu + 1) \left(\frac{1 - \frac{\alpha}{\beta}}{1 - (\frac{\alpha}{\beta})^{\nu+1}} \right).$$

Let $y = \alpha/\beta$ and consider the function $y \rightarrow \frac{1-y}{1-y^{\nu+1}}$. On the interval $y \in [0, 1)$, this function is maximized at $y = 0$ and has a value of 1. Hence, $\alpha = 0$. We then see that (11.6) is equal to $\nu + 1$ on the interval $J = [0, \beta]$ for any $\beta \leq a$, and this is our desired constant. ■

Acknowledgement. We would like to thank Professor Sasha Volberg for his stimulating lectures on the Bellman function technique and for the helpful references and guidance he provided us with.

References

- [1] BASILE, L., D'APUZZO, L. AND SQUILLANTE, M.: The limit class of Gehring type G_∞ . *Boll. Un. Mat. Ital. B (7)* **11** (1997), no. 4, 871–884.
- [2] BASILE, L., D'APUZZO, L. AND SQUILLANTE, M.: The limit class of Gehring type G_∞ in the n -dimensional case. *Rend. Mat. Appl. (7)* **21** (2001), no. 1-4, 207–221.
- [3] COIFMAN, R. R. AND FEFFERMAN, C.: Weighted norm inequalities for maximal functions and singular integrals. *Studia Math.* **51** (1974), no. 3, 241–250.
- [4] CRUZ-URIBE SFO, D. AND NEUGEBAUER, C. J.: The structure of the reverse Hölder classes. *Trans. Amer. Math. Soc.* **347** (1995), no. 8, 2941–2960.
- [5] FEFFERMAN, R., KENIG, C. AND PIPHER, J.: The theory of weights and the Dirichlet problem for elliptic equations. *Ann. of Math. (2)* **134** (1991), no. 1, 65–124.
- [6] GARCÍA-CUERVA, J. AND RUBIO DE FRANCIA, J. L.: *Weighted Norm Inequalities and Related Topics*. North-Holland Mathematics Studies **116**. North-Holland Publishing Co., Amsterdam, 1985.
- [7] GEHRING, F. W.: The L_p -integrability of the partial derivatives of a quasiconformal mapping. *Acta Math.* **130** (1973), 265–277.
- [8] HRUŠČEV, S.: A description of weights satisfying the A_∞ condition of Muckenhoupt. *Proc. Amer. Math. Soc.* **90** (1984), 253–257.
- [9] KINNUNEN, J.: Sharp results on reverse Hölder inequalities. *Ann. Acad. Sci. Fenn. Ser. A I Math. Dissertationes* No. **95**, 1994, 34 pp.
- [10] KORENOVSKII, A. A.: Sharp extension of a reverse Hölder inequality and the Muckenhoupt condition (Russian). *Mat. Zametki* **52** (1992), no. 6, 32–44, 158; translation in *Math. Notes* **52** (1992), no. 5-6, 1192–1201.

- [11] MALAKSIANO, N. A.: On exact inclusions of Gehring classes in Muckenhoupt classes (Russian). *Mat. Zametki* **70** (2001), no. 5, 742–750; translation in *Math. Notes* **70** (2001), no. 5-6, 673–681.
- [12] MUCKENHOUPT, B.: The equivalence of two conditions for weight functions. *Studia Math.* **49** (1973/74), 101–106.
- [13] NAZAROV, F., TREIL, S. AND VOLBERG, A.: Bellman function in stochastic control and harmonic analysis. In *Systems, approximation, singular integral operators, and related topics (Bordeaux, 2000)*, 393–423. Oper. Theory Adv. Appl. **129**. Birkhäuser, Basel, 2001.
- [14] NEUGEBAUER, C. J.: The precise range of indices for the RH_R - and AP -weight classes. Preprint, [arXiv:math.CA/9809162v1](https://arxiv.org/abs/math/9809162v1) 28 Sep 1998, 1–8.
- [15] RIOS, C.: L^p regularity of the Dirichlet problem for elliptic equations with singular drift. *Publ. Mat.* **50** (2006), no. 2, 475–507.
- [16] STEIN, E.: *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. Princeton Mathematical Series **43**. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993.
- [17] VASYUNIN, V.: The exact constant in the inverse Hölder inequality for Muckenhoupt weights. *St. Petersburg Math. J.* **15** (2004), no. 1, 49–79.

Recibido: 2 de enero de 2008

Martin Dindoš
 School of Mathematics
 Maxwell Institute for Mathematical Sciences
 University of Edinburgh
 James Clerk Maxwell Building, Kings Buildings
 Mayfield Road, Edinburgh EH9 3JZ, U.K.
 M.Dindos@ed.ac.uk

Treven Wall
 School of Mathematics
 Maxwell Institute for Mathematical Sciences
 University of Edinburgh
 James Clerk Maxwell Building, Kings Buildings
 Mayfield Road, Edinburgh EH9 3JZ, U.K.
 T.Wall@ed.ac.uk