

Stabilizers of stationary actions of lattices in semisimple groups

Darren Creutz

Abstract. Every stationary action of a strongly irreducible lattice or commensurator of such a lattice in a general semisimple group, with at least one higher-rank connected factor, either has finite stabilizers almost surely or finite index stabilizers almost surely. Consequently, every minimal action of such a lattice on an infinite compact metric space is topologically free.

1. Introduction

Nevo, Stuck and Zimmer [17, 18, 20] proved that every ergodic measure-preserving action of a lattice in a connected semisimple Lie group, all of whose simple factors are higher-rank, either has finite stabilizers almost everywhere or finite index stabilizers almost everywhere.

The author and Peterson [6] extended this to lattices in general semisimple groups, and the author [5] relaxed the higher-rank requirement to a single factor.

However, for nonamenable groups such as lattices, measure-preserving actions is not the correct context as actions on compact metric spaces will not, in general, admit invariant measures.

The natural setting for the study of actions of lattices is that of stationary actions (as stationary measures do always exist). There is a natural measure on a lattice (related to the Poisson boundary of the ambient group), and we say an action of a lattice is stationary when it is stationary with respect to that measure (see Definition 2.2). Corollary 7.5 states the following.

Theorem. *Let $\Gamma < G$ be a strongly irreducible lattice in a semisimple group with finite center and no compact factors, at least one simple factor of higher-rank. (Strong irreducibility meaning the projection of the lattice is dense in every proper subproduct of G .) Every ergodic stationary action of Γ has finite stabilizers almost everywhere or finite index stabilizers almost everywhere.*

We also obtain the same dichotomy for dense commensurators of lattices (Corollary 7.4).

As every action on a compact metric space admits a stationary measure, we obtain that every action of Γ on a minimal compact metric space is either topologically free or the space is finite. Consequently, every ergodic uniformly recurrent subgroup is finite (see [11] for the notion of uniformly recurrent subgroups), answering the generalized form of the question posed in [11, Problem 5.4].

Omitting the requirement of a higher-rank factor but requiring at least two factors, every stationary action either has finite stabilizers or is orbit equivalent to an action of \mathbb{Z} .

Stationary intermediate factor theorems and induced actions. We establish an intermediate factor theorem along the lines of those of Stuck and Zimmer [20] for stationary actions (Theorem 3.3): every intermediate factor between the stationary space and the stationary join of the space with the Poisson boundary (which is the product space in the measure-preserving case) must have certain structure. We generalize the induced action [20] to the stationary setting (Definition 4.1) to apply the factor theorem to actions of lattices.

Projecting actions to the ambient group. The other main tool we introduce is a technique for projecting the action of a lattice or commensurator to the ambient group: Theorems 5.3 and 5.1. These techniques give a very general method for relating the stabilizers of the action of a lattice or commensurator to the stabilizers of an action of the ambient group (on some other space). The technique is summarized as: if Λ is a dense commensurator of a lattice in a locally compact second countable group G and η is a stationary random subgroup of Λ , then there exists a stationary random subgroup of G via the closure map on subgroups, see Corollary 5.2.

Related work. Boutonnet and Houdayer [3] proved an operator-algebraic statement that all stationary characters on a lattice in a connected semisimple group, all simple factors of higher-rank, are trivial or arise from finite-index subrepresentations; in particular, their result implies that stationary actions of such lattices have finite stabilizers or finite index stabilizers. Our result covers a much larger class of lattices, including those where only one simple factor is higher-rank and those with p -adic components.

Extremely recently, Bader, Boutonnet, Houdayer and Peterson [1] independently obtained the result on stationary actions of general lattices as a consequence of an incredibly strong operator-algebraic rigidity statement for lattices. Our result on stationary actions of lattices in products of groups, announced prior to their work,¹ relies on dynamical methods very different from their algebraic methods.

¹See, for example, D. Creutz, Essential freeness of stationary actions of lattices. Talk at Vanderbilt Subfactor Seminar, 2018.

2. Lattices, commensurators and stationary actions

Let G be a locally compact second countable group and $\mu \in P(G)$ an admissible probability measure (nonsingular with respect to Haar measure and with support generating the group). The notation $(G, \mu) \curvearrowright (X, \nu)$ means a quasi-invariant action on a probability space with $\mu * \nu = \nu$.

2.1. Stationary actions of lattices

Definition 2.1. Let $\Gamma < G = G_1 \times \cdots \times G_k$ be a lattice in a product of locally compact second countable groups.

The lattice Γ is strongly irreducible when for every proper subproduct G_0 , the projection map $\text{proj}_{G_0}: G \rightarrow G_0$ has the properties that the image of Γ is dense and the map is faithful on Γ .

Definition 2.2. Let $\Gamma < G = G_1 \times \cdots \times G_k$ be a lattice in a product of locally compact second countable groups.

Let $\kappa = \kappa_1 \times \cdots \times \kappa_k$ be an admissible probability measure on G . When G is a semi-simple group, we take κ_j to be K_j -invariant, where K_j is a maximal compact subgroup if G_j is connected, and K_j is a compact open subgroup if G_j is totally disconnected.

Let μ be the admissible probability measure on Γ such that the Γ action on the (G, κ) -Poisson boundary is μ -stationary [16].

A *stationary action* of Γ is an action which is μ -stationary for some μ coming from a product measure κ on G .

2.2. Stationary actions of commensurators

Definition 2.3. Let $\Gamma < G$ be a lattice in a locally compact second countable group. A countable group $\Lambda < G$ is a *commensurator* of Γ when $\Gamma \subseteq \Lambda$ and for all $\lambda \in \Lambda$, the group $\Gamma \cap \lambda\Gamma\lambda^{-1}$ has finite index in both Γ and $\lambda\Gamma\lambda^{-1}$.

We first establish that dense commensurators are, in a suitable sense, lattices in their own right.

Theorem 2.4. *Let $\Gamma < G$ be a strongly irreducible lattice in a semisimple group G with trivial center of higher-rank (meaning G is either a simple higher-rank connected Lie group or G is a product of at least two simple factors). Let Λ be a dense commensurator of Γ . Then there exists a locally compact totally disconnected nondiscrete group H such that Λ can be embedded as a strongly irreducible lattice in $G \times H$.*

Proof. The group H will be the relative profinite completion $\Lambda // \Gamma$ (see, e.g., [6, Section 6] for details, we indicate here the facts needed for the proof). Define the map $\tau: \Lambda \rightarrow \text{Symm}(\Lambda/\Gamma)$ sending λ to its action on Λ/Γ as a symmetry of Λ/Γ . In [6, Theorem 6.3], it states that $H = \overline{\tau(\Lambda)}$ is a locally compact totally disconnected group and $K = \tau(\Gamma)$ is a compact open subgroup.

Let $N = \ker(\tau)$ be the kernel of τ . Then $N \cap \Gamma \triangleleft \Gamma$ and $N \triangleleft \Lambda$. By Margulis' normal subgroup theorem, $N \cap \Gamma$ is either finite or has finite index in Γ . Since G is center-free, if N is finite, it is trivial. Suppose it has finite index. As $N \triangleleft \Lambda$, then $\bar{N} \triangleleft \bar{\Lambda} = G$. The only normal subgroups of G are finite or are proper subproducts. Since N contains a finite index subgroup of Γ , it is infinite, but then N must be contained in a proper subproduct, so a finite index subgroup of Γ would be contained in a proper subproduct, contradicting that Γ is strongly irreducible. So the kernel of τ is trivial.

In [6, Proposition 6.1.2], it states that H will be discrete if and only if Λ normalizes a finite index subgroup of Γ . If Γ_0 is finite index in Γ and $\Gamma_0 \triangleleft \Lambda$, then $\Gamma_0 \triangleleft G$ as Γ_0 is discrete and Λ is dense, but this is impossible. So H is nondiscrete.

Clearly, we can embed $\Lambda \rightarrow G \times H$ diagonally and faithfully (as the kernel of τ is trivial). Likewise, we can embed Γ diagonally, and we can identify both with their images.

Let U be an open neighborhood of the identity in G such that $\Gamma \cap U = \{e\}$. Then $\Lambda \cap U \times K = \Gamma \cap U \times K = \{e\}$. So Λ is discrete in $G \times H$. Let F be a fundamental domain for Γ in G , that is, $\text{Haar}_G(F) < \infty$ and $F\Gamma = G$. Then $(F \times K)\Lambda = G \times H$ (as the projection to H of Λ is dense) and $\text{Haar}_{G \times H}(F \times K) < \infty$ since K is compact. Hence Λ is a lattice in $G \times H$.

By construction, Λ projects densely to H . As Λ is dense in G by hypothesis, it projects densely to G from $G \times H$. For any proper subproduct G_0 of G , Λ projects densely to G_0 since Γ does. Since Γ projects densely to $G_0 \times K$, we have that Λ projects densely to $G_0 \times H$. Thus Λ is strongly irreducible. ■

This allows us to define the following.

Definition 2.5. Let $\Gamma < G$ be a lattice and $\Lambda < G$ a dense commensurator. Let $\kappa = \kappa_G \times \kappa_H$ be an admissible probability measure on $G \times H$. Then there is a probability measure μ on Λ so that the Λ action on the $(G \times H, \kappa)$ -Poisson boundary is μ -stationary.

An action of Λ is *stationary* when it is stationary for some μ coming from such a κ (i.e., when it is a stationary action if we treat Λ as a lattice).

We also need the converse of the above theorem; lattices can be treated as commensurators.

Proposition 2.6. Let $\Lambda < G \times H$ be a strongly irreducible lattice, G and H locally compact second countable groups with H totally disconnected and $K < H$ a compact open subgroup. Let $\Lambda \curvearrowright (X, \nu)$ be a stationary action. Set $\Gamma = \text{proj}_G(\Lambda \cap G \times K)$. Then Γ is a strongly irreducible lattice in G and $\Gamma \curvearrowright (X, \nu)$ is a stationary action.

Proof. Let $L = \Lambda \cap G \times K$. Then L is discrete since K is open. As K is compact, L has finite covolume in $G \times H$. Since K is compact, $\Gamma = \text{proj}_G L$ is discrete, hence is a lattice in G .

Let $\kappa = \kappa_G \times \kappa_H$ be the admissible measure on $G \times H$ and μ_Λ the probability measure on Λ such that the $(G \times H, \kappa)$ -Poisson boundary is μ_Λ -stationary under the Λ -action and $\Lambda \curvearrowright (X, \nu)$ is μ_Λ -stationary.

Since κ is a product measure, $PB(G \times H, \kappa_G \times \kappa_H) = (B_G, \beta_G) \times (B_H, \beta_H)$ is the product of the (G, κ_G) - and (H, κ_H) -boundaries. The boundary map $b: B_G \times B_H \rightarrow P(X)$ has the property that $\text{bar } b_*(\beta_G \times \beta_H) = \nu$.

Define μ_Γ to be the probability measure on Γ so that (B_G, β_G) is μ_Γ -stationary under the Γ -action. Treating $\Gamma < \Lambda < G \times H$, we have that $\text{proj}_H \Gamma$ leaves β_H invariant since κ_H is chosen to be K -invariant. Since $\mu_\Gamma * \beta_G = \beta_G$, this means that $\mu_\Gamma * \beta_G \times \beta_H = \beta_G \times \beta_H$.

Since the boundary map B is Λ -equivariant, it is Γ -equivariant, and so we conclude that

$$\mu_\Gamma * \nu = \mu_\Gamma * \text{bar } b_*(\beta_G \times \beta_H) = \text{bar } b_*(\mu_\Gamma * \beta_G \times \beta_H) = \text{bar } b_*(\beta_G \times \beta_H) = \nu,$$

which means the action is stationary for Γ . ■

3. Stationary factor theorems

The intermediate factor theorems of Stuck–Zimmer [20] and Bader–Shalom [2], as well as those of the author [5, 6], all make assertions about the structure of an intermediate factor A between a measure-preserving G -space (X, ν) and the product of it with the Poisson boundary $(B \times X, \beta \times \nu)$. We now establish such a factor theorem for stationary actions.

3.1. The stationary joining

For a stationary G -space (X, ν) , the product space $(B \times X, \beta \times \nu)$ will not in general be stationary (and indeed will only be when X is measure-preserving). Our factor theorem employs the stationary joining of Furstenberg–Glasner [9] in its place. The reader is referred to Glasner [10] for a detailed exposition on joinings.

Definition 3.1 ([9]). Let G be a locally compact second countable group and μ an admissible measure on G . Let (X, ν) and (Y, η) be (G, μ) -spaces. Let ν_ω and η_ω , for $\omega \in G^\mathbb{N}$, denote the conditional measures (see, e.g., [2, Theorem 2.10]) which exist $\mu^\mathbb{N}$ -almost surely. Define the probability measure on $X \times Y$ by $\rho = \int_{G^\mathbb{N}} \nu_\omega \times \eta_\omega d\mu^\mathbb{N}(\omega)$. Then $(X \times Y, \rho)$ is a joining of (X, ν) and (Y, η) which is μ -stationary.

This joining is called the μ -stationary joining of the two systems and written $(X, \nu) \vee (Y, \eta)$.

3.2. The invariant products functor

To formulate the intermediate factor theorem for general stationary actions, we recall the invariant products functor, introduced in [2].

Definition 3.2. Let $G = G_1 \times \cdots \times G_k$ be a product of locally compact second countable groups. For each j , write $\tilde{G}_j = \prod_{i \neq j} G_i$ for the subproduct excluding G_j .

For a G -space (Y, η) , write (Y_j, η_j) for the \check{G}_j -ergodic components. When $G \curvearrowright (Y, \eta)$ is ergodic, $G_j \curvearrowright (Y_j, \eta_j)$ is ergodic.

The G -map $(Y, \eta) \rightarrow (Y_1, \eta_1) \times \cdots \times (Y_k, \eta_k)$ is the *invariant product functor*.

3.3. The stationary intermediate factor theorem for product groups

Theorem 3.3. *Let $G = G_1 \times \cdots \times G_k$ be a product of at least two locally compact second countable groups and $\mu = \mu_1 \times \cdots \times \mu_k$ be an admissible probability measure on G . Let (X, ν) be an ergodic (G, μ) -space. Let (B, β) be the Poisson boundary for (G, μ) . Let (A, α) be an intermediate factor*

$$(B, \beta) \vee (X, \nu) \rightarrow (A, \alpha) \rightarrow (X, \nu),$$

where the maps are G -maps that compose to the natural projection map. Then A is isomorphic to the relative independent joining of $A_1 \times \cdots \times A_k$ and X over their common factor $X_1 \times \cdots \times X_k$ (where the $A_1 \times \cdots \times A_k$ and $X_1 \times \cdots \times X_k$ are the invariant product functors on A and X , respectively).

Proof. The invariant product functor (see [5, Section 2.12]) mapping a G -space to the product of the spaces of G_j -ergodic components is relatively measure-preserving for stationary actions ([5, Proposition 2.12.1]; see also [2, Proposition 1.10]).

The \check{G}_j ergodic components of $B \vee X$ are $B_j \vee X_j$ (the reader may verify this straightforward fact). Thus we have the commuting diagram of G -maps

$$\begin{array}{ccccc} B \vee X & \longrightarrow & A & \longrightarrow & X \\ \downarrow & & \downarrow & & \downarrow \\ (B_1 \vee X_1) \times \cdots \times (B_k \vee X_k) & \longrightarrow & A_1 \times \cdots \times A_k & \longrightarrow & X_1 \times \cdots \times X_k. \end{array}$$

Since B is a contractive space, the map $B \vee X \rightarrow X$ is relatively contractive, hence so is the map $A \rightarrow X$. As the downward maps are relatively measure-preserving, [5, Theorem 2.41] implies that A is isomorphic to the relative independent joining of $A_1 \times \cdots \times A_k$ and X over $X_1 \times \cdots \times X_k$. ■

Corollary 3.4. *Let $G = G_1 \times \cdots \times G_k$ be a product of locally compact second countable groups and $\mu = \mu_1 \times \cdots \times \mu_k$ be an admissible probability measure on G . Let (X, ν) be an ergodic (G, μ) -space such that $\text{proj}_j \text{stab}(x)$ is dense in G_j for all j almost everywhere. Then (X, ν) is measure-preserving and weakly amenable.*

Proof. The map $s_j: X \rightarrow S(G_j)$ by $x \mapsto \overline{\text{proj}_{G_j} \text{stab}(x)}$ sending each x to a closed subgroup of G_j gives rise to a random subgroup of G_j ; realizing that stationary random subgroup, via [5, Theorem 3.3], as the stabilizers of an action $G_j \curvearrowright (Z_j, \zeta_j)$ gives us the (G, μ) -stationary space $Z = Z_1 \times \cdots \times Z_k$.

By construction, the stabilizers of Z contain $\overline{\text{proj}_1 \text{stab}(x)} \times \cdots \times \overline{\text{proj}_k \text{stab}(x)} = G_1 \times \cdots \times G_k$. As Z is ergodic (since X is), this means Z is the trivial space.

According to [5, Theorem 4.12], $X \rightarrow Z$ is a relatively measure-preserving extension (since X is stationary and $Z = Z_1 \times \cdots \times Z_k$ is the product random subgroup functor of [5] applied to X), hence we conclude that (X, ν) is in fact measure-preserving.

Now consider the factor theorem when A is an affine space over (X, ν) following the approach pioneered by Stuck and Zimmer [20] and used in [2, 5, 6], etc. The reader is referred to [5, Section 2] for details on affine spaces and their relation to strong and weak amenability of actions.

For any affine space A over (X, ν) , as the action of G on the Poisson boundary is strongly amenable [23, Section 4.3], there are G -maps $B \curvearrowright X \rightarrow A \rightarrow X$ which compose to the projection map. In particular, the pushforward of the stationary measure on $B \times X$ endows A with the structure of a (G, μ) -space.

We consider the case when A is orbital over X (has the same stabilizer subgroups). Since the projections of the stabilizers are dense, the stabilizers of X_j are G_j almost surely meaning that each X_j is trivial. Therefore, A is isomorphic to the independent joining of X and $A_1 \times \cdots \times A_k$.

Let $a \in A$. Since A is orbital over X , $\text{stab}(a) = \text{stab}(x)$ for the $x \in X$ such that a maps to x so almost every a has a stabilizer that projects densely to each G_j . As each A_j is a G_j -space, this means that the stabilizers for A_j are G_j almost everywhere. So the A_j are all trivial, which means that A is isomorphic to X .

As this holds for all affine orbital A , it follows that X is weakly amenable. ■

4. Inducing stationary actions

Let Γ be an irreducible lattice in a locally compact second countable group G . Let F be a fundamental domain for G/Γ with associated cocycle $\alpha: G \times F \rightarrow \Gamma$ given by $gf\alpha(g, f) \in F$. Write m for the Haar measure on G restricted to F and normalized to be a probability measure.

4.1. The classical induced action

The classical induced action, due to Zimmer (e.g., [20]), constructs a measure-preserving G -action from a measure-preserving Γ -action as follows: define $G \curvearrowright F \times X$ by

$$g \cdot (f, x) = (gf\alpha(g, f), \alpha(g, f)^{-1}x),$$

and then the measure $m \times \nu$ is preserved by the G -action.

Much of the issue in finding a generalization to stationary actions lies in the fact that there is exactly one G -quasi-invariant measure on $F \times X$ which projects to ν on X , and this measure is not stationary (unless ν is preserved by Γ), so our induced action will not have the property that its projection to X is ν . However, the projection will be in the same measure class.

4.2. The stationary induced action

Let κ be an admissible probability measure on G , and let (B, β) be the Poisson boundary for (G, κ) . Write μ for the probability measure on Γ so that (B, β) is μ -stationary. Since the action of Γ on (B, β) is strongly amenable (as the action of G is and Γ is closed in G), there exists the boundary Γ -map $\pi: B \rightarrow P(X)$ (Zimmer [23]; see, e.g., [2, Theorem 2.14]).

Let $\text{bar}: P(P(X)) \rightarrow P(X)$ be the barycenter map, $\text{bar } \eta = \int_{P(X)} \nu d\eta(\nu)$ (see [8]). Since $\pi_*: P(B) \rightarrow P(P(X))$ is the pushforward map, we have a Γ -map $\tilde{\pi} = \text{bar} \circ \pi_*: P(B) \rightarrow P(X)$ such that $\tilde{\pi}(\beta) = \nu$.

Let (X, ν) be a (Γ, μ) -space. Define the pointwise action $G \curvearrowright F \times X$ as in the classical case. Define the probability measure ρ on $F \times X$ by

$$\rho = \int_F \delta_f \times \tilde{\pi}(f^{-1}\beta) dm(f).$$

Observe that, since m is G -invariant,

$$\begin{aligned} \kappa * \rho &= \int_G \int_F g \cdot (\delta_f \times \tilde{\pi}(f^{-1}\beta)) dm(f) d\kappa(g) \\ &= \int_G \int_F \delta_{gf\alpha(g,f)} \times \alpha(g,f)^{-1} \tilde{\pi}(f^{-1}\beta) dm(f) d\kappa(g) \\ &= \int_G \int_F \delta_{gf\alpha(g,f)} \times \tilde{\pi}(\alpha(g,f)^{-1} f^{-1} g^{-1} g\beta) dm(f) d\kappa(g) \\ &= \int_G \int_F \delta_{f_0} \times \tilde{\pi}(f_0^{-1} g\beta) dm(f_0) d\kappa(g) \\ &= \int_F \delta_{f_0} \times \tilde{\pi}(f_0^{-1} \kappa * \beta) dm(f_0) \\ &= \int_F \delta_{f_0} \times \tilde{\pi}(f_0^{-1} \beta) dm(f_0) = \rho, \end{aligned}$$

which means that $(F \times X, \rho)$ is a (G, κ) -space, i.e., $\kappa * \rho = \rho$.

Definition 4.1. The (G, κ) -space just defined is the *induced stationary action* from the (Γ, μ) -space (X, ν) . We will write $G \times_\Gamma X$ for this space.

4.3. Properties of the induced action

Proposition 4.2. *The measure on the induced stationary space projects to F as m and projects to X as a measure in the same class as ν (when ν is measure-preserving, it projects to ν itself).*

Proof. Observe that $\text{proj}_X \rho = \int_F \tilde{\pi}(f^{-1}\beta) dm(f) = \tilde{\pi}(\check{m} * \beta)$, where \check{m} is the symmetric opposite of m . So $\text{proj}_X \rho$ is in the same measure class as ν since $\check{m} * \beta$ is in the same measure class as β . That $\text{proj}_F \rho = m$ is immediate. ■

Proposition 4.3. *The induced stationary action is weakly amenable if and only if the action on (X, ν) is weakly amenable.*

Proof. This is [5, Proposition 2.6.1]. While that proposition is stated as holding for measure-preserving actions, the result it relies on, stated as [5, Theorem 2.25], is actually a result of Zimmer [22] which holds for quasi-invariant actions. ■

Proposition 4.4. *If the induced stationary action is measure-preserving so is (X, ν) .*

Proof. Since $g \cdot \rho = \int \delta_{f_0} \times \tilde{\pi}(f_0^{-1}g\beta) dm(f_0)$, the induced action being measure-preserving implies that $\tilde{\pi}(f^{-1}\beta) = \tilde{\pi}(f^{-1}g\beta)$ for all $f \in F$ and $g \in G$ meaning that $\tilde{\pi}(\beta)$ is Γ -invariant. ■

4.4. The stationary intermediate factor theorem for lattices

Corollary 4.5. *Let $\Gamma < G = G_1 \times \cdots \times G_k$ be a strongly irreducible lattice in a product of at least two locally compact second countable groups. Let $\Gamma \curvearrowright (X, \nu)$ be an ergodic stationary action such that $\text{proj}_j \text{stab}(x)$ is dense in G_j for all j almost everywhere. Then (X, ν) is measure-preserving and weakly amenable.*

Proof. Let (Y, η) be the induced stationary action. Since $\text{stab}(f, x) = f \text{stab}(x) f^{-1}$, we have that for almost every y and all j , $\overline{\text{proj}_j \text{stab}(y)} = G_j$. Then Y is measure-preserving by Corollary 3.4, hence X is by Proposition 4.4. ■

5. Projecting actions to the ambient group

Theorem 5.1. *Let $\Gamma < G$ be a lattice in a locally compact second countable group and $\Lambda < G$ a dense commensurator of Γ . Let $\Lambda \curvearrowright (X, \nu)$ be an ergodic stationary action. Let $s: X \rightarrow S(G)$ be a Λ -equivariant map. Then there exists an ergodic stationary G -space (Y, η) such that $\text{stab}_* \eta = s_* \nu$.*

Proof. The measure-preserving case, used in [6], is an easy consequence of density. The stationary case requires more care. Treat (X, ν) as a Γ -space, and let

$$\pi: (B, \beta) \rightarrow (P(X), \pi_* \beta)$$

be the boundary Γ -map, where (B, β) is the G -Poisson boundary. Consider the Γ -map $s_* \circ \pi: B \rightarrow P(S(G))$ which has the property that $\text{bar}_{s_* \pi_* \beta} = s_* \nu$.

Since Λ acts on (B, β) and $(P(S(G)), s_* \pi_* \beta)$, and since (B, β) is a contractive Γ -space, [7, Theorem 2.8] states that $s_* \circ \pi$ is in fact a Λ -equivariant map and there is a G -space (Z, ζ) that is Λ -isomorphic to $(P(S(G)), s_* \pi_* \beta)$ via a Λ -map φ .

Since Λ is dense and $G \curvearrowright P(S(G))$ continuously, then $s_* \pi_* \beta$ is in fact G -stationary as for every $g \in G$, there exist $\lambda_n \in \Lambda$ with $\lambda_n \rightarrow g$, and so $g s_* \pi_* \beta = \lim \lambda_n s_* \pi_* \beta = \lim \lambda_n \varphi_* \zeta = \lim \varphi_*(\lambda_n \zeta) = \varphi_*(g \zeta)$.

As the barycenter map is G -equivariant, then $\text{bar } s_{**}\pi_*\beta$ is G -stationary and in particular there is a quasi-invariant action of G on $(S(G), \text{bar } s_{**}\pi_*\beta)$. Then [5, Theorem 3.3] gives an action $G \curvearrowright (Y, \eta)$ such that $\text{stab}_*\eta = \text{bar } s_{**}\pi_*\beta = s_*\nu$ which may be taken to be ergodic since $\Lambda \curvearrowright (X, \nu)$ is ergodic. ■

Corollary 5.2. *Let $\Gamma < G$ be a lattice in a locally compact second countable group and $\Lambda < G$ a dense commensurator of Γ . Every stationary random subgroup of Λ becomes a stationary random subgroup of G under the closure map.*

Theorem 5.3. *Let $G = \prod_j G_j$ be a product of at least two simple locally compact second countable groups, and let $\Gamma < G$ be a strongly irreducible lattice. Let $\Gamma \curvearrowright (X, \nu)$ be an ergodic stationary action. Let $s: X \rightarrow S(G_0)$ be a Γ -equivariant map to the space of closed subgroups of some proper subproduct $G_0 \triangleleft G$. Then there exists an ergodic G_0 -space (Y, η) such that $\text{stab}_*\eta = s_*\nu$.*

Lemma 5.4 ([19, Lemma 5.1]). *Let $\Gamma < G \times H$ be an irreducible lattice, $U \subseteq G$ an open neighborhood of the identity, (B_H, β) the (H, κ) -Poisson boundary, $E \subseteq B_H$ a positive measure set and $\varepsilon > 0$. Then there exists $\gamma \in \Gamma$ such that $\text{proj}_G \gamma \in U^{-1}U$ and $\beta((\text{proj}_H \gamma)E) > 1 - \varepsilon$.*

Proof. Since Γ is irreducible, H acts ergodically on $((G \times H)/\Gamma, m)$, where m is the unique invariant probability measure. Let $K \subseteq H$ be compact with nonempty interior. Then $m((U \times K)\Gamma) > 0$. The random ergodic theorem [14] tells us for m -almost every $z \in (G \times H)/\Gamma$ and $\kappa^{\mathbb{N}}$ -almost every $(\omega_n) \in H^{\mathbb{N}}$ that $\frac{1}{N} \sum_{n=1}^N \mathbb{1}_{(U \times K)\Gamma}(\omega_n^{-1} \cdots \omega_1^{-1}z) \rightarrow m((U \times K)\Gamma) > 0$. So there exists $(u, k) \in U \times K$ such that $\omega_n^{-1} \cdots \omega_1^{-1}(u, k) \in (U \times K)\Gamma$ infinitely often for almost every (ω_n) .

The conditional measures $\beta_{(\omega_n)} = \lim \omega_1 \cdots \omega_n \beta$ are point masses almost surely and $\beta = \int \beta_{(\omega_n)} d\kappa^{\mathbb{N}}((\omega_n))$ so, as $\beta(kE) > 0$ as $\beta(E) > 0$, there exists $(\omega_n) \in H^{\mathbb{N}}$ with $\omega_n^{-1} \cdots \omega_1^{-1}(u, k) \in (U \times K)\Gamma$ infinitely often and $\beta_{(\omega_n)}(kE) = 1$. So there exists $\{n_j\}$ such that $\omega_{n_j}^{-1} \cdots \omega_1^{-1}(u, k) = (u_j, k_j)\gamma_j$ for some $u_j \in U$, $k_j \in K$, and $\gamma_j \in \Gamma$, and we have $\beta(\omega_{n_j}^{-1} \cdots \omega_1^{-1}kE) \rightarrow 1$.

Since $\gamma_j = (u_j, k_j)^{-1}\omega_{n_j}^{-1} \cdots \omega_1^{-1}(u, k) = (u_j^{-1}u, k_j^{-1}\omega_{n_j}^{-1} \cdots \omega_1^{-1}k)$, we have that $\text{proj}_G \gamma_j \in U^{-1}U$ and $\beta(k_j(\text{proj}_H \gamma_j)E) \rightarrow 1$. Since K is compact, there is $\{j_\ell\}$ so that $k_{j_\ell} \rightarrow k_\infty \in K$. We know the sets $E_\ell = k_{j_\ell}(\text{proj}_H \gamma_{j_\ell})E$ have $\beta(E_\ell) \rightarrow 1$ and since $k_\infty \beta$ is in the same measure class as β , then $\beta(k_{j_\ell}^{-1}E_\ell) \rightarrow 1$. So there exists ℓ such that $1 - \varepsilon < \beta(k_{j_\ell}^{-1}E_\ell) = \beta((\text{proj}_H \gamma_{j_\ell})E)$. ■

Proof of Theorem 5.3. Let (B, β) be the Poisson boundary for G , and let $\pi: (B, \beta) \rightarrow (P(X), \pi_*\beta)$ be the boundary map. Consider the Γ -map $s_* \circ \pi: B \rightarrow P(S(G_0))$. Note the barycenter is $\text{bar } s_{**}\pi_*\beta = s_*\nu$.

Write $G = G_0 \times \tilde{G}$, where \tilde{G} is the simple factors of G not in G_0 . Then $(B, \beta) = (B_0, \beta_0) \times (\tilde{B}, \tilde{\beta})$ is the product of the Poisson boundaries of G and G_0 (we treat each as a G -space by letting \tilde{G} act trivially on B_0 and G_0 act trivially on \tilde{B}).

The intermediate factor theorem for products [2, Theorem 1.7] gives a Γ -isomorphism from $(P(S(G_0)), s_{**}\pi_*\nu)$ to a G -space (Z, ζ) via a Γ -map φ and that there is a G -map $p: (B, \beta) \rightarrow (Z, \zeta)$ such that $p = \varphi \circ s_*\pi$. Moreover, $(Z, \zeta) = (Z_0, \zeta_0) \times (\tilde{Z}, \tilde{\zeta})$, where G_0 acts trivially on \tilde{Z} and p splits as $p_0: (B_0, \beta_0) \rightarrow (Z_0, \zeta_0)$ and $\tilde{p}: (\tilde{B}, \tilde{\beta}) \rightarrow (\tilde{Z}, \tilde{\zeta})$.

Write $\text{proj}_0: \Gamma \rightarrow G_0$ for the (faithful) projection map. If $\gamma_n \in \Gamma$ such that $\text{proj}_0\gamma_n \rightarrow e$ in G_0 , then $s(\gamma_n x) \rightarrow s(x)$ as G_0 acts continuously on $S(G_0)$. Then $\gamma_n \tilde{\zeta} = \varphi_*\gamma_n s_{**}\pi_*\beta \rightarrow \varphi_* s_{**}\pi_*\beta = \tilde{\zeta}$ whenever $\text{proj}_0\gamma_n \rightarrow e$ in G_0 .

Let E be a positive measure subset of \tilde{Z} . By Lemma 5.4, there exist $\gamma_n \in \Gamma$ such that $\text{proj}_0\gamma_n \rightarrow e$ in G_0 and $\tilde{\beta}(\gamma_n \tilde{p}^{-1}(E)) \rightarrow 1$ (taking U_n to be a decreasing sequence of open neighborhoods of e and ε to be $1/n$). Then $\tilde{\zeta}(\gamma_n E) \rightarrow 1$ but also $\gamma_n^{-1}\tilde{\zeta} \rightarrow \tilde{\zeta}$, so we conclude that $\tilde{\zeta}(E) = 1$. So $(\tilde{Z}, \tilde{\zeta})$ is trivial, i.e., $(P(S(G_0)), s_{**}\pi_*\beta)$ is Γ -isomorphic to the G_0 -space (Z_0, ζ_0) .

This means that if $\text{proj}_0\gamma_n \rightarrow g_0$ in G_0 , then $\gamma_n \zeta \rightarrow g_0 \zeta$ which is in the same measure class as ζ . So $g_0 s_{**}\pi_*\beta = g_0 \varphi_*^{-1}\tilde{\zeta} = \lim \gamma_n \varphi_*^{-1}\tilde{\zeta} = \lim \varphi_*^{-1}\gamma_n \zeta = \varphi_* g_0 \zeta$ is in the same class as $s_{**}\pi_*\beta$.

As Γ projects densely to G_0 , this means $g_0 s_{**}\pi_*\beta$ is in the same measure class as $s_{**}\pi_*\beta$ for every $g_0 \in G_0$, i.e., that $s_{**}\pi_*\beta$, and therefore also $s_*\nu$, is G_0 -quasi-invariant (and in fact stationary), so the result follows using [5, Theorem 3.3]. ■

6. Stabilizers of lattices in connected groups

Theorem 6.1. *Let $G = G_1 \times \cdots \times G_k$ be a product of simple connected Lie groups, $k \geq 2$. Let $\Gamma < G$ be a strongly irreducible lattice. Let $\Gamma \curvearrowright (X, \nu)$ be an ergodic stationary action. Then either $\text{stab}_\Gamma(x)$ projects densely to each simple G_k almost everywhere or $\text{stab}_\Gamma(x)$ is finite almost everywhere.*

The proof will occupy this section. We begin with an observation of Vershik [21].

Proposition 6.2. *Let Λ be a countable group and $\Lambda \curvearrowright (X, \nu)$ a quasi-invariant action. For $\lambda \in \Lambda$ and for a subgroup $Q < \Lambda$, write*

$$\text{Fix } \lambda = \{x : \lambda x = x\}, \quad \text{Fix } Q = \{x : qx = x \text{ for all } q \in Q\}.$$

For almost every x , the stabilizer $\text{stab}(x)$ satisfies $\text{stab}(x) \subseteq \{\lambda : \nu(\text{Fix } \lambda) > 0\}$. In particular, if $\nu(\text{Fix } \lambda) = 0$ for all $\lambda \neq e$ then the action is essentially free.

Proof. Write $L = \{\lambda : \nu(\text{Fix } \lambda) > 0\}$. Let $Z = \bigcup_{\lambda \notin L} \text{Fix } \lambda$. Then Z is a countable union of measure zero sets and hence is measure zero. For all $x \notin Z$, we then have $\text{stab}(x) \subseteq L$ since $x \notin \text{Fix } \lambda$ for $\lambda \notin L$ meaning that $\lambda x \neq x$ for $\lambda \notin L$. ■

6.1. Howe–Moore groups

Recall that a group is Howe–Moore when for every unitary representation without invariant vectors, the matrix coefficients vanish at infinity [13].

Lemma 6.3. *Let G be a locally compact second countable group with the Howe–Moore property and $G \curvearrowright (Y, \eta)$ a quasi-invariant action on a probability space. If there exists a subgroup Q with $\eta(\text{Fix } Q) > 0$ and \bar{Q} noncompact, then $\text{Fix } Q = \text{Fix } G$, i.e., almost every point in $\text{Fix } Q$ is fixed by all of G .*

Proof. First, observe that for any $f \in L^2(Y, \eta)$ and $q \in Q$,

$$\int_{\text{Fix } Q} f(qy) d\eta(y) = \int_{\text{Fix } Q} f(y) d\eta(y),$$

and therefore $\frac{dq\eta}{d\eta}(y) = 1$ for $y \in \text{Fix } Q$ and $q \in Q$.

Let π be the unitary representation of G on $L^2(Y, \eta)$: $\pi(g)f(y) = f(g^{-1}y)\sqrt{\frac{dg\eta}{d\eta}(y)}$. Let P be the orthogonal projection from $L^2(Y, \eta)$ to the closed G -invariant subspace \mathcal{I} of π -invariant vectors. Then π restricted to $L^2(Y, \eta) \ominus \mathcal{I}$ has no invariant vectors.

Let f be the characteristic function of some positive measure subset $E \subseteq \text{Fix } Q$. Then $f - Pf \in L^2(Y, \eta) \ominus \mathcal{I}$ so, as G has the Howe–Moore property,

$$\langle \pi(g)(f - Pf), (f - Pf) \rangle \rightarrow 0 \quad \text{as } g \text{ leaves compact sets.}$$

For $q \in Q$ and $y \in E \subseteq \text{Fix } Q$, we have that $\pi(q)f(y) = f(q^{-1}y)\sqrt{\frac{dq\eta}{d\eta}(y)} = 1 = f(y)$. For $y \notin E$, we have that $qy \notin E$ (since $qy \in E$ gives $q^{-1}qy = y \in E$), therefore $\pi(q)f(y) = f(q^{-1}y)\sqrt{\frac{dq\eta}{d\eta}(y)} = 0 = f(y)$.

As Pf is π -invariant, then

$$\langle \pi(q)(f - Pf), (f - Pf) \rangle = \|f - Pf\|^2.$$

As \bar{Q} is noncompact, this means that $\|f - Pf\| = 0$, i.e., f is in fact π invariant.

Suppose E is not a G -invariant set. Then there exists $g \in G$ with $\eta(gE \setminus E) > 0$. Writing $\mathbb{1}_E$ for the characteristic function, we have

$$\mathbb{1}_E(y) = (\mathbb{1}_E(y))^2 = (f(y))^2 = (\pi(g)f(y))^2 = \mathbb{1}_{gE}(y)\frac{dg\eta}{d\eta}(y).$$

Therefore, for $y \in gE \setminus E$ we have $0 = \frac{dg\eta}{d\eta}(y)$ contradicting that the G action is quasi-invariant. Therefore, E is G -invariant as a set.

Since every subset of $\text{Fix } Q$ is a G -invariant set, G fixes almost every point in it. ■

6.2. Connected Lie groups

Lemma 6.4. *Let G be a simple connected Lie group and $G \curvearrowright (Z, \xi)$ be a quasi-invariant (not necessarily ergodic) action. Let $C(z)$ be the connected component of $\text{stab}(z)$. Let $d \geq 0$ be a nonnegative integer. If $\dim C(z) \leq d$ almost everywhere, then one of the following holds:*

- $\text{stab}(z) = G$ on a positive measure set;
- $\text{stab}(z)$ is compact almost everywhere;

- $\text{stab}(z)$ is discrete almost everywhere;
- the diagonal action $G \curvearrowright (Z \times Z, \zeta \times \zeta)$ has $\dim C(z_1, z_2) \leq d - 1$ for almost every $z_1, z_2 \in Z$ which have noncompact and nondiscrete stabilizers.

Proof. Suppose $\zeta \times \zeta(\{(x, y) \in Z \times Z : C(x) = C(y) \text{ are noncompact}\}) > 0$. Fubini's theorem tells us that for some x , there is a positive measure set of y so that $\zeta(\{x : C(x) = C(y)\}) > 0$. This means that $Q = C(y)$ is a noncompact closed subgroup of G with $\zeta(\text{Fix } Q) > 0$. Lemma 6.3 then says that $\text{Fix } Q = \text{Fix } G$, so in fact $C(x) = G$ on that positive measure set, and we are in the first case.

Proceed now assuming we are not in the first case. Let

$$M = \{z \in Z : \text{stab}(z) \text{ is not compact and not discrete}\}.$$

For almost every $x, y \in M$, we have that $C(x) \neq C(y)$ by the above and that $\dim C(x) > 0$ and $\dim C(y) > 0$ since they are nondiscrete. Then $\dim C(x) \cap C(y) < \dim C(x) \leq d$. ■

Proposition 6.5. *Let G be a connected simple Lie group. Let $G \curvearrowright (Y, \eta)$ be an ergodic nontrivial G -space. Then there exists a positive integer m such that the diagonal action $G \curvearrowright (Y^m, \eta^m)$ has the property that for almost every $z \in Y^m$, the stabilizer subgroup $\text{stab}(z)$ is discrete or compact.*

Proof. Write $C(y)$ for the connected component of $\text{stab}(y)$. Set $d = \sup \dim C(y) \leq \dim G$.

Suppose $\eta(\{y \in Y : \text{stab}(y) \text{ is not compact and not discrete}\}) > 0$. Apply Lemma 6.4 to conclude that either $\eta(\{y \in Y : \text{stab}(y) = G\}) > 0$ or that $G \curvearrowright (Y^2, \eta^2)$ has the property that for $\eta \times \eta$ -almost every $y_2 \in Y^2$, it holds that $\text{stab}(y_2)$ is at least one of compact, discrete, all of G , or has dimension $\leq d - 1$. Repeating this process by applying Lemma 6.4 to (Y^2, η^2) , since d is finite, we conclude that there is some finite number t of applications so that for $m = 2^t$, we have that the diagonal action $G \curvearrowright (Y^m, \eta^m)$ has the property that almost every stabilizer is at least one of compact, discrete or all of G .

If $\eta^m(\{y_m \in Y^m : \text{stab}(y_m) = G\}) > 0$, then, writing $y_m = (x_1, \dots, x_m)$ for $x_j \in Y$, we have

$$G = \text{stab}(y_m) = \text{stab}(x_1) \cap \text{stab}(x_2) \cap \dots \cap \text{stab}(x_m),$$

so $\text{stab}(x) = G$ for a positive η -measure set of $x \in Y$. By ergodicity then this holds almost everywhere but that contradicts that (Y, η) is nontrivial. ■

Proposition 6.6. *Let G be a semisimple connected Lie group with trivial center and $\Lambda < G$ a countable subgroup and $G \curvearrowright (Y, \eta)$ a quasi-invariant action. If the G -stabilizers are discrete almost everywhere, then $\Lambda \curvearrowright (Y, \eta)$ is essentially free.*

Proof. Suppose the Λ -action is not essentially free. Then there exists $\lambda \in \Lambda$, $\lambda \neq e$, with $\nu(\text{Fix } \lambda) > 0$. Since G is connected and $\lambda \notin Z(G)$, the centralizer subgroup of λ does

not contain an open neighborhood of the identity in G . So there exist $g_n \rightarrow e$ in G such that $g_n \lambda g_n^{-1} \neq \lambda$ for all n . Note that, writing $E = \text{Fix } \lambda$, we have $\eta(g_n E \Delta E) \rightarrow 0$. Take a subsequence along which $\eta(g_n E \Delta E) < 2^{-n-1} \eta(E)$. Then

$$\begin{aligned} \eta\left(E \cap \bigcap_n g_n E\right) &= \eta(E) - \eta\left(E \Delta \bigcap_n g_n E\right) \geq \eta(E) - \sum_{n=1}^{\infty} \eta(E \Delta g_n E) \\ &> \frac{1}{2} \eta(E) > 0. \end{aligned}$$

For $y \in E \cap (\bigcap_n g_n E) = \text{Fix } \lambda \cap \bigcap_n \text{Fix } g_n \lambda g_n^{-1}$, we have that $\lambda y = y$ and $g_n \lambda g_n^{-1} y = y$ for all n , hence $g_n \lambda g_n^{-1} \in \text{stab}_G(y)$ and $\lambda \in \text{stab}_G(y)$. But $g_n \lambda g_n^{-1} \neq \lambda$ and $g_n \lambda g_n^{-1} \rightarrow \lambda$ contradicting that $\text{stab}_G(y)$ is discrete almost everywhere. ■

6.3. Miscellany

Lemma 6.7. *Let G be a locally compact second countable group and K a compact group. Let $Q < G \times K$ be a closed discrete subgroup. Then the projection of Q to G is discrete in G .*

Proof. Let V be an open set in G with compact closure. Then $Q \cap \bar{V} \times K$ is finite since it is closed, hence compact, and discrete. So $Q \cap V \times K$ is finite meaning that $(\text{proj}_G Q) \cap V$ is finite. So for every $q \in \text{proj}_G Q$, we can find an open U with $U \cap \text{proj}_G Q = \{q\}$. ■

Lemma 6.8. *Let H be a group and $L < H$ a subgroup. If τ is a random subgroup of H that is supported on subgroups of L , then τ is supported on subgroups of N for some $N \triangleleft H$ with $N \subseteq L$.*

Proof. Since τ is supported on $S(L)$, for τ -almost every $Q \in S(H)$ in fact $Q \subseteq L$. As τ is H -quasi-invariant, for all h and τ -almost every Q we have $hQh^{-1} \subseteq L$ as well. Then $Q \subseteq \bigcap_h h^{-1}Lh$ for almost every Q . Set $N = \bigcap_h h^{-1}Lh$ which is normal in H and contained in L . Then $Q \subseteq N$ almost everywhere as claimed. ■

Recall that a group is *locally finite* when every finitely generated subgroup of it is finite.

Proposition 6.9. *Let Γ be a virtually torsion-free countable discrete group and $\Gamma \curvearrowright (X, \nu)$ a quasi-invariant action with locally finite stabilizers almost everywhere. Then the stabilizers are finite almost everywhere.*

Proof. Since Γ is virtually torsion-free, there is a finite upper bound ℓ for the length of every strictly increasing chain of finite subgroups.

As $\text{stab}(x)$ is locally finite, if it is infinite, then it contains an infinite abelian subgroup A (Hall and Kulatilaka [12]). But A must be of the form $\bigoplus_{n=1}^{\infty} F_n$ for some nontrivial finite subgroups F_n , and then $A_N = \bigoplus_{n=1}^N F_n$ would be a strictly increasing chain of finite subgroups of infinite length. Therefore, $\text{stab}(x)$ must in fact be finite. ■

6.4. Lattices and projected actions

Proposition 6.10. *Let $G = G_1 \times \cdots \times G_k$ be a product of simple connected Lie groups, $k \geq 2$. Let $\Gamma < G$ be a strongly irreducible lattice. Let $\Gamma \curvearrowright (X, \nu)$ be an ergodic stationary action. Then either $\text{stab}_\Gamma(x)$ is locally finite almost everywhere or $\text{stab}_\Gamma(x)$ has dense projections to each simple G_j almost everywhere.*

Proof. For each $j \in \{1, 2, \dots, k\}$, the projection map $\text{proj}_{G_j}: G \rightarrow G_j$ has the property that the image of Γ is dense in G_j and the map is faithful on Γ . Define the map $s_j: X \rightarrow S(G_j)$ by $\overline{s_j}(x) = \overline{\text{proj}_{G_j} \text{stab}_\Gamma(x)}$.

By Theorem 5.3, there is an ergodic G_j -space (Y_j, η_j) such that $(\text{stab}_{G_j})_* \eta_j = (s_j)_* \nu$. Define the sets

$$J_d = \{j \in \{1, \dots, k\} : (Y_j, \eta_j) \text{ is trivial}\}, \quad J_c = \{1, \dots, k\} \setminus J_d.$$

Note that for $j \in J_d$, we have that $\text{stab}_* \eta_j$ is a point mass δ_Q for some $Q \in S(G_j)$. As $\text{stab}_* \eta_j$ is Γ -invariant, this means $\gamma Q \gamma^{-1} = Q$ for all γ . As Γ is dense in G_j , this means Q is normal in G , so $Q = G$ or $Q = \{e\}$. If $Q = \{e\}$, then $\Gamma \curvearrowright (X, \nu)$ is essentially free since the projection map is faithful, in which case the proof is complete. So we proceed with the premise that $\text{proj}_{G_j} \text{stab}(x)$ is dense in G_j almost everywhere for $j \in J_d$.

From here on, we assume that $J_c \neq \emptyset$ since if it is empty, then the proof is complete as the projections of the stabilizers are dense to each G_j almost everywhere.

Define the groups

$$G_c = \prod_{j \in J_c} G_j \quad \text{and} \quad G_d = \prod_{j \in J_d} G_j$$

so that $G = G_c \times G_d$ (after appropriate rearrangement).

For $j \in J_c$, Proposition 6.5 says there exists a positive integer m_j so that for the diagonal action, $G_j \curvearrowright (Y_j^{m_j}, \eta_j^{m_j})$ has the property that almost every stabilizer is compact or is discrete.

Set $m = \max\{m_j : j \in J_c\}$. Then almost every stabilizer of $(Y_j, \eta_j)^m$ is discrete or compact. Hence, for almost every $(y_1, \dots, y_m) \in Y_j^m$, we have that $\bigcap_{i=1}^m \text{stab}(y_i)$ is discrete or compact. Then $\bigcap_{i=1}^m \overline{\text{proj}_{G_j} \text{stab}_\Gamma(x_i)}$ is discrete or compact for ν^m -almost every $(x_1, \dots, x_m) \in X^m$.

So for ν^m -almost every (x_1, \dots, x_m) , we have $\overline{\text{proj}_{G_j} \bigcap_{i=1}^m \text{stab}_\Gamma(x_i)}$ is discrete or compact.

Discrete stabilizers. The set $D = \{z \in Y_j^m : \text{stab}(z) \text{ is discrete}\}$ is G_j -invariant, and if it is positive measure, we may consider $G_j \curvearrowright (D, (\eta^m)^*)$, where $(\eta^m)^*$ is η^m restricted to D and normalized to be a probability measure. Applying Proposition 6.6, we determine that $\text{stab}(z) \cap \text{proj}_{G_j} \Gamma = \{e\}$ for almost every $z \in D$. Now if $\text{proj}_{G_j} \text{stab}_\Gamma(x_1, \dots, x_m) \cap \text{proj}_{G_j} \Gamma = \{e\}$, then $\text{proj}_{G_j} \text{stab}_\Gamma(x_1, \dots, x_m) = \{e\}$ and, as the projection is faithful on Γ , then $\text{stab}_\Gamma(x_1, \dots, x_m) = \{e\}$.

Compact stabilizers. For $z \notin D$, the corresponding $\text{proj}_{G_j} \text{stab}_\Gamma(x_1, \dots, x_m)$ is contained in a compact subgroup. Therefore, $\text{proj}_{G_j} \text{stab}_\Gamma(x_1, \dots, x_m)$ is contained in a compact subgroup of G_j a.e. for all $j \in J_c$.

The diagonal action $\Gamma \curvearrowright (X, \nu)^m$. Since $J_c \neq \emptyset$, by Theorem 5.3 there is a G_d -space (Z, ζ) such that $\text{stab}_* \zeta = s_* \nu$, where

$$s(x_1, \dots, x_m) = \overline{\text{proj}_{G_d} \text{stab}_\Gamma(x_1, \dots, x_m)}.$$

Since $\text{stab}_\Gamma(x_1, \dots, x_m)$ is a discrete subgroup of $G_c \times G_d$ and is contained in $K \times G_d$ for some compact subgroup K of G_c , Lemma 6.7 gives that $\text{proj}_{G_d} \text{stab}_\Gamma(x_1, \dots, x_m)$ is discrete in G_d .

The case when $J_d \neq \emptyset$. In this case, $\text{stab}_* \zeta$ is a stationary random subgroup of G_d which is supported on subgroups of $\text{proj}_{G_d} \Gamma$. But this means $\text{stab}_* \zeta = \delta_{\{e\}}$ by Lemma 6.8 since the only normal subgroup of G_d that is contained in Γ is the trivial group (as G_d is nontrivial when $J_d \neq \emptyset$). So we have that $\Gamma \curvearrowright (X, \nu)^m$ is essentially free. For any $\gamma \in \Gamma$ with $\nu(\text{Fix } \gamma) > 0$, also $\nu^m(\text{Fix}_{X^m} \gamma) > 0$. So we have that $\Gamma \curvearrowright (X, \nu)$ is essentially free when $J_d \neq \emptyset$.

The case when $J_d = \emptyset$. In this case, $\text{stab}_\Gamma(x_1, \dots, x_m)$ is contained in a compact subgroup of G almost everywhere. Then for any $\gamma \in \Gamma$ with $\nu(\text{Fix } \gamma) > 0$, since also $\nu^m(\text{Fix } \gamma) > 0$, we have that the group generated by γ is discrete in G but contained in a compact group hence is torsion. If $\text{stab}_\Gamma(x)$ contains a finitely generated infinite subgroup on a positive measure set, then, as there are only countably many finitely generated subgroups of Γ , there is some infinite group A with $A \subseteq \text{stab}_\Gamma(x)$ on a positive measure set. Then $\nu(\text{Fix } A) > 0$ and so $(\eta_1 \times \dots \times \eta_k)^m(\text{Fix } A) > 0$ but then A must be contained in a compact group, which, since A is discrete in G , contradicts that it is infinite. Therefore, when $J_d = \emptyset$, the Γ -stabilizers on X are locally finite. ■

Proof of Theorem 6.1. Proposition 6.10 tells us that either the stabilizers project densely to every proper subproduct or else the stabilizers are locally finite. Selberg's lemma states that Γ , being a lattice in a semisimple Lie group, is virtually torsion-free, so there exists a finite index $\Gamma_0 < \Gamma$ which is torsion-free. Proposition 6.9 then gives that the stabilizers are finite. ■

7. Stationary actions of lattices and commensurators

7.1. Actions of lattices in connected groups

Theorem 7.1. *Let $\Gamma < G = G_1 \times \dots \times G_k$ be a strongly irreducible lattice in a product of at least two connected simple Lie groups. Let $\Gamma \curvearrowright (X, \nu)$ be an ergodic stationary action. Then either $\text{stab}_\Gamma(x)$ is finite almost everywhere or the action is weakly amenable and measure-preserving.*

Proof. Theorem 6.1 tells us that either the stabilizers are finite or they project densely to each simple factor. When the projections are dense, Corollary 4.5 tells us the action is weakly amenable and measure-preserving. ■

Note that weak amenability equates to orbit equivalence to a \mathbb{Z} -action (Connes–Feldman–Weiss [4]).

Theorem 7.2. *Let $\Gamma < G = G_1 \times \cdots \times G_k$ be a strongly irreducible lattice in a product of connected simple Lie groups, at least two simple factors with one of higher-rank. Let $\Gamma \curvearrowright (X, \nu)$ be an ergodic stationary action. Then either $\text{stab}_\Gamma(x)$ is finite almost everywhere or $\text{stab}_\Gamma(x)$ is finite index almost everywhere.*

Proof. By [5, Corollary 8.6], if the action weakly amenable and measure-preserving, then the stabilizers are finite index almost everywhere. ■

7.2. Actions of dense commensurators

Theorem 7.3. *Let $\Gamma < G$ be a strongly irreducible lattice in a connected semisimple Lie groups, at least one of higher-rank. Let $\Lambda < G$ be a dense commensurator of Γ such that the relative profinite completion $\Lambda // \Gamma$ is a product of simple uncountable totally disconnected locally compact groups with the Howe–Moore property. Let $\Lambda \curvearrowright (X, \nu)$ be an ergodic stationary action. Then either $\text{stab}_\Lambda(x)$ is finite almost everywhere or $\text{stab}_\Lambda(x)$ is finite index almost everywhere.*

Proof. By Proposition 2.6, the restriction of the action to Γ is stationary. By Theorem 7.2, the restriction of the action to Γ has stabilizers which are finite or finite index almost everywhere (on each Γ -ergodic component one or the other must occur). For $\lambda \in \Lambda$, we know $\Gamma \cap \lambda\Gamma\lambda^{-1}$ is finite index in both Γ and $\lambda\Gamma\lambda^{-1}$. If $[\Gamma : \text{stab}_\Gamma(x)] < \infty$, then $[\Gamma : \text{stab}(x) \cap \Gamma] < \infty$ and so $[\lambda\Gamma\lambda^{-1} : \lambda\text{stab}(x)\lambda^{-1} \cap \lambda\Gamma\lambda^{-1}] < \infty$. Using the fact that $\Gamma \cap \lambda\Gamma\lambda^{-1}$ is finite index, we have $[\Gamma \cap \lambda\Gamma\lambda^{-1} : \lambda\text{stab}(x)\lambda^{-1} \cap \lambda\Gamma\lambda^{-1} \cap \Gamma] < \infty$, and so $[\Gamma : \lambda\text{stab}(x)\lambda^{-1} \cap \Gamma] < \infty$. But this just says $[\Gamma : \text{stab}_\Gamma(\lambda x)] < \infty$. So the set where the Γ -stabilizers are finite index is Λ -invariant, hence by Λ -ergodicity the Γ -stabilizers are either finite almost everywhere or finite index almost everywhere.

Finite Γ -stabilizers. Let $H = \Lambda // \Gamma$ be the relative profinite completion. Then $\Lambda < G \times H$ is a strongly irreducible lattice (Theorem 2.4).

When $\text{stab}_\Gamma(x)$ is finite, since $\text{proj}_H \text{stab}_\Lambda(x) \cap K = \text{proj}_H \text{stab}_\Lambda(x) \cap \text{proj}_H \Gamma$ by [6, Proposition 6.1.3],

$$\text{proj}_H \text{stab}_\Lambda(x) \cap K = \text{proj}_H (\text{stab}_\Lambda(x) \cap \Gamma) = \text{proj}_H \text{stab}_\Gamma(x)$$

is finite. So $\text{proj}_H \text{stab}_\Lambda(x)$ is discrete in H as K is open. In particular, $\overline{\text{proj}_H \text{stab}_\Lambda(x)} = \text{proj}_H \text{stab}_\Lambda(x) \subseteq \text{proj}_H \Lambda$ almost everywhere.

Treating Λ as an irreducible lattice in $G \times H$, Theorem 5.3 states that there is an ergodic H -space (Y, η) such that $\text{stab}_* \eta = s_* \nu$, where $s: X \rightarrow S(H)$ is the Λ -map $s(x) = \overline{\text{proj}_H \text{stab}_\Lambda(x)}$.

Then $\text{stab}_*\eta$ is a random subgroup of H which is supported on subgroups of $\text{proj}_H \Lambda$. Lemma 6.8 then says it is supported on subgroups of N for some $N \triangleleft H$ with $N \subseteq \text{proj}_H \Lambda$.

Since $H = \prod H_j$ is a product of simple groups, N is a proper subproduct. But $N \subseteq \text{proj}_H \Lambda$, so if N is nontrivial, then there is some simple H_j which is contained in $\text{proj}_H \Lambda$, meaning that H_j is countable, contradicting our hypothesis.

So N is trivial, meaning that $\text{proj}_H \text{stab}_\Lambda(x) = \{e\}$ almost everywhere. The kernel of proj_H is contained in Γ hence is finite (by Margulis' normal subgroup theorem, if it is infinite, then it is finite index; but that would make H discrete hence countable). Therefore, $\text{stab}_\Lambda(x)$ is finite almost everywhere.

Finite index Γ -stabilizers. If $\text{stab}_\Gamma(x)$ is finite index in Γ , we have that $\text{proj}_H \text{stab}_\Lambda(x)$ contains a finite index subgroup of $\text{proj}_H \Gamma$, and therefore $\overline{\text{proj}_H \text{stab}_\Lambda(x)}$ contains a finite index subgroup of $\overline{\text{proj}_H \Gamma} = K$.

As K is a compact open subgroup, then $\overline{\text{proj}_H \text{stab}_\Lambda(x)}$ contains a compact open subgroup and hence is open almost everywhere.

Write $H = \prod H_j$, then $\text{proj}_{H_j} \text{stab}_\Lambda(x)$ is an open subgroup of H_j almost everywhere for each j (as the projection maps are open maps). Since each H_j has the Howe–Moore property, the only proper open subgroups are compact. Write

$$J_c = \{j : \overline{\text{proj}_{H_j} \text{stab}_\Lambda(x)} \text{ is compact a.e.}\},$$

$$J_d = \{j : j \notin J_c\} = \{j : \overline{\text{proj}_{H_j} \text{stab}_\Lambda(x)} = H_j \text{ a.e.}\}$$

and define the groups

$$H_c = \prod_{j \in J_c} H_j, \quad H_d = \prod_{j \in J_d} H_j.$$

If $J_c \neq \emptyset$, then $\text{stab}_\Lambda(x) < G \times H_d \times H_c$ is discrete and $\text{stab}_\Lambda(x) < G \times H_d \times K_x$ for some compact K_x , so Lemma 6.7 says that $\text{proj}_{G \times H_d} \text{stab}_\Lambda(x)$ is discrete in $G \times H_d$.

Since Λ is strongly irreducible as a lattice in $G \times H$, and since H_d is a proper subproduct of H , Λ is a dense commensurator of a lattice in $G \times H_d$ (Proposition 2.6) so Theorem 5.1 tells us there is an ergodic $G \times H_d$ -space (Y, η) such that $\text{stab}_*\eta = s_*\nu$, where $s(x) = \overline{\text{proj}_{G \times H_d} \text{stab}_\Lambda(x)}$ is the Λ -map $s: X \rightarrow S(G \times H_d)$.

Then $\text{stab}_*\eta$ is a random subgroup of $G \times H_d$ supported on subgroups of $\text{proj}_{G \times H_d} \Lambda$. Lemma 6.8 then says it is supported on subgroups of N for some $N \triangleleft G$ with $N \subseteq \text{proj}_G \Lambda$. But no proper subproduct of $G \times H_d$ can be contained in a countable group, so N is trivial. This means that $\text{proj}_{G \times H_d} \text{stab}_\Lambda(x) = \{e\}$ contradicting that the Γ -stabilizers are finite index.

So $J_c = \emptyset$ and $H_d = H$. Then $\text{stab}_\Lambda(x)$ projects densely into each H_j almost everywhere. Since it also projects densely into each simple factor of G (as the Γ -stabilizers do), Corollary 4.5 then tells us that $\Lambda \curvearrowright (X, \nu)$ is weakly amenable and measure-preserving, so by [5, Corollary 8.6] (every simple H_j acts ergodically on the induced space as the projections of the stabilizers are dense), it is essentially transitive meaning that $\text{stab}_\Lambda(x)$ is finite index in Λ a.e. ■

Corollary 7.4. *Let G be a semisimple connected Lie group with finite center, with at least one simple factor of higher-rank, and $\Gamma < G$ be a finitely generated, strongly irreducible lattice. Let Λ be a dense commensurator of Γ . Then every stationary action of Λ has finite stabilizers almost everywhere or has finite index stabilizers almost everywhere.*

Proof. By Margulis' arithmeticity theorem [15], Γ is the \mathbb{Z} -points of some semisimple algebraic group \mathbf{G} (up to finite index and finite kernels). The commensurator of $\mathbf{G}(\mathbb{Z})$ is $\mathbf{G}(\mathbb{Q})$. The relative profinite completion $\Lambda // \Gamma$ is therefore of the form $\prod \mathbf{G}(\mathbb{Q}_p)$ by [6, Theorem 10.2]. Since simple p -adic groups are Howe–Moore, Theorem 7.3 gives the conclusion. ■

7.3. Actions of lattices in general semisimple groups

Corollary 7.5. *Let $\Gamma < G$ be a strongly irreducible lattice in a semisimple group with finite center and no compact factors, with at least one connected higher-rank simple factor. Every stationary action of Γ either has finite stabilizers or finite index stabilizers.*

Proof. Write $G = G_0 \times H$, where G_0 is connected and H is totally disconnected. If H is trivial, then Theorem 7.2 gives the conclusion; if not, then by Proposition 2.6 the projection of Γ to G is a dense commensurator of a strongly irreducible lattice $\Gamma_0 < G_0$ and any stationary action of Γ is stationary for Γ_0 , so Corollary 7.4 then gives the result (the case when G is a simple connected higher-rank group is covered by [3]). ■

Acknowledgments. The author would like to thank R. Boutonnet for numerous helpful discussions, and particularly for pointing out some flaws in early drafts.

Funding. This work was partially supported by a Naval Academy Research Council grant.

References

- [1] U. Bader, R. Boutonnet, C. Houdayer, and J. Peterson, [Charmenability of arithmetic groups of product type](#). *Invent. Math.* **229** (2022), no. 3, 929–985 Zbl 07569645 MR 4462622
- [2] U. Bader and Y. Shalom, [Factor and normal subgroup theorems for lattices in products of groups](#). *Invent. Math.* **163** (2006), no. 2, 415–454 Zbl 1085.22005 MR 2207022
- [3] R. Boutonnet and C. Houdayer, [Stationary characters on lattices of semisimple Lie groups](#). *Publ. Math. Inst. Hautes Études Sci.* **133** (2021), 1–46 Zbl 1504.22009 MR 4292738
- [4] A. Connes, J. Feldman, and B. Weiss, [An amenable equivalence relation is generated by a single transformation](#). *Ergodic Theory Dynam. Systems* **1** (1981), no. 4, 431–450 Zbl 0491.28018 MR 662736
- [5] D. Creutz, [Stabilizers of actions of lattices in products of groups](#). *Ergodic Theory Dynam. Systems* **37** (2017), no. 4, 1133–1186 Zbl 1369.37003 MR 3645514
- [6] D. Creutz and J. Peterson, [Stabilizers of ergodic actions of lattices and commensurators](#). *Trans. Amer. Math. Soc.* **369** (2017), no. 6, 4119–4166 Zbl 1375.37007 MR 3624404

- [7] D. Creutz and Y. Shalom, [A normal subgroup theorem for commensurators of lattices](#). *Groups Geom. Dyn.* **8** (2014), no. 3, 789–810 Zbl [1306.22006](#) MR [3267524](#)
- [8] A. Furman, [Random walks on groups and random transformations](#). In *Handbook of dynamical systems, Vol. 1A*, pp. 931–1014, North-Holland, Amsterdam, 2002 Zbl [1053.60045](#) MR [1928529](#)
- [9] H. Furstenberg and E. Glasner, [Stationary dynamical systems](#). In *Dynamical numbers—interplay between dynamical systems and number theory*, pp. 1–28, Contemp. Math. 532, American Mathematical Society, Providence, RI, 2010 Zbl [1218.22004](#) MR [2762131](#)
- [10] E. Glasner, [Ergodic theory via joinings](#). Math. Surveys Monogr. 101, American Mathematical Society, Providence, RI, 2003 Zbl [1038.37002](#) MR [1958753](#)
- [11] E. Glasner and B. Weiss, [Uniformly recurrent subgroups](#). In *Recent trends in ergodic theory and dynamical systems*, pp. 63–75, Contemp. Math. 631, American Mathematical Society, Providence, RI, 2015 Zbl [1332.37014](#) MR [3330338](#)
- [12] P. Hall and C. R. Kulatilaka, [A property of locally finite groups](#). *J. Lond. Math. Soc.* **39** (1964), 235–239 Zbl [0136.27903](#) MR [161907](#)
- [13] R. E. Howe and C. C. Moore, [Asymptotic properties of unitary representations](#). *J. Funct. Anal.* **32** (1979), no. 1, 72–96 Zbl [0404.22015](#) MR [533220](#)
- [14] S. Kakutani, [Random ergodic theorems and Markoff processes with a stable distribution](#). In *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability, 1950*, pp. 247–261, University California Press, Berkeley, 1951 Zbl [0044.33901](#) MR [44773](#)
- [15] G. A. Margulis, [Finiteness of quotient groups of discrete subgroups](#). *Funct. Anal. Appl.* **13** (1979), no. 3, 178–187 Zbl [0434.22013](#) MR [545365](#)
- [16] G. A. Margulis, [Discrete subgroups of semisimple Lie groups](#). *Ergeb. Math. Grenzgeb.* (3) **17**, Springer, Berlin, 1991 Zbl [0732.22008](#) MR [1090825](#)
- [17] A. Nevo and R. J. Zimmer, [Homogenous projective factors for actions of semi-simple Lie groups](#). *Invent. Math.* **138** (1999), no. 2, 229–252 Zbl [0936.22007](#) MR [1720183](#)
- [18] A. Nevo and R. J. Zimmer, [A generalization of the intermediate factors theorem](#). *J. Anal. Math.* **86** (2002), 93–104 Zbl [1015.22002](#) MR [1894478](#)
- [19] J. Peterson, [Character rigidity for lattices in higher-rank groups](#). 2014, <https://math.vanderbilt.edu/peters10/rigidity.pdf>, visited on 29 February 2024
- [20] G. Stuck and R. J. Zimmer, [Stabilizers for ergodic actions of higher rank semisimple groups](#). *Ann. of Math. (2)* **139** (1994), no. 3, 723–747 Zbl [0836.22018](#) MR [1283875](#)
- [21] A. M. Vershik, [Nonfree actions of countable groups and their characters](#). *J. Math. Sci. (N.Y.)* **174** (2011), no. 1, 1–6 Zbl [1279.37004](#) MR [2749291](#)
- [22] R. J. Zimmer, [Hyperfiniteness and amenable ergodic actions](#). *Invent. Math.* **41** (1977), no. 1, 23–31 Zbl [0361.46061](#) MR [470692](#)
- [23] R. J. Zimmer, [Ergodic theory and semisimple groups](#). Monogr. Math. 81, Birkhäuser, Basel, 1984 Zbl [0571.58015](#) MR [776417](#)

Received 20 October 2021.

Darren Creutz

Mathematics Department, United States Naval Academy, 572C Holloway Road, Annapolis, MD 21402, USA; creutz@usna.edu