

# Sobolev algebras on Lie groups and noncommutative geometry

Cédric Arhancet

**Abstract.** We show that there exists a quantum compact metric space which underlies the setting of each Sobolev algebra associated to a subelliptic Laplacian  $\Delta = -(X_1^2 + \cdots + X_m^2)$  on a compact connected Lie group  $G$  if  $p$  is large enough, more precisely under the (sharp) condition  $p > \frac{d}{\alpha}$ , where  $d$  is the local dimension of  $(G, X)$  and where  $0 < \alpha \leq 1$ . We also provide locally compact variants of this result and generalizations for real second-order subelliptic operators. We also introduce a compact spectral triple (= noncommutative manifold) canonically associated to each subelliptic Laplacian on a compact group. In addition, we show that its spectral dimension is equal to the local dimension of  $(G, X)$ . Finally, we prove that the Connes spectral pseudo-metric allows us to recover the Carnot–Carathéodory distance.

## 1. Introduction

Suppose that  $1 < p < \infty$ . If  $\Delta_p: \text{dom } \Delta_p \subset L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$  is the (positive) Laplacian and if  $\alpha \in \mathbb{R}$ , we can consider the fractional powers  $\Delta_p^{\frac{\alpha}{2}}$  and  $(\text{Id} + \Delta_p)^{\frac{\alpha}{2}}$ . If  $\alpha < 0$ , these operators are the Riesz potential and the Bessel potential of order  $-\alpha$ . The last one was independently introduced by Aronszajn and Smith [9] and Calderón [17], which is nowadays a classical notion in harmonic analysis, see [78, p. 131] and [44, Definition 1.2.4, p. 13].

Strichartz proved in [79, Theorem 2.1, Chapter 2] that the Bessel potential space

$$L_\alpha^p(\mathbb{R}^n) \stackrel{\text{def}}{=} \{f \in L^p(\mathbb{R}^n) : \text{there exists } g \in L^p(\mathbb{R}^n) \text{ such that } f = (\text{Id} + \Delta_p)^{-\frac{\alpha}{2}} g\}$$

is an algebra for the pointwise product for any  $1 < p < \infty$  and any  $\alpha > 0$  such that  $\alpha p > n$ . Note that by [64, Theorem 12.3.4, p. 301], we have

$$L_\alpha^p(\mathbb{R}^n) = \text{dom } \Delta_p^{\frac{\alpha}{2}}.$$

Indeed, for any  $1 < p < \infty$  and any  $\alpha > 0$ , Kato and Ponce showed in their work [54, Lemma X.4, p. 906] on Navier–Stokes equations that

$$L_\alpha^p(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$$

---

2020 *Mathematics Subject Classification.* Primary 58B34; Secondary 46L87.

*Keywords.* Sobolev algebra, Carnot–Carathéodory distance, subelliptic Laplacian on Lie groups, quantum locally compact metric space, spectral triples, spectral dimension.

is an algebra for the pointwise product (see also [47, Theorem 2.2.12, p. 81]). This is a stronger result since by the Sobolev embedding theorem [2, Theorem 1.2.4 (c), p. 14] we have a continuous inclusion

$$L^p_\alpha(\mathbb{R}^n) \subset L^\infty(\mathbb{R}^n) \quad \text{if } \alpha p > n.$$

The proof is a simple consequence of the inequality

$$\|fg\|_{L^p_\alpha(\mathbb{R}^n)} \lesssim_{\alpha,p} \|f\|_{L^p_\alpha(\mathbb{R}^n)} \|g\|_{L^\infty(\mathbb{R}^n)} + \|f\|_{L^\infty(\mathbb{R}^n)} \|g\|_{L^p_\alpha(\mathbb{R}^n)} \tag{1.1}$$

for any  $f, g \in L^p_\alpha(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ , where we use the graph norm of the closed operator  $\Delta^{\frac{\alpha}{2}}_p$

$$\|f\|_{L^p_\alpha(\mathbb{R}^n)} \stackrel{\text{def}}{=} \|f\|_{L^p(\mathbb{R}^n)} + \|\Delta^{\frac{\alpha}{2}}_p(f)\|_{L^p(\mathbb{R}^n)} \approx \|(\text{Id} + \Delta_p)^{\frac{\alpha}{2}}(f)\|_{L^p(\mathbb{R}^n)}.$$

The motivation of this result was the estimate of

$$\|(\text{Id} + \Delta_p)^{\frac{\alpha}{2}}(fg) - f(\text{Id} + \Delta_p)^{\frac{\alpha}{2}}(g)\|_{L^p(\mathbb{R}^n)}$$

for any Schwartz functions  $f$  and  $g$ . This commutator estimate is needed in the study of some nonlinear partial differential equations. We refer to [61] and references therein for a comprehensive view of the state of the art of this kind of inequalities and to [64, Section 12.3] for several equivalent definitions of the Banach space  $L^p_\alpha(\mathbb{R}^n)$ .

In 1996, in their study of Schrödinger semigroups, Gulisashvili and Kon considered in [46] the homogeneous Sobolev space  $\dot{L}^p_\alpha(\mathbb{R}^n)$ , which is the completion of the space  $\text{dom } \Delta^{\frac{\alpha}{2}}_p$  with respect to the norm

$$\|f\|_{\dot{L}^p_\alpha(\mathbb{R}^n)} \stackrel{\text{def}}{=} \|\Delta^{\frac{\alpha}{2}}_p(f)\|_{L^p(\mathbb{R}^n)}.$$

Note that there exist several definitions of this abstract space. We refer to [66] for more information. In this paper, we only use functions of this space belonging to  $\text{dom } \Delta^{\frac{\alpha}{2}}_p$ . If  $\alpha > 0$ , Gulisashvili and Kon observed that  $\dot{L}^p_\alpha(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  is also an algebra for the pointwise product. This result is again a consequence of the Leibniz’s rule [46, Theorem 1.4]

$$\|fg\|_{\dot{L}^p_\alpha(\mathbb{R}^n)} \lesssim_{\alpha,p} \|f\|_{\dot{L}^p_\alpha(\mathbb{R}^n)} \|g\|_{L^\infty(\mathbb{R}^n)} + \|f\|_{L^\infty(\mathbb{R}^n)} \|g\|_{\dot{L}^p_\alpha(\mathbb{R}^n)}. \tag{1.2}$$

Coulhon, Russ, and Tardivel–Nachef [27] extended this result to the case of a unimodular connected Lie group  $G$  with polynomial volume growth by replacing the Laplacian by a subelliptic Laplacian  $-X^2_1 - \dots - X^2_m$ , where  $X \stackrel{\text{def}}{=} (X_1, \dots, X_m)$  is a family of left-invariant Hörmander vector fields. Replacing  $\mathbb{R}^n$  by  $G$ , they obtained generalizations

$$\|fg\|_{L^p_\alpha(G)} \lesssim_{\alpha,p} \|f\|_{L^p_\alpha(G)} \|g\|_{L^\infty(G)} + \|f\|_{L^\infty(G)} \|g\|_{L^p_\alpha(G)}, \quad \alpha > 0 \tag{1.3}$$

for any  $f, g \in L^p_\alpha(G) \cap L^\infty(G)$  and

$$\|fg\|_{\dot{L}^p_\alpha(G)} \lesssim_{\alpha,p} \|f\|_{\dot{L}^p_\alpha(G)} \|g\|_{L^\infty(G)} + \|f\|_{L^\infty(G)} \|g\|_{\dot{L}^p_\alpha(G)}, \quad 0 < \alpha \leq 1 \tag{1.4}$$

( $\alpha > 0$  if  $G$  is nilpotent) of the Leibniz’s rules (1.1) and (1.2). Furthermore, they obtained algebras  $L_\alpha^p(G) \cap L^\infty(G)$  and  $\dot{L}_\alpha^p(G) \cap L^\infty(G)$  called Sobolev algebras. Note that Bohnke has previously proved in [13, Theorem 1] that  $L_\alpha^p(G)$  is an algebra for the pointwise product if  $G$  is a stratified Lie group and if  $\alpha p > d$  where  $d$  is the local dimension of the group. With the help of Sobolev embedding theorem  $L_\alpha^p(G) \subset L^\infty(G)$  under the condition  $\alpha p > d$ , we can see that this is a particular case of the results of [27]. See [16, 72] for the more complicated case of non-unimodular Lie groups.

The concept of quantum compact metric space has its origins in Connes’ paper [22] in 1989, in which he showed that we can recover the geodesic distance  $\text{dist}$  of a compact oriented Riemannian spin manifold  $M$  using the Dirac operator  $\mathcal{D}$  by the formula

$$\text{dist}(x, y) = \sup_{f \in C(M), \|\llbracket \mathcal{D}, f \rrbracket\| \leq 1} |f(x) - f(y)|, \quad x, y \in M, \tag{1.5}$$

where the supremum is taken on all the continuous functions such that the commutator

$$\llbracket \mathcal{D}, f \rrbracket \stackrel{\text{def}}{=} \mathcal{D}f - f\mathcal{D}$$

extends to a contractive operator. Recall that  $\mathcal{D}$  is an unbounded operator acting on the Hilbert space of  $L^2$ -spinors and that the functions of  $C(M)$  act on the same Hilbert space by multiplication operators. Indeed, it is well known that the commutator  $\llbracket \mathcal{D}, f \rrbracket$  induces a bounded operator if and only if  $f$  is a Lipschitz function, and in this case, the Lipschitz constant of  $f$  is equal to the norm  $\|\llbracket \mathcal{D}, f \rrbracket\|$ . Moreover, this space of functions is norm dense in the space  $C(M)$  of continuous functions. See [23, Chapter 6] for more information, and we refer to [86] for a complete proof. If we identify the points  $x, y$  as pure states  $\omega_x$  and  $\omega_y$  on the unital  $C^*$ -algebra  $C(M)$ , we can see this formula as

$$\text{dist}(\omega_x, \omega_y) = \sup_{f \in C(M), \|\llbracket \mathcal{D}, f \rrbracket\| \leq 1} |\omega_x(f) - \omega_y(f)|, \quad x, y \in M.$$

After many years, Rieffel [75] axiomatized this formula replacing the algebra  $C(M)$  by a unital  $C^*$ -algebra  $A$ ,  $f \mapsto \|\llbracket \mathcal{D}, f \rrbracket\|$  by a seminorm  $\|\cdot\|$  defined on a dense subspace of  $A$  and  $\omega_x, \omega_y$  by arbitrary states of  $A$  obtaining essentially the formula (4.2) below and giving rise to a theory of *quantum* compact metric spaces. With this notion, Rieffel was able to define a quantum analogue of Gromov–Hausdorff distance and to give a meaning to many approximations found in the physics literature, as the case of matrix algebras converging to a sphere. Moreover, as research in noncommutative metric geometry progressed, some additional conditions are often added as Leibniz’s rules

$$\|ab\| \lesssim \|a\| \|b\|_A + \|a\|_A \|b\|, \quad a, b \in \text{dom}\|\cdot\|,$$

which look like (1.1) or (1.2).

If  $G$  is a compact connected Lie group  $G$  and if  $\alpha p > d$ , we show that

$$(C(G), \|\cdot\|_{L_\alpha^p(G)})$$

is a quantum compact metric space which underlies the setting of each Leibniz’s rule (1.4) associated to a subelliptic Laplacian  $\Delta$  on a compact connected Lie group  $G$ . Here,  $C(G)$  is the algebra of continuous function on  $G$ . We also provide locally compact variants for non-compact groups in Section 5 with the help of seminorms  $\|\cdot\|_{L^p_\alpha(G)}$ . Finally, our approach is flexible and should be adaptable to different contexts. The present work gets its inspiration from the papers [8, 50, 74].

Spectral triples are generalizations of the setting of Hodge–Dirac operators and Dirac operators on compact oriented Riemannian (spin) manifolds. This notion has emerged as a mean to encode geometric information of spaces in operator and spectral theory. It is at the heart of noncommutative geometry and used to describe quantum spaces, providing efficient tools for such an analysis. Remarkably, this notion also provides a framework for the study of classical spaces as fractals (e.g., [21]) or orbifolds. We refer to [24] for an extensive list of examples.

We introduce a spectral triple associated to each subelliptic Laplacian on a compact connected Lie group  $G$  and we show that the spectral dimension is equal to the local dimension of  $(G, X)$ , where  $X \stackrel{\text{def}}{=} (X_1, \dots, X_m)$  is the family of left-invariant Hörmander vector fields which defines the subelliptic Laplacian (3.9). In retrospect, our proof of this computation is quite simple. However, a variant shows a link between what we call the *local Couhlon–Varopoulos dimension* of a suitable (symmetric) Markovian semigroup  $(T_t)_{t \geq 0}$  acting on  $L^\infty(\Omega)$ , where  $\Omega$  is a finite measure space, and the spectral dimension of the spectral triple defined by an associated canonical Hodge–Dirac operator. We will investigate this more general setting in a future publication [7], also providing generalizations to suitable Markovian semigroups acting on von Neumann algebras.

Recall that this dimension is defined as the infimum of positive real numbers  $d$  such that

$$\|T_t\|_{L^1(\Omega) \rightarrow L^\infty(\Omega)} \lesssim \frac{1}{t^{\frac{d}{2}}}, \quad 0 < t \leq 1; \tag{1.6}$$

see [25] and [28, p. 187] (see also [19] for a related work). Note that the terminology *ultracontractivity* is equally used in [5, Section 7.3.2] and in [6]. This notion is also referred to in [89] under the more suitable term *local ultracontractivity*. We warn the reader that really different definitions of ultracontractivity coexist in the literature; see, e.g., [26, 29], [31, p. 89], and [45]. In the case of a connected Lie group  $G$  equipped with a family  $X$  of left-invariant Hörmander vector fields, the inequality (1.6) is satisfied for the heat semigroup whose generator is the opposite  $-\Delta$  of the subelliptic Laplacian and the local dimension  $d$  of  $(G, X)$ .

**Structure of the paper.** The paper is organized as follows. Section 2 gives background on operator theory. The aim of Section 3 is to describe our setting related to Lie groups and to prove some preliminary useful results. In Section 4, we show the existence of our quantum compact metric spaces. Section 5 is devoted to give *locally* compact variants of these quantum compact metric spaces. In Section 6, we introduce new compact spectral triples and we describe some properties. In particular, we compute the spectral dimension.

In Section 7, we investigate the links between the Connes spectral pseudo-distance and the Carnot–Carathéodory distance but also with the intrinsic pseudo-distance associated to some Dirichlet form. We prove an analog of formula (1.5) for the Carnot–Carathéodory distance. Finally, we state in Section 8 two natural conjectures on the functional calculus of subelliptic Laplacians and their associated Hodge–Dirac operators.

## 2. Preliminaries on operator theory

**Minkowski’s inequality.** Suppose that  $(\Omega_1, \mu_1)$  and  $(\Omega_2, \mu_2)$  are two  $\sigma$ -finite measure spaces and consider a measurable function  $f: \Omega_1 \times \Omega_2 \rightarrow \mathbb{C}$ . We will use the following classical inequality [78, Section A.1, p. 271]:

$$\begin{aligned} & \left[ \int_{\Omega_2} \left| \int_{\Omega_1} f(x, y) \, d\mu_1(x) \right|^p \, d\mu_2(y) \right]^{\frac{1}{p}} \\ & \leq \int_{\Omega_1} \left( \int_{\Omega_2} |f(x, y)|^p \, d\mu_2(y) \right)^{\frac{1}{p}} \, d\mu_1(x). \end{aligned} \tag{2.1}$$

**Dunford–Pettis theorem.** Let  $\Omega$  be a  $\sigma$ -finite measure space such that  $L^1(\Omega)$  is separable. A particular case of Dunford–Pettis theorem, e.g., [76, p. 528] and [45, Section 3], says that if  $T: L^1(\Omega) \rightarrow L^\infty(\Omega)$  is a bounded operator, then there exists a function  $K \in L^\infty(\Omega \times \Omega)$  such that for any  $f \in L^1(\Omega)$ , we have

$$(Tf)(x) = \int_{\Omega} K(x, y) f(y) \, dy$$

for almost all  $x \in \Omega$ . Moreover, we have

$$\|T\|_{L^1(\Omega) \rightarrow L^\infty(\Omega)} = \|K\|_{L^\infty(\Omega \times \Omega)}. \tag{2.2}$$

Conversely, such a function  $K$  defines a bounded operator  $T: L^1(\Omega) \rightarrow L^\infty(\Omega)$ . We also have a similar result for a bounded operator  $T: L^2(\Omega) \rightarrow L^\infty(\Omega)$ . In this case, the equality (2.2) is replaced by

$$\|T\|_{L^2(\Omega) \rightarrow L^\infty(\Omega)} = \|K\|_{L^\infty(\Omega, L^2(\Omega))}. \tag{2.3}$$

**Operator theory.** Recall the characterization of the domain of the closure  $\overline{T}$  of a closable unbounded operator  $T: \text{dom } T \subset Y \rightarrow Z$  between Banach spaces  $Y$  and  $Z$ . We have

$$\begin{aligned} x \in \text{dom } \overline{T} \text{ iff there exists } (x_n) \subset \text{dom } T \text{ such that } x_n \rightarrow x \\ \text{and } T(x_n) \rightarrow y \text{ for some } y. \end{aligned} \tag{2.4}$$

The following is [84, Corollary 5.6, p. 144].

**Theorem 2.1.** *Let  $T$  be a closed densely defined operator on a Hilbert space  $H$ . Then, the operator  $T^*T$  on  $(\text{Ker } T)^\perp$  is unitarily equivalent to the operator  $TT^*$  on  $(\text{Ker } T^*)^\perp$ .*

If  $T$  is densely defined, by [53, Problem 5.27, p. 168], we have

$$\text{Ker } T^* = (\text{Ran } T)^\perp. \tag{2.5}$$

We will also use the following classical equalities [51, Exercise 2.8.45, p. 171]:

$$\text{Ran } T^*T = \text{Ran } T^* \quad \text{and} \quad \text{Ker } T^*T = \text{Ker } T. \tag{2.6}$$

If  $A$  is sectorial operator on a reflexive Banach space  $Y$ , we have by [48, Proposition 2.1.1 (h)] a decomposition

$$Y = \text{Ker } A \oplus \overline{\text{Ran } A}.$$

**Fractional powers.** See [48, 56] for more information on fractional powers. Let  $A$  be a sectorial operator on a Banach space  $Y$ . If  $A$  is densely defined and if  $\alpha$  is a complex number with  $0 < \text{Re } \alpha < n$ , where  $n$  is an integer, then the space  $\text{dom } A^n$  is a core of  $A^\alpha$  by [48, p. 62], i.e.,  $\text{dom } A^n$  is dense in  $\text{dom } A^\alpha$  for the graph norm of  $A^\alpha$ , and we have

$$A^\alpha(x) = \frac{\Gamma(n)}{\Gamma(\alpha)\Gamma(n-\alpha)} \int_0^\infty t^{\alpha-1} (A(t+A)^{-1})^n x \, dt, \quad x \in \text{dom } A^n. \tag{2.7}$$

For any complex numbers  $\alpha, \beta$  with  $\text{Re } \alpha, \text{Re } \beta > 0$ , we have  $A^\alpha A^\beta = A^{\alpha+\beta}$ . By [48, p. 62] and [48, Corollary 3.1.11], for any  $\alpha \in \mathbb{C}$  with  $\text{Re } \alpha > 0$ , we have

$$\text{Ran } A^\alpha \subset \overline{\text{Ran } A} \quad \text{and} \quad \text{Ker } A^\alpha = \text{Ker } A. \tag{2.8}$$

If  $A$  is a sectorial operator on a Banach space  $Y$  and if  $\text{Re } \alpha > 0$ , then by [37, p. 137] the graph norms of the operators  $A^\alpha$  and  $(\text{Id} + A)^\alpha$  are equivalent; i.e.,

$$\|A^\alpha x\|_Y + \|x\|_Y \approx \|(\text{Id} + A)^\alpha x\|_Y, \quad x \in \text{dom } A^\alpha. \tag{2.9}$$

The proof uses [68, p. 28], the equality  $\text{dom}(\text{Id} + A)^\alpha = \text{dom } A^\alpha$  of [48, Proposition 3.1.9, p. 65], and the boundedness of the operator  $(\text{Id} + A)^{-\alpha}$ . See [56, Lemma 15.22, p. 294] and [48, Lemma 6.3.2, p. 148] for the particular case where  $A$  is injective.

**Compactness of fractional powers.** Let  $\Omega$  be a finite measure space. Consider a weak\* continuous semigroup  $(T_t)_{t \geq 0}$  of selfadjoint positive unital contractions on  $L^\infty(\Omega)$  with weak\* (negative) generator  $A_\infty$ . A classical argument shows that each operator  $T_t$  is integral preserving. Such a semigroup induces a strongly continuous semigroup  $(T_{t,p})_{t \geq 0}$  on  $L^p(\Omega)$ , and its generator  $A_p$  is sectorial if  $1 < p < \infty$ .

There exists a weak\* continuous conditional expectation  $\mathbb{E}: L^\infty(\Omega) \rightarrow L^\infty(\Omega)$  on the fixed subalgebra  $\{f \in L^\infty(\Omega) : T_t(f) = f \text{ for all } t \geq 0\}$ . This subset is equal to  $\text{Ker } A_\infty$ . We sketch the argument. By [39, Proposition 3.1.4, p. 120], the induced semigroup  $(T_{t,1})_{t \geq 0}$  on the space  $L^1(\Omega)$  is mean ergodic. In particular, we have a bounded

projection  $Q: L^1(\Omega) \rightarrow L^1(\Omega)$  onto  $\text{Ker } A_1$  along  $\overline{\text{Ran } A_1}$ , which is clearly contractive, satisfying  $Q(1) = 1$ . By [1, Corollary 5.52, p. 222],  $Q$  is a conditional expectation. We conclude by duality that the suitable conditional expectation  $\mathbb{E}$  exists see [38, Exercise 9, p. 159].

If  $\{f \in L^\infty(\Omega) : T_t(f) = f \text{ for all } t \geq 0\} = \mathbb{C}1$ , the conditional expectation is given by

$$\mathbb{E}(f) = \left( \int_{\Omega} f \right) 1.$$

We use the notation  $L_0^p(\Omega)$  for the subspace  $\text{Ker } \mathbb{E}_p$  of  $L^p(\Omega)$ . It is the space of functions with mean 0. We have

$$L_0^p(\Omega) = \overline{\text{Ran } A_p}.$$

Finally, for  $1 \leq p \leq q \leq \infty$ , consider the property

$$\|T_t\|_{L_0^p(\Omega) \rightarrow L_0^q(\Omega)} \lesssim \frac{1}{t^{\frac{n}{2}(\frac{1}{p} - \frac{1}{q})}}, \quad 0 < t \leq 1, \tag{2.10}$$

which is a *local* version of the property [50,  $(R_n^{pq})$ , p. 619]

$$\|T_t\|_{L_0^p(\Omega) \rightarrow L_0^q(\Omega)} \lesssim \frac{1}{t^{\frac{n}{2}(\frac{1}{p} - \frac{1}{q})}}, \quad t > 0.$$

By an interpolation argument similar to the one of [50, Lemma 1.1.2], each of these properties holds for one pair  $1 \leq p < q \leq \infty$  if and only if it holds for all  $1 \leq p \leq q \leq \infty$ . See also [5, Section 7.3.2, p. 65] for a variant.

Recall the following result [30, Theorem 9] (see also [52, Theorem 5.5]) which allows one to obtain compactness via complex interpolation.

**Theorem 2.2.** *Suppose that  $(X_0, X_1)$  and  $(Y_0, Y_1)$  are Banach couples and that  $X_0$  is a UMD-space. Let  $T: X_0 + X_1 \rightarrow Y_0 + Y_1$  such that its restriction  $T_0: X_0 \rightarrow Y_0$  is compact and such that  $T_1: X_1 \rightarrow Y_1$  is bounded. Then, for any  $0 < \theta < 1$ , the map  $T: (X_0, X_1)_\theta \rightarrow (Y_0, Y_1)_\theta$  is compact.*

The following is [50, Theorem 1.1.7]. Note that the proof of this result uses [50, Lemma 1.1.6] whose proof unfortunately seems false in light of the classical problem [52, Problem 5.4]. However, [50, Lemma 1.1.6] can be replaced by Theorem 2.2.

**Proposition 2.3.** *Let  $\Omega$  be a finite measure space. Let  $(T_t)_{t \geq 0}$  be a weak\* continuous semigroup of selfadjoint positive contractions on  $L^\infty(\Omega)$  with  $\{x \in L^\infty(\Omega) : T_t(x) = x \text{ for all } t \geq 0\} = \mathbb{C}1$ , satisfying*

$$\|T_t\|_{L_0^1(\Omega) \rightarrow L^\infty(\Omega)} \lesssim \frac{1}{t^{\frac{n}{2}}}$$

for some  $n$  and such that  $A^{-1}$  is compact on  $L_0^2(\Omega)$ . Then, for all  $1 \leq p < q \leq \infty$  such that  $\frac{2\text{Re } z}{n} > \frac{1}{p} - \frac{1}{q}$ , the operator

$$A^{-z}: L_0^p(\Omega) \rightarrow L_0^q(\Omega)$$

is compact.

### 3. Background and preliminaries results on Lie groups

**Convolution.** If  $G$  is a *unimodular* locally compact group equipped with a Haar measure  $\mu_G$ , recall that the convolution product of two functions  $f$  and  $g$  is given, when it exists, by

$$(f * g)(s) \stackrel{\text{def}}{=} \int_G f(r)g(r^{-1}s) d\mu_G(r) = \int_G f(sr^{-1})g(r) d\mu_G(r). \tag{3.1}$$

**Carnot–Carathéodory distances on connected Lie groups.** Let  $G$  be a connected Lie group with neutral element  $e$ . We consider a finite sequence  $X \stackrel{\text{def}}{=} (X_1, \dots, X_m)$  of left invariant vector fields which generate the Lie algebra  $\mathfrak{g}$  of the group  $G$  such that the vectors  $X_1(e), \dots, X_m(e)$  are linearly independent. We say that it is a family of left-invariant Hörmander vector fields.

Let  $\gamma: [c, d] \rightarrow G$  be an absolutely continuous path such that  $\dot{\gamma}(t)$  belongs to the subspace  $\text{span}\{X_1|_{\gamma(t)}, \dots, X_m|_{\gamma(t)}\}$  for almost all  $t \in [c, d]$ . If  $\dot{\gamma}(t) = \sum_{k=1}^m \dot{\gamma}_k(t) X_k|_{\gamma(t)}$  for almost all  $t \in [c, d]$ , where  $\dot{\gamma}_k(t) \in \mathbb{R}$  and where each  $\dot{\gamma}_k$  is measurable, we can define the  $p$ -length of  $\gamma$  by

$$\ell_p(\gamma) \stackrel{\text{def}}{=} \int_c^d \left( \sum_{k=1}^m |\dot{\gamma}_k(t)|^p \right)^{\frac{1}{p}} dt, \tag{3.2}$$

which belongs to  $[0, \infty]$ . For any  $s, s' \in G$ , there exists such a path  $\gamma: [0, 1] \rightarrow G$  with finite length with  $\gamma(0) = s$  and  $\gamma(1) = s'$ . If  $s, s' \in G$  and  $1 < p < \infty$ , then we define the real number  $\text{dist}_{\text{CC}}^p(s, s')$  between  $s$  and  $s'$  to be the infimum of the length of all such paths with  $\gamma(0) = s$  and  $\gamma(1) = s'$ :

$$\text{dist}_{\text{CC}}^p(s, s') \stackrel{\text{def}}{=} \inf_{\gamma(0)=s, \gamma(1)=s'} \ell_p(\gamma). \tag{3.3}$$

See [87, p. 39] and [34, p. 22] if  $p = 2$ . In this case, we recover the Carnot–Carathéodory distance  $\text{dist}_{\text{CC}}(s, s')$ . We refer also to [67].

If  $f: G \rightarrow \mathbb{C}$  is a smooth function and if  $\gamma: [0, 1] \rightarrow G$  is an absolutely continuous path with tangents in the subspace spanned by  $X_1, \dots, X_m$ , then by [83, p. 64], we have

$$\frac{d}{dt} f(\gamma(t)) = \sum_{k=1}^m \dot{\gamma}_k(t) (X_k f)(\gamma(t)) \quad \text{a.e. } t \in [0, 1], \tag{3.4}$$

i.e.,

$$\frac{d}{dt} f(\gamma(t)) = \langle \dot{\gamma}(t), (Xf)(\gamma(t)) \rangle \quad \text{a.e. } t \in [0, 1].$$

We will need the following elementary inequality. In the following statement, each  $\text{dom } X_{k,p}$  is the domain of  $X_k$  on  $L^p(G)$ ; i.e.,

$$X_{k,p}: \text{dom } X_{k,p} \subset L^p(G) \rightarrow L^p(G).$$



Recall that connected Lie groups are  $\sigma$ -finite under Haar measure. Let  $\lambda: G \rightarrow B(L^p(G))$ ,  $s \mapsto (f \mapsto f(s^{-1}\cdot))$  be the left regular representation of  $G$ .

The following result is a variant of [87, Lemma VIII.1.1, p. 106].

**Lemma 3.1.** *Suppose that  $1 < p < \infty$  and  $\frac{1}{p} + \frac{1}{p^*} = 1$ . Then, for any  $s \in G$  and any  $f$  belonging to  $\text{dom } X_{1,p} \cap \dots \cap \text{dom } X_{m,p}$ , we have*

$$\|(\text{Id} - \lambda_s)f\|_{L^p(G)} \leq \text{dist}_{\text{CC}}^{p^*}(s, e) \left( \sum_{k=1}^m \|X_{k,p}f\|_{L^p(G)}^p \right)^{\frac{1}{p}}. \tag{3.5}$$

*Proof.* Let  $f \in C_c^\infty(G)$ . Let  $s \in G$ , and let  $\gamma: [0, 1] \mapsto G$  be an absolutely continuous path from  $e$  to  $s^{-1}$ . For any  $s' \in G$ , we have

$$\begin{aligned} ((\text{Id} - \lambda_s)f)(s') &= f(s') - (\lambda_s f)(s') = f(s') - f(s^{-1}s') \\ &= - \int_0^1 \frac{d}{dt} f(\gamma(t)s') dt \stackrel{(3.4)}{=} - \int_0^1 \sum_{k=1}^m \dot{\gamma}_k(t) (X_k f)(\gamma(t)s') dt. \end{aligned}$$

Consequently, using Hölder’s inequality, we obtain

$$|((\text{Id} - \lambda_s)f)(s')| \leq \int_0^1 \left( \sum_{k=1}^m \dot{\gamma}_k(t)^p \right)^{\frac{1}{p}} \left( \sum_{k=1}^m [(X_k f)(\gamma(t)s')]^{p^*} \right)^{\frac{1}{p^*}} dt. \tag{3.6}$$

Using Minkowski’s inequality (2.1) and left invariance in the last equality, we deduce that

$$\begin{aligned} \|(\text{Id} - \lambda_s)f\|_{p^*} &= \left[ \int_G |((\text{Id} - \lambda_s)f)(s')|^{p^*} d\mu_G(s') \right]^{\frac{1}{p^*}} \\ &\stackrel{(3.6)}{\leq} \left[ \int_G \left( \int_0^1 \left( \sum_{k=1}^m \dot{\gamma}_k(t)^p \right)^{\frac{1}{p}} \left( \sum_{k=1}^m [(X_k f)(\gamma(t)s')]^{p^*} \right)^{\frac{1}{p^*}} dt \right)^{p^*} d\mu_G(s') \right]^{\frac{1}{p^*}} \\ &\stackrel{(2.1)}{\leq} \int_0^1 \left[ \int_G \left( \sum_{k=1}^m \dot{\gamma}_k(t)^p \right)^{\frac{p^*}{p}} \left( \sum_{k=1}^m [(X_k f)(\gamma(t)s')]^{p^*} \right) d\mu_G(s') \right]^{\frac{1}{p^*}} dt \\ &= \int_0^1 \left( \sum_{k=1}^m \dot{\gamma}_k(t)^p \right)^{\frac{1}{p}} \left[ \sum_{k=1}^m \int_G [(X_k f)(\gamma(t)s')]^{p^*} d\mu_G(s') \right]^{\frac{1}{p^*}} dt \\ &= \int_0^1 \left( \sum_{k=1}^m \dot{\gamma}_k(t)^p \right)^{\frac{1}{p}} \left[ \sum_{k=1}^m \|X_k f\|_{L^{p^*}(G)}^{p^*} \right]^{\frac{1}{p^*}} dt. \end{aligned}$$

Hence, by taking the infimum over all possible paths and observing that

$$\text{dist}_{\text{CC}}^{p^*}(e, s^{-1}) = \text{dist}_{\text{CC}}^{p^*}(s, e),$$

we obtain (3.5) with (3.3). We conclude by using an approximation argument as in the proof of Proposition 3.4 for a general  $f$ . ■

**Growth of volume and dimensions.** Let  $G$  be a connected Lie group equipped with a family  $X \stackrel{\text{def}}{=} (X_1, \dots, X_m)$  of left-invariant Hörmander vector fields and a left Haar measure  $\mu_G$ . For any  $r > 0$  and any  $x \in G$ , we denote by  $B(x, r)$  the open ball with respect to the Carnot–Carathéodory metric centered at  $x$  and of radius  $r$ , and by  $V(r) \stackrel{\text{def}}{=} \mu_G(B(x, r))$  the Haar measure of any ball of radius  $r$ . It is well known, e.g., [87, p. 124], that there exist  $d \in \mathbb{N}^*$ ,  $c, C > 0$  such that

$$cr^d \leq V(r) \leq Cr^d, \quad r \in ]0, 1[. \tag{3.7}$$

The integer  $d$  is called the local dimension of  $(G, X)$ .

When  $r \geq 1$ , only two situations may occur, independently of the choice of  $X$  (see, e.g., [34, p. 26]): either  $G$  has polynomial volume growth, which means that there exist  $D \in \mathbb{N}$  and  $c', C' > 0$  such that

$$c'r^D \leq V(r) \leq C'r^D, \quad r \geq 1, \tag{3.8}$$

or  $G$  has exponential volume growth, which means that there exist  $c_1, C_1, c_2, C_2 > 0$  such that

$$c_1e^{c_2r} \leq V(r) \leq C_1e^{C_2r}, \quad r \geq 1.$$

When  $G$  has polynomial volume growth, the integer  $D$  in (3.8) is called the dimension at infinity of  $G$ . Note that, contrary to  $d$ , it only depends on  $G$  and not on  $X$ ; see [87, Chapter 4].

By [34, Proposition II.4.5, p. 26] or [76, p. 381], each connected Lie group of polynomial growth is unimodular. By [76, pp. 256–257] and [34, p. 26], a connected compact Lie group has polynomial volume growth with  $D = 0$ . Recall finally that connected nilpotent Lie groups have polynomial volume growth by [34, p. 28].

**Example 3.2.** The local dimension of the abelian compact Lie group  $\mathbb{T}^n$  is  $n$  by [76, p. 274], and its dimension at infinity is of 0 since  $\mathbb{T}^n$  is compact.

**Example 3.3.** The local dimension and the dimension at infinity of a stratified Lie group are equal by [34, Proposition II.4.15, p. 29]. The three-dimensional Heisenberg group  $\mathbb{H}_3$  (equipped with its canonical stratification) is a stratified group, and its dimensions are equal to 4 by [34, Example II.4.16].

Let  $G$  be a unimodular connected Lie group endowed with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields, and let  $\mu_G$  be a Haar measure. We consider the subelliptic Laplacian  $\Delta$  on  $G$  defined by

$$\Delta \stackrel{\text{def}}{=} - \sum_{k=1}^m X_k^2. \tag{3.9}$$

For  $1 \leq p < \infty$ , let  $\Delta_p: \text{dom } \Delta_p \subset L^p(G) \rightarrow L^p(G)$  be the smallest closed extension of the closable unbounded operator  $\Delta|_{C_c^\infty(G)}$  to  $L^p(G)$ . Note that the domain  $\text{dom } \Delta_p^{\frac{\alpha}{2}}$  is closed under the adjoint operation  $f \mapsto \bar{f}$ .

We denote by  $(T_t)_{t \geq 0}$  the associated weak\* continuous semigroup of selfadjoint unital (i.e.,  $T_t(1) = 1$ ) positive contractive operators on  $L^\infty(G)$ ; see [87, pp. 20–21], [34, p. 21], and [76, p. 301]. By [76, Proposition 4.13, p. 323] and [34, Proposition 11.3.1, p. 20], for any  $t > 0$ , the operator  $T_t: L^p(G) \rightarrow L^p(G)$  is a convolution operator by a positive function  $K_t$  of  $L^1(G)$ .

Suppose that  $1 < p < \infty$  and that the Lie group  $G$  has polynomial volume growth. By [3, Theorem 2] and [27, p. 339], for any  $f \in C_c^\infty(G)$ , we have

$$\|\Delta_p^{\frac{1}{2}}(f)\|_{L^p(G)} \approx_p \sum_{k=1}^m \|X_k(f)\|_{L^p(G)}. \tag{3.10}$$

Since  $\text{dom } \Delta_p$  is a core of  $\Delta_p^{\frac{1}{2}}$ , a classical argument [68, p. 29] reveals that the subspace  $C_c^\infty(G)$  is a core of the operator  $\Delta_p^{\frac{1}{2}}$ .

The following observation is a natural complement of the equivalences (3.10). Since it is always written in the literature without proof, we give an argument. Note that the subspace  $\text{dom } X_{1,p} \cap \dots \cap \text{dom } X_{m,p}$  is considered in the paper [10, p. 194] and in the book [34, p. 15] and respectively denoted by  $W'_{1,2}(G)$  and  $L'_{2,1}(G)$ .

**Proposition 3.4.** *Let  $G$  be a unimodular connected Lie group with polynomial volume growth. Suppose that  $1 < p < \infty$ . We have*

$$\text{dom } \Delta_p^{\frac{1}{2}} = \text{dom } X_{1,p} \cap \dots \cap \text{dom } X_{m,p}.$$

Moreover, for any  $f \in \text{dom } \Delta_p^{\frac{1}{2}}$ , we have (3.10).

*Proof.* Let  $f \in \text{dom } \Delta_p^{\frac{1}{2}}$ . The subspace  $C_c^\infty(G)$  is dense in  $\text{dom } \Delta_p^{\frac{1}{2}}$  equipped with the graph norm. Hence, we can find a sequence  $(f_n)$  of  $C_c^\infty(G)$  such that

$$f_n \rightarrow f \quad \text{and} \quad \Delta_p^{\frac{1}{2}}(f_n) \rightarrow \Delta_p^{\frac{1}{2}}(f) \quad \text{in } L^p(G).$$

For any integers  $n, l$  and any  $1 \leq k \leq m$ , we obtain

$$\begin{aligned} & \|f_n - f_l\|_{L^p(G)} + \|X_k(f_n) - X_k(f_l)\|_{L^p(G)} \\ & \stackrel{(3.10)}{\lesssim_p} \|f_n - f_l\|_{L^p(G)} + \|\Delta_p^{\frac{1}{2}}(f_n) - \Delta_p^{\frac{1}{2}}(f_l)\|_{L^p(G)}, \end{aligned}$$

which shows that  $(f_n)$  is a Cauchy sequence in  $\text{dom } X_{k,p}$ . By the closedness of  $X_{k,p}$  we infer that this sequence converges to some  $g \in \text{dom } X_{k,p}$  equipped with the graph norm. Since  $\text{dom } X_{k,p}$  equipped with the graph norm is continuously embedded into  $L^p(G)$ , we have  $f_n \rightarrow g$  in  $L^p(G)$ , and therefore,  $f = g$  since  $f_n \rightarrow f$ . It follows that  $f \in \text{dom } X_{k,p}$ . This proves the inclusion

$$\text{dom } \Delta_p^{\frac{1}{2}} \subset \text{dom } X_{k,p}.$$

Moreover, for any integer  $n$ , we have

$$\|X_k(f_n)\|_{L^p(G)} \stackrel{(3.10)}{\lesssim_p} \|\Delta_p^{\frac{1}{2}}(f_n)\|_{L^p(G)}.$$

Since  $f_n \rightarrow f$  in  $\text{dom } X_{k,p}$  and in  $\text{dom } \Delta_p^{\frac{1}{2}}$  both equipped with the graph norm, we conclude that

$$\|X_k(f)\|_{L^p(G)} \lesssim_p \|\Delta_p^{\frac{1}{2}}(f)\|_{L^p(G)}.$$

The proofs of the reverse inclusion and estimate are similar. ■

Suppose that  $1 \leq p < \infty$  and  $\alpha > 0$ . When  $f \in \text{dom } \Delta_p^{\frac{\alpha}{2}}$ , we let

$$\|f\|_{\dot{L}_\alpha^p(G)} \stackrel{\text{def}}{=} \|\Delta_p^{\frac{\alpha}{2}}(f)\|_{L^p(G)}. \tag{3.11}$$

It is related to Sobolev towers; see [40, Section II.5] and [56, Section 15.E]. Note that  $\|\cdot\|_{\dot{L}_\alpha^p(G)}$  is a seminorm on the subspace  $\dot{L}_\alpha^p(G)$  of  $L^p(G)$  (and even a norm if  $G$  is not compact). In this paper, we have no intention to define or use a Banach space  $\dot{L}_\alpha^p(G)$ .

Assume that the unimodular connected Lie group  $G$  has *polynomial volume growth*. For any  $\alpha \in [0, 1]$  and any  $p \in ]1, +\infty[$ , by [27, Theorem 3], the space  $\dot{L}_\alpha^p(G) \cap L^\infty(G)$  is an algebra under pointwise product. In [27, pp. 289–290], the authors give a simple proof of the case  $\alpha = 1$ . More precisely for all  $f, g \in \dot{L}_\alpha^p(G) \cap L^\infty(G)$ , we have  $fg \in \dot{L}_\alpha^p(G) \cap L^\infty(G)$  and (1.4). If  $G$  is in addition nilpotent, the conclusion holds for all  $\alpha \geq 0$ . See also [46, Theorem 1.4] for the particular case  $G = \mathbb{R}^n$  with some generalizations.

In the following statement, the seminorm  $\|\cdot\|_{\dot{L}_\alpha^p(G)}$  is defined on

$$\text{dom}\|\cdot\|_{\dot{L}_\alpha^p(G)} \stackrel{\text{def}}{=} C_0(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}}, \tag{3.12}$$

where  $C_0(G)$  is the Banach space of complex-valued continuous functions on  $G$  that vanish at infinity. Recall that  $C_0(G)$  is equipped with the restriction of the norm  $\|\cdot\|_{L^\infty(G)}$ . If the group  $G$  is compact, we have of course the equality  $C_0(G) = C(G)$ , where  $C(G)$  is the Banach space of complex-valued continuous functions on  $G$ .

**Lemma 3.5.** *Let  $G$  be a connected unimodular Lie group. Suppose that  $1 < p < \infty$  and  $\alpha > 0$ .*

- (1) *The  $\mathbb{C}$ -subspace  $\text{dom}\|\cdot\|_{\dot{L}_\alpha^p(G)}$  is dense in the Banach space  $C_0(G)$ .*
- (2) *The subspace  $\text{dom}\|\cdot\|_{\dot{L}_\alpha^p(G)}$  is closed under the adjoint operation  $f \mapsto \bar{f}$ .*
- (3) *If  $G$  is compact, we have*

$$\{f \in \text{dom}\|\cdot\|_{\dot{L}_\alpha^p(G)} : \|f\|_{\dot{L}_\alpha^p(G)} = 0\} = \mathbb{C}1_{C(G)}.$$

*If  $G$  has polynomial volume growth and is non-compact, we have*

$$\{f \in \text{dom}\|\cdot\|_{\dot{L}_\alpha^p(G)} : \|f\|_{\dot{L}_\alpha^p(G)} = 0\} = \{0\}.$$

- (4) *If  $G$  is compact, the seminorm  $\|\cdot\|_{\dot{L}_\alpha^p(G)}$  is lower semicontinuous.*

*Proof.* (1) Recall that the space  $\text{dom } \Delta_p^n$  is a core of the operator  $\Delta_p^{\frac{\alpha}{2}}$  if  $\frac{\alpha}{2} < n$ . Consequently, the domain

$$\text{dom} \|\cdot\|_{L^\alpha_p(G)} \stackrel{(3.12)}{=} C_0(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}}$$

contains the subspace  $C_c^\infty(G)$  of  $C_0(G)$ . Note that this subspace is dense in  $C_0(G)$  by regularization by [63, Theorem 2.11]. We infer that the  $\mathbb{C}$ -subspace  $\text{dom} \|\cdot\|_{L^\alpha_p(G)}$  is dense in the Banach space  $C_0(G)$ .

(2) Note that the space  $C_0(G)$  is obviously closed under the adjoint operation  $f \mapsto \bar{f}$ . We will show that  $\text{dom} \|\cdot\|_{L^\alpha_p(G)}$  is equally closed under the same operation.

Let  $f \in \text{dom } \Delta_p^{\frac{\alpha}{2}}$ . We know that the subspace  $\text{dom } \Delta_p^n$  is core of  $\Delta_p^{\frac{\alpha}{2}}$ . Hence, there exists a sequence  $(f_j)$  of  $\text{dom } \Delta_p^n$  such that  $f_j \rightarrow f$  and  $\Delta_p^{\frac{\alpha}{2}}(f_j) \rightarrow \Delta_p^{\frac{\alpha}{2}}(f)$ . We have

$$\overline{f_j} \rightarrow \bar{f} \quad \text{and} \quad \Delta_p^{\frac{\alpha}{2}}(\overline{f_j}) = \overline{\Delta_p^{\frac{\alpha}{2}}(f_j)} \rightarrow \overline{\Delta_p^{\frac{\alpha}{2}}(f)},$$

where the equality can be seen with (2.7). By (2.4), we conclude that  $\bar{f} \in \text{dom } \Delta_p^{\frac{\alpha}{2}}$  and that

$$\Delta_p^{\frac{\alpha}{2}}(\bar{f}) = \overline{\Delta_p^{\frac{\alpha}{2}}(f)}.$$

We conclude that  $\text{dom} \|\cdot\|_{L^\alpha_p(G)}$  is closed under the adjoint operation  $f \mapsto \bar{f}$ .

(3) We have

$$\Delta_p(1) \stackrel{(3.9)}{=} - \sum_{k=1}^m X_k^2(1) = 0.$$

Hence, the constant function 1 belongs to  $\text{Ker } \Delta_p \stackrel{(2.8)}{=} \text{Ker } \Delta_p^{\frac{\alpha}{2}}$ . We conclude that

$$\|1\|_{L^\alpha_p(G)} = \|\Delta_p^{\frac{\alpha}{2}}(1)\|_p = 0.$$

In the other direction, if  $\|f\|_{L^\alpha_p(G)} = 0$ , we have

$$\|\Delta_p^{\frac{\alpha}{2}}(f)\|_p = 0.$$

Hence,  $f$  belongs to  $\text{Ker } \Delta_p^{\frac{\alpha}{2}}$ . By (2.8), we deduce that  $\|\Delta_p^{\frac{1}{2}}(f)\|_p = 0$ . Then, according to Proposition 3.4 and (3.10), we have  $\|X_k(f)\|_p = 0$  for any  $k$ . By Lemma 3.1, we infer that  $\lambda_s(f) = f$  for any  $s \in G$ . If  $G$  is compact, we conclude that the function  $f$  is constant; that is,  $f \in \mathbb{C}1$ , and that  $f = 0$  if  $G$  is not compact.

(4) Let  $f \in C(G)$  and  $(f_n)$  be a sequence of elements of  $C(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}}$  such that  $(f_n)$  converges to  $f$  for the norm topology of  $C(G)$  and  $\|f_n\|_{L^\alpha_p(G)} \leq 1$  for any  $n$ ; that is,

$$\|\Delta_p^{\frac{\alpha}{2}}(f_n)\|_{L^p(G)} \leq 1$$

by (3.11). We have to prove that  $f$  belongs to  $\text{dom } \Delta_p^{\frac{\alpha}{2}}$  and that  $\|f\|_{L^\alpha_p(G)} \leq 1$ .

Since  $\|\cdot\|_{L^p(G)} \leq \|\cdot\|_{C(G)}$ , the sequence  $(f_n)$  converges to  $f$  for the norm topology of  $L^p(G)$ , hence for the weak topology of  $L^p(G)$ . Note that the sequence  $\Delta_p^{\frac{\alpha}{2}}(f_n)$  is bounded in the Banach space  $L^p(G)$ . Since bounded sets are weakly relatively compact by [65, Theorem 2.8.2], there exists a weakly convergent subnet  $(\Delta_p^{\frac{\alpha}{2}} f_{n_j})$ . Then,  $(f_{n_j}, \Delta_p^{\frac{\alpha}{2}} f_{n_j})$  is a weakly convergent net in the graph of the closed operator  $\Delta_p^{\frac{\alpha}{2}}$ . Note that this graph is closed and convex, hence weakly closed by [65, Theorem 2.5.16]. Thus, the limit of  $(f_{n_j}, \Delta_p^{\frac{\alpha}{2}} f_{n_j})$  belongs again to the graph and is of the form  $(g, \Delta_p^{\frac{\alpha}{2}} g)$  for some  $g \in \text{dom } \Delta_p^{\frac{\alpha}{2}}$ . In particular,  $(f_{n_j})$  converges weakly to  $g$  and  $\Delta_p^{\frac{\alpha}{2}}(f_{n_j})$  converges weakly to  $\Delta_p^{\frac{\alpha}{2}}(g)$ . We infer that  $f = g$ . We conclude that  $f$  belongs to  $\text{dom } \Delta_p^{\frac{\alpha}{2}}$ . Moreover, using the weakly lower semicontinuity of the norm [65, Theorem 2.5.21], we obtain

$$\|f\|_{L^p(G)} \stackrel{(3.11)}{=} \|\Delta_p^{\frac{\alpha}{2}}(f)\|_{L^p(G)} \leq \liminf_j \|\Delta_p^{\frac{\alpha}{2}}(f_{n_j})\|_{L^p(G)} \leq 1. \quad \blacksquare$$

### 4. Quantum compact metric spaces

**Lipschitz pairs and quantum compact metric spaces.** Following [59, Definition 2.3], a Lipschitz pair  $(A, \|\cdot\|)$  is a  $C^*$ -algebra  $A$  equipped with a seminorm  $\|\cdot\|$  defined on a dense subspace  $\text{dom}\|\cdot\|$  of the selfadjoint part  $(uA)_{\text{sa}}$  such that

$$\{a \in \text{dom}\|\cdot\| : \|a\| = 0\} = \mathbb{R}1_{uA}, \tag{4.1}$$

where  $uA$  is the unitization of the algebra  $A$ . If  $A$  is in addition unital, we say that  $(A, \|\cdot\|)$  is a unital Lipschitz pair.

Recall that a state of a  $C^*$ -algebra  $A$  is a positive linear form  $\varphi$  on  $A$  with  $\|\varphi\| = 1$ . If  $X$  is a compact topological space and if  $A = C(X)$ , a state is the integral associated to a regular Borel measure of probability on  $X$ .

A pair  $(A, \|\cdot\|)$  is a quantum compact metric space when

- (1)  $(A, \|\cdot\|)$  is a unital Lipschitz pair;
- (2) the Monge–Kantorovich metric on the set  $S(A)$  of the states of  $A$ , defined for any two states  $\varphi, \psi \in S(A)$  by

$$\text{dist}_{\text{MK}}(\varphi, \psi) \stackrel{\text{def}}{=} \sup\{|\varphi(a) - \psi(a)| : a \in \text{dom}\|\cdot\| \text{ and } \|a\| \leq 1\}, \tag{4.2}$$

induces the weak\* topology on  $S(A)$ .

In this case, we say that  $\|\cdot\|$  is a Lip-norm. We refer to the nice surveys [59, 75] and references therein for more information.

**Example 4.1.** If  $(X, \text{dist})$  is a compact metric space, a fundamental example [59, Example 2.6] and [58, Example 2.9] is given by  $(C(X), \text{Lip})$ , where  $C(X)$  is the commutative

$C^*$ -algebra of continuous functions on  $X$  and where  $\text{Lip}$  is the Lipschitz seminorm, defined for any Lipschitz function  $f: X \rightarrow \mathbb{C}$  by

$$\text{Lip}(f) \stackrel{\text{def}}{=} \sup \left\{ \frac{|f(x) - f(y)|}{\text{dist}(x, y)} : x, y \in X, x \neq y \right\}.$$

The set of real Lipschitz functions is norm-dense in  $C(X)_{\text{sa}}$  by the Stone–Weierstrass theorem. Indeed,  $\text{Lip}(X)$  contains the constant functions. Moreover,  $\text{Lip}(X)$  separates points in  $X$ . If  $x_0, y_0 \in X$  with  $x_0 \neq y_0$ , we can use the Lipschitz function  $f: X \rightarrow \mathbb{R}$ ,  $x \mapsto \text{dist}(x, y_0)$  since we have  $f(x_0) > 0 = f(y_0)$ . Moreover, it is immediate that a function  $f$  has zero Lipschitz constant if and only if it is constant on  $X$ ; i.e., (4.1) is satisfied.

In the case of  $(C(X), \text{Lip})$ , the equality (4.2) gives the dual formulation of the classical Kantorovich–Rubinstein metric [88, Remark 6.5] for Borel probability measures  $\mu$  and  $\nu$  on  $X$

$$\text{dist}(\mu, \nu) \stackrel{\text{def}}{=} \sup \left\{ \left| \int_X f \, d\mu - \int_X f \, d\nu \right| : f \in C(X)_{\text{sa}}, \text{Lip}(f) \leq 1 \right\}, \tag{4.3}$$

which is a basic concept in optimal transport theory [88]. Considering the Dirac measures  $\delta_x$  and  $\delta_y$  at points  $x, y \in X$  instead of  $\mu$  and  $\nu$ , we recover the distance  $\text{dist}(x, y)$  with the formula (4.3).

**Characterizations of quantum compact metric spaces.** The compatibility of Monge–Kantorovich metric with the weak\* topology is hard to check directly in general. Fortunately, there exists a condition which is more practical. This condition is inspired by the fact that Arzéla–Ascoli’s theorem shows that for any  $x \in X$  the set

$$\{f \in C(X)_{\text{sa}} : \text{Lip}(f) \leq 1, f(x) = 0\}$$

is norm relatively compact, and it is known that this property implies that (4.3) metrizes the weak\* topology on the space of Borel probability measures on  $X$ . Now, we give sufficient conditions in order to obtain quantum compact metric spaces [59, Theorem 2.43]. See also [69, Proposition 1.3].

**Proposition 4.2.** *Let  $(A, \|\cdot\|)$  be a unital Lipschitz pair. The following assertions are equivalent:*

- (a)  $(A, \|\cdot\|)$  is a quantum compact metric space;
- (b) there exists a state  $\mu \in S(A)$  such that the set

$$\{a \in A_{\text{sa}} : \|a\| \leq 1, \mu(a) = 0\}$$

is relatively compact in  $A$  for the topology of the norm of  $A$ ;

- (c) for all states  $\mu \in S(A)$ , the set

$$\{a \in A_{\text{sa}} : \|a\| \leq 1, \mu(a) = 0\}$$

is relatively compact in  $A$  for the topology of the norm of  $A$ .

**Quasi-Leibniz quantum compact metric space.** The Lipschitz seminorm  $\text{Lip}$  of Example 4.1 associated to a compact metric space  $(X, \text{dist})$  enjoys a natural property with respect to the multiplication of functions in  $C(X)$ , called the Leibniz property for any Lipschitz functions  $f, g: X \rightarrow \mathbb{C}$ :

$$\text{Lip}(fg) \leq \|f\|_{C(X)} \text{Lip}(g) + \text{Lip}(f)\|g\|_{C(X)}. \tag{4.4}$$

Moreover, the Lipschitz seminorm is lower-semicontinuous with respect to the norm of the algebra  $C(X)$ , i.e., the uniform convergence norm on  $X$ .

We want to have these additional properties for a quantum compact metric space  $(A, \|\cdot\|)$ . Unfortunately, because of the difficulties with Lipschitz seminorms, Latrémolière has not chosen a direct generalization of (4.4) in this work on quantum compact metric spaces. He introduced the following definition by considering the Jordan–Lie algebra of selfadjoint elements. We say that a quantum compact metric space  $(A, \|\cdot\|)$  is a  $(C, 0)$ -quasi-Leibniz quantum compact metric space if  $\|\cdot\|$  is Jordan–Lie subalgebra of  $A$  and if for any  $a, b \in \text{dom}\|\cdot\|$ , we have

$$\|a \circ b\| \leq C [\|a\| \|b\|_A + \|a\|_A \|b\|] \quad \text{and} \quad \|\{a, b\}\| \leq C [\|a\| \|b\|_A + \|a\|_A \|b\|] \tag{4.5}$$

for some constant  $C > 0$ , where we use the Jordan product

$$a \circ b \stackrel{\text{def}}{=} \frac{1}{2}(ab + ba)$$

and the Lie product

$$\{a, b\} \stackrel{\text{def}}{=} \frac{1}{2i}(ab - ba)$$

and if  $\|\cdot\|$  is lower semicontinuous, i.e.,

$$\{x \in \text{dom}\|\cdot\| : \|x\| \leq 1\} \tag{4.6}$$

is closed for the topology of the norm of  $A$ .

The following is essentially [58, Proposition 2.17] and Proposition 4.2. It is our main tool for checking the definition of quasi-Leibniz quantum compact metric spaces.

**Proposition 4.3.** *Let  $A$  be a unital  $C^*$ -algebra, and let  $\|\cdot\|$  be a seminorm defined on a dense  $\mathbb{C}$ -subspace  $\text{dom}\|\cdot\|$  of  $A$  such that*

- (1)  $\text{dom}\|\cdot\|$  is closed under the adjoint operation;
- (2)  $\{a \in \text{dom}\|\cdot\| : \|a\| = 0\} = \mathbb{C}1_A$ ;
- (3) there exists a constant  $C > 0$  such that for all  $a, b \in \text{dom}\|\cdot\|$ , we have

$$\|ab\| \leq C [\|a\|_A \|b\| + \|a\| \|b\|_A];$$

- (4) there exists a state  $\mu \in S(A)$  such that the set  $\{a \in \text{dom}\|\cdot\| : \|a\| \leq 1, \mu(a) = 0\}$  is relatively compact in  $A$  for the topology of the norm of  $A$ ;
- (5)  $\|\cdot\|$  is lower semicontinuous.



If  $\|\cdot\|_{\text{sa}}$  is the restriction of  $\|\cdot\|$  to  $A_{\text{sa}} \cap \text{dom}\|\cdot\|$ , then  $(A_{\text{sa}}, \|\cdot\|_{\text{sa}})$  is a  $(C, 0)$ -quasi-Leibniz quantum compact metric space.

**New quantum compact metric spaces.** Let  $G$  be a connected Lie group equipped with a family  $X = (X_1, \dots, X_m)$  of left-invariant Hörmander vector fields and a left Haar measure  $\mu_G$ . In this section, we suppose that  $G$  is compact. For the introduction of new quantum compact metric spaces, we need some preliminary results related to some estimates of the heat kernel. For any  $s \in G$ , a particular case of [87, Theorem V.4.3] gives

$$0 \leq K_t(s) \lesssim \frac{1}{t^{\frac{d}{2}}}, \quad 0 < t \leq 1, \tag{4.7}$$

where the local dimension  $d$  is defined in (3.7).

The following is essentially [76, pp. 339–341]. Since a point of [76, pp. 339–341] is misleading and since it is fundamental for us, we give an argument relying on the same nice ideas.

**Lemma 4.4.** *The operator  $\Delta_2: \text{dom}(\Delta_2) \subset L^2(G) \rightarrow L^2(G)$  has compact resolvent, and we have the estimate*

$$\|T_t\|_{L^1(G) \rightarrow L^\infty(G)} \lesssim \frac{1}{t^{\frac{d}{2}}}, \quad 0 < t \leq 1. \tag{4.8}$$

*Proof.* Note that for any  $0 < t \leq 1$ , we have

$$\|K_t\|_{L^2(G)} \lesssim \|K_t\|_{L^\infty(G)} \stackrel{(4.7)}{\lesssim} \frac{1}{t^{\frac{d}{2}}}. \tag{4.9}$$

By translation invariance of the normalized Haar measure of  $G$ , we deduce that

$$\begin{aligned} \int_{G \times G} |K_t(sr^{-1})|^2 ds dr &= \int_G \left( \int_G |K_t(sr^{-1})|^2 ds \right) dr \\ &= \int_G \left( \int_G |K_t(s)|^2 ds \right) dr \stackrel{(4.9)}{\leq} \frac{1}{t^d}. \end{aligned}$$

For any  $t > 0$ , we deduce by [51, Exercise 2.8.38, p. 170] and (3.1) that  $T_t: L^2(G) \rightarrow L^2(G)$  is a Hilbert–Schmidt operator. By [40, Theorem 4.29, p. 119], we conclude that the operator  $\Delta_2$  has compact resolvent. Finally, for any  $t > 0$ , we have

$$\|T_t\|_{L^1(G) \rightarrow L^\infty(G)} \stackrel{(2.2)}{=} \text{esssup}_{s,r \in G} |K_t(sr^{-1})| \stackrel{(4.7)}{\lesssim} \frac{1}{t^{\frac{d}{2}}}. \quad \blacksquare$$

**Lemma 4.5.** *The operator  $\Delta_2^{-1}: L_0^2(G) \rightarrow L_0^2(G)$  is compact.*

*Proof.* By Lemma 4.4, the operator  $\Delta_2: \text{dom} \Delta_2 \subset L^2(G) \rightarrow L^2(G)$  has compact resolvent. Note that  $\text{Ker} \Delta_2$  is an eigenspace, hence a reducing subspace for the selfadjoint operator  $\Delta_2$ . So, for any  $\lambda$  in the resolvent subset  $\rho(\Delta_2)$ , we have a well-defined operator

$(\lambda - \Delta_2)^{-1}: (\text{Ker } \Delta_2)^\perp \rightarrow (\text{Ker } \Delta_2)^\perp$  which is compact by composition. By the resolvent identity [48, p. 273], we deduce that  $\Delta_2^{-1}: (\text{Ker } \Delta_2)^\perp \rightarrow (\text{Ker } \Delta_2)^\perp$  is also compact by [36, p. 3]. Recall that  $(\text{Ker } \Delta_2)^\perp = L_0^2(G)$ . We conclude that  $\Delta_2^{-1}: L_0^2(G) \rightarrow L_0^2(G)$  is compact. ■

For the next statement, the domain of  $\|\cdot\|_{L_\alpha^p(G)}$  is defined as in (3.12).

**Theorem 4.6.** *Let  $G$  be a compact connected Lie group equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields. Suppose that  $0 < \alpha \leq 1$  (or  $0 < \alpha$  if  $G$  is nilpotent) and  $\frac{d}{\alpha} < p < \infty$ , where  $d$  is the local dimension defined in (3.7). Then,  $(C(G), \|\cdot\|_{L_\alpha^p(G)})$  defines a  $(C_{\alpha,p}, 0)$ -quasi-Leibniz quantum compact metric space for some constant  $C_{\alpha,p} > 0$ .*

*Proof.* We will prove the assumptions of Proposition 4.3. The third point of Proposition 4.3 is satisfied by (1.4) and the first two points by Lemma 3.5.

Since the normalized integral  $\int_G: C(G) \rightarrow \mathbb{C}$  is a state of the unital  $C^*$ -algebra  $C(G)$ , it suffices to show that

$$\left\{ f \in \text{dom}\|\cdot\|_{L_\alpha^p(G)} : \|f\|_{L_\alpha^p(G)} \leq 1, \int_G f = 0 \right\} \tag{4.10}$$

is relatively compact in  $C(G)$ .

Note that [34, p. 38] contains a proof of the existence of  $\omega > 0$  such that

$$\|T_t\|_{L_0^1(G) \rightarrow L^\infty(G)} \lesssim e^{-\omega t}, \quad t \geq 1. \tag{4.11}$$

Combined with (4.8), we deduce the estimate

$$\|T_t\|_{L_0^1(G) \rightarrow L^\infty(G)} \lesssim \frac{1}{t^{\frac{d}{2}}}, \quad t > 0.$$

With Lemma 4.5, we conclude that the assumptions of Proposition 2.3 are satisfied. Using this result with  $z = \frac{\alpha}{2}$  and  $q = \infty$ , the operator  $\Delta^{-\frac{\alpha}{2}}: L_0^p(G) \rightarrow L_0^\infty(G)$  is compact if  $p > \frac{d}{\alpha}$ . So, the image  $\mathcal{I}$  by  $\Delta^{-\frac{\alpha}{2}}$  of the closed unit ball  $\{g \in \overline{\text{Ran } \Delta_p} : \|g\|_{L^p(G)} \leq 1\}$  of  $L_0^p(G) = \overline{\text{Ran } \Delta_p}$  is relatively compact. Note that  $\text{Ran } \Delta_p^{\frac{\alpha}{2}} \subset \overline{\text{Ran } \Delta_p}$  by (2.8). Hence, the subset (write  $f = \Delta^{-\frac{\alpha}{2}} \Delta_p^{\frac{\alpha}{2}} f$ )

$$\{f \in C(G)_0 \cap \text{dom } \Delta_p^{\frac{\alpha}{2}} : \|\Delta_p^{\frac{\alpha}{2}}(f)\|_{L^p(G)} \leq 1\}$$

of  $\mathcal{I}$  is relatively compact in  $C(G)$ , where  $C(G)_0$  is the subspace of continuous functions with null integral. Since we have

$$\begin{aligned} & \{f \in C(G)_0 \cap \text{dom } \Delta_p^{\frac{\alpha}{2}} : \|\Delta_p^{\frac{\alpha}{2}}(f)\|_{L^p(G)} \leq 1\} \\ & \stackrel{(3.11)(3.12)}{=} \left\{ f \in \text{dom}\|\cdot\|_{L_\alpha^p(G)} : \|f\|_{L_\alpha^p(G)} \leq 1, \int_G f = 0 \right\}, \end{aligned}$$

we deduce that the subset (4.10) is also relatively compact in  $C(G)$ . The proof is complete. ■

**Remark 4.7.** The result is sharp. Consider the abelian compact group  $G = \mathbb{T}^2$  and the Laplacian

$$\begin{aligned} \Delta_2: \text{dom } \Delta_2 \subset L^2(\mathbb{T}^2) &\rightarrow L^2(\mathbb{T}^2), \\ e^{ni\cdot} \otimes e^{mi\cdot} &\mapsto -(n^2 + m^2)e^{ni\cdot} \otimes e^{mi\cdot} \end{aligned}$$

and  $\alpha = 1$ . By Example 3.2, the local dimension  $d$  of  $\mathbb{T}^2$  is 2. In [8, Remark 5.3] and its proof, it is showed that the set

$$\{f \in C(\mathbb{T}^2)_0 \cap \text{dom } \Delta_2^{\frac{1}{2}} : \|\Delta_2^{\frac{1}{2}}(f)\|_{L^2(\mathbb{T}^2)} \leq 1\}$$

is not bounded, in particular, not relatively compact. With the notation (3.11), this set can be written as

$$\left\{f \in \text{dom } \|\cdot\|_{L^2_1(\mathbb{T}^2)} : \|f\|_{L^2_1(\mathbb{T}^2)} \leq 1, \int_{\mathbb{T}^2} f = 0\right\}.$$

By Proposition 4.2, we conclude that we do not have in general a quantum compact metric space under the critical condition  $p = \frac{d}{\alpha}$ .

**Remark 4.8.** The inequality (1.4) is open if  $\alpha > 1$ . It would be interesting to find a counter-example.

**Remark 4.9.** We can replace the subelliptic Laplacian  $\Delta$  of (3.9) by a real second-order subelliptic operator

$$H \stackrel{\text{def}}{=} - \sum_{i,j=1}^m c_{ij} X_i X_j, \quad \text{where } c_{ij} \in \mathbb{R},$$

satisfying the condition

$$\frac{1}{2}(C + C^T) \geq \mu I \quad \text{for some } \mu > 0 \text{ and } C = [c_{ij}].$$

The  $L^p$ -realization  $H_p$  of this operator is a closed operator with domain  $\text{dom } H_p = L'_{p,2}$ . See [34, Chapter II] for more information. In the case of a compact connected Lie group  $G$ , the boundedness of Riesz transforms is proved in [34, p. 39]. Moreover, for any  $f \in L'_{2,1}$ , we have by [34, pp. 16–17]

$$\text{Re}\langle f, H_2 f \rangle_{L^2(G)} \geq \mu \sum_{k=1}^m \|X_k f\|_{L^2(G)}^2.$$

In particular,  $H_2 f = 0$  if and only if for any  $k \in \{1, \dots, m\}$ , we have  $X_k f = 0$ . This observation is useful for obtaining a suitable generalization of the third point of Lemma 3.5 (unfortunately, this argument only works in the case  $p \geq 2$ ).

The generalization of the Leibniz’s rule (1.4) for these operators for  $\alpha \in ]0, 1[$  is an open question.

**Remark 4.10.** It may be worthy to study the family of the quantum compact metric spaces  $(C(G), \|\cdot\|_{L^\alpha_\alpha(G)})$  when  $p \rightarrow \frac{d}{\alpha}$  from the perspective of the quantum Gromov–Hausdorff distance. What can be said about the “limit”?

**Remark 4.11.** It is unclear if there exists a formula for the restriction of the Monge–Kantorovich metric (4.2) on the subset of pure states of  $C(G)$ , i.e., the map

$$(s, s') \mapsto \sup\{|f(s) - f(s')| : f \in C(G, \mathbb{R}), \|f\|_{L^\alpha_\alpha(G)} \leq 1\}$$

on  $G \times G$ . It would be interesting to understand this quantity to equivalence with respect to a constant. The question is natural when we compare to the next situation of Theorem 7.4.

### 5. Quantum locally compact metric spaces

**Quantum locally compact metric spaces.** The basic reference is [57]. A topography on a  $C^*$ -algebra  $A$  is an abelian  $C^*$ -subalgebra  $\mathfrak{M}$  of  $A$  containing an approximate identity for  $A$ . A topographic quantum space  $(A, \mathfrak{M})$  is an ordered pair of a  $C^*$ -algebra  $A$  and a topography  $\mathfrak{M}$  on  $A$ . Let  $(A, \mathfrak{M})$  be a topographic quantum space. A state  $\varphi \in S(A)$  is local when there exists a compact  $K$  of the Gelfand spectrum of  $\mathfrak{M}$  such that  $\varphi(1_K) = 1$ . A Lipschitz triple  $(A, \|\cdot\|, \mathfrak{M})$  is a triple where  $(A, \|\cdot\|)$  is a Lipschitz pair and  $\mathfrak{M}$  is a topography on  $A$ .

Let  $(A, \|\cdot\|, \mathfrak{M})$  be a Lipschitz triple. The definition of quantum locally compact quantum metric spaces of [57] is equivalent to saying that  $(A, \|\cdot\|, \mathfrak{M})$  is a quantum locally compact quantum metric space if and only if for any compactly supported element  $g, h \in \mathfrak{M}$  and for some local state  $\mu$  of  $A$ , the set

$$\{gah : a \in (uA)_{\text{sa}}, \|a\| \leq 1, \mu(a) = 0\}$$

is relatively compact for the topology associated to  $\|\cdot\|_A$ . Here we identify  $\mu$  with its unique extension  $a + \lambda 1 \mapsto \mu(a) + \lambda$  as a state of the unital  $C^*$ -algebra  $uA$ .

**Quasi-Leibniz quantum locally compact metric space.** Unfortunately, Latrémolière did not generalize the notion of definition of quasi-Leibniz quantum compact metric spaces of Section 4 to the locally compact case. We make an attempt by saying that a quantum locally compact quantum metric space  $(A, \|\cdot\|, \mathfrak{M})$  is a quasi-Leibniz quantum locally compact metric space if the restriction of  $\|\cdot\|$  on  $A_{\text{sa}}$  satisfies the points (4.5) and (4.6), which is slightly less general than [8, Section 5.5].

**Criterion of relative compactness.** The following is a locally compact group generalization [15, Exercise 26, p. VIII.72] [32, Problem 4, p. 283] (see also [34, Theorem A.4.1] for a particular case) of the classical Fréchet–Kolmogorov theorem on relative compactness.

**Theorem 5.1.** *Let  $G$  be a locally compact group equipped with a left Haar measure. Suppose that  $1 \leq p < \infty$ . Let  $\mathcal{F}$  be a subset of the Banach space  $L^p(G)$ . Then,  $\mathcal{F}$  is*

relatively compact if and only if there exists  $M > 0$  such that

$$\limsup_{s \rightarrow e} \sup_{f \in \mathcal{F}} \|\lambda_s f - f\|_{L^p(G)} = 0,$$

$$\sup_{f \in \mathcal{F}} \|f\|_{L^p(G)} \leq M \quad \text{and} \quad \lim_{r \rightarrow \infty} \sup_{f \in \mathcal{F}} \int_{G-B(e,r)} |f(s)|^p d\mu_G(s) = 0.$$

Now, consider a connected Lie group  $G$  equipped with a family  $X$  of left-invariant Hörmander vector fields with polynomial volume growth and local dimension  $d$ . We suppose that  $G$  is *not* compact. Let  $K$  be a compact subset of  $G$ . We denote by  $C_K(G)$  the space of continuous functions on  $G$  with support in  $K$ .

Suppose that  $1 < p < \infty$  and  $\alpha \geq 0$ . Following essentially [27, p. 287], we define the subspace

$$L^p_\alpha(G) \stackrel{\text{def}}{=} \text{dom } \Delta_p^{\frac{\alpha}{2}}$$

of  $L^p(G)$ . If  $f \in L^p_\alpha(G)$ , we will use the notation

$$\|f\|_{L^p_\alpha(G)} \stackrel{\text{def}}{=} \|\Delta_p^{\frac{\alpha}{2}}(f)\|_{L^p(G)} + \|f\|_{L^p(G)} \stackrel{(2.9)}{\approx} \|(\text{Id} + \Delta_p)^{\frac{\alpha}{2}}(f)\|_{L^p(G)}. \tag{5.1}$$

We refer to [40, Section II.5] and [56, Section 15.E] for the link with Sobolev towers. If  $\alpha p > d$ , we have by [27, p. 287] and [16, Theorem 4.4 (c)] a Sobolev embedding  $L^p_\alpha(G) \subset L^\infty(G)$ :

$$\|f\|_{L^\infty(G)} \lesssim \|f\|_{L^p_\alpha(G)}, \quad f \in \text{dom } \Delta_p^{\frac{\alpha}{2}}. \tag{5.2}$$

Note that by [48, Proposition 3.2.3], the Bessel potential  $(\text{Id} + \Delta)^{-\alpha}$  is a *bounded* operator on the Banach space  $L^p(G)$  for any  $\alpha \in \mathbb{C}$  with  $\text{Re } \alpha > 0$ . Consequently, if  $0 < \alpha \leq \beta$ , it is obvious to check with (2.9) that

$$\|f\|_{L^p_\beta(G)} \lesssim \|f\|_{L^p_\alpha(G)}. \tag{5.3}$$

A contractive inclusion for the case  $G = \mathbb{R}^n$  is proved in [78, p. 135] with a different argument. A *contractive* version of (5.3) is stated without proof in the inequality following [16, equation (3.1)], but it is a mistake confirmed by the authors of this paper.

**Proposition 5.2.** *Let  $G$  be a non-compact connected Lie group equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields. Suppose that  $G$  has polynomial volume growth. Let  $\alpha \geq 1$  and  $\max\{1, \frac{d}{\alpha}\} < p < \infty$ . If  $g: G \rightarrow \mathbb{C}$  is a compactly supported continuous function, then the subset*

$$g\{f \in C_0(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}} : \|f\|_{L^p_\alpha(G)} \leq 1, f(e) = 0\} \tag{5.4}$$

is relatively compact in  $L^\infty(G)$ .

*Proof.* Let  $K$  be a compact subset of  $G$ . For any  $M \geq 0$ , consider the subset

$$E_{K,p,M} \stackrel{\text{def}}{=} \{f \in C_K(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}} : \|f\|_{L^p_\alpha(G)} \leq M\} \tag{5.5}$$

of the Banach space  $L^p(G)$ . If  $f \in E_{K,p,M}$ , using the Sobolev embedding  $L_\alpha^p(G) \subset L^\infty(G)$ , we obtain

$$\|f\|_{L^p(G)} \lesssim_{K,p} \|f\|_{L^\infty(G)} \stackrel{(5.2)}{\lesssim} \|f\|_{L_\alpha^p(G)} \leq M.$$

Consequently, the subset  $E_{K,p,M}$  is bounded in  $L^p(G)$ . Moreover, using Lemma 3.1, we have for any function  $f \in E_{K,p,M}$  and any  $s \in G$

$$\begin{aligned} \|(\text{Id} - \lambda_s)f\|_{L^p(G)} &\stackrel{(3.5)}{\leq} \text{dist}_{\text{CC}}^{p*}(s, e) \left( \sum_{k=1}^m \|X_{k,p}(f)\|_{L^p(G)}^p \right)^{\frac{1}{p}} \\ &\approx_p \text{dist}_{\text{CC}}^{p*}(s, e) \sum_{k=1}^m \|X_{k,p}(f)\|_{L^p(G)} \\ &\stackrel{(3.10)}{\lesssim_p} \text{dist}_{\text{CC}}^{p*}(s, e) \|\Delta_p^{\frac{1}{2}}(f)\|_{L^p(G)} \stackrel{(5.1)}{\leq} \text{dist}_{\text{CC}}^{p*}(s, e) \|f\|_{L_1^p(G)} \\ &\stackrel{(5.3)}{\lesssim} \text{dist}_{\text{CC}}^{p*}(s, e) \|f\|_{L_\alpha^p(G)} \stackrel{(5.5)}{\leq} M \text{dist}_{\text{CC}}^{p*}(s, e). \end{aligned}$$

With Theorem 5.1, we obtain the relative compactness of the subset  $E_{K,p,M}$  in  $L^p(G)$  and of its subset

$$F_{K,p,M} \stackrel{\text{def}}{=} \{f \in C_K(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}} : \|f\|_{L_\alpha^p(G)} \leq M, f(e) = 0\}.$$

The operator  $(\text{Id} + \Delta)^{-\frac{\alpha}{2}} : L^p(G) \rightarrow \overline{\text{Ran } \Delta_\infty}$  is bounded by (5.2) and (5.1) (hence uniformly continuous). Applying this operator to the previous subset by writing

$$f = (\text{Id} + \Delta)^{-\frac{\alpha}{2}} (\text{Id} + \Delta_p)^{\frac{\alpha}{2}} f,$$

we obtain that the set  $F_{K,p,M}$  is relatively compact in  $L^\infty(G)$ , hence in  $C_0(G)$ . Note that if  $f$  belongs to  $C_0(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}}$  and satisfies  $\|f\|_{L_\alpha^p(G)} \leq 1$  and if  $g \in C_c(G)$ , we have

$$\begin{aligned} \|gf\|_{L_\alpha^p(G)} &\stackrel{(1.3)}{\lesssim_p} \|g\|_{L_\alpha^p(G)} \|f\|_{L^\infty(G)} + \|g\|_{L^\infty(G)} \|f\|_{L_\alpha^p(G)} \\ &\stackrel{(5.2)}{\lesssim} \|f\|_{L_\alpha^p(G)} [\|g\|_{L^\infty(G)} + \|g\|_{L_\alpha^p(G)}] \leq \|g\|_{L^\infty(G)} + \|g\|_{L_\alpha^p(G)}. \end{aligned}$$

Consequently, if  $\text{supp } g \subset K$ , we obtain that the subset (5.4) is included in some subset  $F_{K,p,M}$  with  $M \stackrel{\text{def}}{=} \|g\|_{L^\infty(G)} + \|g\|_{L_\alpha^p(G)}$ . ■

**Remark 5.3.** In [78, Proposition 3, p. 138], it is proved that a measurable function  $f$  belongs to the space  $L_1^p(\mathbb{R}^n)$  if and only if

$$\|(\text{Id} - \lambda_s)f\|_{L^p(\mathbb{R}^n)} = O(|s|).$$

So, we are not confident in a possible generalization of Proposition 5.2 to the case  $0 < \alpha < 1$ . Note also that in [78, Exercise 6.1, p. 159], it is stated that a measurable function  $f$  belongs to the space  $L_1^p(\mathbb{R}^n)$  if and only if  $f$  belongs to  $L^p(\mathbb{R}^n)$ ,  $f$  is absolutely continuous, and the partial derivatives  $\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n}$  belong to  $L^p(\mathbb{R}^n)$ .

**Remark 5.4.** It is transparent for the author that some form of local ultracontractivity [45, Definition 2.11] can be used to give some variants or generalizations of the previous proof to other contexts.

We also define the seminorm  $\|\cdot\|_{L^p_\alpha(G)}$  on the space  $(C_0(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}}) \oplus \mathbb{C}1$  by letting  $\|f\|_{L^p_\alpha(G)} \stackrel{\text{def}}{=} \|f_0\|_{L^p_\alpha(G)}$  for any element  $f = f_0 + \lambda 1$  of the space  $(C_0(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}}) \oplus \mathbb{C}1$ . Note that the latter space is a subspace of the unitization  $C_0(G) \oplus \mathbb{C}1$  of the non-unital algebra  $C_0(G)$ .

**Lemma 5.5.** *The restriction of the seminorm  $\|\cdot\|_{L^p_\alpha(G)}$  on the subspace  $(C_0(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}})$  is lower semicontinuous.*

*Proof.* Let  $f \in C_0(G)$  and  $(f_n)$  be a sequence of elements of  $C_0(G) \cap \text{dom } \Delta_p^{\frac{\alpha}{2}}$  such that  $(f_n)$  converges to  $f$  for the norm topology of  $C_0(G)$  and  $\|f_n\|_{L^p_\alpha(G)} \leq 1$  for any  $n$ ; that is,

$$\|\Delta_p^{\frac{\alpha}{2}}(f_n)\|_{L^p(G)} + \|f_n\|_{L^p(G)} \leq 1$$

by (5.1). Note that in particular that the sequences  $(f_n)$  and  $(\Delta_p^{\frac{\alpha}{2}}(f_n))$  are bounded in the Banach space  $L^p(G)$ . We have to prove that the function  $f$  belongs to  $\text{dom } \Delta_p^{\frac{\alpha}{2}}$  and that  $\|f\|_{L^p_\alpha(G)} \leq 1$ .

First, we show that the sequence  $(f_n)$  converges to  $f$  for the weak topology of the Banach space  $L^p(G)$ . Indeed, for any function  $g \in C_c(G)$ , we have

$$\left| \int_G (f_n - f)g \, d\mu_G \right| \leq \|f_n - f\|_{L^\infty(G)} \int_G |g| \, d\mu_G \xrightarrow{n \rightarrow +\infty} 0.$$

Using the boundedness of the sequence  $(f_n)$  in  $L^p(G)$ , we obtain the claim with [65, Exercise 2.71, p. 234] since we have the convergence with any function  $g$  of the dense subspace  $C_c(G)$  of the Banach space  $L^{p^*}(G)$ .

Since the sequence  $(\Delta_p^{\frac{\alpha}{2}} f_n)$  is bounded in the Banach space  $L^p(G)$  and since bounded sets are weakly relatively compact by [65, Theorem 2.8.2], there exists a weakly convergent subnet  $(\Delta_p^{\frac{\alpha}{2}} f_{n_j})$ . Then,  $(f_{n_j}, \Delta_p^{\frac{\alpha}{2}} f_{n_j})$  is a weakly convergent net in the graph of the closed operator  $\Delta_p^{\frac{\alpha}{2}}$ . Note that this graph is closed and convex, hence weakly closed by [65, Theorem 2.5.16]. Thus, the limit of  $(f_{n_j}, \Delta_p^{\frac{\alpha}{2}} f_{n_j})$  belongs again to the graph and is of the form  $(g, \Delta_p^{\frac{\alpha}{2}} g)$  for some  $g \in \text{dom } \Delta_p^{\frac{\alpha}{2}}$ . In particular, the net  $(f_{n_j})$  converges weakly to  $g$  and  $\Delta_p^{\frac{\alpha}{2}}(f_{n_j})$  converges weakly to  $\Delta_p^{\frac{\alpha}{2}}(g)$ . We infer that  $f = g$ . We conclude that  $f$  belongs to  $\text{dom } \Delta_p^{\frac{\alpha}{2}}$ . Moreover, using the weakly lower semicontinuity of the norm [65, Theorem 2.5.21], we obtain

$$\begin{aligned} \|f\|_{L^p_\alpha(G)} &\stackrel{(5.1)}{=} \|\Delta_p^{\frac{\alpha}{2}}(f)\|_{L^p(G)} + \|f\|_{L^p(G)} \\ &\leq \liminf_j [\|\Delta_p^{\frac{\alpha}{2}}(f_{n_j})\|_{L^p(G)} + \liminf_j \|f_{n_j}\|_{L^p(G)}] \leq 1. \quad \blacksquare \end{aligned}$$

**Corollary 5.6.** *Let  $G$  be a non-compact connected Lie group equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields. Suppose that  $G$  has polynomial volume growth. Let  $\alpha \geq 1$  and  $\max\{1, \frac{d}{\alpha}\} < p < \infty$ , where  $d$  is the local dimension defined in (3.7). Then*

$$(C_0(G), \|\cdot\|_{L^\alpha_p(G)}, C_0(G))$$

*defines a  $(C_{\alpha,p}, 0)$ -quasi-Leibniz quantum locally compact metric space for some constant  $C_{\alpha,p} > 0$ .*

*Proof.* Parts (1) and (2) of Lemma 3.5 say that

$$\text{dom}\|\cdot\|_{L^\alpha_p(G)} \stackrel{(3.12)}{=} C_0(G) \cap \text{dom} \Delta_p^{\frac{\alpha}{2}}$$

is closed under the adjoint operation and dense in the space  $C_0(G)$ . Consequently,

$$(C_0(G) \cap \text{dom} \Delta_p^{\frac{\alpha}{2}}) \oplus \mathbb{C}1$$

is also closed under the adjoint operation of the algebra  $C_0(G) \oplus \mathbb{C}1$  and dense in  $C_0(G) \oplus \mathbb{C}1$ .

Let  $f = f_0 + \lambda 1$  be an element of the direct sum  $(C_0(G) \cap \text{dom} \Delta_p^{\frac{\alpha}{2}}) \oplus \mathbb{C}1$ . Suppose that  $\|f\|_{L^\alpha_p(G)} = 0$ . Then, by definition,

$$\|\Delta_p^{\frac{\alpha}{2}}(f_0)\|_{L^p(G)} + \|f_0\|_{L^p(G)} \stackrel{(5.1)}{=} \|f_0\|_{L^\alpha_p(G)} = 0.$$

Hence,  $\|f_0\|_{L^p(G)} = 0$  and finally  $f_0 = 0$ . We conclude that  $f = \lambda 1$ . So, (4.1) is satisfied. So, we have a Lipschitz pair

$$(C_0(G), \|\cdot\|_{L^\alpha_p(G)}).$$

This Dirac measure  $\delta_e$  is clearly a local state since it is supported by the compact  $\{e\}$ . The Leibniz rule is given by (1.3). The lower semicontinuity is given by Lemma 5.5. We conclude with Proposition 5.2. ■

In the end of this section, we will investigate what happens when we replace the operator  $\text{Id} + \Delta_p$  by the subelliptic Laplacian  $\Delta_p$  in one case. The obtained result of Proposition 5.7 is a bit different. Indeed, it is obvious that the addition of the identity to the operator  $\Delta$  removes global phenomenons.

Suppose that the connected Lie group  $G$  is equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields and has polynomial growth with  $d < D$ . Such group is not compact. For example, by [76, p. 273], this condition is satisfied if  $G$  is simply connected, nilpotent with  $G \not\cong \mathbb{R}^d$ . Consider some  $1 < p < \infty$  and some  $\alpha > 0$ . If  $d < \alpha p < D$ , it is stated in [27, p. 288] and [28, p. 197] that

$$\|f\|_{L^\infty(G)} \lesssim \|f\|_{L^\alpha_p(G)}, \quad f \in C_c^\infty(G). \tag{5.6}$$

Now, we prove an analog of Proposition 5.2.



**Proposition 5.7.** *Let  $G$  be a connected Lie group equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields. Suppose that  $G$  has polynomial growth with  $d < D$ . Assume that  $d < p < D$ . If  $g: G \rightarrow \mathbb{C}$  is a compactly supported continuous function, then the subset*

$$g\{f \in C_0(G) \cap \text{dom } \Delta_p^{\frac{1}{2}} : \|f\|_{L^p_1(G)} \leq 1, f(e) = 0\} \tag{5.7}$$

is relatively compact in  $L^\infty(G)$ .

*Proof.* Let  $K$  be a compact subset of  $G$ . For any  $M \geq 0$ , consider the subset

$$E_{K,p,M} \stackrel{\text{def}}{=} \{f \in C_K(G) \cap \text{dom } \Delta_p^{\frac{1}{2}} : \|f\|_{L^p_1(G)} \leq M\} \tag{5.8}$$

of the space  $L^p(G)$ . If  $f \in E_{K,p,M}$ , using the Sobolev embedding  $L^p_1(G) \subset L^\infty(G)$  of (5.6), we obtain

$$\|f\|_{L^p(G)} \lesssim_{K,p} \|f\|_{L^\infty(G)} \stackrel{(5.6)}{\lesssim} \|f\|_{L^p_1(G)} \leq M.$$

We infer that the subset  $E_{K,p,M}$  is bounded in  $L^p(G)$ . Furthermore, using Lemma 3.1, we have for any function  $f \in E_{K,p,M}$

$$\begin{aligned} \|(\text{Id} - \lambda_s)f\|_{L^p(G)} &\stackrel{(3.5)}{\leq} \text{dist}_{\text{CC}}^{p^*}(s, e) \left( \sum_{k=1}^m \|X_{k,p}(f)\|_{L^p(G)}^p \right)^{\frac{1}{p}} \\ &\approx_p \text{dist}_{\text{CC}}^{p^*}(s, e) \sum_{k=1}^m \|X_{k,p}(f)\|_{L^p(G)} \\ &\stackrel{(3.10)}{\lesssim_p} \text{dist}_{\text{CC}}^{p^*}(s, e) \|\Delta_p^{\frac{1}{2}}(f)\|_{L^p(G)} \\ &\stackrel{(3.11)}{=} \text{dist}_{\text{CC}}^{p^*}(s, e) \|f\|_{L^p_1(G)} \stackrel{(5.8)}{\leq} M \text{dist}_{\text{CC}}^{p^*}(s, e). \end{aligned}$$

By Theorem 5.1, the subset  $E_{K,p,M}$  is relatively compact in  $L^p(G)$ . Hence, its subset

$$F_{K,p,M} \stackrel{\text{def}}{=} \{f \in C_K(G) \cap \text{dom } \Delta_p^{\frac{1}{2}} : \|f\|_{L^p_1(G)} \leq M, f(e) = 0\}$$

is also relatively compact in  $L^p(G)$ . The operator  $\Delta^{-\frac{1}{2}}: L^p(G) \rightarrow \overline{\text{Ran } \Delta_\infty}$  is bounded by (5.6), hence uniformly continuous. Applying this operator to the previous subset by writing  $f = \Delta^{-\frac{1}{2}} \Delta_p^{\frac{1}{2}} f$ , we obtain that the set  $F_{K,p,M}$  is relatively compact in  $L^\infty(G)$ , hence in the space  $C_0(G)$ . Note that we have

$$\begin{aligned} \|gf\|_{L^p_1(G)} &\stackrel{(1.4)}{\lesssim_p} \|g\|_{L^p_1(G)} \|f\|_{L^\infty(G)} + \|g\|_{L^\infty(G)} \|f\|_{L^p_1(G)} \\ &\stackrel{(5.6)}{\lesssim} \|f\|_{L^p_1(G)} [\|g\|_{L^\infty(G)} + \|g\|_{L^p_1(G)}] \stackrel{(5.8)}{\lesssim} \|g\|_{L^\infty(G)} + \|g\|_{L^p_1(G)}. \end{aligned}$$

Consequently, if  $\text{supp } g \subset K$ , we obtain that the subset (5.7) is included in some subset  $F_{K,p,M}$  with  $M \stackrel{\text{def}}{=} \|g\|_{L^\infty(G)} + \|g\|_{L^p_1(G)}$ . ■

Similarly to Corollary 5.6, we can obtain the following result where the seminorm  $\|\cdot\|_{L^\alpha_p(G)}$  is defined on

$$\text{dom}\|\cdot\|_{L^p_1(G)} \stackrel{(3.12)}{=} C_0(G) \cap \text{dom}\Delta^{\frac{1}{p}}.$$

Unfortunately, we are not able to demonstrate the lower semicontinuity of the seminorm  $\|\cdot\|_{L^p_1(G)}$ . So, we cannot make the statement that we have a *quasi-Leibniz* quantum locally compact metric space.

**Corollary 5.8.** *Let  $G$  be a connected Lie group equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields. Suppose that  $G$  has polynomial growth with  $d < D$ . Then, the triple  $(C_0(G), \|\cdot\|_{L^p_1(G)}, C_0(G))$  defines a quantum locally compact metric space.*

## 6. Compact spectral triples and spectral dimension

**Possibly kernel-degenerate compact spectral triples.** Consider a triple  $(A, Y, \mathcal{D})$  constituted of the following data: a Banach space  $Y$ , a closed unbounded operator  $\mathcal{D}$  on  $Y$  with dense domain  $\text{dom}\mathcal{D} \subset Y$ , and an algebra  $A$  equipped with a homomorphism  $\pi: A \rightarrow B(Y)$ . In this case, we define the Lipschitz algebra

$$\text{Lip}_{\mathcal{D}}(A) \stackrel{\text{def}}{=} \{a \in A : \pi(a) \cdot \text{dom}\mathcal{D} \subset \text{dom}\mathcal{D} \text{ and the unbounded operator } [\mathcal{D}, \pi(a)]: \text{dom}\mathcal{D} \subset Y \rightarrow Y \text{ extends to an element of } B(Y)\}. \quad (6.1)$$

We say that  $(A, Y, \mathcal{D})$  is a (possibly kernel-degenerate) compact spectral triple if in addition  $Y$  is a Hilbert space  $H$ ,  $A$  is a  $C^*$ -algebra,  $D$  is a selfadjoint operator on  $Y$ , and if we have the following:

- (1)  $\mathcal{D}^{-1}$  is a compact operator on  $\overline{\text{Ran}\mathcal{D}} \stackrel{(2.5)}{=} (\text{Ker}\mathcal{D})^\perp$ ;
- (2) the subset  $\text{Lip}_{\mathcal{D}}(A)$  is dense in  $A$ .

We essentially follow [21, Definition 2.1] and [8, Definition 5.10]. Note that there exist different variations of this definition in the literature, see; e.g., [36, Definition 1.1]. Moreover, we can replace  $\mathcal{D}^{-1}$  by  $|\mathcal{D}|^{-1}$  in the first point by an elementary functional calculus argument.

We equally refer to [8, Definition 5.10] for the notion of compact Banach spectral triple which is a generalization for the case of an operator  $\mathcal{D}$  acting on a Banach space  $Y$  instead of a Hilbert space  $H$ .

**Example 6.1.** If  $M$  is a compact oriented Riemannian manifold  $M$ , we can associate the spectral triple  $(C(M), L^2(\wedge T^*M), D)$ , where  $L^2(\wedge T^*M)$  is the Hilbert space of square-integrable complex-valued forms on  $M$  and where  $D$  is the Hodge–Dirac operator (also called Hodge–de Rham operator). If  $M$  is in addition a spin manifold, we can also consider the spectral triple  $(C(M), L^2(M, S), \mathcal{D})$  obtained by using the Hilbert space  $L^2(M, S)$ , the space of square-integrable spinors on  $M$ , and the Dirac operator  $\mathcal{D}$ . In both cases, the functions of  $C(M)$  act on the Hilbert space by multiplication operators.

**Spectral dimension.** Let  $(A, H, \mathcal{D})$  be a compact spectral triple. By [71, Proposition 5.3.38], we have  $\text{Ker } |\mathcal{D}| = \text{Ker } \mathcal{D}$ . Moreover, the operator  $|\mathcal{D}|^{-1}$  is well defined on

$$\overline{\text{Ran } \mathcal{D}} \stackrel{(2.5)}{=} (\text{Ker } \mathcal{D})^\perp.$$

Furthermore, we can extend it by letting  $|\mathcal{D}|^{-1} = 0$  on  $\text{Ker } \mathcal{D}$ . Following [36, p. 4], we say that a compact spectral triple  $(A, H, \mathcal{D})$  is  $\alpha$ -summable for some  $\alpha > 0$  if  $\text{Tr } |\mathcal{D}|^{-\alpha} < \infty$ , that is if the operator  $|\mathcal{D}|^{-1}$  belongs to the Schatten class  $S^\alpha(H)$ . In this case, the spectral dimension of the spectral triple is defined by

$$\dim(A, H, \mathcal{D}) \stackrel{\text{def}}{=} \inf\{\alpha > 0 : \text{Tr } |\mathcal{D}|^{-\alpha} < \infty\}. \tag{6.2}$$

See also [43, p. 450] and [18, p. 38 and Definition 6.2, p. 47] for a variation of this definition.

We will use the following lemma which is a slight variation of [43, Lemma 10.8, p. 450].

**Lemma 6.2.** *If  $|\mathcal{D}|^{-\alpha}$  is trace-class then for any  $t > 0$  the operator  $e^{-t\mathcal{D}^2}$  is trace-class and we have*

$$\text{Tr } e^{-t\mathcal{D}^2} \lesssim \frac{1}{t^{\frac{\alpha}{2}}}, \quad t > 0.$$

*Proof.* Note that here the operator  $|\mathcal{D}|^{-\alpha}$  is defined and bounded on  $\overline{\text{Ran } \mathcal{D}}$ . However, we can extend it by letting  $|\mathcal{D}|^{-\alpha} = 0$  on  $\text{Ker } \mathcal{D}$ . For any  $t > 0$ , we have

$$e^{-t\mathcal{D}^2} = |\mathcal{D}|^\alpha e^{-t\mathcal{D}^2} |\mathcal{D}|^{-\alpha}.$$

An elementary study of the function  $f: \lambda \mapsto \lambda^\alpha e^{-t\lambda^2}$  on  $\mathbb{R}^+$  shows that

$$f'(\lambda) = \lambda^{\alpha-1} e^{-t\lambda^2} (\alpha - 2\lambda^2 t) \quad \text{for any } \lambda \geq 0$$

and consequently that  $f$  is bounded and that its maximum is  $(\frac{\alpha}{2t})^{\frac{\alpha}{2}} e^{-\frac{\alpha}{2}}$  in  $\lambda = \sqrt{\frac{\alpha}{2t}}$ . We conclude by functional calculus that the operator  $|\mathcal{D}|^\alpha e^{-t\mathcal{D}^2}$  is bounded and that

$$\text{Tr } e^{-t\mathcal{D}^2} = \|e^{-t\mathcal{D}^2}\|_{S^1(H)} \leq \| |\mathcal{D}|^\alpha e^{-t\mathcal{D}^2} \|_{B(H)} \| |\mathcal{D}|^{-\alpha} \|_{S^1(H)} \lesssim \frac{1}{t^{\frac{\alpha}{2}}}. \quad \blacksquare$$

**Hodge–Dirac operator.** Let  $G$  be a unimodular connected Lie group equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields and consider a Haar measure  $\mu_G$  on  $G$ . Suppose  $1 \leq p \leq \infty$ . Recall that we have a canonical isometry  $\ell_m^p(L^p(G)) = L^p(G, \ell_m^p)$ . We define the unbounded closed operator  $\nabla_p$  from  $L^p(G)$  into the space  $L^p(G, \ell_m^p)$  by

$$\text{dom } \nabla_p = \text{dom } X_{1,p} \cap \dots \cap \text{dom } X_{m,p}$$

and

$$\nabla_p f \stackrel{\text{def}}{=} (X_{1,p} f, \dots, X_{m,p} f), \quad f \in \text{dom } \nabla_p. \tag{6.3}$$

If  $1 < p < \infty$ , note that  $\text{dom } \nabla_p = \text{dom } \Delta_p^{\frac{1}{2}}$  by Proposition 3.4. For any functions  $f, g$  of  $\text{dom } \nabla_p \cap L^\infty(G)$ , then  $fg$  belongs to  $\text{dom } \nabla_p \cap L^\infty(G)$ , and we have

$$\nabla_p(fg) = g \cdot \nabla_p(f) + f \cdot \nabla_p(g), \tag{6.4}$$

where

$$f \cdot (h_1, \dots, h_m) \stackrel{\text{def}}{=} (fh_1, \dots, fh_m).$$

See [27, p. 289] for a generalization.

If  $1 < p < \infty$ , we introduce the unbounded closed operator

$$\mathcal{D}_p \stackrel{\text{def}}{=} \begin{bmatrix} 0 & (\nabla_p^*)^* \\ \nabla_p & 0 \end{bmatrix} \tag{6.5}$$

on the Banach space  $L^p(G) \oplus_p L^p(G, \ell_m^p)$  defined by

$$\mathcal{D}_p(f, g) \stackrel{\text{def}}{=} ((\nabla_p^*)^*(g), \nabla_p(f)), \quad f \in \text{dom } \nabla_p, g \in \text{dom}(\nabla_p^*)^*. \tag{6.6}$$

We call it the Hodge–Dirac operator of the subelliptic Laplacian  $\Delta = -(X_1^2 + \dots + X_m^2)$ . These operators are related by the computation

$$\mathcal{D}_p^2 \stackrel{(6.5)}{=} \begin{bmatrix} 0 & (\nabla_p^*)^* \\ \nabla_p & 0 \end{bmatrix}^2 = \begin{bmatrix} (\nabla_p^*)^* \nabla_p & 0 \\ 0 & \nabla_p (\nabla_p^*)^* \end{bmatrix} = \begin{bmatrix} \Delta_p & 0 \\ 0 & \nabla_p (\nabla_p^*)^* \end{bmatrix}. \tag{6.7}$$

The operator  $\mathcal{D}_2$  is identical to the operator  $\Pi$  of [10, proof of Theorem 1.2] with  $b = \text{Id}$ .

We will use just the following lemma which describes a tractable subspace for the adjoint operator  $(\nabla_p^*)^*$ . If  $(\varphi_j)$  is a Dirac net of functions of  $C_c^\infty(G)$  and if  $h = (h_1, \dots, h_m)$ , we will use the notation

$$\text{Reg}_j h \stackrel{\text{def}}{=} (h_1 * \varphi_j, \dots, h_m * \varphi_j)$$

as soon as it makes sense.

**Lemma 6.3.** *Suppose that  $1 < p < \infty$ . The subspace  $C_c^\infty(G) \otimes \ell_m^p$  is a core of the unbounded operator  $(\nabla_p^*)^*$ .*

*Proof.* It is easy to check (use [53, Problem 5.24, p. 168]) that  $C_c^\infty(G) \otimes \ell_m^p$  is a subset of  $\text{dom}(\nabla_p^*)^*$ . We consider a Dirac net  $(\varphi_j)$  of functions of  $C_c^\infty(G)$ . Let  $h = (h_1, \dots, h_m)$  be an element of  $\text{dom}(\nabla_p^*)^*$ . Then

$$\text{Reg}_j h = (h_1 * \varphi_j, \dots, h_m * \varphi_j)$$

belongs to  $C_c^\infty(G) \otimes \ell_m^p$ . It remains to show that  $(\text{Reg}_j h)$  converges to  $h$  in the graph norm of  $(\nabla_p^*)^*$ . By [15, Proposition 20, p. VIII.44], the net  $(\text{Reg}_j h)$  converges to  $h$  in  $L^p(G, \ell_m^p)$ . For any  $1 \leq k \leq m$ , we put

$$a_k \stackrel{\text{def}}{=} X_k(e).$$

If  $f \in C_c^\infty(G)$ , using [77, equation (9.19)] and the equalities  $X_k = d\lambda(a_k)$  [34, p. 14] in the second equality and [63, Proposition 3.14] in the third equality, we have

$$\begin{aligned} \nabla \operatorname{Reg}_j f &\stackrel{(6.3)}{=} (X_1(f * \varphi_j), \dots, X_m(f * \varphi_j)) \\ &= (d\lambda(a_1)(\lambda_f(\varphi_j)), \dots, d\lambda(a_m)(\lambda_f(\varphi_j))) \\ &= (\lambda_{X_1 f}(\varphi_j), \dots, \lambda_{X_m f}(\varphi_j)) = ((X_1 f) * \varphi_j, \dots, (X_m f) * \varphi_j) \\ &= \operatorname{Reg}_j(X_1 f, \dots, X_m f) = \operatorname{Reg}_j(\nabla f), \end{aligned} \tag{6.8}$$

where  $\lambda_f(g) \stackrel{\text{def}}{=} f * g$  and where  $d\lambda$  is the derived representation [63, Definition 3.12] of the left regular representation  $\lambda$ . Moreover, for any  $g \in C_c^\infty(G)$ , we have used [32, Theorem (14.10.9)] in the second and the last inequalities

$$\begin{aligned} \langle (\nabla_{p^*})^* \operatorname{Reg}_j h, g \rangle_{L^p(G), L^{p^*}(G)} &= \langle \operatorname{Reg}_j h, \nabla_{p^*} g \rangle_{L^p(G, \ell_m^p), L^{p^*}(G, \ell_m^{p^*})} = \langle h, \operatorname{Reg}_j(\nabla_{p^*} g) \rangle \\ &\stackrel{(6.8)}{=} \langle h, \nabla_{p^*}(\operatorname{Reg}_j g) \rangle = \langle (\nabla_{p^*})^*(h), \operatorname{Reg}_j g \rangle = \langle \operatorname{Reg}_j(\nabla_{p^*})^*(h), g \rangle_{L^p(G), L^{p^*}(G)}, \end{aligned}$$

where here we use the bracket

$$\langle f, g \rangle_{L^p(G), L^{p^*}(G)} = \int_G f(s)g(s^{-1}) ds.$$

Note that the use of the inversion map  $G \rightarrow G, s \mapsto s^{-1}$  in the bracket simplifies [32, Theorem (14.10.9)]. By density and duality, we infer that  $(\nabla_{p^*})^* \operatorname{Reg}_j h = \operatorname{Reg}_j((\nabla_{p^*})^* h)$  which converges to  $(\nabla_{p^*})^*(h)$  in  $L^p(G)$ . ■

If  $f \in L^\infty(G)$ , we define the bounded operator

$$\pi(f): L^p(G) \oplus_p L^p(G, \ell_m^p) \rightarrow L^p(G) \oplus_p L^p(G, \ell_m^p)$$

by

$$\pi(f) \stackrel{\text{def}}{=} \begin{bmatrix} M_f & 0 \\ 0 & \tilde{M}_f \end{bmatrix}, \quad f \in L^\infty(G), \tag{6.9}$$

where the linear map  $M_f: L^p(G) \rightarrow L^p(G), g \mapsto fg$  is the multiplication operator by the function  $f$  and where

$$\tilde{M}_f \stackrel{\text{def}}{=} \operatorname{Id}_{\ell_m^p} \otimes M_f: \ell_m^p(L^p(G)) \rightarrow \ell_m^p(L^p(G)), \quad (h_1, \dots, h_m) \mapsto (fh_1, \dots, fh_m)$$

is also a multiplication operator (by the function  $(f, \dots, f)$  of  $\ell_m^\infty(L^\infty(G))$ ). Using [40, Proposition 4.10, p. 31], it is (really) easy to check that  $\pi: L^\infty(G) \rightarrow B(L^p(G) \oplus_p L^p(G, \ell_m^p))$  is an isometric homomorphism. Moreover, it is obviously continuous when the algebra  $L^\infty(G)$  is equipped with the weak\* topology and when the space  $B(L^p(G) \oplus_p L^p(G, \ell_m^p))$  is equipped with the weak operator topology. Note  $B(L^p(G) \oplus_p L^p(G, \ell_m^p))$  is a dual Banach space whose predual is the projective tensor product

$$(L^p(G) \oplus_p L^p(G, \ell_m^p)) \hat{\otimes} (L^{p^*}(G) \oplus_{p^*} L^{p^*}(G, \ell_m^{p^*})).$$

Using [12, Theorem A.2.5 (2)], it is not difficult to prove that  $\pi$  is *even* weak\* continuous when we equip the space  $B(L^p(G) \oplus_p L^p(G, \ell_m^p))$  with the weak\* topology.

**Proposition 6.4.** *Let  $G$  be a unimodular connected Lie group equipped with a family  $X$  of left-invariant Hörmander vector fields. Consider a Haar measure  $\mu_G$  on  $G$ . Suppose that  $1 < p < \infty$ .*

- (1) *We have  $(\mathcal{D}_p)^* = \mathcal{D}_p^*$ . In particular, the unbounded operator  $\mathcal{D}_2$  is selfadjoint.*
- (2) *We have*

$$\text{dom } \nabla_\infty \subset \text{Lip}_{\mathcal{D}_p}(\mathbb{L}^\infty(G)). \tag{6.10}$$

- (3) *For any  $f \in \text{dom } \nabla_\infty$ , we have*

$$\|[\mathcal{D}_p, \pi(f)]\|_{\mathbb{L}^p(G) \oplus_p \mathbb{L}^p(G, \ell_m^p) \rightarrow \mathbb{L}^p(G) \oplus_p \mathbb{L}^p(G, \ell_m^p)} = \|\nabla_\infty(f)\|_{\mathbb{L}^\infty(G, \ell_m^p)}. \tag{6.11}$$

*Proof.* (1) By definition, an element  $(z, t)$  of the Banach space  $\mathbb{L}^p(G) \oplus_p \mathbb{L}^p(G, \ell_m^p)$  belongs to  $\text{dom}(\mathcal{D}_p)^*$  if and only if there exists  $(h, k) \in \mathbb{L}^p(G) \oplus_p \mathbb{L}^p(G, \ell_m^p)$  such that for any  $(f, g) \in \text{dom } \nabla_p \oplus \text{dom}(\nabla_p^*)^*$ , we have

$$\left\langle \begin{bmatrix} 0 & (\nabla_p^*)^* \\ \nabla_p & 0 \end{bmatrix} \begin{bmatrix} f \\ g \end{bmatrix}, \begin{bmatrix} z \\ t \end{bmatrix} \right\rangle = \left\langle \begin{bmatrix} f \\ g \end{bmatrix}, \begin{bmatrix} h \\ k \end{bmatrix} \right\rangle;$$

that is,

$$\langle (\nabla_p^*)^*(g), z \rangle + \langle \nabla_p(f), t \rangle = \langle g, k \rangle + \langle f, h \rangle. \tag{6.12}$$

If  $z \in \text{dom } \nabla_p^*$  and if  $t \in \text{dom}(\nabla_p)^*$ , the latter holds with  $k = \nabla_p^*(z)$  and  $h = (\nabla_p)^*(t)$ . This proves that  $\text{dom } \nabla_p^* \oplus \text{dom}(\nabla_p)^* \subset \text{dom}(\mathcal{D}_p)^*$  and that

$$(\mathcal{D}_p)^*(z, t) = ((\nabla_p)^*(t), \nabla_p^*(z)) = \begin{bmatrix} 0 & (\nabla_p)^* \\ \nabla_p^* & 0 \end{bmatrix} \begin{bmatrix} z \\ t \end{bmatrix} \stackrel{(6.6)}{=} \mathcal{D}_p^*(z, t).$$

Conversely, if  $(z, t) \in \text{dom}(\mathcal{D}_p)^*$ , choosing  $g = 0$  in (6.12), we obtain  $t \in \text{dom}(\nabla_p)^*$ , and taking  $f = 0$ , we obtain  $z \in \text{dom } \nabla_p^*$ .

- (2) Let  $f \in C_c^\infty(G)$ . A standard calculation shows that

$$\begin{aligned} [\mathcal{D}_p, \pi(f)] &\stackrel{(6.5)(6.9)}{=} \begin{bmatrix} 0 & (\nabla_p^*)^* \\ \nabla_p & 0 \end{bmatrix} \begin{bmatrix} M_f & 0 \\ 0 & \tilde{M}_f \end{bmatrix} - \begin{bmatrix} M_f & 0 \\ 0 & \tilde{M}_f \end{bmatrix} \begin{bmatrix} 0 & (\nabla_p^*)^* \\ \nabla_p & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & (\nabla_p^*)^* \tilde{M}_f \\ \nabla_p M_f & 0 \end{bmatrix} - \begin{bmatrix} 0 & M_f (\nabla_p^*)^* \\ \tilde{M}_f \nabla_p & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & (\nabla_p^*)^* \tilde{M}_f - M_f (\nabla_p^*)^* \\ \nabla_p M_f - \tilde{M}_f \nabla_p & 0 \end{bmatrix}. \end{aligned}$$

We calculate the two non-zero entries of the commutator. For the lower left corner, if  $g \in C_c^\infty(G)$ , we have

$$\begin{aligned} (\nabla_p M_f - \tilde{M}_f \nabla_p)(g) &= \nabla_p M_f(g) - \tilde{M}_f \nabla_p(g) = \nabla_p(fg) - f \cdot \nabla_p(g) \\ &\stackrel{(6.4)}{=} g \cdot \nabla_p(f) = (gX_{1,m}(f), \dots, gX_{m,p}(f)) = M_{\nabla f} J(g), \end{aligned}$$

where  $J: L^p(G) \rightarrow \ell_m^p(L^p(G))$ ,  $g \mapsto (g, \dots, g)$  and where  $M_{\nabla(f)}$  is the multiplication operator on the  $L^p$ -space  $\ell_m^p(L^p(G))$  by  $\nabla(f)$ . For the upper right corner, note that for any  $h \in C_c^\infty(G) \otimes \ell_m^p$  and any  $g \in C_c^\infty(G)$ , we have

$$\begin{aligned} & \langle ((\nabla_{p^*})^* \tilde{M}_f - M_f(\nabla_{p^*})^*)(h), g \rangle \\ &= \langle (\nabla_{p^*})^* \tilde{M}_f(h), g \rangle - \langle M_f(\nabla_{p^*})^*(h), g \rangle \\ &= \langle \tilde{M}_f(h), \nabla_{p^*}(g) \rangle - \langle (\nabla_{p^*})^*(h), M_f(g) \rangle = \langle h, \tilde{M}_f \nabla_{p^*}(g) \rangle - \langle h, \nabla_{p^*} M_f(g) \rangle \\ &= \langle h, \tilde{M}_f \nabla_{p^*}(g) - \nabla_{p^*} M_f(g) \rangle = \langle h, f \cdot \nabla_{p^*}(g) - \nabla_{p^*}(fg) \rangle \\ &\stackrel{(6.4)}{=} -\langle h, g \cdot \nabla(f) \rangle = \langle h, -M_{\nabla f} J(g) \rangle = \langle h, M_{\nabla f} J(g) \rangle \\ &= \langle M_{\nabla f}(h), J(g) \rangle = \langle J^* M_{\nabla f}(h), g \rangle_{L^p(G), L^{p^*}(G)}. \end{aligned}$$

We conclude that

$$((\nabla_{p^*})^* \tilde{M}_f - M_f(\nabla_{p^*})^*)(h) = J^* M_{\nabla f}(h), \quad h \in C_c^\infty(G) \otimes \ell_m^p.$$

The two non-zero components of the commutator are bounded linear operators on  $C_c^\infty(G)$  and on  $C_c^\infty(G) \otimes \ell_m^p$ . We deduce that  $[\mathcal{D}_p, \pi(f)]$  is bounded on the core  $(C_c^\infty(G) \oplus \ell_m^p) \otimes C_c^\infty(G)$  of the unbounded operator  $\mathcal{D}_p$  (here we use Lemma 6.3). By [8, Proposition 26.5], this operator extends to a bounded operator on the Banach space  $L^p(G) \oplus_p L^p(G, \ell_m^p)$ . Hence,  $C_c^\infty(G)$  is a subset of  $\text{Lip}_{\mathcal{D}_p}(L^\infty(G))$ .

If  $(g, h) \in \text{dom } \mathcal{D}_p$  and  $f \in C_c^\infty(G)$ , we have in addition

$$\begin{aligned} \|M_{\nabla f} J\|_{L^p(G) \rightarrow \ell_m^p(L^p(G))} &= \sup_{\|g\|_{L^p(G)}=1} \|M_{\nabla(f)} J(g)\|_{\ell_m^p(L^p(G))} \\ &= \sup_{\|g\|_{L^p(G)}=1} \|(X_{1,p}(f)g, \dots, X_{m,p}(f)g)\|_{\ell_m^p(L^p(G))} \\ &= \sup_{\|g\|_{L^p(G)}=1} \|(X_{1,p}(f)g, \dots, X_{m,p}(f)g)\|_{L^p(G, \ell_m^p)} \\ &= \sup_{\|g\|_{L^p(G)}=1} \left( \int_G |(X_{1,p}f)(s)g(s)|^p + \dots + |(X_{m,p}f)(s)g(s)|^p \, d\mu_G(s) \right)^{\frac{1}{p}} \\ &= \sup_{\|g\|_{L^p(G)}=1} \left( \int_G [|(X_{1,p}f)(s)|^p + \dots + |(X_{m,p}f)(s)|^p] |g(s)|^p \, d\mu_G(s) \right)^{\frac{1}{p}} \\ &= \sup_{\|h\|_{L^1(G)}=1, h \geq 0} \left( \int_G [|(X_{1,p}f)(s)|^p + \dots + |(X_{m,p}f)(s)|^p] h(s) \, d\mu_G(s) \right)^{\frac{1}{p}} \\ &= \left( \sup_{\|h\|_{L^1(G)}=1, h \geq 0} \langle \|\nabla f\|_{\ell_m^p}^p, h \rangle_{L^\infty(G), L^1(G)} \right)^{\frac{1}{p}} \\ &= \|\nabla f\|_{\ell_m^p}^p \|1\|_{L^\infty(G)}^{\frac{1}{p}} = \|\nabla f\|_{L^\infty(G, \ell_m^p)}. \end{aligned}$$

By duality, the second non-null entry of the commutator has the same norm. So we have proved (6.11) in the case where  $f \in C_c^\infty(G)$ .

Let  $f \in \text{dom } \nabla_\infty$ . Since  $C_c^\infty(G)$  is a weak\* core of the operator  $\nabla_\infty$ , we can consider a net  $(f_j)$  in  $C_c^\infty(G)$  such that  $f_j \rightarrow f$  and  $\nabla_\infty(f_j) \rightarrow \nabla_\infty(f)$  both for the weak\* topology of  $L^\infty(G)$ . By [8, Lemma 1.6], we can suppose that the nets  $(f_j)$  and  $(\nabla_\infty(f_j))$  are bounded. By [8, Proposition 5.11 (4)], we deduce that  $f \in \text{Lip}_{\mathcal{D}_p}(L^\infty(G))$ . By continuity of  $\pi$ , note that  $\pi(f_j) \rightarrow \pi(f)$  for the weak operator topology. For any  $\xi \in \text{dom } \mathcal{D}_p$  and any  $\zeta \in \text{dom}(\mathcal{D}_p)^*$ , we have

$$\begin{aligned} & \langle [\mathcal{D}_p, \pi(f_j)]\xi, \zeta \rangle_{L^p(G) \oplus_p L^p(G, \ell_m^p), L^p(G) \oplus_p L^p(G, \ell_m^p)} = \langle (\mathcal{D}_p \pi(f_j) - \pi(f_j) \mathcal{D}_p)\xi, \zeta \rangle \\ & = \langle \mathcal{D}_p \pi(f_j)\xi, \zeta \rangle - \langle \pi(f_j) \mathcal{D}_p \xi, \zeta \rangle = \langle \pi(f_j)\xi, (\mathcal{D}_p)^*\zeta \rangle - \langle \pi(f_j) \mathcal{D}_p \xi, \zeta \rangle \\ & \xrightarrow{j} \langle \pi(f)\xi, (\mathcal{D}_p)^*\zeta \rangle - \langle \pi(f) \mathcal{D}_p \xi, \zeta \rangle = \langle [\mathcal{D}_p, \pi(f)]\xi, \zeta \rangle. \end{aligned}$$

The net  $([\mathcal{D}_p, \pi(f_j)])$  is bounded since

$$\|[\mathcal{D}_p, \pi(f_j)]\|_{p \rightarrow p} \stackrel{(6.11)}{=} \|\nabla_\infty(f_j)\|_{L^\infty(G, \ell_m^p)} \lesssim_{m,p} \|\nabla_\infty(f_j)\|_{L^\infty(G, \ell_m^\infty)} \lesssim 1.$$

We deduce that the net  $([\mathcal{D}_p, \pi(f_j)])$  converges to  $[\mathcal{D}_p, \pi(f)]$  for the weak operator topology by a “net version” of [53, Lemma 3.6, p. 151]. Furthermore, it is (really) easy to check that  $M_{\nabla_\infty(f_j)}J \rightarrow M_{\nabla_\infty(f)}J$  and  $-\mathbb{E}M_{\nabla_\infty(f_j)} \rightarrow -\mathbb{E}M_{\nabla_\infty(f)}$  both for the weak operator topology. Indeed, recall that the composition of operators is separately continuous for the weak operator topology. By uniqueness of the limit, we deduce that the commutator is given by the same formula that in the case of elements of  $C_c^\infty(G)$ . From here, we obtain (6.11) as before. ■

**Remark 6.5.** The inclusion (6.10) is probably an equality. We leave this intriguing question open. We sketch an incomplete proof. Let  $f$  an element of  $\text{Lip}_{\mathcal{D}_p}(L^\infty(G))$ . We consider a Dirac net  $(\varphi_j)$  of functions of  $C^\infty(G)$ . For any  $j$ , we let  $f_j \stackrel{\text{def}}{=} \text{Reg}_j f$ . By an obvious “net version” of [32, Theorem (14.11.1)], the net  $(f_j)$  converges to  $f$  in  $L^\infty(G)$  for the weak\* topology. The point is to prove that  $(\nabla_\infty f_j)$  is a bounded net. If it is true, using Banach–Alaoglu theorem, we can suppose that  $\nabla_\infty(f_j) \rightarrow g$  for the weak\* topology for some function  $g \in L^\infty(G)$ . Since the graph of the unbounded operator  $\nabla_\infty$  is weak\* closed, we would conclude that  $f$  belongs to the subspace  $\text{dom } \nabla_\infty$ .

For the proof of the boundedness of the net, the writing

$$\|\nabla_\infty f_j\|_{L^\infty(G, \ell_m^\infty)} \approx_{m,p} \|\nabla_\infty f_j\|_{L^\infty(G, \ell_m^p)} \stackrel{(6.11)}{=} \sup_{\|\xi\| \leq 1, \|\eta\| \leq 1} |([\mathcal{D}_p, \pi(f_j)]\xi, \eta)|$$

and maybe [32, Theorem (14.10.9)] could be useful.

**Theorem 6.6.** *Let  $G$  be a compact connected Lie group equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields and consider the normalized Haar measure  $\mu_G$  on  $G$ . The triple  $(C(G), L^2(G) \oplus_2 L^2(G, \ell_m^2), \mathcal{D}_2)$  is a compact spectral triple.*



*Proof.* Here, we use the notation  $\nabla \stackrel{\text{def}}{=} \nabla_2$ . On the Hilbert space  $(\text{Ker } \nabla^* \nabla)^\perp \oplus_2 (\text{Ker } \nabla \nabla^*)^\perp$ , we have

$$|\not{D}_2|^{-1} = (\not{D}_2^2)^{-\frac{1}{2}} \stackrel{(6.7)}{=} \begin{bmatrix} \nabla^* \nabla & 0 \\ 0 & \nabla \nabla^* \end{bmatrix}^{-\frac{1}{2}} = \begin{bmatrix} (\nabla^* \nabla)^{-\frac{1}{2}} & 0 \\ 0 & (\nabla \nabla^*)^{-\frac{1}{2}} \end{bmatrix}.$$

By Theorem 2.1, we know that the operators  $\nabla^* \nabla|_{(\text{Ker } \nabla)^\perp}$  and  $\nabla \nabla^*|_{(\text{Ker } \nabla^*)^\perp}$  are unitarily equivalent. Moreover, we have

$$(\text{Ker } \nabla)^\perp \stackrel{(2.5)}{=} \overline{\text{Ran } \nabla^*} \stackrel{(2.6)}{=} \overline{\text{Ran } \nabla^* \nabla} \stackrel{(2.5)}{=} (\text{Ker } \nabla^* \nabla)^\perp$$

and

$$(\text{Ker } \nabla \nabla^*)^\perp \stackrel{(2.6)}{=} (\text{Ker } \nabla^*)^\perp.$$

Consequently,

$$(\nabla^* \nabla)^{-\frac{1}{2}}|_{(\text{Ker } \nabla^* \nabla)^\perp} \quad \text{and} \quad (\nabla \nabla^*)^{-\frac{1}{2}}|_{(\text{Ker } \nabla \nabla^*)^\perp}$$

are also unitarily equivalent. By Lemma 4.5, the operator

$$(\nabla^* \nabla)^{-\frac{1}{2}} = \Delta_2^{-\frac{1}{2}} : \overline{\text{Ran } \Delta_2} \rightarrow \overline{\text{Ran } \Delta_2}$$

is compact (the square root does not change the compactness by [85, Lemma 9.3]) on  $\overline{\text{Ran } \Delta_2} \stackrel{(2.5)}{=} (\text{Ker } \nabla^* \nabla)^\perp$ . Hence, the operator  $(\nabla \nabla^*)^{-\frac{1}{2}}|_{(\text{Ker } \nabla \nabla^*)^\perp}$  is also compact. We conclude that the operator  $|\not{D}_2|^{-1}$  is compact. ■

**Remark 6.7.** Recall that a compact spectral triple  $(A, H, \not{D})$  is even if there exists a selfadjoint unitary operator  $\gamma: H \rightarrow H$  such that  $\gamma \not{D} = -\not{D} \gamma$  and  $\gamma \pi(a) = \pi(a) \gamma$  for any  $a \in A$ . Note that the spectral triple  $(C(G), L^2(G) \oplus_2 L^2(G, \ell_m^2), \not{D})$  is even. Indeed, the Hodge–Dirac operator  $\not{D}_p$  anti-commutes with the involution

$$\gamma_p \stackrel{\text{def}}{=} \begin{bmatrix} -\text{Id}_{L^p(G)} & 0 \\ 0 & \text{Id}_{L^p(G, \ell_m^p)} \end{bmatrix} : L^p(G) \oplus_p L^p(G, \ell_m^p) \rightarrow L^p(G) \oplus_p L^p(G, \ell_m^p)$$

(which is selfadjoint if  $p = 2$ ), since

$$\begin{aligned} \not{D}_p \gamma_p + \gamma_p \not{D}_p &\stackrel{(6.5)}{=} \begin{bmatrix} 0 & (\nabla_p^*)^* \\ \nabla_p & 0 \end{bmatrix} \begin{bmatrix} -\text{Id} & 0 \\ 0 & \text{Id} \end{bmatrix} + \begin{bmatrix} -\text{Id} & 0 \\ 0 & \text{Id} \end{bmatrix} \begin{bmatrix} 0 & (\nabla_p^*)^* \\ \nabla_p & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & (\nabla_p^*)^* \\ -\nabla_p & 0 \end{bmatrix} + \begin{bmatrix} 0 & -(\nabla_p^*)^* \\ \nabla_p & 0 \end{bmatrix} = 0. \end{aligned}$$

Moreover, for any  $f \in L^\infty(G)$ , we have

$$\begin{aligned} \gamma_p \pi(f) &\stackrel{(6.9)}{=} \begin{bmatrix} -\text{Id} & 0 \\ 0 & \text{Id} \end{bmatrix} \begin{bmatrix} M_f & 0 \\ 0 & \tilde{M}_f \end{bmatrix} = \begin{bmatrix} -M_f & 0 \\ 0 & \tilde{M}_f \end{bmatrix} \\ &= \begin{bmatrix} M_f & 0 \\ 0 & \tilde{M}_f \end{bmatrix} \begin{bmatrix} -\text{Id} & 0 \\ 0 & \text{Id} \end{bmatrix} \stackrel{(6.9)}{=} \pi(f) \gamma_p. \end{aligned}$$

In the sequel, if  $1 \leq p < \infty$  and if  $H$  is a Hilbert space, we use the notation  $S^p(H)$  for the space of the compact operators  $T: H \rightarrow H$  such that

$$\|T\|_{S^p(H)} \stackrel{\text{def}}{=} (\text{Tr} |T|^p)^{\frac{1}{p}} < \infty.$$

Moreover, recall that the local dimension  $d$  of  $(G, X)$  is defined in (3.7).

**Proposition 6.8.** *Assume that  $G$  is compact. If  $\alpha > d$ , the operator  $|\mathcal{D}|^{-\alpha}$  is trace-class.*

*Proof.* We consider the canonical projection  $Q: L^2(G) \rightarrow L^2(G)$  on  $L^2_0(G)$ . We have  $Q = \text{Id} - \mathbb{E}$ , where the conditional expectation  $\mathbb{E}$  is defined in Section 2 by  $\mathbb{E}(f) = (\int_G f)1$ . For any  $t > 0$  and any function  $f \in L^2(G)$ , we have

$$T_t Qf = T_t(\text{Id} - \mathbb{E})f = T_t f - T_t \mathbb{E}f = K_t * f - \int_G f. \tag{6.13}$$

For almost all  $s \in G$ , we infer that using the translation invariance of the Haar measure

$$\begin{aligned} T_t Qf(s) &\stackrel{(6.13)}{=} (K_t * f)(s) - \int_G f \stackrel{(3.1)}{=} \int_G K_t(r) f(r^{-1}s) \, d\mu_G(r) - \int_G f(r) \, d\mu_G(r) \\ &= \int_G [K_t(r) - 1] f(r^{-1}s) \, d\mu_G(r). \end{aligned}$$

We conclude that  $T_t Q: L^2(G) \rightarrow L^2(G)$  is a convolution operator by the function  $K_t - 1$ .

For any  $t > 0$ , we have used [51, Exercise 2.8.38 (ii), p. 170]

$$\begin{aligned} \|T_t\|_{S^1(L^2_0(G))} &= \|T_{\frac{t}{2}}\|_{S^1(L^2_0(G))}^2 = \|T_{\frac{t}{2}}\|_{S^2(L^2_0(G))}^2 = \|T_{\frac{t}{2}} Q\|_{S^2(L^2(G))}^2 \\ &= \|K_{\frac{t}{2}} - 1\|_{L^2(G)}^2 = \int_G |(K_{\frac{t}{2}} - 1)(r)|^2 \, d\mu_G(r) \end{aligned} \tag{6.14}$$

for the third equality, consider an orthonormal basis  $(e_i)_{i \in I}$  of the Hilbert space  $L^2(G)$  adapted to the closed subspace  $L^2_0(G)$  and use the equality

$$\|T\|_{S^2(L^2(G))} = \left( \sum_{i \in I} \|T(e_i)\|_{L^2(G)}^2 \right)^{\frac{1}{2}}.$$

Now, by translation invariance of the Haar measure and Dunford–Pettis theorem, we obtain

$$\begin{aligned} \|T_t\|_{S^1(L^2_0(G))} &= \text{esssup}_{s \in G} \int_G |(K_{\frac{t}{2}} - 1)(sr^{-1})|^2 \, d\mu_G(r) \\ &\stackrel{(2.3)}{=} \|T_{\frac{t}{2}} Q\|_{L^2(G) \rightarrow L^\infty(G)}^2 = \|T_{\frac{t}{2}}\|_{L^2_0(G) \rightarrow L^\infty(G)}^2. \end{aligned}$$

By interpolation, we have by (2.10) combined with (4.8) the estimate

$$\|T_{\frac{t}{2}}\|_{L^2_0(G) \rightarrow L^\infty(G)} \lesssim \frac{1}{t^{\frac{d}{4}}}, \quad 0 < t \leq 1.$$

We infer that

$$\|T_t\|_{S^1(L^2_0(G))} \lesssim \frac{1}{t^{\frac{d}{2}}}, \quad 0 < t \leq 1. \tag{6.15}$$

Now, if  $t \geq 1$ , we have since  $G$  is compact

$$\begin{aligned} \|T_t\|_{S^1(L^2_0(G))} &\stackrel{(6.14)}{=} \|K_{\frac{t}{2}} - 1\|_{L^2(G)}^2 \leq \|K_{\frac{t}{2}} - 1\|_{L^\infty(G)}^2 \\ &\stackrel{(2.2)}{=} \|T_{\frac{t}{2}}\mathcal{Q}\|_{L^1(G) \rightarrow L^\infty(G)}^2 = \|T_{\frac{t}{2}}\|_{L^1_0(G) \rightarrow L^\infty(G)}^2. \end{aligned}$$

With the estimate (4.11), we conclude that

$$\|T_t\|_{S^1(L^2_0(G))} \lesssim e^{-2\omega t}, \quad t \geq 1. \tag{6.16}$$

Observe that the map is  $\mathbb{R}^+ \mapsto B(L^2(G))$ ,  $t \mapsto T_t$  is strong operator continuous hence, weak operator continuous. Moreover, if  $\alpha > d$ , we have

$$\begin{aligned} \int_0^\infty t^{\frac{\alpha}{2}-1} \|T_t\|_{S^1(L^2_0(G))} dt &= \int_0^1 t^{\frac{\alpha}{2}-1} \|T_t\|_{S^1(L^2_0(G))} dt + \int_1^\infty t^{\frac{\alpha}{2}-1} \|T_t\|_{S^1(L^2_0(G))} dt \\ &\stackrel{(6.15)}{=} \int_0^1 t^{\frac{\alpha}{2}-1-\frac{d}{2}} dt + \int_1^\infty t^{\frac{\alpha}{2}-1} e^{-2\omega t} dt < \infty. \end{aligned}$$

By [82, Lemma 2.3.2], we deduce that the operator

$$\int_0^\infty t^{\frac{\alpha}{2}-1} T_t dt$$

acting on the Hilbert space  $L^2_0(G)$  is well defined and trace-class. Furthermore, we have

$$\|T_t\|_{L^2_0(G) \rightarrow L^2_0(G)} \stackrel{(6.16)}{\lesssim} e^{-2\omega t} \quad \text{if } t \geq 1,$$

that means that  $(T_t)_{t \geq 0}$  is an exponentially stable semigroup on  $L^2_0(G)$ . Consequently, we know by [48, Corollary 3.3.6] that

$$\Delta^{-\frac{\alpha}{2}} = \frac{1}{\Gamma(\frac{\alpha}{2})} \int_0^\infty t^{\frac{\alpha}{2}-1} T_t dt.$$

We obtain that if  $\alpha > d$ , then the operator  $\Delta^{-\frac{\alpha}{2}}$  is trace-class. The operator  $(\nabla\nabla^*)^{-\alpha}$  is also trace-class since it is unitarily equivalent to  $\Delta^{-\frac{\alpha}{2}}$ , as observed in the proof of Theorem 6.6. Finally, note that

$$|\mathcal{D}_2|^{-\alpha} \stackrel{(6.7)}{=} \begin{bmatrix} \Delta_2^{-\frac{\alpha}{2}} & 0 \\ 0 & (\nabla\nabla^*)^{-\frac{\alpha}{2}} \end{bmatrix}.$$

We conclude that the operator  $|\mathcal{D}_2|^{-\alpha}$  is trace-class if  $\alpha > d$ . ■

**Theorem 6.9.** *Assume that  $G$  is compact. The spectral dimension (6.2) of the spectral triple  $(C(G), L^2(G) \oplus L^2(G, \ell_m^2), \mathcal{D}_2)$  is equal to the local dimension  $d$  of  $(G, X)$ .*

*Proof.* Note Proposition 6.8. Now, suppose that the operator  $|\mathcal{D}|^{-\alpha}$  is trace-class. By Lemma 6.2, we have

$$\text{Tr } e^{-t\mathcal{D}^2} \lesssim \frac{1}{t^{\frac{\alpha}{2}}} \text{ for any } t > 0.$$

Using [34, Proposition II.3.1, p. 20], the relation  $K_t = \check{K}_t$  in the second equality, [51, Exercise 2.8.38 (ii), p. 170] in the fourth equality, the selfadjointness of  $T_t$ , and finally (6.7) in the last inequality, we deduce that

$$\begin{aligned} K_t(e) &= \int_G K_{\frac{t}{2}}(r) K_{\frac{t}{2}}(r^{-1}) \, d\mu_G(r) = \int_G K_{\frac{t}{2}}^2 = \|K_{\frac{t}{2}}\|_{L^2(G)}^2 = \|T_{\frac{t}{2}}\|_{S^2(L^2(G))}^2 \\ &= \|T_{\frac{t}{2}}\|_{S^1(L^2(G))}^2 = \|T_t\|_{S^1(L^2(G))} = \text{Tr } T_t = \text{Tr } e^{-t\Delta_2} \leq \frac{1}{t^{\frac{\alpha}{2}}}. \end{aligned}$$

By [87, (3), p. 113] (see also [34, p. 174] for a more general statement for selfadjoint subelliptic operators), we have

$$\frac{1}{t^{\frac{\alpha}{2}}} \stackrel{(3.7)}{\approx} \frac{1}{V(\sqrt{t})} \lesssim K_t(e), \quad 0 < t \leq 1.$$

We conclude that  $\frac{1}{t^{\frac{\alpha}{2}}} \lesssim \frac{1}{t^{\frac{\alpha}{2}}}$  for any  $0 < t \leq 1$  and consequently  $\alpha \geq d$ . ■

### 7. Some remarks on Carnot–Carathéodory distances

Let  $G$  be a connected unimodular Lie group equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields. In this section, we will show in Theorem 7.4 that the Connes spectral pseudo-distance associated to our Hodge–Dirac operator allows us to recover the Carnot–Carathéodory distance. If  $(A, Y, \mathcal{D})$  is a triple as that precedes (6.1), recall that it is defined by

$$\text{dist}_{\mathcal{D}}(\varphi, \psi) \stackrel{\text{def}}{=} \sup\{|\varphi(a) - \psi(a)| : a \in \text{Lip}_{\mathcal{D}}(A) \text{ and } \|[\mathcal{D}, \pi(a)]\| \leq 1\}, \quad (7.1)$$

where  $\varphi$  and  $\psi$  are two states of the algebra  $A$  and where  $\text{Lip}_{\mathcal{D}}(A)$  is defined in (6.1). The term pseudo-metric is used since  $\text{dist}(\varphi, \psi)$  is not necessarily finite. In general,  $\text{Lip}_{\mathcal{D}}(A)$  is unknown, and we replace this space by a dense subset of  $A$  (or a weak\* dense subset if  $A$  is a dual space) which is contained in  $\text{Lip}_{\mathcal{D}}(A)$ . See [60, 70, 73] for more information.

In [76, Lemma 2.3, p. 265] and [34, p. 24], the following formula is stated for the Carnot–Carathéodory distance, i.e., the case  $p = 2$  of (3.3). For any  $s, s' \in G$ , it is written that

$$\text{dist}_{\text{CC}}(s, s') \stackrel{\text{def}}{=} \sup\{|f(s) - f(s')| : f \in C_c^\infty(G), \|\nabla f\|_{L^\infty(G, \ell_m^2)} \leq 1\}. \quad (7.2)$$

We will see that this formula is strongly related to the distance (7.1) in our setting. Unfortunately, we are unable to understand the sketched proof. The writings “Therefore” and

“by a slight modification of the  $\psi_n$  one can arrange” of [76, Lemma 2.3, p. 265] are obscure to us. Moreover, the same reference [34, p. 24] says without proof that we can replace the space  $C_c^\infty(G)$  by the subspace  $C_c^\infty(G, \mathbb{R})$  of real-valued compactly supported continuous functions in this formula.

Indeed, we can use the following elementary argument. We fix  $s, s' \in G$ . We write

$$f(s) - f(s') = |f(s) - f(s')|e^{i\theta} \quad \text{for some } \theta \in \mathbb{R}.$$

We consider the real-valued function  $\tilde{f} \stackrel{\text{def}}{=} \frac{1}{2}[f e^{-i\theta} + \bar{f} e^{i\theta}]$ . We have

$$\|\nabla \tilde{f}\|_{L^\infty(G, \ell_m^2)} = \frac{1}{2} \|e^{-i\theta} \nabla(f) + e^{-i\theta} \nabla(\bar{f})\|_{L^\infty(G, \ell_m^2)} \leq \|\nabla f\|_{L^\infty(G, \ell_m^2)}$$

and

$$\begin{aligned} |\tilde{f}(s) - \tilde{f}(s')| &= \frac{1}{2} |f(s)e^{-i\theta} + \bar{f}(s)e^{i\theta} - f(s')e^{-i\theta} - \bar{f}(s')e^{i\theta}| \\ &= \frac{1}{2} |e^{-i\theta}[f(s) - f(s')] + e^{i\theta}[\bar{f}(s) - \bar{f}(s')]| \\ &= \frac{1}{2} (|f(s) - f(s')| + |\overline{f(s) - f(s')}|) = |f(s) - f(s')|. \end{aligned}$$

Now, we introduce the following definition.

**Definition 7.1.** Suppose that  $1 < p < \infty$ . Let  $f: G \rightarrow \mathbb{C}$  be a function. The number

$$\text{Lip}_{\text{CC}}^p(f) \stackrel{\text{def}}{=} \sup \left\{ \frac{|f(s) - f(s')|}{\text{dist}_{\text{CC}}^p(s, s')} : s, s' \in G, s \neq s' \right\} \tag{7.3}$$

of  $[0, \infty]$  is called the  $p$ -Carnot–Carathéodory–Lipschitz constant of  $f$ . If  $\text{Lip}_{\text{CC}}^p(f)$  is finite, we call  $f$  a  $p$ -Carnot–Carathéodory–Lipschitz function.

In [10, Proposition 2.5 (i)], it is stated that the domain  $\text{dom } \nabla_\infty$  is the space of the equivalence classes of bounded 2-Carnot–Carathéodory-Lipschitz function on  $G$ . Here, we complete this fact, and we give a variant of [10, Proposition 2.5].

**Lemma 7.2.** Suppose that  $1 < p < \infty$ . Then, an essentially bounded function  $f: G \rightarrow \mathbb{C}$  is a  $p$ -Carnot–Carathéodory-Lipschitz function if and only if its equivalence class belongs to the space  $\text{dom } \nabla_\infty$ . In this case, we have

$$\text{Lip}_{\text{CC}}^p(f) = \|\nabla f\|_{L^\infty(G, \ell_m^{p*})}.$$

*Proof.* Suppose that  $f \in C_c^\infty(G)$ . Let  $s, s' \in G$ , and let  $\gamma: [0, 1] \mapsto G$  be an absolutely continuous path from  $s$  to  $s'$ . We have

$$\begin{aligned} f(s) - f(s') &= f(\gamma(0)) - f(\gamma(1)) \\ &= - \int_0^1 \frac{d}{dt} f(\gamma(t)) dt \stackrel{(3.4)}{=} - \int_0^1 \sum_{k=1}^m \dot{\gamma}_k(t) (X_k f)(\gamma(t)) dt. \end{aligned}$$

Consequently, using Hölder’s inequality, we obtain

$$\begin{aligned}
 |f(s) - f(s')| &\leq \int_0^1 \left| \sum_{k=1}^m \dot{\gamma}_k(t)(X_k f)(\gamma(t)) \right| dt \\
 &\leq \int_0^1 \left( \sum_{k=1}^m |\dot{\gamma}_k(t)|^p \right)^{\frac{1}{p}} \left( \sum_{k=1}^m |(X_k f)(\gamma(t))|^{p^*} \right)^{\frac{1}{p^*}} dt.
 \end{aligned}$$

We deduce that

$$\begin{aligned}
 |f(s) - f(s')| &\leq \int_0^1 \left( \sum_{k=1}^m |\dot{\gamma}_k(t)|^p \right)^{\frac{1}{p}} dt \| (X_1 f, \dots, X_m f) \|_{L^\infty(G, \ell_m^{p^*})} \\
 &\stackrel{(3.2)}{=} \| \nabla f \|_{L^\infty(G, \ell_m^{p^*})} \ell_p(\gamma).
 \end{aligned}$$

Passing to the infimum, we obtain

$$|f(s) - f(s')| \leq \| \nabla f \|_{L^\infty(G, \ell_m^{p^*})} \text{dist}_{\mathbb{C}\mathbb{C}}^p(s, s')$$

by (3.3). Consequently, we have the inequality

$$\text{Lip}_{\mathbb{C}\mathbb{C}}^p(f) \leq \| \nabla f \|_{L^\infty(G, \ell_m^{p^*})}.$$

We conclude with a regularization argument for the general case of a function  $f$  of  $\text{dom } \nabla_\infty$ .

Now, we prove the reverse inequality. Suppose that the function  $f: G \rightarrow \mathbb{C}$  is a  $p$ -Carnot–Carathéodory–Lipschitz function. Let  $\xi \in \ell_m^p$  with  $\|\xi\|_{\ell_m^p} = 1$ . For any  $1 \leq k \leq m$ , we put  $a_k \stackrel{\text{def}}{=} X_k(e)$ . Consider some  $s_0 \in G$  and the path

$$\gamma(t) \stackrel{\text{def}}{=} s_0 \exp \left( t \sum_{k=1}^m \xi_k a_k \right), \quad t \in \mathbb{R}. \tag{7.4}$$

By [33, Remark 19.8.11], for any  $t \in \mathbb{R}$ , we have

$$\dot{\gamma}(t) = \sum_{k=1}^m \xi_k X_k(\gamma(t)). \tag{7.5}$$

We infer that  $\dot{\gamma}(t)$  belongs to the subspace  $\text{span}\{X_1|_{\gamma(t)}, \dots, X_m|_{\gamma(t)}\}$  for all  $t \in \mathbb{R}$ . Moreover, the  $p$ -length of the restriction  $\gamma|[c, d]$  is given by

$$\ell_p(\gamma|[c, d]) \stackrel{(3.2)}{=} \int_c^d \left( \sum_{k=1}^m |\dot{\gamma}_k(t)|^p \right)^{\frac{1}{p}} dt \stackrel{(7.5)}{=} \int_c^d \|\xi\|_{\ell_m^p} dt = |d - c| \|\xi\|_{\ell_m^p} = |d - c|.$$

By (3.3), we deduce that

$$\text{dist}_{\mathbb{C}\mathbb{C}}^p(\gamma(c), \gamma(d)) \leq |d - c|. \tag{7.6}$$

We consider the function  $g: \mathbb{R} \rightarrow \mathbb{R}, t \mapsto f(\gamma(t))$ . Since  $f$  is a  $p$ -Carnot–Carathéodory–Lipschitz function, we have

$$\begin{aligned} |g(c) - g(d)| &= |f(\gamma(c)) - f(\gamma(d))| \\ &\leq \text{Lip}_{\text{CC}}^p(f) \text{dist}_{\text{CC}}^p(\gamma(c), \gamma(d)) \stackrel{(7.6)}{\leq} \text{Lip}_{\text{CC}}^p(f) |d - c|. \end{aligned}$$

Hence, the function  $g$  is a Lipschitz function on  $\mathbb{R}$ , hence differentiable almost everywhere. It is left to the reader to show that  $X_j f$  exists almost everywhere on  $G$ .

If  $t > 0$ , using the notation

$$s_t \stackrel{\text{def}}{=} \exp \left( t \sum_{k=1}^m \xi_k a_k \right),$$

we deduce that

$$\text{Lip}_{\text{CC}}^p(f) \stackrel{(7.3)}{\geq} \frac{|f(\gamma(t)) - f(s_0)|}{\text{dist}_{\text{CC}}^p(\gamma(t), s_0)} \stackrel{(7.6)}{\geq} \frac{|f(\gamma(t)) - f(s_0)|}{t} = \left| \frac{1}{t} ((\rho_{s_t} - \text{Id})f)(s_0) \right|,$$

where  $\rho$  is the right regular representation of  $G$ . Consequently, for any  $t > 0$ , we obtain

$$\left\| \frac{1}{t} (\rho_{s_t} - \text{Id})f \right\|_{L^\infty(G)} \leq \text{Lip}_{\text{CC}}^p(f). \tag{7.7}$$

Now,

$$\left| \frac{1}{t} ((\rho_{s_t} - \text{Id})f)(s_0) \right| = \frac{|f(\gamma(t)) - f(s_0)|}{t}$$

converges almost everywhere when  $t \rightarrow 0$ . Using dominated convergence theorem, we conclude that  $\frac{1}{t} (\rho_{s_t} - \text{Id})f$  converges in  $L^\infty(G)$  for the weak\* topology.

Using [34, p. 14], we infer that the class of  $f$  belongs to  $\text{dom } \nabla_\infty$  and that

$$\frac{1}{t} (\rho_{s_t} - \text{Id})f \rightarrow \sum_{k=1}^m \xi_k Y_{k,\infty} f \quad \text{when } t \rightarrow 0$$

for the weak\* topology of  $L^\infty(G)$ , where  $Y_k$  is the right invariant vector field associated to the element  $a_k$ . Passing to the limit in (7.7) when  $t \rightarrow 0$ , using the weak\* lower semicontinuity of the norm [65, Theorem 2.6.14, p. 227], we obtain

$$\left\| \sum_{k=1}^m \xi_k Y_{k,\infty} f \right\|_{L^\infty(G)} \leq \text{Lip}_{\text{CC}}^p(f).$$

Since  $G$  is unimodular, we have

$$\left\| \sum_{k=1}^m \xi_k X_{k,\infty} f \right\|_{L^\infty(G)} = \left\| \sum_{k=1}^m \xi_k Y_{k,\infty} f \right\|_{L^\infty(G)}$$

(if  $I: G \rightarrow G, s \mapsto s^{-1}$  is the inversion map and  $I_*$  is its associated push-forward map on vector fields, we have  $I_* X_k = -Y_k$ ). We conclude by duality that

$$\|\nabla f\|_{L^\infty(G, \ell_m^*)} = \|(X_1 f, \dots, X_m f)\|_{L^\infty(G, \ell_m^*)} \leq \text{Lip}_{\text{CC}}^p(f). \quad \blacksquare$$

**Lemma 7.3.** *Suppose that  $1 < p < \infty$ . For any  $s, s' \in G$ , we have*

$$\text{dist}_{\text{CC}}^p(s, s') = \sup\{|f(s) - f(s')| : f \in \text{dom } \nabla_\infty, \text{Lip}_{\text{CC}}^p(f) \leq 1\}.$$

Moreover, we can replace  $\text{dom } \nabla_\infty$  by the space  $C_c^\infty(G)$ .

*Proof.* Let  $f \in \text{dom } \nabla_\infty$  with  $\text{Lip}_{\text{CC}}^p(f) \leq 1$ . For any  $s, s' \in G$ , we have by definition

$$|f(s) - f(s')| \leq \text{Lip}_{\text{CC}}^p(f) \cdot \text{dist}_{\text{CC}}^p(s, s') \leq \text{dist}_{\text{CC}}^p(s, s').$$

We deduce that

$$\sup\{|f(s) - f(s')| : f \in \text{dom } \nabla_\infty, \text{Lip}_{\text{CC}}^p(f) \leq 1\} \leq \text{dist}_{\text{CC}}^p(s, s').$$

Now, we prove the reverse inequality. We fix  $s \in G$ . We consider the function  $h: G \rightarrow \mathbb{R}$ ,  $s' \mapsto \text{dist}_{\text{CC}}^p(s, s')$ . Since  $\text{dist}_{\text{CC}}^p$  is a distance on  $G$ , we have for any  $s'' \in G$

$$|h(s') - h(s'')| = |\text{dist}_{\text{CC}}^p(s, s') - \text{dist}_{\text{CC}}^p(s, s'')| \leq \text{dist}_{\text{CC}}^p(s', s'').$$

We infer that  $h$  is  $p$ -Carnot–Carathéodory–Lipschitz function (hence its class belongs to  $\text{dom } \nabla_\infty$  by Lemma 7.2) with  $\text{Lip}_{\text{CC}}^p(h) \leq 1$ . Since

$$|h(s) - h(s')| = \text{dist}_{\text{CC}}^p(s, s'),$$

we obtain

$$\text{dist}_{\text{CC}}^p(s, s') \leq \sup\{|f(s) - f(s')| : f \in \text{dom } \nabla_\infty, \text{Lip}_{\text{CC}}^p(f) \leq 1\}.$$

For the last assertion, we use a regularization argument. We consider a Dirac net  $(\varphi_j)$  of functions of  $C^\infty(G)$ , satisfying in particular

$$\int_G \varphi_j \, d\mu_G = 1.$$

For any  $j$ , we let  $f_j \stackrel{\text{def}}{=} \varphi_j * f$ . Using the left invariance of the distance  $\text{dist}_{\text{CC}}^p$  in the third equality, we have

$$\begin{aligned} \text{Lip}_{\text{CC}}^p(f_j) &\stackrel{(7.3)}{=} \sup_{s \neq s'} \left| \frac{f_j(s) - f_j(s')}{\text{dist}_{\text{CC}}^p(s, s')} \right| \\ &\stackrel{(3.1)}{=} \sup_{s \neq s'} \left| \int_G \frac{f(t^{-1}s)\varphi_j(t) - f(t^{-1}s')\varphi_j(t)}{\text{dist}_{\text{CC}}^p(s, s')} \, d\mu_G(t) \right| \\ &= \sup_{s \neq s'} \left| \int_G \frac{f(t^{-1}s) - f(t^{-1}s')}{\text{dist}_{\text{CC}}^p(t^{-1}s, t^{-1}s')} \varphi_j(t) \, d\mu_G(t) \right| \\ &\leq \int_G \text{Lip}_{\text{CC}}^p(f) \varphi_j(t) \, d\mu_G(t) \\ &= \text{Lip}_{\text{CC}}^p(f) \int_G \varphi_j(t) \, d\mu_G(t) \leq \text{Lip}_{\text{CC}}^p(f). \end{aligned}$$

Since  $f$  is left uniformly continuous, the net  $(f_j)$  converges uniformly by [41, Proposition 2.44, p. 58] to  $f$ . The conclusion is obvious. ■



With the terminology of [60, Definition 1.8], we can interpret the end of the following result by saying that  $(C(G), L^2(G) \oplus_2 L^2(G, \ell_m^2), \mathcal{D}_2)$  is a metric spectral triple.

**Theorem 7.4.** *Let  $G$  be a connected unimodular Lie group equipped with a finite family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields. Suppose that  $1 < p < \infty$ . For any  $s, s' \in G$ , we have*

$$\begin{aligned} \text{dist}_{\text{CC}}^p(s, s') &= \sup\{|f(s) - f(s')| : f \in \text{dom } \nabla_\infty, \|\nabla f\|_{L^\infty(G, \ell_m^{p*})} \leq 1\} \\ &= \sup\{|f(s) - f(s')| : f \in \text{dom } \nabla_\infty, \|[\mathcal{D}_p, \pi(f)]\|_{p^* \rightarrow p^*} \leq 1\}. \end{aligned} \tag{7.8}$$

Moreover, we can replace  $\text{dom } \nabla_\infty$  by the space  $C_c^\infty(G)$ .

Finally, if  $G$  is in addition compact, letting

$$\|f\|_{\mathcal{D}_p} \stackrel{\text{def}}{=} \|[\mathcal{D}_p, \pi(f)]\|_{p \rightarrow p} \quad \text{for any } f \in \text{dom } \nabla_\infty,$$

then the pair  $(C(G), \|\cdot\|_{\mathcal{D}_p})$  is a Leibniz quantum compact metric space.

*Proof.* Combining Lemma 7.2 and Lemma 7.3, we obtain the first equality. The second equality is a consequence of (6.11).

Now, we prove the last sentence. By [8, Proposition 5.11 (2)], note that  $\|\cdot\|_{\mathcal{D}_p}$  is a seminorm on  $\text{Lip}_{\mathcal{D}_p}(L^\infty(G))$ , hence on the subspace  $\text{dom } \nabla_\infty$ . We put

$$\text{dom } \|\cdot\|_{\mathcal{D}_p} \stackrel{\text{def}}{=} \text{dom } \nabla_\infty \quad \text{and} \quad A \stackrel{\text{def}}{=} C(G).$$

We check the properties of Proposition 4.3. Note that with a positive answer to the question raised in Remark 6.5, we could use [8, Proposition 5.11 (3) and Remark 5.7] for some assertions.

- (1) The domain  $\text{dom } \|\cdot\|_{\mathcal{D}_p} = \text{dom } \nabla_\infty$  is clearly closed under  $f \mapsto \bar{f}$ .
- (2) Let  $f \in \text{dom } \nabla_\infty$  with  $\nabla_\infty f = 0$ . For any  $s \in G$ , we have

$$\begin{aligned} \|(\text{Id} - \lambda_s)f\|_{L^p(G)} &\stackrel{(3.5)}{\leq} \text{dist}_{\text{CC}}^{p*}(s, e) \left( \sum_{k=1}^m \|X_k f\|_{L^p(G)}^p \right)^{\frac{1}{p}} \\ &\lesssim_{m,p} \text{dist}_{\text{CC}}^{p*}(s, e) \|\nabla_\infty f\|_{L^\infty(G, \ell_m^p)} = 0. \end{aligned}$$

We deduce that the function  $f$  is constant on  $G$ . The converse is obvious. Hence, we have the equality

$$\{f \in \text{dom } \|\cdot\|_{\mathcal{D}_p} : \|f\|_{\mathcal{D}_p} = 0\} = \mathbb{C}1_{C(G)}.$$

- (3) By [8, Proposition 5.11 (1)], for any  $f, g \in \text{Lip}_{\mathcal{D}_p}(L^\infty(G))$ , we have

$$fg \in \text{Lip}_{\mathcal{D}_p}(L^\infty(G))$$

and

$$\|fg\|_{\mathcal{D}_p} \leq \|f\|_{C(G)} \|g\|_{\mathcal{D}_p} + \|f\|_{\mathcal{D}_p} \|g\|_{C(G)}.$$

(4) Let  $s_0 \in G$  be any point. The Dirac probability measure  $\delta_{s_0}$  is supported on the compact  $\{s_0\}$ . So, it is a local state. We consider the subset  $\{f \in \text{dom}\|\cdot\|_{\mathcal{D}_p} : \|f\|_{\mathcal{D}_p} \leq 1, f(s_0) = 0\}$ . By (7.8) and Lemma 7.2, this subset is equicontinuous. Furthermore, it is pointwise bounded since

$$|f(s)| = |f(s) - f(s_0)| \leq \text{dist}_{\text{CC}}^{p^*}(s_0, s) \lesssim 1, \quad s \in G$$

by continuity of the function  $s \mapsto \text{dist}_{\text{CC}}^{p^*}(s_0, s)$  on the compact  $G$ . By Arzèla–Ascoli theorem, we conclude that it is relatively compact in the space  $C(G)$ .

(5) Suppose that the net  $(f_j)$  converges to  $f$  in the space  $L^\infty(G)$  and that  $\|f_j\|_{\mathcal{D}_p} \leq 1$ ; that is,

$$\|\nabla f_j\|_{L^\infty(G, \ell_m^{p^*})} \leq 1.$$

This net converges for the weak\* topology. Consequently,  $(\nabla_\infty f_j)$  is a bounded net of  $L^\infty(G, \ell_m^{p^*})$ . Using the Banach–Alaoglu theorem, we can suppose that  $\nabla_\infty f_j \rightarrow g$  for the weak\* topology for some function  $g \in L^\infty(G, \ell_m^{p^*})$ . Since the graph of the unbounded operator  $\nabla_\infty$  is weak\* closed, we conclude that  $f$  belongs to the subspace  $\text{dom } \nabla_\infty$  and that  $g = \nabla_\infty f$ . The weak\* lower semicontinuity of the norm [65, Theorem 2.6.14, p. 227] reveals that

$$\|\nabla_\infty f\|_{L^\infty(G, \ell_m^{p^*})} \leq \liminf_j \|\nabla_\infty f_j\|_{L^\infty(G, \ell_m^{p^*})} \leq 1. \quad \blacksquare$$

**Remark 7.5.** If  $G$  is a unimodular Lie group, it seems apparent that the seminorm  $\|\cdot\|_{\mathcal{D}_p}$  can be used to define quantum *locally* compact metric spaces in the spirit of the ones of Section 5. The proof is left to the reader as an exercise.

We finish the paper by connecting our setting to the vast topic of Dirichlet forms. We refer to the books [14, 42, 62] for more information on Dirichlet forms and also to the papers [11, 20, 55, 80, 81] which are connected to our setting. Let  $\Omega$  be a connected second countable Hausdorff locally compact space, and let  $\mu$  be a positive Radon measure with support  $\Omega$ . We denote by  $\mathcal{M}(\Omega)$  the collection of all signed Radon measures on  $\Omega$ .

Recall that a Dirichlet form  $\mathcal{E}$  on  $L^2(\Omega)$  is a closed positive definite symmetric bilinear form defined on  $\text{dom } \mathcal{E} \times \text{dom } \mathcal{E}$ , where  $\text{dom } \mathcal{E}$  is a dense linear subspace of the Hilbert space  $L^2(\Omega)$ .

Beurling and Deny showed that if  $\mathcal{E}$  has no killing measure and no jumping measure, it can be written as

$$\mathcal{E}(f, g) = \int_\Omega d\Gamma(f, g), \quad f, g \in \text{dom } \mathcal{E}$$

for an  $\mathcal{M}(\Omega)$ -valued positive definite symmetric bilinear form  $\Gamma$  defined by the formula

$$\int_\Omega h d\Gamma(f, g) \stackrel{\text{def}}{=} \frac{1}{2}[\mathcal{E}(f, hg) + \mathcal{E}(g, hf) - \mathcal{E}(fg, h)] \tag{7.9}$$

for all  $f, g \in \text{dom } \mathcal{E} \cap L^\infty(\Omega)$  and  $h \in \text{dom } \mathcal{E} \cap C_c(\Omega)$ . The form  $\Gamma$  is called the carré du champ associated to  $\mathcal{E}$ . The Radon–Nikodym derivative  $\frac{d\Gamma(f, f)}{d\mu}(x)$  plays (if it exists) the role of the square of the length of the gradient of  $f \in \text{dom } \mathcal{E}$  at  $x \in \Omega$ .

An intrinsic pseudo-distance on  $X$  associated to  $\mathcal{E}$  is defined in [80, (4.1)] by

$$\text{dist}_{\mathcal{E}}(x, y) \stackrel{\text{def}}{=} \sup \left\{ |f(x) - f(y)| : f \in \text{dom } \mathcal{E} \cap C_c(\Omega), \frac{d\Gamma(f, f)}{d\mu} \leq 1 \right\}. \quad (7.10)$$

Here  $\frac{d\Gamma(f, f)}{d\mu} \leq 1$  means that  $\Gamma(f, f)$  is absolutely continuous with respect to  $\mu$  and that  $\frac{d\Gamma(f, f)}{d\mu} \leq 1$  almost everywhere. We warn the reader that there exist several variants of this distance; see [80] and [81, p. 236].

Returning to the setting of Lie groups, we can consider the symmetric bilinear form

$$\mathcal{E}(f, g) = \int_G \langle \nabla f(s), \nabla g(s) \rangle_{\ell_m^2} d\mu_G(s) \quad (7.11)$$

whose domain is the subspace

$$\text{dom } \mathcal{E} = \text{dom } X_{1,2} \cap \cdots \cap \text{dom } X_{m,2} = \text{dom } \nabla_2$$

of the Hilbert space  $L^2(G)$ . This subspace is considered in the paper [10] and the book [34] and denoted respectively by  $W'_{1,2}(G)$  and  $L'_{2,1}(G)$  (and equipped with a suitable norm). A simple computation for any  $f, g \in \text{dom } \mathcal{E} \cap L^\infty(G)$  and any  $h \in \text{dom } \mathcal{E} \cap C_c(G)$  gives

$$\begin{aligned} & \frac{1}{2} [\mathcal{E}(f, hg) + \mathcal{E}(g, hf) - \mathcal{E}(uv, h)] \\ & \stackrel{(7.11)}{=} \frac{1}{2} \int_G [\langle \nabla f(s), \nabla(hg)(s) \rangle + \langle \nabla g(s), \nabla(hf)(s) \rangle - \langle \nabla(fg)(s), \nabla h(s) \rangle] d\mu_G(s) \\ & \stackrel{(6.4)}{=} \frac{1}{2} \int_G [h(s) \langle \nabla f(s), \nabla g(s) \rangle + g(s) \langle \nabla f(s), \nabla h(s) \rangle + h(s) \langle \nabla g(s), \nabla f(s) \rangle \\ & \quad + f(s) \langle \nabla g(s), \nabla h(s) \rangle - f(s) \langle \nabla g(s), \nabla h(s) \rangle - g(s) \langle \nabla f(s), \nabla h(s) \rangle] d\mu_G(s) \\ & = \int_G h(s) \langle \nabla f(s), \nabla g(s) \rangle d\mu_G(s). \end{aligned}$$

By (7.9), we conclude (with no surprise) that

$$\frac{d\Gamma(f, g)}{d\mu_G}(s) = \langle \nabla f(s), \nabla g(s) \rangle$$

almost everywhere on  $G$ . In particular, we have the equality

$$\frac{d\Gamma(f, f)}{d\mu_G}(s) = \|\nabla f(s)\|_{\ell_m^2}^2$$

almost everywhere. In this case, the intrinsic pseudo-distance (7.10) is given by

$$\begin{aligned} \text{dist}_{\mathcal{E}}(s, s') & \stackrel{(7.10)}{=} \sup \{ |f(s) - f(s')| : f \in W'_{1,2}(G) \cap C_c(G), \|\nabla f(s)\|_{\ell_m^2} \leq 1 \text{ a.e.} \} \\ & = \sup \{ |f(s) - f(s')| : f \in W'_{1,2}(G) \cap C_c(G), \|\nabla f\|_{L^\infty(G, \ell_m^2)} \leq 1 \}, \end{aligned}$$

where  $s, s' \in G$ . Using an approximation procedure similar to the one of the proof of Lemma 7.3 left to the reader, we can conclude that

$$\text{dist}_g(s, s') = \sup\{|f(s) - f(s')| : f \in C_c^\infty(G), \|\nabla f\|_{L^\infty(G, \ell_m^p)} \leq 1\} \stackrel{(7.2)}{=} \text{dist}_{\text{CC}}(s, s');$$

i.e., we obtain the Carnot–Carathéodory distance.

**Remark 7.6.** It is possible that the result [55, Corollary 2.1] can be used to recover a part of the case  $p = 2$  of Lemma 7.2 with a very different argument.

### 8. Some open problems on functional calculus

Let  $G$  be a connected Lie group of polynomial growth (hence unimodular) equipped with a family  $(X_1, \dots, X_m)$  of left-invariant Hörmander vector fields, and consider a Haar measure  $\mu_G$  on  $G$ . We refer to [48, 49] for more information on functional calculus. Here we use the bisector

$$\Sigma_\theta^\pm \stackrel{\text{def}}{=} \Sigma_\theta \cup (-\Sigma_\theta), \quad \text{where } \Sigma_\theta^+ \stackrel{\text{def}}{=} \{z \in \mathbb{C} \setminus \{0\} : |\arg z| < \theta\}$$

for any angle  $\theta \in (0, \frac{\pi}{2})$ . We will explain why the following conjecture is very natural.

**Conjecture 8.1.** Suppose that  $1 < p < \infty$  with  $p \neq 2$ . The unbounded operator  $\mathcal{D}_p$  is bisectorial and admits a bounded  $H^\infty(\Sigma_\theta^\pm)$  functional calculus on a bisector  $\Sigma_\theta^\pm$  for some  $0 < \theta < \frac{\pi}{2}$  on the Banach space  $L^p(G) \oplus_p L^p(G, \ell_m^p)$ .

The case  $p = 2$  is of course obvious since  $\mathcal{D}_2$  is selfadjoint. The boundedness of the  $H^\infty(\Sigma_\theta^\pm)$  functional calculus of the unbounded operator  $\mathcal{D}_p$  implies the boundedness of the Riesz transforms, and this result may be thought of as a strengthening of the equivalence (3.10). Indeed, consider the function  $\text{sgn} \in H^\infty(\Sigma_\theta^\pm)$  defined by

$$\text{sgn}(z) \stackrel{\text{def}}{=} 1_{\Sigma_\theta^+}(z) - 1_{\Sigma_\theta^-}(z).$$

If the operator  $\mathcal{D}_p$  has a bounded  $H^\infty(\Sigma_\theta^\pm)$  functional calculus on  $L^p(G) \oplus_p L^p(G, \ell_m^p)$ , the operator  $\text{sgn}(\mathcal{D}_p)$  is bounded. Moreover, we have

$$|\mathcal{D}_p| = \text{sgn}(\mathcal{D}_p)\mathcal{D}_p \quad \text{and} \quad \mathcal{D}_p = \text{sgn}(\mathcal{D}_p)|\mathcal{D}_p|. \tag{8.1}$$

For any element  $\xi$  of the space  $\text{dom } \mathcal{D}_p = \text{dom } |\mathcal{D}_p|$ , we deduce that

$$\begin{aligned} \|\mathcal{D}_p(\xi)\|_{L^p(G) \oplus_p L^p(G, \ell_m^p)} &\stackrel{(8.1)}{=} \|\text{sgn}(\mathcal{D}_p)|\mathcal{D}_p(\xi)\|_{L^p(G) \oplus_p L^p(G, \ell_m^p)} \\ &\lesssim_p \||\mathcal{D}_p|(\xi)\|_{L^p(G) \oplus_p L^p(G, \ell_m^p)}, \end{aligned}$$

and similarly,

$$\begin{aligned} \||\mathcal{D}_p|(\xi)\|_{L^p(G) \oplus_p L^p(G, \ell_m^p)} &\stackrel{(8.1)}{=} \|\text{sgn}(\mathcal{D}_p)\mathcal{D}_p(\xi)\|_{L^p(G) \oplus_p L^p(G, \ell_m^p)} \\ &\lesssim_p \|\mathcal{D}_p(\xi)\|_{L^p(G) \oplus_p L^p(G, \ell_m^p)}. \end{aligned}$$

Recall that on  $L^p(G) \oplus_p L^p(G, \ell_m^p)$ , we have

$$|\not{D}_p| \stackrel{(6.7)}{=} \begin{bmatrix} \Delta_p^{\frac{1}{2}} & 0 \\ 0 & * \end{bmatrix}.$$

Using (6.5) and by restricting to elements  $\xi$  of the form  $(f, 0)$  with  $f \in \text{dom } \Delta_p^{\frac{1}{2}}$ , we obtain the desired equivalence (3.10).

**Remark 8.2.** With a positive answer to Conjecture 8.1, it is not difficult to show in the case where  $G$  is compact that the triples  $(C(G), L^p(G) \oplus_p L^p(G, \ell_m^p), \not{D}_p)$  give new examples of compact Banach spectral triples in the sense of [8, Definition 5.10].

We finish with another related conjecture.

**Conjecture 8.3.** Suppose that  $1 < p < \infty$ . If  $Y$  is an UMD Banach space, the unbounded operator  $\Delta_p \otimes \text{Id}_Y$  is sectorial and admits a bounded  $H^\infty(\Sigma_\theta)$  functional calculus with  $0 < \theta < \frac{\pi}{2}$  on the Bochner space  $L^p(G, Y)$ .

This is true for the classical Laplacian on  $\mathbb{R}^d$  by [49, Theorem 10.2.25, p. 391]. The scalar case  $Y = \mathbb{C}$  seems true by [35, Theorem 3.4]. The very interesting case where  $Y = S^p$  is a Schatten class could have applications in quantum information theory. It is apparent that [4] is related to this problem.

**Acknowledgments.** The author wishes to thank Tom ter Elst for a short discussion at the very early beginning of this work. I will also thank Frédéric Latrémolière, Jochen Glück, Peer Kunstmann, Bruno Iochum, Tommaso Bruno, and Marco Peloso for short communications.

**Funding.** This research was supported by a grant from the French National Research Agency ANR-18-CE40-0021 (HASCON Project).

## References

- [1] Y. A. Abramovich and C. D. Aliprantis, *An invitation to operator theory*. Grad. Stud. Math. 50, American Mathematical Society, Providence, RI, 2002 Zbl 1022.47001 MR 1921782
- [2] D. R. Adams and L. I. Hedberg, *Function spaces and potential theory*. Grundlehren Math. Wiss. 314, Springer, Berlin, 1996 Zbl 0834.46021 MR 1411441
- [3] G. Alexopoulos, *An application of homogenization theory to harmonic analysis: Harnack inequalities and Riesz transforms on Lie groups of polynomial growth*. *Canad. J. Math.* **44** (1992), no. 4, 691–727 Zbl 0792.22005 MR 1178564
- [4] G. Alexopoulos, *Spectral multipliers on Lie groups of polynomial growth*. *Proc. Amer. Math. Soc.* **120** (1994), no. 3, 973–979 Zbl 0794.43003 MR 1172944
- [5] W. Arendt, *Semigroups and evolution equations: Functional calculus, regularity and kernel estimates*. In *Evolutionary equations. Vol. I*, pp. 1–85, Handb. Differ. Equ., North-Holland, Amsterdam, 2004 Zbl 1082.35001 MR 2103696

- [6] W. Arendt and A. F. M. ter Elst, [Ultracontractivity and eigenvalues: Weyl's law for the Dirichlet-to-Neumann operator](#). *Integral Equations Operator Theory* **88** (2017), no. 1, 65–89 Zbl [06763768](#) MR [3655946](#)
- [7] C. Arhancet, [Spectral triples, Couhlon–Varopoulos dimension and heat kernel estimates](#). 2023, arXiv:[2209.12263](#)
- [8] C. Arhancet and C. Kriegl, [Riesz transforms, Hodge–Dirac operators and functional calculus for multipliers](#). *Lecture Notes in Math.* 2304, Springer, Cham, 2022 Zbl [1493.42001](#) MR [4424941](#)
- [9] N. Aronszajn and K. T. Smith, [Theory of Bessel potentials. I](#). *Ann. Inst. Fourier (Grenoble)* **11** (1961), 385–475 Zbl [0102.32401](#) MR [143935](#)
- [10] L. Bandara, A. F. M. ter Elst, and A. McIntosh, [Square roots of perturbed subelliptic operators on Lie groups](#). *Studia Math.* **216** (2013), no. 3, 193–217 Zbl [1291.35036](#) MR [3092423](#)
- [11] F. Bernicot and D. Frey, [Sobolev algebras through a “carré du champ” identity](#). *Proc. Edinb. Math. Soc. (2)* **61** (2018), no. 4, 1041–1054 Zbl [1427.46022](#) MR [3866517](#)
- [12] D. P. Blecher and C. Le Merdy, [Operator algebras and their modules—an operator space approach](#). *London Math. Soc. Monogr. Ser.* 30, Oxford University Press, Oxford, 2004 Zbl [1061.47002](#) MR [2111973](#)
- [13] G. Bohnke, [Algèbres de Sobolev sur certains groupes nilpotents](#). *J. Funct. Anal.* **63** (1985), no. 3, 322–343 Zbl [0574.22008](#) MR [808266](#)
- [14] N. Bouleau and F. Hirsch, [Dirichlet forms and analysis on Wiener space](#). *De Gruyter Stud. Math.* 14, Walter de Gruyter, Berlin, 1991 Zbl [0748.60046](#) MR [1133391](#)
- [15] N. Bourbaki, [Integration. II. Chapters 7–9](#). *Elem. Math*, Springer, Berlin, 2004 Zbl [1095.28002](#) MR [2098271](#)
- [16] T. Bruno, M. M. Peloso, A. Tabacco, and M. Vallarino, [Sobolev spaces on Lie groups: embedding theorems and algebra properties](#). *J. Funct. Anal.* **276** (2019), no. 10, 3014–3050 Zbl [1466.46024](#) MR [3944287](#)
- [17] A.-P. Calderón, [Lebesgue spaces of differentiable functions and distributions](#). In *Proc. Sympos. Pure Math., Vol. IV*, pp. 33–49, American Mathematical Society, Providence, RI, 1961 Zbl [0195.41103](#) MR [143037](#)
- [18] A. L. Carey, J. Phillips, and A. Rennie, [Spectral triples: examples and index theory](#). In *Noncommutative geometry and physics: renormalisation, motives, index theory*, pp. 175–265, *ESI Lect. Math. Phys.*, European Mathematical Society, Zürich, 2011 Zbl [1252.58004](#) MR [2839056](#)
- [19] E. A. Carlen, S. Kusuoka, and D. W. Stroock, [Upper bounds for symmetric Markov transition functions](#). *Ann. Inst. H. Poincaré Probab. Statist.* **23** (1987), no. 2, 245–287 Zbl [0634.60066](#) MR [898496](#)
- [20] F. Cipriani, [Noncommutative potential theory: a survey](#). *J. Geom. Phys.* **105** (2016), 25–59 Zbl [1341.31008](#) MR [3504150](#)
- [21] F. Cipriani, D. Guido, T. Isola, and J.-L. Sauvageot, [Spectral triples for the Sierpinski gasket](#). *J. Funct. Anal.* **266** (2014), no. 8, 4809–4869 Zbl [1316.28006](#) MR [3177323](#)
- [22] A. Connes, [Compact metric spaces, Fredholm modules, and hyperfiniteness](#). *Ergodic Theory Dynam. Systems* **9** (1989), no. 2, 207–220 Zbl [0718.46051](#) MR [1007407](#)
- [23] A. Connes, [Noncommutative geometry](#). *Academic Press*, San Diego, CA, 1994 Zbl [0818.46076](#) MR [1303779](#)
- [24] A. Connes and M. Marcolli, [A walk in the noncommutative garden](#). In *An invitation to noncommutative geometry*, pp. 1–128, *World Scientific*, Hackensack, NJ, 2008 Zbl [1145.14005](#) MR [2408150](#)

- [25] T. Coulhon, Dimension à l'infini d'un semi-groupe analytique. *Bull. Sci. Math.* **114** (1990), no. 4, 485–500 Zbl [0738.47032](#) MR [1077272](#)
- [26] T. Coulhon and S. Meda, Subexponential ultracontractivity and  $L^p$ - $L^q$  functional calculus. *Math. Z.* **244** (2003), no. 2, 291–308 Zbl [1058.47036](#) MR [1992540](#)
- [27] T. Coulhon, E. Russ, and V. Tardivel-Nachef, Sobolev algebras on Lie groups and Riemannian manifolds. *Amer. J. Math.* **123** (2001), no. 2, 283–342 Zbl [0990.43003](#) MR [1828225](#)
- [28] T. Coulhon and L. Saloff-Coste, Semi-groupes d'opérateurs et espaces fonctionnels sur les groupes de Lie. *J. Approx. Theory* **65** (1991), no. 2, 176–199 Zbl [0745.47030](#) MR [1104158](#)
- [29] M. Cowling and S. Meda, Harmonic analysis and ultracontractivity. *Trans. Amer. Math. Soc.* **340** (1993), no. 2, 733–752 Zbl [0798.47032](#) MR [1127154](#)
- [30] M. Cwikel and N. J. Kalton, Interpolation of compact operators by the methods of Calderón and Gustavsson–Peetre. *Proc. Edinburgh Math. Soc. (2)* **38** (1995), no. 2, 261–276 Zbl [0821.46090](#) MR [1335873](#)
- [31] E. B. Davies, *Heat kernels and spectral theory*. Cambridge Tracts in Math. 92, Cambridge University Press, Cambridge, 1989 Zbl [0699.35006](#) MR [990239](#)
- [32] J. Dieudonné, *Éléments d'analyse. Tome II: Chapitres XII à XV*. Cahiers Scientifiques, Fasc. XXXI, Gauthier-Villars, Paris, 1968 Zbl [0189.05502](#) MR [235946](#)
- [33] J. Dieudonné, *Éléments d'analyse. Tome IV. Chapitres XVIII à XX. nouveau tirage*. Cahiers Scientifiques, Fasc. 34, Gauthier-Villars, Paris, 1977 Zbl [0217.00101](#) MR [467780](#)
- [34] N. Dungey, A. F. M. ter Elst, and D. W. Robinson, *Analysis on Lie groups with polynomial growth*. Progr. Math. 214, Birkhäuser, Boston, MA, 2003 Zbl [1041.43003](#) MR [2000440](#)
- [35] X. T. Duong and D. W. Robinson, Semigroup kernels, Poisson bounds, and holomorphic functional calculus. *J. Funct. Anal.* **142** (1996), no. 1, 89–128 Zbl [0932.47013](#) MR [1419418](#)
- [36] M. Eckstein and B. Iochum, *Spectral action in noncommutative geometry*. SpringerBriefs Math. Phys. 27, Springer, Cham, 2018 Zbl [1416.81008](#) MR [3887635](#)
- [37] M. Egert, *On Kato's conjecture and mixed boundary conditions*. Ph.D. thesis, TU Darmstadt, 2015 Zbl [1336.35002](#)
- [38] T. Eisner, B. Farkas, M. Haase, and R. Nagel, *Operator theoretic aspects of Ergodic theory*. Grad. Texts in Math. 272, Springer, Cham, 2015 Zbl [1353.37002](#) MR [3410920](#)
- [39] E. Y. Emel'yanov, *Non-spectral asymptotic analysis of one-parameter operator semigroups*. Oper. Theory Adv. Appl. 173, Birkhäuser, Basel, 2007 Zbl [1117.47001](#) MR [2302602](#)
- [40] K.-J. Engel and R. Nagel, *One-parameter semigroups for linear evolution equations*. Grad. Texts in Math. 194, Springer, New York, 2000 Zbl [0952.47036](#) MR [1721989](#)
- [41] G. B. Folland, *A course in abstract harmonic analysis*. 2nd edn., Textb. Math., CRC Press, Boca Raton, FL, 2016 Zbl [1342.43001](#) MR [3444405](#)
- [42] M. Fukushima, Y. Oshima, and M. Takeda, *Dirichlet forms and symmetric Markov processes*. extended edn., De Gruyter Stud. Math. 19, Walter de Gruyter, Berlin, 2011 Zbl [1227.31001](#) MR [2778606](#)
- [43] J. M. Gracia-Bondía, J. C. Várilly, and H. Figueroa, *Elements of noncommutative geometry*. Birkhäuser Adv. Texts Basler Lehrbücher, Birkhäuser, Boston, MA, 2001 Zbl [0958.46039](#) MR [1789831](#)
- [44] L. Grafakos, *Modern Fourier analysis*. 3rd edn., Grad. Texts in Math. 250, Springer, New York, 2014 Zbl [1304.42002](#) MR [3243741](#)
- [45] A. Grigor'yan and A. Telcs, Two-sided estimates of heat kernels on metric measure spaces. *Ann. Probab.* **40** (2012), no. 3, 1212–1284 Zbl [1252.35148](#) MR [2962091](#)
- [46] A. Gulisashvili and M. A. Kon, Exact smoothing properties of Schrödinger semigroups. *Amer. J. Math.* **118** (1996), no. 6, 1215–1248 Zbl [0864.47019](#) MR [1420922](#)

- [47] B. Guo, X. Pu, and F. Huang, *Fractional partial differential equations and their numerical solutions*. World Scientific, Hackensack, NJ, 2015 Zbl 1335.35001 MR 3380802
- [48] M. Haase, *The functional calculus for sectorial operators*. Oper. Theory Adv. Appl. 169, Birkhäuser, Basel, 2006 Zbl 1101.47010 MR 2244037
- [49] T. Hytönen, J. van Neerven, M. Veraar, and L. Weis, *Analysis in Banach spaces. Vol. II: Probabilistic methods and operator theory*. Ergeb. Math. Grenzgeb. (3) 67, Springer, Cham, 2017 Zbl 1402.46002 MR 3752640
- [50] M. Junge and T. Mei, *Noncommutative Riesz transforms—a probabilistic approach*. *Amer. J. Math.* **132** (2010), no. 3, 611–680 Zbl 1200.46056 MR 2666903
- [51] R. V. Kadison and J. R. Ringrose, *Fundamentals of the theory of operator algebras. Vol. I: Elementary theory*. Grad. Stud. Math. 15, American Mathematical Society, Providence, RI, 1997 Zbl 0888.46039 MR 1468229
- [52] N. Kalton and S. Montgomery-Smith, *Interpolation of Banach spaces*. In *Handbook of the geometry of Banach spaces, Vol. 2*, pp. 1131–1175, North-Holland, Amsterdam, 2003 Zbl 1041.46012 MR 1999193
- [53] T. Kato, *Perturbation theory for linear operators*. 2nd edn., Grundlehren Math. Wiss., Band 132, Springer, Berlin, 1976 Zbl 0342.47009 MR 407617
- [54] T. Kato and G. Ponce, *Commutator estimates and the Euler and Navier–Stokes equations*. *Comm. Pure Appl. Math.* **41** (1988), no. 7, 891–907 Zbl 0671.35066 MR 951744
- [55] P. Koskela and Y. Zhou, *Geometry and analysis of Dirichlet forms*. *Adv. Math.* **231** (2012), no. 5, 2755–2801 Zbl 1253.53035 MR 2970465
- [56] P. C. Kunstmann and L. Weis, *Maximal  $L_p$ -regularity for parabolic equations, Fourier multiplier theorems and  $H^\infty$ -functional calculus*. In *Functional analytic methods for evolution equations*, pp. 65–311, Lecture Notes in Math. 1855, Springer, Berlin, 2004 Zbl 1097.47041 MR 2108959
- [57] F. Latrémolière, *Quantum locally compact metric spaces*. *J. Funct. Anal.* **264** (2013), no. 1, 362–402 Zbl 1262.46049 MR 2995712
- [58] F. Latrémolière, *The quantum Gromov–Hausdorff propinquity*. *Trans. Amer. Math. Soc.* **368** (2016), no. 1, 365–411 Zbl 1334.46056 MR 3413867
- [59] F. Latrémolière, *Quantum metric spaces and the Gromov–Hausdorff propinquity*. In *Noncommutative geometry and optimal transport*, pp. 47–133, Contemp. Math. 676, American Mathematical Society, Providence, RI, 2016 Zbl 1368.46069 MR 3578737
- [60] F. Latrémolière, *The Gromov–Hausdorff propinquity for metric spectral triples*. *Adv. Math.* **404** (2022), Part A, article no. 108393, 56 Zbl 1500.46055 MR 4411527
- [61] D. Li, *On Kato–Ponce and fractional Leibniz*. *Rev. Mat. Iberoam.* **35** (2019), no. 1, 23–100 Zbl 1412.35261 MR 3914540
- [62] Z. M. Ma and M. Röckner, *Introduction to the theory of (nonsymmetric) Dirichlet forms*. Universitext, Springer, Berlin, 1992 Zbl 0826.31001 MR 1214375
- [63] Z. Magyar, *Continuous linear representations*. North-Holland Math. Stud. 168, North-Holland, Amsterdam, 1992 Zbl 0793.22007 MR 1150050
- [64] C. Martínez Carracedo and M. Sanz Alix, *The theory of fractional powers of operators*. North-Holland Math. Stud. 187, North-Holland, Amsterdam, 2001 Zbl 0971.47011 MR 1850825
- [65] R. E. Megginson, *An introduction to Banach space theory*. Grad. Texts in Math. 183, Springer, New York, 1998 Zbl 0910.46008 MR 1650235
- [66] A. Monguzzi, M. M. Peloso, and M. Salvatori, *Fractional Laplacian, homogeneous Sobolev spaces and their realizations*. *Ann. Mat. Pura Appl. (4)* **199** (2020), no. 6, 2243–2261 Zbl 1473.46043 MR 4165680



- [67] A. Nagel, Analysis and geometry on Carnot–Carathéodory spaces. 2005, <https://people.math.wisc.edu/~nagel/2005Book.pdf>, visited on 23 September 2023
- [68] E. M. Ouhabaz, *Analysis of heat equations on domains*. London Math. Soc. Monogr. Ser. 31, Princeton University Press, Princeton, NJ, 2005 Zbl 1082.35003 MR 2124040
- [69] N. Ozawa and M. A. Rieffel, Hyperbolic group  $C^*$ -algebras and free-product  $C^*$ -algebras as compact quantum metric spaces. *Canad. J. Math.* **57** (2005), no. 5, 1056–1079 Zbl 1101.46047 MR 2164594
- [70] B. Pavlović, Defining metric spaces via operators from unital  $C^*$ -algebras. *Pacific J. Math.* **186** (1998), no. 2, 285–313 Zbl 0943.46036 MR 1663810
- [71] G. K. Pedersen, *Analysis now*. Grad. Texts in Math. 118, Springer, New York, 1989 Zbl 0668.46002 MR 971256
- [72] M. M. Peloso and M. Vallarino, Sobolev algebras on nonunimodular Lie groups. *Calc. Var. Partial Differential Equations* **57** (2018), no. 6, article no. 150, 34 Zbl 1406.46026 MR 3858833
- [73] M. A. Rieffel, Metrics on states from actions of compact groups. *Doc. Math.* **3** (1998), 215–229 Zbl 0993.46043 MR 1647515
- [74] M. A. Rieffel, Group  $C^*$ -algebras as compact quantum metric spaces. *Doc. Math.* **7** (2002), 605–651 Zbl 1031.46082 MR 2015055
- [75] M. A. Rieffel, Compact quantum metric spaces. In *Operator algebras, quantization, and non-commutative geometry*, pp. 315–330, Contemp. Math. 365, American Mathematical Society, Providence, RI, 2004 Zbl 1084.46060 MR 2106826
- [76] D. W. Robinson, *Elliptic operators and Lie groups*. Oxford Math. Monogr., Oxford University Press, New York, 1991 Zbl 0747.47030 MR 1144020
- [77] K. Schmüdgen, *An invitation to unbounded representations of  $*$ -algebras on Hilbert space*. Grad. Texts in Math. 285, Springer, Cham, 2020 Zbl 1458.47002 MR 4180683
- [78] E. M. Stein, *Singular integrals and differentiability properties of functions*. Princeton Math. Ser., No. 30, Princeton University Press, Princeton, NJ, 1970 Zbl 0207.13501 MR 290095
- [79] R. S. Strichartz, Multipliers on fractional Sobolev spaces. *J. Math. Mech.* **16** (1967), 1031–1060 Zbl 0145.38301 MR 215084
- [80] K.-T. Sturm, Analysis on local Dirichlet spaces. I. Recurrence, conservativeness and  $L^p$ -Liouville properties. *J. Reine Angew. Math.* **456** (1994), 173–196 Zbl 0806.53041 MR 1301456
- [81] K.-T. Sturm, On the geometry defined by Dirichlet forms. In *Seminar on Stochastic Analysis, Random Fields and Applications (Ascona, 1993)*, pp. 231–242, Progr. Probab. 36, Birkhäuser, Basel, 1995 Zbl 0834.58039 MR 1360279
- [82] F. Sukochev and D. Zanin, The Connes character formula for locally compact spectral triples. 2018, arXiv:1803.01551
- [83] A. F. M. ter Elst and D. W. Robinson, Subelliptic operators and Lie groups. In *Geometric analysis and applications (Canberra, 2000)*, pp. 67–84, Proc. Centre Math. Appl. Austral. Nat. Univ. 39, Australian National University, Canberra, 2001 Zbl 1129.43004 MR 1852696
- [84] B. Thaller, *The Dirac equation*. Texts Monogr. Phys., Springer, Berlin, 1992 Zbl 0765.47023 MR 1219537
- [85] J. van Neerven, *Functional analysis*. Cambridge Stud. Adv. Math. 201, Cambridge University Press, Cambridge, 2022 Zbl 1505.46002 MR 4425670
- [86] J. C. Varilly, Dirac operators and spectral geometry. Lecture notes on noncommutative geometry and quantum groups edited by P. M. Hajac, <https://www.mimuw.edu.pl/~pwit/toknotes/>, visited on 23 September 2023

- [87] N. T. Varopoulos, L. Saloff-Coste, and T. Coulhon, *Analysis and geometry on groups*. Cambridge Tracts in Math. 100, Cambridge University Press, Cambridge, 1992 Zbl [0744.43006](#) MR [1218884](#)
- [88] C. Villani, *Optimal transport: Old and new*. Grundlehren Math. Wiss. 338, Springer, Berlin, 2009 Zbl [1156.53003](#) MR [2459454](#)
- [89] X. Xiong, *Noncommutative harmonic analysis on semigroups and ultracontractivity*. *Indiana Univ. Math. J.* **66** (2017), no. 6, 1921–1947 Zbl [1392.46047](#) MR [3744814](#)

Received 12 March 2022.

**Cédric Arhancet**

Lycée Lapérouse, 2 Lices Georges Pompidou, 81000 Albi, France;  
[cedric.arhancet@protonmail.com](mailto:cedric.arhancet@protonmail.com)