

Thomas Finn · Alexandre Stauffer

Non-equilibrium multi-scale analysis and coexistence in competing first passage percolation

Received October 18, 2020; revised March 22, 2021

Abstract. The main contribution of this paper is the development of a novel approach to multi-scale analysis that we believe can be used to analyse processes with non-equilibrium dynamics. Our approach will be referred to as *multi-scale analysis with non-equilibrium feedback* and will be used to analyse a natural random growth process with competition on \mathbb{Z}^d called *first passage percolation in a hostile environment* (FPPHE) that consists of two first passage percolation processes FPP_1 and FPP_λ that compete for the occupancy of sites. Initially, FPP_1 occupies the origin and spreads through the edges of \mathbb{Z}^d at rate 1, while FPP_λ is initialised at sites called *seeds* that are distributed according to a product of Bernoulli measures of parameter $p \in (0, 1)$, where a seed remains dormant until FPP_1 or FPP_λ attempts to occupy it before then spreading through the edges of \mathbb{Z}^d at rate $\lambda > 0$. Two fundamental challenges of FPPHE that our approach is able to handle are the absence of monotonicity (for instance, adding seeds could be beneficial to FPP_1 instead of FPP_λ) and its non-equilibrium dynamics; such characteristics, for example, prevent the application of a more standard multi-scale analysis. As a consequence of our main result for FPPHE, we establish a coexistence phase for $d \geq 3$, answering an open question of Sidoravicius and Stauffer (2019). This exhibits a rare situation where a natural random competition model on \mathbb{Z}^d observes coexistence for processes with *different* speeds. Moreover, we are able to establish the stronger result that FPP_1 and FPP_λ can both occupy a *positive density* of sites with positive probability, which is in stark contrast with other competition processes.

Keywords. Competition, first passage percolation, multi-scale analysis

1. Introduction

In this paper we consider competing random spatial growth on \mathbb{Z}^d and study the phenomena of *coexistence*, in which the two types simultaneously grow without bound. Our focus is competing first passage percolation which has been studied for over twenty years, but

Thomas Finn: Department of Mathematical Sciences, University of Bath,
Bath BA2 7AY, UK; t.j.finn@bath.ac.uk

Alexandre Stauffer: Dipartimento di Matematica e Fisica, Università Roma Tre, Roma, Italy; and
Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, UK;
a.stauffer@bath.ac.uk

Mathematics Subject Classification (2020): Primary 60K35; Secondary 82C22, 82C26

many intriguing questions remain open (see Section 1.5 for more details). Understanding for what choice of parameters coexistence can occur is a fundamental question and is often difficult due to intrinsic lack of monotonicity (both processes have to grow indefinitely without blocking each other) and long-range dependencies in such models.

To overcome difficulties inherent in competing first passage percolation, we introduce a novel approach to a widely used technique known as multi-scale analysis (or multi-scale renormalisation). We refer to our approach as *multi-scale analysis with non-equilibrium feedback* and apply it to analyse a challenging growth process known as *first passage percolation in a hostile environment* (FPPHE) [27], in which two types concurrently grow inside \mathbb{Z}^d , occupying the sites of \mathbb{Z}^d while competing with each other for space. As a consequence of our main result, we establish the regime of coexistence for FPPHE on \mathbb{Z}^d that unlike similar models occurs when the two types grow at *different* rates.

Multi-scale analysis is a powerful technique that has been used to analyse a wide range of difficult processes, especially those with *slow decay of correlations*; recent examples include random interacements, dependent percolation processes, and interacting particle systems [9, 16, 22–24, 26, 28]. Despite its power, multi-scale analyses usually rely strongly on certain crucial properties, such as *stationarity* (that is, the process is *in equilibrium*) and *monotonicity*. Such properties are present in all the aforementioned works, but are not satisfied by FPPHE and several other important processes, one additional example being a non-equilibrium model for the spread of an infection among random walk particles, whose analysis is highlighted as a fundamental challenge in [23, 24]; a similar question was recently raised in [18]. We believe our idea could be applicable to analyse this and other processes with non-equilibrium dynamics.

Below we first introduce FPPHE and discuss our results for that model, and then turn to presenting the idea behind our multi-scale analysis with non-equilibrium feedback.

1.1. First passage percolation in a hostile environment (FPPHE)

FPPHE is a natural random growth process with competition introduced by Sidoravicius and Stauffer [27] to analyse a notoriously challenging random aggregation model called *multiparticle diffusion limited aggregation* (MDLA). Nonetheless, FPPHE is an interesting model to analyse in its own right, showing several phase transitions and a wide range of behaviour. In fact, most of the work in [27] is devoted to studying FPPHE, from which results about MDLA can be deduced.

In FPPHE, there are two first passage percolation processes FPP_1 and FPP_λ that compete for the occupancy of sites on \mathbb{Z}^d . Initially the FPP_1 process only occupies the origin, while, following a product of Bernoulli measures of parameter $p \in (0, 1)$ on all other sites of \mathbb{Z}^d , we place *seeds* from which the FPP_λ will eventually attempt to grow. From these initial conditions, FPPHE evolves in time as follows. The FPP_1 process spreads through the edges of \mathbb{Z}^d at rate 1. This means that, for any given edge e , when the FPP_1 process occupies for the first time one of the endpoints of e , then after a random amount of time (that is distributed as an exponential random variable of rate 1), the FPP_1 process attempts to occupy the other endpoint of e , with the attempt being successful if and only

if that endpoint is not already occupied by either process. The FPP_λ process, in turn, does not initially spread but remains dormant. When either the FPP_1 or FPP_λ process attempts to occupy a site that has a seed, the attempt fails and the seed is *activated*, meaning that the FPP_λ process then spreads from that seed through the edges of \mathbb{Z}^d at rate $\lambda > 0$. The other seeds remain dormant until they are activated. Once a site is occupied by either process, it is occupied by that process henceforth and thus FPP_1 and FPP_λ are in competition for the occupancy of sites of \mathbb{Z}^d .

Initially FPP_1 has the advantage as it is able to spread from time 0, while FPP_λ remains dormant in seeds until activated by either process. On the other hand, FPP_λ has the advantage of occupying infinitely many seeds initially and so the spread of FPP_λ from any of these seeds could allow FPP_λ to block the spread of FPP_1 . Understanding how this delicate balance is exhibited under different choices of λ and p is fundamental in the analysis of FPPHE.

Figure 1 illustrates a simulation of FPPHE in two dimensions for two different values of p and λ . We highlight that in Figure 1 (a) the small value of p and relatively large value of λ makes FPP_1 grow ‘almost radially’ from the origin, but even if λ is not very close to 1, FPP_λ manages to conquer a quite large region, highlighting the strong dependences in the process. On the other hand, when p is close to $1 - p_c^{\text{site}}$ (where $p_c^{\text{site}} = p_c^{\text{site}}(d)$ is the critical probability for independent site percolation on \mathbb{Z}^d), as in Figure 1 (b), FPP_1 gives rise to a more irregular shape as it needs to deviate from possible blocks that FPP_λ may create as it eventually occupies part of the thin set of non-seed sites. (We refer the reader to [17] for a background on percolation.) This makes regions of \mathbb{Z}^d that are close




Fig. 1. Simulation of FPPHE in two dimensions. The coloured areas represent the sites occupied by FPP_1 , with different colors representing different epochs of the evolution of FPP_1 . The white area represents the sites occupied by FPP_λ , while the light gray boundary area represents unoccupied sites, including inactive seeds.

to one another be occupied by the process at times that are far apart. For example, in Figure 1 (b), FPP_λ blocked the growth of FPP_1 to the left of the origin (the origin is close to the middle of the picture, where the black region is), forcing FPP_1 to make a long detour to reach regions to the left of the origin.

We say that FPP_1 (resp. FPP_λ) *survives* if in the limit as time goes to infinity, FPP_1 (resp. FPP_λ) occupies an infinite *connected* component of \mathbb{Z}^d . Note that FPP_λ initially has infinitely many seeds, so the requirement of a connected component is natural. If a process does not survive, we say that it *dies out*. Hence, if FPP_λ dies out, then FPP_λ is an infinite collection of connected regions, each of which is almost surely of finite size. When both processes survive, we say that *coexistence* occurs or that FPP_1 and FPP_λ *coexist*.

As at least one process must survive, there are three possible phases that FPPHE can observe. The first is the *extinction phase* where FPP_1 dies out almost surely (and thus FPP_λ survives almost surely). Note that there is always a positive probability of FPP_1 dying out,¹ thus the almost sure criterion is essential in the definition of the extinction phase. Moreover, if $p > 1 - p_c^{\text{site}}$, then for any $\lambda > 0$, FPPHE is in the extinction phase as the origin is almost surely contained in a finite cluster of non-seeds. Consequently, FPPHE is most interesting when $p < 1 - p_c^{\text{site}}$. Similarly, FPP_1 dies out almost surely if $\lambda \geq 1$ (see Section 1.4 for more details) and so we focus on the case $\lambda < 1$. The second phase is when, with positive probability, FPP_1 survives and FPP_λ dies out, which we refer to as the *strong survival phase*. Ultimately, if with positive probability both FPP_1 and FPP_λ survive, we say that FPPHE is in the *coexistence phase*. For $d = 1$ it is immediate that neither the strong survival or coexistence phase can exist.

It seems natural to believe that, as p or λ is increased, we observe a monotone transition from a regime of strong survival to a regime of extinction (possibly passing, on the way, through a regime of coexistence). The phase diagram of FPPHE, as observed via simulations, is illustrated in Figures 2(a) and 2(b). However, as counter-intuitive as it may appear, FPPHE is not a monotone process at all. To explain this, note that once the locations of the seeds and the values of the passage times² have been sampled, the evolution of FPPHE becomes deterministic. The lack of monotonicity in FPPHE corresponds to the fact that it is possible to choose the locations of the seeds and the values of the passage times such that adding a single seed can be beneficial to FPP_1 (instead of FPP_λ). This may sound counter-intuitive at first, but the idea is that the addition of a seed may slow down FPP_1 *locally*, when that seed is activated, but this slow down could have the additional effect of delaying the activation of other seeds that are further away along that direction. It is this delay on activating some seeds that could be helpful for FPP_1 , aiding its growth along nearby directions. By properly choosing the passage times, the addition of a single seed could even make FPP_1 switch from non-survival to survival. We further

¹For example, with probability $p^{2d} > 0$, every site neighbouring the origin hosts a seed, and consequently FPP_1 dies out.

²For an edge (u, v) first occupied by FPP_1 (resp., FPP_λ) at u , the passage time is the time that FPP_1 (resp., FPP_λ) then takes to attempt to occupy v from u .




Fig. 2. Possible phase diagrams for the behaviour of FPPHE, as observed via simulations, split into the cases $d = 2$ and $d \geq 3$, and the known phase diagram for $d \geq 3$. In Figures 2(a) and 2(b), the orange regions represent where we expect there to be strong survival while the white regions represent where we expect extinction. The purple region in Figure 2(b) represents the regime of coexistence. The blue regions represent the cases where the transition between the phases is not clear even from simulations. In particular, for $d = 2$, it is unclear whether there is a regime of coexistence at all, while for $d \geq 3$ it is expected that the purple region extends to the left of p'_c , though simulations are not at all conclusive. In Theorem 1.1, we prove that for $d \geq 3$ and $p < 1 - p_c^{\text{site}}$, if λ is small enough, then FPP_1 survives. A finer description of the phases is illustrated in Figure 2(c); the orange region represents the strong survival phase established in Theorem 1.3 and [27], while the purple region represents the coexistence regime given by Corollary 1.2. The white region is the extinction regimes discussed in Sections 1.1 and 1.4, while the phases in the blue region are not known.

note that in a very recent work [8], a stronger notion of non-monotonicity is established; it is proved that there are graphs for which *the probability that FPP_1 survives is not a monotone function of p and λ* (with several phase transitions taking place). Even though it is not known whether FPPHE on \mathbb{Z}^d is monotone according to the latter notion, the lack of the former notion already creates several problems for the development of a standard multi-scale analysis. The approach we develop is able to circumvent this issue, analyzing FPPHE without relying on monotonicity (in particular, we do not use sprinkling ideas).

1.2. Our results for FPPHE

Our main result for FPPHE is that if $p < 1 - p_c^{\text{site}}$ and λ is small enough, then with positive probability, FPP_1 occupies a positive density of sites, immediately giving that FPP_1 survives. We note that this result is also true in dimension 2. This result is similar in spirit to that of Sidoravicius–Stauffer [27], but the latter establishes survival of FPP_1 by first fixing λ and then setting p small enough. The difference in the order at which the parameters are chosen does make a difference in the overall behaviour of FPPHE, as illustrated in Figure 1 and more thoroughly explained in Section 1.4.

Theorem 1.1. *Suppose $d \geq 2$ and $p < 1 - p_c^{\text{site}}$. There exists a constant $\tilde{\lambda} = \tilde{\lambda}(d, p) > 0$ such that if $\lambda \in (0, \tilde{\lambda})$, then with positive probability, FPP_1 occupies a positive density of sites.*

From Theorem 1.1, coexistence in dimension $d \geq 3$ follows almost immediately. It is well known that $p_c^{\text{site}} < 1/2$ for $d \geq 3$ (cf. Campanino and Russo [6, Theorem 4.1]), and so the inequality $p_c^{\text{site}} < 1 - p_c^{\text{site}}$ holds. In this scenario, if $p \in (p_c^{\text{site}}, 1 - p_c^{\text{site}})$, then there is almost surely an infinite connected component of seeds as well as an infinite connected component of non-seeds. An infinite connected component of seeds guarantees survival for FPP_λ regardless of the value of λ , while FPP_λ has a positive density of sites for all $p > 0$. With this in mind, for $p \in (p_c^{\text{site}}, 1 - p_c^{\text{site}})$, one can use Theorem 1.1 to set λ small enough so that FPP_1 and FPP_λ coexist and both occupy a positive density of sites.

In fact, the situation is more subtle. Through an enhancement argument [2, 5], we can deduce a regime of coexistence just below p_c^{site} . More precisely, after placing seeds with density p , we label as *filled seeds* the set of seeds and sites that are disconnected from infinity by seeds. Let $\{o \xrightarrow{f} \infty\}$ be the event that a neighbour of the origin is contained in an infinite connected component of filled seeds. Define the critical probability of percolation for the process given by filled seeds as

$$p'_c = \sup \{p \in [0, 1] : \mathbb{P}_{1-p}(o \xrightarrow{f} \infty) = 0\},$$

where \mathbb{P}_{1-p} is the probability measure³ induced by the placing seeds with density p . Note that such a critical probability exists as the event of filled seeds percolating is monotone in p . Clearly $p'_c \leq p_c^{\text{site}}$ and if $p > p'_c$, then FPP_λ occupies an infinite connected component of sites almost surely. It is known that $p'_c < p_c^{\text{site}}$ for $d = 2, 3$ by [2, 5] and this strict inequality is conjectured to hold for higher dimensions.

Corollary 1.2. *Suppose $d \geq 3$, $p \in (p'_c, 1 - p_c^{\text{site}})$ and $\tilde{\lambda}$ is as given in Theorem 1.1. If $\lambda \in (0, \tilde{\lambda})$, then with positive probability, FPP_1 and FPP_λ coexist and both occupy a positive density of sites.*

We believe the above result is the first example of a random competition model on \mathbb{Z}^d where processes of different speeds coexist.⁴ In fact, we prove the stronger result that two processes of different speeds can both occupy a *positive density* of sites with positive probability. This is in stark contrast to other natural random competition models on \mathbb{Z}^d , such as the widely studied two-type Richardson model, and highlights the rich behaviour of FPPHE. (We discuss results for other models more thoroughly in Section 1.5.) We remark that the coexistence regime we establish is a bit particular, in the sense that FPP_λ is guaranteed to survive already at time 0. An interesting open problem is to establish the regime of coexistence for some $p < p'_c$; we discuss further open questions in Section 6.

Corollary 1.2 leads one to consider the behaviour of FPPHE when $p < p'_c$ and our second result is a regime of strong survival in this direction. For $M \geq 0$, let $\Lambda_M = [-M/2, M/2]^d$ and $\partial\Lambda_M$ denote the set of sites in Λ_M that have a neighbour not con-

³The choice of subscript $1 - p$ is to remain consistent with later notation, where the emphasis is placed on non-seeds.

⁴We remark that Sidoravicius and Stauffer did prove coexistence for a simplified version of FPPHE, where passage times are deterministic [27]. However, this adaptation renders the model simple enough so that coexistence can be derived from basic percolation results.

tained in Λ_M . Let $\{o \xrightarrow{f} \partial\Lambda_M\}$ be the event that there is a path of filled seeds from a neighbour of the origin to $\partial\Lambda_M$. Define the critical probability

$$p_c'' = \sup \{p \in [0, 1] : \exists a > 0 \text{ such that for all large enough } M, \\ \mathbb{P}_{1-p}(o \xrightarrow{f} \partial\Lambda_M) \leq \exp(-M^a)\}. \quad (1)$$

It is immediate that $p_c'' \leq p_c'$ and we believe there is equality, but proving this is beyond the scope of this article. We prove that if $p < \min\{p_c'', 1 - p_c^{\text{site}}\}$, then for a sufficiently small λ , there is strong survival. We note that we can extend this result to $d = 2$.

Theorem 1.3. *Suppose $d \geq 2$ and $p < \min\{p_c'', 1 - p_c^{\text{site}}\}$. There exists a constant $\tilde{\lambda}_1 = \tilde{\lambda}_1(d, p) > 0$ such that if $\lambda \in (0, \tilde{\lambda}_1)$, then there is strong survival.*

We relate our results to known results for FPPHE in Section 1.4, after we discuss our novel approach to multi-scale analysis.

1.3. Multi-scale analysis with non-equilibrium feedback

The development of multi-scale analysis has arisen as a means of understanding the macroscopic behaviour of complex systems and has proven to be a powerful tool across many mathematical fields. A standard multi-scale construction usually goes as follows. The underlying space is covered by boxes of a fixed size that are labelled as either *good* or *bad* according to (local) events that are measurable with respect to that box. One would choose such events so that large clusters of good boxes imply some desired behaviour at a mesoscopic or macroscopic scale. In some cases this single-scale renormalization procedure will suffice but often one will need to define boxes at infinitely many scales of increasing size and define good and bad appropriately at higher scales. The main challenge of multi-scale constructions is finding a sensible way of defining good and bad boxes so that the desired properties of the system can be captured while good boxes occur with sufficiently high probability at all scales. If one is able to do this, then this robust framework is able to handle systems with strong correlations that otherwise are not tractable to analysis. Examples of such an approach range from random interlacements and dependent percolation [9, 26, 28] to interacting particle systems [16, 22–24].

A common feature of systems that are amenable to a standard multi-scale analysis as described above is that they are in *equilibrium*, thus for example it is possible to assess how good or bad a configuration is at any given moment in time and space without having to observe global information about the system. Another important feature is *monotonicity*, which allows for sprinkling arguments to be developed in order to “decouple” space-time boxes that are sufficiently far from one another. Hence if one can find a suitable definition of good and bad boxes, and can control how they are distributed, then a standard multi-scale analysis can be potent in understanding the macroscopic structure of the system in question.

The main contribution of this paper is to develop a framework, which we refer to as *multi-scale analysis with non-equilibrium feedback*, that we believe can be used to

perform multi-scale analysis on some non-equilibrium systems. As with standard multi-scale analysis, our framework is not a “black box” that is ready to be applied, but rather an approach (or a strategy) to be followed and tailored to each particular problem. Here we provide a heuristic and overall explanation of the main ideas behind our framework, and we give a more detailed account later in Section 4, where we develop it to analyze FPPHE.

For concreteness, henceforth we consider the case of FPPHE, but the reasoning below could be adapted to a more general non-equilibrium process. We would like to characterise some desired macroscopic behaviour of FPPHE such as survival of FPP_1 . We could perform a standard multi-scale analysis by constructing good and bad boxes at infinitely many scales so that a good box implies that the passage times and the distribution of the seeds inside the box is such that if there were only FPP_1 outside the box then FPP_1 could “spread well” throughout the box. Such events could be defined in a local way, but the problem lies in the part “if there were only FPP_1 outside the box”. What if it is FPP_λ that enters the box first? Even worse, what if FPP_λ growing from outside the box ends up taking the whole boundary of the box?

Our approach is to distinguish between two classes of good boxes, which we refer to as being of *positive feedback* or *negative feedback*. For this discussion, we could simplistically say that a box of positive feedback is a box which FPP_1 arrives to it first, before FPP_λ does.⁵ Note that whether a box is of positive or negative feedback is an event that is *not local*, and depends on the whole evolution of the process up to the time it reaches the box. Considering also the fact that FPPHE is not a monotone process, controlling the probability that a box has positive or negative feedback seems to be rather difficult. But the way we proceed is to analyze positive and negative feedback boxes in a completely different manner; in particular, we will *not check* whether boxes have positive or negative feedback, instead we will use this information and assess its consequences. This motivated the choice of the word *feedback*, as we regard this information as a feedback that the box receives from the whole system (that is, it is regarded as information that is given to us, instead of information whose validity we check or whose probability we estimate).

The overall idea is to tune the definition of positive and negative feedback so that the following two properties are obtained:

- (i) a positive feedback box implies the well spread of FPP_1 inside that box (that is, it implies the desired macroscopic behaviour of the process),
- (ii) a box that has negative feedback can be associated to a bad box (called its progenitor) in such a way that we can control how far away the progenitor of a negative feedback box is.

Thus since bad boxes (via a standard multi-scale analysis) can be shown to be rare, (ii) allows us to control the influence and location of boxes with negative feedback. Note

⁵We will need to develop a much more subtle notion of positive feedback, but this simple version is enough to give the overall idea behind our approach.

that (ii) allows us to show that negative feedback boxes are rare by controlling bad boxes only, without having to estimate the probability that a box is of negative feedback.

To give a better idea regarding (ii), the progenitor of a negative feedback box Q is the box that *is responsible* for Q ending up having a negative feedback; in the case of FPPHE, one could imagine that the progenitor is the bad box from which a seed of FPP_λ is activated and grew all the way until entering Q , doing so before FPP_1 . We could generalize (ii) so that a negative feedback box is associated to more than one progenitor, as long as we can control the region inside which such progenitors may be found, but in the present paper it is enough to have a single progenitor for each negative feedback box. We also remark that bad boxes do not need to be classified as a positive feedback or negative feedback. So whenever we refer to a positive or negative feedback box, we assume that the box is good.

Our approach is then naturally split into two parts. The first part consists of a standard multi-scale analysis where we control how the process will behave inside a box given that the behaviour of the process outside the box is nice enough. Then, in the second part we define positive and negative feedback, and derive properties (i) and (ii) above. We divided the bulk of our proof for FPPHE into two sections exactly to better highlight the role of each of these parts, with the first part (the standard multi-scale analysis) being developed in Section 3 and the second part (the actual multi-scale analysis with non-equilibrium feedback) being developed in Section 4.

We believe that the idea behind the multi-scale analysis with positive feedback is robust enough to be applied to other systems with non-equilibrium dynamics. In particular, systems that could be suitable to such type of approach are the ones which either contain a stationary component (like the passage times of FPPHE, which are i.i.d., or systems with a suitable graphical representation involving i.i.d. random variables) or can be approximated by a stationary process. Such a feature could be used in a two-part proof, as described above, with a standard multi-scale analysis being constructed on the stationary component of the process and a positive/negative feedback mechanism being developed to leverage the multi-scale analysis to control the aspects of the process that is not in equilibrium.

1.4. Known results for FPPHE

The first result for FPPHE was proving the existence of a strong survival phase for $d \geq 2$ by Sidoravicius and Stauffer [27, Theorem 1.3]. More precisely, for $d \geq 2$ they proved that for any $\lambda \in (0, 1)$, there exists $p_0 = p_0(\lambda, d) > 0$ such that if $p \in (0, p_0)$, then there is strong survival. The difference between Theorem 1.3 and the strong survival phase proven in [27] is that we fix p first, and then take λ small enough, whereas in [27] it is λ that is fixed first and p that is taken small enough. As noticed in Figure 1, such a change makes FPPHE behave rather differently.

The proof in [27] relied on a multi-scale argument that handled the long-range dependencies in FPPHE in the following manner. One can restrict the spread of FPP_λ from a seed to a (possibly arbitrarily large) region of \mathbb{Z}^d by showing that FPP_1 is able to grow around

(encapsulating) FPP_λ inside this region. Outside of this “bad region”, FPP_λ originating from this seed no longer affects how FPPHE evolves. If such a bad region does not intersect bad regions from other seeds (a fact that ends up being obtained by setting p small enough), then a multi-scale argument that accounts for all bad regions implies that the sites that are not contained in any bad region percolate in \mathbb{Z}^d . This implies the strong survival phase.

In the coexistence phase, however, FPP_1 cannot encapsulate FPP_λ , and in particular there is a seed that gives rise to an infinite component of FPP_λ that FPP_1 must survive against. Hence we had to find a different way to control how much effect a seed may have in the growth of FPP_λ . Drawing a parallel between the two works, we notice that [27] develops a multi-scale analysis that is particularly tailored to the analysis of that specific question and also crucially relies on the fact that outside of bad regions one only finds FPP_1 . This last property allowed for an essentially standard multi-scale analysis to be developed, as local events alone were enough to guarantee the encapsulation of FPP_λ . In particular, the technique in [27] is ineffective to handle the more challenging coexistence phase. On the other hand, not only our proof can handle the coexistence regime, but it also consists in the development of a novel approach to multi-scale analysis, which gives a mechanism that we believe is suitable to be adapted to other questions and models. In particular, in Section 6, we briefly explain how our approach can be tuned to give a cleaner proof of the result in [27].

We end this section with further remarks about FPPHE. It is intuitive that if $\lambda \geq 1$, then FPP_1 dies out almost surely. In particular, adapting a result by van den Berg and Kesten [29] regarding strict inequalities for first passage percolation (in particular, the extension to the limit shape given in [27, Proposition 6.4]), we find that there exists an $\varepsilon > 0$ which does not depend on λ (just on p and d) so that FPP_1 spreads in a way that is strictly slower than first passage percolation of rate $1 - \varepsilon$. This is simply because FPP_1 must at least deviate from a density of seeds. Then if $\lambda > 1 - \varepsilon$, some seed of FPP_λ will manage to grow fast enough to eventually encapsulate FPP_1 inside a finite region. We refer to [12, Theorem 4.1] for details. The known behaviour for FPPHE on \mathbb{Z}^d for $d \geq 3$ can be seen in Figure 2(c).

FPPHE can be readily generalised to any infinite connected graph. For example, if G is a rooted tree and $p < 1 - p_c^{\text{site}}(G)$, then coexistence is equivalent to the root being contained in an infinite cluster of non-seed sites, where the root has the same role as the origin on \mathbb{Z}^d . Such cases are considered trivial as coexistence is an artefact of a more classically understood process. The first non-trivial coexistence phase was established in Candellero and Stauffer [7, Corollary 1.3] when G is a vertex-transitive, hyperbolic and non-amenable graph. First, in [7, Theorem 1.1], they prove that for any $\lambda > 0$, there exists a $p_0 = p_0(G, \lambda)$ such that if $p \in (0, p_0)$, then the FPP_1 process survives with positive probability. Though this is analogous to the result in [27] for \mathbb{Z}^d , there is a difference in behaviour as in hyperbolic and non-amenable graphs, FPP_1 can survive even if $\lambda \geq 1$. Secondly, in [7, Theorem 1.2], they prove that for any $p \in (0, 1)$, and any $\lambda > 0$, the FPP_λ process survives almost surely. In other words, the strong survival phase does not occur for any choice of parameters. We remark that a result analogous to our theorem on

strong survival (Theorem 1.3), where p is fixed first and then λ is set small enough, has not been established in [7] for hyperbolic, non-amenable graphs, the main issue being that such a case requires a more delicate analysis which would involve detailed control of the geometry of non-seeds.

1.5. Related growth models and the search for coexistence

The study of random growth processes with competition is a classical area of research with particular interest in the circumstances in which both processes can coexist with one another, especially on \mathbb{Z}^d . A notable example is the *two-type Richardson model*, in which two types (which we call FPP_1 and FPP_λ) grow throughout \mathbb{Z}^d as follows: FPP_1 starts from the origin and spreads throughout the edges of \mathbb{Z}^d at rate 1 (as a first passage percolation process), whereas FPP_λ starts from a neighbour of the origin and spreads in the same way as FPP_1 , but at a rate λ . Whenever a site is occupied by one of the processes, it remains so forever. Unlike the model FPPHE we study in this paper, in the two-type Richardson model both FPP_1 and FPP_λ start spreading already from time 0.

It is conjectured that when $d \geq 2$ coexistence in the two-type Richardson model can occur with positive probability if and only if $\lambda = 1$ (that is, both processes spread at the same rate). Coexistence for $\lambda = 1$ was established on \mathbb{Z}^2 by Häggström and Pemantle [19] before being generalised to higher dimensions by Garet and Marchand [14] and Hoffman [21]. The other side of the conjecture remains open, but Häggström and Pemantle [20] proved that the set of values for λ for which coexistence can occur with positive probability is at most a countable subset of $(0, \infty)$. At first, it may sound incredible that this does not imply that coexistence can only occur for $\lambda = 1$, but the issue is that the event of coexistence is not monotone in λ . Nonetheless, the two-type Richardson model is a monotone process in the sense that there is a straightforward coupling under which if λ is increased, then the set of sites occupied by FPP_λ can only increase. But the event of coexistence is more subtle since increasing (resp., decreasing) λ could cause the transition from a regime of coexistence to a regime where FPP_λ (resp., FPP_1) is the only one to survive. Recently, the conjecture for the two-type Richardson model was solved for the half-plane by Ahlberg, Deijfen and Hoffman [1], exploiting the planarity of \mathbb{Z}^2 and the boundary of the half-plane. We remark that, as we explained previously, FPPHE is *not* monotone at all, which highlights the challenges that the intrinsic strong dependences in FPPHE can give rise to.

Garet and Marchand [15] proved that in the two-type Richardson model with $\lambda \neq 1$, both processes cannot simultaneously occupy a positive density of sites almost surely. In fact, in [15] they extend this result to a broad class of passage times beyond the context of the two-type Richardson model. This is in stark contrast to the behaviour of FPPHE, as in Corollary 1.2 we see that FPPHE can have both processes occupying a positive density of sites and coexisting with positive probability.

In a somewhat different context, Deijfen, Hirschler and Lopes [10] studied a two-type variant of a particle system known as the *frog model*. In this model, we start two *active* particles from the origin of \mathbb{Z}^d , one particle of type 1 and the other of type λ . In addition,

at each site that is not the origin, we place an independent random number of *sleeping* particles. Then, from time 0 active particles move as independent simple random walks. with type 1 particles jumping at rate 1 and type λ particles jumping at rate λ . Sleeping particles, in turn, do not jump. However, they become active whenever a particle jumps on their site, and at that time acquire the type of the particle that jumped onto them. A central question is to understand which type manages to conquer an infinite number of particles, and in particular whether coexistence is possible (that is, whether both types can simultaneously conquer an infinite number of particles).

In [10], it is shown (for a discrete-time variant of the aforementioned process) that coexistence is possible if the two types have equal jump rates (i.e., $\lambda = 1$). They conjecture that when the jump probabilities are different, coexistence is possible only if the initial distribution of the sleeping particles is especially heavy-tailed. We note that the two-type frog model is also a monotone process. Many other one-type interacting particle systems could also be generalized to two-type competition versions, and references for these can be found in [10].

1.6. Outline of paper

In Section 2 we provide a rigorous construction of FPPHE and fix some useful notation in regard to first passage percolation. In Section 3 we perform a standard multi-scale analysis of FPPHE, as described in Section 1.3. In Section 4 we introduce multi-scale analysis with non-equilibrium feedback and derive all the desired properties. In Section 5 we establish Theorems 1.1 and 1.3 using the results of the previous sections. In Section 6 we conclude with some open questions and final remarks.

2. Preliminaries

In this section we give a formal construction of FPPHE on \mathbb{Z}^d . Initially we place seeds according to i.i.d. Bernoulli random variables of parameter $p \in (0, 1)$ on all sites except the origin. Fix $\lambda > 0$. To define the passage times for FPP_1 and FPP_λ we construct two sets of passage times $\{t_e\}_{e \in E}$ and $\{t_e^\lambda\}_{e \in E}$, where E is the edge set of \mathbb{Z}^d , such that $\{t_e\}_{e \in E}$ (resp. $\{t_e^\lambda\}_{e \in E}$) is a collection of i.i.d. exponentially distributed random variables of rate 1 (resp. λ).

At time $t = 0$ all seeds are inactive and FPP_1 only occupies the origin. The dynamics for $t \geq 0$ are as follows. If x is first occupied by the FPP_1 process at time t and y is a neighbour of x connected by an edge $e_{x,y}$, then FPP_1 attempts to occupy y at time $t + t_{e_{x,y}}$. The occupation is successful if y is not occupied by FPP_1 or FPP_λ before time $t + t_{e_{x,y}}$ and y is not a seed. If y is occupied by FPP_1 or FPP_λ by time $t + t_{e_{x,y}}$, then the occupation from x is unsuccessful. If y is an inactive seed for all time before time $t + t_{e_{x,y}}$, then y becomes an active seed at time $t + t_{e_{x,y}}$, meaning that it is occupied by the FPP_λ process at time $t + t_{e_{x,y}}$. The dynamics if x was instead occupied by the FPP_λ process are precisely the same except with passage times given by $\{t_e^\lambda\}_{e \in E}$.

Remark 2.1. In our analysis we never appeal to the memoryless property of exponentials and so more general passage times with reasonable conditions could be considered instead. For example, distributions with exponential tails would satisfy our analysis. We choose to restrict our attention to exponential passage times for ease of exposition and to avoid cumbersome notation.

To remove possible ambiguities when speaking about the occupancy of sites by FPP_1 and FPP_λ , we introduce the following notation. We let η_t be the *configuration* for FPPHE on \mathbb{Z}^d at time t , meaning that $\eta_t : \mathbb{Z}^d \rightarrow \{-1, 0, 1, 2\}$ where $\{-1, 0, 1, 2\}$ is a set of labels giving the state of occupancy of a site in the following manner. For a fixed site $x \in \mathbb{Z}^d$ and time $t \geq 0$, we have:

- if $\eta_t(x) = -1$, then x is a non-activated seed at time t ,
- if $\eta_t(x) = 0$, then x is not a seed and is not occupied by FPP_1 or FPP_λ at time t ,
- if $\eta_t(x) = 1$, then x is occupied by FPP_1 at time t ,
- if $\eta_t(x) = 2$, then x is either an activated seed or a non-seed occupied by FPP_λ at time t .

Using the above notation, η_0 is defined as follows. We set $\eta_0(0) = 1$ and for $x \in \mathbb{Z}^d \setminus \{0\}$, with probability p we have $\eta_0(x) = -1$, otherwise $\eta_0(x) = 0$.

From the set of passage times $\{t_e\}_{e \in E}$ and $\{t_e^\lambda\}_{e \in E}$ given in the construction of FPPHE, it will be useful to recall some standard definitions in first passage percolation. The interested reader can find an introduction to first passage percolation in the monograph [4] by Auffinger, Damron and Hanson.

Let $\gamma = (v_0, v_1, \dots, v_{n-1}, v_n)$ be a finite simple path of sites on \mathbb{Z}^d , meaning that $v_i \neq v_j$ if $i \neq j$ and $\|v_i - v_{i-1}\|_1 = 1$ for all $i \in \{1, \dots, n\}$. We define the *passage time of γ with respect to $\{t_e\}_{e \in E}$* as the random variable

$$T_1(\gamma) = \sum_{i=1}^n t_{e_i}$$

where e_i is the edge between v_{i-1} and v_i . Similarly, we define the *passage time of γ with respect to $\{t_e^\lambda\}_{e \in E}$* as the random variable

$$T_\lambda(\gamma) = \sum_{i=1}^n t_{e_i}^\lambda.$$

For $x, y \in \mathbb{Z}^d$, let $\Gamma(x \rightarrow y)$ be the set of all finite simple paths from x to y . We define the *passage time between x and y* as the random variable

$$T_\#(x \rightarrow y) = \inf_{\gamma \in \Gamma(x \rightarrow y)} T_\#(\gamma),$$

where the label $\# \in \{1, \lambda\}$ indicates which set of passage times are being referred to. If U is a subgraph of \mathbb{Z}^d , we let $T_\#(x \rightarrow y; U)$ be the infimum of all passage times over paths

that do not exit U , so that

$$T_{\#}(x \rightarrow y; U) = \inf_{\gamma \in \Gamma(x \rightarrow y; U)} T_{\#}(\gamma),$$

where $\Gamma(x \rightarrow y; U)$ is the set of all finite simple paths from x to y that do not exit U .

3. Standard multi-scale analysis

In order to better explain the non-equilibrium feedback idea we introduce, we split our multi-scale analysis in the next two sections. In this section, we describe the part of our analysis that follows a more standard (“in equilibrium”) multi-scale analysis approach, which will be based only on *local* events on the sets of passage times $\{t_e\}_{e \in E}$ and $\{t_e^\lambda\}_{e \in E}$.

Let L_1 be some large integer to be set later. For $k \geq 2$, set

$$L_k = k^2 L_{k-1}^d. \tag{2}$$

Each L_k gives us the side lengths of the k -boxes we define below. We refer to k as the *scale*.

At each scale $k \geq 1$ we partition \mathbb{Z}^d into boxes of side-length $L_k/3$, producing a collection $\{Q_k^{\text{core}}(i)\}_{i \in \mathbb{Z}^d}$ of disjoint boxes that we call *k-cores*,

$$Q_k^{\text{core}}(i) = (L_k/3)i + [-L_k/6, L_k/6]^d.$$

For each $k \geq 1$, define *k-boxes* as a collection $\{Q_k(i)\}_{i \in \mathbb{Z}^d}$ of overlapping boxes of length L_k which are centred on the *k-cores* by

$$Q_k(i) = (L_k/3)i + [-L_k/2, L_k/2]^d.$$

3.1. Good boxes at scale 1

The first step in this multi-scale analysis is defining local events to 1-boxes that will distinguish between *good* and *bad* 1-boxes. Roughly speaking, we would like a good 1-box to imply that FPP_1 has the opportunity to spread with fast passage times throughout the 1-box while the spread of FPP_λ is inhibited. Moreover, we would like to prove that good 1-boxes occur with high probability for a suitable choice of large L_1 and small λ . In this section we make these notions rigorous.

One condition for a 1-box to be good is that the non-seed sites *percolate* well inside the 1-box, giving FPP_1 the opportunity to spread to nearby 1-boxes. To define this condition rigorously, let \mathcal{G} be the random subgraph of \mathbb{Z}^d induced by removing all FPP_λ seeds from \mathbb{Z}^d , noting that we drop the dependence on p for ease of notation. Let $d_{\mathcal{G}}(\cdot, \cdot)$ be the natural graph metric induced by \mathcal{G} and let $V(\mathcal{G})$ be the vertex set of \mathcal{G} . Fix $i \in \mathbb{Z}^d$ and consider the 1-box $Q_1(i)$. Let $\partial Q_1(i)$ be the set of sites of $Q_1(i)$ that have a neighbour not contained in $Q_1(i)$. Enumerate the clusters of $V(\mathcal{G}) \cap Q_1(i)$ where we remove edges between sites in $\partial Q_1(i)$ as

$$\mathcal{C}_1(i), \dots, \mathcal{C}_{n_i}(i),$$

where n_i is the number of clusters and $|\mathcal{C}_1(i)| \geq \dots \geq |\mathcal{C}_{n_i}(i)|$. Note that this ordering may not be unique, but this fact will not alter our arguments. The removal of edges between sites in $\partial Q_1(i)$ is a non-restrictive condition that will simplify arguments later. We let $\mathcal{C}_1^-(i) = \mathcal{C}_1(i) \setminus \partial Q_1(i)$ denote the cluster $\mathcal{C}_1(i)$ with sites in $\partial Q_1(i)$ removed.

Recall we assume $p < 1 - p_c^{\text{site}}$ so that the non-seed sites are supercritical. With this assumption in mind, we have

$$\theta(1-p) = \mathbb{P}_{1-p}(o \text{ is contained in an infinite cluster of } \mathcal{G}) > 0,$$

where \mathbb{P}_{1-p} is the measure given by placements of seeds with parameter p in the construction of FPPHE and we write o for the origin of \mathbb{Z}^d . We write $1-p$ instead of p to emphasise that it is the non-seed sites that we are considering and to remain consistent with standard notation in percolation theory.

For a fixed $\varepsilon > 0$, define the event

$$E_1(i) = \left\{ \begin{array}{l} \text{for all } i' \text{ with } Q_1^{\text{core}}(i') \subset Q_1(i) \text{ the set } \mathcal{C}_1(i) \cap Q_1^{\text{core}}(i') \text{ touches all} \\ \text{faces of } Q_1^{\text{core}}(i') \text{ and } |\mathcal{C}_1(i) \cap Q_1^{\text{core}}(i')| \geq (1-\varepsilon)\theta(1-p)(L_1/3)^d \end{array} \right\} \quad (3)$$

where the dependence on ε is left implicit for ease of notation. Roughly speaking, the event $E_1(i)$ guarantees that within each 1-core contained in $Q_1(i)$, the large component of non-seeds in $Q_1(i)$ percolates well within each 1-core. Deuschel and Pisztora [11, Theorem 1.1] proved large deviation estimates for such events, and in particular there exists a constant $b_1 > 0$ such that

$$\mathbb{P}(E_1(i)) > 1 - \exp(-b_1 L_1^{d-1}).$$

It could be the case that the large components of non-seeds within each core are not contained in the same cluster within a 1-box. To handle this, we consider the event

$$E_2(i) = \{|\mathcal{C}_2(i)| \leq \log^2 |Q_1(i)|\},$$

which is the event that components of non-seeds that are not in the largest component in $Q_1(i)$ are logarithmically small. Penrose and Pisztora [25, Theorem 5] provide large deviation estimates for such events and show that there exists a constant $b_2 > 0$ such that

$$\mathbb{P}(E_2(i)) > 1 - \exp(-b_2 \log^2 L_1).$$

If $Q_1(i)$ and $Q_1(j)$ are neighbours in the sense that i and j are neighbours on \mathbb{Z}^d , there is an overlap of side length at least $L_k/3$. If $E_1(i)$, $E_1(j)$, $E_2(i)$ and $E_2(j)$ all hold, then for large enough L_1 , the intersection of the large components $\mathcal{C}_1(i) \cap \mathcal{C}_1(j)$ of the respective 1-boxes must be non-empty, as the smaller clusters are only logarithmically small.

The event $E_1(i)$ only tells us that the non-seeds percolate well in a 1-box but it could be the case that the paths of non-seeds within the box are superlinear in the graph distance of the sites which would greatly inhibit the spread of FPP_1 . To remedy this, we recall the notion of the *chemical distance* between two sites on a percolated graph, which is their

graph distance induced by the graph under percolation. Let $\mathcal{G}(i)$ be \mathcal{G} restricted to $Q_1(i)$ with edges between sites within $\partial Q_1(i)$ not included. Set $d_{\mathcal{G}(i)}(\cdot, \cdot)$ to be the natural graph distance metric on $\mathcal{G}(i)$.

For a constant $c_1 \geq 1$, define the event

$$E_3(i) = \{\forall x, y \in \mathcal{C}_1(i), d_{\mathcal{G}(i)}(x, y) < c_1 \max \{\|x - y\|_1, \log^2 L_1\}\},$$

that states the chemical distance between two sites in the large component of non-seed sites within $Q_1(i)$ is linear with a logarithmic term. For $E_3(i)$ we can appeal to large deviation estimates for the chemical distance for supercritical Bernoulli percolation by Antal and Pisztor [3, Corollary 1.3]. As long as c_1 is large enough with respect to p and d , as given in [3, Theorem 1.1], we may deduce that for all large enough choices of L_1 ,

$$\mathbb{P}(E_3(i)) > 1 - \exp(-c_1 \log^2 L_1).$$

We observe two technical points to justify the above bound. First, the main results in Antal and Pisztor [3] are stated for *bond* percolation, but also work for site percolation; see the remark following [3, Theorem 1.2]. Second, the event $E_3(i)$ concerns paths restricted to the 1-box $Q_1(i)$, while Antal and Pisztor [3] consider unrestricted paths. However, the same proof in the truncated case follows by performing the coarse graining scheme of Antal and Pisztor, say with subboxes of side-length N partitioning $Q_1(i)$, and then setting c_1 and L_1 large enough with respect to N . More precisely, each subbox is classified as either good or bad according to the presence of a unique large component of non-seeds within the subbox (similar to our events E_1 and E_2). As the probability that a subbox is good can be made large, we are in the domain of highly supercritical percolation. Thus to construct a path from x to y that are both in the large component of $Q_1(i)$, it suffices to fix a deterministic path of subboxes from the subbox containing x to the subbox containing y , and then go around bad clusters of subboxes using an argument of Fontes and Newman [13] as done in [3].

For a large constant $c_2 > 1$, define the event

$$E_4(i) = \{\forall x, y \in \mathcal{C}_1(i), T_1(x \rightarrow y; \mathcal{G}(i)) < c_2 \max \{\|x - y\|_1, \log^2 L_1\}\},$$

which states that the passage times between two non-seed sites in the large component of $Q_1(i)$ are linear in their graph distance. By considering Chernoff bounds for sums of exponentials, one can then deduce that $E_4(i)$ holds with high probability for a large enough c_2 through similar arguments made for $E_3(i)$.

Let $\mathcal{E}(i)$ be the set of edges of \mathbb{Z}^d where both endpoints are contained in $Q_1(i)$ and $E_5(i)$ be the event that all edges in $\mathcal{E}(i)$ have passage time at least $1/\sqrt{\lambda}$ with respect to $\{t_e^\lambda\}_{e \in E}$, so that

$$E_5(i) = \{\forall e \in \mathcal{E}(i), t_e^\lambda \geq 1/\sqrt{\lambda}\}.$$

Recall that if X is an exponentially distributed random variable of rate μ , then $\mathbb{P}(X < x) = 1 - e^{-x\mu}$ for any $x \geq 0$. Hence by taking the union bound over each edge $e \in \mathcal{E}(i)$, we have

$$\mathbb{P}(E_5(i)^c) \leq (1 - e^{-\sqrt{\lambda}})|\mathcal{E}(i)| \leq \sqrt{\lambda} |\mathcal{E}(i)| \tag{4}$$

where $|\cdot|$ denotes cardinality and we use the inequality $e^{-x} \geq 1 - x$ for all $x \in \mathbb{R}$. The right-hand side of (4) may be made arbitrarily small by setting λ small enough.

Definition 3.1 (Good box at scale 1). Let $i \in \mathbb{Z}^d$ and fix any $\varepsilon > 0$ and large enough constants $c_1, c_2 > 1$. Define the event

$$G_1(i) = \bigcap_{j=1}^5 E_j(i).$$

If $G_1(i)$ holds, then we define $Q_1(i)$ to be *good*. Otherwise we define $Q_1(i)$ to be *bad*.

Note that a 1-box is measurable with respect to events concerning the sites and edges inside the 1-box, and thus disjoint 1-boxes being good are independent events. The probability of a 1-box being good is invariant under translations and so we may define ρ_1 to be the probability that an arbitrary 1-box is bad by setting

$$\rho_1 = 1 - \mathbb{P}(G_1(o)).$$

Lemma 3.2. For any $\varepsilon > 0$, suitably large choices of $c_1, c_2 > 1$ and $B > 0$, by setting L_1 large enough and then λ small enough, we have

$$\rho_1 < L_1^{-B}.$$

Proof. The result is an immediate consequence of the bounds derived for the probabilities of the events $E_j(i)$ above. ■

3.2. Good boxes at higher scales

In this section we generalise the notion of good boxes to higher scales. The idea is that for $k \geq 2$, a k -box is good if it contains no more than A disjoint bad $(k - 1)$ -boxes, where A is a fixed constant that we set later.

Definition 3.3 (Good box at scale $k \geq 2$). Fix a constant $A > 0$. Let $k \geq 2$ and consider the k -box $Q_k(i)$. Define $G_k(i)$ to be the event that $Q_k(i)$ contains no more than A disjoint bad $(k - 1)$ -boxes. If $G_k(i)$ holds, then we define $Q_k(i)$ to be *good*. Otherwise we define $Q_k(i)$ to be *bad*.

From the definition of good k -box for $k \geq 2$, we deduce that disjoint k -boxes being good are independent events. Later in Lemma 3.4 we will see that choosing $A > d$ will suffice so long as L_1 is large enough. The intuition behind this definition is that knowing $Q_k(i)$ is good means that there are at most A disjoint bad $(k - 1)$ -boxes, which will simplify controlling the spread of FPP_λ within $Q_k(i)$. There may be many other bad boxes at lower scales, but our multi-scale framework will allow us to consider bad boxes on a scale-by-scale basis. With at most A disjoint bad $(k - 1)$ -boxes in a good k -box, the total number of bad $(k - 1)$ -boxes in that k -box is bounded above by a constant that only depends on A and d .

Fix $k \geq 1$. Similar to the scale 1 case, the probability of a k -box being good is invariant under translations, so we can define ρ_k to be the probability that an arbitrary k -box is bad by setting

$$\rho_k = 1 - \mathbb{P}(G_k(o)).$$

Lemma 3.4. *Let $A > d$. If we choose L_1 sufficiently large and then λ small enough, then for all $k \geq 1$ we have*

$$\rho_k \leq \rho_1^{A^{k-1}}.$$

Proof. We proceed by induction, noting that the case $k = 1$ is immediate. Now let $k > 1$ and assume Lemma 3.4 holds up to scale $k - 1$. We can first bound ρ_k by noticing that

$$\rho_k \leq \mathbb{P}(\Pi_{A+1}),$$

where Π_{A+1} is the event that there are $A + 1$ disjoint bad $(k - 1)$ -boxes in the k -box. We can find an upper bound on the number of $(k - 1)$ -boxes in a k -box by counting the number of $(k - 1)$ -cores contained in a k -box, which equals

$$\left(\frac{L_k}{L_{k-1}/3} \right)^d = 3^d k^{2d} L_{k-1}^{d(d-1)}.$$

Hence the number of possible ways to choose $A + 1$ disjoint bad $(k - 1)$ -boxes is bounded above by $(3^d k^{2d} L_{k-1}^{d(d-1)})^{A+1}$ and by taking a union bound over these configurations we see that

$$\begin{aligned} \rho_k &\leq (3^d k^{2d} L_{k-1}^{d(d-1)} \rho_{k-1})^{A+1}, \\ &\leq (3^d k^{2d} L_{k-1}^{d(d-1)})^{A+1} \rho_1^{(A+1)A^{k-2}}, \end{aligned} \tag{5}$$

where in the last line we appeal to the inductive hypothesis. The result follows from (5) if we prove that

$$(3^d k^{2d} L_{k-1}^{d(d+1)})^{A+1} \leq \rho_1^{-A^{k-2}}, \tag{6}$$

by setting L_1 large enough and then λ small enough. We first note that by (2) we have

$$L_k = \left(\prod_{j=2}^k j^{2d^{k-j}} \right) L_1^{d^{k-1}}.$$

Hence there exists a constant $c > 0$ such that if L_1 is sufficiently large, then

$$3^d k^{2d} L_{k-1}^{d(d+1)} \leq L_1^{d^{k+c}}$$

for all $k \geq 2$. Consequently,

$$(3^d k^{2d} L_{k-1}^{d(d+1)})^{A+1} \leq L_1^{(A+1)d^{k+c}} \leq L_1^{A^{k+c+2}}, \tag{7}$$

where the last inequality follows as $A > d$. Henceforth fix c such that (7) holds. By Lemma 3.2, for any constant $B > 0$, if L_1 is large enough and λ is small enough then

$$L_1 < \rho_1^{-1/B}.$$

In this case, from (7) we deduce that

$$(3^d k^{2d} L_{k-1}^{d(d+1)})^{A+1} < \rho_1^{-A^k+c+2/B} < \rho_1^{-A^{k-2}},$$

where the last inequality follows for all $k \geq 2$ by setting B large enough. Thus, we established (6) and the result follows by induction. ■

It is worth noting that in many applications of such multi-scale analysis constructions, *disjoint* boxes do not necessarily mean *independent* boxes. To deduce an analogue of Lemma 3.4 in such cases, one will typically use a decay of correlation or decoupling inequalities which despite adding some terms (that are proven to be of smaller order) still lead to a similar recursion as in the proof of Lemma 3.4. In our case, disjointness does imply independence and so our recursion argument simplifies to just controlling a combinatorial term.

4. Multi-scale analysis with non-equilibrium feedback

In this section we enhance the multi-scale construction of Section 3 in a way that will allow us to handle the non-equilibrium dynamics of FPPHE; we call this approach *multi-scale analysis with non-equilibrium feedback*. It will be useful to have in mind the intuitive explanation of our idea, which is given in Section 1.3. We start this section by adding some further details to this high-level discussion, and then we move to the rigorous construction.

4.1. High-level description

As explained in Section 1.3, we will introduce a notion of *timeliness* to the entrance of a good box, so that each *good* k -box will be further classified as having either *positive feedback* or *negative feedback*, whereas *bad* k -boxes are not further classified. Roughly speaking, a k -box will be called of *positive feedback* if FPP_1 is able to occupy a site away from the boundary of the k -box sufficiently fast in comparison to the time the box is first visited by either FPP_1 or FPP_λ . Moreover, each k -box Q will be associated to another k -box, called its *parent*. The parent of Q is the box that contains inside its core the site of Q that is first visited by either FPP_1 or FPP_λ . The above definitions will be later given in a fully precise manner, and will be tuned so as to imply the following properties:

- (CAS_k) A k -box with positive feedback is mostly occupied by $(k - 1)$ -boxes of positive feedback, which is a type of *cascading* property of positive feedback boxes.
- (Prog_k) The parent of a negative feedback k -box is either of negative feedback or a bad k -box, which will give rise to a so-called *progenitor structure* discussed below.
- (Fast_k) A good k -box is entered quickly if there is a nearby k -box of positive feedback, giving that positive feedback propagates *fast* enough.
- (Del_k^C) A negative feedback k -box has an entrance time that is sufficiently after the entrance of its parent (if its parent is also of negative feedback), giving that negative feedback k -boxes have a *delayed* entrance time.

The words emphasized above are the terms that we chose to represent each property and that give the property its label.

One could keep in mind from (Prog_k) that the parent of Q witnesses the fact that Q is of negative feedback. So, if the parent of Q is also (a good box) of negative feedback, then by (Prog_k) one finds that it must have a parent itself that is of negative feedback or bad. Applying (Prog_k) inductively, one finds that any negative feedback box can be associated to a path (or a *trail*) of negative feedback boxes (following each box's parent) that culminates into a *bad* box with a good parent, which will be called the *progenitor* of Q . (We remark that the trail of a negative feedback box can contain in its interior both negative feedback boxes and bad boxes with a negative feedback parent, and culminates into the progenitor which is the first bad box with positive feedback parent found along the path of parents.) We will refer to this as a progenitor structure, which will be the crucial property that will allow us to control where negative feedback boxes may occur. This is precisely the property (ii) described in Section 1.3. We emphasise (Prog_k) is not immediate from the definitions and requires λ to be sufficiently small (cf. Lemmas 4.6 and 4.22).

Properties (Fast_k) and (Del_k^C) guarantee that FPPHE gets delayed when traversing a trail of negative feedback boxes, while FPP_1 can quickly traverse through positive feedback boxes nearby. But note that no delay can be guaranteed when FPPHE passes from a bad box to a negative feedback box (or vice versa). Nonetheless, the conclusion we can obtain is that there cannot be long trails composed of large sequences of negative feedback boxes, which together with a control showing that bad boxes are rare will allow us to confine the trail of a negative feedback box to a small region around its progenitor.

Putting the properties together will allow us to conclude the following positive property from positive feedback boxes:

$$\text{positive feedback implies most non-seed sites in a box are occupied by } FPP_1. \quad (8)$$

This is precisely property (i) described in Section 1.3. We note that we cannot guarantee that FPP_1 occupies the *whole* of the box for several reasons. For example, in scale 1, the box will typically contain a density of seeds, and in higher scale it may contain bad boxes of smaller scale inside which we cannot give any guarantee for the spread of FPP_1 . Moreover, just by assuming a box is of positive feedback does not give us much information to control the spread of FPP_1 on sites that are close to the boundary of the box. This last issue will be solved, however, using the fact that boxes have a large overlap. So, the sites that are near the boundary of a box will be contained in the inner part of other boxes, so the spread of FPP_1 on those sites will be controlled by asserting whether those other boxes are of positive feedback.

The proof will then proceed as follows. We will start in Section 4.2 with a brief clarification on how L_1 and λ will be set, and then define positive feedback for scale 1 boxes in Section 4.3, and introduce the parent structure and the progenitor of a box in Section 4.4. Then our proof continues through three major steps: first we show that (CAS_1) holds, then we show that, for all k , property (CAS_k) implies (Prog_k) , (Fast_k) and (Del_k^C) , and ultimately we show that if all properties hold up to scale $k - 1$ then (CAS_k) holds. The three steps together inductively guarantee that all properties hold for all k . The details of the

proofs will be carried out in a few sections, as we need to distinguish the case of scale 1 from higher scales. The proof for scale 1 is derived in Section 4.3 for (CAS₁), Section 4.4 for (Prog₁) and Section 4.5 for (Fast₁) and (Del₁^C). The other sections are reserved to the definitions and the proofs for boxes of higher scales.

4.2. Setting L_1 and λ

Recall that the results of Section 3 hold (in particular, Lemma 3.4) for all L_1 large enough and λ small enough. That is, there exists a value \bar{L} , which depends on d , p and on the constant A from the definition of good boxes, and $\bar{\lambda} = \bar{\lambda}(L_1, d, p, A) > 0$ such that for all $L_1 \geq \bar{L}$, if $\lambda \in (0, \bar{\lambda})$ then the above results hold. Henceforth we fix L_1 larger than \bar{L} and such that $L_1 \geq 5000 \log^2 L_1$; the value 5000 being an arbitrary choice, which just allows us to guarantee that L_1 is sufficiently larger than $\log^2 L_1$. We will also require L_1 to be large enough with respect to some conditions on A , d and p that we only specify in later sections for ease of exposition. Now that L_1 has been fixed, we shall not write any condition on L_1 in the lemmas from this section.

Regarding λ , we will require that λ is not only smaller than $\bar{\lambda}$ but also smaller than another constant. To encapsulate the condition on λ in a simpler form, for any constant $x > 0$, we define

$$\lambda_x = \min \left\{ \bar{\lambda}, \frac{1}{(x + r_1)^2 L_1^2} \right\}, \quad (9)$$

where r_1 is a constant that only depends on d and p that we set later in Fact 4.1. Then by properly choosing x in the lemmas below we will enforce proper conditions on λ . Note that λ_x depends on L_1 , but we omit this from the notation since L_1 is regarded as fixed from now onwards.

4.3. Positive feedback at scale 1

In this section we define what it means for a good 1-box to have positive feedback and prove that a natural consequence of a 1-box having positive feedback is that *most* of the non-seed sites in that box are occupied by FPP_1 , in a sense that is made rigorous in Lemma 4.3.

Before we define positive feedback at scale 1, we first introduce some notation that will be useful for the remainder of the paper. Given $v \in \mathbb{Z}^d$, define the *entrance time for v* , written as $\tau(v)$, as the earliest time that either FPP_1 or FPP_λ occupies v , so that

$$\tau(v) = \inf \{ t \geq 0 : \eta_t(v) \in \{1, 2\} \}.$$

For $k \geq 1$ and $i \in \mathbb{Z}^d$, define the *entrance time for $Q_k(i)$* , written as $\tau_{k,i}$, as the earliest time any site in $Q_k(i)$ is occupied by either FPP_1 or FPP_λ , so that

$$\tau_{k,i} = \inf \{ \tau(v) : v \in Q_k(i) \}. \quad (10)$$

Fact 4.1. Suppose $Q_1(i)$ is a good 1-box. Recall $\mathcal{C}_1^-(i)$ is the component of non-seeds $\mathcal{C}_1(i)$ with sites in $\partial Q_1(i)$ removed. In later arguments it will be useful to have an upper

bound on the passage times between any two sites in $\mathcal{C}_1^-(i)$. Indeed, as a direct consequence of the definition of a good 1-box, for any $x, y \in \mathcal{C}_1^-(i)$, there exists a path γ from x to y such that

$$\gamma \subset \mathcal{C}_1^-(i) \quad \text{and} \quad T_1(\gamma) < c_2 \max \{\|x - y\|_1, \log^2 L_1\} \leq dc_2 L_1,$$

where we recall that c_2 is a large enough constant set in the definition of a good 1-box. It will be useful to set $r_1 = dc_2$ to be consistent with notation we set later.

Definition 4.2 (Positive feedback at scale 1). Let r_1 be as given in Fact 4.1 and suppose that $Q_1(i)$ is good. Define $Q_1(i)$ to have *positive feedback* if there exists a site $v \in \mathcal{C}_1^-(i)$ occupied by the FPP_1 process such that

$$\tau(v) \leq \tau_{1,i} + r_1 L_1.$$

If $Q_1(i)$ is good but does not have positive feedback, then we define it to have *negative feedback*.

From positive feedback we now define a property that will be a key component in a cascading argument in the proof of Theorem 1.1:

(CAS₁) If $Q_1(i)$ has positive feedback and $v \in \mathcal{C}_1^-(i)$, then v is occupied by FPP_1 .

If $Q_1(i)$ is good and $v \in \mathcal{C}_1^-(i)$ is occupied by FPP_λ , then

$$\tau(v) \geq \tau_{1,i} + 1/\sqrt{\lambda},$$

by definition of good 1-box as FPP_λ must cross at least one edge whose endpoints are both contained in $Q_1(i)$. The intuition is that by setting λ to be small enough, we can guarantee that (CAS₁) holds, as given in the following lemma. Recall the definition of λ_x from (9).

Lemma 4.3. *If $\lambda \in (0, \lambda_{r_1})$, then (CAS₁) holds.*

Proof. Suppose $Q_1(i)$ has positive feedback. Let $v \in \mathcal{C}_1^-(i)$ and suppose for contradiction that v is occupied by FPP_λ . Without losing generality, we may assume that v is such a site with earliest entrance time in $\mathcal{C}_1^-(i)$, so that

$$\tau(v) = \inf \{ \tau(w) : w \in \mathcal{C}_1^-(i) \text{ and } w \text{ is occupied by } FPP_\lambda \}.$$

As $v \in \mathcal{C}_1^-(i)$, we note that v is not a seed, and so by definition of good 1-box we have

$$\tau(v) \geq \tau_{1,i} + 1/\sqrt{\lambda}. \tag{11}$$

By definition of positive feedback, there exists a site $u \in \mathcal{C}_1^-(i)$ that is occupied by the FPP_1 process by time $\tau_{1,i} + r_1 L_1$. Recalling Fact 4.1, there exists a path γ from u to v such that $\gamma \subset \mathcal{C}_1^-(i)$ and $T_1(\gamma) < r_1 L_1$. Thus

$$\tau(v) < \tau(u) + r_1 L_1 < \tau_{1,i} + 2r_1 L_1. \tag{12}$$

If λ is small enough so that $2r_1 L_1 < 1/\sqrt{\lambda}$, then (12) contradicts (11), which establishes the result. ■

4.4. Parent and progenitor structure

In this section we provide the fundamental structure that will allow us to control entrance times for boxes at all scales. First we introduce the notion of a *parent*. Recall that for all $k \geq 1$, k -cores give a natural partition of \mathbb{Z}^d and can be used to uniquely identify from where a k -box is entered first by either FPP_1 or FPP_λ .

Definition 4.4 (Parent). Fix $k \geq 1$ and $i \in \mathbb{Z}^d$. Let $e_{k,i} \in \mathbb{Z}^d$ be the almost surely unique site in $Q_k(i)$ that is occupied by either FPP_1 or FPP_λ at time $\tau_{k,i}$, as given in (10). We define the *parent* of $Q_k(i)$ as the k -box $Q_k(i^{\text{par}})$ satisfying $e_{k,i} \in Q_k^{\text{core}}(i^{\text{par}})$.




Fig. 3. A k -box $Q_k(i)$ and its parent $Q_k(i^{\text{par}})$ where squares represent the cores tessellating \mathbb{Z}^d . The arrow represents the spread of FPP_1 or FPP_λ that is first to enter $Q_k(i)$ through $e_{k,i}$.

From the definition of parent, there is a *parent structure* that encodes entrance times as a directed tree, where the vertices are k -cores, and the root of the tree is the k -core containing the origin. See Figure 3 for an illustration of the parent structure. This parent structure is useful as it gives us information about where FPP_1 or FPP_λ first enters a box and if $Q_k(i)$ is a k -box with parent $Q_k(i^{\text{par}})$, then we know that $\tau_{k,i^{\text{par}}} \leq \tau_{k,i}$.

A key element to our method is deducing that all bad boxes and boxes with negative feedback can be traced to a unique bad box, which in some sense *caused* them to be bad or of negative feedback. To make this notion precise, we first define the *progenitor* of a bad box or box with negative feedback at all scales. We remark that we have not yet defined positive or negative feedback at higher scales but for ease of exposition we provide the definition of progenitor now.

Before we define progenitor rigorously, it will be useful to fix some notation. Given any k -box, we can trace through successive parents as far back as we desire, all the way back to $Q_k(o)$, which is the unique k -box that is its own parent. For an arbitrary k -box $Q_k(i_1)$, we set

$$\mathcal{P}_k^{i_1 \leftarrow i_N} = \{Q_k(i_1), \dots, Q_k(i_N)\}$$

to be a collection of k -boxes such that $Q_k(i_n) = Q_k(i_{n-1}^{\text{par}})$ for all $n \in \{2, \dots, N\}$. We call $\mathcal{P}_k^{i_1 \leftarrow i_N}$ the *path of parents from $Q_k(i_1)$ to $Q_k(i_N)$* .

Definition 4.5 (Progenitor). Let $k \geq 1$ and suppose $Q_k(i)$ is bad or has negative feedback. Without losing generality, we may assume that the k -box whose core contains the origin has positive feedback. We let $Q_k(i^{\text{prog}})$ be the unique k -box such that the path of parents $\mathcal{P}_k^{i \leftarrow i^{\text{prog}}}$ contains only bad k -boxes or k -boxes with negative feedback and the parent of $Q_k(i^{\text{prog}})$ has positive feedback. We call $Q_k(i^{\text{prog}})$ the *progenitor* of $Q_k(i)$.

For $k \geq 1$, if $Q_k(o)$ is good and $o \in \mathcal{C}_1(o)$ then it will follow that $Q_k(o)$ has positive feedback. This is clear from the definition of positive feedback at scale 1 and will follow once we define positive feedback at higher scales. Hence if $Q_k(o)$ has positive feedback and $Q_k(i)$ is bad or has negative feedback, then $\mathcal{P}_k^{i \leftarrow o}$ contains at least one k -box with positive feedback, and thus the progenitor is well-defined.

The definition of progenitor does not on its own provide a useful structure to better understand the structure of boxes with negative feedback. Indeed, it could be the case that a box with negative feedback is its own progenitor. The utility of the definition of progenitor becomes apparent by introducing the following property that we refer to as a *progenitor structure*:

(Prog_k) If $Q_k(i)$ has negative feedback, then $Q_k(i^{\text{par}})$ is either bad or has negative feedback.

If there is a progenitor structure at scale k , then every k -box with negative feedback can be traced back to its progenitor, which must be a bad k -box. Moreover, all k -boxes on the path of parents from a k -box with negative feedback to its progenitor must be bad or have negative feedback. These are key properties that will allow us to control the entrance times of boxes with negative feedback. In Figure 4, we see how to implement the progenitor structure to find the progenitor of a bad box or negative feedback box from the parent structure. The arrows provide the parent structure, and by backwards traversing the arrows from a pink or red box, we arrive at a unique red (bad) box that is the progenitor. We note that it is possible for the parent of a box of positive feedback to have negative feedback or be bad.




Fig. 4. An illustration of the progenitor structure from a bad box. Red boxes represent cores of bad boxes, pink boxes represent the cores of boxes of negative feedback and the white boxes represent cores of boxes of positive feedback. The arrows point to the core of a box from the core of its parent. The unique box with a parent of positive feedback (the leftmost red box) is the progenitor of all the other pink and red boxes.

The following lemma shows that if λ is small enough so that (CAS_1) holds, then (Prog_1) holds, providing the desired progenitor structure at scale 1.

Lemma 4.6. *If $\lambda \in (0, \lambda_{r_1})$, then (Prog_1) holds.*

Proof. Assume $Q_k(i)$ has negative feedback, so that there is no site within $\mathcal{C}_1^-(i)$ occupied by FPP_1 at time $\tau_{1,i} + r_1 L_1$, and let $\lambda \in (0, \lambda_{r_1})$. Suppose for contradiction that $Q_1(i^{\text{par}})$ has positive feedback. Recall that $e_{1,i}$ is the almost surely unique site such that $\tau_{1,i} = \tau(e_{1,i})$ and by definition of parent, $e_{1,i} \in Q_1^{\text{core}}(i^{\text{par}})$. Let $u \notin Q_1(i)$ be the neighbour of $e_{1,i}$ that transmitted either FPP_1 or FPP_λ to $e_{1,i}$ at time $\tau_{1,i}$.

First consider the case where u is occupied by FPP_λ and so $e_{1,i}$ is also occupied by FPP_λ . It follows immediately from the definition of good box that

$$\tau_{1,i} = \tau(e_{1,i}) \geq \tau(u) + 1/\sqrt{\lambda} > \tau_{1,i^{\text{par}}} + 1/\sqrt{\lambda}. \tag{13}$$

As $Q_1(i^{\text{par}})$ is assumed to have positive feedback, there exists some site $z \in \mathcal{C}_1^-(i^{\text{par}})$ such that

$$\tau(z) < \tau_{1,i^{\text{par}}} + r_1 L_1.$$

Fix a site $z_0 \in \mathcal{C}_1^-(i) \cap \mathcal{C}_1^-(i^{\text{par}})$. By Fact 4.1, there exists a path γ_0 from z to z_0 such that

$$\gamma_0 \subset \mathcal{C}_1^-(i^{\text{par}}) \quad \text{and} \quad T_1(\gamma_0) < r_1 L_1.$$

As $Q_1(i^{\text{par}})$ has positive feedback, every site on γ_0 is occupied by FPP_1 by (CAS_1) (cf. Lemma 4.3). Hence

$$\tau_{1,i} \leq \tau(z_0) < \tau(z) + r_1 L_1 < \tau_{1,i^{\text{par}}} + 2r_1 L_1. \tag{14}$$

As $\lambda \in (0, \lambda_{r_1})$, (13) contradicts (14) and so u is instead occupied by FPP_1 . As u neighbours a site in $Q_1^{\text{core}}(i^{\text{par}})$, the distance from u to $\partial Q_1(i^{\text{par}})$ is at least $L_1/6$. By definition of good 1-box we have

$$|\mathcal{C}_2(i^{\text{par}})| \leq \log^2 |Q_1(i^{\text{par}})|$$

and so long as L_1 is large enough, it must be the case that $u \in \mathcal{C}_1^-(i^{\text{par}})$.

Arbitrarily fix a site $v \in \mathcal{C}_1^-(i) \cap \mathcal{C}_1^-(i^{\text{par}})$. As $Q_1(i^{\text{par}})$ is good, by Fact 4.1 there exists a path γ_1 from u to v such that

$$\gamma_1 \subset \mathcal{C}_1^-(i^{\text{par}}) \quad \text{and} \quad T_1(\gamma_1) < r_1 L_1. \tag{15}$$

By Lemma 4.3, as $Q_1(i^{\text{par}})$ has positive feedback, every site on γ_1 is occupied by FPP_1 . Moreover, by (15) we have

$$\tau(v) < \tau(u) + r_1 L_1 < \tau(e_{1,i}) + r_1 L_1 = \tau_{1,i} + r_1 L_1,$$

contradicting the negative feedback of $Q_1(i)$. Thus $Q_1(i^{\text{par}})$ is bad or has negative feedback, establishing the result. ■

4.5. Establishing the fundamental properties at scale 1

As a consequence of Lemma 4.3, it is intuitive that a 1-box having positive feedback induces fast entrance times to nearby good 1-boxes due to the large overlap between neighbouring 1-boxes. We make this notion precise in the following property that gives this control for FPP_1 :

(Fast₁) If $Q_1(i)$ is good and $Q_1(j)$ has positive feedback with $\|i - j\|_\infty = 1$, then $\tau_{1,i} < \tau_{1,j} + 2r_1L_1$.

In the following lemma we prove that if λ is so small that (CAS₁) holds, then (Fast₁) holds as well.

Lemma 4.7. *If $\lambda \in (0, \lambda_{r_1})$, then (Fast₁) holds.*

Proof. Let i, j be such that $Q_1(i)$ is good, $Q_1(j)$ has positive feedback and $\|i - j\|_\infty = 1$. By definition of positive feedback, there exists a site $u \in \mathcal{C}_1^-(j)$ such that u is occupied by FPP_1 and $\tau(u) < \tau_{1,j} + r_1L_1$. Arbitrarily fix a site $v \in \mathcal{C}_1^-(j) \cap \mathcal{C}_1^-(i)$. By Fact 4.1 there exists a path γ from u to v such that $\gamma \subset \mathcal{C}_1^-(j)$ and $T_1(\gamma) < r_1L_1$. As $\lambda \in (0, \lambda_{r_1})$, by Lemma 4.3, all sites on γ are occupied by FPP_1 and thus

$$\tau_{1,i} \leq \tau(v) < \tau(u) + r_1L_1 < \tau_{1,j} + 2r_1L_1. \quad \blacksquare$$

Note that in (Fast₁), we cannot deduce that $Q_1(i)$ has positive feedback. It may be the case that it was entered much earlier and actually has negative feedback but Lemma 4.7 gives us that it is impossible that $Q_1(i)$ is entered significantly later than any positive feedback neighbour.

For any constant $C > 0$, define the property

(Del₁^C) If both $Q_1(i)$ and $Q_1(i^{\text{par}})$ have negative feedback, then $\tau_{1,i} > \tau_{1,i^{\text{par}}} + CL_1$.

In other words, (Del₁^C) guarantees that there is a *delay* of time at least CL_1 in the entrance times of a negative feedback 1-box and its parent if the parent also has negative feedback. In the lemma below, we prove that for any constant $C > 0$, (Del₁^C) holds if λ is sufficiently small, providing us the control on FPP_λ at scale 1 that we will ultimately require.

Lemma 4.8. *For any $C > 0$, if $\lambda \in (0, \lambda_C)$, then (Del₁^C) holds.*

Proof. Suppose that $Q_1(i)$ and $Q_1(i^{\text{par}})$ both have negative feedback and $\lambda \in (0, \lambda_C)$. Recall that $e_{1,i}$ is the almost surely unique site such that $\tau(e_{1,i}) = \tau_{1,i}$. By definition of parent, we have

$$e_{1,i} \in Q_1^{\text{core}}(i^{\text{par}}) \subset Q_1(i).$$

Let $u \notin Q_1(i)$ be the neighbouring site of $e_{1,i}$ that transmitted either FPP_1 or FPP_λ to $e_{1,i}$ at time $\tau_{1,i}$. First note that by construction, u is occupied by either FPP_1 or FPP_λ before $Q_1(i)$ is first entered by either process and so

$$\tau_{1,i} > \tau(u). \tag{16}$$

If u was occupied by FPP_λ , then it follows from the definition of good 1-box that

$$\tau_{1,i} - \tau_{1,i^{\text{par}}} \geq 1/\sqrt{\lambda} > CL_1,$$

where the final inequality follows as $\lambda \in (0, \lambda_C)$.

Suppose instead that u was occupied by the FPP_1 process and in this case, by construction of good 1-box and parent, we have $u \in \mathcal{C}_1^-(i^{\text{par}})$. Fix a site $v \in \mathcal{C}_1^-(i) \cap \mathcal{C}_1^-(i^{\text{par}})$. By Fact 4.1, there exists a path γ from u to v such that

$$\gamma \subset \mathcal{C}_1^-(i^{\text{par}}) \quad \text{and} \quad T_1(\gamma) < r_1L_1.$$

By definition of $Q_1(i)$ having negative feedback, there is no site in $\mathcal{C}_1^-(i)$ occupied by FPP_1 at time $\tau_{1,i} + r_1L_1$. However, if every site on γ was occupied by FPP_1 , then Fact 4.1 would imply that $\tau(v) < \tau_{1,i} + r_1L_1$, giving that $Q_1(i)$ has positive feedback, which is a contradiction. Thus there exists a site $v_0 \in \gamma$ that is occupied by FPP_λ with

$$\tau(v_0) < \tau(u) + r_1L_1. \tag{17}$$

By definition of good 1-box and as v_0 is occupied by FPP_λ , we have

$$\tau(v_0) \geq \tau_{1,i^{\text{par}}} + 1/\sqrt{\lambda}. \tag{18}$$

Putting together (16)–(18), we observe that

$$\tau_{1,i} + r_1L_1 > \tau(u) + r_1L_1 > \tau(v_0) \geq \tau_{1,i^{\text{par}}} + 1/\sqrt{\lambda}.$$

By rearranging the above inequality, we see that

$$\tau_{1,i} - \tau_{1,i^{\text{par}}} \geq 1/\sqrt{\lambda} - r_1L_1 > CL_1,$$

where the final inequality holds as $\lambda \in (0, \lambda_C)$. ■

4.6. Definition of clusters and positive feedback at higher scales

In this section we define positive feedback at higher scales. For this we must specify what it means for a good box to be well-separated from bad boxes. In this direction we introduce the idea of *clusters* of $(k - 1)$ -boxes in a k -box.

Definition 4.9 (Clusters). Let $k \geq 2$ and consider the k -box $Q_k(i)$. We will define the clusters of $Q_k(i)$, which will be subsets of the $(k - 1)$ -boxes contained in $Q_k(i)$. Start by taking the set of all $(k - 1)$ -boxes that have a non-empty intersection with the *bad* $(k - 1)$ -boxes from $Q_k(i)$. Note that the bad $(k - 1)$ -boxes are contained in this set. Next, add to this set all the $(k - 1)$ -boxes that are disconnected from infinity by it. This new set may not be connected, but it can be naturally split into its connected components, which we refer to as the *clusters* of $Q_k(i)$. Let \mathcal{A} be such a cluster. We define the *outer-boundary* of \mathcal{A} , denoted by $\partial\mathcal{A}$, as the set of $(k - 1)$ -boxes $Q_{k-1}(\ell) \notin \mathcal{A}$ for which there exists $Q_{k-1}(t) \in \mathcal{A}$ with $\|\ell - t\|_\infty = 1$. We will abuse notation and sometimes let $\partial\mathcal{A}$ refer to the vertices contained in the $(k - 1)$ -boxes in $\partial\mathcal{A}$, but the meaning will be clear from the context. See Figure 5 for an illustration of clusters.

Fix $k \geq 2$ and suppose $Q_k(i)$ is a good k -box with cluster \mathcal{A} . As there are at most A disjoint bad $(k - 1)$ -boxes in $Q_k(i)$, with the d -dimensional isoperimetric inequality we deduce that there exists a constant $\sigma = \sigma(A, d) \in (0, \infty)$ such that

$$\text{the number of } (k - 1)\text{-boxes contained in } \mathcal{A} \cup \partial\mathcal{A} \leq \sigma.$$

Moreover, since there are at most A clusters in $Q_k(i)$, the total number of $(k - 1)$ -boxes contained in some cluster of $Q_k(i)$ is bounded above by $A\sigma$, which is a constant that only depends on A and d .

With the definition of clusters of a k -box at hand, we have a notion of bad $(k - 1)$ -boxes in a good k -box being *well-separated* from one another; in particular, boxes that belong to different clusters are sufficiently far apart. It will be useful to also have a global notion of well-separated, meaning that a site is sufficiently far from bad boxes at all scales. We make this notion precise in the following definition.

Definition 4.10 (Flawless). Let $x \in \mathbb{Z}^d$, and take i to be such that $x \in Q_1^{\text{core}}(i)$. Define x to be *flawless* if it is not contained in a cluster of a box at any scale and is contained in $\mathcal{C}_1(i)$; recall the definition of $\mathcal{C}_1(i)$ from the beginning of Section 3.1.

Our main results concern certain events occurring with positive probability, namely survival of FPP_1 in Theorem 1.1, and survival of FPP_1 and non-survival of FPP_λ in Theorem 1.3. It transpires that the origin being flawless is the specific event we utilise to prove these results. In the following lemma, we prove that such an event occurs with positive probability for large enough L_1 and small enough λ .

Lemma 4.11. *Let $x \in \mathbb{Z}^d$ and $A > d$. If we choose L_1 large enough and then λ small enough, then the point x is flawless with positive probability.*

Proof. For $k \geq 1$, let Ω_k^x be the collection of k -boxes that contain x . Let $i_0 \in \mathbb{Z}^d$ be such that $x \in Q_1^{\text{core}}(i_0)$. We may write the event that x is not flawless as

$$\{x \text{ is not flawless}\} = \bigcup_{k \geq 1} \bigcup_{Q_k(i) \in \Omega_k^x} H_k(i) \cup \{x \notin \mathcal{C}_1(i_0)\}, \tag{19}$$

where $H_k(i)$ is the event that $Q_k(i)$ is in a cluster of some box in Ω_{k+1}^x . As each cluster in a good $(k + 1)$ -box contains no more than σ k -boxes, we deduce that

$$\begin{aligned} \mathbb{P}(H_k(i)) &\leq \mathbb{P}(\text{there exists a bad } k\text{-box } Q_k(j) \text{ such that } \|i - j\|_\infty \leq 10\sigma) \\ &\leq c\rho_k, \end{aligned} \tag{20}$$

where $c = c(A, d) > 0$ and ρ_k is the probability that an arbitrary k -box is bad. Note that

$$\begin{aligned} \mathbb{P}\left(\bigcap_{k \geq 1} \bigcap_{Q_k(i) \in \Omega_k^x} H_k^c(i) \cap \{x \notin \mathcal{C}_1(i_0)\}\right) &\leq \mathbb{P}(x \notin \mathcal{G}) \\ &\leq 1 - \theta(1 - p). \end{aligned} \tag{21}$$

From (19), (20) and the union bound, we have

$$\begin{aligned} \mathbb{P}(x \text{ is not flawless}) &\leq 1 - \theta(1 - p) + \sum_{k \geq 1} \sum_{Q_k(i) \in \Omega_k^x} c\rho_k \\ &\leq 1 - \theta(1 - p) + c' \sum_{k \geq 1} \rho_k, \end{aligned} \tag{22}$$

where $c' = c'(A, d) > 0$ and $\theta(1 - p) > 0$ as $p < 1 - p_c^{\text{site}}$. By recalling Lemma 3.4, the sum in (22) can be made arbitrarily small by setting ρ_1 small enough in Lemma 3.2, establishing the result. ■

As L_1 and λ in Lemma 4.11 only depend on A, d and p , we may add this dependence to how we set L_1 and λ in Section 4.2.

There is an issue of a cluster being too close to the boundary of a k -box, as it may be part of a much larger cluster outside of that k -box. To address this, we introduce the notion of *boundary clusters*.

Definition 4.12 (Boundary and inner clusters). Let $k \geq 2$, let $Q_k(i)$ be good and let \mathcal{A} be a cluster in $Q_k(i)$. We define \mathcal{A} to be a *boundary cluster* in $Q_k(i)$ if the outer boundary of \mathcal{A} has a non-empty intersection with the boundary of $Q_k(i)$, so that

$$\partial\mathcal{A} \cap \partial Q_k(i) \neq \emptyset.$$

Otherwise, we define \mathcal{A} to be an *inner cluster* in $Q_k(i)$.

In Figure 5, the two uppermost clusters intersect the boundary clusters and the rest are inner clusters.

We let $\mathcal{W}(k, i)$ be the set of $(k - 1)$ -boxes in a good k -box $Q_k(i)$ that are not contained in any clusters and call such $(k - 1)$ -boxes *wonderful*. In other words, if $Q_k(i)$ is good and $\mathcal{A}_1, \dots, \mathcal{A}_n$ are all the clusters of bad $(k - 1)$ -boxes in $Q_k(i)$, then

$$\mathcal{W}(k, i) = \left\{ Q_{k-1}(\ell) : Q_{k-1}(\ell) \subset Q_k(i) \text{ and } Q_{k-1}(\ell) \not\subset \bigcup_{j=1}^n \mathcal{A}_j \right\}.$$

Before we give the next definition, it will be useful to also define a collection of boxes. For all $k \geq 1$, let

$$\{Q_k^{\text{inn}}(i)\}_{i \in \mathbb{Z}^d} \quad \text{with} \quad Q_k^{\text{inn}}(i) = (L_k/3)i + [-499L_k/1000, 499L_k/1000]^d, \tag{23}$$

where we refer to $Q_k^{\text{inn}}(i)$ as the *inner part* of $Q_k(i)$. In some cases we will abuse notation and instead mean that $Q_k^{\text{inn}}(i)$ refers to all the $(k - 1)$ -boxes contained in the set given by (23) rather than the sites but the context will always make it unambiguous. Intuitively, the inner part of a k -box contains all sites that are sufficiently far from the boundary of the k -box. Indeed, so long as the distance from sites in the inner part to the boundary is of the order L_k , our analysis will work and so the choice of $L_k/1000$ is purely for concreteness.




Fig. 5. A k -box $Q_k(i)$ with clusters given in red and their respective outer boundary in pink.

Definition 4.13 ($\mathcal{W}^{\text{inn}}(k, i)$). Let $k \geq 2$ and let $Q_k(i)$ be good. Let $\mathfrak{A}(k, i)$ be the collection of all clusters in $Q_k(i)$ whose outer boundary has a non-empty intersection with $Q_k^{\text{inn}}(i)$, so that if $\mathcal{A} \in \mathfrak{A}(k, i)$, then

$$\partial\mathcal{A} \cap Q_k^{\text{inn}}(i) \neq \emptyset.$$

Define $\mathcal{W}^{\text{inn}}(k, i)$ to be the union of all wonderful $(k - 1)$ -boxes in $Q_k^{\text{inn}}(i)$ along with the outer boundary of clusters in $\mathfrak{A}(k, i)$, so that

$$\mathcal{W}^{\text{inn}}(k, i) = (\mathcal{W}(k, i) \cap Q_k^{\text{inn}}(i)) \cup \bigcup_{\mathcal{A} \in \mathfrak{A}(k, i)} \partial\mathcal{A}.$$

It transpires that $\mathcal{W}^{\text{inn}}(k, i)$ is what we choose to replace $C_1^-(i)$ at higher scales. Roughly speaking, we will tune the definition of positive feedback at higher scale so that if there is a $(k - 1)$ -box with positive feedback in $\mathcal{W}^{\text{inn}}(k, i)$ at a relatively early time compared to when either FPP_1 or FPP_λ first enters $Q_k(i)$, then all $(k - 1)$ -boxes in $\mathcal{W}^{\text{inn}}(k, i)$ will have positive feedback. This is made rigorous later in Lemma 4.17.

Remark 4.14. It is worth examining why we choose to define $\mathcal{W}^{\text{inn}}(k, i)$ as we did in Definition 4.13 rather than just setting $\mathcal{W}^{\text{inn}}(k, i)$ to be $\mathcal{W}(k, i) \cap Q_k^{\text{inn}}(i)$. The issue is that $\mathcal{W}(k, i) \cap Q_k^{\text{inn}}(i)$ may not be a single connected component of wonderful $(k - 1)$ -boxes in $Q_k(i)$. It will be essential in later proofs that we can find paths of $(k - 1)$ -boxes through $\mathcal{W}^{\text{inn}}(k, i)$. To be explicit, for $k \geq 1$, we say that $\pi_k = \{Q_k(i_1), \dots, Q_k(i_n)\}$ is a *path of k -boxes* if $\|i_{\ell+1} - i_\ell\|_\infty = 1$ for all $1 \leq \ell < n$, and we write $|\pi_k| = n$. If $\mathcal{W}^{\text{inn}}(k, i)$ is not a single connected component, then this may not be possible without exiting $\mathcal{W}^{\text{inn}}(k, i)$. By including the outer boundaries of any clusters that intersect $Q_k^{\text{inn}}(i)$, we guarantee that $\mathcal{W}^{\text{inn}}(k, i)$ is a single connected component. Hence, if

$$Q_{k-1}(j_0), Q_{k-1}(j_1) \in \mathcal{W}^{\text{inn}}(k, i),$$

then there exists a path π_{k-1} of $(k - 1)$ -boxes from $Q_{k-1}(j_0)$ to $Q_{k-1}(j_1)$ such that

$$\pi_{k-1} \subset \mathcal{W}^{\text{inn}}(k, i) \quad \text{and} \quad |\pi_{k-1}| \leq \frac{3dL_k}{L_{k-1}} + A\sigma.$$

Definition 4.15 (Positive feedback at higher scales). We define positive and negative feedback at higher scales inductively. Let $k \geq 2$, suppose $Q_k(i)$ is good, and positive and negative feedbacks are defined at scale $k - 1$. Define

$$r_k = r_1 \prod_{j=2}^k \left(1 + \frac{a_1}{j^2 L_{j-1}^{d-1}} \right), \tag{24}$$

where r_1 is as given in Fact 4.1 and $a_1 = 1000A\sigma/3$. Note that a_1 depends only on A and d . Recall the definition of entrance time for a k -box as given in (10). If there exists $Q_{k-1}(\ell) \in \mathcal{W}^{\text{inn}}(k, i)$ with positive feedback such that

$$\tau_{k-1,\ell} < \tau_{k,i} + \frac{3}{500} r_k L_k,$$

then we define $Q_k(i)$ to have *positive feedback*. If $Q_k(i)$ does not have positive feedback, then we say it has *negative feedback*.

Remark 4.16. The constant r_k is the analogue of r_1 at scale k , and (24) is justified when we generalise (Fast₁) to higher scales. From (24), we deduce that $\{r_k\}_{k \geq 1}$ is an increasing sequence such that

$$r = \lim_{k \rightarrow \infty} r_k < \infty.$$

Moreover, as long as L_1 is sufficiently large, we have $r < 2r_1$. We assume this is the case, and note that this constraint for L_1 depends on a_1 , which only depends on A and d . Hence, we can add this constraint to how we set L_1 in Section 4.2.

4.7. Fundamental properties at higher scales

In this section we generalise the fundamental properties (CAS₁), (Fast₁) and (Del₁^C) to higher scales; recall that we have already defined the fundamental property (Prog_k) at all scales. We will also introduce a fifth fundamental property that will give information about how FPP_λ spreads through clusters in a good box.

Recall that (CAS₁) states that all sites in the large component of non-seeds in a good 1-box away from the boundary are occupied by FPP_1 and that this property follows from positive feedback by setting λ small enough, as stated in Lemma 4.3. The higher scale version of (CAS₁) is given below. For $k \geq 2$ define the following property:

(CAS_k) If $Q_k(i)$ has positive feedback, then for all i_0 such that $Q_{k-1}(i_0) \in \mathcal{W}^{\text{inn}}(k, i)$, $Q_{k-1}(i_0)$ has positive feedback.

This section contains two main results: the first one, stated below, shows that by setting λ small enough, (CAS_k) holds for all $k \geq 2$. The second main result will be given later, in Lemma 4.22.

Lemma 4.17. *There exists a constant $C^* > 0$ such that if $\lambda \in (0, \lambda_{C^*})$, then (CAS_k) holds for all $k \geq 1$.*

By Lemma 4.3 we know that we may set λ so that (CAS_1) holds but the issue is that if λ is such that (CAS_k) holds, it is not immediately clear how one should set λ to ensure that (CAS_{k+1}) holds. Indeed if this procedure requires that λ tends to 0 as $k \rightarrow \infty$ then this is not sufficient to prove Lemma 4.17. Hence the proof of Lemma 4.17 will require careful consideration of how (CAS_k) implies (CAS_{k+1}) and choosing λ in such a way that it does not depend on the scale k .

The first step in proving Lemma 4.17 is generalising $(Fast_1)$ and (Del_1^C) to higher scales. Then we introduce the notion of the *confinement* of bad clusters to account for the existence of bad boxes in good boxes at higher scales. Once these ideas have been formalised, we prove that if λ is such that (CAS_k) holds and is below a certain constant we define later that does not depend on k , then $(Prog_k)$, $(Fast_k)$ and (Del_k^C) all hold; this will be the content of the second main result of this section, Lemma 4.22. Finally, we show that if all such properties hold up to scale k , then (CAS_{k+1}) holds for a possible smaller value of λ . Noting that the required upper bound on λ converges to a value that is bounded away from 0 as k goes to infinity establishes Lemma 4.17.

In order to establish the fundamental properties at scale $k \geq 2$, we need to pay careful attention to the presence of bad boxes at lower scales, in which we cannot control either FPP_1 or FPP_λ . For example, it could happen that FPP_1 is *slowed down* in some sense as it has to deviate around bad boxes to spread, whereas FPP_λ may be *sped up* by being able to spread through bad boxes at no cost. If the slowing down or speeding up of the respective processes can accumulate in a non-trivial way, then we may lose control of both the processes. The idea is that there are relatively few bad $(k - 1)$ -boxes in a good k -box, so any slowing down or speeding up at scale $(k - 1)$ is minuscule relative to scale k .

For $k \geq 2$ we define the following property that generalises $(Fast_1)$ to higher scales:

$(Fast_k)$ If $Q_k(i)$ is good and $Q_k(j)$ has positive feedback with $\|i - j\|_\infty = 1$, then

$$\tau_{k,i} < \tau_{k,j} + 2r_k L_k,$$

where r_k is as defined in (24).

Next we generalise (Del_1^C) to higher scales. For $k \geq 2$ and any constant $C > 0$, define the following property:

(Del_k^C) If both $Q_k(i)$ and $Q_k(i^{par})$ have negative feedback, then $\tau_{1,i} > \tau_{1,i^{par}} + C\omega_k L_k$, where

$$\omega_k = \prod_{m=2}^k \left(1 - \frac{a_2}{m^2 L_{m-1}^{d-1}} \right), \tag{25}$$

and $a_2 > 0$ is a constant that only depends on A, d and p that we set later in the proof of Lemma 4.22.

Remark 4.18. From (25), we deduce that $\{\omega_k\}_{k \geq 1}$ is a decreasing sequence with

$$\omega = \lim_{k \rightarrow \infty} \omega_k > 0.$$

The sequence $\{\omega_k\}_{k \geq 1}$ accounts for the possible increase of speed that FPP_λ gains at high scales due to the existence of bad boxes at lower scales. From (25), we deduce that if L_1 is sufficiently large, then $\omega > 1/2$. As in Remark 4.16, since a_2 depends only on A , d and p , we add this constraint on L_1 to how we set it in Section 4.2.

We now introduce the concept of *confinement* of a cluster, which will allow us to control the spread of FPP_λ from clusters in a good box at high scales. In particular, the confinement of clusters in a good box is a crucial ingredient in proving that (Del_k^C) holds for $k \geq 2$. For this we will use the notion of inner clusters that was introduced in Definition 4.12.

Definition 4.19 (Successful confinement of inner clusters). Let $k \geq 2$, $Q_k(i)$ be good and \mathcal{A} be an inner cluster in $Q_k(i)$. If there exists a $(k-1)$ -box $Q_{k-1}(j)$ with negative feedback such that

$$Q_{k-1}(j) \in \partial\mathcal{A} \quad \text{and} \quad Q_{k-1}(j^{\text{prog}}) \in \mathcal{A},$$

then we define \mathcal{A} to be *poorly confined*. If \mathcal{A} is not poorly confined, we define it to be *successfully confined*.

Note that, because of the progenitor structure (that is, if (Prog_{k-1}) holds), then we deduce that the existence of any $(k-1)$ -box $Q_{k-1}(t) \notin \mathcal{A}$ of negative feedback (even with $Q_{k-1}(t) \notin \partial\mathcal{A}$) for which $Q_{k-1}(t^{\text{prog}}) \in \mathcal{A}$ implies that \mathcal{A} is poorly confined.

Remark 4.20. Let $k \geq 2$ and suppose $Q_k(i)$ has positive feedback. If (CAS_k) holds then we deduce that if \mathcal{A} is an inner cluster of $Q_k(i)$ that intersects $Q_k^{\text{inn}}(i)$ then it must be successfully confined, because all $(k-1)$ -boxes in $\partial\mathcal{A}$ have positive feedback as $\partial\mathcal{A} \subset \mathcal{W}^{\text{inn}}(k, i)$. But our proof will proceed in the opposite direction, that is, we will use successful confinement of clusters inside a k -box to establish (CAS_k) .

In high-level terms, if a cluster \mathcal{A} is successfully confined, then \mathcal{A} contains all $(k-1)$ -boxes of negative feedback that have a progenitor in \mathcal{A} . In some sense, \mathcal{A} being successfully confined “contains” the effect of the bad boxes in \mathcal{A} . A key point of our analysis is understanding the conditions under which an inner cluster can be poorly confined.

We now provide a different perspective on poorly confined inner clusters. The idea is that any poorly confined inner cluster must have a box of negative feedback in its boundary with a progenitor in *another* cluster. This will allow us to show that a poorly confined cluster must cause a sequence of poorly confined clusters; so a poorly confined cluster inside a k -box will have an effect that is in some sense proportional to its distance to the boundary of the k -box.

We define the *source* of a poorly confined inner cluster as the box of negative feedback in the outer boundary with earliest entrance time (regardless of where its progenitor is located). The source serves as a “witness” to the poor confinement of the cluster.

Definition 4.21 (Source of a poorly confined inner cluster). Let $k \geq 1$ and \mathcal{A} be a poorly confined inner cluster of a good $(k+1)$ -box. Set $Q_k(s)$ be the k -box in $\partial\mathcal{A}$ with negative

feedback that has the smallest entrance time, so that

$$\tau_{k,s} = \inf \{ \tau_{k,h} : Q_k(h) \in \partial\mathcal{A} \text{ and } Q_k(h) \text{ has negative feedback} \},$$

noting that this infimum is over a non-empty set as \mathcal{A} is poorly confined.

As we will see later, it is desirable that the progenitor of the source of a poorly confined inner cluster is not contained in that cluster and is entered relatively soon after the outer boundary of the cluster is entered for the first time. These properties are given below:

(Conf_k) If $Q_{k+1}(i)$ is good and \mathcal{A} is a poorly confined inner cluster in $Q_{k+1}(i)$ with source $Q_k(s) \in \partial\mathcal{A}$, then $Q_k(s^{\text{prog}}) \notin \mathcal{A}$ and $\tau_{k,s} < \tau_{k,b} + 2\sigma r_k L_k$, where $\tau_{k,b} = \inf \{ \tau_{k,b'} : Q_k(b') \in \partial\mathcal{A} \}$.

To prove that the fundamental properties hold at higher scales for an appropriate choice of λ it will be useful to introduce the property (\mathbf{P}_k^C) at scale k , which states that for a constant $C > 0$ all the fundamental properties hold at all scales up to and including k :

(\mathbf{P}_k^C) Properties (CAS_j), (Prog_j), (Fast_j), (Del_j^C) and (Conf_j) hold for all $j \in \{1, \dots, k\}$.

The reason we introduce (\mathbf{P}_k^C) is that to prove some fundamental property at higher scales, we will need to use other fundamental properties at that scale and lower.

Recall that in Lemma 4.17 we wish to set C large enough so that if $\lambda \in (0, \lambda_C)$, then we may deduce that (CAS_k) holds for all $k \geq 1$. The issue is that in choosing C large enough so that if $\lambda \in (0, \lambda_C)$ then (CAS_k) holds, it is not immediate that (CAS_{k+1}) holds too. It transpires that some of the fundamental properties require C to be sufficiently large to deduce that they hold at higher scales. The key point is that we may choose C so that it does not depend on k or L_1 . In fact, we now define a threshold that C needs to be above to allow for the first half of the inductive argument for Lemma 4.17. Define

$$C_0 = 8\sigma r_1 a_3, \tag{26}$$

where $a_3 > 1$ is a constant that only depends on A and d that we set later in the proof of Claim 4.27. The reason for this choice of C_0 will only become apparent in later proofs but the reader should note that C_0 only depends on A , d and p .

The next lemma is the second main result of this section, essentially giving that (CAS_k) and a proper choice of λ imply the fundamental properties at scale k .

Lemma 4.22. *Let $k \geq 2$ and $C > C_0$. If λ is such that (\mathbf{P}_{k-1}^C) and (CAS_k) hold, then (\mathbf{P}_k^C) holds.*

Lemma 4.22 provides a key step in the inductive argument in the proof of Lemma 4.17. One subtlety is that we have not yet proved that (Conf₁) holds but the proof at higher scales transpires to be completely analogous. To streamline later proofs, we now prove that (Conf_k) holds assuming that other fundamental properties at that scale hold.

Lemma 4.23. *Let $k \geq 1$ and $C > C_0$. If λ is such that (Prog_k), (Fast_k) and (Del_k^C) hold, then (Conf_k) holds.*

Proof. Let \mathcal{A} be a poorly confined inner cluster of a good $(k + 1)$ -box with source $Q_k(s)$. Let $Q_k(b)$ be the k -box in $\partial\mathcal{A}$ with the smallest entrance time, so that $\tau_{k,b} = \inf \{ \tau_{k,b'} : Q_k(b') \in \partial\mathcal{A} \}$. As λ is such that (Fast_k) holds, we have

$$\tau_{k,s} < \tau_{k,b} + 2\sigma r_k L_k, \tag{27}$$

because there must exist a k -box with negative feedback within $\partial\mathcal{A}$ by time $\tau_{k,b} + 2\sigma r_k L_k$. Indeed, if no such k -box existed, then every k -box in $\partial\mathcal{A}$ would have positive feedback as (Fast_k) holds and the number of k -boxes in $\partial\mathcal{A}$ is bounded above by σ . This would give a contradiction as \mathcal{A} is assumed to be poorly confined.

It remains to prove that $Q_k(s^{\text{prog}}) \notin \mathcal{A}$. Suppose for contradiction that $Q_k(s^{\text{prog}}) \in \mathcal{A}$ and set $Q_k(j)$ to be the bad k -box in \mathcal{A} with earliest entrance time, so that

$$\tau_{k,j} = \inf \{ \tau_{k,j'} : Q_k(j') \in \mathcal{A} \text{ and } Q_k(j') \text{ is bad} \}.$$

Clearly $\tau_{k,j} \leq \tau_{k,s^{\text{prog}}}$ as we assume that $Q_k(s^{\text{prog}})$ is a bad k -box in \mathcal{A} . By construction we have

$$\tau_{k,b} < \tau_{k,j} \leq \tau_{k,s^{\text{prog}}} < \tau_{k,s}.$$

Moreover, as λ is such that (Prog_k) and (Del_k^C) hold, we have

$$\tau_{k,s} > \tau_{k,s^{\text{prog}}} + C\omega_k L_k > \tau_{k,b} + C\omega_k L_k. \tag{28}$$

If C is large enough so

$$C > 2\sigma r_k / \omega_k, \tag{29}$$

then (28) contradicts (27) and thus $Q_k(s^{\text{prog}}) \notin \mathcal{A}$. By Remarks 4.16 and 4.18, for a large enough L_1 we have $r < 2r_1$ and $\omega > 1/2$ and so if $C > 8\sigma r_1$ then (29) holds for all $k \geq 1$. As this follows immediately from the fact that $C > C_0$, the result is established. ■

Before turning to the proof of Lemma 4.17, we must outline an important consequence that can be deduced from the fundamental properties that provide a lower bound on the entrance time for wonderful boxes that have negative feedback in a good box. The idea is that the poor confinement of an inner cluster in a good box can be traced through the parent structure until reaching the boundary or a boundary cluster of that box. By carefully considering the progenitor structure given by (Prog_k) and also (Conf_k) , we deduce a lower bound on entrance times by the number of boxes of negative feedback on this parent structure. We make this idea rigorous below.

Assume $C > C_0$ and λ is such that (P_k^C) holds. Let $Q_{k+1}(i)$ be good and \mathcal{A}_1 be a poorly confined inner cluster in $Q_{k+1}(i)$ with source $Q_k(s_1)$ that is the witness of the poor confinement of \mathcal{A}_1 . As λ is such that (Conf_k) holds, $Q_k(s_1^{\text{prog}})$ is not contained in \mathcal{A}_1 . If $Q_k(s_1^{\text{prog}})$ is contained in another inner cluster of $Q_{k+1}(i)$, say \mathcal{A}_2 , then \mathcal{A}_2 is also poorly confined since λ is such that (Prog_k) holds. The inner cluster \mathcal{A}_2 has source $Q_k(s_2)$ say. From this point onwards, we can search for the progenitor of $Q_k(s_2)$ and iterate this procedure until the progenitor of a source is either outside of $Q_{k+1}(i)$ or is contained in a boundary cluster in $Q_{k+1}(i)$. This procedure is illustrated in Figure 6 in the case where it terminates when the progenitor of a source is not contained in $Q_{k+1}(i)$.




Fig. 6. An illustration of the path of jumps construction. The pink regions represents poorly confined inner clusters in $Q_{k+1}(i)$. The black arrows represent the path of parents from the sources of the poorly confined inner clusters to their progenitor. The vertical black line represents a piece of the boundary of $Q_{k+1}(i)$, whose interior is to the left of the vertical line.

From this construction we discover a sequence of poorly confined inner clusters $\mathcal{A}_1, \dots, \mathcal{A}_n$ with respective sources $Q_k(s_1), \dots, Q_k(s_n)$ where $Q_k(s_n^{\text{prog}})$ is contained outside of $Q_{k+1}(i)$ or is contained in a boundary cluster of $Q_{k+1}(i)$. Call the collection of paths of parents

$$\mathcal{P}_k^{s_1 \leftarrow s_1^{\text{prog}}}, \dots, \mathcal{P}_k^{s_n \leftarrow s_n^{\text{prog}}},$$

a *path of jumps*; we refer to a *jump* when we concatenate s_i^{prog} with s_{i+1} for each $i \in \{1, \dots, n - 1\}$. Hence, for any poorly confined inner cluster in a good box we can construct a path of jumps through the progenitor structure that terminates when it discovers a progenitor not contained in an inner cluster. Utilising the path of jumps construction and (Del_k^C) we may deduce a useful lower bound on the entrance for wonderful boxes of negative feedback in a good box as given in Lemma 4.24.

Before we state Lemma 4.24 we introduce the following notation. For $B, B' \subset \mathbb{Z}^d$, we define

$$\text{dist}(B, B') = \inf_{x \in B, y \in B'} \|x - y\|_1.$$

If $Q_k(i)$ and $Q_k(i')$ are k -boxes such that $Q_k(i) \subset B$, $Q_k(i') \subset B'$ and the path of parents $\mathcal{P}_k^{i \leftarrow i'}$ is well-defined, then it is easy to verify that

$$\text{the number of } k\text{-boxes in } \mathcal{P}_k^{i \leftarrow i'} \geq \frac{3}{L_k} \text{dist}(B, B').$$

Lemma 4.24. *Let $k \geq 1$ and $C > C_0$. Suppose λ is such that (P_k^C) holds. If $Q_{k+1}(i)$ is good and $Q_k(j_0) \in \mathcal{W}(k + 1, i)$ has negative feedback, then*

$$\tau_{k,j_0} > \tau_{k+1,i} + C\omega_k L_k \left(\frac{3}{L_k} \text{dist}(Q_k(j_0), \partial Q_{k+1}(i)) - a_4 \right), \tag{30}$$

where $a_4 > 0$ is a constant that only depends on A and d .

Proof. Suppose $Q_{k+1}(i)$ is good and $Q_k(j_0) \in \mathcal{W}(k + 1, i)$ has negative feedback. First consider the case where $Q_k(j_0^{\text{prog}})$ is contained in an inner cluster of $Q_{k+1}(i)$, say \mathcal{A}_1 . As λ is such that (P_k^C) holds, \mathcal{A}_1 must be poorly confined and so by the path of jumps construction above, there exists a sequence of poorly confined inner clusters in $Q_{k+1}(i)$ enumerated as $\mathcal{A}_1, \dots, \mathcal{A}_n$ with respective sources $Q_k(s_1), \dots, Q_k(s_n)$ so that the following properties hold:

- $Q_k(j_0^{\text{prog}}) \in \mathcal{A}_1$,
- $Q_k(s_n^{\text{prog}})$ is not contained in an inner cluster in $Q_{k+1}(i)$,
- $Q_k(s_m^{\text{prog}}) \in \mathcal{A}_{m+1}$ for $1 \leq m < n$.

By definition of source, no cluster is repeated in this construction and $n \leq A$ as there are at most A clusters in a good k -box.

From the path of jumps we outline the following procedure to deduce a lower bound on the entrance time for $Q_k(j_0)$. Start following the path of parents from $Q_k(j_0)$ to $Q_k(j_0^{\text{prog}})$ and stop once a k -box in $\partial \mathcal{A}_1$ is discovered, say $Q_k(j_0^*)$, which must have negative feedback as λ is such that (Prog_k) holds. By definition of source and Lemma 4.23, we have $\tau_{k,s_1} < \tau_{k,j_0^*}$. Then jump to $Q_k(s_1)$ and follow its path of parents to $Q_k(s_1^{\text{prog}})$, stopping once a k -box in $\partial \mathcal{A}_2$ is discovered, say $Q_k(s_1^*)$, which again must have negative feedback with $\tau_{k,s_2} < \tau_{k,s_1^*}$. Next, jump to $Q_k(s_2)$ and proceed in this way until we reach $Q_k(s_n)$, from which we follow its path of parents until hitting the boundary of $Q_{k+1}(i)$ or $\partial \mathcal{A}_n$, which in this latter case must be a boundary cluster of $Q_{k+1}(i)$ and we set $Q_{k-1}(s_n^*)$ as before. In the former case we set $Q_k(s_n^*)$ to be the first k -box on the path of parents starting from $Q_k(s_n)$ whose parent is not contained in $Q_{k+1}(i)$. In either case it is clear that $Q_{k+1}(i)$ is entered before $Q_k(s_n^*)$. Through this construction we deduce

$$\tau_{k,s_1} < \tau_{k,j_0^*}, \quad \tau_{k+1,i} \leq \tau_{k,s_n^*} \quad \text{and} \quad \tau_{k,s_{m+1}} < \tau_{k,s_m^*} \quad \text{for all } 1 \leq m < n. \tag{31}$$

Note that in the construction above there are at most A jumps, since each jump occurs in a bad cluster in $Q_{k+1}(i)$ and clusters cannot be repeated due to the ordering of the entrance times of the sources as given in (31). Note also that each jump has length at most σ , since the jump is between two k -boxes contained in the outer boundary of the same cluster. Hence the number of k -boxes on the paths constructed above is bounded below by

$$\frac{3}{L_k} \text{dist}(Q_k(j_0), \partial Q_{k+1}(i)) - a, \tag{32}$$

where $a > 0$ is a constant that depends only on A and d .

Every k -box on these paths is either bad or has negative feedback as λ is such that (Prog_k) holds and the number of such bad boxes is bounded above by a constant that only depends on A and d . For every pair of boxes with negative feedback on these paths where one is the parent of the other, we deduce that there is a delay of at least $C\omega_k L_k$ in the respective entrance times of the boxes as λ is such that (Del_k^C) holds. From (32), we deduce the number of such pairs is bounded below by

$$\frac{3}{L_k} \text{dist}(Q_k(j_0), \partial Q_{k+1}(i)) - a', \tag{33}$$

where $a' \geq a$ is a constant that depends only on A and d .

Combining (33) and (31) with the delay of at least $C\omega_k L_k$ between boxes with negative feedback that have a negative feedback parent, we deduce

$$\begin{aligned} \tau_{k,j_0} &> \tau_{k,s_n^*} + C\omega_k L_k \left(\frac{3}{L_k} \text{dist}(Q_k(j_0), \partial Q_{k+1}(i)) - a' \right), \\ &\geq \tau_{k+1,i} + C\omega_k L_k \left(\frac{3}{L_k} \text{dist}(Q_k(j_0), \partial Q_{k+1}(i)) - a' \right). \end{aligned}$$

Now consider the case where $Q_k(j_0^{\text{prog}})$ is either not contained in $Q_{k+1}(i)$ or contained in a boundary cluster of $Q_{k+1}(i)$, \mathcal{B} say. In this scenario set $Q_k(j_0^*)$ to be the first k -box on the path of parents $\mathcal{P}_k^{j_0 \leftarrow o}$ whose parent is not contained in $Q_{k+1}(i)$ or the first box that is contained in \mathcal{B} . In this case, the result follow by an analogous argument to the previous case where we instead consider the path of parents $\mathcal{P}_k^{j_0 \leftarrow j_0^*}$, noting that no jumps need be considered. The lemma follows by setting $a_4 = a'$. ■

To conclude the analysis of the fundamental properties, we are left with the task of proving Lemmas 4.17 and 4.22. This will be done in the next two sections.

4.8. Proof of the cascading lemma (Lemma 4.17)

In this section, we give the proof for Lemma 4.17 that provides the cascading argument required in the proof of Theorem 1.1. For this proof we assume that Lemma 4.22 holds, which will be established in the next section. The main step of the proof is given in the following claim, from which Lemma 4.17 readily follows.

Claim 4.25. *There exists a constant $C_1 > 0$ that only depends on p and d such that the following holds. For all $k \geq 1$ and $C > \max\{C_0, C_1\}$, if λ is such that (P_k^C) holds, then (CAS_{k+1}) holds.*

Proof of Lemma 4.17. Assuming Claim 4.25 holds, let $C^* = \max\{r_1, C_0, C_1\}$, $C > C^*$ and $\lambda \in (0, \lambda_C)$. Then Lemmas 4.3, 4.6, 4.7, 4.8 and 4.23 give us that (P_1^C) holds. Claim 4.25 immediately implies that (CAS_2) also holds. Lemma 4.22 then gives us for this choice of C and λ that (P_2^C) also holds. The choice of C and λ does not depend on k and so by repeatedly using Claim 4.25 and Lemma 4.22, we see that if $C > C^*$ and $\lambda \in (0, \lambda_C)$, then (CAS_k) holds for all $k \geq 1$. ■

Proof of Claim 4.25. Let $k \geq 1$ and $C > C_0$. Assume that λ is such that (P_k^C) holds and $Q_{k+1}(i)$ has positive feedback. Suppose for contradiction that there exists some $Q_k(j_0) \in \mathcal{W}^{\text{inn}}(k+1, i)$ that has negative feedback. Without losing generality, we may assume that $Q_k(j_0)$ is the k -box in $\mathcal{W}^{\text{inn}}(k+1, i)$ of negative feedback with the smallest entrance time, so that

$$\tau_{k,j_0} = \inf \{ \tau_{k,h} : Q_k(h) \in \mathcal{W}^{\text{inn}}(k+1, i) \text{ and } Q_k(h) \text{ has negative feedback} \}.$$

As we assume $Q_{k+1}(i)$ has positive feedback, there exists $Q_k(i_0) \in \mathcal{W}^{\text{inn}}(k+1, i)$ such that

$$\tau_{k,i_0} < \tau_{k+1,i} + \frac{3}{500} r_{k+1} L_{k+1}.$$

By Remark 4.14, we may fix a path of k -boxes Ξ_k from $Q_k(i_0)$ to $Q_k(j_0)$ so that $\Xi_k \subset \mathcal{W}^{\text{inn}}(k+1, i)$ and

$$|\Xi_k| \leq 3dL_{k+1}/L_k + A\sigma.$$

If all k -boxes on Ξ_k have positive feedback with the exception of $Q_k(j_0)$, then (Fast $_k$) implies that

$$\begin{aligned} \tau_{k,j_0} &< \tau_{k,i_0} + 2r_k L_k |\Xi_k|, \\ &\leq \tau_{k,i_0} + 2r_k L_k (3dL_{k+1}/L_k + A\sigma). \end{aligned}$$

Recalling that $Q_{k+1}(i)$ has positive feedback and $\tau_{k,i_0} < \tau_{k+1,i} + \frac{3}{500} r_{k+1} L_{k+1}$, from (24) and the above we deduce that

$$\begin{aligned} \tau_{k,j_0} &< \tau_{k+1,i} + \frac{3}{500} r_{k+1} L_{k+1} + 6dr_k L_{k+1} \left(1 + \frac{A\sigma}{3d(k+1)^2 L_k^{d-1}} \right), \\ &< \tau_{k+1,i} + \left(\frac{3}{500} + 6d \right) r_{k+1} L_{k+1}. \end{aligned} \tag{34}$$

With an upper bound for τ_{k,j_0} established in (34), our aim is to establish a corresponding lower bound for τ_{k,j_0} that will give a contradiction.

By definition of $\mathcal{W}^{\text{inn}}(k+1, i)$, if $Q_k(q) \in \Xi_k$, then

$$\frac{3}{L_k} \text{dist}(Q_k(q), \partial Q_{k+1}(i)) \geq \frac{3L_{k+1}}{1000L_k} - \sigma. \tag{35}$$

By (35), (Del $_k^C$) and Lemma 4.24, for all $Q_k(q) \in \Xi_k$, we have

$$\tau_{k,q} > \tau_{k+1,i} + C\omega_k L_k \left(\frac{3L_{k+1}}{1000L_k} - a_5 \right), \tag{36}$$

where $a_5 > 0$ is a constant that only depends on A and d . From the above inequality we deduce

$$\begin{aligned} \tau_{k,q} &= \tau_{k+1,i} + \frac{3}{1000} C\omega_k L_{k+1} \left(1 - \frac{1000a_5}{3(k+1)^2 L_k^{d-1}} \right) \\ &> \tau_{k+1,i} + \frac{1}{1000} C\omega_{k+1} L_{k+1}, \end{aligned} \tag{37}$$

where the final inequality will hold so long as L_1 is sufficiently large and by recalling the definition of ω_{k+1} in (25). This requirement on L_1 only depends on a_5 and thus can be added to the conditions on setting L_1 in Section 4.2.

For $k \geq 1$, set

$$\tilde{C}_{k+1} = 6(1 + 1000d)r_{k+1}/\omega_{k+1}.$$

If $C > \max \{C_0, \tilde{C}_{k+1}\}$ then all k -boxes on Ξ_k are necessarily positive feedback as otherwise (34) contradicts (37). However, this leads to a contradiction, as then $Q_{k-1}(j_0)$ must also have positive feedback. Hence if $C > \max \{C_0, \tilde{C}_{k+1}\}$ and λ is such that (P_k^C) holds, then (CAS_{k+1}) also holds.

To remove the dependence on k in setting λ , we desire that

$$C > 6(1 + 1000d)r/\omega, \tag{38}$$

where r is as given in Remark 4.16 and ω as in Remark 4.18. Recall that so long as L_1 is sufficiently large, we have $r < 2r_1$ and $\omega > 1/2$. Thus, if we set

$$C_1 = 24(1 + 1000d)r_1,$$

and let $C > C_1$, then (38) holds. Consequently, if $C > C_1$, then $C > \tilde{C}_{k+1}$ for all $k \geq 1$, establishing the claim. ■

4.9. Proofs of fundamental properties at higher scales (Lemma 4.22)

To ease exposition, we split the proof into four parts, one for each fundamental property to be established. We also note that for this proof we shall not assume Lemma 4.17, as the proof of Lemma 4.17 uses Lemma 4.22.

Proof of Lemma 4.22 for (Prog_k) . Let $k \geq 2$ and $C > C_0$. Assume that λ is such that (P_{k-1}^C) and (CAS_k) hold. Suppose that $Q_k(i)$ has negative feedback. Hence, if $Q_{k-1}(\ell) \in \mathcal{W}^{\text{inn}}(k, i)$ has positive feedback, then

$$\tau_{k-1,\ell} \geq \tau_{k,i} + \frac{3}{500}r_k L_k. \tag{39}$$

Suppose for contradiction that $Q_k(i^{\text{par}})$ has positive feedback. The strategy of this proof is to construct a path of wonderful $(k - 1)$ -boxes from where FPP_1 enters $Q_k(i)$ to $\mathcal{W}^{\text{inn}}(k, i)$ and use the fact that all $(k - 1)$ -boxes on this path must have positive feedback by the positive feedback of $Q_k(i^{\text{par}})$ and (CAS_k) . This will imply that $Q_k(i)$ has positive feedback, leading to the desired contradiction. To begin, we make the following claim that we prove later.

Claim 4.26. *Suppose the assumptions above hold. There exist $(k - 1)$ -boxes $Q_{k-1}(i_1^*)$ and $Q_{k-1}(i_0^*)$ such that*

- (i) $Q_{k-1}(i_0^*) \in \mathcal{W}^{\text{inn}}(k, i^{\text{par}})$,
- (ii) $\tau_{k-1,i_0^*} \leq \tau_{k,i}$,
- (iii) $Q_{k-1}(i_1^*) \in \mathcal{W}^{\text{inn}}(k, i) \cap \mathcal{W}^{\text{inn}}(k, i^{\text{par}})$,

and there exists a path of $(k - 1)$ -boxes π_{k-1} from $Q_{k-1}(i_0^*)$ to $Q_{k-1}(i_1^*)$ such that

$$\pi_{k-1} \subset \mathcal{W}^{\text{inn}}(k, i^{\text{par}}) \quad \text{and} \quad |\pi_{k-1}| \leq \frac{3L_k}{1000L_{k-1}} + A\sigma. \quad (40)$$

Assuming Claim 4.26 holds and $Q_k(i^{\text{par}})$ has positive feedback, we deduce that all $(k - 1)$ -boxes on π_{k-1} have positive feedback as λ is such that (CAS_k) holds. Moreover, as λ is such that (PC_{k-1}^C) holds, then (Fast_{k-1}) holds. Thus

$$\tau_{k-1, i_1^*} < \tau_{k-1, i_0^*} + 2r_{k-1}L_{k-1}|\pi_{k-1}|.$$

By (40) we deduce

$$\begin{aligned} \tau_{k-1, i_1^*} &< \tau_{k-1, i_0^*} + 2r_{k-1}L_{k-1} \left(\frac{3L_k}{1000L_{k-1}} + A\sigma \right) \\ &\leq \tau_{k, i} + 2r_{k-1}L_{k-1} \left(\frac{3L_k}{1000L_{k-1}} + A\sigma \right). \end{aligned}$$

Recalling the definition of r_k in (24) where $a_1 = 1000A\sigma/3$, we have

$$\begin{aligned} \tau_{k-1, i_1^*} &< \tau_{k, i} + \frac{3}{500}r_{k-1}L_k \left(1 + \frac{a_1}{k^2L_{k-1}^{d-1}} \right) \\ &= \tau_{k, i} + \frac{3}{500}r_kL_k. \end{aligned} \quad (41)$$

As $Q_{k-1}(i_1^*) \in \mathcal{W}^{\text{inn}}(k, i)$, (41) contradicts (39). Hence it must be the case that $Q_k(i^{\text{par}})$ is either bad or has negative feedback. ■

Proof of Claim 4.26. Let $Q_{k-1}(i_{\text{ent}})$ be such that $e_{k, i} \in Q_{k-1}^{\text{core}}(i_{\text{ent}})$. If $Q_{k-1}(i_{\text{ent}}) \in \mathcal{W}^{\text{inn}}(k, i^{\text{par}})$ then the first two conditions in Claim 4.26 follow immediately by setting $i_0^* = i_{\text{ent}}$. The other possibility is where $Q_{k-1}(i_{\text{ent}}) \in \mathcal{A}$, where \mathcal{A} is an inner cluster in $Q_k(i^{\text{par}})$. In this case we let $Q_{k-1}(i_0^*)$ be the $(k - 1)$ -box in $\partial\mathcal{A}$ with the smallest entrance time and the first two conditions of Claim 4.26 follow. Note that $Q_{k-1}(i_{\text{ent}})$ cannot be contained in a boundary cluster of $Q_k(i^{\text{par}})$, as by definition of parent, we have $Q_{k-1}^{\text{core}}(i_{\text{ent}}) \subset Q_k^{\text{core}}(i^{\text{par}})$.

The final part of the claim follows by setting π_{k-1} to be the shortest path of $(k - 1)$ -boxes from $Q_{k-1}(i_0^*)$ to $\mathcal{W}^{\text{inn}}(k, i)$ that is completely contained in $\mathcal{W}^{\text{inn}}(k, i^{\text{par}})$. Since the distance between $\partial Q_k(i)$ and $\mathcal{W}^{\text{inn}}(k, i)$ is at most $\frac{3L_k}{1000}$, by further accounting for possible deviations around clusters of $Q_k(i)$ through their respective outer boundaries, we obtain the desired conditions. ■

Proof of Lemma 4.22 for (Fast_k) . Let $k \geq 2$ and $C > C_0$. Assume that λ is such that (PC_{k-1}^C) and (CAS_k) hold. Suppose $Q_k(i)$ is good and $Q_k(j)$ has positive feedback so that $\|i - j\|_\infty = 1$. As $Q_k(j)$ has positive feedback, there exists a $(k - 1)$ -box $Q_{k-1}(j_0) \in \mathcal{W}^{\text{inn}}(k, j)$ that has positive feedback such that

$$\tau_{k-1, j_0} < \tau_{k, j} + \frac{3}{500}r_kL_k.$$

Fix a $(k - 1)$ -box $Q_{k-1}(j_1) \in \mathcal{W}^{\text{inn}}(k, j) \cap \mathcal{W}(k, i)$ such that there is a path π_{k-1} from $Q_{k-1}(j_0)$ to $Q_{k-1}(j_1)$ with

$$\pi_{k-1} \subset \mathcal{W}^{\text{inn}}(k, j) \quad \text{and} \quad |\pi_{k-1}| \leq \left(1 - \frac{3}{1000}\right) \frac{L_k}{L_{k-1}} + A\sigma.$$

As λ is such that (PC_{k-1}^C) holds, by (Fast_{k-1}) we deduce that

$$\begin{aligned} \tau_{k,i} &\leq \tau_{k-1,j_1} < \tau_{k-1,j_0} + 2r_{k-1}L_{k-1} \left(\left(1 - \frac{3}{1000}\right) \frac{L_k}{L_{k-1}} + A\sigma \right) \\ &= \tau_{k-1,j_0} + 2r_{k-1}L_k \left(1 - \frac{3}{1000} + \frac{A\sigma}{k^2L_{k-1}^{d-1}} \right) \\ &< \tau_{k,j} + 2r_{k-1}L_k \left(1 - \frac{3}{1000} + \frac{A\sigma}{k^2L_{k-1}^{d-1}} \right) + \frac{3}{500}r_kL_k. \end{aligned}$$

By taking the final term above inside the bracket, we deduce that

$$\begin{aligned} \tau_{k,i} &< \tau_{k,j} + 2r_{k-1}L_k \left(1 + \frac{3}{1000} \left(\frac{r_k}{r_{k-1}} - 1 \right) + \frac{A\sigma}{k^2L_{k-1}^{d-1}} \right), \\ &= \tau_{k,j} + 2r_{k-1}L_k \left(1 + \frac{2A\sigma}{k^2L_{k-1}^{d-1}} \right) \\ &< \tau_{k,j} + 2r_kL_k \end{aligned}$$

where in the first two lines we recall (24) and $a_1 = 1000A\sigma/3$. From the above inequality, we deduce (Fast_k) holds. ■

Proof of Lemma 4.22 for (Del_k^C) . Let $k \geq 2$, $C > C_0$ and assume that λ is such that (PC_{k-1}^C) and (CAS_k) hold. Suppose that $Q_k(i)$ and $Q_k(i^{\text{par}})$ both have negative feedback.

Claim 4.27. *Suppose the assumptions above hold. There exists a $(k - 1)$ -box*

$$Q_{k-1}(\hat{i}_0) \subset Q_k(i^{\text{par}})$$

with negative feedback such that

- (1) $\text{dist}(Q_{k-1}(\hat{i}_0), \partial Q_k(i^{\text{par}})) \geq \frac{1}{3}(L_k - a_6L_{k-1})$,
- (2) $\tau_{k-1,\hat{i}_0} < \tau_{k,i} + a_7r_{k-1}L_{k-1}$,

where $a_6, a_7 > 0$ are constants that depend only on A and d .

Let $Q_{k-1}(\hat{i}_0)$ be as given in the claim above. As λ is such that (PC_{k-1}^C) holds, by (Del_{k-1}^C) , Lemma 4.24 and Claim 4.27, we deduce that

$$\begin{aligned} \tau_{k-1,\hat{i}_0} &> \tau_{k,i^{\text{par}}} + C\omega_{k-1}L_{k-1} \left(\frac{3}{L_{k-1}} \text{dist}(Q_{k-1}(\hat{i}_0), \partial Q_k(i^{\text{par}})) - a_4 \right), \\ &\geq \tau_{k,i^{\text{par}}} + C\omega_{k-1}L_k \left(1 - \frac{a_4 + a_6}{k^2L_{k-1}^{d-1}} \right). \end{aligned} \tag{42}$$

Moreover, by Claim 4.27 and (42),

$$\begin{aligned} \tau_{k,i} &> \tau_{k-1,\hat{i}_0} - a_7 r_{k-1} L_{k-1}, \\ &> \tau_{k,i^{\text{par}}} + C\omega_{k-1} L_k \left(1 - \frac{a_4 + a_6 + a_7 r_{k-1}}{k^2 L_{k-1}^{d-1}}\right). \end{aligned} \tag{43}$$

If we set

$$a_2 = a_4 + a_6 + 2a_7 r_1,$$

then $a_2 > a_4 + a_6 + a_7 r_{k-1}$ for all $k \geq 2$ by Remark 4.16 for a sufficiently large L_1 , and only depends on A, d and p . In particular, by (43) we have

$$\tau_{k,i} > \tau_{k,i^{\text{par}}} + C\omega_{k-1} \left(1 - \frac{a_2}{k^2 L_{k-1}^{d-1}}\right),$$

establishing the result. ■

Proof of Claim 4.27. Let $Q_{k-1}(i_{\text{ent}})$ be a $(k - 1)$ -box such that $e_{k,i} \in Q_{k-1}^{\text{core}}(i_{\text{ent}})$. If $Q_{k-1}(i_{\text{ent}}) \in \mathcal{W}(k, i^{\text{par}})$ has negative feedback, then set $\hat{i}_0 = i_{\text{ent}}$. Another simple case is when $Q_{k-1}(i_{\text{ent}})$ is contained in some poorly confined inner cluster of $Q_k(i^{\text{par}})$ which must have a negative feedback $(k - 1)$ -box in its outer boundary; the one with the smallest entrance time can be chosen to be $Q_{k-1}(\hat{i}_0)$ as λ is such that (Conf_k) holds. Note that $Q_{k-1}(i_{\text{ent}})$ cannot be contained in a boundary cluster of $Q_k(i^{\text{par}})$, as by definition of parent we have $Q_{k-1}^{\text{core}}(i_{\text{ent}}) \subset Q_k^{\text{core}}(i^{\text{par}})$.

The remaining two cases to consider are when $Q_{k-1}(i_{\text{ent}}) \in \mathcal{W}(k, i^{\text{par}})$ has positive feedback, and when $Q_{k-1}(i_{\text{ent}})$ is contained in a successfully confined inner cluster of $Q_k(i^{\text{par}})$. The idea of the proof for these cases is to find a $(k - 1)$ -box $Q_{k-1}(i_0) \in \mathcal{W}(k, i^{\text{par}})$ with positive feedback near $Q_{k-1}(i_{\text{ent}})$ whose entrance time is not much larger than $\tau_{k-1,i_{\text{ent}}}$. From $Q_{k-1}(i_0)$ we fix a path of $(k - 1)$ -boxes π_{k-1} to $\mathcal{W}^{\text{inn}}(k, i)$, so that

$$\pi_{k-1} \subset \mathcal{W}^{\text{core}}(k, i^{\text{par}}) \quad \text{and} \quad |\pi_{k-1}| \leq \frac{3L_k}{1000L_{k-1}} + A\sigma, \tag{44}$$

where $\mathcal{W}^{\text{core}}(k, i^{\text{par}})$ is the set of wonderful $(k - 1)$ -boxes in $Q_k(i^{\text{par}})$ that are contained either in $Q_k^{\text{core}}(i^{\text{par}})$ or in the outer boundary of an inner cluster that intersects $Q_k^{\text{core}}(i^{\text{par}})$. As λ is such that (Fast_{k-1}) holds, if all $(k - 1)$ -boxes on π_{k-1} have positive feedback then $Q_k(i)$ must have positive feedback, which is a contradiction. Hence there exists some $(k - 1)$ -box on π_{k-1} with negative feedback and we prove that this box is necessarily close to $Q_{k-1}(i_{\text{ent}})$, completing the proof.

Now we implement the idea above rigorously. If $Q_{k-1}(i_{\text{ent}}) \in \mathcal{W}(k, i^{\text{par}})$ has positive feedback then we simply set $i_0 = i_{\text{ent}}$. Otherwise, if $Q_{k-1}(i_{\text{ent}})$ is contained in a successfully confined inner cluster of $Q_k(i^{\text{par}})$, \mathcal{A} say, then we may assume that all $(k - 1)$ -boxes within $\partial\mathcal{A}$ have positive feedback as otherwise we may set \hat{i}_0 to correspond to one of these boxes with negative feedback. Note that \mathcal{A} intersects $Q_k^{\text{core}}(i^{\text{par}})$. Hence we may choose $Q_{k-1}(i_0)$ to be the $(k - 1)$ -box in $\partial\mathcal{A}$ with earliest entrance time. Through this construction, we have

$$\tau_{k-1,i_0} \leq \tau_{k,i}. \tag{45}$$

Fix a geodesic path of $(k - 1)$ -boxes from $Q_{k-1}(i_0)$ to $\mathcal{W}_k^{\text{inn}}(i) \cap Q_k^{\text{core}}(i^{\text{par}})$, that is, the shortest path of $(k - 1)$ -boxes from $Q_{k-1}(i_0)$ to some $(k - 1)$ -box contained in $\mathcal{W}_k^{\text{inn}}(i) \cap Q_k^{\text{core}}(i^{\text{par}})$. Let π_{k-1} be the augmentation of this path that deviates through the outer boundary of clusters in $Q_k(i^{\text{par}})$ it intersects, so that π_{k-1} ends at some $(k - 1)$ -box $Q_{k-1}(j) \in \mathcal{W}_k^{\text{inn}}(k, i) \cap Q_k^{\text{core}}(i^{\text{par}})$ and π_{k-1} satisfies (44). This construction is possible as $Q_{k-1}(i_0) \in \partial\mathcal{A} \subset \mathcal{W}^{\text{core}}(k, i^{\text{par}})$ and we may always choose the deviations so that such a $Q_{k-1}(j)$ exists.

If $\pi_{k-1} \subset \mathcal{W}(k, i)$, then define $\pi_{k-1}^{\text{end}} = \pi_{k-1}$ and $\pi_{k-1}^{\text{start}} = \emptyset$. Otherwise, let π_{k-1}^{end} be the subpath of π_{k-1} given by starting from $Q_{k-1}(j)$ and backwards traversing π_{k-1} until there is a $(k - 1)$ -box not contained in $Q_k^{\text{core}}(i^{\text{par}})$, and terminating at the $(k - 1)$ -box immediately before this. Hence, if $Q_{k-1} \in \pi_{k-1}^{\text{end}}$, then $Q_{k-1} \subset Q_k^{\text{core}}(i^{\text{par}})$ and $Q_{k-1} \in \mathcal{W}(k, i)$. Let $\pi_{k-1}^{\text{start}} = \pi_{k-1} \setminus \pi_{k-1}^{\text{end}}$ and note that the number of $(k - 1)$ -boxes in π_{k-1}^{start} is bounded above by a constant that depends only on A and d . This is because in the construction above, π_{k-1} will only need to be outside $Q_k^{\text{core}}(i^{\text{par}})$ to deviate around inner clusters in $Q_k(i^{\text{par}})$.

As $Q_k(i)$ has negative feedback, there must exist some $Q_{k-1}(j_0) \in \pi_{k-1}$ that has negative feedback and without losing generality we may assume that it is the $(k - 1)$ -box on π_{k-1} with negative feedback that is first discovered by traversing π_{k-1} from $Q_{k-1}(i_0)$. Indeed, if such a $Q_{k-1}(j_0)$ did not exist, then (Fast $_{k-1}$) and (44) would imply that $Q_k(i)$ has positive feedback. Let $\pi_{k-1}^{j_0}$ be the subpath of π_{k-1} from $Q_{k-1}(i_0)$ to $Q_{k-1}(j_0)$.

If $Q_{k-1}(j_0) \in \pi_{k-1}^{\text{start}}$, then $\pi_{k-1}^{j_0} \subset \pi_{k-1}^{\text{start}}$. In this case, there exist constants $a_8, a_9 > 0$ that depend only on A and d such that

$$\frac{3}{L_{k-1}} \text{dist}(Q_{k-1}(j_0), \partial Q_k(i)) \leq a_8 \quad \text{and} \quad |\pi_{k-1}^{j_0}| \leq a_9. \tag{46}$$

If $Q_{k-1}(j_0) \in \pi_{k-1}^{\text{end}}$, then $Q_{k-1}(j_0) \subset Q_k^{\text{core}}(i^{\text{par}}) \subset Q_k(i)$. As $C > C_0$ and λ is such that (PC $_{k-1}^C$) holds, by Lemma 4.24, then

$$\tau_{k-1, j_0} > \tau_{k, i} + C\omega_{k-1}L_{k-1} \left(\frac{3}{L_{k-1}} \text{dist}(Q_{k-1}(j_0), \partial Q_k(i)) - a_4 \right). \tag{47}$$

In other words, if $Q_{k-1}(j_0)$ is contained in $Q_k(i)$ but is not close to the boundary of $Q_k(i)$, we can use Lemma 4.24 to deduce a non-trivial lower bound on the difference between τ_{k-1, j_0} and $\tau_{k, i}$. In this case,

$$|\pi_{k-1}^{j_0}| \leq \frac{3}{L_{k-1}} \text{dist}(Q_{k-1}(j_0), \partial Q_k(i)) + A\sigma. \tag{48}$$

By construction, every $(k - 1)$ -box on $\pi_{k-1}^{j_0}$ has positive feedback except for $Q_{k-1}(j_0)$. Thus, as λ is such that (Fast $_{k-1}$) holds, by (45) we deduce that

$$\begin{aligned} \tau_{k-1, j_0} &< \tau_{k-1, i_0} + 2r_{k-1}L_{k-1}|\pi_{k-1}^{j_0}| \\ &\leq \tau_{k, i} + 2r_{k-1}L_{k-1} \left(\frac{3}{L_{k-1}} \text{dist}(Q_{k-1}(j_0), \partial Q_k(i)) + A\sigma \right). \end{aligned} \tag{49}$$

If $\frac{3}{L_{k-1}} \text{dist}(Q_{k-1}(j_0), \partial Q_k(i)) > a_4$ and $C > 0$ is large enough so that

$$C\omega_{k-1} \left(\frac{3}{L_{k-1}} \text{dist}(Q_{k-1}(j_0), \partial Q_k(i)) - a_4 \right) > 2r_{k-1} \left(\frac{3}{L_{k-1}} \text{dist}(Q_{k-1}(j_0), \partial Q_k(i)) + A\sigma \right), \quad (50)$$

then (47) contradicts (49). Rearranging the above inequality, we deduce that (50) holds if

$$C > \frac{2\sigma r_{k-1}}{\omega_{k-1}} a_3,$$

where $a_3 > 1$ is a constant that only depends on A and d which is exactly the choice of a_3 in the definition of C_0 in (26). Moreover, if $C > C_0$ and L_1 is large enough so that $r < 2r_1$ and $\omega > 1/2$, then the above inequality implies that (50) follows for all $k \geq 2$. Consequently,

$$\frac{3}{L_{k-1}} \text{dist}(Q_{k-1}(j_0), \partial Q_k(i)) \leq a_4. \quad (51)$$

The claim follows from (46), (48), (51) and (Fast $_k$) by setting $\hat{i}_0 = j_0$, $a_6 = \max\{a_4, a_8\}$ and $a_7 = \max\{a_4 + A\sigma, a_9\}$. ■

Proof of Lemma 4.22 for (Conf $_k$). This is an immediate consequence of the proofs for the three fundamental properties above and Lemma 4.23. ■

5. Proofs of Theorems 1.1 and 1.3

In this section we prove Theorems 1.1 and 1.3. First we state and prove the following lemma.

Lemma 5.1. *If L_1 is large enough and λ small enough, then with positive probability, $Q_k(o)$ has positive feedback for all $k \geq 1$.*

Proof. By Lemma 4.11 it suffices to prove that if the origin is flawless, then $Q_k(o)$ has positive feedback for all $k \geq 1$. We proceed by induction and note that the case $k = 1$ is immediate from the definition of positive feedback at scale 1. Now suppose $k > 1$ and that the result holds up to scale $k - 1$. As a consequence of the origin being flawless, we have

$$Q_{k-1}(o) \in \mathcal{W}^{\text{inn}}(k, o),$$

and by the inductive hypothesis, $Q_{k-1}(o)$ has positive feedback. Moreover, as $\tau_{k-1,o} = \tau_{k,o} = 0$ and $\eta_0(o) = 1$, we see that $Q_k(o)$ has positive feedback at scale k . ■

Proof of Theorem 1.1. Fix L_1 large enough and then λ small enough so that Lemmas 4.17 and 5.1 hold. Assume that $Q_\kappa(o)$ has positive feedback for all $\kappa \geq 1$. Fix a large scale k and consider $Q_k(o)$. By Lemma 4.17, every $(k - 1)$ -box contained in $\mathcal{W}^{\text{inn}}(k, o)$ has

positive feedback. For ease of argument, let us focus on $(k - 1)$ -cores in the intersection $Q_k^{\text{core}}(o) \cap \mathcal{W}^{\text{inn}}(k, o)$. The number of such $(k - 1)$ -cores is at least

$$\left(\frac{L_k}{L_{k-1}}\right)^d - A\sigma = \left(\frac{L_k}{L_{k-1}}\right)^d \left(1 - \frac{A\sigma}{k^{2d} L_{k-1}^{d(d-1)}}\right),$$

where the $A\sigma$ term counts all $(k - 1)$ -boxes contained in clusters in $Q_k(o)$. Through analogous reasoning, in each of these $(k - 1)$ -cores in $Q_k^{\text{core}}(o) \cap \mathcal{W}^{\text{inn}}(k, o)$, the number of $(k - 2)$ -cores that are not contained in a cluster of bad $(k - 2)$ -boxes in a good $(k - 1)$ -box is bounded below by

$$\left(\frac{L_{k-1}}{L_{k-2}}\right)^d - A\sigma = \left(\frac{L_{k-1}}{L_{k-2}}\right)^d \left(1 - \frac{A\sigma}{(k - 1)^{2d} L_{k-2}^{d(d-1)}}\right).$$

Iterating this argument all the way to $k = 1$ allows us to deduce that the number of 1-cores contained in $Q_k^{\text{core}}(o) \cap \mathcal{W}^{\text{inn}}(k, o)$ that are not contained in a cluster of any good box up to scale k is bounded below by

$$\left(\frac{L_k}{L_1}\right)^d \left(\prod_{j=1}^{k-1} \zeta_j\right) \quad \text{where} \quad \zeta_j = 1 - \frac{A\sigma}{(j + 1)^{2d} L_j^{d(d-1)}}.$$

From ζ_j given above, we deduce that $\{\prod_{j=1}^{k-1} \zeta_j\}_{k \geq 1}$ is a decreasing sequence that is uniformly bounded away from 0, so that

$$\zeta = \lim_{k \rightarrow \infty} \prod_{j=1}^{k-1} \zeta_j > 0, \tag{52}$$

where ζ depends only on A and d .

If $Q_1(g) \subset Q_k^{\text{core}}(o) \cap \mathcal{W}^{\text{inn}}(k, o)$ and is not contained in a cluster of any good box up to scale k , then $Q_1(g)$ has positive feedback as $Q_k(o)$ has positive feedback. Recall $\mathcal{C}_1^-(g)$ is the largest component of non-seeds in $Q_1(g)$ with sites in the boundary $\partial Q_1(g)$ removed. By Lemma 4.3, all sites in $\mathcal{C}_1^-(g)$ are occupied by FPP_1 , and in particular, all sites in $\mathcal{C}_1(g) \cap Q_1^{\text{core}}(g)$. As $Q_1(g)$ is good, for ε we fix in (3) we have

$$|\mathcal{C}_1(g) \cap Q_1^{\text{core}}(g)| \geq (1 - \varepsilon)\theta(1 - p)(L_1/3)^d, \tag{53}$$

where we recall that $\theta(1 - p) > 0$ as $p < 1 - p_c^{\text{site}}$. As 1-cores partition \mathbb{Z}^d , we deduce from (52) and (53) that the number of sites that FPP_1 occupies in $Q_k(o)$ is bounded below by

$$3^{-d} \zeta (1 - \varepsilon)\theta(1 - p)L_k^d. \tag{54}$$

By dividing (54) by L_k^d , we see that a lower bound of the density of sites occupied by FPP_1 in $Q_k(o)$ is a positive constant that does not depend on k . The result follows by letting $k \rightarrow \infty$. ■

Recall the critical probability p_c'' defined in (1). To prove Theorem 1.3, we augment the definition of a good 1-box as we will need to exploit the fact that $p < p_c''$. Consider the 1-box $Q_1(i)$. Enumerate the connected components of filled seeds in $Q_1(i)$ as

$$\mathcal{S}_1(i), \dots, \mathcal{S}_{m_i}(i),$$

so that $|\mathcal{S}_1(i)| \geq \dots \geq |\mathcal{S}_{m_i}(i)|$, where m_i is the number of connected components of filled seeds in $Q_1(i)$. Define the event

$$E_6(i) = \{|\mathcal{S}_1(i)| \leq L_1/100\}.$$

By setting $M = L_1/100$ in (1), we observe that we can include $E_6(i)$ in the definition of $Q_1(i)$ being good and all the previous results in our multi-scale analysis follow in the same manner for $p < p_c''$.

The proof of Theorem 1.3 also relies on extending the notion of parent and progenitor to sites, which one can view as *scale 0* boxes. We define a site to be good if it does not host a seed and bad otherwise. A good site has positive feedback if it is occupied by FPP_1 and has negative feedback if it is occupied by FPP_λ . The parent of a site v is the neighbouring site from which FPP_1 or FPP_λ propagated to that site at time $\tau(v)$. Any bad or negative feedback site v (i.e. a site occupied by FPP_λ) can be traced through the parent structure to a unique seed that was activated by FPP_1 , which we call the progenitor in an analogous manner to scales $k \geq 1$. Hence both v^{prog} and $\mathcal{P}_0^{v \leftarrow v^{\text{prog}}}$ are well-defined. With these considerations, we are now in a position to prove Theorem 1.3.

Proof of Theorem 1.3. Suppose $d \geq 2$, $p < \min\{p_c'', 1 - p_c^{\text{site}}\}$ and L_1 is large enough and λ is small enough so that Lemmas 4.17 and 5.1 and Theorem 1.1 hold. Assume that $Q_\kappa(o)$ has positive feedback for all $\kappa \geq 1$. In this scenario, FPP_1 survives with positive probability, and so strong survival will be established if we prove all connected components of FPP_λ are almost surely finite.

Let S be an arbitrary component of seeds and S' be the set of sites whose progenitor is contained in S . Fix a scale k large enough so that $S \subset Q_k^{\text{core}}(o)$. We note that such a k exists as $p < p_c'' \leq p_c^{\text{site}}$, and so all components of seeds are almost surely finite.

To set up the proof, we introduce the following definition. For $\ell \geq 1$ and $B \subset \mathbb{Z}^d$, we call a finite set \mathcal{O}_ℓ of ℓ -boxes an *annulus of ℓ -boxes separating B from infinity* if any infinite self-avoiding path from B intersects the core of an ℓ -box from \mathcal{O}_ℓ , and moreover we can find two disjoint connected sets Z_1 and Z_2 such that

$$\mathbb{Z}^d \setminus \bigcup_{Q_\ell \in \mathcal{O}_\ell} Q_\ell^{\text{core}} = Z_1 \cup Z_2, \tag{55}$$

where Z_1 is finite and contains B while Z_2 is unbounded.

Let $\mathcal{O}_{k-1} \subset \mathcal{W}^{\text{inn}}(k, o)$ be an annulus of $(k - 1)$ -boxes separating $Q_k^{\text{core}}(o)$ from infinity. As λ is such that (CAS $_k$) holds, all $(k - 1)$ -boxes in \mathcal{O}_{k-1} have positive feedback. Let \mathcal{O}_{k-2} be an annulus of $(k - 2)$ -boxes separating $Q_k^{\text{core}}(o)$ from infinity such that

$$\mathcal{O}_{k-2} \subset \bigcup_{Q_{k-1}(j) \in \mathcal{O}_{k-1}} \mathcal{W}^{\text{inn}}(k - 1, j).$$

As all $(k - 1)$ -boxes in \mathcal{O}_{k-1} have positive feedback and λ is such that (CAS_{k-1}) holds, all $(k - 2)$ -boxes in \mathcal{O}_{k-2} also have positive feedback. We continue this procedure until we construct an annulus of 1-boxes that separates $Q_k^{\text{core}}(o)$ from infinity, \mathcal{O}_1 say, such that all 1-boxes in \mathcal{O}_1 have positive feedback.

Let Z_2 be the unbounded connected set induced by \mathcal{O}_1 as given in (55), noting that $\mathbb{Z}^d \setminus Z_2$ must be a finite set that contains S . If S' is infinite, then by the definition of annulus there must exist a box $Q_1(j) \in \mathcal{O}_1$ whose core contains a site from S' . Hence, $\mathcal{C}_1^-(j)$ must intersect S' , which contradicts the fact that $Q_1(j)$ is of positive feedback. ■

6. Concluding remarks and open questions

To establish the coexistence phase in Corollary 1.2, we require that the FPP_λ survives almost surely by letting $p \in (p'_c, 1 - p_c^{\text{site}})$, which is only possible if $d \geq 3$. This simplifies our analysis as we know that FPP_λ survives almost surely in such situations, so by setting λ small enough we can prove FPP_1 survives with positive probability as outlined in Theorem 1.1. The most natural question is whether or not there is a coexistence phase without the almost sure survival of FPP_λ .

Question 6.1. For $d \geq 3$, does there exist a coexistence phase for all $p \in (0, 1 - p_c^{\text{site}})$?

The coexistence phase established in Corollary 1.2 leads one to wonder if there could be coexistence when $p < p'_c$, but the case $d = 2$ is wide open in the sense that it is not known if there is coexistence for any choice of parameters.

Question 6.2. Does there exist a coexistence phase for FPPHE when $d = 2$?

The main difficulty in approaching these two questions is controlling for the survival of both processes simultaneously. By Theorem 1.1, we see that for all $p < 1 - p_c^{\text{site}}$, if λ is small enough then FPP_1 survives with positive probability but there is no information about the survival of FPP_λ unless we assume that the seeds are supercritical or use an enhancement argument with filled seeds. For a fixed p it could be the case that for some values of λ there is coexistence, but once λ is sufficiently small, there is strong survival. This issue is compounded as the strong survival and coexistence may not be disjoint phases, in that there could be a choice of λ and p for which both occur with positive probability. This in fact happens when the graph G is composed of two copies of \mathbb{Z}^2 that are connected by a single edge between their respective origins. By Sidoravicius and Stauffer [27, Theorem 1.3], there is a phase of strong survival on \mathbb{Z}^2 . If λ and p are such that there is strong survival on \mathbb{Z}^2 , then strong survival and coexistence both have a positive probability of occurring on G . Nonetheless, we expect that, on \mathbb{Z}^d , strong survival and coexistence are indeed disjoint phases.

Question 6.3. For what graphs are the strong survival and coexistence phases distinct?

Even if we could assume that the strong survival and coexistence phases were distinct, distinguishing between them is still challenging due to the lack of monotonicity in

FPPHE. As discussed in Section 1.5, one can help the spread of FPP_1 by strategically planting additional seeds or increasing passage times and one can even construct graphs where increasing p can increase the probability that FPP_1 survives. However, one would expect that if the underlying graph was transitive, then the probability of FPP_1 surviving would be a non-increasing function of λ and p . Establishing for what graphs this property holds is an interesting question.

Question 6.4. For what graphs is the probability of FPP_1 surviving a non-increasing function of λ and p ?

In Corollary 1.2, a coexistence phase is established in which both FPP_1 and FPP_λ occupy a positive density of sites. It is possible that coexistence occurs with FPP_1 surviving while only occupying a set of zero density, a regime we refer to as *weak coexistence*. It is not difficult to prove there is a phase of weak coexistence on (non-amenable) trees, but establishing a phase of weak coexistence on \mathbb{Z}^d remains an open problem.

Question 6.5. For $d \geq 2$, does there exist a regime of weak coexistence on \mathbb{Z}^d ?

One may consider an alternative definition of survival of FPP_λ where the spread of FPP_λ from a single seed gives rise to an infinite connected component of FPP_λ . More precisely, we construct an infinite directed forest on \mathbb{Z}^d encoding the spread of FPP_1 and FPP_λ as follows. If FPP_1 (resp. FPP_λ) spreads from x to successfully occupy y , we attach a directed edge from x to y . When a seed is activated by the attempted occupation of FPP_1 , we attach no directed edge with endpoint at that seed. We say there is *infinite geodesic survival* if there exists at least one infinite connected component of FPP_λ in the directed forest above. Clearly infinite geodesic survival implies survival considered in this paper. A natural question is whether there can be coexistence with this definition.

Question 6.6. For $d \geq 2$, can FPP_1 and FPP_λ observe infinite geodesic survival simultaneously with positive probability?

We can use our multi-scale analysis with non-equilibrium feedback to recover the strong survival phase established in [27], in which for $\lambda \in (0, 1)$, there is strong survival for sufficiently small p . We briefly indicate how this can be done now.

Fix $\lambda \in (0, 1)$. The aim is to prove that FPP_1 survives while all connected components of FPP_λ are finite for a small enough choice of p . The standard multi-scale construction proceeds in an almost identical manner except we alter the definition of good 1-box. We define a 1-box to be good if FPP_1 can readily spread in the box and there are no seeds. By setting L_1 large enough and p small enough, the standard multi-scale construction then follows readily, so that good boxes at all scales occur with sufficiently high probability.

The idea of the proof is that in a good k -box, all bad boxes at lower scales (away from the boundary) are well-separated and one can construct an encapsulation argument to control the spread of FPP_λ from these bad boxes. Indeed, as FPP_λ must spread from bad boxes, it is reasonable to construct a parent and progenitor structure analogous to the one in Section 4.4 to identify the good boxes whose sites FPP_λ manages to occupy. For this, we tune the definition of positive and negative feedback so that FPP_1 occupies all sites in

a good box that are sufficiently far from bad boxes at lower scales. For example, a 1-box has positive feedback if FPP_1 occupies a site away from the boundary at a relatively fast time after the box was first entered. Through a cascading argument, FPP_λ is confined to a collection of regions in this k -box that can no longer spread while FPP_1 occupies every other site. By taking the limit $k \rightarrow \infty$, FPP_1 survives while FPP_λ fails to observe an infinite connected component. The spirit of this proof is close to the one given in [27], but our multi-scale analysis with non-equilibrium feedback offers a more systematic approach through the introduction of positive and negative feedback, leading to a clearer and more streamlined proof.

Acknowledgements. A. Stauffer is grateful to Ioan Manolescu for useful discussions. Part of this work was completed during a visit by T. Finn to the University of Roma Tre, for which he is grateful for their generous hospitality.

Funding. T. Finn was supported by a scholarship from the EPSRC Centre for Doctoral Training in Statistical Applied Mathematics at Bath (SAMBa), under the project EP/L015684/1 and EB/MA1206A.

A. Stauffer was supported by EPSRC Fellowship EP/N004566/1.

References

- [1] Ahlberg, D., Deijfen, M., Hoffman, C.: The two-type Richardson model in the half-plane. *Ann. Appl. Probab.* **30**, 2261–2273 (2020) Zbl [1457.60140](#) MR [4149528](#)
- [2] Aizenman, M., Grimmett, G.: Strict monotonicity for critical points in percolation and ferromagnetic models. *J. Statist. Phys.* **63**, 817–835 (1991) MR [1116036](#)
- [3] Antal, P., Pisztor, A.: On the chemical distance for supercritical Bernoulli percolation. *Ann. Probab.* **24**, 1036–1048 (1996) Zbl [0871.60089](#) MR [1404543](#)
- [4] Auffinger, A., Damron, M., Hanson, J.: 50 Years of First-Passage Percolation. University Lecture Ser. 68, Amer. Math. Soc., Providence, RI (2017) Zbl [1452.60002](#) MR [3729447](#)
- [5] Balister, P., Bollobás, B., Riordan, O.: Essential enhancements revisited. arXiv:[1402.0834](#) (2014)
- [6] Campanino, M., Russo, L.: An upper bound on the critical percolation probability for the three-dimensional cubic lattice. *Ann. Probab.* **13**, 478–491 (1985) Zbl [0567.60096](#) MR [781418](#)
- [7] Candellero, E., Stauffer, A.: Coexistence of competing first passage percolation on hyperbolic graphs. *Ann. Inst. Henri Poincaré Probab. Statist.* **57**, 2128–2164 (2021) Zbl [1492.60268](#) MR [4330845](#)
- [8] Candellero, E., Stauffer, A.: First passage percolation in a hostile environment is not monotone. arXiv:[2110.05821](#) (2021)
- [9] Candellero, E., Teixeira, A.: Percolation and isoperimetry on roughly transitive graphs. *Ann. Inst. Henri Poincaré Probab. Statist.* **54**, 1819–1847 (2018) Zbl [1417.60078](#) MR [3865659](#)
- [10] Deijfen, M., Hirscher, T., Lopes, F.: Competing frogs on \mathbb{Z}^d . *Electron. J. Probab.* **24**, art. 146, 17 pp. (2019) Zbl [1428.60137](#) MR [4049082](#)
- [11] Deuschel, J.-D., Pisztor, A.: Surface order large deviations for high-density percolation. *Probab. Theory Related Fields* **104**, 467–482 (1996) Zbl [0842.60023](#) MR [1384041](#)
- [12] Finn, T.: Topics in random growth models. PhD thesis, Univ. of Bath (2021)
- [13] Fontes, L., Newman, C. M.: First passage percolation for random colorings of \mathbb{Z}^d . *Ann. Appl. Probab.* **3**, 746–762 (1993) Zbl [0780.60101](#) MR [1233623](#)

- [14] Garet, O., Marchand, R.: Coexistence in two-type first-passage percolation models. *Ann. Appl. Probab.* **15**, 298–330 (2005) Zbl [1080.60092](#) MR [2115045](#)
- [15] Garet, O., Marchand, R.: First-passage competition with different speeds: positive density for both species is impossible. *Electron. J. Probab.* **13**, no. 70, 2118–2159 (2008) Zbl [1191.60111](#) MR [2461538](#)
- [16] Gracar, P., Stauffer, A.: Multi-scale Lipschitz percolation of increasing events for Poisson random walks. *Ann. Appl. Probab.* **29**, 376–433 (2019) Zbl [1408.60038](#) MR [3910007](#)
- [17] Grimmett, G.: *Percolation*. 2nd ed., Grundlehren Math. Wiss. 321, Springer, Berlin (1999) Zbl [926.60004](#) MR [1707339](#)
- [18] Grimmett, G. R., Li, Z.: Brownian snails with removal: epidemics in diffusing populations. *Electron. J. Probab.* **27**, art. 78, 31 pp. (2022) Zbl [1495.60091](#) MR [4441145](#)
- [19] Häggström, O., Pemantle, R.: First passage percolation and a model for competing spatial growth. *J. Appl. Probab.* **35**, 683–692 (1998) Zbl [0920.60085](#) MR [1659548](#)
- [20] Häggström, O., Pemantle, R.: Absence of mutual unbounded growth for almost all parameter values in the two-type Richardson model. *Stochastic Process. Appl.* **90**, 207–222 (2000) Zbl [1047.60099](#) MR [1794536](#)
- [21] Hoffman, C.: Coexistence for Richardson type competing spatial growth models. *Ann. Appl. Probab.* **15**, 739–747 (2005) Zbl [1067.60098](#) MR [2114988](#)
- [22] Kesten, H., Sidoravicius, V.: Branching random walk with catalysts. *Electron. J. Probab.* **8**, art. 5, 51 pp. (2003) Zbl [1064.60196](#) MR [1961167](#)
- [23] Kesten, H., Sidoravicius, V.: The spread of a rumor or infection in a moving population. *Ann. Probab.* **33**, 2402–2462 (2005) Zbl [1111.60074](#) MR [2184100](#)
- [24] Kesten, H., Sidoravicius, V.: A phase transition in a model for the spread of an infection. *Illinois J. Math.* **50**, 547–634 (2006) Zbl [1101.92040](#) MR [2247840](#)
- [25] Penrose, M. D., Pisztor, A.: Large deviations for discrete and continuous percolation. *Adv. Appl. Probab.* **28**, 29–52 (1996) Zbl [0853.60085](#) MR [1372330](#)
- [26] Peres, Y., Sinclair, A., Sousi, P., Stauffer, A.: Mobile geometric graphs: detection, coverage and percolation. *Probab. Theory Related Fields* **156**, 273–305 (2013) Zbl [1273.82060](#) MR [3055260](#)
- [27] Sidoravicius, V., Stauffer, A.: Multi-particle diffusion limited aggregation. *Invent. Math.* **218**, 491–571 (2019) Zbl [1491.60183](#) MR [4011705](#)
- [28] Sznitman, A.-S.: Vacant set of random interlacements and percolation. *Ann. of Math. (2)* **171**, 2039–2087 (2010) Zbl [1202.60160](#) MR [2680403](#)
- [29] van den Berg, J., Kesten, H.: Inequalities for the time constant in first-passage percolation. *Ann. Appl. Probab.* **3**, 56–80 (1993) Zbl [0771.60092](#) MR [1202515](#)