

Geometric pushforward in Hodge filtered complex cobordism and secondary invariants

Knut Bjarte Haus and Gereon Quick

Abstract. We construct a functorial pushforward homomorphism in geometric Hodge filtered complex cobordism along proper holomorphic maps between arbitrary complex manifolds. This significantly improves previous results on such transfer maps and is a much stronger result than the ones known for differential cobordism of smooth manifolds. This enables us to define and provide a concrete geometric description of Hodge filtered fundamental classes for all proper holomorphic maps. Moreover, we give a geometric description of a cobordism analog of the Abel–Jacobi invariant for nullbordant maps which is mapped to the classical invariant under the Hodge filtered Thom morphism. For the latter we provide a new construction in terms of geometric cycles.

Contents

1. Introduction	457
2. Currential geometric Hodge filtered cobordism	461
3. Hodge filtered MU -orientations	471
4. Pushforward along proper Hodge filtered MU -oriented maps	475
5. A canonical Hodge filtered MU -orientation for holomorphic maps	488
6. Fundamental classes and secondary cobordism invariants	494
7. Hodge filtered Thom morphism	499
8. Image and kernel for compact Kähler manifolds	506
References	508

1. Introduction

The study of the analytic submanifolds of a given compact Kähler manifold is a central theme in complex geometry. Fundamental classes provide important invariants for this study. For a classical example, let X be a compact Kähler manifold and $Z \subset X$ a submanifold of codimension p . The Poincaré dual of the pushforward of the fundamental class of Z along the inclusion defines a cohomology class $[Z]$ in $H^{2p}(X; \mathbb{Z})$. In fact, $[Z]$ lies in the subgroup $\text{Hdg}^{2p}(X) = H^{2p}(X; \mathbb{Z}) \cap H^{p,p}(X; \mathbb{C})$ of integral classes of Hodge type

Mathematics Subject Classification 2020: 55N22 (primary); 14F43, 58J28, 19E15, 32C35 (secondary).
Keywords: cobordism, differential cohomology theories, Hodge filtered orientation, Thom morphism, Abel–Jacobi map.

(p, p) . This induces a homomorphism from the free abelian group $\mathcal{Z}^p(X)$ generated by submanifolds of codimension p of X to $\text{Hdg}^{2p}(X)$. This map lifts to a homomorphism to Deligne cohomology $H_{\mathcal{D}}^{2p}(X; \mathbb{Z}(p))$. The latter group fits in the short exact sequence

$$0 \rightarrow J^{2p-1}(X) \rightarrow H_{\mathcal{D}}^{2p}(X; \mathbb{Z}(p)) \rightarrow \text{Hdg}^{2p}(X) \rightarrow 0, \tag{1.1}$$

where $J^{2p-1}(X)$ denotes Griffiths’ intermediate Jacobian (see for example [24, Section 12]). On the subgroup $\mathcal{Z}_{\text{hom}}^p(X)$ of submanifolds whose fundamental class is homologically trivial sequence (1.1) induces the Abel–Jacobi map $\mathcal{Z}_{\text{hom}}^p(X) \rightarrow J^{2p-1}(X)$. As described in [24, Section 12.1] this map has a concrete geometric description via evaluating integrals over singular cycles in X , and one may consider it as a secondary cohomology invariant. In [19, 20] Karoubi constructed an analog of Deligne cohomology for complex K -theory over complex manifolds in which secondary invariants for vector bundles can be defined (see also [9]). In [16], Hopkins and Quick show that there is a bigraded analog of Deligne cohomology $E_{\mathcal{D}}$ for every rationally even cohomology theory E . If X is a compact Kähler manifold, there is a short exact sequence

$$0 \rightarrow J_E^{2p-1}(X) \rightarrow E_{\mathcal{D}}^{2p}(p)(X) \rightarrow \text{Hdg}_E^{2p}(X) \rightarrow 0$$

generalizing sequence (1.1). Let X be a smooth projective complex algebraic variety and $\tilde{\mathcal{M}}^p(X)$ be the free abelian group generated by isomorphism classes $[f]$ of projective smooth morphisms $f: Y \rightarrow X$ of codimension p between complex algebraic varieties. Based on the work of Levine and Morel [21] on algebraic cobordism, it is shown in [16] that for $E = MU$ there is a natural homomorphism $\hat{\varphi}: \tilde{\mathcal{M}}^p(X) \rightarrow MU_{\mathcal{D}}^{2p}(p)(X)$ where X also denotes the underlying complex manifold of complex points of X . On the subgroup $\tilde{\mathcal{M}}^p(X)_{\text{top}}$ of topologically cobordant maps this induces an Abel–Jacobi type homomorphism $\text{AJ}: \tilde{\mathcal{M}}^p(X)_{\text{top}} \rightarrow J_{MU}^{2p-1}(X)$. This homomorphism has been studied in more detail in [22]. However, both $\hat{\varphi}$ and AJ are only defined for complex algebraic varieties and are not induced by a geometric procedure as their classical analogs, but by a rather abstract machinery.

In [14], the authors define for every complex manifold X and integers n and p , geometric Hodge filtered complex cobordism groups $MU^n(p)(X)$ recalled below. The main result of [14] is that there is a natural isomorphism of Hodge filtered cohomology groups

$$MU_{\mathcal{D}}^n(p)(X) \cong MU^n(p)(X). \tag{1.2}$$

The aim of the present paper is to construct pushforward homomorphisms along proper holomorphic maps for geometric Hodge filtered cobordism. This will allow us to give a concrete description of the Hodge filtered fundamental classes of holomorphic maps $f: Y \rightarrow X$ for any complex manifold X and of the Abel–Jacobi invariant AJ for topologically trivial cobordism cycles on compact Kähler manifolds. We note that the results of the present paper are independent of the comparison isomorphism (1.2) of [14]. The only results from [14] that we assume here are the verification of some natural properties of the groups $MU^n(p)(X)$.

We will now briefly describe the construction of the groups $MU^n(p)(X)$ of [14] which we also recall in more detail in Section 2 and will then describe our main results in more detail. Consider the genus $\phi: MU_* \rightarrow \mathcal{V}_* := MU_* \otimes_{\mathbb{Z}} \mathbb{C}$ given by multiplication by $(2\pi i)^n$ in degree $2n$. By Thom’s theorem, MU_n is the bordism group of n -dimensional almost complex manifolds Z . Hirzebruch showed that any genus $\phi: MU_* \rightarrow \mathcal{V}_*$ is of the form

$$\phi(Z) = \int_Z (K^\phi(TZ))^{-1}$$

for a multiplicative sequence K^ϕ , where TZ denotes the tangent bundle of Z . This yields a \mathcal{V}_* -valued characteristic class of complex vector bundles. For $p \in \mathbb{Z}$, we consider the characteristic class $K^p = (2\pi i)^p \cdot K^\phi$. If ∇ is a connection on a complex vector bundle E , Chern–Weil theory gives a form $K^p(\nabla)$ representing $K^p(E)$. Given a form ω on Z and a proper oriented map $f: Z \rightarrow X$, we consider the pushforward current $f_*\omega$, which acts on compactly supported forms on X by $\sigma \mapsto \int_Z \omega \wedge f^*\sigma$. We define Hodge filtered cobordism cycles as triples (f, ∇, h) where f is a proper complex-oriented map $f: Z \rightarrow X$, ∇ is a connection on the complex stable normal bundle of f and h is a current on X such that

$$f_*K^p(\nabla) - dh \text{ is a smooth form in } F^p\mathcal{A}^n(X; \mathcal{V}_*).$$

After defining a suitable Hodge filtered bordism relation we obtain the group $MU^n(p)(X)$ of Hodge filtered cobordism classes. The main new technical contribution of the present paper is the construction of pushforwards for geometric Hodge filtered complex cobordism.

Theorem 1.1. *Let $g: X \rightarrow Y$ be a proper holomorphic map between complex manifolds of complex codimension $d = \dim_{\mathbb{C}} Y - \dim_{\mathbb{C}} X$. Then there is a pushforward homomorphism of $MU^*(*)(Y)$ -modules*

$$g_*: MU^n(p)(X) \rightarrow MU^{n+2d}(p+d)(Y)$$

which is functorial for proper holomorphic maps and compatible with pullbacks.

In [16, Section 7], the authors show that there is an $MU_{\mathcal{D}}$ -pushforward along projective morphisms between smooth projective complex varieties. In fact, they show that there are pushforward maps for a logarithmically refined version of $MU_{\mathcal{D}}$ for quasi-projective smooth complex varieties, as a rather formal consequence of the projective bundle formula. This theory coincides with $MU_{\mathcal{D}}$ for projective smooth complex varieties. Hence, assuming the comparison isomorphism (1.2) of [14], Theorem 1.1 extends the existence of pushforwards to a significantly larger class of maps than the one in [16]. Since pushforwards in cohomology $g_*: MU^n(X) \rightarrow MU^{n+2d}(Y)$ only exist for proper and complex oriented continuous maps of (real) codimension d , the class of proper holomorphic maps is the largest possible subclass of holomorphic maps for which a pushforward with good properties may exist.

The construction of g_* in Theorem 1.1 is similar to the one of pushforwards for differential cobordism for smooth manifolds in [5]. However, the pushforward in differential cobordism exists only for proper submersions with a choice of a smooth MU -orientation. We will now explain why the pushforward for Hodge filtered cobordism exists for all proper holomorphic maps. In Section 3 we first define the group of a Hodge filtered MU -orientation as a Grothendieck group of triples (E, ∇, σ) where E is a complex vector bundle with connection ∇ and σ is a form on X such that $K(\nabla) - d\sigma \in F^0 \mathcal{A}^0(X; \mathcal{V}_*)$. The relations involve a Chern–Simons transgression form associated to the multiplicative sequence K . A Hodge filtered MU -oriented map is then a holomorphic map with a lift of the stable normal bundle to the group of Hodge filtered MU -orientations. Then we show in Section 4 that there is a pushforward along every proper Hodge filtered MU -oriented map. Finally, we show in Section 5 that there is a canonical choice of a Hodge filtered MU -orientation for every proper holomorphic map. The key idea is a variation of a result of Karoubi’s [19, Theorem 6.7] which establishes a mapping of virtual holomorphic vector bundles to the group of Hodge filtered MU -orientations by picking a Bott connection.¹ Applying this result to the virtual normal bundle of a holomorphic map defines a canonical orientation which we call the *Bott orientation*.

Another crucial point for the construction of pushforwards is that there is a currential version of Hodge filtered cobordism which we introduce in Section 2.3. A key difference to differential cohomology theories on smooth manifolds, such as differential cobordism or differential K -theory, is that, for Hodge filtered cobordism, the currential description and the one using forms are canonically isomorphic. This is not the case for differential theories as explained for differential K -theory in [10], where particularly the exact sequences [10, (2.20)] and [10, (2.29)] make it clear that the smooth and currential differential K -theory groups are different in general. The main reason is that the space of closed currents $\mathcal{D}^n(X)_{cl}$ is strictly larger than the space of closed forms $\mathcal{A}^n(X)_{cl}$. In the Hodge filtered context, however, we use $H^n(X; F^p \mathcal{A}^*)$ and $H^n(X; F^p \mathcal{D}^*)$, and $H^n(X; \mathcal{A}^*/F^p)$ and $H^n(X; \mathcal{D}^*/F^p)$ instead of $\mathcal{A}^n(X)/\text{Im}(d)$ and $\mathcal{D}^n(X)/\text{Im}(d)$. Since the Dolbeault–Grothendieck lemma holds both for currents and forms, in both cases the canonical map from the first to the second group is an isomorphism (see Lemma 2.2 and Theorem 2.16).

We will now describe the remaining content and results of the paper. In Section 4, we show that the pushforward is functorial, compatible with pullbacks and satisfies a projection formula. In Section 6, we introduce the Hodge filtered fundamental class $[f] = f_*(1) \in MU^{2p}(p)(X)$ associated to a holomorphic map $f: Y \rightarrow X$ of codimension p as the pushforward of the unit element along f . If X is a compact Kähler manifold and f is a nullbordant proper holomorphic map, then $[f]$ has image in the subgroup $J_{MU}^{2p-1}(X)$ which has the structure of a complex torus. In this case we also write $\text{AJ}(f) := [f]$ for the

¹The notion of Bott connection in [19, Section 6] is a generalization of Bott connections in foliation theory. For complex manifolds, Bott connections are connections compatible with the holomorphic structure.

class of f in $J_{MU}^{2p-1}(X)$. Since the pushforward f_* has a geometric construction, we are able to give a geometric description of the secondary invariant $AJ(f)$ in Section 6. The main ingredient in the formulas are certain Chern–Simons transgression forms as in [6] mediating between an arbitrary connection on the normal bundle N_f and Bott connections on the corresponding tangent bundles. In Section 7 we present a cycle model for Deligne cohomology inspired by but slightly simpler than the one of Gillet and Soulé in [11]. The main difference is that we use currents of integration instead of integral currents in the sense of geometric measure theory (see also [13]). The new construction may be of independent interest and useful for other applications. This enables us to give a cycle description of the Hodge filtered Thom morphism $MU_{\mathcal{D}}^n(p)(X) \rightarrow H_{\mathcal{D}}^n(X; \mathbb{Z}(p))$ for every complex manifold X and integers n and p . In Section 8, we report on our knowledge of the current status of examples and phenomena related to the kernel and image of the Hodge filtered Thom morphism for compact Kähler manifolds.

2. Currential geometric Hodge filtered cobordism

First we briefly recall some facts about currents and the construction of geometric Hodge filtered complex cobordism groups from [14]. Then we introduce a currential version of Hodge filtered cobordism.

2.1. Currents

Let X be a smooth manifold and let Λ_X denote the orientation bundle of X . Let $\mathcal{A}_c^*(X; \Lambda_X)$ be the space of compactly supported smooth forms on X with values in Λ_X . Let $\mathcal{D}^*(X)$ denote the space of currents on X , defined as the topological dual of $\mathcal{A}_c^*(X; \Lambda_X)$. Given a form $\omega \in \mathcal{A}^*(X)$ and a current $T \in \mathcal{D}^*(X)$, their product acts by

$$T \wedge \omega(\sigma) = T(\omega \wedge \sigma).$$

There is an injective map $\mathcal{A}^*(X) \hookrightarrow \mathcal{D}^*(X)$ given by

$$\omega \mapsto T_\omega = \left(\sigma \mapsto \int_X \omega \wedge \sigma, \sigma \in \mathcal{A}_c^*(X; \Lambda_X) \right).$$

We equip \mathcal{D}^* with a grading so that this injection preserves degrees. That is, $\mathcal{D}^k(X)$ consists of the currents which vanish on a homogeneous Λ_X -valued form σ , unless possibly if $\deg \sigma = \dim_{\mathbb{R}} X - k$. We will not always distinguish ω from T_ω in our notation.

If X is a manifold without boundary, Stokes’ theorem implies for $\omega \in \mathcal{A}^k(X)$:

$$T_{d\omega}(\sigma) = (-1)^{k+1} T_\omega(d\sigma).$$

Hence the exterior differential can be extended to a map $d: \mathcal{D}^k(X) \rightarrow \mathcal{D}^{k+1}(X)$ by

$$dT(\sigma) = (-1)^{k+1} T(d\sigma).$$

For a vector space V we set $\mathcal{D}^*(X; V) = \mathcal{D}^*(X) \otimes V$, and for an evenly graded complex vector space $\mathcal{V}_* = \bigoplus_j \mathcal{V}_{2j}$ we set

$$\mathcal{D}^n(X; \mathcal{V}_*) = \bigoplus_j \mathcal{D}^{n+2j}(X; \mathcal{V}_{2j}).$$

An orientation of a map $f: Z \rightarrow X$ is equivalent to an isomorphism $\Lambda_Z \cong f^* \Lambda_X$. If f is proper and oriented, we therefore get a map

$$f_*: \mathcal{A}_c^*(X; \Lambda_X) \rightarrow \mathcal{A}_c^*(Z; \Lambda_Z)$$

which induces a homomorphism

$$f_*: \mathcal{D}^*(Z) \rightarrow \mathcal{D}^{*+d}(X),$$

where $d = \text{codim } f = \dim X - \dim Z$. We also denote by f_* the homomorphism

$$\mathcal{D}^*(Z; \mathcal{V}_*) \rightarrow \mathcal{D}^*(X; \mathcal{V}_*)$$

induced by tensoring f_* with the identity of the various \mathcal{V}_{2j} . We then have the identity

$$d \circ f_* = (-1)^d f_* \circ d.$$

Remark 2.1. In the case of a submersion $\pi: W \rightarrow X$ the pushforward π_* preserves smoothness. We thus obtain the *integration over the fiber* map

$$\int_{W/X} : \mathcal{A}^*(W) \rightarrow \mathcal{A}^{*+d}(X)$$

defined by the equation

$$T_{\int_{W/X} \omega} = \pi_* T_\omega.$$

Now we assume that X is a complex manifold. Then the space of currents is bigraded as follows. We write $\mathcal{D}^{p,q}(X)$ for the subgroup of those currents which vanish on compactly supported (p', q') -forms unless $p' + p = \dim_{\mathbb{C}} X = q' + q$. Then $\mathcal{A}^{*,*}(X) \rightarrow \mathcal{D}^{*,*}(X)$, $\omega \mapsto T_\omega$, is a morphism of double complexes. The Hodge filtration on currents is defined by

$$F^p \mathcal{D}^n(X) = \bigoplus_{i \geq p} \mathcal{D}^{i, n-i}(X).$$

For an evenly graded complex vector space \mathcal{V}_* we set

$$F^p \mathcal{D}^n(X; \mathcal{V}_*) := \bigoplus_j F^{p+j} \mathcal{D}^{n+2j}(X; \mathcal{V}_{2j}),$$

$$\frac{\mathcal{D}^n}{F^p}(X; \mathcal{V}_*) := \bigoplus_j \frac{\mathcal{D}^{n+2j}(X; \mathcal{V}_{2j})}{F^{p+j} \mathcal{D}^{n+2j}(X; \mathcal{V}_{2j})}.$$

With similar notation for $F^p \mathcal{A}^*(X; \mathcal{V}_*)$ and $\frac{\mathcal{A}^*}{F^p}(X; \mathcal{V}_*)$ we get the following result which will be crucial for the proof of Theorem 2.16.

Lemma 2.2. *Let X be a complex manifold. For every p , the maps of complexes of sheaves*

$$F^p \mathcal{A}^*(\mathcal{V}_*) \rightarrow F^p \mathcal{D}^*(\mathcal{V}_*) \quad \text{and} \quad \frac{\mathcal{A}^*}{F^p}(\mathcal{V}_*) \rightarrow \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*)$$

are quasi-isomorphisms on the site of open subsets of X .

Proof. It suffices to prove the assertions for $\mathcal{V}_* = \mathbb{C}$. Let Ω^p be the sheaf of holomorphic p forms on X . The maps of complexes

$$\Omega^p \rightarrow \mathcal{A}^{p,*} \rightarrow \mathcal{D}^{p,*}$$

are quasi-isomorphisms by the Dolbeault–Grothendieck lemma as formulated and proven in [12, pp. 382–385] (see also [8, Lemma 3.29, p. 28]), where we consider Ω^p as a complex concentrated in a single degree. The sheaves $\mathcal{A}^{p,q}$ and $\mathcal{D}^{p,q}$, being modules over \mathcal{A}^0 , are *fine*. Therefore, it follows from [24, Lemma 8.5] that the solid inclusions

$$\begin{array}{ccc} & \Omega^{*\geq p} & \\ \swarrow & & \searrow \\ F^p \mathcal{A}^* & \cdots \cdots \cdots \rightarrow & F^p \mathcal{D}^* \end{array}$$

are quasi-isomorphisms. This implies that the dotted arrow is a quasi-isomorphism and proves the first assertion. By the same argument we have de Rham’s theorem and the inclusion $\mathcal{A}^* \rightarrow \mathcal{D}^*$ is a quasi-isomorphism. Hence the induced map between the cokernels of the maps $\mathcal{A}^* \rightarrow \mathcal{D}^*$ and $F^p \mathcal{A}^* \rightarrow F^p \mathcal{D}^*$ is a quasi-isomorphism as well. This proves the second assertion. ■

2.2. Geometric Hodge filtered cobordism groups

We briefly recall the construction of the geometric Hodge filtered cobordism groups of [14]. For further details we refer to [14, Section 2]. Let $\mathbf{Man}_{\mathbb{C}}$ denote the category of complex manifolds with holomorphic maps. For $X \in \mathbf{Man}_{\mathbb{C}}$ let $f: Z \rightarrow X$ be a proper complex-oriented map, and let N_f be a complex vector bundle which represents the stable normal bundle of f . Let ∇_f be a connection on N_f . We call the triple $\tilde{f} = (f, N_f, \nabla_f)$ a *geometric cycle* over X . We let $\widetilde{ZMU}^n(X)$ denote the abelian group generated by isomorphism classes, in the obvious sense, of geometric cycles over X of codimension n with the relations

$$\tilde{f}_1 + \tilde{f}_2 = \tilde{f}_1 \sqcup \tilde{f}_2.$$

Let MU_* be the graded ring with $MU_n = MU^{-n}(\text{pt})$. A map of rings $MU_* \rightarrow R$ for R an integral domain over \mathbb{Q} is called a complex genus. Complex genera may be constructed in the following way. For each $i \in \mathbb{N}$, let x_i be an indeterminate of degree i . Let $Q \in R[[y]]$ be a formal power series in the variable y of degree 2. Let σ_i denote the i -th elementary symmetric function in x_1, x_2, \dots . We may then define a sequence of

polynomials K_i^Q satisfying

$$K^Q(\sigma_1, \sigma_2, \dots) = 1 + K_2^Q(\sigma_1) + K_4^Q(\sigma_1, \sigma_2) + \dots = \prod_{i=1}^{\infty} Q(x_i)$$

since the right-hand side is symmetric in the x_i . Then we get a characteristic class K^Q defined on a complex vector bundle $E \rightarrow X$ of dimension n by

$$K^Q(E) := K^Q(c_1(E), \dots, c_n(E)) \in H^*(X; R),$$

where $c_i(E)$ denotes the i -th Chern class of E . In fact, by [15, Section 1.8], all genera are of the form

$$\phi^Q([X]) = \int_X K^Q(N_X),$$

where N_X denotes the complex vector bundle representing the stable normal bundle of X obtained from the complex orientation of $X \rightarrow \text{pt}$. From now on we set $\mathcal{V}_* := MU_* \otimes_{\mathbb{Z}} \mathbb{C}$. We assume that the power series $Q(y) = 1 + r_1 y + r_2 y^2 + \dots$ has total degree 0. This is equivalent to assuming ϕ^Q to be a degree-preserving genus. Then $K^Q(E)$ has total degree 0. By [5, Lemma 3.26], ϕ^Q extends to a morphism of multiplicative cohomology theories

$$\phi^Q: MU^n(X) \rightarrow H^n(X; \mathcal{V}_*)$$

by

$$\phi^Q([f]) = f_* K^Q(N_f).$$

Here $H^n(X; \mathcal{V}_*) \cong \bigoplus_j H^{n+2j}(X; \mathcal{V}_{2j})$, so that in particular $H^{-2j}(\text{pt}; \mathcal{V}_*) \cong \mathcal{V}_{2j}$. Now we fix the multiplicative natural transformation

$$\phi: MU^*(X) \rightarrow H^*(X; \mathcal{V}_*)$$

characterized by restricting to multiplication with $(2\pi i)^k$ on $MU_{2k} \rightarrow MU_{2k} \otimes \mathbb{C}$. Let

$$K = 1 + K_2(\sigma_1) + K_4(\sigma_1, \sigma_2) + \dots$$

be the multiplicative sequence satisfying $\phi([f]) = f_* K(N_f)$. For $p \in \mathbb{Z}$ we set $K^p = (2\pi i)^p \cdot K$ and

$$\phi^p([f]) = f_* K^p(N_f).$$

Let $f: Z \rightarrow X$ be a proper complex-oriented map, and let ∇_f be a connection on N_f . By Chern–Weil theory there is a well-defined form $c(\nabla_f) \in \mathcal{A}^*(Z)$ representing the total Chern class $c(N_f)$. In fact, with respect to local coordinates, we have

$$c(\nabla_f) = 1 + c_1(\nabla_f) + c_2(\nabla_f) + \dots = \det\left(I - \frac{1}{2\pi i} F^{\nabla_f}\right),$$

where F^{∇_f} denotes the curvature of ∇_f . Then the form

$$K(\nabla_f) := K(c_1(\nabla_f), c_2(\nabla_f), \dots) \in \mathcal{A}^0(Z; \mathcal{V}_*)$$

represents the cohomology class $K(N_f)$.

Definition 2.3. For a geometric cycle $\tilde{f} \in \widetilde{ZMU}^n(X)$ we define, using the orientation of f induced by its complex orientation, the current

$$\phi^p(\tilde{f}) = f_* K^p(\nabla_f) \in \mathcal{D}^n(X; \mathcal{V}_*).$$

Note that $\phi^p(\tilde{f})$ is a closed current representing the cohomology class

$$\phi^p([f]) = f_* K^p(N_f) \in H^n(X; \mathcal{V}_*).$$

By de Rham’s work [7, Theorem 14] we can always find a current $h \in \mathcal{D}^{n-1}(X; \mathcal{V}_*)$ such that

$$\phi^p(\tilde{f}) - dh = f_* K^p(\nabla_f) - dh \text{ is a form, i.e., lies in } \mathcal{A}^n(X; \mathcal{V}_*).$$

Definition 2.4. Let X be a complex manifold and n, p integers. The group of *Hodge filtered cycles* of degree (n, p) on X is defined as the subgroup

$$ZMU^n(p)(X) \subset (\widetilde{ZMU}^n(X) \times \mathcal{D}^{n-1}(X; \mathcal{V}_*) / d\mathcal{D}^{n-2}(X; \mathcal{V}_*))$$

consisting of pairs $\gamma = (\tilde{f}, h)$ satisfying

$$f_* K^p(\nabla_f) - dh \in F^p \mathcal{A}^n(X; \mathcal{V}_*).$$

Remark 2.5. To simplify the notation, we will often write ϕ and K instead of ϕ^p and K^p , respectively. We may sometimes consider a Hodge filtered cobordism cycle as a triple

$$\gamma = (\tilde{f}, \omega, h) \in \widetilde{ZMU}^n(X) \times F^p \mathcal{A}^n(X; \mathcal{V}_*) \times \mathcal{D}^{n-1}(X; \mathcal{V}_*) / d\mathcal{D}^{n-2}(X; \mathcal{V}_*),$$

where $(\tilde{f}, h) \in ZMU^n(p)(X)$ and the form $\omega := \phi(\tilde{f}) - dh = f_* K(\nabla_f) - dh$.

Next we introduce the cobordism relation. The group of geometric bordism data over X is the subgroup of elements $\tilde{b} \in \widetilde{ZMU}^n(\mathbb{R} \times X)$, with underlying maps of the form $b = (c_b, f): W \rightarrow \mathbb{R} \times X$ such that 0 and 1 are regular values for c_b . Then $W_t = c_b^{-1}(t)$ is a closed manifold for $t = 0, 1$, and $f_t = f|_{W_t}$ is a geometric cycle. We define

$$\partial \tilde{b} := \tilde{f}_1 - \tilde{f}_0 \in \widetilde{ZMU}^n(X)$$

and, setting $W_{[0,1]} = c_b^{-1}([0, 1])$, we define the current

$$\psi^p(\tilde{b}) := (-1)^n (f|_{W_{[0,1]}})_* (K^p(\nabla_b)). \tag{2.1}$$

We will often write ψ instead of ψ^p to simplify the notation. By [14, Proposition 2.17], a geometric bordism datum \tilde{b} over X satisfies

$$\phi^p(\partial \tilde{b}) - d\psi^p(\tilde{b}) = 0.$$

Hence we consider $(\partial\tilde{b}, \psi^p(\tilde{b}))$ as a Hodge filtered cycle of degree $(\text{codim } b, p)$. We call such cycles *nullbordant* and let $BMU_{\text{geo}}^n(p)(X) \subset ZMU^n(p)(X)$ denote the subgroup they generate. We follow Karoubi in [19, Section 4.1] and denote

$$\tilde{F}^p \mathcal{A}^{n-1}(X; \mathcal{V}_*) := F^p \mathcal{A}^{n-1}(X; \mathcal{V}_*) + d \mathcal{A}^{n-2}(X; \mathcal{V}_*). \tag{2.2}$$

We define the map

$$a: d^{-1}(F^p \mathcal{A}^n(X; \mathcal{V}_*))^{n-1} \rightarrow ZMU^n(p)(X), \quad a(h) := (0, h), \tag{2.3}$$

where $d^{-1}(F^p \mathcal{A}^n(X; \mathcal{V}_*))^{n-1}$ denotes the subset of elements in $\mathcal{A}^{n-1}(X; \mathcal{V}_*)$ which are sent to the subgroup $F^p \mathcal{A}^n(X; \mathcal{V}_*)$ under $d: \mathcal{A}^{n-1}(X; \mathcal{V}_*) \rightarrow \mathcal{A}^n(X; \mathcal{V}_*)$. The group of *Hodge filtered cobordism relations* is defined as

$$BMU^n(p)(X) = BMU_{\text{geo}}^n(p)(X) + a(\tilde{F}^p \mathcal{A}^{n-1}(X; \mathcal{V}_*)).$$

Definition 2.6. Let $X \in \mathbf{Man}_{\mathbb{C}}$ and let n and p be integers. The *geometric Hodge filtered cobordism group* of X of degree (n, p) is defined as the quotient

$$MU^n(p)(X) := \frac{ZMU^n(p)(X)}{BMU^n(p)(X)}.$$

We denote the Hodge filtered cobordism class of the cycle $\gamma = (\tilde{f}, h) = (f, N_f, \nabla_f, h)$ by $[\gamma] = [\tilde{f}, h] = [f, N_f, \nabla_f, h]$.

We define maps R and I on the level of cycles as follows:

$$\begin{aligned} R: ZMU^n(p)(X) &\rightarrow F^p \mathcal{A}^n(X; \mathcal{V}_*)_{\text{cl}}, & R(\tilde{f}, h) &= f_* K(\nabla_f) - dh, \\ I: ZMU^n(p)(X) &\rightarrow ZMU^n(X), & I(\tilde{f}, h) &= f. \end{aligned} \tag{2.4}$$

Note that the maps R, I , and a above induce well-defined homomorphisms on cohomology by [14, Proposition 2.19].

Remark 2.7. Note that $[R(\gamma)] = \phi(I(\gamma))$. In that sense, R refines the topological information of I with Hodge filtered differential geometric content. It is shown in [14] that R and I fit in a homotopy pullback in a suitable model category which can be used to construct Hodge filtered cobordism.

For the following theorem we let $\bar{\phi}$ denote the composition of ϕ with the homomorphism induced by reducing the coefficients modulo F^p .

Theorem 2.8. *For every $p \in \mathbb{Z}$, the assignment $X \mapsto MU^*(p)(X)$ has the following properties:*

- For every $X \in \mathbf{Man}_{\mathbb{C}}$ there is the following long exact sequence:

$$\begin{array}{ccccccc} \dots & \longrightarrow & H^{n-1}(X; \frac{\mathcal{A}^*}{F^p}(\mathcal{V}_*)) & \xrightarrow{a} & MU^n(p)(X) & \xrightarrow{I} & MU^n(X) \\ & & \xrightarrow{\bar{\phi}} & & H^n(X; \frac{\mathcal{A}^*}{F^p}(\mathcal{V}_*)) & \xrightarrow{a} & MU^{n+1}(p)(X) \longrightarrow \dots \end{array}$$

- For every holomorphic map $g: Y \rightarrow X$ and every n there is a homomorphism

$$g^*: MU^n(p)(X) \rightarrow MU^n(p)(Y).$$

Hence $MU^n(p)$ is a contravariant functor on $\mathbf{Man}_{\mathbb{C}}$.

- For every $X \in \mathbf{Man}_{\mathbb{C}}$, there is a structure of a bigraded ring on

$$MU^*(*)(X) = \bigoplus_{n,p} MU^n(p)(X).$$

Proof. The first assertion is proven in [14, Section 2.6] and follows from a direct verification of the exactness. The second and third assertions are proven in [14, Section 2.7] and [14, Section 2.8], respectively. We will, however, recall the construction of the pullback and of the ring structure in Section 4. ■

For later purposes we now show how the Hodge filtered cobordism class depends on the connection on the representative of the normal bundle.

Definition 2.9. Let X be a smooth manifold, and let

$$\mathcal{E} = (0 \rightarrow E_1 \rightarrow E_2 \rightarrow E_3 \rightarrow 0)$$

be a short exact sequence of complex vector bundles over X with connections ∇^{E_i} on E_i . Let $\pi: [0, 1] \times X \rightarrow X$ denote the projection. Let $\nabla^{\pi^*E_2}$ be a connection on π^*E_2 which equals $\pi^*\nabla^{E_2}$ near $\{1\} \times X$ and equals $\pi^*(\nabla^{E_1} \oplus \nabla^{E_3})$ near $\{0\} \times X$. The Chern–Simons transgression form of the short exact sequence \mathcal{E} associated to the multiplicative sequence K is given by

$$\begin{aligned} CS_K(\mathcal{E}) &= CS_K(\nabla^{E_1}, \nabla^{E_2}, \nabla^{E_3}) \\ &= \int_{[0,1] \times X / X} K(\nabla^{\pi^*E_2}) \in \mathcal{A}^{-1}(X; \mathcal{V}_*) / \text{Im}(d). \end{aligned}$$

Remark 2.10. The construction of $CS_K(\mathcal{E})$ requires choosing a section $s: E_3 \rightarrow E_2$ as well as a connection $\nabla^{\pi^*E_2}$. However, the form $CS_K(\mathcal{E})$ is independent of these choices in the quotient $\mathcal{A}^{-1}(X; \mathcal{V}_*) / \text{Im}(d)$. By Stokes’ theorem, the derivative of the Chern–Simons form $CS_K(\nabla^{E_1}, \nabla^{E_2}, \nabla^{E_3})$ satisfies

$$\begin{aligned} d CS_K(\nabla^{E_1}, \nabla^{E_2}, \nabla^{E_3}) &= K(\nabla^{E_2}) - K(\nabla^{E_1} \oplus \nabla^{E_3}) \\ &= K(\nabla^{E_2}) - K(\nabla^{E_1}) \wedge K(\nabla^{E_3}). \end{aligned}$$

Remark 2.11. We will often consider the following special case. Let E be a complex vector bundle over the smooth manifold X . Let ∇_0 and ∇_1 be two connections on E . We can form the short exact sequence

$$\mathcal{E} = (0 \rightarrow E \xrightarrow{\text{id}} E \rightarrow 0 \rightarrow 0)$$

and define $CS_K(\nabla_0, \nabla_1) := CS_K(\mathcal{E})$. This Chern–Simons transgression form can be expressed as

$$CS_K(\nabla_1, \nabla_0) = \int_{[0,1] \times X/X} K(t \cdot \pi^* \nabla_1 + (1-t) \cdot \pi^* \nabla_0)$$

and its derivative satisfies

$$d CS_K(\nabla_1, \nabla_0) = K(\nabla_1) - K(\nabla_0).$$

Lemma 2.12. *Let $\tilde{f}_0 = (f, N, \nabla_0)$, and $\tilde{f}_1 = (f, N, \nabla_1) \in \widetilde{ZMU}^n(X)$ be two geometric cycles over X with the same underlying complex-oriented map $f: Z \rightarrow X$. Then there is a geometric bordism \tilde{b} with $\partial\tilde{b} = \tilde{f}_1 - \tilde{f}_0$ and*

$$\psi(\tilde{b}) = (-1)^n f_* CS_K(\nabla_0, \nabla_1).$$

Proof. Let $b = \text{id}_{\mathbb{R}} \times f: \mathbb{R} \times Z \rightarrow \mathbb{R} \times X$, and let $\pi_Z: \mathbb{R} \times Z \rightarrow Z$ denote the projection. With the product complex orientation, using $N_{\text{id}_{\mathbb{R}}} = 0$, we have $N_b = N_{\text{id}_{\mathbb{R}} \times f} = \pi_Z^* N_f$. On $\pi_Z^* N_f$ we consider the connection

$$\nabla_b := t \cdot \pi_Z^* \nabla_0 + (1-t) \cdot \pi_Z^* \nabla_1,$$

where t is the \mathbb{R} -coordinate. We can then promote b to a geometric bordism

$$\tilde{b} = (b, N_b, \nabla_b).$$

Then we have $\partial\tilde{b} = \tilde{f}_1 - \tilde{f}_0$. Using $f \circ \pi_Z = \pi_X \circ b$ the assertion follows from

$$f_* CS_K(\nabla_0, \nabla_1) = f_* \circ (\pi_Z|_{[0,1] \times Z})_* K(\nabla_b) = ((\pi_X \circ b)|_{[0,1] \times Z})_* K(\nabla_b). \quad \blacksquare$$

2.3. Currential Hodge filtered complex cobordism

Now we introduce a new and alternative description of Hodge filtered cobordism groups by considering the Hodge filtration on currents instead of forms. The difference to the previous definition may seem negligible but turns out to be crucial for the construction of a general pushforward later.

Definition 2.13. Let X be a complex manifold and n, p integers. We define the group of *currential* Hodge filtered cycles $ZMU_{\delta}^n(p)(X)$ as the subgroup

$$ZMU_{\delta}^n(p)(X) \subset (\widetilde{ZMU}^n(X) \times \mathcal{D}^{n-1}(X; \mathcal{V}_*) / d\mathcal{D}^{n-2}(X; \mathcal{V}_*))$$

consisting of pairs (\tilde{f}, h) such that

$$\phi(\tilde{f}) - dh = f_* K(\nabla_f) - dh \in F^p \mathcal{D}^n(X; \mathcal{V}_*). \tag{2.5}$$

We will sometimes write a currential Hodge filtered cycle (\tilde{f}, h) as a triple (\tilde{f}, T, h) with $T = \phi(\tilde{f}) - dh$. Let a_{δ} denote the map

$$a_{\delta}: d^{-1}(F^p \mathcal{D}^n(X; \mathcal{V}_*))^{n-1} \rightarrow ZMU^n(p)(X), \quad a_{\delta}(h) := (0, h),$$

where $d^{-1}(F^p \mathcal{D}^n(X; \mathcal{V}_*))^{n-1}$ denotes the subset of elements in $\mathcal{D}^{n-1}(X; \mathcal{V}_*)$ which are sent to the subgroup $F^p \mathcal{D}^n(X; \mathcal{V}_*)$ under

$$d: \mathcal{D}^{n-1}(X; \mathcal{V}_*) \rightarrow \mathcal{D}^n(X; \mathcal{V}_*).$$

We define the group of *currential cobordism relations* by

$$BMU_\delta^n(p)(X) := BMU_{\text{geo}}^n(X) + a_\delta(\tilde{F}^p \mathcal{D}^*(X; \mathcal{V}_*)).$$

Definition 2.14. For $X \in \text{Man}_\mathbb{C}$ and integers n, p , we define the *currential Hodge filtered cobordism groups* by

$$MU_\delta^n(p)(X) := ZMU_\delta^n(p)(X) / BMU_\delta^n(p)(X).$$

Similar to (2.4), we define maps on the level of currential cycles as follows:

$$\begin{aligned} R_\delta: ZMU_\delta^n(p)(X) &\rightarrow F^p \mathcal{D}^n(X; \mathcal{V}_*), & R_\delta(\tilde{f}, h) &= f_* K(\nabla_f) - dh, \\ I_\delta: ZMU_\delta^n(p)(X) &\rightarrow ZMU^n(X), & I_\delta(\tilde{f}, h) &= f. \end{aligned} \tag{2.6}$$

By slight abuse of notation, we also denote by the symbols R_δ, I_δ and a_δ the corresponding induced homomorphisms on cohomology groups. When the context is clear, we will often drop the subscript δ from the notation. For the next statement let $\bar{\phi}_\delta$ denote the composition of ϕ with the homomorphism induced by reducing the coefficients modulo F^p .

Proposition 2.15. *Let X be a complex manifold. There is a long exact sequence*

$$\begin{array}{ccccccc} \dots & \xrightarrow{\bar{\phi}_\delta} & H^{n-1}(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*)) & \xrightarrow{a_\delta} & MU_\delta^n(p)(X) & \xrightarrow{I_\delta} & MU^n(X) \\ & & \xrightarrow{\bar{\phi}_\delta} & H^n(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*)) & \xrightarrow{a_\delta} & MU_\delta^{n+1}(p)(X) & \xrightarrow{I_\delta} \dots \end{array}$$

Proof. The proof follows that of [14, Theorem 2.21] closely. We provide the details of the proof for the convenience of the reader. We start with exactness at $MU_\delta^n(p)(X)$. By definition of a_δ and I_δ we have

$$I_\delta(a_\delta([h])) = I_\delta([0, dh, h]) = 0.$$

To show the converse we work at the level of cycles. Let $\gamma = (\tilde{f}, h) \in ZMU_\delta^n(p)(X)$ and suppose $I_\delta(\gamma) = 0$. That means $f = \partial b$ for some bordism datum b . We may extend the geometric structure of \tilde{f} over b and obtain a geometric bordism datum \tilde{b} such that $\partial \tilde{b} = \tilde{f}$. We then have

$$(\tilde{f}, h) - (\partial \tilde{b}, \psi(\tilde{b})) = (0, h') = a_\delta(h').$$

The last equality follows from the observation that, since $(0, h') \in ZMU^n(p)(X)$ is a currential Hodge filtered cycle, we must have $dh' \in F^p \mathcal{D}^n(X; \mathcal{V}_*)$. Hence we know $\gamma \in BMU_\delta^n(p)(X)$.

Next we show exactness at $MU^n(X)$. The vanishing $\bar{\phi}_\delta \circ I_\delta = 0$ follows from the following commutative diagram, where the bottom row is exact:

$$\begin{array}{ccccc}
 MU_\delta^n(p)(X) & \xrightarrow{I_\delta} & MU^n(X) & & \\
 R_\delta \downarrow & & \phi \downarrow & \searrow \bar{\phi}_\delta & \\
 H^n(X; F^p \mathcal{D}^*(\mathcal{V}_*)) & \xrightarrow{\text{inc}_*} & H^n(X; \mathcal{V}_*) & \longrightarrow & H^n(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*)).
 \end{array}$$

Conversely, suppose $\bar{\phi}_\delta([f]) = 0$. Then we can find $\tilde{\omega} \in F^p \mathcal{D}^n(X; \mathcal{V})$ such that

$$\phi([f]) = \text{inc}_*([\tilde{\omega}]).$$

Let ∇_f be a connection on N_f so that we get a geometric cycle \tilde{f} with $I_\delta(\tilde{f}) = f$. Then $\phi(\tilde{f})$ is a current representing $\phi([f])$. Hence $\phi(\tilde{f})$ and $\tilde{\omega}$ are cohomologous, i.e., there is a current $h \in \mathcal{D}^{n-1}(X; \mathcal{V}_*)$ such that $\phi(\tilde{f}) - dh = \tilde{\omega}$. Then $\gamma := (\tilde{f}, h)$ is a currential Hodge filtered cycle with $I_\delta(\gamma) = f$.

Now we show exactness at $H^n(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*))$. Let $f: Z \rightarrow X$ be a bordism cycle on X . We will show

$$a_\delta(\bar{\phi}_\delta([f])) = 0 \in MU_\delta^{n+1}(p)(X).$$

Lifting f to a geometric cycle $\tilde{f} \in \widetilde{ZMU}^n(X)$, we can write

$$a_\delta(\bar{\phi}_\delta([f])) = [0, \phi(\tilde{f})].$$

We may build from \tilde{f} a geometric bordism datum \tilde{b} with underlying map

$$Z \xrightarrow{(\frac{1}{2}, f)} \mathbb{R} \times X$$

where $\frac{1}{2}$ denotes the constant map with value $\frac{1}{2}$. We have $\partial \tilde{b} = 0$ and $\psi(\tilde{b}) = (-1)^n \phi(\tilde{f})$. Hence

$$(\partial \tilde{b}, \psi(\tilde{b})) = (0, (-1)^n \phi(\tilde{f})) \in BMU_\delta^n(p)(X)$$

and we conclude that $a_\delta(\bar{\phi}_\delta([f])) = 0$.

Conversely, suppose that $h \in (d^{-1} F^p \mathcal{D}^n(X; \mathcal{V}_*))^{n-1}$ is such that $a_\delta(h) = (0, h)$ represents 0 in $MU_\delta^n(p)(X)$. Then there is a geometric bordism datum \tilde{b} with underlying map $(c_b, f_b): W \rightarrow \mathbb{R} \times X$, and a current $h' \in \tilde{F}^p \mathcal{D}^{n-1}(X; \mathcal{V}_*)$ such that

$$(0, h) = (\partial \tilde{b}, \psi(\tilde{b}) + h').$$

Note that $\tilde{F}^p \mathcal{D}^{n-1}(X; \mathcal{V}_*)$ is the group of relations for $H^{n-1}(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*))$, where we therefore have

$$[h] = [\psi(\tilde{b})].$$

Since $\partial \tilde{b} = 0$, we have

$$f := f_b|_{c_b^{-1}(\{0,1\})} \in ZMU^n(X)$$

is a bordism cycle. By definition of ψ , we have $\psi(\tilde{b}) = (-1)^n \phi(\tilde{f})$ where \tilde{f} is the obvious geometric cycle over f . Hence

$$[h] = [\psi(\tilde{b})] = (-1)^n \bar{\phi}_\delta([f]) \in \text{Im}(\bar{\phi}_\delta).$$

This finishes the proof. ■

There is a natural homomorphism

$$\tau: ZMU^n(p)(X) \rightarrow ZMU_\delta^n(p)(X), \quad (\tilde{f}, h) \mapsto (\tilde{f}, h)$$

which forgets that $\phi(\tilde{f}) - dh$ is a form and not just a current. Since τ sends $BMU^n(p)(X)$ to $BMU_\delta^n(p)(X)$, it follows that there is an induced natural homomorphism

$$\tau: MU^n(p)(X) \rightarrow MU_\delta^n(p)(X).$$

Theorem 2.16. *For every $X \in \mathbf{Man}_\mathbb{C}$ and all integers n and p , the natural homomorphism $\tau: MU^n(p)(X) \rightarrow MU_\delta^n(p)(X)$ is an isomorphism.*

Proof. The long exact sequences of Theorem 2.8 and Proposition 2.15 fit into the commutative diagram

$$\begin{array}{ccccccc} \dots & \longrightarrow & H^{n-1}(X; \frac{\mathcal{A}^*}{F^p}(\mathcal{V}_*)) & \xrightarrow{a} & MU^n(p)(X) & \xrightarrow{I} & MU^n(X) \xrightarrow{\bar{\phi}} \dots \\ & & \cong \downarrow & & \downarrow & & \downarrow \text{id} \\ \dots & \longrightarrow & H^{n-1}(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*)) & \xrightarrow{a_\delta} & MU_\delta^n(p)(X) & \xrightarrow{I_\delta} & MU^n(X) \xrightarrow{\bar{\phi}_\delta} \dots \end{array}$$

That the left-most vertical arrow is an isomorphism, is a corollary of the fact that the Dolbeault–Grothendieck lemma holds for currents as well as forms, see Lemma 2.2 for the full argument. The right-most arrow is the identity. The assertion now follows from the five-lemma. ■

Remark 2.17. In [14], $MU^n(p)(-)$ is defined on the larger category \mathbf{Man}_F of manifolds with a filtration of \mathcal{A}^* . We note that Theorem 2.16 does not extend to that context, since its proof uses that there is a Hodge filtration for currents which extends that of forms and that the inclusion $F^p \mathcal{A}^* \rightarrow F^p \mathcal{D}^*$ is a quasi-isomorphism.

Remark 2.18. As we discussed in the introduction, Theorem 2.16 reflects an important difference between Hodge filtered cohomology and differential cohomology.

3. Hodge filtered MU -orientations

We will now define the notion of a Hodge filtered MU -orientation of a holomorphic map in two steps: First as a type of Hodge filtered K -theory class with \mathcal{V}_* -coefficients. Then we apply this to the normal bundle of a holomorphic map. Recall from (2.2) the notation

$$\tilde{F}^0 \mathcal{A}^{-1}(X; \mathcal{V}_*) = F^0 \mathcal{A}^{-1}(X; \mathcal{V}_*) + \text{Im}(d) \subset \mathcal{A}^{-1}(X; \mathcal{V}_*).$$

Definition 3.1. Let X be a complex manifold. We define the group $K_{MU_{\mathcal{D}}}(X)$ of *Hodge filtered MU-orientations*, *MU $_{\mathcal{D}}$ -orientations* for short, to be the quotient of the free abelian group generated by triples $\varepsilon := (E, \nabla, \sigma)$ where E is a complex vector bundle on X , ∇ is a connection on E and $\sigma \in \mathcal{A}^{-1}(X; \mathcal{V}_*) / \tilde{F}^0 \mathcal{A}^{-1}(X; \mathcal{V}_*)$ such that

$$\mathcal{K}(\varepsilon) = \mathcal{K}(E, \nabla, \sigma) := K(\nabla) - d\sigma \in F^0 \mathcal{A}^0(X; \mathcal{V}_{*cl}) \tag{3.1}$$

is a form in filtration step F^0 modulo the subgroup generated by $(\underline{\mathbb{C}}_X, d, 0)$ for the trivial bundle $\underline{\mathbb{C}}_X$ on X with the canonical connection and by

$$(E_2, \nabla_2, \sigma_2) - (E_1, \nabla_1, \sigma_1) - (E_3, \nabla_3, \sigma_3)$$

whenever there is a short exact sequence of the form

$$0 \rightarrow E_1 \rightarrow E_2 \rightarrow E_3 \rightarrow 0$$

with the identity

$$\sigma_2 = \sigma_1 \wedge \mathcal{K}(\varepsilon_3) + \sigma_3 \wedge K(\nabla_1) + CS_K(\nabla_1, \nabla_2, \nabla_3) \tag{3.2}$$

in $\mathcal{A}^{-1}(X; \mathcal{V}_*) / \tilde{F}^0 \mathcal{A}^{-1}(X; \mathcal{V}_*)$. We denote the image of (E, ∇, σ) in the quotient by $[E, \nabla, \sigma]$.

Remark 3.2. For $q \in \mathbb{Z}$ we could modify the above definition and define an $MU_{\mathcal{D}}$ -orientation of *filtration q* to be a triple (E, ∇, σ) as above such that $\mathcal{K}(\varepsilon) = K(\nabla) - d\sigma \in F^q \mathcal{A}^0(X; \mathcal{V}_*)$. Using appropriate relations, the addition on $K_{MU_{\mathcal{D}}}(X)$ extends to the direct limit of pointed sets over all $q \in \mathbb{Z}$. The group $K_{MU_{\mathcal{D}}}(X)$ of Definition 3.1 is the subgroup of orientations of filtration 0. Since we do not know of applications to support the additional generality and complexity, we only consider orientations of filtration 0 in this paper.

We will now discuss the group $K_{MU_{\mathcal{D}}}(X)$ in more detail.

Lemma 3.3. *The addition in $K_{MU_{\mathcal{D}}}(X)$ is given by*

$$[\varepsilon_1] + [\varepsilon_2] = [E_1, \nabla_1, \sigma_1] + [E_2, \nabla_2, \sigma_2] = [E_1 \oplus E_2, \nabla_1 \oplus \nabla_2, \sigma_{12}],$$

where

$$\sigma_{12} = \sigma_1 \wedge \mathcal{K}(\varepsilon_2) + \sigma_2 \wedge (K(\nabla_1)).$$

Proof. We consider the sequence

$$0 \rightarrow E_1 \rightarrow E_1 \oplus E_2 \rightarrow E_2 \rightarrow 0.$$

Since the connections split, $CS_K(\nabla_1, \nabla_1 \oplus \nabla_2, \nabla_2) = 0$. Hence condition (3.2) for

$$(E_1 \oplus E_2, \nabla_1 \oplus \nabla_2, \sigma_{12}) - (E_1, \nabla_1, \sigma_1) - (E_2, \nabla_2, \sigma_2)$$

to be a relation reduces to

$$\sigma_{12} = \sigma_1 \wedge \mathcal{K}(\varepsilon_2) + \sigma_2 \wedge K(\nabla_1). \quad \blacksquare$$

Remark 3.4. It follows from Lemma 3.3 that the identity element of $K_{MU_{\mathcal{D}}}(X)$ is represented by the triple $(0, d, 0)$ where the first 0 denotes the zero-dimensional trivial bundle.

Remark 3.5. Suppose we have two triples $\varepsilon_1 = (E, \nabla_1, \sigma_1)$ and $\varepsilon_2 = (E, \nabla_2, \sigma_2)$ with the same underlying bundle E . By Remark 3.4, the triple $0 = (0, d, 0)$ represents the identity element in $K_{MU_{\mathcal{D}}}(X)$. Since $K(d) = 1$, considering $E \xrightarrow{\text{id}} E$ as a short exact sequence as in Remark 2.11, we get a relation $[\varepsilon_2] - [\varepsilon_1] - [0]$ for $K_{MU_{\mathcal{D}}}$, i.e., we have $[\varepsilon_1] = [\varepsilon_2]$ in $K_{MU_{\mathcal{D}}}(X)$, if and only if

$$\sigma_2 = \sigma_1 + \text{CS}_K(\nabla_1, \nabla_2).$$

Remark 3.6. There is a hidden symmetry in equation (3.2) of Definition 3.1: Since σ_1 and σ_3 are of odd degree, we have

$$\sigma_1 \wedge d\sigma_3 = \sigma_3 \wedge d\sigma_1 \pmod{\text{Im}(d)}.$$

Hence, modulo $\text{Im}(d)$, we have

$$\begin{aligned} \sigma_1 \wedge (K(\nabla_3) - d\sigma_3) + \sigma_3 \wedge K(\nabla_1) &= \sigma_1 \wedge K(\nabla_3) - \sigma_1 \wedge d\sigma_3 + \sigma_3 \wedge K(\nabla_1) \\ &= \sigma_1 \wedge K(\nabla_3) + \sigma_3 \wedge K(\nabla_1) - \sigma_3 \wedge d\sigma_1 \\ &= \sigma_1 \wedge K(\nabla_3) + \sigma_3 \wedge (K(\nabla_1) - d\sigma_1). \end{aligned}$$

Using the map \mathcal{K} we can rewrite this relation as

$$\sigma_1 \wedge \mathcal{K}(\varepsilon_3) + \sigma_3 \wedge K(\nabla_1) = \sigma_1 \wedge K(\nabla_3) + \sigma_3 \wedge \mathcal{K}(\varepsilon_1) \pmod{\text{Im}(d)}.$$

We now discuss further properties of the assignment

$$\varepsilon = (E, \nabla, \sigma) \mapsto \mathcal{K}(\varepsilon) = K(\nabla) - d\sigma$$

defined in (3.1). We note that there is a certain similarity in the behaviors and roles of the forms $R(\tilde{f}, h)$ (respectively current $R_{\delta}(\tilde{f}, h)$) and $\mathcal{K}(\varepsilon)$. Both $R_{\delta}(\tilde{f}, h)$ and $\mathcal{K}(\varepsilon)$ contribute to the construction of the pushforward along holomorphic maps in Section 4 (see also Lemma 4.2, Remark 4.5, Proposition 4.10 and Remark 4.15). Moreover, while the current $f_*K(\nabla_f)$ is not a cobordism invariant, the class of the difference

$$R(\tilde{f}, h) = f_*K(\nabla_f) - dh$$

is indeed invariant. Similarly, we will show in Proposition 3.8 that $\mathcal{K}(\varepsilon) = K(\nabla) - d\sigma$ is an invariant of the equivalence class $[\varepsilon]$ in $K_{MU_{\mathcal{D}}}(X)$ while $K(\nabla)$ is not. In fact, we will show that \mathcal{K} respects the group structure on $K_{MU_{\mathcal{D}}}(X)$. This result will be used in several of our main results and their proofs. In particular, \mathcal{K} plays a key role in the proof of the functoriality of the pushforward in Theorem 4.11.

Lemma 3.7. For representatives $(E_i, \nabla_i, \sigma_i)$ with $i = 1, 2$ of generators in $K_{MU_{\mathcal{D}}}(X)$ we have

$$\mathcal{K}(\varepsilon_1 + \varepsilon_2) = \mathcal{K}(\varepsilon_1) \wedge \mathcal{K}(\varepsilon_2).$$

Proof. To prove the assertion we use Lemma 3.3. Both $d\sigma_i$ and $K(\nabla_i)$ are closed and lie in $\mathcal{A}^0(X_i; \mathcal{V}_*)_{\text{cl}}$. In particular, this means that they are in the center of the ring $\mathcal{A}^*(X_i; \mathcal{V}_*)$. Using this fact we get

$$\begin{aligned} \mathcal{K}(\varepsilon_1 + \varepsilon_2) &= K(\nabla_1 \oplus \nabla_2) - d(\sigma_1 \wedge (K(\nabla_2) - d\sigma_2) + \sigma_2 \wedge (K(\nabla_1))) \\ &= K(\nabla_1) \wedge K(\nabla_2) - d\sigma_1 \wedge \mathcal{K}(\varepsilon_2) - d\sigma_2 \wedge K(\nabla_1) \\ &= K(\nabla_1) \wedge \mathcal{K}(\varepsilon_2) - d\sigma_1 \wedge \mathcal{K}(\varepsilon_2) \\ &= \mathcal{K}(\varepsilon_1) \wedge \mathcal{K}(\varepsilon_2). \end{aligned}$$

Proposition 3.8. *The map \mathcal{K} descends to a morphism of monoids*

$$\mathcal{K}: (K_{MU_{\mathcal{D}}}(X), +) \rightarrow (F^0\mathcal{A}^0(X; \mathcal{V}_*)_{\text{cl}}, \wedge).$$

Proof. Since the triple $(\underline{\mathbb{C}}_X^N, d, 0)$ represents the identity element in $K_{MU_{\mathcal{D}}}(X)$, we see that \mathcal{K} sends the identity element to the identity. The fact that \mathcal{K} descends to a map on $K_{MU_{\mathcal{D}}}(X)$ and respects the monoid structure then follows from Lemma 3.7 and the defining relations of $K_{MU_{\mathcal{D}}}(X)$. ■

Remark 3.9. Let $\varepsilon = (E, \nabla, \sigma)$ in $K_{MU_{\mathcal{D}}}(X)$ be a representative of a generator in $K_{MU_{\mathcal{D}}}(X)$. We may consider

$$\mathcal{K}(\varepsilon) = K(\nabla) - d\sigma$$

as a power series over the commutative ring $\mathcal{A}^{2*}(X)$ in the generators of \mathcal{V}_* . Having leading term 1, $\mathcal{K}(\varepsilon)$ is an invertible power series.

Remark 3.10. The triple $(\underline{\mathbb{C}}_X^N, d, 0)$ represents the identity element in $K_{MU_{\mathcal{D}}}(X)$. Given a generator $\varepsilon = (E, \nabla, \sigma)$ for $K_{MU_{\mathcal{D}}}(X)$, we can construct a class $[\varepsilon']$ which satisfies $[\varepsilon] + [\varepsilon'] = 0$ in $K_{MU_{\mathcal{D}}}(X)$ as follows: Since X is a finite-dimensional manifold, we can find a complex vector bundle E' and an isomorphism $E \oplus E' \cong \underline{\mathbb{C}}_X^N$ for some N . We equip E' with the connection ∇' induced from d by the direct sum decomposition $E \oplus E' = \underline{\mathbb{C}}_X^N$. Using Remark 3.9 we let

$$\sigma' = -\sigma \wedge K(\nabla') \wedge \mathcal{K}(\varepsilon)^{-1}.$$

To check that $\varepsilon' = (E', \nabla', \sigma')$ satisfies $[\varepsilon] + [\varepsilon'] = 0$ we use Lemma 3.3 to write

$$[\varepsilon] + [\varepsilon'] = [\underline{\mathbb{C}}_X^N, d, \sigma''],$$

where

$$\begin{aligned} \sigma'' &= \sigma \wedge K(\nabla') + \sigma' \wedge \mathcal{K}(\varepsilon) \\ &= \sigma \wedge K(\nabla') - \sigma \wedge K(\nabla') \wedge \mathcal{K}(\varepsilon)^{-1} \wedge \mathcal{K}(\varepsilon) \\ &= 0 \end{aligned}$$

which proves the claim.

Remark 3.11. Assume we have a short exact sequence of complex vector bundles

$$0 \rightarrow E_1 \rightarrow E_2 \rightarrow E_3 \rightarrow 0$$

and that we have orientations $\nu_i = [E_i, \nabla_i, \sigma_i]$ involving two of the three bundles. Then it follows from the defining relations in $K_{MU_{\mathcal{D}}}(X)$ and Remark 3.9 that for any connection ∇_j on the remaining bundle E_j we can find a form σ_j such that $\nu_j = [E_j, \nabla_j, \sigma_j]$ is an orientation and such that $\nu_1 + \nu_3 = \nu_2$.

Definition 3.12. Let $f: X \rightarrow Y$ be a holomorphic map. Since the defining relations are compatible with pullbacks of bundles and connections, there is a well-defined pullback of orientations

$$f^*: K_{MU_{\mathcal{D}}}(Y) \rightarrow K_{MU_{\mathcal{D}}}(X)$$

defined by $f^*[E, \nabla, \sigma] = [f^*E, f^*\nabla, f^*\sigma]$.

Next we define the notion of a Hodge filtered orientation of a holomorphic map. We will use this notion in the following section to define the pushforward along a holomorphic map. In Section 5 we show that every holomorphic map has a canonical choice of a Hodge filtered orientation.

Definition 3.13. Let $g: X \rightarrow Y$ be a holomorphic map. We define a *Hodge filtered MU -orientation of g* , or an $MU_{\mathcal{D}}$ -orientation of g for short, to be a class

$$\nu = [N_g, \nabla_g, \sigma_g] \in K_{MU_{\mathcal{D}}}(X),$$

where N_g represents the complex stable normal bundle associated with g as a complex-oriented map.

Definition 3.14. Let $X_1 \xrightarrow{g_1} X_2 \xrightarrow{g_2} X_3$ be proper holomorphic maps of complex codimension d_1 and d_2 , respectively. Let $\nu_i \in K_{MU_{\mathcal{D}}}(X_i)$ be $MU_{\mathcal{D}}$ -orientations of g_i for $i = 1, 2$. We define the *composed Hodge filtered MU -orientation of $g_2 \circ g_1$* to be

$$\nu_1 + g_1^* \nu_2 \in K_{MU_{\mathcal{D}}}(X_1).$$

Remark 3.15. With the notation of Definition 3.14, we recall that the stable normal bundle of $g_2 \circ g_1$ is isomorphic to the sum of the stable normal bundle of g_1 and the pullback of the stable normal bundle of g_2 along g_1 . Hence $\nu_1 + g_1^* \nu_2$ is, in fact, a Hodge filtered MU -orientation of $g_2 \circ g_1$ in the sense of Definition 3.13.

4. Pushforward along proper Hodge filtered MU -oriented maps

We will now define a pushforward homomorphism for proper $MU_{\mathcal{D}}$ -oriented maps and show that it is functorial. Then we show that the pushforward is compatible with pullback and cup product.

Let $g: X \rightarrow Y$ be a holomorphic map and let \circ be an orientation of g . We write g° for g together with the orientation \circ and refer to g° as an $MU_{\mathcal{D}}$ -oriented holomorphic map. If we want to specify the representative $\varepsilon = (N_g, \nabla_g, \sigma_g)$ of \circ , we write $g^\varepsilon = (g, N_g, \nabla_g, \sigma_g)$, and we write \tilde{g}^ε for the underlying geometric cycle $\tilde{g}^\varepsilon = (g, N_g, \nabla_g)$. We will now define the pushforward of a Hodge filtered cycle along an oriented proper holomorphic map.

Definition 4.1. Let $g: X \rightarrow Y$ be a proper holomorphic map of complex codimension d . Let $\varepsilon = (N_g, \nabla_g, \sigma_g)$ be a representative of an orientation class of g in $KMU_{\mathcal{D}}(X)$. If $\tilde{f} = (f, N_f, \nabla_f)$ is a geometric cycle on X , we write

$$\tilde{g}^\varepsilon \circ \tilde{f} = (g \circ f, N_f \oplus f^*N_g, \nabla_f \oplus f^*\nabla_g)$$

for the composed geometric cycle on Y . We define the *pushforward homomorphism* on currential Hodge filtered cycles by

$$g_*^\varepsilon(\tilde{f}, h) = (\tilde{g}^\varepsilon \circ \tilde{f}, g_*(K(\nabla_g) \wedge h + \sigma_g \wedge R_\delta(\tilde{f}, h))),$$

where g_* denotes the pushforward of currents along g and $R_\delta(\tilde{f}, h) = f_*K(\nabla_f) - dh$ is defined as in (2.6).

We will explain the choices made in Definition 4.1 further in Remarks 4.4 and 4.5 below. But first we need to check that the construction is well-defined, i.e., we have to show that $g_*^\varepsilon(\tilde{f}, h)$ actually is a currential Hodge filtered cycle. We will achieve this in two steps as follows:

Lemma 4.2. *For every $(\tilde{f}, h) \in ZMU_\delta^n(p)(X)$ we have*

$$R_\delta(g_*^\varepsilon(\tilde{f}, h)) = g_*(\mathcal{K}(\varepsilon) \wedge R_\delta(\tilde{f}, h)).$$

Proof. We check this claim by applying the definition of R_δ and then rewrite the current as follows:

$$\begin{aligned} R_\delta(g_*^\varepsilon(\tilde{f}, h)) &= g_*f_*K(\nabla_f \oplus f^*\nabla_g) - dg_*(K(\nabla_g) \wedge h + \sigma_g \wedge R_\delta(\tilde{f}, h)) \\ &= g_*(f_*K(\nabla_f) \wedge K(\nabla_g)) - g_*(K(\nabla_g) \wedge dh + d\sigma_g \wedge R_\delta(\tilde{f}, h)) \\ &= g_*(K(\nabla_g) \wedge (f_*K(\nabla_f) - dh) - d\sigma_g \wedge R_\delta(\tilde{f}, h)) \\ &= g_*((K(\nabla_g) - d\sigma_g) \wedge R_\delta(\tilde{f}, h)) \\ &= g_*(\mathcal{K}(\varepsilon) \wedge R_\delta(\tilde{f}, h)). \end{aligned}$$

■

We can now use this observation to show that $g_*^\varepsilon(\tilde{f}, h)$ is a currential Hodge filtered cycle.

Proposition 4.3. *For every $(\tilde{f}, h) \in ZMU_\delta^n(p)(X)$ we have*

$$g_*^\varepsilon(\tilde{f}, h) \in ZMU_\delta^{n+2d}(p+d)(Y).$$

Proof. It follows from the definition that $\tilde{g}^\varepsilon \circ \tilde{f}$ is a geometric cycle. Hence, by definition of currential Hodge filtered cycles in Definition 2.13, it remains to check that the current

$$R_\delta(g_*^\varepsilon(\tilde{f}, h)) = \phi(\tilde{g}^\varepsilon \circ \tilde{f}) - dh$$

satisfies condition (2.5) on the filtration step of a current in a Hodge filtered cycle, i.e., we have to show that $R_\delta(g_*^\varepsilon(\tilde{f}, h))$ lies in $F^{p+d}\mathcal{D}^{n+2d}(Y; \mathcal{V}_*)$. By Lemma 4.2 we know $R_\delta(g_*^\varepsilon(\tilde{f}, h)) = g_*(\mathcal{K}(\varepsilon) \wedge R_\delta(\tilde{f}, h))$. Hence it suffices to observe that

$$\mathcal{K}(\varepsilon) = K(\nabla_g) - d\sigma_g \in F^0\mathcal{A}^0(X; \mathcal{V}_*)$$

and

$$R_\delta(\tilde{f}, h) = f_*K(\nabla_f) - dh \in F^p\mathcal{D}^n(X; \mathcal{V}_*).$$

Since g is holomorphic of codimension d , it follows that

$$g_*(\mathcal{K}(\varepsilon) \wedge R_\delta(\tilde{f}, h)) \in F^{p+d}\mathcal{D}^{n+2d}(Y; \mathcal{V}_*)$$

as required. ■

Remark 4.4. One might arrive at the formula for the current in the definition of $g_*^\varepsilon(\tilde{f}, h)$ as follows: If $\sigma_g = 0$, then $g_*(K(\nabla_g) \wedge h)$ is the only natural candidate, and it does satisfy the desirable formulas. Having made that choice, consider next the case $\mathfrak{o} = [N_g, \nabla_g, \sigma_g]$ such that there is a connection ∇'_g with $\mathfrak{o} = [N_g, \nabla'_g, 0]$ in $KMU_{\mathcal{D}}(X)$. Then the rest of the formula can be derived using Lemma 2.12 and the relations in $KMU_{\mathcal{D}}(X)$.

Remark 4.5. Let $\mathfrak{o} = [N_g, \nabla_g, \sigma_g] \in KMU_{\mathcal{D}}(X)$. Since σ_g is of degree -1 , we have $d(\sigma_g \wedge h) = d\sigma_g \wedge h - \sigma_g \wedge dh$. Hence, modulo $\text{Im}(d)$, we have $\sigma_g \wedge dh = d\sigma_g \wedge h$, and it follows that modulo $\text{Im}(d)$ we have

$$K(\nabla_g) \wedge h + \sigma_g \wedge (f_*K(\nabla_f) - dh) = (K(\nabla_g) - d\sigma_g) \wedge h + \sigma_g \wedge f_*K(\nabla_f). \quad (4.1)$$

Using the maps \mathcal{K} and R_δ we can rewrite this relation as

$$K(\nabla_g) \wedge h + \sigma_g \wedge R_\delta(\tilde{f}, h) = \mathcal{K}(\mathfrak{o}) \wedge h + \sigma_g \wedge f_*K(\nabla_f).$$

We will now show that the map g_*^ε of Definition 4.1 induces a well-defined pushforward homomorphism on Hodge filtered cobordism. We first show that g_*^ε sends Hodge filtered bordism data to Hodge filtered bordisms in Lemma 4.6.

Lemma 4.6. *We have*

$$g_*^\varepsilon(BMU_\delta^n(p)(X)) \subset BMU_\delta^{n+2d}(p+d)(Y).$$

Proof. Let $h \in \tilde{F}^p\mathcal{D}^{n-1}(X; \mathcal{V}_*)$. By definition of the map a in (2.3), we have $a(h) = (0, h)$. Using relation (4.1) we get

$$g_*(K(\nabla_g) \wedge h - \sigma_g \wedge dh) = g_*((K(\nabla_g) - d\sigma_g) \wedge h).$$

Since $g_*((K(\nabla) - d\sigma_g) \wedge h) \in \tilde{F}^{p+d} \mathcal{D}^{n+2d}(Y; \mathcal{V}_*)$, we conclude that

$$g_*^\varepsilon(a(h)) \in BMU_\delta^{n+2d}(p+d)(Y).$$

It remains to show

$$g_*^\varepsilon(BMU_{\text{geo}}^n(X)) \subset BMU_{\text{geo}}^{n+2d}(Y).$$

This follows from [5, Lemma 4.35]. We provide a proof for the reader's convenience. Let $\tilde{b} \in \widetilde{ZMU}^n(\mathbb{R} \times X)$ be a geometric bordism datum on X . Let \tilde{e} denote the geometric cycle $\text{id}_{\mathbb{R}} \times g: \mathbb{R} \times X \rightarrow \mathbb{R} \times Y$ with the obvious geometric structure. Then $\tilde{e} \circ \tilde{b}$ is a geometric bordism datum over Y .

By definition of $\psi(\tilde{b})$ in (2.1), the fact that g is of even real codimension implies

$$\psi(\tilde{e} \circ \tilde{b}) = g_*(K(\nabla_g) \wedge \psi(\tilde{b})).$$

This shows that we have

$$(\partial(\tilde{e} \circ \tilde{b}), \psi(\tilde{e} \circ \tilde{b})) = (\tilde{g} \circ \partial\tilde{b}, g_*(K(\nabla_g) \wedge \psi(\tilde{b}))).$$

By definition of g_*^ε , we have

$$g_*^\varepsilon(\partial\tilde{b}, \psi(\tilde{b})) = (\tilde{g} \circ \partial\tilde{b}, g_*(K(\nabla_g) \wedge \psi(\tilde{b}) + \sigma_g \wedge R_\delta(\partial\tilde{b}, \psi(\tilde{b}))).$$

By [14, Proposition 2.17], geometric bordism data satisfy $R(\partial\tilde{b}, \psi(\tilde{b})) = 0$ and hence also $R_\delta(\partial\tilde{b}, \psi(\tilde{b})) = 0$. Thus, we get

$$g_*^\varepsilon(\partial\tilde{b}, \psi(\tilde{b})) = (\tilde{g} \circ \partial\tilde{b}, g_*(K(\nabla_g) \wedge \psi(\tilde{b}))).$$

Hence in total we have shown that

$$g_*^\varepsilon(\partial\tilde{b}, \psi(\tilde{b})) = (\partial(\tilde{e} \circ \tilde{b}), \psi(\tilde{e} \circ \tilde{b})).$$

This shows $g_*^\varepsilon(BMU_{\text{geo}}^n(X)) \subset BMU_{\text{geo}}^{n+2d}(Y)$ and finishes the proof. ■

Next we show that the equivalence class of $g_*^\varepsilon(\tilde{f}, h)$ does not depend on the choice of a representative of the $MU_{\mathcal{D}}$ -orientation on g .

Lemma 4.7. *Let $\varepsilon = (N_g, \nabla, \sigma)$ and $\varepsilon' = (N_g, \nabla', \sigma')$ be two representatives of the $MU_{\mathcal{D}}$ -orientation \circ of $g: X \rightarrow Y$. Then, for each $\gamma \in ZMU_\delta^n(p)(X)$, we have*

$$[g_*^\varepsilon \gamma] = [g_*^{\varepsilon'} \gamma] \text{ in } MU_\delta^{n+2d}(p+d)(Y).$$

Proof. Let $\gamma = (\tilde{f}, h)$ be a currential cycle. By the definition of $g_*^\varepsilon \gamma$ we have

$$[g_*^\varepsilon(\tilde{f}, h)] = [\tilde{g}^\varepsilon \circ \tilde{f}, g_*(K(\nabla) \wedge h + \sigma \wedge R_\delta(\tilde{f}, h))].$$

Using (4.1), we get

$$[g_*^\varepsilon(\tilde{f}, h)] = [\tilde{g}^\varepsilon \circ \tilde{f}, g_*(\mathcal{K}(\varepsilon) \wedge h + \sigma \wedge f_* K(\nabla_f))]. \tag{4.2}$$

Similarly, for the representative ε' , we get

$$[g_*^{\varepsilon'}(\tilde{f}, h)] = [\tilde{g}^{\varepsilon'} \circ \tilde{f}, g_*(\mathcal{K}(\varepsilon') \wedge h + \sigma' \wedge f_*K(\nabla_f))]. \tag{4.3}$$

We need to show that the two cycles in (4.2) and (4.3), respectively, are connected by a Hodge filtered bordism. By Proposition 3.8, we know

$$\mathcal{K}(\varepsilon) = K(\nabla) - d\sigma = K(\nabla') - d\sigma' = \mathcal{K}(\varepsilon'). \tag{4.4}$$

By Remark 3.5 we can assume $\sigma - \sigma' = \text{CS}_K(\nabla, \nabla')$. Hence we get

$$\sigma \wedge f_*K(\nabla_f) = \sigma' \wedge f_*K(\nabla_f) + \text{CS}_K(\nabla, \nabla') \wedge f_*K(\nabla_f).$$

Since $f_*K(\nabla_f)$ is of degree n and $\text{CS}_K(\nabla, \nabla')$ is of degree -1 , switching factor on the right-hand side yields

$$\sigma \wedge f_*K(\nabla_f) = \sigma' \wedge f_*K(\nabla_f) + (-1)^n f_*K(\nabla_f) \wedge \text{CS}_K(\nabla, \nabla'). \tag{4.5}$$

The projection formula $f_*(T \wedge f^*\omega) = (f_*T) \wedge \omega$ applied to the current $T = K(\nabla_f)$ and the form $\omega = \text{CS}_K(\nabla, \nabla')$ implies

$$f_*K(\nabla_f) \wedge \text{CS}_K(\nabla, \nabla') = f_*(K(\nabla_f) \wedge f^*\text{CS}_K(\nabla, \nabla')). \tag{4.6}$$

The connections of $\tilde{g}^\varepsilon \circ \tilde{f}$ and $\tilde{g}^{\varepsilon'} \circ \tilde{f}$ are $\nabla_f \oplus f^*\nabla$ and $\nabla_f \oplus f^*\nabla'$, respectively. The Chern–Simons form for these two connections satisfies

$$\text{CS}_K(\nabla_f \oplus f^*\nabla, \nabla_f \oplus f^*\nabla') = K(\nabla_f) \wedge f^*\text{CS}_K(\nabla, \nabla').$$

Together with (4.6) this implies

$$f_*K(\nabla_f) \wedge \text{CS}_K(\nabla, \nabla') = f_*\text{CS}_K(\nabla_f \oplus f^*\nabla, \nabla_f \oplus f^*\nabla'). \tag{4.7}$$

Hence, identities (4.4), (4.5), and (4.7) together with $g_* \circ f_* = (g \circ f)_*$ on currents show that

$$\begin{aligned} &g_*(\mathcal{K}(\varepsilon) \wedge h + \sigma \wedge f_*K(\nabla_f)) \\ &= g_*((\mathcal{K}(\varepsilon') \wedge h + \sigma' \wedge f_*K(\nabla_f)) \\ &\quad + (-1)^{n+2d}(g \circ f)_*\text{CS}_K(\nabla_f \oplus f^*\nabla, \nabla_f \oplus f^*\nabla')). \end{aligned} \tag{4.8}$$

Since $\nabla_f \oplus f^*\nabla$ and $\nabla_f \oplus f^*\nabla'$ are the connections of $\tilde{g}^\varepsilon \circ \tilde{f}$ and $\tilde{g}^{\varepsilon'} \circ \tilde{f}$, respectively, Lemma 2.12 and (4.8) imply that the difference of the cycles $g_*^\varepsilon(\tilde{f}, h)$ and $g_*^{\varepsilon'}(\tilde{f}, h)$ lies in $BMU_\delta^{n+2d}(p+d)(Y)$. This proves the assertion of the lemma. ■

From now on we will use the canonical isomorphism $\tau: MU^n(p)(X) \rightarrow MU_\delta^n(p)(X)$ of Theorem 2.16 to identify $MU^n(p)(X)$ with $MU_\delta^n(p)(X)$. Putting the previous results together we have shown the following result.

Theorem 4.8. Let $g^{\circ}: X \rightarrow Y$ be a proper $MU_{\mathcal{D}}$ -oriented holomorphic map with $\mathfrak{o} = [\varepsilon] \in K_{MU_{\mathcal{D}}}(X)$. The assignment

$$[\tilde{f}, h] \mapsto [g_*^{\varepsilon}(\tilde{f}, h)]$$

induces a well-defined homomorphism

$$g_*^{\circ}: MU^n(p)(X) \rightarrow MU^{n+2d}(p+d)(Y).$$

We refer to g_*° as the pushforward along g° .

Remark 4.9. Following Remark 3.2 we could consider an orientation \mathfrak{o}_q of filtration q for $q \in \mathbb{Z}$. Then we would get a pushforward homomorphism

$$g_*^{\mathfrak{o}_q}: MU^n(p)(X) \rightarrow MU^{n+2d}(p+q+d)(Y)$$

with an additional shift by q . Since we are mainly interested in the orientation of Definition 5.9 which is of filtration 0 in this terminology, we decided to skip the additional level of generality. We note, however, that all the computations in this section could be modified accordingly.

The following result shows how the pushforward of Theorem 4.8 relates to the pushforwards of complex cobordism and sheaf cohomology.

Proposition 4.10. Let $g^{\circ}: X \rightarrow Y$ be a proper $MU_{\mathcal{D}}$ -oriented holomorphic map of complex codimension d , with $\mathfrak{o} = [N_g, \nabla_g, \sigma_g] \in K_{MU_{\mathcal{D}}}(X)$. Recall that we write $\mathcal{K}(\mathfrak{o}) = K(\nabla_g) - d\sigma_g$. Then the following diagrams commute:

$$\begin{array}{ccccc}
 H^{n-1}(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*)) & \xrightarrow{a} & MU^n(p)(X) & \xrightarrow{I} & MU^n(X) \\
 \downarrow g_*(\mathcal{K}(\mathfrak{o}) \wedge -) & & \downarrow g_*^{\circ} & & \downarrow g_* \\
 H^{n-1+2d}(Y; \frac{\mathcal{D}^*}{F^{p+d}}(\mathcal{V}_*)) & \xrightarrow{a} & MU^{n+2d}(p+d)(Y) & \xrightarrow{I} & MU^{n+2d}(Y)
 \end{array} \tag{4.9}$$

$$\begin{array}{ccc}
 MU^n(p)(X) & \xrightarrow{R} & H^n(X; F^p \mathcal{D}^*(X; \mathcal{V}_*)) \\
 \downarrow g_*^{\circ} & & \downarrow g_*(\mathcal{K}(\mathfrak{o}) \wedge -) \\
 MU^{n+2d}(p+d)(Y) & \xrightarrow{R} & H^{n+2d}(X; F^{p+d} \mathcal{D}^*(Y; \mathcal{V}_*)).
 \end{array} \tag{4.10}$$

Proof. For $[h] \in H^{n-1}(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*))$ we have

$$\begin{aligned}
 g_*^{\circ}(a[h]) &= [0, g_*(\mathcal{K}(\mathfrak{o}) \wedge h)] \\
 &= a(g_*(\mathcal{K}(\mathfrak{o}) \wedge h))
 \end{aligned}$$

which proves that the left-hand square in (4.9) commutes. That the right-hand square in (4.9) commutes follows from the observation that the underlying complex-oriented

map of a composition of geometric cycles, is the composition of the underlying complex-oriented maps. Hence we have

$$g_*(I[\tilde{f}, h]) = g_*[f, N_f] = [g \circ f, N_f \oplus f^* N_g] = I(g_*^{\circ}[f, h]).$$

Finally, by Lemma 4.2 we have

$$R(g_*^{\circ}(\gamma)) = g_*(\mathcal{K}(\circ) \wedge R(\gamma))$$

which shows that square (4.10) commutes as well. ■

We will now show that the pushforward is functorial.

Theorem 4.11. *Let the composition of proper holomorphic maps*

$$g_2 \circ g_1: X_1 \xrightarrow{g_1} X_2 \xrightarrow{g_2} X_3$$

be endowed with the composed $MU_{\mathcal{D}}$ -orientation $\circ_1 + g_1^ \circ_2$. Then we have*

$$(g_{2*}^{\circ_2}) \circ (g_{1*}^{\circ_1}) = (g_2 \circ g_1)_*^{\circ_1 + g_1^* \circ_2}$$

as homomorphisms

$$MU^n(p)(X_1) \rightarrow MU^{n+2d_1+2d_2}(p + d_1 + d_2)(X_3).$$

Proof. Let $\gamma = (\tilde{f}, h) \in ZMU_{\mathcal{D}}^n(p)(X_1)$. For $i = 1, 2$, let $\varepsilon_i = (N_i, \nabla_i, \sigma_i)$ represent \circ_i . We use the representative of $\circ_1 + g_1^* \circ_2$ suggested by Lemma 3.3,

$$\varepsilon_{12} = (N_1 \oplus g_1^* N_2, \nabla_1 \oplus g_1^* \nabla_2, \sigma_{12})$$

with

$$\sigma_{12} := \sigma_1 \wedge g_1^* \mathcal{K}(\circ_2) + g_1^* \sigma_2 \wedge K(\nabla_1). \tag{4.11}$$

Observe that the underlying geometric cycle of $g_{12}^{\varepsilon_{12}}$, which we denote by \tilde{g}_{12} , is the composed geometric cycle

$$\tilde{g}_{12} = \tilde{g}_2 \circ \tilde{g}_1.$$

Therefore, we know that the underlying geometric cycles of

$$(g_{12}^{\varepsilon_{12}})_* \gamma = (\tilde{g}_{12} \circ \tilde{f}, h_{12})$$

and

$$(g_2^{\varepsilon_2})_* \circ (g_1^{\varepsilon_1})_* \gamma = (\tilde{g}_2 \circ \tilde{g}_1 \circ \tilde{f}, h_{\circ})$$

coincide. It remains to show that $h_{12} = h_{\circ}$ modulo $\text{Im}(d)$. By definition of the pushforward and Remark 4.5 we have

$$h_{12} = (g_2 \circ g_1)_* [\mathcal{K}(\circ_1 + g_1^* \circ_2) \wedge h + \sigma_{12} \wedge f_* K(\nabla_f)].$$

On the other hand, applying Remark 4.5 to h_\circ yields

$$h_\circ = g_{2*}[\mathcal{K}(\nu_2) \wedge g_{1*}(\mathcal{K}(\nu_1) \wedge h + \sigma_1 \wedge f_*K(\nabla_f)) + \sigma_2 \wedge (g_1 \circ f)_*K(\nabla_{g_1 \circ f})].$$

Now we use the projection formula $f_*(T \wedge f^*\omega) = (f_*T) \wedge \omega$ for a current T and a form ω . Since we have $K(\nabla_{g_1 \circ f}) = K(\nabla_f) \wedge f^*K(\nabla_1)$, the projection formula for $T = K(\nabla_f)$ and $\omega = K(\nabla_1)$ implies

$$f_*K(\nabla_{g_1 \circ f}) = f_*(K(\nabla_f) \wedge f^*K(\nabla_1)) = f_*K(\nabla_f) \wedge K(\nabla_1).$$

Hence we can rewrite h_\circ as

$$h_\circ = g_{2*}[\mathcal{K}(\nu_2) \wedge g_{1*}(\mathcal{K}(\nu_1) \wedge h + \sigma_1 \wedge f_*K(\nabla_f)) + \sigma_2 \wedge g_{1*}(f_*K(\nabla_f) \wedge K(\nabla_1))].$$

We apply again the projection formula to the pushforward along g_1 , once with $T = \mathcal{K}(\nu_1) \wedge h + \sigma_1 \wedge f_*K(\nabla_f)$ and $\omega = \mathcal{K}(\nu_2)$, and once with $T = f_*K(\nabla_f) \wedge K(\nabla_1)$ and $\omega = \sigma_2$. Since $g_1^*\mathcal{K}(\nu_2)$ and $K(\nabla_1)$ lie in $\mathcal{A}^0(X_1; \mathcal{V}_*)$, and hence in the center of the ring $\mathcal{A}^*(X_1; \mathcal{V}_*)$, we then get

$$h_\circ = (g_2 \circ g_1)_*[g_1^*\mathcal{K}(\nu_2) \wedge (\mathcal{K}(\nu_1) \wedge h + \sigma_1 \wedge f_*K(\nabla_f)) + g_1^*\sigma_2 \wedge K(\nabla_1) \wedge f_*K(\nabla_f)].$$

Next we collect the terms that are wedged with $f_*K(\nabla_f)$ and obtain:

$$h_\circ = (g_2 \circ g_1)_*[g_1^*\mathcal{K}(\nu_2) \wedge \mathcal{K}(\nu_1) \wedge h + (g_1^*\mathcal{K}(\nu_2) \wedge \sigma_1 + g_1^*\sigma_2 \wedge K(\nabla_1)) \wedge f_*K(\nabla_f)].$$

By Proposition 3.8 we have $\mathcal{K}(\nu_1 + g_1^*\nu_2) = \mathcal{K}(\nu_1) \wedge \mathcal{K}(g_1^*\nu_2)$ which implies

$$h_\circ = (g_2 \circ g_1)_*[\mathcal{K}(\nu_1 + g_1^*\nu_2) \wedge h + (g_1^*\mathcal{K}(\nu_2) \wedge \sigma_1 + g_1^*\sigma_2 \wedge K(\nabla_1)) \wedge f_*K(\nabla_f)].$$

Finally, by formula (4.11) for σ_{12} , we get

$$h_\circ = (g_2 \circ g_1)_*[\mathcal{K}(\nu_1 + g_1^*\nu_2) \wedge h + \sigma_{12} \wedge f_*K(\nabla_f)].$$

This shows $h_\circ = h_{12}$ and finishes the proof. ■

Remark 4.12. Let $g: X \rightarrow Y$ and $q: W \rightarrow Y$ be transverse proper holomorphic maps of codimensions d and d' , respectively. Let $\pi: W \times_Y X \rightarrow Y$ be the map induced by the following cartesian diagram in $\mathbf{Man}_{\mathbb{C}}$

$$\begin{CD} W \times_Y X @>q'>> X \\ @Vg'VV @VVgV \\ W @>q>> Y. \end{CD}$$

(Note: A dashed arrow labeled π also points from $W \times_Y X$ to Y .)

Let $\nu_g \in K_{MU_{\mathcal{D}}}(X)$ and $\nu_q \in K_{MU_{\mathcal{D}}}(W)$ be $MU_{\mathcal{D}}$ -orientations of g and q , respectively. We then have natural isomorphisms of stable normal bundles $(g')^*N_q = N_{q'}$, $(q')^*N_g = N_{g'}$, and

$$N_\pi = (g')^*N_q \oplus N_{g'} = (q')^*N_g \oplus N_{q'}.$$

Hence $(g')^* \circ_q + (q')^* \circ_g$ is an orientation of π , and Theorem 4.11 implies that we have the following identity

$$g_*^{\circ_g} \circ (q')_*^{(g')^* \circ_q} = \pi_*^{(g')^* \circ_q + (q')^* \circ_g} = q_*^{\circ_q} \circ (g')_*^{(q')^* \circ_g}$$

of homomorphisms $MU^n(p)(W \times_Y X) \rightarrow MU^{n+2d+2d'}(p+d+d')(Y)$.

Next we will show that the pushforward is compatible with pullbacks. First we briefly recall the construction of pullback homomorphisms in $MU^*(p)(-)$ from [14, Theorem 2.22]. For further details we refer to [14, Section 2.7] and the references therein. Let $k: Y' \rightarrow Y$ be a holomorphic map. We consider the following cartesian diagram of manifolds

$$\begin{array}{ccc} Z' & \xrightarrow{k_Z} & Z \\ k^* f \downarrow & & \downarrow f \\ Y' & \xrightarrow{k} & Y \end{array}$$

where k and f are transverse, and $\tilde{f} = (f, N_f, \nabla_f)$ is a geometric cycle on Y . By transversality we get that $k^* f$ is complex-oriented with $N_{k_Z} = k_Z^* N_f$. We define $k^* \tilde{f}$ by

$$k^* \tilde{f} = (k^* f, k_Z^* N_f, k_Z^* \nabla_f).$$

For a cycle $(\tilde{f}, h) \in ZMU^n(p)(Y)$, it remains to define the pullback of the current h . Since the pullback of an arbitrary current is not defined, this requires to restrict to the subgroup $ZMU_k^n(p)(Y) \subset ZMU^n(p)(Y)$ consisting of those $\gamma = (\tilde{f}, h)$ satisfying

$$WF(h) \cap N(k) = \emptyset \quad \text{and} \quad k \pitchfork f$$

where $WF(h)$ denotes the wave-front set of h and $N(k)$ is the normal set of f as defined in [17, Section 8.1]. For $\gamma = (\tilde{f}, h) \in ZMU_k^n(p)(Y)$, we then have a well-defined pullback

$$k^* \gamma = k^*(\tilde{f}, h) = (k^* f, k_Z^* N_f, k_Z^* \nabla_f, k^* h)$$

where $k^* h$ is well-defined by [17, Theorem 8.2.4]. By [14, Theorem 2.25], this induces a pullback homomorphism

$$k^*: MU^n(p)(Y) \rightarrow MU^n(p)(Y').$$

Theorem 4.13. *Suppose we have a cartesian diagram in $\mathbf{Man}_{\mathbb{C}}$*

$$\begin{array}{ccc} X' & \xrightarrow{k'} & X \\ g' \downarrow & & \downarrow g \\ Y' & \xrightarrow{k} & Y \end{array} \tag{4.12}$$

with k transverse to g , and g proper of codimension d . Let \circ be an $MU_{\mathcal{D}}$ -orientation of g . We equip g' with the pullback orientation $\circ' := k'^*\circ$. Then we have

$$k^*g_*^{\circ} = (g')_*^{\circ'}k'^*: MU^n(p)(X) \rightarrow MU^{n+2d}(p+d)(Y').$$

Proof. Let $\gamma = (\tilde{f}, h) = (f, N_f, \nabla_f, h) \in ZMU^n(p)(X)$ be a cycle. Since transversality is generic we can assume f to be transverse with k' . Let $k'_Z: Z' \rightarrow Z$ be the induced map in the top cartesian rectangle in

$$\begin{array}{ccc} Z' & \xrightarrow{k'_Z=k_Z} & Z \\ k'^*f \downarrow & & \downarrow f \\ X' & \xrightarrow{k'} & X \\ g' \downarrow & & \downarrow g \\ Y' & \xrightarrow{k} & Y. \end{array}$$

Since both rectangles are cartesian, the outer rectangle is cartesian as well. Hence the map $k_Z: Z' \rightarrow Z$ induced by the outer cartesian diagram agrees with k'_Z . We write

$$k^*g_*^{\circ}(\tilde{f}, h) =: (f_{\lrcorner}, h_{\lrcorner}) \quad \text{and} \quad (g')_*^{\circ'}k'^*(\tilde{f}, h) =: (f^{\lrcorner}, h^{\lrcorner}).$$

Let $\varepsilon = (N_g, \nabla_g, \sigma_g)$ be a representative of the orientation \circ of g . Then we have

$$f_{\lrcorner} = (k^*(g \circ f), k_Z^*N_f \oplus k_Z^*f^*N_g, k_Z^*\nabla_f \oplus k_Z^*f^*\nabla_g).$$

The pullback orientation $k'^*\circ$ is represented by $\varepsilon' = (k'^*N_g, k'^*\nabla_g, k'^*\sigma_g)$. Since diagram (4.12), is cartesian we have

$$f_{\lrcorner} = ((g' \circ k'^*f), N_{k'^*f} \oplus (k'^*f)^*N_g, \nabla_{k'^*f} \oplus (k'^*f)^*\nabla_g) = f^{\lrcorner}.$$

Now we check the effect on the current h using that we have $k^*g_* = g'_*k'^*$ by [14, Theorem 2.27] whenever the involved maps are defined:

$$\begin{aligned} h_{\lrcorner} &= k^*(g_*(K(\nabla_g) \wedge h + \sigma_g \wedge (f_*K(\nabla_f) - dh))) \\ &= g'_*(k'^*K(\nabla_g) \wedge k'^*h + k'^*\sigma_g \wedge k'^*(f_*K(\nabla_f) - dh)) \\ &= h^{\lrcorner}. \end{aligned}$$

We recall from [14, Section 2.8] that there is a natural product of the form

$$MU^{n_1}(p_1)(X) \times MU^{n_2}(p_2)(X) \rightarrow MU^{n_1+n_2}(p_1+p_2)(X) \tag{4.13}$$

turning $MU^*(*)(X)$ into a ring. The product of two classes $[\gamma_1]$ and $[\gamma_2]$ is denoted by $[\gamma_1] \cdot [\gamma_2]$ and is induced by the following construction: We consider the operation

$$\otimes: \mathcal{D}^{n_1}(X_1; \mathcal{V}_*) \times \mathcal{D}^{n_2}(X_2; \mathcal{V}_*) \rightarrow \mathcal{D}^{n_1+n_2}(X_1 \times X_2; \mathcal{V}_*) \tag{4.14}$$

satisfying $T_1 \otimes T_2 = \pi_1^* T_1 \wedge \pi_2^* T_2$. Since K is multiplicative, we have

$$K^{p_1+p_2}(\nabla_1 \oplus \nabla_2) = K^{p_1}(\nabla_1) \otimes K^{p_2}(\nabla_2).$$

We then define the symbol \times , and refer to it as the external product of Hodge filtered cycles by potential slight abuse of terminology, by

$$\gamma_1 \times \gamma_2 := (\tilde{f}_1 \times \tilde{f}_2, h_1 \otimes R(\gamma_2) + (-1)^{n_1} f_{1*} K(\nabla_{f_1}) \otimes h_2). \tag{4.15}$$

The product in (4.13) is then defined as the pullback along the diagonal map $\Delta_X: X \rightarrow X \times X$:

$$[\gamma_1] \cdot [\gamma_2] = \Delta_X^*([\gamma_1 \times \gamma_2]).$$

The following theorem shows that g_* is a homomorphism of $MU^*(*)(Y)$ -modules.

Theorem 4.14. *Let $g: X \rightarrow Y$ be a proper holomorphic map of codimension d and let \circ be an $MU_{\mathcal{D}}$ -orientation of g . Then, for all integers n, p, m, q , and all elements $x \in MU^n(p)(X)$ and $y \in MU^m(q)(Y)$, we have the following projection formula*

$$g_*^{\circ}(g^* y \cdot x) = y \cdot g_*^{\circ} x \text{ in } MU^{n+m+2d}(p+q+d)(Y).$$

Proof. Since the product is defined by pulling back an exterior product along the diagonal, we consider the following commutative diagram

$$\begin{array}{ccccc} X & \xrightarrow{(g, \text{id}_X)} & Y \times X & \xrightarrow{\pi_X} & X \\ g \downarrow & & G = \text{id}_Y \times g \downarrow & & \downarrow g \\ Y & \xrightarrow{\Delta_Y} & Y \times Y & \xrightarrow{\text{pr}_2} & Y. \end{array}$$

We denote by $\pi_Y: Y \times X \rightarrow Y$ and $\pi_X: Y \times X \rightarrow X$, and by $\text{pr}_1: Y \times Y \rightarrow Y$ and $\text{pr}_2: Y \times Y \rightarrow Y$ the projections onto the first and second factors, respectively. We endow the map $G := \text{id}_Y \times g$ with the pullback $MU_{\mathcal{D}}$ -orientation $\circ' := \pi_X^* \circ$. We claim that in order to prove the assertion of the theorem it suffices to show the identity

$$G_*^{\circ'}(y \times x) = y \times g_*^{\circ}(x). \tag{4.16}$$

To prove that it suffices to show (4.16), we observe that (4.16) implies that

$$\Delta_Y^* G_*^{\circ'}(y \times x) = \Delta_Y^*(y \times g_*^{\circ}(x)) = y \cdot g_*^{\circ} x$$

by definition of the cup product on $MU^*(*)(Y)$. Hence it remains to show that

$$\Delta_Y^* G_*^{\circ'}(y \times x) = g_*^{\circ}(g^* y \cdot x).$$

To do so we consider the following diagram

$$\begin{array}{ccccc} X & \xrightarrow{\Delta_X} & X \times X & \xrightarrow{g \times \text{id}_X} & Y \times X \\ g \downarrow & & g \times g \downarrow & & \downarrow G \\ Y & \xrightarrow{\Delta_Y} & Y \times Y & \xrightarrow{\text{id}_Y \times Y} & Y \times Y. \end{array} \tag{4.17}$$

Since the outer diagram in (4.17) is cartesian and since G and Δ_Y are transverse, we can apply Theorem 4.13 to get

$$\begin{aligned} \Delta_Y^* G_*^{o'}(y \times x) &= g_*^o((g \times \text{id}_X) \circ \Delta_X)^*(y \times x) \\ &= g_*^o \Delta_X^*((g \times \text{id}_X)^*(y \times x)) \\ &= g_*^o \Delta_X^*((g^* y \times x)) \\ &= g_*^o(g^* y \cdot x), \end{aligned}$$

where the last equality uses the definition of the cup product on $MU^*(*)(X)$. This proves the claim.

We will now show that identity (4.16) holds by proving the corresponding formula on the level of cycles. Let $\varepsilon = (N_g, \nabla_g, \sigma_g)$ be a representative of \circ . Then the triple $\varepsilon' = (\pi_X^* N_g, \pi_X^* \nabla_g, \pi_X^* \sigma_g)$ represents $\circ' = \pi_X^* \circ$. Let (\tilde{f}_x, h_x) and (\tilde{f}_y, h_y) be cycles such that $x = [\tilde{f}_x, h_x]$ and $y = [\tilde{f}_y, h_y]$. We write $h_{y \times x}$ for the current defined by (4.15) such that $y \times x = [\tilde{f}_y \times \tilde{f}_x, h_{y \times x}]$. The theorem will then follow once we have proven the identity of cycles

$$G_*^{\varepsilon'}(\tilde{f}_y \times \tilde{f}_x, h_{y \times x}) = (\tilde{f}_y, h_y) \times g_*^\varepsilon(\tilde{f}_x, h_x). \tag{4.18}$$

Formula (4.18) can be checked separately on the level of geometric cycles and on the level of currents. To simplify the notation we denote the cycle $G_*^{\varepsilon'}(\tilde{f}_y \times \tilde{f}_x, h_{y \times x})$ by (\tilde{f}_G, h_G) . We write $(\tilde{f}_{g_*(x)}, h_{g_*(x)})$ for the cycle $g_*^\varepsilon(\tilde{f}_x, h_x)$, and $(\tilde{f}_{y \times g_*(x)}, h_{y \times g_*(x)})$ for the cycle $(\tilde{f}_y, h_y) \times g_*^\varepsilon(\tilde{f}_x, h_x)$. For the geometric cycles the formula $\tilde{f}_G = \tilde{f}_{y \times g_*(x)}$ follows directly from the definition of the pushforward and the definition of the map $G = \text{id}_Y \times g$.

Now we show that (4.18) holds for the corresponding currents. Recall that we use the notation $\phi(\gamma) = (f_\gamma)_* K(\nabla_{f_\gamma})$ and $R(\gamma) = \phi(\gamma) - dh_\gamma$ for a cycle $\gamma = (f_\gamma, h_\gamma)$. We then have by definition of the exterior product \times

$$h_{y \times x} = h_y \otimes R(x) + (-1)^m \phi(y) \otimes h_x. \tag{4.19}$$

By definition of the pushforward we have

$$h_G = G_*(\pi_X^* K(\nabla_g) \wedge h_{y \times x} + \pi_X^* \sigma_g \wedge R(y \times x)).$$

Using formula (4.19) and the formula $R(y \times x) = R(y) \otimes R(x)$, which is verified in [14, p. 26], we can rewrite h_G as

$$h_G = G_*(\pi_X^* K(\nabla_g) \wedge (h_y \otimes R(x) + (-1)^m \phi(y) \otimes h_x) + \pi_X^* \sigma_g \wedge (R(y) \otimes R(x))).$$

By definition of \otimes in (4.14) and the fact that $R(y)$ is of degree m we then get

$$\begin{aligned} h_G &= G_*(h_y \otimes (K(\nabla_g) \wedge R(x)) + (-1)^m \phi(y) \otimes (K(\nabla_g) \wedge h_x) \\ &\quad + (-1)^m R(y) \otimes (\sigma_g \wedge R(x))). \end{aligned}$$

Now we use the definition of G as $G = \text{id}_Y \times g$ to get:

$$h_G = h_Y \otimes g_*(K(\nabla_g) \wedge R(x)) + (-1)^m \phi(y) \otimes g_*(K(\nabla_g) \wedge h_x) + (-1)^m R(y) \otimes g_*(\sigma_g \wedge R(x)).$$

On the other hand we compute

$$\begin{aligned} h_{Y \times g_*^\varepsilon(x)} &= h_Y \otimes R(g_*^\varepsilon(\tilde{f}_x, h_x)) + (-1)^m \phi(y) \otimes h_{g_*^\varepsilon(x)} \\ &= h_Y \otimes g_*((K(\nabla_g) - d\sigma_g) \wedge R(x)) + (-1)^m \phi(y) \otimes g_*(K(\nabla_g) \wedge h_x + \sigma_g \wedge R(x)) \\ &= h_Y \otimes g_*(K(\nabla_g) \wedge R(x)) - h_Y \otimes g_*(d\sigma_g \wedge R(x)) \\ &\quad + (-1)^m \phi(y) \otimes g_*(K(\nabla_g) \wedge h_x) + (-1)^m \phi(y) \otimes g_*(\sigma_g \wedge R(x)). \end{aligned}$$

Comparing the expressions for h_G and $h_{Y \times g_*^\varepsilon(x)}$ it remains to show

$$(-1)^m R(y) \otimes g_*(\sigma_g \wedge R(x)) + h_Y \otimes g_*(d\sigma_g \wedge R(x)) = (-1)^m \phi(y) \otimes g_*(\sigma_g \wedge R(x))$$

modulo $\text{Im}(d)$. Since, by definition of R in (2.4), $R(x)$ is a closed form, we have

$$d(\sigma_g \wedge R(x)) = d\sigma_g \wedge R(x) - \sigma_g \wedge dR(x) = d\sigma_g \wedge R(x).$$

Since h_Y is of degree m , we therefore get

$$d(h_Y \otimes g_*(\sigma_g \wedge R(x))) = dh_Y \otimes g_*(d\sigma_g \wedge R(x)) + (-1)^m h_Y \otimes g_*(d\sigma_g \wedge R(x)).$$

Hence, modulo image of d , we get the following identity

$$h_Y \otimes g_*(d\sigma_g \wedge R(x)) = (-1)^m dh_Y \otimes g_*(d\sigma_g \wedge R(x)) \quad \text{modulo } \text{Im}(d).$$

Since $R(y) = \phi(y) - dh_Y$ by definition, we can thus conclude

$$\begin{aligned} (-1)^m R(y) \otimes g_*(\sigma_g \wedge R(x)) + h_Y \otimes g_*(d\sigma_g \wedge R(x)) &= (-1)^m R(y) \otimes g_*(\sigma_g \wedge R(x)) + (-1)^m dh_Y \otimes g_*(\sigma_g \wedge R(x)) \\ &= (-1)^m \phi(y) \otimes g_*(\sigma_g \wedge R(x)) \end{aligned}$$

modulo $\text{Im}(d)$. This shows (4.18) and finishes the proof. ■

We end this section with a further observation on the relationship of the maps R and \mathcal{K} .

Remark 4.15. As in the proof of Proposition 4.10, we can express the identity shown in Lemma 4.2 as

$$R(g_*^\circ(\gamma)) = g_*(\mathcal{K}(\circ) \wedge R(\gamma))$$

for every element $[\gamma]$ and proper holomorphic map $g: X \rightarrow Y$ with $MU_{\mathcal{D}}$ -orientation \circ . For the special case that γ is the identity element 1_X of the ring $MU^*(\circ)(X)$, i.e., for $[\gamma] = 1_X = [\text{id}_X, d, 0]$, we get

$$R(g_*^\circ(1_X)) = g_*(\mathcal{K}(\circ)).$$

5. A canonical Hodge filtered MU -orientation for holomorphic maps

We will now show that for every holomorphic map there is a natural choice for an $MU_{\mathcal{D}}$ -orientation. The key result is Theorem 5.6 which provides us with a canonical choice of a class of connections. The existence of a canonical choice of a class of orientation and Theorem 5.12 may be seen as justification for defining $MU_{\mathcal{D}}$ -orientations as a K -group and not just as a set. We recall from [19, Section 6.3] the following terminology.

Definition 5.1. Let X be a complex manifold and let D be a smooth connection on a holomorphic vector bundle E over X . Then with respect to local coordinates (U_i, g_i) , D acts as $d + \Gamma_i$, where $\Gamma_i = (\Gamma_i^{jk})$ is a matrix of 1-forms. Recall that we have

$$\Gamma_i = g_{ji}^{-1} dg_{ji} + g_{ji}^{-1} \Gamma_j g_{ji},$$

where the g_{ij} denote the transition functions. Conversely any such cocycle $\{\Gamma_i\}$ defines a connection. Then D is called a *Bott connection* if for each i, j, k we have

$$\Gamma_i^{jk} \in F^1 \mathcal{A}^1(X).$$

Remark 5.2. As noted in the introduction, Bott connections are more commonly referred to as connections compatible with the holomorphic structure. Here we follow Karoubi, who uses the terminology in [19] in a context where Bott connections generalize both connections compatible with a holomorphic structure and Bott connections of foliation theory. Since Bott connections are frequently used in what follows, we adopt *Bott connection* as a convenient terminology.

Remark 5.3. Every holomorphic vector bundle on a complex manifold admits a Bott connection. In fact, the Chern connection on a holomorphic bundle with a hermitian metric is defined as the unique Bott connection which is compatible with the hermitian structure. By [18, Proposition 4.1.4] every complex vector bundle admits a hermitian metric. By [18, Proposition 4.2.14] every holomorphic bundle with a hermitian structure has a Chern connection. Alternatively, one can show the existence of Bott connections as in [19, Section 6] using a local trivialization of the bundle and a partition of unity.²

Remark 5.4. If D is a Bott connection, then the curvature of D , which in local coordinates is represented by the matrix

$$d\Gamma_i + \Gamma_i \wedge \Gamma_i,$$

belongs to $F^1 \mathcal{A}^2(X; \text{End}(E))$. This implies the following key fact about Bott connections:

$$K(D) \in F^0 \mathcal{A}^0(X; \mathcal{V}_*). \tag{5.1}$$

²We emphasize again that a Bott connection does not have to be holomorphic, but is merely required to be smooth. Hence one may use a partition of unity for the construction as explained in [19, Section 6].

Remark 5.5. Let D be a Bott connection on E . Then (5.1) implies that the triple $(E, D, 0)$ defines an element in $K_{MU_D}(X)$. The following result, inspired by [19, Theorem 6.7], shows that the associated orientation class $[E, D, 0]$ is independent of the choice of Bott connection D .

We will now prove the key technical result of this section.

Theorem 5.6. *For every $X \in \mathbf{Man}_{\mathbb{C}}$, there is a natural homomorphism*

$$B: K_{\text{hol}}^0(X) \rightarrow K_{MU_D}(X)$$

induced by

$$B[E] = [E, D, 0],$$

for each holomorphic vector bundle E where D is any Bott connection on E .

Proof. The existence of a Bott connection was pointed out in Remark 5.3. The assertion of the theorem then follows from the following two lemmas. ■

As a first step we analyze the Chern–Simons form of two Bott connections on a given holomorphic vector bundle and show that they lead to the same orientation class.

Lemma 5.7. *Let D and D' be two Bott connections for a holomorphic vector bundle $E \rightarrow X$. Then $\text{CS}_K(D, D') \in F^0\mathcal{A}^0(X; \mathcal{V}_*)$, so $[E, D, 0] = [E, D', 0]$ in $K_{MU_D}(X)$.*

Proof. Let Γ_i and Γ'_i be the connection matrices of D and D' , respectively, with respect to local holomorphic coordinates z_1, \dots, z_l on U_i . Let $I = [0, 1]$ be the unit interval and let $\pi: I \times X \rightarrow X$ denote the projection. Then consider the connection $D'' = t \cdot \pi^* D + (1 - t) \cdot \pi^* D'$ on $\pi^* E$. Its connection matrix on $I \times U_i$ is

$$\Gamma'' = t\Gamma_i + (1 - t)\Gamma'_i.$$

The curvature of D'' is given on $I \times U_i$ by

$$\begin{aligned} \Omega''_i &= d\Gamma''_i + \Gamma''_i \wedge \Gamma''_i \\ &= dt \wedge \Gamma_i + t \cdot d\Gamma_i - dt \wedge \Gamma'_i + (1 - t)d\Gamma'_i + t^2\Gamma_i \wedge \Gamma_i \\ &\quad + (1 - t)^2\Gamma'_i \wedge \Gamma'_i + t(1 - t)\Gamma_i \wedge \Gamma'_i. \end{aligned}$$

Each term is of filtration 1 in the sense that at least one of the dz_j s appears in each term of each entry. Hence the Chern form $c_k(D'')$ has at least k many dz_j s appearing in its local expression, and in that sense belongs to $F^k\mathcal{A}^*([0, 1] \times X)$. Integrating out dt maps this filtration step $F^k\mathcal{A}^*([0, 1] \times X)$ to the Hodge filtration $F^k\mathcal{A}^*(X)$. This implies

$$\pi_* K(D'') \in F^0\mathcal{A}^{-1}(X; \mathcal{V}_*).$$

Since $\text{CS}_K(D, D') = \pi_* K(D'')$, this proves

$$[E, D, 0] = [E, D', 0] \in K_{MU_D}(X). \quad \blacksquare$$

Next, we show that all the defining relations of $K_{\text{hol}}^0(X)$ and $K_{MU_{\mathcal{D}}}(X)$ are respected by B .

Lemma 5.8. *Let E be a holomorphic bundle over X and D be a Bott connection on E . The assignment $E \mapsto (E, D, 0)$ induces a map $B: K_{\text{hol}}^0(X) \rightarrow K_{MU_{\mathcal{D}}}(X)$.*

Proof. The first part of this proof follows [19, Proof of Theorem 6.7]. Let

$$0 \rightarrow E_1 \xrightarrow{\alpha} E_2 \xrightarrow{\beta} E_3 \rightarrow 0$$

be a short exact sequence of holomorphic vector bundles, and let D_i be a Bott connection on E_i . By the defining relations for $K_{MU_{\mathcal{D}}}(X)$ we need to establish

$$CS_K(D_1, D_2, D_3) \in \tilde{F}^0 \mathcal{A}^{-1}(X; \mathcal{V}_*)$$

where we recall that the notation \tilde{F} has been introduced in (2.2). Let $\gamma: E_3 \rightarrow E_2$ be a smooth splitting. This yields a smooth isomorphism of bundles

$$u = (\alpha, \gamma): E_1 \oplus E_3 \rightarrow E_2.$$

The inverse u^{-1} has the form

$$u^{-1} = \begin{pmatrix} \sigma \\ \beta \end{pmatrix},$$

where σ is a left-inverse of α . We choose holomorphic coordinates for each E_j over an open $U_i \subset X$. We then get the following equations of matrix valued forms:

$$u_i \cdot u_i^{-1} = \begin{pmatrix} \alpha_i & \gamma_i \end{pmatrix} \cdot \begin{pmatrix} \sigma_i \\ \beta_i \end{pmatrix} = 1$$

and

$$\begin{pmatrix} \sigma_i \\ \beta_i \end{pmatrix} \begin{pmatrix} \alpha_i & \gamma_i \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Since D_2 is a Bott connection, it is represented by a matrix Γ_i^2 with coefficients in F^1 . Note that, since γ and σ may not be holomorphic, $\Delta^2 := u^* D_2$ may not be a Bott connection. However, locally on U_i , Δ^2 takes the form

$$\Delta_i^2 = u_i^{-1} du_i + u_i^{-1} \Gamma_i^2 u_i.$$

We have $u_i^{-1} \Gamma_i^2 u_i \in F^1$, since Γ_i^2 is in F^1 . The matrix $u_i^{-1} du_i$ expands as

$$\begin{pmatrix} \sigma_i \\ \beta_i \end{pmatrix} \begin{pmatrix} d\alpha_i & d\gamma_i \end{pmatrix} = \begin{pmatrix} \sigma_i d\alpha_i & \sigma_i d\gamma_i \\ \beta_i d\alpha_i & \beta_i d\gamma_i \end{pmatrix}.$$

Since $\beta_i \gamma_i = 1$, we have $\beta_i d\gamma_i = -d\beta_i \gamma_i \in F^1$. Since $d\alpha_i \in F^1$, we see that $u_i^{-1} du_i$ is upper triangular modulo F^1 .

Now let $\pi: [0, 1] \times X \rightarrow X$ be the projection. Let

$$\nabla = t \cdot \pi^* \Delta^2 + (1 - t) \cdot \pi^*(D_1 \oplus D_3),$$

and let θ_i be the connection matrix of ∇ with respect to local coordinates on $U_i \subset X$. We continue to use the notion of filtration on $\mathcal{A}^*(\pi^*(E_1 \oplus E_3))$, $\mathcal{A}^*(\pi^*(E_2))$, and $\mathcal{A}^*([0, 1] \times X; \mathcal{V}_*)$ as in the proof of Lemma 5.7. We know that $(1 - t) \cdot \pi^*(D_1 \oplus D_3)$ is in F^1 , and we have just shown that $t \cdot \pi^* \Delta^2$ is upper triangular modulo F^1 . Thus, θ_i is upper triangular modulo F^1 as well. Hence the local curvature form of ∇ , i.e., $\Omega_i = d\theta_i + \theta_i \wedge \theta_i$, is upper triangular modulo F^1 as well. This implies that $c_i(\nabla) \in F^i \mathcal{A}^{2i}([0, 1] \times X)$ and hence $K(\nabla) \in F^0 \mathcal{A}^0([0, 1] \times X; \mathcal{V}_*)$. Now we note that we defined the Chern–Simons form $CS_K(D_1, D_2, D_3)$ as the integral of $K(\nabla')$, and not $K(\nabla)$, for the connection ∇' on $[0, 1] \times \pi^* E_2$ given by

$$\nabla' = (u^{-1})^* \nabla = t \cdot \pi^* D_2 + (1 - t) \cdot \pi^*((u^{-1})^*(D_1 \oplus D_3)).$$

Locally we can express the curvature of ∇' as

$$\Omega'_i = u_i^{-1} \Omega_i u_i.$$

Thus ∇ and ∇' have identical Chern–Weil forms. In particular, this implies that $K(\nabla') \in F^0 \mathcal{A}^0([0, 1] \times X; \mathcal{V}_*)$. Thus, again since integrating out dt sends $F^0 \mathcal{A}^0$ to $F^0 \mathcal{A}^{-1}$, we have shown

$$\pi_* K(\nabla') = CS_K(D_1, D_2, D_3) \in \tilde{F}^0 \mathcal{A}^{-1}(X; \mathcal{V}_*)$$

which finishes the proof of the lemma and of Theorem 5.6. ■

A key application of Theorem 5.6 is that it allows us to make a canonical choice of a Hodge filtered MU -orientation for each holomorphic map:

Definition 5.9. Let $g: X \rightarrow Y$ be a holomorphic map, and let

$$\mathcal{N}_g := [g^*TY] - [TX] \in K_{\text{hol}}^0(X)$$

denote the virtual holomorphic normal bundle of g . We define the *Bott $MU_{\mathcal{D}}$ -orientation of g* , or *Bott orientation of g* for short, to be $B(\mathcal{N}_g) \in K_{MU_{\mathcal{D}}}(X)$, i.e., the image of \mathcal{N}_g under $B: K_{\text{hol}}^0(X) \rightarrow K_{MU_{\mathcal{D}}}(X)$.

The next lemma shows that the Bott orientation is functorial, i.e., it is compatible with pullbacks in the following way.

Lemma 5.10. Assume we have a pullback diagram in $\mathbf{Man}_{\mathbb{C}}$

$$\begin{array}{ccc} X' & \xrightarrow{f'} & X \\ g' \downarrow & & \downarrow g \\ Y' & \xrightarrow{f} & Y \end{array} \tag{5.2}$$

with f transverse to g . Let \mathcal{N}_g and $\mathcal{N}_{g'}$ be the virtual holomorphic normal bundles of g and g' , respectively. Then we have

$$f'^* B(\mathcal{N}_g) = B(\mathcal{N}_{g'}) \text{ in } K_{MU_{\mathcal{D}}}(X').$$

Proof. Since f is transverse to g , we have $f'^* \mathcal{N}_g = \mathcal{N}_{g'}$ in $K_{\text{hol}}^0(X')$. Since the choice of Bott connection does not matter for B by Theorem 5.6, this induces the desired identity $f'^* B(\mathcal{N}_g) = B(\mathcal{N}_{g'})$ in $K_{MU_{\mathcal{D}}}(X')$. ■

Remark 5.11. Let $X_1 \xrightarrow{g_1} X_2 \xrightarrow{g_2} X_3$ be proper holomorphic maps. Since the map

$$K_{\text{hol}}^0(X) \rightarrow K_{MU_{\mathcal{D}}}(X)$$

is a homomorphism of groups, we have

$$B(\mathcal{N}_{g_2 \circ g_1}) = B(\mathcal{N}_{g_1} \oplus g_1^* \mathcal{N}_{g_2}) = B(\mathcal{N}_{g_1}) + g_1^* B(\mathcal{N}_{g_2}).$$

Hence the Bott orientation of $g_2 \circ g_1$ is the composed $MU_{\mathcal{D}}$ -orientation of the Bott orientations of g_1 and g_2 , respectively. Together with Lemma 5.10 this may justify to call the Bott orientation a *canonical* Hodge filtered MU -orientation for a holomorphic map.

Applying Theorems 4.8, 4.11, 4.13, and 4.14 with the Bott orientation together with Remark 5.11 yields the following result.

Theorem 5.12. Let X and Y be complex manifolds, and let $g: X \rightarrow Y$ be a proper holomorphic map of codimension d . We equip g with its Bott orientation $\circ := B(\mathcal{N}_g)$. Then $g_* := g_*^{\circ}$ defines a functorial pushforward map

$$g_*: MU^n(p)(X) \rightarrow MU^{n+2d}(p+d)(Y).$$

This is a homomorphism of $MU^*(*) (Y)$ -modules in the sense that, for all integers n, p, m, q , and all elements $x \in MU^n(p)(X)$ and $y \in MU^m(q)(Y)$, we have

$$g_*(g^* y \cdot x) = y \cdot g_* x \text{ in } MU^{n+m+2d}(p+q+d)(Y).$$

Furthermore, if $f: Y' \rightarrow Y$ is holomorphic and transversal to g , letting f' and g' denote the induced maps as in (5.2), the following formula holds

$$f'^* \circ g_* = g'_* \circ f'^*.$$

In the remainder of this section we further reflect on the Bott orientation class $B(\mathcal{N}_g)$. We note that $[\mathcal{N}_g] = [g^*TY] - [TX]$ merely is a virtual bundle and, in general, there may not be a holomorphic bundle N_g over X which represents $[g^*TY] - [TX]$ in $K_{\text{hol}}^0(X)$. We can, however, obtain a representative of the orientation class $B(\mathcal{N}_g)$ in $K_{MU_{\mathcal{D}}}(X)$ as follows: Let $g: X \rightarrow Y$ be a holomorphic map and $i: X \rightarrow \mathbb{C}^k$ a smooth embedding. We then get a short exact sequence of complex vector bundles of the form

$$0 \rightarrow TX \xrightarrow{D(g,i)} g^*TY \oplus \underline{\mathbb{C}}_X^k \rightarrow N_{(g,i)} \rightarrow 0. \tag{5.3}$$

Proposition 5.13. *Let X be a Stein manifold, Y any complex manifold and $g: X \rightarrow Y$ a holomorphic map. Then we can represent the virtual normal bundle of g , i.e., the element $[g^*TY] - [TX] \in K^0(X)$, by a holomorphic vector bundle on X .*

Proof. Since X is Stein, we can assume i in (5.3) to be holomorphic. Hence $N_{(g,i)}$ admits a holomorphic structure. ■

For general X , however, we cannot expect $N_{(g,i)}$ to be holomorphic. In particular, $N_{(g,i)}$ does not, in general, represent the difference $[g^*TY] - [TX]$ in $K_{\text{hol}}^0(X)$. Yet we have the following result which follows from the defining relations in $K_{MU_{\mathcal{D}}}(X)$ (see also Remark 3.11).

Proposition 5.14. *With the above notation, let D_X be a Bott connection for TX , and D_Y a Bott connection for TY . Let $\nabla_{(g,i)}$ be a connection on $N_{(g,i)}$. We set*

$$\circ_g := [N_{(g,i)}, \nabla_{(g,i)}, -\text{CS}_K(D_X, g^*D_Y \oplus d, \nabla_{(g,i)})] \in K_{MU_{\mathcal{D}}}(X).$$

Then we have

$$B(\mathcal{N}_g) = \circ_g \quad \text{in } K_{MU_{\mathcal{D}}}(X). \quad \blacksquare$$

For a projective complex manifold we can represent the canonical $MU_{\mathcal{D}}$ -orientation in the following way.

Proposition 5.15. *Let $g: X \rightarrow Y$ be a proper holomorphic map. Assume that X is a projective complex manifold. Then there is a holomorphic vector bundle N on X and a Bott connection D on N such that $(N, D, 0)$ is an $MU_{\mathcal{D}}$ -orientation of g and $B(\mathcal{N}_g) = [N, D, 0]$ in $K_{MU_{\mathcal{D}}}(X)$.*

Proof. Recall the Euler sequence

$$0 \rightarrow \underline{\mathbb{C}} \rightarrow \gamma_1^{\oplus(n+1)} \rightarrow T\mathbb{C}\mathbb{P}^n \rightarrow 0$$

where $\gamma_1 \rightarrow \mathbb{C}\mathbb{P}^n$ is the tautological line bundle. There is a canonical inclusion $\gamma_1 \rightarrow \underline{\mathbb{C}}^{n+1}$, and we denote the quotient by γ_1^\perp . Hence $-\gamma_1 = [\gamma_1^\perp] - [\underline{\mathbb{C}}^{n+1}]$. Thus we obtain the identity

$$-[T\mathbb{C}\mathbb{P}^n] = (n + 1) \cdot ([\gamma_1^\perp] - [\underline{\mathbb{C}}^{n+1}]) + [\underline{\mathbb{C}}] = (n + 1) \cdot [\gamma_1^\perp] - [\underline{\mathbb{C}}^{n^2+2n}]$$

in $K_{\text{hol}}^0(\mathbb{C}\mathbb{P}^n)$. Now let X be a projective manifold and let $\iota: X \hookrightarrow \mathbb{C}\mathbb{P}^n$ denote a holomorphic embedding. We have a short exact sequence of holomorphic vector bundles over X

$$0 \rightarrow TX \rightarrow \iota^*T\mathbb{C}\mathbb{P}^n \rightarrow NX \rightarrow 0.$$

In $K_{\text{hol}}^0(X)$ this implies the identities

$$-[TX] = [NX] - \iota^*[T\mathbb{C}\mathbb{P}^n] = [NX] + (n + 1)\iota^*[\gamma_1^\perp] - [\underline{\mathbb{C}}_X^{n^2+2n}]$$

and hence

$$\mathcal{N}_g = [g^*TY] - [TX] = [g^*TY] + [NX] + (n + 1)\iota^*[\gamma_1^\perp] - [\mathbb{C}_X^{n^2+2n}].$$

We define the holomorphic bundle

$$N := g^*TY \oplus NX \oplus \iota^*(\gamma_1^\perp)^{\oplus(n+1)}.$$

Since $B(\mathbb{C}^{n^2+2n}) = 0$, we then get the identity $B(\mathcal{N}_g) = B(N)$ in $K_{MU,d}(X)$. Thus we have $B(\mathcal{N}_g) = [N, D, 0]$ for any Bott connection D on N . ■

6. Fundamental classes and secondary cobordism invariants

The existence of pushforwards along proper holomorphic maps allows us to define special types of Hodge filtered cobordism classes. In particular, we can define fundamental classes as follows:

Definition 6.1. Let $f: Y \rightarrow X$ be a proper holomorphic map of codimension d . Let $1_Y \in MU^0(0)(Y)$ be the identity element of the graded commutative ring $MU^*(*)(Y)$. We endow f with its Bott orientation. We then refer to the element $[f] := f_*(1_Y) \in MU^{2d}(d)(X)$ as the *fundamental class of f* . If the context of f and X is clear, we may also write $[Y]$ for $[f]$ and call it the *fundamental class of Y* .

Let $f: Y \rightarrow X$ be a proper holomorphic map of codimension d . Let $i: Y \rightarrow \mathbb{C}^k$ be a smooth embedding. We then get a short exact sequence of the form

$$0 \rightarrow TY \rightarrow f^*TX \oplus \mathbb{C}_Y^k \rightarrow N_{(f,i)} \rightarrow 0.$$

With this notation, we have the following result.

Proposition 6.2. *The fundamental class $[f]$ of f in $MU^{2p}(p)(X)$ is given by*

$$f_*[1_Y] = [\tilde{f}, f_*\sigma_{(f,i)}] = [f, N_{(f,i)}, \nabla_{(f,i)}, f_*\sigma_{(f,i)}],$$

where $\nabla_{(f,i)}$ is any connection on $N_{(f,i)}$ and $\sigma_{(f,i)} = -CS_K(D_Y, f^*D_X \oplus d, \nabla_{(f,i)})$ for Bott connections D_X on TX and D_Y on TY .

Proof. This follows directly from the description of the Bott orientation in Proposition 5.14 and the definition of the pushforward map using $1_Y = [id_Y, d, 0]$. ■

Next we show that the fundamental class is compatible with products in the following sense.

Lemma 6.3. *Let $f: Y \rightarrow X$ and $g: Z \rightarrow X$ be proper holomorphic maps of codimension d and d' , respectively. Let π denote the map induced by the following cartesian diagram*

in $\mathbf{Man}_{\mathbb{C}}$

$$\begin{array}{ccc}
 Y \times_X Z & \xrightarrow{f'} & Z \\
 g' \downarrow & \searrow \pi & \downarrow g \\
 Y & \xrightarrow{f} & X.
 \end{array}$$

Assume that f and g are transverse. Then we have

$$[f] \cdot [g] = [\pi] \quad \text{in } MU^{2d+2d'}(d+d')(X).$$

Proof. Since f and g are transverse, we can apply Theorem 4.13 to get

$$f^* g_* = g'_* f'^*.$$

Since $\pi = g' \circ f$ by definition, Theorem 4.11 implies

$$f_* f^* g_* = f_* g'_* f'^* = \pi_* f'^*.$$

We apply this to $1_Z \in MU^0(0)(Z)$ and use that $f'^*(1_Z) = 1_{Y \times_X Z}$ to get

$$[\pi] = \pi_* f'^*(1_Z) = f_* f^* g_*(1_Z) = f_* f^*[g].$$

Now we apply Theorem 4.14 to $y = [g]$ and $x = 1_Y$ to conclude

$$[\pi] = \pi_*(1_{Y \times_X Z}) = f_* f^*[g] = [g] \cdot f_*(1_Y) = [g] \cdot [f].$$

Finally, we note that the product in the subring consisting of elements of even cohomological degrees $MU^{2*}(\mathbb{C})(X)$ is commutative to conclude the proof. ■

Remark 6.4. If $f: Y \hookrightarrow X$ is the embedding of a complex *submanifold* of codimension d , then the normal bundle N_f is a holomorphic bundle. Hence, in this case, the Bott orientation of f is given by $B(N_f) = (N_f, D_f, 0)$ with a Bott connection D_f on N_f , and we have $[f] = [f, N_f, D_f, 0]$ in $MU^{2d}(d)(X)$.

Remark 6.5. Let $f_0: Y_0 \rightarrow X$ and $f_1: Y_1 \rightarrow X$ be two embeddings of complex submanifolds of codimension d . By Remark 6.4, we can write the associated fundamental classes as $[f_0] = [f_0, N_{f_0}, D_{f_0}, 0]$ and $[f_1] = [f_1, N_{f_1}, D_{f_1}, 0]$. Now assume that f_0 and f_1 are cobordant, i.e., they represent the same element in $MU^{2d}(X)$. Then we can find a geometric bordism \tilde{b} with $\partial \tilde{b} = \tilde{f}_1 - \tilde{f}_0$. The bordism \tilde{b} is, in general, not sufficient to show $[f_0] = [f_1]$ in $MU^{2d}(d)(X)$, since the associated current $\psi(\tilde{b})$ defined in (2.1) may not vanish. In fact, \tilde{b} defines a Hodge filtered bordism datum between f_0 and f_1 if and only if

$$\psi(\tilde{b}) \in \tilde{F}^d \mathcal{D}^{2d-1}(X; \mathcal{V}_*) = F^d \mathcal{D}^{2d-1}(X; \mathcal{V}_*) + d \mathcal{D}^{2d-2}(X; \mathcal{V}_*).$$

In particular, two homotopic maps f_0 and f_1 do not define the same class in Hodge filtered cobordism in general (see also Lemma 2.12 and [14, Lemma 5.9]). This shows that the current $\psi(\tilde{b})$ contains information that is not detected by $MU^{2d}(X)$.

Following Remark 6.5 we will now study the case of a *topologically* cobordant fundamental class in more detail. For the rest of this section we assume that X is a *compact Kähler* manifold. Then we can split the long exact sequence of Proposition 2.15 into a short exact sequence as follows. Let $\text{Hdg}_{\mathbb{S}MU}^{2p}(X) = I(MU^{2p}(p)(X))$. We write

$$J_{MU}^{2p-1}(X) = \frac{H^{2p-1}(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*))}{\phi(MU^{2p-1}(X))}.$$

Then we get a short exact sequence

$$0 \rightarrow J_{MU}^{2p-1}(X) \rightarrow MU^{2p}(p)(X) \rightarrow \text{Hdg}_{\mathbb{S}MU}^{2p}(X) \rightarrow 0. \tag{6.1}$$

Remark 6.6. Note that, since X is compact Kähler, we have an isomorphism

$$H^{2p-1}\left(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*)\right) \cong \frac{H^{2p-1}(X; \mathcal{V}_*)}{F^p H^{2p-1}(X; \mathcal{V}_*)}.$$

Thus we can rewrite $J_{MU}^{2p-1}(X)$ as

$$J_{MU}^{2p-1}(X) = \frac{H^{2p-1}(X; \mathcal{V}_*)}{F^p H^{2p-1}(X; \mathcal{V}_*) + \phi(MU^{2p-1}(X))}.$$

Remark 6.7. As noted in [16, Remark 4.12], it follows from the Hodge decomposition that $J_{MU}^{2p-1}(X)$ is isomorphic to the group $MU^{2p-1}(X) \otimes \mathbb{R}/\mathbb{Z}$. This implies that, as a *real* Lie group, $J_{MU}^{2p-1}(X)$ is a homotopy invariant of X , while as a *complex* Lie group $J_{MU}^{2p-1}(X)$ depends on the complex structure of X .

Definition 6.8. Assume we have an element $[\gamma]$ in $MU^{2p}(p)(X)$ such that $I([\gamma])$ vanishes in $MU^{2p}(X)$. Then sequence (6.1) shows that we may use $J_{MU}^{2p-1}(X)$ as the target for *secondary cobordism invariants*. For example, let $f: Y \rightarrow X$ be a proper holomorphic map of codimension p . Assume that the fundamental class of f in $MU^{2p}(X)$, given as the pushforward of $1_Y \in MU^0(Y)$ along f , vanishes. Then the fundamental class of f in $MU^{2p}(p)(X)$ has image in the subgroup $J_{MU}^{2p-1}(X)$. Because of the similarity to the Abel–Jacobi map of Deligne–Griffiths (see e.g. [24, Section 12]) we will denote the image of f in the subgroup $J_{MU}^{2p-1}(X)$ by $\text{AJ}(f)$ and will refer to $\text{AJ}(f)$ as the *Abel–Jacobi invariant* of f .

Let $f: Y \rightarrow X$ be a proper holomorphic map of codimension p . We will now describe $\text{AJ}(f)$ in more detail. Let $[\gamma_f] := [f, N_{(f,i)}, \nabla_{(f,i)}, f_*\sigma_{(f,i)}]$ be as in Proposition 6.2. We assume that $f_*(1_Y) = 0$ in $MU^{2p}(X)$. Then there is a topological bordism datum $b: W \rightarrow \mathbb{R} \times X$ such that $\partial b = f$. Let N_b be the associated normal bundle. We can extend the connection $\nabla_{(f,i)}$ on $N_{(f,i)}$ to get a connection ∇_b on N_b , and obtain a geometric cobordism datum \tilde{b} . Then we have

$$\gamma_f - (\partial\tilde{b}, \psi(\tilde{b})) = (0, f_*\sigma_{(f,i)} - \psi(\tilde{b})) = (0, f_*\sigma_{(f,i)} - (\pi_X \circ b|_{W_{[0,1]}})_*(K^p(\nabla_b)))$$

by definition of $\psi(\tilde{b})$ in (2.1). Hence we get

$$[\gamma_f] = a[f_*\sigma_{(f,i)} - \psi(\tilde{b})]$$

under the homomorphism

$$a: H^{2p-1}\left(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*)\right) \rightarrow MU^{2p-1}(p)(X)$$

which is induced by the map defined on the level of forms in (2.3). The class $[f_*\sigma_f - \psi(\tilde{b})]$ in $H^{2p-1}(X; \frac{\mathcal{D}^*}{F^p}(\mathcal{V}_*))$ may depend on the choice of \tilde{b} . However, if \tilde{b}' is a different bordism datum, then we have

$$a[\psi(\tilde{b}) - \psi(\tilde{b}')] \in \phi(MU^{2p-1}(X)) \subset H^{2p-1}(X; \mathcal{V}_*).$$

Thus, after taking the quotient, we get a well-defined class. We summarise these observations in the following theorem.

Theorem 6.9. *With the above assumptions on f and X , the fundamental class of f in $MU^{2p}(p)(X)$ is the image of*

$$AJ(f) = [f_*\sigma_f - \psi(\tilde{b})] \in \frac{H^{2p-1}(X; \mathcal{V}_*)}{F^p H^{2p-1}(X; \mathcal{V}_*) + \phi(MU^{2p-1}(X))} = J_{MU}^{2p-1}(X). \quad \blacksquare$$

Now we give an alternative description of $AJ(f)$. Let \mathcal{V}'_* be the \mathbb{C} -dual graded algebra with homogeneous components

$$\mathcal{V}'_j = (\mathcal{V}_{-j})' = \text{Hom}_{\mathbb{C}}(\mathcal{V}_{-j}, \mathbb{C}).$$

Then the canonical pairing given by evaluation

$$\text{ev}: \mathcal{V}'_* \otimes \mathcal{V}_* \rightarrow \mathbb{C}$$

has degree 0, if \mathbb{C} is interpreted as a graded vector space concentrated in degree 0. Let $n = \dim_{\mathbb{C}} X$. Poincaré duality and the fact that all vector spaces involved are finite-dimensional imply that the pairing

$$H^k(X; \mathcal{V}_*) \times H^{2n-k}(X; \mathcal{V}'_*) \rightarrow \mathbb{C} \tag{6.2}$$

$$([\eta], [\omega]) = \text{ev}\left(\int_X \eta \wedge \omega\right)$$

is perfect. Here $\eta \wedge \omega$ is interpreted as a $\mathcal{V}_* \otimes \mathcal{V}'_*$ -valued form. We may thus identify $H^{2p-1}(X; \mathcal{V}_*)$ with $(H^{2n-2p+1}(X; \mathcal{V}'_*))'$. Hodge symmetry and Serre duality then imply that, under this identification, the subspace $F^p H^{2p-1}(X; \mathcal{V}_*)$ corresponds to the subspace $(F^{n-p+1} H^{2n-2p+1}(X; \mathcal{V}'_*))^\perp$. This implies that there is a natural isomorphism

$$\frac{H^{2p-1}(X; \mathcal{V}_*)}{F^p H^{2p-1}(X; \mathcal{V}_*)} \cong (F^{n-p+1} H^{2n-2p+1}(X; \mathcal{V}'_*))'.$$

Now we let ϕ' denote the composition of $\phi: MU^k(X) \rightarrow H^k(X; \mathcal{V}_*)$ followed by the identification under pairing (6.2), i.e., ϕ' maps the element $[f: Z \rightarrow X] \in MU^k(X)$ to $\phi'(f)$ in $(H^{2n-k}(X; \mathcal{V}'_*))'$ defined by

$$\phi'(f)([\omega]) := \text{ev} \left(\int_Z K(\nabla_f) \wedge f^* \omega \right),$$

where ∇_f is a connection on the normal bundle N_f . We note that, since Y is closed, it follows from Stokes' theorem that this pairing is independent of the choice of representative of $[\omega]$ and of the choice of connection. In fact, it is independent of the choice of form which represents the class $K(N_f)$. Then we conclude from the above arguments that there is a natural isomorphism

$$J_{MU}^{2p-1}(X) \cong \frac{(F^{n-p+1} H^{2n-2p+1}(X; \mathcal{V}'_*))'}{\phi'(MU^{2p-1}(X))}. \tag{6.3}$$

Now let $f: Y \rightarrow X$ be a proper holomorphic map of codimension p such that $[f] = 0$ in $MU^{2p}(X)$. Let $\tilde{b} = (b, N_b, \nabla_b)$ be a geometric bordism datum over

$$b = (c_b, f_b): W \rightarrow \mathbb{R} \times X.$$

We set $W_{[0,1]} := c_b^{-1}([0, 1])$, and $w := f_b|_{W_{[0,1]}}$.

Theorem 6.10. *With the above notation, the image of $\text{AJ}(f)$ under isomorphism (6.3) is represented by the functional in $(F^{n-p+1} H^{2n-2p+1}(X; \mathcal{V}'_*))'$ defined by*

$$[\omega] \mapsto \text{ev} \left(\int_Y \sigma_f \wedge f^* \omega + \int_{W_{[0,1]}} K(\nabla_b) \wedge w^* \omega \right).$$

Proof. We recall from Theorem 6.9 that $\text{AJ}(f) = [f_* \sigma_f - \psi(\tilde{b})] \in J_{MU}^{2p-1}(X)$. Let ω be a closed form in $F^{n-p+1} \mathcal{A}^{2n-2p+1}(X; \mathcal{V}'_*)$. Since the codimension of f_b is odd, we have $\psi(\tilde{b}) = -w_* K(\nabla_b)$. Then the interaction of pushforwards and pullbacks with integrals and Stokes' theorem yield:

$$\begin{aligned} \int_X (f_* \sigma_f - \psi(\tilde{b})) \wedge \omega &= \int_X f_* \sigma_f \wedge \omega - \int_X \psi(\tilde{b}) \wedge \omega \\ &= \int_Y \sigma_f \wedge f^* \omega + \int_{W_{[0,1]}} K(\nabla_b) \wedge w^* \omega. \end{aligned}$$

By construction of isomorphism (6.3), the image of $\text{AJ}(f)$ is the homomorphism that sends $[\omega]$ to the class given by evaluating the above sum of integrals.

It remains to show that this evaluation yields a well-defined element in the group $(F^{n-p+1} H^{2n-2p+1}(X; \mathcal{V}'_*))'$. Assume $\omega = d\psi$. Then

$$\int_{W_{[0,1]}} K(\nabla_b) \wedge w^*(d\psi) = \int_Y K(\nabla_b)|_Y \wedge f^* \psi \tag{6.4}$$

by Stokes' theorem. Since f is holomorphic, we have $K(N_b)|_Y = K(N_f) \in H^{0,0}(Y; \mathcal{V}_*)$ and thus $K(\nabla_b)|_Y \in F^0 \mathcal{A}^0(Y; \mathcal{V}_*)$. Since Hodge theory implies the vanishing of the subspace $F^{n-p+1} H^{2n-2p}(Y; \mathcal{V}_* \otimes \mathcal{V}'_*) = 0$, integral (6.4) vanishes.

For the other integral we note that by Stokes' theorem we have

$$\int_Y \sigma_f \wedge f^*(d\psi) = \int_Y d\sigma_f \wedge f^*\psi. \tag{6.5}$$

We recall from Proposition 6.2 that $\sigma_f = -CS_K(D_Y, f^*D_X \oplus d, \nabla_f)$ for Bott connections D_X on TX and D_Y on TY , and an arbitrary connection ∇_f on the normal bundle. The derivative of σ_f satisfies

$$d\sigma_f = K(f^*D_X \oplus d) - K(D_Y) - K(\nabla_f).$$

Since K is multiplicative and $K(d) = 1$, we have $K(f^*D_X \oplus d) = K(f^*D_X)$. Since D_X and D_Y are Bott connections, we know that $K(f^*D_X)$ and $K(D_Y)$ are in $F^0 \mathcal{A}^0(Y; \mathcal{V}_*)$. This implies again for reasons of type that the integrals

$$\int_Y K(f^*D_X) \wedge f^*\psi \quad \text{and} \quad \int_Y K(D_Y) \wedge f^*\psi$$

both vanish. The remaining term to analyse is the integral $\int_Y K(\nabla_f) \wedge f^*\psi$ which we already have shown to vanish. Thus integral (6.5) vanishes and the functional is well-defined. Finally, we note that integral (6.5) is independent of the chosen bordism datum, while the difference between the integrals (6.4) corresponding to two different bordism data is an element in $\phi'(MU^{2p-1}(X))$. ■

Remark 6.11. The formula in Theorem 6.10 simplifies if the orientation \mathfrak{o}_f admits a representative of the form $(N, \nabla, 0)$. If f is projective, we obtain such a representative from Proposition 5.15, and if f is a holomorphic embedding, $B(f^*TX/TY)$ will do. We do not know if such representatives exist for the canonical orientations of general holomorphic maps.

7. Hodge filtered Thom morphism

We will now define a Thom morphism from Hodge filtered cobordism to Deligne cohomology. In order to define a map on the level of cycles we will first construct a new cycle model for Deligne cohomology. Our construction is similar to that of Gillet–Soulé in [11] (see also [13]). However, our construction is more elementary than the one in [11] in the sense that it avoids the use of geometric measure theory.

Let X be a complex manifold and $U \subseteq X$ an open subset. For an integer $p \geq 0$, let $\mathbb{Z}(p)$ denote $(2\pi i)^p \cdot \mathbb{Z}$ and let $\mathbb{Z}_{\mathcal{D}}(p)$ be the complex of sheaves

$$0 \rightarrow \mathbb{Z}(p) \rightarrow \mathcal{O}_X \rightarrow \Omega_X^1 \rightarrow \cdots \rightarrow \Omega_X^{p-1} \rightarrow 0,$$

where $\mathbb{Z}(p)$ is placed in degree 0. Then the Deligne cohomology group $H_{\mathbb{D}}^q(X; \mathbb{Z}(p))$ may be defined as the q -th hypercohomology of the complex $\mathbb{Z}_{\mathbb{D}}(p)$. We recall the group of smooth relative chains defined as the quotient

$$C_{\dim X - k}^{\text{diff}}(X, X \setminus \bar{U}; \mathbb{Z}(p)) = \frac{C_{\dim X - k}^{\text{diff}}(X; \mathbb{Z}(p))}{C_{\dim X - k}^{\text{diff}}(X \setminus \bar{U}; \mathbb{Z}(p))}.$$

Let \bar{C}^k denote the presheaf

$$U \mapsto \bar{C}^k(U) := C_{\dim X - k}^{\text{diff}}(X, X \setminus \bar{U}; \mathbb{Z}(p)).$$

The restriction maps of \bar{C}^k are induced by quotienting out the appropriate additional chains. The presheaf \bar{C}^k is very close to being a sheaf since it satisfies the sheaf condition for coverings of X . However, it does not satisfy the sheaf condition for general collections of open subsets of X . Hence let C^k be the sheafification of \bar{C}^k . The sheaf C^k is not fine, but it is homotopically fine, meaning that its endomorphism sheaf admits a homotopy partition of unity. We refer to [4, p. 172], from which we also recall the implication that $H^*(H^j(\bar{C}^*(U))) = 0$ for $j > 0$. Hence the hypercohomology spectral sequence degenerates on the E_2 -page, past which only the row $H^0(C^*(U))$ survives. On stalks the sheaf C^k coincides with the presheaf \bar{C}^k . Let U be a small contractible open subset of X . By excision we have

$$H^k(\bar{C}^*(U)) = H_{\dim X - k}(\mathbb{R}^{\dim X}, \mathbb{R}^{\dim X} \setminus \mathbb{D}; \mathbb{Z}(p)),$$

for \mathbb{D} the closed unit disc. Hence we get

$$H^j(\bar{C}^*(U)) = \begin{cases} 0 & j > 0, \\ \mathbb{Z}(p) & j = 0. \end{cases}$$

This proves the following result.

Lemma 7.1. *The complex C^* is an acyclic resolution of the constant sheaf $\mathbb{Z}(p)$ as sheaves on X . ■*

By [4, Appendix B, Solution I.12], we also have the following fact.

Lemma 7.2. *The canonical map $\bar{C}^* \rightarrow C^*$ induces an isomorphism of cohomology groups on global sections $H^k(\bar{C}^*(X)) = H^k(C^*(X))$. ■*

In other words, the sheaf cohomology $H^k(X; \mathbb{Z}(p))$ can be computed as the cohomology of the complex $\bar{C}^*(X)$. Now we consider the map of complexes

$$T: \bar{C}^*(X) \rightarrow \mathcal{D}^*(X)$$

induced by integration. Let $\mathcal{D}_{\mathbb{Z}}^*(X)$ be the image of T in $\mathcal{D}^*(X)$. Since T is a map of chain complexes, it follows that $\mathcal{D}_{\mathbb{Z}}^*(X)$ is a complex as well.

Proposition 7.3. *The map $T: \bar{C}^*(X) \rightarrow \mathcal{D}_{\mathbb{Z}}^*(X)$ induces an isomorphism on cohomology.*

Proof. By Whitehead’s triangulation theorem, we may pick a smooth triangulation of X , i.e., a set $S = \{f_i: \Delta^{k_i} \rightarrow X\}$ such that each f_i is a continuous embedding which extends to a smooth mapping of a neighborhood of $\Delta^k \subset \mathbb{R}^k$, and each $x \in X$ is in the interior of a unique cell $S_i = \text{Im}(f_i)$. It is well-known that the inclusion $C_*(S; \mathbb{Z}(p)) \rightarrow C_*(X; \mathbb{Z}(p))$ of cellular chains is a quasi-isomorphism. Hence it suffices to show that T restricts to a quasi-isomorphism on the cellular chains of S . Since each point $x \in X$ is contained in the interior of a unique cell of S , we can show that T is injective on cellular chains as follows. We can construct for each i a form $\omega_i \in \mathcal{A}^{k_i}(X)$ such that $\int_{\Delta^{k_i}} f_i^* \omega_i \neq 0$, and such that the only k_i -cell intersecting the support of ω_i is S_i . Suppose $T(c) = 0$ for $c = \sum a_i f_i$. Then $T(c)(\omega_i) = a_i T(f_i)(\omega_i)$ is a nonzero multiple of a_i , and we get $a_i = 0$ for all i . To see that the map induced by T from cellular homology is injective, we first note that the inclusion of cellular chains into singular chains is a deformation retract since it is a quasi-isomorphism between complexes of projective modules. Let r be a retraction onto the cellular chains. Now let c be a cellular cycle with $T(c) = dT(\alpha)$ for α an arbitrary integral chain $\alpha \in C^*(X)$. Then we have $T(c) = dT(\alpha) = T(\partial\alpha)$ and thus $T(c) = T(r(c)) = T(r(\partial\alpha)) = T(\partial r(\alpha))$. Since T is injective on cellular chains, we get $c = \partial r(\alpha)$. Hence c represents 0 in cellular homology, and the map induced by T on cellular homology is injective. It remains to see that T restricted to cellular chains is surjective on homology.

By definition of $\mathcal{D}_{\mathbb{Z}}^*(X)$ as the image of T , every element of $\mathcal{D}_{\mathbb{Z}}^*(X)$ is of the form $\sum_i T(a_i \cdot g_i)$ where g_i are smooth maps $\Delta^k \rightarrow X$. Assume that $\sum_i T(a_i \cdot g_i)$ is a cycle and hence represents a class in $H^k(X; \mathcal{D}_{\mathbb{Z}})$. To simplify the notation, we write

$$g := \sum_i a_i \cdot g_i.$$

By assumption, we have $dT(g) = 0$. Since r is a deformation retraction, there is a homotopy h of the cellular chains such that

$$\partial h + h\partial = 1 - r.$$

By applying r , we define a cellular chain $f := r(g)$. Omitting the inclusion from cellular chains into chains from the notation we then have the identity of chains

$$g' := f - h\partial(g) = \partial(h(g)) + g.$$

Applying r again defines a cellular chain $r(g')$ such that

$$dT(r(g')) = dT(r(\partial(h(g)) + g)) = T(\partial\partial(h(g))) + dT(g) = 0,$$

where we use the assumption $dT(g) = 0$. Hence we get

$$T(\partial(r(g'))) = dT(r(g')) = 0.$$

Since T is injective on cellular chains, this implies $\partial r(g') = 0$, i.e., that $f' := r(g')$ is a cellular cycle. Since $T(f') - T(g) = T(\partial(h(g))) = dT(h(g))$ is an exact current, we have found a cellular cycle f' whose homology class is mapped to the homology class of g under T . This completes the proof. ■

We are now ready to give our presentation of Deligne cohomology. Let

$$i_F: F^p \mathcal{A}^* \rightarrow \mathcal{D}^*$$

be the map of sheaves induced by T , and let $i_c: \mathcal{D}_{\mathbb{Z}}^*(X) \rightarrow \mathcal{D}^*(X)$ be the inclusion. We will show that the following cochain complex

$$C_{\mathcal{D}}^*(p)(X) = \text{cone}(\mathcal{D}_{\mathbb{Z}}^*(X) \oplus F^p \mathcal{A}^*(X) \xrightarrow{i_c - i_F} \mathcal{D}^*(X))$$

computes the Deligne cohomology of X . In degree k we have the group

$$C_{\mathcal{D}}^k(p)(X) = \mathcal{D}_{\mathbb{Z}}^k(X) \oplus F^p \mathcal{A}^k(X) \oplus \mathcal{D}^{k-1}(X).$$

The differential is defined by

$$d(T, \omega, h) = (dT, d\omega, i_c(T) - dh + i_F(\omega)).$$

Theorem 7.4. *The cohomology of the cochain complex $C_{\mathcal{D}}^*(p)(X)$ is naturally isomorphic to Deligne cohomology.*

To prove the theorem we will use multicomplexes, which are more flexible than bicomplexes. We recall from [3] that a multicomplex of abelian groups consists of the data of a bigraded abelian group, $E^{s,t}$, and differentials $d_r^{s,t}: E^{s,t} \rightarrow E^{s+r,t-r+1}$ such that

$$\sum_{i+j=k} d_j^{s+i,t-i+1} \circ d_i^{s,t} = 0: E^{s,t} \rightarrow E^{s+k,t-k+2}.$$

One can consider multicomplexes of objects in any abelian category. We are considering here multicomplexes of abelian sheaves.

Proof of Theorem 7.4. We will construct a series of quasi-isomorphisms of complexes of sheaves

$$\mathbb{Z}_{\mathcal{D}}(p) \simeq C_{\mathcal{D}}^*(p) \simeq \text{Tot}(M)$$

and a quasi-isomorphism of complexes of abelian groups $\text{Tot}(M)(X) \rightarrow C_{\mathcal{D}}^*(X)$, where M is the following multicomplex of sheaves on X :

$$M^{s,t} = \begin{cases} C^t & s = 0, \\ \mathcal{D}^{s-1,t} & 0 < s < p, \\ F^p \mathcal{A}^{s,t} \oplus \mathcal{D}^{s-1,t} & p \leq s. \end{cases}$$

To define the differentials let $\Pi^{s,k-s}: \mathcal{D}^k \rightarrow \mathcal{D}^{s,k-s}$ be the projection. For $s > 0$, there is only d_0 and d_1 . The differentials of M are

$$d_0^{s,t} = \begin{cases} d: C^t \rightarrow C^{t+1} & s = 0, \\ -\bar{\partial}: \mathcal{D}^{s-1,t} \rightarrow \mathcal{D}^{s-1,t+1} & 0 < s < p, \\ (\bar{\partial}, i_F - \bar{\partial}): F^p \mathcal{A}^{s,t} \oplus \mathcal{D}^{s-1,t} \rightarrow F^p \mathcal{A}^{s,t+1} \oplus \mathcal{D}^{s-1,t+1} & s \geq p, \end{cases}$$

$$d_1^{s,t} = \begin{cases} \Pi^{0,t} \circ i_c: C^t \rightarrow \mathcal{D}^{0,t} & s = 0, \\ -\partial: \mathcal{D}^{s-1,t} \rightarrow \mathcal{D}^{s,t} & 0 < s < p, \\ (\partial, i_F - \partial): F^p \mathcal{A}^{s,t} \oplus \mathcal{D}^{s-1,t} \rightarrow F^p \mathcal{A}^{s+1,t} \oplus \mathcal{D}^{s,t} & s \geq p, \end{cases}$$

$$d_r^{0,t} = \Pi^{r,t-r} \circ i_c: C^t \rightarrow \mathcal{D}^{r,t-r}.$$

The total complex of M is given by

$$\text{Tot}^*(M) = \text{cone} \left(C^* \oplus F^p \mathcal{A}^* \xrightarrow{i_F - i_c} \mathcal{D}^* \right).$$

There is therefore a natural map $\text{Tot}^*(M(X)) \rightarrow C_{\mathcal{D}}^*(p)(X)$ defined by

$$\text{Tot}^*(M(X)) \ni (c, \omega, h) \mapsto (aT(c), \omega, h) \in C_{\mathcal{D}}^*(p)(X),$$

where we write aT for the sheafified map induced by T . This map of complexes induces an isomorphism on cohomology since each of the maps

$$\text{id}: F^p \mathcal{A}^*(X) \rightarrow F^p \mathcal{A}^*(X), \quad T: \bar{C}^*(X) \rightarrow \mathcal{D}_{\mathbb{Z}}^*(X), \quad \text{and} \quad \text{id}: \mathcal{D}^*(X) \rightarrow \mathcal{D}^*(X)$$

is a quasi-isomorphism. We define yet another complex of sheaves

$$C_{\mathcal{D}}'^*(p) = (\mathbb{Z}(p) \rightarrow \Omega^0 \xrightarrow{d} \dots \xrightarrow{d} \Omega^{p-2} \xrightarrow{(0,d)} \Omega^p \oplus \Omega^{p-1} \xrightarrow{\delta_p} \Omega^{p+1} \oplus \Omega^p \xrightarrow{\delta_{p+1}} \dots)$$

with $\delta_i(\omega, \tau) = (d\omega, \omega - d\tau)$ for $i \geq p$. There is a map $f: \mathbb{Z}_{\mathcal{D}}(p)(X) \rightarrow C_{\mathcal{D}}'^*(p)(X)$ given by

$$\begin{array}{ccccccccccccccc} \mathbb{Z}(p) & \longrightarrow & \Omega^0 & \xrightarrow{d} & \Omega^1 & \xrightarrow{d} & \dots & \xrightarrow{d} & \Omega^{p-2} & \xrightarrow{d} & \Omega^{p-1} & \longrightarrow & 0 \\ \text{id} \downarrow & & \text{id} \downarrow & & \text{id} \downarrow & & & & \text{id} \downarrow & & \alpha \downarrow & & \\ \mathbb{Z}(p) & \longrightarrow & \Omega^0 & \xrightarrow{d} & \Omega^1 & \longrightarrow & \dots & \xrightarrow{d} & \Omega^{p-2} & \xrightarrow{(0,d)} & \Omega^p \oplus \Omega^{p-1} & \xrightarrow{\delta_p} & \dots \end{array}$$

with $\alpha(\omega) = (d\omega, \omega)$. We claim that this is a quasi-isomorphism of complexes of sheaves. This is clear in degrees $< p$, and in degrees $> p$ it follows from the fact that $C_{\mathcal{D}}'^*(p)$ is exact in that range. In degree p we need to show that f induces an isomorphism on cohomology of stalks. Let U be a polydisc. Then

$$H_{\mathcal{D}}^p(U; \mathbb{Z}(p)) = \frac{\Omega^{p-1}(U)}{\text{Im } d},$$

and

$$H^p(U; C_{\mathcal{D}}^*(p)) = \frac{\{(\omega, \tau) \in \Omega^p(U) \oplus \Omega^{p-1}(U) : d\tau = \omega\}}{\text{Im}(0, d)}.$$

It is clear that the map induced by f , which can be described as $[\tau] \mapsto [d\tau, \tau]$, is an isomorphism. Hence f is a quasi-isomorphism as claimed. Next there is a natural map $C_{\mathcal{D}}^*(p) \rightarrow M$ given by

$$\begin{array}{ccccccccccc} \mathbb{Z}(p) & \longrightarrow & \Omega^0 & \longrightarrow & \dots & \longrightarrow & \Omega^{p-2} & \longrightarrow & \Omega^p \oplus \Omega^{p-1} & \longrightarrow & \dots \\ \downarrow \varepsilon & & \downarrow & & & & \downarrow & & \downarrow & & \\ C^0 & \longrightarrow & \mathcal{D}^{0,0} & \longrightarrow & \dots & \longrightarrow & \mathcal{D}^{p-2,0} & \longrightarrow & \mathcal{A}^p \oplus \mathcal{D}^{p-1} & \longrightarrow & \dots \end{array}$$

where ε is the quasi-isomorphism $\mathbb{Z}(p) \rightarrow C^*$. The column $M^{i,*}$ is a resolution of the sheaf $C_{\mathcal{D}}^i(p)$ by Lemma 7.1 and the arguments in [12, pp. 382–385]. Hence the natural map

$$C_{\mathcal{D}}^*(p) \rightarrow M$$

is a quasi-isomorphism. This concludes the proof. ■

Remark 7.5. If we choose a smooth triangulation of Z , then by summing up the top cells we get a smooth singular cycle c_Z representing the fundamental class $[Z] \in H_{\dim Z}(Z; \mathbb{Z})$. We have $T(c_Z) = 1 \in \mathcal{D}^0(Z)$, and so

$$f_* 1 = f_* T(c_Z) = T(f_* c_Z) \in \mathcal{D}_{\mathbb{Z}}^*(X).$$

The advantage of using \mathcal{D}_Z is that no choice of triangulation is needed in order to get the current $f_* 1$.

Let τ_0 be the map

$$\mathcal{D}^*(X; \mathcal{V}_*) \rightarrow \mathcal{D}^*(X; \mathbb{C})$$

induced by the map on coefficients

$$\mathcal{V}_* = MU_* \otimes \mathbb{C} \rightarrow \mathbb{C}$$

determined by the additive formal group law over \mathbb{C} . Then τ_0 is a chain map and it preserves the Hodge filtration. Now we are ready to define our Hodge filtered Thom morphism on the level of cycles:

$$\begin{aligned} \tau_Z: ZMU^n(p)(X) &\rightarrow C_{\mathcal{D}}^n(p)(X), \\ \gamma = (\tilde{f}, h) &\mapsto (f_* 1, \tau_0(R(\gamma)), \tau_0(h)). \end{aligned}$$

Lemma 7.6. *We have $\tau_0(f_* K(\nabla_f)) = f_* 1$.*

Proof. This follows from the definition of τ_0 and the fact $K_0 = 1$ since K is a multiplicative sequence. ■

Theorem 7.7. *For every $X \in \mathbf{Man}_{\mathbb{C}}$, the map $\tau_{\mathbb{Z}}$ induces a natural homomorphism*

$$\hat{\tau}_{\mathbb{Z}}: MU^n(p)(X) \rightarrow H_{\mathcal{D}}^n(X; \mathbb{Z}(p))$$

which fits into a morphism of long exact sequences

$$\begin{CD} \cdots @>>> H^{n-1}(X; \frac{\mathcal{A}^*}{\tilde{F}^p}(\mathcal{V}_*)) @>>> MU^n(p)(X) @>>> MU^n(X) @>>> \cdots \\ @. @V \tau_0 VV @V \hat{\tau}_{\mathbb{Z}} VV @V \tau VV @. \\ \cdots @>>> H^{n-1}(X; \frac{\mathcal{A}^*}{\tilde{F}^p}(\mathbb{C})) @>>> H_{\mathcal{D}}^n(X; \mathbb{Z}(p)) @>>> H^n(X; \mathbb{Z}) @>>> \cdots \end{CD} \tag{7.1}$$

Proof. It is clear that $\tau_{\mathbb{Z}}$ is a group homomorphism. We need to prove that, for a cycle $\gamma = (\tilde{f}, h) \in ZMU^n(p)(X)$, we have

$$d\tau_{\mathbb{Z}}(\gamma) = 0 \quad \text{and} \quad \tau_{\mathbb{Z}}(BMU^n(p)(X)) \subset dC_{\mathcal{D}}^{n-1}(p)(X).$$

We begin with the former. We have

$$\begin{aligned} d\tau_{\mathbb{Z}}(\tilde{f}, h) &= d(f_*1, \tau_0(R(\gamma)), \tau_0(h)) \\ &= (df_*1, \tau_0(dR(\gamma)), \tau_0(dh) + f_*1 - \tau_0(R(\gamma))). \end{aligned}$$

Since f_*1 is a closed current, and $R(\gamma)$ is a closed form, we deduce $d\tau_{\mathbb{Z}}(\gamma) = 0$ from Lemma 7.6. Now let \tilde{b} be a geometric bordism datum. Then

$$\hat{\tau}_{\mathbb{Z}}(\partial\tilde{b}, \psi(\tilde{b})) = (\tau_0\phi(\partial\tilde{b}), 0, \tau_0\psi(\tilde{b})) = (\tau_0d\psi(\tilde{b}), 0, \tau_0\psi(\tilde{b})) = d(\tau_0\psi(\tilde{b}), 0, 0).$$

Next let $h \in \tilde{F}^p\mathcal{A}^{n-1}(X; \mathcal{V}_*)$. Then $\tau_0(h) \in \tilde{F}^p\mathcal{A}^{n-1}(X)$, so that

$$(0, \tau_0(h), 0) \in C_{\mathcal{D}}^{n-1}(p)(X).$$

We have

$$\tau_{\mathbb{Z}}(a(h)) = \tau_{\mathbb{Z}}(0, h) = (0, \tau_0(dh), \tau_0(h)) = d(0, \tau_0(h), 0)$$

which finishes the proof that $\tau_{\mathbb{Z}}$ induces a homomorphism. The second assertion follows directly from the construction of $\hat{\tau}_{\mathbb{Z}}$. ■

Let X be a compact Kähler manifold. Let $f: Y \rightarrow X$ be the inclusion of a complex submanifold of codimension p such that its fundamental class in $MU^{2p}(X)$ vanishes. The latter condition implies that the fundamental class of f in $H^{2p}(X; \mathbb{Z})$ vanishes as well. Hence both the classical Abel–Jacobi invariant $AJ_H(f)$ of Deligne–Griffiths (see e.g. [24, Section 12]) and the invariant $AJ(f)$ of Theorems 6.9 and 6.10 are defined.

Theorem 7.8. *With the above notation and assumptions, we have*

$$\tau_0(AJ(f)) = AJ_H(f).$$

Proof. By Theorem 6.10 the invariant $AJ(f)$ may be represented by the functional

$$[\omega] \mapsto \text{ev} \left(\int_Y \sigma_f \wedge f^* \omega + \int_{W_{[0,1]}} (K(\nabla_b)) \wedge w^* \omega \right).$$

The image of the Chern–Simons form σ_f under $\hat{\tau}_{\mathbb{Z}}$ and τ_0 is zero since σ_f is a form in degree -1 . By Lemma 7.6, $K(\nabla_b)$ is mapped to 1. Thus, τ_0 maps $AJ(f)$ to the class of the functional in $F^{n-p+1}H^{2n-2p+1}(X; \mathbb{C})'$ defined by

$$[\omega] \mapsto \text{ev} \int_{W_{[0,1]}} w^* \omega.$$

This corresponds to the characterization of $AJ_H(f)$ in [24, Section 12.1.2, p. 294]. ■

8. Image and kernel for compact Kähler manifolds

We assume again that X is a compact Kähler manifold. Then the morphism of long exact sequences (7.1) induces a map of short exact sequences

$$\begin{array}{ccccccc}
 0 & \longrightarrow & J_{MU}^{2p-1}(X) & \longrightarrow & MU^{2p}(p)(X) & \longrightarrow & \text{Hdg}_{MU}^{2p}(X) \longrightarrow 0 \\
 & & \downarrow \tau_J & & \downarrow \hat{\tau}_{\mathbb{Z}} & & \downarrow \tau \\
 0 & \longrightarrow & J^{2p-1}(X) & \longrightarrow & H_{\mathcal{D}}^{2p}(X; \mathbb{Z}(p)) & \longrightarrow & \text{Hdg}^{2p}(X) \longrightarrow 0.
 \end{array} \tag{8.1}$$

Let $\mathcal{M}^p(X)$ be the free abelian group generated by isomorphism classes $[f]$ of proper holomorphic maps $f: Y \rightarrow X$ of codimension p . For a proper holomorphic map $f: Y \rightarrow X$ of codimension p we denote its fundamental class in $MU^{2p}(p)(X)$ by $\hat{\varphi}(f)$ and its fundamental class in $MU^{2p}(X)$ by $\varphi(f)$. This defines homomorphisms of abelian groups

$$\hat{\varphi}: \mathcal{M}^p(X) \rightarrow MU^{2p}(p)(X) \quad \text{and} \quad \varphi: \mathcal{M}^p(X) \rightarrow MU^{2p}(X).$$

We denote the kernel of φ by $\mathcal{M}^p(X)_{\text{top}}$. Then the Abel–Jacobi invariant of Definition 6.8 defines a homomorphism

$$AJ: \mathcal{M}^p(X)_{\text{top}} \rightarrow J_{MU}^{2p-1}(X).$$

Note that every element in $\mathcal{M}^p(X)_{\text{top}}$ is homologically equivalent to zero and therefore has a well-defined image in $J^{2p-1}(X)$. By Theorems 7.7 and 7.8 composition with the respective maps of diagram (8.1) produces the classical invariants. Diagram (8.1) shows that studying the kernel and image of $\hat{\tau}_{\mathbb{Z}}$ is equivalent to analysing the kernel and image of τ_J and τ , respectively. We expect the maps $\hat{\varphi}$ and AJ to be useful to discover new phenomena and examples that the classical invariants with values in Deligne cohomology are not able to detect. We will now briefly report on some results in this direction.

First we look at the image of $\widehat{\tau}_{\mathbb{Z}}$. Let X be a smooth projective complex algebraic variety. In [23], Totaro showed that an element in $H^{2*}(X(\mathbb{C}); \mathbb{Z})$ which is not in the image of $\tau: MU^{2*}(X(\mathbb{C})) \rightarrow H^{2*}(X(\mathbb{C}); \mathbb{Z})$ cannot be algebraic. This is a refinement of the obstruction induced by the Atiyah–Hirzebruch spectral sequence (see also [2]). It follows from [16, Corollary 7.12] that an algebraic class in $H^{2*}(X(\mathbb{C}); \mathbb{Z})$ has to be in the subgroup $\tau(\text{Hdg}_{MU}^{2*}(X(\mathbb{C})))$. In [1, Section 3.4], Benoist shows that this obstruction to algebraicity of cohomology classes is in fact finer than the one of [23].

Now we consider the kernel of τ_J . Since τ_0 is an epimorphism of vector spaces, the map τ_J is surjective, and the snake lemma implies that there is a short exact sequence

$$0 \rightarrow \ker \tau_J \rightarrow \ker \widehat{\tau}_{\mathbb{Z}} \rightarrow \ker \tau \rightarrow 0.$$

Hence $\ker \widehat{\tau}_{\mathbb{Z}}$ contains information on the failure of the Thom morphism τ to be injective on Hodge classes, and on the failure of τ_J to be injective. We have a further short exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & MU^{2p-1}(X)_{mt} & \xrightarrow{\phi_{mt}} & \frac{H^{2p-1}(X; \mathbb{V}_*)}{F^p H^{2p-1}(X; \mathbb{V}_*)} & \longrightarrow & J_{MU}^{2p-1}(X) \longrightarrow 0 \\ & & \downarrow \tau_{mt} & & \downarrow \tau_{\bar{J}} & & \downarrow \tau_J \\ 0 & \longrightarrow & H^{2p-1}(X; \mathbb{Z})_{mt} & \xrightarrow{i} & \frac{H^{2p-1}(X; \mathbb{C})}{F^p H^{2p-1}(X; \mathbb{C})} & \longrightarrow & J^{2p-1}(X) \longrightarrow 0 \end{array}$$

where the subscript mt means modulo torsion. Again, since τ_0 is onto, it follows that $\tau_{\bar{J}}$ is onto. Then the snake lemma places $\ker \tau_J$ in the exact sequence

$$0 \rightarrow \ker \tau \rightarrow \ker \tau_{\bar{J}} \rightarrow \ker \tau_J \rightarrow \text{coker } \tau_{mt} \rightarrow 0.$$

This indicates two methods to construct elements in $\ker \tau_J$: as elements coming from $\ker \tau_{\bar{J}}$ or as elements coming from $\text{coker } \tau_{mt}$. We will now briefly describe both these methods.

The arguments in [16, Section 7.3] show how to construct elements in $\ker \tau_{\bar{J}}$. We note that even though we have not shown that $MU^{2*}(\ast)(-)$ receives a map from algebraic cobordism for algebraic varieties, we can adjust the arguments as follows. Let \mathbb{P}^1 be the complex projective line, and let $[\mathbb{P}^1]$ denote corresponding element in MU^{-2} . Let $f: Y \rightarrow X$ be a proper holomorphic map of codimension p . Let $\mathbb{P}_X^1 \rightarrow X$ denote the pullback of \mathbb{P}^1 to X . By Lemma 6.3 we get a well-defined homomorphism

$$\mathcal{M}^p(X) \rightarrow MU^{2p-2}(p-1)(X)$$

induced by sending $[Y]$ to $[Y] \cdot [\mathbb{P}_X^1]$. Since X is compact, there is an isomorphism

$$MU^*(X) \otimes_{\mathbb{Z}} \mathbb{Q} \cong H^*(X; \mathbb{Q}) \otimes_{\mathbb{Z}} MU^*.$$

This implies that the sum $\bigoplus_{p \in \mathbb{Z}} J_{MU}^{2p-1}(X) \otimes \mathbb{Q}$ is a flat MU^* -module. Thus, for $\gamma \in \mathcal{M}^p(X)$, if $\text{AJ}(\gamma)$ is non-zero in $J_{MU}^{2p-1}(X) \otimes \mathbb{Q}$, then $\text{AJ}(\gamma) \cdot [\mathbb{P}^1]$ is non-zero in

$J_{MU}^{2p-3}(X) \otimes \mathbb{Q}$ and therefore non-zero in $J_{MU}^{2p-3}(X)$. Now we can take an element $\gamma \in \mathcal{M}^p(X)$ such that $\varphi(\gamma) = 0$ and the image of γ in $J^{2p-1}(X)$ is non-torsion. Then the above argument shows that $\text{AJ}(\gamma) \cdot [\mathbb{P}^1]$ is non-zero in $J_{MU}^{2p-3}(X)$. However, the image $\tau_J(\text{AJ}(\gamma) \cdot [\mathbb{P}^1])$ vanishes in $J^{2p-3}(X)$ since τ_0 sends $[\mathbb{P}^1]$ to zero. Examples of this situation where X is a projective smooth complex algebraic variety are described in [16, Examples 7.15 and 7.16].

Finally, we look at $\text{coker } \tau$. The most interesting case is that of a non-torsion element in $\text{coker } \tau$ which induces an element in $\ker \tau_J$ that remains non-trivial after taking the tensor product with \mathbb{R}/\mathbb{Z} over MU^* . For certain complex Lie groups, for example $SO(5)$, we can show that there are such elements in $\text{coker } \tau$. However, we are so far not able to produce such elements for X being compact or even projective.

Acknowledgments. Parts of the present paper grew out of the work on the first-named author's doctoral thesis. Both authors would like to thank the Department of Mathematical Sciences at NTNU for the continuous support during the work on the thesis and this paper. We thank the anonymous referee for many valuable comments, suggestions and questions which helped improving the paper.

Funding. The second author was partially supported by the RCN Project No. 313472.

References

- [1] O. Benoist, Steenrod operations and algebraic classes. 2022, arXiv:2209.03685
- [2] O. Benoist and J. C. Ottem, *Two coniveau filtrations*. *Duke Math. J.* **170** (2021), no. 12, 2719–2753 Zbl 1478.14022 MR 4305380
- [3] J. M. Boardman, *Conditionally convergent spectral sequences*. In *Homotopy invariant algebraic structures (Baltimore, MD, 1998)*, pp. 49–84, Contemp. Math. 239, American Mathematical Society, Providence, RI, 1999 Zbl 0947.55020 MR 1718076
- [4] G. E. Bredon, *Sheaf theory*. 2nd edn., Grad. Texts in Math. 170, Springer, New York, 1997 Zbl 0874.55001 MR 1481706
- [5] U. Bunke, T. Schick, I. Schröder, and M. Wiethaup, *Landweber exact formal group laws and smooth cohomology theories*. *Algebr. Geom. Topol.* **9** (2009), no. 3, 1751–1790 Zbl 1181.55006 MR 2550094
- [6] S. S. Chern and J. Simons, *Characteristic forms and geometric invariants*. *Ann. of Math. (2)* **99** (1974), 48–69 Zbl 0283.53036 MR 353327
- [7] G. de Rham, *Differentiable manifolds*. Grundlehren Math. Wiss. 266, Springer, Berlin, 1984 Zbl 0534.58003 MR 760450
- [8] J. P. Demailly, *Complex analytic and differential geometry*. 2012, online book draft
- [9] H. Esnault, *Characteristic classes of flat bundles*. *Topology* **27** (1988), no. 3, 323–352 Zbl 0699.32016 MR 963635
- [10] D. S. Freed and J. Lott, *An index theorem in differential K-theory*. *Geom. Topol.* **14** (2010), no. 2, 903–966 Zbl 1197.58007 MR 2602854

- [11] H. Gillet and C. Soulé, [Arithmetic Chow groups and differential characters](#). In *Algebraic K-theory: connections with geometry and topology (Lake Louise, AB, 1987)*, pp. 29–68, NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci. 279, Kluwer Academic Publishers, Dordrecht, 1989 Zbl [0719.14003](#) MR [1045844](#)
- [12] P. Griffiths and J. Harris, *Principles of algebraic geometry*. Wiley Classics Lib., John Wiley & Sons, New York, 1994 Zbl [0836.14001](#) MR [1288523](#)
- [13] R. Harvey, B. Lawson, and J. Zweck, [The de Rham–Federer theory of differential characters and character duality](#). *Amer. J. Math.* **125** (2003), no. 4, 791–847 Zbl [1060.58004](#) MR [1993742](#)
- [14] K. B. Haus and G. Quick, [Geometric Hodge filtered complex cobordism](#). *Adv. Math.* **431** (2023), article no. 109244 Zbl [07741093](#) MR [4628822](#)
- [15] F. Hirzebruch, T. Berger, and R. Jung, *Manifolds and modular forms*. Aspects Math., E 20, Springer Fachmedien Wiesbaden, Braunschweig, 1992 Zbl [0767.57014](#) MR [1189136](#)
- [16] M. J. Hopkins and G. Quick, [Hodge filtered complex bordism](#). *J. Topol.* **8** (2015), no. 1, 147–183 Zbl [1349.32009](#) MR [3335251](#)
- [17] L. Hörmander, *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis*. Classics Math., Springer, Berlin, 2003; Reprint of the second (1990) edition Zbl [1028.35001](#) MR [1996773](#)
- [18] D. Huybrechts, *Complex geometry*. Universitext, Springer, Berlin, 2005 Zbl [1055.14001](#) MR [2093043](#)
- [19] M. Karoubi, [Théorie générale des classes caractéristiques secondaires](#). *K-Theory* **4** (1990), no. 1, 55–87 Zbl [0716.57018](#) MR [1076525](#)
- [20] M. Karoubi, [Classes caractéristiques de fibrés feuilletés, holomorphes ou algébriques](#). In *Proceedings of Conference on Algebraic Geometry and Ring Theory in honor of Michael Artin, Part II (Antwerp, 1992)*. *K-Theory* **8** (1994), no. 2, 153–211 Zbl [0833.57012](#) MR [1273841](#)
- [21] M. Levine and F. Morel, *Algebraic cobordism*. Springer Monogr. Math., Springer, Berlin, 2007 Zbl [1188.14015](#) MR [2286826](#)
- [22] G. Quick, [An Abel–Jacobi invariant for cobordant cycles](#). *Doc. Math.* **21** (2016), 1645–1668 Zbl [1355.14016](#) MR [3603931](#)
- [23] B. Totaro, [Torsion algebraic cycles and complex cobordism](#). *J. Amer. Math. Soc.* **10** (1997), no. 2, 467–493 Zbl [0989.14001](#) MR [1423033](#)
- [24] C. Voisin, *Hodge theory and complex algebraic geometry. I*. Cambridge Stud. Adv. Math. 76, Cambridge University Press, Cambridge, 2002 Zbl [1005.14002](#) MR [1967689](#)

Communicated by Otmar Venjakob

Received 2 April 2023; revised 6 December 2023.

Knut Bjarte Haus

Department of Mathematical Sciences, Norwegian University of Science and Technology, 7491 Trondheim, Norway; knut.b.haus@gmail.com

Gereon Quick

Department of Mathematical Sciences, Norwegian University of Science and Technology, 7491 Trondheim, Norway; gereon.quick@ntnu.no