

# Liouville-type theorems for double-phase problems involving the Grushin operator

Quang Thanh Khuat

**Abstract.** In this paper, we are concerned with the double-phase problem involving the Grushin operator in the whole space  $\mathbb{R}^N = \mathbb{R}^{N_1} \times \mathbb{R}^{N_2}$

$$-\operatorname{div}_G(|\nabla_G u|^{p-2} \nabla_G u + w(z)|\nabla_G u|^{q-2} \nabla_G u) = f(z)|u|^{r-1} u,$$

where  $\nabla_G$  is the Grushin gradient,  $\Delta_G$  is the Grushin operator,  $q \geq p \geq 2$ ,  $r > q - 1$  and  $w, f \in L^1_{\text{loc}}(\mathbb{R}^N)$  are two nonnegative functions satisfying some growth conditions at infinity. Our purpose is to establish some Liouville-type theorems for stable weak solutions or for weak solutions which are stable outside a compact set of the equation above.

## 1. Introduction and main results

The Liouville-type theorems, which are concerned with the nonexistence of nontrivial solutions of PDEs, have been intensively developed in the past few decades since they have emerged as a fundamental tool for many applications to the qualitative properties of solutions of partial differential equations; see, e.g., [5, 6, 16, 27, 28]. In this paper, we study Liouville-type theorems for the double-phase problem involving the Grushin operator in the whole space  $\mathbb{R}^N = \mathbb{R}^{N_1} \times \mathbb{R}^{N_2}$

$$-\operatorname{div}_G(|\nabla_G u|^{p-2} \nabla_G u + w(z)|\nabla_G u|^{q-2} \nabla_G u) = f(z)|u|^{r-1} u, \quad (1.1)$$

where  $\nabla_G = (\nabla_x, |x|^\gamma \nabla_y)$ ,  $\Delta_G = \operatorname{div}_G \circ \nabla_G = \Delta_x + |x|^{2\gamma} \Delta_y$ ,  $q \geq p \geq 2$ , and  $r > q - 1$ . In this paper, we assume that  $w, f \in L^1_{\text{loc}}(\mathbb{R}^N)$  are two nonnegative functions verifying the following condition: there exist  $R_0, C_1, C_2 > 0$  and  $a, b \in \mathbb{R}$  such that

$$w(z) \leq C_1 |z|_G^a \quad (1.2)$$

and

$$f(z) \geq C_2 |z|_G^b \quad (1.3)$$

---

*Mathematics Subject Classification 2020:* 35B53 (primary); 35B35, 35J60 (secondary).

*Keywords:* Liouville-type theorems, double-phase problem, Grushin operator, stable solutions.

for all  $|z|_G > R_0$ . Here,  $z = (x, y) \in \mathbb{R}^{N_1} \times \mathbb{R}^{N_2}$  and

$$|z|_G = [ |x|^{2(\gamma+1)} + (\gamma + 1)^2 |y|^2 ]^{\frac{1}{2(\gamma+1)}}.$$

Note that, there is a group of dilations  $\{\delta_t\}_{t>0}$

$$\delta_t : \mathbb{R}^N \rightarrow \mathbb{R}^N, \quad \delta_t(z) = \delta_t(x, y) = (tx, t^{\gamma+1}y)$$

such that the operator  $\Delta_G = \operatorname{div}_G \circ \nabla_G = \Delta_x + |x|^{2\gamma} \Delta_y$  is  $\delta_t$ -homogeneous of degree two. This operator is elliptic when  $|x| \neq 0$  and degenerates on  $\{0\} \times \mathbb{R}^{N_2}$ .

Let us first recall that, in the case  $\gamma = 0, w = 0, p = 2$ , the left-hand side of (1.1) becomes  $-\Delta u$ . The most well-known Liouville-type result for nonlinear elliptic equations involving the Laplace operator was given in the celebrated article [16], in which the authors proved that the Lane–Emden equation

$$-\Delta u = u^r \quad \text{in } \mathbb{R}^N$$

has no positive solution if and only if  $r < \frac{N+2}{N-2}$ . About the class of stable solutions of the Lane–Emden equation, Farina [13] obtained an optimal Liouville-type theorem; see also [30].

Next, in the case  $\gamma = 0, w = 0, p \geq 2$ , equation (1.1) becomes

$$-\Delta_p u = f(z)|u|^{r-1}u \quad \text{in } \mathbb{R}^N.$$

The reader is referred to [29] for the nonexistence result of positive solutions and to [7, 9] for the nonexistence result of stable solutions of this equation.

Considering (1.1) in the case  $\gamma = 0, w \geq 0, q \geq p \geq 2$ , the operator of the left-hand side becomes the double-phase operator, and we have the equation

$$-\operatorname{div}(|\nabla u|^{p-2} \nabla u + w(z)|\nabla u|^{q-2} \nabla u) = f(z)|u|^{r-1}u \quad \text{in } \mathbb{R}^N. \tag{1.4}$$

Following the ideas in the papers [9, 14], Phuong Le [24] established the nonexistence of nontrivial stable solutions in  $W_{\text{loc}}^{1,H}(\mathbb{R}^N)$  of (1.4) under the condition

$$N < \min \left\{ \frac{p(\alpha_0 + r) + b(\alpha_0 + p - 1)}{r - p + 1}, \frac{(q - a)(\alpha_0 + r) + b(\alpha_0 + q - 1)}{r - q + 1} \right\},$$

where

$$\alpha_0 := \frac{2r - q + 1 + 2\sqrt{r(r - q + 1)}}{q - 1}.$$

We now consider the general case  $\gamma \geq 0$ . In recent years, there has been an increasing interest in Liouville-type theorems for elliptic equations involving the operator  $\Delta_G = \operatorname{div}_G \circ \nabla_G = \Delta_x + |x|^{2\gamma} \Delta_y$ ; see, e.g., [11, 12, 23, 31, 33]. Currently, this operator is usually named Grushin operator. The operators of this kind were first introduced and studied by Franchi and Lanconelli [15]. Recently, they were named by Kogoj and Lanconelli [19]

$\Delta_\lambda$ -Laplacians, under the additional assumption that the operators are homogeneous of degree two with respect to a group of dilations; see also [1, 10, 20–22, 25, 31, 32]. The operator considered by Grushin [17] is a very particular case of the  $\Delta_\lambda$ -Laplacians; Grushin studied this operator by adding lower terms with complex coefficients; see also [3]. For the equation

$$-\Delta_G u = u^r \quad \text{in } \mathbb{R}^N = \mathbb{R}^{N_1} \times \mathbb{R}^{N_2},$$

the Liouville-type theorem was proved by Monticelli [26] for nonnegative classical solutions and by Yu [33] for nonnegative weak solutions. The optimal condition on the range of the exponent is

$$r < \frac{Q + 2}{Q - 2},$$

where

$$Q = N_1 + (\gamma + 1)N_2 \tag{1.5}$$

is called the homogeneous dimension of  $\mathbb{R}^N$  associated to the Grushin operator. The Liouville results for the nonlinear elliptic equations involving  $p$ -Laplace-type Grushin operator

$$-\operatorname{div}_G(w(z)|\nabla_G u|^{p-2}\nabla_G u) = f(z)g(u) \quad \text{in } \mathbb{R}^N = \mathbb{R}^{N_1} \times \mathbb{R}^{N_2}$$

was studied in [31], where  $g(u) = e^u$  or  $g(u) = -u^{-p}$ . However, to our best knowledge, there has not any work treating the double-phase problem (1.1) involving the Grushin operator. The purpose of this paper is to establish some nonexistence results of nontrivial stable solutions of (1.1). Notice that double-phase differential operators and their corresponding energy functionals appear in nonlinear elasticity theory, strongly anisotropic materials, Lavrentiev’s phenomenon, and so on; see, e.g., [2, 4, 8, 18, 34–36].

We next recall some notations which will be used in the sequel. Let  $\Omega \subset \mathbb{R}^N$  be an open domain, and let  $H : \Omega \times [0, \infty) \rightarrow [0, \infty)$  be the function  $(z, t) \mapsto t^p + w(z)t^q$ . Put

$$\rho_H(u) = \int_\Omega H(z, |u|) = \int_\Omega (|u|^p + w(z)|u|^q)$$

and

$$L^H(\Omega) = \{u : \Omega \rightarrow \mathbb{R} \mid u \text{ is measurable and } \rho_H(u) < \infty\},$$

with the norm

$$\|u\|_H = \inf\left\{\tau > 0 \mid \rho_H\left(\frac{u}{\tau}\right) \leq 1\right\}.$$

Define

$$W^{1,H}(\Omega) = \{u \in L^H(\Omega) \mid |\nabla_G u| \in L^H(\Omega)\},$$

with the norm

$$\|u\|_{1,H} = \|\nabla_G u\|_H + \|u\|_H.$$

Denote by  $W_0^{1,H}(\Omega)$  the closure of  $C_c^1(\Omega)$  with respect to the  $\|\cdot\|_{1,H}$  norm and

$$W_{\text{loc}}^{1,H}(\Omega) = \{u : \Omega \rightarrow \mathbb{R} \mid u\varphi \in W_0^{1,H}(\Omega) \text{ for all } \varphi \in C_c^1(\Omega)\}.$$

In order to state our results, we need the following definitions.

**Definition 1.1.** A function  $u \in W_{\text{loc}}^{1,H}(\mathbb{R}^N)$  is said to be a weak solution of equation (1.1) if  $f(z)|u|^r \in L_{\text{loc}}^1(\mathbb{R}^N)$ , and for all  $\varphi \in C_c^1(\mathbb{R}^N)$  we have

$$\int_{\mathbb{R}^N} (|\nabla_G u|^{p-2} \nabla_G u + w(z)|\nabla_G u|^{q-2} \nabla_G u) \cdot \nabla_G \varphi = \int_{\mathbb{R}^N} f(z)|u|^{r-1} u \varphi. \tag{1.6}$$

Moreover,  $u$  is called a finite energy solution if

$$\int_{\mathbb{R}^N} (|\nabla_G u|^p + w(z)|\nabla_G u|^q + f(z)|u|^{r+1}) < \infty.$$

**Definition 1.2.** We say that a weak solution  $u$  of (1.1)

- is stable if for all  $\varphi \in C_c^1(\mathbb{R}^N)$  we have

$$\begin{aligned} L_u(\varphi) := & \int_{\mathbb{R}^N} [|\nabla_G u|^{p-2} |\nabla_G \varphi|^2 + (p-2)|\nabla_G u|^{p-4} (\nabla_G u \cdot \nabla_G \varphi)^2] \\ & + \int_{\mathbb{R}^N} w(z) [|\nabla_G u|^{q-2} |\nabla_G \varphi|^2 + (q-2)|\nabla_G u|^{q-4} (\nabla_G u \cdot \nabla_G \varphi)^2] \\ & - r \int_{\mathbb{R}^N} f(z)|u|^{r-1} \varphi^2 \geq 0, \end{aligned} \tag{1.7}$$

- is stable outside a compact set  $K \subset \mathbb{R}^N$  if  $L_u(\varphi) \geq 0$  only holds for  $\varphi \in C_c^1(\mathbb{R}^N \setminus K)$ ,
- has Morse index equal to  $k \geq 1$  if  $k$  is the maximal dimension of a subspace  $V \subset C_c^1(\mathbb{R}^N)$  such that  $L_u(\varphi) < 0$  for all  $\varphi \in V \setminus \{0\}$ .

Notice that the stability condition (1.7) is nothing but the fact that the second variation at  $u$  of the energy functional

$$E(u) = \int_{\mathbb{R}^N} \left( \frac{|\nabla_G u|^p}{p} + \frac{w(z)|\nabla_G u|^q}{q} - \frac{f(z)|u|^{r+1}}{r+1} \right)$$

is nonnegative. Therefore, all the local minima of the functional are stable solutions of (1.1). Besides, we also know that every finite Morse index solution is stable outside a compact set. Indeed, there exist  $m_0 \geq 1$  and  $V_0 := \text{Span}\{\varphi_1, \dots, \varphi_{m_0}\} \subset C_c^1(\mathbb{R}^N)$  such that  $L_u(\varphi) < 0$  for any  $\varphi \in V_0 \setminus \{0\}$ . Therefore,  $L_u(\varphi) \geq 0$  for every  $\varphi \in C_c^1(\mathbb{R}^N \setminus K)$ , where  $K := \bigcup_{j=1}^{m_0} \text{supp}(\varphi_j)$ .

Our first result concerning the classification of stable weak solutions of (1.1) is as follows.

**Theorem 1.3.** Let  $q \geq p \geq 2, r > q - 1$ . Suppose that  $w, f \in L_{\text{loc}}^1(\mathbb{R}^N; [0, \infty))$  satisfying (1.2) and (1.3). Under the condition

$$Q < \min \left\{ \frac{p(\alpha_0 + r) + b(\alpha_0 + p - 1)}{r - p + 1}, \frac{(q - a)(\alpha_0 + r) + b(\alpha_0 + q - 1)}{r - q + 1} \right\}, \tag{1.8}$$

where

$$\alpha_0 := \frac{2r - q + 1 + 2\sqrt{r(r - q + 1)}}{q - 1},$$

equation (1.1) has only the trivial stable weak solution. Here, the homogeneous dimension  $Q$  is given in (1.5).

Next, we consider a special case of  $f$  and  $w$  in (1.1), i.e.,

$$-\operatorname{div}_G(|\nabla_G u|^{p-2} \nabla_G u + |z|_G^a |\nabla_G u|^{q-2} \nabla_G u) = |z|_G^b |u|^{r-1} u \quad \text{in } \mathbb{R}^N = \mathbb{R}^{N_1} \times \mathbb{R}^{N_2}. \tag{1.9}$$

We next give some conditions ensuring the nonexistence of nontrivial finite energy solutions of (1.9). Our second result is as follows.

**Theorem 1.4.** *Let  $q > p \geq 2, r > q - 1$ . Suppose that  $u$  is a finite energy solution of (1.9) satisfying  $|\nabla_G u|^{p-2} \nabla_G u + |z|_G^a |\nabla_G u|^{q-2} \nabla_G u \in W_{\text{loc}}^{1,2}(\mathbb{R}^N, \mathbb{R}^N)$ . If*

$$\frac{Q + b}{r + 1} > \max \left\{ \frac{Q - p}{p}, \frac{Q - q + a}{q} \right\},$$

then  $u$  must be trivial, i.e.,  $u = 0$ .

Under the same assumptions in Theorem 1.4, we will prove that any weak solution, which is stable outside a compact set, is also a finite energy solution. Therefore, we obtain the following theorem.

**Theorem 1.5.** *Let  $q > p \geq 2, r > q - 1$ . Suppose that  $u$  is a weak solution of (1.9) which is stable outside a compact set and  $|\nabla_G u|^{p-2} \nabla_G u + |z|_G^a |\nabla_G u|^{q-2} \nabla_G u \in W_{\text{loc}}^{1,2}(\mathbb{R}^N, \mathbb{R}^N)$ . If*

$$\frac{Q + b}{r + 1} > \max \left\{ \frac{Q - p}{p}, \frac{Q - q + a}{q} \right\},$$

then  $u$  must be trivial, i.e.,  $u = 0$ .

In particular, our results generalize those in [24] from the Laplace operator to the Grushin operator. Our results are also extensions of those in [31] to the double-phase problem. Inspired by [11, 24, 31], our approach in this paper is also based on the energy method and a Pohozaev identity involving the Grushin operator established below.

This paper is organized as follows. In Section 2, we prove Theorem 1.3. The proof of Theorem 1.4 is given in Section 3. The last section is devoted to the proof of Theorem 1.5.

## 2. Proof of Theorem 1.3

We begin this section by fixing some notations. Let  $\Omega_R = B_1(0, R) \times B_2(0, R^{\gamma+1})$ , where  $B_1(0, R) \subset \mathbb{R}^{N_1}$  and  $B_2(0, R^{\gamma+1}) \subset \mathbb{R}^{N_2}$  are the Euclidean balls.

First, we prove the following a priori estimate for stable weak solutions of (1.1).

**Proposition 2.1.** *Let  $u$  be a stable weak solution of (1.1). Suppose that  $\alpha \geq 1, r > \frac{(q-1)(\alpha+1)^2}{4\alpha}$  and  $m \geq 1$  so that*

$$\min \left\{ \frac{(mq - p)(\alpha + r)}{\alpha + p - 1}, \frac{(mq - q)(\alpha + r)}{\alpha + q - 1} \right\} \geq mq. \tag{2.1}$$

Then, for all  $\eta \in C_c^1(\mathbb{R}^N; [0, 1])$  and  $\nabla_G \eta = 0$  in  $\Omega_{R_0}$ , there is a positive constant  $C$  depending on  $p, q, r, \alpha, m$  such that

$$\begin{aligned} & \int_{\mathbb{R}^N} (|\nabla_G u|^p |u|^{\alpha-1} + w(z) |\nabla_G u|^q |u|^{\alpha-1} + f(z) |u|^{\alpha+r}) \eta^{qm} \\ & \leq C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} f(z)^{-\frac{\alpha+p-1}{r-p+1}} |\nabla_G \eta|^{\frac{p(\alpha+r)}{r-p+1}} \\ & \quad + C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} w(z)^{\frac{\alpha+r}{r-q+1}} f(z)^{-\frac{\alpha+q-1}{r-q+1}} |\nabla_G \eta|^{\frac{q(\alpha+r)}{r-q+1}}. \end{aligned}$$

*Proof.* For each  $k \in \mathbb{N}$ , we define

$$a_k(t) = \begin{cases} |t|^{\frac{\alpha-1}{2}} t, & |t| < k, \\ k^{\frac{\alpha-1}{2}} t, & |t| \geq k, \end{cases} \quad \text{and} \quad b_k(t) = \begin{cases} |t|^{\alpha-1} t, & |t| < k, \\ k^{\alpha-1} t, & |t| \geq k. \end{cases}$$

By direct calculations, we have

$$a'_k(t) = \begin{cases} \frac{\alpha+1}{2} |t|^{\frac{\alpha-1}{2}}, & |t| < k, \\ k^{\frac{\alpha-1}{2}}, & |t| \geq k, \end{cases} \quad \text{and} \quad b'_k(t) = \begin{cases} \alpha |t|^{\alpha-1}, & |t| < k, \\ k^{\alpha-1}, & |t| \geq k. \end{cases}$$

It follows that

$$\begin{aligned} a_k(t)^2 = t b_k(t), \quad a'_k(t)^2 & \leq \frac{(\alpha+1)^2}{4\alpha} b'_k(t), \\ |a_k(t)|^s a'_k(t)^{2-s} + |b_k(t)|^s b'_k(t)^{1-s} & \leq C_{\alpha,s} |t|^{\alpha+s-1} \end{aligned} \tag{2.2}$$

for all  $t \in \mathbb{R}$  and  $s \geq 2$ .

By density arguments, we observe that (1.6) holds for all  $\varphi \in W_0^{1,H}(\mathbb{R}^N)$ . Moreover, one may easily check that if  $u \in W_{\text{loc}}^{1,H}(\mathbb{R}^N)$ , then  $a_k(u), b_k(u) \in W_{\text{loc}}^{1,H}(\mathbb{R}^N)$  for any  $k \in \mathbb{N}$ . This implies that  $a_k(u) \psi^{\frac{\beta}{2}} \in W_0^{1,H}(\mathbb{R}^N)$  and  $b_k(u) \psi^\beta \in W_0^{1,H}(\mathbb{R}^N)$  for all  $\psi \in C_c^1(\mathbb{R}^N)$ ,  $k \in \mathbb{N}$ , and  $\beta \geq q$ .

Let  $\beta \geq q$ ,  $\varepsilon \in (0, 1)$ , and  $\psi \in C_c^1(\mathbb{R}^N)$  satisfying  $0 \leq \psi \leq 1$ . Using

$$\varphi = b_k(u) \psi^\beta$$

as a test function in (1.6) gives us

$$\begin{aligned} & \int_{\mathbb{R}^N} |\nabla_G u|^p b'_k(u) \psi^\beta + \beta \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} b_k(u) \psi^{\beta-1} \nabla_G u \cdot \nabla_G \psi \\ & + \int_{\mathbb{R}^N} w(z) |\nabla_G u|^q b'_k(u) \psi^\beta + \beta \int_{\mathbb{R}^N} w(z) |\nabla_G u|^{q-2} b_k(u) \psi^{\beta-1} \nabla_G u \cdot \nabla_G \psi \\ & = \int_{\mathbb{R}^N} f(z) |u|^{r-1} u b_k(u) \psi^\beta. \end{aligned}$$

Applying Young’s inequality, we obtain

$$\begin{aligned}
 & -\beta \int_{\mathbb{R}^N} w(z) |\nabla_G u|^{q-2} b_k(u) \psi^{\beta-1} \nabla_G u \cdot \nabla_G \psi \\
 & \leq \beta \int_{\mathbb{R}^N} w(z) |\nabla_G u|^{q-1} |b_k(u)| \psi^{\beta-1} |\nabla_G \psi| \\
 & \leq \int_{\mathbb{R}^N} \left\{ \varepsilon \left( w(z)^{\frac{q-1}{q}} |\nabla_G u|^{q-1} b'_k(u)^{\frac{q-1}{q}} \psi^{\frac{(q-1)\beta}{q}} \right)^{\frac{q}{q-1}} \right. \\
 & \quad \left. + C_\varepsilon \left( w(z)^{\frac{1}{q}} |b_k(u)| b'_k(u)^{\frac{1-q}{q}} \psi^{\frac{\beta-q}{q}} |\nabla_G \psi| \right)^q \right\} \\
 & = \varepsilon \int_{\mathbb{R}^N} w(z) |\nabla_G u|^q b'_k(u) \psi^\beta \\
 & \quad + C_\varepsilon \int_{\mathbb{R}^N} w(z) |b_k(u)|^q b'_k(u)^{1-q} \psi^{\beta-q} |\nabla_G \psi|^q.
 \end{aligned}$$

In the same way, we have

$$\begin{aligned}
 & -\beta \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} b_k(u) \psi^{\beta-1} \nabla_G u \cdot \nabla_G \psi \\
 & \leq \varepsilon \int_{\mathbb{R}^N} |\nabla_G u|^p b'_k(u) \psi^\beta + C_\varepsilon \int_{\mathbb{R}^N} |b_k(u)|^p b'_k(u)^{1-p} \psi^{\beta-p} |\nabla_G \psi|^p.
 \end{aligned}$$

Employing the two estimates above and (2.2), we get that

$$\begin{aligned}
 & (1 - \varepsilon) \int_{\mathbb{R}^N} (|\nabla_G u|^p + w(z) |\nabla_G u|^q) b'_k(u) \psi^\beta \\
 & \leq \int_{\mathbb{R}^N} f(z) |u|^{r-1} u b_k(u) \psi^\beta \\
 & \quad + C_\varepsilon \int_{\mathbb{R}^N} |b_k(u)|^p b'_k(u)^{1-p} \psi^{\beta-p} |\nabla_G \psi|^p \\
 & \quad + C_\varepsilon \int_{\mathbb{R}^N} w(z) |b_k(u)|^q b'_k(u)^{1-q} \psi^{\beta-q} |\nabla_G \psi|^q \\
 & \leq \int_{\mathbb{R}^N} f(z) |u|^{r-1} u b_k(u) \psi^\beta + C_\varepsilon \int_{\mathbb{R}^N} |u|^{\alpha+p-1} \psi^{\beta-p} |\nabla_G \psi|^p \\
 & \quad + C_\varepsilon \int_{\mathbb{R}^N} w(z) |u|^{\alpha+q-1} \psi^{\beta-q} |\nabla_G \psi|^q. \tag{2.3}
 \end{aligned}$$

Applying Schwartz’s inequality for the stability condition (1.7), we obtain

$$\begin{aligned}
 & (p - 1) \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} |\nabla_G \varphi|^2 + (q - 1) \int_{\mathbb{R}^N} w(z) |\nabla_G u|^{q-2} |\nabla_G \varphi|^2 \\
 & \geq r \int_{\mathbb{R}^N} f(z) |u|^{r-1} \varphi^2 \tag{2.4}
 \end{aligned}$$

for all  $\varphi \in C_c^1(\mathbb{R}^N)$ . By density arguments, we have that (2.4) holds true for all  $\varphi \in W_0^{1,H}(\mathbb{R}^N)$ . Next, using  $\varphi = a_k(u)\psi^{\frac{\beta}{2}} \in W_0^{1,H}(\mathbb{R}^N)$  as a test function in (2.4), we obtain

$$\begin{aligned} & (p-1) \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} \left| a'_k(u)\psi^{\frac{\beta}{2}} \nabla_G u + \frac{\beta}{2} a_k(u)\psi^{\frac{\beta-2}{2}} \nabla_G \psi \right|^2 \\ & + (q-1) \int_{\mathbb{R}^N} w(z) |\nabla_G u|^{q-2} \left| a'_k(u)\psi^{\frac{\beta}{2}} \nabla_G u + \frac{\beta}{2} a_k(u)\psi^{\frac{\beta-2}{2}} \nabla_G \psi \right|^2 \\ & \geq r \int_{\mathbb{R}^N} f(z) |u|^{r-1} a_k(u)^2 \psi^\beta. \end{aligned}$$

Applying the inequality

$$|z_1 + z_2|^2 \leq (1 + \delta)|z_1|^2 + C_\delta |z_2|^2 \quad \text{for } z_1, z_2 \in \mathbb{R}^N, \delta > 0,$$

we arrive at

$$\begin{aligned} & r \int_{\mathbb{R}^N} f(z) |u|^{r-1} a_k(u)^2 \psi^\beta \\ & \leq \left( p-1 + \frac{\varepsilon}{2} \right) \int_{\mathbb{R}^N} |\nabla_G u|^p a'_k(u)^2 \psi^\beta + A_\varepsilon \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} a_k(u)^2 \psi^{\beta-2} |\nabla_G \psi|^2 \\ & \quad + \left( q-1 + \frac{\varepsilon}{2} \right) \int_{\mathbb{R}^N} w(z) |\nabla_G u|^q a'_k(u)^2 \psi^\beta \\ & \quad + B_\varepsilon \int_{\mathbb{R}^N} w(z) |\nabla_G u|^{q-2} a_k(u)^2 \psi^{\beta-2} |\nabla_G \psi|^2. \end{aligned} \tag{2.5}$$

In the case  $q > 2$ , employing Young’s inequality, we get that

$$\begin{aligned} & B_\varepsilon \int_{\mathbb{R}^N} w(z) |\nabla_G u|^{q-2} a_k(u)^2 \psi^{\beta-2} |\nabla_G \psi|^2 \\ & \leq \int_{\mathbb{R}^N} \left\{ \frac{\varepsilon}{2} \left( w(z)^{\frac{q-2}{q}} |\nabla_G u|^{q-2} a'_k(u)^{\frac{2(q-2)}{q}} \psi^{\frac{(q-2)\beta}{q}} \right)^{\frac{q}{q-2}} \right. \\ & \quad \left. + C_\varepsilon \left( w(z)^{\frac{2}{q}} a_k(u)^2 a'_k(u)^{\frac{2(2-q)}{q}} \psi^{\frac{2(\beta-q)}{q}} |\nabla_G \psi|^2 \right)^{\frac{q}{2}} \right\} \\ & = \frac{\varepsilon}{2} \int_{\mathbb{R}^N} w(z) |\nabla_G u|^q a'_k(u)^2 \psi^\beta + C_\varepsilon \int_{\mathbb{R}^N} w(z) |a_k(u)|^q a'_k(u)^{2-q} \psi^{\beta-q} |\nabla_G \psi|^q. \end{aligned}$$

We see that this inequality is also true in the case  $q = 2$ . By the same argument, we have

$$\begin{aligned} & A_\varepsilon \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} a_k(u)^2 \psi^{\beta-2} |\nabla_G \psi|^2 \\ & \leq \frac{\varepsilon}{2} \int_{\mathbb{R}^N} |\nabla_G u|^p a'_k(u)^2 \psi^\beta + C_\varepsilon \int_{\mathbb{R}^N} |a_k(u)|^p a'_k(u)^{2-p} \psi^{\beta-p} |\nabla_G \psi|^p. \end{aligned}$$



Putting these two estimates back into (2.5) gives

$$\begin{aligned} & r \int_{\mathbb{R}^N} f(z)|u|^{r-1}a_k(u)^2\psi^\beta \\ & \leq (p-1+\varepsilon) \int_{\mathbb{R}^N} |\nabla_G u|^p a'_k(u)^2\psi^\beta + (q-1+\varepsilon) \int_{\mathbb{R}^N} w(z)|\nabla_G u|^q a'_k(u)^2\psi^\beta \\ & \quad + C_\varepsilon \int_{\mathbb{R}^N} |a_k(u)|^p a'_k(u)^{2-p}\psi^{\beta-p}|\nabla_G \psi|^p \\ & \quad + C_\varepsilon \int_{\mathbb{R}^N} w(z)|a_k(u)|^q a'_k(u)^{2-q}\psi^{\beta-q}|\nabla_G \psi|^q. \end{aligned}$$

Using the fact that  $q \geq p$  and (2.2), we derive

$$\begin{aligned} & r \int_{\mathbb{R}^N} f(z)|u|^{r-1}a_k(u)^2\psi^\beta \\ & \leq (q-1+\varepsilon) \int_{\mathbb{R}^N} (|\nabla_G u|^p + w(z)|\nabla_G u|^q)a'_k(u)^2\psi^\beta \\ & \quad + C_\varepsilon \int_{\mathbb{R}^N} |u|^{\alpha+p-1}\psi^{\beta-p}|\nabla_G \psi|^p + C_\varepsilon \int_{\mathbb{R}^N} w(z)|u|^{\alpha+q-1}\psi^{\beta-q}|\nabla_G \psi|^q \\ & \leq \frac{(q-1+\varepsilon)(\alpha+1)^2}{4\alpha} \int_{\mathbb{R}^N} (|\nabla_G u|^p + w(z)|\nabla_G u|^q)b'_k(u)\psi^\beta \\ & \quad + C_\varepsilon \int_{\mathbb{R}^N} |u|^{\alpha+p-1}\psi^{\beta-p}|\nabla_G \psi|^p + C_\varepsilon \int_{\mathbb{R}^N} w(z)|u|^{\alpha+q-1}\psi^{\beta-q}|\nabla_G \psi|^q. \end{aligned} \tag{2.6}$$

Combining (2.3) and (2.6), we obtain

$$\begin{aligned} & r \int_{\mathbb{R}^N} f(z)|u|^{r-1}a_k(u)^2\psi^\beta \\ & \leq \frac{(q-1+\varepsilon)(\alpha+1)^2}{4\alpha(1-\varepsilon)} \int_{\mathbb{R}^N} f(z)|u|^{r-1}ub_k(u)\psi^\beta \\ & \quad + C_\varepsilon \int_{\mathbb{R}^N} |u|^{\alpha+p-1}\psi^{\beta-p}|\nabla_G \psi|^p + C_\varepsilon \int_{\mathbb{R}^N} w(z)|u|^{\alpha+q-1}\psi^{\beta-q}|\nabla_G \psi|^q. \end{aligned}$$

Using (2.2) gives us

$$\begin{aligned} & r \int_{\mathbb{R}^N} f(z)|u|^{r-1}a_k(u)^2\psi^\beta \\ & \leq \frac{(q-1+\varepsilon)(\alpha+1)^2}{4\alpha(1-\varepsilon)} \int_{\mathbb{R}^N} f(z)|u|^{r-1}a_k(u)^2\psi^\beta \\ & \quad + C_\varepsilon \int_{\mathbb{R}^N} |u|^{\alpha+p-1}\psi^{\beta-p}|\nabla_G \psi|^p + C_\varepsilon \int_{\mathbb{R}^N} w(z)|u|^{\alpha+q-1}\psi^{\beta-q}|\nabla_G \psi|^q. \end{aligned}$$

Therefore,

$$\begin{aligned} & D_\varepsilon \int_{\mathbb{R}^N} f(z)|u|^{r-1}a_k(u)^2\psi^\beta \\ & \leq C_\varepsilon \int_{\mathbb{R}^N} |u|^{\alpha+p-1}\psi^{\beta-p}|\nabla_G \psi|^p + C_\varepsilon \int_{\mathbb{R}^N} w(z)|u|^{\alpha+q-1}\psi^{\beta-q}|\nabla_G \psi|^q, \end{aligned}$$

where

$$D_\varepsilon := r - \frac{(q-1+\varepsilon)(\alpha+1)^2}{4\alpha(1-\varepsilon)}.$$

Since  $\lim_{\varepsilon \rightarrow 0^+} D_\varepsilon = r - \frac{(q-1)(\alpha+1)^2}{4\alpha} > 0$ , we can fix some  $\varepsilon$  sufficiently close to zero such that  $D_\varepsilon > 0$ . We also fix  $\beta = q$ . Hence,

$$\begin{aligned} \int_{\mathbb{R}^N} f(z)|u|^{r-1}a_k(u)^2\psi^q &\leq C \int_{\mathbb{R}^N} |u|^{\alpha+p-1}\psi^{q-p}|\nabla_G\psi|^p \\ &\quad + C \int_{\mathbb{R}^N} w(z)|u|^{\alpha+q-1}|\nabla_G\psi|^q. \end{aligned}$$

Combining this with (2.3) and using (2.2), we can also bound gradient terms as follows:

$$\begin{aligned} \int_{\mathbb{R}^N} (|\nabla_G u|^p + w(z)|\nabla_G u|^q)b'_k(u)\psi^q &+ \int_{\mathbb{R}^N} f(z)|u|^{r-1}a_k(u)^2\psi^q \\ &\leq C \int_{\mathbb{R}^N} |u|^{\alpha+p-1}\psi^{q-p}|\nabla_G\psi|^p + C \int_{\mathbb{R}^N} w(z)|u|^{\alpha+q-1}|\nabla_G\psi|^q. \end{aligned}$$

Using Fatou’s lemma when letting  $k \rightarrow \infty$ , we get that

$$\begin{aligned} \int_{\mathbb{R}^N} (|\nabla_G u|^p|u|^{\alpha-1} + w(z)|\nabla_G u|^q|u|^{\alpha-1} + f(z)|u|^{\alpha+r})\psi^q \\ \leq C \int_{\mathbb{R}^N} |u|^{\alpha+p-1}\psi^{q-p}|\nabla_G\psi|^p + C \int_{\mathbb{R}^N} w(z)|u|^{\alpha+q-1}|\nabla_G\psi|^q. \end{aligned} \tag{2.7}$$

Next, take  $\psi = \eta^m$  in (2.7) and apply the Young inequality to find that

$$\begin{aligned} &\int_{\mathbb{R}^N} (|\nabla_G u|^p|u|^{\alpha-1} + w(z)|\nabla_G u|^q|u|^{\alpha-1} + f(z)|u|^{\alpha+r})\eta^{qm} \\ &\leq C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} |u|^{\alpha+p-1}|\nabla_G \eta|^p \eta^{mq-p} + C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} w(z)|u|^{\alpha+q-1}|\nabla_G \eta|^q \eta^{q(m-1)} \\ &\leq \int_{\mathbb{R}^N \setminus \Omega_{R_0}} \left\{ \frac{1}{4} \left( f(z)^{\frac{\alpha+p-1}{\alpha+r}} |u|^{\alpha+p-1} \eta^{mq-p} \right)^{\frac{\alpha+r}{\alpha+p-1}} + C \left( f(z)^{-\frac{\alpha+p-1}{\alpha+r}} |\nabla_G \eta|^p \right)^{\frac{\alpha+r}{r-p+1}} \right\} \\ &\quad + \int_{\mathbb{R}^N \setminus \Omega_{R_0}} \left\{ \frac{1}{4} \left( f(z)^{\frac{\alpha+q-1}{\alpha+r}} |u|^{\alpha+q-1} \eta^{q(m-1)} \right)^{\frac{\alpha+r}{\alpha+q-1}} \right. \\ &\quad \left. + C \left( w(z) f(z)^{-\frac{\alpha+q-1}{\alpha+r}} |\nabla_G \eta|^q \right)^{\frac{\alpha+r}{r-q+1}} \right\} \\ &\leq \frac{1}{4} \int_{\mathbb{R}^N \setminus \Omega_{R_0}} f(z)|u|^{\alpha+r} \eta^{qm} + C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} f(z)^{-\frac{\alpha+p-1}{r-p+1}} |\nabla_G \eta|^{\frac{p(\alpha+r)}{r-p+1}} \\ &\quad + \frac{1}{4} \int_{\mathbb{R}^N \setminus \Omega_{R_0}} f(z)|u|^{\alpha+r} \eta^{qm} + C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} w(z)^{\frac{\alpha+r}{r-q+1}} f(z)^{-\frac{\alpha+q-1}{r-q+1}} |\nabla_G \eta|^{\frac{q(\alpha+r)}{r-q+1}}. \end{aligned}$$

Therefore,

$$\begin{aligned} & \int_{\mathbb{R}^N} (|\nabla_G u|^p |u|^{\alpha-1} + w(z) |\nabla_G u|^q |u|^{\alpha-1} + f(z) |u|^{\alpha+r}) \eta^{qm} \\ & \leq C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} f(z)^{-\frac{\alpha+p-1}{r-p+1}} |\nabla_G \eta|^{\frac{p(\alpha+r)}{r-p+1}} \\ & \quad + C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} w(z)^{\frac{\alpha+r}{r-q+1}} f(z)^{-\frac{\alpha+q-1}{r-q+1}} |\nabla_G \eta|^{\frac{q(\alpha+r)}{r-q+1}}. \end{aligned}$$

The proof is complete. ■

Let  $R > 0$ ,  $\Omega_{2R} = B_1(0, 2R) \times B_2(0, 2R^{\gamma+1})$ , where

$$B_1(0, 2R) \subset \mathbb{R}^{N_1}, \quad B_2(0, 2R^{\gamma+1}) \subset \mathbb{R}^{N_2}$$

are the Euclidean balls. Define

$$\eta_{1,R}(x) = \eta_1\left(\frac{|x|}{R}\right), \quad \eta_{2,R}(y) = \eta_2\left(\frac{|y|}{R^{\gamma+1}}\right),$$

where  $\eta_1, \eta_2 \in C_c^\infty([0, +\infty))$ ,  $0 \leq \eta_1, \eta_2 \leq 1$ ,

$$\eta_i(t) = \begin{cases} 1 & \text{in } [0, 1], \\ 0 & \text{in } [2, +\infty), \end{cases}$$

and for some constant  $C > 0$  and  $\eta_{1,R}, \eta_{2,R}$  satisfy

$$|\nabla_x \eta_{1,R}| \leq \frac{C}{R}, \quad |\nabla_y \eta_{2,R}| \leq \frac{C}{R^{\gamma+1}}.$$

**Lemma 2.2.** *The following assertions hold true.*

(i) *There exists a constant  $C > 0$  independent of  $R$  such that*

$$|\nabla_G \eta_R| \leq \frac{C}{R}, \quad \forall z \in \Omega_{2R},$$

where  $\eta_R(z) = \eta_{1,R}(x)\eta_{2,R}(y)$ .

(ii) *There exists a constant  $C > 0$  independent of  $R$  such that if  $z \in \Omega_{2R}$ , then*

$$|z|_G \leq CR.$$

(iii) *If  $z \notin \Omega_R$ , then  $|z|_G > R$ .*

*Proof.* Proof of (i). We have

$$\nabla_G \eta_R = (\nabla_x \eta_R, |x|^\gamma \nabla_y \eta_R) = (\eta_{2,R} \nabla_x \eta_{1,R}, |x|^\gamma \eta_{1,R} \nabla_y \eta_{2,R}).$$

For any  $z = (x, y) \in \Omega_{2R}$ , we have  $x \in B_1(0, 2R)$ . This implies

$$|x| \leq 2R.$$

Combining this with the hypothesis about functions  $\eta_{i,R}$ ,  $i = 1, 2$ , there exists a constant  $C > 0$  independent of  $R$  such that

$$|\nabla_G \eta_R|^2 = \eta_{2,R}^2 |\nabla_x \eta_{1,R}|^2 + |x|^{2\gamma} \eta_{1,R}^2 |\nabla_y \eta_{2,R}|^2 \leq \frac{C}{R^2}, \quad \forall z \in \Omega_{2R}.$$

Therefore, there is a constant  $C > 0$  independent of  $R$  such that

$$|\nabla_G \eta_R| \leq \frac{C}{R}, \quad \forall z \in \Omega_{2R}.$$

Proof of (ii). For any  $z = (x, y) \in \Omega_{2R}$ , we have  $x \in B_1(0, 2R)$ ,  $y \in B_2(0, 2R^{\gamma+1})$ . This implies

$$|x| \leq 2R \quad \text{and} \quad |y| \leq 2R^{\gamma+1}.$$

Then, we get

$$\begin{aligned} |z|_G &= [|x|^{2(\gamma+1)} + (\gamma + 1)^2 |y|^2]^{\frac{1}{2(\gamma+1)}} \\ &\leq [(2R)^{2(\gamma+1)} + (\gamma + 1)^2 (2R^{\gamma+1})^2]^{\frac{1}{2(\gamma+1)}}. \end{aligned}$$

By direct calculation, we obtain

$$|z|_G \leq CR,$$

where  $C$  is a positive constant independent of  $R$ .

Proof of (iii). For any  $z = (x, y) \notin \Omega_R$ , we have  $x \notin B_1(0, R)$ ,  $y \notin B_2(0, R^{\gamma+1})$ . This implies

$$|x| > R \quad \text{and} \quad |y| > R^{\gamma+1}.$$

We deduce that

$$|z|_G > [R^{2(\gamma+1)} + (\gamma + 1)^2 (R^{\gamma+1})^2]^{\frac{1}{2(\gamma+1)}}.$$

By direct calculation, we have  $|z|_G > R$ . ■

*Proof Theorem 1.3.* Observe that

$$\alpha \geq 1 \quad \text{and} \quad r > \frac{(q-1)(\alpha+1)^2}{4\alpha}$$

are equivalent to  $1 \leq \alpha < \alpha_0$ . Besides, it follows from the assumption  $r > q - 1 \geq p - 1$  that there is  $m \geq 1$  satisfying (2.1). Clearly, the standard cut-off function  $\eta_R$  is chosen as in Lemma 2.2 satisfying  $\eta_R \in C_c^1(\mathbb{R}^N)$ ,  $0 \leq \eta_R \leq 1$  in  $\mathbb{R}^N$  and

$$\begin{cases} \eta_R = 1 & \text{in } \Omega_R, \\ \eta_R = 0 & \text{in } \mathbb{R}^N \setminus \Omega_{2R}, \\ |\nabla_G \eta_R| \leq \frac{C}{R} & \text{in } \Omega_{2R} \setminus \Omega_R, \end{cases}$$

where  $R > R_0$ . For all  $R > R_0$ , applying the Proposition 2.1 with  $\alpha \in [1, \alpha_0)$  and the function  $\eta_R$  as in Lemma 2.2, there is a constant  $C$  independent of  $R$  such that

$$\begin{aligned} & \int_{\Omega_R} (|\nabla_G u|^p |u|^{\alpha-1} + w(z)|\nabla_G u|^q |u|^{\alpha-1} + f(z)|u|^{\alpha+r}) \\ & \leq C \int_{\Omega_{2R} \setminus \Omega_R} f(z)^{-\frac{\alpha+p-1}{r-p+1}} |\nabla_G \eta_R|^{\frac{p(\alpha+r)}{r-p+1}} \\ & \quad + C \int_{\Omega_{2R} \setminus \Omega_R} w(z)^{\frac{\alpha+r}{r-q+1}} f(z)^{-\frac{\alpha+q-1}{r-q+1}} |\nabla_G \eta_R|^{\frac{q(\alpha+r)}{r-q+1}}. \end{aligned}$$

Since  $|z|_G > R > R_0, \forall z \notin \Omega_R$ , and using the hypothesis about  $w(z), f(z), \eta_R$ , we obtain

$$\begin{aligned} & \int_{\Omega_R} (|\nabla_G u|^p |u|^{\alpha-1} + w(z)|\nabla_G u|^q |u|^{\alpha-1} + f(z)|u|^{\alpha+r}) \\ & \leq C \int_{\Omega_{2R} \setminus \Omega_R} |z|_G^{-\frac{b(\alpha+p-1)}{r-p+1}} R^{-\frac{p(\alpha+r)}{r-p+1}} + C \int_{\Omega_{2R} \setminus \Omega_R} |z|_G^{\frac{a(\alpha+r)}{r-q+1}} |z|_G^{-\frac{b(\alpha+q-1)}{r-q+1}} R^{-\frac{q(\alpha+r)}{r-q+1}}. \end{aligned}$$

Combining this with  $|z|_G > R, \forall z \notin \Omega_R$  and  $|z|_G \leq CR, \forall z \in \Omega_{2R}$ , we have

$$\int_{\Omega_R} (|\nabla_G u|^p |u|^{\alpha-1} + w(z)|\nabla_G u|^q |u|^{\alpha-1} + f(z)|u|^{\alpha+r}) \leq CR^\theta, \tag{2.8}$$

where

$$\theta = Q - \min \left\{ \frac{p(\alpha+r) + b(\alpha+p-1)}{r-p+1}, \frac{(q-a)(\alpha+r) + b(\alpha+q-1)}{r-q+1} \right\}.$$

By assumption (1.8), we may choose  $\alpha$  sufficiently close to  $\alpha_0$  such that  $\theta < 0$ . Then, by letting  $R \rightarrow \infty$  in (2.8), we get the conclusion of the theorem. ■

### 3. Proof of Theorem 1.4

We begin this section with the following Pohozaev-type identity involving the Grushin operator.

**Proposition 3.1.** *Suppose that  $u$  is a finite energy solution of equation (1.9) and*

$$|\nabla_G u|^{p-2} \nabla_G u + |z|_G^a |\nabla_G u|^{q-2} \nabla_G u \in W_{\text{loc}}^{1,2}(\mathbb{R}^N; \mathbb{R}^N).$$

Then, we have

$$\frac{Q-p}{p} \int_{\mathbb{R}^N} |\nabla_G u|^p + \frac{Q-q+a}{q} \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^q = \frac{Q+b}{r+1} \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1}.$$

*Proof.* By density arguments and the regularity provided, we may use  $v_R = \eta_R z^* \cdot \nabla_G u$  as a test function in (1.6), where  $\eta_R$  is chosen as in Lemma 2.2 and  $z^* = (x, \frac{\gamma+1}{|x|^\gamma} y)$ , we have

$$\int_{\mathbb{R}^N} (|\nabla_G u|^{p-2} \nabla_G u \cdot \nabla_G v_R + |z|_G^a |\nabla_G u|^{q-2} \nabla_G u \cdot \nabla_G v_R) = \int_{\mathbb{R}^N} |z|_G^b |u|^{r-1} u v_R. \tag{3.1}$$

Using the divergence theorem, we obtain

$$\begin{aligned} & \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} \nabla_G u \cdot \nabla_G v_R \\ &= \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} (\nabla_G u \cdot \nabla_G \eta_R) (z^* \cdot \nabla_G u) \\ & \quad + \int_{\mathbb{R}^N} |\nabla_G u|^p \eta_R + \int_{\mathbb{R}^N} \eta_R z^* \cdot \nabla_G \left( \frac{|\nabla_G u|^p}{p} \right) \\ &= \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} (\nabla_G u \cdot \nabla_G \eta_R) (z^* \cdot \nabla_G u) \\ & \quad + \int_{\mathbb{R}^N} |\nabla_G u|^p \eta_R - \int_{\mathbb{R}^N} \nabla_G \eta_R \cdot z^* \frac{|\nabla_G u|^p}{p} - \int_{\mathbb{R}^N} \eta_R (\nabla_G \cdot z^*) \frac{|\nabla_G u|^p}{p} \\ &= \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} (\nabla_G u \cdot \nabla_G \eta_R) (z^* \cdot \nabla_G u) \\ & \quad + \int_{\mathbb{R}^N} |\nabla_G u|^p \eta_R - \int_{\mathbb{R}^N} \nabla_G \eta_R \cdot z^* \frac{|\nabla_G u|^p}{p} - Q \int_{\mathbb{R}^N} \eta_R \frac{|\nabla_G u|^p}{p}. \end{aligned}$$

It follows from  $\int_{\mathbb{R}^N} |\nabla_G u|^p < \infty$  and the dominated convergence theorem that

$$\lim_{R \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} \nabla_G u \cdot \nabla_G v_R = -\frac{Q-p}{p} \int_{\mathbb{R}^N} |\nabla_G u|^p. \tag{3.2}$$

Similarly,

$$\begin{aligned} & \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^{q-2} \nabla_G u \cdot \nabla_G v_R \\ &= \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^{q-2} (\nabla_G u \cdot \nabla_G \eta_R) (z^* \cdot \nabla_G u) \\ & \quad + \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^q \eta_R + \int_{\mathbb{R}^N} |z|_G^a \eta_R z^* \cdot \nabla_G \left( \frac{|\nabla_G u|^q}{q} \right) \\ &= \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^{q-2} (\nabla_G u \cdot \nabla_G \eta_R) (z^* \cdot \nabla_G u) + \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^q \eta_R \\ & \quad - a \int_{\mathbb{R}^N} |z|_G^a \eta_R \frac{|\nabla_G u|^q}{q} - \int_{\mathbb{R}^N} |z|_G^a \nabla_G \eta_R \cdot z^* \frac{|\nabla_G u|^q}{q} - Q \int_{\mathbb{R}^N} |z|_G^a \eta_R \frac{|\nabla_G u|^q}{q}. \end{aligned}$$

It follows from  $\int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^q < \infty$  and the dominated convergence theorem that

$$\lim_{R \rightarrow \infty} \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^{q-2} \nabla_G u \cdot \nabla_G v_R = -\frac{Q-q+a}{q} \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^q. \tag{3.3}$$

Now, we take  $\nu_R = \eta_R z^* \cdot \nabla_G u$  on the right-hand side of (3.1) and employing the divergence theorem to find

$$\begin{aligned} & \int_{\mathbb{R}^N} |z|_G^b |u|^{r-1} u \nu_R \\ &= \int_{\mathbb{R}^N} |z|_G^b |u|^{r-1} u \eta_R z^* \cdot \nabla_G u \\ &= \int_{\mathbb{R}^N} |z|_G^b \eta_R z^* \cdot \nabla_G \left( \frac{|u|^{r+1}}{r+1} \right) \\ &= - \int_{\mathbb{R}^N} \nabla_G (|z|_G^b) \cdot z^* \eta_R \frac{|u|^{r+1}}{r+1} - \int_{\mathbb{R}^N} |z|_G^b \nabla_G \eta_R \cdot z^* \frac{|u|^{r+1}}{r+1} \\ &\quad - \int_{\mathbb{R}^N} |z|_G^b \eta_R (\nabla_G \cdot z^*) \frac{|u|^{r+1}}{r+1} \\ &= -b \int_{\mathbb{R}^N} |z|_G^b \eta_R \frac{|u|^{r+1}}{r+1} - \int_{\mathbb{R}^N} |z|_G^b \nabla_G \eta_R \cdot z^* \frac{|u|^{r+1}}{r+1} - Q \int_{\mathbb{R}^N} |z|_G^b \eta_R \frac{|u|^{r+1}}{r+1}. \end{aligned}$$

Since  $\int_{\mathbb{R}^N} |z|_G^b |u|^{r+1} < \infty$ , we deduce from the dominated convergence theorem that

$$\lim_{R \rightarrow \infty} \int_{\mathbb{R}^N} |z|_G^b |u|^{r-1} u \nu_R = -\frac{Q+b}{r+1} \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1}. \tag{3.4}$$

The conclusion of the proposition follows from (3.1)–(3.4). ■

*Proof of Theorem 1.4.* To prove Theorem 1.4, first, we prove the following identity:

$$\int_{\mathbb{R}^N} |\nabla_G u|^p + |z|_G^a |\nabla_G u|^q = \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1}.$$

Indeed, using  $\varphi = u \eta_R$  as a test function in (1.6), where  $\eta_R$  is chosen as in Lemma 2.2, we obtain

$$\begin{aligned} & \int_{\mathbb{R}^N} |\nabla_G u|^p \eta_R + \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} u \nabla_G u \cdot \nabla_G \eta_R \\ &+ \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^q \eta_R + \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^{q-2} u \nabla_G u \cdot \nabla_G \eta_R \\ &= \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1} \eta_R. \end{aligned} \tag{3.5}$$

By means of Holder’s inequality, we have

$$\begin{aligned} & \left| \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} u \nabla_G u \cdot \nabla_G \eta_R \right| \\ & \leq \int_{\mathbb{R}^N} |\nabla_G u|^{p-1} |u| |\nabla_G \eta_R| \\ & \leq \left( \int_{\mathbb{R}^N} |\nabla_G u|^p \right)^{\frac{p-1}{p}} \left( \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1} \right)^{\frac{1}{r+1}} \left( \int_{\mathbb{R}^N} |z|_G^{-\frac{bp}{r-p+1}} |\nabla_G \eta_R|^{\frac{p(r+1)}{r-p+1}} \right)^{\frac{r-p+1}{p(r+1)}}. \end{aligned}$$

Using the hypothesis about  $\eta_R$  and Lemma 2.2, we have

$$\begin{aligned} & \left| \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} u \nabla_G u \cdot \nabla_G \eta_R \right| \\ & \leq C \left( \int_{\mathbb{R}^N} |\nabla_G u|^p \right)^{\frac{p-1}{p}} \left( \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1} \right)^{\frac{1}{r+1}} \left( \int_{\Omega_{2R} \setminus \Omega_R} |z|_G^{-\frac{bp}{r-p+1}} R^{-\frac{p(r+1)}{r-p+1}} \right)^{\frac{r-p+1}{p(r+1)}} \\ & \leq CR^{\frac{Q(r-p+1)}{p(r+1)} - \frac{r+b+1}{r+1}} \left( \int_{\mathbb{R}^N} |\nabla_G u|^p \right)^{\frac{p-1}{p}} \left( \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1} \right)^{\frac{1}{r+1}}. \end{aligned}$$

Using the condition  $\frac{Q+b}{r+1} > \frac{Q-p}{p}$ , we see that  $\frac{Q(r-p+1)}{p(r+1)} - \frac{r+b+1}{r+1} < 0$ . Therefore, by letting  $R \rightarrow \infty$ , we get that

$$\lim_{R \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla_G u|^{p-2} u \nabla_G u \cdot \nabla_G \eta_R = 0. \tag{3.6}$$

In the same way, we have

$$\begin{aligned} & \left| \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^{q-2} u \nabla_G u \cdot \nabla_G \eta_R \right| \\ & \leq \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^{q-1} |u| |\nabla_G \eta_R| \\ & \leq \left( \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^q \right)^{\frac{q-1}{q}} \left( \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1} \right)^{\frac{1}{r+1}} \\ & \quad \times \left( \int_{\mathbb{R}^N} |z|_G^{\frac{a(r+1)}{r-q+1}} |z|_G^{-\frac{bq}{r-q+1}} |\nabla_G \eta_R|^{\frac{q(r+1)}{r-q+1}} \right)^{\frac{r-q+1}{q(r+1)}} \\ & \leq CR^{\frac{Q(r-q+1)}{q(r+1)} + \frac{a}{q} - \frac{r+b+1}{r+1}} \left( \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^q \right)^{\frac{q-1}{q}} \left( \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1} \right)^{\frac{1}{r+1}}. \end{aligned}$$

By  $\frac{Q+b}{r+1} > \frac{Q-q+a}{q}$ , there holds  $\frac{Q(r-q+1)}{q(r+1)} + \frac{a}{q} - \frac{r+b+1}{r+1} < 0$ . Therefore, letting  $R \rightarrow \infty$ , we get that

$$\lim_{R \rightarrow \infty} \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^{q-2} u \nabla_G u \cdot \nabla_G \eta_R = 0. \tag{3.7}$$

Combining (3.5)–(3.7) leads to

$$\int_{\mathbb{R}^N} (|\nabla_G u|^p + |z|_G^a |\nabla_G u|^q) = \int_{\mathbb{R}^N} |z|_G^b |u|^{r+1}.$$

Substituting this identity into Proposition 3.1, we obtain

$$\left( \frac{Q+b}{r+1} - \frac{Q-p}{p} \right) \int_{\mathbb{R}^N} |\nabla_G u|^p + \left( \frac{Q+b}{r+1} - \frac{Q-q+a}{q} \right) \int_{\mathbb{R}^N} |z|_G^a |\nabla_G u|^q = 0.$$

Since  $\frac{Q+b}{r+1} > \max\{\frac{Q-p}{p}, \frac{Q-q+a}{q}\}$ , then it follows that  $u$  is constant and hence must be zero. ■



### 4. Proof of Theorem 1.5

The following a priori estimate holds for weak solutions which are stable outside a compact set.

**Proposition 4.1.** *Let  $u$  be a weak solution of (1.1) which is stable outside a compact set  $K \subset \mathbb{R}^N$ . Suppose that  $\alpha \geq 1$ ,  $r > \frac{(q-1)(\alpha+1)^2}{4\alpha}$ , and  $m \geq 1$  verifying*

$$\min \left\{ \frac{(mq - p)(\alpha + r)}{\alpha + p - 1}, \frac{(mq - q)(\alpha + r)}{\alpha + q - 1} \right\} \geq mq.$$

*Then, for all  $\eta \in C_c^1(\mathbb{R}^N \setminus (K \cup \Omega_{R_0}); [0, 1])$  and  $\nabla_G \eta = 0$  in  $\Omega_{R_0}$ , there is a constant  $C$  depending on  $p, q, r, \alpha, m$  such that*

$$\begin{aligned} & \int_{\mathbb{R}^N} (|\nabla_G u|^p |u|^{\alpha-1} + w(z) |\nabla_G u|^q |u|^{\alpha-1} + f(z) |u|^{\alpha+r}) \eta^{qm} \\ & \leq C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} f(z)^{-\frac{\alpha+p-1}{r-p+1}} |\nabla_G \eta|^{\frac{p(\alpha+r)}{r-p+1}} \\ & \quad + C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} w(z)^{\frac{\alpha+r}{r-q+1}} f(z)^{-\frac{\alpha+q-1}{r-q+1}} |\nabla_G \eta|^{\frac{q(\alpha+r)}{r-q+1}}. \end{aligned}$$

The proof of Proposition 4.1 follows closely to that of Proposition 2.1, and so, it will be omitted.

*Proof of Theorem 1.5.* Let  $u$  be a weak solution of (1.9) which is stable outside a compact set  $K$ . We fix  $R_1 > R_0$  so that  $K \subset \Omega_{R_1}$ . For any  $R > R_1 + 1$ , we consider  $\xi_R \in C_c^1(\mathbb{R}^N)$  such that  $0 \leq \xi_R \leq 1$  in  $\mathbb{R}^N$  and

$$\xi_R = \begin{cases} 0 & \text{in } \Omega_{R_1}, \\ 1 & \text{in } \Omega_R \setminus \Omega_{R_1+1}, \\ 0 & \text{in } \mathbb{R}^N \setminus \Omega_{2R}. \end{cases}$$

We may assume furthermore that  $|\nabla_G \xi_R| \leq C$  in  $\Omega_{R_1+1} \setminus \Omega_{R_1}$  and  $|\nabla_G \xi_R| \leq \frac{C}{R}$  in  $\Omega_{2R} \setminus \Omega_R$ , where  $C$  is independent of  $R$ . Then, applying Proposition 4.1 with  $\alpha = 1$ ,  $w(z) = |z|_G^a$ ,  $f(z) = |z|_G^b$ , and  $\eta = \xi_R$ , we obtain

$$\begin{aligned} & \int_{\mathbb{R}^N} (|\nabla_G u|^p + |z|_G^a |\nabla_G u|^q + |z|_G^b |u|^{r+1}) \xi_R^{qm} \\ & \leq C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} |z|_G^{-\frac{bp}{r-p+1}} |\nabla_G \xi_R|^{\frac{p(r+1)}{r-p+1}} \\ & \quad + C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} |z|_G^{\frac{a(r+1)}{r-q+1}} |z|_G^{-\frac{bq}{r-q+1}} |\nabla_G \xi_R|^{\frac{q(r+1)}{r-q+1}}. \end{aligned} \tag{4.1}$$

Using the hypothesis about  $\xi_R$ , we have

$$\begin{aligned} & \int_{\Omega_R \setminus \Omega_{R_{1+1}}} (|\nabla_G u|^p + |z|_G^a |\nabla_G u|^q + |z|_G^b |u|^{r+1}) \\ & \leq \int_{\mathbb{R}^N} (|\nabla_G u|^p + |z|_G^a |\nabla_G u|^q + |z|_G^b |u|^{r+1}) \xi_R^{qm}. \end{aligned}$$

We also obtain

$$\begin{aligned} & C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} |z|_G^{-\frac{bp}{r-p+1}} |\nabla_G \xi_R|^{\frac{p(r+1)}{r-p+1}} \\ & = C \int_{\Omega_{R_{1+1}} \setminus \Omega_{R_1}} |z|_G^{-\frac{bp}{r-p+1}} |\nabla_G \xi_R|^{\frac{p(r+1)}{r-p+1}} + C \int_{\Omega_{2R} \setminus \Omega_R} |z|_G^{-\frac{bp}{r-p+1}} |\nabla_G \xi_R|^{\frac{p(r+1)}{r-p+1}} \\ & \leq C_0 + C \int_{\Omega_{2R} \setminus \Omega_R} |z|_G^{-\frac{bp}{r-p+1}} |\nabla_G \xi_R|^{\frac{p(r+1)}{r-p+1}} \\ & \leq C_0 + C_1 R^{Q - \frac{p(r+b+1)}{r-p+1}}. \end{aligned}$$

Similarly,

$$C \int_{\mathbb{R}^N \setminus \Omega_{R_0}} |z|_G^{\frac{a(r+1)}{r-q+1}} |z|_G^{-\frac{bq}{r-q+1}} |\nabla_G \xi_R|^{\frac{q(r+1)}{r-q+1}} \leq C_0 + C_2 R^{Q - \frac{q(r+b+1)-a(r+1)}{r-q+1}}.$$

Substituting these three estimates into (4.1), we have

$$\begin{aligned} & \int_{\Omega_R \setminus \Omega_{R_{1+1}}} (|\nabla_G u|^p + |z|_G^a |\nabla_G u|^q + |z|_G^b |u|^{r+1}) \\ & \leq C_0 + C_1 R^{Q - \frac{p(r+b+1)}{r-p+1}} + C_2 R^{Q - \frac{q(r+b+1)-a(r+1)}{r-q+1}}. \end{aligned} \tag{4.2}$$

The condition

$$\frac{Q+b}{r+1} > \max \left\{ \frac{Q-p}{p}, \frac{Q-q+a}{q} \right\}$$

implies that

$$Q < \frac{p(r+b+1)}{r-p+1}$$

and

$$Q < \frac{q(r+b+1)-a(r+1)}{r-q+1}.$$

Therefore, letting  $R \rightarrow \infty$  in (4.2), we receive

$$|\nabla_G u|^p + |z|_G^a |\nabla_G u|^q + |z|_G^b |u|^{r+1} \in L^1(\mathbb{R}^N).$$

This and Theorem 1.4 complete the proof. ■

**Acknowledgments.** The author is grateful to the referee for the careful reading and helpful suggestions to improve the presentation of the paper.

**Funding.** Quang Thanh Khuat is supported by the Academy of Finance, Hanoi, Vietnam.

## References

- [1] C. T. Anh and B. K. My, [Liouville-type theorems for elliptic inequalities involving the  \$\Delta\_\lambda\$ -Laplace operator](#). *Complex Var. Elliptic Equ.* **61** (2016), no. 7, 1002–1013 Zbl 1346.35030 MR 3500512
- [2] A. Bahrouni, V. D. Rădulescu, and D. D. Repovš, [Double phase transonic flow problems with variable growth: nonlinear patterns and stationary waves](#). *Nonlinearity* **32** (2019), no. 7, 2481–2495 Zbl 1419.35056 MR 3957220
- [3] M. S. Baouendi, [Sur une classe d’opérateurs elliptiques dégénérés](#). *Bull. Soc. Math. France* **95** (1967), 45–87 Zbl 0179.19501 MR 0228819
- [4] V. Benci, P. D’Avenia, D. Fortunato, and L. Pisani, [Solitons in several space dimensions: Derrick’s problem and infinitely many solutions](#). *Arch. Ration. Mech. Anal.* **154** (2000), no. 4, 297–324 Zbl 0973.35161 MR 1785469
- [5] M.-F. Bidaut-Véron and L. Véron, [Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations](#). *Invent. Math.* **106** (1991), no. 3, 489–539 Zbl 0755.35036 MR 1134481
- [6] I. Birindelli, I. Capuzzo Dolcetta, and A. Cutri, [Liouville theorems for semilinear equations on the Heisenberg group](#). *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **14** (1997), no. 3, 295–308 Zbl 0876.35033 MR 1450950
- [7] C. Chen, [Liouville type theorem for stable solutions of  \$p\$ -Laplace equation in  \$\mathbb{R}^N\$](#) . *Appl. Math. Lett.* **68** (2017), 62–67 Zbl 1365.35045 MR 3614279
- [8] L. Cherfils and Y. Il’yasov, [On the stationary solutions of generalized reaction diffusion equations with  \$p\$ - \$q\$ -Laplacian](#). *Commun. Pure Appl. Anal.* **4** (2005), no. 1, 9–22 Zbl 1210.35090 MR 2126276
- [9] L. Damascelli, A. Farina, B. Sciunzi, and E. Valdinoci, [Liouville results for  \$m\$ -Laplace equations of Lane–Emden–Fowler type](#). *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **26** (2009), no. 4, 1099–1119 Zbl 1172.35405 MR 2542716
- [10] A. T. Duong, T. H. Giang, P. Le, and T. H. A. Vu, [Classification results for a sub-elliptic system involving the  \$\Delta\_\lambda\$ -Laplacian](#). *Math. Methods Appl. Sci.* **44** (2021), no. 5, 3615–3629 Zbl 1470.35086 MR 4227948
- [11] A. T. Duong and N. T. Nguyen, [Liouville type theorems for elliptic equations involving Grushin operator and advection](#). *Electron. J. Differential Equations* (2017), article no. 108 Zbl 1370.35125 MR 3651905
- [12] A. T. Duong and Q. H. Phan, [Liouville type theorem for nonlinear elliptic system involving Grushin operator](#). *J. Math. Anal. Appl.* **454** (2017), no. 2, 785–801 Zbl 1371.35078 MR 3658799
- [13] A. Farina, [On the classification of solutions of the Lane–Emden equation on unbounded domains of  \$\mathbb{R}^N\$](#) . *J. Math. Pures Appl. (9)* **87** (2007), no. 5, 537–561 Zbl 1143.35041 MR 2322150
- [14] A. Farina, [Stable solutions of  \$-\Delta u = e^u\$  on  \$\mathbb{R}^N\$](#) . *C. R. Math. Acad. Sci. Paris* **345** (2007), no. 2, 63–66 Zbl 1325.35048 MR 2343553
- [15] B. Franchi and E. Lanconelli, [Une métrique associée à une classe d’opérateurs elliptiques dégénérés](#). *Rend. Sem. Mat. Univ. Politec. Torino* (1983), no. Special Issue, 105–114 (1984); Conference on linear partial and pseudodifferential operators (Torino, 1982) Zbl 0553.35033 MR 0745979

- [16] B. Gidas and J. Spruck, [Global and local behavior of positive solutions of nonlinear elliptic equations](#). *Comm. Pure Appl. Math.* **34** (1981), no. 4, 525–598 Zbl 0465.35003 MR 0615628
- [17] V. V. Grushin, [On a class of elliptic pseudo differential operators degenerate on a submanifold](#). *Math. USSR Sb.* **13** (1971), no. 2, article no. 155
- [18] V. V. Jikov, S. M. Kozlov, and O. A. Oleĭnik, [Homogenization of differential operators and integral functionals](#). Springer, Berlin, 1994; Translated from the Russian by G. A. Yosifian [G. A. Iosif yan] Zbl 0838.35001 MR 1329546
- [19] A. E. Kogoj and E. Lanconelli, [On semilinear  \$\Delta\_\lambda\$ -Laplace equation](#). *Nonlinear Anal.* **75** (2012), no. 12, 4637–4649 Zbl 1260.35020 MR 2927124
- [20] A. E. Kogoj and E. Lanconelli, [Linear and semilinear problems involving  \$\Delta\_\lambda\$ -Laplacians](#). In *Proceedings of the International Conference “Two nonlinear days in Urbino 2017”*, pp. 167–178, Electron. J. Differ. Equ. Conf. 25, Texas State University–San Marcos, Department of Mathematics, San Marcos, TX, 2018 Zbl 1400.35130 MR 3883635
- [21] A. E. Kogoj and S. Sonner, [Attractors for a class of semi-linear degenerate parabolic equations](#). *J. Evol. Equ.* **13** (2013), no. 3, 675–691 Zbl 1286.35046 MR 3089799
- [22] A. E. Kogoj and S. Sonner, [Hardy type inequalities for  \$\Delta\_\lambda\$ -Laplacians](#). *Complex Var. Elliptic Equ.* **61** (2016), no. 3, 422–442 Zbl 1362.35107 MR 3454116
- [23] P. Le, [Liouville theorems for stable weak solutions of elliptic problems involving Grushin operator](#). *Commun. Pure Appl. Anal.* **19** (2020), no. 1, 511–525 Zbl 1427.35029 MR 4025955
- [24] P. Le, [Liouville results for double phase problems in  \$\mathbb{R}^N\$](#) . *Qual. Theory Dyn. Syst.* **21** (2022), no. 3, article no. 59 Zbl 1489.35022 MR 4412564
- [25] D. T. Luyen and N. M. Tri, [Existence of solutions to boundary-value problems for similinear  \$\Delta\_\gamma\$  differential equations](#). *Math. Notes* **97** (2015), no. 1-2, 73–84 Zbl 1325.35051 MR 3394492
- [26] D. D. Monticelli, [Maximum principles and the method of moving planes for a class of degenerate elliptic linear operators](#). *J. Eur. Math. Soc. (JEMS)* **12** (2010), no. 3, 611–654 Zbl 1208.35068 MR 2639314
- [27] Q. H. Phan and P. Souplet, [Liouville-type theorems and bounds of solutions of Hardy–Hénon equations](#). *J. Differential Equations* **252** (2012), no. 3, 2544–2562 Zbl 1233.35093 MR 2860629
- [28] P. Poláčik and P. Quittner, [Liouville type theorems and complete blow-up for indefinite super-linear parabolic equations](#). In *Nonlinear elliptic and parabolic problems*, pp. 391–402, Progr. Nonlinear Differential Equations Appl. 64, Birkhäuser, Basel, 2005 Zbl 1093.35037 MR 2185228
- [29] J. Serrin and H. Zou, [Cauchy–Liouville and universal boundedness theorems for quasilinear elliptic equations and inequalities](#). *Acta Math.* **189** (2002), no. 1, 79–142 Zbl 1059.35040 MR 1946918
- [30] C. Wang and D. Ye, [Some Liouville theorems for Hénon type elliptic equations](#). *J. Funct. Anal.* **262** (2012), no. 4, 1705–1727 Zbl 1246.35092 MR 2873856
- [31] Y. Wei, C. Chen, Q. Chen, and H. Yang, [Liouville-type theorem for nonlinear elliptic equations involving  \$p\$ -Laplace-type Grushin operators](#). *Math. Methods Appl. Sci.* **43** (2020), no. 1, 320–333 Zbl 1445.35161 MR 4044241
- [32] Y. Wei, C. Chen, and H. Yang, [Liouville-type theorem for Kirchhoff equations involving Grushin operators](#). *Bound. Value Probl.* (2020), article no. 13 Zbl 1489.35119 MR 4055450

- [33] X. Yu, [Liouville type theorem for nonlinear elliptic equation involving Grushin operators](#). *Commun. Contemp. Math.* **17** (2015), no. 5, article no. 1450050, 12 pp. Zbl [1326.35139](#) MR [3404744](#)
- [34] V. V. Zhikov, Averaging of functionals of the calculus of variations and elasticity theory. *Izv. Akad. Nauk SSSR Ser. Mat.* **50** (1986), no. 4, 675–710, 877 Zbl [0599.49031](#) MR [0864171](#)
- [35] V. V. Zhikov, On Lavrentiev’s phenomenon. *Russian J. Math. Phys.* **3** (1995), no. 2, 249–269 Zbl [0910.49020](#) MR [1350506](#)
- [36] V. V. Zhikov, On some variational problems. *Russian J. Math. Phys.* **5** (1997), no. 1, 105–116 (1998) Zbl [0917.49006](#) MR [1486765](#)

Received 21 September 2023.

**Quang Thanh Khuat**

Faculty of Fundamental Science, Academy of Finance, 58 Le Van Hien, Duc Thang, Bac Tu Liem, 10000 Hanoi, Vietnam; [khuatquangthanh@hvtc.edu.vn](mailto:khuatquangthanh@hvtc.edu.vn)