

# Tangential contact between free and fixed boundaries for variational solutions to variable-coefficient Bernoulli-type free boundary problems

Diego Moreira and Harish Shrivastava

**Abstract.** In this paper, we show that, given appropriate boundary data, the free boundaries of minimizers of functionals of type  $J(v; A, \lambda_+, \lambda_-, \Omega) = \int_{\Omega} (A(x)\nabla v, \nabla v) + \Lambda(v) dx$  and the fixed boundary touch each other in a tangential fashion. We extend the results of Karakhanyan, Kenig, and Shahgholian [Calc. Var. Partial Differential Equations 28 (2007), 15–31] to the case of variable coefficients. We prove this result via classification of the global profiles, as per Karakhanyan, Kenig, and Shahgholian [Calc. Var. Partial Differential Equations 28 (2007), 15–31].

## 1. Introduction

The objective of this paper is to study the behavior of free boundary near the fixed boundary of domain for minimizers of Bernoulli-type functionals with Hölder continuous coefficients

$$J(v; A, \lambda_+, \lambda_-, \Omega) = \int_{\Omega} ((A(x)\nabla v, \nabla v) + \Lambda(v)) dx, \quad (1.1)$$

where  $A$  is an elliptic matrix with Hölder continuous entries, and  $\Lambda(v) = \lambda_+ \chi_{\{v>0\}} + \lambda_- \chi_{\{v\leq 0\}}$ . We prove that if the value of boundary data and its derivative at a point are equal to zero (i.e., it satisfies the (DPT) condition mentioned below), then the contact of free boundary and the fixed boundary is tangential.

Several authors have extended the works of Alt, Caffarelli, and Friedman [2] on free boundary problems with constant coefficients to the case of variable coefficients. For example, the works of Argiolas and Ferrari [4, 14] which extend the seminal works [8, 9] to the case of  $x$  dependent coefficients. See also the recent work of Ferrari and Lederman [15] for the case of variable exponents. For the non-homogenous case with constant coefficients, one can refer to the work of De Silva, Ferrari, and Salsa [10].

Boundary interactions of free boundaries have gained significant attention in recent years. Whenever there are two media involved, the interactions of their respective diffusions can be modeled by free boundary problems. Often, free boundary of solution and

---

2020 *Mathematics Subject Classification.* Primary 49J05; Secondary 35B65, 35Q92, 35Q35.

*Keywords.* Variational calculus, Bernoulli-type free boundary problems, boundary behavior, Alt–Caffarelli–Friedman-type minimizers.

fixed boundary of set come in contact. In applications, the dam problem [3] and jets, wakes, and cavities [5] model phenomena which involve understanding of free boundary and fixed boundary.

Very recently, the works of Indrei [19, 20] study the interactions of free boundaries and fixed boundaries for fully non-linear obstacle problems. We refer to [17], where the authors shed more light into the angle of contact between fixed boundary and free boundary for one-phase Bernoulli problem.

We refer the reader to the work of Kenig, Karakhanyan, and Shahgholian [22, 23] where they deal with constant-coefficient Bernoulli-type free boundary problems. The case of variable coefficients in this paper brings forward some new difficulties (for example, the proof of Proposition 3.9). As it is common by now, our strategy in this article is to classify blowups of minimizers. We prove that the blowups and also their positivity sets converge to a global solution in  $\mathcal{P}_\infty$  (cf. Definition 2.5). In Section 2, we list the assumptions and set some notations, and then in Section 3, we prove that blowups of minimizers converge to those of global solutions (cf. Definition 2.6). In the last section, we prove our main result.

## 2. Setting up the problem

We consider the following class of functions which we denote as  $\mathcal{P}_r(\alpha, M, \lambda, \mathcal{D}, \mu)$ . Before definition, we set the following notations:

$$B_R^+ := \{x \in B_R \text{ such that } x_N > 0\},$$

$$B'_R := \{x \in B_R \text{ such that } x_N = 0\}.$$

For  $x \in \mathbb{R}^N$ , we denote  $x' \in \mathbb{R}^{N-1}$  as the projection of  $x$  on the plane  $\{x_N = 0\}$ ; we denote the tangential gradient  $\nabla'$  as follows:

$$\nabla' u := \left( \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_{N-1}} \right).$$

We define the affine space set  $H_\phi^1(B_R^+)$  as follows:

$$H_\phi^1(B_R^+) = \{v \in H^1(B_R^+) : v - \phi \in H_0^1(B_R^+)\}.$$

For a given function  $v \in H^1(B_2^+)$ , we denote  $F(v)$  as

$$F(v) := \partial\{v > 0\},$$

and  $\text{Id}$  is the notation for  $N \times N$  identity matrix. For a given function  $v$ , we define  $v^\pm$  as

$$v^+ := \max(0, v),$$

$$v^- := \max(0, -v).$$

**Definition 2.1.** A function  $u \in H^1(B_{2/r}^+)$  is said to belong to the class  $\mathcal{P}_r(\alpha, M, \lambda_{\pm}, \mathcal{D}, \mu)$  if there exist symmetric matrix coefficients  $A \in C^\infty(B_{2/r}^+)^{N \times N}$ ,  $\phi \in C^{1,\alpha}(B_{2/r}^+)$ ,  $\lambda_{\pm} > 0$ ,  $0 < \mu < 1$ , and  $\mathcal{D} > 0$  such that the following hold:

(P1)  $\|A\|_{L^\infty(B_{2/r}^+)} \leq M$ ,  $\|\nabla\phi\|_{L^\infty(B_{2/r}^+)} \leq M$ ,  $[A]_{C^\alpha(B_{2/r}^+)}, [\nabla\phi]_{C^{1,\alpha}(B_{2/r}^+)} \leq r^\alpha M$ , and  $|\phi(x')| \leq Mr^{1+\alpha}|x'|^{1+\alpha}$  ( $x' \in B'_{2/r}$ ).  $\phi$  satisfies the following Degenerate Phase Transition condition (DPT):

$$\forall x' \in B'_{2/r} \text{ such that } \phi(x') = 0; \text{ then } |\nabla'\phi(x')| = 0. \quad (\text{DPT})$$

(P2)  $A(0) = \text{Id}$ ,  $\mu|\xi|^2 \leq \langle A(x)\xi, \xi \rangle \leq \frac{1}{\mu}|\xi|^2$  for all  $x \in B_{2/r}^+$  and  $\xi \in \mathbb{R}^N$ .

(P3)  $0 < \lambda_- < \lambda_+$ .

(P4)  $u$  minimizes  $J(\cdot; A, \lambda_+, \lambda_-, B_{2/r}^+)$  (cf. (1.1)); that is, for every  $u - v \in H_0^1(B_{2/r}^+)$ ,

$$\int_{B_{2/r}^+} ((A(x)\nabla u, \nabla u) + \Lambda(u)) dx \leq \int_{B_{2/r}^+} ((A(x)\nabla v, \nabla v) + \Lambda(v)) dx$$

$$(\Lambda(s) = \lambda_+\chi_{\{s>0\}} + \lambda_-\chi_{\{s\leq 0\}}) \quad \text{and} \quad 0 \in F(u) \cap \overline{B_{2/r}^+}.$$

(P5)  $u \in H_\phi^1(B_{2/r}^+)$ .

(P6) There exists  $0 < r_0$  such that for all  $0 < \rho \leq r_0$  we have

$$\frac{|B_\rho^+(0) \cap \{u > 0\}|}{|B_\rho^+(0)|} > \mathcal{D}. \quad (2.1)$$

**Remark 2.2.** In fact, the functions  $u \in \mathcal{P}_r(\alpha, M, \lambda_{\pm}, \mathcal{D}, \mu)$  carry more regularity than being only a Sobolev function. They are Hölder continuous in  $B_{2/r}^+$  (cf. Lemma 3.2).

**Remark 2.3.** We have assumed the coefficient matrices  $A$  to be of the class  $C^\infty(B_2^+)$ . However, all the estimates in this paper depend only on the  $C^\alpha$  norms of  $A$ . Primarily, the reason we imposed the assumption  $A \in C^\infty(B_2^+)^{N \times N}$  is to ensure that we can represent solutions to PDEs  $-\text{div}(A(s)\nabla u) = 0$  in terms of co-normal derivatives of Green’s function (cf. (3.26) and (3.35)) which follows from [21, equation (1.12)]. This representation (in (3.26) and (3.35)) is also true for the case  $A \in C^\alpha(B_2^+)^{N \times N}$ ; a formal proof of this fact will be present in our forthcoming articles. This way, one can replace the assumption  $A \in C^\infty(B_2^+)^{N \times N}$  (in Definition 2.1) by  $A \in C^\alpha(B_2^+)^{N \times N}$ .

**Remark 2.4.** The assumption (P2), i.e.,  $A(0) = \text{Id}$ , does not compromise the generality of the problem. Indeed, assume that  $A(0) \neq \text{Id}$ . Since  $A(x)$  is a symmetric matrix for all  $x \in B_R^+$ , we have

$$J(u; B_R^+) = \int_{B_R^+} ((A(x)\nabla u, \nabla u) + \Lambda(u)) dx$$

$$= \int_{B_R^+} (|A^{1/2}(x)\nabla u(x)|^2 + \Lambda(u)(x)) dx.$$

Under the change of variables  $y \rightarrow A(0)^{-1/2}x$ , the functional  $J(\cdot; B_R^+)$  is transformed into

$$J(u; B_R^+) = |\det(A(0))|^{1/2} \int_{B_R^+} (|\mathcal{A}^{1/2}(y)\nabla v|^2 + \Lambda(v)) dy,$$

where

$$\begin{aligned} \mathcal{A}^{1/2}(y) &:= A^{1/2}(A(0)^{1/2}y)A(0)^{-1/2} \quad (\text{note that } \mathcal{A}(0) = \text{Id}), \\ v(y) &:= u(A(0)^{1/2}y), \\ B_R^+ &:= A(0)^{-1/2}B_R^+. \end{aligned}$$

Then, we can reformulate the minimization problem to minimize the functional

$$\mathcal{J}(v; B_R^+) := \int_{B_R^+} (|\mathcal{A}^{1/2}(y)\nabla v|^2 + \Lambda(v)) dy = \int_{B_R^+} (\langle \mathcal{A}(y)\nabla v, \nabla v \rangle + \Lambda(v)) dy.$$

Moreover, under the (linear) transformation  $x \rightarrow A(0)^{-1/2}x$ , all the assumptions (P1)–(P6) remain structurally unchanged.

In the absence of ambiguity on values of  $\alpha, M, \lambda_{\pm}, \mathcal{D}$ , and  $\mu$ , we use the notation  $\mathcal{P}_r$  in place of  $\mathcal{P}_r(\alpha, M, \lambda_{\pm}, \mathcal{D}, \mu)$ . If  $\phi \in C^{1,\alpha}(B_2^+)$  satisfies (DPT), from [7, Lemma 10.1], we know that  $\phi^{\pm}|_{B'_2} \in C^{1,\alpha}(B'_2)$  and also

$$\|\phi^{\pm}\|_{C^{1,\alpha}(B'_2)} \leq \|\phi\|_{C^{1,\alpha}(B'_2)}.$$

Given  $v \in H^1(B_{R/r}^+)$  and  $r > 0$ , we define the blowup  $v_r \in H^1(B_{R/r}^+)$  as follows:

$$v_r(x) := \frac{1}{r}v(rx). \tag{2.2}$$

For the coefficient matrix  $A$ ,  $A^r(x)$  is defined as follows:

$$A^r(x) := A(rx).$$

One can check that if  $u \in \mathcal{P}_1(\alpha, M, \lambda_{\pm}, \mathcal{D}, \mu)$ , then  $u_r \in \mathcal{P}_r(\alpha, M, \lambda_{\pm}, \mathcal{D}, \mu)$ . Indeed if  $u \in \mathcal{P}_1$  and  $u$  is a minimizer of the functional  $J$  (cf. (P4))

$$\begin{aligned} J(v; A, \lambda_+, \lambda_-, B_2^+) &:= \int_{B_2^+} (\langle A(x)\nabla v, \nabla v \rangle + \Lambda(v)) dx, \\ (\Lambda(s)) &= \lambda_+\chi_{\{s>0\}} + \lambda_-\chi_{\{s\leq 0\}} \end{aligned}$$

with boundary data  $\phi \in C^{1,\alpha}(B_2^+)$  (i.e.,  $u \in H^1_{\phi}(B_2^+)$ ), then, by simple change of variables, we can check that  $u_r \in H^1_{\phi_r}(B_{2/r}^+)$  (this verifies (P5)) and  $u_r$  minimizes

$$\begin{aligned} J(v; A^r, \lambda_+, \lambda_-, B_{2/r}^+) &:= \int_{B_{2/r}^+} (\langle A^r(x)\nabla v, \nabla v \rangle + \Lambda(v)) dx, \\ (\Lambda(s)) &= \lambda_+\chi_{\{s>0\}} + \lambda_-\chi_{\{s\leq 0\}}. \end{aligned}$$

Moreover, if  $A$  and  $\phi$  satisfy conditions (P1) and (P2) for  $r = 1$ , then  $A^r$  and  $\phi_r$  satisfy (P1) and (P2) for  $r$ . (P3) and (P6) remain invariant under the change of variables. Therefore,  $u_r \in \mathcal{P}_r$ .

In order to study the blowup limits ( $\lim_{r \rightarrow 0} u_r$ ) of functions  $u \in \mathcal{P}_1(\alpha, M, \lambda_{\pm}, \mathcal{D}, \mu)$ , we define a class of global solutions  $\mathcal{P}_{\infty}(C, \lambda_{\pm})$ . Let us set the following notation before giving the definition:

$$\Pi := \{x \in \mathbb{R}^N : x_N = 0\}.$$

**Definition 2.5** (Global solution). We say that  $u \in H^1(\mathbb{R}_+^N)$  belongs to the class of functions  $\mathcal{P}_{\infty}(C, \lambda_{\pm})$ ; that is,  $u$  is a global solution if there exist  $C > 0$  and  $0 < \lambda_+ < \lambda_-$  such that

- (G1)  $|u(x)| \leq C|x|$  for all  $x \in \mathbb{R}_+^N$ ,
- (G2)  $u$  is continuous up to the boundary  $\Pi$ ,
- (G3)  $u = 0$  on  $\Pi$ ,
- (G4) and for every ball  $B_r(x_0)$ ,  $u$  is a minimizer of  $J(\cdot; \text{Id}, \lambda_+, \lambda_-, B_r(x_0) \cap \mathbb{R}_+^N)$  (cf. (1.1)); that is,

$$\int_{B_r(x_0) \cap \mathbb{R}_+^N} (|\nabla u|^2 + \Lambda(u)) \, dx \leq \int_{B_r(x_0) \cap \mathbb{R}_+^N} (|\nabla v|^2 + \Lambda(v)) \, dx.$$

Here,  $(\Lambda(s) = \lambda_+ \chi_{\{s>0\}} + \lambda_- \chi_{\{s \leq 0\}})$  and for every  $v \in H^1(B_r(x_0) \cap \mathbb{R}_+^N)$  such that  $u - v \in H_0^1(B_r(x_0) \cap \mathbb{R}_+^N)$ .

Our main result intends to show that for a minimizer  $u$  of  $J(\cdot; A, \lambda_+, \lambda_-, B_2^+)$  with  $A, \lambda_{\pm}$ , and  $u$  satisfying the properties (P1)–(P6), the free boundary of every such minimizer touches the flat part of fixed boundary tangentially at the origin. For this, we prove that as we approach closer and closer to the origin, the free boundary points cannot lie completely inside any cone which is perpendicular to the flat boundary and has its tip at the origin. The main result in this paper is stated below.

**Theorem 2.6.** *There exist a constant  $\rho_0$  and a modulus of continuity  $\sigma$  such that if*

$$u \in \mathcal{P}_1(\alpha, M, \lambda_{\pm}, \mathcal{D}, \mu),$$

then

$$F(u) \cap B_{\rho_0}^+ \subset \{x : x_N \leq \sigma(|x|)|x|\}.$$

Here,  $\sigma$  depends only on  $\alpha, M, \lambda_{\pm}, \mathcal{D}, \mu$ .

### 3. Blowup analysis

The following is a classical result (cf. [1, Remark 4.2]); we present the proof for the case of variable coefficients.

**Lemma 3.1.** *Given a bounded and strictly elliptic matrix  $A(x)$  and a non-negative continuous function  $w$  such that  $\operatorname{div}(A(x)\nabla w) = 0$  in  $\{w > 0\} \cap B_2^+$  in weak sense (i.e., in the distributional sense in the context of Sobolev spaces), then  $w \in H_{\text{loc}}^1(B_2^+)$  and  $\operatorname{div}(A(x)\nabla w) \geq 0$  in weak sense in  $B_2^+$ .*

*Proof.* Let  $D \Subset B_2^+$  and  $\eta \in C_c^\infty(B_2^+)$  be cutoff function for  $D$ . That is,  $\eta \in C_c^\infty(B_2^+)$  be such that

$$\eta(x) = \begin{cases} 1 & \text{in } D, \\ 0 & \text{on } \partial B_2^+. \end{cases}$$

Since  $\operatorname{div}(A(x)\nabla w) = 0$  in  $\{w > 0\} \cap B_2^+$ , we have

$$\begin{aligned} 0 &= \int_{B_2^+} \langle A(x)\nabla w, \nabla((w - \varepsilon)^+ \eta^2) \rangle dx \\ &= \int_{B_2^+ \cap \{w > \varepsilon\}} \eta^2 \langle A(x)\nabla w, \nabla w \rangle dx + \int_{B_2^+ \cap \{w > \varepsilon\}} w \langle A(x)\nabla w, \nabla \eta^2 \rangle dx \\ &\quad - \varepsilon \int_{B_2^+ \cap \{w > \varepsilon\}} \langle A(x)\nabla w, \nabla \eta^2 \rangle dx \end{aligned}$$

which implies

$$\begin{aligned} \int_{B_2^+ \cap \{w > \varepsilon\}} \langle A(x)\nabla w, \nabla w \rangle \eta^2 dx &\leq \int_{B_2^+ \cap \{w > \varepsilon\}} |w \langle A(x)\nabla w, \nabla \eta^2 \rangle| dx \\ &\quad + \varepsilon \int_{B_2^+ \cap \{w > \varepsilon\}} |\langle A(x)\nabla w, \nabla \eta^2 \rangle| dx. \end{aligned}$$

By the choice of  $\eta$  and ellipticity of the matrix  $A$ , we obtain, using Young’s inequality,

$$\begin{aligned} &\mu \int_{B_2^+ \cap \{w > \varepsilon\}} |\nabla w|^2 \eta^2 dx \\ &\leq \frac{1}{\mu} \left| \int_{B_2^+ \cap \{w > \varepsilon\}} \eta w |\nabla w| |\nabla \eta| dx \right| + \frac{\varepsilon}{\mu} \left| \int_{B_2^+ \cap \{w > \varepsilon\}} \eta |\nabla w| |\nabla \eta| dx \right| \\ &\leq C_1(\mu) \left[ \frac{1}{\delta} \int_{B_2^+ \cap \{w > \varepsilon\}} w^2 |\nabla \eta|^2 dx + \delta \int_{B_2^+ \cap \{w > \varepsilon\}} \eta^2 |\nabla w|^2 dx \right. \\ &\quad \left. + \delta \varepsilon \int_{B_2^+ \cap \{w > \varepsilon\}} \eta^2 |\nabla w|^2 dx + \frac{\varepsilon}{\delta} \int_{B_2^+ \cap \{w > \varepsilon\}} |\nabla \eta|^2 dx \right]. \end{aligned}$$

After choosing  $\delta > 0$  very small and rearranging the terms in the equation above, since  $\eta = 1$  in  $D$ , we finally get

$$\begin{aligned} \int_{D \cap \{w > \varepsilon\}} |\nabla w|^2 dx &\leq \int_{B_2^+ \cap \{w > \varepsilon\}} |\nabla w|^2 \eta^2 dx \\ &\leq C(\mu) \left[ \int_{B_2^+ \cap \{w > \varepsilon\}} w^2 |\nabla \eta|^2 dx + \int_{B_2^+ \cap \{w > \varepsilon\}} |\nabla \eta|^2 dx \right]. \end{aligned}$$

As  $\varepsilon \rightarrow 0$ , we obtain

$$\begin{aligned} \int_D |\nabla w|^2 dx &= \int_{\{w>0\} \cap D} |\nabla w|^2 dx = \lim_{\varepsilon \rightarrow 0} \int_{\{w>\varepsilon\} \cap D} |\nabla w|^2 \\ &\leq C(\mu) \lim_{\varepsilon \rightarrow 0} \left[ \int_{B_2^+ \cap \{w>\varepsilon\}} w^2 |\nabla \eta|^2 dx + \int_{B_2^+ \cap \{w>\varepsilon\}} |\nabla \eta|^2 dx \right] \\ &\leq C(\mu, D) \left[ \int_{B_2^+ \cap \text{supp}(\eta)} w^2 dx + 1 \right]. \end{aligned}$$

Since  $w \in C(B_2^+)$ , therefore  $w$  is uniformly bounded in  $\text{supp}(\eta)$ , and therefore,

$$\int_D (|\nabla w|^2 + |w^2|) dx \leq C(\mu, D) \left[ \int_{B_2^+ \cap \text{supp}(\eta)} w^2 dx + 1 \right] \leq C(\mu, D, \|w\|_{L^\infty(\text{supp}(\eta))}).$$

Now, for  $0 \leq \varphi \in C_c^\infty(B_2^+)$ , consider the test function

$$\begin{aligned} v &= \varphi \left( 1 - \left( \min \left( 2 - \frac{w}{\varepsilon}, 1 \right) \right)^+ \right) \\ \int_{B_2^+} \langle A(x) \nabla w, \nabla \varphi \rangle dx &= \int_{B_2^+} \left\langle A(x) \nabla w, \nabla \left( \varphi \left( \left( 2 - \frac{w}{\varepsilon} \right) \wedge 1 \right)^+ \right) \right\rangle dx. \end{aligned}$$

We can easily check that  $v \geq 0$  in  $B_2^+$  and  $v \in H_0^1(B_2^+)$ ; in particular,

$$\varphi \left( \left( 2 - \frac{w}{\varepsilon} \right) \wedge 1 \right)^+ = \begin{cases} \varphi, & x \in \{w \leq \varepsilon\}, \\ \varphi \cdot \left( 2 - \frac{w}{\varepsilon} \right), & x \in \{\varepsilon < w \leq 2\varepsilon\}, \\ 0, & x \in \{w > 2\varepsilon\}. \end{cases}$$

Therefore, we have

$$\begin{aligned} &\int_{B_2^+} \langle A(x) \nabla w, \nabla \varphi \rangle dx \\ &= \int_{B_2^+} \left\langle A(x) \nabla w, \nabla \left( \varphi \left( \left( 2 - \frac{w}{\varepsilon} \right) \wedge 1 \right)^+ \right) \right\rangle dx \\ &= \int_{B_2^+ \cap \{w \leq \varepsilon\}} \langle A(x) \nabla w, \nabla \varphi \rangle dx + \int_{B_2^+ \cap \{\varepsilon < w \leq 2\varepsilon\}} \left\langle A(x) \nabla w, \nabla \left( \varphi \left( 2 - \frac{w}{\varepsilon} \right) \right) \right\rangle dx \\ &= \int_{B_2^+ \cap \{w \leq \varepsilon\}} \langle A(x) \nabla w, \nabla \varphi \rangle dx + 2 \int_{B_2^+ \cap \{\varepsilon < w \leq 2\varepsilon\}} \langle A(x) \nabla w, \nabla \varphi \rangle dx \\ &\quad - \frac{2}{\varepsilon} \int_{B_2^+ \cap \{\varepsilon < w \leq 2\varepsilon\}} \langle A(x) \nabla w, \nabla(w\varphi) \rangle dx \\ &\leq C(\mu) \int_{B_2^+ \cap \{\varepsilon < w \leq 2\varepsilon\}} |\nabla w| |\nabla \varphi| dx + \frac{2}{\varepsilon} \int_{B_2^+ \cap \{\varepsilon < w \leq 2\varepsilon\}} w \langle A(x) \nabla w, \nabla \varphi \rangle dx \\ &\quad - \frac{2}{\varepsilon} \int_{B_2^+ \cap \{\varepsilon < w \leq 2\varepsilon\}} \varphi \langle A(x) \nabla w, \nabla w \rangle dx \\ &\leq C(\mu) \int_{B_2^+ \cap \{\varepsilon < w \leq 2\varepsilon\}} |\nabla w| |\nabla \varphi| dx. \end{aligned}$$

The last term goes to zero as  $\varepsilon \rightarrow 0$ . Therefore, we can say that

$$\int_{B_2^+} \langle A(x)\nabla w, \nabla\varphi \rangle dx \leq 0.$$

We conclude the proof from [13, Theorem 1.39] (see also [24, Theorem 6.22]). ■

**Lemma 3.2** (Hölder continuity). *If  $u \in \mathcal{P}_1$ , then  $u \in C^{0,\alpha_0}(\overline{B_2^+})$  for some  $0 < \alpha_0 < 1$ . In fact,*

$$\|u\|_{C^{\alpha_0}(B_2^+)} \leq C(\mu, \lambda_{\pm})\|u\|_{L^\infty(B_2^+)}.$$

*Proof.* The functional  $J(\cdot; A, \lambda_+, \lambda_-, B_2^+)$  satisfies the hypothesis of [16, Theorem 7.3] and  $\phi \in C^{1,\alpha}(\overline{B_2^+})$ ; therefore, Lemma 3.2 follows from the arguments in [16, Section 7.8], where the boundary regularity is treated. ■

**Remark 3.3.** Since every function  $u \in \mathcal{P}_1$  is continuous, therefore the positivity set  $\{u > 0\}$  is an open set.

**Corollary 3.4.** *If  $u \in \mathcal{P}_1$ , then  $u^\pm$  are  $A$ -subharmonic.*

*Proof.* The claim follows directly from Lemmas 3.2 and 3.1. ■

**Lemma 3.5.** *If  $u \in \mathcal{P}_1$ , then*

$$|u(x)| \leq C(\mu, [A]_{C^\alpha(B_2^+)}, N)M|x| \quad \text{in } B_1^+. \tag{3.1}$$

*Proof.* Let  $w$  be such that

$$\begin{cases} \operatorname{div}(A(x)w) = 0 & \text{in } B_2^+, \\ w = \phi^+ & \text{in } \partial B_2^+. \end{cases}$$

Since  $u$  is  $A$ -subharmonic in  $B_2^+$  (cf. Corollary 3.4), by maximum principle, if  $x \in B_1^+$  we have

$$\begin{aligned} u^+(x) &\leq w(x) \leq w(x) - w(x') + w(x') \\ &\leq \|\nabla w\|_{L^\infty(B_1^+)}|x - x'| + |\phi^+(x')| \\ &= \|\nabla w\|_{L^\infty(B_1^+)}x_N + |\phi^+(x')| \\ &\leq (\|\nabla w\|_{L^\infty(B_1^+)} + M)|x| \quad \forall x \in B_1^+. \end{aligned} \tag{3.2}$$

In the last inequality, we have used (P1). Now, we prove that the term  $\|\nabla w\|_{L^\infty(B_1^+)}$  is uniformly bounded.

From [6, Theorem 2], we have

$$\|\nabla w\|_{L^\infty(B_1^+)} \leq C(\mu, [A]_{C^\alpha(B_2^+)}, N)[\|w\|_{L^\infty(B_2^+)} + \|\phi^+\|_{C^{1,\alpha}(B_2^+)}]. \tag{3.3}$$



By comparison principle,

$$\|w\|_{L^\infty(B_2^+)} = \|w\|_{L^\infty(\partial B_2^+)} = \|\phi^+\|_{L^\infty(\partial B_2^+)} \leq M.$$

Plugging this information in (3.3), we get

$$\|\nabla w\|_{L^\infty(B_1^+)} \leq C(\mu, [A]_{C^\alpha(B_2^+)}, N)M,$$

and then using (3.2), we obtain

$$u^+(x) \leq C(\mu, [A]_{C^\alpha(B_2^+)}, N)M|x| \quad \forall x \in B_1^+. \tag{3.4}$$

And analogously,

$$u^-(x) \leq C(\mu, [A]_{C^\alpha(B_2^+)}, N)M|x| \quad \forall x \in B_1^+. \tag{3.5}$$

By adding (3.4) and (3.5), we prove (3.1). ■

**Remark 3.6.** We can check that, for every  $u \in \mathcal{P}_1$ ,  $u_r \in \mathcal{P}_r$  and  $u_r$  is  $A^r$ -subharmonic and satisfies (3.1) in  $B_{1/r}^+$ . That is,

$$|u_r(x)| \leq C(\mu, [A]_{C^\alpha(B_2^+)}, N)M|x|, \quad x \in B_{1/r}^+.$$

**Lemma 3.7** (Uniform  $H^1$  bounds). *Let  $u \in \mathcal{P}_1$ . Then, for  $R > 0$  such that  $2R \leq \frac{2}{r}$ , we have*

$$\int_{B_R^+} |\nabla u_r|^2 dx \leq C(N, \lambda, \mu, R, M).$$

*Proof.* Since  $u_r \in \mathcal{P}_r$ , from (P4), we can say that  $u_r$  is a minimizer of  $J(\cdot; A^r, \lambda_\pm, B_{2R}^+)$  with boundary data  $\phi_r$ . Here,  $A^r$  and  $\phi_r$  satisfy conditions (P1) and (P2). Precisely speaking,  $u_r$  is the minimizer of the following functional:

$$J(v; A^r, \lambda_\pm, B_{2R}^+) := \int_{B_{2R}^+} (\langle A^r(x)\nabla v, \nabla v \rangle + \Lambda(v)) dx.$$

Here,  $(\Lambda(v) = \lambda_+\chi_{\{v>0\}} + \lambda_-\chi_{\{v\leq 0\}})$ . Consider  $h \in H^1(B_{2R}^+)$  as a harmonic replacement

$$\begin{cases} \operatorname{div}(A^r(x)\nabla h) = 0 & \text{in } B_{2R}^+ \\ h - u_r \in H_0^1(B_{2R}^+). \end{cases}$$

In other words,  $h$  is the minimizer of  $\int_{B_{2R}^+} \langle A^r(x)\nabla h, \nabla h \rangle dx$  in the set  $H_{u_r}^1(B_{2R}^+)$ .

From the minimality of  $u_r$  and the choice of  $h$ , we have

$$\begin{aligned} & \int_{B_{2R}^+} \langle A^r(x)\nabla(u_r - h), \nabla(u_r - h) \rangle dx \\ &= \int_{B_{2R}^+} \langle A^r(x)\nabla(u_r - h), \nabla(u_r + h - 2h) \rangle dx \end{aligned}$$

$$\begin{aligned}
 &= \int_{B_{2R}^+} \langle A^r(x) \nabla(u_r - h), \nabla(u_r + h) \rangle dx - 2 \int_{B_{2R}^+} \langle A^r(x) \nabla(u_r - h), \nabla h \rangle dx \\
 &= \int_{B_{2R}^+} ((A^r(x) \nabla u_r, \nabla u_r) - \langle A^r(x) \nabla h, \nabla h \rangle) dx \quad (\text{since } h \text{ is } A^r\text{-harmonic in } B_R^+) \\
 &\leq \int_{B_{2R}^+} (\Lambda(h) - \Lambda(u_r)) dx \leq C(N, \lambda, R).
 \end{aligned}$$

Using the ellipticity of  $A$ , we get

$$\int_{B_R^+} |\nabla(u_r - h)|^2 dx \leq \int_{B_{2R}^+} |\nabla(u_r - h)|^2 dx \leq C(N, \lambda, \mu, R).$$

Expanding the left-hand side, we get

$$\begin{aligned}
 \int_{B_R^+} |\nabla u_r|^2 dx &\leq \int_{B_R^+} |\nabla u_r|^2 dx + \int_{B_R^+} |\nabla h|^2 dx \\
 &\leq C(N, \lambda, \mu, R) + 2 \int_{B_R^+} \nabla u_r \cdot \nabla h dx \\
 &\leq C(N, \lambda, \mu, R) + \varepsilon \int_{B_R^+} |\nabla u_r|^2 dx + \frac{1}{\varepsilon} \int_{B_R^+} |\nabla h|^2 dx.
 \end{aligned}$$

By choosing  $\varepsilon = \frac{1}{8}$ , we are left with

$$\int_{B_R^+} |\nabla u_r|^2 dx \leq C(N, \lambda, \mu, R) \left( 1 + \int_{B_R^+} |\nabla h|^2 dx \right).$$

To close the argument, we need to show that  $\int_{B_R^+} |\nabla h|^2 dx$  is uniformly bounded, indeed from [6, Theorem 2],  $\|\nabla h\|_{L^\infty(B_R^+)} \leq C(\mu, M)$ . ■

**Lemma 3.8** (Compactness). *Let  $r_j \rightarrow 0^+$ , and a sequence  $\{v_j\} \in \mathcal{P}_1$ . Then, the blowups  $u_j := (v_j)_{r_j}$  (as defined in (2.2)) converge (up to subsequence) uniformly in  $B_R^+$  and weakly in  $H^1(B_R^+)$  to some limit for any  $R > 0$ . Moreover, if  $u_0$  is such a limit of  $u_j$  in the above-mentioned topologies, then  $u_0$  belongs to  $\mathcal{P}_\infty$ .*

*Proof.* We fix  $R > 0$ ; since  $v_j \in \mathcal{P}_1$ , therefore  $u_j \in \mathcal{P}_{r_j}$ , and as argued in the proof of previous lemma, the functions  $u_j$  are minimizers of the functional  $J(\cdot; A_j, \lambda_\pm, B_R^+)$  for  $j$  sufficiently large that  $R < \frac{1}{r_j}$ . We set the notation for the functional  $J_j$  as

$$J_j(v) := J(v; A_j, \lambda_\pm, B_R^+) = \int_{B_R^+} ((A_j(x) \nabla v, \nabla v) + \Lambda(v)) dx, \quad v \in H_{u_j}^1(B_R^+).$$

We require to rescale the boundary data in the same way as we do to  $v_j$ , we also denote the boundary values for  $u_j \in \mathcal{P}_{r_j}$  as  $\phi_j$ . Here, the sequences  $A_j \in C^\alpha(B_{2/r_j}^+)^{N \times N}$  and  $\phi_j \in C^{1,\alpha}(B_{2/r_j})$  satisfy conditions (P1) and (P2) with

$$r = r_j, \quad \Lambda(v) := \lambda_+ \chi_{\{v>0\}} + \lambda_- \chi_{\{v \leq 0\}}.$$

We set the following notation for the functional  $J_0$ :

$$J_0(v; B_R^+) := \int_{B_R^+} (|\nabla v|^2 + \Lambda(v)) \, dx. \tag{3.6}$$

From Lemma 3.2, we know that  $u_j \in C^{\alpha_0}(\overline{B_{2/r_j}^+})$  which implies  $C^{\alpha_0}(\overline{B_R^+})$ . In particular,  $\|u_j\|_{C^{0,\alpha_0}(B_R^+)} \leq C(\mu, \lambda_{\pm})$ . Hence,  $u_j$  is a uniformly bounded and equicontinuous sequence in  $\overline{B_R^+}$ ; we can apply Arzela–Ascoli theorem to show that  $u_j$  uniformly converges to a function  $u_0 \in C^{0,\alpha_0}(\overline{B_R^+})$ .

Since  $u_j = \phi_j$  on  $B'_R$ , from (P1), we have  $|\phi_j(x)| \leq M r_j^{1+\alpha} |x|^{1+\alpha}$  for  $x \in B'_R$ ; therefore,  $|\phi_j(x)| \leq C(M, \alpha) r_j^{1+\alpha} R^{1+\alpha}$ . Hence,  $\phi_j \rightarrow 0$  uniformly on  $B'_R$ . We have

$$u_0 = \lim_{j \rightarrow \infty} u_j = \lim_{j \rightarrow \infty} \phi_j = 0 \quad \text{on } B'_R.$$

Thus,  $u_0$  satisfies (G2) and (G3) inside the domain  $\overline{B_R^+}$ . Also, from Lemma 3.7, we have

$$\int_{B_R^+} |\nabla u_j|^2 \, dx \leq C(N, \lambda_{\pm}, \mu, R, M, \alpha). \tag{3.7}$$

Then, by the linear growth condition (cf. Remark 3.6),  $u_j$  also satisfies

$$|u_j(x)| \leq C(\mu, \alpha) M |x| \quad x \in B_R^+.$$

Hence, passing to the limit, we have  $|u_0(x)| \leq C(\mu, \alpha) M |x|, \forall x \in B_R^+$ . In other words,  $u_0$  satisfies (G1) in  $\overline{B_R^+}$ . Moreover, we have

$$\int_{B_R^+} |u_j|^2 \, dx \leq C(\mu, \alpha, M) \int_{B_R^+} |x|^2 \, dx \leq C(\mu, \alpha, M, N, R). \tag{3.8}$$

Thus, (3.7) and (3.8) imply that  $u_j$  is a bounded sequence in  $H^1(B_R^+)$ . Hence, up to a subsequence,  $u_j \rightharpoonup u_0$  weakly in  $H^1(B_R^+)$ . We rename the subsequence again as  $u_j$ .

We have found a blowup limit up to a subsequence  $u_0$  and have shown that  $u_0$  satisfies (G1), (G2), and (G3) in  $\overline{B_R^+}$ . In order to show that  $u_0 \in \mathcal{P}_{\infty}$ , it only remains to verify that  $u_0$  satisfies (G4); i.e.,  $u_0$  is a local minimizer of  $J_0(\cdot; B_R^+)$  for all  $R > 0$  (cf. (3.6)). For that, we first claim that

$$\int_{B_R^+} (|\nabla u_0|^2 + \Lambda(u_0)) \, dx \leq \liminf_{j \rightarrow \infty} \int_{B_R^+} (\langle A_j(x) \nabla u_j, \nabla u_j \rangle + \Lambda(u_j)) \, dx. \tag{3.9}$$

Indeed, let us look separately at the term  $J_j(u_j)$  on the right-hand side of the above equation

$$J_j(u_j) = \int_{B_R^+} (\langle A_j(x) \nabla u_j, \nabla u_j \rangle + \lambda_+ \chi_{\{u_j > 0\}} + \lambda_- \chi_{\{u_j \leq 0\}}) \, dx.$$

We rewrite the first term as follows:

$$\int_{B_R^+} \langle A_j(x) \nabla u_j, \nabla u_j \rangle dx = \int_{B_R^+} \langle (A_j(x) - \text{Id}) \nabla u_j, \nabla u_j \rangle dx + \int_{B_R^+} |\nabla u_j|^2 dx. \tag{3.10}$$

From (P1) and (P2), we have for all  $x \in B_R^+$

$$\|A_j(x) - \text{Id}\|_{L^\infty(B_R^+)} \leq M r_j^\alpha |x|^\alpha \leq C(M, R, \alpha) r_j^\alpha \rightarrow 0 \quad \text{as } j \rightarrow \infty.$$

Therefore,  $A_j \rightarrow \text{Id}$  uniformly and  $\|\nabla u_j\|_{L^2(B_2^+)}$  is bounded (cf. (3.7)). Hence, the first term on the right-hand side of (3.10) tends to zero as  $j \rightarrow \infty$ . Thus, from (3.10) and by weak lower semi-continuity of  $H^1$  norm, we have

$$\int_{B_R^+} |\nabla u_0|^2 dx \leq \liminf_{j \rightarrow \infty} \int_{B_R^+} |\nabla u_j|^2 dx = \liminf_{j \rightarrow \infty} \int_{B_R^+} \langle A_j(x) \nabla u_j, \nabla u_j \rangle dx. \tag{3.11}$$

For the second term, we claim that

$$\int_{B_R^+} \lambda_+ \chi_{\{u_0 > 0\}} + \lambda_- \chi_{\{u_0 \leq 0\}} dx \leq \liminf_{j \rightarrow \infty} \int_{B_R^+} \lambda_+ \chi_{\{u_j > 0\}} + \lambda_- \chi_{\{u_j \leq 0\}} dx. \tag{3.12}$$

To see this, we first show that, for almost every  $x \in B_R^+$ , we have

$$\lambda_+ \chi_{\{u_0 > 0\}}(x) + \lambda_- \chi_{\{u_0 \leq 0\}}(x) \leq \liminf_{j \rightarrow \infty} (\lambda_+ \chi_{\{u_j > 0\}}(x) + \lambda_- \chi_{\{u_j \leq 0\}}(x)). \tag{3.13}$$

Indeed, let  $x_0 \in B_R^+ \cap (\{u_0 > 0\} \cup \{u_0 < 0\})$ . Then, by the uniform convergence of  $u_j$  to  $u_0$ , we can easily see that  $u_j(x_0)$  attains the sign of  $u_0(x_0)$  for sufficiently large value of  $j$ . Hence, (3.13) holds in  $\{u_0 > 0\} \cup \{u_0 < 0\}$ .

Now, assume that  $x_0 \in B_R^+ \cap \{u_0 = 0\}$ . Then, the left-hand side of (3.13) is equal to

$$\lambda_+ \chi_{\{u_0 > 0\}}(x_0) + \lambda_- \chi_{\{u_0 \leq 0\}}(x_0) = \lambda_-.$$

Regarding the right-hand side of (3.13), we see that

$$\lambda_+ \chi_{\{u_j > 0\}}(x_0) + \lambda_- \chi_{\{u_j \leq 0\}}(x_0) = \begin{cases} \lambda_+ & \text{if } u_j(x_0) > 0, \\ \lambda_- & \text{if } u_j(x_0) \leq 0. \end{cases}$$

Since  $\lambda_- < \lambda_+$  (cf. (P3)), the right-hand side in the equation above is always greater than or equal to  $\lambda_-$ . Then,

$$\lambda_+ \chi_{\{u_0 > 0\}}(x_0) + \lambda_- \chi_{\{u_0 \leq 0\}}(x_0) = \lambda_- \leq \liminf_{j \rightarrow \infty} (\lambda_+ \chi_{\{u_j > 0\}}(x_0) + \lambda_- \chi_{\{u_j \leq 0\}}(x_0)).$$

Thus, (3.13) is proven for all  $x \in B_R^+$ , and hence, (3.12) holds by Fatou’s lemma.

By adding (3.11) and (3.12) and [12, Theorem 3.127], we obtain (3.9). Now, we will use (3.9) to prove the minimality of  $u_0$  for the functional  $J_0(\cdot; B_R^+)$  (cf. 3.6).




Figure 1. Curvy line represents the free boundary of  $w$ .

Pick any  $w \in H^1(B_R^+)$  such that  $u_0 - w \in H_0^1(B_R^+)$ . We construct an admissible competitor  $w_j^\delta$  to compare the minimality of  $u_j$  for the functional  $J_j(\cdot; B_R^+)$ . Then, we intend to use (3.9).

In this direction, we define two cutoff functions  $\eta_\delta : \mathbb{R}^N \rightarrow \mathbb{R}$  and  $\theta : \mathbb{R} \rightarrow \mathbb{R}$  as follows:

$$\eta_\delta(x) := \begin{cases} 1, & x \in B_{R-\delta}, \\ 0, & x \in \mathbb{R}^N \setminus B_R, \end{cases} \quad \theta(t) := \begin{cases} 1, & |t| \leq 1/2, \\ 0, & |t| \geq 1. \end{cases}$$

We can take  $|\nabla \eta_\delta| \leq \frac{C(N)}{\delta}$ . We define  $\theta_j(x) = \theta(\frac{x_N}{d_j})$  for a sequence  $d_j \rightarrow 0$ , which will be suitably chosen in later steps of the proof. Let  $w_j^\delta$  be a test function defined as

$$w_j^\delta := w + (1 - \eta_\delta)(u_j - u_0) + \eta_\delta \theta_j \phi_j.$$

Since the function  $w_j^\delta - u_j$  is continuous in  $\overline{B_R^+}$  and is pointwise equal to zero on  $\partial B_R^+$ , which is a Lipschitz surface in  $\mathbb{R}^N$ , therefore  $u_j - w_j^\delta \in H_0^1(B_R^+)$ . For further steps, the reader can refer to Figure 1.

Let  $\Omega_{\delta,j} = B_R^+ \cap \{\theta_j = 0\} \cap \{\eta_\delta = 1\}$ , and  $\mathcal{R}_{\delta,j} = B_R^+ \setminus \Omega_{\delta,j}$ ; by observing that  $w_j^\delta = w$  on  $\Omega_{\delta,j}$ , we see that

$$\begin{aligned} |\{w_j^\delta > 0\} \cap B_R^+| &= |\{w_j^\delta > 0\} \cap \Omega_{\delta,j}| + |\{w_j^\delta > 0\} \cap \mathcal{R}_{\delta,j}| \\ &= |\{w > 0\} \cap \Omega_{\delta,j}| + |\{w_j^\delta > 0\} \cap \mathcal{R}_{\delta,j}| \\ &= |\{w > 0\} \cap (B_R^+ \setminus \mathcal{R}_{\delta,j})| + |\{w_j^\delta > 0\} \cap \mathcal{R}_{\delta,j}| \\ &= |\{w > 0\} \cap B_R^+| - |\{w > 0\} \cap \mathcal{R}_{\delta,j}| + |\{w_j^\delta > 0\} \cap \mathcal{R}_{\delta,j}|. \end{aligned}$$

From the above discussions, we have

$$|\{w > 0\} \cap B_R^+| - |\mathcal{R}_{\delta,j}| \leq |\{w_j^\delta > 0\} \cap B_R^+| \leq |\{w > 0\} \cap B_R^+| + |\mathcal{R}_{\delta,j}|.$$

Since we know that  $\lim_{\delta \rightarrow 0} (\lim_{j \rightarrow \infty} |\mathcal{R}_{\delta,j}|) = 0$ , we have

$$\lim_{\delta \rightarrow 0} \left( \lim_{j \rightarrow \infty} |\{w_j^\delta > 0\} \cap B_R^+| \right) = |\{w > 0\} \cap B_R^+|, \tag{3.14}$$

and similarly,

$$\lim_{\delta \rightarrow 0} \left( \lim_{j \rightarrow \infty} |\{w_j^\delta \leq 0\} \cap B_R^+| \right) = |\{w \leq 0\} \cap B_R^+|. \tag{3.15}$$

Given  $u_j \in \mathcal{P}_{r_j}$  and  $w_j^\delta - u_j \in H_0^1(B_R^+)$ , from the minimality of  $u_j$  for the functional  $J_j$ , we have

$$\begin{aligned} & \int_{B_R^+} \left( \langle A_j(x) \nabla u_j, \nabla u_j \rangle + \lambda_+ \chi_{\{u_j > 0\}} + \lambda_- \chi_{\{u_j \leq 0\}} \right) dx \\ & \leq \int_{B_R^+} \left( \langle A_j(x) \nabla w_j^\delta, \nabla w_j^\delta \rangle + \lambda_+ \chi_{\{w_j^\delta > 0\}} + \lambda_- \chi_{\{w_j^\delta \leq 0\}} \right) dx, \end{aligned}$$

and from (3.9), we obtain

$$\begin{aligned} & \int_{B_R^+} \left( |\nabla u_0|^2 + \lambda_+ \chi_{\{u_0 > 0\}} + \lambda_- \chi_{\{u_0 \leq 0\}} \right) dx \\ & \leq \liminf_{j \rightarrow \infty} \int_{B_R^+} \left( \langle A_j(x) \nabla w_j^\delta, \nabla w_j^\delta \rangle + \lambda_+ \chi_{\{w_j^\delta > 0\}} + \lambda_- \chi_{\{w_j^\delta \leq 0\}} \right) dx \\ & \leq \limsup_{j \rightarrow \infty} \int_{B_R^+} \left( \langle A_j(x) \nabla w_j^\delta, \nabla w_j^\delta \rangle + \lambda_+ \chi_{\{w_j^\delta > 0\}} + \lambda_- \chi_{\{w_j^\delta \leq 0\}} \right) dx. \end{aligned} \tag{3.16}$$

From the same reasoning as for the justification of (3.11), we have

$$\limsup_{j \rightarrow \infty} \int_{B_R^+} \langle A_j(x) \nabla w_j^\delta, \nabla w_j^\delta \rangle dx = \limsup_{j \rightarrow \infty} \int_{B_R^+} |\nabla w_j^\delta|^2 dx.$$

Therefore, rewriting (3.16) as

$$\begin{aligned} & \int_{B_R^+} \left( |\nabla u_0|^2 + \lambda_+ \chi_{\{u_0 > 0\}} + \lambda_- \chi_{\{u_0 \leq 0\}} \right) dx \\ & \leq \limsup_{j \rightarrow \infty} \int_{B_R^+} \left( |\nabla w_j^\delta|^2 dx + \lambda_+ \chi_{\{w_j^\delta > 0\}} + \lambda_- \chi_{\{w_j^\delta \leq 0\}} \right) dx, \end{aligned} \tag{3.17}$$

we claim that

$$\lim_{\delta \rightarrow 0} \left( \limsup_{j \rightarrow \infty} \int_{B_R^+} |\nabla w_j^\delta|^2 dx \right) = \int_{B_R^+} |\nabla w|^2 dx. \tag{3.18}$$

To obtain the claim above, we prove that

$$\lim_{\delta \rightarrow 0} \left( \limsup_{j \rightarrow \infty} \int_{B_R^+} |\nabla(w_j^\delta - w)|^2 dx \right) = 0.$$

From the definition of  $w_j^\delta$ , we know that

$$w_j^\delta - w = (1 - \eta_\delta)(u_j - u_0) + \eta_\delta \theta_j \phi_j.$$

Therefore, we have

$$\begin{aligned}
 & \int_{B_R^+} |\nabla(w_j^\delta - w)|^2 dx \\
 & \leq C \left( \int_{B_R^+} |\nabla((1-\eta_\delta)(u_j - u_0))|^2 dx + \int_{B_R^+} |\nabla(\theta_j \eta_\delta \phi_j)|^2 dx \right) \\
 & \leq C(N) \left( \int_{B_R^+} (1-\eta_\delta)^2 |\nabla(u_j - u_0)|^2 dx + \frac{1}{\delta^2} \int_{B_R^+} |u_j - u_0|^2 dx \right. \\
 & \quad \left. + \int_{B_R^+} |\nabla(\theta_j \eta_\delta \phi_j)|^2 dx \right). \tag{3.19}
 \end{aligned}$$

Let us consider the first term on the right-hand side. We know that  $\int_{B_R^+} |\nabla(u_j - u_0)|^2 dx$  is uniformly bounded in  $j \in \mathbb{N}$  (cf. (3.7)). Therefore,

$$\begin{aligned}
 & \lim_{\delta \rightarrow 0} \left( \limsup_{j \rightarrow \infty} \int_{B_R^+} (1 - \eta_\delta)^2 |\nabla(u_j - u_0)|^2 dx \right) \\
 & = \left( \lim_{\delta \rightarrow 0} \|1 - \eta_\delta\|_{L^\infty(B_R^+)}^2 \right) \left( \limsup_{j \rightarrow \infty} \int_{B_R^+} |\nabla(u_j - u_0)|^2 dx \right) \\
 & \leq C(N, \lambda, \mu, R, M, \alpha) \lim_{\delta \rightarrow 0} \|1 - \eta_\delta\|_{L^\infty(B_R^+)} = 0. \tag{3.20}
 \end{aligned}$$

Regarding the second term, since  $|u_j - u_0|$  tends to zero in  $L^2(B_R^+)$  as  $j \rightarrow \infty$ , therefore the second term also tends to zero as  $j \rightarrow \infty$ . We write

$$\lim_{\delta \rightarrow 0} \left( \lim_{j \rightarrow \infty} \frac{1}{\delta^2} \int_{B_R^+} |u_j - u_0|^2 dx \right) = 0. \tag{3.21}$$

Lastly, we claim that

$$\lim_{j \rightarrow \infty} \int_{B_R^+} |\nabla(\theta_j \eta_\delta \phi_j)|^2 dx = 0. \tag{3.22}$$

Indeed, we have

$$\begin{aligned}
 \int_{B_R^+} |\nabla(\theta_j \eta_\delta \phi_j)|^2 dx & \leq C \left( \int_{B_R^+} |\nabla \eta_\delta|^2 (\theta_j \phi_j)^2 dx + \int_{B_R^+} |\nabla \phi_j|^2 (\eta_\delta \theta_j)^2 dx \right. \\
 & \quad \left. + \int_{B_R^+} |\nabla \theta_j|^2 (\eta_\delta \phi_j)^2 dx \right). \tag{3.23}
 \end{aligned}$$

Since  $\eta_\delta, \theta_j \leq 1$ ,  $|\nabla \eta_\delta| \leq \frac{C(N)}{\delta}$ , and  $\|\nabla \phi_j\|_{L^\infty(B_R^+)} \leq M$  (cf. (P1)), we obtain

$$\begin{aligned}
 & \int_{B_R^+} |\nabla \eta_\delta|^2 (\theta_j \phi_j)^2 dx + \int_{B_R^+} |\nabla \phi_j|^2 (\eta_\delta \theta_j)^2 dx \\
 & \leq C(N) |\{\theta_j \neq 0\} \cap B_R^+| \left( \frac{1}{\delta^2} + M^2 \right). \tag{3.24}
 \end{aligned}$$

We know that  $|\{\theta_j \neq 0\} \cap B_R^+| \rightarrow 0$  as  $j \rightarrow \infty$ ; hence, from (3.24), the first and second terms in (3.23) tend to zero as  $j \rightarrow \infty$ . The last term in (3.23) also tends to zero as  $j \rightarrow \infty$ ; indeed from (P1), we have

$$[\nabla \phi_j]_{C^\alpha(B_R^+)} \leq r_j^\alpha M.$$

Since  $\phi_j(0) = 0$ , therefore we have  $|\phi_j| \leq R^\alpha [\nabla \phi_j]_{B_R^+} \leq R^\alpha r_j^\alpha M$  in  $B_R^+$ . Also, observing that  $|\nabla \theta_j| \leq \frac{1}{d_j}$ ,  $\eta_\delta \leq 1$  in  $B_R^+$ , we have

$$\int_{B_R^+} |\nabla \theta_j|^2 (\eta_\delta \phi_j)^2 dx \leq MR^{2\alpha} \frac{r_j^{2\alpha}}{d_j^2} |B_R^+|.$$

If we choose a sequence  $d_j \rightarrow 0^+$  such that we also have

$$\frac{r_j^\alpha}{d_j} \rightarrow 0,$$

the third term in (3.23) tends to zero as  $j \rightarrow \infty$ . Plugging in the estimates above (3.20), (3.21), and (3.22) in (3.19), we obtain the claim (3.18).

From equations (3.14), (3.15), and (3.18), we obtain that the right-hand side of (3.17) is equal to  $J_0(w; B_R^+)$ ; therefore,  $u_0$  is a minimizer of  $J_0(\cdot; B_R^+)$ . That is,

$$J_0(u_0; B_R^+) \leq J_0(w; B_R^+)$$

for every  $w \in H^1(B_R^+)$  such that  $u_0 - w \in H_0^1(B_R^+)$ . Since the inequality above (which corresponds to (G4)) and other verified properties of  $u_0$  (i.e., (G1), (G2), and (G3) in  $B_R^+$ ) hold for every  $R > 0$ , therefore  $u_0 \in H_{loc}^1(\mathbb{R}_+^N)$  satisfies all the properties in Definition 2.5. Hence,  $u_0 \in \mathcal{P}_\infty$ . ■

After proving that the (subsequential) limits of blowups are global solutions, we proceed to show that the positivity sets (and hence the free boundaries) of blowups converge in certain sense to that of blowup limit. For this, we will need to establish that the minimizers  $u \in \mathcal{P}_r$  are non-degenerate near the free boundary. In the proof below, we adapt the ideas from [2].

**Proposition 3.9** (Non-degeneracy near the free boundary). *Let  $u \in \mathcal{P}_{r_0}$  for some  $r_0 > 0$  and  $x_0 \in B_{2/r_0}^+$ . Then, for every  $0 < \kappa < 1$ , there exists a constant  $c(\mu, N, \kappa, \lambda_\pm) > 0$  such that, for all  $B_r(x_0) \subset B_{\frac{2}{r_0}}^+$ , we have*

$$\frac{1}{r} \int_{\partial B_r(x_0)} u^+ d\mathcal{H}^{N-1}(x) < c(\mu, N, \kappa, \lambda_\pm) \implies u^+ = 0 \quad \text{in } B_{\kappa r}(x_0).$$

*Proof.* We fix  $x_0 \in \{u > 0\} \cap B_{2/r_0}^+$  and  $r > 0$  such that  $B_r(x_0) \subset B_{2/r_0}^+$ . We denote  $\gamma := \frac{1}{r} \int_{B_r(x_0)} u^+ dx$ . We know from Lemma 3.2 that the set  $\{u > 0\}$  is open. Also, since  $u \in \mathcal{P}_{r_0}$ , there exists  $A \in C^\infty(B_{2/r_0}^+)^{N \times N}$ ,  $\varphi \in C^{1,\alpha}(B_{2/r_0}^+)$ ,  $\lambda_\pm$  satisfying (P1)–(P6).






Figure 2. Graph of  $v_\varepsilon$ .

Therefore,  $u$  solves the PDE  $\operatorname{div}(A(x)\nabla u) = 0$  in  $\{u > 0\} \cap B_{2/r_0}^+$ . By elliptic regularity theory,  $u$  is locally smooth,  $(C_{\text{loc}}^{1,\alpha}(\{u > 0\} \cap B_{2/r_0}^+))$ . Then, for almost every  $\varepsilon > 0$ , (smooth)  $B_r \cap \partial\{u > \varepsilon\}$  is a  $C^{1,\alpha}$  surface. Pick one such small  $\varepsilon > 0$ , and we consider the test function  $v_\varepsilon$  given by

$$\begin{cases} \operatorname{div}(A(x)\nabla v_\varepsilon) = 0 & \text{in } (B_r(x_0) \setminus B_{\kappa r}(x_0)) \cap \{u > \varepsilon\}, \\ v_\varepsilon = u & \text{in } B_r(x_0) \cap \{u \leq \varepsilon\}, \\ v_\varepsilon = \varepsilon & \text{in } B_{\kappa r}(x_0) \cap \{u > \varepsilon\}, \\ v_\varepsilon = u & \text{on } \partial B_r(x_0). \end{cases}$$

Refer to Figure 2 for a pictorial understanding of definition  $v_\varepsilon$ . The function  $v_\varepsilon$  defined above belongs to  $H^1(B_r(x_0))$ ; thanks to [11, Theorem 3.44] ([11, Theorem 3.44] is proven for  $C^1$  domains, but the proof can also be adapted for Lipschitz domains [13, Theorem 4.6]). We intend to show that  $v_\varepsilon$  is bounded in  $H^1(B_r(x_0))$ . This ensures the existence of limit  $\lim_{\varepsilon \rightarrow 0} v_\varepsilon$  exists in weak sense in  $H^1(B_r(x_0))$  and strong sense in  $L^2(B_r(x_0))$ . Let  $G$  be the Green function for  $L(v) = \operatorname{div}(A(x)\nabla v)$  in the ring  $B_r(x_0) \setminus B_{\kappa r}(x_0)$ . Then, if there is a function  $w$  such that

$$\begin{cases} \operatorname{div}(A(x)\nabla w) = 0 & \text{in } B_r(x_0) \setminus B_{\kappa r}(x_0), \\ w = u & \text{on } \partial B_r(x_0) \cap \{u > \varepsilon\}, \\ w = \varepsilon & \text{elsewhere on } \partial(B_r(x_0) \setminus B_{\kappa r}(x_0)), \end{cases} \tag{3.25}$$

we can also write that  $w - \varepsilon = (u - \varepsilon)^+$  on  $\partial B_r(x_0)$  and  $w - \varepsilon = 0$  on  $\partial B_{\kappa r}(x_0)$ . From boundary elliptic regularity (cf. [6, Theorem 2]),  $(w - \varepsilon) \in C^{1,\alpha}(\overline{B_{r-\eta}(x_0)} \setminus \overline{B_{\kappa r}(x_0)})$  for every  $\eta \in (0, (1 - \kappa)r)$ . Consider  $\bar{x} \in \partial B_{\kappa r}(x_0)$  and any sequence  $\{x_k\} \subset B_r(x_0) \setminus B_{\kappa r}(x_0)$  such that  $x_k \rightarrow \bar{x} \in \partial B_{\kappa r}(x_0)$ . We have

$$\nabla(w - \varepsilon)(\bar{x}) = \lim_{k \rightarrow \infty} \nabla(w - \varepsilon)(x_k).$$

By Green representation formulae for  $w - \varepsilon$  in (3.25) (cf. [21, equation (1.12)]), we have

$$(w - \varepsilon)(x_k) = \int_{\partial(B_r(x_0) \setminus B_{\kappa r}(x_0))} (u - \varepsilon)^+(y)(A(y)\nabla_y G(x_k, y)) \cdot \nu_y \, d\sigma(y). \quad (3.26)$$

Therefore,

$$\begin{aligned} & \nabla(w - \varepsilon)(\bar{x}) \\ &= \lim_{k \rightarrow \infty} \nabla(w - \varepsilon)(x_k) \\ &= \lim_{k \rightarrow \infty} \nabla_x \left( \int_{\partial(B_r(x_0) \setminus B_{\kappa r}(x_0))} (u - \varepsilon)^+(y)(A(y)\nabla_y G(x, y)) \cdot \nu_y \, d\sigma(y) \right) \Big|_{x=x_k} \\ &= \lim_{k \rightarrow \infty} \int_{\partial B_r(x_0) \cap \{u > \varepsilon\}} (u - \varepsilon)^+(y)(A(y)\nabla_x \nabla_y G(x_k, y)) \cdot \nu_y \, d\sigma(y) \quad \forall \bar{x} \in \partial B_{\kappa r}(x_0), \end{aligned} \quad (3.27)$$

where  $\nu_y$  is the unit outer normal vector at a point  $y$  on the boundary. Computations in (3.27) are justified by the regularity and estimates given in [26, Theorem 3] (see also the classical paper by Grüter and Widman [18, Theorem 3.3 (vi)]). As a matter of fact, we can proceed further to obtain for  $\bar{x} \in \partial B_{\kappa r}(x_0)$

$$\begin{aligned} & |\nabla w(\bar{x})| \\ & \leq C(\mu, N) \lim_{k \rightarrow \infty} \int_{\partial B_r(x_0) \cap \{u > \varepsilon\}} (u - \varepsilon)^+(y) |\nabla_x \nabla_y G(x_k, y)| \, d\mathcal{H}^{N-1}(y) \\ & \leq \frac{C(\mu, N)}{(1 - \kappa)^N} \frac{1}{r} \int_{\partial B_r} (u - \varepsilon)^+ \, d\mathcal{H}^{N-1}(y) \leq C(\mu, N, \kappa)\gamma \quad \text{on } \partial B_{\kappa r}(x_0). \end{aligned} \quad (3.28)$$

We can easily check by the respective definitions that  $w \geq v_\varepsilon$  on  $\partial(B_r(x_0) \setminus B_{\kappa r}(x_0))$ . Moreover, since  $w \in C(\overline{B_r(x_0) \setminus B_{\kappa r}(x_0)})$  and  $w \geq \varepsilon$  on  $\partial(B_r(x_0) \setminus B_{\kappa r}(x_0))$  by the maximum principle (recall  $\operatorname{div}(A(x)\nabla w) = 0$  in  $B_r(x_0) \setminus B_{\kappa r}(x_0)$ ), we conclude that  $w \geq v_\varepsilon$  in  $B_r(x_0) \setminus B_{\kappa r}(x_0)$ . In particular,  $w \geq v_\varepsilon$  on  $\partial D_\varepsilon$ , where

$$D_\varepsilon := (B_r(x_0) \setminus B_{\kappa r}(x_0)) \cap \{u > \varepsilon\}.$$

By comparison principle, we know that  $w \geq v_\varepsilon$  in  $D_\varepsilon$ , and since  $w = v_\varepsilon = \varepsilon$  on  $\partial B_{\kappa r}(x_0) \cap \{u > \varepsilon\}$ , hence from (3.28)

$$|\nabla v_\varepsilon| \leq |\nabla w| \leq C(\mu, N, \kappa)\gamma \quad \text{on } \partial B_{\kappa r}(x_0) \cap \{u > \varepsilon\}. \quad (3.29)$$

Given that  $\operatorname{div}(A(x)\nabla v_\varepsilon) = 0$  in  $D_\varepsilon$ , we have, by divergence theorem and (3.29),

$$\begin{aligned} \int_{D_\varepsilon} (A(x)\nabla v_\varepsilon) \cdot \nabla(v_\varepsilon - u) \, dx &= \int_{\partial B_{\kappa r}(x_0) \cap \{u > \varepsilon\}} (u - v_\varepsilon)(A(x)\nabla v_\varepsilon) \cdot \nu(y) \, d\mathcal{H}^{N-1}(y) \\ &\leq C(\mu) \int_{\partial B_{\kappa r}(x_0) \cap \{u > \varepsilon\}} |u - \varepsilon| |\nabla v_\varepsilon| \, d\mathcal{H}^{N-1}(y) \\ &\leq C(\mu, N, \kappa)\gamma \int_{\partial B_{\kappa r}(x_0) \cap \{u > \varepsilon\}} |u - \varepsilon| \, d\mathcal{H}^{N-1}(y) \\ &=: M_0(u). \end{aligned}$$

The justification for the use of divergence theorem in  $D_\varepsilon$  can be found in [2, equation (3.4)]. From the calculations above, we can write

$$\begin{aligned} & \int_{D_\varepsilon} (A(x)\nabla v_\varepsilon) \cdot \nabla (v_\varepsilon - u) \, dx \leq M_0 \\ \Rightarrow & \int_{D_\varepsilon} (A(x)\nabla v_\varepsilon) \cdot \nabla v_\varepsilon \, dx \leq M_0 + \int_{D_\varepsilon} (A(x)\nabla v_\varepsilon) \cdot \nabla u \, dx \\ \Rightarrow & \mu \int_{D_\varepsilon} |\nabla v_\varepsilon|^2 \, dx \leq M_0 + \frac{1}{\mu} \int_{D_\varepsilon} |\nabla v_\varepsilon| |\nabla u| \, dx \\ \Rightarrow & \mu \int_{D_\varepsilon} |\nabla v_\varepsilon|^2 \, dx \leq M_0 + \frac{\varepsilon_0}{2\mu} \int_{D_\varepsilon} |\nabla v_\varepsilon|^2 \, dx + \frac{1}{2\varepsilon_0\mu} \int_{D_\varepsilon} |\nabla u|^2 \, dx. \end{aligned}$$

Putting very small  $\varepsilon_0 > 0$  in the last inequality, we have

$$\int_{D_\varepsilon} |\nabla v_\varepsilon|^2 \, dx \leq M_0 + C(\mu) \int_{D_\varepsilon} |\nabla u|^2 \, dx =: M_1(u).$$

Since  $v_\varepsilon = \varepsilon$  in  $B_{\kappa r}(x_0) \cap \{u > \varepsilon\}$  which implies that  $\nabla v_\varepsilon = 0$  in  $B_{\kappa r}(x_0) \cap \{u > \varepsilon\}$  and  $v_\varepsilon = u$  in  $B_r(x_0) \setminus D_\varepsilon$ ,

$$\int_{B_r(x_0)} |\nabla v_\varepsilon|^2 \, dx = \int_{D_\varepsilon} |\nabla v_\varepsilon|^2 \, dx + \int_{B_r(x_0) \setminus D_\varepsilon} |\nabla u|^2 \, dx =: M_2(u). \tag{3.30}$$

By the definition of  $v_\varepsilon$ ,  $0 < v_\varepsilon \leq u$  on  $\partial D_\varepsilon$  and  $\operatorname{div}(A(x)\nabla v_\varepsilon) = \operatorname{div}(A(x)\nabla u) = 0$  in  $D_\varepsilon$ , therefore, by comparison principle,  $0 < v_\varepsilon < u$  in  $D_\varepsilon$ . In the set  $B_{\kappa r}(x_0) \cap \{u > \varepsilon\}$ , we have  $v_\varepsilon = \varepsilon < u$  and  $v_\varepsilon = u$  in  $B_r(x_0) \cap \{u \leq \varepsilon\}$ . Overall, we have  $0 < v_\varepsilon \leq u$  in  $B_r(x_0) \cap \{u > \varepsilon\}$ . Therefore,

$$\begin{aligned} \int_{B_r(x_0)} |v_\varepsilon|^2 \, dx & \leq \int_{B_r(x_0) \cap \{u \leq \varepsilon\}} |u|^2 \, dx + \int_{B_r(x_0) \cap \{u > \varepsilon\}} |u|^2 \, dx \\ & = \int_{B_r(x_0)} |u|^2 \, dx. \end{aligned} \tag{3.31}$$

Hence, from (3.30) and (3.31),  $v_\varepsilon$  is bounded in  $H^1(B_r(x_0))$ . Therefore, up to a subsequence, there exists a limit  $v = \lim_{\varepsilon \rightarrow 0} v_\varepsilon$  in weak  $H^1$  sense such that  $v$  satisfies the following:

$$\begin{cases} \operatorname{div}(A(x)\nabla v) = 0 & \text{in } (B_r(x_0) \setminus B_{\kappa r}(x_0)) \cap \{u > 0\}, \\ v = u & \text{in } B_r(x_0) \cap \{u \leq 0\}, \\ v = 0 & \text{in } B_{\kappa r}(x_0) \cap \{u > 0\}, \\ v = u & \text{on } \partial B_r(x_0). \end{cases} \tag{3.32}$$

We verify the above properties (3.32) of  $v$  at the end of this proof.

Let us use the function  $v$  as a test function with respect to minimality condition on  $u$  in  $B_r(x_0)$ ; we have

$$\int_{B_r(x_0)} (\langle A(x)\nabla u, \nabla u \rangle + \Lambda(u)) \, dx \leq \int_{B_r(x_0)} (\langle A(x)\nabla v, \nabla v \rangle + \Lambda(v)) \, dx.$$

Since  $v = u$  in  $\{u \leq 0\}$  and  $\{v > 0\} \subset \{u > 0\}$ , the integration in the set  $\{u \leq 0\}$  gets canceled from both sides, and we are left with the terms mentioned below.

Setting  $D_0 := (B_r(x_0) \setminus B_{kr}(x_0)) \cap \{u > 0\}$ , we have

$$\begin{aligned} & \int_{B_r(x_0) \cap \{u > 0\}} (\langle A(x) \nabla u, \nabla u \rangle - \langle A(x) \nabla v, \nabla v \rangle) dx \\ & \leq \int_{B_r(x_0) \cap \{u > 0\}} (\Lambda(v) - \Lambda(u)) dx \\ & = \int_{B_{kr}(x_0) \cap \{u > 0\}} (\Lambda(v) - \Lambda(u)) dx \\ & = \lambda_0 |B_{kr}(x_0) \cap \{u > 0\}| \quad (\lambda_0 := -(\lambda_+ - \lambda_-)). \end{aligned}$$

We have the second equality above because

$$\chi_{\{v > 0\}} = \chi_{\{u > 0\}} \quad \text{in } D_0.$$

Since  $v = 0$  in  $\{u > 0\} \cap B_{kr}$ , we have

$$\begin{aligned} & \int_{B_{kr}(x_0) \cap \{u > 0\}} \langle A(x) \nabla u, \nabla u \rangle dx \\ & + \int_{D_0} (\langle A(x) \nabla u, \nabla u \rangle - \langle A(x) \nabla v, \nabla v \rangle) dx \leq \lambda_0 |B_{kr}(x_0) \cap \{u > 0\}|. \end{aligned}$$

Using the ellipticity of  $A$  and shuffling the terms in the above equation, we obtain

$$\begin{aligned} & \int_{B_{kr}(x_0) \cap \{u > 0\}} (\mu |\nabla u|^2 - \lambda_0) dx \\ & \leq \int_{D_0} (\langle A(x) \nabla v, \nabla v \rangle - \langle A(x) \nabla u, \nabla u \rangle) dx \\ & = \int_{D_0} (\langle A(x) \nabla(v - u), \nabla(v + u) \rangle) dx \\ & = \int_{D_0} (\langle A(x) \nabla(v - u), \nabla(u - v + 2v) \rangle) dx \\ & \leq 2 \int_{D_0} \langle A(x) \nabla v, \nabla(v - u) \rangle dx \\ & \leq 2 \liminf_{\varepsilon \rightarrow 0} \int_{D_0} \langle A(x) \nabla v_\varepsilon, \nabla(v_\varepsilon - u) \rangle \\ & = 2 \liminf_{\varepsilon \rightarrow 0} \int_{D_\varepsilon} \langle A(x) \nabla v_\varepsilon, \nabla(v_\varepsilon - u) \rangle dx \quad (\text{since } v_\varepsilon = u \text{ in } D_\varepsilon \setminus D_0) \\ & = 2 \liminf_{\varepsilon \rightarrow 0} \int_{\partial B_{kr}(x_0) \cap \{u > \varepsilon\}} (u - \varepsilon) \langle A(x) \nabla v_\varepsilon, \nu \rangle dx \\ & \leq \frac{2}{\mu} \liminf_{\varepsilon \rightarrow 0} \int_{\partial B_{kr}(x_0) \cap \{u > \varepsilon\}} (u - \varepsilon) |\nu \cdot \nabla v_\varepsilon| dx := M. \end{aligned} \tag{3.33}$$

The second-to-last equality in the above calculation is obtained from integration by parts; its justification can be found in [2, equation (3.4)]. From (3.33) and (3.29), and using the trace inequality in  $H^1(B_{\kappa r})$ , we have (for some different constant  $C(\kappa)$ )

$$\begin{aligned}
 M &\leq C(\mu, N, \kappa)\gamma \int_{\partial B_{\kappa r}(x_0)} u^+ d\mathcal{H}^{N-1}(x) \\
 &\leq C(\mu, N, \kappa)\gamma \int_{B_{\kappa r}(x_0)} \left( |\nabla u^+| + \frac{1}{r}u^+ \right) dx \\
 &\leq C(\mu, N, \kappa)\gamma \left[ |B_{\kappa r}(x_0) \cap \{u > 0\}|^{1/2} \left( \int_{B_{\kappa r}(x_0)} |\nabla u^+|^2 dx \right)^{1/2} \right. \\
 &\quad \left. + \frac{1}{r} \sup_{B_{\kappa r}(x_0)} (u^+) |\{B_{\kappa r}(x_0) \cap \{u > 0\}\}| \right] \\
 &\leq C(\mu, N, \kappa)\gamma \left[ \frac{1}{2\sqrt{-\lambda_0}} \int_{B_{\kappa r}(x_0) \cap \{u > 0\}} |\nabla u^+|^2 dx \right. \\
 &\quad \left. + \frac{\sqrt{-\lambda_0}}{2} |B_{\kappa r}(x_0) \cap \{u > 0\}| + \frac{1}{r} \sup_{B_{\kappa r}(x_0)} (u^+) \int_{B_{\kappa r}(x_0) \cap \{u > 0\}} 1 dx \right] \\
 &= \frac{C(\mu, N, \kappa)\gamma}{2\sqrt{-\lambda_0}} \left( \int_{B_{\kappa r}(x_0) \cap \{u > 0\}} |\nabla u^+|^2 - \lambda_0 dx \right) \\
 &\quad + \frac{C(\mu, N, \kappa)\gamma}{\lambda_0 r} \sup_{B_{\kappa r}(x_0)} (u^+) \int_{B_{\kappa r}(x_0) \cap \{u > 0\}} \lambda_0 dx. \tag{3.34}
 \end{aligned}$$

We have used Hölder’s inequality and then Young’s inequality above. From Lemma 3.1,  $u^+$  is  $A$ -subharmonic in  $B_r(x_0)$ . If  $G'$  is the Green’s function for

$$L'(v) = \operatorname{div}(A(x)\nabla v) \quad \text{in } B_r(x_0),$$

then by comparison principle and Green’s representation (cf. [21, equation (1.12)])

$$u^+(x) \leq \int_{\partial B_r(x_0)} u^+(y) (A(y)\nabla_y G'(x, y)) \cdot \nu_y d\mathcal{H}^{N-1}(y) \quad \forall x \in B_{\kappa r}(x_0). \tag{3.35}$$

Since for all  $y \in \partial B_r(x_0)$  and  $x \in B_{\kappa r}(x_0)$ , we have

$$\frac{1}{|x - y|^{N-1}} \leq \frac{C(\kappa)}{r^{N-1}},$$

then, using the Green’s function estimates (cf. [18, Theorem 3.3 (v)]), we get

$$\begin{aligned}
 \sup_{B_{\kappa r}(x_0)} u^+ &\leq C(\mu) \int_{\partial B_r(x_0)} \frac{u^+(y)}{|x - y|^{N-1}} d\mathcal{H}^{N-1}(y) \\
 &\leq C(\mu, \kappa, N) \int_{\partial B_r(x_0)} u^+ d\mathcal{H}^{N-1}(y) = C(\mu, \kappa, N)\gamma r. \tag{3.36}
 \end{aligned}$$

Using (3.33) and (3.36) in (3.34), we have

$$\begin{aligned} & \mu \int_{B_{\kappa r}(x_0) \cap \{u > 0\}} (|\nabla u|^2 - \lambda_0) dx \\ & \leq \frac{C(\mu, \kappa, N)\gamma}{2\sqrt{-\lambda_0}} \int_{B_{\kappa r}(x_0) \cap \{u > 0\}} (|\nabla u^+|^2 - \lambda_0) dx \\ & \quad + \frac{C(\mu, \kappa, N)\gamma}{\lambda_0 r} \sup_{B_{\kappa r}(x_0)} (u^+) \int_{B_{\kappa r}(x_0) \cap \{u > 0\}} \lambda_0 dx \\ & \leq \frac{C(\mu, \kappa, N)\gamma}{\mu\sqrt{-\lambda_0}} \left(1 + \frac{C(\kappa)\gamma}{\sqrt{-\lambda_0}}\right) \mu \int_{B_{\kappa r}(x_0) \cap \{u > 0\}} (|\nabla u|^2 - \lambda_0) dx. \end{aligned}$$

If  $\gamma$  is small enough, then

$$\int_{B_{\kappa r}(x_0) \cap \{u > 0\}} (|\nabla u|^2 - \lambda_0) dx = 0.$$

In particular,  $|\{u > 0\} \cap B_{\kappa r}(x_0)| = 0$ , that is,  $u^+ = 0$  almost everywhere in  $B_{\kappa r}(x_0)$ .

It remains to verify the properties of  $v$  in (3.32). Before looking at the proof, we observe that, for a given  $\varphi \in C_c^\infty(D_0)$ , there exists  $\varepsilon_0 > 0$  such that  $\varphi \in C_c^\infty(D_\varepsilon)$  for all  $\varepsilon < \varepsilon_0$ . Indeed, since  $\text{supp}(\varphi)$  is a compact set and  $\bigcup_{\varepsilon > 0} D_\varepsilon$  is a cover of  $\text{supp}(\varphi)$ , then for a finite set  $\{\varepsilon_1, \dots, \varepsilon_n\}$  we have  $\text{supp}(\varphi) \subset \bigcup_{i=1}^n D_{\varepsilon_i} \subset D_{\varepsilon_{\max}}$ , where  $\varepsilon_{\max} = \max(\varepsilon_1, \dots, \varepsilon_n)$ . Therefore,  $\varphi \in C_c^\infty(D_\varepsilon)$  for all  $\varepsilon < \varepsilon_{\max}$ .

Let us first verify that  $\text{div}(A(x)\nabla v) = 0$  in  $D_0$ . For this, let  $\varphi \in C_c^\infty(D_0)$ ; then from the continuity of  $u$ , there exists a  $\varepsilon_0 > 0$  such that  $\text{supp}(\varphi) \subset D_\varepsilon$  for all  $\varepsilon < \varepsilon_0$ ; also, we have

$$\int_{D_0} \langle A\nabla v, \nabla \varphi \rangle dx = \int_{\text{supp}(\varphi)} \langle A\nabla v, \nabla \varphi \rangle dx \tag{3.37}$$

since  $\text{supp}(\varphi) \subset D_\varepsilon$ ; from the definition of  $v_\varepsilon$ , we have

$$\int_{\text{supp}(\varphi)} \langle A\nabla v_\varepsilon, \nabla \varphi \rangle dx = 0,$$

and we know that  $v$  is a weak limit of  $v_\varepsilon$  in  $H^1(B_r(x_0))$ ; therefore, from (3.37), we have

$$\int_{D_0} \langle A\nabla v, \nabla \varphi \rangle dx = \int_{\text{supp}(\varphi)} \langle A\nabla v, \nabla \varphi \rangle dx = \lim_{\varepsilon \rightarrow 0} \int_{\text{supp}(\varphi)} \langle \nabla v_\varepsilon, \nabla \varphi \rangle dx = 0.$$

Hence, we show that  $\text{div}(A(x)\nabla v) = 0$  in  $D_0$ . To show that  $v = 0$  in  $B_{\kappa r}(x_0) \cap \{u > 0\}$ , we now take the function  $\varphi \in C_c^\infty(B_{\kappa r}(x_0) \cap \{u > 0\})$ . From the same reasoning as above, we know that there exists an  $\varepsilon_0 > 0$  such that  $\text{supp}(\varphi) \subset B_{\kappa r}(x_0) \cap \{u > \varepsilon\}$  for all  $\varepsilon < \varepsilon_0$ . From the definition of  $v_\varepsilon$ , we have

$$\int_{\text{supp}(\varphi)} v_\varepsilon \varphi dx = \int_{\{u > \varepsilon\} \cap B_{\kappa r}(x_0)} v_\varepsilon \varphi dx = \varepsilon \int_{\{u > \varepsilon\} \cap B_{\kappa r}(x_0)} \varphi dx,$$

and in limit  $\varepsilon \rightarrow 0$ , from the above equation, we have

$$\int_{\text{supp}(\varphi)} v\varphi \, dx = \lim_{\varepsilon \rightarrow 0} \int_{\text{supp}(\varphi)} v_\varepsilon\varphi \, dx = 0,$$

and therefore,  $v = 0$  a.e. in  $B_{\kappa r}(x_0) \cap \{u > 0\}$ . To prove that  $v = u$  in  $\{u \leq 0\}$ , we observe that  $\{u \leq 0\} \subset \{u \leq \varepsilon\}$ ; hence, from the definition of  $v_\varepsilon$ , we have

$$v_\varepsilon = u \quad \text{in } \{u \leq 0\}.$$

Since the weak limits maintain the equality (cf. [25, Lemma 3.14]), the claim follows in the limit  $\varepsilon \rightarrow 0$ . Apart from that, since  $v_\varepsilon = u$  on  $\partial B_r(x_0)$ , therefore, from the conservation of traces in weak convergence, it follows that  $v = u$  on  $\partial B_r(x_0)$ . This completes the proof of Proposition 3.9. ■

**Remark 3.10.** In the proposition above, the constant is local in nature; this means that the value of the constant depends on the choice of compact set  $K \subset\subset B_2^+$ , where  $x_0 \in K$ .

**Lemma 3.11.** *Let  $u_0$  and  $u_j$  be as in Lemma 3.8. Then, for a subsequence of  $u_j$ , for any  $R > 0$ , we have*

$$\chi_{\{u_k > 0\} \cap B_R^+} \rightarrow \chi_{\{u_0 > 0\} \cap B_R^+} \quad \text{a.e. in } B_R^+. \tag{3.38}$$

This in turn implies

$$\chi_{\{u_j > 0\} \cap B_R^+} \rightarrow \chi_{\{u_0 > 0\} \cap B_R^+} \quad \text{in } L^1(B_R^+). \tag{3.39}$$

*Proof.* From Lemma 3.8, we can consider a subsequence of  $u_j$  such that  $u_j \rightarrow u_0$  in  $L^\infty(B_R^+)$ . Let  $x \in B_R^+$ . If  $x \in \{u_0 > 0\} \cap B_R^+$  (or  $\chi_{\{u_0 > 0\} \cap B_R^+}(x) = 1$ ), then  $u_j(x) > \frac{u(x)}{2} > 0$  (or  $\chi_{\{u_j > 0\} \cap B_R^+}(x) = 1$ ) for  $k$  sufficiently large. Thus, we conclude that

$$\chi_{\{u_j > 0\} \cap B_R^+}(x) \rightarrow \chi_{\{u_0 > 0\} \cap B_R^+}(x) \text{ as } j \rightarrow \infty \text{ for all } x \in \{u_0 > 0\} \cap B_R^+.$$

If  $x \in \{u_0 \leq 0\} \cap B_R^+$  (or  $\chi_{\{u_0 > 0\} \cap B_R^+}(x) = 0$ ), then there exists  $\delta > 0$  such that

$$B_\delta(x) \subset \{u_0 \leq 0\} \cap B_R^+.$$

Thus, we have

$$\frac{1}{\delta} \int_{\partial B_\delta(x)} u_0^+ \, d\mathcal{H}^{N-1} = 0.$$

Again, by the uniform convergence of  $u_j$  to  $u_0$  in  $B_R^+$  (cf. Lemma 3.8), we obtain

$$\frac{1}{\delta} \int_{\partial B_\delta(x)} u_j^+ \, d\mathcal{H}^{N-1} \leq \frac{1}{2} c(\mu, N, \lambda_\pm) \quad \text{for } j \text{ sufficiently large.} \tag{3.40}$$

Here,  $c(\mu, N, \lambda_\pm)$  is as in Proposition 3.9. This implies that  $u_j \leq 0$  in  $B_{\frac{\delta}{2}}(x)$  (cf. Proposition 3.9). In particular,  $\chi_{\{u_j(x) \leq 0\}}(x) = 0$  for  $j$  sufficiently large. This way, we obtain

$$\chi_{\{u_j > 0\} \cap B_R^+}(x) \rightarrow \chi_{\{u_0 > 0\} \cap B_R^+}(x) \text{ as } j \rightarrow \infty \text{ for all } x \in \{u_0 \leq 0\} \cap B_R^+. \tag{3.41}$$

From the representation theorem [2, Theorem 7.3], we know that

$$|\partial\{u_0 > 0\} \cap B_R^+| = 0.$$

From (3.40), (3.41), and the fact that

$$|\partial\{u_0 > 0\} \cap B_R^+| = 0,$$

we obtain the claim (3.38). Since

$$|\chi_{\{u_j > 0\} \cap B_R^+}| \leq 1,$$

the claim (3.39) follows from Lebesgue’s dominated convergence theorem. ■

### 4. The main result

We rephrase the notion of the tangential touch of the free boundary to the fixed boundary, which is equivalent to the tangential touch condition mentioned in the statement of Theorem 2.6.

In the proof of our main result, we will show that, given  $u \in \mathcal{P}_1$ , for every  $\varepsilon > 0$  there exists  $\rho_\varepsilon > 0$  such that

$$\partial\{u > 0\} \cap B_\rho^+ \subset B_\rho^+ \setminus K_\varepsilon \quad \forall 0 < \rho \leq \rho_\varepsilon,$$

where

$$K_\varepsilon := \{x \in \mathbb{R}_+^N : x_N \geq \varepsilon \sqrt{x_1^2 + \dots + x_{N-1}^2}\}.$$

*Proof of Theorem 2.6.* We assume, by contradiction, that the free boundaries of functions in  $\mathcal{P}_1$  do not touch the origin in a tangential fashion to the plane. Then, there exist  $\varepsilon > 0$  and sequences  $v_j \in \mathcal{P}_1$  and  $x_j \in F(v_j) \cap K_\varepsilon$  such that  $|x_j| \rightarrow 0$  as  $j \rightarrow \infty$ . Let  $r_j = |x_j|$ , and we consider the blowups  $u_j := (v_j)_{r_j}$ .

Let  $u_0 := \lim_{j \rightarrow \infty} u_j$  as in Lemma 3.8. Also, let  $x_0 \in \partial B_1^+ \cap K_\varepsilon$  be a limit up to a subsequence (still called  $x_j$ ) such that

$$x_0 = \lim_{j \rightarrow \infty} \frac{x_j}{|x_j|}.$$

Since  $x_j \in F(v_j)$ , we have  $v_j(x_j) = 0$ . Therefore, on rescaling,

$$u_j\left(\frac{x_j}{r_j}\right) = \frac{1}{r_j} v_j(x_j) = 0.$$

In the limit as  $j \rightarrow \infty$ , we have

$$u_0(x_0) = \lim_{j \rightarrow \infty} u_j\left(\frac{x_j}{r_j}\right) = 0.$$



From the density assumption that  $u_j$  satisfy condition (2.1) and Lemma 3.11, we have for any given  $R > 0$

$$\begin{aligned} \frac{|\{u_0 > 0\} \cap B_R^+|}{|B_R^+|} &= \int_{B_R^+} \chi_{\{u_0 > 0\}} dx = \lim_{j \rightarrow \infty} \int_{B_R^+} \chi_{\{u_j > 0\}} dx \\ &= \lim_{j \rightarrow \infty} \frac{1}{|B_{Rr_j}^+|} \int_{B_{Rr_j}^+} \chi_{\{v_j > 0\}} dx \\ &= \lim_{j \rightarrow \infty} \frac{|\{v_j > 0\} \cap B_{Rr_j}^+|}{|B_{Rr_j}^+|} > \mathcal{D}. \end{aligned} \tag{4.1}$$

We can see that the computations done in (4.1), in fact, show that the density property remains invariant under the blowup of any function  $v$ . This way, we conclude that the function  $(u_0)_0$  which is the blowup limit of  $(u_0)$  (in particular,  $(u_0)_0 := \lim_{r \rightarrow 0} (u_0)_r$ ) also satisfies

$$\frac{|\{(u_0)_0 > 0\} \cap B_R^+|}{|B_R^+|} > \mathcal{D} \quad \forall R > 0. \tag{4.2}$$

Now, we note that from Lemma 3.8  $u_0 \in \mathcal{P}_\infty$ . Moreover, from (4.1),  $u_0 \not\equiv 0$ , and from [22, Theorem 4.2, Lemma 4.3], we have  $u_0 \geq 0$ ; also, from (4.2), we conclude that  $(u_0)_0 \not\equiv 0$ . This way, again by [22, Theorem 4.9], we have  $u_0(x) = c x_N^+$  for all  $x \in \mathbb{R}_+^N$  for some constant  $c > 0$ .

Hence, the function  $u_0$  cannot be equal to zero at any point in  $\mathbb{R}_+^N$ . But we have  $x_0 \in \partial B_1^+ \cap K_\varepsilon$  and  $u_0(x_0) = 0$ . This leads to a contradiction. In order to construct the modulus of continuity  $\sigma$ , for  $x \in \mathbb{R}_+^N$ , we assign the value  $\sigma(|x|)$  to be the maximum opening  $\eta$  of cone  $K_\eta$  such that  $F(u) \cap B_r^+ \subset K_\eta^c$  for all  $r \leq |x|$ . In other words,

$$\begin{aligned} \sigma(|x|) &= \inf\{\eta \mid F(u) \cap B_r^+ \subset K_\eta^c, r \leq |x|\} \\ &= \sup\left\{\frac{x_N}{|x'|} \mid x \in F(u) \cap B_r^+ \setminus \{0\}, r \leq |x|\right\}. \end{aligned} \quad \blacksquare$$

## References

[1] H. W. Alt and L. A. Caffarelli, Existence and regularity for a minimum problem with free boundary. *J. Reine Angew. Math.* **325** (1981), 105–144 Zbl [0449.35105](#) MR [618549](#)

[2] H. W. Alt, L. A. Caffarelli, and A. Friedman, Variational problems with two phases and their free boundaries. *Trans. Amer. Math. Soc.* **282** (1984), no. 2, 431–461 Zbl [0844.35137](#) MR [732100](#)

[3] H. W. Alt and G. Gilardi, The behavior of the free boundary for the dam problem. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **9** (1982), no. 4, 571–626 Zbl [0521.76092](#) MR [693780](#)

[4] R. Argiolas and F. Ferrari, Flat free boundaries regularity in two-phase problems for a class of fully nonlinear elliptic operators with variable coefficients. *Interfaces Free Bound.* **11** (2009), no. 2, 177–199 Zbl [1179.35349](#) MR [2511639](#)

- [5] G. Birkhoff and E. H. Zarantonello, *Jets, wakes, and cavities*. Academic Press, New York, 1957 Zbl 0077.18703 MR 88230
- [6] M. V. Borsuk, [Dini-continuity of first derivatives of solutions of the Dirichlet problem for second-order linear elliptic equations in a nonsmooth domain](#). *Sibirsk. Mat. Zh.* **39** (1998), no. 2, 261–280 Zbl 0902.35031 MR 1631768
- [7] J. E. M. Braga and D. R. Moreira, [Up to the boundary gradient estimates for viscosity solutions to nonlinear free boundary problems with unbounded measurable ingredients](#). *Calc. Var. Partial Differential Equations* **61** (2022), no. 5, article no. 197, 65 pp. Zbl 1496.35129 MR 4469486
- [8] L. A. Caffarelli, [A Harnack inequality approach to the regularity of free boundaries. I. Lipschitz free boundaries are  \$C^{1,\alpha}\$](#) . *Rev. Mat. Iberoamericana* **3** (1987), no. 2, 139–162 Zbl 0676.35085 MR 990856
- [9] L. A. Caffarelli, [A Harnack inequality approach to the regularity of free boundaries. II. Flat free boundaries are Lipschitz](#). *Comm. Pure Appl. Math.* **42** (1989), no. 1, 55–78 Zbl 0676.35086 MR 973745
- [10] D. De Silva, F. Ferrari, and S. Salsa, [Free boundary regularity for fully nonlinear non-homogeneous two-phase problems](#). *J. Math. Pures Appl. (9)* **103** (2015), no. 3, 658–694 Zbl 1342.35457 MR 3310271
- [11] F. Demengel and G. Demengel, *Functional spaces for the theory of elliptic partial differential equations*. Universitext, Springer, London; EDP Sciences, Les Ulis, 2012 Zbl 1239.46001 MR 2895178
- [12] C. Dunn, *Introduction to analysis*. Textb. Math., CRC Press, Boca Raton, FL, 2017 Zbl 1375.26001
- [13] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*. Revised edn., Textb. Math., CRC Press, Boca Raton, FL, 2015 Zbl 1310.28001 MR 3409135
- [14] F. Ferrari, [Two-phase problems for a class of fully nonlinear elliptic operators. Lipschitz free boundaries are  \$C^{1,\gamma}\$](#) . *Amer. J. Math.* **128** (2006), no. 3, 541–571 Zbl 1142.35108 MR 2230916
- [15] F. Ferrari and C. Lederman, [Regularity of flat free boundaries for a  \$p\(x\)\$ -Laplacian problem with right hand side](#). *Nonlinear Anal.* **212** (2021), article no. 112444, 25 pp. Zbl 1472.35464 MR 4273843
- [16] E. Giusti, *Direct methods in the calculus of variations*. World Scientific, River Edge, NJ, 2003 Zbl 1028.49001 MR 1962933
- [17] G. Gravina and G. Leoni, [On the behavior of the free boundary for a one-phase Bernoulli problem with mixed boundary conditions](#). *Commun. Pure Appl. Anal.* **19** (2020), no. 10, 4853–4878 Zbl 1460.35397 MR 4147334
- [18] M. Grüter and K.-O. Widman, [The Green function for uniformly elliptic equations](#). *Manuscripta Math.* **37** (1982), no. 3, 303–342 Zbl 0485.35031 MR 657523
- [19] E. Indrei, [Boundary regularity and nontransversal intersection for the fully nonlinear obstacle problem](#). *Comm. Pure Appl. Math.* **72** (2019), no. 7, 1459–1473 Zbl 1429.35087 MR 3957397
- [20] E. Indrei, [Free boundary regularity near the fixed boundary for the fully nonlinear obstacle problem](#). In *Advances in harmonic analysis and partial differential equations*, pp. 147–156, Contemp. Math. 748, American Mathematical Society, Providence, RI, 2020 Zbl 1436.35329 MR 4085452
- [21] D. S. Jerison and C. E. Kenig, [The Dirichlet problem in nonsmooth domains](#). *Ann. of Math. (2)* **113** (1981), no. 2, 367–382 Zbl 0434.35027 MR 607897

- [22] A. L. Karakhanyan, C. E. Kenig, and H. Shahgholian, [The behavior of the free boundary near the fixed boundary for a minimization problem](#). *Calc. Var. Partial Differential Equations* **28** (2007), no. 1, 15–31 Zbl [1111.35135](#) MR [2267752](#)
- [23] A. L. Karakhanyan and H. Shahgholian, [Analysis of a free boundary at contact points with Lipschitz data](#). *Trans. Amer. Math. Soc.* **367** (2015), no. 7, 5141–5175 Zbl [1348.35331](#) MR [3335413](#)
- [24] E. Lieb and M. Loss, *Analysis*. CRM Proc. Lecture Notes, American Mathematical Society, Providence, RI, 2001
- [25] D. Moreira and H. Shrivastava, [Optimal regularity for variational solutions of free transmission problems](#). *J. Math. Pures Appl. (9)* **169** (2023), 1–49 Zbl [1504.35105](#) MR [4523460](#)
- [26] A. V. Vähäkangas, [On regularity and extension of Green’s operator on bounded smooth domains](#). *Potential Anal.* **37** (2012), no. 1, 57–77 Zbl [1250.35009](#) MR [2928238](#)

Received 3 July 2022; revised 15 March 2023.

**Diego Moreira**

Departamento de Matemática, Universidade Federal do Ceará, 60.455-760 Fortaleza, Brazil;  
[dmoreira@mat.ufc.br](mailto:dmoreira@mat.ufc.br)

**Harish Shrivastava**

Gandhi Institute of Technology and Management (GITAM), 560010 Bangalore, India;  
[hshrivas@gitam.edu](mailto:hshrivas@gitam.edu)