

# On minimal non-supersoluble groups

Adolfo Ballester-Bolinches and Ramón Esteban-Romero

*Dedicated to the memory of Klaus Doerk (1939–2004)*

## Abstract

The aim of this paper is to classify the finite minimal non- $p$ -supersoluble groups,  $p$  a prime number, in the  $p$ -soluble universe.

## 1. Introduction

All groups considered in this paper are finite.

Given a class  $\mathfrak{X}$  of groups, we say that a group  $G$  is a *minimal non- $\mathfrak{X}$ -group* or an  *$\mathfrak{X}$ -critical group* if  $G \notin \mathfrak{X}$ , but all proper subgroups of  $G$  belong to  $\mathfrak{X}$ . It is rather clear that detailed knowledge of the structure of  $\mathfrak{X}$ -critical groups could help to give information about what makes a group belong to  $\mathfrak{X}$ .

Minimal non- $\mathfrak{X}$ -groups have been studied for various classes of groups  $\mathfrak{X}$ . For instance, Miller and Moreno [10] analysed minimal non-abelian groups, while Schmidt [14] studied minimal non-nilpotent groups. These groups are now known as *Schmidt groups*. Rédei classified completely the minimal non-abelian groups in [12] and the Schmidt groups in [13]. More precisely,

**Theorem 1** ([12]). *The minimal non-abelian groups are of one of the following types:*

1.  $G = [V_q]C_{r^s}$ , where  $q$  and  $r$  are different prime numbers,  $s$  is a positive integer, and  $V_q$  is an irreducible  $C_{r^s}$ -module over the field of  $q$  elements with kernel the maximal subgroup of  $C_{r^s}$ ,
2. the quaternion group of order 8,
3.  $G_{II}(q, m, n) = \langle a, b \mid a^{q^m} = b^{q^n} = 1, a^b = a^{1+q^{m-1}} \rangle$ , where  $q$  is a prime number,  $m \geq 2$ ,  $n \geq 1$ , of order  $q^{m+n}$ , and

---

2000 Mathematics Subject Classification: 20D10, 20F16.

Keywords: Finite groups, supersoluble groups, critical groups.

4.  $G_{III}(q, m, n) = \langle a, b \mid a^{q^m} = b^{q^n} = [a, b]^q = [a, b, a] = [a, b, b] = 1 \rangle$ ,  
 where  $q$  is a prime number,  $m \geq n \geq 1$ , of order  $q^{m+n+1}$ .

We must note that there is a misprint in the presentation of the last type of groups in Huppert's book [7; Aufgabe III.22].

**Theorem 2** ([13], see also [2]). *Schmidt groups fall into the following classes:*

1.  $G = [P]Q$ , where  $Q = \langle z \rangle$  is cyclic of order  $q^r > 1$ , with  $q$  a prime not dividing  $p - 1$  and  $P$  an irreducible  $Q$ -module over the field of  $p$  elements with kernel  $\langle z^q \rangle$  in  $Q$ .
2.  $G = [P]Q$ , where  $P$  is a non-abelian special  $p$ -group of rank  $2m$ , the order of  $p$  modulo  $q$  being  $2m$ ,  $Q = \langle z \rangle$  is cyclic of order  $q^r > 1$ ,  $z$  induces an automorphism in  $P$  such that  $P/\Phi(P)$  is a faithful irreducible  $Q$ -module, and  $z$  centralises  $\Phi(P)$ . Furthermore,  $|P/\Phi(P)| = p^{2m}$  and  $|P'| \leq p^m$ .
3.  $G = [P]Q$ , where  $P = \langle a \rangle$  is a normal subgroup of order  $p$ ,  $Q = \langle z \rangle$  is cyclic of order  $q^r > 1$ , with  $q$  dividing  $p - 1$ , and  $a^z = a^i$ , where  $i$  is the least primitive  $q$ -th root of unity modulo  $p$ .

Here  $[K]H$  denotes the semidirect product of  $K$  with  $H$ , where  $H$  acts on  $K$ .

Itô [8] considered the minimal non- $p$ -nilpotent groups for a prime  $p$ , which turn out to be Schmidt groups.

Doerk [5] was the first author in studying the minimal non-supersoluble groups. Later, Nagrebeckiĭ [11] classified them.

Let  $p$  be a prime number. A group  $G$  is said to be  $p$ -supersoluble whenever  $G$  is  $p$ -soluble and all  $p$ -chief factors of  $G$  are cyclic groups of order  $p$ .

Kontorovič and Nagrebeckiĭ [9] studied the minimal non- $p$ -supersoluble groups for a prime  $p$  with trivial Frattini subgroup. Tuccillo [15] tried to classify all minimal non- $p$ -supersoluble groups in the soluble case, and gave results about non-soluble minimal non- $p$ -supersoluble groups. Unfortunately, there is a gap in his paper and some groups are missing from his classification.

**Example 3.** The extraspecial group  $N = \langle a, b \rangle$  of order  $41^3$  and exponent 41 has automorphisms  $y$  of order 5 and  $z$  of order 8, given by  $a^y = a^{10}$ ,  $b^y = b^{37}$ , and  $a^z = b^{19}$ ,  $b^z = a^{35}$ , satisfying  $y^z = y^{-1}$ . The semidirect product  $G$  of  $N$  by  $\langle x, y \rangle$  is a minimal non-supersoluble group such that the Frattini subgroup  $\Phi(N)$  of  $N$  is not a central subgroup of  $G$ . This is a minimal non-41-supersoluble group not appearing in any type of Tuccillo's result.

**Example 4.** The extraspecial group  $N = \langle a, b \rangle$  of order  $17^3$  and exponent 17 has an automorphism  $z$  of order 32 given by  $a^z = b$ ,  $b^z = a^3$ . The semidirect product  $G = [N]\langle z \rangle$  is a minimal non-17-supersoluble group. It is clear that  $[a, b]^z = [a, b]^{14}$  and so  $[a, b]$  does not belong to the centre of  $G$ . This is another group missing in Tuccillo's work.

**Example 5.** The automorphism group of the extraspecial group of order  $7^3$  and exponent 7 has a subgroup isomorphic to the symmetric group  $\Sigma_3$  of degree 3. The corresponding semidirect product is a minimal non-7-supersoluble group not corresponding to any case of Tuccillo's work.

**Example 6.** Let  $E = \langle x_1, x_2 \rangle$  be an extraspecial group of order 125 and exponent 5. This group has two automorphisms  $\alpha$  and  $\beta$  given by  $x_1^\alpha = x_2^4$ ,  $x_2^\alpha = x_1$ ,  $x_1^\beta = x_1^2$ , and  $x_2^\beta = x_2^3$  generating a quaternion group  $H$  of order 8 such that the corresponding semidirect product  $[E]H$  is a minimal non-5-supersoluble group. This group is also missing in [15].

**Example 7.** With the same notation as in Example 6, the automorphisms  $\beta$  and  $\gamma$  defined by  $x_1^\gamma = x_2$ ,  $x_2^\gamma = x_1$  generate a dihedral group  $D$  of order 8. The corresponding semidirect product  $[E]D$  is a minimal non-5-supersoluble group not appearing in [15].

By looking at these examples, we see that the classification of minimal non- $p$ -supersoluble groups given in [15] is far from being complete. In our examples, the Frattini subgroup of the Sylow  $p$ -subgroup is not a central subgroup, contrary to the claim in [15; 1.7].

The aim of this paper is to give the complete classification of minimal non- $p$ -supersoluble groups in the  $p$ -soluble universe. This restriction is motivated by the following result.

**Proposition 8.** *Let  $G$  be a minimal non- $p$ -supersoluble group. Then either  $G/\Phi(G)$  is a simple group of order divisible by  $p$ , or  $G$  is  $p$ -soluble.*

Our main theorem is the following:

**Theorem 9.** *The minimal  $p$ -soluble non- $p$ -supersoluble groups for a prime  $p$  are exactly the groups of the following types:*

**Type 1:** *Let  $q$  be a prime number such that  $q$  divides  $p - 1$ . Let  $C$  be a cyclic group of order  $p^s$ , with  $s \geq 1$ , and let  $M$  be an irreducible  $C$ -module over the field of  $q$  elements with kernel the maximal subgroup of  $C$ . Consider a group  $E$  with a normal  $q$ -subgroup  $F$  contained in the Frattini subgroup of  $E$  and  $E/F$  isomorphic to the semidirect product  $[M]C$ . Let  $N$  be an irreducible  $E$ -module over the field of  $p$*

elements with kernel the Frattini subgroup of  $E$ . Let  $G = [N]E$  be the corresponding semidirect product. In this case,  $\Phi(G)_p$ , the Sylow  $p$ -subgroup of  $\Phi(G)$ , which coincides with the Frattini subgroup of a Sylow  $p$ -subgroup of  $E$ , is a central subgroup of  $G$  and  $\Phi(G)_q$ , the Sylow  $q$ -subgroup of  $\Phi(G)$ , is equal to  $\Phi(E)$ , which coincides with the Frattini subgroup of a Sylow  $q$ -subgroup of  $E$  and centralises  $N$ .

**Type 2:**  $G = [P]Q$ , where  $Q = \langle z \rangle$  is cyclic of order  $q^r > 1$ , with  $q$  a prime not dividing  $p - 1$ , and  $P$  is an irreducible  $Q$ -module over the field of  $p$  elements with kernel  $\langle z^q \rangle$  in  $Q$ .

**Type 3:**  $G = [P]Q$ , where  $P$  is a non-abelian special  $p$ -group of rank  $2m$ , the order of  $p$  modulo  $q$  being  $2m$ ,  $q$  is a prime,  $Q = \langle z \rangle$  is cyclic of order  $q^r > 1$ ,  $z$  induces an automorphism in  $P$  such that  $P/\Phi(P)$  is a faithful and irreducible  $Q$ -module, and  $z$  centralises  $\Phi(P)$ . Furthermore,  $|P/\Phi(P)| = p^{2m}$  and  $|P'| \leq p^m$ .

**Type 4:**  $G = [P]Q$ , where  $P = \langle a_0, a_1, \dots, a_{q-1} \rangle$  is an elementary abelian  $p$ -group of order  $p^q$ ,  $Q = \langle z \rangle$  is cyclic of order  $q^r$ , with  $q$  a prime such that  $q^f$  is the highest power of  $q$  dividing  $p - 1$  and  $r > f \geq 1$ . Define  $a_j^z = a_{j+1}$  for  $0 \leq j < q - 1$  and  $a_{q-1}^z = a_0^i$ , where  $i$  is a primitive  $q^f$ -th root of unity modulo  $p$ .

**Type 5:**  $G = [P]Q$ , where  $P = \langle a_0, a_1 \rangle$  is an extraspecial group of order  $p^3$  and exponent  $p$ ,  $Q = \langle z \rangle$  is cyclic of order  $2^r$ , with  $2^f$  the largest power of 2 dividing  $p - 1$  and  $r > f \geq 1$ . Define  $a_1 = a_0^z$  and  $a_1^z = a_0^i x$ , where  $x \in \langle a_0, a_1 \rangle$  and  $i$  is a primitive  $2^f$ -th root of unity modulo  $p$ .

**Type 6:**  $G = [P]E$ , where  $E$  is a 2-group with a normal subgroup  $F$  such that  $F \leq \Phi(E)$  and  $E/F$  is isomorphic to a quaternion group of order 8 and  $P$  is an irreducible module for  $E$  with kernel  $F$  over the field of  $p$  elements of dimension 2, where  $4 \mid p - 1$ .

**Type 7:**  $G = [P]E$ , where  $E$  is a 2-group with a normal subgroup  $F$  such that  $F \leq \Phi(E)$  and  $E/F$  is isomorphic to a quaternion group of order 8,  $P$  is an extraspecial group of order  $p^3$  and exponent  $p$ , where  $4 \mid p - 1$ , and  $P/\Phi(P)$  is an irreducible module for  $E$  with kernel  $F$  over the field of  $p$  elements.

**Type 8:**  $G = [P]E$ , where  $E$  is a  $q$ -group for a prime  $q$  with a normal subgroup  $F$  such that  $F \leq \Phi(E)$  and  $E/F$  is isomorphic to a group  $G_{\Pi}(q, m, 1)$  of Theorem 1,  $P$  is an irreducible  $E$ -module of dimension  $q$  over the field of  $p$  elements with kernel  $F$ , and  $q^m$  divides  $p - 1$ .

**Type 9:**  $G = [P]E$ , where  $E$  is a 2-group with a normal subgroup  $F$  such that  $F \leq \Phi(E)$  and  $E/F$  is isomorphic to a group  $G_{\Pi}(2, m, 1)$  of Theorem 1,  $P$  is an extraspecial group of order  $p^3$  and exponent  $p$  such

that  $P/\Phi(P)$  is an irreducible  $E$ -module of dimension 2 over the field of  $p$  elements with kernel  $F$ , and  $2^m$  divides  $p - 1$ .

**Type 10:**  $G = [P]E$ , where  $E$  is a  $q$ -group for a prime  $q$  with a normal subgroup  $F$  such that  $F \leq \Phi(E)$  and  $E/F$  is isomorphic to an extraspecial group of order  $q^3$  and exponent  $q$ , with  $q$  odd,  $P$  is an irreducible  $E$ -module over the field of  $p$  elements with kernel  $F$  and dimension  $q$ , and  $q$  divides  $p - 1$ .

**Type 11:**  $G = [P]MC$ , where  $C$  is a cyclic subgroup of order  $r^{s+t}$ , with  $r$  a prime number and  $s$  and  $t$  integers such that  $s \geq 1$  and  $t \geq 0$ , normalising a Sylow  $q$ -subgroup  $M$  of  $G$ ,  $M/\Phi(M)$  is an irreducible  $C$ -module over the field of  $q$  elements,  $q$  a prime, with kernel the subgroup  $D$  of order  $r^t$  of  $C$ , and  $P$  is an irreducible  $MC$ -module over the field of  $p$  elements, where  $q$  and  $r^s$  divide  $p - 1$ . In this case,  $\Phi(G)_{p'}$ , the Hall  $p'$ -subgroup of  $\Phi(G)$ , coincides with  $\Phi(M) \times D$  and centralises  $P$ .

**Type 12:**  $G = [P]MC$ , where  $C$  is a cyclic subgroup of order  $2^{s+t}$ , with  $s$  and  $t$  integers such that  $s \geq 1$  and  $t \geq 0$ , normalising a Sylow  $q$ -subgroup  $M$  of  $G$ ,  $q$  a prime,  $M/\Phi(M)$  is an irreducible  $C$ -module over the field of  $q$  elements with kernel the subgroup  $D$  of order  $2^t$  of  $C$ , and  $P$  is an extraspecial group of order  $p^3$  and exponent  $p$  such that  $P/\Phi(P)$  is an irreducible  $MC$ -module over the field of  $p$  elements, where  $q$  and  $2^s$  divide  $p - 1$ . In this case,  $\Phi(G)_{p'}$ , the Hall  $p'$ -subgroup of  $\Phi(G)$ , is equal to  $\Phi(M) \times D$  and centralises  $P$ .

From Proposition 8 and Theorem 9 we deduce immediately that a minimal non- $p$ -supersoluble group is either a Frattini extension of a non-abelian simple group of order divisible by  $p$ , or a soluble group.

As a consequence of Theorem 9, bearing in mind that minimal non-supersoluble groups are soluble by [5] and minimal non- $p$ -supersoluble groups for a prime  $p$ , we obtain the classification of minimal non-supersoluble groups:

**Theorem 10.** *The minimal non-supersoluble groups are exactly the groups of Types 2 to 12 of Theorem 9, with  $r$  dividing  $q - 1$  in the case of groups of Type 11.*

The classification of minimal non- $p$ -supersoluble groups can be applied to get some new criteria for supersolubility. A well-known theorem of Buckley [4] states that if a group  $G$  has odd order and all its subgroups of prime order are normal, then  $G$  is supersoluble. The next generalisation follows easily from our classification:

**Theorem 11.** *Let  $G$  be a group whose subgroups of prime order permute with all Sylow subgroups of  $G$  and no section of  $G$  is isomorphic to the quaternion group of order 8. Then  $G$  is supersoluble.*

As a final remark, we mention that Tuccillo [15] also gave some partial results for Frattini extensions of non-abelian simple groups of order divisible by  $p$ . Looking at the results of Section 4 of that paper, it seems that the classification of minimal non- $p$ -supersoluble groups in the general finite universe is a hard task.

## 2. Preliminary results

First we gather the main properties of a minimal non-supersoluble group. They appear in Doerk's paper [5].

**Theorem 12.** *Let  $G$  be a minimal non-supersoluble group. We have:*

1.  $G$  is soluble.
2.  $G$  has a unique normal Sylow subgroup  $P$ .
3.  $P/\Phi(P)$  is a minimal normal subgroup of  $G/\Phi(P)$ .
4. The Frattini subgroup  $\Phi(P)$  of  $P$  is supersolubly embedded in  $G$ , i. e., there exists a series  $1 = N_0 \leq N_1 \leq \dots \leq N_m = \Phi(P)$  such that  $N_i$  is a normal subgroup of  $G$  and  $|N_i/N_{i-1}|$  is prime for  $1 \leq i \leq m$ .
5.  $\Phi(P) \leq Z(P)$ ; in particular,  $P$  has class at most 2.
6. The derived subgroup  $P'$  of  $P$  has at most exponent  $p$ , where  $p$  is the prime dividing  $|P|$ .
7. For  $p > 2$ ,  $P$  has exponent  $p$ ; for  $p = 2$ ,  $P$  has exponent at most 4.
8. Let  $Q$  be a complement to  $P$  in  $G$ . Then  $Q \cap C_G(P/\Phi(P)) = \Phi(G) \cap \Phi(Q) = \Phi(G) \cap Q$ .
9. If  $\overline{Q} = Q/(Q \cap \Phi(G))$ , then  $\overline{Q}$  is a minimal non-abelian group or a cyclic group of prime power order.

In [6; VII, 6.18], some properties of critical groups for a saturated formation in the soluble universe are given. This result has been extended to the general finite universe by the first author and Pedraza-Aguilera. Recall that if  $\mathfrak{F}$  is a formation, the  $\mathfrak{F}$ -residual of a group  $G$ , denoted by  $G^{\mathfrak{F}}$ , is the smallest normal subgroup of  $G$  such that  $G/G^{\mathfrak{F}}$  belongs to  $\mathfrak{F}$ .

**Lemma 13** ([3; Theorem 1 and Proposition 1]). *Let  $\mathfrak{F}$  be a saturated formation.*

1. *Assume that  $G$  is a group such that  $G$  does not belong to  $\mathfrak{F}$ , but all its proper subgroups belong to  $\mathfrak{F}$ . Then  $F'(G)/\Phi(G)$  is the unique minimal normal subgroup of  $G/\Phi(G)$ , where  $F'(G) = \text{Soc}(G \text{ mod } \Phi(G))$ , and  $F'(G) = G^{\mathfrak{F}}\Phi(G)$ . In addition, if the derived subgroup of  $G^{\mathfrak{F}}$  is a proper subgroup of  $G^{\mathfrak{F}}$ , then  $G^{\mathfrak{F}}$  is a soluble group. Furthermore, if  $G^{\mathfrak{F}}$  is soluble, then  $F'(G) = F(G)$ , the Fitting subgroup of  $G$ . Moreover  $(G^{\mathfrak{F}})' = T \cap G^{\mathfrak{F}}$  for every maximal subgroup  $T$  of  $G$  such that  $G/\text{Core}_G(T) \notin \mathfrak{F}$  and  $F'(G)T = G$ .*
2. *Assume that  $G$  is a group such that  $G$  does not belong to  $\mathfrak{F}$  and there exists a maximal subgroup  $M$  of  $G$  such that  $M \in \mathfrak{F}$  and  $G = MF(G)$ . Then  $G^{\mathfrak{F}}/(G^{\mathfrak{F}})'$  is a chief factor of  $G$ ,  $G^{\mathfrak{F}}$  is a  $p$ -group for some prime  $p$ ,  $G^{\mathfrak{F}}$  has exponent  $p$  if  $p > 2$  and exponent at most 4 if  $p = 2$ . Moreover, either  $G^{\mathfrak{F}}$  is elementary abelian or  $(G^{\mathfrak{F}})' = Z(G^{\mathfrak{F}}) = \Phi(G^{\mathfrak{F}})$  is an elementary abelian group.*

It is clear that the class  $\mathfrak{F}$  of all  $p$ -supersoluble groups for a given prime  $p$  is a saturated formation [7; VI, 8.3]. Thus Lemma 13 applies to this class.

The following series of lemmas is also needed in the proof of Theorem 9.

**Lemma 14.** *Let  $N$  be a non-abelian special normal  $p$ -subgroup of a group  $G$ ,  $p$  a prime, such that  $N/\Phi(N)$  is a minimal normal subgroup of  $G/\Phi(N)$ . Assume that there exists a series  $1 = N_0 \trianglelefteq N_1 \trianglelefteq \dots \trianglelefteq N_t = \Phi(N)$  with  $N_i$  normal in  $G$  for all  $i$  and cyclic factors  $N_i/N_{i-1}$  of order  $p$  for  $1 \leq i \leq t$ . Then  $N/\Phi(N)$  has order  $p^{2m}$  for an integer  $m$ .*

**Proof.** The result holds if  $N$  is extraspecial by [6; A, 20.4]. Assume that  $N$  is not extraspecial. Let  $T = N_1$  be a minimal normal subgroup of  $G$  contained in  $\Phi(N)$ , then  $T$  has order  $p$ . It is clear that  $(N/T)' = N'/T$  and  $\Phi(N/T) = \Phi(N)/T$ . Consequently  $(N/T)' = \Phi(N/T)$ . On the other hand,  $\Phi(N/T) = \Phi(N)/T = Z(N)/T \leq Z(N/T)$ . If  $\Phi(N/T) \neq Z(N/T)$ , then  $Z(N/T) = N/T$  because  $N/\Phi(N)$  is a chief factor of  $G$ , but this implies that  $N/T$  is abelian, in particular,  $T = N'$  and  $N$  is extraspecial, a contradiction. Therefore  $G/T$  satisfies the hypothesis of the lemma and  $N/T$  is non-abelian. By induction,  $(N/T)/\Phi(N/T) \cong N/\Phi(N)$  has order  $p^{2m}$ . ■

**Lemma 15.** *Let  $G$  be a group, and let  $N$  be a normal subgroup of  $G$  contained in  $\Phi(G)$ . If  $p$  is a prime and  $G$  is a minimal non- $p$ -supersoluble group, then  $G/N$  is a minimal non- $p$ -supersoluble group.*

*Conversely, if  $G/N$  is a minimal non- $p$ -supersoluble group,  $N \leq \Phi(G)$ , and there exists a series  $1 = N_0 \trianglelefteq N_1 \trianglelefteq \dots \trianglelefteq N_t = N$  with  $N_i$  normal in  $G$*

for all  $i$  and whose factors  $N_i/N_{i-1}$  are either cyclic of order  $p$  or  $p'$ -groups for  $1 \leq i \leq t$ , then  $G$  is a minimal non- $p$ -supersoluble group.

**Proof.** Assume that  $G$  is a minimal non- $p$ -supersoluble group and  $N \leq \Phi(G)$ . If  $M/N$  is a proper subgroup of  $G/N$ , then  $M$  is a proper subgroup of  $G$ . Hence  $M$  is  $p$ -supersoluble, and so is  $M/N$ . If  $G/N$  were  $p$ -supersoluble, since  $N \leq \Phi(G)$ ,  $G$  would be  $p$ -supersoluble, a contradiction. Therefore  $G/N$  is minimal non- $p$ -supersoluble.

Conversely, assume that  $G/N$  is a minimal non- $p$ -supersoluble group,  $N \leq \Phi(G)$ , and that there exists a series  $1 = N_0 \triangleleft N_1 \triangleleft \cdots \triangleleft N_t = N$  with  $N_i$  normal in  $G$  for all  $i$  and factors  $N_i/N_{i-1}$  cyclic of order  $p$  or  $p'$ -groups for  $1 \leq i \leq t$ . It is clear that  $G$  cannot be  $p$ -supersoluble. Let  $M$  be a maximal subgroup of  $G$ . Since  $N \leq \Phi(G)$ ,  $N \leq M$ . Thus  $M/N$  is  $p$ -supersoluble. On the other hand, it is clear that every chief factor of  $M$  below  $N$  is either a  $p'$ -group or a cyclic group of order  $p$ . Consequently,  $M$  is  $p$ -supersoluble. ■

**Lemma 16** ([1]). *Let  $A$  be a group, and let  $B$  be a normal subgroup of  $A$  of prime index  $r$  dividing  $p - 1$ ,  $p$  a prime. If  $M$  is an irreducible and faithful  $A$ -module over  $\text{GF}(p)$  of dimension greater than 1 and the restriction of  $M$  to  $B$  is a sum of irreducible  $B$ -modules of dimension 1, then  $M$  has dimension  $r$ . In this case,  $M$  is isomorphic to the induced module of one of the direct summands of  $M_B$  from  $B$  up to  $A$ .*

In the rest of the paper,  $\mathfrak{F}$  will denote the formation of all  $p$ -supersoluble groups,  $p$  a prime.

**Lemma 17.** *Let  $G$  be a minimal non- $p$ -supersoluble group whose  $p$ -supersoluble residual  $N = G^{\mathfrak{F}}$  is normal Sylow  $p$ -subgroup. Then a Hall  $p'$ -subgroup  $R/\Phi(G)$  of  $G/\Phi(G)$  is either cyclic of prime power order or a minimal non-abelian group.*

**Proof.** By Lemma 15, we can assume without loss of generality that  $\Phi(G) = 1$ . Then, by Lemma 13,  $G$  is a primitive group and  $C_G(N) = N$ . In particular, for each subgroup  $X$  of  $G$ , we have that  $O_{p',p}(XN) = N$ . Let  $M$  be a maximal subgroup of  $R$ . Then  $MN$  is a  $p$ -supersoluble group and so  $MN/O_{p',p}(MN) = MN/N$  is abelian of exponent dividing  $p - 1$ . Therefore if  $R$  is non-abelian, then it is a minimal non-abelian group. Suppose that  $R$  is abelian. If  $R$  has a unique maximal subgroup, then  $R$  is cyclic of prime power order. Assume now that  $R$  has at least two different maximal subgroups. Then  $R$  is a product of two subgroups of exponent dividing  $p - 1$ . Consequently  $R$  has exponent  $p - 1$  and so  $N$  is a cyclic group of order  $p$  by [6; B, 9.8], a contradiction. Therefore if  $R$  is not cyclic of prime power order,  $R$  must be a minimal non-abelian group and the lemma is proved. ■



**Lemma 18.** *Let  $G$  be a minimal non- $p$ -supersoluble group with a normal Sylow  $p$ -subgroup  $N$  such that  $G/\Phi(N)$  is a Schmidt group. Then  $G$  is a Schmidt group.*

**Proof.** Let  $G$  be a minimal non- $p$ -supersoluble group with a normal Sylow  $p$ -subgroup  $N$  such that  $G/\Phi(N)$  is a Schmidt group. Then  $G = NQ$ , for a Hall  $p'$ -subgroup  $Q$  of  $G$ . Moreover, since  $G$  is not  $p$ -supersoluble and  $G/\Phi(N)$  is a Schmidt group, we have that  $Q$  is a cyclic  $q$ -group for a prime  $q$  and  $q$  does not divide  $p - 1$  by Theorem 2. Let  $M$  be a maximal subgroup of  $G$ . If  $N$  is not contained in  $M$ , then a conjugate of  $Q$  is contained in  $M$  and so we can assume without loss of generality that  $M = \Phi(N)Q$ . Since  $q$  does not divide  $p - 1$  and  $M$  is  $p$ -supersoluble, we have that  $Q$  centralises all chief factors of a chief series of  $M$  passing through  $\Phi(N)$ . But by [6; A, 12.4], it follows that  $Q$  centralises  $\Phi(N)$  by and so  $M$  is nilpotent. If  $N$  is contained in  $M$ , then  $M$  is a normal subgroup of  $G$  such that  $M/\Phi(N)$  is nilpotent. By [7; III, 3.5], it follows that  $M$  is nilpotent. This completes the proof. ■

### 3. Proof of the main theorems

**Proof of Proposition 8.** By Lemma 13,  $G/\Phi(G)$  has a unique minimal normal subgroup  $T/\Phi(G)$  and  $T = G^{\mathfrak{S}}\Phi(G)$ . It follows that  $T/\Phi(G)$  must have order divisible by  $p$ . Assume that  $T/\Phi(G)$  is a direct product of non-abelian simple groups. We note that, since  $G/\Phi(G)$  is a minimal non- $p$ -supersoluble group by Lemma 15,  $T/\Phi(G) = G/\Phi(G)$  and so  $G/\Phi(G)$  is a simple non-abelian group.

Assume now that  $T/\Phi(G)$  is a  $p$ -group. By Lemma 13, we have that  $G^{\mathfrak{S}}$  is a  $p$ -group. In this case,  $T/\Phi(G)$  is complemented by a maximal subgroup  $M/\Phi(G)$  of  $G/\Phi(G)$ . Since  $M$  is  $p$ -supersoluble, so is  $M/\Phi(G)$ . Therefore  $G/\Phi(G)$  is  $p$ -soluble. It follows that  $G$  is  $p$ -soluble. ■

**Proof of Theorem 9.** Assume that  $G$  is a  $p$ -soluble minimal non- $p$ -supersoluble group. By Lemma 13 and Proposition 8,  $N = G^{\mathfrak{S}}$  is a  $p$ -group.

Assume first that  $N$  is not a Sylow subgroup of  $G$ . By Lemma 13,  $N/\Phi(N)$  is non-cyclic.

Assume that  $\Phi(G) = 1$ . Then  $N$  is the unique minimal normal subgroup of  $G$ , which is an elementary abelian  $p$ -group, and it is complemented by a subgroup,  $R$  say. Moreover,  $N$  is self-centralising in  $G$ . This implies that  $O_{p',p}(G) = N = O_p(G)$ . Since  $N$  is not a Sylow  $p$ -subgroup of  $G$ , we have that  $p$  divides the order of  $R$ . Consider a maximal normal subgroup  $M$  of  $R$ . Observe that  $NM$  is a  $p$ -supersoluble group and  $O_{p',p}(NM) = O_p(NM) = N$  because  $O_p(M)$  is contained in  $O_p(R) = 1$ . Therefore  $M \cong MN/O_{p',p}(MN)$

is abelian of exponent dividing  $p - 1$ . It follows that  $M$  is a normal Hall  $p'$ -subgroup of  $R$  and  $|R : M| = p$  because  $p$  divides  $|R|$ . In particular,  $M$  is the only maximal normal subgroup of  $R$ . Moreover, if  $C$  is a Sylow  $p$ -subgroup of  $R$ , then  $C$  is a cyclic group of order  $p$ .

Let  $M_0$  be a normal subgroup of  $R$  such that  $M/M_0$  is a chief factor of  $R$ . Let  $X = NM_0C$ . Since  $X$  is a proper subgroup of  $G$ , we have that  $X$  is  $p$ -supersoluble. Hence  $X/O_{p',p}(X)$  is an abelian group of exponent dividing  $p - 1$ . It follows that  $C \leq O_{p',p}(X)$ . In particular,  $C = M_0C \cap O_{p',p}(X)$  is a normal subgroup of  $M_0C$  which intersects trivially  $M_0$ . We conclude that  $C$  centralises  $M_0$ . If  $M_1$  is another normal subgroup of  $R$  such that  $M/M_1$  is a chief factor of  $R$ , then  $M = M_0M_1$ . The same argument shows that  $C$  centralises  $M_1$  and so  $C$  centralises  $M$  as well, a contradiction because in this case  $C \leq Z(R)$  and then  $C \leq O_p(R) = 1$ . Consequently  $M_0$  is the unique such normal subgroup. Since  $M$  is abelian, we have that  $M_0 \leq Z(R)$ .

Now  $R$  has an irreducible and faithful module  $N$  over  $\text{GF}(p)$ . By [6; B, 9.4],  $Z(R)$  is cyclic. In particular,  $M_0$  is cyclic. We will prove next that  $M_0 = 1$ . In order to do so, assume, by way of contradiction, that  $M$  is not a minimal normal subgroup of  $R$ . First of all, if  $M$  is not a  $q$ -group for a prime  $q$ , then  $M$  is a direct product of its Sylow subgroups, but all of them should be contained in  $M_0$ , a contradiction. Therefore,  $M$  is a  $q$ -group for a prime  $q$ . Since  $M$  has exponent dividing  $p - 1$ , we have that  $q$  divides  $p - 1$ . If  $\text{Soc}(M)$  is a proper normal subgroup of  $M$ , then  $\text{Soc}(M) \leq M_0$ . Since  $M_0$  is cyclic, we have that  $M$  is an abelian group with a cyclic socle. Therefore  $M$  is cyclic. But since  $q$  divides  $p - 1$ , we have that  $C$  centralises  $M$  and so  $C \leq O_p(R) = 1$ , a contradiction. Consequently  $M = \text{Soc}(M)$ , and  $M$  is a  $C$ -module over  $\text{GF}(q)$ . If  $M$  is not irreducible as  $C$ -module, then  $M$  can be expressed as a direct sum of proper  $C$ -modules over  $\text{GF}(q)$ . Hence  $M$  has at least two maximal  $C$ -submodules, which yield two different chief factors  $M/M_1$  and  $M/M_2$  of  $R$ , a contradiction. Therefore  $M$  is a minimal normal subgroup of  $R$ ,  $R = MC$ , and  $C_R(M) = M$ . On the other hand,  $N$  is a faithful and irreducible  $R$ -module over  $\text{GF}(p)$ . By Clifford's theorem [6; B, 7.3], the restriction of  $N$  to  $M$  is a direct sum of  $|R : T|$  homogeneous components, where  $T$  is the inertia subgroup of one of the irreducible components of  $N$  when regarded as an  $M$ -module. Moreover, by [6; B, 8.3], we have that each of these homogeneous components  $N_i$  is irreducible. Therefore they have dimension 1 because  $N_iM$  is supersoluble for every  $i$ . Since  $N$  is not cyclic, we have that  $|R : T| > 1$ . Since  $M \leq T \leq R$ , we have that  $M = T$  and so  $N$  has order  $p^p$ .

Assume now that  $\Phi(G) \neq 1$ . In this case,  $\overline{G} = G/\Phi(G)$  is a minimal non- $p$ -supersoluble group by Lemma 15 and  $\Phi(\overline{G}) = 1$ . We observe that  $N\Phi(G)/\Phi(G)$  cannot be a Sylow  $p$ -subgroup of  $G/\Phi(G)$ , because otherwise

$NH$ , where  $H$  is a Hall  $p'$ -subgroup of  $G$ , would be a proper supplement to  $\Phi(G)$  in  $G$ , which is impossible. In particular, if  $T$  is a normal subgroup of  $G$  contained in  $\Phi(G)$ , then the  $p$ -supersoluble residual  $NT/T$  of  $G/T$  is not a Sylow  $p$ -subgroup of  $G/T$ . Therefore  $\overline{G}$  has the above structure. Since

$$N\Phi(G) = F(G), \quad F(G/\Phi(G)) = F(G)/\Phi(G), \quad \text{and} \quad \Phi(F(G)/\Phi(G)) = 1,$$

we have that  $\overline{N} = (\overline{G})^{\mathfrak{F}} = N\Phi(G)/\Phi(G)$  satisfies

$$\begin{aligned} \overline{N}/\Phi(\overline{N}) &= (N\Phi(G)/\Phi(G))/\Phi(N\Phi(G)/\Phi(G)) \\ &= (F(G)/\Phi(G))/\Phi(F(G)/\Phi(G)), \end{aligned}$$

which is isomorphic to  $F(G)/\Phi(G) = N\Phi(G)/\Phi(G)$ , and the latter is  $G$ -isomorphic to  $N/(N \cap \Phi(G)) = N/\Phi(N)$  by Lemma 13. Assume that  $\Phi(N) \neq 1$ . By Lemma 14, we have that  $N/\Phi(N)$  has square order. But this order is equal to  $|\overline{N}/\Phi(\overline{N})| = p^2$ , which implies that  $p = 2$ . This contradicts the fact that  $q$  divides  $p - 1$ . Therefore  $\Phi(N) = 1$ . Now we will prove that  $\Phi(G)_p$ , the Sylow  $p$ -subgroup of  $\Phi(G)$ , is a central cyclic subgroup of  $G$ . Assume first that  $\Phi(G)_{p'}$ , the Hall  $p'$ -subgroup of  $\Phi(G)$ , is trivial. We have that  $G/\Phi(G) = \overline{N}\overline{M}\overline{C}$ , where  $\overline{C}$  is a cyclic group of order  $p$ ,  $\overline{M}$  is an irreducible and faithful module for  $\overline{C}$  over  $\text{GF}(q)$ ,  $q$  a prime dividing  $p - 1$ , and  $\overline{N}$  is an irreducible and faithful module for  $\overline{M}\overline{C}$  over  $\text{GF}(p)$  of dimension  $p$ . Let  $N$ ,  $M$ , and  $C$  be, respectively, preimages of  $\overline{N}$ ,  $\overline{M}$ , and  $\overline{C}$  by the canonical epimorphism from  $G$  to  $G/T$ . We can assume that  $N = G^{\mathfrak{F}}$  and  $M$  is a Sylow  $q$ -subgroup of  $G$ . Since  $\overline{C}$  is cyclic of order  $p$ , we can find a cyclic subgroup  $C$  of  $G$  such that  $\overline{C} = C\Phi(G)/\Phi(G)$ . Consider now a chief factor  $H/K$  of  $G$  contained in  $\Phi(G)_p$ . Then  $G/C_G(H/K)$  is an abelian group of exponent dividing  $p - 1$  and  $H/K$  is centralised by a Sylow  $p$ -subgroup of  $G/K$ ; in particular,  $G/C_G(H/K)$  is isomorphic to a factor group of a group with a unique normal subgroup of index  $p$ . It follows that  $C_G(H/K) = G$ , that is,  $H/K$  is a central factor of  $G$ . Now  $N$  centralises  $\Phi(G)$  because  $\Phi(N) = 1 = N \cap \Phi(G)$  and  $M$  is a  $q$ -group stabilising a series of  $\Phi(G)$ . By [6; A, 12.4],  $M$  centralises  $\Phi(G)$ . Moreover  $C$  normalises  $M$  because  $M\Phi(G) = M \times \Phi(G)$  is normalised by  $C$ . In particular,  $MC$  is a subgroup of  $G$ . Since  $G = N(MC)$  and  $N$  is a minimal normal subgroup of  $G$ , it follows that  $MC$  is a maximal subgroup of  $G$ . Hence  $\Phi(G)$  is contained in  $MC$  and so in  $C$ . This implies that  $\Phi(C) \leq Z(G)$ . In the general case, we have that

$$\Phi(G)/\Phi(G)_{p'} \leq Z(G/\Phi(G)_{p'}).$$

Then  $[G, \Phi(G)_p] \leq \Phi(G)_{p'}$ . Therefore  $\Phi(G)_p \leq Z(G)$ . On the other hand, it is clear that  $\Phi(G)_p$  is a proper subgroup of  $C$ . Thus  $\Phi(G)_p \leq \Phi(C)$  and so  $\Phi(G)_p \leq \Phi(MC)$ . Now  $\Phi(G)_{p'} = \Phi(G)_q$ , the Sylow  $q$ -subgroup of  $\Phi(G)$ ,

is contained in  $M$  and  $M/\Phi(G)_{p'}$  is elementary abelian. Hence  $\Phi(M) \leq \Phi(G)_{p'}$ . Moreover, by Maschke's theorem [6; A, 11.4], the elementary abelian group  $M/\Phi(M)$  admits a decomposition

$$M/\Phi(M) = \Phi(G)_{p'}/\Phi(M) \times A/\Phi(M),$$

where  $A$  is normalised by  $C$ . In this case,  $R = MC = A(C\Phi(G)_{p'})$ . Since  $C$  normalises  $A$ , we have that  $AC$  is a subgroup of  $G$ . Therefore  $N(AC)$  is a subgroup of  $G$  and so  $G = (NAC)\Phi(G)_{p'}$ . We conclude that  $G = NAC$ . By order considerations, we have that  $M = A$  and so  $\Phi(M) = \Phi(G)_{p'}$ .

Now let  $G$  be a minimal non- $p$ -supersoluble group such that  $N$  is a Sylow  $p$ -subgroup of  $G$ . Let  $Q$  be a Hall  $p'$ -subgroup of  $G$ . Then  $G = NQ$ . Denote with bars the images in  $\overline{G} = G/\Phi(G)$ . By Lemma 13,  $\overline{N} = N\Phi(G)/\Phi(G)$  is a minimal normal subgroup of  $\overline{G} = G/\Phi(G)$  and either  $N$  is elementary abelian, or  $N' = Z(N) = \Phi(N)$ . Note that  $\Phi(N) = \Phi(G)_p$ , the Sylow  $p$ -subgroup of  $\Phi(G)$ , because  $\Phi(N)$  is contained in  $\Phi(G)_p$  and  $\overline{N}$  is a chief factor of  $G$ . Assume that  $\Phi(G)_{p'}$ , the Hall  $p'$ -subgroup of  $\Phi(G)$ , is not contained in  $\Phi(Q)$ . Then there exists a maximal subgroup  $A$  of  $Q$  such that  $Q = A\Phi(G)_{p'}$ . In this case,  $G = NQ = NA\Phi(G)_{p'}$  and so  $G = NA$ . It follows that  $A = Q$  by order considerations, a contradiction. Therefore  $\Phi(G)_{p'} \leq \Phi(Q)$ . We also note that since

$$\overline{Q} = Q\Phi(G)/\Phi(G) \cong Q/\Phi(G)_q,$$

where  $\Phi(G)_q$  is the Sylow  $q$ -subgroup of  $\Phi(G)$ , has an irreducible and faithful module  $\overline{N} = N/\Phi(N)$  over  $\text{GF}(p)$ , we have that  $Z(\overline{Q})$  is cyclic by [6; B, 9.4].

By Lemma 17 we have that the Hall  $p'$ -subgroup  $\overline{Q}$  of  $\overline{G}$  is either a cyclic group of prime power order or a minimal non-abelian group.

Suppose that  $\overline{Q} = \langle \bar{z} \rangle$  is a cyclic group of order a power of a prime number,  $q$  say. Since this group is isomorphic to  $Q/\Phi(G)_q$  and  $\Phi(G)_q \leq \Phi(Q)$ , we have that  $Q$  is a cyclic group of  $q$ -power order,  $Q = \langle z \rangle$  say.

Suppose that the order of  $\bar{z}$  is  $q^f$ . Then  $q^{f-1}$  divides  $p-1$ . If  $\bar{z}^q = 1$ , then  $\overline{G}$  is a Schmidt group. By Lemma 18,  $G$  is a Schmidt group. By Theorem 2,  $G$  is a group of Type 2 if  $\Phi(N) = 1$ , or 3 if  $\Phi(N) \neq 1$ .

Assume now that  $f \geq 2$ . In this case,  $q$  divides  $p-1$  and, by Lemma 16, we have that  $\overline{N}$  has order  $p^q$ . Let  $a_0 \in \overline{N} \setminus 1$ . Let  $a_i = a_0^{z^i}$  for  $1 \leq i \leq q-1$ , then  $a_0^{z^q} = a_0^i$ , where  $i$  is a  $q^{f-1}$ -root of unity modulo  $p$ . It follows that

$$(a_0^{z^{q^{f-1}}}) = a_0^{i^{q^{f-2}}}.$$

If  $i$  is not a primitive  $q^{f-1}$ -th root of unity modulo  $p$ , we have that  $i^{q^{f-2}} \equiv 1 \pmod{p}$ . In particular,  $a_0^{z^{q^{f-1}}} = a_0$ , which contradicts the fact that the order

of  $\bar{z}$  is  $q^f$ . If  $\Phi(N) = 1$ , then we obtain a group of Type 4. If  $\Phi(N) \neq 1$ , then  $\bar{N}$  has square order by Lemma 14 and so  $q = 2$ . Hence  $N$  is an extraspecial group of order  $p^3$  and exponent 3, and  $G$  is a group of Type 5.

Assume now that  $\bar{Q}$  is not cyclic. In this case,  $\bar{Q}$  is a minimal non-abelian group by Lemma 17. Let  $x$  be an element of  $\bar{Q}$ . Since  $\bar{N}\langle x \rangle$  is a  $p$ -supersoluble group, we have that the order of  $x$  divides  $p-1$ . It follows that the exponent of  $\bar{Q}$  divides  $p-1$ . Since  $\bar{N} = N/\Phi(N)$  is an irreducible and faithful  $\bar{Q}$ -module over  $\text{GF}(p)$  of dimension greater than 1 and the restriction of  $\bar{N}$  to every maximal subgroup of  $\bar{Q}$  is a sum of irreducible modules of dimension 1, we have that  $\bar{N}$  has order  $p^q$  by Lemma 16.

Suppose that  $\bar{Q}$  is a  $q$ -group for a prime  $q$ . By Theorem 1,

$$\text{either } \bar{Q} \cong Q_8, \quad \text{or } \bar{Q} \cong G_{\text{II}}(q, m, n), \quad \text{or } \bar{Q} \cong G_{\text{III}}(q, m, n).$$

Suppose that  $\bar{Q}$  is isomorphic to a quaternion group  $Q_8$  of order 8. In this case,  $q = 2$ ,  $|\bar{N}| = p^2$  and  $\exp(\bar{Q}) = 4$  divides  $p-1$ . If  $\Phi(N) = 1$ , then we have a group of Type 6. Assume that  $\Phi(N) \neq 1$ . In this case,  $N$  is an extraspecial group of order  $p^3$  and exponent  $p$  and so  $G$  is a group of Type 7.

Suppose that  $\bar{Q}$  is isomorphic to

$$G_{\text{II}}(q, m, n) = \langle a, b \mid a^{q^m} = b^{q^n} = 1, a^b = a^{1+q^{m-1}} \rangle,$$

where  $m \geq 2$ ,  $n \geq 1$ , of order  $q^{m+n}$ . Since  $\bar{Q}$  has an irreducible and faithful module  $\bar{N}$ , we have that  $Z(\bar{Q})$  is cyclic by [6; B, 9.4]. Since  $\langle a^p, b^p \rangle \leq Z(\bar{Q})$  and  $m \geq 2$ , we have that  $b^p = 1$  and so  $n = 1$ . Hence  $q^m$  divides  $p-1$ . If  $\Phi(N) = 1$ , then we obtain a group of Type 8. If  $\Phi(N) \neq 1$ , then  $N$  is non-abelian and so  $|\bar{N}|$  is a square by Lemma 14. It follows that  $q = 2$  and  $G$  is a group of Type 9.

Suppose now that  $\bar{Q}$  is isomorphic to

$$G_{\text{III}}(q, m, n) = \langle a, b \mid a^{q^m} = b^{q^n} = [a, b]^q = [a, b, a] = [a, b, b] = 1 \rangle,$$

where  $m \geq n \geq 1$ , of order  $q^{m+n+1}$ . Since  $G_{\text{III}}(2, 1, 1) \cong G_{\text{II}}(2, 2, 1)$ , we can assume that  $(q, m, n) \neq (2, 1, 1)$ .

As before,  $Z(\bar{Q})$  is cyclic. Consider  $\langle a^q, b^q, [a, b] \rangle$ , which is contained in  $Z(\bar{Q})$ . If  $m \geq 2$ , then  $\langle a^q, [a, b] \rangle$  is cyclic. Since  $[a, b]$  has order  $p$ , we have that  $[a, b] = a^{qt}$  for a natural number  $t$ . But hence  $a^b = a^{1+qt}$  and so  $\langle a \rangle$  is a normal subgroup of  $G$ . Therefore  $|\bar{Q}| = |\langle a, b \rangle| = |\langle a \rangle \langle b \rangle| \leq q^{m+n}$ , a contradiction. Consequently  $m = 1$ . It follows that  $\bar{Q}$  is an extraspecial group of order  $q^3$  and exponent  $q$ . If  $\Phi(N) \neq 1$ , then  $\bar{N}$  has square order, but this implies that  $q = 2$ , a contradiction. Consequently,  $\Phi(N) = 1$  and we have a group of Type 10.

Assume now that  $\overline{Q}$  is a minimal non-abelian group which is not a  $q$ -group for any prime  $q$ . Then  $\overline{Q}$  is isomorphic to  $[V_q]C_{r^s}$ , where  $q$  and  $r$  are different primes numbers,  $s$  is a positive integer, and  $V_q$  is an irreducible  $C_{r^s}$ -module over the field of  $q$  elements with kernel the maximal subgroup of  $C_{r^s}$ . Since  $\overline{NV}_q$  is a  $p$ -supersoluble subgroup, it follows that the restriction of  $\overline{N}$  to  $V_q$  can be expressed as a direct sum of irreducible modules of dimension 1. By Lemma 16, we have that  $\overline{N}$  has dimension  $r$ . We know that  $\Phi(G)_{p'} \leq \Phi(Q)$  and  $\Phi(G)_p = \Phi(N)$ . Since  $\overline{Q}$  is isomorphic to  $Q/\Phi(G)_{p'}$ , and this group is  $r$ -nilpotent,  $Q$  is  $r$ -nilpotent. Consequently  $Q$  has a normal Sylow  $q$ -subgroup  $M$ . On the other hand,  $\Phi(G)_q$ , the Sylow  $q$ -subgroup of  $\Phi(G)$ , is contained in  $M$  and  $M/\Phi(G)_q$  is elementary abelian. This implies that  $\Phi(M)$  is contained in  $\Phi(G)_q$ . Let  $C$  be a Sylow  $r$ -subgroup of  $G$ . Then, by Maschke's theorem [6; A, 11.4],

$$M/\Phi(M) = \Phi(G)_q/\Phi(M) \times A/\Phi(M)$$

for a subgroup  $A$  of  $M$  normalised by  $C$ . Then  $Q = (AC)\Phi(G)_q = AC$  and so  $A = M$ . Consequently  $\Phi(M) = \Phi(G)_q$ . Now the Sylow  $r$ -subgroup  $\Phi(G)_r$  of  $\Phi(G)$  is contained in  $C$ . If  $\Phi(G)_r$  were not contained in  $\Phi(C)$ , there would exist a maximal subgroup  $T$  of  $C$  such that  $C = T\Phi(G)_r$ . This would imply  $Q = MT$  and  $T = C$ , a contradiction. Hence  $\Phi(G)_r$  is contained in  $\Phi(C)$  and  $C$  is cyclic. Moreover  $\Phi(G)_r$  centralises  $M$ .

If  $\Phi(N) = 1$ , then we have a group of Type 11. If  $\Phi(N) \neq 1$ , then  $r = 2$  and  $N$  is an extraspecial group of order  $p^3$  and exponent  $p$ . This is a group of Type 12.

Conversely, it is clear that the groups of Types 1 to 12 are minimal non- $p$ -supersoluble. ■

**Proof of Theorem 10.** It is clear that all groups of the statement of the theorem are minimal non-supersoluble. Conversely, assume that a group is minimal non-supersoluble. Hence it is soluble, and so its  $p$ -supersoluble residual is a  $p$ -group by Proposition 8. Note that groups of Type 1 in Theorem 9 are not minimal non-supersoluble. On the other hand, groups of Type 11 are not minimal non-supersoluble when  $r$  does not divide  $q - 1$ , because in this case the subgroup  $MC$  is not supersoluble. ■

**Proof of Theorem 11.** Assume that the result is false. Choose for  $G$  a counterexample of least order. Since the property of the statement is inherited by subgroups, it is clear that  $G$  must be a minimal non-supersoluble group, and so a minimal non- $p$ -supersoluble group for a prime  $p$ . In particular, the  $p$ -supersoluble residual  $N = G^{\mathfrak{F}}$  of  $G$  is a  $p$ -group. Suppose that  $N$  has exponent  $p$ . The hypothesis implies that every subgroup of  $N$  is normalised by  $O^p(G)$ . In particular,  $N/\Phi(N)$  is cyclic, a contradiction.

Consequently  $p = 2$  and the exponent of  $N$  is 4. By Theorem 9, the only group with  $\mathfrak{F}$ -residual of exponent 4 is a group of Type 3. But in this case either  $N/\Phi(N)$  has order 4 and  $N$  must be isomorphic to the quaternion group of order 8, because the dihedral group of order 8 does not have any automorphism of odd order, or  $N/\Phi(N)$  has order greater than 4. In the last case,  $N$  has an extraspecial quotient, which has a section isomorphic to a quaternion group of order 8, final contradiction. ■

### Acknowledgement

The authors are indebted to the referee for his/her helpful suggestions.

### References

- [1] BALLESTER-BOLINCHES, A. AND COSSEY, J.: On finite groups whose subgroups are either supersoluble or subnormal. Preprint.
- [2] BALLESTER-BOLINCHES, A., ESTEBAN-ROMERO, R. AND ROBINSON, D. J. S.: On finite minimal non-nilpotent groups. *Proc. Amer. Math. Soc.* **133** (2005), no. 12, 3455–3462 (electronic).
- [3] BALLESTER-BOLINCHES, A. AND PEDRAZA-AGUILERA, M. C.: On minimal subgroups of finite groups. *Acta Math. Hungar.* **73** (1996), 335–342.
- [4] BUCKLEY, J. Finite groups whose minimal subgroups are normal. *Math. Z.* **116** (1970), 15–17.
- [5] DOERK, K.: Minimal nicht überauflösbare, endliche Gruppen. *Math. Z.* **91** (1966), 198–205.
- [6] DOERK, K. AND HAWKES, T.: *Finite Soluble Groups*. De Gruyter Expositions in Mathematics **4**. Walter de Gruyter, Berlin, 1992.
- [7] HUPPERT, B.: *Endliche Gruppen. I*. Die Grundlehren der Mathematischen Wissenschaften **134**. Springer-Verlag, Berlin-New York, 1967.
- [8] ITÔ, N.: Note on  $(LM)$ -groups of finite orders. *Kōdai Math. Sem. Rep.* **3** (1951), 1–6.
- [9] KONTOROVICH, N. P. AND NAGREBECKII, V. T.: Finite minimal not  $p$ -supersolvable groups. *Ural. Gos. Univ. Mat. Zap.* **9** (1975), 53–59, 134–135.
- [10] MILLER, G. A. AND MORENO, H. C.: Non-abelian groups in which every subgroup is abelian. *Trans. Amer. Math. Soc.* **4** (1903), 398–404.
- [11] NAGREBECKII, V. T.: Finite minimal non-supersolvable groups. In *Finite groups (Proc. Gomel Sem., 1973/74)* (Russian), 104–108, 229. Izdat. “Nauka i Tehnika”, Minsk, 1975.
- [12] RÉDEI, L.: Das “schiefe Produkt” in der Gruppentheorie mit Anwendung auf die endlichen nichtkommutativen Gruppen mit lauter kommutativen echten Untergruppen und die Ordnungszahlen, zu denen nur kommutative Gruppen gehören. *Comment. Math. Helv.* **20** (1947), 225–264.

- [13] RÉDEI, L.: Die endlichen einstufig nichtnilpotenten Gruppen. *Publ. Math. Debrecen* **4** (1956), 303–324.
- [14] SCHMIDT, O. J.: Über Gruppen, deren sämtliche Teiler spezielle Gruppen sind. *Mat. Sbornik* **31** (1924), 366–372.
- [15] TUCCILLO, F.: On finite minimal non- $p$ -supersoluble groups. *Colloq. Math.* **63** (1992), 119–131.

*Recibido:* 15 de febrero de 2005

*Revisado:* 4 de mayo de 2005

Adolfo Ballester-Bolinchés  
Departament d'Àlgebra  
Universitat de València  
Dr. Moliner, 50  
E-46100 Burjassot, València, Spain  
Adolfo.Ballester@uv.es

Ramón Esteban-Romero  
Departament de Matemàtica Aplicada  
Institut de Matemàtica Pura i Aplicada  
Universitat Politècnica de València  
Camí de Vera, s/n  
E-46022 València, Spain.  
resteban@mat.upv.es