

Infinite metacyclic subgroups of the mapping class group

Pankaj Kapari, Kashyap Rajeevsarathy, and Apeksha Sanghi

Abstract. For $g \geq 2$, let $\text{Mod}(S_g)$ be the mapping class group of the closed orientable surface S_g of genus g . In this paper, we provide necessary and sufficient conditions for a pair of elements in $\text{Mod}(S_g)$ to generate an infinite metacyclic subgroup. In particular, we provide necessary and sufficient conditions under which a pseudo-Anosov mapping class generates an infinite metacyclic subgroup of $\text{Mod}(S_g)$ with a nontrivial periodic mapping class. As applications of our main results, we establish the existence of infinite metacyclic subgroups of $\text{Mod}(S_g)$ isomorphic to $\mathbb{Z} \rtimes \mathbb{Z}_m$, $\mathbb{Z}_n \rtimes \mathbb{Z}$, and $\mathbb{Z} \rtimes \mathbb{Z}$. Furthermore, we derive bounds on the order of a nontrivial periodic generator of an infinite metacyclic subgroup of $\text{Mod}(S_g)$ that are realized. Finally, we show that the centralizer of an irreducible periodic mapping class F is either $\langle F \rangle$ or $\langle F \rangle \times \langle i \rangle$, where i is a hyperelliptic involution.

1. Introduction

Let $\text{Mod}(S_g)$ be the mapping class group of the closed orientable surface S_g of genus $g \geq 2$. A metacyclic group is an extension of a cyclic group by a cyclic group. Given $F, G \in \text{Mod}(S_g)$, it is natural to ask the following question: Can one derive necessary and sufficient conditions under which F and G generate a metacyclic subgroup of $\text{Mod}(S_g)$? Ivanov (see [14, Theorem 7.5A]) derived necessary and sufficient conditions under which two pure mapping classes commute in $\text{Mod}(S_g)$. Subsequently, the finite abelian subgroups of $\text{Mod}(S_g)$ have been extensively studied [4, 11, 21]. Furthermore, in [5, 6, 30], the question (posed earlier) has been answered in the affirmative for finite metacyclic subgroups of $\text{Mod}(S_g)$ up to conjugacy of their generators. Moreover, it was shown in [6] that for $g \geq 5$, $\text{Mod}(S_g)$ has an infinite metacyclic subgroup generated by a bounding pair map and an involution. Taking inspiration from these works, in this paper, we settle this question for infinite metacyclic subgroups of $\text{Mod}(S_g)$.

A *multicurve* in S_g is a nonempty collection of isotopy classes of pairwise disjoint essential simple closed curves. A left-handed (or positive) Dehn twist about a simple closed curve c will be denoted by T_c . Given a multicurve $C = \{c_1, c_2, \dots, c_\ell\}$ in S_g and nonzero integers q_i , for $1 \leq i \leq \ell$, a mapping class of the form $T_{c_1}^{q_1} T_{c_2}^{q_2} \cdots T_{c_\ell}^{q_\ell}$ is said to be a *multitwist* about C . The Nielsen–Thurston classification [33] asserts that each mapping class in $\text{Mod}(S_g)$ is either periodic, reducible, or pseudo-Anosov. Furthermore,

Mathematics Subject Classification 2020: 57M60 (primary); 57K20 (secondary).

Keywords: surface, pseudo-periodic mapping class, pseudo-Anosov mapping class, metacyclic group.

a pseudo-Anosov mapping class is neither periodic nor reducible. The intersection of all maximal reduction systems of a reducible mapping class F is called its *canonical reduction system*, which we denote by $\mathcal{C}(F)$.

Let $F \in \text{Mod}(S_g)$ be an infinite order reducible mapping class. Let $\mathcal{C}(F) = \{c_1, c_2, \dots, c_\ell\}$ be the canonical reduction system for F and N an F -invariant closed regular neighborhood of $\mathcal{C}(F)$. Let n be the least positive integer such that F^n fixes each path component of $S_g \setminus \overline{N} \cup N$. Then, as a consequence of the Nielsen–Thurston classification [33], there exist $s \in \mathbb{N} \cup \{0\}$ and $q_i \in \mathbb{Z} \setminus \{0\}$ such that

$$F^n = T_{c_1}^{q_1} T_{c_2}^{q_2} \cdots T_{c_\ell}^{q_\ell} \eta_1(F_1) \eta_2(F_2) \cdots \eta_s(F_s) \quad (1)$$

with $F_i \in \text{Mod}(R_i)$ is either periodic or pseudo-Anosov, where R_i is a path component of $S_g \setminus \overline{N}$ and $\eta_i : \text{Mod}(R_i) \rightarrow \text{Mod}(S_g)$ is the natural inclusion map. For $1 \leq j \leq s$, F_j 's (or $\eta_j(F_j)$'s) will be called the *canonical components* of F . The product $T_{c_1}^{q_1} T_{c_2}^{q_2} \cdots T_{c_\ell}^{q_\ell}$ appearing in (1) will be called the *multitwist component* of F . The decomposition of the form (1) will be called the *canonical decomposition (or the Nielsen decomposition)* of F . Without loss of generality, we assume that $F_1, F_2, \dots, F_{s'}$ are periodic canonical components, where $s' \leq s$. The integer $n \cdot \text{lcm}(|F_1|, |F_2|, \dots, |F_{s'}|)$ will be called the *degree* of F . For a multicurve C , the cut surface obtained by capping the boundary components of $S_g \setminus \overline{N}$ by marked disks will be denoted by $S_g(C)$, where N is a closed regular neighborhood of C .

Suppose that $F, G \in \text{Mod}(S_g)$ generate an infinite metacyclic subgroup of $\text{Mod}(S_g)$ such that $\langle F \rangle \triangleleft \langle F, G \rangle$. Then it follows that F and G satisfy the relation $G^{-1}FG = F^k$, for some nonzero integer k . Hence, the group $\langle F, G \rangle$ is a semidirect product of $\langle F \rangle$ and $\langle G \rangle$ and will be denoted by $\langle F \rangle \rtimes_k \langle G \rangle$. In Section 3, we derive the main results of this paper. To begin with, in Section 3.1, we derive necessary and sufficient conditions for the existence of infinite metacyclic subgroups of $\text{Mod}(S_g)$ with a pseudo-Anosov generator depending upon the Nielsen–Thurston type of the other generator (see Theorem 3.1). We achieve this by analyzing its invariant foliations and the dilatation homomorphism (see [23]). In particular, we have given necessary and sufficient conditions under which a pseudo-Anosov mapping class F forms an infinite metacyclic subgroup $\langle F, G \rangle$ with a nontrivial periodic mapping class G such that $\langle F \rangle \triangleleft \langle F, G \rangle$. Furthermore, for other types of G , we have the following main result.

Theorem 1. *For $g \geq 2$, consider nontrivial mapping classes $F, G \in \text{Mod}(S_g)$. Let $\langle F, G \rangle$ be metacyclic with $\langle F \rangle \triangleleft \langle F, G \rangle$. Then the following statements hold.*

- (i) *If F is a pseudo-Anosov, then G cannot be an infinite order reducible mapping class.*
- (ii) *If F and G are pseudo-Anosov, then $\langle F, G \rangle$ is abelian. Furthermore, either $\langle F, G \rangle \cong \mathbb{Z}$ or $\langle F, G \rangle \cong \mathbb{Z}_n \times \mathbb{Z}$ for some $n \in \mathbb{N}$.*
- (iii) *Let G be pseudo-Anosov and $\langle F, G \rangle$ is non-abelian. Then F is a reducible mapping class of finite order.*

In Section 3.2, by decomposing each reducible generator into its canonical components, we obtain necessary and sufficient conditions under which two reducible elements of $\text{Mod}(S_g)$ form an infinite metacyclic subgroup. In this direction, we have our second main result (see Theorem 3.12) which generalizes a result of Ivanov (see [14, Theorem 7.5A]).

Theorem 2. *For $g \geq 2$, let $F, G \in \text{Mod}(S_g)$ be two nontrivial mapping classes such that at least one of F or G is of infinite order and neither F nor G is pseudo-Anosov. Assume that F, G have degrees n, m , with multitwist components*

$$T_{c_1}^{q_1} T_{c_2}^{q_2} \dots T_{c_\ell}^{q_\ell} \quad \text{and} \quad T_{c'_1}^{q'_1} T_{c'_2}^{q'_2} \dots T_{c'_{\ell'}}^{q'_{\ell'}},$$

respectively, where $q_i, q'_i \in \mathbb{Z} \setminus \{0\}$, $\mathcal{C}(F) = \{c_1, c_2, \dots, c_\ell\}$, and $\mathcal{C}(G) = \{c'_1, c'_2, \dots, c'_{\ell'}\}$. Then $\langle F, G \rangle$ is an infinite metacyclic subgroup with $\langle F \rangle \triangleleft \langle F, G \rangle$ if and only if the following conditions hold.

- (i) $\mathcal{C}(F) \cup \mathcal{C}(G)$ is a multicurve.
- (ii) If F is periodic with $G^{-1}FG = F^k$, then $k^m \equiv 1 \pmod{n}$.
- (iii) Define $A_i := \{c_j \in \mathcal{C}(F) \mid q_j = q_i\}$, $B_i := \{c_j \in \mathcal{C}(F) \mid q_j = kq_i\}$, and $C_i := \{c'_j \in \mathcal{C}(G) \mid q'_j = q'_i\}$. Then $G(A_i) = B_i$, $G(B_i) = A_i$, and $F(C_i) = C_i$ for every i .
- (iv) For every path component R of $S_g(\mathcal{C}(F) \cup \mathcal{C}(G))$, then $G_r^{-1}F_rG_r = F_r^{k^{p_r}}$, where $G_r, F_r \in \text{Mod}(R)$ are induced by G, F , respectively, and p_r is the size of orbit of R under G .
- (v) For two path components R, S of $S_g(\mathcal{C}(F) \cup \mathcal{C}(G))$ such that $G(R) = S$, then F_r^k is conjugate to F_s , where $F_r \in \text{Mod}(R)$, $F_s \in \text{Mod}(S)$ are induced by F .

The following result is a direct consequence of Theorem 2.

Corollary 1. *For $g \geq 2$, let $F, G \in \text{Mod}(S_g)$ be two nontrivial mapping classes such that at least one of F or G is of infinite order and neither F nor G is pseudo-Anosov. Let $\langle F, G \rangle$ be an infinite metacyclic subgroup of $\text{Mod}(S_g)$ with $\langle F \rangle \triangleleft \langle F, G \rangle$. Then the following statements hold.*

- (i) F and G are reducible mapping classes.
- (ii) If F, G are of infinite order such that G is of odd degree, then $\langle F, G \rangle$ is abelian.
- (iii) If G is of infinite order of degree 1, then $\langle F, G \rangle$ is abelian.

By applying our main theorems, we have shown that infinite metacyclic subgroups of $\text{Mod}(S_g)$ are abundant. In general, we have established that $\text{Mod}(S_g)$ has infinite metacyclic subgroups isomorphic to $\mathbb{Z}_n \rtimes_k \mathbb{Z}$, $\mathbb{Z} \rtimes_k \mathbb{Z}_n$, and $\mathbb{Z} \rtimes_k \mathbb{Z}$. We have constructed several explicit examples (see Sections 3–4) of such subgroups.

In Section 4, we derive several other applications of our main results. In Section 4.1, we obtain the following characterization of the infinite metacyclic subgroups of level m subgroups $\text{Mod}(S_g)[m]$ of $\text{Mod}(S_g)$ for $m \geq 3$.

Proposition 1. *For $g \geq 2$ and $m \geq 3$, let $F, G \in \text{Mod}(S_g)[m]$ be two nontrivial mapping classes. Then $\langle F, G \rangle$ is metacyclic with $\langle F \rangle \triangleleft \langle F, G \rangle$ if and only if the following hold.*

- (i) F and G are infinite order reducible mapping classes that commute.
- (ii) $\mathcal{C}(F) \cup \mathcal{C}(G)$ is a multicurve.
- (iii) The nontrivial canonical components of F and G are pseudo-Anosov mapping classes.
- (iv) The nontrivial canonical components of F and G with the same support generate a cyclic group.

Moreover, when $g \geq 3$, we show the existence of non-abelian infinite metacyclic subgroups in $\text{Mod}(S_g)[2]$. The following construction is motivated by a family of Penner-type pseudo-Anosov mapping classes described in [2].

Corollary 2. *For $g \geq 3$, there is an infinite metacyclic subgroup of $\langle F, G \rangle < \text{Mod}(S_g)[2]$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$, where F is a Penner-type pseudo-Anosov and G is a hyperelliptic involution.*

In Section 4.2, we have derived bounds on the order of a nontrivial periodic generator of an infinite metacyclic subgroup of $\text{Mod}(S_g)$ that are realized (see Proposition 4.5). In particular, we have the following result.

Proposition 2. *For $g \geq 2$, let $F, G \in \text{Mod}(S_g)$ be two nontrivial mapping classes such that $\langle F, G \rangle$ is an infinite metacyclic subgroup with $\langle F \rangle \triangleleft \langle F, G \rangle$.*

- (i) *Let F be a pseudo-Anosov mapping class and G a periodic mapping class.*
 - (a) *If $\langle F, G \rangle$ is abelian, then $2 \leq |G| \leq 2g$.*
 - (b) *If $\langle F, G \rangle$ is non-abelian, then $2 \leq |G| \leq 4g$.*
- (ii) *Let F be a reducible mapping class of infinite order and G a periodic mapping class.*
 - (a) *If $\langle F, G \rangle$ is abelian, then $2 \leq |G| \leq 2g + 2$.*
 - (b) *If $\langle F, G \rangle$ is non-abelian, then $2 \leq |G| \leq 2g$.*
- (iii) *If F is periodic and $\langle F, G \rangle$ is non-abelian, then $3 \leq |F| \leq 2g + 2$.*

Moreover, all of the above bounds are realized.

In Section 4.3, we describe pseudo-Anosovs in $\text{Mod}(S_g)$ which can be written as a product of two nontrivial periodic mapping classes of the same order.

Corollary 3. *Let $\langle F, G \rangle < \text{Mod}(S_g)$ be a non-abelian infinite metacyclic subgroup with $\langle F \rangle \triangleleft \langle F, G \rangle$, where F is a pseudo-Anosov and G is nontrivial periodic. Then, for integers i, j such that i is odd and j is even, $G^i F^j$ is conjugate to G^i . In particular, GF^2 is conjugate to G , and therefore, F^2 can be written as a product of two nontrivial periodic mapping classes of the same order.*

As a final application to our theory, in Section 4.4, we analyze the centralizers of irreducible periodic mapping classes in $\text{Mod}(S_g)$ (see Proposition 4.13). In particular, we have the following result.

Corollary 4. *Let $F \in \text{Mod}(S_g)$ be an irreducible periodic mapping class. Then the centralizer of F in $\text{Mod}(S_g)$ is either $\langle F \rangle$ or $\langle F \rangle \times \langle i \rangle$, where i is a hyperelliptic involution.*

2. Preliminaries

For $g \geq 2$, let S_g be the connected closed orientable surface of genus g . The *mapping class group* of S_g is the group of path components of $\text{Homeo}^+(S_g)$, and it will be denoted by $\text{Mod}(S_g)$. The elements of $\text{Mod}(S_g)$ are called *mapping classes*. The Nielsen–Thurston classification [33] asserts that each mapping class in $\text{Mod}(S_g)$ is either periodic, reducible, or pseudo-Anosov.

2.1. Periodic mapping classes

In view of the Nielsen–Kerckhoff theorem [18], a periodic mapping class $F \in \text{Mod}(S_g)$ of order n has a representative \mathcal{F} of the same order (known as a *Nielsen representative*) which induces a \mathbb{Z}_n -action on S_g via isometries. The *corresponding orbifold* of F is the quotient orbifold $\mathcal{O}_F := S_g / \langle \mathcal{F} \rangle$ (see [32, Chapter 13]), which is homeomorphic to S_{g_0} , where g_0 is the *orbifold genus* of \mathcal{O}_F . The \mathbb{Z}_n -action induces a branched covering $p : S_g \rightarrow \mathcal{O}_F$ with k branch points (or *cone points*) x_1, \dots, x_k in \mathcal{O}_F of orders n_1, \dots, n_k , respectively. The *order* of a cone point x_i is the order of the stabilizer subgroup of any point in the preimage of x_i . From orbifold covering space theory, the branch covering $p : S_g \rightarrow \mathcal{O}_F$ corresponds to an exact sequence

$$1 \rightarrow \pi_1(S_g) \xrightarrow{p_*} \pi_1^{\text{orb}}(\mathcal{O}_F) \xrightarrow{\phi} \mathbb{Z}_n \rightarrow 1.$$

Moreover, $\pi_1^{\text{orb}}(\mathcal{O}_F)$ is a Fuchsian group [17, 20] that has the following presentation:

$$\left\langle \alpha_1, \beta_1, \dots, \alpha_{g_0}, \beta_{g_0}, \gamma_1, \dots, \gamma_k \mid \gamma_1^{n_1} = \dots = \gamma_k^{n_k} = \prod_{i=1}^k \gamma_i \prod_{i=1}^{g_0} [\alpha_i, \beta_i] = 1 \right\rangle.$$

The epimorphism $\phi : \pi_1^{\text{orb}}(\mathcal{O}_F) \rightarrow \mathbb{Z}_n$ (classically known as a *surface kernel map*) is order-preserving on torsion elements and is given by $\phi(\gamma_i) = \mathcal{F}^{(n/n_i)d_i}$, where $\text{gcd}(d_i, n_i) = 1$, for $1 \leq i \leq k$. The tuple $(g_0; n_1, \dots, n_k)$ is called the *signature* of the quotient orbifold \mathcal{O}_F which we denote by $\Gamma(\mathcal{O}_F)$. Each cone point x_i of order n_i in \mathcal{O}_F lifts under p to an orbit of size n/n_i on S_g and the *local rotation* induced by \mathbb{Z}_n -action in this orbit is given by $2\pi d_i^{-1}/n_i$, where $\text{gcd}(d_i, n_i) = 1$. Thus, the orbit data of a cyclic action along with the structure of its corresponding orbifold can be compactly encoded as a tuple of integers.

Definition 2.1. For $n \geq 2$, $g_0 \geq 0$, and $0 \leq r \leq n - 1$, a *cyclic data set of degree n* , denoted by $n(D)$, is a tuple of the form

$$D = (n, g_0, r; (d_1, n_1), \dots, (d_k, n_k))$$

with the following conditions.

- (i) $r > 0$ if and only if $k = 0$, and when $r > 0$, then $\gcd(r, n) = 1$.
- (ii) $n_i \geq 2$, $n_i \mid n$, $\gcd(d_i, n_i) = 1$, for all i .
- (iii) $\text{lcm}(n_1, \dots, \widehat{n_i}, \dots, n_k) = \text{lcm}(n_1, \dots, n_k)$, for all i .
- (iv) If $g_0 = 0$, then $\text{lcm}(n_1, \dots, n_k) = n$.
- (v) $\sum_{i=1}^k \frac{n}{n_i} d_i \equiv 0 \pmod{n}$.
- (vi) $\frac{2g-2}{n} = 2g_0 - 2 + \sum_{i=1}^k (1 - \frac{1}{n_i})$ (Riemann-Hurwitz equation).

The number g determined by the Riemann–Hurwitz equation is the *genus* of the data set and will be denoted by $g(D)$.

The quantity r (in Definition 2.1) will be nonzero if and only if D represents a free rotation of S_g by $2\pi r/n$. We will not include r in the notation of a data set, whenever $r = 0$. The significance of the cyclic data set is given in the following proposition due to Nielsen [25] (see also [31, Theorem 3.9]).

Proposition 2.2. *Cyclic data sets of degree n and genus g are in one-to-one correspondence with conjugacy classes of periodic mapping classes of order n in $\text{Mod}(S_g)$.*

From here on, a periodic mapping class F and its associated cyclic action \mathcal{F} up to conjugacy will be represented by its corresponding data set, which we denote by D_F and $D_{\mathcal{F}}$, respectively. The corresponding orbifold of F will also be denoted by \mathcal{O}_F .

We now state some results concerning nontrivial periodic mapping classes which will be used later. The following result due to Gilman [10] characterizes irreducible periodic mapping classes $F \in \text{Mod}(S_g)$ based on the corresponding orbifold \mathcal{O}_F .


Theorem 2.3. *For $g \geq 2$, let $F \in \text{Mod}(S_g)$ be a nontrivial periodic mapping class. Then F is irreducible if and only if \mathcal{O}_F is a sphere with 3 cone points.*

We will now state a useful lemma [16, Theorem 4.1] due to Kasahara.


Lemma 2.4. *For $g \geq 2$, let $F \in \text{Mod}(S_g)$ be a nontrivial reducible periodic mapping class. Then $|F| \leq 2g + 2$. The upper bound is realized if and only if g is even and $\Gamma(\mathcal{O}_F) = (0; 2, 2, g + 1, g + 1)$. Furthermore, when $|F| < 2g + 2$, we have $|F| \leq 2g$. Equivalently, if either $|F| = 2g + 1$ or $|F| > 2g + 2$, then F is irreducible.*

Finally, we state the following assertion which follows from a result of Kulkarni [19].

Lemma 2.5. *There are no periodic mapping classes of order $4g + 1$ in $\text{Mod}(S_g)$.*



(a) The sum of local rotation angles about the two fixed points of cyclic action $D = (5, 0; (4, 5), (3, 5), (3, 5))$ associated with the pair $(3, 5)$ is $-2\pi/5$ modulo 2π . Hence, the action of D can be extended to a pseudo-periodic mapping class $F \in \text{Mod}(S_3)$ such that $F^5 = T_c^{-1}$, where c is a non-separating curve.



(b) The sum of local rotation angles about the fixed points of cyclic actions $D_1 = (8, 0; (1, 2), (5, 8), (7, 8))$ and $D_2 = (10, 0; (1, 2), (2, 5), (1, 10))$ associated with the pairs $(7, 8)$ and $(1, 10)$, respectively, is $-2\pi/40$ modulo 2π . Since $\text{lcm}(8, 10) = 40$, a pseudo-periodic $F \in \text{Mod}(S_4)$ can be constructed from D_1 and D_2 such that $F^{40} = T_c^{-1}$, where c is a separating curve.

Figure 1. Construction of pseudo-periodic mapping classes.

2.2. Pseudo-periodic mapping classes

Let $F \in \text{Mod}(S_g)$ be an infinite order reducible mapping class. From here on, we will use the notions of *canonical decomposition* and the *degree* of F as defined in Section 1. A mapping class is said to be *pseudo-periodic* if it is either a nontrivial periodic or of infinite order reducible with only periodic canonical components. Thus, a nontrivial periodic mapping class F will be considered as pseudo-periodic with $\mathcal{C}(F) = \emptyset$, degree $|F|$, and multitwist component equal to identity. We observe that multitwists are pseudo-periodic mapping classes having trivial periodic canonical components.

In the following example, we construct some infinite order pseudo-periodic mapping classes whose power is a Dehn twist about a simple closed curve.

Example 2.6. Let $F \in \text{Mod}(S_g)$ be a pseudo-periodic mapping class such that $F^n = T_c$. Then F is represented by an $\mathcal{F} \in \text{Homeo}^+(S_g)$ such that $\mathcal{F}(N) = N$, where N is a closed annular neighborhood of c . Thus, \mathcal{F} induces a \mathbb{Z}_n -action on $S_g(c)$ with two fixed points. Moreover, the sum of induced rotation angles about these fixed points is $2\pi/n$ modulo 2π . Conversely, given nontrivial periodic mapping classes having a (two, in case c is nonseparating) distinguished fixed point such that the sum of induced rotation angles about these fixed points is $2\pi/n$ modulo 2π , one can reverse this process to recover F . (We refer the reader to [24, 28, 29, 31] for details.) We illustrate this construction of roots of Dehn twists in Figure 1.

The angle sum condition in Example 2.6 (in the construction of pseudo-periodic) generalizes to a formal “compatibility condition” between pairs of orbits of one or more cyclic actions (see [15] for more details).

Definition 2.7. For $i = 1, 2$, let O_i be an orbit of cyclic action D_i such that $|O_1| = |O_2|$. Let k be an integer such that $0 \leq |k| \leq n/2$, where $n = \text{lcm}(n(D_1), n(D_2))$.

- (i) We say that O_1 and O_2 are *trivially n -compatible* if $|O_1| = |O_2| = n$ (in this case $n(D_1) = n(D_2)$).
- (ii) Let the pair (d_i, n_i) correspond to the orbit O_i in the data set D_i , where we assume that $(d_i, n_i) = (0, 1)$ if $|O_i| = n(D_i)$. We say that the orbits O_1 and O_2 are *$|O_i|$ -compatible with twist factor k* if

$$\frac{2\pi d_1^{-1}}{n_1} + \frac{2\pi d_2^{-1}}{n_2} \equiv \frac{2\pi k}{n} \pmod{2\pi}. \quad (2)$$

When the twist factor associated with the compatibility of the D_i is 0, we simply say that the D_i are *$|O_i|$ -compatible*.

2.3. Metacyclic groups

A group H is said to be a *metacyclic group* if there is a short exact sequence

$$1 \rightarrow N \rightarrow H \rightarrow L \rightarrow 1, \quad (3)$$

where N and L are cyclic groups. If a metacyclic group H fits into an exact sequence as in (3) that splits, then we say that H is a *split metacyclic group*. Thus, the split metacyclic group H is isomorphic to the semidirect product $N \rtimes L$. Given integers $u, n \in \mathbb{N}$, a finite metacyclic group H of order $u \cdot n$ admits the following presentation:

$$H = \langle \mathcal{F}, \mathcal{G} \mid \mathcal{F}^n = 1, \mathcal{F}^r = \mathcal{G}^u, \mathcal{G}^{-1} \mathcal{F} \mathcal{G} = \mathcal{F}^k \rangle, \quad (4)$$

where $r \in \mathbb{N}$, $k \in \mathbb{Z}_n^\times$ such that $r \mid n$, $k^u \equiv 1 \pmod{n}$, and $r(k-1) \equiv 0 \pmod{n}$. For integers $m, n \in \mathbb{N}$ and $k \in \mathbb{Z}_n^\times$, a split metacyclic group admits the following presentation:

$$H = \langle \mathcal{F}, \mathcal{G} \mid \mathcal{F}^n = 1, \mathcal{G}^m = 1, \mathcal{G}^{-1} \mathcal{F} \mathcal{G} = \mathcal{F}^k \rangle \cong \mathbb{Z}_n \rtimes_k \mathbb{Z}_m.$$

Metacyclic groups have been completely classified by Hempel in [12].

An *infinite metacyclic group* is a metacyclic group of infinite order. It is known [12, Chapter 7] that an infinite metacyclic group admits exactly one of the following presentations:

$$\begin{aligned} \langle \mathcal{F}, \mathcal{G} \mid \mathcal{G}^{-1} \mathcal{F} \mathcal{G} = \mathcal{F}^k \rangle &\cong \mathbb{Z} \rtimes_k \mathbb{Z}, & \text{for } k = \pm 1, \\ \langle \mathcal{F}, \mathcal{G} \mid \mathcal{G}^{2m} = 1, \mathcal{G}^{-1} \mathcal{F} \mathcal{G} = \mathcal{F}^k \rangle &\cong \mathbb{Z} \rtimes_k \mathbb{Z}_{2m}, & \text{for } k = -1, m \in \mathbb{N}, \\ \langle \mathcal{F}, \mathcal{G} \mid \mathcal{F}^n = 1, \mathcal{G}^{-1} \mathcal{F} \mathcal{G} = \mathcal{F}^k \rangle &\cong \mathbb{Z}_n \rtimes_k \mathbb{Z}, & \text{for } k \in \mathbb{Z}_n^\times, n \in \mathbb{N}. \end{aligned} \quad (5)$$

Throughout this paper, we will only consider non-cyclic (i.e., two-generator) infinite metacyclic groups. As a consequence of the relation $\mathcal{G}^{-1} \mathcal{F} \mathcal{G} = \mathcal{F}^k$ in a metacyclic group $H = \langle \mathcal{F}, \mathcal{G} \rangle$, we have the following elementary lemma.

Lemma 2.8. *Let $H = \langle \mathcal{F}, \mathcal{G} \rangle$ be a metacyclic group, where $\mathcal{G}^{-1} \mathcal{F} \mathcal{G} = \mathcal{F}^k$. For integers i, j , we have*

- (i) $\mathcal{F}^i \mathcal{G}^j = \mathcal{G}^j \mathcal{F}^{ik^j}$ and
- (ii) $(\mathcal{G}^i \mathcal{F}^j)^\ell = \mathcal{G}^{i\ell} \mathcal{F}^{j(1+k^i+k^{2i}+\dots+k^{i(\ell-2)}+k^{i(\ell-1)})}$.

2.4. Induced orbifold automorphisms

Let $\langle \mathcal{F}, \mathcal{G} \rangle$ be a metacyclic subgroup of $\text{Homeo}^+(S_g)$, where \mathcal{F} has finite order. Each cone point $[x] \in \mathcal{O}_{\mathcal{F}}$ corresponds to a unique pair of the form (c_x, n_x) in the data set $D_{\mathcal{F}}$ corresponding to \mathcal{F} . If $[x] \in \mathcal{O}_{\mathcal{F}}$ is not a cone point, then we take $(c_x, n_x) = (0, 1)$. As $\langle \mathcal{F} \rangle \triangleleft H$, it is known [34] that \mathcal{G} would induce a $\bar{\mathcal{G}} \in \text{Homeo}^+(\mathcal{O}_{\mathcal{F}})$ that preserves the set of cone points in $\mathcal{O}_{\mathcal{F}}$ along with their orders. We will call $\bar{\mathcal{G}}$, *the induced automorphism on $\mathcal{O}_{\mathcal{F}}$ by \mathcal{G}* , and we formalize this notion in the following definition.

Definition 2.9. Let $\mathcal{F} \in \text{Homeo}^+(S_g)$ be a finite order map such that $|F| = n$. We say a $\bar{\mathcal{G}} \in \text{Homeo}^+(\mathcal{O}_{\mathcal{F}})$ is an *automorphism of $\mathcal{O}_{\mathcal{F}}$* if for $[x], [y] \in \mathcal{O}_{\mathcal{F}}$, $k \in \mathbb{Z}_n^\times$, and $\bar{\mathcal{G}}([x]) = [y]$, we have

- (i) $n_x = n_y$,
- (ii) $c_x = kc_y$.

We denote the group of automorphisms of $\mathcal{O}_{\mathcal{F}}$ by $\text{Aut}_k(\mathcal{O}_{\mathcal{F}})$. When $k = 1$, we simply write $\text{Aut}(\mathcal{O}_{\mathcal{F}})$ instead of $\text{Aut}_1(\mathcal{O}_{\mathcal{F}})$. In the following lemma, we state some basic properties of induced automorphisms.

Lemma 2.10 ([6, Lemma 2.9]). *Let $\mathcal{F} \in \text{Homeo}^+(S_g)$ be a map of order n and $\mathcal{G} \in \text{Homeo}^+(S_g)$ a map such that $\mathcal{G}^{-1} \mathcal{F} \mathcal{G} = \mathcal{F}^k$. Then \mathcal{G} induces a $\bar{\mathcal{G}} \in \text{Aut}_k(\mathcal{O}_{\mathcal{F}})$ such that*

$$\mathcal{O}_{\mathcal{F}} / \langle \bar{\mathcal{G}} \rangle = S_g / \langle \mathcal{F}, \mathcal{G} \rangle.$$

Furthermore, \mathcal{G} has infinite order if and only if $\bar{\mathcal{G}}$ has infinite order. If $|\mathcal{G}| = m$, then

- (i) $|\bar{\mathcal{G}}|$ divides $|\mathcal{G}|$, and
- (ii) $|\bar{\mathcal{G}}| < m$ if and only if $\mathcal{F}^r = \mathcal{G}^u$, for some $0 < r < n$ and $0 < u < m$.

We refer the reader to [5, 6] for further details on induced orbifold automorphisms.

2.5. Pseudo-Anosov mapping classes

For $g \geq 2$, let $F \in \text{Mod}(S_g)$ be *pseudo-Anosov* mapping class. We will now describe a well-known construction of pseudo-Anosov mapping classes due to Penner [27, Theorem 3.1].

Theorem 2.11. *Let $A = \{a_1, a_2, \dots, a_m\}$ and $B = \{b_1, b_2, \dots, b_n\}$ be multicurves in S_g such that $A \cup B$ fills S_g . Any product of positive powers of T_{a_i} and negative powers of T_{b_j} , where each a_i and each b_j appear at least once, is a pseudo-Anosov mapping class.*

Note that a collection of simple closed curves C in S_g is said to *fill* S_g if $\overline{S_g \setminus C}$ is a union of closed disks. Let $F, G \in \text{Mod}(S_g)$ be nontrivial mapping classes, where F is pseudo-Anosov with stretch factor $\lambda > 1$ satisfying the relation $G^{-1}FG = F^k$. Let (\mathfrak{F}_s, μ_s) and (\mathfrak{F}_u, μ_u) be the stable and unstable singular measured foliations of F . We will require the following result due to McCarthy [23, Lemma 1] in proving our results.

Lemma 2.12. *Let $F, G \in \text{Mod}(S_g)$ such that F is a pseudo-Anosov mapping class satisfying $G^{-1}FG = F^k$. Then $k = \pm 1$ and there exists a positive real number ρ such that the following conditions hold:*

- (i) *if $k = 1$, then $G(\mathfrak{F}_s, \mu_s) = (\mathfrak{F}_s, \rho^{-1}\mu_s)$ and $G(\mathfrak{F}_u, \mu_u) = (\mathfrak{F}_u, \rho\mu_u)$,*
- (ii) *if $k = -1$, then $G(\mathfrak{F}_s, \mu_s) = (\mathfrak{F}_u, \rho^{-1}\mu_u)$ and $G(\mathfrak{F}_u, \mu_u) = (\mathfrak{F}_s, \rho\mu_s)$.*

Remark 2.13. Let $\mathcal{H} = \{G \in \text{Mod}(S_g) : G(\mathfrak{F}_s) = \mathfrak{F}_s \text{ and } G(\mathfrak{F}_u) = \mathfrak{F}_u\}$ and let \mathbb{R}_+ be the group of positive real numbers under multiplication. There is a homomorphism $\lambda : \mathcal{H} \rightarrow \mathbb{R}_+$ such that $\lambda(G) = \lambda_G$ with $G(\mathfrak{F}_u, \mu_u) = (\mathfrak{F}_u, \lambda_G\mu_u)$ and $G(\mathfrak{F}_s, \mu_s) = (\mathfrak{F}_s, \lambda_G^{-1}\mu_s)$ (see [23]). This homomorphism is known as the *dilatation homomorphism*.

For a singular point p of \mathfrak{F}_u , let \mathcal{H}_p be the subgroup of \mathcal{H} consisting of mapping classes that fix p . Let \mathcal{L}_p be the set of all singular leaves of \mathfrak{F}_u originating at the singular point p . The action of \mathcal{H}_p on \mathcal{L}_p induces a homomorphism $\phi_p : \mathcal{H}_p \rightarrow \Sigma_{|\mathcal{L}_p|}$, where $\Sigma_{|\mathcal{L}_p|}$ is the permutation group on $|\mathcal{L}_p|$ letters.

The image and kernel of the dilatation homomorphism λ have also been described in [23, Lemmas 2-3].

Lemma 2.14. *For the dilatation homomorphism $\lambda : \mathcal{H} \rightarrow \mathbb{R}_+$, we have that $\lambda(\mathcal{H})$ is infinite cyclic and $\ker \lambda$ is a finite group.*

3. Infinite metacyclic subgroups of mapping class group

For $g \geq 2$ and two nontrivial periodic mapping classes $F, G \in \text{Mod}(S_g)$, the necessary and sufficient number-theoretic conditions under which conjugates F', G' (of F, G resp.) generate a finite metacyclic group have been derived in [5, 6, 30]. In this section, we analyze the infinite metacyclic subgroups of $\text{Mod}(S_g)$. From here on, for $F, G \in \text{Mod}(S_g)$, we will assume that if $\langle F, G \rangle$ is a metacyclic group, then $\langle F \rangle \triangleleft \langle F, G \rangle$, which implies that $G^{-1}FG = F^k$ for some nonzero integer k .

3.1. Metacyclic subgroups with pseudo-Anosov generators

Let $F \in \text{Mod}(S_g)$ be a pseudo-Anosov mapping class with stretch factor $\lambda > 1$. Let (\mathfrak{F}_s, μ_s) and (\mathfrak{F}_u, μ_u) be the stable and unstable singular measured foliations for F , respectively. We will now prove our first main result concerning infinite metacyclic subgroups of $\text{Mod}(S_g)$ with at least one pseudo-Anosov generator. The homomorphism ϕ_p in the statement of the following theorem has been defined in Remark 2.13.

Theorem 3.1 (Main Theorem 1). *For $g \geq 2$, consider nontrivial mapping classes $F, G \in \text{Mod}(S_g)$.*

- (i) *Let $\langle F, G \rangle$ be metacyclic with $\langle F \rangle \triangleleft \langle F, G \rangle$. Then the following statements hold.*
 - (a) *If F is a pseudo-Anosov, then G cannot be an infinite order reducible mapping class.*
 - (b) *If F and G are pseudo-Anosov, then $\langle F, G \rangle$ is abelian. Furthermore, either $\langle F, G \rangle \cong \mathbb{Z}$ or $\langle F, G \rangle \cong \mathbb{Z}_n \times \mathbb{Z}$ for some $n \in \mathbb{N}$.*
 - (c) *Let G be pseudo-Anosov and $\langle F, G \rangle$ is non-abelian. Then F is a reducible mapping class of finite order.*
- (ii) *Let F be pseudo-Anosov and G is either periodic or pseudo-Anosov. Then $\langle F, G \rangle$ is an abelian metacyclic subgroup if and only if*
 - (a) $G(\mathfrak{F}_u, \mu_u) = (\mathfrak{F}_u, \lambda_G \mu_u)$, $G(\mathfrak{F}_s, \mu_s) = (\mathfrak{F}_s, \lambda_G^{-1} \mu_s)$, and
 - (b) *there exists a singular point p of \mathfrak{F}_u such that $G^{-1} F G F^{-1} \in \ker \phi_p$.*
- (iii) *Let F be pseudo-Anosov and let G be periodic. Then $\langle F, G \rangle$ is a non-abelian metacyclic subgroup with $\langle F \rangle \triangleleft \langle F, G \rangle$ if and only if*
 - (a) $G(\mathfrak{F}_u, \mu_u) = (\mathfrak{F}_s, \mu_s)$, $G(\mathfrak{F}_s, \mu_s) = (\mathfrak{F}_u, \mu_u)$, and
 - (b) *there exists a singular point p of \mathfrak{F}_u such that $G^{-1} F G F \in \ker \phi_p$.*

Proof. Let F be a pseudo-Anosov mapping class with (\mathfrak{F}_u, μ_u) and (\mathfrak{F}_s, μ_s) as its unstable and stable invariant singular measured foliations, respectively.

To begin with, we consider the case when $\langle F, G \rangle$ is metacyclic with $\langle F \rangle \triangleleft \langle F, G \rangle$. Let F be pseudo-Anosov and G an infinite order reducible mapping class. Consider the dilatation homomorphism $\lambda : \mathcal{H} \rightarrow \mathbb{R}_+$ (see Remark 2.13). Since $G^{-1} F G = F^k$, where $k = \pm 1$, G^2 commutes with F . By Lemma 2.12, $G^2 \in \mathcal{H}$, and since G is not pseudo-Anosov, $G^2 \in \ker \lambda$. This is impossible since $\ker \lambda$ is finite and G^2 has infinite order.

Next, we consider the case when F, G are pseudo-Anosov and $\langle F, G \rangle$ is a non-abelian metacyclic subgroup. If $G^{-1} F G = F^{-1}$, then G^2 commutes with F . By Lemma 2.12, it follows that G^2 preserves (\mathfrak{F}_u, μ_u) and (\mathfrak{F}_s, μ_s) . Thus, F and G keep (\mathfrak{F}_u, μ_u) and (\mathfrak{F}_s, μ_s) invariant, which contradicts Lemma 2.12. Therefore, $\langle F, G \rangle$ is abelian, and from Lemma 2.12, we have $\langle F, G \rangle \subset \mathcal{H}$. Since $\ker \lambda$ is a finite group, if $\ker \lambda|_{\langle F, G \rangle} \neq 1$, then $\langle F, G \rangle \cong \mathbb{Z}_n \times \mathbb{Z}$ for some $n \in \mathbb{N}$. Furthermore, if $\ker \lambda|_{\langle F, G \rangle} = 1$, then $\langle F, G \rangle \cong \mathbb{Z}$.

Next, we assume that G is a pseudo-Anosov mapping class and $\langle F, G \rangle$ is non-abelian metacyclic subgroup. As discussed above, F can not be pseudo-Anosov. Let F be an infinite order reducible mapping class. Since $\mathcal{C}(G^{-1} F G) = G^{-1}(\mathcal{C}(F))$, $\mathcal{C}(F^{-1}) = \mathcal{C}(F)$ and $G^{-1} F G = F^{-1}$, it follows that $G(\mathcal{C}(F)) = \mathcal{C}(F)$. But as $\mathcal{C}(F) \neq \emptyset$ and G is irreducible, this contradicts our assumption. Now, assume that F is periodic. If F is irreducible, then, by Theorem 2.3, $\mathcal{O}_F \approx S_{0,3}$. Since $G F G^{-1} = F^k$, G induces an infinite order mapping class in $\text{Mod}(S_{0,3})$ which is not possible. Hence, F must be reducible periodic.

Finally, we consider the case when F is pseudo-Anosov and G is either periodic or pseudo-Anosov. Consider the homomorphism ϕ_p defined in Remark 2.13. From Lemma 2.12, if $G^{-1}FG = F$, then $G(\mathfrak{F}_u, \mu_u) = (\mathfrak{F}_u, \lambda_G \mu_u)$ and $G(\mathfrak{F}_s, \mu_s) = (\mathfrak{F}_s, \lambda_G^{-1} \mu_s)$. Furthermore, it is apparent that

$$G^{-1}FGF^{-1} \in \ker \phi_p.$$

Conversely, we assume that $G(\mathfrak{F}_u, \mu_u) = (\mathfrak{F}_u, \lambda_G \mu_u)$, $G(\mathfrak{F}_s, \mu_s) = (\mathfrak{F}_s, \lambda_G^{-1} \mu_s)$, and $G^{-1}FGF^{-1} \in \ker \phi_p$ for some singular point p of \mathfrak{F}_u . Let \mathcal{F} and \mathcal{G} be representatives of F and G , respectively. Since $G^{-1}FGF^{-1} \in \ker \phi_p$, we have $\mathcal{G}^{-1}\mathcal{F}\mathcal{G}\mathcal{F}^{-1}(L) = L$, where L is a leaf of \mathfrak{F}_u originating at the singular point p . Since $\lambda(\mathcal{G}^{-1}\mathcal{F}\mathcal{G}\mathcal{F}^{-1}) = 1$, $\mathcal{G}^{-1}\mathcal{F}\mathcal{G}\mathcal{F}^{-1}$ fixes L pointwise. Since L is dense in S_g [8, Theorem 9.6], we must have $\mathcal{G}^{-1}\mathcal{F}\mathcal{G}\mathcal{F}^{-1} = 1$, and hence $G^{-1}FG = F$. By a similar argument, (iii) follows. ■

We address the case when G is a pseudo-Anosov mapping class and F is a nontrivial periodic mapping class in the following remark.

Remark 3.2. For $g \geq 2$, let $F, G \in \text{Mod}(S_g)$ be such that G is a pseudo-Anosov mapping class and F is a nontrivial periodic mapping class. By Birman–Hilden theory [3, 22], it follows that $\langle F, G \rangle$ is metacyclic with $\langle F \rangle \triangleleft \langle F, G \rangle$ if and only if there exists a pseudo-Anosov mapping class $\bar{G} \in \text{Mod}(\mathcal{O}_F)$ such that \bar{G} lifts to G under the branched cover $p : S_g \rightarrow \mathcal{O}_F$. By removing branch points and their preimages, p can be considered an unbranched cover between punctured surfaces. Since p is an abelian cover, a \bar{G} lifts under p if and only if the induced isomorphism $\bar{G}_\# \in \text{Aut}(H_1(\mathcal{O}_F, \mathbb{Z}))$ leaves the subgroup of $H_1(\mathcal{O}_F, \mathbb{Z})$ corresponding to the cover p invariant. This homological criterion is often straightforward to compute (see [1, 9]).

Now, we construct several infinite metacyclic subgroups of $\text{Mod}(S_g)$ with a pseudo-Anosov generator. In the following example, we describe a non-abelian infinite metacyclic subgroup having a nontrivial periodic and a pseudo-Anosov generator.

Example 3.3. For $g \geq 1$, let G be a rotation of S_{4g} by $2\pi/4$ as shown in Figure 2. By considering the multicurves $A = \{a_1, a_2, \dots, a_{4g+1}\}$ and $B = \{b_1, b_2, \dots, b_{4g+1}\}$, we see that the curves in $A \cup B$ fill S_{4g} . From Theorem 2.11, it follows that

$$F = \prod_{i=1}^{4g+1} T_{a_i} \prod_{j=1}^{4g+1} T_{b_j}^{-1}$$

is a pseudo-Anosov mapping class. For $1 \leq i \leq 3g$ and $1 \leq i' \leq g$, we have $G(a_i) = b_{g+i}$, $G(b_i) = a_{g+i}$, $G(a_{3g+i'}) = b_{i'}$, $G(b_{3g+i'}) = a_{i'}$, and G exchanges the curves a_{4g+1} and b_{4g+1} . Therefore, $G^{-1}FG = F^{-1}$, and so we have $\langle F, G \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}_4$. We observe that

$$\langle F, G^2 \rangle \cong \mathbb{Z} \times \mathbb{Z}_2.$$

Note that the construction described in Example 3.3 generalizes to any even integer $n \geq 4$, where n is the order of the periodic generator. For even genera, we will now provide an example of a metacyclic subgroup of $\text{Mod}(S_g)$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$.




Figure 2. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_{4g})$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_4$ generated by a periodic mapping class G of order 4 and a pseudo-Anosov mapping class F .




Figure 3. An infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_{2h})$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$, generated by an involution G and a pseudo-Anosov F .

Example 3.4. For an integer $h \geq 1$, let $A = \{a_1, a_2, \dots, a_{2h}, d\}$ and $B = \{b_1, b_2, \dots, b_{2h}, c\}$ be two multicurves in S_{2h} , and let $G \in \text{Mod}(S_{2h})$ be an involution, as shown in Figure 3. Since $A \cup B$ fills S_{2h} , by Theorem 2.11, the mapping class

$$F = \prod_{i=1}^{2h} T_{a_i} T_d \prod_{j=1}^{2h} T_{b_j}^{-1} T_c^{-1}$$

is pseudo-Anosov. For $1 \leq i \leq 2h$, G maps a_i to b_{2h+1-i} , b_i to a_{2h+1-i} , and c to d . Therefore, we have $G^{-1}FG = F^{-1}$, and so it follows that $\langle F, G \rangle < \text{Mod}(S_{2h})$ is isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$.

Examples 3.3–3.4 together generalize to the following corollary.




Figure 4. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_{4g+1})$ isomorphic to $\mathbb{Z} \times \mathbb{Z}_4$ generated by a periodic mapping class G of order 4 and a pseudo-Anosov mapping class F .

Corollary 3.5. *For an even positive integer $n \mid g$, there is an infinite metacyclic subgroup of $\langle F, G \rangle < \text{Mod}(S_g)$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_n$, where F is a Penner-type pseudo-Anosov and G is periodic with $D_G = (n, g/n; (1, n), (n-1, n))$.*

For positive integer m , let $\langle F, G \mid G^{2m} = 1, G^{-1}FG = F^{-1} \rangle$ be an infinite metacyclic subgroup of $\text{Mod}(S_g)$, where F is a pseudo-Anosov mapping class and G is a periodic mapping class. Then it is easily seen that $\langle F, G^2 \rangle$ is abelian (as in Example 3.3). However, a natural question is whether every infinite abelian metacyclic subgroup of $\text{Mod}(S_g)$ arises this way. The following example shows that this is not true in general.

Example 3.6. For $g \geq 1$, let G be a free rotation of S_{4g+1} by $2\pi/4$ as shown in Figure 4. We observe that the multicurves $A = \{a_1, a_2, \dots, a_{4g+1}\}$ and $B = \{b_1, b_2, \dots, b_{4g+4}\}$ fill S_{4g+1} . From Theorem 2.11, the mapping class

$$F = \prod_{i=1}^{4g+1} T_{a_i} \prod_{j=1}^{4g+4} T_{b_j}^{-1}$$

is pseudo-Anosov. For $1 \leq i \leq 3g$, $1 \leq i' \leq g$, and $1 \leq j \leq 3$, we have that $G(a_i) = a_{i+g}$, $G(b_i) = b_{i+g}$, $G(a_{3g+i'}) = a_{i'}$, $G(b_{3g+i'}) = b_{i'}$, $G(a_{4g+1}) = a_{4g+1}$, $G(b_{4g+j}) = b_{4g+j+1}$, and $G(b_{4g+4}) = b_{4g+1}$. By construction, we have $GF = FG$, and so it follows that $\langle F, G \rangle \cong \mathbb{Z} \times \mathbb{Z}_4$.




Figure 5. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_{4g})$ isomorphic to $\mathbb{Z} \times \mathbb{Z}_4$ generated by a periodic mapping class G of order 4 and a pseudo-Anosov mapping class F .

For $g = 1$, assume that $\langle F, G \rangle$ is a subgroup of a non-abelian infinite metacyclic subgroup $\langle F, G' \rangle$, where $(G')^2 = G$. It follows from [5, Corollary 5.7] that G is primitive. Therefore, such a G' cannot exist.

In Example 3.6, the periodic generator was represented by a nontrivial free rotation, but in the following example, the periodic generator is represented by a nontrivial non-free rotation.

Example 3.7. For $g \geq 1$, let G be a rotation of S_{4g} by $2\pi/4$ as shown in Figure 5. We observe that the multicurves $A = \{a_1, a_2, \dots, a_{4g+1}\}$ and $B = \{b_1, b_2, \dots, b_{4g}\}$ fill S_{4g} . From Theorem 2.11, the mapping class

$$F = \prod_{i=1}^{4g+1} T_{a_i} \prod_{j=1}^{4g} T_{b_j}^{-1}$$

is pseudo-Anosov. For $1 \leq i \leq 3g$, $1 \leq i' \leq g$, and $1 \leq j \leq 3$, we have that $G(a_i) = a_{i+g}$, $G(b_i) = b_{i+g}$, $G(a_{3g+i'}) = a_{i'}$, $G(b_{3g+i'}) = b_{i'}$, and $G(a_{4g+1}) = a_{4g+1}$. By construction, we have $GF = FG$, and so it follows that $\langle F, G \rangle \cong \mathbb{Z} \times \mathbb{Z}_4$.

Examples 3.6–3.7 can be generalized to the following.

Corollary 3.8. For any positive integer $n \geq 2$ such that $n \mid g$ (resp. $n \mid g - 1$), there is an infinite metacyclic subgroup of $\langle F, G \rangle < \text{Mod}(S_g)$ isomorphic to $\mathbb{Z} \times \mathbb{Z}_n$, where F is

a Penner-type pseudo-Anosov and G is periodic with $D_G = (n, g/n; (1, n), (n-1, n))$ (resp. $D_G = (n, (g+n-1)/n, 1; -)$).

Remark 3.9. An infinite metacyclic subgroup of $\langle F, G \rangle < \text{Mod}(S_g)$ generated by a Penner-type pseudo-Anosov mapping class F can also have pseudo-Anosovs of non-Penner-type. In Examples 3.3 and 3.6, it can be seen that every pseudo-Anosov mapping class in $\langle F, G \rangle$ which is not a power of F is a non-Penner-type pseudo-Anosov. In fact, F can also be replaced with a non-Penner-type pseudo-Anosov generator. In Example 3.6, for $i \not\equiv 0 \pmod{|G|}$ and $j \neq 0$, the mapping class $G^i F^j$ is a non-Penner-type pseudo-Anosov, while in Example 3.3, $G^i F^j$, where i is an even positive integer such that $i \not\equiv 0 \pmod{|G|}$ and $j \neq 0$, is a non-Penner-type pseudo-Anosov. Furthermore, in each case, taking $j = \pm 1$ would yield elements that are possible generators of $\langle F, G \rangle$ in place of F .

Remark 3.10. Let F be a pseudo-Anosov generator of a metacyclic subgroup of $\text{Mod}(S_g)$. Then there is no upper bound on the stretch factor $\lambda(F)$ of F . This follows from the simple fact that if $\langle F, G \rangle$ is a metacyclic subgroup of $\text{Mod}(S_g)$, then $\langle F^n, G \rangle$ is also a metacyclic subgroup for all $n > 1$, where $\lambda(F^n) = \lambda(F)^n$.

3.2. Metacyclic subgroups with reducible generators of infinite order

We begin this subsection with the following lemma which provides necessary and sufficient conditions under which two multitwists are equal.

Lemma 3.11 ([7, Lemma 3.17]). *Let $A = \{a_1, \dots, a_n\}$ and $B = \{b_1, \dots, b_m\}$ be two multicurves in S_g . Let p_i and q_i be nonzero integers. If*

$$T_{a_1}^{p_1} \dots T_{a_n}^{p_n} = T_{b_1}^{q_1} \dots T_{b_m}^{q_m}$$

in $\text{Mod}(S_g)$, then $m = n$ and the sets $\{T_{a_i}^{p_i}\}, \{T_{b_i}^{q_i}\}$ are equal.

We will now establish our second main result that gives necessary and sufficient conditions under which two mapping classes which are not pseudo-Anosov form an infinite metacyclic subgroup of $\text{Mod}(S_g)$.

Theorem 3.12 (Main Theorem 2). *For $g \geq 2$, let $F, G \in \text{Mod}(S_g)$ be two nontrivial mapping classes such that at least one of F or G is of infinite order and neither F nor G is pseudo-Anosov. Assume that F, G have degrees n, m , with multitwist components*

$$T_{c_1}^{q_1} T_{c_2}^{q_2} \dots T_{c_\ell}^{q_\ell} \quad \text{and} \quad T_{c'_1}^{q'_1} T_{c'_2}^{q'_2} \dots T_{c'_{\ell'}}^{q'_{\ell'}},$$

respectively, where $q_i, q'_i \in \mathbb{Z} \setminus \{0\}$, $\mathcal{C}(F) = \{c_1, c_2, \dots, c_\ell\}$, and $\mathcal{C}(G) = \{c'_1, c'_2, \dots, c'_{\ell'}\}$. Then $\langle F, G \rangle$ is an infinite metacyclic subgroup with $\langle F \rangle \triangleleft \langle F, G \rangle$ if and only if the following conditions hold.

- (i) $\mathcal{C}(F) \cup \mathcal{C}(G)$ is a multicurve.
- (ii) If F is periodic with $G^{-1}FG = F^k$, then $k^m \equiv 1 \pmod{n}$.

- (iii) Define $A_i := \{c_j \in \mathcal{C}(F) \mid q_j = q_i\}$, $B_i := \{c_j \in \mathcal{C}(F) \mid q_j = kq_i\}$, and $C_i := \{c'_j \in \mathcal{C}(G) \mid q'_j = q'_i\}$. Then $G(A_i) = B_i$, $G(B_i) = A_i$, and $F(C_i) = C_i$ for every i .
- (iv) For every path component R of $S_g(\mathcal{C}(F) \cup \mathcal{C}(G))$, $G_r^{-1}F_rG_r = F_r^{k p_r}$, where $G_r, F_r \in \text{Mod}(R)$ are induced by G, F , respectively, and p_r is the size of orbit of R under G .
- (v) For two path components R, S of $S_g(\mathcal{C}(F) \cup \mathcal{C}(G))$ such that $G(R) = S$, F_r^k is conjugate to F_s , where $F_r \in \text{Mod}(R)$, $F_s \in \text{Mod}(S)$ are induced by F .

Proof. Let $\langle F, G \rangle$ be an infinite metacyclic subgroup of $\text{Mod}(S_g)$. First, we assume that F has infinite order. Since $G^{-1}FG = F^k$, where $k = \pm 1$, we have $G^{-1}F^nG = F^{kn}$, and so their multitwist components are equal; that is,

$$T_{G^{-1}(c_1)}^{q_1} T_{G^{-1}(c_2)}^{q_2} \cdots T_{G^{-1}(c_\ell)}^{q_\ell} = T_{c_1}^{kq_1} T_{c_2}^{kq_2} \cdots T_{c_\ell}^{kq_\ell}.$$

By Lemma 3.11, it follows that

$$\{T_{G^{-1}(c_i)}^{q_i} \mid 1 \leq i \leq \ell\} = \{T_{c_j}^{kq_j} \mid 1 \leq j \leq \ell\},$$

and so $G(A_i) = B_i$ and $G(B_i) = A_i$ for every i . Hence, $G(\mathcal{C}(F)) = \mathcal{C}(F)$. Since $k = \pm 1$, G^2 commutes with F . By comparing the multitwist components in $FG^2F^{-1} = G^2$, we have $F(C_i) = C_i$ for every i . As $\mathcal{C}(G)$ is the intersection of all maximal reduction system of G , $\mathcal{C}(G)$ is contained in the maximal reduction system of G containing $\mathcal{C}(F)$. Therefore, it follows that $\mathcal{C}(F) \cup \mathcal{C}(G)$ is a multicurve. The same conclusion holds trivially for the case when F is periodic.

Suppose that G has infinite order and F is periodic. Since $G^{-1}FG = F^k$, we have $G^{-ma}FG^{ma} = F^{k^{ma}} = F$, where $a = |k|$. By comparing the multitwist components in $FG^{ma}F^{-1} = G^{ma}$, it follows that

$$T_{F(c'_1)}^{aq'_1} T_{F(c'_2)}^{aq'_2} \cdots T_{F(c'_\ell)}^{aq'_\ell} = T_{c'_1}^{aq'_1} T_{c'_2}^{aq'_2} \cdots T_{c'_\ell}^{aq'_\ell}.$$

By Lemma 3.11, we have $F(C_i) = C_i$ for each i , and so $F(\mathcal{C}(G)) = \mathcal{C}(G)$. Since $F(\mathcal{C}(G)) = \mathcal{C}(G)$ and $G^{-m}(FG^mF^{-1}) = F^{k^m-1}$, it follows that $F^{k^m-1} = 1$. Therefore, $k^m \equiv 1 \pmod{n}$, and we have established (i)–(iii).

Restricting the relation $G^{-1}FG = F^k$ to a path component R of $S_g(\mathcal{C}(F) \cup \mathcal{C}(G))$ gives $(G^{-p_r}FG^{p_r})|_R = F_r^{k p_r}|_R$, where p_r is the size of the orbit of R under G . Therefore, $G_r^{-1}F_rG_r = F_r^{k p_r}$, where $G_r, F_r \in \text{Mod}(R)$ are induced by G, F , respectively. For two distinct path components R, S of $S_g(\mathcal{C}(F) \cup \mathcal{C}(G))$ such that $G(R) = S$, restricting the relation $G^{-1}FG = F^k$ to R , it follows that F_s is conjugate to F_r^k , where $F_r \in \text{Mod}(R)$, $F_s \in \text{Mod}(S)$ are induced by F . This completes the argument for (iv)–(v).

Conversely, we assume that F and G satisfies (i)–(v). Since the relation $G_r^{-1}F_rG_r = F_r^{k p_r}$ holds in $\text{Mod}(R)$ for every path component R of $S_g(\mathcal{C}(F) \cup \mathcal{C}(G))$, it follows from conditions (i)–(iii), (v) that the relation $G^{-1}FG = F^k$ holds in $\text{Mod}(S_g)$. Hence, $\langle F, G \rangle$ is an infinite metacyclic subgroup. \blacksquare




Figure 6. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_3)$ isomorphic to $\mathbb{Z} \times \mathbb{Z}_2$ generated by F such that $F^3 = T_c$ and a free involution G .

We have the following direct consequence of Theorem 3.12.

Corollary 3.13. *For $g \geq 2$, let $F, G \in \text{Mod}(S_g)$ be two nontrivial mapping classes such that at least one of F or G is of infinite order and neither F nor G is pseudo-Anosov. Let $\langle F, G \rangle$ be an infinite metacyclic subgroup of $\text{Mod}(S_g)$ with $\langle F \rangle \triangleleft \langle F, G \rangle$. Then the following statements hold.*

- (i) F and G are reducible mapping classes.
- (ii) If F, G are of infinite order such that G is of odd degree, then $\langle F, G \rangle$ is abelian.
- (iii) If G is of infinite order of degree 1, then $\langle F, G \rangle$ is abelian.

Proof. By Theorem 3.12 (i), (iii), F and G preserve the multicurve $\mathcal{C}(F) \cup \mathcal{C}(G)$. Therefore, F and G are reducible mapping classes. Let n, m denote the degrees of F, G , respectively, where m is odd, and assume that F, G are of infinite order. Since $\mathcal{C}(F) \cup \mathcal{C}(G)$ is a multicurve, comparing the multitwist components in $G^{-m} F^n G^m = F^{nk^m}$, it follows that $k^m = 1$. As m is odd, $k = 1$, which implies that $\langle F, G \rangle$ is abelian. Finally, when F is periodic and G is an infinite order reducible mapping class of degree 1, by Theorem 3.12 (ii), we have that $\langle F, G \rangle$ is abelian. ■

Now, we give several examples of infinite metacyclic subgroups of $\text{Mod}(S_g)$ involving reducible generators. In the following example, we use the n -compatibility of cyclic actions to construct an infinite metacyclic subgroup of $\text{Mod}(S_g)$ generated by a nontrivial periodic and a pseudo-periodic mapping class of infinite order.

Example 3.14. Let $\tilde{F}, \tilde{G} \in \text{Mod}(S_2)$ be periodic mapping classes with

$$D_{\tilde{F}} = (3, 0; (1, 3), (1, 3), (2, 3)_1, (2, 3)_1) \quad \text{and} \quad D_{\tilde{G}} = (2, 1; (1, 2)_2, (1, 2)_2),$$

respectively, as in Figure 6. From the theory developed in [5], there exist conjugates F' and G' of \tilde{F} and \tilde{G} , respectively, that commute in $\text{Mod}(S_g)$. We observe that the orbits corresponding to the cone points of $D_{\tilde{F}}$ (resp. $D_{\tilde{G}}$) with the same suffix are 1-compatible with twist factor 1 (resp. 1-compatible). Hence, F' and G' extend to a pseudo-periodic F and a periodic G (represented by \mathcal{G}), respectively, in $\text{Mod}(S_3)$ such that $F^3 = T_c$ and $D_G = (2, 2, 1; -)$. Since $\langle F', G' \rangle$ is abelian, from Theorem 3.12, it follows that $\langle F, G \rangle \cong \mathbb{Z} \times \mathbb{Z}_2$.




Figure 7. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_{2g+1})$ isomorphic to $\mathbb{Z}_{2g+1} \rtimes_{-1} \mathbb{Z}$ generated by a periodic mapping class F of order $2g+1$ and G such that $G^2 = T_a^2 T_b^{-2}$.

The following corollary is a direct generalization of Example 3.14.

Corollary 3.15. For $g \geq 2$, let $F \in \text{Mod}(S_g)$ be a nontrivial periodic mapping class with

$$D_F = (n, g_0; (a, n), (b, n), (c_1, n_1), \dots, (c_\ell, n_\ell)).$$

For $1 < m < n$ and $m \mid n$ such that $\gcd(m, n/m) = 1$, there is an infinite metacyclic subgroup of $\text{Mod}(S_{g+1})$ isomorphic to $\mathbb{Z} \times \mathbb{Z}_m$ if the following conditions hold.

- (i) $a + b \equiv 0 \pmod{m}$.
- (ii) $a^{-1} + b^{-1} \equiv k \pmod{n/m}$, where $k \in \mathbb{Z}_{n/m} \setminus \{0\}$.

In the following example, we construct a non-abelian infinite metacyclic subgroup $\langle F, G \rangle$, where F is a nontrivial reducible periodic mapping class.

Example 3.16. For $g \geq 1$, let $F_1, F_2 \in \text{Mod}(S_g)$ be two periodic mapping classes (see Figure 7) with

$$\begin{aligned} D_{F_1} &= (2g+1, 0; (1, 2g+1), (g, 2g+1)_1, (g, 2g+1)_2), \\ D_{F_2} &= (2g+1, 0; (2g, 2g+1), (g+1, 2g+1)_1, (g+1, 2g+1)_2). \end{aligned}$$

Since the orbits corresponding to the cone points with the same suffix are 1-compatible, a periodic mapping class $F \in \text{Mod}(S_{2g+1})$ can be constructed from F_1, F_2 with

$$D_F = (2g+1, 1; (1, 2g+1), (2g, 2g+1)).$$

Let $G' \in \text{Mod}(S_{2g+1})$ be an involution represented by \mathcal{G}' as shown in the figure with $D_{G'} = (2, g+1, 1; -)$. From the theory developed in [6], we have $G'^{-1} F G' = F^{-1}$. Now, consider $G \in \text{Mod}(S_{2g+1})$ such that $G = G' T_a T_b^{-1}$. Since $F(a) = a$ and $F(b) = b$, it follows that $G^{-1} F G = F^{-1}$. Hence, $\langle F, G \rangle \cong \mathbb{Z}_{2g+1} \rtimes_{-1} \mathbb{Z}$.

In [6, Example 4.19], an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ was constructed, where F was an infinite-order pseudo-periodic and G was a nontrivial periodic




Figure 8. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_{2g+1})$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$ generated by an F such that $F^{2g+1} = T_a^3 T_b^{-3}$ and an involution G .

mapping class such that $\langle \mathcal{G} \rangle$ acted non-transitively on the path components of $S_g(\mathcal{C}(F))$. We now provide an example in which the action of $\langle \mathcal{G} \rangle$ on $S_g(\mathcal{C}(F))$ is transitive.

Example 3.17. For $g \geq 2$, let $F_1, F_2 \in \text{Mod}(S_g)$ be periodic mapping classes (see Figure 8) with

$$\begin{aligned} D_{F_1} &= (2g+1, 0; (g, 2g+1)_1, (1, 2g+1)_2, (g, 2g+1)), \\ D_{F_2} &= (2g+1, 0; (2g, 2g+1)_1, (g+1, 2g+1)_2, (g+1, 2g+1)). \end{aligned}$$

Here, the orbits corresponding to cone points with the same suffix are 1-compatible with twist factor ± 3 . Thus, there exists a pseudo-periodic $F \in \text{Mod}(S_{2g+1})$ with F_1, F_2 as its canonical components such that

$$F^{2g+1} = T_a^3 T_b^{-3},$$

where $\mathcal{C}(F) = \{a, b\}$ is a bounding pair. Let $G \in \text{Mod}(S_{2g+1})$ be represented by a free involution \mathcal{G} as shown in the figure with $D_G = (2, g+1, 1; -)$. From Theorem 3.12, it follows that $GFG^{-1} = F^{-1}$, and hence, $\langle F, G \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$.

The constructions in Examples 3.16 and 3.17 easily generalize to the following.

Corollary 3.18. For $g \geq 2$, let $F \in \text{Mod}(S_g)$ be a periodic mapping class with

$$D_F = (n, g_0; (a, n), (b, n), (c_1, n_1), \dots, (c_\ell, n_\ell)), \quad \text{where } 3 \leq n \leq 4g.$$

Then the following statements hold.

- (i) If $a = b$, then there is an infinite metacyclic subgroup of $\text{Mod}(S_{2g+1})$ isomorphic to $\mathbb{Z}_n \rtimes_{-1} \mathbb{Z}$.
- (ii) If $a \neq b$, then there is an infinite metacyclic subgroup of $\text{Mod}(S_{2g+1})$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$.

So far, we have only constructed infinite metacyclic subgroups with nontrivial periodic elements. In the next couple of examples, we construct infinite metacyclic subgroups that do not have any nontrivial periodic element.




Figure 9. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ isomorphic to $\mathbb{Z} \times \mathbb{Z}$ and $\mathbb{Z} \rtimes_{-1} \mathbb{Z}$ generated by two pseudo-periodic mapping classes F and G .




Figure 10. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ isomorphic to $\mathbb{Z} \times \mathbb{Z}$ and $\mathbb{Z} \rtimes_{-1} \mathbb{Z}$ generated by two pseudo-periodic mapping classes F and G .

Example 3.19. For an odd integer $g > 1$, let $G_1, G_2 \in \text{Mod}(S_g)$ be represented by a free involution \mathcal{E}_1 and a hyperelliptic involution \mathcal{E}_2 as in Figure 9. We observe that \mathcal{E}_1 and \mathcal{E}_2 commute. Consider $F_1, F_2, G \in \text{Mod}(S_g)$ such that

$$F_1 = G_2 T_a T_b, \quad F_2 = G_2 T_a T_b^{-1} T_e T_f^{-1}, \quad \text{and} \quad G = G_1 T_c.$$

Since $G^2 = T_c^2$, $F_1^2 = T_a T_b T_e T_f$, and $F_2^2 = T_a^2 T_b^{-2} T_e^2 T_f^{-2}$, F_1, F_2 , and G are pseudo-periodic mapping classes. Now, it can be verified that $G^{-1} F_1 G = F_1$ and $G^{-1} F_2 G = F_2^{-1}$. Thus, we have $\langle F_1, G \rangle \cong \mathbb{Z} \times \mathbb{Z}$ and $\langle F_2, G \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}$.

Example 3.20. For an even integer $g > 2$, let $G_1, G_2 \in \text{Mod}(S_g)$ be represented by an involution \mathcal{E}_1 and a hyperelliptic involution \mathcal{E}_2 as in Figure 10. We observe that G_1 and G_2 commute. Consider $F, G \in \text{Mod}(S_g)$ such that $F = G_2 T_c T_d^{-1}$ and $G = G_1 T_a T_e$. Since $F^2 = T_c^2 T_d^{-2}$ and $G^2 = T_a T_b T_e T_f$, F and G are pseudo-periodic mapping classes. It can be verified that $G^{-1} F G = F^{-1}$, and hence we have $\langle F, G \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}$. Considering $F' \in \text{Mod}(S_g)$ such that $F' = G_2 T_c T_d$, it can be seen that $\langle F', G \rangle \cong \mathbb{Z} \times \mathbb{Z}$.

Taking inspiration from [6, Example 4.20], where a non-abelian infinite metacyclic subgroup was constructed with a nontrivial periodic generator, we will now describe an example where both generators are pseudo-periodics of infinite order.

Example 3.21. Let $F, G' \in \text{Mod}(S_{13})$ such that $F^3 = T_{c_1} T_{c_2}^{-1} T_{c_3} T_{c_4}^{-1}$ and G' represented by \mathcal{E}' with $D_{\mathcal{E}'} = (4, 4, 1; -)$ (see Figure 11). In [6, Example 4.20], it was shown that $G' F G'^{-1} = F^{-1}$, and therefore $\langle F, G' \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}_4 < \text{Mod}(S_{13})$. Now, we consider




Figure 11. Realization of an infinite metacyclic subgroup of $\text{Mod}(S_{13})$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}$.

$G \in \text{Mod}(S_{13})$ such that $G = G' T_{c_1}$. Since $G^4 = T_{c_1} T_{c_2} T_{c_3} T_{c_4}$, the G is pseudo-periodic of degree 4. As $F(c_1) = c_1$ and

$$G^{-1} F G = T_{c_1}^{-1} G'^{-1} F G' T_{c_1} = T_{c_1}^{-1} F^{-1} T_{c_1} = F^{-1},$$

we have $\langle F, G \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}$.

In the preceding examples, we saw infinite metacyclic subgroups with pseudo-periodic generators. In the following examples, we construct infinite metacyclic subgroups with an infinite order reducible generator with canonical components that are nontrivial periodic and pseudo-Anosov.

Example 3.22. For $g \geq 3$, consider the collection of curves as shown in Figure 12 and the mapping class

$$F = T_{b_1} T_{b_2} T_{a_1}^{-1} T_{a_2}^{-1} T_{b_3} \prod_{i=3}^g T_{a_i} T_{b_{i+1}}.$$

Since $F(b_3) = b_3$, F is a reducible mapping class of infinite order with pseudo-Anosov canonical component $T_{b_1} T_{b_2} T_{a_1}^{-1} T_{a_2}^{-1}$ and periodic canonical component $\prod_{i=3}^g T_{a_i} T_{b_{i+1}}$. Let G be the hyperelliptic involution as shown in the figure. Since $G^{-1} F G = F$, we have $\langle F, G \rangle \cong \mathbb{Z} \times \mathbb{Z}_2$.




Figure 12. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ isomorphic to $\mathbb{Z} \times \mathbb{Z}_2$ generated by a hyperelliptic involution G and a reducible mapping class F of infinite order.




Figure 13. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_6)$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$ generated by an involution G and a reducible mapping class F of infinite order.

The construction in Example 3.22 generalizes to the following assertion.

Corollary 3.23. *For $g \geq 2$, there is an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ isomorphic to $\mathbb{Z} \times \mathbb{Z}_2$ generated by a hyperelliptic involution G and a reducible mapping class of infinite order containing at least one pseudo-Anosov and one nontrivial periodic canonical component.*

Example 3.24. Consider the collection of curves in S_6 as shown in Figure 13 and the mapping class

$$F = (T_{b_1} T_{a_1} T_{b_2})(T_{c_1} T_{c_2}^{-1})(T_{b_3} T_{d_1} T_{a_4} T_{a_3}^{-1} T_{d_2}^{-1} T_{b_5}^{-1})(T_{c_3} T_{c_4}^{-1})(T_{b_6}^{-1} T_{a_6}^{-1} T_{b_7}^{-1}).$$

Since $F(\{c_1, c_2, c_3, c_4\}) = \{c_1, c_2, c_3, c_4\}$, F is a reducible mapping class of infinite order with two nontrivial periodic canonical components and one pseudo-Anosov canonical component. Let $G \in \text{Mod}(S_6)$ be an involution as shown in the figure. Since $G^{-1} F G = F^{-1}$, $\langle F, G \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$.

A direct generalization of Example 3.24 is the following result.

Corollary 3.25. *For an even integer $g \geq 4$, there is an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$ generated by an involution G with $D_G = (2, g/2; (1, 2), (1, 2))$ and a reducible mapping class of infinite order containing at least one pseudo-Anosov and one nontrivial periodic canonical component.*

4. Applications

In this section, we derive some applications of the theory developed in this paper.

4.1. Infinite metacyclic subgroups of the level m subgroup of $\text{Mod}(S_g)$

The action of $\text{Mod}(S_g)$ on $H_1(S_g, \mathbb{Z})$ affords a surjective representation [7, Chapter 6] $\Psi : \text{Mod}(S_g) \rightarrow \text{Sp}(2g, \mathbb{Z})$. The subgroup $\ker \Psi$ is known as the *Torelli group* and is denoted by $\mathcal{I}(S_g)$. Further, for an integer $m \geq 2$, the *level m congruence subgroup* is the kernel of the composition

$$\text{Mod}(S_g) \rightarrow \text{Sp}(2g, \mathbb{Z}) \rightarrow \text{Sp}(2g, \mathbb{Z}_m),$$

denoted by $\text{Mod}(S_g)[m]$. By definition $\mathcal{I}(S_g) \subset \text{Mod}(S_g)[m]$ for every m . For $m \geq 3$, it is known [7, Chapter 6] that $\text{Mod}(S_g)[m]$ is torsion-free and that an infinite order reducible in $\text{Mod}(S_g)[m]$ has degree 1 [13, Corollary 1.8]. The only torsion elements of $\text{Mod}(S_g)[2]$ are the hyperelliptic involutions. The following result follows immediately from Theorems 3.1, 3.12, and Corollary 3.13.

Proposition 4.1. *For $g \geq 2$ and $m \geq 3$, let $F, G \in \text{Mod}(S_g)[m]$ be two nontrivial mapping classes. Then $\langle F, G \rangle$ is metacyclic with $\langle F \rangle \triangleleft \langle F, G \rangle$ if and only if the following hold.*

- (i) *F and G are infinite order reducible mapping classes that commute.*
- (ii) *$\mathcal{C}(F) \cup \mathcal{C}(G)$ is a multicurve.*
- (iii) *The nontrivial canonical components of F and G are pseudo-Anosov mapping classes.*
- (iv) *The nontrivial canonical components of F and G with the same support generate a cyclic group.*

In the following examples, we construct infinite metacyclic subgroups of $\text{Mod}(S_g)[2]$ with a pseudo-Anosov generator. Since the hyperelliptic involution of $\text{Mod}(S_2)$ lies in the center, we will assume $g \geq 3$.

Example 4.2. Consider the multicurves $A = \{a_1, a_2, \dots, a_g\}$ and $B = \{b_1, b_2, \dots, b_{g+1}\}$ as shown in Figure 14. Since the curves of $A \cup B$ fill S_g , by Theorem 2.11, the mapping class

$$F = \prod_{i=1}^g T_{a_i}^2 \prod_{i=i}^{g+1} T_{b_i}^{-2}$$

is pseudo-Anosov. Let G be the hyperelliptic involution as shown in Figure 14. Since $G(c) = c$ for every $c \in A \cup B$, we have $G^{-1}FG = F$. As $F, G \in \text{Mod}(S_g)[2]$, $\langle F, G \rangle < \text{Mod}(S_g)[2]$ isomorphic to $\mathbb{Z} \times \mathbb{Z}_2$.

The following example draws inspiration from a family of Penner-type pseudo-Anosov mapping classes described in [2].

Example 4.3. For $g \geq 3$, we construct a non-abelian metacyclic subgroup of $\text{Mod}(S_g)[2]$ generated by a pseudo-Anosov element in $\mathcal{I}(S_g)$. First we describe a filling collection of curves C in S_g which is a disjoint union of two multicurves A and B . Consider the surfaces S, S' , and S'' with curves and arcs as shown in Figure 15. We construct a closed




Figure 14. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)[2]$ isomorphic to $\mathbb{Z} \times \mathbb{Z}_2$ generated by a hyperelliptic involution G and a pseudo-Anosov F .




Figure 15. The surface S'' on the left, S in center, and S' on the right used to construct a filling system of curves in S_g .




Figure 16. A filling system of curves in S_4 and $S_7 = S + S' + S$.

surface by combining multiple copies of S , S' , and S'' as follows. For $X_i \in \{S, S', S''\}$, we write $X_1 + X_2 \cdots + X_n$ for the surface obtained by gluing X_i end to end and capping the remaining boundary components after gluing. For $m \geq 1$, we write mS for $S + S + \cdots + S$. For $m \geq 1$, we can write $S_g = mS$ if $g = 3m$, $S_g = S + S' + mS$ if $g = 3m + 4$, $S_g = S + S' + mS + S' + S$ if $g = 3m + 8$, $S_5 = S''$, and $S_8 = S'' + S'$. The multicurves A and B are drawn with red and blue color, respectively.

Let F be a product of positive (left-handed) Dehn twists about the curves in A and negative Dehn twists about the curves in B , where each Dehn twist is taken exactly once. We observe that for each curve $a \in A$, there exist a unique curve $b \in B$ such that $\{a, b\}$ bounds a subsurface of S_g and vice-versa. Since $A \cup B$ fills S_g , by Theorem 2.11, F is a pseudo-Anosov. It can be seen that $F \in \mathcal{I}(S_g)$. Let G be the hyperelliptic involution shown in Figure 16. Since G exchanges multicurves A and B , we have

$$G^{-1}FG = F^{-1}.$$

Hence, $\langle F, G \rangle < \text{Mod}(S_g)[2]$ is isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_2$.

4.2. Bounds on the order of a periodic generator of an infinite metacyclic group

In this subsection, we derive bounds on the order of a nontrivial periodic generator of an infinite metacyclic subgroup of $\text{Mod}(S_g)$ which are realized. We will require the following remark.

Remark 4.4. For an even integer g , let $F \in \text{Mod}(S_g)$ be a nontrivial reducible periodic mapping class of order $2g + 2$. From the theory developed in [26], it follows that F arises as a 1-compatibility between fixed points of F' and F'^{-1} , where $F' \in \text{Mod}(S_{g/2})$ is a periodic mapping class of order $4(g/2) + 2 = 2g + 2$. Hence, F has a unique maximal reduction system containing a single separating curve.

In the following proposition, we obtain bounds on the order of a nontrivial periodic generator which are realized.

Proposition 4.5. *For $g \geq 2$, let $F, G \in \text{Mod}(S_g)$ be two nontrivial mapping classes such that $\langle F, G \rangle$ is an infinite metacyclic subgroup with $\langle F \rangle \triangleleft \langle F, G \rangle$.*

- (i) *Let F be a pseudo-Anosov mapping class and G a periodic mapping class.*
 - (a) *If $\langle F, G \rangle$ is non-abelian, then $2 \leq |G| \leq 4g$.*
 - (b) *If $\langle F, G \rangle$ is abelian, then $2 \leq |G| \leq 2g$.*
- (ii) *Let F be a reducible mapping class of infinite order and G a periodic mapping class.*
 - (a) *If $\langle F, G \rangle$ is abelian, then $2 \leq |G| \leq 2g + 2$.*
 - (b) *If $\langle F, G \rangle$ is non-abelian, then $2 \leq |G| \leq 2g$.*
- (iii) *If F is periodic and $\langle F, G \rangle$ is non-abelian, then $3 \leq |F| \leq 2g + 2$.*

Moreover, all of the above bounds are realized.

Proof. (i) Suppose that $\langle F, G \rangle$ is non-abelian. Example 4.6 shows that an order $4g$ periodic mapping class can form a non-abelian metacyclic subgroup with F . Since it is known that $|G| \leq 4g + 2$ [11] and there is no periodic mapping class of order $4g + 1$ (Lemma 2.5), it suffices to show that $|G| \neq 4g + 2$. If $|G| = 4g + 2$, then by Lemma 2.4, G^2 is irreducible. Furthermore, from Theorem 2.3, it follows that $\mathcal{O}_{G^2} \approx S_{0,3}$. Since G^2 commutes with F , by Lemma 2.10, F induces an infinite order mapping class in the finite group $\text{Mod}(S_{0,3})$, which is impossible. Thus, it follows that $|G| \leq 4g$.

Next, we consider the case when $\langle F, G \rangle$ is abelian. The preceding argument shows that G is a reducible mapping class. By Lemma 2.4, it follows that $|G| \leq 2g + 2$. Moreover, Example 4.6 shows that an order $2g$ periodic mapping class can form an infinite abelian metacyclic subgroup with F . Since G is reducible, it suffices to show that $|G| \neq 2g + 2$. Let G be a reducible periodic mapping class of order $2g + 2$ that commutes with F . By Remark 4.4, G has a unique maximal reduction system containing a single separating curve, say, c . Since $GF = FG$, we have $GF(c) = F(c)$, and so $F(c) = c$, which is not possible as F is irreducible. Thus, it follows that $|G| \leq 2g$. Examples 4.2–4.3 show that the lower bounds are realized.

(ii) Let $\langle F, G \rangle$ be abelian. From Corollary 3.13, it follows that G is reducible, and from Lemma 2.4, we have $|G| \leq 2g + 2$. As before, a periodic mapping class of order $2g + 2$ has a unique maximal reduction system $\{c\}$. Taking $F = T_c$, we have G commutes with F . This shows that the upper bound $2g + 2$ is realized when $\langle F, G \rangle$ is abelian.




Figure 17. Realization of an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_2)$ isomorphic to $\mathbb{Z} \rtimes_{-1} \mathbb{Z}_8$ generated by an irreducible periodic mapping class G of order 8 and a pseudo-Anosov mapping class F .

Let $\langle F, G \rangle$ be non-abelian. If $|G| = 2g + 2$, then from Theorem 3.12, it follows that $G(\mathcal{C}(F)) = \mathcal{C}(F)$. Since G has a unique maximal reduction system $\{c\}$, it follows that $\mathcal{C}(F) = \{c\}$ (as $\mathcal{C}(F) \neq \emptyset$). Therefore, the multitwist component of F^n is T_c^q , for some $n \in \mathbb{N}$ and $q \in \mathbb{Z} \setminus \{0\}$. By comparing multitwist components in $G^{-1}F^nG = F^{-n}$, we have $G^{-1}T_c^qG = T_c^{-q}$. This is impossible since G commutes with T_c . Hence, $|G| \leq 2g$ and Example 4.7 shows that this upper bound is realized. Examples 3.17 and 4.7 show that the lower bounds are realized.

(iii) Since $\langle F, G \rangle$ is non-abelian and $\langle F \rangle \triangleleft \langle F, G \rangle$, we have $k \in \mathbb{Z}_n^\times \setminus \{1\}$, where $n = |F|$. Hence, $n \geq 3$, and by Theorem 3.1 and Corollary 3.13, F is reducible. From Lemma 2.4, we have $n \leq 2g + 2$. Thus, the assertion follows, and by Corollaries 3.18 and 4.9, it follows that the bounds are realized. ■

We will now provide examples demonstrating that the upper bound on the order of the periodic generator G of the group $\langle F, G \rangle$ obtained in Proposition 4.5 is realized.

Example 4.6. For $g \geq 2$, let $G \in \text{Mod}(S_g)$ be a periodic mapping class of order $4g$ realized as $2\pi/4g$ -rotation of a $4g$ -gon with side-pairing $a_1a_2 \cdots a_{2g}a_1^{-1}a_2^{-1} \cdots a_{2g}^{-1}$ as shown in Figure 17 (for $g = 2$). For $1 \leq i \leq 2g - 1$, let $c_i = a_i a_{i+1}$, $c_{2g} = a_{2g} a_1^{-1}$, $A = \{c_1, c_3, \dots, c_{2g-1}\}$, and $B = \{c_2, c_4, \dots, c_{2g}\}$. We note that c_i is homotopic to the concatenation of a_i and a_{i+1} . We observe that A and B are multicurves such that the curves in $A \cup B$ fill S_g . By Theorem 2.11, the mapping class

$$F = \prod_{k=1}^g T_{c_{2k-1}} \prod_{k=1}^g T_{c_{2k}}^{-1}$$

is pseudo-Anosov. Since $G(c_i) = c_{i+1}$, where $1 \leq i \leq 2g - 1$ and $G(c_{2g}) = c_1$, we have $G^{-1}FG = F^{-1}$. Hence, $\langle F, G \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}_{4g}$. Furthermore, since $G^{-2}FG^2 = F$, we have $\langle F, G^2 \rangle \cong \mathbb{Z} \times \mathbb{Z}_{2g}$.




Figure 18. An infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_2)$ isomorphic to $\mathbb{Z}_6 \rtimes_{-1} \mathbb{Z}$ generated by an order 6 mapping class F and a pseudo-periodic mapping class G .

Example 4.7. For even integer $g \geq 2$, let $G \in \text{Mod}(S_g)$ be a periodic of order $2g$ realized as the square of $2\pi/4g$ -rotation of a $4g$ -gon with side-pairing $a_1 a_2 \cdots a_{2g} a_1^{-1} a_2^{-1} \cdots a_{2g}^{-1}$ as shown in Figure 17 (for $g = 2$). Consider the multicurve $\mathcal{C} = \{c_{2i-1} := a_{2i-1} a_{2i} \mid 1 \leq i \leq g\}$, where c_{2i-1} is homotopic to the concatenation of a_{2i-1} and a_{2i} . Define the multitwist

$$F = \prod_{i=1}^{g/2} T_{c_{4i-3}} T_{c_{4i-1}}^{-1}.$$

For $1 \leq i \leq g-1$, as $G(c_{2i-1}) = c_{2i+1}$ and $G(c_{2g-1}) = c_1$, we have $G^{-1}FG = F^{-1}$ and $G^{-g}FG^g = F$. Therefore, it follows that $\langle F, G \rangle \cong \mathbb{Z} \rtimes_{-1} \mathbb{Z}_{2g}$ and $\langle F, G^g \rangle \cong \mathbb{Z} \times \mathbb{Z}_2$.

The following example shows that the upper bound on the order of the periodic generator F of a non-abelian metacyclic subgroup $\langle F, G \rangle$ obtained in Proposition 4.5 is realized when G is reducible of infinite order.

Example 4.8. Let $F_1, F_2 \in \text{Mod}(S_1)$ be periodic with

$$D_{F_1} = (6, 0; (1, 2), (1, 3), (1, 6)_1) \text{ and } D_{F_2} = (6, 0; (1, 2), (2, 3), (5, 6)_1).$$

Since the orbits corresponding to cone points with the same suffix are 1-compatible, a periodic mapping class $F \in \text{Mod}(S_2)$ can be constructed from F_1, F_2 with

$$D_F = (6, 0; (1, 2), (1, 2), (1, 3), (2, 3)).$$

Let $G' \in \text{Mod}(S_2)$ be an involution represented by \mathcal{G}' as shown in Figure 18 with $D_{G'} = (2, 1; (1, 2), (1, 2))$. From the theory developed in [6], it follows that $G'^{-1}FG' = F^{-1}$. Now, consider $G \in \text{Mod}(S_2)$ such that $G = G'T_c$. Since $F(c) = c$, we have $G^{-1}FG = T_c^{-1}G'^{-1}FG'T_c = F^{-1}$, and hence, $\langle F, G \rangle \cong \mathbb{Z}_6 \rtimes_{-1} \mathbb{Z}$.

Example 4.8 generalizes to the following corollary.

Corollary 4.9. For an even integer $g \geq 2$, there is an infinite metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ isomorphic to $\mathbb{Z}_{2g+2} \rtimes_{-1} \mathbb{Z}$, where F is a nontrivial periodic mapping class and G is a pseudo-periodic mapping class of infinite order.

4.3. Types of elements in an infinite metacyclic group

Let $\langle F, G \rangle < \text{Mod}(S_g)$ be an infinite metacyclic subgroup with $\langle F \rangle \triangleleft \langle F, G \rangle$. In this subsection, we determine the Nielsen–Thurston type of the elements in $\langle F, G \rangle$ depending upon the Nielsen–Thurston type of F, G .

Lemma 4.10. *For $g \geq 2$, consider a (non-cyclic) metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ that admits the presentation $\langle F, G \mid G^{-1}FG = F^k \rangle$, where $k = \pm 1$. Then every nontrivial element of $\langle F, G \rangle$ is a reducible mapping class of infinite order.*

Proof. From Lemma 2.8, $F^i G^j = G^j F^{ik^j}$, so every element of $\langle F, G \rangle$ is of the form $G^i F^j$ for some integers i, j . For $G^i F^j \in \langle F, G \rangle$, from Lemma 2.8, we have

$$(G^i F^j)^\ell = G^{i\ell} F^{j(1+k^i+k^{2i}+\dots+k^{i(\ell-2)}+k^{i(\ell-1)})}.$$

It follows that every nontrivial element $G^i F^j \in \langle F, G \rangle$ has infinite order. Furthermore, by Theorem 3.1, neither G nor F can be pseudo-Anosov mapping classes. When G and F are infinite order reducibles, from Theorem 3.12, it follows that every nontrivial element of $\langle F, G \rangle$ preserves the multicurve $\mathcal{C}(F) \cup \mathcal{C}(G)$ and hence is a reducible mapping class. ■

Lemma 4.11. *For $g \geq 2$, consider a metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ that admits the presentation $\langle F, G \mid F^n = 1, G^{-1}FG = F^k \rangle$, where $n \geq 3$ and $k \in \mathbb{Z}_n^\times \setminus \{1\}$. Every nontrivial element of $\langle F, G \rangle$, except the powers of F , is of the same Nielsen–Thurston type as G .*

Proof. We note that G can be either pseudo-Anosov or reducible of infinite order. When G is reducible, from Theorem 3.12, we have $F(\mathcal{C}(G)) = \mathcal{C}(G)$. Moreover, for $i \neq 0$, we consider $G^i F^j \in \langle F, G \rangle$ and set $\ell = |F||k|$. Then from Lemma 2.8, we have

$$\begin{aligned} (G^i F^j)^\ell &= G^{i\ell} F^{j(1+k^i+k^{2i}+\dots+k^{i(\ell-2)}+k^{i(\ell-1)})} \\ &= G^{i\ell} F^{j|F|(1+k^i+k^{2i}+\dots+k^{i(|k|-2)}+k^{i(|k|-1)})} = G^{i\ell}. \end{aligned}$$

Hence, every nontrivial element of $\langle F, G \rangle$, except the powers of F , has the same Nielsen–Thurston type as G . ■

Lemma 4.12. *For $g, m \geq 2$, consider a metacyclic subgroup $\langle F, G \rangle < \text{Mod}(S_g)$ that admits the presentation $\langle F, G \mid G^m = 1, G^{-1}FG = F^k \rangle$, $k = \pm 1$ (for $k = -1$, m is even).*

- (i) *If $\langle F, G \rangle$ is abelian, then every nontrivial element of $\langle F, G \rangle$, except the powers of G , has the same Nielsen–Thurston type as F .*
- (ii) *If $\langle F, G \rangle$ is non-abelian, then for integers i, j , where $j \neq 0$, $G^i F^j$ has the same Nielsen–Thurston type as F if i is even, and $G^i F^j$ is periodic of order $|G^i|$ when i is odd. Furthermore, for i odd and j even, $G^i F^j$ is conjugate to G^i .*

Proof. When $\langle F, G \rangle$ is abelian, it follows that every nontrivial element of $\langle F, G \rangle$, except the powers of G , is of infinite order of the same Nielsen–Thurston type as of F . We now consider the case when $\langle F, G \rangle$ is non-abelian. Since G^2 commutes with F , $G^{2i} F^j$ has the same Nielsen–Thurston type as F , where $j \neq 0$. By Lemma 2.8,

$$(G^i F^j)^2 = G^{2i} F^{(1+(-1)^i)} = G^{2i}$$

if and only if i is odd. Thus, it follows that, for $j \neq 0$, $G^i F^j$ is periodic of order $|G^i|$ if and only if i is odd. When j is even and i is odd, we have $F^{j/2}(G^i F^j)F^{-j/2} = F^{j/2}G^i F^{j/2} = G^i F^{(j[1+(-1)^i])/2} = G^i$. Therefore, $G^i F^j$ is conjugate to G^i . ■

4.4. Centralizers of irreducible periodic mapping classes

In this subsection, we describe the centralizers of irreducible periodic mapping classes in $\text{Mod}(S_g)$.

Proposition 4.13. *For $g \geq 2$, let $F \in \text{Mod}(S_g)$ be an irreducible periodic mapping class with $D_F = (n, 0; (c_1, n_1), (c_2, n_2), (c_3, n_3))$. Let H be the centralizer of F in $\text{Mod}(S_g)$.*

- (i) *If either $n > 2g + 2$, or the (c_i, n_i) are all distinct for $i = 1, 2, 3$, then $H = \langle F \rangle$.*
- (ii) *If $n \leq 2g + 2$ and $(c_i, n_i) = (c_j, n_j)$ for some $i, j \in \{1, 2, 3\}$ and $i \neq j$, then $H = \langle F \rangle \times \langle i \rangle$, where i is a hyperelliptic involution.*

Proof. Since F is irreducible, by Theorem 2.3, $\mathcal{O}_F \approx S_{0,3}$. By Definition 2.1 (vi), we have $n \geq 2g + 1$. For $G \in H \setminus \langle F \rangle$, by Lemma 2.10, there exists $\bar{\mathcal{G}} \in \text{Aut}(\mathcal{O}_F)$ induced by \mathcal{G} . Since $\text{Mod}(S_{0,3}) \cong \Sigma_3$ (where Σ_3 is the permutation group on three letters), it follows that $|G| = 2$ or 3 .

First, we consider the case when $|G| = 3$. Since $\bar{\mathcal{G}} \in \text{Aut}(\mathcal{O}_F)$, $\bar{\mathcal{G}}$ permutes the three cone points of \mathcal{O}_F , which implies that $(c_i, n_i) = (c_j, n_j)$ for all $i, j \in \{1, 2, 3\}$. By Definition 2.1 (iv), we have $(c_i, n_i) = (c_1, n)$ for every i . Moreover, by Definition 2.1 (v), we have $3c_1 \equiv 0 \pmod{n}$, which is impossible (as $n \geq 4$).

We now consider the case when $|G| = 2$. Since $\bar{\mathcal{G}} \in \text{Aut}(\mathcal{O}_F)$, $\bar{\mathcal{G}}$ permutes two cone points of \mathcal{O}_F and fixes the third one. By Definition 2.1 (iv), we have $(c_i, n_i) = (c_j, n_j) = (c_i, n)$, for some $i, j \in \{1, 2, 3\}$ and $i \neq j$. We note that such a $\bar{G} \in \Sigma_3$ is uniquely determined. Since $G \in H$, we have $\langle F, G \rangle \cong \mathbb{Z}_n \times \mathbb{Z}_2$. By a result of Maclachlan [21], it follows that $|\langle F, G \rangle| \leq 4g + 4$, and this implies that $n \leq 2g + 2$. Thus, (i) follows.

When $n = 2g + 1$, by Definition 2.1 (vi), we have $n_i = 2g + 1$ for every i . Since $\Gamma(S_g / \langle \mathcal{F}, \mathcal{G} \rangle) = (0; 2, 2g + 1, 4g + 2)$, it follows that $\langle \mathcal{F}, \mathcal{G} \rangle$ is cyclic. From the theory developed in [5], it follows that \mathcal{G} is a hyperelliptic involution. When $n = 2g + 2$, by Definition 2.1 (vi), we have $\Gamma(\mathcal{O}_F) = (0; g + 1, 2g + 2, 2g + 2)$. Since $\bar{\mathcal{G}} \in \text{Aut}(\mathcal{O}_F)$, it follows that $\Gamma(S_g / \langle \mathcal{F}, \mathcal{G} \rangle) = (0; 2, 2g + 2, 2g + 2)$. Again, from the theory developed in [5], the possible data sets for G are

- (a) $D_{G_1} = (2, 0; \underbrace{(1, 2), (1, 2), \dots, (1, 2)}_{(2g+2) \text{ times}})$,

(b) $D_{G_2} = (2, g/2; (1, 2), (1, 2))$ when g is even, and

(c) $D_{G_3} = (2, (g + 1)/2, 1; -)$ when g is odd.

Furthermore, it can be shown that $F^{g+1}G_1$ is conjugate to G_2 (resp. G_3) when g is even (resp. odd). Further, we note that lifts of \bar{G} are GF^j , where $1 \leq j \leq n$. This concludes our argument for (ii). ■

Acknowledgments. The authors are grateful to the referees for helpful suggestions that have significantly improved the exposition in this manuscript.

Funding. The first author was supported by the Prime Minister Research Fellowship (PMRF) scheme instituted by the Ministry of Education, India.

References

- [1] N. Agarwal, S. Dey, N. K. Dhanwani, and K. Rajeevsarathy, Lifiable mapping class groups of regular cyclic covers. *Houston J. Math.* **47** (2021), no. 1, 223–243 Zbl 1489.57016 MR 4447971
- [2] I. Agol, C. J. Leininger, and D. Margalit, Pseudo-Anosov stretch factors and homology of mapping tori. *J. Lond. Math. Soc. (2)* **93** (2016), no. 3, 664–682 Zbl 1388.37033 MR 3509958
- [3] J. S. Birman and H. M. Hilden, On isotopies of homeomorphisms of Riemann surfaces. *Ann. of Math. (2)* **97** (1973), 424–439 Zbl 0237.57001 MR 0325959
- [4] S. A. Broughton and A. Wootton, Finite abelian subgroups of the mapping class group. *Algebr. Geom. Topol.* **7** (2007), 1651–1697 Zbl 1126.14038 MR 2366175
- [5] N. K. Dhanwani and K. Rajeevsarathy, Commuting conjugates of finite-order mapping classes. *Geom. Dedicata* **209** (2020), 69–93 Zbl 1451.57012 MR 4163393
- [6] N. K. Dhanwani, K. Rajeevsarathy, and A. Sanghi, Split metacyclic actions on surfaces. *New York J. Math.* **28** (2022), 617–649 Zbl 07530325 MR 4401493
- [7] B. Farb and D. Margalit, *A primer on mapping class groups*. Princeton Math. Ser. 49, Princeton University Press, Princeton, NJ, 2012 Zbl 1245.57002 MR 2850125
- [8] A. Fathi, F. Laudenbach, and V. Poénaru, *Thurston’s work on surfaces*. Math. Notes 48, Princeton University Press, Princeton, NJ, 2012 Zbl 1244.57005 MR 3053012
- [9] T. Ghaswala and R. R. Winarski, The liftable mapping class group of balanced superelliptic covers. *New York J. Math.* **23** (2017), 133–164 Zbl 1402.57003 MR 3611078
- [10] J. Gilman, Structures of elliptic irreducible subgroups of the modular group. *Proc. London Math. Soc. (3)* **47** (1983), no. 1, 27–42 Zbl 0522.30035 MR 0698926
- [11] W. J. Harvey, Cyclic groups of automorphisms of a compact Riemann surface. *Quart. J. Math. Oxford Ser. (2)* **17** (1966), 86–97 Zbl 0156.08901 MR 0201629
- [12] C. E. Hempel, *Metacyclic groups*. Ph.D. thesis, Australian National University, 1998, https://openresearch-repository.anu.edu.au/bitstream/1885/144742/2/b20383460_hempel_e_j_c.pdf visited on 28 March 2024
- [13] N. V. Ivanov, *Subgroups of Teichmüller modular groups*. Transl. Math. Monogr. 115, American Mathematical Society, Providence, RI, 1992 Zbl 0776.57001 MR 1195787

- [14] N. V. Ivanov, Mapping class groups. In *Handbook of geometric topology*, pp. 523–633, North-Holland, Amsterdam, 2002 Zbl [1002.57001](#) MR [1886678](#)
- [15] P. Kapari and K. Rajeevsarathy, [General primitivity in the mapping class group](#). *J. Topol. Anal.* (2024), DOI [10.1142/S1793525323500541](#)
- [16] Y. Kasahara, Reducibility and orders of periodic automorphisms of surfaces. *Osaka J. Math.* **28** (1991), no. 4, 985–997 Zbl [0758.57010](#) MR [1152963](#)
- [17] S. Katok, *Fuchsian groups*. Chicago Lectures in Math., University of Chicago Press, Chicago, IL, 1992 Zbl [0753.30001](#) MR [1177168](#)
- [18] S. P. Kerckhoff, [The Nielsen realization problem](#). *Ann. of Math. (2)* **117** (1983), no. 2, 235–265 Zbl [0528.57008](#) MR [0690845](#)
- [19] R. S. Kulkarni, [Riemann surfaces admitting large automorphism groups](#). In *Extremal Riemann surfaces (San Francisco, CA, 1995)*, pp. 63–79, Contemp. Math. 201, American Mathematical Society, Providence, RI, 1997 Zbl [0863.30050](#) MR [1429195](#)
- [20] A. M. Macbeath, *Discontinuous groups and birational transformations: Summer School, Queen's College Dundee, University of St. Andrews*. Department of Mathematics, Queen's College, 1961
- [21] C. Maclachlan, [Abelian groups of automorphisms of compact Riemann surfaces](#). *Proc. London Math. Soc. (3)* **15** (1965), 699–712 Zbl [0156.08902](#) MR [0179348](#)
- [22] C. Maclachlan and W. J. Harvey, [On mapping-class groups and Teichmüller spaces](#). *Proc. London Math. Soc. (3)* **30** (1975), no. part, part 4, 496–512 Zbl [0303.32020](#) MR [0374414](#)
- [23] J. D. McCarthy, Normalizers and centralizers of pseudo-Anosov mapping classes. 1994, <https://users.math.msu.edu/users/mccarthy/publications/normcent.pdf> visited on 28 March 2024
- [24] D. McCullough and K. Rajeevsarathy, [Roots of Dehn twists](#). *Geom. Dedicata* **151** (2011), 397–409 Zbl [1219.57019](#) MR [2780759](#)
- [25] J. Nielsen, Die Struktur periodischer Transformationen von Flächen. *Danske Vid. Selsk. Math.-Fys. Medd.* **15** (1937), no. 1, 1–77 Zbl [0017.13302](#)
- [26] S. Parsad, K. Rajeevsarathy, and B. Sanki, [Geometric realizations of cyclic actions on surfaces](#). *J. Topol. Anal.* **11** (2019), no. 4, 929–964 Zbl [1432.57047](#) MR [4040017](#)
- [27] R. C. Penner, [A construction of pseudo-Anosov homeomorphisms](#). *Trans. Amer. Math. Soc.* **310** (1988), no. 1, 179–197 Zbl [0706.57008](#) MR [0930079](#)
- [28] K. Rajeevsarathy, [Roots of Dehn twists about separating curves](#). *J. Aust. Math. Soc.* **95** (2013), no. 2, 266–288 Zbl [1284.57018](#) MR [3142359](#)
- [29] K. Rajeevsarathy, [Fractional powers of Dehn twists about nonseparating curves](#). *Glasg. Math. J.* **56** (2014), no. 1, 197–210 Zbl [1286.57017](#) MR [3137860](#)
- [30] K. Rajeevsarathy and A. Sanghi, [Metacyclic actions on surfaces](#). 2022, arXiv:[2201.09602](#)
- [31] K. Rajeevsarathy and P. Vaidyanathan, [Roots of Dehn twists about multicurves](#). *Glasg. Math. J.* **60** (2018), no. 3, 555–583 Zbl [1409.57022](#) MR [3830536](#)
- [32] W. P. Thurston, [The geometry and topology of three manifolds](#). <https://library.slmath.org/books/gt3m/> visited on 28 March 2024
- [33] W. P. Thurston, [On the geometry and dynamics of diffeomorphisms of surfaces](#). *Bull. Amer. Math. Soc. (N.S.)* **19** (1988), no. 2, 417–431 Zbl [0674.57008](#) MR [0956596](#)
- [34] T. W. Tucker, [Finite groups acting on surfaces and the genus of a group](#). *J. Combin. Theory Ser. B* **34** (1983), no. 1, 82–98 Zbl [0521.05027](#) MR [0701174](#)

Received 19 September 2022.

Pankaj Kapari

Department of Mathematics, Indian Institute of Science Education and Research Bhopal, Bhopal Bypass Road, Bhauri Bhopal, Madhya Pradesh 462 066, India; pankajkapri02@gmail.com

Kashyap Rajeevsarathy

Department of Mathematics, Indian Institute of Science Education and Research Bhopal, Bhopal Bypass Road, Bhauri Bhopal, Madhya Pradesh 462 066, India; kashyap@iiserb.ac.in

Apeksha Sanghi

Department of Mathematics, Indian Institute of Science Education and Research Bhopal, Bhopal Bypass Road, Bhauri Bhopal, Madhya Pradesh 462 066, India; apekshasanghi93@gmail.com