

Wreath products of groups acting with bounded orbits

Paul-Henry LEEMANN and Grégoire SCHNEEBERGER

Abstract. If \mathbf{S} is a subcategory of metric spaces, we say that a group G has property \mathbf{BS} if any isometric action on an \mathbf{S} -space has bounded orbits. Examples of such subcategories include metric spaces, affine real Hilbert spaces, $\text{CAT}(0)$ cube complexes, connected median graphs, trees or ultra-metric spaces. The corresponding properties \mathbf{BS} are respectively Bergman's property, property \mathbf{FH} (which, for countable groups, is equivalent to the celebrated Kazhdan's property (\mathbf{T})), property \mathbf{FW} (both for $\text{CAT}(0)$ cube complexes and for connected median graphs), property \mathbf{FA} and uncountable cofinality. Historically many of these properties were defined using the existence of fixed points.

Our main result is that for many subcategories \mathbf{S} , the wreath product $G \wr_X H$ has property \mathbf{BS} if and only if both G and H have property \mathbf{BS} and X is finite. On one hand, this encompasses in a general setting previously known results for properties \mathbf{FH} and \mathbf{FW} . On the other hand, this also applies to the Bergman's property. Finally, we also obtain that $G \wr_X H$ has uncountable cofinality if and only if both G and H have uncountable cofinality and H acts on X with finitely many orbits.

Mathematics Subject Classification 2020: 20E22 (primary); 20F65 (secondary).

Keywords: wreath product, action with bounded orbits, fixed-point properties, property (\mathbf{T}) , property \mathbf{FW} , Bergman's property.

1. Introduction

When working with group properties, it is natural to ask if they are stable under “natural” group operations. One such operation, of great use in geometric group theory, is the wreath product, see Section 2 for all the relevant definitions.

An \mathbf{S} -space is a metric space with an “additional structure” and we will say that a group G has *property* \mathbf{BS} if every action by isometries which preserves the structure on an \mathbf{S} -space has bounded orbits. Formally, this means that \mathbf{S} is a subcategory of the category of metric spaces, and that the actions are by \mathbf{S} -automorphisms. We note that having one bounded orbit implies that all the orbits are bounded.

In the context of properties defined by actions with bounded orbits, the first result concerning wreath products, due to Cherix, Martin and Valette and later refined by Neuhauser, concerns Kazhdan's property (T).

Theorem 1.1 ([9, 23]). *Let G and H be two discrete groups with G non-trivial and let X be a set on which H acts. The wreath product $G \wr_X H$ has property (T) if and only if G and H have property (T) and X is finite.*

For countable groups (and more generally for σ -compact locally-compact topological groups), property (T) is equivalent, by the Delorme–Guichardet theorem, to property FH (every action on an affine real Hilbert space has bounded orbits), see [2, Theorem 2.12.4]. Hence, Theorem 1.1 can also be viewed, for countable groups, as a result on property FH.

The corresponding result for property FA (every action on a tree has bounded orbits) and property FR (every action on a real tree has bounded orbits), is a little more convoluted and was obtained a few years later by Cornulier and Kar.

Theorem 1.2 ([13]). *Let G and H be two groups with G non-trivial and let X be a set on which H acts with finitely many orbits and without fixed points. Then $G \wr_X H$ has property FA (respectively property FR) if and only if H has property FA (respectively property FR), G has no quotient isomorphic to \mathbf{Z} and cannot be written as a countable increasing union of proper subgroups.*

Observe that our statement of Theorem 1.2 differs of the original statement of [13]. Indeed, where we ask G to have uncountable cofinality and no quotient isomorphic to \mathbf{Z} , the authors of [13] ask G to have uncountable cofinality and finite abelianization. However, these two sets of conditions are equivalent. One implication is trivial, as finite abelian groups do not project onto \mathbf{Z} . For the other implication, suppose that G has uncountable cofinality but infinite abelianization $G/[G, G]$. The group $G/[G, G]$ being an infinite abelian group, it has a countably infinite quotient A – a classical fact, of which Y. de Cornulier kindly reminded us; see [18, Section 16.11.c] for a proof. The quotient A has uncountable cofinality, see Lemma 3.2, and is therefore an infinite finitely generated abelian group, which hence projects onto \mathbf{Z} .

Finally, we have an analogous of Theorem 1.1 for property FW (every action on a CAT(0) cube complex has bounded orbits).

Theorem 1.3 ([11, 20]). *Let G and H be two groups with G non-trivial and let X be a set on which H acts. Suppose that all three of G , H and $G \wr_X H$ are finitely generated. Then the wreath product $G \wr_X H$ has property FW if and only if G and H have property FW and X is finite.*

It is straightforward to prove that the wreath product $G \wr_X H$ is finitely generated if and only if both G and H are finitely generated and the number of orbits of the action of H on X is finite.

Theorem 1.3 was first proved, using *cardinal definite functions*, for arbitrary groups by Cornulier [11, Propositions 5.B.3 and 5.B.4], but without the implication “if $G \wr_X H$ has property FW, then G has property FW.” The authors then gave an elementary proof of it via *Schreier graphs* for the specific case of finitely generated groups [20]. Y. Stalder has let us know (private communication) that, using *space with walls* instead of Schreier graphs, the arguments of [20] can be adapted to replace the finite generation hypothesis of Theorem 1.3 by the condition that all three of G , H and X are at most countable. Finally, A. Genevois published in [14] a proof of Theorem 1.3 for wreath products of the form $G \wr_H H$, based on his *diadem product of spaces*.

The above results on properties FH, FW and FA were obtained with distinct methods even if the final results have a common flavor. At the same time, all three properties FH, FW and FA can be characterized by the fact that any isometric action on a suitable metric space (respectively, affine real Hilbert space, connected median graph and tree) has bounded orbits; see Definition 2.4. But more group properties can be characterized in terms of actions with bounded orbits. This is, for example, the case of the Bergman’s property (actions on metric spaces), the property FB_r (actions on reflexive real Banach spaces) or of uncountable cofinality (actions on ultrametric spaces).

By adopting the point of view of actions with bounded orbits, we obtain a unified proof of the following result; see also Theorem 3.1 for the general (and more technical) statement.

Theorem A. *Let BS be any one of the following properties: Bergman’s property, property FB_r , property FH or property FW. Let G and H be two groups with G non-trivial and let X be a set on which H acts. Then the wreath product $G \wr_X H$ has property BS if and only if G and H have property BS and X is finite.*

With a little twist, we also obtain a similar result for groups with uncountable cofinality.

Proposition B. *Let G and H be two groups with G non-trivial and let X be a set on which H acts. Then the wreath product $G \wr_X H$ has uncountable cofinality if and only if G and H have uncountable cofinality and H acts on X with finitely many orbits.*

A crucial ingredient of our proofs is that the spaces under consideration admit a natural notion of Cartesian product. In particular, some of our results do not work for trees and property FA, nor do they for real trees and the corresponding property FR. Nevertheless, we are still able to show that if $G \wr_X H$ has property FA, then H acts on X

with finitely many orbits. Combining this with Theorem 1.2 we obtain the following theorem.

Theorem C. *Let G and H be two groups with G non-trivial and X a set on which H acts. Suppose that H acts on X without fixed points. Then $G \wr_X H$ has property FA (respectively, has property FR) if and only if H has property FA (respectively has property FR), H acts on X with finitely many orbits, G has no quotient isomorphic to \mathbf{Z} and cannot be written as a countable increasing union of proper subgroups.*

2. Definitions and examples

This section contains all the definitions, as well as some useful preliminary facts and some examples.

2.1. Wreath products. Let X be a set and G a group. We view $\bigoplus_X G$ as the set of functions from X to G with finite support:

$$\bigoplus_X G = \{\varphi: X \rightarrow G \mid \varphi(x) = 1 \text{ for all but finitely many } x\}.$$

This is naturally a group, where multiplication is taken componentwise.

If H is a group acting on X , then it naturally acts on $\bigoplus_X G$ by

$$(h.\varphi)(x) = \varphi(h^{-1}.x).$$

This leads to the following standard definition.

Definition 2.1. Let G and H be groups and X be a set on which H acts. The (restricted¹) wreath product $G \wr_X H$ is the group $(\bigoplus_X G) \rtimes H$.

A prominent particular case of wreath products is of the form $G \wr_H H$, where H acts on itself by left multiplication. They are sometimes called *standard wreath products* or simply *wreath products*, while general $G \wr_X H$ are sometimes called *permutational wreath products*. Best known example of wreath product is the so called *lamplighter group* $(\mathbf{Z}/2\mathbf{Z}) \wr_{\mathbf{Z}} \mathbf{Z}$. Other (trivial) examples of wreath products are direct products $G \times H$ which correspond to wreath products over a singleton $G \wr_{\{*\}} H$.

2.2. Classical actions with bounded orbits. In this subsection we discuss some classical group properties, which are defined by actions with bounded orbits on various metric spaces.

¹There exists an unrestricted version of this product where the direct sum is replaced by a direct product.

Median graphs. For u and v two vertices of a connected² graph \mathcal{G} , we define the total interval $[u, v]$ as the set of vertices that lie on some shortest path between u and v . A connected graph \mathcal{G} is *median* if for any three vertices u, v, w , the intersection

$$[u, v] \cap [v, w] \cap [u, w]$$

consists of a unique vertex, denoted $m(u, v, w)$. A graph is *median* if each of its connected components is median. For more on median graphs and spaces, see [1, 6, 19]. If X and Y are both (connected) median graphs, then their Cartesian product is also a (connected) median graph. The class of median graphs was introduced by Nebeský in 1971 [22] and Gerasimov [15, 16], Roller [27] and Chepoi [8] realized independently that this class coincides with the class of 1-skeleta of CAT(0) cube complexes. Trees are the simplest examples of connected median graphs, while the ensuing classical example show that any power set can be endowed with a median graph structure.

Example 2.2. Let X be a set and let $\mathcal{P}(X) = 2^X$ be the set of all subsets of X . Define a graph structure on $\mathcal{P}(X)$ by putting an edge between E and F if and only if

$$\#(E \Delta F) = 1,$$

where Δ is the symmetric difference. Therefore, the distance between two subsets E and F is $\#(E \Delta F)$ and the connected component of E is the set of all subsets F with $E \Delta F$ finite. For E and F in the same connected component, $[E, F]$ consists of all subsets of X that both contain $E \cap F$ and are contained in $E \cup F$. In particular, $\mathcal{P}(X)$ is a median graph, with $m(D, E, F)$ being the set of all elements belonging to at least two of D, E and F . In other words,

$$m(D, E, F) = (D \cap E) \cup (D \cap F) \cup (E \cap F).$$

These graphs are useful due to the following fact. Any action of a group G on a set X naturally extends to an action of G on $\mathcal{P}(X)$ by graph homomorphisms:

$$g.Y := \{g.y \mid y \in Y\}$$

for $Y \subset X$. Note that the action of G on $\mathcal{P}(X)$ may exchange the connected components. In fact, the connected component of $E \in \mathcal{P}(X)$ is stabilized by G if and only if E is *commensurated* by G , that is if for every $g \in G$ the set $E \Delta g.E$ is finite.

²We will always assume that our connected graphs are non-empty. This is coherent with the definition that a connected graph is a graph with exactly one connected component.

Uncountable cofinality. Recall that a metric space (X, d) is *ultrametric* if d satisfies the strong triangular inequality:

$$d(x, y) \leq \max\{d(x, z), d(z, y)\}$$

for any x, y and z in X . A group G has *uncountable cofinality* if every action on ultrametric spaces has bounded orbits. The following characterization of groups of countable cofinality can be extracted from [10] and we include a proof only for the sake of completeness. It implies in particular that a countable group has uncountable cofinality if and only if it is finitely generated.

Lemma 2.3. *Let G be a group. Then the following are equivalent:*

- (1) G can be written as a countable increasing union of proper subgroups;
- (2) G does not have uncountable cofinality, i.e., there exists an ultrametric space X on which G acts with an unbounded orbit;
- (3) There exists a G -invariant (for the action by left multiplication) ultrametric d on G such that $G \curvearrowright G$ has an unbounded orbit.

Proof. It is clear that the third item implies the second.

Let (X, d) be an ultrametric space on which G acts with an unbounded orbit $G.x_0$. For any $n \in \mathbf{N}$, let H_n be the subset of G defined by

$$H_n := \{g \in G \mid d(x_0, g.x_0) \leq n\}.$$

Then G is the union of the (countably many) H_n , which are subgroups of G . Indeed, H_n is trivially closed under taking the inverse, and is also closed under taking products as we have

$$\begin{aligned} d(x_0, gh.x_0) &\leq \max\{d(x_0, g.x_0), d(g.x_0, gh.x_0)\} \\ &= \max\{d(x_0, g.x_0), d(x_0, h.x_0)\}. \end{aligned}$$

As $G.x_0$ is unbounded, they are proper subgroups. Since they are proper subgroups and $H_n \leq H_{n+1}$, we can extract an increasing subsequence $(H_{r_n})_n$ that still satisfies $G = \bigcup_n H_{r_n}$.

Finally, suppose that $G = \bigcup_{n \in \mathbf{N}} H_n$, where the H_n form an increasing sequence of proper subgroups. It is always possible to suppose that $H_0 = \{1\}$. Define d on G by

$$d(g, h) := \min\{n \mid g^{-1}h \in H_n\}.$$

One easily verifies that d is a G -invariant ultrametric. Moreover, the orbit of 1 contains all of G and is hence unbounded. ■

Some classical group properties. We now discuss the bounded orbits properties for actions on various classes of metric spaces.

Definition 2.4. Let G be a group. It is said to have the following properties:

- *Bergman's property* if any action by isometries on a metric space has bounded orbits;
- *Property FB_r* if any action by affine isometries on a reflexive real Banach space has bounded orbits;
- *Property FH* if any action by affine isometries on a real Hilbert space has bounded orbits;
- *Property FW* if any action by graph isomorphisms on a connected median graph has bounded orbits;
- *Property FR* if any action by isometries on a real tree has bounded orbits;
- *Property FA* if any action by graph isomorphisms on a tree has bounded orbits;
- *Uncountable cofinality* if any action by isometries on an ultrametric space has bounded orbits.

In the above, we insisted on the fact that actions are supposed to preserve the structure of the metric space under consideration, or in other words to be by automorphisms of the considered category. This is sometimes automatic, as for example any isometry of a real Banach or Hilbert space is affine by the Mazur–Ulam theorem. However, for cube complexes, for example, this is not the case; a 2-regular tree has automorphism group $\mathbf{Z} \times (\mathbf{Z}/2\mathbf{Z})$, while its isometry group is $\mathbf{R} \times (\mathbf{Z}/2\mathbf{Z})$. In other words, an isometry of a cube complex is not necessarily a cube complex isomorphism; see also Example 2.8.

In the following, we will often do a slight abuse of notation and simply speak of a *group action* on a space X , without always specifying by which kinds of maps the group acts, which should always be clear from the context.

The names FB_r , FH, FW, FR and FA come from the fact that these properties admit a description in terms of (and were first studied in the context of) the existence of a fixed point for actions on reflexive Banach spaces, on Hilbert spaces, on spaces with walls (or equivalently on CAT(0) cube complexes, see [7, 25]), on real trees and on trees (*arbres*, in French). However this is equivalent with the bounded orbit property, see Proposition 2.16 and the discussion below it. Observe that spaces with walls admit a natural pseudo-metric on them, which is not necessarily a metric.


The Bergman's property can also be characterized via length function, see for example the beginning of [29].

For a survey on property FB_r , see [26] and the references therein.

For countable groups (and more generally for σ -compact locally compact groups), property FH is equivalent, by the Delorme–Guichardet theorem, to the celebrated Kazhdan’s property (T), but this is not true in general. Indeed by [3] symmetric groups over infinite sets are uncountable discrete group with Bergman’s property which, as we will see just below, implies property FH. Such groups cannot have property (T) as, for discrete groups, it implies finite generation.

A classical result of the Bass–Serre theory of groups acting on trees [28], is that a group G has property FA if and only if it satisfies the following three conditions: G has uncountable cofinality, G has no quotient isomorphic to \mathbf{Z} and G is not a non-trivial amalgam. In view of this characterization, Theorem 1.2 says that property FA almost behaves well under wreath products.

Proposition 2.5. *There are the following implications between the properties of Definition 2.4:*



Moreover, except maybe for the implication [Bergman’s property \implies FB_r], all implications are strict.

Proof. The implications [Bergman’s property \implies $\text{FB}_r \implies$ FH] and [FW \implies FA \iff FR] trivially follow from the fact that Hilbert spaces are reflexive Banach spaces, which are themselves metric spaces and that trees are both real trees and connected median graphs.

The implication [FH \implies FW] follows from the fact that a group G has property FW if and only if any affine action on a real Hilbert space which preserves integral points has bounded orbits [11, Proposition 7.I.3].

The implication [FH \implies FR] follows from the fact that real trees are median metric spaces, and that such spaces can be embedded into L^1 -spaces (see, for instance, [30, Theorem V.2.4]).

Finally, the implication [FA \implies uncountable cofinality] is due to Serre [28]: If G is an increasing union of subgroups G_i , then $\bigsqcup G/G_i$ admits a tree structure by joining any $gG_i \in G/G_i$ to $gG_{i+1} \in G/G_{i+1}$. The action of G by multiplication on $\bigsqcup G/G_i$ is by graph isomorphisms and with unbounded orbits.

We now present some examples demonstrating the strictness of the implications. Countable groups with property FB_r are finite by [5], while infinite finitely generated

groups with property (T), e.g., $SL_3(\mathbf{Z})$, have property FH. The group $SL_2(\mathbf{Z}[\sqrt{2}])$ has property FW but not FH; see [12]. If G is a non-trivial finite group and H is an infinite group with property FA (respectively, property FR), then $G \wr_H H$ has property FA (respectively property FR) by Theorem C, but does not have property FW by Theorem A. The group \mathbf{Z} has uncountable cofinality, while it acts by translations and with unbounded orbits on the infinite 2-regular tree. Finally, Minasyan constructed examples of groups with FA but without FR in [21]. ■

The reader familiar with triangle groups,

$$\Delta(l, m, n) = \langle a, b, c \mid a^2 = b^2 = c^2 = (ab)^l = (bc)^m = (ca)^n = 1 \rangle$$

with $l, m, n \in \{1, 2, \dots\} \cup \{\infty\}$, will be happy to observe that they provide explicit examples of groups with property FA but not property FW. Indeed, if l, m and n are all three integers, then $\Delta(l, m, n)$ has property FA by Serre [28, Section 6.5, Corollaire 2]. And if

$$\kappa(l, m, n) := \frac{1}{l} + \frac{1}{m} + \frac{1}{n} \leq 1,$$

then $\Delta(l, m, n)$ is the infinite symmetric group of a tiling of the Euclidean plane (if $\kappa(l, m, n) = 1$) or of the hyperbolic plane (if $\kappa(l, m, n) < 1$), and hence acts on a space with walls without fixed point, which implies that it does not have property FW.

In view of Proposition 2.5, two questions remain open:

is the implication [Bergman’s property $\implies FB_r$] strict, and does property FW imply property FR?

It is possible to consider relative versions of the properties appearing in Definition 2.4. Let \mathbf{S} be any classes of metric spaces considered in Definition 2.4 and let BS be the corresponding group property. If G is a group and H a subgroup of G , we say that the pair (G, H) has *relative property BS* if for every G -action on an \mathbf{S} -space, the H orbits are bounded. A group G has property BS if and only if for every subgroup H the pair (G, H) has relative property BS, and if and only if for every overgroup L the pair (L, G) has relative property BS.

2.3. Groups acting with bounded orbits on S-spaces. It is possible to define other properties in the spirit of Definition 2.4 for any “subclass of metric spaces”, or more precisely for any subcategory of pseudo-metric spaces. A reader not familiar with category theory and interested only in one specific subclass of metric spaces may forget all these general considerations and only verify that the arguments of Section 3 apply for their favorite subclass of metric spaces.

Definition 2.6. A *pseudo-metric space* is a set X with a map $d: X \times X \rightarrow \mathbf{R}_{\geq 0}$, called a *pseudo-distance*, such that

- (1) $d(x, x) = 0$ for all $x \in X$;
- (2) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (3) $d(x, z) \leq d(x, y) + d(y, z)$.

If moreover $d(x, y) \neq 0$ for $x \neq y$, the map d is a *distance* and (X, d) is a *metric space*. On the other hand, an *ultra-pseudo-metric space* is a pseudo-metric space (X, d) such that d satisfies the strong triangular inequality:

$$d(x, z) \leq \max\{d(x, y), d(y, z)\}.$$

A *morphism* (or *short map*) between two pseudo-metric spaces (X_1, d_1) and (X_2, d_2) is a distance non-increasing map $f: X_1 \rightarrow X_2$, that is

$$d_2(f(x), f(y)) \leq d_1(x, y)$$

for any x and y in X_1 . If f is bijective and distance preserving, then it is an *isomorphism* (or *isometry*). Pseudo-metric spaces with short maps form a category **PMet**, of which the category of metric spaces (with short maps) **Met** is a full subcategory.

If (X, d) is a pseudo-metric space, we have a natural notion of the *diameter* of a subset $Y \subset X$ with values in $[0, \infty]$, defined by

$$\text{diam}(Y) := \sup\{d(x, y) \mid x, y \in Y\},$$

and we say that Y is *bounded* if it has finite diameter.

Remind that a *subcategory of PMet*, is a category **S** whose objects are pseudo-metric spaces, and whose morphisms are short maps. The subcategory **S** is *full* if given two **S**-objects X and Y , any short map from X to Y is a **S**-morphism. A *G-action on an S-space X* is simply a group homomorphism $\alpha: G \rightarrow \text{Aut}_{\mathbf{S}}(X)$.

In practice, a lot of examples of (full) subcategories of **PMet** are already subcategories of **Met**. Obvious examples of full subcategories of **Met** include metric spaces and ultrametric spaces (with short maps). Affine real Hilbert and Banach spaces and more generally normed vector spaces (with affine maps) are also full subcategories of **Met** if we restrict ourselves to morphisms that do not increase the norm (that is, such that $\|f(x)\| \leq \|x\|$). In particular, for us isomorphisms of Hilbert and Banach spaces will always be affine isometries.

For connected graphs (and hence for its full subcategories of connected median graphs and of trees), one looks at the category **Graph** where objects are connected simple graphs $G = (V, E)$ and where a morphism $f: (V, E) \rightarrow (V', E')$ is a function between the vertex sets such that if (x, y) is an edge, then either $f(x) = f(y)$

or $(f(x), f(y))$ is an edge. There are two natural ways to see **Graph** as a subcategory of **Met**. The first one consists to look at the vertex set V endowed with the shortest-path metric: $d(v, w)$ is the minimal number of edges on a path between v and w . The second one, consists at looking at the so called geometric realization of (V, E) , where each edge is seen as an isometric copy of the segment $[0, 1]$. Similarly to what happens for cube complexes (see the discussion after Definition 2.4), the geometric realization of a graph gives an embedding **Graph** \hookrightarrow **Met** which is not full. Nevertheless, for our purpose, the particular choice of one of the above two embeddings **Graph** \hookrightarrow **Met** will make no difference.

We can now formally define the group property **BS**.

Definition 2.7. Let **S** be a subcategory of **PMet**. A group G has *property BS* if every G -action by **S**-automorphisms on an **S**-space has all its orbits bounded. A pair (G, H) of a group and a subgroup has *relative property BS* if for every G -action by **S**-automorphisms on an **S**-space, the H -orbits are bounded.

Observe that a group G has property **BS** if and only if any G -action on an **S**-space has at least one bounded orbit.

All the properties of Definition 2.4 are of the form **BS**. Another example of property of the form **BS** can be found in [17, Definition 6.22]: a group has property $(\text{FHyp}_{\mathbb{C}})$ if any action on a real or complex hyperbolic space of finite dimension has bounded orbits. This property is implied by property **FH**, but does not imply property **FA** [17, Corollary 6.23 and Example 6.24]. One can also want to look at the category of all Banach spaces (the corresponding property **BB** hence stands between the Bergman's property and property FB_r), or the category of L^p -spaces for p fixed [4] (if $p \notin \{1, \infty\}$, then BL^p is implied by FB_r).

Another interesting example of a property of the form **BS** is the fact to have no quotient isomorphic to \mathbf{Z} ; see Example 2.8. The main interest for us of this example is that property **FA** is the conjunction of three properties, two of them (uncountable cofinality and having no quotient isomorphic to \mathbf{Z}) still being of the form **BS**.

Example 2.8. Let Z be the 2-regular tree, or in other words the Cayley graph of \mathbf{Z} for the standard generating set. Then $\text{Aut}_{\text{Graph}}(Z) = \mathbf{Z} \times (\mathbf{Z}/2\mathbf{Z})$ is the infinite dihedral group and its subgroup of orientation preserving isomorphisms is isomorphic to \mathbf{Z} . Let **S** be the category with one object Z and with morphisms the orientation preserving isomorphisms. Hence, we obtain that a group G has no quotient isomorphic to \mathbf{Z} if and only if every G -action on **S**-space has bounded orbits. Let us denote by **BZ** this property.

Since Z is a tree, property **FA** implies property **BZ**. This implication is strict as demonstrated by **Q**. In fact, the counterexample **Q** shows that **BZ** does not imply

uncountable cofinality, while **Z** demonstrates that uncountable cofinality does not imply **BZ**.

An example of an uninteresting property **BS** is given by taking **S** to be the category of metric spaces of finite diameter (together with short maps), in which case any group has **BS**. On the other hand, if **S** is the category of extended pseudo-metric spaces (d takes values in $\mathbf{R}_+ \cup \{\infty\}$), only the trivial group has **BS**. Indeed, one can put the extended metric $d(x, y) = \infty$ if $x \neq y$ on G and then the action by left multiplication of G on (G, d) is transitive and with an unbounded orbit as soon as G is non-trivial.

The category **PMet** has the advantage (over **Met**) of behaving more nicely with respect to categorical constructions and quotients. However, we have the following lemma.

Lemma 2.9. *A group G has Bergman's property (respectively uncountable cofinality) if and only if any isometric G -action on a pseudo-metric (respectively ultra-pseudo-metric) space has bounded orbits.*

Proof. One direction is trivial. For the other direction, let (X, d) be a pseudo-metric space on which G acts by isometries. Let $X' := X / \sim$ be the quotient of X for the relation $x \sim y$ if $d(x, y) = 0$ and let d' be the quotient of d . Then (X', d') is a metric space, the action of G passes to the quotient and $G.x$ is d -bounded if and only if $G.[x]$ is d' -bounded. Finally, if d satisfies the strong triangular inequality, then so does d' . ■

On the other hand, the following result is perhaps more surprising.

Lemma 2.10. *A group G has Bergman's property if and only if any G -action by graph automorphisms on a connected graph has bounded orbits.*

Proof. The left-to-right implication is clear.

For the other direction, let X be a metric space. Let $G(X)$ denotes the graph obtained from a vertex-set X by applying the following process: for any two $x, y \in X$ add a path of length $\lfloor d(x, y) \rfloor + 1$ between x and y . $G(X)$ is connected and the obvious inclusion $\iota: X \rightarrow G(X)$ is a quasi-isometric embedding. Moreover, the construction is canonical, so every group action on X extends to a group action on $G(X)$, making ι equivariant. So if a group satisfying the bounded orbit property on connected graphs acts on a metric space X , then its induced action on $G(X)$ has bounded orbits, which implies that its orbits in X are bounded. ■

Observe that an alternative proof of the above lemma can be easily deduced from the following characterization of Bergman's property due to Cornulier [10]: A group G has Bergman's property if and only if it has uncountable cofinality and for every generating set T of G , the Cayley graph $\text{Cay}(G; T)$ is bounded. Details are left to the reader.

While we will be able to obtain some results for a general subcategory of **PMet**, we will sometimes need to restrict ourselves to subcategories satisfying good properties. In particular, we will use three axioms: one on the existence of non-trivial G -action, one on the existence of finite Cartesian powers and one on infinite Cartesian powers. Our Cartesian powers will need to be in some sense compatible with the bornology, but the conditions for finite and infinite powers will not be the same. A summary of which axioms are satisfied by the above mentioned subcategories of **PMet** can be found in Table 1.

Definition 2.11 (Axiom (A1)). A subcategory **S** of **PMet** has *non-trivial group actions* if for every non-trivial group G there exists an **S**-space X and an action $G \curvearrowright X$ by **S**-automorphisms moving at least one point.

Examples of categories **S** with non-trivial group actions include (ultra-) metric spaces and metric spaces of finite diameter (with the action by multiplication of G on itself, endowed with the discrete metric), (reflexive) Banach spaces and Hilbert spaces (with $X = l^2(G)$ and the permutation action of G), L^p spaces (G acting by permutation on $\ell^p(G)$) and finally connected median graphs and (real) trees (X has one central vertex to which we attach an edge for every $g \in G$ and the action of G is by left multiplication). On the opposite side, both Z from Example 2.8 and real and complex finite-dimensional hyperbolic spaces do not have non-trivial group actions. Indeed, a group acts non-trivially on Z if and only if it projects onto \mathbf{Z} . For hyperbolic spaces, a group G acts on a hyperbolic space of dimension n if and only if it projects onto a subgroup of $SO(n, 1)$ or $SU(n, 1)$. In particular, if the action is non-trivial, then G projects onto a non-trivial subgroup of $GL_n(\mathbf{C})$, whose all finitely generated subgroups are residually finite. We conclude that a finitely generated infinite simple group G does not admit a non-trivial action on a real or complex hyperbolic space of finite dimension.

Before introducing the axioms about Cartesian powers, let us recall the definition and some properties of the product distances d_p .

Definition 2.12. For a real $p \geq 1$ and a collection of non-empty metric spaces, which we denote by $(X_i, d_i)_{i \in I}$, we have the maps

$$d_p: \prod_{i \in I} X_i \times \prod_{i \in I} X_i \rightarrow \mathbf{R},$$

$$(f, g) \mapsto \left(\sum_{i \in I} d_i(f(i), g(i))^p \right)^{\frac{1}{p}}$$

and

$$d_\infty: \prod_{i \in I} X_i \times \prod_{i \in I} X_i \rightarrow \mathbf{R},$$

$$(f, g) \mapsto \sup_{i \in I} d_i(f(i), g(i)),$$

with the convention that a sum with uncountably non-zero summands is infinite.

If I is finite, then d_p is a distance on $\prod_{i \in I} X_i$. Moreover, it is compatible with the bornology in the sense that if $E \subseteq X$ is unbounded, then the diagonal $\text{diag}(E) \subset X^n$ is also unbounded. The following definition generalizes this comportment to other distances.

Definition 2.13 (Axiom (A2)). A subcategory \mathbf{S} of \mathbf{PMet} satisfies *axiom (A2)* if for any \mathbf{S} -space X and any integer n , there exists an \mathbf{S} -object, called a *n th Cartesian power of X* and written X^n , such that:

- (1) as a set, X^n is the n th Cartesian power of X ;
- (2) the canonical image of $\text{Aut}_{\mathbf{S}}(X)^n \rtimes \text{Sym}(n)$ in $\text{Bij}(X^n)$ lies in $\text{Aut}_{\mathbf{S}}(X^n)$;
- (3) if $E \subset X$ is unbounded, then the diagonal $\text{diag}(E) \subset X^n$ is unbounded.

A good heuristic is that your favorite subcategory of \mathbf{PMet} would satisfy axiom (A2) in the above sense if and only if it already has a classical operation, which is called *Cartesian product*. An \mathbf{S} -object satisfying the first two properties of Definition 2.13 will be called a *finite Cartesian power*.

For metric spaces (of finite diameter), the categorical product (corresponding to the metric $d_\infty = \max\{d_X, d_Y\}$) works fine, but any product metric of the form

$$d_p = (d_X^p + d_Y^p)^{\frac{1}{p}}$$

for $p \in [1, \infty]$ works as well. For ultra-metric spaces, the categorical product d_∞ works fine. For L^p spaces (and hence for Hilbert spaces), we take the usual Cartesian product (which is also the categorical product), which corresponds to the metric d_p . For (reflexive) Banach spaces, any product metric of the form d_p works. For connected graphs, the usual Cartesian product, which corresponds to $d_1 = d_X + d_Y$ works well, but one can also take the strong product (which is the categorical product in \mathbf{Graph}), that is the distance d_∞ . For connected median graphs, only the usual Cartesian product with d_1 works.³ On the other hand, (real) trees, Z from Example 2.8 and hyperbolic spaces do not have finite Cartesian powers and hence cannot satisfy axiom (A2).

³The category of median graphs does not have categorical products.

As the above examples illustrate, there can be multiple non-isomorphic spaces playing the role of X^n . As our results will not depend on a particular choice of a Cartesian power, we will sometimes make a slight abuse of language and speak of *the* Cartesian power X^n .

The situation for infinite products is more complicated. Indeed, if I is infinite then the map d_p is not necessary a distance on $\prod_{i \in I} X_i$ as it can take infinite values. The solution consists of looking at the subset of $\prod_{i \in I} X_i$ on which d_p takes finite values. Formally we first need to chose a base-point x_i in X_i for each $i \in I$, which gives us an element $f_0 \in \prod_{i \in I} X_i$ defined by $f_0(i) = x_i$. We can then define

$$\bigoplus_{i \in I} X_i := \left\{ f \in \prod_{i \in I} X_i \mid f(i) = f_0(i) \text{ for all but finitely many } i \right\},$$

$$\bigoplus_{i \in I}^p X_i := \left\{ f \in \prod_{i \in I} X_i \mid \sum_i d_i(f(i), f_0(i))^p < \infty \right\}.$$

A priori, the above definitions depend on the choice of the x_i . However, since our results will not depend on a particular choice of base-points, we will omit to specify it. Moreover, if $\text{Aut}_{\mathbf{S}}(X)$ acts transitively on X , then $\bigoplus_{i \in I}^p X$ will not depend on the choice of $x \in X$. If all the X_i are equal to \mathbf{R} with $x_i = 0$, then $\bigoplus_{i \in I}^p X_i$ is the classical Banach space $\ell^p(I)$ while $\bigoplus_{i \in I} X_i$ is the (non-complete) sequence space $c_{00}(I)$.

It is straightforward to verify that

$$\bigoplus_{i \in I} X_i \subseteq \bigoplus_{i \in I}^p X_i \subseteq \prod_{i \in I} X_i,$$

and that the first inclusion is an equality if the X_i are uniformly discrete with a uniform lower bound on their packing radius. Moreover, the restriction of the map d_p to $\bigoplus_{i \in I}^p X_i$ is a distance. While $(\bigoplus_{i \in I} X_i, d_p)$ is a metric space, it is in general not complete even when the X_i are complete, which is why we needed to define $\bigoplus_{i \in I}^p X_i$; this will be important for Banach and Hilbert spaces.

One common feature of the product distances d_p for $p \neq \infty$ is that, in some rough way, they are able to detect the number of coordinates on which f differs from f_0 . Our last axiom will generalize this comportment.

Definition 2.14 (Axiom (A3)). A subcategory \mathbf{S} of \mathbf{PMet} satisfies *axiom (A3)* if for any \mathbf{S} -space X , any element x_0 of X and any infinite set I , there exists an \mathbf{S} -object, called the *I*th Cartesian power of X and written $\bigoplus_I^{\mathbf{S}} X$, such that:

- (1) as sets we have the inclusions $\bigoplus_I X \subseteq \bigoplus_I^{\mathbf{S}} X \subseteq \prod_I X$;
- (2) the canonical image of $\text{Aut}_{\mathbf{S}}(X) \wr_I \text{Sym}(I)$ in $\text{Bij}(\prod_I X)$ lies in $\text{Aut}_{\mathbf{S}}(\bigoplus_I^{\mathbf{S}} X)$;

(3) for any y in X with $d(y, x_0) > 0$, the following set has infinite diameter:

$$\left\{ f \in \bigoplus_I X \mid f(i) = y \text{ for finitely many } i \text{ and otherwise } f(i) = x_0 \right\}.$$

An object satisfying the first two items of Definition 2.14 will be called an infinite Cartesian power.

In practice, the above definition is often easy to verify. Indeed, in most cases when \mathbf{S} has finite Cartesian powers it is for some product metric of the form d_p . Then the metric space $(\bigoplus_I^p X, d_p)$ will usually be an infinite Cartesian power in \mathbf{S} and, if $p \neq \infty$, it will satisfy (A3). In particular, (A3) holds in the following categories: metric spaces, (reflexive) Banach spaces (with d_p for $1 < p < \infty$), Hilbert spaces and L^p spaces if $p \neq \infty$, connected (median) graphs. On the other hand, (real) trees, hyperbolic spaces and Z from Example 2.8 do not have a sensible notion of infinite Cartesian powers. Finally, while ultra-metric spaces, L^∞ spaces and spaces of finite diameter have infinite Cartesian powers (for d_∞), axiom (A3) does not hold as the diameter of the set appearing in Definition 2.14 is $d(y, x_0)$.

Finally, we introduce one last definition

Definition 2.15. A subcategory \mathbf{S} of \mathbf{PMet} has *bornological Cartesian powers* if it satisfies both axiom (A2) and (A3).

2.4. Variations and generalizations. This subsection is devoted to variations and generalizations of property BS. It is intended as a note for the interested reader, and can be skipped without any harm.

Groups acting with fixed point on S-spaces. Some of the properties that are of interest for us have been historically defined via the existence of a fixed point for some action. More generally, we say that a group G has *property FS* if any G -action on an \mathbf{S} -space has a fixed point.

Since our actions are by isometries, property FS implies property BS. The other implication holds as soon as we have a suitable notion of the center of a (non-empty) bounded subset X . For a large class of metric spaces, this is provided by the following result of Bruhat and Tits:

Proposition 2.16 ([17, Chapter 3.b]). *Let (X, d) be a complete metric space such that the following two conditions are satisfied:*

(1) *for all x and y in X , there exists a unique $m \in X$ (the middle of $[x, y]$) such that*

$$d(x, m) = d(y, m) = \frac{1}{2}d(x, y);$$

(2) for all x, y and z in X , if m is the middle of $[y, z]$ we have the median's inequality

$$2d(x, m)^2 + \frac{1}{2}d(y, z)^2 \leq d(x, y)^2 + d(x, z)^2.$$

Then if G is a group acting by isometries on X with a bounded orbit, it has a fixed point.

Examples of complete metric spaces satisfying Proposition 2.16 include among others: Hilbert spaces, Bruhat-Tits Buildings, Hadamard spaces (i.e., complete CAT(0) spaces and in particular CAT(0) cube complexes which are either finite-dimensional or locally finite), trees and \mathbf{R} -trees; with the caveat that for trees and \mathbf{R} -trees, the fixed point is either a vertex or the middle of an edge. See [17, Chapter 3.b] and the references therein for more on this subject. On the other hand, [2, Lemma 2.2.7] gives a simple proof of the existence of a center for bounded subsets of Hilbert spaces, and more generally of reflexive Banach spaces, but this also directly follows from the Ryll-Nardzewski fixed-point theorem. Finally, properties FS and BS are equivalent for the category of separable uniformly convex Banach spaces by the existence of the Chebyshev center of a (non-empty) bounded set.

For action on metric spaces or on connected median graphs, FS is strictly stronger than BS. Indeed, this trivially follows from the action by rotation of $\mathbf{Z}/4\mathbf{Z}$ on the square graph. However, by [16, 27] if a group G acts on a connected median graph with a bounded orbit, then it has a finite orbit. For actions on an ultra-metric spaces FS is strictly stronger than BS since the finite group $\mathbf{Z}/2\mathbf{Z}$ acts without fixed point on the Cantor space

$$X \subset [0, 1] \quad \text{by } x \mapsto 1 - x.$$

Property FS is also strictly stronger than BS for \mathbf{S} the category of all Banach spaces. Indeed, by [24] any infinite discrete group admits an action without fixed point on some Banach space and hence does not have property FS, while there exists infinite groups with the Bergmann's property which implies property BS.

Actions with uniformly bounded orbits. One might wonder what happens if in Definition 2.4 we replace the requirement of having bounded orbits by having uniformly bounded orbits. It turns out that this is rather uninteresting, as a group G is trivial if and only if any G -action on a metric space (respectively on an Hilbert space, on a connected median graph, on a tree or on an ultrametric space) has uniformly bounded orbits. Indeed, if G is non-trivial, then, for the action of G on the Hilbert space $\ell^2(G)$ the orbit of $n \cdot \delta_g$ has diameter $n\sqrt{2}$. For a tree (and hence also for a connected median graph), one may look at the tree T obtained by taking a root r on which we glue an infinite ray for each element of G . Then G naturally acts on T by permuting the rays.

The orbits for this action are the

$$\mathcal{L}_n = \{v \mid d(v, r) = n\},$$

which have diameter $2n$. Finally, it is possible to put an ultradistance on the vertices of T by

$$d_\infty(x, y) := \max\{d(x, r), d(y, r)\}$$

if $x \neq y$. Then the orbits are still the \mathcal{L}_n , but this time with diameter n .

Topological groups. One can wonder what happens for topological groups. While, the wreath product of topological groups is not in general a topological group, this is the case if G is discrete and X is a discrete set endowed with a continuous H -action. In this particular context, Theorem 3.1, as well as its proof, remains true. The details are left to the interested reader.

Categorical generalizations. In the above, we defined property **BS** for \mathbf{S} a subcategory of \mathbf{PMet} . It is possible to generalize this definition to more general categories. We are not aware of any example of the existence of a group property arising in this general context that is not equivalent to a property **BS** in the sense of Definition 2.7, but still mention it for the curious reader.

On one hand, we can replace \mathbf{PMet} with a more general category. For example, one can look at the category \mathbf{M} of sets X endowed with a map $d: X \times X \rightarrow \mathbf{R}_{\geq 0}$ satisfying the triangle inequality. That is, d is a pseudo-distance, except that it is not necessary symmetric and $d(x, x)$ may be greater than 0. All the statements and the proofs remain true for \mathbf{S} a subcategory of \mathbf{M} .

On the other hand, we can define property **BS** for any category \mathbf{S} over \mathbf{PMet} , that is for any category \mathbf{S} endowed with a faithful functor $F: \mathbf{S} \rightarrow \mathbf{PMet}$. Such a couple $(\mathbf{S}, F: \mathbf{S} \rightarrow \mathbf{PMet})$ is sometimes called a *structure over \mathbf{PMet}* , and F is said to be *forgetful*. In this context, we need to be careful to define Cartesian powers (Definitions 2.13 and 2.14) using F , but apart for that all the statements and all the proofs remain unchanged. An example of such an \mathbf{S} that cannot be expressed as a subcategory of \mathbf{PMet} is the category of edge-labeled graphs, where the morphisms are graph morphisms that induce a permutation on the set of labels. However, in this case the property **BS** is equivalent to the Bergman's property.

One can also combine the above two examples and look at couples $(\mathbf{S}, F: \mathbf{S} \rightarrow \mathbf{M})$, with F faithful.

Finally, in view of Definitions 2.7, 2.13 and 2.14, the reader might ask why we are working in \mathbf{PMet} or \mathbf{M} instead of \mathbf{Born} , the category of bornological spaces together with bounded maps. The reason behind this is the forthcoming Lemma 3.3 and its corollaries, which fail for general bornological spaces. In fact, all the statements and

Category \mathbf{S}	Corresponding group property	Satisfies axiom		
		(A1)	(A2)	(A3)
Metric spaces	Bergmann's property	✓	✓	✓
Banach spaces	BB	✓	✓	✓
Reflexive Banach spaces	FB_r	✓	✓	✓
L^p spaces (p fixed)	BL^p	✓	✓	iff $p \neq \infty$
Hilbert spaces	FH	✓	✓	✓
\mathbf{R} and \mathbf{C} hyperbolic spaces	$\text{FHyp}_{\mathbf{C}}$	✗	✗	✗
Median graphs	FW	✓	✓	✓
Real trees	FR	✓	✗	✗
Trees	FA	✓	✗	✗
Ultrametric spaces	uncountable cofinality	✓	✓	✗
$Z = 2$ -regular tree with Isom^+	BZ	✗	✗	✗
Spaces of finite diameter	hold for all groups	✓	✓	✗

TABLE 1
 Axioms for category \mathbf{S} . (A1) = non-trivial group actions (Definition 2.11), (A2) = Definition 2.13, (A3) = Definition 2.14, bornological Cartesian powers = (A2) + (A3).

the proofs remain true for a general $(\mathbf{S}, F: \mathbf{S} \rightarrow \mathbf{Born})$ as soon as \mathbf{S} satisfies Lemma 3.3. Here is an example of such an \mathbf{S} which does not appear as a category over \mathbf{M} . Take κ to be an infinite cardinal and let \mathbf{S}_κ be the subcategory of \mathbf{Born} where a subset E of a \mathbf{S}_κ -space is bounded if and only if $|E| < \kappa$. A group G has property BS_κ if and only if $|G| < \kappa$.

3. Proofs of the main results

Throughout this section, \mathbf{S} will denote a subcategory of \mathbf{PMet} and BS the group property *every action by \mathbf{S} -automorphisms on an \mathbf{S} -space has bounded orbits*. In Section 2.3, we defined three axioms that \mathbf{S} might satisfy. Axiom (A1) simply states that a non-trivial group acts non-trivially on some \mathbf{S} -space. Axioms (A2) and (A3) guarantee the existence of finite and infinite Cartesian powers, which should be compatible in some sense with the bornology. Finally, \mathbf{S} has bornological Cartesian powers if it satisfy both axioms (A2) and (A3). In Table 1 we present a short reminder on whenever these axioms are satisfied for some subcategories of \mathbf{PMet} that were mentioned in Sections 1 and 2.

The main result of this section is the following theorem that implies Theorem A.

Theorem 3.1. *Suppose that \mathbf{S} has non-trivial group actions and bornological Cartesian powers. Let G and H be two groups with G non-trivial and let X be a set on which H acts. Then the wreath product $G \wr_X H$ has property BS if and only if G and H have property BS and X is finite.*

Theorem 3.1 is a direct consequence of the forthcoming Corollary 3.6 and Lemmas 3.10 and 3.12. The conclusion of Theorem 3.1 remains true if the hypothesis on \mathbf{S} are replaced by “ \mathbf{S} satisfies (A2) and property BS implies property FW”, see the discussion after Lemma 3.10 for more details.

We now state two elementary but useful results.

Lemma 3.2. *Let G be a group and H be a quotient. If G has property BS, then so has H .*

Proof. If H acts on some \mathbf{S} -space X with an unbounded orbit, then the surjection $G \twoheadrightarrow H$ gives us a G -action on X , with the same orbits as the H -action. ■

Lemma 3.3. *Let G be a group and A and B be two subgroups such that $G = AB$. If both (G, A) and (G, B) have relative property BS, then G has property BS.*

Proof. Let X be an \mathbf{S} -space on which G acts and let x be an element of X . Let D_1 be the diameter of $A.x$ and D_2 the diameter of $B.x$. By assumption, they both are finite. Since A acts by isometries, all the $a.Bx$ have diameter D_2 . Let y be an element of $G.x$. There exists $a \in A$ such that y belongs to $a.Bx$. Since 1 belongs to B , y is at distance at most D_2 of $a.x$ and hence at distance at most $D_1 + D_2$ of x . Therefore, the diameter of $G.x$ is finite. ■

By combining Lemmas 3.2 and 3.3, we obtain the following three corollaries on direct, semi-direct and wreath products.

Corollary 3.4. *Let G and H be two groups. Then $G \times H$ has property BS if and only if both G and H have property BS.*

Corollary 3.5. *Let $N \rtimes H$ be a semi-direct product. Then,*

- (1) *if $N \rtimes H$ has property BS, then so has H ;*
- (2) *if both N and H have property BS, then $N \rtimes H$ also has property BS.*

Corollary 3.6. *Let G and H be two groups and X a set on which H acts. Then,*

- (1) *if $G \wr_X H$ has property BS, then so has H ;*
- (2) *if both G and H have property BS and X is finite, then $G \wr_X H$ has property BS.*

When \mathbf{S} has a suitable notion of quotients (by a group of isometries), it is possible to obtain a strong version of Lemmas 3.2 and 3.3. Here is the corresponding result for Bergman’s property and uncountable cofinality.

Proposition 3.7. *Let BS be either Bergman’s property or the property of having uncountable cofinality. Let $1 \rightarrow N \rightarrow G \rightarrow H \rightarrow 1$ be a group extension. Then G has property BS if and only if H has property BS and the pair (G, N) has relative BS property.*

Proof. One direction is simply Lemma 3.2 and the definition of relative property BS.

On the other hand, let (X, d) be a pseudo-metric space on which G acts by isometries and let x be an element of X . Let $\{g_i \mid i \in I\}$ be a transversal for N , that is,

$$H \cong \{g_i N \mid i \in I\}$$

with the quotient multiplication. By assumption, $N.x$ is bounded of diameter D_1 and for any $i \in I$ the subset $g_i N.x$ of X has also diameter D_1 . Since N is a subgroup of isometries of X , the map $d': X/N \times X/N \rightarrow \mathbf{R}$ defined by

$$d'([x], [y]) := \inf\{d(x', y') \mid x' \in N.x, y' \in N.y\}$$

is the quotient pseudo-distance on X/N . Indeed, while the map d' might not satisfies the triangle inequality for a generic quotient X/\sim , this is the case if the quotient is by a subgroup of isometries; details are left to the reader. Moreover, if d satisfies the strong triangle inequality, then so does d' . The quotient action of $H \cong G/N$ on X/N is by isometries and the diameter of $H.xN$ is bounded, say by D_2 . In particular, for any i and j in I , the distance between the subsets $g_i N.x$ and $g_j N.x$ of X is bounded by D_2 . Since this distance is an infimum, there exist actual elements of $g_i N.x$ and $g_j N.x$ at distance less than $D_2 + 1$. Altogether, we obtain that any y in $G.x$ is at distance at most $D_1 + D_2 + 1$ of x . Hence, the orbit $G.x$ is bounded. ■

Since the triangle graph, which is not median, is a quotient of the 2-regular infinite tree by a subgroup of isometries, the proof of Proposition 3.7 does not carry over for properties FW and FA. Similarly, the quotient of \mathbf{R} by the action of $\mathbf{Z}/2\mathbf{Z}$ given by $x \mapsto -x$ is not a Banach space and hence the proof of Proposition 3.7 does not apply to properties FH and FB_r . However, the statement of Proposition 3.7 (stability under extensions) remains true for properties FH, FB_r , FW and FA. For properties FH and FB_r , this follows from the fixed-point definition and the fact that a non-empty closed subset of an Hilbert space (respectively, of a reflexive Banach space) is an Hilbert space (respectively, a reflexive Banach space) itself. For properties FW and FA, see [11, Proposition 5.B.3] and [28].

We now state a result on infinite direct sums.

Lemma 3.8. *Let G and $(G_x)_{x \in X}$ be non-trivial groups and let H be a group acting on X . Then,*

- (1) $\bigoplus_{x \in X} G_x$ has uncountable cofinality if and only if all the G_x have uncountable cofinality and X is finite;
- (2) if $G \wr_X H$ has uncountable cofinality, then H acts on X with finitely many orbits.

It is of course possible to prove Lemma 3.8 using the characterization of uncountable cofinality in terms of subgroups, in which case the proof is a short exercise left to the reader. However, we find enlightening to prove it using the characterization in terms of actions on ultrametric spaces.

Proof of Lemma 3.8. One direction of the first assertion is simply Corollary 3.4 and holds for any property of the form BS. For the other direction, for any \mathbf{S} , if $\bigoplus_{x \in X} G_x$ has property BS then all its quotients, and hence all the G_x , have property BS. Hence, we have to prove that an infinite direct sum of non-trivial groups does not have uncountable cofinality. If X is infinite, there exists a countable subset $Y \subset X$. Let $Z := X \setminus Y$, thus we have $X = Y \sqcup Z$. We can decompose the direct sum as

$$\bigoplus_{x \in X} G_x = \left(\bigoplus_{x \in Y} G_x \right) \times \left(\bigoplus_{x \in Z} G_x \right)$$

and then, by Corollary 3.4, if $\bigoplus_{x \in Y} G_x$ does not have uncountable cofinality, then neither does $\bigoplus_{x \in X} G_x$. So let us fix an enumeration of Y and let $K := \bigoplus_{i \geq 1} G_i$ and for each i , choose $g_i \neq 1$ in G_i . Let $d_{\max}(f, g) := \max\{i \mid f(i) \neq g(i)\}$. This is a K -invariant ultra-metric for the action by left multiplication of G on itself. Then for every integer n , the orbit $K \cdot \{1, 1, \dots\}$ contains $\{g_1, \dots, g_n, 1, \dots\}$, which is at distance n of $\{1, 1, \dots\}$ for d_{\max} if the g_i are not equal to 1.

The second assertion is a simple variation on the first. Indeed, we have

$$G \wr_X H \cong \left(\bigoplus_{Y \in X/H} L_Y \right) \rtimes H \quad \text{with } L_Y \cong \bigoplus_{y \in Y} G_y,$$

where X/H is the set of H -orbits. The important fact for us is that H fixes the decomposition into L_Y factors: for all Y we have $H.L_Y = L_Y$. Up to regrouping some of the L_Y together, we hence have

$$G \wr_X H \cong \left(\bigoplus_{i \geq 1} L_i \right) \rtimes H$$

with $H.L_i = L_i$ for all i . Now, we have an ultradistance d_{\max} on $L := \bigoplus_{i \geq 1} L_i$ as above and we can put the discrete distance d on H . Then $d'_{\max} := \max\{d_{\max}, d\}$ is an

ultradistance on $(\bigoplus_{i \geq 1} L_i) \rtimes H$, which is $(\bigoplus_{i \geq 1} L_i) \rtimes H$ -invariant (for the action by left multiplication). From a practical point of view, we have

$$d'_\infty((\varphi, h), (\varphi', h')) := \max\{i \mid \varphi(i) \neq \varphi'(i)\}$$

if $\varphi \neq \varphi'$, and

$$d'_{\max}((\varphi, h), (\varphi, h')) = 1$$

if $h \neq h'$. Since the action of L on itself has an unbounded orbit for d_{\max} , the action of $(\bigoplus_{i \geq 1} L_i) \rtimes H$ on itself has an unbounded orbit for d'_{\max} . ■

We directly obtain the following corollary.

Corollary 3.9. *Suppose that BS implies having uncountable cofinality. Let G and $(G_x)_{x \in X}$ be non-trivial groups and let H be a group acting on X . Then,*

- (1) $\bigoplus_{x \in X} G_x$ has property BS if and only if all the G_x have property BS and X is finite;
- (2) if $G \wr_X H$ has property BS, then H acts on X with finitely many orbits.

While the statement (and the proof) of Corollary 3.9 (1) is expressed in terms of uncountable cofinality, it is also possible to state it and prove it for a subcategory **S** of **PMet** without a priori knowing if BS is stronger than having uncountable cofinality. The main idea is to find a “natural” **S**-space on which $G = \bigoplus_{i \geq 1} G_i$ acts. For example, for (reflexive) Banach, Hilbert and L^p spaces, one can take $\bigoplus_{i \geq 1} \ell^p(G_i)$. For connected median graphs, one takes the connected component of $\{1_{G_1}, 1_{G_2}, \dots\}$ in $\mathcal{P}(\bigsqcup_{i \geq 1} G_i)$. For (real) trees, it is possible to put a forest structure on $\mathcal{P}(\bigsqcup_{i \geq 1} G_i)$ in the following way. For $E \in \mathcal{P}(\bigsqcup_{i \geq 1} G_i)$, and for each i such that $E \cap G_j$ is empty for all $j \leq i$, add an edge from E to $E \cup \{g\}$ for each $g \in G_i$. The graph obtained this way is a G -invariant subforest of the median graph on $\mathcal{P}(\bigsqcup_{i \geq 1} G_i)$. For Corollary 3.9 (2), we also need that the corresponding structure is invariant by the action of H , which is the case of the above examples, except for the tree structure.

It is also possible to give a proof of Corollary 3.9 (1) using axioms similar to (A1)–(A3). More precisely, we need a variation of (A3) for countable Cartesian products (for morphisms we only ask that $\bigoplus_{i \in \mathbb{N}} \text{Aut}_{\mathbf{S}}(X_i) \subseteq \text{Aut}_{\mathbf{S}}(\bigoplus_{i \in \mathbb{N}} X_i)$) and a strong version of (A1) saying that there exists a universal bound M such that any non-trivial group G acts on some **S**-space moving a point at distance at least M . The axiomatization of Corollary 3.9 (2) is a little more complex. However, since in the following we will use Corollary 3.9 only for (real) trees, which do not have Cartesian powers, we will not elaborate on the details and let the proof to the interested reader. Instead, we will give an axiomatic proof of the following variation of Corollary 3.9 (2).

Lemma 3.10. *Suppose that \mathbf{S} has non-trivial group actions and satisfies axiom (A3). Let G and H be two groups with G non-trivial and let X be a set on which H acts. If $G \wr_X H$ has BS, then X is finite.*

Proof. We will prove that if X is infinite, then $G \wr_X H$ does not have property BS. Suppose that X is infinite. By non-trivial groups actions, there exists an \mathbf{S} -space Y on which G acts non-trivially by moving some element y_0 to another element $z_0 \neq y_0$. Let $\bigoplus_X Y$ be the corresponding Cartesian power and f_0 be the constant function $f_0(x) = y_0$. By assumption, the natural action of $G \wr_X H$ on $\bigoplus_X Y$ is by \mathbf{S} -automorphisms. Since X is infinite, it contains a countable subset $I = \{i_1, i_2, \dots\}$. For every integer n , the function

$$f_n(x) := \begin{cases} z_0 & \text{if } x = i_m, m \leq n, \\ y_0 & \text{otherwise} \end{cases}$$

is in the $G \wr_X H$ -orbit of f_0 . By axiom (A3) this orbit is unbounded and $G \wr_X H$ does not have property BS. ■

As a direct corollary, we obtain that if property BS implies property BS' for some S' with non-trivial group actions and (A3) (example: BS' = FW), then the conclusion of Lemma 3.10 holds even if \mathbf{S} might not satisfy its premises. Conversely, it follows from Theorem 1.2 and Proposition B that Lemma 3.10 do not holds for property FR, property FA or having uncountable cofinality.

We now turn our attention to properties that behave well under finite Cartesian products in the sense of axiom (A2). We first describe the comportment of property BS under finite index subgroups.

Lemma 3.11. *Let G be a group and let H be a finite index subgroup. Then,*

- (1) *if H has property BS, then so has G ;*
- (2) *if \mathbf{S} satisfies (A2) and G has property BS, then H has property BS.*

Proof. Suppose that G does not have BS and let X be an \mathbf{S} -space on which G acts with an unbounded orbit \mathcal{O} . Then H acts on X and \mathcal{O} is a union of at most $[G : H]$ orbits. This directly implies that H has an unbounded orbit and therefore does not have BS.

On the other hand, suppose that $H \leq G$ is a finite index subgroup of G without property BS. Let $\alpha: H \curvearrowright X$ be an action of H on an \mathbf{S} -space (X, d_X) such that there is an unbounded orbit \mathcal{O} . Similarly to the classical theory of representations of finite groups, we have the induced action $\text{Ind}_H^G(\alpha): G \curvearrowright X^{G/H}$ on the set $X^{G/H}$. Since H has finite index, $X^{G/H}$ is an \mathbf{S} -space and the action is by \mathbf{S} -automorphisms. On the other hand, the subgroup $H \leq G$ acts diagonally on $X^{G/H}$, which implies that $\text{diag}(\mathcal{O})$ is contained in a G -orbit. Since $\text{diag}(\mathcal{O})$ is unbounded, G does not have property BS.

For readers that are not familiar with representations of finite groups, here is the above argument in more details. Let $(f_i)_{i=1}^n$ be a transversal for G/H . The natural action of G on G/H gives rise to an action of G on $\{1, \dots, n\}$. Hence, for any g in G and i in $\{1, \dots, n\}$ there exists a unique $h_{g,i}$ in H such that $gf_i = f_{g.i}h_{g,i}$. That is, $h_{g,i} = f_{g.i}^{-1}gf_i$. We then define

$$g.(x_1, \dots, x_n) := (h_{g,g^{-1}.1}.x_{g^{-1}.1}, \dots, h_{g,g^{-1}.n}.x_{g^{-1}.n}).$$

This is indeed an action by \mathbf{S} -automorphisms on $X^{G/H}$ by Definition 2.13 (2). Moreover, every element $h \in H$ acts diagonally by $h.(x_1, \dots, x_n) = (h.x_1, \dots, h.x_n)$. In particular, this G -action has an unbounded orbit. ■

We now prove one last lemma that will be necessary for the proof of Theorem A.

Lemma 3.12. *Suppose that \mathbf{S} satisfies (A2). If X is finite and $G \wr_X H$ has property BS, then G has property BS.*

Proof. Suppose that G does not have BS and let (Y, d_Y) be an \mathbf{S} -space on which G acts with an unbounded orbit $G.y$. Then (Y^X, d) is an \mathbf{S} -space and we have the primitive action of the wreath product $G \wr_X H$ on Y^X :

$$((\varphi, h).\psi)(x) = \varphi(h^{-1}.x).\psi(h^{-1}.x).$$

By Definition 2.13 (2), this action is by \mathbf{S} -automorphisms. The orbit $G.y$ embeds diagonally and hence $\text{diag}(G.y)$ is an unbounded subset of some $G \wr_X H$ -orbit, which implies that $G \wr_X H$ does not have property BS. ■

It is also possible to derive Lemma 3.12 directly from Lemma 3.11 (2), with a more algebraic proof. Indeed, using the notation and hypothesis of Lemma 3.12, let H' be the kernel of the action of H on X and $\pi: G \wr_X H \rightarrow H$ be the canonical projection. Then $\pi^{-1}(H') \cong \bigoplus_X G \oplus H'$ is a finite index subgroup of $G \wr_X H$ and hence has property BS. Since G is a quotient of $\bigoplus_X G \oplus H'$, we conclude that it also has property BS.

We now proceed to prove Proposition B. As for Lemma 3.8, it is also possible to prove it using the characterization of uncountable cofinality in terms of subgroups, in which case it is an easy exercise, but we will only give a proof using the characterization in terms of actions on ultrametric spaces.

Proof of Proposition B. By Corollary 3.6 and Lemma 3.8 we already know that if $G \wr_X H$ has uncountable cofinality, then H has uncountable cofinality and it acts on X with finitely many orbits. We will now prove that if $G \wr_X H$ has uncountable cofinality, so does G . Let us suppose that G has countable cofinality. By Lemma 2.3, there exists

an ultrametric d on G such that the action of G on itself by left multiplication has an unbounded orbit. But then we have the primitive action of the wreath product $G \wr_X H$ on $G^X \cong \prod_X G$, which preserves $\bigoplus_X G$. It is easy to check that the map $d_\infty: \bigoplus_X G \times \bigoplus_X G \rightarrow \mathbf{R}$ defined by

$$d_\infty(\psi_1, \psi_2) := \max\{d(\psi_1(x), \psi_2(x)) \mid x \in X\}$$

is a $G \wr_X H$ -invariant ultrametric. Finally, let $g_0 \in G$ be an element of unbounded G -orbit for d and let x_0 be any element of X . For g in G and x in X , we define the following analog of Kronecker's delta functions

$$\delta_x^g(y) := \begin{cases} g & \text{if } y = x, \\ 1 & \text{if } y \neq x. \end{cases}$$

Then we have $(\delta_{x_0}^g, 1) \cdot \delta_{x_0}^{g_0} = \delta_{x_0}^{gg_0}$, and hence $d_\infty(\delta_{x_0}^{g_0}, \delta_{x_0}^{gg_0}) = d(g_0, gg_0)$ is unbounded.

Suppose now that both G and H have uncountable cofinality and that H acts on X with finitely many orbits. We want to prove that $G \wr_X H$ has uncountable cofinality.

Let (Y, d) be an ultrametric space on which $G \wr_X H$ acts. Then H and all the G_x act on Y with bounded orbits. Let $H.x_1, \dots, H.x_n$ be the H -orbits on X and let y be any element of Y . Then $H.y$ has finite diameter D_0 while $G_{x_i}.y$ has finite diameter D_i . For any $x \in X$, there exists $1 \leq i \leq n$ and $h \in H$ such that $x = h.x_i$. We have

$$\begin{aligned} d((\delta_{x_i}^g, h^{-1}).y, y) &\leq \max\{d((\delta_{x_i}^g, h^{-1}).y, (\delta_{x_i}^g, 1).y), d((\delta_{x_i}^g, 1).y, y)\} \\ &= \max\{d((1, h^{-1}).y, y), d((\delta_{x_i}^g, 1).y, y)\} \\ &\leq \max\{D_0, D_i\}, \end{aligned}$$

which implies that the diameter of $G_{x_i}h^{-1}.y$ is bounded by $\max\{D_0, D_i\}$. However, $G_{x_i}h^{-1}.y$ has the same diameter as $hG_{x_i}h^{-1}.y = G_{h.x_i}.y = G_x.y$.

On the other hand, the diameter of $\bigoplus_X G.y$ is bounded by the supremum of the diameters of the $G_x.y$, and hence bounded by $\max\{D_0, D_1, \dots, D_n\}$. Finally, for (φ, h) in $G \wr_X H$, we have

$$\begin{aligned} d(y, (\varphi, h).y) &\leq \max\{d(y, (\varphi, 1).y), d((\varphi, 1).y, (\varphi, h).y)\} \\ &= \max\{d(y, (\varphi, 1).y), d(y, (1, h).y)\} \\ &\leq \max\{\max\{D_0, D_1, \dots, D_n\}, D_0\}. \end{aligned}$$

That is, the diameter of $G \wr_X H.y$ is itself bounded by $\max\{D_0, D_1, \dots, D_n\}$, which finishes the proof. ■

While the fact that trees do not have Cartesian powers is an obstacle to our methods, we still have a weak version of Theorem 3.1 for properties **FA** and **FR**. Before stating it, remind that we already know, by Proposition B, the behavior of uncountable cofinality under wreath products. On the other hand, we have the following result.

Lemma 3.13. *The group $G \wr_X H$ has no quotient isomorphic to \mathbf{Z} if and only if both G and H have no quotient isomorphic to \mathbf{Z} .*

Proof. The desired result follows from $(G \wr_X H)^{\text{ab}} \cong \bigoplus_{X/H} (G^{\text{ab}}) \times H^{\text{ab}}$ and the claim that a direct sum $\bigoplus_{y \in Y} K_y$ has a quotient isomorphic to \mathbf{Z} if and only if at least one of the factor has a quotient isomorphic to \mathbf{Z} . Indeed, one direction of the claim is trivial. For the other direction, remind that K does not project onto \mathbf{Z} if and only if any action of K by orientation preserving isomorphisms on \mathbf{Z} , the 2-regular tree, has bounded orbits. But the only possibility for such an action to have bounded orbits is to be trivial. If none of the K_y projects onto \mathbf{Z} , all their actions on Z are trivial and so is any action of $\bigoplus_{y \in Y} K_y$, which can therefore not project onto \mathbf{Z} . ■

By Corollary 3.6, Proposition B and Lemma 3.13, we directly obtain the following partial version of Theorem 1.2.

Proposition 3.14. *Let G and H be two groups with G non-trivial and X a set on which H acts. Then,*

- (1) *if $G \wr_X H$ has property FA (respectively, property FR), then H has property FA (respectively, property FR), H acts on X with finitely many orbits, G has no quotient isomorphic to \mathbf{Z} and G has uncountable cofinality;*
- (2) *if both G and H have no quotient isomorphic to \mathbf{Z} , have uncountable cofinality and H acts on X with finitely many orbits, then $G \wr_X H$ has no quotient isomorphic to \mathbf{Z} and has uncountable cofinality;*
- (3) *if both G and H have property FA (respectively, property FR) and X is finite, then $G \wr_X H$ has property FA (respectively, property FR).*

Moreover, by using Corollary 3.9 we can get ride of the *finitely many orbits* hypothesis in Theorem 1.2 in order to obtain Theorem C.

Acknowledgments. The authors would like to thank P. de la Harpe, A. Genevois, T. Nagnibeda, and A. Valette for helpful comments and to the NSF for its support. They also gratefully thank the anonymous referees for their valuable remarks.

Funding. P.-H. Leemann was supported by grant 200021_188578 of the Swiss National Fund for Scientific Research.

References

- [1] H.-J. BANDEL and V. CHEPOI, [Metric graph theory and geometry: A survey](#). In *Surveys on discrete and computational geometry*, pp. 49–86, Contemp. Math. 453, Amer. Math. Soc., Providence, RI, 2008. Zbl [1169.05015](#) MR [2405677](#)

- [2] B. BEKKA, P. DE LA HARPE and A. VALETTE, *Kazhdan's property (T)*. New Math. Monogr. 11, Cambridge University Press, Cambridge, 2008. Zbl 1146.22009 MR 2415834
- [3] G. M. BERGMAN, *Generating infinite symmetric groups*. *Bull. London Math. Soc.* **38** (2006), no. 3, 429–440. Zbl 1103.20003 MR 2239037
- [4] M. BOURDON, *Un théorème de point fixe sur les espaces L^p* . *Publ. Mat.* **56** (2012), no. 2, 375–392. Zbl 1334.31006 MR 2978328
- [5] N. BROWN and E. GUENTNER, *Uniform embeddings of bounded geometry spaces into reflexive Banach space*. *Proc. Amer. Math. Soc.* **133** (2005), no. 7, 2045–2050. Zbl 1069.46003 MR 2137870
- [6] I. CHATTERJI, C. DRUȚU and F. HAGLUND, *Kazhdan and Haagerup properties from the median viewpoint*. *Adv. Math.* **225** (2010), no. 2, 882–921. Zbl 1271.20053 MR 2671183
- [7] I. CHATTERJI and G. NIBLO, *From wall spaces to CAT(0) cube complexes*. *Internat. J. Algebra Comput.* **15** (2005), no. 5-6, 875–885. Zbl 1107.20027 MR 2197811
- [8] V. CHEPOI, *Graphs of some CAT(0) complexes*. *Adv. in Appl. Math.* **24** (2000), no. 2, 125–179. Zbl 1019.57001 MR 1748966
- [9] P.-A. CHERIX, F. MARTIN and A. VALETTE, *Spaces with measured walls, the Haagerup property and property (T)*. *Ergodic Theory Dynam. Systems* **24** (2004), no. 6, 1895–1908. Zbl 1068.43007 MR 2106770
- [10] Y. DE CORNULIER, *Strongly bounded groups and infinite powers of finite groups*. *Comm. Algebra* **34** (2006), no. 7, 2337–2345. Zbl 1125.20023 MR 2240370
- [11] Y. CORNULIER, *Group actions with commensurated subsets, wallings and cubings*. [v1] 2013, [v2] 2016, arXiv:1302.5982v2.
- [12] — *Irreducible lattices, invariant means, and commensurating actions*. *Math. Z.* **279** (2015), no. 1-2, 1–26. Zbl 1311.43001 MR 3299841
- [13] Y. CORNULIER and A. KAR, *On property (FA) for wreath products*. *J. Group Theory* **14** (2011), no. 1, 165–174. Zbl 1228.20027 MR 2764930
- [14] A. GENEVOIS, *Lamplighter groups, median spaces and Hilbertian geometry*. *Proc. Edinb. Math. Soc. (2)* **65** (2022), no. 2, 500–529. Zbl 1514.20150 MR 4449680
- [15] V. N. GERASIMOV, *Semi-splittings of groups and actions on cubings*. In *Algebra, geometry, analysis and mathematical physics (Novosibirsk, 1996)*, pp. 91–109, Institut Matematiki im. S. L. Soboleva, Novosibirsk, 1997. Zbl 0906.20025 MR 1624115
- [16] — *Fixed-point-free actions on cubings*. *Siberian Adv. Math.* **8** (1998), no. 3, 36–58. Zbl 0912.20028 MR 1663779
- [17] P. DE LA HARPE and A. VALETTE, *La propriété (T) de Kazhdan pour les groupes localement compacts (avec un appendice de Marc Burger)*. Astérisque 175, Société Mathématique de France, Paris, 1989. Zbl 0759.22001 MR 1023471

- [18] E. HEWITT and K. A. ROSS, *Abstract harmonic analysis. Vol. I*. Second edn., Grundlehren Math. Wiss. 115, Springer, Berlin-New York, 1979. Zbl [0416.43001](#) MR [0551496](#)
- [19] S. KLAVŽAR and H. M. MULDER, Median graphs: Characterizations, location theory and related structures. *J. Combin. Math. Combin. Comput.* **30** (1999), 103–127. Zbl [0931.05072](#) MR [1705337](#)
- [20] P.-H. LEEMANN and G. SCHNEEBERGER, [Property FW and wreath products of groups: A simple approach using Schreier graphs](#). *Expo. Math.* **40** (2022), no. 4, 1261–1270. Zbl [1525.20023](#) MR [4518652](#)
- [21] A. MINASYAN, [New examples of groups acting on real trees](#). *J. Topol.* **9** (2016), no. 1, 192–214. Zbl [1365.20021](#) MR [3465847](#)
- [22] L. NEBESKÝ, Median graphs. *Comment. Math. Univ. Carolinae* **12** (1971), 317–325. Zbl [0215.34001](#) MR [0286705](#)
- [23] M. NEUHAUSER, [Relative property \(T\) and related properties of wreath products](#). *Math. Z.* **251** (2005), no. 1, 167–177. Zbl [1139.22002](#) MR [2176470](#)
- [24] L. NGUYEN VAN THÉ and V. G. PESTOV, [Fixed point-free isometric actions of topological groups on Banach spaces](#). *Bull. Belg. Math. Soc. Simon Stevin* **17** (2010), no. 1, 29–51. Zbl [1207.22002](#) MR [2656670](#)
- [25] B. NICA, [Cubulating spaces with walls](#). *Algebr. Geom. Topol.* **4** (2004), 297–309. Zbl [1131.20030](#) MR [2059193](#)
- [26] P. W. NOWAK, Group actions on Banach spaces. In *Handbook of group actions. Vol. II*, pp. 121–149, Adv. Lect. Math. (ALM) 32, International Press, Somerville, MA, 2015. MR [3382026](#)
- [27] M. ROLLER, Poc sets, median algebras and group actions. PhD thesis, Universität Regensburg, 1998.
- [28] J.-P. SERRE, *Arbres, amalgames, SL_2* . Astérisque 46, Société Mathématique de France, Paris, 1977. Zbl [0369.20013](#) MR [0476875](#)
- [29] R. TESSERA and A. VALETTE, [Locally compact groups with every isometric action bounded or proper](#). *J. Topol. Anal.* **12** (2020), no. 2, 267–292. Zbl [1444.22006](#) MR [4119107](#)
- [30] E. R. VERHEUL, *Multimedians in metric and normed spaces*. CWI Tract 91, Stichting Mathematisch Centrum, Centrum voor Wiskunde en Informatica, Amsterdam, 1993. Zbl [0790.46008](#) MR [1244813](#)

(Reçu le 25 juin 2021)

Paul-Henry LEEMANN, Institut de Mathématiques, Université de Neuchâtel,
11 rue Emile-Argand, 2000 Neuchâtel, Switzerland; e-mail: paulhenry.leemann@xjtlu.edu.cn

Grégoire SCHNEEBERGER, Section de mathématiques, Université de Genève,
7-9 rue du Conseil Général, 1205 Genève, Switzerland;
e-mail: gregoire.schneeberger@unige.ch