

Levinson's theorem for two-dimensional scattering systems: It was a surprise, it is now topological!

Angus Alexander, Thanh Duc Nguyen, Adam Rennie, and Serge Richard

Abstract. We prove a general Levinson's theorem for Schrödinger operators in two dimensions with threshold obstructions at zero energy. Our results confirm and simplify earlier seminal results of Bollé, Gesztesy et al., while providing an explicit topological interpretation. We also derive explicit formulas for the wave operators, and so show that they are elements of a C^* -algebra introduced by Cordes. As a consequence of our approach, we provide an evaluation of the spectral shift function at zero in the presence of p -resonances.

1. Introduction

We re-examine scattering theory for two-dimensional Schrödinger operators, a challenging subject that has been the focus of many studies. The aim of the present paper is threefold. We firstly confirm the results obtained in [6] for Levinson's theorem. Secondly, we recast their proof in an updated framework with more powerful tools. Thirdly, we restore the topological nature of Levinson's theorem by exhibiting it as an index theorem in scattering theory. In doing so, we reach our main result and recover the analytic formula for the number of bound states stated in [6, Theorem 6.3], and below in (1.2). Our general approach to index theorems in scattering theory has been described in the review paper [36] and illustrated in several examples [2, 16, 17, 24–29, 33, 35, 37].

With H_0 the free Hamiltonian and V a suitable potential, it is known that the 0-energy behaviour of the self-adjoint operator $H = H_0 + V$ is fairly complicated and plays a crucial role. Namely, the possible coexistence of 0-energy bound states and of two types of 0-energy resonances has an impact on propagation properties of the evolution group and on boundedness of the wave operators in various spaces. For concreteness, let us immediately recall from [21, Theorem 6.2] that for a two-dimensional Schrödinger operator H , a solution $\Psi \neq 0$ of the equation $H\Psi = 0$ is an s -resonance if $\Psi \in L^\infty(\mathbb{R}^2)$ but $\Psi \notin L^p(\mathbb{R}^2)$ for any $p < \infty$, Ψ is a p -resonance

Mathematics Subject Classification 2020: 81U05 (primary); 35P25 (secondary).

Keywords: Schrödinger operators, wave operators, resonances, topological index theorem.

if $\Psi \in L^p(\mathbb{R}^2)$ for any $p > 2$ but $\Psi \notin L^2(\mathbb{R}^2)$, and Ψ is an eigenfunction of H if $\Psi \in L^2(\mathbb{R}^2)$. Note that, in the case of a spherically symmetric potential V , the attributes s or p match with the s - or p -orbitals in atomic physics, see also [30, Table 1]. A short review of the corresponding literature is provided below.

The intricate 0-energy behaviour of two-dimensional Schrödinger operators has also a tremendous impact on the so-called Levinson's theorem, see (1.2). It was shown in [7], and later confirmed in [6], that an s -resonance does not play any role in this context while any p -resonance leads to a contribution of 1, similar to bound states. These properties are in sharp contrast from the one-dimensional or three-dimensional situation, in which resonances give a contribution of $\pm \frac{1}{2}$, and for that reason it was announced as *a surprise* in the first of the two mentioned papers. Unfortunately, the proofs of the results of [6] are based on double asymptotic expansions of the resolvent, which make them strenuous to follow. Note that the expression *half-bound states* for resonances in dimension 1 and 3 has been coined because of the mentioned contribution $\pm \frac{1}{2}$, but this expression is no more meaningful in the present context.

Before presenting our results in more detail, let us provide a brief (and non-exhaustive) description of the literature related to our work. A decade after the two surprising papers [6, 7], a renewed interest in the two-dimensional case has been triggered by the work [48] and then [22] on the L^p -boundedness of the wave operators. However, these works were conducted under the assumption that 0-energy bound states and 0-energy resonances are absent (the so-called regular or generic case). The next breakthrough came with the derivation in [21] of a simplified resolvent expansion, no longer given as a two parameter expansion, but in terms of powers of a single parameter. Subsequently, numerous works took advantage of this simplified resolvent expansion, as for example [4, 13, 38, 42] in which the assumption of the absence 0-energy bound states and 0-energy resonances remains. In other works, it was assumed that 0-energy bound states and p -resonances are absent, as for example in [45], or that only the p -resonances are absent, as in [11]. The first results on the behaviour of the Schrödinger evolution in the general case appeared then in [12]. More recently, two-dimensional Schrödinger operators with point interactions have been investigated: the boundedness of the wave operators in L^p -spaces in the regular case has been discussed in [9], while a full picture has been provided in [49]. Simultaneously, results on the scattering operator in the general setting have been exhibited in [40], together with an analysis of the wave operators in the absence of p -resonance. In particular, this paper contains the confirmation of the 0-energy behaviour of the scattering matrix, namely $\lim_{\lambda \searrow 0} S(\lambda) = 1$, obtained in [6] based on the double asymptotic expansion of the resolvent. Finally, building on [49], L^p -boundedness for more general Schrödinger operators with threshold obstructions has been fully investigated in [50].

Our approach for obtaining Levinson's theorem as an index theorem in scattering theory is based on a detailed study of the wave operators. Let us recall their definition, and refer to Section 2 for more details. We consider the scattering system given by the pair of operators (H, H_0) , where H_0 is the Laplacian in the Hilbert space $L^2(\mathbb{R}^2)$ and $H := H_0 + V$ with V a real potential decaying rapidly at infinity. Under quite general conditions on V it is known that the wave operators

$$W_{\pm} := \text{s-lim}_{t \rightarrow \pm\infty} e^{itH} e^{-itH_0}$$

exist and are complete. In particular, they are Fredholm operators with no kernel and with a cokernel given by the subspace spanned by the eigenfunctions of H . This subspace is of finite dimension for sufficiently fast decaying potential. Another important operator in this context is the scattering operator defined by $S := W_{+}^* W_{-}$. This operator is unitary in $L^2(\mathbb{R}^2)$. Since S strongly commutes with H_0 , the operator S decomposes in the spectral representation of H_0 . Thus, if we denote by \mathcal{F}_0 the unitary map from $L^2(\mathbb{R}^2)$ to $L^2(\mathbb{R}_{+}; L^2(\mathbb{S}))$ satisfying $(\mathcal{F}_0 H_0 f)(\lambda) = \lambda(\mathcal{F}_0 f)(\lambda)$ for any f in the domain of H_0 , then one has $\mathcal{F}_0 S \mathcal{F}_0^* = \{S(\lambda)\}_{\lambda \in \mathbb{R}_{+}}$, meaning that S is unitarily equivalent to a family of unitary operators $\{S(\lambda)\}_{\lambda \in \mathbb{R}_{+}}$ in $L^2(\mathbb{S})$. For historical reasons, the operator $S(\lambda)$ is called the *scattering matrix at energy λ* , even though it acts on an infinite-dimensional Hilbert space $L^2(\mathbb{S}) =: \mathfrak{h}$.

By using the stationary representation of the wave operators, our first result is a new formula for the wave operator W_{-} . More precisely, for V decaying fast enough, we show that the following equality holds:

$$\mathcal{F}_0(W_{-} - 1)\mathcal{F}_0^* = \left(\frac{1}{2}(1 - \tanh(\pi A_{+})) \otimes 1_{\mathfrak{h}}\right)(S(L) - 1) - N_2 \Xi B + K,$$

where A_{+} corresponds to the generator of the dilation group in $L^2(\mathbb{R}_{+})$, $S(L)$ denotes the operator of multiplication by the function $\lambda \mapsto S(\lambda)$, and K is a compact operator. So far, the product $N_2 \Xi B$ of three bounded operators is not really meaningful, but let us stress that this term is non-compact whenever H admits one or two p -resonances at 0. The precise definitions of these three operators are given in (3.4), (3.10), and (3.12); for now, it is enough to mention that N_2 and B are operator valued multiplication operators. Let us emphasise that this formula for W_{-} is at the root of the topological version of Levinson's theorem, and similar formulas have been obtained in several contexts, see for example [5, 15, 18, 19, 32, 34, 39, 43].

In order to get a better understanding of the new term $N_2 \Xi B$, a new representation is better suited. By conjugation with a suitable unitary rescaling, the wave operator W_{-} can be realised on $L^2(\mathbb{R}; L^2(\mathbb{S}))$, and by the decomposition into even and odd functions on \mathbb{R} , we end up studying the wave operator in the Hilbert space $L^2(\mathbb{R}_{+}; L^2(\mathbb{S}))^2$.

In this representation, the operator W_- takes the form

$$\begin{aligned} & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 1 & \tanh(\frac{\pi}{2}\sqrt{-\Delta_N})\phi(A_+) \\ \tilde{\phi}(A_+) \tanh(\frac{\pi}{2}\sqrt{-\Delta_N}) & 1 \end{pmatrix} \begin{pmatrix} \tilde{S}_e(L)-1 & \tilde{S}_o(L) \\ \tilde{S}_o(L) & \tilde{S}_e(L)-1 \end{pmatrix} \\ & + \begin{pmatrix} (\tilde{N}_2)_e(L) & (\tilde{N}_2)_o(L) \\ (\tilde{N}_2)_o(L) & (\tilde{N}_2)_e(L) \end{pmatrix} \begin{pmatrix} \frac{2}{1+i2A_+} & 0 \\ 0 & \frac{2}{1+i2A_+} \end{pmatrix} \begin{pmatrix} \tilde{B}_e(L) & \tilde{B}_o(L) \\ \tilde{B}_o(L) & \tilde{B}_e(L) \end{pmatrix} + K, \end{aligned} \quad (1.1)$$

with $\phi(A_+) := -\tanh(\pi A_+) + i \cosh(\pi A_+)^{-1}$, the indices e and o for the even or odd part of a function defined on \mathbb{R} , and the tilde functions meaning a rescaling, as for example $\tilde{S}(x) := S(e^{-2x})$ for any $x \in \mathbb{R}$. Again, the operator K is a compact operator. Let us emphasise some of the specific features of the previous formulas. It involves functions of three natural operators acting on $L^2(\mathbb{R}_+)$, namely the Neumann Laplacian $-\Delta_N$, the operator L of multiplication by the variable, and the generator A_+ of the unitary dilation group. In addition, it is shown in the following sections that all functions involved in this expression are continuous functions having limits either at $\pm\infty$, or at 0 and $+\infty$.

Obtaining formula (1.1) involves purely analytical tools, starting from the asymptotic expansion of the resolvent provided by [21], and using various analytical tricks for studying the stationary formula for the wave operators. Note that some of these tricks have been suggested by [50], even if the aims and the methods are different. These investigations are presented in Section 3 and in the first half of Section 4. The next key observation is that a C^* -algebra \mathcal{E} generated by functions of the three generators mentioned above has been thoroughly studied in [8, Chapter 5]. In particular, a precise description of the quotient of this algebra by the set of compact operators is provided: the quotient consists of continuous functions defined on the edges of a hexagon (this hexagon is illustrated in Section 5). By considering $M_2(\mathcal{E})$, the set of 2×2 matrices with values in \mathcal{E} , enlarging this algebra by a tensor product with $\mathcal{K}(L^2(\mathbb{S}))$, and adding a unit, one ends up with a C^* -algebra in which the expression (1.1) for the wave operator is natural.

Once in this framework, the rest of the investigation is more algebraic, and is presented in the second half of Section 4 and in Section 5. It firstly consists in computing the image of (1.1) in the quotient algebra. Since W_- is a Fredholm operator, this image is given by a continuous function Γ defined on the edges of the hexagon and taking unitary values in $\mathbb{C} + M_2(\mathcal{K}(L^2(\mathbb{S})))$. This function is provided in Proposition 4.4 and in Lemma 4.7. The operators \tilde{N}_2 and \tilde{B} take a much more explicit and interesting form in the quotient algebra: together they are the image of a projection P_p which is directly linked with the p -resonance of H , see (4.9) for a precise definition of P_p and Remark 4.6 for a discussion about the specific form of this projection.

Secondly, using a K -theoretic argument, the function Γ can be linked to the projection $E_p(H)$ on the subspace spanned by the eigenvectors of H . This construction is presented in Section 5 and is based on a description of the index map borrowed

from [41, Proposition 9.2.4.(ii)]. Thirdly, by applying traces, one infers a numerical equality. This equality involves the index of a Fredholm operator W_S defined by

$$W_S - 1 = \left(\frac{1}{2}(1 - \tanh(\pi A_+)) \otimes 1_{\mathfrak{h}} \right) (S(L) - 1),$$

and the operator trace of the bound state projection $E_p(H)$. It results in the following equality:

$$\text{Index}(W_S) + \dim(P_p) = -\#\sigma_p(H),$$

where $\#\sigma_p(H)$ corresponds to the cardinality of the set of eigenvalues of H , multiplicity counted. Note that the dimension of P_p corresponds to the number of p -resonances, and is 0, 1, or 2. By taking care of the high energy behaviour of the scattering matrix, one finally deduces our main result:

$$\frac{1}{2\pi i} \int_0^\infty \text{tr}(S(\lambda)^* S'(\lambda)) d\lambda + \frac{1}{4\pi} \int_{\mathbb{R}^2} V(x) dx + \dim(P_p) = -\#\sigma_p(H). \quad (1.2)$$

Simultaneously, we also determine the value of the spectral shift function at zero in the presence of p -resonances. We refer to Section 5 for more precise statements.

Equality (1.2) confirms that each p -resonance provides a contribution of 1 to Levinson's theorem, while the s -resonance does not provide any contribution. For comparison, let us recall the version of Levinson's theorem obtained in [6, Theorem 6.3] under the assumption of exponential decay of the potential and the condition $\int_{\mathbb{R}^2} V(x) dx \neq 0$. In the framework of [6], Levinson's theorem is expressed as

$$\int_0^\infty \text{Im}((H - \lambda - i0)^{-1} - (H_0 - \lambda - i0)^{-1}) d\lambda = -\pi N_- + \pi \Delta_{-1,-1} - \frac{1}{4} \int_{\mathbb{R}^2} V(x) dx, \quad (1.3)$$

where N_- is the number of strictly negative eigenvalues of H and $-\Delta_{-1,-1}$ is equal to the number of 0-energy eigenvalues and p -resonances. Taking into account the formal identity [6, equation (6.45)],

$$\text{Im} \text{tr}((H - \lambda - i0)^{-1} - (H_0 - \lambda - i0)^{-1}) = -\frac{i}{2} \frac{d}{d\lambda} \text{tr}(\ln(S(\lambda))),$$

it follows that (1.3) corresponds to (1.2).

Let us finally mention one main difference between (1.2) and (1.3): the contribution of the p -resonance is not on the same side of the equality, and the same remark holds for the expression involving the integral of V . Our right-hand side term contains only (minus) the trace of $E_p(H)$, which corresponds to the Fredholm index of W_- . In our approach, the contribution of the p -resonance projection P_p is coming

from the function Γ mentioned above, which describes the image of the wave operator W_- under the quotient map. In that respect, the p -resonance data has to stay on the same side as the scattering operator, which is also coming from Γ . On the other hand, the term involving the integral of V is due to a regularization process for the computation of $\text{Index}(W_S)$. For that reason, it also has to stay on the left-hand side of equality (1.2). Altogether, these contributions coming from scattering theory are equal to the contribution due to index theory, namely (minus) the trace on $E_p(H)$. Even though $\dim(P_p)$ is an integer, moving it to the other side of the equality sign would remove the topological character of this equality. As said in the title: it was a surprise, it is now topological!

Notations. $\mathbb{N} := \{0, 1, 2, \dots\}$ is the set of natural numbers, \mathcal{S} the Schwartz space on \mathbb{R}^2 , $\mathbb{R}_+ := (0, \infty)$, and $\langle \cdot \rangle := \sqrt{1 + |\cdot|^2}$. The sets \mathcal{H}_t^s are the weighted Sobolev spaces over \mathbb{R}^2 with index $s \in \mathbb{R}$ for derivatives and index $t \in \mathbb{R}$ for decay at infinity [3, Section 4.1], and with shorthand notations $\mathcal{H}^s := \mathcal{H}_0^s$, $\mathcal{H}_t := \mathcal{H}_t^0$, and $\mathcal{H} := \mathcal{H}_0^0 = L^2(\mathbb{R}^2)$. For any $s, t \in \mathbb{R}$, the 2-dimensional Fourier transform \mathcal{F} is a topological isomorphism of \mathcal{H}_t^s onto \mathcal{H}_s^t , and the scalar product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ (antilinear in the first argument) extends continuously to a duality $\langle \cdot, \cdot \rangle_{\mathcal{H}_t^s, \mathcal{H}_t^{-s}}$ between \mathcal{H}_t^s and \mathcal{H}_t^{-s} . Given two Banach spaces \mathcal{E}_1 and \mathcal{E}_2 , $\mathcal{B}(\mathcal{E}_1, \mathcal{E}_2)$ (resp. $\mathcal{K}(\mathcal{E}_1, \mathcal{E}_2)$) denotes the set of bounded (resp. compact) operators from \mathcal{E}_1 to \mathcal{E}_2 , with shorthand notation $\mathcal{B}(\mathcal{E}_1) := \mathcal{B}(\mathcal{E}_1, \mathcal{E}_1)$ (resp. $\mathcal{K}(\mathcal{E}_1) := \mathcal{K}(\mathcal{E}_1, \mathcal{E}_1)$). Finally, \otimes stands for the closed tensor product of Hilbert spaces or the spatial tensor product of operators.

2. Preliminaries

In this section, we briefly recall some notations and preliminary results introduced in [40, Section 2].

2.1. Free operator

Set $\mathfrak{h} := L^2(\mathbb{S})$ and $\mathcal{H} := L^2(\mathbb{R}_+; \mathfrak{h})$, and let H_0 be the (positive) self-adjoint operator in $\mathcal{H} = L^2(\mathbb{R}^2)$ given by minus the Laplacian $-\Delta$ on \mathbb{R}^2 . Then, the unitary operator $\mathcal{F}_0: \mathcal{H} \rightarrow \mathcal{H}$ defined by

$$((\mathcal{F}_0 f)(\lambda))(\omega) = 2^{-1/2}(\mathcal{F}f)(\sqrt{\lambda}\omega), \quad f \in \mathcal{S}, \lambda \in \mathbb{R}_+, \omega \in \mathbb{S}, \quad (2.1)$$

is a spectral transformation for H_0 in the sense that

$$(\mathcal{F}_0 H_0 f)(\lambda) = \lambda(\mathcal{F}_0 f)(\lambda) = (L\mathcal{F}_0 f)(\lambda), \quad f \in \mathcal{H}^2, \text{ a.e. } \lambda \in \mathbb{R}_+,$$

with L the maximal multiplication operator by the variable $\lambda \in \mathbb{R}_+$ in \mathcal{H} . Moreover, for each $\lambda \in \mathbb{R}_+$, the operator $\mathcal{F}_0(\lambda): \mathcal{S} \rightarrow \mathfrak{h}$ given by $\mathcal{F}_0(\lambda)f := (\mathcal{F}_0 f)(\lambda)$ extends

to an element of $\mathcal{B}(\mathcal{H}_t^s, \mathfrak{h})$ for any $s \in \mathbb{R}$ and $t > 1/2$, and the function $\mathbb{R}_+ \ni \lambda \mapsto \mathcal{F}_0(\lambda) \in \mathcal{B}(\mathcal{H}_t^s, \mathfrak{h})$ is continuous.

The asymptotic expansion of $\mathcal{F}_0(\lambda)$ as $\lambda \searrow 0$ plays an important role. By expanding the exponential $e^{-i\sqrt{\lambda}\omega \cdot x}$ in a Taylor series, one gets

$$\mathcal{F}_0(\lambda) = \gamma_0 + \sqrt{\lambda}\gamma_1 + \lambda\gamma_2 + o(\lambda), \quad \lambda \in \mathbb{R}_+, \quad (2.2)$$

with $\gamma_j: \mathcal{S} \rightarrow \mathfrak{h}$ ($j = 0, 1, 2$) the operator given by

$$(\gamma_j f)(\omega) := \frac{(-i)^j}{2^{3/2}\pi(j!)} \int_{\mathbb{R}^2} dx (\omega \cdot x)^j f(x), \quad f \in \mathcal{S}, \omega \in \mathcal{S}.$$

The operator γ_j extends to an element of $\mathcal{B}(\mathcal{H}_t^s, \mathfrak{h})$ for any $s \in \mathbb{R}$ and $t > j + 1$, which implies that the expansion (2.2) holds in $\mathcal{B}(\mathcal{H}_t^s, \mathfrak{h})$ as $\lambda \searrow 0$ for any $s \in \mathbb{R}$ and $t > 3$. We shall sometimes use the abbreviated notation $\gamma_2(\lambda)$, or $O(\lambda)$, for the sum $\lambda\gamma_2 + o(\lambda)$ in (2.2).

2.2. Perturbed operator

Let us now consider a potential $V \in L^\infty(\mathbb{R}^2; \mathbb{R})$ satisfying for some $\rho > 1$ the bound

$$|V(x)| \leq \text{Const.} \langle x \rangle^{-\rho}, \quad \text{a.e. } x \in \mathbb{R}^2. \quad (2.3)$$

Then, the perturbed Hamiltonian $H := H_0 + V$ is a short range perturbation of H_0 , and it is known that the corresponding wave operators

$$W_\pm := \text{s-lim}_{t \rightarrow \pm\infty} e^{itH} e^{-itH_0} \quad (2.4)$$

exist and are complete. As a consequence, the scattering operator $S := W_+^* W_-$ is unitary in \mathcal{H} . Now, define for $z \in \mathbb{C} \setminus \mathbb{R}$ the resolvents of H_0 and H

$$R_0(z) := (H_0 - z)^{-1} \quad \text{and} \quad R(z) := (H - z)^{-1}.$$

In order to recall properties of $R_0(z)$ and $R(z)$ as z approaches the real axis, it is convenient to decompose the potential V according to the following rule: for a.e. $x \in \mathbb{R}^2$ set

$$v(x) := |V(x)|^{1/2} \quad \text{and} \quad u(x) := \begin{cases} +1 & \text{if } V(x) \geq 0, \\ -1 & \text{if } V(x) < 0, \end{cases} \quad (2.5)$$

so that u is self-adjoint and unitary and $V = uv^2$. Then, using the fact that H does not have any positive eigenvalues [23, Section 1] and that a limiting absorption principle holds for H_0 and H [1, Theorem 4.2], we infer that the limits

$$vR_0(\lambda \pm i0)v := \lim_{\varepsilon \searrow 0} vR_0(\lambda \pm i\varepsilon)v \quad \text{and} \quad vR(\lambda \pm i0)v := \lim_{\varepsilon \searrow 0} vR(\lambda \pm i\varepsilon)v,$$

exist in $\mathcal{B}(\mathcal{H})$ and are continuous in the variable $\lambda \in \mathbb{R}_+$. This, together with the relation

$$u - uvR(\lambda \pm i\varepsilon)vu = (u + vR_0(\lambda \pm i\varepsilon)v)^{-1}, \quad \lambda \in \mathbb{R}_+, \varepsilon > 0,$$

implies the existence and the continuity of the function $\mathbb{R}_+ \ni \lambda \mapsto (u + vR_0(\lambda \pm i0)v)^{-1} \in \mathcal{B}(\mathcal{H})$. Furthermore, one has $\lim_{\lambda \rightarrow \infty} (u + vR_0(\lambda \pm i0)v)^{-1} = u$ in $\mathcal{B}(\mathcal{H})$, since $\lim_{\lambda \rightarrow \infty} vR_0(\lambda \pm i0)v = 0$ in $\mathcal{B}(\mathcal{H})$ [47, Proposition 7.1.2]. On the other hand, the existence in $\mathcal{B}(\mathcal{H})$ of the limits $\lim_{\lambda \searrow 0} (u + vR_0(\lambda \pm i0)v)^{-1}$ depends on the presence or absence of eigenvalues or resonances at 0-energy. This problem has been studied in detail in [21] in dimensions 1 and 2. We recall here the main result in dimension 2 [21, Theorem 6.2 (ii)]. Take $\kappa \in \mathbb{C}^*$ with $\operatorname{Re}(\kappa) \geq 0$, let $\eta := 1/\ln(\kappa)$ (with \ln the principal value of the complex logarithm), and set

$$\mathbf{M}(\kappa) := u + vR_0(-\kappa^2)v.$$

Then, if V satisfies (2.3) with $\rho > 11$ and if $0 < |\kappa| < \kappa_0$ with $\kappa_0 > 0$ small enough, the operator $\mathbf{M}(\kappa)^{-1}$ admits an expansion

$$\mathbf{M}(\kappa)^{-1} = I_1(\kappa) - g(\kappa)I_2(\kappa) - \frac{g(\kappa)\eta}{\kappa^2}I_3(\kappa), \quad (2.6)$$

with

$$\begin{aligned} I_1(\kappa) &:= (\mathbf{M}(\kappa) + S_1)^{-1}, \\ I_2(\kappa) &:= (\mathbf{M}(\kappa) + S_1)^{-1}S_1(M_1(\kappa) + S_2)^{-1}S_1(\mathbf{M}(\kappa) + S_1)^{-1}, \\ I_3(\kappa) &:= (\mathbf{M}(\kappa) + S_1)^{-1}S_1(M_1(\kappa) + S_2)^{-1} \\ &\quad \cdot S_2(T_3m(\kappa)^{-1}T_3 - T_3m(\kappa)^{-1}b(\kappa)d(\kappa)^{-1}S_3 - S_3d(\kappa)^{-1}c(\kappa)m(\kappa)^{-1}T_3 \\ &\quad \quad + S_3d(\kappa)^{-1}c(\kappa)m(\kappa)^{-1}b(\kappa)d(\kappa)^{-1}S_3 + S_3d(\kappa)^{-1}S_3)S_2 \\ &\quad \cdot (M_1(\kappa) + S_2)^{-1}S_1(\mathbf{M}(\kappa) + S_1)^{-1}, \end{aligned}$$

and where $S_1 \geq S_2 \geq S_3$ are finite-dimensional orthogonal projections in \mathcal{H} , $T_3 := S_2 - S_3$, $g: \mathbb{C} \rightarrow \mathbb{C}$ satisfies $g(\kappa) = O(\eta^{-1})$ for $0 < |\kappa| < \kappa_0$, $m: \mathbb{C} \rightarrow \mathcal{B}(\mathcal{H})$ satisfies $m(\kappa) = O(\eta^{-1})$ for $0 < |\kappa| < \kappa_0$, and all other factors are operator-valued functions having limits in $\mathcal{B}(\mathcal{H})$ as $\kappa \rightarrow 0$. As proved in [21, Theorem 6.2], the dimension of $T_2 := S_1 - S_2$ is at most 1 and this projection is related to s -resonance, the dimension of T_3 is at most 2 and this projection is related to p -resonances, and the projection S_3 is related to the possible 0-energy bound state(s) with its dimension equal to the number of linearly independent 0-energy eigenfunctions.

One of the initial tasks in [40] has been to provide the expansion near 0 of an operator related to $\mathbf{M}(\kappa)$ which plays an important role for the stationary expression of the wave operators. The statement is recalled below, with the convention that $\lambda > 0$, $\kappa := -i\sqrt{\lambda}$ which means that $\eta = \frac{1}{\ln(\lambda)/2 - i\pi/2}$.

Theorem 2.1 (Theorem 4.7 of [40]). *If V satisfies (2.3) with $\rho > 11$, then one has as $\lambda \searrow 0$*

$$\begin{aligned}
 & (u + vR_0(\lambda + i0)v)^{-1}v\mathcal{F}_0(\lambda)^* \\
 &= \frac{g(\kappa)\eta}{\sqrt{\lambda}}(T_3 - S_3d(\kappa)^{-1}c(\kappa))m(\kappa)^{-1}T_3v\gamma_1^* + \frac{1}{\eta}S_3O(1) + O(1) \\
 &= \frac{\eta}{\sqrt{\lambda}}S_2(T_3 - S_3d(\kappa)^{-1}c(\kappa))g(\kappa)m(\kappa)^{-1}T_3v\gamma_1^* + \frac{1}{\eta}S_3O(1) + O(1) \\
 &= S_2\left(\frac{\eta}{\sqrt{\lambda}}(T_3 - S_3d(\kappa)^{-1}c(\kappa))g(\kappa)m(\kappa)^{-1}T_3v\gamma_1^* + \frac{1}{\eta}S_3O(1) + O(1)\right) \\
 &\quad + S_2^\perp O(1), \tag{2.7}
 \end{aligned}$$

where $S_3O(1)$ denotes a family of bounded operators with their range in the subspace defined by the projection S_3 , and similarly for $S_2O(1)$ and $S_2^\perp O(1)$.

In part of the analysis performed in [40] the assumption $T_3 = 0$ was imposed. In this case, the main singularity in this expansion disappears, and only the milder singular term $\frac{1}{\eta}S_3O(1) + O(1)$ remains. Note that the factor S_3 is associated with 0-energy bound states, while the projection T_3 is related to the so-called p -resonances. In the sequel, we shall remove the assumption that $T_3 = 0$.

3. Stationary expression for the wave operators

In this section, we start by recalling the stationary expression for the wave operator W_- . We then decompose this expression into smaller pieces, which will be analysed separately. This analysis is taking place in the spectral representation of H_0 , namely in the space \mathcal{H} .

Since the subsequent developments are based on the asymptotic expansions provided in (2.6) and in (2.7), which hold under the assumption (2.3) with $\rho > 11$, we shall assume this decay in the rest of the paper, without repeating it. Under this assumption, the wave operators (2.4) obtained by the time-dependent approach and those described by the time-independent approach coincide [46, Theorem 5.3.6]. For suitable $\varphi, \psi \in \mathcal{H} = L^2(\mathbb{R}_+; \mathfrak{h})$, we recall the stationary expression for W_- , namely

$$\begin{aligned}
 & \langle \mathcal{F}_0(W_- - 1)\mathcal{F}_0^*\varphi, \psi \rangle_{\mathcal{H}} \\
 &= - \int_{\mathbb{R}} d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu)v\mathbf{1}(u + vR_0(\lambda + i\varepsilon)v)^{-1}v\mathcal{F}_0^*\delta_\varepsilon(L - \lambda)\varphi, \\
 &\quad (\mu - \lambda + i\varepsilon)^{-1}\psi(\mu) \rangle_{\mathfrak{h}}, \tag{3.1}
 \end{aligned}$$

where

$$\delta_\varepsilon(L - \lambda) := \frac{\varepsilon}{\pi} (L - \lambda + i\varepsilon)^{-1} (L - \lambda - i\varepsilon)^{-1}.$$

Note that we have artificially inserted an identity operator $\mathbf{1}$ in the above expression. Indeed, this identity operator can be rewritten as

$$\mathbf{1} = S_2 + S_2^\perp,$$

which then provides two expressions from (3.1). The one with S_2^\perp does not contain any singularity at 0-energy, and has been thoroughly studied in [40].

We shall now carefully decouple the low energy and the high energy parts of this expression. For that purpose, let us fix $\varrho \in C(\mathbb{R}_+; [0, 1])$ with

$$\varrho(\lambda) = \begin{cases} 0, & \lambda < \frac{1}{4}, \\ 1, & \lambda > \frac{3}{4}. \end{cases} \quad (3.2)$$

The function $1 - \varrho: \mathbb{R}_+ \rightarrow [0, 1]$ is denoted by ϱ^\perp . We shall also use two auxiliary functions $\varrho_1 \in C(\mathbb{R}_+; [0, 1])$ and $\varrho_0 \in C(\mathbb{R}_+; [0, 1])$ with

$$\varrho_1(\lambda) = \begin{cases} 0, & \lambda < \frac{1}{8}, \\ 1, & \lambda > \frac{1}{4}, \end{cases} \quad \text{and} \quad \varrho_0(\lambda) = \begin{cases} 1, & \lambda < \frac{3}{4}, \\ 0, & \lambda > \frac{7}{8}. \end{cases} \quad (3.3)$$

As a consequence, $\varrho\varrho_1 = \varrho$ and that $\varrho^\perp\varrho_0 = \varrho^\perp$. Finally, for $\varphi \in \mathcal{H}$ and $\varepsilon > 0$ we also set

$$\tilde{\varphi}_\varepsilon(\lambda) := S_2(u + vR_0(\lambda + i\varepsilon)v)^{-1} v\mathcal{F}_0^* \delta_\varepsilon(L - \lambda) \varrho^\perp(L)\varphi.$$

Then the term with the factor S_2 can be rewritten as

$$\begin{aligned} & \int_{\mathbb{R}} d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu) v (\mu - \lambda - i\varepsilon)^{-1} S_2(u + vR_0(\lambda + i\varepsilon)v)^{-1} \\ & \quad \cdot v\mathcal{F}_0^* \delta_\varepsilon(L - \lambda) \varphi, \psi(\mu) \rangle_{\mathfrak{H}} \\ &= \int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu) v S_2(\mu - \lambda - i\varepsilon)^{-1} S_2(u + vR_0(\lambda + i\varepsilon)v)^{-1} \\ & \quad \cdot v\mathcal{F}_0^* \delta_\varepsilon(L - \lambda) \varrho(L) \varphi, \psi(\mu) \rangle_{\mathfrak{H}} \\ & \quad + \int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu) v S_2 \varrho^\perp(\mu) \frac{1}{\sqrt{\mu}} (\sqrt{\mu} - \sqrt{\lambda} + \sqrt{\lambda}) \\ & \quad \cdot (\mu - \lambda - i\varepsilon)^{-1} \tilde{\varphi}_\varepsilon(\lambda), \psi(\mu) \rangle_{\mathfrak{H}} \\ & \quad + \int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu) v S_2 \varrho(\mu) \frac{1}{\mu} (\mu - \lambda + \lambda) \\ & \quad \cdot (\mu - \lambda - i\varepsilon)^{-1} \tilde{\varphi}_\varepsilon(\lambda), \psi(\mu) \rangle_{\mathfrak{H}} \\ &= R_0 + R_1 + R_2 + R_3 + R_4 \end{aligned}$$

where

$$\begin{aligned}
 R_0 &:= \int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu) v S_2(\mu - \lambda - i\varepsilon)^{-1} S_2(u + vR_0(\lambda + i\varepsilon)v)^{-1} \\
 &\quad \cdot v\mathcal{F}_0^* \delta_\varepsilon(L - \lambda) \varrho(L) \varphi, \psi(\mu) \rangle_{\mathfrak{h}}, \\
 R_1 &:= \int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu) v S_2 \varrho^\perp(\mu) \frac{1}{\sqrt{\mu}} (\sqrt{\mu} - \sqrt{\lambda}) \\
 &\quad \cdot (\mu - \lambda - i\varepsilon)^{-1} \tilde{\varphi}_\varepsilon(\lambda), \psi(\mu) \rangle_{\mathfrak{h}}, \\
 R_2 &:= \int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu) v S_2 \varrho^\perp(\mu) \frac{1}{\sqrt{\mu}} (\mu - \lambda - i\varepsilon)^{-1} \sqrt{\lambda} \tilde{\varphi}_\varepsilon(\lambda), \psi(\mu) \rangle_{\mathfrak{h}}, \\
 R_3 &:= \int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu) v S_2 \varrho(\mu) \frac{1}{\mu} (\mu - \lambda) (\mu - \lambda - i\varepsilon)^{-1} \tilde{\varphi}_\varepsilon(\lambda), \psi(\mu) \rangle_{\mathfrak{h}}, \\
 R_4 &:= \int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \mathcal{F}_0(\mu) v S_2 \varrho(\mu) \frac{1}{\mu} (\mu - \lambda - i\varepsilon)^{-1} \lambda \tilde{\varphi}_\varepsilon(\lambda), \psi(\mu) \rangle_{\mathfrak{h}}.
 \end{aligned}$$

These various terms will be treated below. For now, let us observe that for any $\varphi \in C_c(\mathbb{R}_+; \mathfrak{h})$ and any $\lambda > 0$, the following limit exists in \mathcal{H} , as a consequence of [39, Lemma 2.3]:

$$\text{s-}\lim_{\varepsilon \searrow 0} \tilde{\varphi}_\varepsilon(\lambda) = \varrho^\perp(\lambda) S_2(u + vR_0(\lambda + i0)v)^{-1} v\mathcal{F}_0(\lambda)^* \varphi(\lambda).$$

It is then natural to consider the subsequent operator-valued function of λ and study its behaviour for $\lambda \searrow 0$. The following statement is mainly a consequence of the expansion (2.7) and the properties of the function ϱ^\perp .

Lemma 3.1. *The following map is continuous and bounded:*

$$\mathbb{R}_+ \ni \lambda \mapsto B(\lambda) := \varrho^\perp(\lambda) \sqrt{\lambda} \ln(\lambda) S_2[(u + vR_0(\lambda + i0)v)^{-1} v\mathcal{F}_0(\lambda)^*] \in \mathcal{K}(\mathfrak{h}, \mathcal{H}). \quad (3.4)$$

The multiplication operator

$$B: C_c(\mathbb{R}_+; \mathfrak{h}) \rightarrow L^2(\mathbb{R}_+; \mathcal{H})$$

given, for $\varphi \in C_c(\mathbb{R}_+; \mathfrak{h})$ and $\lambda \in \mathbb{R}_+$, by

$$(B\varphi)(\lambda) := B(\lambda)\varphi(\lambda)$$

extends continuously to an element of $\mathcal{B}(\mathcal{H}, L^2(\mathbb{R}_+; \mathcal{H}))$.

Proof. The continuity of the functions $\lambda \mapsto B(\lambda)$ follows from the continuities already mentioned in Section 2. For the behavior of this function near 0, we can use Theorem 2.1 and replace the term inside the square bracket by the right-hand side of (2.7). Then, by recalling that $g(\kappa) = O(\eta^{-1})$ and $m(\kappa)^{-1} = O(\eta)$ as $\lambda \searrow 0$, one infers that the first term behaves as $O((\sqrt{\lambda} \ln(\lambda))^{-1})$ in the limit $\lambda \searrow 0$. The factor $\sqrt{\lambda} \ln(\lambda)$ makes this product bounded in the limit $\lambda \searrow 0$. For $\lambda > \frac{3}{4}$, the function $\lambda \mapsto B(\lambda)$ vanishes, since the factor $\varrho^\perp(\lambda)$ has this property. The rest of the statement is a direct consequence of boundedness of the map $\lambda \mapsto B(\lambda)$. ■

Let us now provide the analysis of the terms R_0 to R_4 which appear in our study of the stationary expression for the wave operator W_- . An important role is played by the generator of dilations, which we now describe. We recall that the dilation group $\{U_t^+\}_{t \in \mathbb{R}}$ in $L^2(\mathbb{R}_+)$, with self-adjoint generator A_+ , is given by $(U_t^+ \varphi)(\lambda) := e^{t/2} \varphi(e^t \lambda)$ for $\varphi \in C_c(\mathbb{R}_+)$, $\lambda \in \mathbb{R}_+$ and $t \in \mathbb{R}$. Then, by standard functional calculus for unitary groups one has

$$[\varphi(A_+)f](x) = (2\pi)^{-\frac{1}{2}} \int_{-\infty}^{\infty} (\mathcal{F}_1^* \varphi)(t) [U_t^+ f](x) dt, \quad (3.5)$$

with \mathcal{F}_1 the usual Fourier transform on $L^2(\mathbb{R})$ and $\varphi, f \in C_c^\infty(\mathbb{R})$. We shall also use the function $\vartheta: \mathbb{R} \rightarrow \mathbb{R}$ given for any $s \in \mathbb{R}$ by

$$\vartheta(s) := \frac{1}{2}(1 - \tanh(\pi s)). \quad (3.6)$$

Lemma 3.2. *The term R_0 can be rewritten as $\langle 2\pi i N_0(\vartheta(A_+) \otimes 1_{\mathcal{H}}) B_0 \varphi, \psi \rangle_{\mathcal{H}}$, with N_0 and B_0 defined below in (3.7) and in (3.8) respectively.*

Proof. Let us firstly define the map

$$\mathbb{R}_+ \ni \mu \mapsto N_0(\mu) := \mathcal{F}_0(\mu) v S_2 \in \mathcal{K}(\mathcal{H}, \mathfrak{h}). \quad (3.7)$$

It is easy to check that this function is continuous, admits a limit as $\mu \searrow 0$, and vanishes as $\mu \rightarrow \infty$, see [40, Lemma 4.9 (a)] and its proof for a similar statement. The multiplication operator

$$N_0: C_c(\mathbb{R}_+; \mathcal{H}) \rightarrow \mathcal{H}$$

given, for $\xi \in C_c(\mathbb{R}_+; \mathcal{H})$ and $\mu \in \mathbb{R}_+$, by

$$(N_0 \xi)(\mu) := N_0(\mu) \xi(\mu)$$

extends then continuously to an element of $\mathcal{B}(L^2(\mathbb{R}_+; \mathcal{H}), \mathcal{H})$.

Secondly, similar arguments to those of Lemma 3.1 show that the map

$$\mathbb{R}_+ \ni \lambda \mapsto B_0(\lambda) := \varrho(\lambda) S_2(u + v R_0(\lambda + i0)v)^{-1} v \mathcal{F}_0(\lambda)^* \in \mathcal{K}(\mathfrak{h}, \mathcal{H}) \quad (3.8)$$

is continuous and bounded. As a consequence, the multiplication operator

$$B_0: C_c(\mathbb{R}_+; \mathfrak{h}) \rightarrow L^2(\mathbb{R}_+; \mathcal{H})$$

given, for $\varphi \in C_c(\mathbb{R}_+; \mathfrak{h})$ and $\lambda \in \mathbb{R}_+$, by

$$(B_0\varphi)(\lambda) := B_0(\lambda)\varphi(\lambda)$$

extends continuously to an element of $\mathcal{B}(\mathcal{H}, L^2(\mathbb{R}_+; \mathcal{H}))$.

Then, by considering $\varphi, \psi \in C_c^\infty(\mathbb{R}_+) \odot C(\mathbb{S})$, we can prove as in [38, Theorem 2.5] that the expression R_0 given by

$$\int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu (\mathcal{F}_0(\mu) v S_2(\mu - \lambda - i\varepsilon)^{-1} S_2(u + v R_0(\lambda + i\varepsilon)v)^{-1} \cdot v \mathcal{F}_0^* \delta_\varepsilon(L - \lambda) \varrho(L) \varphi, \psi(\mu))_{\mathfrak{h}}$$

reduces to

$$\langle 2\pi i N_0(\vartheta(A_+) \otimes 1_{\mathcal{H}}) B_0\varphi, \psi \rangle_{\mathcal{H}}.$$

Note that the key argument in the proof is an application of Lebesgue's dominated convergence theorem, as shown in the proof of [38, Theorem 2.5]. \blacksquare

For the next statement, recall that ϱ is a localization near ∞ while ϱ_0 is a localization near 0, see (3.2) and (3.3).

Lemma 3.3. *The term R_3 can be rewritten as $\langle K\varphi, \psi \rangle_{\mathcal{H}}$, with $K \in \mathcal{K}(\mathcal{H})$.*

Proof. For $\mu, \lambda > 0$ and $\varepsilon > 0$ we consider the kernel

$$\Theta_\varepsilon(\mu, \lambda) := \varrho(\mu) \frac{1}{\mu} (\mu - \lambda)(\mu - \lambda - i\varepsilon)^{-1} \frac{1}{\sqrt{\lambda \ln(\lambda)}} \varrho_0(\lambda).$$

This function defines a Hilbert–Schmidt operator in $L^2(\mathbb{R}_+)$ with Hilbert–Schmidt norm satisfying

$$\begin{aligned} \|\Theta_\varepsilon\|_{\text{HS}}^2 &= \int_0^\infty d\mu \int_0^\infty d\lambda |\Theta_\varepsilon(\mu, \lambda)|^2 \\ &\leq \int_{\frac{1}{4}}^\infty d\mu \left(\frac{\varrho(\mu)}{\mu}\right)^2 \int_0^{\frac{7}{8}} d\lambda \left(\frac{\varrho_0(\lambda)}{\sqrt{\lambda \ln(\lambda)}}\right)^2 \end{aligned}$$

with an upper bound independent of ε . By an application of the dominated convergence theorem, one then infers that Θ_ε converges in the Hilbert–Schmidt norm to Θ_0 defined for $\mu, \lambda > 0$ by $\Theta_0(\mu, \lambda) := \frac{\varrho(\mu)}{\mu} \frac{\varrho_0(\lambda)}{\sqrt{\lambda \ln(\lambda)}}$.

By an application of the Lebesgue dominated convergence theorem, one obtains as in the previous proof that for $\varphi, \psi \in C_c^\infty(\mathbb{R}_+) \odot C(\mathbb{S})$, the expression R_3 given by

$$\int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \left\langle \mathcal{F}_0(\mu) v S_2 \varrho(\mu) \frac{1}{\mu} (\mu - \lambda) (\mu - \lambda - i\varepsilon)^{-1} \tilde{\varphi}_\varepsilon(\lambda), \psi(\mu) \right\rangle_{\mathfrak{H}}$$

reduces to

$$\langle N_0 (1_{L^2(\mathbb{R}_+)} \otimes S_2) (\Theta_0 \otimes 1_{\mathcal{H}}) B \varphi, \psi \rangle_{\mathcal{H}},$$

with N_0 defined in (3.7) and with B defined in (3.4). As S_2 is finite rank, the product $(1_{L^2(\mathbb{R}_+)} \otimes S_2) (\Theta_0 \otimes 1_{\mathcal{H}})$ corresponds to a compact operator on $L^2(\mathbb{R}_+; \mathcal{H})$. By multiplying this factor by the bounded operators B on the right and N_0 on the left, one obtains an element of $\mathcal{K}(\mathcal{H})$. ■

Lemma 3.4. *The term R_4 can be rewritten as*

$$\langle 2\pi i N_4 (\vartheta(A_+) \otimes 1_{\mathcal{H}}) (M_4 \otimes 1_{\mathcal{H}}) B \varphi, \psi \rangle_{\mathcal{H}},$$

with B defined in (3.4), and M_4 and N_4 bounded multiplication operators defined in proof.

Proof. For $\mu > 0$ we set

$$N_4(\mu) := N_0(\mu) \varrho(\mu) \frac{1}{\mu}. \quad (3.9)$$

This operator valued function is bounded, vanishes as $\mu \searrow 0$, and satisfies the limit $\lim_{\mu \rightarrow \infty} N_4(\mu) = 0$. For $\lambda > 0$ let us also set

$$M_4(\lambda) := \varrho_0(\lambda) \sqrt{\lambda} \frac{1}{\ln(\lambda)}.$$

This scalar function is bounded, vanishes for $\lambda > \frac{7}{8}$, and satisfies $\lim_{\lambda \searrow 0} N_4(\lambda) = 0$. The corresponding bounded multiplication operators are denoted by N_4 and M_4 , respectively.

Then, by considering $\varphi, \psi \in C_c^\infty(\mathbb{R}_+) \odot C(\mathbb{S})$, we can prove as in [38, Theorem 2.5] that the expressions R_4 given by

$$\int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \left\langle \mathcal{F}_0(\mu) v S_2 \varrho(\mu) \frac{1}{\mu} (\mu - \lambda - i\varepsilon)^{-1} \lambda \tilde{\varphi}_\varepsilon(\lambda), \psi(\mu) \right\rangle_{\mathfrak{H}}$$

reduces to

$$\langle 2\pi i N_4(\vartheta(A_+) \otimes 1_{\mathcal{H}})(M_4 \otimes 1_{\mathcal{H}})B\varphi, \psi \rangle_{\mathcal{H}},$$

with B defined in (3.4). ■

Lemma 3.5. *The term R_2 can be rewritten as*

$$\langle 2\pi i N_2(\vartheta(A_+) \otimes 1_{\mathcal{H}})(M_2 \otimes 1_{\mathcal{H}})B\varphi, \psi \rangle_{\mathcal{H}}$$

with B defined in (3.4), and M_2 and N_2 bounded multiplication operators defined in proof.

Proof. For $\lambda > 0$, let us set $M_2(\lambda) := \varrho_0(\lambda) \frac{1}{\ln(\lambda)}$. This scalar function is clearly bounded, vanishes for $\lambda > \frac{7}{8}$, and satisfies $\lim_{\lambda \searrow 0} M_2(\lambda) = 0$. The corresponding bounded multiplication operator in $L^2(\mathbb{R}_+)$ are denoted by M_2 . Let us also define the map

$$\mathbb{R}_+ \ni \mu \mapsto N_2(\mu) := \mathcal{F}_0(\mu) \varrho^\perp(\mu) \frac{1}{\sqrt{\mu}} v S_2 \in \mathcal{B}(\mathcal{H}, \mathfrak{h}). \quad (3.10)$$

Clearly, this map vanishes as $\mu \rightarrow \infty$. By the expansion (2.2) together with the algebraic cancellation $\gamma_0 v S_2 = 0$, as shown in [40, Lemma 3.2 (b)], one infers that this map admits a limit as $\mu \searrow 0$. As a consequence, the operator valued multiplication operator $N_2: C_c(\mathbb{R}_+; \mathcal{H}) \rightarrow \mathcal{H}$ given by $(N_2 \xi)(\mu) := N_2(\mu) \xi(\mu)$ for $\xi \in C_c(\mathbb{R}_+; \mathcal{H})$ and $\mu \in \mathbb{R}_+$, extends then continuously to an element of $\mathcal{B}(L^2(\mathbb{R}_+; \mathcal{H}), \mathcal{H})$.

Then, by considering $\varphi, \psi \in C_c^\infty(\mathbb{R}_+) \odot C(\mathbb{S})$, we can prove as in [38, Theorem 2.5] that the expressions R_2 given by

$$\int_0^\infty d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \left\langle \mathcal{F}_0(\mu) v S_2 \varrho^\perp(\mu) \frac{1}{\sqrt{\mu}} (\mu - \lambda - i\varepsilon)^{-1} \sqrt{\lambda} \tilde{\varphi}_\varepsilon(\lambda), \psi(\mu) \right\rangle_{\mathfrak{h}}$$

reduces to

$$\langle 2\pi i N_2(\vartheta(A_+) \otimes 1_{\mathcal{H}})(M_2 \otimes 1_{\mathcal{H}})B\varphi, \psi \rangle_{\mathcal{H}}$$

with B defined in (3.4). ■

For the study of the term R_1 we need some preparatory results. For that purpose, let us consider the unitary transformation $\mathcal{U}: L^2(\mathbb{R}_+) \rightarrow L^2(\mathbb{R})$ defined for $f \in L^2(\mathbb{R}_+)$ and $x \in \mathbb{R}$ by

$$[\mathcal{U}f](x) := \sqrt{2} e^{-x} f(e^{-2x}). \quad (3.11)$$

We also introduce the integral operator $\Xi: L^2(\mathbb{R}_+) \rightarrow L^2(\mathbb{R}_+)$ with kernel given by

$$\Xi(\mu, \lambda) := \varrho_0(\mu) \frac{1}{\sqrt{\mu} + \sqrt{\lambda}} \frac{1}{\sqrt{\lambda} \ln(\lambda)} \varrho_0(\lambda). \quad (3.12)$$

We can then express Ξ in terms of dilations and position operators.

Lemma 3.6. *The following equality holds in $L^2(\mathbb{R})$:*

$$\mathcal{U}\Xi\mathcal{U}^* = -\varrho_0(e^{-2X})\frac{2}{1+i2A}\varrho_0(e^{-2X}) + K_1$$

with A the generator of dilation in $L^2(\mathbb{R})$, X the operator by multiplication by the variable in $L^2(\mathbb{R})$, with remainder $K_1 \in \mathcal{K}(L^2(\mathbb{R}))$.

Proof. By a direct computation on any $f \in L^2(\mathbb{R})$ one has

$$\begin{aligned} [\mathcal{U}\Xi\mathcal{U}^*f](x) &= -\varrho_0(e^{-2x}) \int_{\mathbb{R}} \frac{dy}{y} \frac{1}{1+e^{x-y}} \varrho_0(e^{-2y}) f(y) \\ &= -\varrho_0(e^{-2x}) \int_{\mathbb{R}} \frac{dy}{y} \left[\frac{1}{1+e^{x-y}} - \chi_-(x-y) \right] \varrho_0(e^{-2y}) f(y) \end{aligned} \quad (3.13)$$

$$- \varrho_0(e^{-2x}) \int_{\mathbb{R}} \frac{dy}{y} \chi_-(x-y) \varrho_0(e^{-2y}) f(y), \quad (3.14)$$

where χ_- denotes the characteristic function on \mathbb{R}_- . Observe now that the function

$$\mathbb{R} \ni s \mapsto \frac{1}{1+e^s} - \chi_-(s) \in \mathbb{R}$$

belongs to $L^1(\mathbb{R})$. Then, since the function $a: \mathbb{R} \ni x \mapsto \varrho_0(e^{-2x}) \in \mathbb{R}$ belongs to $C_0((-\infty, \infty])$ and since the function $c: \mathbb{R} \ni y \mapsto \frac{1}{y} \varrho_0(e^{-2y}) \in \mathbb{R}$ belongs to $C_0(\mathbb{R})$, one infers that (3.13) defines a compact operator K_1 of the form $a(X)b(D)c(X)$, with $b \in C_0(\mathbb{R})$ and where (X, D) are the canonically conjugate position and momentum operators on $L^2(\mathbb{R})$.

For (3.14), observe that for $x > 0$ one has

$$\begin{aligned} & -\varrho_0(e^{-2x}) \int_{\mathbb{R}} \frac{dy}{y} \chi_-(x-y) \varrho_0(e^{-2y}) f(y) \\ &= -\varrho_0(e^{-2x}) \int_x^\infty \frac{dy}{y} [\varrho_0(e^{-2X})f](y) \\ &= -\varrho_0(e^{-2x}) \int_0^\infty ds [\varrho_0(e^{-2X})f](e^s x) \\ &= -\varrho_0(e^{-2x}) \int_{\mathbb{R}} ds \chi_+(s) e^{-s/2} (U_s[\varrho_0(e^{-2X})f])(x), \end{aligned}$$

where χ_+ denotes the characteristic function of \mathbb{R}_+ and $\{U_s\}_{s \in \mathbb{R}}$ corresponds to the dilation group in $L^2(\mathbb{R})$. Note that we have assumed $x > 0$ in the above computation,

since for $x < 0$ the factor $\varrho_0(e^{-2x})$ already vanishes. Now, since the function $h: \mathbb{R} \ni s \mapsto \chi_+(s) e^{-s/2} \in \mathbb{R}$ belongs to $L^1(\mathbb{R})$, one infers that (3.14) defines an operator of the form $a(X)b(A)a(X)$, with the function a introduced above, with the function b also belonging to $C_0(\mathbb{R})$, and with A the generator of dilation in $L^2(\mathbb{R})$. In fact, the function b can be explicitly computed by taking the Fourier transform of h and using an analogue of (3.5), from which we obtain $b(s) := \frac{2}{1+i2s}$. ■

We now consider the term R_1 .

Lemma 3.7. *The term R_1 can be rewritten as $\langle (N_2 \Xi B + K)\varphi, \psi \rangle_{\mathcal{H}}$ with N_2 introduced in (3.10), Ξ introduced in (3.12), B introduced in (3.4), and $K \in \mathcal{K}(\mathcal{H})$.*

Proof. First, we show that for a dense set of $\varphi, \psi \in \mathcal{H}$ and for a.e. $\lambda \in \mathbb{R}_+$ we have

$$\begin{aligned} & \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \langle \tilde{\varphi}_\varepsilon(\lambda), (\sqrt{\mu} - \sqrt{\lambda})(\mu - \lambda + i\varepsilon)^{-1} N_2(\mu)^* \psi(\mu) \rangle_{\mathcal{H}} \\ &= \int_0^\infty d\mu \left\langle \varrho^\perp(\lambda) S_2(u + vR_0(\lambda + i0)v)^{-1} v \mathcal{F}_0(\lambda)^* \varphi(\lambda), \right. \\ & \quad \left. \frac{1}{\sqrt{\mu} + \sqrt{\lambda}} N_2(\mu)^* \psi(\mu) \right\rangle_{\mathcal{H}} \end{aligned}$$

where $N_2(\mu)$ has been defined in (3.10). This can be obtained by a straightforward application of Lebesgue's dominated convergence theorem by choosing two functions $\varphi, \psi \in C_c(\mathbb{R}_+; \mathfrak{h})$, and by observing that

$$\begin{aligned} & |\langle \tilde{\varphi}_\varepsilon(\lambda), (\sqrt{\mu} - \sqrt{\lambda})(\mu - \lambda + i\varepsilon)^{-1} N_2(\mu)^* \psi(\mu) \rangle_{\mathcal{H}}| \\ & \leq \|\tilde{\varphi}_\varepsilon(\lambda)\|_{\mathcal{H}} \left| \frac{\sqrt{\mu} - \sqrt{\lambda}}{\mu - \lambda + i\varepsilon} \right| \|N_2(\mu)^* \psi(\mu)\|_{\mathcal{H}} \\ & \leq \text{Const.}(\lambda) \frac{1}{\sqrt{\mu} + \sqrt{\lambda}} \| \|N_2(\cdot)^* \psi(\cdot)\|_{\mathcal{H}} \|_\infty \chi_{\text{supp } \psi}(\mu), \end{aligned}$$

which is clearly $L^1(\mathbb{R}_+)$ as a function of μ . Note that we have used the strong convergence of the map $\varepsilon \mapsto \tilde{\varphi}_\varepsilon(\lambda)$ in \mathcal{H} to infer the uniform bound on $\|\tilde{\varphi}_\varepsilon(\lambda)\|_{\mathcal{H}}$.

The rest of the proof is straightforward. Only for the term K it is necessary to observe that this term is compact since $\mathcal{U}^* K_1 \mathcal{U} \otimes S_2 \in \mathcal{K}(L^2(\mathbb{R}_+) \otimes \mathcal{H})$ with $K_1 \in \mathcal{K}(L^2(\mathbb{R}))$ obtained from Lemma 3.6. ■

Let us now consider the final term R_5 , namely the one with the factor S_2^\perp inserted in (3.1). For that purpose, we introduce the map

$$\mathbb{R}_+ \ni \lambda \mapsto N_5(\lambda) := \mathcal{F}_0(\lambda) v S_2^\perp \in \mathcal{B}(\mathcal{H}, \mathfrak{h}), \quad (3.15)$$

which is continuous, admits a limit as $\lambda \searrow 0$, and vanishes as $\lambda \rightarrow \infty$, see for example [40, Lemma 4.8]. The corresponding multiplication operator in $\mathcal{B}(L^2(\mathbb{R}_+; \mathcal{H}), \mathcal{H})$ is denoted by N_5 . Similarly, we define the map

$$\mathbb{R}_+ \ni \lambda \mapsto B_5(\lambda) := S_2^\perp(u + vR_0(\lambda + i0)v)^{-1}v\mathcal{F}_0(\lambda)^* \in \mathcal{B}(\mathfrak{h}, \mathcal{H}).$$

This map is continuous, bounded as $\lambda \searrow 0$ thanks to the expansion provided in Theorem 2.1, and vanishes as $\lambda \rightarrow \infty$. The corresponding multiplication operator in $\mathcal{B}(\mathcal{H}, L^2(\mathbb{R}_+; \mathcal{H}))$ is denoted by B_5 .

Lemma 3.8. *The term*

$$R_5 := \int_{\mathbb{R}} d\lambda \lim_{\varepsilon \searrow 0} \int_0^\infty d\mu \left\langle \mathcal{F}_0(\mu)vS_2^\perp(u + vR_0(\lambda + i\varepsilon)v)^{-1}v\mathcal{F}_0^*\delta_\varepsilon(L - \lambda)\varphi, (\mu - \lambda + i\varepsilon)^{-1}\psi(\mu) \right\rangle_{\mathfrak{h}}$$

can be rewritten as $\langle 2\pi i N_5(\vartheta(A_+) \otimes 1_{\mathcal{H}})B_5\varphi, \psi \rangle_{\mathcal{H}}$.

The proof of this Lemma is very similar to the one of Lemma 3.2 and involves only an application of Lebesgue's dominated convergence theorem, as shown in the proof of [38, Theorem 2.5].

4. Various representations

In this section we provide two new representations for the expressions obtained above. These representations will be useful for considering Levinson's theorem as an index theorem. The first task is to perform some commutations with the function $\vartheta(A_+)$ from (3.6) which has appeared several times, The first statement provides the necessary information for these commutations.

Lemma 4.1. *Let ϑ be the function introduced in (3.6). Then, for $j \in \{0, 2, 4, 5\}$ the following inclusion holds:*

$$N_j(\vartheta(A_+) \otimes 1_{\mathcal{H}}) - (\vartheta(A_+) \otimes 1_{\mathfrak{h}})N_j \in \mathcal{K}(L^2(\mathbb{R}_+; \mathcal{H}), \mathcal{H}),$$

with N_0 introduced in (3.7), N_2 in (3.10), N_4 in (3.9), and N_5 in (3.15).

Proof. This type of result has already been proved in [39, Lemma 2.7] and is based on an argument of Cordes, see for instance [3, Theorem 4.1.10]. The key element is that the functions $\lambda \mapsto N_j(\lambda)$ have limits at 0 and at ∞ , and that the function $s \mapsto \vartheta(s)$ has a limit at $-\infty$ and at ∞ . For ϑ is property is clear, and for N_j these properties have been discussed when these functions have been introduced. ■

For the next statement, recall that the scattering operator is defined by $S := W_+^* W_-$, and that its representation in the spectral representation of H_0 is denoted by $S(L)$, namely $S(L) := \mathcal{F}_0 S \mathcal{F}_0^*$

Corollary 4.2. *The following equality holds:*

$$R_0 + R_2 + R_3 + R_4 + R_5 = -\langle (\vartheta(A_+) \otimes 1_{\mathfrak{h}})(S(L) - 1)\psi \rangle_{\mathcal{H}} + \langle K\varphi, \psi \rangle_{\mathcal{H}},$$

with $K \in \mathcal{K}(\mathcal{H})$.

Proof. Let us firstly observe that for any $\lambda > 0$ one has

$$\begin{aligned} & N_0(\lambda)B_0(\lambda) + N_2(\lambda)B(\lambda) + N_4(\lambda)N_4(\lambda) + N_5(\lambda)B_5(\lambda) \\ &= \mathcal{F}_0(\lambda)v \left[S_2 \varrho(\lambda) + S_2 \varrho^\perp(\lambda) \frac{1}{\sqrt{\lambda}} \cdot \varrho_0(\lambda) \frac{1}{\ln(\lambda)} \cdot \varrho^\perp(\lambda) \sqrt{\lambda} \ln(\lambda) \right. \\ &\quad \left. + S_2 \varrho(\lambda) \frac{1}{\lambda} \cdot \varrho_0(\lambda) \sqrt{\lambda} \frac{1}{\ln(\lambda)} \cdot \varrho^\perp(\lambda) \sqrt{\lambda} \ln(\lambda) + S_2^\perp \right] \\ &\quad \cdot (u + vR_0(\lambda + i0)v)^{-1} v \mathcal{F}_0(\lambda)^* \\ &= \mathcal{F}_0(\lambda)v \left[S_2 (\varrho(\lambda) + \varrho^\perp(\lambda)^2 + \varrho^\perp(\lambda)\varrho(\lambda)) + S_2^\perp \right] \\ &\quad \cdot (u + vR_0(\lambda + i0)v)^{-1} v \mathcal{F}_0(\lambda)^* \\ &= \mathcal{F}_0(\lambda)v (u + vR_0(\lambda + i0)v)^{-1} v \mathcal{F}_0(\lambda)^* \\ &= \frac{1}{-2\pi i} (S(\lambda) - 1) \end{aligned}$$

where $S(\lambda)$ denotes the scattering matrix at energy λ . The last equality can be found for example in [47, Theorem 1.8.1]. It remains then to collect the formulas obtained in Lemmas 3.2, 3.3, 3.4, 3.5, 3.8, together with Lemma 4.1 to obtain the result. ■

Collecting the results obtained so far, we have obtained the equality

$$\mathcal{F}_0(W_- - 1)\mathcal{F}_0^* = \left(\frac{1}{2}(1 - \tanh(\pi A_+)) \otimes 1_{\mathfrak{h}} \right) (S(L) - 1) - N_2 \Xi B + K, \quad (4.1)$$

with $K \in \mathcal{K}(\mathcal{H})$. This equality holds in $\mathcal{H} = L^2(\mathbb{R}_+; \mathfrak{h})$.

As suggested by the analysis of the operator Ξ in Lemma 3.6, we shall firstly look at this equality in the Hilbert space $L^2(\mathbb{R}; \mathfrak{h})$ by using the unitary map \mathcal{U} defined in (3.11). The image of Ξ in this representation has been computed in Lemma 3.6. For any multiplication operator M defined by $\mathbb{R}_+ \ni \lambda \mapsto M(\lambda)$, it is easily seen that for any $\mathfrak{f} \in L^2(\mathbb{R}; \mathfrak{h})$ one has

$$[\mathcal{U}M\mathcal{U}^*\mathfrak{f}](x) = M(e^{-2x})\mathfrak{f}(x) \equiv [M(e^{-2X})\mathfrak{f}](x).$$

Finally, if we consider the dilation group $\{U_t^+\}_{t \in \mathbb{R}}$ we obtain by a straightforward computation that

$$[\mathcal{U}U_t^+\mathcal{U}^*\mathfrak{f}](x) = \mathfrak{f}\left(x - \frac{1}{2}t\right) = [e^{-it\frac{1}{2}D}\mathfrak{f}](x),$$

where $D = -i\frac{d}{dx}$. Summing up this information we find in $L^2(\mathbb{R}; \mathfrak{h})$ the equality

$$\begin{aligned} \mathcal{U}\mathcal{F}_0(W_- - 1)\mathcal{F}_0^*\mathcal{U}^* &= \left(\frac{1}{2}\left(1 - \tanh\left(\frac{\pi}{2}D\right)\right) \otimes 1_{\mathfrak{h}}\right)(\tilde{S}(X) - 1) \\ &\quad + \tilde{N}_2(X)\frac{2}{1 + i2A}\tilde{B}(X) + K, \end{aligned} \quad (4.2)$$

with $K \in \mathcal{K}(L^2(\mathbb{R}, \mathfrak{h}))$ and with the tilde functions given by rescaling the arguments. More precisely, we set $\tilde{S}(X) := S(e^{-2X})$, $\tilde{N}_2(X) := N_2(e^{-2X})$, $\tilde{B}(X) := B(e^{-2X})$. As clearly visible in this formula, the three generators X, D, A are involved in this expression, namely the position operator, the generator of translation, and the generator of dilations. For completeness, let us recall the formal commutations relations between these three generators, namely $[iD, X] = 1$, $[iD, A] = D$ and $[iX, A] = -X$.

It turns out that a C^* -algebra generated by continuous functions of 3 generators has been introduced and studied in [8, Chapter 5]. The algebra is constructed on the Hilbert space $L^2(\mathbb{R}_+)$ while the above expression is taking place on $L^2(\mathbb{R})$. In order to fit into the framework of Cordes, we need to consider one more unitary transformation, namely the decomposition into even and odd functions on $L^2(\mathbb{R})$.

Let us consider $\mathcal{V}: L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}_+; \mathbb{C}^2)$ given by

$$\mathcal{V}\mathfrak{f} := \sqrt{2}\begin{pmatrix} \mathfrak{f}_e \\ \mathfrak{f}_o \end{pmatrix} \quad \text{and} \quad \left[\mathcal{V}^*\begin{pmatrix} \mathfrak{f}_1 \\ \mathfrak{f}_2 \end{pmatrix}\right](x) := \frac{1}{\sqrt{2}}[f_1(|x|) + \text{sgn}(x)f_2(|x|)],$$

for $\mathfrak{f} \in L^2(\mathbb{R})$, $\begin{pmatrix} \mathfrak{f}_1 \\ \mathfrak{f}_2 \end{pmatrix} \in L^2(\mathbb{R}_+; \mathbb{C}^2)$, and $x \in \mathbb{R}$. Here $\mathfrak{f}_e, \mathfrak{f}_o$ denote the even and the odd part of \mathfrak{f} . Then, one observes that if m is a function on \mathbb{R}

$$\mathcal{V}m(X)\mathcal{V}^* = \begin{pmatrix} m_e(L) & m_o(L) \\ m_o(L) & m_e(L) \end{pmatrix} \quad (4.3)$$

while

$$\mathcal{V}m(A)\mathcal{V}^* = \begin{pmatrix} m(A_+) & 0 \\ 0 & m(A_+) \end{pmatrix}. \quad (4.4)$$

In order to consider $\mathcal{V}m(D)\mathcal{V}^*$, let us denote by \mathcal{F}_1 the usual unitary Fourier transform in $L^2(\mathbb{R})$, and let $\mathcal{F}_N, \mathcal{F}_D$ be the unitary cosine and sine transforms on $L^2(\mathbb{R}_+)$, respectively. The subscripts N and D are related to the Neumann Laplacian and the Dirichlet Laplacian in $L^2(\mathbb{R}_+)$, which are diagonalised by \mathcal{F}_N and \mathcal{F}_D , respectively. Note also that these operators correspond to their own inverse. It is then easily checked that

$$\mathcal{V}\mathcal{F}_1\mathcal{V}^* = \begin{pmatrix} \mathcal{F}_N & 0 \\ 0 & i\mathcal{F}_D \end{pmatrix}.$$

In addition, by a straightforward computation one gets

$$\mathcal{V}m(D)\mathcal{V}^* = \mathcal{V}\mathcal{F}_1^*m(X)\mathcal{F}_1\mathcal{V}^* = \begin{pmatrix} \mathcal{F}_N m_e(L)\mathcal{F}_N & -i\mathcal{F}_N m_o(L)\mathcal{F}_D \\ i\mathcal{F}_D m_o(L)\mathcal{F}_N & \mathcal{F}_D m_e(L)\mathcal{F}_D \end{pmatrix}.$$

For the final step, let us recall that the Neumann Laplacian satisfies $-\Delta_N := \mathcal{F}_N L^2 \mathcal{F}_N$, and that

$$i\mathcal{F}_N \mathcal{F}_D = -\tanh(\pi A_+) + i \cosh(\pi A_+)^{-1} =: \phi(A_+).$$

We refer for example to [10, Proposition 4.13] for a proof of the above equality. Then, we end up with

$$\mathcal{V}m(D)\mathcal{V}^* = \mathcal{V}\mathcal{F}_1^*m(X)\mathcal{F}_1\mathcal{V}^* = \begin{pmatrix} m_e(\sqrt{-\Delta_N}) & -m_o(\sqrt{-\Delta_N})\phi(A_+) \\ -\bar{\phi}(A_+)m_o(\sqrt{-\Delta_N}) & \bar{\phi}(A_+)m_e(\sqrt{-\Delta_N})\phi(A_+) \end{pmatrix}. \quad (4.5)$$

Thus, the three equalities (4.3), (4.4), and (4.5) allow us to compute the image of (4.2) into $L^2(\mathbb{R}_+; \mathfrak{h})^2$.

Lemma 4.3. *The expression $\mathcal{V}\mathcal{U}\mathcal{F}_0 W_- \mathcal{F}_0^* \mathcal{U}^* \mathcal{V}^*$ is given by*

$$\begin{aligned} & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 1 & \tanh(\frac{\pi}{2}\sqrt{-\Delta_N})\phi(A_+) \\ \bar{\phi}(A_+)\tanh(\frac{\pi}{2}\sqrt{-\Delta_N}) & 1 \end{pmatrix} \begin{pmatrix} \tilde{\mathcal{S}}_e(L)-1 & \tilde{\mathcal{S}}_o(L) \\ \tilde{\mathcal{S}}_o(L) & \tilde{\mathcal{S}}_e(L)-1 \end{pmatrix} \\ & + \begin{pmatrix} (\tilde{N}_2)_e(L) & (\tilde{N}_2)_o(L) \\ (\tilde{N}_2)_o(L) & (\tilde{N}_2)_e(L) \end{pmatrix} \begin{pmatrix} \frac{2}{1+i2A_+} & 0 \\ 0 & \frac{2}{1+i2A_+} \end{pmatrix} \begin{pmatrix} \tilde{\mathcal{B}}_e(L) & \tilde{\mathcal{B}}_o(L) \\ \tilde{\mathcal{B}}_o(L) & \tilde{\mathcal{B}}_e(L) \end{pmatrix} + K \end{aligned} \quad (4.6)$$

with $K \in \mathcal{K}(L^2(\mathbb{R}_+; \mathfrak{h})^2)$.

Let us now recall the already mentioned construction of Cordes. In [8, Section V.7], the following C^* -subalgebra of $\mathcal{B}(L^2(\mathbb{R}_+))$ is introduced:

$$\mathcal{E} := C^*(a_i(A_+)b_i(L)c_i(-\Delta_N) \mid a_i \in C([-\infty, +\infty]), b_i, c_i \in C([0, +\infty])).$$

It is then shown in [8, Theorem V.7.3] that the quotient algebra $\mathcal{E}/\mathcal{K}(L^2(\mathbb{R}_+))$ is isomorphic to $C(\heptagon)$, the set of continuous functions defined on the edges of a hexagon; see Figure 1. For an operator of the form $a(A_+)b(L)c(\sqrt{-\Delta_N}) \in \mathcal{E}$, its image in the quotient algebra takes the form

$$\Gamma_1(s) := a(s)b(0)c(+\infty), \quad s \in [-\infty, +\infty], \quad (4.7a)$$

$$\Gamma_2(\ell) := a(+\infty)b(\ell)c(+\infty), \quad \ell \in [0, +\infty], \quad (4.7b)$$

$$\Gamma_3(\xi) := a(+\infty)b(+\infty)c(\xi), \quad \xi \in [+\infty, 0], \quad (4.7c)$$

$$\Gamma_4(s) := a(s)b(+\infty)c(0), \quad s \in [+\infty, -\infty], \quad (4.7d)$$

$$\Gamma_5(\xi) := a(-\infty)b(+\infty)c(\xi), \quad \xi \in [0, +\infty], \quad (4.7e)$$

$$\Gamma_6(\ell) := a(-\infty)b(\ell)c(+\infty), \quad \ell \in [+\infty, 0]. \quad (4.7f)$$

Observe that we gave an orientation on the interval on which these functions are defined. As a result, the concatenation map

$$\Gamma \equiv (\Gamma_1, \Gamma_2, \Gamma_3, \Gamma_4, \Gamma_5, \Gamma_6) := \square \rightarrow \mathbb{C}$$

is continuous, even at the vertices of the hexagon.

Our interest in the construction of Cordes comes from the similarity between the elements of \mathcal{E} and the formula (4.6). Indeed, the functions of the three operators L , A_+ , $-\Delta$ are continuous, and have limits either at $-\infty$ and $+\infty$, or at 0 and $+\infty$ (we shall recall these limits below). The only difference is that we have to consider the unital C^* -algebra $(M_2(\mathcal{E}) \otimes \mathcal{K}(\mathfrak{h}))^+$, the 2×2 matrices with values in \mathcal{E} tensor product with the compact operators on \mathfrak{h} , and \mathbb{C} times the identity added. Clearly, this algebra contains the ideal $M_2(\mathcal{K}(L^2(\mathbb{R}_+))) \otimes \mathcal{K}(\mathfrak{h})$, and one has

$$(M_2(\mathcal{E}) \otimes \mathcal{K}(\mathfrak{h}))^+ / M_2(\mathcal{K}(L^2(\mathbb{R}_+))) \otimes \mathcal{K}(\mathfrak{h}) = (M_2(C(\square)) \otimes \mathcal{K}(\mathfrak{h}))^+.$$

One can thus look at the image of (4.6) through the quotient map

$$q := (M_2(\mathcal{E}) \otimes \mathcal{K}(\mathfrak{h}))^+ \rightarrow (M_2(C(\square)) \otimes \mathcal{K}(\mathfrak{h}))^+$$

with kernel $M_2(\mathcal{K}(L^2(\mathbb{R}_+))) \otimes \mathcal{K}(\mathfrak{h})$. In the next statement we provide this image, keeping the convention provided in (4.7) for the enumeration of the 6 components.




Figure 1. Representation of the quotient algebra, with orientation indicated on the edges. The starting point of Γ_1 is located on the lower left corner.

Proposition 4.4. *The operator provided in (4.6) belongs to $(M_2(\mathcal{E}) \otimes \mathcal{K}(\mathfrak{h}))^+$, and its image through the quotient map q consists in the following 6 operator-valued functions:*

$$\begin{aligned}\Gamma_1(s) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2}(S(1) - 1) \begin{pmatrix} 1 & \phi(s) \\ \bar{\phi}(s) & 1 \end{pmatrix}, & s \in [-\infty, +\infty], \\ \Gamma_2(\ell) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2}(S(e^{2\ell}) - 1) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, & \ell \in [0, +\infty], \\ \Gamma_3(\xi) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & \xi \in [+\infty, 0], \\ \Gamma_4(s) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2} \frac{2}{1 + i2s} N_2(0) B(0) \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, & s \in [+\infty, -\infty], \\ \Gamma_5(\xi) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & \xi \in [0, +\infty], \\ \Gamma_6(\ell) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2}(S(e^{-2\ell}) - 1) \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, & \ell \in [+\infty, 0].\end{aligned}$$

Proof. When the various factors of (4.6) were introduced, their continuity properties and the existence of their limits at endpoints have been discussed. The only missing information is about $S(\lambda) - 1$. It is known that for any $\lambda > 0$, one has $S(\lambda) - 1 \in \mathcal{K}(\mathfrak{h})$, and that the map $\lambda \mapsto S(\lambda) - 1$ is continuous, see for example [47, Proposition 8.1.5]. In addition, it has been shown in [40, Theorem 1.1] that $\lim_{\lambda \searrow 0} S(\lambda) = 1$. Since $\lim_{\lambda \rightarrow \infty} S(\lambda) = 1$, with the limit taken in $\mathcal{B}(\mathfrak{h})$, the function $\lambda \mapsto S(\lambda) - 1$ belongs to $C_0(\mathbb{R}_+, \mathcal{K}(\mathfrak{h}))$. By inspection of the various factors, one can now infer that the operator provided in (4.6) belongs to $(M_2(\mathcal{E}) \otimes \mathcal{K}(\mathfrak{h}))^+$.

Let us move to the image of this operator in the quotient algebra. By using the formulas proposed in (4.7), the computations are rather straightforward. For Γ_1 , it is necessary to observe that $\tilde{S}_e(0) = S(1)$ while $\tilde{S}_o(0) = 0$. In addition, we have $\lim_{\xi \rightarrow +\infty} \tanh(\frac{\pi}{2}\xi) = 1$. Because of the localization function ϱ^\perp , one also observes that $\tilde{N}_2(0) = 0$ and $\tilde{B}(0) = 0$. For Γ_2 , note that $\lim_{s \rightarrow +\infty} \phi(s) = -1$, and then the expression

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} \tilde{S}_e(\ell) - 1 & \tilde{S}_o(\ell) \\ \tilde{S}_o(\ell) & \tilde{S}_e(\ell) - 1 \end{pmatrix}$$

leads directly to the result. The computation for Γ_6 is very similar, once the equality $\lim_{s \rightarrow -\infty} \phi(s) = 1$ is taken into account. For Γ_3 and for Γ_5 , it is sufficient to remember that $\lim_{\lambda \searrow 0} S(\lambda) = 1$ and that $\lim_{\lambda \rightarrow \infty} S(\lambda) = 1$. These equalities imply that $\lim_{\lambda \rightarrow \pm\infty} \tilde{S}_e(\ell) = 1$ while $\lim_{\lambda \rightarrow \pm\infty} \tilde{S}_o(\ell) = 0$, with these limits taken in $\mathcal{B}(\mathfrak{h})$. Finally, for Γ_4 , it is necessary to observe that $\lim_{\ell \rightarrow \infty} \tilde{N}_2(\ell) = N_2(0)$, $\lim_{\ell \rightarrow \infty} \tilde{B}(\ell) = B(0)$, and then we have

$$\lim_{\ell \rightarrow \infty} (\tilde{N}_2)_e(\ell) = \frac{1}{2} N_2(0) = \lim_{\ell \rightarrow \infty} (\tilde{N}_2)_o(\ell),$$

and

$$\lim_{\ell \rightarrow \infty} (\tilde{B})_e(\ell) = \frac{1}{2}B(0) = \lim_{\ell \rightarrow \infty} (\tilde{B})_o(\ell).$$

This leads us directly to the statement. ■

In order to fully exploit the previous result, it remains to compute the terms appearing in Γ_4 . So, we first determine the expression $B(0) := \lim_{\lambda \searrow 0} B(\lambda)$ explicitly by using the results of [21, Theorem 6.1 and Theorem 6.2]. We denote by X_j the multiplication operator by x_j in $L^2(\mathbb{R}^2)$, and recall from [21, Theorem 6.2 (i)] that $\text{Ran}(T_3)$ is spanned by

$$Q_j = S_2 X_j v$$

for $j = 1, 2$. Note that one or both of the Q_j may vanish or they may be linearly dependent, in which case the dimension of $\text{Ran}(T_3)$ is strictly smaller than 2.

Lemma 4.5. *The following equality holds:*

$$\begin{aligned} B(0) &= 2(T_3 - S_3 d(0)^{-1} c(0)) \left(-\frac{1}{4\pi} \sum_{j=1}^2 \langle Q_j, \cdot \rangle Q_j \right)^{-1} T_3 v \gamma_1^* \\ &= 2(T_3 - S_3 d(0)^{-1} c(0)) \left(-\frac{1}{4\pi} \sum_{j=1}^2 |Q_j\rangle \langle Q_j| \right)^{-1} T_3 v \gamma_1^*, \end{aligned}$$

where the standard bra-ket notation has been introduced for the last expression.

Proof. We begin by recalling some facts about the function $g(\kappa)$ and the operator $m(\kappa)$. Firstly, by [21, equation (6.30)] the scalar-valued function g satisfies

$$g(\kappa) = \eta^{-1} \left(-\frac{\|v\|^2}{2\pi} + \eta h(\kappa) \right)$$

where h is bounded near zero. Secondly, by [21, equation (6.42)], the operator valued function m satisfies

$$m(\kappa) = \eta^{-1} \frac{\|v\|^2}{8\pi^2} \sum_{j=1}^d \langle Q_j, \cdot \rangle Q_j + f(\kappa)$$

where f is bounded. Then one may write

$$\begin{aligned} g(\kappa) m(\kappa)^{-1} &= g(\kappa) \left(\eta^{-1} \frac{\|v\|^2}{8\pi^2} \sum_{j=1}^2 \langle Q_j, \cdot \rangle Q_j + f(\kappa) \right)^{-1} \\ &= \left(\eta^{-1} g(\kappa)^{-1} \frac{\|v\|^2}{8\pi^2} \sum_{j=1}^2 \langle Q_j, \cdot \rangle Q_j + g(\kappa)^{-1} f(\kappa) \right)^{-1}, \end{aligned}$$

and observe that

$$\lim_{\kappa \rightarrow 0} g(\kappa)^{-1} f(\kappa) = \lim_{\kappa \rightarrow 0} \eta \left(-\frac{\|v\|^2}{2\pi} + \eta h(\kappa) \right)^{-1} f(\kappa) = 0,$$

since f is bounded and $\eta \rightarrow 0$ as $\kappa \rightarrow 0$. We also have the limit

$$\lim_{\kappa \rightarrow 0} \eta^{-1} g(\kappa)^{-1} = \lim_{\kappa \rightarrow 0} \left(-\frac{\|v\|^2}{2\pi} + \eta h(\kappa) \right)^{-1} = -\frac{2\pi}{\|v\|^2},$$

since h is bounded and $\eta \rightarrow 0$ as $\kappa \rightarrow 0$. Thus, we find

$$\begin{aligned} \lim_{\kappa \rightarrow 0} g(\kappa) m(\kappa)^{-1} &= \lim_{\kappa \rightarrow 0} \left(\eta^{-1} g(\kappa)^{-1} \frac{\|v\|^2}{8\pi^2} \sum_{j=1}^2 \langle Q_j, \cdot \rangle Q_j + g(\kappa)^{-1} f(\kappa) \right)^{-1} \\ &= \left(-\frac{1}{4\pi} \sum_{j=1}^2 \langle Q_j, \cdot \rangle Q_j \right)^{-1}. \end{aligned}$$

It finally remains to insert the expansion (2.7) into the expression for $B(\lambda)$ and we obtain

$$\begin{aligned} B(0) &= \lim_{\lambda \searrow 0} B(\lambda) = \lim_{\lambda \searrow 0} \varrho^\perp(\lambda) \sqrt{\lambda} \ln(\lambda) S_2 [(u + vR_0(\lambda + i0)v)^{-1} v\mathcal{F}_0(\lambda)^*] \\ &= \lim_{\lambda \searrow 0} \ln(\lambda) \eta S_2 (T_3 - S_3 d(\kappa)^{-1} c(\kappa)) g(\kappa) m(\kappa)^{-1} T_3 v \gamma_1^* \\ &= 2(T_3 - S_3 d(0)^{-1} c(0)) \left(-\frac{1}{4\pi} \sum_{j=1}^2 \langle Q_j, \cdot \rangle Q_j \right)^{-1} T_3 v \gamma_1^* \end{aligned}$$

which leads to the claim. ■

Let us compute still more explicitly these expressions. Recall firstly that

$$[\gamma_1 f](\omega) := \frac{-i}{2^{3/2}\pi} \int_{\mathbb{R}^2} dx(\omega \cdot x) f(x)$$

for suitable $f \in L^2(\mathbb{R}^2)$. Let us also define $\xi_{\pm 1} \in \mathfrak{h}$ with $\xi_{\pm 1}(\theta) := \frac{1}{\sqrt{2\pi}} e^{\pm i\theta}$ and $\|\xi_{\pm 1}\|_{\mathfrak{h}} = 1$. As a consequence, for any $\tau \in \mathfrak{h}$ one has

$$\begin{aligned} [\gamma_1^* \tau](x) &= \frac{i}{2^{3/2}\pi} \int_{\mathbb{S}} d\omega(x \cdot \omega) \tau(\omega) \\ &= \frac{i}{2^{3/2}\pi} \int_0^{2\pi} d\theta (x_1 \cos(\theta) + x_2 \sin(\theta)) \tau((\cos(\theta), \sin(\theta))) \end{aligned}$$

$$\begin{aligned}
&= \frac{i}{4\sqrt{\pi}} \int_0^{2\pi} d\theta [x_1(\xi_1(\theta) + \xi_{-1}(\theta)) - ix_2(\xi_1(\theta) - \xi_{-1}(\theta))] \\
&\quad \cdot \tau((\cos(\theta), \sin(\theta))) \\
&= \frac{i}{4\sqrt{\pi}} [x_1(\langle \xi_{-1} | + \langle \xi_1 |) - ix_2(\langle \xi_{-1} | - \langle \xi_1 |)] \tau,
\end{aligned}$$

where $\langle \xi_{\pm 1} | \tau := \int_0^{2\pi} d\theta \overline{\xi_{\pm 1}(\theta)} \tau((\cos(\theta), \sin(\theta)))$. Since $S_3 v \gamma_1^* = 0$, one infers that

$$\begin{aligned}
T_3 v \gamma_1^* &= S_2 v \gamma_1^* = \frac{i}{4\sqrt{\pi}} S_2 v [X_1(\langle \xi_{-1} | + \langle \xi_1 |) - iX_2(\langle \xi_{-1} | - \langle \xi_1 |)] \\
&= \frac{i}{4\sqrt{\pi}} [|Q_1\rangle \langle \xi_{-1} + \xi_1 | - i|Q_2\rangle \langle \xi_{-1} - \xi_1 |] \\
&= \frac{i}{4\sqrt{\pi}} [|Q_1 - iQ_2\rangle \langle \xi_{-1} | + |Q_1 + iQ_2\rangle \langle \xi_1 |]. \tag{4.8}
\end{aligned}$$

With these expressions at hand, we can finally compute the expression for term Γ_4 of Proposition 4.4. For this, we define an orthogonal projection P_p as follows.

- If $\dim(T_3) = 0$, then

$$P_p := 0;$$

- if $\dim(T_3) = 1$, then

- if $Q_1 = 0$,

$$P_p := \left| \frac{1}{\sqrt{2}}(\xi_{-1} - \xi_1) \right\rangle \left\langle \frac{1}{\sqrt{2}}(\xi_{-1} - \xi_1) \right|, \tag{4.9a}$$

- if $Q_2 = 0$,

$$P_p := \left| \frac{1}{\sqrt{2}}(\xi_{-1} + \xi_1) \right\rangle \left\langle \frac{1}{\sqrt{2}}(\xi_{-1} + \xi_1) \right|, \tag{4.9b}$$

- if $Q_2 = \alpha Q_1$,

$$\begin{aligned}
P_p &:= \left| \frac{1}{\sqrt{2(1 + |\alpha|^2)}}((1 + i\bar{\alpha})\xi_{-1} + (1 - i\bar{\alpha})\xi_1) \right\rangle \\
&\quad \cdot \left\langle \frac{1}{\sqrt{2(1 + |\alpha|^2)}}((1 + i\bar{\alpha})\xi_{-1} + (1 - i\bar{\alpha})\xi_1) \right|; \tag{4.9c}
\end{aligned}$$

- if $\dim(T_3) = 2$, then

$$P_p := (|\xi_{-1}\rangle \langle \xi_{-1}| + |\xi_1\rangle \langle \xi_1|).$$

Remark 4.6. Observe that the projection P_p corresponds to an orthogonal projection in $L^2(\mathbb{S})$, but the exact form of this projection is independent of V . The V -dependence appears only in the conditions $Q_1 = 0$, $Q_2 = 0$, or $Q_2 = \alpha Q_1$. A quite similar feature was already observed for Schrödinger operators in \mathbb{R}^3 , for which a 0-energy

resonance affects the 0-energy scattering matrix $S(0)$ only in the subspace of spherically symmetric functions of $L^2(\mathbb{S}^2)$, see [20, Theorems 5.2 and 5.4]. In particular, let us stress that these projections are not linked to any symmetry of the potential V , since no such requirement is imposed.

Lemma 4.7. *For any $s \in [+\infty, -\infty]$, one has*

$$\Gamma_4(s) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \frac{1}{2} \frac{2}{1+i2s} P_p \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 - \frac{1}{1+i2s} P_p & -\frac{1}{1+i2s} P_p \\ -\frac{1}{1+i2s} P_p & 1 - \frac{1}{1+i2s} P_p \end{pmatrix}.$$

Proof. First of all, observe that

$$\begin{aligned} N_2(0)B(0) &= \gamma_1 v S_2 B(0) \\ &= 2\gamma_1 v (T_3 + S_3)(T_3 - S_3 d(0)^{-1} c(0)) \left(-\frac{1}{4\pi} \sum_{j=1}^2 |Q_j\rangle\langle Q_j| \right)^{-1} T_3 v \gamma_1^* \\ &= 2\gamma_1 v T_3 \left(-\frac{1}{4\pi} \sum_{j=1}^2 |Q_j\rangle\langle Q_j| \right)^{-1} T_3 v \gamma_1^* \end{aligned}$$

where the algebraic equality $\gamma_1 v S_3 = 0$ has been taken into account, see [40, Lemma 3.2(c)].

Then, by using the expression (4.8) for $T_3 v \gamma_1^*$ and for its adjoint, one infers that the following equality holds:

$$\begin{aligned} &2\gamma_1 v T_3 \left(-\frac{1}{4\pi} \sum_{j=1}^2 |Q_j\rangle\langle Q_j| \right)^{-1} T_3 v \gamma_1^* \\ &= -\frac{1}{2} [|\xi_{-1}\rangle\langle Q_1 - iQ_2| + |\xi_1\rangle\langle Q_1 + iQ_2|] T_3 \left(\sum_{j=1}^2 |Q_j\rangle\langle Q_j| \right)^{-1} T_3 \\ &\quad \cdot [|\xi_{-1}\rangle\langle Q_1 - iQ_2| + |\xi_1\rangle\langle Q_1 + iQ_2|]. \end{aligned} \tag{4.10}$$

By a direct computation, one gets that (4.10) is equal to $-P_p$, as defined above. Note that for the case $\dim(T_3) = 2$, we have used a convenient result due to Parra about the inversion of a matrix on its range. This statement and its proof are gathered in Appendix A.

It only remains to observe that

$$\Gamma_4(s) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2} \frac{2}{1+i2s} N_2(0)B(0) \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \frac{1}{2} \frac{2}{1+i2s} P_p \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix},$$

which gives us the statement. ■

5. Topological Levinson's theorem

In this section, we briefly recall the C^* -algebraic framework leading to a topological version of Levinson's theorem, and show that our current investigations fit into this framework. We refer to the survey paper [36] for additional information on this program and for the presentation of several examples.

It has already been shown that the unital C^* -algebra $(M_2(\mathcal{E}) \otimes \mathcal{K}(\mathfrak{h}))^+$ plays the important role of containing the wave operator W_- , once suitable unitary conjugations are applied. In addition, this algebra contains the ideal $M_2(\mathcal{K}(L^2(\mathbb{R}_+))) \otimes \mathcal{K}(\mathfrak{h})$ which is nothing but the algebra $\mathcal{K}(L^2(\mathbb{R}_+; \mathfrak{h})^2)$ of compact operators on this Hilbert space. Then, as a consequence of Cordes' result, one has the short exact sequence of C^* -algebras

$$0 \rightarrow \mathcal{K}(L^2(\mathbb{R}_+; \mathfrak{h})^2) \rightarrow (M_2(\mathcal{E}) \otimes \mathcal{K}(\mathfrak{h}))^+ \xrightarrow{q} (M_2(C(\circ)) \otimes \mathcal{K}(\mathfrak{h}))^+ \rightarrow 0$$

and the corresponding 6 terms exact sequence for the K -theory of these algebras. In particular, one has $K_0(\mathcal{K}(L^2(\mathbb{R}_+; \mathfrak{h})^2)) \cong \mathbb{Z}$ and $K_1((M_2(C(\circ)) \otimes \mathcal{K}(\mathfrak{h}))^+) \cong \mathbb{Z}$.

Since the operator-valued function $\Gamma = (\Gamma_1, \Gamma_2, \Gamma_3, \Gamma_4, \Gamma_5, \Gamma_6)$ exhibited in Proposition 4.4 belongs to $(M_2(C(\circ)) \otimes \mathcal{K}(\mathfrak{h}))^+$ and is invertible, it defines an element $[\Gamma]_1$ in the K_1 -group of this algebra. In addition, since $W_- \in (M_2(\mathcal{E}) \otimes \mathcal{K}(\mathfrak{h}))^+$ is an isometry and a lift for Γ , one directly infers from [41, Proposition 9.2.4.(ii)] that

$$\text{ind}([\Gamma]_1) = [1 - W_-^* W_-]_0 - [1 - W_- W_-^*]_0 = -[E_p(H)]_0, \quad (5.1)$$

with $E_p(H)$ the projection on the subspace spanned by the eigenfunctions of H . Let us emphasise that the equality (5.1) corresponds to the topological version of Levinson's theorem: it is a relation (by the index map) between the equivalence class in K_1 of quantities related to scattering theory, and the equivalence class in K_0 of the projection on the bound states of H . Note that the operator Γ contains the scattering operator in its components Γ_2 and Γ_6 , but also a new contribution related to p -resonance in its component Γ_4 .

The standard formulation of Levinson's theorem is an equality between numbers. Thus, our last task is to extract a numerical equality from (5.1). In a more general setting, we might pair the K_1 class of the scattering matrix with the Chern character of a suitable spectral triple, as in [2]. For this specific case, we proceed in a more elementary way by using the determinant and winding number directly. Thus, on $\mathcal{K}(L^2(\mathbb{R}_+; \mathfrak{h})^2)$, one uses the usual trace (on finite-dimensional projections), and on

$$(M_2(C(\circ)) \otimes \mathcal{K}(\mathfrak{h}))^+ \cong (C(\circ; M_2(\mathcal{K}(\mathfrak{h}))))^+$$

the winding number of the pointwise determinant is the correct notion to be used.

Remark 5.1. When computing the winding number, and pairing the equality (5.1) with traces, a few conventions about signs have to be taken. As introduced in [36, Section 2], we shall turn around the hexagon clockwise, and the increase in the winding number is also counted clockwise. The convention about the path is illustrated in Figure 1, with the starting point of Γ_1 located on the lower left corner. With this convention, the multiplicative factor n , which relates the winding number computed on Γ and the trace applied to $-E_p(H)$, is equal to -1 , see [36, Theorem 4.4] for the details.

For the computation of the pointwise determinant of the components of Γ , let us recall from [47, Corollary 8.1.7] that $S(\lambda) - 1$ is trace class, and that the map $\lambda \mapsto \det(S(\lambda))$ is continuous. Then, based on the following lemma, it will be possible to get simpler expressions for Γ_1 , Γ_2 , and Γ_6 .

Lemma 5.2. *Let \mathcal{H} be a complex Hilbert space and let c be a complex number with $|c| = 1$. For a unitary operator $U \in \mathcal{B}(\mathcal{H})$ with $U - 1$ trace class, define the operator $B \in \mathcal{B}(\mathcal{H} \oplus \mathcal{H})$ by*

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2}(U - 1) \begin{pmatrix} 1 & c \\ \bar{c} & 1 \end{pmatrix}.$$

Then $\sigma(B) \setminus \{1\} = \sigma(U) \setminus \{1\}$, multiplicity counted, and $\det(U) = \det(B)$.

Proof. Let λ be an eigenvalue of B , with non zero eigenvector $\begin{pmatrix} \xi \\ \eta \end{pmatrix}$, namely $B \begin{pmatrix} \xi \\ \eta \end{pmatrix} = \lambda \begin{pmatrix} \xi \\ \eta \end{pmatrix}$. This equation is equivalent to the two equations

$$\frac{1}{2}(U - 1)(\xi + c\eta) = (\lambda - 1)\xi, \quad \frac{1}{2}(U - 1)(\bar{c}\xi + \eta) = (\lambda - 1)\eta.$$

By multiplying the second line by c , we infer the relation $(\lambda - 1)\xi = (\lambda - 1)c\eta$. For $\lambda \neq 1$, it follows that $\eta = \bar{c}\xi$. By inserting this in the first equation, we get

$$\frac{1}{2}(U - 1)(\xi + c\eta) = (U - 1)\xi = (\lambda - 1)\xi,$$

implying that $U\xi = \lambda\xi$. Note that $\xi \neq 0$, otherwise the eigenvector of B would be the 0 vector.

Conversely, if $\xi \neq 0$ satisfies $U\xi = \lambda\xi$, then one easily checks that the vector $\begin{pmatrix} \xi \\ \bar{c}\xi \end{pmatrix}$ is an eigenvector of B associated with the eigenvalue λ . ■

Corollary 5.3. *One has*

$$\begin{aligned} \det(\Gamma_1(s)) &= \det(S(1)), \\ \det(\Gamma_2(\ell)) &= \det(S(e^{2\ell})), \\ \det(\Gamma_6(\ell)) &= \det(S(e^{-2\ell})). \end{aligned}$$

Since one trivially gets $\det(\Gamma_3(\xi)) = 1$ for any $\xi \in [+\infty, 0]$, and $\det(\Gamma_5(\xi)) = 1$ for any $\xi \in [0, +\infty]$, it only remains to compute $\det(\Gamma_4(s))$ for $s \in [+\infty, -\infty]$. However, based on the content of Lemma 4.7 and since P_p is a finite-dimensional projection, this computation is easy. By using again Lemma 5.2, one infers that

$$\det(\Gamma_4(s)) = \left(\frac{i2s - 1}{i2s + 1} \right)^{\dim(P_p)}. \quad (5.2)$$

Before the explicit computation of the winding number of the pointwise determinant, it is useful to divide the computation of the Fredholm index of W_- into two contributions. For that purpose, we define the operator $W_S \in \mathcal{B}(\mathcal{H})$ by the equality

$$W_S - 1 := \left(\frac{1}{2} \left(1 - \tanh(\pi A_+) \right) \otimes 1_{\mathfrak{h}} \right) (S(L) - 1). \quad (5.3)$$

We then directly obtain its main properties.

Lemma 5.4. *The operator W_S is a Fredholm operator.*

Proof. It is sufficient to observe that the operator W_{S^*} defines an inverse for W_S , up to compact operators. Indeed, this can be easily checked by firstly recalling that $[\vartheta(A_+) \otimes 1_{\mathfrak{h}}, S(L)] \in \mathcal{K}(L^2(\mathbb{R}_+; \mathfrak{h}))$, with ϑ defined in (3.6). In addition, since $S(0) = \lim_{\lambda \rightarrow \infty} S(\infty) = 1$ and since $\vartheta - \vartheta^2$ vanishes at $\pm\infty$, operators of the form

$$(\vartheta(A_+) \otimes 1_{\mathfrak{h}} - \vartheta^2(A_+) \otimes 1_{\mathfrak{h}})(S(L) - 1)$$

or

$$(\vartheta(A_+) \otimes 1_{\mathfrak{h}} - \vartheta^2(A_+) \otimes 1_{\mathfrak{h}})(S^*(L) - 1)$$

belong to $\mathcal{K}(L^2(\mathbb{R}_+; \mathfrak{h}))$. ■

For a Fredholm operator W , let us denote by $\text{Index}(W)$ its Fredholm index. Then the following statement holds.

Proposition 5.5. *If V satisfies (2.3) with $\rho > 11$, then the following equality holds:*

$$\text{Index}(W_S) + \dim(P_p) = -\#\sigma_p(H).$$

Proof. Let W_S be as in (5.3) and define $W_p := 1 - N_2 \otimes B$. It follows from (4.1) that

$$\mathcal{F}_0 W_- \mathcal{F}_0^* = W_S + (W_p - 1) + K$$

for a compact operator K . By construction, both the operator $\mathcal{V}UW_S\mathcal{U}^*\mathcal{V}^*$ and the operator $\mathcal{V}U(W_p - 1)\mathcal{U}^*\mathcal{V}^*$ belong to $(M_2(\mathcal{E}) \otimes \mathcal{K}(\mathfrak{h}))^+$. For $j \in \{1, 2, 3, 4, 5, 6\}$ let $\Gamma_{S,j}$ and $\Gamma_{p,j}$ respectively denote the components of the images $q(\mathcal{V}UW_S\mathcal{U}^*\mathcal{V}^*)$

and $q(\mathcal{V}\mathcal{U}(W_p - 1)\mathcal{U}^*\mathcal{V}^*)$ in the quotient algebra. Then a proof similar to Proposition 4.4 and Lemma 4.7 leads to

$$\begin{aligned}\Gamma_{S,1}(s) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2}(S(1) - 1) \begin{pmatrix} 1 & \phi(s) \\ \bar{\phi}(s) & 1 \end{pmatrix}, & s \in [-\infty, +\infty], \\ \Gamma_{S,2}(\ell) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2}(S(e^{2\ell}) - 1) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, & \ell \in [0, +\infty], \\ \Gamma_{S,3}(\xi) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & \xi \in [+ \infty, 0], \\ \Gamma_{S,4}(s) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & s \in [+ \infty, -\infty], \\ \Gamma_{S,5}(\xi) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & \xi \in [0, +\infty], \\ \Gamma_{S,6}(\ell) &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{1}{2}(S(e^{-2\ell}) - 1) \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, & \ell \in [+ \infty, 0],\end{aligned}$$

and to

$$\begin{aligned}\Gamma_{p,j} &:= \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, & j \in \{1, 2, 3, 5, 6\}, \\ \Gamma_{p,4}(s) &:= -\frac{1}{2} \frac{2}{1 + i2s} P_p \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, & s \in [+ \infty, -\infty].\end{aligned}$$

Explicit computation shows that $\Gamma_{S,j}^* \Gamma_{p,j} = \Gamma_{p,j} = 0$ for $j \neq 4$ and $\Gamma_{S,4}^* \Gamma_{p,4} = \Gamma_{p,4}$. Thus, we find $q(\mathcal{V}\mathcal{U}W_S^*(W_p - 1)\mathcal{U}^*\mathcal{V}^*) = q(\mathcal{V}\mathcal{U}(W_p - 1)\mathcal{U}^*\mathcal{V}^*)$. Since their image under the quotient map agrees, we have $1 + W_S^*(W_p - 1) = W_p + K$ for some compact operator K . Now, we note the equalities

$$\mathcal{F}_0 W_- \mathcal{F}_0^* = W_S + (W_p - 1) + K = W_S(1 + W_S^*(W_p - 1)) + K',$$

for some compact operators K and K' , which lead to

$$\text{Index}(W_-) = \text{Index}(W_S) + \text{Index}(1 + W_S^*(W_p - 1)) = \text{Index}(W_S) + \text{Index}(W_p).$$

Clearly, one has $\text{Index}(W_-) = -\#\sigma_p(H)$. On the other hand, the value $\text{Index}(W_p)$ can be computed with the winding number of the pointwise determinant of $1 + \Gamma_{p,4}$, as mentioned in Remark 5.1. More precisely, one has

$$\text{Index}(W_p) = -\text{Wind}(\det(1 + \Gamma_{p,4})) = -\text{Wind}(\det(\Gamma_4))$$

with $\det(\Gamma_4)$ provided in (5.2). However, since $s \mapsto \det(\Gamma_4(s))$ has to be computed from $+\infty$ to $-\infty$, and that on this path the increase of the argument is anti-clockwise, one gets $\text{Index}(W_p) = \dim(P_p)$, leading directly to the statement. ■

Our next aim is to compute $\text{Index}(W_-)$ in terms of Γ_S , as introduced in the proof of Proposition 5.5. Due to the high energy behaviour of the scattering matrix, some regularization is necessary to obtain an analytic formula. Indeed, even though the map

$\lambda \mapsto S(\lambda)$ converges to 1 in the norm on $\mathcal{B}(\mathfrak{h})$ as $\lambda \rightarrow \infty$, the map $\lambda \mapsto \det(S(\lambda))$ does not converge to 1 as $\lambda \rightarrow \infty$. A more precise statement is provided in Lemma 5.9.

For $\lambda \in \mathbb{R}_+$, we define the self-adjoint operator $A(\lambda)$ in $\mathcal{B}(\mathfrak{h})$ by

$$A(\lambda) = 4 \tan^{-1}(\lambda) \mathcal{F}_0(\lambda) V \mathcal{F}_0(\lambda)^*.$$

The main properties of this operator are gathered in the following statement.

Lemma 5.6. *For each $\lambda \in \mathbb{R}_+$, the operator $A(\lambda)$ is self-adjoint and trace class.*

Proof. Since V is real-valued, the self-adjointness property is clear. Based on the definition of \mathcal{F}_0 given in (2.1), we can write explicitly the integral kernel of $A(\lambda)$ as

$$A(\lambda, \omega, \omega') = 2 \tan^{-1}(\lambda) (2\pi)^{-2} \int_{\mathbb{R}^2} e^{-i\sqrt{\lambda}(\omega-\omega') \cdot x} V(x) dx.$$

Integrating along the diagonal shows that the trace of $A(\lambda)$ is

$$\mathrm{tr}(A(\lambda)) = \frac{1}{\pi} \tan^{-1}(\lambda) \int_{\mathbb{R}^2} V(x) dx. \quad (5.4)$$

The computation is justified by writing

$$4 \tan^{-1}(\lambda) \mathcal{F}_0(\lambda) v u v \mathcal{F}_0(\lambda)^* \quad (5.5)$$

with u, v introduced in (2.5), and by observing that (5.5) contains two factors which are Hilbert–Schmidt. ■

By the properties of the map $\lambda \mapsto \mathcal{F}_0(\lambda)$ exhibited in [40, Lemma 4.8], one infers that the operator-valued map $\lambda \mapsto A(\lambda) \in \mathcal{B}(\mathfrak{h})$ is continuous and has norm limits $\lim_{\lambda \searrow 0} A(\lambda) = 0$ and $\lim_{\lambda \rightarrow \infty} A(\lambda) = 0$. As a consequence of (5.4), one also observes that the map $\lambda \mapsto \mathrm{tr}(A(\lambda))$ is continuous on \mathbb{R}_+ and satisfies $\lim_{\lambda \searrow 0} \mathrm{tr}(A(\lambda)) = 0$ and $\lim_{\lambda \rightarrow \infty} \mathrm{tr}(A(\lambda)) = \frac{1}{2} \int_{\mathbb{R}^2} V(x) dx$.

Based on these observations, let us now define the unitary operator in $\mathcal{B}(\mathfrak{h})$

$$\beta(\lambda) := \exp(iA(\lambda))$$

which clearly satisfies $\det(\beta(\lambda)) = e^{i \mathrm{tr}(A(\lambda))}$ for all $\lambda \in \mathbb{R}_+$. We also define the operator $W_\beta \in \mathcal{B}(\mathcal{H})$ by the equality

$$W_\beta - 1 = \left(\frac{1}{2} \left(1 - \tanh(\pi A_+) \right) \otimes 1_{\mathfrak{h}} \right) (\beta(L) - 1).$$

Our main interest for this operator is related to the properties shown in the next statement.

Lemma 5.7. *The operator W_β is a Fredholm operator satisfying $\text{Index}(W_\beta) = 0$.*

Proof. Observe firstly that we have $\lim_{\lambda \searrow 0} \beta(\lambda) = 1$ and that $\lim_{\lambda \rightarrow \infty} \beta(\lambda) = 1$, both limits in the norm sense. Thus, by the same argument provided in the proof of Lemma 5.4 one gets that the operator W_{β^*} defines an inverse for W_β , up to compact operators. It directly follows that W_β is a Fredholm operator.

It remains to show that $\text{Index}(W_\beta) = 0$. To see this we consider, for fixed $\lambda \in \mathbb{R}_+$, the map $[0, 1] \ni t \mapsto A_t(\lambda)$ with $A_t(\lambda)$ defined by

$$A_t(\lambda) = 4 \tan^{-1}((1-t)\lambda) \mathcal{F}_0(\lambda) V \mathcal{F}_0(\lambda)^*.$$

The map $A_t(\lambda)$ defines a norm continuous path in $\mathcal{B}(\mathfrak{h})$ from $A(\lambda)$ to 0. Defining the path $A_t = A_t(L) \in \mathcal{B}(\mathcal{H})$ we then obtain a norm continuous path in $\mathcal{B}(\mathcal{H})$ from A to 0. As a consequence, $\beta_t = \exp(iA_t)$ defines a norm continuous path of unitary operators in $\mathcal{B}(\mathcal{H})$ from β to 1. Hence, the path W_{β_t} defines a norm continuous path in $\mathcal{B}(\mathcal{H})$ from W_β to the identity, along which the Fredholm index is constant, and so equal to 0. ■

Lemma 5.8. *The Fredholm operators W_S and W_β satisfy $W_S W_\beta - W_{S\beta} \in \mathcal{K}(\mathcal{H})$ and*

$$\text{Index}(W_S) = \text{Index}(W_{S\beta}).$$

Proof. The equality $W_S W_\beta = W_{S\beta}$ up to compact operators follows from one more commutator computation as provided in the proof of Lemmas 5.4 and 5.7. The index claim follows from the fact that $\text{Index}(W_\beta) = 0$ and the composition rule for Fredholm index. ■

The next statement shows that the operator β provides the correct regularization for the operator S , and consequently W_β will provide the correct regularization to the operator W_S . The proof is using some properties of the spectral shift function developed in [47, Chapter 9].

Lemma 5.9. *The map $\lambda \mapsto \det(S(\lambda)) \det(\beta(\lambda))$ satisfies*

$$\lim_{\lambda \searrow 0} \det(S(\lambda)) \det(\beta(\lambda)) = 1$$

and

$$\lim_{\lambda \rightarrow \infty} \det(S(\lambda)) \det(\beta(\lambda)) = 1.$$

Proof. Let us firstly recall the Birman–Kreĭn formula linking the scattering operator and the spectral shift function, namely $\det(S(\lambda)) = e^{-2\pi i \xi(\lambda)}$. By [47, Theorem 9.1.14], there exists a continuous function ξ_2 (the regularised spectral shift

function) such that

$$\xi(\lambda) = \xi_2(\lambda) + \frac{1}{4\pi} \int_{\mathbb{R}^2} V(x) \, dx,$$

with $\lim_{\lambda \rightarrow \infty} \xi_2(\lambda) = 0$. In addition, since $S(0) = 1$, it follows that $\lim_{\lambda \searrow 0} \xi(\lambda) \in \mathbb{Z}$. We can thus consider the function $\lambda \mapsto f(\lambda)$ with

$$\begin{aligned} f(\lambda) &:= -2\pi i \xi(\lambda) + i \operatorname{tr}(A(\lambda)) \\ &= -2\pi i \left[\xi_2(\lambda) + \frac{1}{4\pi} \left(1 - \frac{2}{\pi} \tan^{-1}(\lambda) \right) \int_{\mathbb{R}^2} V(x) \, dx \right], \end{aligned}$$

which satisfies $\lim_{\lambda \rightarrow \infty} f(\lambda) = 0$ and $\lim_{\lambda \searrow 0} f(\lambda) \in (-2\pi i)\mathbb{Z}$. It finally remains to observe that

$$\det(S(\lambda)) \det(\beta(\lambda)) = e^{-2\pi i \xi(\lambda)} e^{i \operatorname{tr}(A(\lambda))} = e^{f(\lambda)}$$

and so the map $\lambda \mapsto \det(S(\lambda)) \det(\beta(\lambda))$ satisfies the properties stated. \blacksquare

We finally recall from [47, equation (9.1.22)] that the spectral shift function ξ satisfies for $\lambda > 0$ the equality

$$\operatorname{tr}(S(\lambda)^* S'(\lambda)) = -2\pi i \xi'(\lambda), \quad (5.6)$$

with the differentiability of ξ being guaranteed by [47, Theorem 9.1.18]. We can thus state the main result of this section.

Proposition 5.10. *The following equality holds:*

$$\operatorname{Index}(W_S) = \frac{1}{2\pi i} \int_0^\infty \operatorname{tr}(S(\lambda)^* S'(\lambda)) \, d\lambda + \frac{1}{4\pi} \int_{\mathbb{R}^2} V(x) \, dx.$$

Proof. By Lemma 5.8, one has $\operatorname{Index}(W_S) = \operatorname{Index}(W_{S\beta})$. Note that, by [44, Theorem 3.5 (a)], we have $\det(S\beta) = \det(S) \det(\beta)$. By Lemma 5.9, $\det(S\beta)$ defines a loop, and using Gohberg–Kreĭn theory (cf. [14] and [31, Theorem 4.9]) we can compute the index of $W_{S\beta}$ as

$$\begin{aligned} \operatorname{Index}(W_{S\beta}) &= \operatorname{Wind}(\det(S\beta)) \\ &= \frac{1}{2\pi i} \int_0^\infty \frac{\frac{d}{d\lambda} [\det(S(\lambda)) \det(\beta(\lambda))]}{\det(S(\lambda)) \det(\beta(\lambda))} \, d\lambda \\ &= \frac{1}{2\pi i} \int_0^\infty \frac{d}{d\lambda} (-2\pi i \xi(\lambda) + i \operatorname{tr}(A(\lambda))) \, d\lambda \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2\pi i} \int_0^\infty (-2\pi i \xi'(\lambda)) \, d\lambda + \frac{1}{2\pi} [\operatorname{tr}(A(\infty)) - \operatorname{tr}(A(0))] \\
 &= \frac{1}{2\pi i} \int_0^\infty \operatorname{tr}(S(\lambda)^* S'(\lambda)) \, d\lambda + \frac{1}{4\pi} \int_{\mathbb{R}^2} V(x) \, dx,
 \end{aligned}$$

as claimed. ■

By collecting the content of Proposition 5.5 and of Proposition 5.10, we can now confirm the statement of [6, Theorem 6.3].

Theorem 5.11. *If V satisfies (2.3) with $\rho > 11$, then the following equality holds:*

$$\frac{1}{2\pi i} \int_0^\infty \operatorname{tr}(S(\lambda)^* S'(\lambda)) \, d\lambda + \frac{1}{4\pi} \int_{\mathbb{R}^2} V(x) \, dx + \dim(P_p) = -\#\sigma_p(H).$$

We can then complement the content of [47, Theorem 9.1.14] about the spectral shift function.

Corollary 5.12. *If V satisfies (2.3) with $\rho > 11$, then the spectral shift function for the pair (H, H_0) satisfies*

$$\lim_{\varepsilon \searrow 0} \xi(\varepsilon) = -\#\sigma_p(H) - \dim(P_p).$$

Proof. By [47, Theorem 9.1.14] we have

$$\xi(\infty) = \frac{1}{4\pi} \int_{\mathbb{R}^2} V(x) \, dx.$$

By taking the equality (5.6) into account, we observe that

$$\begin{aligned}
 \lim_{\varepsilon \searrow 0} \xi(\varepsilon) &= - \int_0^\infty \xi'(\lambda) \, d\lambda + \xi(\infty) \\
 &= \frac{1}{2\pi i} \int_0^\infty \operatorname{tr}(S(\lambda)^* S'(\lambda)) \, d\lambda + \frac{1}{4\pi} \int_{\mathbb{R}^2} V(x) \, dx.
 \end{aligned}$$

The result now follows from Theorem 5.11. ■

A. Appendix

In the following statement, the standard bra-ket notation is freely used.

Lemma A.1. *Let \mathcal{H} be a complex Hilbert space, and let $\varphi, \psi \in \mathcal{H}$ be linearly independent. Consider $c \in \mathbb{C}$ with $|c| = 1$, define $T: \mathcal{H} \rightarrow \mathcal{H}$ by*

$$T := |\varphi\rangle\langle\varphi| + c|\psi\rangle\langle\psi|,$$

and set

$$k := \|\varphi\|^2\|\psi\|^2 - |\langle\varphi, \psi\rangle|^2 > 0.$$

Then, the operator T^\dagger defined by

$$\begin{aligned} T^\dagger := & \frac{1}{ck^2} [(c\|\psi\|^4 + |\langle\varphi, \psi\rangle|^2)|\varphi\rangle\langle\varphi| - (c\|\psi\|^2\langle\varphi, \psi\rangle + \|\varphi\|^2\langle\varphi, \psi\rangle)|\varphi\rangle\langle\psi| \\ & - (c\|\psi\|^2\langle\psi, \varphi\rangle + \|\varphi\|^2\langle\psi, \varphi\rangle)|\psi\rangle\langle\varphi| + (\|\varphi\|^4 + c|\langle\varphi, \psi\rangle|^2)|\psi\rangle\langle\psi|] \end{aligned}$$

satisfies $TT^\dagger = T^\dagger T = P_{\text{Ran}(T)}$, the projection on the range of T . Furthermore, the following equalities hold:

$$\langle\varphi, T^\dagger\varphi\rangle = 1, \quad \langle\psi, T^\dagger\psi\rangle = \bar{c}, \quad \langle\varphi, T^\dagger\psi\rangle = 0, \quad \langle\psi, T^\dagger\varphi\rangle = 0.$$

Proof. We first observe that

$$T\varphi = \|\varphi\|^2\varphi + c\langle\psi, \varphi\rangle\psi, \quad T\psi = \langle\varphi, \psi\rangle\varphi + c\|\psi\|^2\psi.$$

We can also compute

$$\begin{aligned} T^\dagger\varphi &= \frac{1}{ck^2} [((c\|\psi\|^4 + |\langle\varphi, \psi\rangle|^2)\|\varphi\|^2 - (c\|\psi\|^2\langle\varphi, \psi\rangle + \|\varphi\|^2\langle\varphi, \psi\rangle)\langle\psi, \varphi\rangle)\varphi \\ & \quad + (-c\|\psi\|^2\langle\psi, \varphi\rangle + \|\varphi\|^2\langle\psi, \varphi\rangle)\|\varphi\|^2 \\ & \quad + (\|\varphi\|^4 + c|\langle\varphi, \psi\rangle|^2)\langle\psi, \varphi\rangle\psi] \\ &= \frac{1}{ck^2} [c\|\psi\|^2(\|\psi\|^2\|\varphi\|^2 - |\langle\psi, \varphi\rangle|^2)\varphi + c\langle\psi, \varphi\rangle(|\langle\varphi, \psi\rangle|^2 - \|\psi\|^2\|\varphi\|^2)\psi] \\ &= \frac{1}{k} [\|\psi\|^2\varphi - \langle\psi, \varphi\rangle\psi], \end{aligned}$$

and similarly one gets

$$\begin{aligned} T^\dagger\psi &= \frac{1}{ck^2} [((c\|\psi\|^4 + |\langle\varphi, \psi\rangle|^2)\langle\varphi, \psi\rangle - (c\|\psi\|^2\langle\varphi, \psi\rangle + \|\varphi\|^2\langle\varphi, \psi\rangle)\|\psi\|^2)\varphi \\ & \quad + (-c\|\psi\|^2\langle\psi, \varphi\rangle + \|\varphi\|^2\langle\psi, \varphi\rangle)\langle\varphi, \psi\rangle \\ & \quad + (\|\varphi\|^4 + c|\langle\varphi, \psi\rangle|^2)\|\psi\|^2\psi] \\ &= \frac{1}{ck^2} [\langle\varphi, \psi\rangle(|\langle\varphi, \psi\rangle|^2 - \|\psi\|^2\|\varphi\|^2)\varphi + \|\varphi\|^2(\|\psi\|^2\|\varphi\|^2 - |\langle\psi, \varphi\rangle|^2)\psi] \\ &= \frac{1}{ck} [-\langle\varphi, \psi\rangle\varphi + \|\varphi\|^2\psi]. \end{aligned}$$

From these we obtain that

$$\begin{aligned} T^\dagger T \varphi &= \|\varphi\|^2 T^\dagger \varphi + c \langle \psi, \varphi \rangle T^\dagger \psi \\ &= \frac{\|\varphi\|^2}{k} [\|\psi\|^2 \varphi - \langle \psi, \varphi \rangle \psi] + \frac{c \langle \psi, \varphi \rangle}{ck} [-\langle \varphi, \psi \rangle \varphi + \|\varphi\|^2 \psi] = \varphi, \end{aligned}$$

and

$$\begin{aligned} T T^\dagger \varphi &= \frac{1}{k} [\|\psi\|^2 T \varphi - \langle \psi, \varphi \rangle T \psi] \\ &= \frac{1}{k} [\|\psi\|^2 (\|\varphi\|^2 \varphi + c \langle \psi, \varphi \rangle \psi) - \langle \psi, \varphi \rangle (\langle \varphi, \psi \rangle \varphi + c \|\psi\|^2 \psi)] = \varphi. \end{aligned}$$

By a similar computation, one also gets $T^\dagger T \psi = \psi$ and $T T^\dagger \psi = \psi$. The remaining equalities can also be obtained straightforwardly. ■

Acknowledgements. We thank the anonymous referee for a careful reading and helpful suggestions.

Funding. A. Alexander is supported by an Australian Government RTP scholarship. A. Rennie is supported by ARC Discovery grant DP220101196. S. Richard is supported by JSPS Grant-in-Aid for scientific research C no 21K03292.

References

- [1] S. Agmon, Spectral properties of Schrödinger operators and scattering theory. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **2** (1975), no. 2, 151–218 Zbl [0315.47007](#) MR [0397194](#)
- [2] A. Alexander and A. Rennie, [Levinson's theorem as an index pairing](#). *J. Funct. Anal.* **286** (2024), no. 5, article no. 110287 Zbl [07787696](#) MR [4682457](#)
- [3] W. O. Amrein, A. Boutet de Monvel, and V. Georgescu, *C_0 -groups, commutator methods and spectral theory of N -body Hamiltonians*. Progr. Math. 135, Birkhäuser, Basel, 1996 Zbl [0962.47500](#) MR [1388037](#)
- [4] M. Beceanu, [Decay estimates for the wave equation in two dimensions](#). *J. Differential Equations* **260** (2016), no. 6, 5378–5420 Zbl [1339.35037](#) MR [3448782](#)
- [5] J. Bellissard and H. Schulz-Baldes, [Scattering theory for lattice operators in dimension \$d \geq 3\$](#) . *Rev. Math. Phys.* **24** (2012), no. 8, article no. 1250020 Zbl [1256.35040](#) MR [2974082](#)
- [6] D. Bollé, F. Gesztesy, and C. Danneels, Threshold scattering in two dimensions. *Ann. Inst. H. Poincaré Phys. Théor.* **48** (1988), no. 2, 175–204 Zbl [0696.35040](#) MR [0952661](#)
- [7] D. Bollé, F. Gesztesy, C. Danneels, and S. F. J. Wilk, Threshold behavior and Levinson's theorem for two-dimensional scattering systems: a surprise. *Phys. Rev. Lett.* **56** (1986), 900–903
- [8] H. O. Cordes, *Elliptic pseudodifferential operators—an abstract theory*. Lecture Notes in Math. 756, Springer, Berlin, 1979 Zbl [0417.35004](#) MR [0551619](#)

- [9] H. D. Cornean, A. Michelangeli, and K. Yajima, [Two-dimensional Schrödinger operators with point interactions: threshold expansions, zero modes and \$L^p\$ -boundedness of wave operators](#). *Rev. Math. Phys.* **31** (2019), no. 4, article no. 1950012 Zbl 1423.35274 MR 3939663
- [10] J. Dereziński and S. Richard, [On Schrödinger operators with inverse square potentials on the half-line](#). *Ann. Henri Poincaré* **18** (2017), no. 3, 869–928 Zbl 1370.81070 MR 3611018
- [11] M. B. Erdoğan, M. Goldberg, and W. R. Green, [On the \$L^p\$ boundedness of wave operators for two-dimensional Schrödinger operators with threshold obstructions](#). *J. Funct. Anal.* **274** (2018), no. 7, 2139–2161 Zbl 1516.35132 MR 3762098
- [12] M. B. Erdoğan and W. R. Green, [Dispersive estimates for Schrödinger operators in dimension two with obstructions at zero energy](#). *Trans. Amer. Math. Soc.* **365** (2013), no. 12, 6403–6440 Zbl 1282.35143 MR 3105757
- [13] M. B. Erdoğan and W. R. Green, [A weighted dispersive estimate for Schrödinger operators in dimension two](#). *Comm. Math. Phys.* **319** (2013), no. 3, 791–811 Zbl 1272.35053 MR 3040376
- [14] I. C. Gohberg and M. G. Kreĭn, The basic propositions on defect numbers, root numbers and indices of linear operators. *Amer. Math. Soc. Transl. (2)* **13** (1960), 185–264 Zbl 0089.32201 MR 0113146
- [15] H. Inoue, [Explicit formula for Schrödinger wave operators on the half-line for potentials up to optimal decay](#). *J. Funct. Anal.* **279** (2020), no. 7, article no. 108630 Zbl 1481.47010 MR 4103875
- [16] H. Inoue and S. Richard, [Index theorems for Fredholm, semi-Fredholm, and almost periodic operators: all in one example](#). *J. Noncommut. Geom.* **13** (2019), no. 4, 1359–1380 Zbl 1440.81039 MR 4059823
- [17] H. Inoue and S. Richard, [Topological Levinson’s theorem for inverse square potentials: complex, infinite, but not exceptional](#). *Rev. Roumaine Math. Pures Appl.* **64** (2019), no. 2-3, 225–250 Zbl 1449.81019 MR 4012604
- [18] H. Inoue and N. Tsuzu, [Schrödinger wave operators on the discrete half-line](#). *Integral Equations Operator Theory* **91** (2019), no. 5, article no. 42 Zbl 07123543 MR 4013752
- [19] H. Isozaki and S. Richard, [On the wave operators for the Friedrichs–Faddeev model](#). *Ann. Henri Poincaré* **13** (2012), no. 6, 1469–1482 Zbl 1251.81091 MR 2966469
- [20] A. Jensen and T. Kato, [Spectral properties of Schrödinger operators and time-decay of the wave functions](#). *Duke Math. J.* **46** (1979), no. 3, 583–611 Zbl 0448.35080 MR 0544248
- [21] A. Jensen and G. Nenciu, [A unified approach to resolvent expansions at thresholds](#). *Rev. Math. Phys.* **13** (2001), no. 6, 717–754 Zbl 1029.81067 MR 1841744
- [22] A. Jensen and K. Yajima, [A remark on \$L^p\$ -boundedness of wave operators for two-dimensional Schrödinger operators](#). *Comm. Math. Phys.* **225** (2002), no. 3, 633–637 Zbl 1057.47011 MR 1888876
- [23] T. Kato, [Growth properties of solutions of the reduced wave equation with a variable coefficient](#). *Comm. Pure Appl. Math.* **12** (1959), 403–425 Zbl 0091.09502 MR 0108633

- [24] J. Kellendonk, K. Pankrashkin, and S. Richard, [Levinson's theorem and higher degree traces for Aharonov–Bohm operators](#). *J. Math. Phys.* **52** (2011), no. 5, article no. 052102 Zbl 1317.81106 MR 2839058
- [25] J. Kellendonk and S. Richard, [Levinson's theorem for Schrödinger operators with point interaction: a topological approach](#). *J. Phys. A* **39** (2006), no. 46, 14397–14403 Zbl 1105.81033 MR 2276222
- [26] J. Kellendonk and S. Richard, On the structure of the wave operators in one dimensional potential scattering. *Math. Phys. Electron. J.* **14** (2008), article no. 3 Zbl 1175.47010 MR 2486691
- [27] J. Kellendonk and S. Richard, Topological boundary maps in physics. In *Perspectives in operator algebras and mathematical physics*, pp. 105–121, Theta Ser. Adv. Math. 8, Theta, Bucharest, 2008 Zbl 1199.81029 MR 2433030
- [28] J. Kellendonk and S. Richard, [The topological meaning of Levinson's theorem, half-bound states included](#). *J. Phys. A* **41** (2008), no. 29, article no. 295207 Zbl 1149.81325 MR 2455280
- [29] J. Kellendonk and S. Richard, [On the wave operators and Levinson's theorem for potential scattering in \$\mathbb{R}^3\$](#) . *Asian-Eur. J. Math.* **5** (2012), no. 1, article no. 1250004 Zbl 1253.81124 MR 2931291
- [30] M. Klaus and B. Simon, [Coupling constant thresholds in nonrelativistic quantum mechanics. I. Short-range two-body case](#). *Ann. Physics* **130** (1980), no. 2, 251–281 Zbl 0455.35112 MR 0610664
- [31] M. Lesch, On the index of the infinitesimal generator of a flow. *J. Operator Theory* **26** (1991), no. 1, 73–92 Zbl 0784.46041 MR 1214921
- [32] H. S. Nguyen, S. Richard, and R. Tiedra de Aldecoa, [Discrete Laplacian in a half-space with a periodic surface potential I: Resolvent expansions, scattering matrix, and wave operators](#). *Math. Nachr.* **295** (2022), no. 5, 912–949 Zbl 07747276 MR 4453943
- [33] F. Nicoleau, D. Parra, and S. Richard, [Does Levinson's theorem count complex eigenvalues?](#) *J. Math. Phys.* **58** (2017), no. 10, article no. 102101 Zbl 1373.81215 MR 3707598
- [34] K. Pankrashkin and S. Richard, [Spectral and scattering theory for the Aharonov–Bohm operators](#). *Rev. Math. Phys.* **23** (2011), no. 1, 53–81 Zbl 1230.81054 MR 2774603
- [35] K. Pankrashkin and S. Richard, [One-dimensional Dirac operators with zero-range interactions: spectral, scattering, and topological results](#). *J. Math. Phys.* **55** (2014), no. 6, article no. 062305 Zbl 1298.81080 MR 3390663
- [36] S. Richard, [Levinson's theorem: an index theorem in scattering theory](#). In *Spectral theory and mathematical physics*, pp. 149–203, Oper. Theory Adv. Appl. 254, Birkhäuser, Cham, 2016 Zbl 06981835 MR 3526450
- [37] S. Richard and R. Tiedra de Aldecoa, [New formulae for the wave operators for a rank one interaction](#). *Integral Equations Operator Theory* **66** (2010), no. 2, 283–292 Zbl 1206.47013 MR 2595658
- [38] S. Richard and R. Tiedra de Aldecoa, [Explicit formulas for the Schrödinger wave operators in \$\mathbb{R}^2\$](#) . *C. R. Math. Acad. Sci. Paris* **351** (2013), no. 5-6, 209–214 Zbl 1268.81152 MR 3089680

- [39] S. Richard and R. Tiedra de Aldecoa, [New expressions for the wave operators of Schrödinger operators in \$\mathbb{R}^3\$](#) . *Lett. Math. Phys.* **103** (2013), no. 11, 1207–1221
Zbl [1273.81219](#) MR [3095143](#)
- [40] S. Richard, R. Tiedra de Aldecoa, and L. Zhang, [Scattering operator and wave operators for 2D Schrödinger operators with threshold obstructions](#). *Complex Anal. Oper. Theory* **15** (2021), no. 6, article no. 106 Zbl [1476.81136](#) MR [4308889](#)
- [41] M. Rørdam, F. Larsen, and N. Laustsen, *An introduction to K -theory for C^* -algebras*. London Math. Soc. Stud. Texts 49, Cambridge University Press, Cambridge, 2000
Zbl [0967.19001](#) MR [1783408](#)
- [42] W. Schlag, [Dispersive estimates for Schrödinger operators in dimension two](#). *Comm. Math. Phys.* **257** (2005), no. 1, 87–117 Zbl [1134.35321](#) MR [2163570](#)
- [43] H. Schulz-Baldes, [The density of surface states as the total time delay](#). *Lett. Math. Phys.* **106** (2016), no. 4, 485–507 Zbl [1341.82094](#) MR [3474898](#)
- [44] B. Simon, *Trace ideals and their applications*. Second edn., Math. Surveys Monogr. 120, American Mathematical Society, Providence, RI, 2005 Zbl [1074.47001](#) MR [2154153](#)
- [45] E. Toprak, [A weighted estimate for two dimensional Schrödinger, matrix Schrödinger, and wave equations with resonance of the first kind at zero energy](#). *J. Spectr. Theory* **7** (2017), no. 4, 1235–1284 Zbl [1383.35069](#) MR [3737892](#)
- [46] D. R. Yafaev, *Mathematical scattering theory*. General Theory. Transl. Math. Monogr. 105, American Mathematical Society, Providence, RI, 1992 Zbl [0761.47001](#)
MR [1180965](#)
- [47] D. R. Yafaev, *Mathematical scattering theory*. Math. Surveys Monogr. 158, American Mathematical Society, Providence, RI, 2010 Zbl [1197.35006](#) MR [2598115](#)
- [48] K. Yajima, [\$L^p\$ -boundedness of wave operators for two-dimensional Schrödinger operators](#). *Comm. Math. Phys.* **208** (1999), no. 1, 125–152 Zbl [0961.47004](#) MR [1729881](#)
- [49] K. Yajima, [\$L^p\$ -boundedness of wave operators for 2D Schrödinger operators with point interactions](#). *Ann. Henri Poincaré* **22** (2021), no. 6, 2065–2101 Zbl [1467.35121](#)
MR [4264892](#)
- [50] K. Yajima, [The \$L^p\$ -boundedness of wave operators for two dimensional Schrödinger operators with threshold singularities](#). *J. Math. Soc. Japan* **74** (2022), no. 4, 1169–1217
Zbl [1523.35084](#) MR [4499832](#)

Received 16 November 2023; revised 12 February 2024.

Angus Alexander

School of Mathematics and Applied Statistics, University of Wollongong, Northfields Ave,
Wollongong, NSW 2522, Australia; angusa@uow.edu.au

Thanh Duc Nguyen

Department of Physics, Graduate School of Science, Nagoya University, Furo-cho,
Chikusa-ku, Nagoya 464-8602, Japan; thanhnghuyen@st.phys.nagoya-u.ac.jp

Adam Rennie

School of Mathematics and Applied Statistics, University of Wollongong, Northfields Ave,
Wollongong, NSW 2522, Australia; renniea@uow.edu.au

Serge Richard

Institute for Liberal Arts and Sciences & Graduate School of Mathematics, Nagoya University,
Furo-cho, Chikusa-ku, Nagoya 464-8601, Japan; richard@math.nagoya-u.ac.jp