

Factoring strongly irreducible group shift actions onto full shifts of lower entropy

Dawid Huczek and Sebastian Kopacz

Abstract. We show that if G is a countable amenable group G with the comparison property and X is a strongly irreducible G -shift satisfying certain aperiodicity conditions, then X factors onto the full G -shift on N symbols, so long as the logarithm of N is less than the topological entropy of G .

1. Introduction

A well-known result in the study of symbolic dynamical systems states that any subshift of finite type (SFT) with the action of \mathbb{Z} and entropy greater than or equal to $\log N$ factors onto the full shift over N symbols—this was proven in [9] and [1] for the cases of equal and unequal entropy, respectively. Extending these results for actions of other groups has been difficult, and it is known that a factor map onto a full shift of equal entropy may not exist, even in the case of \mathbb{Z}^d , where $d > 1$ (see [3]). Johnson and Madden [7] showed that any SFT with the action of \mathbb{Z}^d , which has entropy greater than $\log N$ and satisfies an additional mixing condition (known as corner gluing), has an extension which is finite-to-one (hence of equal entropy) and maps onto the full shift over N symbols. This result was later improved by Desai in [4] to show that such a system factors directly onto the full shift, without the intermediate extension, and then by Boyle, Pavlov, and Schraudner [2] to replace the corner gluing by a weaker mixing condition (block gluing).

In this paper, we use similar methods to adapt these constructions to symbolic dynamical systems with actions of amenable groups. Our approach requires three assumptions: that the group G has the comparison property (satisfied, for instance, for all countable amenable groups with subexponential growth), that the system X is strongly irreducible (which replaces the corner gluing condition and allows the construction to be valid without assuming that the underlying system be an SFT), and that it has nonperiodic blocks for all possible sets of periods. With these assumptions, we prove that X can be factored onto any full shift of smaller entropy (i.e., a full shift over N symbols, where $\log N < h(X)$). We note that our method does not apply for the case of equal entropy ($\log N = h(X)$); as

Mathematics Subject Classification 2020: 37B10 (primary); 37B40, 43A07 (secondary).

Keywords: countable amenable group, (dynamical) tiling, free action, topological entropy, comparison property.

we mentioned earlier, Boyle and Schraudner [3] have shown that there exist \mathbb{Z}^d -shifts of finite type which do not factor onto full shifts of equal entropy.

2. Preliminaries

In this section, we establish the definitions, notation, and standard facts we will use in this paper. Since this is mainly standard material, we omit most proofs and references.

2.1. Amenability, Følner sets, and invariance

Throughout this paper, G will denote a countable amenable group and (F_n) will denote a fixed Følner sequence, that is, a sequence of finite subsets of G such that for every $g \in G$ the sequence $\frac{|gF_n \Delta F_n|}{|F_n|}$ tends to 0 as n goes to infinity. Multiplication involving sets will always be understood element wise, so gF_n is the set $\{gf : f \in F_n\}$, and KF in the following definition denotes the set $\{kf : k \in K, f \in F\}$.

Definition 2.1. Let K, F be finite subsets of G and let $\varepsilon > 0$. We say that F is (K, ε) -invariant if $|KF \Delta F| < \varepsilon|F|$.

The defining property of the Følner sequence can be equivalently (and usefully) stated as follows: For every finite $K \subset G$ and every $\varepsilon > 0$, there exists an N such that for every $n \geq N$ the set F_n is (K, ε) -invariant.

Definition 2.2. Let D, F be finite subsets of G . Let $F_D = \{g \in F : Dg \subset F\}$. We will refer to F_D as the D -interior of F .

A straightforward computation shows that for any $\varepsilon > 0$, if F is $(D, \frac{\varepsilon}{|D|})$ -invariant, then $|F_D| \geq (1 - \varepsilon)|F|$ (we will describe such a relation in cardinalities by saying that F_D is a $(1 - \varepsilon)$ -subset of F). Also observe that $F_{DK} = (F_D)_K$.

2.2. Symbolic dynamical systems

Let Λ be a finite set (referred to as the *alphabet*). The *full G -shift* over Λ is the product set Λ^G with the product topology (induced by the discrete topology on Λ) endowed with the right-shift action of G :

$$(gx)(h) = x(hg).$$

By a symbolic dynamical system, a shift space, or subshift, we understand any closed, shift-invariant subset X of Λ^G .

Definition 2.3. For a finite $T \subset G$, by a block with domain T we understand a function $B : T \rightarrow \Lambda$. If X is a symbolic dynamical system over Λ , and $x \in X$, then we say that B occurs in x (at position g) if for every $t \in T$ we have $x(tg) = B(t)$, which we denote more concisely as $x|_{Tg} = B$. We say that B occurs in X if it occurs in some $x \in X$, and we denote the set of all possible blocks with domain T which occur in X by $\mathcal{B}_T(X)$.

Definition 2.4. We say that a symbolic dynamical system X is *strongly irreducible* (with irreducibility distance D , where D is a finite subset of G containing the identity element e) if for any blocks B_1, B_2 (with domains T_1, T_2) which occur in X , and any $g_1, g_2 \in G$ such that $DT_1g_1 \cap T_2g_2 = \emptyset$, there exists an $x \in X$ such that $x|_{T_1g_1} = B_1$ and $x|_{T_2g_2} = B_2$.

Definition 2.5. If P is a finite subset of G , we say that a block B with domain T is P -aperiodic if for every $p \in P$ there exists a $t \in T$ such that tp is also in T , and $B(tp) \neq B(t)$. We will say that a symbolic dynamical system X is aperiodic if it has a P -aperiodic block for every finite $P \subset G$.

Definition 2.6. For a fixed Følner sequence (F_n) , let $\mathcal{N}_{F_n}(X)$ be the cardinality of the set $\mathcal{B}_{F_n}(X)$. The topological entropy of a symbolic dynamical system X is defined as

$$h(X) = \lim_{n \rightarrow \infty} \frac{1}{|F_n|} \log \mathcal{N}_{F_n}(X).$$

(In this paper, we use logarithms with base 2, although as usual the theorems remain true if one defines entropy using any other base, so long as the choice remains consistent throughout.) It is a standard fact that the obtained value of $h(X)$ does not depend on the choice of the Følner sequence. In fact, the relation between entropy and the number of blocks holds for any sufficiently invariant domain, as per the following theorem [8].

Theorem 2.7. *For any $\varepsilon > 0$, there exists an $N > 0$ and $\delta > 0$ such that if T is an (F_n, δ) -invariant set for some $n > N$, then $\mathcal{N}_T(X) > 2^{(h(X) - \varepsilon)|T|}$.*

2.3. Quasitilings and tilings

Definition 2.8. A *quasitiling* of a countable amenable group G is any collection \mathcal{T} of finite subsets of G (referred to as *tiles*) such that there exists a finite collection \mathcal{S} of finite subsets of G (referred to as *shapes*) such that every $T \in \mathcal{T}$ has a unique representation $T = Sc$ for some $S \in \mathcal{S}$ and some $c \in G$. We refer to such a c as the *center* of T . If the tiles of \mathcal{T} are disjoint and their union is all of G , then we refer to \mathcal{T} as a *tiling*.

Remark 2.9. If we enumerate the set of shapes of a quasitiling \mathcal{T} as $\{S_1, S_2, \dots, S_r\}$, then we can identify \mathcal{T} with an element $x_{\mathcal{T}}$ of the set $\{0, 1, \dots, r\}^G$, letting $x_{\mathcal{T}}(g) = j$ if $S_j g$ is a shape of \mathcal{T} and $x_{\mathcal{T}}(g) = 0$ otherwise. This in turn induces (via orbit closure) a subshift $X_{\mathcal{T}}$, and any element of $X_{\mathcal{T}}$ in turn corresponds to a quasitiling of G which has the same disjointness, invariance, and density properties as \mathcal{T} . This allows us to discuss some properties of quasitilings using the notions of topological dynamics (in particular, it makes sense to consider entropy and factorizations), by interpreting these notions as applied to the corresponding subshifts.

We will use several theorems which guarantee the existence of quasitilings and/or tilings satisfying certain properties. The first is proven in [5], and we will invoke it

when constructing subsystems with specified entropy (in Section 3) and marker blocks (in Section 4).

Theorem 2.10. *For any finite $K \subset G$ and any $\varepsilon > 0$, there exists a tiling \mathcal{T} of G such that all tiles of \mathcal{T} are (K, ε) -invariant and $h(\mathcal{T}) = 0$.*

When we construct the factor map onto the full shift, we will rely on combining two other results. The first one originally appears in the seminal paper by Ornstein and Weiss [10], although in [5] it is stated and proven in an equivalent form closer to the one stated here (the main difference being that the original version does not explicitly use the notion of lower Banach density).

Theorem 2.11. *For any $\delta > 0$ and $N > 0$, there exists a quasitiling \mathcal{T} of G such that:*

- (1) *The shapes of \mathcal{T} are all Følner sets, that is, $\mathcal{S} = \{F_{n_1}, F_{n_2}, \dots, F_{n_r}\}$, where $N \leq n_1 < n_2 < \dots < n_r$, and r depends only on δ .*
- (2) *\mathcal{T} is δ -disjoint, that is, every tile $T \in \mathcal{T}$ has a subset T° such that $|T^\circ| > (1 - \delta)|T|$ and these subsets are pairwise disjoint.*
- (3) *\mathcal{T} is $(1 - \delta)$ -covering, that is, the lower Banach density of the union of \mathcal{T} is greater than $1 - \delta$.*

The statement of the next theorem involves the comparison property. We will restrict ourselves to recalling the definition of comparison property ([6, Definition 6.1]) and formulating the result we are going to invoke; for more details we refer the reader to the cited paper.

Definition 2.12. A countable amenable group G is said to have the comparison property if for every action of G on a zero-dimensional compact metric space X , and every pair of clopen subsets $A, B \subset X$, such that $\mu(A) \leq \mu(B)$ for every G -invariant Borel measure μ on X , there exists a finite partition $A = \bigcup_{i=1}^k A_i$ into clopen sets, and elements $g_1, g_2, \dots, g_k \in G$ such that $g_1(A_1), g_2(A_2), \dots, g_k(A_k)$ are disjoint subsets of B .

Per [6, Theorem 6.33], every group of subexponential growth has the comparison property. In fact, it seems to be still an open question whether countable amenable groups without the comparison property exist at all.

The following theorem is established as a step in the proof of [6, Theorem 7.5]; we will state it as a stand-alone result.

Theorem 2.13. *Suppose G is a countable amenable group with the comparison property, and K is a finite subset of G containing the neutral element e . For every $\varepsilon > 0$, there exist $\delta > 0$ and $N > 0$ such that every $(1 - \delta)$ -covering quasitiling \mathcal{T} whose tiles are pairwise disjoint and shapes are (K, δ) -invariant and have cardinality larger than N can be modified to a tiling \mathcal{T}° whose shapes are (K, ε) -invariant. Moreover, the sets of centers of \mathcal{T}*

and \mathcal{T}° are identical, and there exists a finite $H \subset G$ such that for any $g \in G$ we can determine the tile of \mathcal{T}° to which it belongs, so long as we know the set $Hg \cap \mathcal{T}$ (i.e., using the language of topological dynamics, \mathcal{T}° is a factor of \mathcal{T}).

For convenience, we will combine Theorems 2.11 and 2.13 into the following form (which will be used in the main construction).

Theorem 2.14. *If G is a countable amenable group with the comparison property, then for every $\varepsilon > 0$, $N > 0$, and every finite $K \subset G$, there exists a quasitiling \mathcal{T}' of G such that:*

- (1) *The shapes of \mathcal{T}' are all Følner sets, that is, $\mathcal{S} = \{F_{n_1}, F_{n_2}, \dots, F_{n_r}\}$, where $N \leq n_1 < n_2 < \dots < n_r$, and r depends only on ε .*
- (2) *\mathcal{T}' factors onto an exact tiling \mathcal{T}'' of G whose shapes are (K, ε) -invariant.*

3. Subsystem entropy

The result below seems classical, but for the convenience of the reader, we include the elementary proof.

Theorem 3.1. *If X is a strongly irreducible subshift of positive entropy h , then the set of topological entropies of subshifts of X is dense in $(0, h)$.*

Proof. Let a and b be such that $0 < a < b < h$. We will show that there exists a subshift $Y \subset X$ such that $a \leq h(Y) \leq b$ (which is an equivalent formulation of the theorem). Fix a positive η smaller than $(b - a)/2$ and note that for every sufficiently large L , there exists a positive integer n such that $\frac{1}{L} \log n$ is in the interval $(a + \eta, b - \eta)$. Combined with Theorem 2.7, this lets us state the following.

Fact. *There exist $N > 0$, $\delta > 0$, and $M > 0$ such that if T is an (F_n, δ) -invariant set for some $n > N$, and $|T| \geq M$, then $\mathcal{B}_T(X)$ has a subset \mathcal{B}_T^* such that $a + \eta < \frac{1}{|T|} \log |\mathcal{B}_T^*| < b - \eta$.*

Let D denote the irreducibility distance of X . By Theorem 2.14, there exists a tiling \mathcal{T} of G such that $h(\mathcal{T}) = 0$ and the shapes of \mathcal{T} can have arbitrarily good invariance properties, which we will specify within the next few sentences. Enumerate the shapes of \mathcal{T} by S_1, S_2, \dots, S_J . There exist some δ^*, δ^{**} , and n such that if the shapes of \mathcal{T} are (D, δ^*) and (F_n, δ^{**}) -invariant, then for every j we can choose a family of blocks $\mathcal{B}_j \subset \mathcal{B}_{(S_j)_D}(X)$ such that if we denote $N_j = |\mathcal{B}_j|$, then we have $a + \eta < \frac{1}{|(S_j)_D|} \log N_j < b - \eta$. In addition, since for small enough δ^* the relative difference between $|(S_j)_D|$ and $|S_j|$ can be arbitrarily small, we can even write

$$a + \eta < \frac{1}{|S_j|} \log N_j < b - \eta.$$

Now, let Y be the orbit closure of the set of all points $x \in X$ such that for every $T \in \mathcal{T}$ $x|_{T_D} \in \mathcal{B}_j$, where j is such that S_j is the shape of T . Observe that strong irreducibility, combined with the fact that T_D is disjoint from all other tiles of \mathcal{T} , means (via a standard compactness argument) that we can choose the blocks $x|_{T_D}$ independently of each other, and any such choice will yield a valid element of Y .

We will estimate the entropy of Y by considering the number of blocks with domain F_n as n increases to infinity, beginning with estimating this number of blocks described above. Fix some large n and note that every block B with domain F_n that occurs in Y has the property that there exists some right-translate \mathcal{T}' of \mathcal{T} such that for every tile T of \mathcal{T}' such that $T \subset F_n$ the block $B|_{T_D}$ belongs to \mathcal{B}_j , where j is such that S_j is the shape of T . Let H_n be the number of ways in which the right-translates of \mathcal{T} can intersect F_n , or equivalently, the number of ways in which the right-translates of F_n can intersect \mathcal{T} . Since \mathcal{T} has entropy 0, for large enough n we have $\frac{1}{|F_n|} \log H_n < \frac{\eta}{2}$.

For any such right-translate \mathcal{T}' let l_j be the number of tiles of \mathcal{T}' with shape S_j that are subsets of F_n . Note that if n is large enough, then $\sum_{j=1}^J l_j |S_j|$ is almost equal to $|F_n|$. Since for every tile of \mathcal{T}' the block $B|_{T_D}$ is a block from \mathcal{B}_j , and thus one of the N_j possible blocks, the D -interiors of the tiles of \mathcal{T}' can be filled in at most $(\prod_{j=1}^J N_j^{l_j})$ ways. We have no control over the symbols outside these interiors, but we know that there are at most $|F_n| - \sum_{j=1}^J l_j |(S_j)_D|$ such symbols. It follows that the number of blocks associated with \mathcal{T}' is at most

$$\left(\prod_{j=1}^J N_j^{l_j} \right) \cdot |\Lambda|^{|F_n| - \sum_{j=1}^J l_j |(S_j)_D|}.$$

Taking logarithms and dividing by $|F_n|$, we obtain

$$\frac{1}{|F_n|} \sum_{j=1}^J l_j \log N_j + \frac{|F_n| - \sum_{j=1}^J l_j |(S_j)_D|}{|F_n|} \log |\Lambda|.$$

If δ^* was small enough so that the D -interiors of the tiles of \mathcal{T}' form a $(1 - \frac{\eta}{2|\Lambda|})$ -covering quasitiling (which we can safely assume since D , η , and Λ were known before we started the construction), then for large enough n the second term will not exceed $\frac{\eta}{2}$. As for the first term, we can further estimate it as follows:

$$\frac{1}{|F_n|} \sum_{j=1}^J l_j \log N_j = \frac{1}{|F_n|} \sum_{j=1}^J l_j |S_j| \frac{1}{|S_j|} \log N_j < (b - \eta) \frac{1}{|F_n|} \sum_{j=1}^J l_j |S_j| < b - \eta.$$

It follows that for any right-translate \mathcal{T}' of \mathcal{T} , the number of blocks in Y with domain F_n that have blocks from \mathcal{B}_j in every tile of \mathcal{T}' with shape S_j does not exceed $2^{(b - \frac{\eta}{2})|F_n|}$, and thus the cardinality of $\mathcal{B}_{F_n}(Y)$ does not exceed $H_n 2^{(b - \frac{\eta}{2})|F_n|}$. This lets us estimate that

$$\frac{1}{|F_n|} \log |\mathcal{B}_{F_n}(Y)| \leq \frac{1}{|F_n|} \log H_n + b - \frac{\eta}{2} < b,$$

and hence,

$$h(Y) < b.$$

The lower bound for entropy is much simpler—for any n the number of blocks with domain F_n which occur in Y is equal to at least $(\prod_{j=1}^J N_j^{l_j})$, where l_j is the number of tiles of \mathcal{T} with shape S_j which are subsets of F_n (this time we do not even need to consider possible translates of \mathcal{T}). Consequently,

$$\begin{aligned} \frac{1}{|F_n|} \log |\mathcal{B}_{F_n}(Y)| &\geq \frac{1}{|F_n|} \sum_{j=1}^J l_j \log N_j = \frac{1}{|F_n|} \sum_{j=1}^J l_j |S_j| \frac{1}{|S_j|} \log N_j \\ &> (a + \eta) \frac{\sum_{j=1}^J l_j |S_j|}{|F_n|}, \end{aligned}$$

which for large enough n will be greater than a , and therefore $h(Y) > a$, which concludes the proof. ■

4. Marker blocks

According to Theorem 2.7, if $h(X) > \log N$, it is possible to define a surjective block-to-block map from $\mathcal{B}_{F_n}(X)$ onto $\{0, 1, \dots, N-1\}^{F_n}$, for sufficiently large n . This allows us, with the use of a tiling, to build a map Π from X onto full G -shift over N symbols, by defining it on a tile-by-tile basis. However, without some extra care such a map might not commute with the group action, that is, $\Pi(gx)$ might not be equal to $g\Pi(x)$. Therefore, we need to work with the subshift $X_{\mathcal{T}}$ rather than directly with the tiling \mathcal{T} , and we have to find a suitable way of “decoding” a tiling based on the content of $x \in X$. To achieve this, we will use special blocks called *marker blocks* or, more concisely, *markers*. Each tile will have its own marker block which we will place inside, for example, in the center, and outside the marker, we will leave enough space to apply the block-to-block map. That way the factor map will involve first locating the marker block to determine the tiling and then using the block-to-block map to determine the content of the image within each tile. The procedure is possible due to strong irreducibility: First, when we put marker blocks in tiles (strong irreducibility gives us a lot of freedom in specifying the content of a tile except within a small “neighborhood” of the marker blocks and near the boundary of the tile) and again, when we “glue” the contents of tiles together to get final point (i.e., we will specify the symbols in the part of the tile that is suitably “far from the boundary,” and strong irreducibility will ensure that any combination of two tiles with such specified content will occur somewhere in X). A very important element of the construction (and a major source of difficulties) will be to ensure that we will have only one marker block per tile: Once the marker blocks are placed, *a priori* there is the risk that the other symbols might accidentally create a marker block in a different position—either within the areas which we will use for the actual block-to-block code or in the “gaps” whose contents we

will have to leave mostly uncontrolled in order to take advantage of strong irreducibility. The same problem occurs in the case of \mathbb{Z}^d , but the solution is more complicated for general amenable groups, because the lack of commutativity greatly increases the number of possible scenarios we need to consider. Thus, our main goal in this section will be to prove the following theorem, which shows how to build the marker blocks.

Theorem 4.1. *Let X be an aperiodic, strongly irreducible symbolic dynamical system (with the irreducibility distance D) over alphabet Λ , with action of a countable amenable group G . Let Y be a proper subshift of X . Then there exists a block M with shape K , and a tiling \mathcal{T} (with a family of shapes \mathcal{S}), satisfying the following conditions:*

- * *For every $T \in \mathcal{T}$, any $t \in T$, and any $x \in X$, if $x|_{Kt} = M$, then the set $T_D \cap Kt$ is not empty, and the block $x|_{T_D \cap Kt}$ does not occur in Y .*
- ** *For two different $g_1, g_2 \in G$ and any $x \in X$, if $x|_{Kg_1} = x|_{Kg_2} = M$, then $Kg_2 \setminus DKg_1 \neq \emptyset$, and $x|_{Kg_2 \setminus DKg_1}$ is a block which does not appear in Y .*

Proof. Before we proceed, let us briefly discuss the intuition and meaning behind our two conditions and especially how a block M satisfying such conditions is used in the context of marker blocks. We know that there exists a block B such that B does not occur in Y . The marker block M will contain many occurrences of B (and some extra aperiodic blocks) to ensure that certain subblocks of M also cannot be blocks occurring in Y . In the main construction, inside the D -interior of each tile of \mathcal{T} we will want to mostly put blocks from Y , other than the one instance of the marker block M . The fact that M (and some of its specific subblocks) cannot occur in Y helps us control where in a tile it might appear. The first property states that if M occurs outside the D -interior of a tile of \mathcal{T} , then a large part of it would still overlap the D -interior of some tile, and the overlapping part would be a block that is forbidden in Y . Therefore, if we ensure that the D -interiors of tiles (outside of the explicitly placed markers) are blocks from Y , this will not be possible in our construction. To accomplish that, we will “enlarge” M , adding some copies of the block B in a specific way. The second property says that any two occurrences of M within one point cannot overlap too much, and in addition, the nonoverlapping parts form a block that cannot occur in Y . This lets us ensure that a new marker block M will not be accidentally created between the original marker block and rest of the tile: In $x|_{Kg_1}$ we put the marker block M , and we can be sure that we will not find any other marker blocks within coordinates “close” to DKg_1 . Here, the proof differs from the case \mathbb{Z}^d ; since our group G need not be Abelian, there might exist elements g_1 and g_2 such that $Kg_1 = Kg_2$, greatly complicating the ways in which an overlap between two occurrences might be produced. The core idea we use to avoid such problems is simple: We add two more copies of the block B to the marker block, in a way that prevents all possible conflicts. Unfortunately, verifying this involves checking many conditions on how the overlaps might be produced, but there are still finitely many of such conditions, each of them disallowing a finite number of places where we could put these copies of B and finally letting us obtain the desired marker block M .

For clarity, we will separate the proof into stages:

- (1) Since Y is a proper subshift of X , there exists some block B which occurs in X but not in Y . Let Z denote the shape of B .
- (2) Let $P_e = Z^{-1}DZ$ and for any $g \in G$ let $P_g = g^{-1}P_e g$. Observe that for any g , P_g is a finite set (and all of these sets have equal cardinality), and $e \in P_g$ (because we assumed that $e \in D$). Let P'_g be the largest (in terms of cardinality) subset of P_g such that $P'_g \subset P_h$ for infinitely many h (if there is more than one such subset of equal cardinality, we can choose any of them). Choose g_0 so that P'_{g_0} has maximal cardinality among all the P_g s. Set $P = P'_{g_0}$ and $G_P = \{g \in G : P \subset P_g\}$. Observe that if any $h \in G$ occurs in infinitely many P_g for $g \in G_P$, then due to the maximality of P we necessarily have $h \in P$. It follows that for any $E \subset G$ we have $E \cap P_g \subset P$ for all but finitely many $g \in G_P$.
- (3) By our assumption, there exists in X a block M_0 with domain K_0 which is P -aperiodic, that is, for every $g \in P$ there exists a $k \in K_0$ such that k and kg are both elements of K_0 and $M_0(kg) \neq M_0(k)$. By strong irreducibility, we can also require that M_0 has B as a subblock, and thus M_0 cannot occur in Y .
- (4) Let \mathcal{T}' be a tiling of G , such that the shapes of \mathcal{T}' are supersets of DZ , let S' denote the union of all shapes of \mathcal{T}' , and let \mathcal{T} be another tiling, whose shapes are (S', δ) -invariant, where δ is so small that for every shape S of \mathcal{T} , and every $s \in G$ the set $S_D s^{-1}$ contains an entire tile of \mathcal{T}' . Let S be the union of all shapes of \mathcal{T} . By the invariance property established earlier, for any $s \in S \setminus S_{DK_0}$ the set $S_D s^{-1}$ contains an entire tile of \mathcal{T}' , which allows us to choose a finite set C' (consisting of centers of such tiles) such that for every $s \in S \setminus S_{DK_0}$ the set $S_D s^{-1}$ is a superset of Zc' for some $c' \in C'$.
 Let $K_1 = K_0 \cup \bigcup_{c' \in C'} DZc'$. Strong irreducibility yields the existence of a block M_1 with domain K_1 , such that $M_1|_{K_0} = M_0$, and for every $c' \in C'$ we have $M_1|_{Zc'} = B$. Observe that for any tile $T \in \mathcal{T}$, if $g \in T$, then either $K_0 g \subset T_D$ or for at least one $c' \in C'$ we have $Zc'g \subset T_D$. Indeed, T has the form Sc , where S is one of the shapes of \mathcal{T} . Let $s = gc^{-1}$. If $s \in S_{DK_0}$, then $DK_0 s \subset S$, and thus $DK_0 g = DK_0 s c \subset Sc = T$. Otherwise, we know that for some $c' \in C'$ we have $Zc' \subset S_D s^{-1}$. Therefore, $Zc' s \subset S_D$, which in turn yields $Zc' g = Zc' s c \subset S_D c = T_D$.
 That way we have obtained a block M_1 with domain K_1 that satisfies the first property stated in our theorem. Note that any block that has M_1 as a subblock will retain this property.
- (5) Now choose c_2 as an element of G_P which satisfies the following conditions (each of them is satisfied for all but finitely many elements of the group, and G_P is infinite, which makes such a choice possible). The significance of these conditions is certainly not obvious at first, but will become clear later:
 - (a) $c_2 \notin Z^{-1}D^{-1}K_1$,

- (b) $c_2^{-1} \notin K_1^{-1}DK_1K_1^{-1}D^{-1}Z$,
(c) $c_2^{-1} \notin K_1^{-1}DZZ^{-1}D^{-1}Z$.

Also choose c_3 as an element of G_P satisfying the following conditions (which is again possible because each of the following is true for all but finitely many elements of G_P):

- (d) $c_3 \notin Z^{-1}D^{-1}(K_1 \cup Zc_2)$,
(e) $c_3 \notin Z^{-1}D^{-1}K_1c_2^{-1}Z^{-1}DK_1$,
(f) $c_3 \notin Z^{-1}D^{-1}K_1c_2^{-1}Z^{-1}DZc_2$,
(g) $c_3 \notin Z^{-1}D^{-1}Zc_2K_1^{-1}DK_1$,
(h) $c_3 \notin Z^{-1}D^{-1}Zc_2K_1^{-1}DZc_2$,
(i) $P_{c_3} \cap P_{c_2} = P$,
(j) $c_3^{-1} \notin K_1^{-1}DK_1K_1^{-1}D^{-1}Z$,
(k) $c_3^{-1} \notin c_2^{-1}Z^{-1}DZZ^{-1}D^{-1}Z$,
(l) $P_{c_3} \cap K_1^{-1}DZc_2 = P$,
(m) $c_3^{-1} \notin K_1^{-1}DZc_2K_1^{-1}D^{-1}Z$,
(n) $c_3^{-1} \notin K_1^{-1}DZZ^{-1}D^{-1}Z$.

Finally, we replace K' with $K = K_1 \cup Zc_2 \cup Zc_3$, and we choose a block M such that $M|_{K_1} = M_1$, $M|_{Zc_2} = M|_{Zc_3} = B$. Such an M exists, because conditions (5a) and (5d) imply that DZc_2 is disjoint from K_1 and DZc_3 is disjoint from $K_1 \cup Zc_2$, which allows us to invoke the strong irreducibility of X . Now, assume that for some $x \in X$ and $g \in G$, we have $x|_{Kg} = x|_K = M$. We will show that in such a situation, at least one of the sets K_1g , Zc_2g , and Zc_3g is disjoint with DK . This will require some laborious and repetitive computations, which we will separate into the following steps:

- $x|_{Kg} = x|_K = M$ implies that for every $k \in K_1$ such that kg is also in K_1 , we have $M_1(k) = M_1(kg)$. Since M_1 is P -aperiodic, g cannot belong to P .
- Observe that for any $g \in G$ each of the sets DK_1 , DZc_2 , and DZc_3 is intersected by at most one of the sets K_1g and Zc_2g . Indeed:
 - Suppose that $DK_1 \cap K_1g$ and $DK_1 \cap Zc_2g$ are both nonempty sets. This implies that $g \in K_1^{-1}DK_1$ and $g \in c_2^{-1}Z^{-1}DK_1$, thus we obtain $K_1^{-1}DK_1 \cap c_2^{-1}Z^{-1}DK_1 \neq \emptyset$. This, however, would imply that $c_2^{-1} \in K_1^{-1}DK_1K_1^{-1}D^{-1}Z$, contradicting condition (5b).
 - Similarly, if $DZc_2 \cap K_1g$ and $DZc_2 \cap Zc_2g$ are both nonempty, a similar reasoning tells us that the intersection $K_1^{-1}DZc_2 \cap c_2^{-1}Z^{-1}DZc_2$ is nonempty, thus so is the set $K_1^{-1}DZ \cap c_2^{-1}Z^{-1}DZ$. This would mean that $c_2^{-1} \in K_1^{-1}DZZ^{-1}D^{-1}Z$, contradicting condition (5c).

- In the case where $DZc_3 \cap K_1g$ and $DZc_3 \cap Zc_2g$ are both nonempty, we can similarly conclude that the set $K_1^{-1}DZc_3 \cap c_2^{-1}Z^{-1}DZc_3$ is nonempty, thus so is $K_1^{-1}DZ \cap c_2^{-1}Z^{-1}DZ$, which is exactly the same conclusion as above (which means it also cannot hold).
- As established above, even if K_1g and Zc_2g both intersect DK , they intersect different “components” of DK . We will now show that if either of K_1g or Zc_2g intersects DZc_3 , then the other one is disjoint from DK . Indeed, suppose $K_1g \cap DZc_3$ is nonempty, which is equivalent to the condition $g \in K_1^{-1}DZc_3$.
 - We have already established that in this scenario Zc_2g cannot intersect DZc_3 .
 - If $Zc_2g \cap DK_1$ is nonempty, then we have $g \in c_2^{-1}Z^{-1}DK_1$. It follows that the set $c_2^{-1}Z^{-1}DK_1 \cap K_1^{-1}DZc_3$ is nonempty, that is, $c_3 \in Z^{-1}D^{-1}K_1c_2^{-1}Z^{-1}DK_1$, contradicting condition (5e).
 - If $Zc_2g \cap DZc_2$, then we may conclude that $g \in c_2^{-1}Z^{-1}DZc_2$, so that $c_2^{-1}Z^{-1}DZc_2 \cap K_1^{-1}DZc_3$ is nonempty. Consequently, we have $c_3 \in Z^{-1}D^{-1}K_1c_2^{-1}Z^{-1}DZc_2$, contradicting (5f).

On the other hand, if $Zc_2g \cap DZc_3$ is nonempty, then $g \in c_2^{-1}Z^{-1}DZc_3$, and we reason as follows:

- We already know K_1g cannot intersect DZc_3 .
- If $K_1g \cap DK_1 \neq \emptyset$, then we have $g \in K_1^{-1}DK_1$. Thus, $c_2^{-1}Z^{-1}DZc_3 \cap K_1^{-1}DK_1$ is nonempty, hence $c_3 \in Z^{-1}D^{-1}Zc_2K_1^{-1}DK_1$, contradicting (5g).
- If $K_1g \cap DZc_2$, then $g \in K_1^{-1}DZc_2$. So, $K_1^{-1}DZc_2 \cap c_2^{-1}Z^{-1}DZc_3$ is nonempty, yielding $c_3 \in Z^{-1}D^{-1}Zc_2K_1^{-1}DZc_2$, and this is impossible by virtue of condition (5h).
- There remain only two possible cases where K_1g and Zc_2g both overlap DK , and we will show that in both of those situations Zc_3g must be disjoint from DK . The first case is that $K_1g \cap DK_1$ and $Zc_2g \cap DZc_2$ are both nonempty. This gives us $g \in K_1^{-1}DK_1$ and $g \in c_2^{-1}Z^{-1}DZc_2 (= P_{c_2})$, and we need to again consider three possibilities.
 - Suppose $Zc_3g \cap DZc_3 \neq \emptyset$, and thus $g \in c_3^{-1}Z^{-1}DZc_3 = P_{c_3}$, which means that $g \in P_{c_2} \cap P_{c_3}$, but by virtue of (5i), this means $g \in P$. However, we have already established in step (a) that g cannot be in P , so this situation is not possible.
 - If $Zc_3g \cap DK_1$ is nonempty, then it follows that $g \in c_3^{-1}Z^{-1}DK_1$. Combining that with the fact that $g \in K_1^{-1}DK_1$, we establish that $K_1^{-1}DK_1 \cap c_3^{-1}Z^{-1}DK_1$ is nonempty, and thus $c_3^{-1} \in K_1^{-1}DK_1K_1^{-1}D^{-1}Z$, contradicting (5j).

- If $Zc_3g \cap DZc_2$ is nonempty, then we have $g \in c_3^{-1}Z^{-1}DZc_2$. Combining this with the fact that $g \in c_2^{-1}Z^{-1}DZc_2$, we see that $c_3^{-1}Z^{-1}DZc_2 \cap c_2^{-1}Z^{-1}DZc_2 \neq \emptyset$, and therefore it can be seen that $c_3^{-1}Z^{-1}DZ \cap c_2^{-1}Z^{-1}DZ$ is also nonempty. This gives $c_3^{-1} \in c_2^{-1}Z^{-1}DZZ^{-1}D^{-1}Z$, contradicting (5k).
- The final possibility to exclude is that $K_1g \cap DZc_2$ and $Zc_2g \cap DK_1$ are both nonempty. These conditions translate into $g \in K_1^{-1}DZc_2$ and $g \in c_2^{-1}Z^{-1}DK_1$. For the last time, we need to consider three possible cases for Zc_3g .
 - If $Zc_3g \cap DZc_3 \neq \emptyset$, then $g \in c_3^{-1}Z^{-1}DZc_3 = P_{c_3}$. We also know that $g \in K_1^{-1}DZc_2$, but in light of condition (5l), this means $g \in P$, which is not possible.
 - If $Zc_3g \cap DK_1 \neq \emptyset$, then we obtain $g \in c_3^{-1}Z^{-1}DK_1$. Also, $g \in K_1^{-1}DZc_2$, so we have $c_3^{-1}Z^{-1}DK_1 \cap K_1^{-1}DZc_2 \neq \emptyset$, thus $c_3^{-1} \in K_1^{-1}DZc_2K_1^{-1}D^{-1}Z$, contradicting (5m).
 - If $Zc_3g \cap DZc_2 \neq \emptyset$, it follows that $g \in c_3^{-1}Z^{-1}DZc_2$. We also know that $g \in K_1^{-1}DZc_2$, so $c_3^{-1}Z^{-1}DZc_2 \cap K_1^{-1}DZc_2 \neq \emptyset$, and consequently, $c_3^{-1}Z^{-1}DZ \cap K_1^{-1}DZ$ is also nonempty. Thus, $c_3^{-1} \in K_1^{-1}DZZ^{-1}D^{-1}Z$, contradicting (5n).

We have shown that if $x|_{Kg} = x|_K = M$, then $Kg \setminus DK$ is a superset of at least one out of K_1g , Zc_2g , and Zc_3g . Thus, the block $x|_{Kg \setminus DK}$ has either M_1 or B as a subblock, but neither of those blocks occur in Y , and therefore, $x|_{Kg \setminus DK}$ also does not occur in Y . More generally, if $x|_{Kg_1} = x|_{Kg_2} = M$, then since $x|_{Kg_2} = x|_{Kg_2g_1^{-1}g_1} = (g_1x)|_{Kg_2g_1^{-1}}$, we can write $(g_1x)|_K = (g_1x)|_{Kg_2g_1^{-1}} = M$. It follows that $(g_1x)|_{Kg_2g_1^{-1} \setminus DK}$ is a block which does not occur in Y , but $(g_1x)|_{Kg_2g_1^{-1} \setminus DK} = x|_{Kg_2 \setminus DKg_1}$, so this block also does not occur in Y , as required. ■

Note that the tiling \mathcal{T} constructed above can (and probably will) have shapes smaller than the domain of M . This will not be an obstacle in our construction, but nevertheless we note that one can replace \mathcal{T} by a larger, congruent tiling (i.e., one whose tiles are unions of tiles of \mathcal{T}), the shapes of which can be arbitrarily large and have arbitrarily good invariance properties, and such a replacement will retain the properties specified in the theorem. Thus, we can make the following remark.

Remark 4.2. The tiling \mathcal{T} in Theorem 4.1 can be chosen to be (F, ε) -invariant for any $\varepsilon > 0$ and any finite $F \subset G$.

5. Constructing the extension

Theorem 5.1. *If G is a countable amenable group with the comparison property, and X is an aperiodic, strongly irreducible symbolic dynamical system with the shift action of G ,*

then for every $N \in \mathbb{N}$ such that $\log(N) < h(X)$, there exists a factor map from X onto the full G -shift over N symbols.

Proof. By Theorem 3.1, X has a subsystem Y such that $\log N < h(Y) < h(X)$. Before we delve into the technical details, here is a rough outline of the construction:

- (1) The factor map will determine if an element of X can be tiled (at least within some finite area around the neutral element of G) using a certain collection of large shapes.
- (2) This decision will be made based on the occurrences of a marker block within a certain window.
- (3) If such a local tiling can be found, there is a correspondence between blocks over its tiles and blocks in the full shift, which induces the image under the factor map (in other words, we define the map as a sliding block code). If the contents of x do not induce a local tiling (which is entirely possible, and in fact more likely than not), the code just assigns 0 to the image of the “problematic” coordinates.
- (4) To show that this map is a surjection, we tile every element of the full shift using large shapes and construct an element in X that only has marker blocks at the centers of such shapes, and blocks from Y elsewhere. Since $h(Y) > \log N$, the space not taken up by markers is enough to encode the entire element of the full shift.

We can apply Theorem 4.1 to obtain a block M with domain K and tiling \mathcal{T} with the following properties:

- * If T is a tile of \mathcal{T} and for some $x \in X$ we have $x|_{K_T} = M$, then the (nonempty) block $x|_{T_D \cap K_T}$ does not occur in Y .
- ** For any two different $g_1, g_2 \in G$, if $x|_{K_{g_1}} = x|_{K_{g_2}} = M$, then the (nonempty) block $x|_{K_{g_2} \setminus DK_{g_1}}$ does not occur in Y .

Let $\varepsilon = h(Y) - \log N$ and let E denote the union of the shapes of \mathcal{T} . There exists a δ such that if S is any (E, δ) -invariant set, then the union of tiles of \mathcal{T} which are contained in S is a $(1 - \frac{\varepsilon}{2})$ -subset of S .

We are about to apply Theorem 2.14 (note: we are not yet applying this theorem; we are just discussing one of its parameters) with the parameter δ , so we know it will yield a quasitiling with r_δ shapes $S_1, S_2, \dots, S_{r_\delta}$, where r_δ will depend only on δ . Let $K^* \subset G$ be a finite set such that $|\mathcal{B}_{K^*}(Y)| > r_\delta \cdot |E|$, and thus there exists a surjective function $\psi : \mathcal{B}_{K^*}(Y) \rightarrow \{1, \dots, r_\delta\} \times E$. We can also assume that K^* is disjoint from DK .

We can now apply Theorem 2.14 to obtain a quasitiling \mathcal{T}' such that:

- \mathcal{T}' has r_δ shapes.
- \mathcal{T}' factors onto a tiling \mathcal{T}'' with the same set of centers, and such that all shapes of \mathcal{T}'' are (E, δ) -invariant.

- Every shape S of \mathcal{T}'' is a superset of $DK \cup DK^*$, and in fact, the number of blocks with domain $S_D \setminus (DK \cup DK^*)$ that occur in Y is greater than $N^{|S|}$, hence there exists a surjective map $\Pi : \mathcal{B}_{S_D \setminus (DK \cup DK^*)}(Y) \rightarrow \{0, 1, \dots, N-1\}^S$.

(The latter property easily follows from the fact that if the shapes of \mathcal{T}' are sufficiently large Følner sets, then $S_D \setminus (DK \cup DK^*)$ can have arbitrarily large relative cardinality in S , and the entropy of Y exceeds $\log N$.)

We now have all the objects we need to define the factor map π from X onto the full G -shift over N symbols. We will do so by describing a procedure to determine the symbol $\pi(x)(e)$ based on $x|_{H'}$ for a certain finite $H' \subset G$.

Let H' be large enough that the knowledge of how the set of centers of \mathcal{T}' intersects H' allows us to determine the tile T'' of \mathcal{T}'' such that $e \in T''$. Consider the set of all $c \in H'$ such that $x|_{Kc} = M$. For every such c , if we have $\psi(x|_{K^*c}) = (j, g)$ (for $1 \leq j \leq r_\delta$ and $g \in E$), we obtain a set of the form S_jgc . There are now two possibilities:

- If the set of these shapes is equal to the intersection of H' with some right-translate of \mathcal{T}' , then it determines the intersection of H' with the same right-translate of \mathcal{T}'' , and in particular, it uniquely assigns e to a set of the form Sh for some shape S of \mathcal{T}'' and some $h \in H'$. If $x|_{S_D h \setminus (DK \cup DK^*)h}$ is some block A which occurs in Y , then let $\pi(x)(e) = \Pi(A)(h^{-1})$.
- If uniquely determining $x(e)$ as above is not possible (or the resulting block A does not occur in Y), set $\pi(x)(e) = 0$.

The map defined above is a sliding block code (with window H'), and thus it is a factor map from X onto some subset of the full G -shift over N symbols, and the only nontrivial property left to verify is surjectivity. In other words, we need to show that for every $z \in \{0, 1, \dots, N-1\}^G$ there exists some $x \in X$ such that $\pi(x) = z$. For such a z , we will define its preimage x by prescribing the content of x within D -interiors of disjoint sets (mostly tiles of \mathcal{T}); strong irreducibility means that such an x will exist provided the individual blocks do occur in X . Enumerate the tiles of \mathcal{T}' as T'_1, T'_2, \dots . For every k , the tile T'_k has the form $S'_{j(k)}c'_k$, where $j(k) \in \{1, 2, \dots, r_\delta\}$. The center c'_k belongs to some tile $S_k c_k$ of \mathcal{T} , and thus $c'_k = g_k c_k$ for some $g_k \in E$. The set K^*c_k is a subset of the D -interior of some union of finitely many tiles of \mathcal{T} ; denote this interior by \widehat{K}^*c_k . Let $x|_{Kc_k} = M$, let $x|_{[(S_k)_D \setminus K]c_k}$ be any block from Y , and let $x|_{\widehat{K}^*c_k}$ be a block from Y such that $\psi(x|_{K^*c_k}) = (j(k), g_k)$. Let $\widehat{S}_k c_k$ be the union of all tiles of \mathcal{T} that are disjoint from $DKc_k \cup D\widehat{K}^*c_k$ and contained within $(T'_k)_D$. There exists a block A in Y , with domain $(S_k)_D \setminus (DK \cup DK^*)$, such that $\Pi(A) = z(T'')$, where T'' is the tile of \mathcal{T}'' whose center is c_k (we know there is exactly one such tile), and we can extend A to a block A_k (also occurring with Y) with domain \widehat{S}_k . Set $x|_{\widehat{S}_k c_k} = A_k$. In addition, for any tile T of \mathcal{T} which is not a subset of $(T'_k)_D$ for any k , we can set $x|_{T_D}$ to be any block in Y .

The above construction, together with the properties of M , means that $x|_{Kc} = M$ if and only if $c = c_k$ for some k (this is because all D -interiors of tiles of \mathcal{T} , except the places where we explicitly put the marker, were chosen to be blocks from Y). This means

that for every g we can uniquely determine the tile of \mathcal{T}'' to which g belongs, based on the contents of $x|_{H'g}$, and hence $\pi(x) = z$. ■

References

- [1] M. Boyle, [Lower entropy factors of sofic systems](#). *Ergodic Theory Dynam. Systems* **3** (1983), no. 4, 541–557 Zbl [0529.28014](#) MR [0753922](#)
- [2] M. Boyle, R. Pavlov, and M. Schraudner, [Multidimensional sofic shifts without separation and their factors](#). *Trans. Amer. Math. Soc.* **362** (2010), no. 9, 4617–4653 Zbl [1207.37011](#) MR [2645044](#)
- [3] M. Boyle and M. Schraudner, [\$\mathbb{Z}^d\$ shifts of finite type without equal entropy full shift factors](#). *J. Difference Equ. Appl.* **15** (2009), no. 1, 47–52 Zbl [1161.37021](#) MR [2484416](#)
- [4] A. Desai, [A class of \$\mathbb{Z}^d\$ shifts of finite type which factors onto lower entropy full shifts](#). *Proc. Amer. Math. Soc.* **137** (2009), no. 8, 2613–2621 Zbl [1172.37008](#) MR [2497473](#)
- [5] T. Downarowicz, D. Huczek, and G. Zhang, [Tilings of amenable groups](#). *J. Reine Angew. Math.* **747** (2019), 277–298 Zbl [1411.37017](#) MR [3905135](#)
- [6] T. Downarowicz and G. Zhang, [Symbolic extensions of amenable group actions and the comparison property](#). *Mem. Amer. Math. Soc.* **281** (2023), no. 1390, vi+95p. Zbl [1520.37003](#) MR [4539364](#)
- [7] A. Johnson and K. Madden, [Factoring higher-dimensional shifts of finite type onto the full shift](#). *Ergodic Theory Dynam. Systems* **25** (2005), no. 3, 811–822 Zbl [1140.37304](#) MR [2142947](#)
- [8] E. Lindenstrauss and B. Weiss, [Mean topological dimension](#). *Israel J. Math.* **115** (2000), 1–24 Zbl [0978.54026](#) MR [1749670](#)
- [9] B. Marcus, [Factors and extensions of full shifts](#). *Monatsh. Math.* **88** (1979), no. 3, 239–247 Zbl [0432.54036](#) MR [0553733](#)
- [10] D. S. Ornstein and B. Weiss, [Entropy and isomorphism theorems for actions of amenable groups](#). *J. Analyse Math.* **48** (1987), 1–141 Zbl [0637.28015](#) MR [0910005](#)

Received 20 June 2021.

Dawid Huczek

Faculty of Pure and Applied Mathematics, Wrocław University of Science and Technology, Wybrzeże Wyspiańskiego 27, 50-370 Wrocław, Poland; dawid.huczek@pwr.edu.pl

Sebastian Kopacz

Faculty of Pure and Applied Mathematics, Wrocław University of Science and Technology, Wybrzeże Wyspiańskiego 27, 50-370 Wrocław, Poland; sebastian.kopacz@pwr.edu.pl