

Parabolic fractal geometry of stable Lévy processes with drift

Peter Kern and Leonard Pleschberger

Abstract. We explicitly calculate the Hausdorff dimension of the graph and range of an isotropic stable Lévy process X plus deterministic drift function f . For that purpose we use a restricted version of the genuine Hausdorff dimension, which is called the parabolic Hausdorff dimension. It turns out that covers by parabolic cylinders are optimal for treating self-similar processes, since their distinct non-linear scaling between time and space geometrically matches the self-similarity of the processes. We provide explicit formulas for the Hausdorff dimension of the graph and the range of $X + f$. In sum, the parabolic Hausdorff dimension of the drift term f alone contributes to the Hausdorff dimension of $X + f$. Furthermore, we derive some formulas and bounds for the parabolic Hausdorff dimension.

1. Introduction

Let $X = (X_t)_{t \geq 0}$ be a Lévy process in \mathbb{R}^d that is a stochastic process on some probability space $(\Omega, \mathcal{A}, \mathbb{P})$ with the following properties:

- (i) The process \mathbb{P} -almost surely starts in $0 \in \mathbb{R}^d$.
- (ii) X possesses independent increments, that is, for any $0 \leq t_0 < \dots < t_n$, the random variables $X_{t_0}, X_{t_1} - X_{t_0}, \dots, X_{t_n} - X_{t_{n-1}}$ are independent.
- (iii) X has stationary increments, that is, for all $t, h \geq 0$, the distribution of the increment $X_{t+h} - X_t \stackrel{d}{=} X_h$ does not depend on t , where the symbol $\stackrel{d}{=}$ denotes equality in distribution.
- (iv) X is stochastically continuous, that is, $\mathbb{P}(\|X_{t+h} - X_t\| > \varepsilon) \rightarrow 0$ as $h \rightarrow 0$ for any $t \geq 0$ and $\varepsilon > 0$.

Mathematics Subject Classification 2020: 60G51 (primary); 28A78, 28A80, 60G17, 60G18, 60G52 (secondary).

Keywords: stable Lévy process, drift function, fractal path behavior, parabolic Hausdorff dimension, self-similarity.

Additionally assuming self-similarity, the Lévy process is called stable. In this paper we only deal with the special case of an isotropic α -stable Lévy process, in which case the self-similarity is given by

$$(X_{c \cdot t})_{t \geq 0} \stackrel{\text{fd}}{=} (c^{1/\alpha} \cdot X_t)_{t \geq 0} \quad \text{for all } c > 0, \quad (1.1)$$

where $\stackrel{\text{fd}}{=}$ denotes equality of all finite-dimensional distributions that characterize the stochastic processes in law. In this case the Hurst index $H = 1/\alpha$ is restricted to $H \geq \frac{1}{2}$, that is, $\alpha \in (0, 2]$ and the isotropic α -stable Lévy process is also uniquely determined in law by the characteristic function $\mathbb{E}[e^{i\langle \xi, X_t \rangle}] = e^{-t \cdot C \|\xi\|^\alpha}$ with Lévy exponent $\Psi(\xi) = C \|\xi\|^\alpha$ for some constant $C > 0$. In case of $\alpha = 2$ we obtain Brownian motion. For details on stable Lévy processes, we refer to the monograph [15].

The integrability of $\xi \mapsto \exp(-t \cdot C \|\xi\|^\alpha)$ ensures the applicability of the Fourier inversion formula. Therefore, for any $t > 0$, the random variable X_t possesses the continuous density function

$$x \mapsto p(t, x) := (2\pi)^{-d} \int_{\mathbb{R}^d} e^{-i\langle \xi, x \rangle} e^{-t\Psi(\xi)} d\xi = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{-i\langle \xi, x \rangle} e^{-t \cdot C \|\xi\|^\alpha} d\xi,$$

which for $\alpha \in (0, 2)$ cannot be expressed in simple terms but belongs to $C^\infty(\mathbb{R}^d)$ with all derivatives in $L^1(\mathbb{R}^d) \cap C_0(\mathbb{R}^d)$; see [15]. Furthermore, from the self-similarity property in (1.1), it easily follows that

$$p(t, x) = t^{-d/\alpha} \cdot p\left(1, \frac{x}{t^{1/\alpha}}\right) \quad \text{for all } t > 0 \text{ and } x \in \mathbb{R}^d. \quad (1.2)$$

Thus, we define $p(x) := p(1, x)$ as the density at time $t = 1$ and by [4, Theorem 2.1], we have the tail estimate

$$p(x) = \mathcal{O}(\|x\|^{-d-\alpha}) \quad \text{as } \|x\| \rightarrow \infty. \quad (1.3)$$

This density is bounded and rotationally symmetric, that is, writing $x = ry \neq 0$ with $r = \|x\| > 0$ and $y = x/\|x\| \in S^{d-1}$, the density $p(x) = p(ry)$ does not depend on y and due to unimodality (see [15]), $r \mapsto p(ry)$ is non-increasing.

Our aim is to analyze isotropic stable Lévy processes plus (arbitrary) Borel measurable drift functions by methods from fractal geometry. In particular, we determine formulas for the Hausdorff dimension of the graph and the range of an isotropic α -stable Lévy process plus drift. Starting with Brownian motion, in the past decades much effort has been made to explicitly calculate the Hausdorff dimension of the range and the graph of stable Lévy processes with an even more general self-similarity relation than (1.1) (see, e.g., [1, 3, 5, 7, 10, 14, 17, 19] or the excellent review article [18]). Some of these results were extended to Markov processes, but require additional regularity assumptions for the transition probabilities in the non-homogeneous

situation; see [18] and the references therein for details. Only recently, Peres and Sousi started to deal with Hausdorff dimension results of self-similar processes with an additional drift function by considering Brownian motion [12] and fractional Brownian motion [13]. We will follow the method in [13] to prove corresponding results for isotropic stable Lévy processes. The restriction to isotropic stable Lévy processes is due to rotational symmetry, which is needed in the proof method. Compared to the method in [13], we have to overcome some additional issues:

- (1) An isotropic α -stable Lévy process for $\alpha \in (0, 2)$ is a pure jump process. Hence, we cannot use Hölder continuity of the sample paths to derive upper bounds for the Hausdorff dimension as in case of fractional Brownian motion in [13]. Instead, we will use the covering lemma of Pruitt and Taylor ([14, Lemma 6.1]) for the Lévy process together with a deterministic cover of the drift function.
- (2) The Hurst index $H = 1/\alpha$ of an isotropic α -stable Lévy process is restricted to $H \geq 1/2$, whereas $H \in (0, 1)$ for fractional Brownian motion. It will turn out that the case $H \geq 1$ needs different arguments than the blueprint given for $H \in (0, 1)$ in [13]. The reason is that diameters of parabolic cylinders introduced in (2.1) are determined by the size of the cylinders in the space domain for $H \in (0, 1)$, whereas for $H > 1$, the size of the cylinders in the time domain dominates.
- (3) By (1.3), the tail of the probability density of an isotropic α -stable Lévy process falls off as a power function, whereas the normal density of fractional Brownian motion decreases exponentially fast. The power law tails require more delicate estimates of energy kernels in the potential-theoretic method to derive lower bounds of the Hausdorff dimension.

We introduce a generalized version of the genuine Hausdorff dimension, which is called the α -parabolic Hausdorff dimension in Section 2. We also give a priori upper and lower bounds for the α -parabolic Hausdorff dimension in terms of the genuine Hausdorff dimension. It turns out that covers by α -parabolic cylinders are optimal for treating self-similar processes, since their distinct non-linear scaling between time and space geometrically matches the self-similarity of the processes. We provide explicit formulas for the Hausdorff dimension of the graph and the range of an isotropic α -stable Lévy process plus Borel measurable drift function in Section 3 and defer the proofs to Sections 4–6. In sum, the α -parabolic Hausdorff dimension of the drift term f alone contributes to the Hausdorff dimension of $X + f$. We derive new formulas and estimates for the α -parabolic Hausdorff dimension of constant functions and Hölder continuous functions in Section 7.

2. On parabolic fractal geometry

We start with the definition of the α -parabolic Hausdorff dimension, in which a distinct non-linear scaling between time and space inheres. Throughout this paper the symbol $|\cdot|$ denotes the diameter of a set in real Euclidean space. We use the same symbol for the absolute value of reals, without fear to cause confusion, and denote by $\|\cdot\|$ the Euclidean norm of vectors in \mathbb{R}^d or \mathbb{R}^{1+d} . For real functions f, g the symbol $f \lesssim g$ denotes the existence of a constant $C \in (0, \infty)$ not depending on the variables such that $f \leq C \cdot g$ and $f \asymp g$ is short for $f \lesssim g$ together with $g \lesssim f$.

Definition 2.1. Let $A \subseteq \mathbb{R}^{1+d}$ be any set and $\alpha, \beta \in (0, \infty)$. The α -parabolic β -Hausdorff (outer) measure of A is defined as

$$\mathcal{P}^\alpha - \mathcal{H}^\beta(A) := \lim_{\delta \downarrow 0} \inf \left\{ \sum_{k=1}^{\infty} |\mathbb{P}_k|^\beta : A \subseteq \bigcup_{n=1}^{\infty} \mathbb{P}_k, \mathbb{P}_k \in \mathcal{P}^\alpha, |\mathbb{P}_k| \leq \delta \right\},$$

where the α -parabolic cylinders $(\mathbb{P}_n)_{n \in \mathbb{N}}$ are contained in

$$\mathcal{P}^\alpha := \left\{ [t, t+c] \times \prod_{j=1}^d [x_j, x_j + c^{1/\alpha}] : t, x_j \in \mathbb{R}, c \in (0, 1] \right\}. \quad (2.1)$$

We define the α -parabolic Hausdorff dimension of A as

$$\mathcal{P}^\alpha\text{-dim} A := \inf \{ \beta > 0 : \mathcal{P}^\alpha - \mathcal{H}^\beta(A) = 0 \} = \sup \{ \beta > 0 : \mathcal{P}^\alpha - \mathcal{H}^\beta(A) = \infty \}.$$

The case $\alpha = 1$ equals the genuine Hausdorff dimension, which is simply denoted by the symbol \dim .

Let us compare the α -parabolic Hausdorff dimension to other parabolic Hausdorff dimensions appearing in the literature. For the Gaussian case, Taylor and Watson [16] introduced the *parabolic Hausdorff dimension* $\mathcal{P}\text{-dim}$ in the same way we did for $\alpha = 2$ in order to determine polar sets for the heat equation. On the contrary, for $H \in (0, 1]$, Peres and Sousi [13] defined the *H-parabolic β -Hausdorff content*

$$\Psi_H^\beta(A) := \inf \left\{ \sum_{k=1}^{\infty} c_k^\beta : A \subseteq \bigcup_{k=1}^{\infty} \mathbb{P}_k, \mathbb{P}_k \in \mathcal{P}^{1/H} \right\},$$

where the covering sets

$$\mathbb{P}_k = [t_k, t_k + c_k] \times \prod_{j=1}^d [x_{j,k}, x_{j,k} + c_k^H]$$

are from the class of $1/H$ -parabolic cylinders $\mathcal{P}^{1/H}$ from (2.1) with $|\mathbb{P}_k| \asymp c_k^H$. This results in what they call the H -parabolic Hausdorff dimension

$$\dim_{\Psi, H} A := \inf \{ \beta > 0 : \Psi_H^\beta(A) = 0 \} = \sup \{ \beta > 0 : \Psi_H^\beta(A) > 0 \}.$$

Proposition 2.2. *Let $A \subseteq \mathbb{R}^{1+d}$ be an arbitrary set, let $\alpha = 1/H \in [1, \infty)$, and let $\beta \in (0, \infty)$. Then, one has*

$$\mathcal{P}^\alpha\text{-dim} A = (\dim_{\Psi, H} A)/H, \quad (2.2)$$

that is, Peres and Sousi's H -parabolic Hausdorff dimension differs from our α -parabolic Hausdorff dimension by a constant factor.

Proof. First we introduce the auxiliary α -parabolic β -Hausdorff content

$$\Phi_\alpha^\beta(A) := \inf \left\{ \sum_{k=1}^{\infty} |\mathbb{P}_k|^\beta : A \subseteq \bigcup_{k=1}^{\infty} \mathbb{P}_k, \mathbb{P}_k \in \mathcal{P}^\alpha \right\}.$$

Since for $\alpha \geq 1$ and $\mathbb{P}_k = [t_k, t_k + c_k] \times \prod_{j=1}^d [x_{j,k}, x_{j,k} + c_k^{1/\alpha}] \in \mathcal{P}^\alpha$, the diameter fulfills $|\mathbb{P}_k|^\beta \asymp c_k^{\beta/\alpha} = c_k^{H \cdot \beta}$, one has

$$\Phi_\alpha^\beta(A) = \Psi_H^{\beta \cdot H}(A). \quad (2.3)$$

Following the arguments in [11, Proposition 4.9], one gets $\mathcal{P}^\alpha - \mathcal{H}^\beta(A) = 0$ if and only if $\Phi_\alpha^\beta(A) = 0$, which together with (2.3) shows (2.2). ■

The α -parabolic Hausdorff dimension fulfills the following countable stability property, which easily follows from monotonicity and σ -subadditivity of the α -parabolic Hausdorff measure, as argued for the genuine Hausdorff dimension in [6, page 29]:

Proposition 2.3. *For every $\alpha \in (0, \infty)$ and $(A_k)_{k \in \mathbb{N}} \subseteq \mathbb{R}^{1+d}$, the α -parabolic Hausdorff dimension fulfills the countable stability property*

$$\mathcal{P}^\alpha\text{-dim} \bigcup_{k=1}^{\infty} A_k = \sup_{k \in \mathbb{N}} \mathcal{P}^\alpha\text{-dim} A_k.$$

Moreover, we derive the following a priori estimates for the Hausdorff dimension in terms of the parabolic Hausdorff dimension:

Theorem 2.4. *Let $A \subseteq \mathbb{R}^{1+d}$ be any set. Let $\phi_\alpha = \mathcal{P}^\alpha\text{-dim} A$. Then, one has*

$$\dim A \leq \begin{cases} \phi_\alpha \wedge (\alpha \cdot \phi_\alpha + 1 - \alpha) & \alpha \in (0, 1], \\ \phi_\alpha \wedge \left(\frac{1}{\alpha} \cdot \phi_\alpha + \left(1 - \frac{1}{\alpha}\right) \cdot d \right) & \alpha \in [1, \infty) \end{cases}$$

and

$$\dim A \geq \begin{cases} \phi_\alpha + (1 - \frac{1}{\alpha}) \cdot d & \alpha \in (0, 1], \\ \phi_\alpha + 1 - \alpha & \alpha \in [1, \infty). \end{cases}$$

Proof. (i) Let $\alpha \in (0, \infty)$. By the definition of the α -parabolic β -Hausdorff measure, there are only coverings by \mathcal{P}^α -sets permitted. So, besides \mathcal{P}^α , there could exist more efficient covers of A with respect to their shape. Therefore, $\mathcal{H}^\beta(A) \leq \mathcal{P}^\alpha - \mathcal{H}^\beta(A)$, which implies $\dim A \leq \phi_\alpha$.

(ii) Let $\alpha \in (0, 1]$ and $\varepsilon > 0$ be arbitrary. If $\beta > \alpha \cdot \phi_\alpha + 1 - \alpha$, then $\frac{\beta}{\alpha} + (1 - \frac{1}{\alpha}) > \phi_\alpha$. Hence, we can cover A by the α -parabolic cylinders

$$(\mathbb{P}_k)_{k \in \mathbb{N}} = \left([t_k, t_k + c_k] \times \prod_{j=1}^d [x_{j,k}, x_{j,k} + c_k^{1/\alpha}] \right)_{k \in \mathbb{N}} \subseteq \mathcal{P}^\alpha$$

with $|\mathbb{P}_k| \asymp c_k$ for every $k \in \mathbb{N}$ such that $\sum_{k=1}^{\infty} |\mathbb{P}_k|^{\beta/\alpha + 1 - 1/\alpha} \leq \varepsilon$. Each \mathbb{P}_k can be covered by $\lceil c_k^{1-1/\alpha} \rceil$ hypercubes \square_k with sidelength $c_k^{1/\alpha}$. Hence,

$$\mathcal{H}^\beta(A) \leq \sum_{k=1}^{\infty} \lceil c_k^{1-1/\alpha} \rceil \cdot |\square_k|^\beta \lesssim \sum_{k=1}^{\infty} c_k^{\beta/\alpha + 1 - 1/\alpha} \lesssim \sum_{k=1}^{\infty} |\mathbb{P}_k|^{\beta/\alpha + 1 - 1/\alpha} \leq \varepsilon.$$

Since $\beta > \alpha \cdot \phi_\alpha + 1 - \alpha$ is arbitrary, we have $\dim A \leq \alpha \cdot \phi_\alpha + 1 - \alpha$.

(iii) Let $\alpha \in [1, \infty)$ and $\varepsilon > 0$ be arbitrary. If $\beta > 1/\alpha \cdot \phi_\alpha + (1 - 1/\alpha) \cdot d$, then we have $\alpha\beta + (1 - \alpha) \cdot d > \phi_\alpha$. Hence, a cover of A by α -parabolic cylinders $(\mathbb{P}_k)_{k \in \mathbb{N}}$ as in part (i) now fulfills $|\mathbb{P}_k| \asymp c_k^{1/\alpha}$ such that $\sum_{k=1}^{\infty} |\mathbb{P}_k|^{\alpha\beta + (1-\alpha) \cdot d} \leq \varepsilon$. Each \mathbb{P}_k can be covered by $\lceil c_k^{1/\alpha - 1} \rceil^d$ hypercubes \square_k with sidelength c_k . Then,

$$\begin{aligned} \mathcal{H}^\beta(A) &\leq \sum_{k=1}^{\infty} \lceil c_k^{1/\alpha - 1} \rceil^d \cdot |\square_k|^\beta \lesssim \sum_{k=1}^{\infty} (c_k^{1/\alpha})^{\alpha\beta + (1-\alpha) \cdot d} \lesssim \sum_{k=1}^{\infty} |\mathbb{P}_k^{1/\alpha}|^{\alpha\beta + (1-\alpha) \cdot d} \\ &\leq \varepsilon. \end{aligned}$$

Since $\beta > 1/\alpha \cdot \phi_\alpha + (1 - 1/\alpha) \cdot d$ is arbitrary, we have $\dim A \leq \frac{1}{\alpha} \cdot \phi_\alpha + (1 - \frac{1}{\alpha}) \cdot d$.

(iv) Let $\alpha \in (0, 1]$. Further, let $\beta > \dim A$ and $\varepsilon > 0$ be arbitrary. Then, we can cover A with hypercubes

$$(\square_k)_{k \in \mathbb{N}} = \left([t_k, t_k + c_k] \times \prod_{j=1}^d [x_{j,k}, x_{j,k} + c_k] \right)_{k \in \mathbb{N}} \subseteq \mathcal{P}^1$$

of sidelength $c_k \leq 1$ such that $\sum_{k=1}^{\infty} |\square_k|^\beta \leq \varepsilon$. Each \square_k can be covered by $\lceil c_k^{1-1/\alpha} \rceil^d$ α -parabolic cylinders

$$(\mathbb{P}_k)_{k \in \mathbb{N}} = \left([t_k, t_k + c_k] \times \prod_{j=1}^d [x_{j,k}, x_{j,k} + c_k^{1/\alpha}] \right)_{k \in \mathbb{N}} \subseteq \mathcal{P}^\alpha$$

with $|P_k| \asymp c_k$. By choosing $\gamma = \beta + (1/\alpha - 1) \cdot d$, one has

$$\begin{aligned} \mathcal{P}^\alpha - \mathcal{H}^\gamma(A) &\leq \sum_{k=1}^{\infty} [c_k^{1-1/\alpha}]^d \cdot |P_k|^\gamma \lesssim \sum_{k=1}^{\infty} c_k^{(1-1/\alpha)d+\gamma} = \sum_{k=1}^{\infty} c_k^\beta \lesssim \sum_{k=1}^{\infty} |\square_k|^\beta \\ &\leq \varepsilon. \end{aligned}$$

Since $\beta > \dim A$ is arbitrary, one has $\mathcal{P}^\alpha\text{-dim} A \leq \dim A + (\frac{1}{\alpha} - 1) \cdot d$.

(v) Let $\alpha \in [1, \infty)$. Each \square_k from part (iv) can be covered by $[c_k^{1-\alpha}]$ α -parabolic cylinders

$$(P_k)_{k \in \mathbb{N}} = \left([t_k, t_k + c_k^\alpha] \times \prod_{j=1}^d [x_{j,k}, x_{j,k} + c_k] \right) \subseteq \mathcal{P}^\alpha$$

with $|P_k| \asymp c_k$. By choosing $\gamma = \beta + \alpha - 1$, with similar calculations as above we get $\mathcal{P}^\alpha - \mathcal{H}^\gamma(A) \lesssim \varepsilon$. Since $\beta > \dim A$ is arbitrary, one has $\mathcal{P}^\alpha\text{-dim} A \leq \dim A + \alpha - 1$ and the theorem is proven. ■

3. Main results

So far, our considerations regarding the parabolic Hausdorff dimension were of a purely geometric nature. Now we will apply it to stochastic processes. We unite the cogitations of the following sections and begin with the Hausdorff dimension of the graph $\mathcal{G}_T(X + f) = \{(t, X_t + f(t)) : t \in T\}$ of an isotropic stable Lévy process X plus Borel measurable drift function f .

Theorem 3.1. *Let $T \subseteq \mathbb{R}_+$ be a Borel set and $\alpha \in (0, 2]$. Let $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d . Further, let $f : T \rightarrow \mathbb{R}^d$ be a Borel measurable function. Let $\varphi_\alpha = \mathcal{P}^\alpha\text{-dim} \mathcal{G}_T(f)$. Then, one \mathbb{P} -almost surely has*

$$\dim \mathcal{G}_T(X + f) = \begin{cases} \varphi_1 & \alpha \in (0, 1], \\ \varphi_\alpha \wedge \left(\frac{1}{\alpha} \cdot \varphi_\alpha + \left(1 - \frac{1}{\alpha}\right) \cdot d \right) & \alpha \in [1, 2]. \end{cases}$$

Proof. The Gaussian case where $\alpha = 2$ follows from [13, Theorem 1.2] together with Proposition 2.2. The other cases will follow by the combination of Corollary 4.2 with Theorem 5.7 for drift functions f with $\|f(t) - f(s)\| \leq 1$ for all $s, t \in T$. For the treatment of arbitrary drift functions f , we write $T = \bigcup_{z \in (\mathbb{Z}/2)^d} T_z$, where $T_z := \{t \in T : \|f(t) - z\| \leq \frac{1}{2}\}$. The claim now follows easily by using the countable stability property from Proposition 2.3. ■

The formula for the Hausdorff dimension of the range $\mathcal{R}_T(X + f) = \{X_t + f(t) : t \in T\}$ of an isotropic stable Lévy process X plus Borel measurable drift function f reads as follows:

Theorem 3.2. *Let $T \subseteq \mathbb{R}_+$ be a Borel set and $\alpha \in (0, 2]$. Let $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d and $f : T \rightarrow \mathbb{R}^d$ be a Borel measurable function. Let $\varphi_\alpha = \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f)$. Then, one \mathbb{P} -almost surely has*

$$\dim \mathcal{R}_T(X + f) = \begin{cases} (\alpha \cdot \varphi_\alpha) \wedge d & \alpha \in (0, 1], \\ \varphi_\alpha \wedge d & \alpha \in [1, 2]. \end{cases}$$

Proof. The Gaussian case where $\alpha = 2$ follows from [13, Theorem 1.2]. The rest will follow by Theorems 6.1 and 6.4, analogously to the proof of Theorem 3.1. ■

Our main theorems imply an improvement of Theorem 2.4 for graph sets.

Corollary 3.3. *Let $\varphi_\alpha = \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f)$. In the case of $\alpha \in (0, 1]$, one has*

$$\varphi_1 \geq (\alpha \cdot \varphi_\alpha) \vee \left(\varphi_\alpha + \left(1 - \frac{1}{\alpha}\right) \cdot d \right).$$

Proof. For $\alpha \in (0, 1]$, the combination of Theorems 3.1 and 3.2 directly yields

$$\varphi_1 = \dim \mathcal{G}_T(f) \geq \dim \mathcal{G}_T(X + f) \geq \dim \mathcal{R}_T(X + f) = (\alpha \cdot \varphi_\alpha) \wedge d.$$

Furthermore, we have

$$\alpha \cdot \varphi_\alpha \geq \varphi_\alpha + \left(1 - \frac{1}{\alpha}\right) \cdot d \quad \text{if and only if} \quad \alpha \cdot \varphi_\alpha \leq d$$

and

$$d \geq \varphi_\alpha + \left(1 - \frac{1}{\alpha}\right) \cdot d \quad \text{if and only if} \quad \alpha \cdot \varphi_\alpha \leq d,$$

which proves the claim. ■

4. Graph: Upper bound via geometric measure theory

We calculate an upper bound for the Hausdorff dimension of the graph of an isotropic stable Lévy process X plus drift function by means of an efficient covering.

Theorem 4.1. *Let $T \subseteq \mathbb{R}_+$ be any set and $\alpha \in (0, 2]$. Let $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d and $f : T \rightarrow \mathbb{R}^d$ be any function. Furthermore, let $\varphi_\alpha = \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f)$ be the α -parabolic Hausdorff dimension of the graph of f over T . Then, for $\alpha \in (0, 1]$ one \mathbb{P} -almost surely has*

$$\dim \mathcal{G}_T(X + f) \leq \dim \mathcal{G}_T(f) = \varphi_1,$$

and for $\alpha \in [1, 2]$, one \mathbb{P} -almost surely has

$$\mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(X + f) \leq \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f) = \varphi_\alpha.$$

Proof. (i) Let $\alpha \in (0, 1]$ and $\beta = \varphi_1$ and let $\delta, \varepsilon > 0$ be arbitrary. Then, $\mathcal{G}_T(f)$ can be covered by hypercubes

$$(\square_k)_{k \in \mathbb{N}} = \left([t_k, t_k + c_k] \times \prod_{i=1}^d [x_{i,k}, x_{i,k} + c_k] \right)_{k \in \mathbb{N}} \subseteq \mathcal{P}^1$$

such that $\sum_{k=1}^{\infty} |\square_k|^{\beta+\delta} \lesssim \sum_{k=1}^{\infty} c_k^{\beta+\delta} \leq \varepsilon$. Let $M_k(\omega)$ be the random number of a fixed 2^d -nested collection of hypercubes (see [14, Lemma 6.1] for the definition) with sidelength $c_k^{1/\alpha}$ that the path $t \mapsto X_t(\omega)$ hits at some time $t \in [t_k, t_k + c_k]$. Let $\bigcup_{k \in \mathbb{N}} P_k(\omega) \supseteq \mathcal{G}_T(X(\omega))$ with

$$(P_k(\omega))_{k \in \mathbb{N}} = \left([t_k, t_k + c_k] \times \bigcup_{j=1}^{M_k(\omega)} \prod_{i=1}^d [\xi_{i,j,k}(\omega), \xi_{i,j,k}(\omega) + c_k^{1/\alpha}] \right)_{k \in \mathbb{N}}$$

being a corresponding random parabolic cover of the graph of this path. Then, for all $t \in [t_k, t_k + c_k]$, there exists $j \in \{1, \dots, M_k(\omega)\}$ such that for the i -th component of $X + f$, we have

$$\begin{aligned} \xi_{i,j,k}(\omega) + x_{i,k} &\leq X_t^{(i)}(\omega) + f_i(t) \leq \xi_{i,j,k}(\omega) + x_{i,k} + c_k^{1/\alpha} + c_k \\ &\leq \xi_{i,j,k}(\omega) + x_{i,k} + 2c_k. \end{aligned}$$

Hence, we obtain a random cover $\bigcup_{k \in \mathbb{N}} \widetilde{\square}_k(\omega) \supseteq \mathcal{G}_T(X(\omega) + f)$, where

$$\begin{aligned} \widetilde{\square}_k(\omega) &= [t_k, t_k + c_k] \times \bigcup_{j=1}^{M_k(\omega)} \prod_{i=1}^d ([\xi_{i,j,k}(\omega) + x_{i,k}, \xi_{i,j,k}(\omega) + x_{i,k} + c_k] \\ &\quad \cup [\xi_{i,j,k}(\omega) + x_{i,k} + c_k, \xi_{i,j,k}(\omega) + x_{i,k} + 2c_k]). \end{aligned}$$

This is a union of $M_k(\omega) \cdot 2^d$ sets with diameter $\sqrt{d+1} \cdot c_k$. An application of Pruitt and Taylor's covering lemma (see [14, Lemma 6.1]) and [10, Lemma 3.4] shows that for all $\delta' > 0$, one has

$$\mathbb{E}[M_k] \lesssim \frac{c_k}{\mathbb{E}[T(c_k^{1/\alpha}/3, c_k)]} \lesssim c_k^{-\delta'/\alpha},$$

where $T(c_k^{1/\alpha}/3, c_k)$ is the sojourn time of the process $(X_t)_{t \in [0, c_k]}$ in a ball of radius $c_k^{1/\alpha}/3$ centered at the origin. Hence, we get for $\varepsilon' = \delta + \delta'/\alpha > 0$,

$$\mathbb{E}[\mathcal{H}^{\beta+\varepsilon'}(\mathcal{G}_T(X + f))] \leq \mathbb{E}\left[\sum_{k=1}^{\infty} |\widetilde{\square}_k|^{\beta+\varepsilon'}\right] \lesssim \sum_{k=1}^{\infty} \mathbb{E}[M_k] \cdot c_k^{\beta+\varepsilon'} \lesssim \sum_{k=1}^{\infty} c_k^{\beta+\delta} \leq \varepsilon.$$

Since $\varepsilon, \varepsilon' > 0$ are arbitrary, we get for all $\alpha \in (0, 1]$ and $\beta' > \beta$,

$$\mathbb{E}[\mathcal{H}^{\beta'}(\mathcal{G}_T(X + f))] = 0,$$

which implies $\mathcal{H}^{\beta'}(\mathcal{G}_T(X + f)) = 0$ \mathbb{P} -almost surely. Since $\beta' > \beta$ is arbitrary, we finally get \mathbb{P} -almost surely $\dim \mathcal{G}_T(X + f) \leq \beta = \varphi_1$.

(ii) Let $\alpha \in [1, 2]$ and $\beta = \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f)$ and also let $\varepsilon, \delta > 0$ be arbitrary. Then, $\mathcal{G}_T(f)$ can be covered by α -parabolic cylinders

$$(\mathbb{P}_k)_{k \in \mathbb{N}} = \left([t_k, t_k + c_k] \times \prod_{i=1}^d [x_{i,k}, x_{i,k} + c_k^{1/\alpha}] \right)_{k \in \mathbb{N}} \subseteq \mathcal{P}^\alpha$$

such that $\sum_{k=1}^{\infty} |\mathbb{P}_k|^{\beta+\delta} \lesssim \sum_{k=1}^{\infty} c_k^{(\beta+\delta)/\alpha} \leq \varepsilon$. Let $M_k(\omega)$ be the random number of a fixed 2^d -nested collection of hypercubes with sidelength $c_k^{1/\alpha}$ that the path $t \mapsto X_t(\omega)$ hits at some time $t \in [t_k, t_k + c_k]$. As in part (i), we obtain a random parabolic cover $\bigcup_{k \in \mathbb{N}} \tilde{\mathbb{P}}_k(\omega) \supseteq \mathcal{G}_T(X(\omega) + f)$, where

$$\begin{aligned} \tilde{\mathbb{P}}_k(\omega) = [t_k, t_k + c_k] \times & \bigcup_{j=1}^{M_k(\omega)} \prod_{i=1}^d ([\xi_{i,j,k}(\omega) + x_{i,k}, \xi_{i,j,k}(\omega) + x_{i,k} + c_k^{1/\alpha}] \\ & \cup [\xi_{i,j,k}(\omega) + x_{i,k} + c_k^{1/\alpha}, \xi_{i,j,k}(\omega) + x_{i,k} + 2c_k^{1/\alpha}]). \end{aligned}$$

This is a union of $M_k(\omega) \cdot 2^d$ sets with diameter $|\tilde{\mathbb{P}}_k(\omega)| \lesssim c_k^{1/\alpha}$. As in part (i), we get $\mathbb{E}[M_k] \lesssim c_k^{-\delta'/\alpha}$. Hence, we get for $\varepsilon' = \delta + \delta' > 0$ with the similar calculations as above,

$$\mathbb{E}[\mathcal{P}^\alpha - \mathcal{H}^{\beta+\varepsilon'}(\mathcal{G}_T(X + f))] \lesssim \sum_{k=1}^{\infty} \mathbb{E}[M_k] \cdot c_k^{(\beta+\varepsilon')/\alpha} \lesssim \sum_{k=1}^{\infty} c_k^{(\beta+\delta)/\alpha} \leq \varepsilon.$$

Since $\varepsilon, \varepsilon' > 0$ are arbitrary, as in part (i), we finally get

$$\mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(X + f) \leq \beta = \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f) \quad \mathbb{P}\text{-almost surely,}$$

as claimed. ■

Corollary 4.2. *Let $T \subseteq \mathbb{R}_+$ be any set and $\alpha \in (0, 2]$. Let $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d and $f : T \rightarrow \mathbb{R}^d$ be any function. Furthermore, let $\varphi_\alpha = \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f)$. Then, one \mathbb{P} -almost surely has*

$$\dim \mathcal{G}_T(X + f) \leq \begin{cases} \varphi_1 & \alpha \in (0, 1], \\ \varphi_\alpha \wedge \left(\frac{1}{\alpha} \cdot \varphi_\alpha + \left(1 - \frac{1}{\alpha}\right) \cdot d \right) & \alpha \in [1, 2]. \end{cases}$$

Proof. The Gaussian case $\alpha = 2$ follows from [13, Corollary 2.3] and Proposition 2.2. The rest follows directly from Theorems 2.4 and 4.1. ■

5. Graph: Lower bound via potential theory

Next we want to calculate a lower bound for the Hausdorff dimension of isotropic stable Lévy processes with drift. This will be accomplished by the energy method; see [11, Section 4.3]. This method makes use of the Lebesgue integral. Hence, for the first time we have to impose restrictions on the domain $T \subseteq \mathbb{R}_+$ and the drift function $f : T \rightarrow \mathbb{R}^d$ with regard to their measurability. For a Borel-measurable function, it is well known that the graph is always a Borel set, whereas the range is not necessarily a Borel set, but belongs to the Suslin sets. Suslin sets (also called analytic sets) supersede the Borel sets and can be represented as the image of a Borel set under a continuous mapping. For details on Suslin sets, we refer to [8, Section 39]. We introduce some notions from potential theory in this slightly more general setting, to be also applicable for the range in Section 6.

Definition 5.1. Let $A \subseteq \mathbb{R}^{1+d}$ be a Suslin set and μ be a probability measure supported on A , that is, $\mu \in \mathcal{M}^1(A)$. Further, let $K : \mathbb{R}^{1+d} \rightarrow [0, \infty]$ be a Lebesgue measurable function that is called the *difference kernel*. The K -energy of a probability measure μ is defined to be

$$\mathcal{E}_K(\mu) := \int_A \int_A K(t-s, x-y) d\mu(t, x) d\mu(s, y)$$

and the *equilibrium value* of A is defined as $\mathcal{E}_K^* := \inf_{\mu \in \mathcal{M}^1(A)} \mathcal{E}_K(\mu)$. We define the K -capacity of A as

$$\text{Cap}_K(A) := \frac{1}{\mathcal{E}_K^*}.$$

Whenever the kernel has the form $K(t, x) = \|(t, x)\|^{-\beta}$, we write $\mathcal{E}_\beta(\mu)$ for $\mathcal{E}_K(\mu)$ and $\text{Cap}_\beta(A)$ for $\text{Cap}_K(A)$ and we refer to them as the β -energy of a probability measure μ and the Riesz β -capacity of A , respectively. Next we state Frostman's theorem.

Theorem 5.2. Let $\alpha > 0$. For any Suslin set $A \subseteq \mathbb{R}^{1+d}$, one has

$$\mathcal{P}^\alpha\text{-dim} A \geq \dim A = \sup\{\beta : \text{Cap}_\beta(A) > 0\}.$$

Proof. This follows from the first assertion in the proof of Theorem 2.4 together with [2, Appendix B]. ■

The next lemma shows that we can work with an energy integral where the stable process X is transformed into the kernel.

Lemma 5.3. Let $T \subseteq \mathbb{R}_+$ be a Borel set and $\alpha \in (0, 2]$. Let $X = (X_t)_{t \geq 0}$ be a stochastic process in \mathbb{R}^d with stationary increments and $f : \mathbb{R}_+ \rightarrow \mathbb{R}^d$ be a Borel measurable

function. Define the difference kernel

$$K^\beta(t, x) := \mathbb{E}[\|(t, \text{sign}(t) \cdot X_{|t|} + x)\|^{-\beta}].$$

Then, from $\text{Cap}_{K^\beta}(\mathcal{G}_T(f)) > 0$, it follows that \mathbb{P} -almost surely $\text{Cap}_\beta(\mathcal{G}_T(X + f)) > 0$ holds. Hence, $\mathcal{E}_{K^\beta}(\mu) < \infty$ for some probability measure $\mu \in \mathcal{M}^1(\mathcal{G}_T(f))$ implies

$$\dim \mathcal{G}_T(X + f) \geq \beta \quad \mathbb{P}\text{-almost surely.}$$

Proof. For every $\omega \in \Omega$, the pathwise bijection that $(t, f(t)) \in \mathcal{G}_T(f)$ if and only if $(t, X_t(\omega) + f(t)) \in \mathcal{G}_T(X_t(\omega) + f)$ yields the existence of some random probability measure $\nu_\omega \in \mathcal{M}^1(\mathcal{G}_T(X(\omega) + f))$ with $\nu_\omega(\tilde{A}_\omega) = \mu(A)$ for all Borel sets $A \subseteq \mathcal{G}_T(f)$, where $\tilde{A}_\omega := \{(t, x + X_t(\omega)) : (t, x) \in A\}$. Therefore, Tonelli's theorem and the stationarity of the increments of X yield

$$\begin{aligned} \mathbb{E}[\mathcal{E}_\beta(\nu_\omega)] &= \mathbb{E}\left[\int_{\mathcal{G}_T(X(\omega)+f)} \int_{\mathcal{G}_T(X(\omega)+f)} \|(t-s, x-y)\|^{-\beta} d\nu_\omega(t, x) d\nu_\omega(s, y)\right] \\ &= \mathbb{E}\left[\int_{\mathcal{G}_T(f)} \int_{\mathcal{G}_T(f)} \|(t-s, x + X_t(\omega) - (y + X_s(\omega)))\|^{-\beta} d\mu(t, x) d\mu(s, y)\right] \\ &= \int_{\mathcal{G}_T(f)} \int_{\mathcal{G}_T(f)} \mathbb{E}[\|(t-s, X_t(\omega) - X_s(\omega) + x - y)\|^{-\beta}] d\mu(t, x) d\mu(s, y) \\ &= \int_{\mathcal{G}_T(f)} \int_{\mathcal{G}_T(f)} \mathbb{E}[\|(t-s, \text{sign}(t-s) \cdot X_{|t-s|}(\omega) + x - y)\|^{-\beta}] d\mu(t, x) d\mu(s, y) \\ &= \mathcal{E}_{K^\beta}(\mu) \end{aligned}$$

By assumption, there exists $\mu \in \mathcal{M}^1(\mathcal{G}_T(f))$ such that $\mathcal{E}_{K^\beta}(\mu) < \infty$; therefore, $\mathcal{E}_\beta(\nu_\omega) < \infty$ \mathbb{P} -almost surely. The rest of the claim follows by Frostman's theorem (Theorem 5.2). \blacksquare

Frostman's lemma provides the suitable candidate for the probability measure μ . We give a parabolic version of it.

Theorem 5.4. *Let $A \subseteq \mathbb{R}^{1+d}$ be a Borel set. If $\mathcal{P}^\alpha\text{-dim} A > \beta$, then there exists $\mu \in \mathcal{M}^1(A)$ such that we have*

$$\mu\left([t, t+c] \times \prod_{i=1}^d [x_i, x_i + c^{1/\alpha}]\right) \lesssim \begin{cases} c^\beta & \alpha \in (0, 1], \\ c^{\beta/\alpha} & \alpha \in [1, \infty) \end{cases}$$

for every $c \in (0, 1]$ and $t, x_1, \dots, x_d \in \mathbb{R}$.

Proof. The parabolic case can easily be proven along the lines of the classical case; see, for example, [11, Section 4.4]. \blacksquare

The following lemma is a refinement of [13, (2.7)]:

Lemma 5.5. *Let $t \in \mathbb{R}$ be fixed and $h : \mathbb{R}^d \rightarrow \mathbb{R}$, $h(x) = \|(t, x)\|^{-\beta} = (t^2 + \|x\|^2)^{-\beta/2}$. Then, h is rotationally symmetric and the mapping $r \mapsto h(r \cdot y)$ is non-increasing for $r = \|x\|$ and does not depend on $y = x/\|x\| \in S^{d-1}$. Further, let $p : \mathbb{R}^d \rightarrow \mathbb{R}$ be a rotationally symmetric function such that also $r \mapsto p(r \cdot y)$ is non-increasing for $r = \|x\|$ and $y = x/\|x\| \in S^{d-1}$. Then, for all $u \in \mathbb{R}^d$, we have*

$$\int_{\mathbb{R}^d} h(x+u) \cdot p(x) \, dx \lesssim \int_{\mathbb{R}^d} h(x) \cdot p(x) \, dx,$$

provided that the integrals exist.

Proof. The first part is obvious. Further, by monotonicity, we have

$$\begin{aligned} & \int_{\mathbb{R}^d} h(x+u) \cdot p(x) \, dx \\ &= \int_{\{\|x\| < \|x+u\|\}} \underbrace{h(x+u) \cdot p(x)}_{\leq h(x)} \, dx + \int_{\{\|x\| \geq \|x+u\|\}} h(x+u) \cdot \underbrace{p(x)}_{\leq p(x+u)} \, dx \\ &\leq 2 \int_{\mathbb{R}^d} h(x) \cdot p(x) \, dx, \end{aligned}$$

as claimed. ■

Inspired by [13, Lemma 2.5], we give a priori estimates for the difference kernel K^β from Lemma 5.3 that will later turn out to provide appropriate estimates of the energy integral.

Lemma 5.6. *Let $\alpha \in (0, 2)$ and $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d . Let $\beta \geq 0$, $\tau \in \mathbb{R}$, and $\delta \in \mathbb{R}^d$ be such that $|\tau| \in (0, 1]$, $\|\delta\| \in [0, 1]$. Then, for the difference kernel $K^\beta(\tau, \delta) = \mathbb{E}[\|(\tau, \text{sign}(\tau) \cdot X_{|\tau|} + \delta)\|^{-\beta}]$ from Lemma 5.3, one has*

$$K^\beta(\tau, \delta) \lesssim \begin{cases} |\tau|^{-\beta}, \\ |\tau|^{-\beta/\alpha} & \text{for } \beta < d, \\ |\tau|^{(1-1/\alpha)d-\beta} & \text{for } \beta > d \end{cases} \quad (5.1)$$

and

$$K^\beta(\tau, \delta) \lesssim \begin{cases} \|\delta\|^{-\beta} & \text{for } \alpha \in (0, 1], |\tau| \leq \|\delta\|, \\ \|\delta\|^{-\beta} & \text{for } \alpha \in [1, 2), \beta < d, |\tau| \leq \|\delta\|^\alpha, \\ \|\delta\|^{(\alpha-1)d-\alpha\beta} & \text{for } \alpha \in [1, 2), \beta > d, |\tau| \leq \|\delta\|^\alpha. \end{cases} \quad (5.2)$$

Proof. Let $p(x)$ denote the density function of $X_1 \stackrel{d}{=} |\tau|^{-1/\alpha} X_{|\tau|}$. We define rescaled increments $\tilde{\tau} := \tau/|\tau|^{1/\alpha}$ and $\tilde{\delta} := \delta/|\tau|^{1/\alpha}$. Trivial estimation always yields

$$\mathbb{E}[\|(\tau, \text{sign}(\tau) \cdot X_{|\tau|} + \delta)\|^{-\beta}] \leq |\tau|^{-\beta}.$$

The self-similarity of the stable Lévy process and Lemma 5.5 yield

$$\begin{aligned}
 \mathbb{E}[\|(\tau, \text{sign}(\tau) \cdot X_{|\tau|} + \delta)\|^{-\beta}] &= |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|(\tilde{\tau}, \text{sign}(\tau) \cdot x + \tilde{\delta})\|^{-\beta} \cdot p(x) \, dx \\
 &= |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|(|\tilde{\tau}|, x + \text{sign}(\tau) \cdot \tilde{\delta})\|^{-\beta} \cdot p(x) \, dx \\
 &\lesssim |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|(|\tilde{\tau}|, x)\|^{-\beta} \cdot p(x) \, dx.
 \end{aligned} \tag{5.3}$$

Let $\beta < d$. Then, by (5.3), we get

$$\begin{aligned}
 \mathbb{E}[\|(\tau, \text{sign}(\tau) \cdot X_{|\tau|} + \delta)\|^{-\beta}] &\lesssim |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|x\|^{-\beta} \cdot p(x) \, dx \\
 &\lesssim |\tau|^{-\beta/\alpha} \cdot \mathbb{E}[\|X_1\|^{-\beta}] \lesssim |\tau|^{-\beta/\alpha},
 \end{aligned}$$

since negative moments of order $\beta < d$ exist; see [1, Lemma 3.1].

Let $\beta > d$. Then, by (5.3) one has, using the volume of a ball with radius $\tilde{\tau}$,

$$\begin{aligned}
 \mathbb{E}[\|(\tau, \text{sign}(\tau) \cdot X_{|\tau|} + \delta)\|^{-\beta}] &\lesssim |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|(|\tilde{\tau}|, x)\|^{-\beta} \cdot p(x) \, dx \\
 &\leq |\tau|^{-\beta/\alpha} \left(\int_{\{\|x\| < |\tilde{\tau}|\}} |\tilde{\tau}|^{-\beta} \cdot p(x) \, dx + \int_{\{\|x\| \geq |\tilde{\tau}|\}} \|x\|^{-\beta} \cdot p(x) \, dx \right) \\
 &\leq |\tau|^{-\beta/\alpha} \left(|\tilde{\tau}|^{d-\beta} + \int_{\{|\tilde{\tau}| \leq \|x\| \leq 1\}} \|x\|^{-\beta} \, dx + \int_{\{\|x\| > 1\}} p(x) \, dx \right) \\
 &\leq |\tau|^{-\beta/\alpha} \left(|\tilde{\tau}|^{d-\beta} + \int_{|\tilde{\tau}|}^1 \int_{S^{d-1}} \|ry\|^{-\beta} \cdot r^{d-1} \, dy \, dr + 1 \right) \\
 &\lesssim |\tau|^{-\beta/\alpha} \left(|\tilde{\tau}|^{d-\beta} + \int_{|\tilde{\tau}|}^1 r^{d-\beta-1} \, dr \right) \\
 &\lesssim |\tau|^{-\beta/\alpha} \cdot |\tilde{\tau}|^{d-\beta} = |\tau|^{-\beta/\alpha} \cdot |\tau|^{(1-1/\alpha)(d-\beta)} = |\tau|^{(1-1/\alpha)d-\beta}.
 \end{aligned}$$

This proves (5.1). To prove (5.2), consider the region $\|x\| \leq \tilde{|\delta|}/2$, which yields

$$\|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \geq \| \|x\| - \tilde{|\delta|} \| = \|\tilde{\delta}\| - \|x\| \geq \frac{1}{2} \cdot \|\tilde{\delta}\|.$$

Thus, for the estimates in (5.2), we have

$$\begin{aligned}
 \mathbb{E}[\|(\tau, \text{sign}(\tau) \cdot X_{|\tau|} + \delta)\|^{-\beta}] &= |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|(\tilde{\tau}, \text{sign}(\tau) \cdot x + \tilde{\delta})\|^{-\beta} \cdot p(x) \, dx \\
 &\lesssim \underbrace{\|\delta\|^{-\beta} + |\tau|^{-\beta/\alpha} \int_{\{\|x\| \geq \|\tilde{\delta}\|/2, \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \geq |\tilde{\tau}|\}}}_{=: I_1} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} \cdot p(x) \, dx
 \end{aligned}$$

$$+ \underbrace{|\tau|^{-\beta/\alpha} \int_{\{\|x\| \geq \|\tilde{\delta}\|/2, \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \leq |\tilde{\tau}|\}} \tilde{\tau}^{-\beta} \cdot p(x) dx}_{=: I_2}.$$

Now,

$$\begin{aligned} I_1 &= |\tau|^{-\beta/\alpha} \int_{\{\|x\| \geq \|\tilde{\delta}\|/2, \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \geq |\tilde{\tau}|\}} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} \cdot p(x) dx \\ &= |\tau|^{-\beta/\alpha} \int_{\{\|x\| \geq \|\tilde{\delta}\|/2, \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \geq \max(|\tilde{\tau}|, \|\tilde{\delta}\|)\}} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} \cdot p(x) dx \\ &\quad + |\tau|^{-\beta/\alpha} \int_{\{\|x\| \geq \|\tilde{\delta}\|/2, \|\tilde{\delta}\| \geq \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \geq |\tilde{\tau}|\}} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} \cdot p(x) dx \\ &\leq \|\delta\|^{-\beta} + \underbrace{|\tau|^{-\beta/\alpha} \int_{\{\|x\| \geq \|\tilde{\delta}\|/2, \|\tilde{\delta}\| \geq \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \geq |\tilde{\tau}|\}} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} \cdot p(x) dx}_{=: I_3}. \end{aligned}$$

By using (1.3), we further have

$$\begin{aligned} I_3 &= |\tau|^{-\beta/\alpha} \int_{\{\|x\| \geq \|\tilde{\delta}\|/2, \|\tilde{\delta}\| \geq \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \geq |\tilde{\tau}|\}} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} \cdot p(x) dx \\ &\lesssim |\tau|^{-\beta/\alpha} \cdot \|\tilde{\delta}\|^{-d-\alpha} \int_{\{\|\tilde{\delta}\| \geq \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \geq |\tilde{\tau}|\}} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} dx \\ &= |\tau|^{-\beta/\alpha} \cdot \|\tilde{\delta}\|^{-d-\alpha} \int_{|\tilde{\tau}|}^{\|\tilde{\delta}\|} r^{d-\beta-1} dr. \end{aligned} \tag{5.4}$$

For $\alpha \in [1, 2)$, $\beta < d$, and $|\tau| \leq \|\delta\|^\alpha$, by (5.4) we get

$$I_3 \lesssim |\tau|^{-\beta/\alpha} \cdot \|\tilde{\delta}\|^{-\alpha-\beta} = |\tau| \cdot \|\delta\|^{-\alpha-\beta} \lesssim \|\delta\|^{-\beta},$$

whereas for $\alpha \in (0, 1]$ and $|\tau| \leq \|\delta\|$, one has

$$I_3 \lesssim |\tau| \cdot \|\delta\|^{-\alpha-\beta} \leq \|\delta\|^{1-\alpha} \cdot \|\delta\|^{-\beta} \lesssim \|\delta\|^{-\beta}.$$

For $\alpha \in [1, 2)$, $\beta > d$, and $|\tau| \leq \|\delta\|^\alpha$, by (5.4) one has

$$\begin{aligned} I_3 &\lesssim |\tau|^{-\beta/\alpha} \cdot \|\tilde{\delta}\|^{-d-\alpha} \int_{|\tilde{\tau}|}^{\infty} r^{-\beta} \cdot r^{d-1} dr \lesssim |\tau|^{-\beta/\alpha} \cdot \|\tilde{\delta}\|^{-d-\alpha} \cdot \tilde{\tau}^{d-\beta} \\ &= |\tau|^{d+1-\beta} \cdot \|\delta\|^{-d-\alpha} \leq \|\delta\|^{\alpha(d+1-\beta)} \cdot \|\delta\|^{-d-\alpha} = \|\delta\|^{(\alpha-1)d-\alpha\beta}. \end{aligned}$$

Finally, by using (1.3), we get

$$I_2 = |\tau|^{-\beta/\alpha} \int_{\{\|x\| \geq \|\tilde{\delta}\|/2, \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \leq |\tilde{\tau}|\}} |\tilde{\tau}|^{-\beta} \cdot p(x) dx$$

$$\begin{aligned}
 &\lesssim |\tau|^{-\beta/\alpha} \cdot |\tilde{\tau}|^{-\beta} \int_{\|x + \text{sign}(\tau) \cdot \tilde{\delta}\| \leq |\tilde{\tau}|} \|x\|^{-d-\alpha} dx \\
 &\lesssim |\tau|^{-\beta/\alpha} \cdot |\tilde{\tau}|^{-\beta} \cdot \|\tilde{\delta}\|^{-d-\alpha} \cdot |\tilde{\tau}|^d = |\tau|^{d-\beta+1} \cdot \|\delta\|^{-d-\alpha},
 \end{aligned}$$

using the volume of a ball with radius $|\tilde{\tau}|$ and center $-\text{sign}(\tau) \cdot \tilde{\delta}$. Now, $\alpha \in (0, 1]$, $\beta < d$, and $|\tau| \leq \|\delta\|$ result in

$$I_2 \lesssim |\tau|^{d-\beta+1} \cdot \|\delta\|^{-d-\alpha} \leq \|\delta\|^{1-\alpha-\beta} \leq \|\delta\|^{-\beta}.$$

If $\alpha \in [1, 2)$, $\beta \leq d$, and $|\tau| \leq \|\delta\|^\alpha$ one has

$$I_2 \lesssim |\tau|^{d-\beta+1} \cdot \|\delta\|^{-d-\alpha} \leq \|\delta\|^{(\alpha-1)d-\alpha\beta} \leq \|\delta\|^{-\beta}.$$

If $\alpha \in [1, 2)$, $\beta \geq d$, and $|\tau| \leq \|\delta\|^\alpha$ one has

$$I_2 \lesssim \|\delta\|^{(\alpha-1)d-\alpha\beta} \quad \text{and} \quad \|\delta\|^{-\beta} \leq \|\delta\|^{(\alpha-1)d-\alpha\beta}.$$

Altogether, we have shown

$$\mathbb{E}[\|(\tau, \text{sign}(\tau) \cdot X_{|\tau|} + \delta)\|^{-\beta}] \lesssim \|\delta\|^{-\beta} + I_1 + I_2 \lesssim \|\delta\|^{-\beta} + I_3 + I_2$$

and our upper bounds for I_2 and I_3 directly yield (5.2). \blacksquare

Now, we are able to calculate the lower bound via energy estimates.

Theorem 5.7. *Let $T \subseteq \mathbb{R}_+$ be a Borel set and $\alpha \in (0, 2)$. Let $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d and $f : T \rightarrow \{y \in \mathbb{R}^d : \|y - x\| \leq \frac{1}{2}\}$ for fixed $x \in \mathbb{R}^d$ be a Borel measurable function. Let $\varphi_\alpha = \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f)$. Then, one \mathbb{P} -almost surely has*

$$\dim \mathcal{G}_T(X + f) \geq \begin{cases} \varphi_1 & \alpha \in (0, 1], \\ \varphi_\alpha \wedge \left(\frac{1}{\alpha} \cdot \varphi_\alpha + \left(1 - \frac{1}{\alpha}\right) \cdot d\right) & \alpha \in [1, 2]. \end{cases} \quad (5.5)$$

Proof. We define the increments $\tau := t - s$ and $\delta := f(t) - f(s)$ with $\|\delta\| \in [0, 1]$ and consider the difference kernel $K^\beta(t, x) = \mathbb{E}[\|(t, \text{sign}(t) \cdot X_{|t|} + x)\|^{-\beta}]$. We prove that $\mathcal{E}_{K^\beta}(\mu) < \infty$ holds for $\mu \in \mathcal{M}^1(\mathcal{G}_T(f))$ from the parabolic version of Frostman's lemma in Theorem 5.4 and for every β less than the right-hand side of (5.5). Then, the claim follows due to Lemma 5.3. For the energy integral, we have

$$\begin{aligned}
 \mathcal{E}_{K^\beta}(\mu) &= \int \int_{\mathcal{G}_T(f) \times \mathcal{G}_T(f)} K^\beta(t - s, f(t) - f(s)) d\mu(s, x) d\mu(t, y) \\
 &\leq \int \int_{\{|t-s| \in (0, 1]\}} K^\beta(\tau, \delta) d\mu d\mu + \int \int_{\{|t-s| \in (1, \infty)\}} |t - s|^{-\beta} d\mu d\mu \\
 &\lesssim \int \int_{\{|\tau| \in (0, 1]\}} \mathbb{E}[\|(\tau, \text{sign}(\tau) \cdot X_{|\tau|}(\omega) + \delta)\|^{-\beta}] d\mu d\mu. \quad (5.6)
 \end{aligned}$$

(i) We begin with the case $\alpha \in (0, 1]$ and $\beta = \varphi_1 - 2\varepsilon$ for some arbitrary $\varepsilon > 0$. Due to Lemma 5.6, we have

$$\begin{aligned} \mathcal{E}_{K^\beta}(\mu) &\lesssim \underbrace{\int \int_{\{|\tau| \in (0, 1], \|\delta\| \in [0, |\tau|]\}} |\tau|^{-\beta} d\mu d\mu}_{=: I_1} \\ &\quad + \underbrace{\int \int_{\{|\tau| \in (0, 1], \|\delta\| \in (|\tau|, 1]\}} \|\delta\|^{-\beta} d\mu d\mu}_{=: I_2}. \end{aligned}$$

We get

$$I_1 \lesssim \sum_{k=1}^{\infty} 2^{k\beta} \cdot \mu \otimes \mu\{|\tau| \in (2^{-k}, 2 \cdot 2^{-k}], \|\delta\| \in [0, 2 \cdot 2^{-k}]\}.$$

Further,

$$I_2 \lesssim \sum_{k=1}^{\infty} 2^{k\beta} \cdot \mu \otimes \mu\{|\tau| \in (0, 2 \cdot 2^{-k}], \|\delta\| \in (2^{-k}, 2 \cdot 2^{-k}]\}.$$

Now we have to calculate the expressions $\mu \otimes \mu\{\cdot\}$ for I_1 and I_2 . For each $k \in \mathbb{N}$ we tile $\mathbb{R}_+ \times \mathbb{R}^d$ by disjoint hypercubes of size $2^{-k} \times \dots \times 2^{-k}$ and denote the collection of such hypercubes by \mathcal{D}_k . For every $c \in (0, 1]$, $\gamma = \varphi_1 - \varepsilon$ and $\alpha \in (0, 1]$ Frostman's lemma 5.4 yields

$$\mu\left([t, t+c] \times \prod_{i=1}^d [x_i, x_i+c]\right) \lesssim c^\gamma,$$

in particular we have $\mu(Q') \lesssim 2^{-k\gamma}$ for each $Q' \in \mathcal{D}_k$. In order to estimate I_1 we define the following relation on \mathcal{D}_k . For two hypercubes $Q, Q' \in \mathcal{D}_k$ we write $Q \sim Q'$ if there exists $(s, x) \in Q$ and $(t, y) \in Q'$ such that $|\tau| = |t-s| \in (2^{-k}, 2 \cdot 2^{-k}]$ and $\|\delta\| = \|y-x\| \in [0, 2 \cdot 2^{-k}]$. Thus,

$$I_1 \lesssim \sum_{k=1}^{\infty} 2^{k\beta} \sum_{\substack{Q, Q' \in \mathcal{D}_k \\ Q \sim Q'}} \mu \otimes \mu(Q \times Q') = \sum_{k=1}^{\infty} 2^{k\beta} \sum_{Q, Q' \in \mathcal{D}_k, Q \sim Q'} \mu(Q) \cdot \mu(Q').$$

The number of hypercubes related to a fixed $Q \in \mathcal{D}_k$ via \sim is bounded by a universal constant not depending on k and Q , hence

$$I_1 \lesssim \sum_{k=1}^{\infty} 2^{k\beta} \sum_{Q \in \mathcal{D}_k} \sum_{Q \sim Q'} \mu(Q) \cdot \mu(Q') \lesssim \sum_{k=1}^{\infty} 2^{k(\beta-\gamma)} \cdot \sum_{Q \in \mathcal{D}_k} \mu(Q).$$

Note that $\sum_{Q \in \mathcal{D}_k} \mu(Q) = \mu(\bigcup_{Q \in \mathcal{D}_k} Q) = \mu(\mathbb{R}_+ \times \mathbb{R}^d) = 1$ and we conclude

$$I_1 \lesssim \sum_{k=1}^{\infty} 2^{k(\beta-\gamma)} = \sum_{k=1}^{\infty} 2^{-k\varepsilon} < \infty,$$

since $\beta = \varphi_1 - 2\varepsilon$ and $\gamma = \varphi_1 - \varepsilon$. For the estimation of I_2 we define another relation on \mathcal{D}_k . For two hypercubes $Q, Q' \in \mathcal{D}_k$ we write $Q \approx Q'$ if there exists $(s, x) \in Q$ and $(t, y) \in Q'$ such that $|\tau| \in (0, 2 \cdot 2^{-k}]$ and $\|\delta\| \in (2^{-k}, 2 \cdot 2^{-k}]$. Thus,

$$I_2 \lesssim \sum_{k=1}^{\infty} 2^{k\beta} \sum_{Q, Q' \in \mathcal{D}_k, Q \approx Q'} \mu(Q) \cdot \mu(Q').$$

Again, the number of hypercubes related to some fixed Q via \approx is bounded by a universal constant. Hence the same calculation as for I_1 yields

$$I_2 \lesssim \sum_{k=1}^{\infty} 2^{k(\beta-\gamma)} = \sum_{k=1}^{\infty} 2^{-k\varepsilon} < \infty.$$

(ii) Now we treat the case $\alpha \in [1, 2)$ and $\varphi_\alpha \leq d$. Let $\beta = \varphi_\alpha - 2\alpha \cdot \varepsilon < d$ for some arbitrary $\varepsilon > 0$. Due to Lemma 5.6 we have

$$\begin{aligned} \mathcal{E}_{K^\beta}(\mu) &\lesssim \underbrace{\int \int_{\{|\tau| \in (0, 1], \|\delta\| \in [0, |\tau|^{1/\alpha}]\}} |\tau|^{-\beta/\alpha} d\mu d\mu}_{=: I_3} \\ &\quad + \underbrace{\int \int_{\{|\tau| \in (0, 1], \|\delta\| \in (|\tau|^{1/\alpha}, 1]\}} \|\delta\|^{-\beta} d\mu d\mu}_{=: I_4}. \end{aligned}$$

We get

$$I_3 \lesssim \sum_{k=1}^{\infty} 2^{k\beta/\alpha} \cdot \mu \otimes \mu\{|\tau| \in (2^{-k}, 2 \cdot 2^{-k}], \|\delta\| \in [0, 2^{1/\alpha} \cdot 2^{-k/\alpha}]\}.$$

Further,

$$I_4 \lesssim \sum_{k=1}^{\infty} 2^{k\beta/\alpha} \cdot \mu \otimes \mu\{|\tau| \in (0, 2 \cdot 2^{-k}], \|\delta\| \in (2^{-k/\alpha}, 2^{1/\alpha} \cdot 2^{-k/\alpha}]\}.$$

Now we have to calculate the expressions $\mu \otimes \mu\{\cdot\}$ for I_3 and I_4 . For each $k \in \mathbb{N}$, we tile $\mathbb{R}_+ \times \mathbb{R}^d$ by disjoint α -parabolic cylinders of size $2^{-k} \times 2^{-k/\alpha} \times \dots \times 2^{-k/\alpha}$

and again denote the collection of such cylinders by \mathcal{D}_k . For every $c \in (0, 1]$, $\gamma = \varphi_\alpha - \alpha \cdot \varepsilon$, and $\alpha \in [1, 2)$, Frostman's lemma (Theorem 5.4) yields

$$\mu\left([t, t+c] \times \prod_{i=1}^d [x_i, x_i + c^{1/\alpha}]\right) \lesssim c^{\gamma/\alpha};$$

in particular, we have $\mu(Q') \lesssim 2^{-k\gamma/\alpha}$ for each $Q' \in \mathcal{D}_k$. The same technique as in (i) results in

$$I_3, I_4 \lesssim \sum_{k=1}^{\infty} 2^{k(\beta-\gamma)/\alpha} \leq \sum_{k=1}^{\infty} 2^{-k\varepsilon} < \infty,$$

since $\beta = \varphi_\alpha - 2\alpha \cdot \varepsilon$ and $\gamma = \varphi_\alpha - \alpha \cdot \varepsilon$.

(iii) Finally, we treat the case $\alpha \in [1, 2)$ and $\varphi_\alpha > d$. Let $\beta = (1 - \frac{1}{\alpha}) \cdot d + \frac{1}{\alpha} \cdot \varphi_\alpha - 2\varepsilon > d$ for sufficiently small $\varepsilon > 0$. Due to Lemma 5.6, we have

$$\begin{aligned} \mathcal{E}_{K^\beta}(\mu) &\lesssim \int \int_{\{|\tau| \in (0, 1], \|\delta\| \in [0, |\tau|^{1/\alpha}]\}} |\tau|^{(1-1/\alpha)d-\beta} d\mu d\mu \\ &\quad + \int \int_{\{|\tau| \in (0, 1], \|\delta\| \in (|\tau|^{1/\alpha}, 1]\}} \|\delta\|^{(\alpha-1)d-\alpha\beta} d\mu d\mu \end{aligned}$$

and the same techniques as in (ii) yield the finiteness of this expression when choosing $\gamma = \varphi_\alpha - \varepsilon$ in Frostman's lemma (Theorem 5.4). ■

6. Range: Upper and lower bounds

We give upper and lower bounds for the Hausdorff dimension of the range of a stable Lévy process with drift.

Theorem 6.1. *Let $T \subseteq \mathbb{R}_+$ be any set and $\alpha \in (0, 2]$. Let $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d and $f : T \rightarrow \mathbb{R}^d$ be any function. Define $\varphi_\alpha := \mathcal{P}^\alpha$ - $\dim \mathcal{G}_T(f)$. Then, one \mathbb{P} -almost surely has*

$$\dim \mathcal{R}_T(X + f) \leq \begin{cases} (\alpha \cdot \varphi_\alpha) \wedge d & \alpha \in (0, 1], \\ \varphi_\alpha \wedge d & \alpha \in [1, 2]. \end{cases} \quad (6.1)$$

Proof. The Gaussian case follows from the proof of [13, Theorem 1.2] and Proposition 2.2. Since the Hausdorff dimension of the range never exceeds the topological dimension of the space a function maps to, we always have $\dim \mathcal{R}_T(X + f) \leq d$. In the case of $\alpha \in [1, 2)$ the claim directly follows from Theorems 2.4 and 4.1, which yield

$$\dim \mathcal{R}_T(X + f) \leq \dim \mathcal{G}_T(X + f) \leq \mathcal{P}^\alpha\text{-dim} \mathcal{G}_T(X + f) \leq \mathcal{P}^\alpha\text{-dim} \mathcal{G}_T(f) = \varphi_\alpha.$$

Now, let $\alpha \in (0, 1]$ and $\beta = \alpha \cdot \varphi_\alpha$ and let $\delta, \varepsilon > 0$ be arbitrary. Then, $\mathcal{G}_T(f)$ can be covered by α -parabolic cylinders

$$(\mathbb{P}_k)_{k \in \mathbb{N}} = \left([t_k, t_k + c_k] \times \prod_{i=1}^d [x_{i,k}, x_{i,k} + c_k^{1/\alpha}] \right)_{k \in \mathbb{N}} \subseteq \mathcal{P}^\alpha$$

such that $\sum_{k=1}^{\infty} |\mathbb{P}_k|^{(\beta+\delta)/\alpha} \lesssim \sum_{k=1}^{\infty} c_k^{(\beta+\delta)/\alpha} \leq \varepsilon$. Let $M_k(\omega)$ be the random number of a fixed 2^d -nested collection of hypercubes with sidelength $c_k^{1/\alpha}$ that the path $t \mapsto X_t(\omega)$ hits at some time $t \in [t_k, t_k + c_k]$. As in the proof of Theorem 4.1, we obtain a random parabolic cover $\bigcup_{k \in \mathbb{N}} \tilde{\mathbb{P}}_k(\omega) \supseteq \mathcal{G}_T(X(\omega) + f)$ where

$$\begin{aligned} \tilde{\mathbb{P}}_k(\omega) &= [t_k, t_k + c_k] \times \bigcup_{j=1}^{M_k(\omega)} \prod_{i=1}^d ([\xi_{i,j,k}(\omega) + x_{i,k}, \xi_{i,j,k}(\omega) + x_{i,k} + c_k^{1/\alpha}] \\ &\quad \cup [\xi_{i,j,k}(\omega) + x_{i,k} + c_k^{1/\alpha}, \xi_{i,j,k}(\omega) + x_{i,k} + 2c_k^{1/\alpha}]). \end{aligned}$$

By projection, we get the random cover $\bigcup_{k \in \mathbb{N}} \square_k \supseteq \mathcal{R}_T(X(\omega) + f)$ with

$$\begin{aligned} \square_k(\omega) &= \bigcup_{k=1}^{M_k(\omega)} \prod_{i=1}^d ([\xi_{i,j,k}(\omega) + x_{i,k}, \xi_{i,j,k}(\omega) + x_{i,k} + c_k^{1/\alpha}] \\ &\quad \cup [\xi_{i,j,k}(\omega) + x_{i,k} + c_k^{1/\alpha}, \xi_{i,j,k}(\omega) + x_{i,k} + 2c_k^{1/\alpha}]). \end{aligned}$$

This is a union of $M_k(\omega) \cdot 2^d$ hypercubes with diameter $|\square_k(\omega)| \asymp c_k^{1/\alpha}$. As in the proof of Theorem 4.1 we get $\mathbb{E}[M_k] \lesssim c_k^{-\delta'/\alpha}$. Hence, we get for $\varepsilon' = \delta + \delta' > 0$,

$$\begin{aligned} \mathbb{E}[\mathcal{H}^{\beta+\varepsilon'}(\mathcal{R}_T(X + f))] &\leq \mathbb{E}\left[\sum_{k=1}^{\infty} |\square_k|^{(\beta+\varepsilon')}\right] \lesssim \mathbb{E}\left[\sum_{k=1}^{\infty} M_k \cdot 2^d \cdot c_k^{(\beta+\varepsilon')/\alpha}\right] \\ &\lesssim \sum_{k=1}^{\infty} \mathbb{E}[M_k] \cdot c_k^{(\beta+\varepsilon')/\alpha} \lesssim \sum_{k=1}^{\infty} c_k^{(\beta+\varepsilon'-\delta')/\alpha} = \sum_{k=1}^{\infty} c_k^{(\beta+\delta)/\alpha} \leq \varepsilon. \end{aligned}$$

Since $\varepsilon, \varepsilon' > 0$ are arbitrary, for all $\beta' > \beta$ we get $\mathbb{E}[\mathcal{H}^{\beta'}(\mathcal{R}_T(X + f))] = 0$, and hence,

$$\mathcal{H}^{\beta'}(\mathcal{R}_T(X + f)) = 0 \quad \mathbb{P}\text{-almost surely.}$$

Since $\beta' > \beta$ is also arbitrary, we finally get $\dim \mathcal{R}_T(X + f) \leq \beta = \alpha \cdot \varphi_\alpha$ \mathbb{P} -almost surely, as claimed. \blacksquare

The lower bound is obtained by the energy method. The stable process X is transformed into the kernel of the energy integral.

Lemma 6.2. *Let $T \subseteq \mathbb{R}_+$ be a Borel set and $\alpha \in (0, 2)$. Let $X = (X_t)_{t \geq 0}$ be an isotropic stable Lévy process in \mathbb{R}^d and $f : \mathbb{R}_+ \rightarrow \mathbb{R}^d$ be a Borel measurable function. Define the difference kernel*

$$\kappa^\beta(t, x) := \mathbb{E}[\|\text{sign}(t) \cdot X_{|t|} + x\|^{-\beta}].$$

Then, $\text{Cap}_{\kappa^\beta}(\mathcal{G}_T(f)) > 0$ implies that \mathbb{P} -almost surely $\text{Cap}_\beta(\mathcal{R}_T(X(\omega) + f)) > 0$ holds. Hence, $\mathcal{E}_{\kappa^\beta}(\mu) < \infty$ for some probability measure $\mu \in \mathcal{M}^1(\mathcal{G}_T(f))$ implies that \mathbb{P} -almost surely $\dim \mathcal{R}_T(X + f) \geq \beta$ holds.

Proof. Let $\mu \in \mathcal{M}^1(\mathcal{G}_T(f))$ and π_t denote the projection onto the time component, that is, $\pi_t(t, f(t)) = t$. Define the probability measure $\nu \in \mathcal{M}^1(\mathbb{R}_+)$ as the pushforward measure $\nu(A) = \mu(\pi_T^{-1}(A))$ for Borel sets $A \subseteq \mathbb{R}_+$ and, further, the random probability measure $\tilde{\mu}_\omega(R) = \nu((X(\omega) + f)^{-1}(R))$ for every Borel set $R \subseteq \mathbb{R}^d$. Then, Tonelli's theorem and the stationarity of the increments of X yield

$$\begin{aligned} \mathbb{E}[\mathcal{E}_\beta(\tilde{\mu}_\omega)] &= \mathbb{E}\left[\int_{\mathcal{R}_T(X(\omega)+f)} \int_{\mathcal{R}_T(X(\omega)+f)} \|x - y\|^{-\beta} d\tilde{\mu}_\omega(x) d\tilde{\mu}_\omega(y)\right] \\ &= \mathbb{E}\left[\int_T \int_T \|X_t(\omega) + f(t) - (X_s(\omega) + f(s))\|^{-\beta} d\nu(t) d\nu(s)\right] \\ &= \int_{\mathcal{G}_T(f)} \int_{\mathcal{G}_T(f)} \mathbb{E}[\|X_t(\omega) - X_s(\omega) + x - y\|^{-\beta}] d\mu(t, x) d\mu(s, y) \\ &= \int_{\mathcal{G}_T(f)} \int_{\mathcal{G}_T(f)} \mathbb{E}[\|\text{sign}(t-s) \cdot X_{|t-s|}(\omega) + x - y\|^{-\beta}] d\mu(t, x) d\mu(s, y) \\ &= \mathcal{E}_{\kappa^\beta}(\mu). \end{aligned}$$

By assumption, there exists $\mu \in \mathcal{M}^1(\mathcal{G}_T(f))$ such that $\mathcal{E}_{\kappa^\beta}(\mu) < \infty$ holds. From that one \mathbb{P} -almost surely has $\mathcal{E}_\beta(\tilde{\mu}_\omega) < \infty$ and the final statement immediately follows by Frostman's theorem (Theorem 5.2), since the range of a Borel set under a Borel measurable function is a Suslin set; see [8, Section 11]. \blacksquare

Next we provide estimates for the difference kernel κ^β from Lemma 6.2 that will give appropriate estimates of the energy integral.

Lemma 6.3. *Let $\alpha \in (0, 2)$ and $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d . Let $\beta \in (0, d)$ and $\tau \in \mathbb{R}$, $\delta \in \mathbb{R}^d$ be such that $|\tau| \in (0, 1]$, $\|\delta\| \in [0, 1]$. Then, for the difference kernel $\kappa^\beta(\tau, \delta) = \mathbb{E}[\|\text{sign}(t) \cdot X_{|t|} + x\|^{-\beta}]$ from Lemma 6.2 one has*

$$\kappa^\beta(\tau, \delta) \lesssim \begin{cases} |\tau|^{-\beta/\alpha}, \\ \|\delta\|^{-\beta} \end{cases} \quad \text{for } |\tau| \leq \|\delta\|^\alpha.$$

Proof. Let $p(x)$ denote the density function of $X_1 \stackrel{d}{=} |\tau|^{-1/\alpha} X_{|\tau|}$. We define the rescaled increment $\tilde{\delta} := \delta/|\tau|^{1/\alpha}$. The self-similarity of the stable Lévy process and Lemma 5.5 yield

$$\begin{aligned} \mathbb{E}[\|\text{sign}(\tau) \cdot X_{|\tau|} + \delta\|^{-\beta}] &= |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} \cdot p(x) \, dx \\ &= |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|x + \text{sign}(\tau) \cdot \tilde{\delta}\|^{-\beta} \cdot p(x) \, dx \\ &\lesssim |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|x\|^{-\beta} \cdot p(x) \, dx \lesssim |\tau|^{-\beta/\alpha} \cdot \mathbb{E}[\|X_1\|^{-\beta}] \lesssim |\tau|^{-\beta/\alpha}, \end{aligned}$$

since negative moments of order $\beta < d$ exist; see [1, Lemma 3.1]. Now consider the region $\|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \leq \|\tilde{\delta}\|/2$, which yields

$$\begin{aligned} \|x\| &= \|\text{sign}(\tau) \cdot x + \tilde{\delta} - \tilde{\delta}\| \geq \|\|\text{sign}(\tau) \cdot x + \tilde{\delta}\| - \|\tilde{\delta}\|\| \\ &= \|\tilde{\delta}\| - \|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \geq \frac{1}{2} \cdot \|\tilde{\delta}\|. \end{aligned}$$

Thus, $\beta < d$ and $\tau \leq \|\delta\|^\alpha$ lead to

$$\begin{aligned} \mathbb{E}[\|\text{sign}(\tau) \cdot X_{|\tau|} + \delta\|^{-\beta}] &= |\tau|^{-\beta/\alpha} \int_{\mathbb{R}^d} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} \cdot p(x) \, dx \\ &\lesssim \|\delta\|^{-\beta} + |\tau|^{-\beta/\alpha} \int_{\{\|\text{sign}(\tau) \cdot x + \tilde{\delta}\| \leq \|\tilde{\delta}\|/2\}} \|\text{sign}(\tau) \cdot x + \tilde{\delta}\|^{-\beta} \cdot p(x) \, dx \\ &\lesssim \|\delta\|^{-\beta} + |\tau|^{-\beta/\alpha} \cdot \|\tilde{\delta}\|^{-d-\alpha} \int_0^{\|\tilde{\delta}\|} r^{d-\beta-1} \, dr = \|\delta\|^{-\beta} + |\tau|^{-\beta/\alpha} \cdot \|\tilde{\delta}\|^{-\alpha-\beta} \\ &= \|\delta\|^{-\beta} + |\tau| \cdot \|\delta\|^{-\alpha-\beta} \lesssim \|\delta\|^{-\beta}, \end{aligned}$$

where we have used (1.3) to estimate the tail densities. \blacksquare

We get a lower bound for the Hausdorff dimension of the range of a stable Lévy process with drift.

Theorem 6.4. *Let $T \subseteq \mathbb{R}_+$ be a Borel set and $\alpha \in (0, 2)$. Let $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d and $f : T \rightarrow \{y \in \mathbb{R}^d : \|y - x\| \leq \frac{1}{2}\}$ for fixed $x \in \mathbb{R}^d$ be a Borel measurable function. Let $\varphi_\alpha = \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f)$. Then, one \mathbb{P} -almost surely has*

$$\dim \mathcal{R}_T(X + f) \geq \begin{cases} (\alpha \cdot \varphi_\alpha) \wedge d & \alpha \in (0, 1], \\ \varphi_\alpha \wedge d & \alpha \in [1, 2). \end{cases} \quad (6.2)$$

Proof. We consider the difference kernel $\kappa^\beta(t, x) = \mathbb{E}[\|\text{sign}(t) \cdot X_{|t|}(\omega) + x\|^{-\beta}]$ from Lemma 6.2. Analogously to the proof of Theorem 5.7, we can show that $\mathcal{E}_{\kappa^\beta}(\mu)$ is finite for $\mu \in \mathcal{M}^1(\mathcal{G}_T(f))$ from the parabolic version of Frostman's lemma in

Theorem 5.4 and for every β less than the right-hand side of (6.2). Note that the different lower bounds for the dimension of the range in cases $\alpha \in (0, 1]$ and $\alpha \in [1, 2)$ occur due to the different upper bounds in the parabolic version of Frostman's lemma. \blacksquare

7. Estimates for the parabolic Hausdorff dimension

We first give an estimate for the α -parabolic Hausdorff dimension of a constant function.

Lemma 7.1. *Let $T \subseteq \mathbb{R}$ be any set and $\alpha \in (0, \infty)$. Define the constant function $f_C(x) = C \in \mathbb{R}^d$ for all $x \in T$. Then, one has*

$$\mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f_C) \leq (\alpha \vee 1) \cdot \dim T.$$

Proof. Without loss of generality, let $f_C = f_0 \equiv 0 \in \mathbb{R}^d$.

(i) Let $\alpha \in (0, 1]$, $\beta = \dim T$ and let $\delta, \varepsilon > 0$ be arbitrary. Then, there exists a cover $\bigcup_{k \in \mathbb{N}} T_k \supseteq T$ with $T_k = [t_k, t_k + c_k]$ and $c_k \leq 1$ such that $\sum_{k=1}^{\infty} |T_k|^{\beta+\delta} = \sum_{k=1}^{\infty} c_k^{\beta+\delta} \leq \varepsilon$. Now, $\mathcal{G}_T(f_0)$ can be covered by α -parabolic cylinders

$$(\mathbb{P}_k)_{k \in \mathbb{N}} = \left([t_k, t_k + c_k] \times \prod_{j=1}^d [0, c_k^{1/\alpha}] \right)_{k \in \mathbb{N}} \subseteq \mathcal{P}^\alpha$$

with $|\mathbb{P}_k| \asymp c_k$. Hence,

$$\mathcal{P}^\alpha - \mathcal{H}^{\beta+\delta}(\mathcal{G}_T(f_0)) \leq \sum_{k=1}^{\infty} |\mathbb{P}_k|^{\beta+\delta} \lesssim \sum_{k=1}^{\infty} c_k^{\beta+\delta} \leq \varepsilon.$$

Since $\delta > 0$ is arbitrary, for all $\beta' > \beta$ we have $\mathcal{P}^\alpha - \mathcal{H}^{\beta'}(\mathcal{G}_T(f_0)) < \infty$ and, therefore, one has $\mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f_0) \leq \beta'$. Since $\beta' > \beta$ is also arbitrary, we obtain

$$\mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f_0) \leq \beta = \dim T.$$

(ii) Let $\alpha \in [1, \infty)$ and $\beta = \alpha \cdot \dim T$ and let $\delta, \varepsilon > 0$ be arbitrary. With the cover $\bigcup_{k \in \mathbb{N}} T_k \supseteq T$ from part (i), we get $\sum_{k=1}^{\infty} |T_k|^{(\beta+\delta)/\alpha} = \sum_{k=1}^{\infty} c_k^{(\beta+\delta)/\alpha} \leq \varepsilon$. Then, the cover $\bigcup_{k \in \mathbb{N}} \mathbb{P}_k \supseteq \mathcal{G}_T(f_0)$ from part (i) now fulfills $|\mathbb{P}_k| \asymp c_k^{1/\alpha}$ and it follows that

$$\mathcal{P}^\alpha - \mathcal{H}^{\beta+\delta}(\mathcal{G}_T(f_0)) \leq \sum_{k=1}^{\infty} |\mathbb{P}_k|^{\beta+\delta} \lesssim \sum_{k=1}^{\infty} c_k^{(\beta+\delta)/\alpha} \leq \varepsilon.$$

Since $\delta > 0$ and $\beta' > \beta$ are arbitrary, as in part (i) we get $\mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f_0) \leq \beta = \alpha \cdot \dim T$. \blacksquare

We can calculate the α -parabolic Hausdorff dimension of the graph of an isotropic α -stable Lévy process itself.

Theorem 7.2. *Let $T \subseteq \mathbb{R}_+$ be a Borel set and $\alpha \in (0, 2]$. Let $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process in \mathbb{R}^d . Then, one \mathbb{P} -almost surely has*

$$\mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(X) = (\alpha \vee 1) \cdot \dim T.$$

Proof. By [17, Theorem 3.2], Theorems 2.4 and 4.1, and Lemma 7.1 in the case $\alpha \cdot \dim T \geq 1$, that is, $\alpha \in [1, 2]$ and $f_0 \equiv 0 \in \mathbb{R}^d$, one \mathbb{P} -almost surely has

$$\begin{aligned} \dim T + 1 - 1/\alpha &= \dim \mathcal{G}_T(X) \leq 1/\alpha \cdot \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(X) + 1 - 1/\alpha \\ &\leq 1/\alpha \cdot \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f_0) + 1 - 1/\alpha \leq \dim T + 1 - 1/\alpha. \end{aligned}$$

In the other cases, [17, Theorem 3.1] together with the same theorems as above \mathbb{P} -almost surely yield

$$\begin{aligned} (\alpha \vee 1) \cdot \dim T &= \dim \mathcal{G}_T(X) \leq \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(X) \leq \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f_0) \\ &\leq (\alpha \vee 1) \cdot \dim T \end{aligned}$$

and the claim follows. ■

Remark 7.3. We can also deduce the Hausdorff dimension of the graph of the fractional Brownian motion $B^H = (B_t^H)_{t \geq 0}$ in \mathbb{R}^d of Hurst index $1/\alpha = H \in (0, 1]$. One \mathbb{P} -almost surely has

$$\mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(B^H) = \frac{\dim T}{H} = \alpha \cdot \dim T.$$

This follows from [20, Theorem 2.1], Proposition 2.2, Theorem 2.4, [13, Lemma 2.2], and Lemma 7.1 for $\alpha \cdot \dim T \leq d$ and $f_0 \equiv 0 \in \mathbb{R}^d$, which \mathbb{P} -almost surely yield

$$\alpha \cdot \dim T = \dim \mathcal{G}_T(B^H) \leq \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(B^H) = \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f_0) \leq \alpha \cdot \dim T.$$

In the other cases the same theorems \mathbb{P} -almost surely yield

$$\begin{aligned} \dim T + (1 - 1/\alpha) \cdot d &= \dim \mathcal{G}_T(B^H) \leq \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(B^H)/\alpha + (1 - 1/\alpha) \cdot d \\ &= \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f_0)/\alpha + (1 - 1/\alpha) \cdot d \leq \dim T + (1 - 1/\alpha) \cdot d \end{aligned}$$

and the claim follows.

The calculations in the proof of the previous theorem further show the next result.

Corollary 7.4. *Let $T \subseteq \mathbb{R}_+$ be a Borel set and $\alpha \in (0, \infty)$. Define the constant function $f_C(x) = C \in \mathbb{R}^d$ for all $x \in T$. Then, one has*

$$\mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f_C) = (\alpha \vee 1) \cdot \dim T.$$

As a consequence, we recover a well-known result for the range of an isotropic α -stable Lévy process; see [3] and [10, Theorem 3.1].

Corollary 7.5. *Let $T \subseteq \mathbb{R}_+$ be a Borel set and $\alpha \in (0, 2]$. Let $X = (X_t)_{t \geq 0}$ be an isotropic α -stable Lévy process. One \mathbb{P} -almost surely has*

$$\dim \mathcal{R}_T(X) = (\alpha \cdot \dim T) \wedge d.$$

Proof. From Theorem 3.2 and Corollary 7.4 it follows that

$$\dim \mathcal{R}_T(X) = ((\alpha \wedge 1) \cdot \mathcal{P}^\alpha\text{-dim} \mathcal{G}_T(f_0)) \wedge d = (\alpha \cdot \dim T) \wedge d,$$

as claimed. ■

We can also give some a priori estimates for the α -parabolic Hausdorff dimension of the graph of a function in terms of the genuine Hausdorff dimension.

Theorem 7.6. *Let $T \subseteq \mathbb{R}$ be any set and $f : T \rightarrow \mathbb{R}^d$ be any function. Let $\varphi_\alpha = \mathcal{P}^\alpha\text{-dim} \mathcal{G}_T(f)$. Then, one has*

$$\varphi_\alpha \leq \begin{cases} (\varphi_1 + (\frac{1}{\alpha} - 1) \cdot d) \wedge (d + 1) & \alpha \in (0, 1], \\ (\varphi_1 + \alpha - 1) \wedge (d + 1) & \alpha \in [1, \infty) \end{cases} \quad (7.1)$$

and

$$\varphi_\alpha \geq \begin{cases} \varphi_1 \vee (\frac{1}{\alpha} \cdot \varphi_1 + 1 - \frac{1}{\alpha}) & \alpha \in (0, 1], \\ \varphi_1 \vee (\alpha \cdot \varphi_1 + (1 - \alpha) \cdot d) & \alpha \in [1, \infty). \end{cases} \quad (7.2)$$

Further, if $T \subseteq \mathbb{R}_+$ is a Borel set and $f : T \rightarrow \mathbb{R}^d$ is a Borel measurable function, then we obtain

$$\varphi_\alpha \leq \left(\frac{1}{\alpha} \cdot \varphi_1\right) \wedge \left(\varphi_1 + \left(\frac{1}{\alpha} - 1\right) \cdot d\right) \wedge (d + 1), \quad \alpha \in (0, 1]. \quad (7.3)$$

Proof. This follows immediately by Theorem 2.4 for (7.1) and (7.2) and Corollary 3.3 for (7.3) and the fact that the Hausdorff dimension never exceeds the topological dimension. ■

Next we calculate some bounds for the parabolic Hausdorff dimension of β -Hölder continuous functions. These are functions $f : T \rightarrow \mathbb{R}^d$ that fulfill $\|f(t) - f(s)\| \leq C \cdot |t - s|^\beta$ for all $s, t \in T$ and some $\beta \in (0, 1]$, $C > 0$, denoted by $f \in C^\beta(T, \mathbb{R}^d)$. In the case of $\alpha = 1$, the following theorem is well known; see [9]:

Theorem 7.7. *Let $T \subseteq \mathbb{R}$ be any set, $\alpha \in (0, \infty)$, $\beta \in (0, 1]$, and $f \in C^\beta(T, \mathbb{R}^d)$. Define $\varphi_\alpha := \mathcal{P}^\alpha\text{-dim}\mathcal{G}_T(f)$. Then, one has*

$$\varphi_\alpha \leq \begin{cases} (\dim T + d \cdot (\frac{1}{\alpha} - \beta)) \wedge \frac{\dim T}{\alpha\beta} \wedge (d + 1) & \alpha \in (0, 1], \\ (\alpha \cdot \dim T + d \cdot (1 - \alpha\beta)) \wedge \frac{\dim T}{\beta} \wedge (d + 1) & \alpha \in [1, \frac{1}{\beta}], \\ (\alpha \cdot \dim T) \wedge (d + 1) & \alpha \in [\frac{1}{\beta}, \infty). \end{cases}$$

Proof. Let $\tau > \dim T$ and $\varepsilon > 0$ be arbitrary. Then, we can cover T by intervals $(\mathbb{T}_k)_{k \in \mathbb{N}}$ with sidelength $|\mathbb{T}_k| < 1$ such that $\sum_{k=1}^{\infty} |\mathbb{T}_k|^\tau < \varepsilon$. Since $f \in C^\beta(T, \mathbb{R}^d)$, we can cover $\mathcal{G}_T(f)$ by $(\mathbb{B}_k)_{k \in \mathbb{N}} \subseteq \mathbb{R}^{1+d}$ where

$$\mathbb{B}_k := \mathbb{T}_k \times \prod_{i=1}^d [x_{i,k}, x_{i,k} + C \cdot |\mathbb{T}_k|^\beta] \quad \text{for every } k \in \mathbb{N}.$$

Note that without loss of generality we may assume $C \geq 1$ for the constant in the definition of Hölder continuity.

(i) Let $\alpha \in (0, 1]$. On the one hand, for every $k \in \mathbb{N}$ we can cover \mathbb{B}_k by (several) α -parabolic cylinders with sidelength $|\mathbb{T}_k|$ in time. Since $K \cdot |\mathbb{T}_k|^{1/\alpha} \geq C \cdot |\mathbb{T}_k|^\beta$ iff $K \geq C \cdot |\mathbb{T}_k|^{\beta-1/\alpha}$ for some hypercubes $\square_{k,l}$ with sidelength $|\mathbb{T}_k|^{1/\alpha}$, we find a cover

$$\mathcal{G}_T(f) \subseteq \bigcup_{k=1}^{\infty} \bigcup_{l=1}^{\lceil C \cdot |\mathbb{T}_k|^{\beta-1/\alpha} \rceil} \mathbb{T}_k \times \square_{k,l}$$

with $\mathbb{T}_k \times \square_{k,l} \in \mathcal{P}^\alpha$ and $|\mathbb{T}_k \times \square_{k,l}| \asymp |\mathbb{T}_k|$ for every $k, l \in \mathbb{N}$. Now, for $\gamma = \tau + d \cdot (1/\alpha - \beta)$, we have

$$\mathcal{P}^\alpha - \mathcal{H}^\gamma(\mathcal{G}_T(f)) \lesssim \sum_{k=1}^{\infty} |\mathbb{T}_k|^{d \cdot (\beta-1/\alpha) + \gamma} = \sum_{k=1}^{\infty} |\mathbb{T}_k|^\tau < \varepsilon.$$

Since $\tau > \dim T$ is arbitrary, $\varphi_\alpha \leq \dim T + d \cdot (1/\alpha - \beta)$. On the other hand, for every $k \in \mathbb{N}$, we can cover \mathbb{B}_k by a single α -parabolic cylinder with sidelength $C^\alpha \cdot |\mathbb{T}_k|^{\alpha\beta}$ in time. Then, $\mathcal{G}_T(f) \subseteq \bigcup_{k \in \mathbb{N}} \mathbb{P}_k$ with $|\mathbb{P}_k| \asymp |\mathbb{T}_k|^{\alpha\beta}$. Now, for $\gamma = \tau/(\alpha\beta)$, we have $\mathcal{P}^\alpha - \mathcal{H}^\gamma(\mathcal{G}_T(f)) \lesssim \sum_{k=1}^{\infty} |\mathbb{T}_k|^{\alpha\beta \cdot \gamma} < \varepsilon$. Since $\tau > \dim T$ is arbitrary, this results in $\varphi_\alpha \leq \frac{\dim T}{\alpha\beta}$.

(ii) Let $\alpha \in [1, 1/\beta]$. On the one hand, for every $k \in \mathbb{N}$ we can cover \mathbb{B}_k by (several) α -parabolic cylinders with sidelength $|\mathbb{T}_k|$ in time. The covering sets from part (i) now fulfill $|\mathbb{T}_k \times \square_{k,l}| \asymp |\mathbb{T}_k|^{1/\alpha}$ and for $\gamma = \alpha \cdot \tau + d \cdot (1 - \alpha\beta)$, we have

$$\mathcal{P}^\alpha - \mathcal{H}^\gamma(\mathcal{G}_T(f)) \lesssim \sum_{k=1}^{\infty} |\mathbb{T}_k|^{d \cdot (\beta-1/\alpha) + \gamma/\alpha} = \sum_{k=1}^{\infty} |\mathbb{T}_k|^\tau < \varepsilon.$$

Since $\tau > \dim T$ is arbitrary, this results in $\varphi_\alpha \leq \alpha \cdot \dim T + d \cdot (1 - \alpha\beta)$. On the other hand, as in part (i), for every $k \in \mathbb{N}$ we can cover B_k by a single α -parabolic cylinder with sidelength $C^\alpha \cdot |\tau_k|^{\alpha\beta}$ in time. Then, the cover $\mathcal{G}_T(f) \subseteq \bigcup_{k \in \mathbb{N}} P_k$ now fulfills $|P_k| \asymp |\tau_k|^\beta$ and for $\gamma = \tau/\beta$, we have $\mathcal{P}^\alpha - \mathcal{H}^\gamma(\mathcal{G}_T(f)) \lesssim \sum_{k=1}^\infty |\tau_k|^{\beta \cdot \gamma} < \varepsilon$. Since $\tau > \dim T$ is arbitrary, this results in $\varphi_\alpha \leq \dim T/\beta$.

(iii) Let $\alpha \in [1/\beta, \infty)$. For every $k \in \mathbb{N}$, we can cover B_k by a single α -parabolic cylinder P_k with length $C^\alpha \cdot |\tau_k|$ in time. Then, $\mathcal{G}_T(f) \subseteq \bigcup_{k=1}^\infty P_k$ with $|P_k| \asymp |\tau_k|^{1/\alpha}$ and for $\gamma \geq \alpha \cdot \tau$, we have $\mathcal{P}^\alpha - \mathcal{H}^\gamma(\mathcal{G}_T(f)) \lesssim \sum_{k=1}^\infty |\tau_k|^{\gamma/\alpha} < \varepsilon$. Since $\tau > \dim T$ is arbitrary, this results in $\varphi_\alpha \leq \alpha \cdot \dim T$. ■

Let us inspect the important case $\alpha = 2$, that is, we aim to get a bound for the Hausdorff dimension of the graph of Brownian motion plus β -Hölder continuous drift function over T .

Corollary 7.8. *Let $T \subseteq \mathbb{R}_+$ be any set. Let $B = (B_t)_{t \geq 0}$ denote the d -dimensional Brownian motion and let $f \in C^\beta(T, \mathbb{R}^d)$ for some $\beta \in (0, 1]$. Then, one \mathbb{P} -almost surely has*

$$\dim \mathcal{G}_T(B + f) \leq \begin{cases} d + \frac{1}{2} & \beta \leq \frac{\dim T}{d} \wedge \frac{1}{2} \wedge (\dim T - \frac{1}{2}), \\ \dim T + d \cdot (1 - \beta) & \dim T - \frac{1}{2} \leq \beta \leq (\frac{\dim T}{d} \wedge \frac{1}{2}), \\ \frac{\dim T}{\beta} & \frac{\dim T}{d} \leq \beta \leq \frac{1}{2}, \\ (2 \cdot \dim T) \wedge (\dim T + \frac{d}{2}) & \beta \geq \frac{1}{2}. \end{cases}$$

Moreover, one \mathbb{P} -almost surely has

$$\dim \mathcal{R}_T(B + f) \leq \begin{cases} \frac{\dim T}{\beta} & \frac{\dim T}{d} \leq \beta \leq \frac{1}{2}, \\ (2 \cdot \dim T) \wedge d & \beta \geq \frac{1}{2}, \\ d & \text{else.} \end{cases}$$

Proof. Let $\varphi_2 = \mathcal{P}^2\text{-dim} \mathcal{G}_T(f)$. Corollary 4.2 \mathbb{P} -almost surely yields

$$\dim \mathcal{G}_T(B + f) \leq \varphi_2 \wedge \frac{\varphi_2 + d}{2}$$

and Theorem 6.1 \mathbb{P} -almost surely yields

$$\dim \mathcal{R}_T(B + f) \leq \varphi_2 \wedge d.$$

Finally, by Theorem 7.7, we easily get

$$\varphi_2 \leq \begin{cases} (2 \cdot \dim T + d \cdot (1 - 2\beta)) \wedge (d + 1) & \beta \leq \frac{\dim T}{d} \wedge \frac{1}{2}, \\ \frac{\dim T}{\beta} & \frac{\dim T}{d} \leq \beta \leq \frac{1}{2}, \\ 2 \cdot \dim T & \beta \geq \frac{1}{2} \end{cases}$$

and the claim follows. ■

References

- [1] P. Becker-Kern, M. M. Meerschaert, and H.-P. Scheffler, [Hausdorff dimension of operator stable sample paths](#). *Monatsh. Math.* **140** (2003), no. 2, 91–101 Zbl [1038.60034](#) MR [2017662](#)
- [2] C. J. Bishop and Y. Peres, *Fractals in probability and analysis*. Cambridge Stud. Adv. Math. 162, Cambridge University Press, Cambridge, 2017 Zbl [1390.28012](#) MR [3616046](#)
- [3] R. M. Blumenthal and R. K. Gettoor, [A dimension theorem for sample functions of stable processes](#). *Illinois J. Math.* **4** (1960), 370–375 Zbl [0093.14402](#) MR [121881](#)
- [4] R. M. Blumenthal and R. K. Gettoor, [Some theorems on stable processes](#). *Trans. Amer. Math. Soc.* **95** (1960), 263–273 Zbl [0107.12401](#) MR [119247](#)
- [5] R. M. Blumenthal and R. K. Gettoor, [The dimension of the set of zeros and the graph of a symmetric stable process](#). *Illinois J. Math.* **6** (1962), 308–316 Zbl [0286.60031](#) MR [138134](#)
- [6] K. Falconer, *Fractal geometry*. John Wiley & Sons, Chichester, 1990 Zbl [0689.28003](#) MR [1102677](#)
- [7] W. J. Hendricks, [A dimension theorem for sample functions of processes with stable components](#). *Ann. Probability* **1** (1973), 849–853 Zbl [0269.60036](#) MR [426168](#)
- [8] T. Jech, *Set theory*. Springer Monogr. Math., Springer, Berlin, 2003 Zbl [1007.03002](#) MR [1940513](#)
- [9] J.-P. Kahane, *Some random series of functions*. Second edn., Cambridge Stud. Adv. Math. 5, Cambridge University Press, Cambridge, 1985 Zbl [0776.28003](#) MR [833073](#)
- [10] M. M. Meerschaert and Y. Xiao, [Dimension results for sample paths of operator stable Lévy processes](#). *Stochastic Process. Appl.* **115** (2005), no. 1, 55–75 Zbl [1074.60079](#) MR [2105369](#)
- [11] P. Mörters and Y. Peres, *Brownian motion*. Camb. Ser. Stat. Probab. Math. 30, Cambridge University Press, Cambridge, 2010 Zbl [1243.60002](#) MR [2604525](#)
- [12] Y. Peres and P. Sousi, [Brownian motion with variable drift: 0-1 laws, hitting probabilities and Hausdorff dimension](#). *Math. Proc. Cambridge Philos. Soc.* **153** (2012), no. 2, 215–234 Zbl [1260.60168](#) MR [2981924](#)
- [13] Y. Peres and P. Sousi, [Dimension of fractional Brownian motion with variable drift](#). *Probab. Theory Related Fields* **165** (2016), no. 3–4, 771–794 Zbl [1344.60040](#) MR [3520018](#)
- [14] W. E. Pruitt and S. J. Taylor, [Sample path properties of processes with stable components](#). *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **12** (1969), 267–289 Zbl [0181.21103](#) MR [258126](#)
- [15] K. Sato, *Lévy processes and infinitely divisible distributions*. Cambridge Stud. Adv. Math. 68, Cambridge University Press, Cambridge, 1999 Zbl [0973.60001](#) MR [1739520](#)
- [16] S. J. Taylor and N. A. Watson, [A Hausdorff measure classification of polar sets for the heat equation](#). *Math. Proc. Cambridge Philos. Soc.* **97** (1985), no. 2, 325–344 Zbl [0584.31006](#) MR [771826](#)
- [17] L. Wedrich, [Hausdorff dimension of the graph of an operator semistable Lévy process](#). *J. Fractal Geom.* **4** (2017), no. 1, 21–41 Zbl [1362.60046](#) MR [3631375](#)

- [18] Y. Xiao, [Random fractals and Markov processes](#). In *Fractal geometry and applications: a jubilee of Benoît Mandelbrot, Part 2*, pp. 261–338, Proc. Sympos. Pure Math. 72, American Mathematical Society, Providence, RI, 2004 Zbl [1055.37003](#) MR [2112126](#)
- [19] Y. Xiao and H. Lin, [Dimension properties of sample paths of self-similar processes](#). *Acta Math. Sinica (N.S.)* **10** (1994), no. 3, 289–300 Zbl [0813.60042](#) MR [1415699](#)
- [20] Y. M. Xiao, Dimension results for Gaussian vector fields and index- α stable fields. *Ann. Probab.* **23** (1995), no. 1, 273–291 Zbl [0834.60040](#) MR [1330771](#)

Received 21 December 2023; revised 31 May 2024.

Peter Kern

Mathematical Institute, Faculty of Mathematics and Natural Sciences, Heinrich Heine University Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf, Germany; kern@hhu.de

Leonard Pleschberger

Mathematical Institute, Faculty of Mathematics and Natural Sciences, Heinrich Heine University Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf, Germany;
leonard.pleschberger@hhu.de