

Positive solutions for a p -Laplacian equation with sub-critical singular parametric reaction term

Pasquale Candito, Giuseppe Failla, and Roberto Livrea

Abstract. The existence of at least two smooth positive solutions for a parametric quasilinear elliptic problem driven by a p -Laplacian operator involving a mildly singular non-linearity perturbed with a sub-critical term is established. Although, to get our conclusions, we combine variational and truncation techniques, we do not use the usual trick of C^1 versus Sobolev minimizers. An explicit quantitative estimate from below of the best theoretical parameters considered is furnished.

1. Introduction

In this paper, the following p -Laplacian problem involving a singular non-linearity perturbed with a sub-critical term is studied, namely,

$$\begin{cases} -\Delta_p u = \lambda f(x, u) + \mu g(x, u) & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (P_{\lambda, \mu})$$

where $\Omega \subset \mathbb{R}^N$, ($N \geq 3$) is a bounded domain with a smooth boundary, $\partial\Omega \in C^2$, the driven operator is the usual p -Laplacian,

$$\Delta_p u := \operatorname{div}(|\nabla u|^{p-2} \nabla u)$$

with $p \in (1, N)$, λ and μ are two positive parameters. Furthermore, we assume that $f : \Omega \times (0, +\infty) \rightarrow [0, +\infty)$ and $g : \Omega \times [0, +\infty) \rightarrow [0, +\infty)$ are two Carathéodory functions fulfilling the following conditions:

(Q_1) $\lim_{s \rightarrow 0^+} f(x, s) = +\infty$ uniformly w.r.t. $x \in \Omega$;

(Q_2) there exist positive constants c_i for $i \in \{1, \dots, 4\}$ and some $\gamma \in (0, 1)$ such that

(Q_{21}) $f(x, s) \leq c_1 s^{-\gamma} + c_2$ for a.a. $x \in \Omega \forall s > 0$;

(Q_{22}) $g(x, s) \leq c_3 s^{q-1} + c_4$ for a.a. $x \in \Omega \forall s > 0, q \in (1, p^*)$,

with $p^* = \frac{Np}{N-p}$ being the critical Sobolev exponent.

Mathematics Subject Classification 2020: 35J92 (primary); 35A01, 35J20, 35J75 (secondary).

Keywords: variational methods, quasilinear elliptic equations, p -Laplacian problem, singular reaction term, truncation techniques.

A wide literature on parametric problems has been produced since the pioneering and seminal papers due to Ambrosetti, Brézis, Cerami [1] and García Azorero, Peral, Manfredi [16], where among the others results, in this latter the fundamental result of Brézis, Nirenberg [8] concerning H^1 versus C^1 local minimizers is extended to the p -Laplacian with $p \neq 2$, which is a well-known key point to allow merging sub-super-solution methods with variational ones.

Subsequently, these ideas have been developed in singular settings, overcoming, in an original way, many technical difficulties arising from the presence of the singularity on the non-linearity, as, for instance, the lack of regularity of the energy functional associated with the problem investigated; see Giacomoni, Saoudi [18] and Giacomoni, Schindler, and Takáč [19].

For a general look on p -Laplacian singular problems, we refer the interested reader to the recent survey [20], the papers [28–31] ($\gamma < 1$), [11, 15] for strongly singular problems ($\gamma \geq 1$) and references therein.

The study of this type of singular problems with a p -Laplacian operator is also largely encouraged by a wide range of physical and engineering applications, particularly the so-called non-Newtonian fluids, see [13, Remark 2.2] and references therein for more details.

Our main results, Theorems 3.1 and 3.4, establish the existence of at least one or two smooth positive solutions for $(P_{\lambda,\mu})$, respectively.

Adapting here some arguments introduced and developed in [9, 10], the first solution for $(P_{\lambda,\mu})$ is obtained by applying a local minimum theorem (Theorem 2.12) due to Bonanno [3], which beyond the existence of a solution, allows us to get two additional features:

- (B₁) the first solution is a local minimum of the functional J_λ , regardless of the asymptotic behavior at infinity of the perturbation g ;
- (B₂) a quantitative estimate of the parameters λ, μ , for which $(P_{\lambda,\mu})$ admits a positive solution, i.e.,

$$\lambda \left(\frac{c_1}{1-\gamma} + c_2 \right) + \mu(c_3 + c_4) < \begin{cases} +\infty & \text{if } 1 < q < p, \\ \frac{N}{p} \pi^{\frac{p}{2}} |\Omega|^{-\frac{p}{N}} \left(\frac{N-p}{p-1} \right)^{p-1} \left[\frac{\Gamma(\frac{N}{p})\Gamma(1+N-\frac{N}{p})}{\Gamma(1+\frac{N}{2})\Gamma(N)} \right]^p & \text{if } q = p, \\ \frac{N}{p} \pi^{\frac{p}{2}} |\Omega|^{-\frac{p}{N}} \left(\frac{N-p}{p-1} \right)^{p-1} \left(\frac{q-p}{2p} \right)^{\frac{q-p}{q}} \left[\frac{\Gamma(\frac{N}{p})\Gamma(1+N-\frac{N}{p})}{\Gamma(1+\frac{N}{2})\Gamma(N)} \right]^p & \text{if } p < q < p^*, \end{cases}$$

where Γ is the gamma function.

In a few words, we combine variational and truncation techniques, but we do not use the usual trick of C^1 versus Sobolev minimizers, recalled above, as well as we do not need a priori estimates or to prove the existence of a super-solution. More precisely, we can use the point (B₁) together with the request on g to fulfill the classical unilateral Ambrosetti–Rabinowitz condition (in the short (AR)-condition), see [2, 26], i.e., there exists $\eta > p$

such that

$$\operatorname{ess\,inf}_{x \in \Omega} \int_0^s g(x, t) dt > 0, \quad 0 < \eta \int_{s_1}^s g(x, t) dt \leq g(x, s)s \quad (\text{AR})$$

for a.a. $x \in \Omega$ and for all $s \geq s_1 > 0$. Here, we realize that the functional J_λ satisfies two key ingredients of the mountain pass theorem [2], that are, the so-called mountain pass geometry, see [4, Theorem 2.1], and the Palais–Smale condition.

Roughly speaking, to obtain the second solution in the super-linear case, $p < q < p^*$, as in the above-mentioned papers, we apply the powerful mountain pass theorem of Ambrosetti and Rabinowitz [2] to a C^1 energy functional J_λ associated to an auxiliary problem, whose positive smooth solutions give back solutions of $(P_{\lambda, \mu})$.

We also highlight that the estimates achieved in (B_2) are not the best theoretical ones, as, for example, in [18] or [19], but are completely computable since they involve only the constants c_i given in (Q_2) and an upper bound of the Sobolev embedding of $W_0^{1,p}(\Omega)$ in $L^q(\Omega)$ as in (2.1) below.

To summarize, Section 2 shows some preliminary results and describes the variational setting adopted to solve $(P_{\lambda, \mu})$. Section 3 is devoted to the main results.

Finally, we observe that our results guarantee the existence of at least two solutions for the following class of elliptic problems:

$$\begin{cases} -\Delta_4 u = \frac{\lambda}{\sqrt{u}} + \mu u^8 & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

with $\Omega \subseteq \mathbb{R}^N$ ($5 \leq N \leq 7$).

2. Preliminaries

In this paper, we will use the symbol $\|\cdot\|_p$ to denote the norm in the Lebesgue space $L^p(\Omega)$, i.e.,

$$\|u\|_p = \left(\int_\Omega |u(x)|^p dx \right)^{\frac{1}{p}} \quad \forall u \in L^p(\Omega);$$

in addition, the symbol $\|\cdot\|$ refers to the norm in Sobolev space $W_0^{1,p}(\Omega)$, i.e.,

$$\|u\| = \|\nabla u\|_p \quad \forall u \in W_0^{1,p}(\Omega).$$

We recall that by compact embedding $W_0^{1,p}(\Omega) \hookrightarrow L^q(\Omega)$ for $q \in [1, p^*)$ we obtain

$$\|u\|_q \leq C_q \|u\| \quad \forall u \in W_0^{1,p}(\Omega),$$

where according to Talenti [34] and by Hölder's inequality, we have

$$C_q \leq S_q := \pi^{-\frac{1}{2}} N^{-\frac{1}{p}} |\Omega|^{\frac{p^*-q}{p^*q}} \left(\frac{p-1}{N-p} \right)^{1-\frac{1}{p}} \left[\frac{\Gamma(1 + \frac{N}{2})\Gamma(N)}{\Gamma(\frac{N}{p})\Gamma(1 + N - \frac{N}{p})} \right], \quad (2.1)$$

with Γ being the gamma function. Since our problem involves a singular term, we also need the inequality [27, Theorem 21.3] in the following particular theorem.

Theorem 2.1 (Hardy–Sobolev’s inequality). *Let $p \in (1, N)$, $\tau \in [0, 1]$ and $\frac{1}{r} = \frac{1}{p} - \frac{1-\tau}{N}$. Then, for any $u \in W_0^{1,p}(\Omega)$, we have $ud^{-\tau} \in L^r(\Omega)$ and*

$$\|ud^{-\tau}\|_r \leq D_\tau \|u\|,$$

where D_τ is a positive constant and d denotes the distance function, i.e.,

$$d(x) := \text{dist}(x, \partial\Omega) = \min_{y \in \partial\Omega} |x - y| \quad \forall x \in \Omega.$$

As in [12, Definitions 3.14–3.15], we recall the definitions of sub-solution and weak solution for the problem $(P_{\lambda,\mu})$.

Definition 2.2. The function $\underline{u} \in W_0^{1,p}(\Omega)$ is called a (weak) sub-solution of $(P_{\lambda,\mu})$, if

$$\int_{\Omega} |\nabla \underline{u}|^{p-2} \nabla \underline{u} \nabla v \, dx \leq \lambda \int_{\Omega} f(x, \underline{u}) v \, dx + \mu \int_{\Omega} g(x, \underline{u}) v \, dx \quad \forall v \in W_0^{1,p}(\Omega) \cap L_+^p(\Omega); \quad (2.2)$$

likewise, the function $u \in W_0^{1,p}(\Omega)$ is called a weak solution of problem $(P_{\lambda,\mu})$, if

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla v \, dx = \lambda \int_{\Omega} f(x, u) v \, dx + \mu \int_{\Omega} g(x, u) v \, dx \quad \forall v \in W_0^{1,p}(\Omega); \quad (2.3)$$

in both cases, it is understood that the right-hand side is well posed.

Now, adopting some reasoning as in [9, 10], we prove the existence of a sub-solution for $(P_{\lambda,\mu})$.

Lemma 2.3. *Suppose (Q_1) holds, then, for all $\lambda, \mu > 0$, there exist $0 < \alpha < 1$, $\delta > 0$, $l > 0$ and a sub-solution \underline{u} of problem $(P_{\lambda,\mu})$ such that $\underline{u} \in C^{1,\alpha}(\bar{\Omega})$ and $ld(x) \leq \underline{u} \leq \delta$ for a.a. $x \in \Omega$.*

Proof. Notice that, from (Q_1) , there exists $\delta > 0$ small enough such that

$$f(x, s) \geq 1 \quad \forall (x, s) \in \Omega \times (0, \delta). \quad (2.4)$$

For $\lambda > 0$, we study the auxiliary problem

$$\begin{cases} -\Delta_p u = \lambda & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases} \quad (2.5)$$

Since $\lambda \in W^{-1,p'}(\Omega)$ and negative p -Laplacian is strictly monotone for $1 < p < 2$, strongly monotone for $p = 2$ and uniformly monotone for $p > 2$ (see (2.18) below and [25, Example 2.27 (c)]), by Minty–Browder’s theorem [6, Theorem 5.16] and combining [35, Theorem 5], [24, Theorem 1.1] (see also [23]) and [21, Lemma 3.1], there exists

$e_\lambda \in C^{1,\alpha}(\bar{\Omega})$ with $0 < \alpha < 1$ (see also [22, Theorem 4.8]), a unique positive solution of (2.5). Let $M := \max\{\delta, \max_{\bar{\Omega}} e_\lambda\}$ and choose c such that $cM \leq \delta$. We put $\underline{u} = ce_\lambda$, then, by (2.4),

$$-\Delta_p \underline{u} = c^{p-1}(-\Delta_p e_\lambda) = c^{p-1}\lambda \leq \lambda \leq \lambda f(x, \underline{u}) \leq \lambda f(x, \underline{u}) + \mu g(x, \underline{u}).$$

Therefore, \underline{u} is a sub-solution of the problem $(P_{\lambda,\mu})$. Moreover, from [32, Theorem 5.3.1], there exists a positive constant l such that

$$ld(x) \leq \underline{u} \leq c \max_{\bar{\Omega}} e_\lambda \leq \delta \quad (2.6)$$

and this achieves the proof. \blacksquare

To apply variational tools and to avoid blow up phenomena, we truncate the reaction term as follows, set $f^* : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ and $g^* : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$

$$f^*(x, s) = \begin{cases} f(x, s) & \text{if } s \geq \underline{u}; \\ f(x, \underline{u}) & \text{if } |s| < \underline{u}; \\ f(x, -s) & \text{if } s \leq -\underline{u}; \end{cases} \quad (2.7)$$

and

$$g^*(x, s) = \begin{cases} g(x, s) & \text{if } s \geq \underline{u}; \\ g(x, \underline{u}) & \text{if } |s| < \underline{u}; \\ g(x, -s) & \text{if } s \leq -\underline{u}. \end{cases} \quad (2.8)$$

Notice that f^* and g^* are Carathéodory functions. In particular, by (Q_2) and (2.6), we obtain

$$f^*(x, s) \leq c_1 \underline{u}^{-\gamma}(x) + c_2 \leq \tilde{c}_1 d^{-\gamma}(x) + c_2 \quad \text{for a.a. } x \in \Omega, \forall s \in \mathbb{R}, \quad (2.9)$$

where $\tilde{c}_1 = c_1 l^{-\gamma}$. Analogously, we get that by (2.6),

$$g^*(x, s) \leq c_3 |s|^{q-1} + c_5 \quad \text{for a.a. } x \in \Omega, \forall s \in \mathbb{R}, \quad (2.10)$$

where $c_5 = c_3 \delta^{q-1} + c_4$.

Now, with the aim of applying variational methods for $(P_{\lambda,\mu})$, we consider the following problem:

$$\begin{cases} -\Delta_p u = \lambda h_{\lambda,\mu}^*(x, u) & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (\text{P}^*)$$

where $h_{\lambda,\mu} : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is given, for a.a. $x \in \Omega$, by

$$h_{\lambda,\mu}^*(x, u) := \begin{cases} f^*(x, u) + \frac{\mu}{\lambda} g^*(x, u) & \text{if } |u| \geq \underline{u}; \\ f(x, \underline{u}) + \frac{\mu}{\lambda} g(x, \underline{u}) & \text{if } |u| < \underline{u}. \end{cases}$$

Remark 2.4. We explicitly observe that the weak solution u of (\mathbf{P}^*) is well-defined, i.e., $u \in W_0^{1,p}(\Omega)$ is such that

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla v dx = \lambda \int_{\Omega} f^*(x, u) v dx + \mu \int_{\Omega} g^*(x, u) v dx \quad \forall v \in W_0^{1,p}(\Omega).$$

Indeed, if $u, v \in W_0^{1,p}(\Omega)$, by (2.9), (2.10), Hölder, Sobolev, and Hardy–Sobolev inequalities, it follows that

$$\begin{aligned} \left| \int_{\Omega} h_{\lambda,\mu}^*(x, u) v dx \right| &\leq \int_{\Omega} f^*(x, u) |v| dx + \frac{\mu}{\lambda} \int_{\Omega} g^*(x, u) |v| dx \\ &\leq \int_{\Omega} (\tilde{c}_1 d^{-\gamma}(x) + c_2) |v| dx + \frac{\mu}{\lambda} \int_{\Omega} (c_3 |u|^{q-1} + c_5) |v| dx \\ &\leq \tilde{c}_1 D_{\gamma} \|v\| + c_2 \|v\|_1 + \frac{\mu}{\lambda} (c_3 \|u\|_q^{q-1} \|v\|_q + c_5 \|v\|_1) < \infty. \end{aligned}$$

Now, we set

$$H_{\lambda,\mu}(x, s) := \int_0^s h_{\lambda,\mu}^*(x, t) dt \quad \text{for a.a. } x \in \Omega, \forall s \in \mathbb{R}.$$

Since $h_{\lambda,\mu}^*(x, s)$ is an even function and by (2.6) (we recall that one can choose $\underline{u} \leq \delta < 1$), we point out that for a.a. $x \in \Omega$ and for all $s \in \mathbb{R}$,

$$|H_{\lambda,\mu}(x, s)| \leq \int_0^{|s|} h_{\lambda,\mu}^*(x, t) dt = \int_0^{\underline{u}} h_{\lambda,\mu}^*(x, \underline{u}) dt + \int_{\underline{u}}^{|s|} h_{\lambda,\mu}^*(x, t) dt. \quad (2.11)$$

Moreover, for a.a. $x \in \Omega$, one has

$$\int_0^{\underline{u}} f(x, \underline{u}) dt + \frac{\mu}{\lambda} \int_0^{\underline{u}} g(x, \underline{u}) dt \leq \frac{c_1}{1-\gamma} + c_2 + \frac{\mu}{\lambda} (c_3 + c_4)$$

and

$$\int_{\underline{u}}^{|s|} f^*(x, t) dt + \frac{\mu}{\lambda} \int_{\underline{u}}^{|s|} g^*(x, t) dt \leq \frac{c_1}{1-\gamma} |s|^{1-\gamma} + c_2 |s| + \frac{\mu}{\lambda} (c_3 |s|^q + c_4 |s|),$$

which implies that

$$\int_0^{|s|} h^*(x, t) dt \leq \frac{c_1}{1-\gamma} + c_2 + \frac{\mu}{\lambda} (c_3 + c_4) + \frac{c_1}{1-\gamma} |s|^{1-\gamma} + c_2 |s| + \frac{\mu}{\lambda} (c_3 |s|^q + c_4 |s|).$$

At this point, it follows that

$$H_{\lambda,\mu}(x, |s|) \leq \begin{cases} 2 \left[\frac{c_1}{1-\gamma} + c_2 + \frac{\mu}{\lambda} (c_3 + c_4) \right] & \text{if } |s| \leq 1; \\ \frac{c_1}{1-\gamma} + c_2 + \frac{\mu}{\lambda} (c_3 + c_4) + \left[\frac{c_1}{1-\gamma} + c_2 + \frac{\mu}{\lambda} (c_3 + c_4) \right] |s|^q & \text{if } |s| > 1. \end{cases}$$

So, for a.a. $x \in \Omega$ and for all $s \in \mathbb{R}$, we have

$$H(x, |s|) \leq 2 \left[\frac{c_1}{1-\gamma} + c_2 + \frac{\mu}{\lambda} (c_3 + c_4) \right] + \left[\frac{c_1}{1-\gamma} + c_2 + \frac{\mu}{\lambda} (c_3 + c_4) \right] |s|^q. \quad (2.12)$$

To simplify the notation, for our convenience, let us put

$$A_{\lambda,\mu} := \frac{c_1}{1-\gamma} + c_2 + \frac{\mu}{\lambda}(c_3 + c_4) \quad (2.13)$$

from (2.12), we get

$$|H_{\lambda,\mu}(x, s)| \leq H_{\lambda,\mu}(x, |s|) \leq A_{\lambda,\mu}(2 + |s|^q) \quad \text{for a.a. } x \in \Omega, \forall s \in \mathbb{R}. \quad (2.14)$$

Lemma 2.5. *For all $\lambda, \mu > 0$, every weak solution of (\mathbf{P}^*) is a weak solution of $(P_{\lambda,\mu})$.*

Proof. Let $u \in W_0^{1,p}(\Omega)$ be a solution of (\mathbf{P}^*) . Following the idea in [12, Lemma 3.50], it is enough to show that $u \geq \underline{u}$. Indeed, by (2.7)–(2.8), if $u \geq \underline{u}$ then $h_{\lambda,\mu}^*(x, s) = f(x, s) + \frac{\mu}{\lambda}g(x, s)$ for a.a. $x \in \Omega$, for all $s \in \mathbb{R}$. Since u is a weak solution of (\mathbf{P}^*) , then by (2.3), one has

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla v dx = \lambda \int_{\Omega} f^*(x, u) v dx + \mu \int_{\Omega} g^*(x, u) v dx, \quad v \in W_0^{1,p}(\Omega). \quad (2.15)$$

Therefore, \underline{u} is a sub-solution of (\mathbf{P}^*) , then by (2.2), we have

$$\int_{\Omega} |\nabla \underline{u}|^{p-2} \nabla \underline{u} \nabla v dx \leq \lambda \int_{\Omega} f^*(x, \underline{u}) v dx + \mu \int_{\Omega} g^*(x, \underline{u}) v dx, \quad v \in W_0^{1,p}(\Omega) \cap L_+^p(\Omega). \quad (2.16)$$

Subtracting (2.16) and (2.15) and choosing $v = (\underline{u} - u)^+$,

$$\begin{aligned} & \int_{\Omega} (|\nabla \underline{u}|^{p-2} \nabla \underline{u} - |\nabla u|^{p-2} \nabla u) \nabla (\underline{u} - u)^+ dx \\ & \leq \lambda \int_{\Omega} (f^*(x, \underline{u}) - f^*(x, u)) (\underline{u} - u)^+ dx + \mu \int_{\Omega} (g^*(x, \underline{u}) - g^*(x, u)) (\underline{u} - u)^+ dx. \end{aligned}$$

Notice that in $\{\underline{u} > u\}$, from (2.7)–(2.8), we have that $f \equiv f^*$, $g \equiv g^*$, then

$$\int_{\Omega} (|\nabla \underline{u}|^{p-2} \nabla \underline{u} - |\nabla u|^{p-2} \nabla u) \nabla (\underline{u} - u)^+ dx \leq 0. \quad (2.17)$$

On the other hand, by monotonicity of $(-\Delta_p, W_0^{1,p}(\Omega))$ (see, [25, Example 2.27 (c)] and [12, Example 2.110]), it follows that

$$\langle -\Delta_p u + \Delta_p v, u - v \rangle \geq \begin{cases} c_1(p) (\|u\| + \|v\|)^{p-2} \|u - v\|^2 & \text{if } 1 < p < 2, \\ c_2(p) \|u - v\|^p & \text{if } p \geq 2, \end{cases} \quad (2.18)$$

for suitable positive constants $c_1(p), c_2(p)$. Moreover, by (2.17)–(2.18), there exists a positive constant c such that

$$\begin{aligned} 0 & \leq \|(\underline{u} - u)^+\|^p = \int_{\{\underline{u} > u\}} |\nabla (\underline{u} - u)|^p dx \\ & \leq c \int_{\{\underline{u} > u\}} (|\nabla \underline{u}|^{p-2} \nabla \underline{u} - |\nabla u|^{p-2} \nabla u) \nabla (\underline{u} - u) dx \\ & = c \int_{\Omega} (|\nabla \underline{u}|^{p-2} \nabla \underline{u} - |\nabla u|^{p-2} \nabla u) \nabla (\underline{u} - u)^+ dx \leq 0. \end{aligned}$$

Therefore, $\|(\underline{u} - u)^+\|^p = 0$ and $u \geq \underline{u}$. ■

The energy functional associated to (\mathbf{P}^*) is defined by setting

$$J_\lambda(u) = \Phi(u) - \lambda\Psi(u), \quad (2.19)$$

where

$$\Phi(u) = \frac{1}{p}\|u\|^p, \quad \Psi(u) = \int_{\Omega} H_{\lambda,\mu}(x, u)dx \quad \forall u \in W_0^{1,p}(\Omega). \quad (2.20)$$

Lemma 2.6. *Suppose (Q_1) and (Q_2) hold. Then, Ψ is well-posed, continuously Gâteaux differentiable and Ψ' is completely continuous (i.e., if $x_n \rightharpoonup x$ then $\Psi'(x_n) \rightarrow \Psi'(x)$, see [12, Definition 2.95]).*

Proof. By (2.14) Ψ is well-posed. Now, we compute and show that

$$\langle \Psi'(u), v \rangle = \int_{\Omega} h_{\lambda,\mu}^*(x, u)v dx \quad \forall u, v \in W_0^{1,p}(\Omega).$$

Indeed, fixing $u, v \in W_0^{1,p}(\Omega)$, we get

$$\langle \Psi'(u), v \rangle = \lim_{t \rightarrow 0^+} \frac{\Psi(u + tv) - \Psi(u)}{t} = \lim_{t \rightarrow 0^+} \int_{\Omega} \frac{H_{\lambda,\mu}(x, u + tv) - H_{\lambda,\mu}(x, u)}{t} dx,$$

and Torricelli–Barrow’s theorem ensures that, for each $t \in [0, 1]$ and for a.a. $x \in \Omega$, we get

$$\begin{aligned} H_{\lambda,\mu}(x, u + tv) - H_{\lambda,\mu}(x, u) &= tv \int_0^1 \frac{d}{ds} H_{\lambda,\mu}(x, u + stv) ds \\ &= tv \int_0^1 h_{\lambda,\mu}^*(x, u + stv) ds. \end{aligned}$$

Furthermore, fixed $x \in \Omega$,

$$h_{\lambda,\mu}^*(x, y + z) \leq \tilde{c}_1 d^{-\gamma}(x) + c_2 + \frac{\mu}{\lambda} c_3 |y + z|^{q-1} + \frac{\mu}{\lambda} c_5 \quad \forall y, z \in \Omega.$$

Then, by Fubini’s theorem and Lebesgue’s dominate convergence theorem (see, respectively, [6, Theorems 4.5 and 4.2]), it follows that

$$\begin{aligned} &\lim_{t \rightarrow 0^+} \int_{\Omega} \frac{H_{\lambda,\mu}(x, u + tv) - H_{\lambda,\mu}(x, u)}{t} dx \\ &= \lim_{t \rightarrow 0^+} \int_{\Omega} \left(\int_0^1 \frac{tv h_{\lambda,\mu}^*(x, u + stv)}{t} ds \right) dx \\ &= \lim_{t \rightarrow 0^+} \int_0^1 \left(\int_{\Omega} h_{\lambda,\mu}^*(x, u + stv) v dx \right) ds \\ &= \int_{\Omega} h_{\lambda,\mu}^*(x, u) v dx. \end{aligned}$$

Finally, we prove that Ψ' is completely continuous and $\{u_n\} \subseteq W_0^{1,p}(\Omega)$ is a sequence such that $u_n \rightharpoonup u$ in $W_0^{1,p}(\Omega)$ and let $v \in W_0^{1,p}(\Omega)$,

$$\begin{aligned} & \lim_{n \rightarrow \infty} |\langle \Psi'(u_n) - \Psi(u), v \rangle| \\ &= \lim_{n \rightarrow \infty} \left| \int_{\Omega} (h_{\lambda,\mu}^*(x, u_n) - h_{\lambda,\mu}^*(x, u)) v dx \right| \\ &\leq \lim_{n \rightarrow \infty} \int_{\Omega} |h_{\lambda,\mu}^*(x, u_n) - h_{\lambda,\mu}^*(x, u)| |v| dx \\ &\leq \lim_{n \rightarrow \infty} \left(\int_{\Omega} |f^*(x, u_n) - f^*(x, u)| |v| dx + \frac{\mu}{\lambda} \int_{\Omega} |g^*(x, u_n) - g^*(x, u)| |v| dx \right). \end{aligned}$$

By Rellich–Kondrachov’s embedding theorem [6, Theorem 9.16], $W_0^{1,p}(\Omega) \hookrightarrow L^q(\Omega)$ is compact for $q \in [1, p^*)$. So, in $L^q(\Omega)$, there exists $w \in L^q(\Omega)$ such that $u_n \rightarrow u$ in $L^q(\Omega)$, $u_n \rightarrow u$ a.e. in Ω and $|u_n| \leq w$ [6, Theorem 4.9]. Moreover, from (Q_2) , (2.9)–(2.10) and the Lebesgue’s dominate convergence theorem, we get that Ψ' is completely continuous. ■

Definition 2.7. Let X be a Banach space and $J \in C^1(X, \mathbb{R})$. We say that J satisfies the Palais–Smale condition (briefly, (PS)-condition), if any sequence $\{u_n\} \subseteq X$ such that

- (1) $J(u_n)$ is bounded,
- (2) $\|J'(u_n)\|_{X^*} \rightarrow 0$ as $n \rightarrow +\infty$,

admitting a convergent subsequence.

A more general condition can be found in [3].

Definition 2.8 ([3, Section 2]). Let Φ, Ψ be two continuously Gâteaux differentiable functions; put

$$J = \Phi - \Psi,$$

and fix $r \in [-\infty, +\infty]$; we say that the function J fulfills the Palais–Smale condition cut off upper at r (in short $(PS)^r$ -condition), if any sequence $\{u_n\}$, in addition to (1) and (2) in the previous definition, accomplishing also

- (3) $\Phi(u_n) < r \forall n \in \mathbb{N}$,

possesses a convergent subsequence.

Lemma 2.9. Suppose that (Q_1) , (Q_2) , and (AR) hold. Then, for all $p < q < p^*$, the functional J_λ associated to (P^*) satisfies the (PS)-condition.

Proof. Fix $\lambda, \mu > 0$, let J_λ be as in (2.19) and $\{u_n\}$ a (PS)-sequence in $W_0^{1,p}(\Omega)$, i.e., it fulfills (1) and (2) introduced in the Definition 2.7, that is, for all $n \in \mathbb{N}$, there exists a suitable constant c_b such that

$$J_\lambda(u_n) = \frac{1}{p} \|u_n\|^p - \lambda \int_{\Omega} H_{\lambda,\mu}(x, u_n) dx \leq c_b \quad (2.21)$$

and

$$|\langle J'_\lambda(u_n), v \rangle| = \left| \int_\Omega |\nabla u_n|^{p-2} \nabla u_n \nabla v dx - \lambda \int_\Omega h_{\lambda, \mu}^*(x, u_n) v dx \right| \leq \|v\| \quad \forall v \in W_0^{1,p}(\Omega). \quad (2.22)$$

In order to prove that $\{u_n\}$ is bounded first we verify it on $\{u_n^-\}$. From (2.22), with $v = -u_n^-$, we obtain

$$\|u_n^-\|^p \leq \|u_n^-\|^p + \lambda \int_\Omega h_{\lambda, \mu}^*(x, u_n) u_n^- dx \leq \|u_n^-\|,$$

then $\|u_n^-\| \leq 1$. Now, we prove that also $\{u_n^+\}$ is bounded. Arguing as in (2.14), putting $A_1 := (\frac{c_1}{1-\gamma} + c_2)$, we have

$$\int_0^{|s|} f^*(x, t) dt \leq A_1(2 + |s|) \quad \forall s \in \mathbb{R}.$$

At this point, we use (AR). Let $n \in \mathbb{N}$ for a.a. $x \in \Omega$ and for all $s \in \mathbb{R}$, we compute

$$H_{\lambda, \mu}(x, u_n^+) \leq \int_0^{u_n^+} f^*(x, t) dt + \frac{\mu}{\lambda} \int_0^{u_n^+} g^*(x, t) dt \leq A_1(2 + u_n^+) + \frac{\mu}{\lambda} \int_0^{u_n^+} g^*(x, t) dt.$$

Define

$$\Omega_R := \{x \in \Omega : u_n^+(x) \geq R\}$$

and

$$\Omega'_R = \Omega \setminus \Omega_R.$$

We remember that we can choose $\underline{u} < 1$ and $R \geq \max\{s_1, 1\}$. So, from the previous inequalities, we point out

$$\begin{aligned} & \int_\Omega H_{\lambda, \mu}(x, u_n^+) dx \\ & \leq \int_\Omega \int_0^{u_n^+} f^*(x, t) dt dx + \frac{\mu}{\lambda} \int_\Omega \int_0^{u_n^+} g^*(x, t) dt dx \\ & \leq A_1(2|\Omega| + S_1 \|u_n^+\|) + \frac{\mu}{\lambda} \int_{\Omega_R} \int_0^{u_n^+} g^*(x, t) dt dx + \frac{\mu}{\lambda} \int_{\Omega'_R} \int_0^{u_n^+} g^*(x, t) dt dx. \end{aligned} \quad (2.23)$$

In particular, one has

$$\begin{aligned} & \int_{\Omega'_R} \int_0^{u_n^+} g^*(x, t) dt dx \quad (2.24) \\ & \leq \int_{\Omega'_R} \int_0^R g^*(x, t) dt dx \\ & = \int_{\Omega'_R} \int_0^{\underline{u}} g(x, \underline{u}) dt dx + \int_{\Omega'_R} \int_{\underline{u}}^R g(x, t) dt dx \leq (c_3 + c_4) |\Omega| (1 + R^q). \end{aligned} \quad (2.25)$$

Moreover, we have

$$\begin{aligned} & \int_{\Omega_R} \int_0^{u_n^+} g^*(x, t) dt dx \\ &= \int_{\Omega_R} \int_0^{\underline{u}} g(x, \underline{u}) dt dx + \int_{\Omega_R} \int_{\underline{u}}^R g(x, t) dt dx + \int_{\Omega_R} \int_R^{u_n^+} g(x, t) dt dx, \end{aligned} \quad (2.26)$$

where

$$\int_{\Omega_R} \int_0^{\underline{u}} g(x, \underline{u}) dt dx \leq (c_3 + c_4)|\Omega|, \quad (2.27)$$

$$\int_{\Omega_R} \int_{\underline{u}}^R g(x, t) dt dx \leq (c_3 + c_4)|\Omega|R^q, \quad (2.28)$$

since $\Omega = \Omega_R \cup \Omega'_R$, we have

$$\int_{\Omega_R} \int_R^{u_n^+} g(x, t) dt dx \leq \int_{\Omega} \int_R^{u_n^+} g(x, t) dt dx. \quad (2.29)$$

So, combining (2.23)–(2.29), we get

$$\begin{aligned} & \int_{\Omega} H_{\lambda, \mu}(x, u_n^+) dx \\ & \leq A_1(2|\Omega| + S_1 \|u_n^+\|) + \frac{\mu}{\lambda}(2(c_3 + c_4)|\Omega|(1 + R^q)) + \frac{\mu}{\lambda} \int_{\Omega} \int_R^{u_n^+} g^*(x, t) dt dx. \end{aligned}$$

Furthermore, by (AR)-condition,

$$\begin{aligned} & \int_{\Omega} H_{\lambda, \mu}(x, u_n^+) dx \\ & \leq A_1(2|\Omega| + S_1 \|u_n^+\|) + \frac{\mu}{\lambda}(2(c_3 + c_4)|\Omega|(1 + R^q)) + \frac{\mu}{\lambda\eta} \int_{\Omega} g^*(x, u_n) u_n^+ dx. \end{aligned}$$

By (2.22), with $v = u_n^+$, it follows that

$$\begin{aligned} -\|u_n^+\|^p - \|u_n^+\| & \leq -\lambda \int_{\Omega} f^*(x, u_n^+) u_n^+ dx - \mu \int_{\Omega} g^*(x, u_n^+) u_n^+ dx \\ & \leq -\mu \int_{\Omega} g^*(x, u_n) u_n^+ dx. \end{aligned} \quad (2.30)$$

By (2.21), we arrive at

$$\begin{aligned} \frac{1}{p} \|u_n^+\|^p & \leq c_b + \lambda \int_{\Omega} H_{\lambda, \mu}(x, u_n^+) dx \leq c_b + \lambda A_1(2|\Omega| + S_1 \|u_n^+\|) \\ & \quad + \mu(2(c_3 + c_4)|\Omega|(1 + R^q)) + \frac{\mu}{\eta} \int_{\Omega} g^*(x, u_n) u_n^+ dx, \end{aligned} \quad (2.31)$$

then, dividing (2.30) by η and adding to (2.31), we derive

$$\left(\frac{1}{p} - \frac{1}{\eta}\right)\|u_n^+\|^p - \left(\frac{1}{\eta} + \lambda S_1 A_1\right)\|u_n^+\| \leq c_b + 2|\Omega|[\lambda A_1 + \mu(c_3 + c_4)(1 + R^q)].$$

So, $\{u_n^+\}$ is also bounded and our claim is shown, i.e., $\{u_n\}$ is bounded in $W_0^{1,p}(\Omega)$. By Lemma 2.6, Ψ' is completely continuous, then

$$\lim_{n \rightarrow \infty} \langle \Phi'(u_n), u_n - u \rangle = \lim_{n \rightarrow \infty} \langle J'_\lambda(u_n), u_n - u \rangle + \lambda \lim_{n \rightarrow \infty} \langle \Psi'(u_n), u_n - u \rangle = 0,$$

that is,

$$\limsup_{n \rightarrow \infty} \langle -\Delta_p(u_n), u_n - u \rangle = 0.$$

Thus, by (S_+) property of $(-\Delta_p, W_0^{1,p}(\Omega))$ (see [12, Definition 2.96 and Lemma 2.111]), we have that if $u_n \rightharpoonup u$ then $u_n \rightarrow u$ and J_λ satisfies the (PS)-condition. ■

The following is folklore, but for completeness we prove it also in our setting.

Lemma 2.10. *Suppose (Q_1) , (Q_2) , and (AR) hold. Then, for all $p < q < p^*$, the functional J_λ associated to (P^*) is unbounded from below.*

Proof. Fix $\lambda, \mu > 0$. For all $M \geq 1$, we put $\Omega_M = \{x \in \Omega : M\varphi_1(x) \geq R\}$ and we consider $\Omega'_M = \Omega \setminus \Omega_M$, where $R \geq \max\{s_1, 1\}$ and φ_1 is the first positive eigenfunction of $(-\Delta_p, W_0^{1,p}(\Omega))$, normalized in $L^p(\Omega)$. From (AR), for all $\omega \geq R$,

$$\begin{aligned} \eta \int_{s_1}^\omega g(x, t) dt &\leq g(x, \omega)\omega, \\ \frac{\eta}{\omega} &\leq \frac{g(x, \omega)}{\int_{s_1}^\omega g(x, t) dt} = \frac{\frac{\partial}{\partial \omega} \left(\int_{s_1}^\omega g(x, t) dt \right)}{\int_{s_1}^\omega g(x, t) dt}. \end{aligned}$$

By integrating both sides for $s \geq R$ in $[R, s]$ w.r.t. ω , we obtain

$$\eta \ln\left(\frac{s}{R}\right) \leq \ln\left(\frac{\int_{s_1}^s g(x, t) dt}{\int_{s_1}^R g(x, t) dt}\right);$$

then,

$$\int_{s_1}^s g(x, t) dt \geq s^\eta R^{-\eta} \int_{s_1}^R g(x, t) dt. \quad (2.32)$$

Compute

$$\begin{aligned} J_\lambda(M\varphi_1) &= \frac{1}{p} \|M\varphi_1\|^p - \lambda \int_\Omega \int_0^{M\varphi_1} h_{\lambda, \mu}^*(x, t) dt dx \\ &= \frac{\lambda_1}{p} M^p - \lambda \int_\Omega \int_0^{M\varphi_1} h_{\lambda, \mu}^*(x, t) dt dx, \end{aligned}$$

we focus on the right-hand side term, in particular,

$$\int_{\Omega} \int_0^{M\varphi_1} h_{\lambda, \mu}^*(x, t) dt dx = \int_{\Omega} \int_0^{M\varphi_1} f^*(x, t) dt dx + \frac{\mu}{\lambda} \int_{\Omega} \int_0^{M\varphi_1} g^*(x, t) dt dx.$$

As in (2.14), we get

$$\int_{\Omega} \int_0^{M\varphi_1} f(x, t) dt dx \leq A_1(2|\Omega| + MS_1\|\varphi_1\|) = A_1(2|\Omega| + \lambda_1 S_1 M).$$

While, on the integral of g , we have

$$\int_{\Omega} \int_0^{M\varphi_1} g(x, t) dt dx = \int_{\Omega_M} \int_0^{M\varphi_1} g(x, t) dt dx + \int_{\Omega'_M} \int_0^{M\varphi_1} g(x, t) dt dx,$$

where

$$\int_{\Omega'_M} \int_0^{M\varphi_1} g(x, t) dt dx \leq \int_{\Omega'_M} \int_0^R g(x, t) dt dx \leq (c_3 + c_4)R^q|\Omega|$$

and

$$\int_{\Omega_M} \int_0^{M\varphi_1} g(x, t) dt dx = \int_{\Omega_M} \int_0^R g(x, t) dt dx + \int_{\Omega_M} \int_R^{M\varphi_1} g(x, t) dt dx$$

with

$$\int_{\Omega_M} \int_0^R g(x, t) dt dx \leq (c_3 + c_4)R^q|\Omega|.$$

Thanks to these estimates, we have

$$\begin{aligned} J_{\lambda}(M\varphi_1) &\leq \frac{\lambda_1 M^p}{p} + \lambda A_1(2|\Omega| + S_1 \lambda_1 M) \\ &\quad + 2\mu(c_3 + c_4)R^q|\Omega| - \mu \int_{\Omega_M} \int_R^{M\varphi_1} g(x, t) dt dx. \end{aligned}$$

By (2.32), we get

$$\int_R^s g(x, t) dt \geq \left(\frac{s^\eta}{R^\eta} - 1\right) \int_{s_1}^R g(x, t) dt.$$

We use this inequality to see that

$$\begin{aligned} J_{\lambda}(M\varphi_1) &\leq \frac{\lambda_1 M^p}{p} + \lambda A_1(2|\Omega| + S_1 \lambda_1 M) \\ &\quad + 3\mu(c_3 + c_4)R^q|\Omega| - \mu \frac{M^\eta}{R^\eta} \int_{\Omega_M} \int_{s_1}^R g(x, t) \varphi_1^\eta(x) dt dx; \end{aligned}$$

then, $J_{\lambda}(M\varphi_1) \rightarrow -\infty$ as $M \rightarrow +\infty$. ■

For the reader's convenience, let us recall our two fundamental tools. The first one is the celebrated Ambrosetti–Rabinowitz's mountain pass theorem (see [2, 7]).

Theorem 2.11 (Mountain pass theorem). *Let X be a Banach space. Let $J : X \rightarrow \mathbb{R}$ be a functional such that $J \in C^1(X, \mathbb{R})$. Let $x_0, x_1 \in X$, $r > 0$ such that $\|x_1 - x_0\| > r$ and*

$$\inf_{\|u-x_0\|=r} J(u) > \max \{J(x_0), J(x_1)\}.$$

Let

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} J(\gamma(t)),$$

where $\Gamma = \{\gamma \in C([0, 1], X) : \gamma(0) = x_0, \gamma(1) = x_1\}$. Suppose that J satisfies the Palais–Smale condition. Then, c is a critical value for J .

For the second one, fixing $r > 0$, let $\varphi(r)$ be as follows:

$$\varphi(r) = \inf_{v \in \Phi^{-1}]\!-\infty, r[} \frac{\sup_{u \in \Phi^{-1}]\!-\infty, r[} \Psi(u) - \Psi(v)}{r - \Phi(v)}. \quad (2.33)$$

Theorem 2.12 ([3, Theorem 5.2]). *Let X be a Banach space and let $\Phi, \Psi : X \rightarrow \mathbb{R}$ be two Gâteaux differentiable functions with Φ bounded from below. Fix $r > \inf_X \Phi$ and assume that for each $\lambda \in]0, \frac{1}{\varphi(r)}[$, where φ is given in (2.33), the function $J_\lambda = \Phi - \lambda\Psi$ satisfies $(PS)^r$ -condition. Then, for each $\lambda \in]0, \frac{1}{\varphi(r)}[$ there is $u^* \in \Phi^{-1}]\!-\infty, r[$ such that $J_\lambda(u^*) \leq J_\lambda(u) \forall u \in \Phi^{-1}]\!-\infty, r[$ and $J'_\lambda(u^*) = 0$.*

Remark 2.13. We point out that this type of local minima theorems follows from the ideas introduced by Ricceri in [33], in which the author used the weakly closure of suitable sublevels. In [5], Bonanno and Candito got the result starting from direct methods of calculus of variations and, in [3], Bonanno himself obtained the local minimum via Ekeland variational principle.

3. Main results

In this section, we prove our main results. In particular, we obtain the existence of at least one solution for $1 < q \leq p$ and the existence of at least two solutions for $p < q < p^*$ for problem $(P_{\lambda, \mu})$. In addition, we derive a computable estimate of the parameters λ, μ .

The following positive constants are central in the main results:

$$Z_1 := \frac{N}{p} \pi^{\frac{p}{2}} |\Omega|^{-\frac{p}{N}} \left(\frac{N-p}{p-1}\right)^{p-1} \left[\frac{\Gamma(\frac{N}{p})\Gamma(1 + N - \frac{N}{p})}{\Gamma(1 + \frac{N}{2})\Gamma(N)} \right]^p, \quad (3.1)$$

$$Z_2 := \frac{N}{p} \pi^{\frac{p}{2}} |\Omega|^{-\frac{p}{N}} \left(\frac{N-p}{p-1}\right)^{p-1} \left(\frac{q-p}{2p}\right)^{\frac{q-p}{q}} \left[\frac{\Gamma(\frac{N}{p})\Gamma(1 + N - \frac{N}{p})}{\Gamma(1 + \frac{N}{2})\Gamma(N)} \right]^p. \quad (3.2)$$

Theorem 3.1. *Assume that (Q_1) and (Q_2) hold. In addition, we suppose that*

$$\lambda \left(\frac{c_1}{1-\gamma} + c_2 \right) + \mu(c_3 + c_4) < \begin{cases} +\infty & \text{if } 1 < q < p; \\ Z_1 & \text{if } q = p, \\ Z_2 & \text{if } p < q < p^*, \end{cases}$$

where Z_1, Z_2 are as in (3.1)–(3.2). Then, problem $(P_{\lambda,\mu})$ admits at least one solution $u \in C_0^{1,\alpha}(\bar{\Omega})$, with $\alpha \in (0, 1)$.

Proof. Let J_λ, Φ , and Ψ be as in (2.19)–(2.20). By Lemma 2.6, J_λ is a C^1 -functional. Fix $r > 0$, let $\{u_n\}$ be a $(PS)^r$ -sequence. Since $\Phi(u_n) < r$ and reasoning is as in Lemma 2.9, we have that J_λ satisfies the $(PS)^r$ -condition as in Definition 2.8. In this setting, Theorem 2.12 furnishes a local minimum $u^* = u^*(\lambda, \mu)$ of J_λ for all $\lambda \in]0, \bar{\lambda}[$. So, u^* is a weak solution of (P^*) and then, by Lemma 2.5, a weak solution of $(P_{\lambda,\mu})$. To give an approximation of $\bar{\lambda}$ we need to estimate (2.33). Since $r > \inf_X \Phi = \Phi(0) = \Psi(0) = 0$ and (2.14), we detect that

$$\begin{aligned} \varphi(r) &\leq \frac{1}{r} \sup_{u \in \Phi^{-1}(]-\infty, r])} \Psi(u) \\ &\leq \frac{1}{r} \left(\sup_{\|u\| \leq (pr)^{\frac{1}{p}}} \int_{\Omega} |H_{\lambda,\mu}(x, u)| dx \right) \\ &\leq \frac{1}{r} \sup_{\|u\| \leq (pr)^{\frac{1}{p}}} \int_{\Omega} A_{\lambda,\mu} (2 + |u|^q) dx \\ &\leq \frac{1}{r} A_{\lambda,\mu} (2|\Omega| + S_q^q p^{\frac{q}{p}} r^{\frac{q}{p}}) \quad \forall r > 0. \end{aligned} \tag{3.3}$$

where $A_{\lambda,\mu}$ is defined in (2.13). We put

$$k(r) := \frac{1}{r} (2|\Omega| + S_q^q p^{\frac{q}{p}} r^{\frac{q}{p}}) \quad \text{for all } r > 0.$$

A straightforward calculation shows that

$$\lim_{r \rightarrow 0^+} k(r) = \lim_{r \rightarrow 0^+} \frac{1}{r} (2|\Omega| + S_q^q p^{\frac{q}{p}} r^{\frac{q}{p}}) = +\infty \quad \text{for all } 1 < q < p^*, \tag{3.4}$$

and, for $r \rightarrow +\infty$, three cases arise:

$$\lim_{r \rightarrow +\infty} k(r) = \begin{cases} 0 & \text{if } 1 < q < p; \\ S_p^p p & \text{if } q = p; \\ +\infty & \text{if } p < q < p^*. \end{cases} \tag{3.5}$$

To guarantee that $\varphi(r) < \frac{1}{\lambda}$ for some $r > 0$, by (3.3), it follows that

$$\varphi(r) \leq A_{\lambda,\mu} k(r) < \frac{1}{\lambda}.$$

Considering that $p > 1, q > 1$, then by (3.4)–(3.5), we have the following cases:

- (1) if $1 < q < p$, since $k(r)$ is a continuous function, there exists $\bar{r}_{\lambda,\mu} > 0$ such that

$$k(\bar{r}_{\lambda,\mu}) = \frac{1}{\lambda A_{\lambda,\mu}}.$$

Moreover, from the strictly decreasing of $k(r)$, our claim holds for all $r > \bar{r}_{\lambda,\mu}$;

- (2) if $q = p$, there exists $\bar{r}_{\lambda,\mu} > 0$ such that $\lambda A_{\lambda,\mu} = \frac{1}{k(r)} < \frac{1}{S_p^p} =: Z_1$, then since $k(r)$ is strictly decreasing, the result holds for all $r > \bar{r}_{\lambda,\mu}$;
- (3) if $p < q < p^*$, then $k(r)$ has a global minimum point for $r > 0$. We compute the external point of $k(r)$,

$$\bar{r} = \left(\frac{2p|\Omega|}{S_q^q p^{\frac{q}{p}}(q-p)} \right)^{\frac{p}{q}}, \quad \inf_{r>0} k(\bar{r}) = S_q^p p \left(\frac{2p|\Omega|}{q-p} \right)^{1-\frac{p}{q}}.$$

Thus, by (3.3), it follows that

$$\frac{1}{Z_2} := S_q^p p \left(\frac{2p|\Omega|}{q-p} \right)^{1-\frac{p}{q}} < \frac{1}{\lambda A_{\lambda,\mu}}.$$

So, also, in this case, our statement holds by keeping in mind (2.1).

Notice that our solution belongs to $C_0^{1,\alpha}(\bar{\Omega})$ with $0 < \alpha < 1$. In particular, from the estimates in (2.9) and (2.10), it is sufficient first to apply [17, Theorem 1.5.5] to obtain $u^* \in L^\infty(\Omega)$; finally, from [17, Theorem 1.5.6], we derive u belongs to a Hölder space. ■

Remark 3.2. A careful reading of the proof of Theorem 3.1, gives us, for $p < q < p^*$, an estimate on the Sobolev norm of the solution, i.e.,

$$\|u\| \leq \bar{r} = \left(\frac{2p|\Omega|}{S_q^q p^{\frac{q}{p}}(q-p)} \right)^{\frac{p}{q}}.$$

Remark 3.3. Notice that since $X = W_0^{1,p}(\Omega)$ is a reflexive Banach space (for $1 < p < +\infty$), Ψ is also weakly sequentially continuous [36, Corollary 41.9]. Then, J_λ is a C^1 -functional and it is weakly sequentially lower semi-continuous. For $1 < q \leq p$, we can obtain the existence of a solution for (P^*) as global minimum of J_λ , given that, in this case, the energy functional is also coercive. Indeed, from estimates in (2.14), one has

$$J_\lambda(u) = \frac{1}{p} \|u\|^p - \lambda \int_\Omega H_{\lambda,\mu}(x, u) dx \geq \frac{1}{p} \|u\|^p - A_{\lambda,\mu} (2|\Omega| + S_q^q \|u\|^q),$$

and so, we have the following cases:

- (1) if $1 < q < p$,

$$\lim_{\|u\| \rightarrow +\infty} J(u) = +\infty$$

for all $\lambda, \mu > 0$;

(2) if $q = p$,

$$\lim_{\|u\| \rightarrow +\infty} J(u) = +\infty$$

provided $\frac{1}{p} - A_{\lambda, \mu} S_p^p > 0$, i.e.,

$$\lambda \left(\frac{c_1}{1-\gamma} + c_2 \right) + \mu(c_3 + c_4) < \frac{1}{S_p^p}.$$

Then, by Tonelli–Weierstrass’ theorem [14, Theorem 1.2], there exists a weak solution of (P*). Our approach (Theorem 2.12), allows us to guarantee the existence of a weak solution also for the super-linear case and, in all three cases, without the assumption of weakly lower semi-continuity on J_λ .

Now, we prove the multiplicity result.

Theorem 3.4. *Assume that (Q₁) and (Q₂) hold. Moreover, we suppose that there exist $\eta > p$ and $s_1 > 0$ such that*

$$\operatorname{ess\,inf}_{x \in \Omega} \int_0^s g(x, t) dt > 0, \quad 0 < \eta \int_{s_1}^s g(x, t) dt \leq g(x, s)s. \quad (\text{AR})$$

Then, if $p < q < p^$, the problem (P _{λ, μ}) admits at least two solutions: $u^*(\lambda, \mu)$ a local minimum and $\tilde{u}(\lambda, \mu)$ a mountain pass point of the energy functional J_λ , for all λ, μ such that*

$$\lambda \left(\frac{c_1}{1-\gamma} + c_2 \right) + \mu(c_3 + c_4) < Z_2,$$

with Z_2 being as in (3.2).

Proof. From Lemma 2.5, it is enough to obtain solution of problem (P*). Moreover, the energy functional

$$J_\lambda = \Phi - \lambda\Psi$$

satisfies the (PS)-condition (Lemma 2.9) and it is a C^1 -functional (Lemma 2.6). So, by Theorem 3.1, there exists u^* local minimum of J_λ such that $J_\lambda(u^*) \leq J_\lambda(u)$ for all $u \in \bar{B}(0, r)$ for a suitable $r > 0$ as in Theorem 3.1 (see also Remark 3.2). Since J_λ is unbounded from below (Lemma 2.10), [4, Theorem 2.1] ensures that it satisfies also the mountain pass geometry and the proof is completed, owing the Theorem 2.11. ■

We desire to conclude our work by giving the following example.

Example 3.5. For every $\lambda, \mu > 0$ such that

$$2\lambda + \mu < \frac{5}{54} \sqrt[9]{\frac{3125}{64}} \left(\frac{77\pi}{1024} \right)^4.$$

Theorem 3.4 guarantees the existence of at least two solutions for the problem (1.1) proposed in the Introduction, where $\Omega \subset \mathbb{R}^5$ is a bounded domain with smooth boundary and

$|\Omega| = 1$, $p = 4$, $q = 9$, $4 < \eta \leq 9$ and $\gamma = \frac{1}{2}$. Moreover, taking into account Remark 3.2, the first solution u^* (corresponding to a local minimum of the energy functional) satisfies

$$\|u^*\| \leq \frac{10}{27} \sqrt[9]{\frac{8}{625}} \left(\frac{77\pi}{1024}\right)^4.$$

Acknowledgments. The authors are very grateful to the anonymous referees for their knowledgeable reports, which helped them improve the manuscript.

Funding. The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). This study was partly funded by Research Project of MIUR (Italian Ministry of Education, University and Research) Prin 2022 “Nonlinear differential problems with applications to real phenomena” (Grant no. 2022ZXZTN2).

References

- [1] A. Ambrosetti, H. Brezis, and G. Cerami, [Combined effects of concave and convex nonlinearities in some elliptic problems](#). *J. Funct. Anal.* **122** (1994), no. 2, 519–543 Zbl 0805.35028 MR 1276168
- [2] A. Ambrosetti and P. H. Rabinowitz, [Dual variational methods in critical point theory and applications](#). *J. Functional Analysis* **14** (1973), 349–381 Zbl 0273.49063 MR 0370183
- [3] G. Bonanno, [A critical point theorem via the Ekeland variational principle](#). *Nonlinear Anal.* **75** (2012), no. 5, 2992–3007 Zbl 1239.58011 MR 2878492
- [4] G. Bonanno, [Relations between the mountain pass theorem and local minima](#). *Adv. Nonlinear Anal.* **1** (2012), no. 3, 205–220 Zbl 1277.35170 MR 3034869
- [5] G. Bonanno and P. Candito, [Non-differentiable functionals and applications to elliptic problems with discontinuous nonlinearities](#). *J. Differential Equations* **244** (2008), no. 12, 3031–3059 Zbl 1149.49007 MR 2420513
- [6] H. Brezis, [Functional analysis, Sobolev spaces and partial differential equations](#). Universitext, Springer, New York, 2011 Zbl 1220.46002 MR 2759829
- [7] H. Brezis and L. Nirenberg, [Remarks on finding critical points](#). *Comm. Pure Appl. Math.* **44** (1991), no. 8-9, 939–963 Zbl 0751.58006 MR 1127041
- [8] H. Brezis and L. Nirenberg, [H¹ versus C¹ local minimizers](#). *C. R. Acad. Sci. Paris Sér. I Math.* **317** (1993), no. 5, 465–472 Zbl 0803.35029 MR 1239032
- [9] P. Candito, U. Guarnotta, and R. Livrea, [Existence of two solutions for singular \$\Phi\$ -Laplacian problems](#). *Adv. Nonlinear Stud.* **22** (2022), no. 1, 659–683 Zbl 1505.35192 MR 4517919
- [10] P. Candito, U. Guarnotta, and K. Perera, [Two solutions for a parametric singular \$p\$ -Laplacian problem](#). *J. Nonlinear Var. Anal.* **4** (2020), 455–468 Zbl 1473.35309
- [11] A. Canino, B. Sciunzi, and A. Trombetta, [Existence and uniqueness for \$p\$ -Laplace equations involving singular nonlinearities](#). *NoDEA Nonlinear Differential Equations Appl.* **23** (2016), no. 2, article no. 8 Zbl 1341.35074 MR 3478284
- [12] S. Carl, V. K. Le, and D. Motreanu, [Nonsmooth variational problems and their inequalities: Comparison principles and applications](#). Springer Monogr. Math., Springer, New York, 2007 Zbl 1109.35004 MR 2267795

- [13] S. Carl and K. Perera, Generalized solutions of singular p -Laplacian problems in \mathbb{R}^N . *Nonlinear Stud.* **18** (2011), no. 1, 113–124 Zbl 1217.35083 MR 2814088
- [14] D. G. Costa, *An invitation to variational methods in differential equations*. Birkhäuser, Boston, MA, 2007 Zbl 1123.35001 MR 2321283
- [15] L. Gambera and U. Guarnotta, Strongly singular convective elliptic equations in \mathbb{R}^N driven by a non-homogeneous operator. *Commun. Pure Appl. Anal.* **21** (2022), no. 9, 3031–3054 Zbl 1501.35238 MR 4484115
- [16] J. P. García Azorero, I. Peral Alonso, and J. J. Manfredi, Sobolev versus Hölder local minimizers and global multiplicity for some quasilinear elliptic equations. *Commun. Contemp. Math.* **2** (2000), no. 3, 385–404 Zbl 0965.35067 MR 1776988
- [17] L. Gasiński and N. S. Papageorgiou, *Nonsmooth critical point theory and nonlinear boundary value problems*. Series in Mathematical Analysis and Applications 8, Chapman & Hall/CRC, Boca Raton, FL, 2005 Zbl 1058.58005 MR 2092433
- [18] J. Giacomoni and K. Saoudi, $W_0^{1,p}$ versus C^1 local minimizers for a singular and critical functional. *J. Math. Anal. Appl.* **363** (2010), no. 2, 697–710 Zbl 1193.35067 MR 2564889
- [19] J. Giacomoni, I. Schindler, and P. Takáč, Sobolev versus Hölder local minimizers and existence of multiple solutions for a singular quasilinear equation. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **6** (2007), no. 1, 117–158 Zbl 1181.35116 MR 2341518
- [20] U. Guarnotta, R. Livrea, and S. A. Marano, Some recent results on singular p -Laplacian equations. *Demonstr. Math.* **55** (2022), no. 1, 416–428 Zbl 1498.35312 MR 4467097
- [21] D. D. Hai, On a class of singular p -Laplacian boundary value problems. *J. Math. Anal. Appl.* **383** (2011), no. 2, 619–626 Zbl 1223.35137 MR 2812410
- [22] A. Lê, Eigenvalue problems for the p -Laplacian. *Nonlinear Anal.* **64** (2006), no. 5, 1057–1099 Zbl 1208.35015 MR 2196811
- [23] G. M. Lieberman, The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations. *Comm. Partial Differential Equations* **16** (1991), no. 2-3, 311–361 Zbl 0742.35028 MR 1104103
- [24] G. Marino and P. Winkert, L^∞ -bounds for general singular elliptic equations with convection term. *Appl. Math. Lett.* **107** (2020), article no. 106410 Zbl 1441.35078 MR 4092591
- [25] D. Motreanu, V. V. Motreanu, and N. Papageorgiou, *Topological and variational methods with applications to nonlinear boundary value problems*. Springer, New York, 2014 Zbl 1292.47001 MR 3136201
- [26] D. Mugnai, Addendum to: Multiplicity of critical points in presence of a linking: application to a superlinear boundary value problem, NoDEA. *Nonlinear Differential Equations Appl.* **11** (2004), no. 3, 379–391, and a comment on the generalized Ambrosetti–Rabinowitz condition [mr2090280]. *NoDEA Nonlinear Differential Equations Appl.* **19** (2012), no. 3, 299–301 Zbl 1102.35040 MR 2926299
- [27] B. Opic and A. Kufner, *Hardy-type inequalities*. Pitman Res. Notes Math. Ser. 219, Longman Scientific & Technical, Harlow, 1990 Zbl 0698.26007 MR 1069756
- [28] N. S. Papageorgiou, V. D. Rădulescu, and D. D. Repovš, Pairs of positive solutions for resonant singular equations with the p -Laplacian. *Electron. J. Differential Equations* (2017), article no. 249 Zbl 1372.35089 MR 3711202
- [29] N. S. Papageorgiou, V. D. Rădulescu, and D. D. Repovš, Positive solutions for nonlinear parametric singular Dirichlet problems. *Bull. Math. Sci.* **9** (2019), no. 3, article no. 1950011 Zbl 1471.35168 MR 4045383

- [30] N. S. Papageorgiou and P. Winkert, [Singular \$p\$ -Laplacian equations with superlinear perturbation](#). *J. Differential Equations* **266** (2019), no. 2-3, 1462–1487 Zbl [1417.35050](#) MR [3906221](#)
- [31] N. S. Papageorgiou and P. Winkert, [Positive solutions for weighted singular \$p\$ -Laplace equations via Nehari manifolds](#). *Appl. Anal.* **100** (2021), no. 11, 2436–2448 Zbl [1475.35158](#) MR [4291386](#)
- [32] P. Pucci and J. Serrin, *The maximum principle*. Progr. Nonlinear Differential Equations Appl. 73, Birkhäuser, Basel, 2007 Zbl [1134.35001](#) MR [2356201](#)
- [33] B. Ricceri, [A general variational principle and some of its applications](#). *J. Comput. Appl. Math.* **113** (2000), no. 1-2, 401–410 Zbl [0946.49001](#) MR [1735837](#)
- [34] G. Talenti, [Best constant in Sobolev inequality](#). *Ann. Mat. Pura Appl. (4)* **110** (1976), 353–372 Zbl [0353.46018](#) MR [0463908](#)
- [35] J. L. Vázquez, [A strong maximum principle for some quasilinear elliptic equations](#). *Appl. Math. Optim.* **12** (1984), no. 3, 191–202 Zbl [0561.35003](#) MR [0768629](#)
- [36] E. Zeidler, *Nonlinear functional analysis and its applications: Variational methods and optimization. III*. Springer, New York, 1985 Zbl [0583.47051](#) MR [0768749](#)

Received 6 December 2023; revised 15 May 2024.

Pasquale Candito

Dipartimento di Ingegneria Civile, dell’Energia, dell’Ambiente e dei Materiali, Università degli Studi Mediterranea di Reggio Calabria, via Zehender (Località Feo di Vito), 89122 Reggio Calabria, Italy; pasquale.candito@unirc.it

Giuseppe Failla

Dipartimento di Scienze Matematiche e Informatiche, Scienze Fisiche e Scienze della Terra, Università degli Studi di Messina, Viale Ferdinando Stagno d’Alcontres, 98166 Messina, Italy; giuseppe.failla@studenti.unime.it

Roberto Livrea

Dipartimento di Matematica e Informatica, Università degli Studi di Palermo, via Archirafi, 90123 Palermo, Italy; roberto.livrea@unipa.it