

A remark on solutions to semilinear equations with Robin boundary conditions

Antonio Celentano, Alba Lia Masiello, and Gloria Paoli

Abstract. Symmetry properties of solutions to elliptic quasilinear equations have been widely studied in the context of Dirichlet boundary conditions. We show that, in the context of Robin boundary conditions, the symmetry property á la Gidas, Ni, and Nirenberg does not hold in dimension $n \geq 2$, even for superharmonic functions, and we provide an explicit example.

1. Introduction

The task of proving symmetries of solutions to quasilinear or nonlinear PDEs that reflect the symmetries of the domain has interested many authors. In this context, the classical result by Gidas, Ni, and Nirenberg, contained in the celebrated paper [12, Theorem 1], is stated as follows.

Theorem 1.1 (Gidas–Ni–Nirenberg symmetry result). *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be such that $f = f_1 + f_2$, where f_1 is a locally Lipschitz function and f_2 is non-decreasing. Then, any positive solution $u \in C^2(\bar{B})$ to the problem*

$$\begin{cases} -\Delta u = f(u) & \text{in } B, \\ u = 0 & \text{on } \partial B, \end{cases} \quad (1)$$

where B is a ball of \mathbb{R}^n of radius R , has to be radial and

$$\frac{\partial u}{\partial r} < 0 \quad \text{for } 0 < r < R.$$

In order to prove this result, the authors make use of the method of moving planes, first introduced by Aleksandrov in [1], and then applied by Serrin in [22] in the context of PDEs.

In the case $n = 2$, Lions in [14] gives an alternative proof of Theorem 1.1, which allows one to consider weaker smoothness assumptions on f and u , under the additional hypothesis $f \geq 0$. The technique used in [14] relies on the Schwartz symmetrization,

the isoperimetric inequality, and the Pohozaev identity. These techniques were applied and generalized in [13] to the context of the n -Laplacian, and, eventually, in [21] to the solutions to the p -Laplace equation for any p , in any dimension $n \geq 2$.

The result contained in [12] is a milestone in proving the symmetry of solutions to PDEs: in the linear case, see, for instance, [4, 5, 11, 20] and, for the p -Laplace operator, we refer to [3, 7, 9, 10]. In all the aforementioned papers, it is possible to prove the radiality of positive solution to (1) either under the regularity hypothesis on f stated in Theorem 1.1 (using the moving plane method as in [12]) or under the assumption $f \geq 0$ (with symmetrization techniques as in [13, 14, 21]). For a sign-changing f , the Lipschitz continuity property cannot be relaxed to Hölder continuity, as shown in [8]. Indeed, in this case, the author finds a positive solution to (1) that is not radially symmetric.

Recently, the study of symmetrization techniques for PDEs problem with Robin boundary condition has gained attention; see, for instance, [2, 15, 16]. The aim of the present work is to study the behavior of the solution to the Robin problem

$$\begin{cases} -\Delta u = f(u) & \text{in } B, \\ \frac{\partial u}{\partial \nu} + \beta u = 0 & \text{on } \partial B \end{cases} \quad (2)$$

whenever β is a positive parameter. To our knowledge, in the literature, few results deal with symmetry properties of the solutions to differential equations with Robin boundary conditions. For instance, in [6], the authors consider the following problem:

$$\begin{cases} -\varepsilon^2 \Delta u = f(u) - u & \text{in } B, \\ \varepsilon \frac{\partial u}{\partial \nu} + \beta u = 0 & \text{on } \partial B, \end{cases} \quad (3)$$

where $\varepsilon > 0$, $f : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function of the form

$$f(t) = f_1(t) - f_2(t)$$

for $t \geq 0$, with $f_1, f_2 \geq 0$, satisfying some structural growth conditions (for the precise details, see [6, Section 1]). Under these assumptions, it is possible to prove the existence of positive least-energy solutions to (3) by making use of the Palais–Smale condition and the mountain pass lemma. The authors show that there exists a $\beta_* > 1$ such that, for $\beta > \beta_*$, a least-energy solution has the maximum at the center of the ball, while, for $\beta \leq \beta_*$ and $\varepsilon \rightarrow 0$, a least-energy solution has the maximum near the boundary, and, consequently, the function cannot be radially symmetric. These results regarding the location of the maximum of a least-energy solution to problem (3) are actually obtained in [6] for the more general case of a bounded and smooth domain $\Omega \subset \mathbb{R}^n$.

For the sake of completeness, we recall that the case $\beta = 0$, i.e., the Neumann problem, and the case $\beta = +\infty$, i.e., the Dirichlet problem, have been studied, for instance, respectively, in [17–19].

In Theorem 2.1, we show that in the one-dimensional case the symmetry result for the solution to (2) holds under the standard hypotheses of Gidas, Ni, and Nirenberg and the

additional hypothesis $f \geq 0$. On the other hand, this is not the case for $n \geq 2$, as pointed out in Corollary 3.2:

In dimension $n \geq 2$, there exists a positive superharmonic function φ that is a solution to (2) and that is not radially symmetric.

So, the main novelty of our paper is that we can find a non-radial solution to problem (2) when the nonlinearity f is positive, and, moreover, this solution is explicit (see Theorem 3.1).

2. One-dimensional case: The symmetry holds

We start by analyzing the one-dimensional case.

Theorem 2.1. *Let $R > 0$, and let $I =]-R, R[$ be the open ball of radius R . Let $u \in C^2([-R, R])$ be a solution to*

$$\begin{cases} -u'' = f(u) & \text{in } I, \\ \frac{\partial u}{\partial \nu}(x) + \beta u(x) = 0 & \text{in } x = \pm R, \end{cases}$$

where $\beta > 0$. Let us assume that f satisfies the following assumptions:

- (i) $f \geq 0$ in \mathbb{R} , f is not identically zero in $u(I)$,
- (ii) $f = f_1 + f_2$, where f_1 is locally Lipschitz in \mathbb{R} and f_2 is non-decreasing.

Then, $u(x) = u(-x)$ for all $x \in [-R, R]$. Moreover,

$$u'(x) < 0, \quad x \in [0, R].$$

Proof. We divide the proof into two steps. In the first step, we prove that the function u is strictly positive, and in the second one, we prove that we can apply the result contained in Theorem 1.1.

Step 1. We start by proving that $u > 0$ in $[-R, R]$. Since $u'' \leq 0$, u' is non-increasing in $]-R, R[$, so the minimum of u on $[-R, R]$ is achieved either in $-R$ or in R . Let us denote by x_m the minimum point of u in $[-R, R]$. From the Robin boundary conditions, we have that

$$-\beta u(x_m) = \frac{\partial u}{\partial \nu}(x_m) \leq 0,$$

and, as a consequence, $u \geq 0$ in $[-R, R]$.

Now, we want to prove that $u > 0$ in $[-R, R]$. By contradiction, we assume that $u(x_m) = 0$. If $x_m = -R$, the Robin boundary conditions imply that

$$0 = \beta u(-R) = -\frac{\partial u}{\partial \nu}(-R) = u'(-R) \geq u'(x), \quad \forall x \in I,$$

where we have observed that u' is non-increasing. This implies that also u is a non-increasing function, so $-R$ should be both a minimum and a maximum. This is not possible, since u should be constant and this contradicts the hypothesis $f \not\equiv 0$ in $u(I)$. Therefore, we have that $x_m = R$ and, arguing as before, we have

$$0 = -\beta u(R) = \frac{\partial u}{\partial v}(R) = u'(R) \leq u'(x), \quad \forall x \in I,$$

So, u is a non-decreasing function, the point R is both a minimum and a maximum, and we get a contradiction as before.

Step 2. We prove now that $u(R) = u(-R)$. Let us assume by contradiction that $u(R) \neq u(-R)$. Without loss of generality, we can suppose $u(R) < u(-R)$. As a consequence of Step 1, the function u is strictly increasing in a neighborhood of the point $-R$, so, by the continuity of u , there exists $y \in I$ such that $u(y) = u(-R)$. Therefore, the following quantity is well defined:

$$\lambda := \inf\{t \in I : u(t) = u(-R)\},$$

and $\lambda > -R$. Moreover, the continuity of u also implies $u(\lambda) = u(-R)$ and $u(x) > u(-R)$ in $(-R, \lambda)$.

We define now the function $v := u - u(-R)$, which is a positive solution to

$$\begin{cases} -\Delta v = \tilde{f}(v) & \text{in } (-R, \lambda), \\ v = 0 & \text{in } x = -R, x = \lambda, \end{cases}$$

where $\tilde{f}(v) = f(v + u(R))$. So, we can use Theorem 1.1 in the interval $(-R, \lambda)$. We have that u is symmetric with respect to the line $x = 2^{-1}(\lambda - R)$, and, as a consequence, we get

$$\frac{du}{dv}(-R) = -\frac{du}{dx}(-R) = \frac{du}{dx}(\lambda) < 0. \quad (4)$$

Using (4) and the fact that u' is non-increasing, we obtain

$$\beta u(R) = -\frac{du}{dv}(R) = -\frac{du}{dx}(R) \geq -\frac{du}{dx}(\lambda) = -\frac{du}{dv}(-R) = \beta u(-R),$$

and, therefore,

$$u(R) \geq u(-R),$$

which is a contradiction.

From Steps 1 and 2, we have that the function $v = u - u(R)$ is a positive solution to

$$\begin{cases} -\Delta v = \tilde{f}(v) & \text{in } I, \\ v = 0 & \text{in } x = \pm R. \end{cases}$$

So, we can conclude by using Theorem 1.1. ■

We do not know if the hypotheses (i)–(ii) on the function f are the optimal ones to obtain the symmetry result. Nevertheless, we will show in Remark 3.1, in the next session, that the assumption $f \geq 0$ cannot be removed.




Figure 1. Construction of the function φ .

3. Counterexample: Symmetry breaking in dimension $n \geq 2$

In the following, we will denote by $|\cdot|$ the Euclidean norm in \mathbb{R}^n .

Theorem 3.1. *Let $B_R \subset \mathbb{R}^n$, $n \geq 2$, be the ball centered at the origin with radius R , let β be a positive constant, and let $x_0 \neq 0$ in B_R . Then, there exists a positive function $\varphi \in C^\infty(\overline{B_R})$ that is a non-radial function in B_R (i.e., $\varphi(x) \neq \varphi(|x|)$), and it is a solution to*

$$\begin{cases} -\Delta\varphi = f(\varphi) & \text{in } B_R, \\ \frac{\partial\varphi}{\partial\nu} + \beta\varphi = 0 & \text{on } \partial B_R, \end{cases} \quad (5)$$

where

$$f(t) = c_1 t [c_2 t^{\frac{1}{\beta R}} + c_3 t^{\frac{2}{\beta R}}], \quad (6)$$

with $c_1, c_3 > 0$, $c_2 \in \mathbb{R}$ defined as follows:

$$c_1 = 2\beta R, \quad c_2 = -2(\beta R + 1) + n, \quad c_3 = 2(\beta R + 1)\alpha^2, \quad \alpha^2 = R^2 - |x_0|^2.$$

Proof. We define the following quantities (see Figure 1):

- $a := |x_0|$,
- $\alpha^2 := R^2 - a^2 > 0$.

We show the existence of a positive function $\varphi(x) = \varphi(|x - x_0|) = \varphi(r)$ such that

$$\frac{\partial\varphi}{\partial\nu} + \beta\varphi = 0 \quad \text{on } \partial B_R, \quad (7)$$

where ν is the unit outer normal to ∂B_R . Let us fix $x \in \partial B_R$. Being

$$\nabla(\varphi(x)) = \varphi'(r) \frac{x - x_0}{r}, \quad \nu(x) = \frac{x}{R},$$

the Robin boundary conditions (7) become

$$\varphi'(r) \frac{x - x_0}{r} \cdot \frac{x}{R} + \beta\varphi(r) = 0. \quad (8)$$

Denoting, now, by θ the angle between the vectors x_0 and x , we have the following relation:

$$\cos(\theta) = \frac{R^2 + a^2 - r^2}{2aR}. \quad (9)$$

So, from (8) and (9), recalling that $\alpha^2 = R^2 - a^2$, we have

$$\frac{\varphi'(r)}{rR} (R^2 - Ra \cos(\theta)) + \beta\varphi(r) = \frac{\varphi'(r)}{2rR} (r^2 + \alpha^2) + \beta\varphi(r) = 0,$$

and, therefore,

$$\frac{\varphi'(r)}{\varphi(r)} = -(2\beta R) \frac{r}{r^2 + \alpha^2}. \quad (10)$$

Integrating (10), we get

$$\varphi(r) = \frac{c}{(r^2 + \alpha^2)^{\beta R}}. \quad (11)$$

If we choose $c = 1$, we have

$$\begin{aligned} \varphi'(r) &= -\frac{2\beta R r}{(r^2 + \alpha^2)^{\beta R + 1}}, \\ \varphi''(r) &= -(-\beta R - 1) \frac{4\beta R r^2}{(r^2 + \alpha^2)^{\beta R + 2}} - \frac{2\beta R}{(r^2 + \alpha^2)^{\beta R + 1}}. \end{aligned}$$

and, consequently,

$$\begin{aligned} -\Delta\varphi &= -\varphi''(r) - \frac{n-1}{r}\varphi'(r) \\ &= \frac{2\beta R}{(r^2 + \alpha^2)^{\beta R + 1}} \left[n - 2(\beta R + 1) + (\beta R + 1) \frac{2\alpha^2}{r^2 + \alpha^2} \right] \\ &= 2\beta R \varphi(r)^{\frac{1}{\beta R} + 1} \left[n - 2(\beta R + 1) + 2(\beta R + 1) \alpha^2 \varphi(r)^{\frac{1}{\beta R}} \right] \\ &= f(\varphi(r)), \end{aligned}$$

where f is the function defined in (6). So, we have proved the desired claim, since we have found a non-radial function of the form $\varphi(x) = \varphi(|x - x_0|) = \varphi(r)$, defined in (11), that satisfies (5). \blacksquare

As a consequence of Theorem 3.1, we obtain the following corollary.

Corollary 3.2. *Let $n \geq 2$. There exists a positive superharmonic function φ that is a solution to (2) and that is not radially symmetric.*

Proof. In the case $n = 2$, the right-hand side of (6) becomes

$$f(t) = 4\beta R t \left(-\beta R t^{\frac{1}{\beta R}} + \alpha^2 (1 + \beta R) t^{\frac{2}{\beta R}} \right).$$

We notice that

$$f(t) \geq 0, \quad \text{if } t \geq \left(\frac{\beta R}{\alpha^2 (1 + \beta R)} \right)^{\beta R}, \quad (12)$$

so the function $f \circ \varphi$ is positive if φ satisfies (12) for all $x \in B_R$, and this follows by imposing the following geometric constraint:

$$\beta \leq \frac{R - a}{R(R + a)}.$$

If $n \geq 3$, we can choose the constant $c_2 \geq 0$; by imposing the condition

$$\beta \leq \frac{n - 2}{2R} \quad (13)$$

and, under these assumptions, we have that $f(t) \geq 0$ for $t \geq 0$.

Therefore, we can see that, by imposing the geometrical constraints (12) and (13), respectively, for $n = 2$ and $n \geq 3$, the function φ defined in (11) is an example of positive superharmonic function, which is non-radial and satisfies (5). ■

We conclude with a remark on the one-dimensional case.

Remark 3.1. The function φ defined in Theorem 3.1, in the case $n = 1$, satisfies the problem

$$\begin{cases} -\varphi'' = f(\varphi) & \text{in } (-R, +R), \\ \frac{\partial \varphi}{\partial \nu} + \beta \varphi = 0 & \text{in } x = \pm R. \end{cases}$$

We note that $\varphi \in C^\infty([-R, R])$ and f is a locally Lipschitz function, but f does not satisfy the hypothesis (i), that is, the positiveness. Indeed, by straightforward computations, we obtain that $f \circ \varphi$ is a sign-changing function in $\varphi([-R, +R])$ for every $\beta > 0$ and $R > a > 0$.

Acknowledgments. The authors would like to thank Professor Carlo Nitsch for the useful and insightful discussions.

Funding. All the authors are partially supported by Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of Istituto Nazionale di Alta Matematica (INdAM). During the writing of the present work, Gloria Paoli was supported by the Alexander von Humboldt Foundation through an Alexander von Humboldt Research Fellowship.

Conflicts of interest. There are no conflicts of interest to disclose regarding the publication of this paper.

References

- [1] A. D. Aleksandrov, Uniqueness theorems for surfaces in the large. V. *Amer. Math. Soc. Transl.* (2) **21** (1962), 412–416 Zbl [0119.16603](#) MR [0150710](#)
- [2] A. Alvino, C. Nitsch, and C. Trombetti, [A Talenti comparison result for solutions to elliptic problems with Robin boundary conditions](#). *Comm. Pure Appl. Math.* **76** (2023), no. 3, 585–603 Zbl [1525.35076](#) MR [4544805](#)
- [3] M. Badiale and E. Nabana, [A note on radiality of solutions of \$p\$ -Laplacian equation](#). *Appl. Anal.* **52** (1994), no. 1-4, 35–43 Zbl [0841.35008](#) MR [1380325](#)
- [4] H. Berestycki and L. Nirenberg, [Monotonicity, symmetry and antisymmetry of solutions of semilinear elliptic equations](#). *J. Geom. Phys.* **5** (1988), no. 2, 237–275 Zbl [0698.35031](#) MR [1029429](#)
- [5] H. Berestycki and L. Nirenberg, [On the method of moving planes and the sliding method](#). *Bol. Soc. Brasil. Mat. (N.S.)* **22** (1991), no. 1, 1–37 Zbl [0784.35025](#) MR [1159383](#)
- [6] H. Berestycki and J. Wei, [On singular perturbation problems with Robin boundary condition](#). *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **2** (2003), no. 1, 199–230 Zbl [1121.35008](#) MR [1990979](#)
- [7] F. Brock, [Radial symmetry for nonnegative solutions of semilinear elliptic equations involving the \$p\$ -Laplacian](#). In *Progress in partial differential equations, Vol. 1 (Pont-à-Mousson, 1997)*, pp. 46–57, Pitman Res. Notes Math. Ser. 383, Longman, Harlow, 1998 Zbl [0920.35051](#) MR [1628044](#)
- [8] F. Brock, [Continuous rearrangement and symmetry of solutions of elliptic problems](#). *Proc. Indian Acad. Sci. Math. Sci.* **110** (2000), no. 2, 157–204 Zbl [0965.49002](#) MR [1758811](#)
- [9] L. Damascelli and F. Pacella, [Monotonicity and symmetry results for \$p\$ -Laplace equations and applications](#). *Adv. Differential Equations* **5** (2000), no. 7-9, 1179–1200 Zbl [1002.35045](#) MR [1776351](#)
- [10] L. Damascelli and B. Sciunzi, [Regularity, monotonicity and symmetry of positive solutions of \$m\$ -Laplace equations](#). *J. Differential Equations* **206** (2004), no. 2, 483–515 Zbl [1108.35069](#) MR [2096703](#)
- [11] L. E. Fraenkel, [An introduction to maximum principles and symmetry in elliptic problems](#). Cambridge Tracts in Math. 128, Cambridge University Press, Cambridge, 2000 Zbl [0947.35002](#) MR [1751289](#)
- [12] B. Gidas, W. M. Ni, and L. Nirenberg, [Symmetry and related properties via the maximum principle](#). *Comm. Math. Phys.* **68** (1979), no. 3, 209–243 Zbl [0425.35020](#) MR [0544879](#)
- [13] S. Kesavan and F. Pacella, [Symmetry of positive solutions of a quasilinear elliptic equation via isoperimetric inequalities](#). *Appl. Anal.* **54** (1994), no. 1-2, 27–37 Zbl [0833.35040](#) MR [1382205](#)
- [14] P.-L. Lions, [Two geometrical properties of solutions of semilinear problems](#). *Applicable Anal.* **12** (1981), no. 4, 267–272 Zbl [0445.35043](#) MR [0653200](#)
- [15] A. L. Masiello and G. Paoli, [Rigidity results for the \$p\$ -Laplacian Poisson problem with Robin boundary conditions](#). *J. Optim. Theory Appl.* (2024), DOI [10.1007/s10957-024-02442-1](#)
- [16] A. L. Masiello and G. Paoli, [A rigidity result for the Robin torsion problem](#). *J. Geom. Anal.* **33** (2023), no. 5, article no. 149 Zbl [1512.35176](#) MR [4554054](#)
- [17] W.-M. Ni and I. Takagi, [On the shape of least-energy solutions to a semilinear Neumann problem](#). *Comm. Pure Appl. Math.* **44** (1991), no. 7, 819–851 Zbl [0754.35042](#) MR [1115095](#)
- [18] W.-M. Ni and I. Takagi, [Locating the peaks of least-energy solutions to a semilinear Neumann problem](#). *Duke Math. J.* **70** (1993), no. 2, 247–281 Zbl [0796.35056](#) MR [1219814](#)

- [19] W.-M. Ni and J. Wei, [On the location and profile of spike-layer solutions to singularly perturbed semilinear Dirichlet problems](#). *Comm. Pure Appl. Math.* **48** (1995), no. 7, 731–768
Zbl [0838.35009](#) MR [1342381](#)
- [20] E. Rosset, [An approximate Gidas–Ni–Nirenberg theorem](#). *Math. Methods Appl. Sci.* **17** (1994), no. 13, 1045–1052 Zbl [0806.35041](#) MR [1300801](#)
- [21] J. Serra, [Radial symmetry of solutions to diffusion equations with discontinuous nonlinearities](#). *J. Differential Equations* **254** (2013), no. 4, 1893–1902 Zbl [1263.35117](#) MR [3003296](#)
- [22] J. Serrin, [A symmetry problem in potential theory](#). *Arch. Rational Mech. Anal.* **43** (1971), 304–318 Zbl [0222.31007](#) MR [0333220](#)

Received 26 October 2023; revised 27 June 2024.

Antonio Celentano

Dipartimento di Matematica e Applicazioni “R. Caccioppoli”, Università degli Studi di Napoli Federico II, Via Cintia, Complesso Universitario di Monte S. Angelo, 80126 Napoli, Italy;
antonio.celentano2@unina.it

Alba Lia Masiello

Dipartimento di Matematica e Applicazioni “R. Caccioppoli”, Università degli Studi di Napoli Federico II, Via Cintia, Complesso Universitario di Monte S. Angelo, 80126 Napoli, Italy;
albalia.masiello@unina.it

Gloria Paoli

Dipartimento di Matematica e Applicazioni “R. Caccioppoli”, Università degli Studi di Napoli Federico II, Via Cintia, Complesso Universitario di Monte S. Angelo, 80126 Napoli, Italy;
gloria.paoli@unina.it