

MATHEMATISCHES FORSCHUNGSINSTITUT OBERWOLFACH

Report No. 14/2024

DOI: 10.4171/OWR/2024/14

## **K-Stability, Birational Geometry and Mirror Symmetry**

Organized by  
Thibaut Delcroix, Montpellier  
Liana Heuberger, Bath  
Susanna Zimmermann, Orsay

17 March – 22 March 2024

**ABSTRACT.** The workshop *K-stability, Birational Geometry and Mirror Symmetry* presented recent advances in all three topics, in the form of research level mini-courses, research talks and lightning talk sessions. Deep interactions between the three topics were highlighted, together with applications (e.g. to K-moduli or subgroups of the Cremona groups) and new directions (such as non-Archimedean methods in K-stability and Mirror Symmetry).

*Mathematics Subject Classification (2020):* 14E07, 14E30, 14J33, 14J45, 14B07, 14N35, 14J10, 32Q20, 32Q26.

*License:* Unless otherwise noted, the content of this report is licensed under CC BY SA 4.0.

### **Introduction by the Organizers**

The workshop *K-stability, Birational Geometry and Mirror Symmetry*, organised by Thibaut Delcroix (Montpellier), Liana Heuberger (Bath) and Susanna Zimmermann (Orsay) was well attended with 44 participants. The program of the workshop consisted in three mini-courses of 4 hours each, delivered by pairs of experts each morning from Monday to Thursday, 8 talks, and two sessions of 5 lightning talks of 10 minutes each. In what follows, we describe the main themes covered by the workshop.

The mini-course on birational geometry, with lectures by Julia Schneider and Jeremy Blanc, focused on recent major results on quotients of the Cremona group (the group of birational self-maps of projective space) in various dimensions and over various fields. One of the major means of understanding this classical group (or, rather, the groupoid of birational maps between Mori fibre spaces), expanded upon in detail during the mini-course, is its presentation in terms of Sarkisov links

and elementary relations. Focusing first on surfaces with Julia Schneider's two lectures, and switching to higher dimensions in Jérémy Blanc's talks, surveyed the main ideas from groundbreaking results of Blanc, Lamy, Zimmermann, Schneider, Yasinsky et al. on the quotients of the Cremona group, as well as highlighting key ingredients of their proofs.

The mini-course on K-stability, by Ivan Cheltsov and Elena Denisova, centred around recent techniques in K-stability applied to the explicit study of K-stability for smooth Fano threefolds. They presented the vast program whose results are collected in the book "The Calabi problem for Fano threefolds" by Araujo, Cheltsov et al. The purpose of this study is to determine which smooth Fano threefolds are K-stable, as well as finding applications which study of their behaviour in families, i.e. to determine the properties of K-moduli spaces. Here, a key input is Fujita and Li's valuative criterion for the K-stability of Fano manifolds, together with Abban and Zhuang's method of finding a lower bound of the delta invariant from a geometrically relevant choice of flag of subvarieties. With these tools at hand, the problem essentially reduces to a difficult and subtle, but explicit, study of the birational geometry of Fano threefolds, and Sarkisov links appear again in this setting.

The third mini-course, by Giulia Gugiatti and Andrea Petracci, presented recent advances around the Fano/Landau-Ginzburg (F/LG) correspondence in mirror symmetry. Various incarnations of this correspondence were discussed, culminating in a construction of smoothings of affine singular toric threefolds by Corti-Hacking-Petracci. A nuanced discussion of the limits of currently applicable systematic methods crystallised in the study of Johnsson-Koll ar surfaces, whose anticanonical system is known to be empty. A discussion of the homological mirror symmetry (a stronger version of the mirror theorem than the F/LG correspondence) for del Pezzo surfaces, by Auroux-Katzarkov-Orlov was incorporated in this more general singular setting.

Some of the research talks shared striking similarities, highlighting the interconnectedness and broad impact of the chosen topics. The talks of Anne-Sophie Kaloghiros and Yuchen Liu focussed on the topic of K-moduli spaces, which was partly addressed in the mini-course, and expanded upon it in different directions. The result of Kaloghiros describes certain irreducible components of K-moduli space for Fano threefolds in the simplest possible non-trivial case (i.e. not reduced to a point), namely when it is of dimension one, by combining birational geometry with a technique from mirror symmetry which produces toric smoothings. Liu delivered, on the final day of the workshop, a talk on the moduli continuity method for K-stability. In connection to the mini-course, he explained how to apply this method to a family of smooth Fano threefolds (No. 2-15 in the Mori-Mukai list) to explicitly determine its irreducible component in the K-moduli space using its description as a GIT quotient.

In the last part of the workshop, the talks of S ebastien Boucksom and Enrica Mazzon were hinging toward non-Archimedean techniques in two of the main subjects of the workshop. Boucksom explained how non-Archimedean formalism

allows for a valuative interpretation of K-stability beyond the Fano case, emphasizing that nonetheless explicit methods as presented in the mini-course are currently out of reach in the general polarization case and require further research. Mazzon's talk introduced the SYZ conjecture in mirror symmetry, and explained how the non-Archimedean approach first proposed by Kontsevich and Soibelman is giving rise to exciting new results appearing in work of Yang Li and Hultgren-Jonsson-Mazzon-McCleerey.

The remaining talks provide a wide overview of various recent advances and fields of investigation in algebraic geometry, which we briefly summarize. Hendrik Süß highlighted surprising relations between the notion of normalized volume of singularities, underlying the valuative criterion for K-stability, and convex geometry in the setting of toric singularities. Ana-Maria Castravet's talk introduced the notion of higher Fano manifolds, and presented an overview of known examples and non-examples. Carolina Araujo focused on a problem of Gizatullin, asking when automorphisms of a quartic surface in the projective space are induced by Cremona transformations of the projective space, and presented her results in this direction relating to the Sarkisov program presented in the mini-courses. Finally, Alex Duncan's talk focused, in contrast to the mini-course, on small, finite subgroups of Cremona groups, measuring the complexity of the Cremona group in terms of the representation dimension of its finite subgroups.

We ran two hour-long sessions of 10-minute lightning talks, so that participants could advertise a new result, a question or a key idea. Both junior and senior participants were eager to contribute, resulting in a widely representative, discussion-fuelling workshop. We also welcomed the input of an Oberwolfach Research Fellow which was not originally planned as a participant of the workshop. The lightning talks were given by Andrés Jaramillo Puentes, Jaroslaw Wisniewski, Roland Púček, Egor Yasinsky, King Leung Lee, Eduardo Alves da Silva, Ignacio Barros, Erroxe Etxabbarri Alberdi, Tran Trung Nghiem and Jürgen Hausen.

*Acknowledgement:* The MFO and the workshop organizers would like to thank the National Science Foundation for supporting the participation of junior researchers in the workshop by the grant DMS-2230648, "US Junior Oberwolfach Fellows". Moreover, the MFO and the workshop organizers would like to thank the Simons Foundation for supporting Alexander R. Duncan in the "Simons Visiting Professors" program at the MFO.



## Workshop: K-Stability, Birational Geometry and Mirror Symmetry

### Table of Contents

Jérémy Blanc and Julia Schneider	
<i>Sarkisov links and Cremona groups</i> .....	833
Ivan Cheltsov and Elena Denisova (joint with many people)	
<i>Calabi problem for smooth Fano threefolds</i> .....	836
Giulia Gugiatti and Andrea Petracci	
<i>Minicourse on Mirror Symmetry for Fano varieties</i> .....	844
Hendrik Süß (joint with Joaquín Moraga)	
<i>On the normalised volume of toric singularities</i> .....	847
Ana-Maria Castravet (joint with Carolina Araujo, Roya Beheshti, Kelly Jabbusch, Enrica Mazzon, Svetlana Makarova, Will Reynolds, Libby Taylor, Nivedita Viswanathan)	
<i>Higher Fano Manifolds</i> .....	848
Anne-Sophie Kaloghiros	
<i>On K-moduli spaces of Fano 3-folds</i> .....	852
Carolina Araujo	
<i>On a problem of Gizatullin</i> .....	855
Alexander Duncan (joint with Jason Bailey Heath, Christian Urech)	
<i>Finite subgroups of Cremona groups and representation dimension</i> .....	858
Sébastien Boucksom (joint with Mattias Jonsson)	
<i>A valuative criterion for K-stability: beyond the Fano case</i> .....	860
Yuchen Liu (joint with Chenyang Xu, Junyan Zhao)	
<i>Moduli Continuity method in K-stability</i> .....	861
Enrica Mazzon	
<i>Non-archimedean approach to SYZ conjecture</i> .....	864



## Abstracts

### Sarkisov links and Cremona groups

JÉRÉMY BLANC AND JULIA SCHNEIDER

Let  $K$  be a field. The *Cremona group of rank  $n$  over  $K$*  is the group of birational transformations of the projective space of dimension  $n$ , written  $\text{Bir}_K(\mathbb{P}^n)$ . Non-trivial normal subgroups of  $\text{Bir}_K(\mathbb{P}^2)$  have been constructed using geometric group theory (see [CL13] for algebraically closed fields and [Lon16] for arbitrary fields). The resulting quotients, however, are complicated and remain mysterious. In fact, if  $K$  is algebraically closed, every non-trivial quotient of  $\text{Bir}_K(\mathbb{P}^2)$  contains  $\text{PGL}_3(K)$ ; in particular, there is no finite and no abelian quotients, except the trivial group.

For fields that are not algebraically closed, this is no longer true, as was discovered by Zimmermann who showed that the abelianisation of  $\text{Bir}_{\mathbb{R}}(\mathbb{P}^2)$  is isomorphic to  $\bigoplus_{\mathbb{R}} \mathbb{Z}/2\mathbb{Z}$  [Zim18]. This opened up the approach of studying Cremona groups via their group homomorphisms. For perfect fields  $K$ , quotients of  $\text{Bir}_K(\mathbb{P}^2)$  that are a free product of direct sums of  $\mathbb{Z}/2\mathbb{Z}$  have been constructed in [LZ20, Sch22, Zim22]. For  $n \geq 3$ , quotients of  $\text{Bir}_{\mathbb{C}}(\mathbb{P}^n)$  that are a free product of direct sums of  $\mathbb{Z}/2\mathbb{Z}$  have been constructed in [BLZ21], showing that complex Cremona groups in higher dimensions are not simple. More quotients were constructed in this setting [BY20, Zik23]. Finally, using motivic invariants of birational maps, it was shown that  $\mathbb{Z}$  is a quotient of  $\text{Bir}_{\mathbb{C}}(\mathbb{P}^n)$  for  $n \geq 4$ , as well as of  $\text{Bir}_{\mathbb{Q}}(\mathbb{P}^n)$  for  $n \geq 3$  [LS22]. Hence, these Cremona groups are not generated by elements of finite order, contrasting the case of plane Cremona groups over perfect fields which are generated by involutions [LS24]. Notably, the question whether  $\text{Bir}_{\mathbb{C}}(\mathbb{P}^3)$  is generated by elements of finite order remains open.

### 1. THE RESULTS

Together with Egor Yasinsky, we studied  $\text{Bir}_K(S)$ , where  $S$  is a non-trivial Severi-Brauer surface over a perfect field  $K$ , that is,  $S_{\bar{K}} \simeq \mathbb{P}_{\bar{K}}^2$  and  $S(K) = \emptyset$ .

**Theorem 1.** [BSY23] *Let  $S$  be a non-trivial Severi-Brauer surface. For  $d \in \{3, 6\}$  we denote by  $\mathcal{P}_d$  the set of degree  $d$  points of  $S$  up to the action of  $\text{Aut}_K(S)$ . Then,  $|\mathcal{P}_3| \geq 2$  and for each  $p \in \mathcal{P}_3$ , there is a surjective group homomorphism*

$$\Psi: \text{Bir}_K(S) \rightarrow \bigoplus_{\mathcal{P}_3 \setminus \{p\}} \mathbb{Z}/3\mathbb{Z} * \left( \ast_{\mathcal{P}_6} \mathbb{Z} \right).$$

*In particular,  $\text{Bir}_K(S)$  is not a perfect group (and is thus not simple). Moreover, if  $\mathcal{P}_6 \neq \emptyset$ , then  $\text{Bir}_K(S)$  is not generated by elements of finite order.*

As an application, we study fibrations  $X \rightarrow B$  whose generic fibre is a non-trivial Severi-Brauer surface  $S$  over  $\mathbb{C}(B)$ . Such  $S$  exist only if  $\dim B \geq 2$ , and hence  $\dim X \geq 4$ . For  $n \geq 4$ , we construct a surjective group homomorphism from  $\text{Bir}_{\mathbb{C}}(\mathbb{P}^n)$  to  $F(\mathbb{C}) = \ast_{\mathbb{C}} \mathbb{Z}$ , the free group indexed by  $\mathbb{C}$ . This implies the following:

**Theorem 2.** [BSY23] *Let  $n \geq 4$ , and let  $G$  be any group of cardinality  $|G| \leq |\mathbb{C}|$ . Then  $G$  is a quotient of  $\text{Bir}_{\mathbb{C}}(\mathbb{P}^n)$ .*

2. THE STRATEGY: THE SARKISOV PROGRAM

Instead of just considering  $\text{Bir}(X)$  one can consider a larger class of birational maps, namely the *groupoid*  $\text{BirMori}(X)$  consisting of birational maps between *Mori fibre spaces* birational to  $X$ . The *Sarkisov program* states that  $\text{BirMori}(X)$  is generated by *Sarkisov links* and isomorphisms of Mori fibre spaces, and every relation between Sarkisov links is generated by *elementary relations* and trivial relations (see [Isk96, Cor95, HM13, Kal13, LZ20, BLZ21]). In order to define Mori fibre spaces, Sarkisov links and elementary relations in a uniform way, we introduce rank  $r$  fibrations:

Assume first that  $X$  is a smooth projective surface over a perfect field  $K$ . A surjective morphism  $\pi: X \rightarrow B$  with connected fibres is a *rank  $r$  fibration* if  $B$  is smooth with  $\dim(B) < \dim(X)$ , relative Picard rank  $\rho(X/B) = r$ , and  $-K_X$  is relatively ample. Note that  $X/\text{Spec}(K)$  is a rank  $r$  fibration exactly if  $X$  is a del Pezzo surface of Picard rank  $\rho(X) = r$ . The higher dimensional analogue of rank  $r$  fibrations tries to capture some of the nice properties of del Pezzo surfaces in terms of birational geometry. In particular (see [BLZ21] for the precise definition), one requires that  $X$  is terminal and  $\mathbb{Q}$ -factorial (mild singularities),  $X/B$  is a Mori Dream space (hence one can run any MMP from  $X$  over  $B$ , and there are only finitely many outputs), every output of an MMP from  $X$  over  $B$  is again  $\mathbb{Q}$ -factorial and terminal and  $-K_X$  is big over  $B$ .

For example, note that the Hirzebruch surface  $\mathbb{F}_2$  is a rank 1 fibration over  $\mathbb{P}^1$  but not a rank 2 fibration over  $\text{Spec}(K)$ :  $-\mathbb{K}_{\mathbb{F}_2}$  is big but the contraction of the section with self-intersection  $-2$  produces a singular surface (not terminal).

The observant reader might have realised by now that rank 1 fibrations are exactly Mori fibre spaces. A rank 2 fibration  $Y/B$  dominates exactly two rank 1 fibrations  $X_1/B_1$  and  $X_2/B_2$ , giving rise to a diagram as follows, called *Sarkisov diagram*, where the dotted arrow denotes a pseudo-isomorphism:

$$\begin{array}{ccc}
 Y = Y_1 & \cdots\cdots\cdots & Y_2 \\
 \downarrow & & \downarrow \\
 X_1 & \cdots\cdots\cdots & X_2 \\
 \downarrow & & \downarrow \\
 B_1 & \longrightarrow & B \longleftarrow B_2
 \end{array}$$

A birational map  $\chi: X_1 \dashrightarrow X_2$  between two Mori fibre spaces is called *Sarkisov link* exactly if it can be put into such a diagram. Depending on whether the birational morphisms  $Y_i \rightarrow X_i$  are isomorphisms or divisorial contractions one obtains four types of links; if both of them are divisorial the link is said to be a Sarkisov link of type II.

A rank 3 fibration gives rise to a relation between Sarkisov links. These are the *elementary relations* that generate relations, as explained before.

This allows us to define *groupoid* homomorphisms  $\text{BirMori}(X) \rightarrow G$  as follows. We choose a certain type of Sarkisov links, typically a link of type II, such that



we have a good control over the elementary relations in which they appear. For a Sarkisov link  $\chi: X_1 \dashrightarrow X_2$  of type II, denote by  $\Gamma \subset X_1$  the exceptional locus of  $\chi$ , and  $Y \rightarrow X_1$  the blow-up at  $\Gamma$ . In the case of surfaces, Bertini involutions where  $\Gamma$  is a point of degree 8 were used in [LZ20]: They do not appear in any elementary relation (because  $Y$  is a del Pezzo surface of degree 1) but they do satisfy the trivial relation  $\chi = \chi^{-1}$ , giving a quotient of the form  $*_{\mathcal{P}_8} \mathbb{Z}/2\mathbb{Z}$ .

For Theorem 1 we use the fact that the degree of any point on a non-trivial Severi-Brauer surface  $S$  is divisible by 3, and that any Sarkisov link  $\chi$  from  $S$  goes to the opposite Severi-Brauer surface  $S^{\text{op}}$ . The construction of the groupoid homomorphism goes as follows, depending on the degree  $d$  of the base-point  $\Gamma$  of  $\chi$ . If  $d = 6$ , then  $\chi$  does not appear in any elementary relation, so we send  $\chi$  onto  $1_{[\Gamma]} \in *_{\mathcal{P}_6} \mathbb{Z}$ . If  $d = 3$ , there is an elementary relation involving  $\chi$ ; we send  $\chi$  onto  $1_{[\Gamma]} \in \bigoplus_{\mathcal{P}_3} \mathbb{Z}/3\mathbb{Z}$ .

In higher dimensions, [BLZ21] used links between conic bundles where  $\Gamma$  has large covering gonality. To prove Theorem 2, we use Sarkisov links  $\chi$  between Mori fibre spaces whose generic fibre is a non-trivial Severi-Brauer surface, such that  $\chi$  induces a Sarkisov link  $\hat{\chi}$  between the corresponding Severi-Brauer surfaces, and, moreover, such that  $\Gamma$  has large covering genus.

## REFERENCES

- [BLZ21] J. Blanc, S. Lamy & S. Zimmermann. Quotients of higher dimensional Cremona groups. *Acta Math*, 226(2):211–318, 2021.
- [BY20] J. Blanc & E. Yasinsky. Quotients of groups of birational transformations of cubic del Pezzo fibrations. *J. Éc. Polytech., Math.*, 7:1089–1112, 2020.
- [BSY23] J. Blanc, J. Schneider & E. Yasinsky. Birational maps of severi-brauer surfaces, with applications to cremona groups of higher rank, 2023.
- [CL13] S. Cantat & S. Lamy. Normal subgroups in the Cremona group. *Acta Math.*, 210(1):31–94, 2013. With an appendix by Yves de Cornulier.
- [Cor95] A. Corti. Factoring birational maps of threefolds after Sarkisov. *J. Algebraic Geom.*, 4(2):223–254, 1995.
- [HM13] C. D. Hacon & J. McKernan. The Sarkisov program. *J. Algebraic Geom.*, 22(2):389–405, 2013.
- [Isk96] V. A. Iskovskikh. Factorization of birational maps of rational surfaces from the viewpoint of Mori theory. *Russ. Math. Surv.*, 51(4):585–652, 1996.
- [Kal13] A.-S. Kaloghiros. Relations in the Sarkisov program. *Compos. Math.*, 149(10):1685–1709, 2013.
- [LS24] S. Lamy & J. Schneider. Generating the plane Cremona groups by involutions. *Algebr. Geom.*, 11(1):111–162, 2024.
- [LZ20] S. Lamy & S. Zimmermann. Signature morphisms from the Cremona group over a non-closed field. *J. Eur. Math. Soc. (JEMS)*, 22(10):3133–3173, 2020.
- [LS22] H.-Y. Lin & E. Shinder. Motivic invariants of birational maps, 2022. to appear in *Annals of mathematics*.
- [Lon16] A. Lonjou. Non simplicité du groupe de Cremona sur tout corps. *Annales de l'Institut Fourier*, 66(5):2021–2046, 2016.
- [Sch22] J. Schneider. Relations in the Cremona group over perfect fields. *Annales de l'Institut Fourier*, 72(1):1–42, 2022.
- [Zik23] S. Zikas. Rigid birational involutions of  $\mathbb{P}^3$  and cubic threefolds. *J. Éc. Polytech., Math.*, 10:233–252, 2023.

- [Zim18] S. Zimmermann. The Abelianization of the real Cremona group. *Duke Math. J.*, 167(2):211–267, 2018.
- [Zim22] S. Zimmermann. A remark on Geiser involutions. *European Journal of Mathematics*, 8(3):1291–1306, sep 2022.

## Calabi problem for smooth Fano threefolds

IVAN CHELTSOV AND ELENA DENISOVA

(joint work with many people)

This is an abstract for four lectures given at the workshop. The goal of these lectures were to explain how to prove K-stability of smooth Fano threefolds.

### 1. INDUCTIVE APPROACH TO K-STABILITY

Let  $X$  be a smooth Fano threefold, let  $f: W \rightarrow X$  be a birational morphism such that  $W$  is normal, and let  $E$  be a prime divisor in  $W$ . We say that  $E$  is a divisor over  $X$ , and we denote this as  $E/X$ . We can relate several numbers to  $E$ . One is log-discrepancy  $A_X(E) = 1 + \text{ord}_E(K_{W/X})$ , which is easy to compute. The second one is the pseudo-effective threshold:

$$\tau(E) = \sup \left\{ u \in \mathbb{R}_{>0} \mid f^*(-K_X) - uE \text{ is pseudo-effective} \right\},$$

which is not so easy to compute. The third number is the most difficult one:

$$S_X(E) = \frac{1}{(-K_X)^n} \int_0^\infty \text{vol}(-K_X - uE) du = \frac{1}{(-K_X)^n} \int_0^{\tau(E)} \text{vol}(f^*(-K_X) - uE) du.$$

Let  $\beta(E) = A_X(E) - S_X(E)$ . Then the Fujita–Li valuation criterion [6, 7] says that  $X$  is K-stable  $\Leftrightarrow \beta(E) > 0$  for every prime divisor  $E$  over  $X$ . This criterion can be restated as follows. For every point  $P \in X$ , let

$$\delta_P(X) = \inf_{\substack{E/X \\ P \in C_X(E)}} \frac{A_X(E)}{S_X(E)}$$

where the infimum is taken by all prime divisors over  $X$  whose center on  $X$  contains the point  $P$ . Then  $X$  is K-stable if  $\delta_P(X) > 1$  for every  $P \in X$ .

To show that  $\delta_P(X) > 1$ , we can use Abban–Zhuang theory [1] and Fujita’s formula derived from this theory [2]. Namely, let us do the following:

- (1) choose a surface  $S \subset X$  such that  $P \in S$  and  $S$  has Du Val singularities;
- (2) compute  $\tau = \tau(S) = \sup \{ u \in \mathbb{Q}_{>0} \mid -K_X - uS \text{ is pseudo-effective} \}$
- (3) for  $u \in [0, \tau]$ , compute
  - $P(u)$  = the positive part of the Zariski decomposition of  $-K_X - uS$ ,
  - $N(u)$  = the negative part of the Zariski decomposition of  $-K_X - uS$ .

Then  $S_X(S) = \frac{1}{(-K_X)^3} \int_0^\tau P(u)^3 du$ . If  $S_X(S) \geq 1$ , then  $\delta_P(X) \leq 1$ . Thus, we may assume  $S_X(S) < 1$ . Then Abban–Zhuang theory and the Fujita’s formula give:

$$\delta_P(X) \geq \min \left\{ \frac{1}{S_X(S)}, \delta_P(S, W_{\bullet, \bullet}^S) \right\}$$

for

$$\delta_P(S, W_{\bullet, \bullet}^S) = \inf_{\substack{F/S \\ P \in C_S(F)}} \frac{A_S(F)}{S(W_{\bullet, \bullet}^S; F)},$$

where the infimum is taken by all prime divisors  $F$  over  $S$  such that  $P \in C_S(F)$  and

$$\begin{aligned} S(W_{\bullet, \bullet}^S; F) &= \frac{3}{(-K_X)^3} \int_0^\tau (P(u)^2 \cdot S) \cdot \text{ord}_F(N(u)|_S) du \\ &\quad + \frac{3}{(-K_X)^3} \int_0^\tau \int_0^\infty \text{vol}(P(u)|_S - vF) dv du. \end{aligned}$$

Note that the number  $\delta_P(S, W_{\bullet, \bullet}^S)$  has the same nature as  $\delta_P(X)$ , but it is always easier to compute, because  $S$  is a surface. Moreover, to estimate  $\delta_P(S, W_{\bullet, \bullet}^S)$ , we can apply Abban–Zhuang theory again.

## 2. $\delta$ -INVARIANTS OF POLARIZED SURFACES

Abban–Zhuang theory can be applied to estimate  $\delta$ -invariants of any polarized variety. To illustrate this, let  $S$  be a smooth surface, and let  $D$  be a big and nef  $\mathbb{R}$ -divisor  $D$  on the surface  $S$ . For every prime divisor  $F$  over  $S$ , we set

$$S_D(F) = \frac{1}{D^2} \int_0^\infty \text{vol}(D - vF) dv$$

similar to what we did for smooth Fano threefold in Section 1. Then we let

$$\delta_P(S, D) = \inf_{P \in C_S(F)} \frac{A_S(F)}{S_D(F)}$$

where the infimum is taken by all prime divisors  $F$  over  $S$  whose support contains the point  $P$ .

To apply Abban–Zhuang theory to estimate  $\delta_P(S, D)$ , we do the following:

- (1) choose a smooth curve  $C \subset S$  that passes through  $P$ ;
- (2) compute  $\tau = \sup \left\{ v \in \mathbb{R}_{\geq 0} \mid \text{the divisor } D - vC \text{ is pseudo-effective} \right\}$ ;
- (3) for  $v \in [0, \tau]$ , compute
  - $P(v)$  = the positive part of the Zariski decomposition of  $D - vC$ ,
  - $N(v)$  = the negative part of the Zariski decomposition of  $D - vC$ .

Then  $S_D(C) = \frac{1}{D^2} \int_0^\infty \text{vol}(D - vC) dv = \frac{1}{D^2} \int_0^\tau P(v)^2 dv$ . Thus, we have

$$\delta_P(S, D) \leq \frac{A_S(C)}{S_D(C)} = \frac{1}{S_D(C)}.$$

Set

$$\begin{aligned} S(W_{\bullet, \bullet}^C; P) &= \frac{2}{D^2} \int_0^\tau \text{ord}_P(N(v)|_C)(P(v) \cdot C) dv + \frac{1}{D^2} \int_0^\tau (P(v) \cdot C)^2 dv \\ &= \frac{2}{D^2} \int_0^\tau h(v) dv, \end{aligned}$$

where

$$h(v) = (P(v) \cdot C) \times (N(v) \cdot C)_P + \frac{(P(v) \cdot C)^2}{2}.$$

Then it follows from Abban–Zhuang theory and Fujita’s formula that

$$\delta_P(S, D) \geq \min \left\{ \frac{1}{S_D(C)}, \frac{1}{S(W_{\bullet, \bullet}^C; P)} \right\}.$$

We can apply Abban–Zhuang theory to the exceptional curve of a weighted blow up of  $S$  at  $P$ . For simplicity, let us show how to do this for the usual blow up. Namely, let  $f: \tilde{S} \rightarrow S$  be the blow up of the surface  $S$  at the point  $P$ , and let  $E$  be the  $f$ -exceptional curve. Set

$$\tilde{\tau} = \sup \left\{ v \in \mathbb{R}_{\geq 0} \mid \text{the divisor } f^*(D) - vE \text{ is pseudo-effective} \right\}.$$

For  $v \in [0, \tilde{\tau}]$ , let  $\tilde{P}(v)$  and  $\tilde{N}(v)$  be the positive and negative part of the Zariski decomposition of the divisor  $f^*(D) - vE$ , respectively. Then  $A_S(E) = 2$  and  $S_D(E) = \frac{1}{D^2} \int_0^{\tilde{\tau}} \tilde{P}(v)^2 dv$ . Thus, we have

$$\delta_P(S, D) \leq \frac{2}{S_D(E)}$$

Now, we for every point  $O \in E$ , we set

$$\begin{aligned} S(W_{\bullet, \bullet}^E; O) &= \frac{2}{D^2} \int_0^{\tilde{\tau}} \text{ord}_O(\tilde{N}(v)|_E)(\tilde{P}(v)|_E) dv + \frac{1}{D^2} \int_0^{\tilde{\tau}} (\tilde{P}(v) \cdot E)^2 dv \\ &= \frac{1}{D^2} \int_0^{\tilde{\tau}} h(v) dv, \end{aligned}$$

where

$$h(v) = (\tilde{P}(v) \cdot E) \times (\tilde{N}(v) \cdot E)_P + \frac{(\tilde{P}(v) \cdot E)^2}{2}.$$

Then Abban–Zhuang theory gives

$$\delta_P(S, D) \geq \min \left\{ \frac{2}{S_D(E)}, \inf_{O \in E} \frac{1}{S(W_{\bullet, \bullet}^E; O)} \right\}.$$

3. NEMURRO LEMMA

Now, we go back to smooth Fano threefolds. Let us use all assumptions and notations of Section 1. Set

$$D = D(u) = P(u)|_S.$$

for every  $u \in [0, \tau]$ . Consider the polarized pair  $(S, D)$ . Here, the divisor  $D$  is nef by construction. However, it may not be big in general (especially for  $u = \tau$ ), but everything we described in Section 2 still works. Thus, arguing as in Section 2, we can find an estimate

$$\delta_P(S, D) \geq q(u)$$

for  $u \in [0, \tau]$ , where  $q(u): [0, \tau] \rightarrow \mathbb{R}_{>0}$  is a continuous non-negative function. Following [3], let us show how to use this estimate for  $\delta_P(S, D)$  to estimate  $\delta_P(X)$ . Namely, observe that

$$\begin{aligned} S(W_{\bullet, \bullet}^S; F) &= \frac{3}{(-K_X)^3} \int_0^\tau (P(u)^2 \cdot S) \cdot \text{ord}_F(N(u)|_S) du \\ &\quad + \frac{3}{(-K_X)^3} \int_0^\tau \int_0^\infty \text{vol}(P(u)|_S - vF) dv du \\ &\leq \frac{3}{(-K_X)^3} \int_0^\tau (P(u)^2 \cdot S) \cdot \text{ord}_F(N(u)|_S) du \\ &\quad + \left( \frac{3}{(-K_X)^3} \int_0^\tau \frac{D^2}{q(u)} du \right) A_S(F) \end{aligned}$$

for every prime divisor  $F$  over  $S$ . Thus, if  $P \notin \text{Supp}(N(u))$  for  $u \in [0, \tau]$ , then

$$\delta_P(S, W_{\bullet, \bullet}^S) \geq \frac{1}{\frac{3}{(-K_X)^3} \int_0^\tau \frac{D^2}{q(u)} du}.$$

What if  $P \in \text{Supp}(N(u))$  for some  $u \in [0, \tau]$ ? In this case, we want to find a  $K > 0$  such that

$$\frac{3}{(-K_X)^3} \int_0^\tau (P(u)^2 \cdot S) \cdot \text{ord}_F(N(u)|_S) du \leq K \cdot A_S(F)$$

for every prime divisor  $F$  over  $X$ . How to do this?

**Example 1.** Suppose that there is  $a \in (0, \tau)$  such that

$$N(u) = \begin{cases} 0 & \text{for } u \in [0, a], \\ (u - a)E & \text{for } u \in [a, \tau], \end{cases}$$

where  $E$  is a prime divisor in  $X$  such that  $E \neq S$ . If  $(S, E|_S)$  is log canonical, then

$$\begin{aligned} \frac{3}{(-K_X)^3} \int_0^\tau (P(u)^2 \cdot S) \cdot \text{ord}_F(N(u)|_S) du \\ \leq \left( \frac{3}{(-K_X)^3} \int_a^\tau (P(u)^2 \cdot S)(u - a) du \right) A_S(F) \end{aligned}$$

for every prime divisor  $F$  over  $S$ . This holds if  $S$  is smooth, and  $E|_S$  is a smooth curve.

#### 4. SECOND FUJITA'S FORMULA

Let us use all assumptions and notations of Section 1. Suppose, in addition, that  $S$  is smooth. Let us show a simple way how to estimate  $\delta_P(X)$  using the second Fujita's formula found in [2]. To do this, fix a smooth curve  $C \subset S$  such that  $P \in C$ . For every  $u \in [0, \tau]$ , write

$$N(u)|_S = d(u)C + N'(u),$$

where  $N'(u)$  is effective  $\mathbb{R}$ -divisor such that  $C \not\subset \text{Supp}(N'(u))$ , and

$$d(u) = \text{ord}_C(N(u)|_S).$$

Then, for every  $u \in [0, \tau]$ , compute

$$t(u) = \sup \left\{ v \in \mathbb{R}_{\geq 0} \mid \text{the divisor } P(u)|_S - vC \text{ is pseudo-effective} \right\}.$$

After this, for every  $v \in [0, t(u)]$ , compute

- $P(u, v)$  = positive part of Zariski decomposition of  $P(u)|_S - vC$ ,
- $N(u, v)$  = negative part of Zariski decomposition of  $P(u)|_S - vC$ .

Finally, compute

$$S(W_{\bullet, \bullet, \bullet}^S; C) = \frac{3}{(-K_X)^3} \int_0^\tau d(u)(P(u, 0))^2 du + \frac{3}{(-K_X)^3} \int_0^\tau \int_0^{t(u)} (P(u, v))^2 dv du$$

compute

$$F_P(W_{\bullet, \bullet, \bullet}^{S, C}) = \frac{6}{(-K_X)^3} \int_0^\tau \int_0^{t(u)} (P(u, v) \cdot C) \cdot \text{ord}_P(N'(u)|_C + N(u, v)|_C) dv du,$$

and compute

$$S(W_{\bullet,\bullet,\bullet}^{S,C}; P) = \frac{3}{(-K_X)^3} \int_0^\tau \int_0^{\tilde{t}(u)} (P(u, v) \cdot C)^2 dvdu + F_P(W_{\bullet,\bullet,\bullet}^{S,C}).$$

Then the second Fujita’s formula derived from Abban–Zhuang theory gives

$$\delta_P(X) \geq \min \left\{ \frac{1}{S_X(S)}, \frac{1}{S(W_{\bullet,\bullet,\bullet}^S; C)}, \frac{1}{S(W_{\bullet,\bullet,\bullet}^{S,C}; P)} \right\}.$$

In many cases, this gives us the desired estimate  $\delta_P(X) > 1$ .

However, if this approach does not work, we can blow up the surface  $S$  and apply a similar formula to the exceptional curve [2]. Namely, let  $f: \tilde{S} \rightarrow S$  be the blow up of the point  $P$ , and let  $F$  be the  $f$ -exceptional curve. Write

$$f^*(N(u)|_S) = \tilde{d}(u)F + \tilde{N}'(u),$$

where  $\tilde{N}'(u)$  is the strict transform on  $\tilde{S}$  of  $N(u)|_S$ , and  $\tilde{d}(u) = \text{mult}_P(N(u)|_S)$ . For  $u \in [0, \tau]$ , compute

$$\tilde{t}(u) = \sup \left\{ v \in \mathbb{R}_{\geq 0} \mid \text{the divisor } f^*(P(u)|_S) - vF \text{ is big} \right\}.$$

Then, for every  $v \in [0, \tilde{t}(u)]$ , compute

- $\tilde{P}(u, v)$  = positive part of Zariski decomposition of  $f^*(P(u)|_S) - vF$ ,
- $\tilde{N}(u, v)$  = negative part of Zariski decomposition of  $f^*(P(u)|_S) - vF$ .

After this, we compute

$$S(W_{\bullet,\bullet,\bullet}^S; F) = \frac{3}{(-K_X)^3} \int_0^\tau \tilde{d}(u)(P(u, 0))^2 du + \frac{3}{(-K_X)^3} \int_0^\tau \int_0^{\tilde{t}(u)} (\tilde{P}(u, v))^2 dvdu.$$

Then, for every point  $O \in F$ , compute

$$F_O(W_{\bullet,\bullet,\bullet}^{\tilde{S},F}) = \frac{6}{(-K_X)^3} \int_0^\tau \int_0^{\tilde{t}(u)} (\tilde{P}(u, v) \cdot F) \cdot \text{ord}_O(\tilde{N}'(u)|_F + \tilde{N}(u, v)|_F) dvdu$$

and

$$S(W_{\bullet,\bullet,\bullet}^{\tilde{S},F}; O) = \frac{3}{(-K_X)^3} \int_0^\tau \int_0^{\tilde{t}(u)} (\tilde{P}(u, v) \cdot F)^2 dvdu + F_O(W_{\bullet,\bullet,\bullet}^{\tilde{S},F}).$$

Then we have

$$\delta_P(X) \geq \min \left\{ \frac{1}{S_X(S)}, \frac{2}{S(W_{\bullet,\bullet,\bullet}^S; F)}, \inf_{O \in F} \frac{1}{S(W_{\bullet,\bullet,\bullet}^{\tilde{S},F}; O)} \right\}.$$

5. ONE APPLICATION

Let  $Y = \mathbb{P}^1 \times \mathbb{P}^1$ , and let  $Z$  be a smooth curve in  $Y$  of degree  $(5, 1)$ . Then, choosing appropriate coordinates  $([u : v], [x : y])$  on the surface  $Y$ , we may assume that the curve  $Z$  is given by

$$u(x^5 + a_1x^4y + a_2x^3y^2 + a_3x^2y^3) = v(y^5 + b_1xy^4 + b_2x^2y^3 + b_3x^3y^2)$$

for some  $a_1, a_2, a_3, b_1, b_2, b_3$ . Consider the embedding  $Y \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^2$  given by

$$([u : v], [x : y]) \mapsto ([u : v], [x^2 : xy : y^2]).$$

Identify  $Y$  and  $Z$  with their images in  $\mathbb{P}^1 \times \mathbb{P}^2$ . Let  $\pi : X \rightarrow \mathbb{P}^1 \times \mathbb{P}^2$  be the blow up of the curve  $Z$ . Then  $X$  is a Fano threefold of degree  $-K_X^3 = 20$ .

Let  $\text{pr}_1 : \mathbb{P}^1 \times \mathbb{P}^2 \rightarrow \mathbb{P}^1$  be the projection to the first factor. Set  $\phi_1 = \text{pr}_1 \circ \pi$ . Then  $\phi_1$  is a fibration into del Pezzo surfaces of degree 4.

**Theorem 1** ([4]). *Suppose that every singular fiber of the fibration  $\phi_1$  has singular points of type  $\mathbb{A}_1$ . Then  $X$  is  $K$ -stable.*

Let us briefly explain how to prove this result. Suppose that every singular fiber of the del Pezzo fibration  $\phi_1$  has singular points of type  $\mathbb{A}_1$ . Fix a point  $P \in X$ . To prove Theorem 1, it is enough to show that  $\delta_P(X) > 1$ . Let  $\tilde{Y}$  be the strict transform on  $X$  of the surface  $Y$ .

**Lemma 1** ([2, Lemma 5.68]). Suppose that  $P \in \tilde{Y}$ . Then  $\delta_P(X) > 1$ .

*Proof.* Apply results described in Section 4 with  $S = \tilde{Y}$  and  $C$  being one of the rulings of the smooth surface  $\tilde{Y} \simeq Y \simeq \mathbb{P}^1 \times \mathbb{P}^1$  that passes through  $P$ . □

Thus, we may assume that  $P \notin \tilde{Y}$ . Let us apply results of Section 1 with  $S$  being the fiber of  $\phi_1$  that contains  $P$ . Then  $\tau = 2$ . Moreover, we compute

$$P(u) = \begin{cases} -K_X - uS & \text{for } u \in [0, 1], \\ -K_X - uS - (u - 1)\tilde{Y} & \text{for } u \in [1, 2] \end{cases}$$

and

$$N(u) = \begin{cases} 0 & \text{for } u \in [0, 1], \\ (u - 1)\tilde{Y} & \text{for } u \in [1, 2]. \end{cases}$$

This gives  $S_X(S) = \frac{1}{20} \int_0^2 P(u)^3 du = \frac{69}{80} < 1$ .



Set  $Z = \tilde{Y}|_S$ . For every prime divisor  $F$  over  $S$  whose support on  $S$  contains the point  $P$ , we have

$$\begin{aligned} S(W_{\bullet, \bullet}^S; F) &= \frac{3}{20} \left( \int_0^1 \int_0^\infty \text{vol}(-K_S - vF) dv du \right. \\ &\quad \left. + \int_1^2 \int_0^\infty \text{vol}(-K_S - (u-1)Z - vF) dv du \right) \\ &\leq \frac{3}{20} \left( \int_0^\infty \text{vol}(-K_S - vF) dv + \int_0^\infty \text{vol}(-K_S - vF) dv \right) = \\ &= \frac{3}{10} \left( \int_0^\infty \text{vol}(-K_S - vF) dv \right) = \frac{6}{5} \left( \frac{1}{4} \int_0^\infty \text{vol}(-K_S - vF) dv \right) \\ &= \frac{6}{5} S_S(F) \leq \frac{6}{5} \cdot \frac{A_S(F)}{\delta_P(S)} \end{aligned}$$

Thus, if  $\delta_P(S) > \frac{6}{5}$ , then  $\delta_P(X) > 1$ . If  $S$  is smooth, then  $\delta_P(S) > \frac{6}{5}$  [2], which implies that  $\delta_P(X) > 1$ . Thus, we may assume that  $S$  is singular.

As in Section 3, set  $D = P(u)|_S$  for every  $u \in [0, 2]$ . Then, applying results described in Section 2 to the minimal resolution of singularities of the surface  $S$ , and arguing as in [5], we see that  $\delta_P(S, D) \geq q(u)$  for

$$q(u) = \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} & \text{for } u \in [1, a], \\ \frac{15 - 3u^2}{11 - u^3} & \text{for } u \in [a, 2], \end{cases}$$

where  $a \in [1, 2]$  such that  $3a^3 - 9a^2 + 3a + 5 = 0$ . Thus, for every prime divisor  $F$  over the del Pezzo surface  $S$  whose support on  $S$  contains  $P$ , we have

$$\begin{aligned} S(W_{\bullet, \bullet}^S; F) &= \frac{3}{20} \int_0^2 \int_0^\infty \text{vol}(D - vF) dv du \\ &\leq \frac{3}{20} \cdot \frac{4A_S(F)}{\delta_P(S)} + \left( \frac{3}{20} \int_1^2 \frac{(5 - u^2)}{q(u)} du \right) A_S(F) \leq \frac{99}{100} A_S(F) \end{aligned}$$

which implies that  $\delta_P(S, W_{\bullet, \bullet}^S) \geq \frac{100}{99}$ . Then  $\delta_P(X) > 1$  by the inequality described in Section 1. This shows that  $X$  is K-stable.

## REFERENCES

- [1] H. Abban, Z. Zhuang, *K-stability of Fano varieties via admissible flags*, Forum of Mathematics, Pi **10** (2022), e15.
- [2] C. Araujo, A.-M. Castravet, I. Cheltsov, K. Fujita, A.-S. Kaloghiros, J. Martinez-Garcia, C. Shramov, H. Süß, N. Viswanathan, *The Calabi problem for Fano threefolds*, LMS Lecture Notes in Mathematics **485**, Cambridge University Press, 2023.
- [3] I. Cheltsov, K. Fujita, T. Kishimoto, T. Okada, *K-stable divisors in  $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^2$  of degree  $(1, 1, 2)$* , Nagoya Mathematical Journal **251** (2023), 686–714.
- [4] E. Denisova, *K-stability of Fano 3-folds of Picard rank 3 and degree 20*, to appear in Annali dell' Università di Ferrara.
- [5] E. Denisova,  *$\delta$ -invariant of Du Val del Pezzo surfaces of degree  $\geq 4$* , preprint, ArXiv:2304.11412.
- [6] K. Fujita, *A valuative criterion for uniform K-stability of  $\mathbb{Q}$ -Fano varieties*, J. Reine Angew. Math. **751** (2019), 309–338.
- [7] C. Li, *K-semistability is equivariant volume minimization*, Duke Math. Jour. **166** (2017), 3147–3218.

## Minicourse on Mirror Symmetry for Fano varieties

GIULIA GUGIATTI AND ANDREA PETRACCI

Mirror Symmetry for Fano varieties takes the shape of the Fano/Landau–Ginzburg (LG) correspondence: conjecturally, the mirror of a Fano orbifold of dimension  $n$  is an  $n$ -dimensional LG model  $(Y, w)$ , i.e. a pair formed by a non compact manifold  $Y$  and a complex-valued function  $w$  on  $Y$  called superpotential. Broadly speaking, the correspondence interchanges the symplectic/complex geometry of  $X$  with the complex/symplectic geometry of the critical points of  $w$ .

In this series of lectures we discussed certain aspects of the Fano/LG correspondence.

**Lecture 1 (AP).** In this lecture we provided an introduction to the original formulation of the Fano/LG correspondence, due to Batyrev, Givental, Hori, Vafa, and others. This formulation predicts an identity between two cohomological invariants: the regularised quantum period of  $X$ , which is a generating function for certain Gromov–Witten invariants of  $X$ , and a period of  $(Y, w)$ , encoding information on the variation of cohomology of the smooth fibres of  $w$ .

The lecture primarily focused on a series of conjectures, first laid out by Coates, Corti, Galkin, Golyshev, and Kasprzyk [17, 18, 9], according to which  $\mathbb{Q}$ -Gorenstein (qG) deformation families of Fano orbifolds of dimension  $n$  should be mirror (in the sense of the above-mentioned formulation) to mutation equivalence classes of maximally mutable Laurent polynomials [12] in  $n$  variables. This conjectural framework agrees with classical mirror constructions [15, 22] and finds theoretical ground in the intrinsic mirror symmetry program by Gross–Siebert [19]. However, it notably excludes Fano varieties with empty anticanonical linear system [14, Remark 2.7].

Part of the lecture was devoted to define the quantum period of a Fano orbifold, and to sketch some of the available techniques for its computation [16, 11, 13, 10,

6, 8, 26, 28]. The lecture was supported by some running examples in dimension two.

**Lecture 2 (GG).** The lecture was structured in two parts.

In the first part, we expanded on the complex geometry of LG models. We revisited the notion of periods of a LG model and its specialisation to that of classical period of a Laurent polynomial [9, 29]. We discussed an equivalent formulation of the Fano/LG correspondence in terms of polynomial ordinary differential equations/complex local systems underlying a one-dimensional variation of (pure) Hodge Structure. Throughout the exposition, we revisited the 2-dimensional examples encountered in the first lecture and computed the relative periods and differential operators.

The focus of the second part of the lecture was on Fano varieties with empty anticanonical linear system, currently lying beyond the context of any systematic mirror construction. The simplest instance of such varieties is the series of log del Pezzo surfaces  $X_{8k+4} \subset \mathbb{P}(2, 2k+1, 2k+1, 4k+1)$ ,  $k \in \mathbb{N}_{>0}$ , first studied by Johnson and Kollár [24]. The work of GG with Corti [14] builds the only known mirror series to this series. In the lecture we sketched the main ideas of our mirror construction, which builds upon the hypergeometric nature of the regularised quantum period of the surfaces and the motivic origin of hypergeometric functions [7, 31].

**Lecture 3 (AP).** Since qG-deformations of del Pezzo surfaces are unobstructed [20], it is clear what the general qG-deformation of a toric del Pezzo surface is. In dimension  $\geq 3$  this is no longer true, and so one needs to study deformation theory of Fano varieties more carefully. This has also applications to the moduli theory of Fano varieties.

In this lecture we presented some features of deformation theory of toric Fano varieties. In particular, we showed that there exist toric Fano 3-folds which deform to different smooth Fano 3-folds [23, 30, 25]. This is reflected on the mirror side by the fact that on the same polytope there might be different maximally mutable Laurent polynomials. These examples and results on the deformation theory of toric Fano 3-folds builds on the deformation theory of toric singularities studied by Altmann [4, 3, 2, 1].

**Lecture 4 (GG).** This lecture approached the Fano/LG correspondence from the point of view of Kontsevich's Homological Mirror Symmetry (HMS) conjecture [27]. One formulation of HMS predicts an equivalence between the bounded derived category of coherent sheaves of  $X$  and the analog of the Fukaya category for a symplectic fibration, namely the bounded derived category of Lagrangian vanishing cycles of  $(Y, w)$ . A rigorous definition of this category was proposed by Seidel [32] in the case where  $w$  is a symplectic Lefschetz fibration.

After briefly sketching the construction of the category of Lagrangian vanishing cycles, we focused on the implication of the above formulation of HMS at the numerical level, i.e. at the level of the numerical Grothendieck groups of the two categories [33, 21]. For smooth del Pezzo surfaces we reviewed the homological mirror construction given in [5], and we explained how to recover the outputs of

this construction from the Laurent polynomial mirrors to the surfaces. A special focus was placed on the case of smooth del Pezzo surfaces of degree two. These surfaces appear as the degenerate case ( $k = 0$ ) of the series of log del Pezzo surfaces constructed by Johnson–Kollár [24], for which HMS has not yet been established.

#### REFERENCES

- [1] Klaus Altmann. Computation of the vector space  $T^1$  for affine toric varieties. *J. Pure Appl. Algebra*, 95(3):239–259, 1994.
- [2] Klaus Altmann. Minkowski sums and homogeneous deformations of toric varieties. *Tohoku Math. J. (2)*, 47(2):151–184, 1995.
- [3] Klaus Altmann. Infinitesimal deformations and obstructions for toric singularities. *J. Pure Appl. Algebra*, 119(3):211–235, 1997.
- [4] Klaus Altmann. The versal deformation of an isolated toric Gorenstein singularity. *Invent. Math.*, 128(3):443–479, 1997.
- [5] Denis Auroux, Ludmil Katzarkov, and Dmitri Orlov. Mirror symmetry for del Pezzo surfaces: vanishing cycles and coherent sheaves. *Invent. Math.*, 166(3):537–582, 2006.
- [6] Aaron Bertram, Ionuț Ciocan-Fontanine, and Bumsig Kim. Gromov-Witten invariants for abelian and nonabelian quotients. *J. Algebraic Geom.*, 17(2):275–294, 2008.
- [7] Frits Beukers, Henri Cohen, and Anton Mellit. Finite hypergeometric functions. *Pure Appl. Math. Q.*, 11(4):559–589, 2015.
- [8] Ionuț Ciocan-Fontanine, Bumsig Kim, and Claude Sabbah. The abelian/nonabelian correspondence and Frobenius manifolds. *Invent. Math.*, 171(2):301–343, 2008.
- [9] Tom Coates, Alessio Corti, Sergey Galkin, Vasily Golyshev, and Alexander Kasprzyk. Mirror symmetry and Fano manifolds. In *European Congress of Mathematics*, pages 285–300. (2013).
- [10] Tom Coates, Alessio Corti, Hiroshi Iritani, and Hsian-Hua Tseng. A mirror theorem for toric stacks. *Compos. Math.*, 151(10):1878–1912, 2015.
- [11] Tom Coates and Alexander Givental. Quantum Riemann-Roch, Lefschetz and Serre. *Ann. of Math. (2)*, 165(1):15–53, 2007.
- [12] Tom Coates, Alexander Kasprzyk, Giuseppe Pitton, and Ketil Tveiten. Maximally mutable laurent polynomials. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 477, 10 2021.
- [13] Tom Coates, Yuan-Pin Lee, Alessio Corti, and Hsian-Hua Tseng. The quantum orbifold cohomology of weighted projective spaces. *Acta Math.*, 202(2):139–193, 2009.
- [14] Alessio Corti and Giulia Gugiatti. Hyperelliptic integrals and mirrors of the Johnson-Kollár del Pezzo surfaces. *Trans. Amer. Math. Soc.*, 374(12):8603–8637, 2021.
- [15] Alexander Givental. A mirror theorem for toric complete intersections. In *Topological field theory, primitive forms and related topics (Kyoto, 1996)*, volume 160 of *Progr. Math.*, pages 141–175. Birkhäuser Boston, Boston, MA, 1998.
- [16] Alexander B. Givental. Equivariant Gromov-Witten invariants. *Internat. Math. Res. Notices*, (13):613–663, 1996.
- [17] Vasily Golyshev. Classification problems and mirror duality. In *Surveys in geometry and number theory: reports on contemporary Russian mathematics*, volume 338 of *London Math. Soc. Lecture Note Ser.*, pages 88–121. Cambridge Univ. Press, Cambridge, 2007.
- [18] Vasily Golyshev. Spectra and strains. *arXiv e-prints*, page arXiv:0801.0432, January 2008.
- [19] Mark Gross and Bernd Siebert. The canonical wall structure and intrinsic mirror symmetry. *Invent. Math.*, 229(3):1101–1202, 2022.
- [20] Paul Hacking and Yuri Prokhorov. Smoothable del Pezzo surfaces with quotient singularities. *Compos. Math.*, 146(1):169–192, 2010.
- [21] Andrew Harder and Alan Thompson. Pseudolattices, del Pezzo surfaces, and Lefschetz fibrations. *Trans. Amer. Math. Soc.*, 373(3):2071–2104, 2020.

- [22] Kentaro Hori and Cumrun Vafa. Mirror Symmetry. *arXiv e-prints*, pages hep-th/0002222, 2000.
- [23] Priska Jahnke and Ivo Radloff. Terminal Fano threefolds and their smoothings. *Math. Z.*, 269(3-4):1129–1136, 2011.
- [24] J. M. Johnson and J. Kollár. Kähler-Einstein metrics on log del Pezzo surfaces in weighted projective 3-spaces. *Ann. Inst. Fourier (Grenoble)*, 51(1):69–79, 2001.
- [25] Anne-Sophie Kaloghiros and Andrea Petracci. On toric geometry and K-stability of Fano varieties. *Trans. Amer. Math. Soc. Ser. B*, 8:548–577, 2021.
- [26] Bumsig Kim. Quantum hyperplane section theorem for homogeneous spaces. *Acta Math.*, 183(1):71–99, 1999.
- [27] Maxim Kontsevich. Homological algebra of mirror symmetry. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994)*, pages 120–139. Birkhäuser, Basel, 1995.
- [28] Y.-P. Lee. Quantum Lefschetz hyperplane theorem. *Invent. Math.*, 145(1):121–149, 2001.
- [29] Travis Mandel. Fano mirror periods from the Frobenius structure conjecture, 2019.
- [30] Andrea Petracci. An example of mirror symmetry for Fano threefolds. In *Birational geometry and moduli spaces*, volume 39 of *Springer INdAM Ser.*, pages 173–188. Springer, Cham, 2020.
- [31] David P. Roberts and Fernando Rodriguez Villegas. Hypergeometric motives. *Notices Amer. Math. Soc.*, 69(6):914–929, 2022.
- [32] Paul Seidel. *Fukaya categories and Picard-Lefschetz theory*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [33] S. Sugiyama. On the Fukaya-Seidel categories of surface Lefschetz fibrations. arXiv:1607.02263.

## On the normalised volume of toric singularities

HENDRIK SÜSS

(joint work with Joaquín Moraga)

The notion of normalised volume for log terminal singularities was introduced by Chi Li in [2]. In the special case of the anticanonical cone over a K-semistable Fano variety it coincides with the usual anticanonical volume (aka Fano degree) of the Fano variety.

In general the normalised volume is hard to calculate, but in the case of Gorenstein toric singularities we obtain a nice formula in terms of convex geometry. Given a full-dimensional polytope  $P \subset \mathbb{R}^{d-1}$ , we consider the Gorenstein toric variety  $X = \text{Spec}(k[\sigma^\vee \cap \mathbb{Z}^d])$  corresponding to the cone  $\sigma = \mathbb{R}_{\geq 0} \cdot (P \times \{1\}) \subset \mathbb{R}^d$ . We choose  $x \in X$  to be the (unique) torus fixed point. Then by [3] we have

$$\widehat{\text{vol}}(X, x) = \min \{ \text{vol}((P - v)^*) \mid v \in \text{int}(P) \}.$$

Here,  $(P - v)^*$  denotes the polar dual of the polytope  $P$  after translation by  $-v$ . In convex geometry the (unique) point  $v \in \text{int}(P)$  where the minimum is attained, is known as the *Santaló point* of  $P$ . Moreover, for a convex body  $P$  the product

$$M(P) = \text{vol}(P) \cdot \min \{ \text{vol}((P - v)^*) \mid v \in \text{int}(P) \}$$

is known as the *Mahler volume* of  $P$ , which is an affine invariant of the polytope. The Mahler volume appears in two remarkable inequalities. The first one is known as the *Blaschke-Santaló inequality* and states that the Mahler volume of convex

bodies of fixed dimension is maximised by the unit ball. In [3] we utilised this inequality to show that for a fixed dimension and fixed  $\epsilon > 0$  there are only finitely many toric singularities with normalised volume being at least  $\epsilon$ .

A second, but to this point only conjectural, inequality states that for  $\dim(P) = d - 1$  one has

$$M(P) \geq d^d,$$

where the equality is achieved for simplices. The latter inequality is known as the *non-symmetric Mahler conjecture*. It is natural to ask whether there is a reasonable interpretation of this inequality in terms of algebraic geometry. Hence, we are looking for an interpretation of the volume of  $P$  in terms of algebraic geometry. But this volume is known to coincide with the Euler characteristic of a crepant resolution  $\tilde{X}$  of  $X$  (at least if such a resolution exists). However, even for non-abelian quotient singularities does the corresponding inequality

$$(1) \quad \widehat{\text{vol}}(X, x)\chi(\tilde{X}) \geq d^d$$

not longer hold. In [1] Gulotta suggests to replace  $\chi(\tilde{X})$  by another quantity associated to *non-commutative crepant resolutions* of  $X$  in the sense of [4]. With this adjustment the corresponding inequality of the type (1) does again hold (with equality) for finite quotient singularities and we can also verify it for a range of other examples, such as anticanonical cones over del Pezzo surfaces and  $cA_n$  singularities.

## REFERENCES

- [1] D. R. Gulotta, *Properly ordered dimers, R-charges, and an efficient inverse algorithm*. J. High Energy Phys. 2008, No. 10, Paper No. 014, 31 p.
- [2] C. Li, *Minimizing normalized volumes of valuations* Math. Z. **289** (2018), No. 1–2, 491–513
- [3] J. Moraga and H. Süß, *Bounding toric singularities with normalized volume*, arXiv:2111.01738 [math.AG]
- [4] M. van den Bergh, *Non-commutative crepant resolutions*. in: The legacy of Niels Henrik Abel. Papers from the Abel bicentennial conference, University of Oslo, Oslo, Norway, June 3–8, 2002. Berlin: Springer. 749–770

## Higher Fano Manifolds

ANA-MARIA CASTRAVET

(joint work with Carolina Araujo, Roya Beheshti, Kelly Jabbusch, Enrica Mazzon, Svetlana Makarova, Will Reynolds, Libby Taylor, Nivedita Viswanathan)

A Fano manifold  $X$  is a complex projective manifold with ample first Chern class  $c_1(T_X)$ . This condition has far reaching geometric implications. For instance, any Fano manifold is rationally connected, i.e., there are rational curves connecting any two points [6, 11]. A celebrated result of Graber, Harris and Starr states that proper families of rationally connected complex projective manifolds over smooth curves always admit sections [10]. This generalizes Tsen’s theorem in the case of function fields of curves. A theorem of Tsen and Lang states that a family

$\pi : \mathcal{X} \rightarrow B$  of degree  $d$  hypersurfaces over a  $k$ -dimensional base  $B$  admit (rational) sections if  $d^k \leq n$ . For hypersurfaces of degree  $d$  in  $\mathbb{P}^n$ , being Fano or rationally connected is equivalent to the numerical condition  $d \leq n$ . Hence, for  $k = 1$ , the result of Graber, Harris and Starr replaces the condition of being a hypersurface of degree  $d \leq n$  with the condition of being rationally connected.

There has been quite some effort towards finding suitable geometric conditions on the fibers of  $\pi : \mathcal{X} \rightarrow B$  that generalize the Tsen-Lang theorem for higher dimensional bases  $B$ . More generally one can consider the following problem: Find intrinsic (geometric) conditions  $\mathcal{F}_k$  such that

- Hypersurfaces of degree  $d$  in  $\mathbb{P}^n$  satisfy  $\mathcal{F}_k$  iff  $d^k \leq n$ ,
- Projective manifolds satisfying  $\mathcal{F}_k$  are covered by rational  $k$ -folds,
- The Tsen–Lang Theorem holds for a family  $\pi : \mathcal{X} \rightarrow B$  over a  $k$ -dimensional base  $B$  if the fibers of  $\pi$  satisfy  $\mathcal{F}_k$  (modulo the Brauer obstruction).

For  $k = 1$ , the condition  $\mathcal{F}_1$  can hence be taken as either the condition of being Fano or the condition of being rationally connected.

In a series of papers [7, 8, 9], de Jong and Starr asked if for  $k = 2$  the condition  $\mathcal{F}_2$  can be taken to be *rationally simply connected*, a technical condition taking inspiration from topology which requires (at the very least) that a suitable irreducible component of the space of rational curves through two general points is itself rationally connected. De Jong, He and Starr proved that rational homogeneous spaces and hypersurfaces of degree  $d$  in  $\mathbb{P}^n$  with  $d^2 \leq n$  are rationally simply connected and that they satisfy the Tsen-Lang theorem. However, the condition of being rationally simply connected is hard to verify in practice and it is desirable to have natural geometric conditions that imply it. In this context, 2-manifolds were introduced by de Jong and Starr in [8, 9]. More generally, consider the following:

**Definition.** [9, 1] A complex projective manifold  $X$  is *k-Fano* if  $X$  is Fano and the  $i$ -th graded piece of the Chern character  $ch_i(T_X) \in H^{2i}(X, \mathbb{Z})$  is positive for all  $i \in \{1, \dots, k\}$ , i.e.,  $ch_i(T_X) \cdot Z > 0$  for all  $Z \subset X$  with  $\dim(Z) = i$ .

The following may be easily verified:

- (1) The  $n$ -dimensional projective space  $\mathbb{P}^n$  is  $n$ -Fano.
- (2) Hypersurfaces of degree  $d$  in  $\mathbb{P}^n$  are  $k$ -Fano if and only if  $d^k \leq n$ . In particular, hypersurfaces of degree  $d$  in  $\mathbb{P}^n$  are 2-Fano if and only if  $d^2 \leq n$ .

**Question.** [1] *Can one take the condition  $\mathcal{F}_k$  to be “ $X$  is  $k$ -Fano”?*

Some evidence towards an affirmative answer has been given in [9, 1]. Roughly speaking, we know that 2-Fano manifolds are covered by rational surfaces and 3-Fano manifolds are covered by rational threefolds. The approach taken in [1] is to consider the implications of the  $k$ -Fano condition on the *spaces of minimal rational curves*  $H_x$  through a general point  $x \in X$ . For example, one may compute all the Chern characters of  $H_x$  in terms of the Chern characters of  $X$  and a canonical polarization on  $H_x$  coming from the map  $H_x \rightarrow \mathbb{P}(T_x(X))$  that associates to a rational curve through  $x$  its tangent line at  $x$ . One can prove that there is an inductive structure: If  $X$  is 2-Fano, then  $H_x$  is Fano [9, 1], and similarly, if  $X$  is

3-Fano, then  $H_x$  is 2-Fano [1] (under certain technical assumptions). Results in the same spirit have been obtained in the case when  $X$  is  $k$ -Fano by Nagaoka [12] and Suzuki [13].

There are however few examples of higher Fano manifolds. What is known:

- (1) Complete intersections  $X_{d_1, \dots, d_c} \subset \mathbb{P}(a_0, \dots, a_n)$  are  $k$ -Fano iff  $\sum d_i^k < \sum a_i^k$ .
- (2) The only 2-Fano surface is  $\mathbb{P}^2$ .
- (3) The only 2-Fano threefolds are  $\mathbb{P}^3$  and the quadric hypersurface  $Q \subset \mathbb{P}^4$  [1].
- (4) The only 3-Fano threefold is  $\mathbb{P}^3$  [1].
- (5) The only known examples of 3-Fano manifolds are complete intersections

$$X_{d_1, \dots, d_c} \subset \mathbb{P}(a_0, \dots, a_n), \quad \sum d_i^3 < \sum a_i^3.$$

(6) A classification of 2-Fano manifolds  $X$  of index  $\geq \dim(X) - 2$  was given in [3]: they are either complete intersections  $X_{d_1, \dots, d_c}$  in weighted projective spaces  $\mathbb{P}(a_0, \dots, a_n)$  or certain rational homogeneous spaces of Picard number  $\rho = 1$  or certain complete intersections in them [3]. A classification of 2-Fano manifolds among rational homogeneous spaces with  $\rho = 1$  was also given in [3]:

**Theorem 1.** [3] *Classification of 2-Fano rational homogeneous spaces with  $\rho = 1$ :*

- $A_n/P^k, k = 1, \frac{n}{2}, \frac{n+1}{2}, n$
- $B_n/P^k, k = 1, \frac{2n-1}{3}, n$
- $C_n/P^k, k = 1, \frac{2n+2}{3}, n$
- $D_n/P^k, k = 1, \frac{2n+2}{3}, n-1, n$
- $E_6/P^1, E_6/P^2, E_6/P^3, E_7/P^1, E_7/P^2, E_7/P^7, E_8/P^1, E_8/P^2, E_8/P^8$
- $F_4/P^4$
- $G_2/P^1, G_2/P^2$

(Here we use the Bourbaki labeling of vertices in a Dynkin diagram.)

- (7) Products  $X \times Y$  with  $\dim(X), \dim(Y) \geq 1$  are not 2-Fano.
- (8) Projectivizations  $\mathbb{P}(E)$  of vector bundles  $E$  of rank  $\geq 2$  over a positive dimensional base are not 2-Fano.
- (9) No blow-up of a projective manifold along a smooth subvariety of codimension at least 2 is known to be 2-Fano. One is hence lead to the following:

**Question.** *Do all 2-Fano manifolds have Picard number one?*

**Problem.** Find examples of 3-Fano manifolds other than complete intersections in weighted projective spaces.

**Conjecture.** [3] *Let  $X$  be a  $k$ -Fano manifold of dimension  $n$ . If  $k \geq \lceil \log_2(n+1) \rceil$ , then  $X \cong \mathbb{P}^n$ .*

We now concentrate on the **toric** case:

**Conjecture.** *If  $X$  a toric 2-Fano manifold, then  $X \cong \mathbb{P}^n$ .*



Assume  $X$  a smooth projective toric variety with lattice  $N$  and fan  $\Sigma$ . We denote by  $G(\Sigma) \subset N$  set of primitive vectors generating the rays of  $\Sigma$ .

**Definition.** A set  $P = \{v_0, \dots, v_r\} \subseteq G(\Sigma)$  is called a *primitive collection* if

- $\langle v_0, \dots, v_r \rangle \notin \Sigma$ , but
- $\langle v_0, \dots, \hat{v}_i, \dots, v_r \rangle \in \Sigma$  for all  $i = 0, \dots, r$

If  $\sigma(P) = \langle w_1, \dots, w_s \rangle$  is the smallest cone in  $\Sigma$  containing  $v_0 + \dots + v_r$ , then  $v_0 + \dots + v_r = \mu_1 w_1 + \dots + \mu_s w_s$ , for some  $\mu_1, \dots, \mu_s \in \mathbb{Z}_{>0}$ . We say that the primitive collection  $P = \{v_0, \dots, v_r\}$  is *centered* if  $\sigma(P) = 0$ , i.e.,

$$v_0 + \dots + v_r = 0.$$

Furthermore, as relations between the primitive vectors in  $G(\Sigma)$  correspond to numerical equivalence classes of 1-cycles on  $X$ , via this correspondence primitive relations correspond to effective 1-cycle classes that generate the Mori cone of  $X$ .

A theorem of Batyrev [5] asserts that centered primitive relations always exist. In [4] it is proved that for every centered primitive relation  $v_0 + \dots + v_r = 0$  one may associate a torus invariant open set  $U \subseteq X$  which comes with a  $\mathbb{P}^r$ -bundle structure  $\pi : U \rightarrow W$  and the centered primitive relation corresponds to the classes of lines in the fibers of  $\pi$ . A theorem of Chen-Fu-Hwang states that all minimal covering families of rational curves on  $X$  may be described in this way.

**Definition.** [4] The minimal  $\mathbb{P}$ -dimension  $m(X)$  is the smallest  $r > 0$  such that  $\Sigma$  has a centered primitive collection  $\{v_0, \dots, v_r\}$ .

If  $X$  is a toric projective manifold of dimension  $n$ , it is not difficult to see that  $m(X) = n$  implies that  $X \cong \mathbb{P}^n$ . Furthermore, if  $X$  is Fano and  $m(X) = n - 1$ , one can prove that  $X$  is the blow-up of  $\mathbb{P}^n$  along a linear subspace of codimension 2 (which is not 2-Fano). We prove:

**Theorem 2.** [4] *Assume  $X$  is a Fano toric projective manifold of dimension  $n$ .*

- (1) *If  $m(X) = n - 2$ , then  $\rho \geq 3$  and  $X$  belong to 8 explicit isomorphism classes. In particular  $X$  is not 2-Fano.*
- (2) *If  $m(X) = 1$ , then  $X$  is not 2-Fano.*

One can see from the classification of Fano toric manifolds in small dimensions that the toric Fano varieties  $X$  with  $m(X) = 1$  form the largest class:

dim( $X$ )	# Fanos	$m=1$	$m=2$	$m=3$	$m=4$	$m=5$	$m=6$
4	124	107	15	1	1		
5	866	744	112	8	1	1	
6	7622	6333	1174	105	8	1	1

TABLE 1. The minimal  $\mathbb{P}$ -dimension of toric Fano manifolds of low dimension.

## REFERENCES

- [1] C. Araujo and A.-M. Castravet, *Polarized minimal families of rational curves and higher Fano manifolds*, Amer. J. Math. 134.1 (2012), 87–107.
- [2] C. Araujo and A.-M. Castravet, *Classification of 2-Fano manifolds with high index*. A celebration of algebraic geometry. Vol. 18. Clay Math. Proc. Amer. Math. Soc., Providence, RI, 2013, 1–36.
- [3] C. Araujo, R. Beheshti, A.-M. Castravet, K. Jabbusch, S. Makarova, E. Mazzon, L. Taylor and N. Viswanathan, *Higher Fano manifolds*. Rev. Un. Mat. Argentina 64.1 (2022), pp. 103–125.
- [4] C. Araujo, R. Beheshti, A.-M. Castravet, K. Jabbusch, S. Makarova, E. Mazzon and N. Viswanathan (with an appendix by W. Reynolds), *The minimal projective bundle dimension and toric 2-Fano manifolds*, to appear in Trans. Amer. Math. Soc. arXiv:2301.00883.
- [5] V. V. Batyrev. *On the classification of smooth projective toric varieties*. Tohoku Math. J. (2) 43.4 (1991), 569–585
- [6] F. Campana, *Connexité rationnelle des variétés de Fano*. Ann. Sci. Ecole Norm. Sup. (4) 25.5 (1992), 539–545.
- [7] A. J. de Jong, X. He and J. M. Starr, *Families of rationally simply connected varieties over surfaces and torsors for semisimple groups*. Publ. Math. Inst. Hautes Etudes Sci. 114 (2011), 1–85.
- [8] A. J. de Jong and J. M. Starr, *A note on Fano manifolds whose second Chern character is positive*. arXiv:math/0602644. 2006.
- [9] A. J. de Jong and J. M. Starr, *Higher Fano manifolds and rational surfaces*. Duke Math. J. 139.1 (2007), 173–183
- [10] T. Graber, J. Harris and J. Starr, *Families of rationally connected varieties*. J. Amer. Math. Soc. 16.1 (2003), 57–67.
- [11] J. Kollár, Y. Miyaoka and S. Mori, *Rational connectedness and boundedness of Fano manifolds*. J. Diff. Geom. 36.3 (1992), 765–779.
- [12] T. Nagaoka, *On a sufficient condition for a Fano manifold to be covered by rational  $N$ -folds*. J. Pure Appl. Algebra 223.11 (2019), 4677–4688.
- [13] T. Suzuki, *Higher order minimal families of rational curves and Fano manifolds with nef Chern characters*. J. Math. Soc. Japan 73.3 (2021), 949–964.

On  $K$ -moduli spaces of Fano 3-folds

ANNE-SOPHIE KALOGHIROS

The notion of  $K$ -polystability was introduced to characterise the existence of Kähler–Einstein metrics on Fano manifolds. More precisely, the Yau–Tian–Donaldson conjecture (now a theorem, due to Chen–Donaldson and Sun) states that a Fano manifold is Kähler–Einstein precisely when it is  $K$ -polystable.

Recent advances in the theory of  $K$ -stability have shown that this notion also allows one to construct moduli spaces for Fano varieties. More precisely:

**Theorem 1.** [1] *There is a projective good moduli space  $M_{n,V}^{Kps}$  whose points parametrise  $K$ -polystable  $\mathbb{Q}$ -Fano varieties of dimension  $n$  and volume  $V$ .*

In each dimension, there are finitely many families of smooth manifolds, which have been classified in dimension up to 3. There are 10 families of smooth del Pezzo surfaces and 105 families of smooth Fano 3-folds. For each of these families, we can ask the following questions:

- (A) Is the general member of the family K-polystable? (In other words, is the associated component of  $M_{n,V}^{Kps}$  non-empty?)
- (B) Is every (smooth) member of the family K-polystable?
- (C) What is the compactification of the associated component of  $M_{n,V}^{Kps}$ ? In particular, what are the K-polystable limits of Fano manifolds in the family?

In dimension 2, (A) and (B) were answered by Tian [2] and (C) was answered in degree 4 by Mabuchi and Mukai and in general by Odaka-Spotti-Sun [3].

Following Fujita and Li’s valuative criterion for K-polystability [6, 7], a purely algebraic theory of (K-poly)stability was formulated, and this has led to much progress in recent years. Notably, Abban and Zhuang [4] developed techniques to determine K-stability using flags and Zhuang showed how to exploit symmetries effectively [5]; these have yielded many results in explicit K-stability of Fano 3-folds. We now know:

**Theorem 2.** [9] *Let  $X$  be the general member of one of the 105 deformation families of Fano 3-folds. Then one of:*

- $X$  belongs to family  $MM_{2-26}$ , or
- $X$  belongs to one of the 26 deformation families of K-divisionally unstable Fano 3-folds classified by Fujita [8], or
- $X$  is K-polystable

This answers (A) above in dimension 3; 78 families have smooth K-polystable members, and in some cases the families contain both K-polystable and non-K-polystable smooth Fano 3-folds. The answer to (B) is known for 58 out of 78 families with K-polystable members.

Relatively few known examples of K-moduli spaces of Fano 3-folds are known: Liu and Xu have shown that the K-moduli space of cubic threefolds coincides with the GIT moduli space [10], and a number of recent works have considered specific families.

We could also investigate specifically K-moduli spaces of small dimension. 44 of the 105 deformation families of Fano 3-folds have 0-dimensional moduli, and out of these, 21 yield a non-empty component of the associated K-moduli  $M_{3,V}^{Kps}$ . There are 8 families with 1-dimensional moduli, and 6 of these yield a non-empty component of the associated K-moduli  $M_{3,V}^{Kps}$ . My collaborators and I show:

**Theorem 3.** [11] *All one dimensional components of  $M_{3,V}^{Kps}$  associated to families of smoothable Fano 3-folds are isomorphic to  $\mathbb{P}^1$ .*

As a by-product, we obtain:

**Corollary 1.** [11] *All singular K-polystable limits of Fano 3-folds in families  $MM_{2-22}$ ,  $MM_{2-24}$ ,  $MM_{2-25}$ ,  $MM_{3-12}$ ,  $MM_{3-13}$  and  $MM_{4-13}$  are constructed explicitly.*

Finally, I present a construction of the 3-dimensional component of  $M_{3,24}^{Kps}$  associated to the deformation family  $MM_{4-1}$ . Smooth members of  $MM_{4-1}$  are divisors of multidegree  $(1, 1, 1, 1)$  in  $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ .

**Theorem 4** ([12]). *A  $K$ -polystable limit of members of  $\text{MM}_{4-1}$*

- *either an irreducible divisor  $(1, 1, 1, 1) \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$  of the form*

$$(*) \quad X = \{a(x_1x_2x_3x_4 + y_1y_2y_3y_4) + b(x_1x_2y_3y_4 + x_3x_4y_1y_2) + c(x_1x_3y_2y_4 + x_2x_4y_1y_3) + d(x_1x_4y_2y_3 + x_2x_3y_1y_4) = 0\}$$

*for  $(a : b : c : d) \in \mathbb{P}^3$ .*

- *or an irreducible  $(2, 2) \subset \mathbb{P}(1, 1, 2) \times \mathbb{P}(1, 1, 2)$  of the form*

$$(**) \quad X = \{w_1w_2 + \alpha s_1t_1s_2t_2 + \beta(s_1^2s_2^2 + t_1^2t_2^2) + \gamma(s_1^2t_2^2 + t_1^2s_2^2) = 0\}$$

*for  $(\alpha : \beta : \gamma) \in \mathbb{P}^2$ .*

We show:

**Theorem 5.** *The component of  $K$ -moduli space  $\text{M}_{3,24}^{\text{Kps}}$  associated to family  $\text{MM}_{4-1}$  is the blowup of  $\mathbb{P}(1, 3, 4, 6)$  at the smooth point  $\{[2 : 2 : 0 : 0]\}$  with weights  $(1, 2, 3)$ .*

#### REFERENCES

- [1] Y. Liu, C. Xu and Z. Zhuang, *Finite generation for valuations computing stability thresholds and applications to  $K$ -stability*, Ann.of Math.(2), **196** (2022), 507–566.
- [2] G. Tian, *On Calabi’s conjecture for complex surfaces with positive first Chern class*, Invent. Math.**101** (1990), 101–172.
- [3] Y. Odaka, C. Spotti and S. Sun, *Compact moduli spaces of del Pezzo surfaces and Kähler-Einstein metrics*, J. Differ. Geom., **102** (2016), 127–172.
- [4] H. Abban, Z. Zhuang,  *$K$ -stability of Fano varieties via admissible flags*, Forum of Mathematics Pi **10** (2022), 1–43.
- [5] Z. Zhuang, *Optimal destabilizing centers and equivariant  $K$ -stability*, Invent. Math. **226** (2021), 195–223.
- [6] K. Fujita, *A valuative criterion for uniform  $K$ -stability of  $\mathbb{Q}$ -Fano varieties*, J. Reine Angew. Math. **751** (2019), 309–338.
- [7] C. Li,  *$K$ -semistability is equivariant volume minimization*, Duke Math. J. **166** (2017), 3147–3218.
- [8] K. Fujita, *On  $K$ -stability and the volume functions of  $\mathbb{Q}$ -Fano varieties*, Proc. Lond. Math. Soc. **113** (2016), 541–582.
- [9] C. Araujo, A.-M. Castravet, I. Cheltsov, K. Fujita, A.-S. Kaloghiros, J. Martinez-Garcia, C. Shramov, H. Süß, N. Viswanathan, *The Calabi problem for Fano threefolds*, Lecture Notes in Mathematics, Cambridge University Press, 2023.
- [10] Y. Liu, C. Xu,  *$K$ -stability of cubic threefolds*, Duke Math. J. **168** (2019), 2029–2073.
- [11] H. Abban, I. Cheltsov, E. Denisova, E. Etxabbarri-Alberdi, D. Jiao, A.-S. Kaloghiros, J. Martinez-Garcia and T. Papazachariou *One-dimensional components in the  $K$ -moduli of smooth Fano 3-folds*, arXiv: 2309.12518.
- [12] I. Cheltsov, M. Fedorchuk, K. Fujita and A.-S. Kaloghiros,  *$K$ -moduli of four qubits*, in preparation.

## On a problem of Gizatullin

CAROLINA ARAUJO

### 1. OVERVIEW

Our work is motivated by the general problem of describing automorphisms of smooth hypersurfaces in projective spaces. Let  $X_d \subset \mathbb{P}^{n+1}$  be a smooth complex hypersurface of degree  $d$ . Apart from the two exceptional cases  $(n, d) = (1, 3)$  and  $(n, d) = (2, 4)$ , every automorphism of  $X_d$  is the restriction of a linear automorphism of the ambient space  $\mathbb{P}^{n+1}$  ([8], [4]). It is natural to wonder where the automorphisms of  $X_d$  come from in the two exceptional cases.

For  $(n, d) = (1, 3)$ , that is, when  $C = X_3 \subset \mathbb{P}^2$  is a smooth cubic curve, the automorphism group of  $C$  is well known:

$$\mathrm{Aut}(C) = C \rtimes \mathbb{Z}_m, \text{ for some } m \in \{2, 4, 6\},$$

where  $C$  is viewed as an elliptic curve acting on itself by translation. In this case, every automorphism of  $C$  is the restriction of a Cremona transformation of  $\mathbb{P}^2$  ([10, §2]). To see this, first make a linear change of coordinates to write  $C$  in Weierstrass form. Then write down the expression for the translation by a point on the curve in these coordinates, and check that this expression gives a Cremona transformation of  $\mathbb{P}^2$  inducing the given translation on  $C$ . The automorphisms in the finite factor  $\mathbb{Z}_m$  are easily seen to be restrictions of linear automorphisms of  $\mathbb{P}^2$ . The group of Cremona transformations of  $\mathbb{P}^2$  stabilizing the curve  $C$  is called the *decomposition group* of  $C$  and is denoted by  $\mathrm{Dec}(\mathbb{P}^2, C)$ , while the group of Cremona transformations fixing the curve  $C$  pointwise is called the *inertia group* of  $C$  and is denoted by  $\mathrm{In}(\mathbb{P}^2, C)$ . So we have an exact sequence

$$1 \rightarrow \mathrm{In}(\mathbb{P}^2, C) \rightarrow \mathrm{Dec}(\mathbb{P}^2, C) \rightarrow \mathrm{Aut}(C) \rightarrow 1.$$

Generators of the decomposition group  $\mathrm{Dec}(\mathbb{P}^2, C)$  were given in [11], while the inertia group  $\mathrm{In}(\mathbb{P}^2, C)$  was investigated in [3].

For  $(n, d) = (2, 4)$ ,  $S = X_4 \subset \mathbb{P}^3$  is a K3 surface, and  $\mathrm{Aut}(S)$  can be infinite and fairly complicated. The following question is attributed to Gizatullin:

**Problem** (Gizatullin). Which automorphisms of a smooth quartic surface  $S \subset \mathbb{P}^3$  are restrictions of Cremona transformations of  $\mathbb{P}^3$ ?

In [9], Oguiso constructed a smooth quartic surface  $S \subset \mathbb{P}^3$  with Picard rank  $\rho(S) = 2$ ,  $\mathrm{Aut}(S) \cong \mathbb{Z}$  and trivial decomposition group,  $\mathrm{Dec}(\mathbb{P}^3, S) = \{1\}$ . So no nontrivial automorphism of  $S$  is induced by a Cremona transformation of  $\mathbb{P}^3$ . In [10], Oguiso constructed a smooth quartic surface  $S \subset \mathbb{P}^3$  with  $\rho(S) = 3$ ,  $\mathrm{Aut}(S) \cong \mathbb{Z}_2 * \mathbb{Z}_2 * \mathbb{Z}_2$ , and such that every automorphism of  $S$  is the restriction of a Cremona transformation of  $\mathbb{P}^3$ . Inspired by these examples, Oguiso asked whether every automorphism of finite order of a smooth quartic surface is the restriction of a Cremona transformation of  $\mathbb{P}^3$ . This question was negatively answered by Paiva and Queido in [12], with the construction of a smooth quartic surface  $S \subset \mathbb{P}^3$  with

$\rho(S) = 2$ ,  $\text{Aut}(S) \cong \mathbb{Z}_2 * \mathbb{Z}_2$  (and hence  $\text{Aut}(S)$  is generated by involutions), while  $\text{Dec}(\mathbb{P}^3, S) = \{1\}$ .

In this talk, we present a general framework that can be used to study Gizatullin's problem and to investigate the decomposition and inertia groups of a quartic surface  $S \subset \mathbb{P}^3$ , namely, the birational theory of *Calabi-Yau pairs*. We also report on recent progress on these problems in collaboration with Alessio Corti and Alex Massarenti, and with Daniela Paiva and Sokratis Zikas.

## 2. BIRATIONAL GEOMETRY OF CALABI-YAU PAIRS

Let  $S \subset \mathbb{P}^3$  be a smooth quartic surface. The pair  $(\mathbb{P}^3, S)$  is an example of a *Calabi-Yau pair*, and an element of its decomposition group  $\text{Dec}(\mathbb{P}^3, S)$  is an example of a *volume preserving* birational self-map of  $\mathbb{P}^3$  with respect to the pair  $(\mathbb{P}^3, S)$ . In this section, we introduce these notions in a more general context, and explain how they allow us to use tools from the Minimal Model Program (MMP) in order to investigate Gizatullin's problem.

**Definition.** A *Calabi-Yau (CY) pair* is a pair  $(X, D)$  consisting of a terminal projective variety  $X$  and an effective Weil divisor  $D$  on  $X$  such that  $K_X + D \sim 0$  and  $(X, D)$  has klt singularities.

Let  $(X, D_X)$  and  $(Y, D_Y)$  be CY pairs, and  $f : X \dashrightarrow Y$  a birational map. We say that  $f$  is *volume preserving* if, for every geometric valuation  $E$  with center on both  $X$  and  $Y$ , the discrepancies of  $E$  with respect to the pairs  $(X, D_X)$  and  $(Y, D_Y)$  are equal:  $a(E, K_X + D_X) = a(E, K_Y + D_Y)$ .

The *birational group of a CY pair*  $(X, D)$  is the group  $\text{Bir}(X, D)$  of birational self-maps of  $X$  which are volume preserving with respect to  $(X, D)$ .

**Remark.** The terminology is explained by the following interpretation. Given a CY pair  $(X, D)$ , there is a rational volume form  $\omega$  on  $X$ , unique up to scaling, such that  $D + \text{div}(\omega) = 0$ . A birational self-map of  $X$  is volume preserving with respect to  $(X, D)$  if and only if it preserves the volume form  $\omega$  up to scaling.

When the CY pair  $(X, D)$  has canonical singularities, a birational self-map of  $X$  is volume preserving if and only if it restricts to a birational self-map of  $D$ . This is the case with the pair  $(\mathbb{P}^3, S)$ , where  $S \subset \mathbb{P}^3$  is a quartic surface with at worst rational double points as singularities. In this case, the decomposition group  $\text{Dec}(\mathbb{P}^3, S)$  coincides with the birational group  $\text{Bir}(\mathbb{P}^3, S)$  of the pair  $(\mathbb{P}^3, S)$ , which can be studied with tools of the MMP, as we now explain.

Given a uniruled variety, the MMP produces a *Mori fiber space* that is birationally equivalent to it. In general, there might be several different Mori fiber spaces in the same birational equivalence class. The *Sarkisov program* provides a factorization theorem for birational maps between Mori fiber spaces in terms of simpler birational maps, called *Sarkisov links*. It was established in dimension 3 in [5], and in higher dimensions in [7]. The Sarkisov program has become a powerful tool to investigate the Cremona group, as it allows one to factorize any birational self-map of  $\mathbb{P}^n$  as a composition of Sarkisov links between Mori fiber

spaces. In [6], Corti and Kaloghiros established the following volume preserving version of the Sarkisov program:

**Theorem 1.** *A volume preserving birational map between Mori fibered CY pairs is a composition of volume preserving Sarkisov links.*

### 3. THE DECOMPOSITION GROUP OF QUARTIC SURFACES

We end this report by discussing a few consequences of Theorem 1. In [1], in collaboration with Alessio Corti and Alex Massarenti, we developed a framework to study the birational geometry of CY pairs. By our first main result ([1, Theorem A]), if the quartic surface  $S \subset \mathbb{P}^3$  is very general, then the group  $\text{Bir}(\mathbb{P}^3, S)$  is trivial. More generally:

**Theorem 2.** *Let  $D \subset \mathbb{P}^{n+1}$  be a hypersurface of degree  $n + 1$ . Suppose that  $D$  is terminal and  $\text{Cl}(D) = \mathbb{Z} \cdot [\mathcal{O}_{\mathbb{P}^{n+1}}(1)|_D]$ . Then  $\text{Bir}(\mathbb{P}^{n+1}, D) = \{1\}$ .*

Therefore, if we want to produce interesting subgroups of the Cremona group  $\text{Bir}(\mathbb{P}^3)$  using CY pairs  $(\mathbb{P}^3, S)$ , then the quartic surface  $S \subset \mathbb{P}^3$  must be chosen special. Namely, either  $S$  should be singular, or it must have Picard rank  $\rho(S) \geq 2$ . The singular case was treated in [1, Theorem B]:

**Theorem 3.** *Let  $S \subset \mathbb{P}^3$  be a general singular quartic surface, so that  $S$  has a unique rational double point of type  $A_1$ , and  $\text{Cl}(S) = \mathbb{Z} \cdot [\mathcal{O}_{\mathbb{P}^3}(1)|_S]$ . Then we have a split exact sequence*

$$1 \rightarrow \text{In}(\mathbb{P}^3, S) \rightarrow \text{Bir}(\mathbb{P}^3, S) \xrightarrow{\sim} \text{Bir}(S) \cong \mathbb{Z}_2 \rightarrow 1,$$

and the inertia group  $\text{In}(\mathbb{P}^3, S)$  is a form of  $\mathbb{G}_m$  over  $\mathbb{C}(x, y)$ , i.e.,  $\text{In}(\mathbb{P}^3, S)$  is an algebraic group over the field  $\mathbb{C}(x, y)$  which is isomorphic to  $\mathbb{G}_m$  over the algebraic closure  $\overline{\mathbb{C}(x, y)}$ .

Gizatullin’s problem for smooth quartic surfaces  $S \subset \mathbb{P}^3$  with  $\rho(S) = 2$  was addressed in [12] and in the recent paper [2], in collaboration with Daniela Paiva and Sokratis Zikas. Let  $S \subset \mathbb{P}^3$  be a general smooth quartic surface with  $\rho(S) = 2$ . In a suitable basis for  $\text{Pic}(S) \cong \mathbb{Z}^2$ , the intersection product is given by a matrix of the form

$$\begin{pmatrix} 4 & b \\ b & 2c \end{pmatrix},$$

with  $b, c \in \mathbb{Z}$ . Denote by  $r = b^2 - 8c$  the discriminant of  $S$ . It follows from [12] that  $\text{Dec}(\mathbb{P}^3, S) = \{1\}$  whenever  $r > 233$ . In [2], we determined the image of the restriction homomorphism  $\text{Dec}(\mathbb{P}^3, S) \rightarrow \text{Aut}(S)$  for each value of  $r \leq 233$ . In particular, there are examples of smooth quartic surfaces  $S$  with  $\rho(S) = 2$ ,  $\text{Aut}(S) \cong \mathbb{Z}_2, \mathbb{Z}$  or  $\mathbb{Z}_2 * \mathbb{Z}_2$ , and such that every automorphism of  $S$  is the restriction of a Cremona transformation of  $\mathbb{P}^3$ .

## REFERENCES

- [1] C. Araujo, A. Corti and A. Massarenti, *Birational geometry of Calabi-Yau pairs and 3-dimensional Cremona transformations*, arXiv:2306.00207 (2023).
- [2] C. Araujo, D. Paiva and S. Zikas, *On Gizatullin's Problem for quartic surfaces of Picard rank 2*, in preparation (2024).
- [3] J. Blanc, *On the inertia group of elliptic curves in the Cremona group of the plane*, Michigan Math. J. 56 (2) (2008) 315–330.
- [4] H. C. Chang, *On plane algebraic curves*, Chinese J. Math. 6 (1978) 185–189.
- [5] A. Corti, *Factoring birational maps of threefolds after Sarkisov*, J. Algebraic Geom. 4 (2) (1995) 223–254.
- [6] A. Corti, A.-S. Kaloghiros, *The Sarkisov program for Mori fibred Calabi-Yau pairs*, Algebraic Geometry 3 (3) (2016) 370–384.
- [7] C. D. Hacon and J. McKernan, *The Sarkisov program*, J. Algebraic Geom. 22 (2) (2013) 389–405.
- [8] H. Matsumura and P. Monsky, *On the automorphisms of hypersurfaces*, J. Math. Kyoto Univ. 3 (1964) 347–361.
- [9] K. Oguiso, *Quartic K3 surfaces and Cremona transformations*, Arithmetic and geometry of K3 surfaces and Calabi-Yau threefolds, Fields Inst. Commun., vol. 67, Springer, New York (2013) 455–460.
- [10] K. Oguiso, *Smooth quartic K3 surfaces and Cremona transformations, II*, arxiv:1206.5049 (2012).
- [11] I. Pan, *Sur le sous-groupe de décomposition d'une courbe irrationnelle dans le groupe de Cremona du plan*, Michigan Math. J. 55(2) (2007) 285–298.
- [12] D. Paiva and A. Quedo, *Automorphisms of quartic surfaces and Cremona transformations*, arXiv:2302.09014 (2023).

## Finite subgroups of Cremona groups and representation dimension

ALEXANDER DUNCAN

(joint work with Jason Bailey Heath, Christian Urech)

For a field  $k$  and a positive integer  $n$ , the *Cremona group of degree  $n$  over  $k$* , denoted  $\mathrm{Cr}_n(k)$ , is the group of birational automorphisms of the projective space  $\mathbb{P}_k^n$ . We are interested in understanding how complicated its finite subgroups can be.

We have  $\mathrm{Cr}_1(k) \cong \mathrm{PGL}_2(k)$ , so the finite subgroups are completely classified for all fields when  $n = 1$ . The finite subgroups of  $\mathrm{Cr}_2(\mathbb{C})$  were (almost) completely classified by Dolgachev and Iskovskikh in [2], building on work going back more than a century. Over other fields there has been abundant progress towards full classifications of finite subgroups, but much work still needs to be done. In higher dimensions, there is some hope for  $\mathrm{Cr}_3(k)$ , but a full classification in higher degrees seems out of reach of current techniques.

Rather than a full classification, a coarser approach is to merely bound the complexity of finite subgroups of  $\mathrm{Cr}_n(k)$ . For some “small” fields, such as number fields, one can bound the order of finite subgroup groups as was done by Serre for  $\mathrm{Cr}_2(k)$  in [4]. However, there is no finite bound on order for  $k = \mathbb{C}$  — even for  $n = 1$ . Another alternative is to study the *Jordan constant* of  $\mathrm{Cr}_n(k)$ , which bounds the index of normal abelian subgroups of the finite subgroups. As a consequence of



Birkar’s proof of the BAB Conjecture [1], Prokhorov and Shramov [3] have shown that the Jordan constant is finite for  $\text{Cr}_k(n)$  for all  $n$  and all fields of characteristic 0.

Here we consider another measure of complexity. The *representation dimension* of a finite group  $G$  over  $k$ , denoted  $\text{rdim}_k(G)$ , is the minimal  $N$  such that there is an embedding  $G \hookrightarrow \text{GL}_N(k)$ . For a field  $k$  and a positive integer  $n$ , define

$$c_n(k) := \sup \{ \text{rdim}_k(G) \mid G \text{ finite group such that } G \subseteq \text{Cr}_n(k) \} .$$

We are able to compute  $c_n(k)$  exactly for small  $n$  and arbitrary fields.

$$c_1(k) = \begin{cases} 2 & \text{if char}(k) = 2, \\ 3 & \text{if char}(k) \geq 3, \\ 3 & \text{if char}(k) = 0 \text{ and } -1 \text{ is a sum of two squares,} \\ 2 & \text{otherwise.} \end{cases}$$

$$c_2(k) = \begin{cases} \infty & \text{if char}(k) \neq 0, \\ 8 & \text{if char}(k) = 0 \text{ and } \sqrt{-3} \in k, \\ 6 & \text{otherwise.} \end{cases}$$

In the important special case of the complex space Cremona group, we have the following bounds

$$15 \leq c_3(\mathbb{C}) \leq 62208$$

where we do not believe that the upper bound is sharp.

We also prove that  $c_n(k)$  is infinite for all  $n \geq 2$  and all fields  $k$  of positive characteristic. As a consequence of the finiteness of the Jordan constant, we can show that  $c_n(k)$  is finite for all  $n$  when  $k = \mathbb{C}$  or  $k$  is a number field. Indeed, it is likely that  $c_n(k)$  is finite for all fields of characteristic 0.

Finally, by investigating the automorphism groups of (possibly non-split) toric varieties, we obtain lower bounds for all dimensions  $n$  and all fields  $k$

$n$	1	2	3	4	5	6	$\geq 7$
$c_n(k) \geq$	2	6	12	24	40	72	$2^n$

While explicit *upper* bounds for  $c_n(k)$  for large  $n$  is difficult with current techniques, we do think it is worth trying to find new lower bounds and encourage the community to try to improve the numbers above.

#### REFERENCES

- [1] Caucher Birkar. Singularities of linear systems and boundedness of Fano varieties. *Ann. of Math. (2)*, 193(2):347–405, 2021.
- [2] Igor V. Dolgachev and Vasily A. Iskovskikh. Finite subgroups of the plane Cremona group. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I*, volume 269 of *Progr. Math.*, pages 443–548. Birkhäuser Boston Inc., Boston, MA, 2009.
- [3] Yuri Prokhorov and Constantin Shramov. Jordan property for Cremona groups. *Amer. J. Math.*, 138(2):403–418, 2016.
- [4] Jean-Pierre Serre. A Minkowski-style bound for the order of the finite subgroups of the Cremona group of rank 2 over an arbitrary field. *Moscow Math. J.*, 9:193–208, 2009.

## A valuative criterion for K-stability: beyond the Fano case

SÉBASTIEN BOUCKSOM

(joint work with Mattias Jonsson)

The notion of K-stability was introduced in complex differential geometry as a conjectural—and now partially confirmed—algebro-geometric criterion for the existence of special Kähler metrics. Lately, it has also become a subject in its own respect. The main purpose of this talk is to review a series of two joint papers with Mattias Jonsson [BJ21, BJ22b], in which we show how global pluripotential theory over a trivially valued field, as developed in [BoJ22a], can be used to study K-stability.

Let  $X$  be a projective variety (reduced and irreducible) of dimension  $n \geq 1$  over an algebraically closed field  $k$  of characteristic 0, and  $L$  an ample  $\mathbb{Q}$ -line bundle on  $X$ . The definition of K-stability of the polarized variety  $(X, L)$ , as given by Donaldson, involves the sign of an invariant attached to (ample) test configurations for  $(X, L)$ , which can be interpreted as a non-Archimedean version of the Mabuchi K-energy functional. Filtrations of the section ring of  $(X, L)$  provide another, widely used description of test configurations; more precisely, the latter correspond to  $\mathbb{Z}$ -filtrations of finite type, as first pointed out by Witt Nyström. Chi Li's recent breakthrough on the Yau–Tian–Donaldson conjecture for cscK metrics shows that a stronger form of uniform K-stability, formulated in terms of filtrations, indeed implies the existence of a cscK metric.

To describe this, note that each filtration  $\chi$  can be canonically approximated by a sequence  $\chi_d$  of finitely generated  $\mathbb{Z}$ -filtrations, i.e. test configurations. The proper definition of K-stability for filtrations relies on a detailed study of the non-Archimedean Mabuchi K-energy functional  $M(\chi)$ . For a test configuration, the latter decomposes into ‘energy’ terms  $E(\chi), E^{K_X}(\chi)$ , and an ‘entropy’ term  $\text{Ent}(\chi) := \int A_X \text{MA}(\chi)$ , where  $A_X$  is the log discrepancy function, defined on the set  $X^{\text{div}}$  of divisorial valuations, and the *Monge–Ampère measure*  $\text{MA}(\chi)$  is a *divisorial measure*, i.e. a probability measure with finite support in  $X^{\text{div}}$ . Our first main result is as follows:

**Theorem 1.** *There exists a unique extension of  $E(\chi), E^{K_X}(\chi), \text{MA}(\chi)$  to arbitrary filtrations  $\chi$ , obtained as the limits of the corresponding quantities for the canonical approximants  $\chi_d$ .*

Here  $\text{MA}(\chi)$  is a positive measure on the Berkovich space  $X^{\text{an}}$ , a natural compactification of  $X^{\text{div}}$ . Since  $A_X$  extends to  $X^{\text{an}}$  [BFJ08], this provides a natural extension of  $M(\chi)$  to all filtrations. Our second main result is then:

**Theorem 2.** *The Monge–Ampère operator induces a 1–1 correspondence between the set of divisorial filtrations (up to translation by a constant) and that of divisorial measures. Furthermore, K-stability for filtrations can be tested on the subset of divisorial filtrations.*

## REFERENCES

- [BFJ08] S. Boucksom, C. Favre, M. Jonsson. *Valuations and plurisubharmonic singularities*. Publ. Res. Inst. Math. Sci. **44** (2008), 449–494.
- [BJ21] S. Boucksom, M. Jonsson, *A non-Archimedean approach to K-stability, I: Metric geometry of spaces of test configurations and valuations*, preprint [arXiv:2107.11221](https://arxiv.org/abs/2107.11221), to appear in Ann. Inst. Fourier.
- [BoJ22a] S. Boucksom, M. Jonsson. *Global pluripotential theory over a trivially valued field*. Ann. Fac. Sci. Toulouse. **31** (2022), 647–836.
- [BJ22b] S. Boucksom, M. Jonsson, *A non-Archimedean approach to K-stability, II: A non-Archimedean approach to K-stability, II: divisorial stability and openness*, preprint [arXiv:2206.09492](https://arxiv.org/abs/2206.09492), to appear in Crelle’s journal.

## Moduli Continuity method in K-stability

YUCHEN LIU

(joint work with Chenyang Xu, Junyan Zhao)

This talk is a report on the paper [12] joint with Chenyang Xu and the preprint [14] joint with Junyan Zhao.

It is remarkable that K-stability, a notion introduced by Tian [19] and Donaldson [6] to characterize Kähler–Einstein metrics on Fano varieties, provides the correct condition to construct compact moduli spaces of Fano varieties. As a consequence of about a dozen important recent papers, the K-moduli theorem states that for every fixed dimension  $n$  and volume  $V$ , there exists a projective scheme  $M_{n,V}^K$ , known as the K-moduli space, parameterizes all  $n$ -dimensional K-polystable Fano varieties  $X$  with  $(-K_X)^n = V$ .

Despite the general theory being completed, a natural question remains: can we determine the K-moduli space for explicit Fano varieties? More precisely, the question often asks to describe a certain irreducible component  $M^K$  of  $M_{n,V}^K$  that compactifies a given family of smooth Fano manifolds. A notable approach to this question, called the *moduli continuity method*, is based on the study of singularities and volumes and crucially uses the compactness of K-moduli spaces. The moduli continuity method first appeared in Tian’s solution of the Kähler–Einstein problem for smooth del Pezzo surfaces [18]. Later, Mabuchi–Mukai [15] and Odaka–Spotti–Sun [16] used this method to successfully describe K-moduli spaces of del Pezzo surfaces. In this note, we shall focus on two families of Fano threefolds: cubic threefolds and blow-ups of  $\mathbb{P}^3$  along a genus 4 curve. For further families in higher dimensions, see e.g. [17, 11, 2].

**Theorem 1** ([12]). *The K-moduli space of cubic threefolds is isomorphic to the corresponding GIT moduli space.*

**Theorem 2** ([14]). *The K-moduli space of blow-ups of  $\mathbb{P}^3$  along a genus 4 curve is isomorphic to a variation of GIT moduli space of slope  $\frac{22}{51}$  of (2, 3)-complete intersections in  $\mathbb{P}^3$ .*

Below, we sketch the proofs of these results using the moduli continuity method. Note that the a priori estimate for singularities is good enough to determine the

K-moduli space in the first family. Nonetheless, some new ingredients such as moduli of K3 surfaces and Sarkisov links are needed in the second family.

*Step 0.* We show that the K-moduli space  $M^K$  is non-empty. In other words, this is saying that there exists at least one K-stable member of the given family of Fano varieties. Usually, one finds a member (sometimes singular) with large symmetry and apply the equivariant K-stability criterion [22]. Then, by the openness of K-(semi)stability [3, 20], a general member of the given family is K-stable. For cubic threefolds, we choose the Fermat cubic threefold. For blow-ups of  $\mathbb{P}^3$  along a genus 4 curve, we choose a specific curve with large symmetry; see [1, Proposition 4.33].

*Step 1.* We establish an a priori estimate for singularities that can appear in  $X \in M^K$ . The key estimate is the following local-to-global volume comparison from [10] (after [7]):

$$(1) \quad \frac{(-K_X)^n}{(-K_{\mathbb{P}^n})^n} \leq \frac{\widehat{\text{vol}}(x, X)}{\widehat{\text{vol}}(p, \mathbb{P}^n)}.$$

Here  $n = \dim X$ ,  $p \in \mathbb{P}^n$  is a point, and  $\widehat{\text{vol}}$  stands for the local volume (also known as normalized volume) of a klt singularity introduced by C. Li in [9]. Let us restrict to the case of Fano threefolds. Together with the finite degree formula for local volumes [21] and the ODP Gap Theorem in dimension 3 [12], the inequality (1) implies that as long as  $(-K_X)^3 \geq 20$ , every smoothable  $\mathbb{Q}$ -Cartier Weil divisor  $L$  on  $X$  is Cartier; in particular,  $X$  is Gorenstein canonical.

*Step 2.* We describe the geometry of  $X \in M^K$  by investigating the linear system  $|L|$  for a suitable divisor  $L$ . Let  $\mathcal{X} \rightarrow T$  be a  $\mathbb{Q}$ -Gorenstein smoothing of  $X \cong X_0$  over a pointed smooth curve  $0 \in T$ .

If  $X_t$  is a cubic threefold for  $t \neq 0$ , we take  $L$  to be the degeneration of  $\mathcal{O}_{X_t}(1)$  as a Weil divisor. Then  $L$  is  $\mathbb{Q}$ -Cartier ample as  $-K_X \sim 2L$ . From Step 1, we conclude that  $L$  is Cartier. Thus [8] implies that  $|L|$  is very ample and induces a closed embedding  $X \hookrightarrow \mathbb{P}^4$  as a (possibly singular) cubic hypersurface.

If  $\pi_t : X_t \rightarrow \mathbb{P}^3$  is the blow-up along a genus 4 curve  $C_t$  for  $t \neq 0$ , we take  $\mathcal{L}$  to be the divisor on the total space  $\mathcal{X}$  such that  $\mathcal{L}|_{X_t} = \pi_t^* \mathcal{O}_{\mathbb{P}^3}(1)$ , and let  $L = \mathcal{L}|_{X_0}$ . Then we encounter a major issue as  $\mathcal{L}$  may not be  $\mathbb{Q}$ -Cartier since it is no longer proportional to  $-K_{\mathcal{X}/T}$ . To resolve this, we take a small  $\mathbb{Q}$ -Cartierization  $\tilde{\mathcal{X}} \rightarrow \mathcal{X}$  so that the strict transform  $\tilde{\mathcal{L}}$  of  $\mathcal{L}$  on  $\tilde{\mathcal{X}}$  is  $\mathbb{Q}$ -Cartier and ample over  $\mathcal{X}$ . Then we prove that  $\tilde{\mathcal{L}}$  is indeed big and semiample over  $T$ , and  $X \cong \tilde{X}_0$  is still a blow-up of  $\mathbb{P}^3$ . This is achieved by Reid's technique of general elephants, a delicate analysis of moduli of lattice-polarized K3 surfaces, and the Sarkisov link structure on these Fano threefolds as blow-ups of singular cubic threefolds.

*Step 3.* We show that the K-moduli space  $M^K$  is isomorphic to a suitable GIT moduli space  $M^{\text{GIT}}$ . In summary, the previous steps show that every  $X \in M^K$  belongs to a suitable parameter space  $W$  with an action of a reductive group  $G$ . If the above estimates are strong enough, then we often have that the Picard rank of  $W$  is small, and the CM line bundle  $\lambda_{\text{CM}}$  on  $W$  is ample. Thus we can take the GIT quotient  $M^{\text{GIT}} = W //_{\lambda_{\text{CM}}} G$ . By the Paul–Tian criterion, we have an

injective birational morphism

$$\phi : M^K \rightarrow M^{\text{GIT}}.$$

Since  $M^K$  is proper [4, 13], we conclude that  $\phi$  is an isomorphism by Zariski's main theorem. For cubic threefolds, we take  $W = \mathbb{P}H^0(\mathbb{P}^4, \mathcal{O}(3))$  and  $G = \text{PGL}(5)$ , and Theorem 1 follows. For blow-ups of  $\mathbb{P}^3$  along genus 4 curves, we take  $W$  to be the projective bundle over  $\mathbb{P}^9 = \mathbb{P}H^0(\mathbb{P}^3, \mathcal{O}(2))$  parameterizing  $(2, 3)$ -intersections in  $\mathbb{P}^3$ , and  $G = \text{PGL}(4)$ . Then computations of the CM line bundle show that  $\lambda_{\text{CM}}$  has slope  $\frac{22}{51}$  in the Picard group of  $W$ , which has rank 2. Thus Theorem 2 follows.

Finally, we note that the variation of GIT in Theorem 2 was studied in detail in [5], where it was shown that the VGIT moduli spaces provide models for the Hassett–Keel program of genus 4 curves. Therefore, our K-moduli space  $M^K$  for the second family of Fano threefolds appears as a specific model in the Hassett–Keel program.

#### REFERENCES

- [1] C. Araujo, A. Castravet, I. Cheltsov, K. Fujita, A. Kaloghiros, J. Martinez-Garcia, C. Shramov, H. Süß, and N. Viswanathan, *The Calabi problem for Fano threefolds*, London Math. Soc. Lecture Note Ser., **485** Cambridge University Press, Cambridge, 2023.
- [2] K. Ascher, K. DeVleming, and Y. Liu, *K-stability and birational models of moduli of quartic K3 surfaces*, *Invent. Math.* **232** (2023), no. 2, 471–552.
- [3] H. Blum, Y. Liu, and C. Xu, *Openness of K-semistability for Fano varieties*, *Duke Math. J.* **171** (2022), no. 13, 2753–2797.
- [4] H. Blum and C. Xu, *Uniqueness of K-polystable degenerations of Fano varieties*, *Ann. of Math.* **190** (2019), no. 2, 609–656.
- [5] S. Casalaina-Martin, D. Jensen, and R. Laza, *Log canonical models and variation of GIT for genus 4 canonical curves*, *J. Algebraic Geom.* **23** (2014), no. 4, 727–764.
- [6] S. Donaldson, *Scalar curvature and stability of toric varieties*, *J. Differential Geom.* **62** (2002), no. 2, 289–349.
- [7] K. Fujita, *Optimal bounds for the volumes of Kähler-Einstein Fano manifolds*, *Amer. J. Math.*, **140** (2018), no. 2, 391–414.
- [8] T. Fujita, *On singular del Pezzo varieties*, *Algebraic geometry (L'Aquila, 1988)*, 117–128, *Lecture Notes in Math.*, **1417**, Springer, Berlin, 1990.
- [9] C. Li, *Minimizing normalized volumes of valuations*, *Math. Z.* **289** (2018), no. 1-2, 491–513.
- [10] Y. Liu, *The volume of singular Kähler-Einstein Fano varieties*, *Compos. Math.*, **154** (2018), no. 6, 1131–1158.
- [11] Y. Liu, *K-stability of cubic fourfolds*, *J. Reine Angew. Math.* **786** (2022), 55–77.
- [12] Y. Liu and C. Xu, *K-stability of cubic threefolds*, *Duke Math. J.* **168** (2019), no. 11, 2029–2073.
- [13] Y. Liu, C. Xu, and Z. Zhuang, *Finite generation for valuations computing stability thresholds and applications to K-stability*, *Ann. of Math. (2)* **196** (2022), no. 2, 507–566.
- [14] Y. Liu and J. Zhao, *K-moduli of Fano threefolds and genus four curves*, preprint arXiv:2403.16747 (2024).
- [15] T. Mabuchi and S. Mukai, *Stability and Einstein-Kähler metric of a quartic del Pezzo surface*, *Einstein metrics and Yang-Mills connections (Sanda, 1990)*, 133–160, *Lecture Notes in Pure and Appl. Math.*, **145**, Dekker, New York, 1993.
- [16] Y. Odaka, C. Spotti, and S. Sun, *Compact moduli spaces of del Pezzo surfaces and Kähler-Einstein metrics*, *J. Differential Geom.* **102** (2016), no. 1, 127–172.
- [17] C. Spotti and S. Sun, *Explicit Gromov-Hausdorff compactifications of moduli spaces of Kähler-Einstein Fano manifolds*, *Pure Appl. Math. Q.* **13** (2017), no. 3, 477–515.

- [18] G. Tian, *On Calabi's conjecture for complex surfaces with positive first Chern class*, Invent. Math. **101** (1990), no. 1, 101–172.
- [19] G. Tian, *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. **130** (1997), no. 1, 1–37.
- [20] C. Xu, *A minimizing valuation is quasi-monomial*, Ann. of Math. **191** (2020), no. 3, 1003–1030.
- [21] C. Xu and Z. Zhuang, *Uniqueness of the minimizer of the normalized volume function*, Camb. J. Math. **9** (2021), no. 1, 149–176.
- [22] Z. Zhuang, *Optimal destabilizing centers and equivariant K-stability*, Invent. Math. **226** (2021), no. 1, 195–223.

## Non-archimedean approach to SYZ conjecture

ENRICA MAZZON

Mirror symmetry is a fast-moving research area at the boundary between mathematics and theoretical physics. Originated from observations in string theory, it suggests the existence of a duality between Calabi–Yau (CY) manifolds, complex manifolds with a nowhere vanishing holomorphic form of maximal degree. It predicts that every CY manifold  $X$  has a mirror partner  $\check{X}$ , such that the complex geometry of  $\check{X}$  is equivalent to the symplectic geometry of  $X$ , in some appropriate sense, and vice versa.

Various approaches have been developed to find a rigorous definition of a mirror pair  $(X, \check{X})$ , and methods to construct mirror partners; a geometric explanation was proposed by Strominger, Yau and Zaslow (SYZ) in [SYZ96]. In its current formulation, the SYZ conjecture concerns CY manifolds in certain degenerating families rather than individual manifolds. More precisely, consider a projective family  $(X_t)_t$  of CY varieties of dimension  $n$  over a punctured disk, such that the family is maximally degenerate, i.e. the monodromy operator on the degree  $n$  cohomology of  $X_t$  has a Jordan block of maximal size, that is  $n + 1$ .

**Conjecture 1** (SYZ conjecture). For all sufficiently small  $t$ ,  $X_t$  admits a fibration  $\pi : X_t \rightarrow B$ , whose fibres are special Lagrangian tori, away from a locus  $\Delta$  of codimension 2 in  $B$ . Moreover, the mirror partner  $\check{X}_t$  of  $X_t$  is obtained by dualizing the special Lagrangian toric fibres of  $\pi$  and by suitably compactifying the resulting space.

While some examples of special Lagrangian torus fibrations can be produced, dealing with the general case seems very difficult. The insight of Kontsevich and Soibelman is to replace the above conjecture by an analogous one in the non-archimedean world, and to interpret the latter as an asymptotic limit of the complex phenomenon when  $t \rightarrow 0$ .

More precisely, one can associate to the degenerating family  $X = (X_t)_t$  the Berkovich non-archimedean space  $X^{\text{an}}$ , whose points are valuations defined locally on  $X$ . Given a degeneration  $\mathcal{X}$  of  $X$ , we say that  $\mathcal{X}$  is snc (respectively dlt) if the pair  $(\mathcal{X}, \mathcal{X}_0)$  is strict normal crossing (respectively divisorially log terminal), where  $\mathcal{X}_0$  is fiber over  $t = 0$ ; see [Kol13] for more details. Given any snc or dlt

degeneration, the dual intersection complex  $\mathcal{D}(\mathcal{X}_0)$  is a simplicial complex encoding the combinatorics of the multiple intersections of the components of  $\mathcal{X}_0$ . It admits a canonical embedding in  $X^{\text{an}}$ , whose image is called the skeleton of  $\mathcal{X}$  and denoted  $\text{Sk}(\mathcal{X})$ , and a retraction  $\rho_{\mathcal{X}} : X^{\text{an}} \rightarrow \text{Sk}(\mathcal{X})$ . Among various degenerations, minimal (in the sense of MMP) dlt models  $\mathcal{X}$  of  $X$  determine a canonical skeleton  $\text{Sk}(X) = \text{Sk}(\mathcal{X})$ , called the essential skeleton of  $X$  and independent of the choice of the minimal model; see [MN15, NX16] for more details. The essential skeleton and the retractions  $\rho_{\mathcal{X}} : X^{\text{an}} \rightarrow \text{Sk}(\mathcal{X})$  are of particular relevance in the non-archimedean approach to the SYZ conjecture, as we will see in the sequel.

By the celebrated Yau theorem,  $X_t$  carries a unique Kähler form  $\omega_t$  such that  $[\omega_t] \in c_1(L_t)$  and  $\omega_t^n = C_t \Omega_t \wedge \overline{\Omega}_t$  for a constant  $C_t$ . Finding such form  $\omega_t$  boils down to solving an equation, called complex Monge–Ampère equation. By the works [CL06, BFJ15], non-archimedean Monge–Ampère equations can be defined on  $X^{\text{an}}$  as well, and solved for any measure supported on a subset  $\mathcal{D}(\mathcal{X}_0)$  of  $X^{\text{an}}$ . In particular, let  $\Psi$  be the solution to

$$\text{MA}_{\text{NA}}(\Psi) = d\mu_{\text{Sk}(X)},$$

where  $\text{MA}_{\text{NA}}$  denotes the non-archimedean Monge–Ampère operator, and  $d\mu_{\text{Sk}(X)}$  the Lebesgue measure on  $\text{Sk}(X)$ . Up to fixing a reference metric, we can think of  $\Psi$  as a function on  $X^{\text{an}}$ . In [Li23] Li reduced the SYZ conjecture to a conjecture in non-archimedean geometry about Monge–Ampère metrics

**Theorem 1** ([Li23]). *Let  $X$  be a maximally degenerate family of Calabi–Yau varieties. If there exists a degeneration  $\mathcal{X}$  of  $X$  such that the solution  $\Psi$  is invariant with respect to the retraction  $\rho_{\mathcal{X}}$ , i.e.*

$$\Psi = \Psi \circ \rho_{\mathcal{X}} \quad \text{on} \quad \rho_{\mathcal{X}}^{-1}(\text{Int}(\tau)),$$

*over the interior of any maximal face  $\tau$  of  $\mathcal{D}(\mathcal{X}_0)$ , then an SYZ fibration exists on a large region  $U_t \subseteq X_t$ .*

This approach reduces the construction of SYZ fibrations to a property in non-archimedean geometry. In [HJMM24], Hultgren, Jonsson, McCleerey and I provided first evidence for such conjecture, when  $X$  is not one-dimensional or an abelian variety. More precisely, let  $X = \{z_0 z_1 \dots z_{n+1} + t f(z) = 0\} \subset \mathbb{P}^{n+1}$  be a family of Calabi–Yau hypersurfaces where  $f$  is a generic polynomial of degree  $n + 2$ . In this case, the essential skeleton  $\text{Sk}(X)$  is a sphere and can be identified with the boundary of the standard unit simplex in  $\mathbb{R}^{n+1}$ .

**Theorem 2** ([HJMM24]). *If  $\nu$  is a symmetric measure on  $\text{Sk}(X)$ , then the solution to  $\text{MA}_{\text{NA}}(\cdot) = \nu$  is the restriction of a symmetric toric metric on  $\mathcal{O}_{\mathbb{P}^{n+1}}(n + 2)^{\text{an}}$ , thus is determined by the restriction to  $\text{Sk}(X)$  of a convex function on  $\mathbb{R}^{n+1}$ .*

Applying Theorem 2 to  $\nu = d\mu_{\text{Sk}(X)}$ , we show that the characterization of the solution  $\Psi$  provided by the theorem is sufficient to prove the invariance property of Theorem 1. We conclude therefore that SYZ fibrations exist on large regions of CY hypersurfaces. See

## REFERENCES

- [AH23] R. Andreasson and J. Hultgren. *Solvability of Monge-Ampère equations and tropical affine structures on reflexive polytopes*. [arXiv:2303.05276](#).
- [BFJ15] S. Boucksom, C. Favre, and M. Jonsson. *Solution to a non-Archimedean Monge-Ampère equation*. *Journal of the American Mathematical Society* **3** (2015), 617–667.
- [CL06] A. Chambert-Loir. *Mesures et équidistribution sur les espaces de Berkovich*. *Journal für die Reine und Angewandte Mathematik* **595** (2006), 215–235.
- [HJMM24] J. Hultgren, M. Jonsson, E. Mazzon and N. McCleerey. *Tropical and non-Archimedean Monge-Ampère equations for a class of Calabi-Yau hypersurfaces*. *Adv. Math.* **439** (2024), 109494.
- [Kol13] J. Kollár. *Singularities of the minimal model program*. Cambridge University Press, Cambridge, 200 (2013).
- [KS06] M. Kontsevich and Y. Soibelman. *Affine structures and non-Archimedean analytic spaces*. In *The unity of mathematics*. *Progr. Math.* 244, 321–385. Birkhäuser Boston, Boston, MA, 2006.
- [Li24] Y. Li. *Metric SYZ conjecture for certain toric Fano hypersurfaces*. *Cambridge J. Math.* **12** (2024), 223–252.
- [Li23] Y. Li. *Metric SYZ conjecture and non-archimedean geometry*. *Duke Math. J.* **172** (2023), 3227–3255.
- [MN15] M. Mustață and J. Nicaise. *Weight functions on non-archimedean analytic spaces and the Kontsevich–Soibelman skeleton*. *Algebr. Geom.* **2** (2015), 365–404.
- [NX16] J. Nicaise and C. Xu. *The essential skeleton of a degeneration of algebraic varieties*. *American Journal of Mathematics* **138** (2016), 1645–1667.
- [SYZ96] A. Strominger, S.-T. Yau, E. Zaslov. *Mirror Symmetry is T-duality*. *Nucl. Phys.* B479, 243–259, 1996.



## Participants

**Dr. Eduardo Alves da Silva**

Laboratoire de Mathématique d'Orsay  
Université Paris-Saclay  
Bâtiment 307, rue Michel Magat  
91405 Orsay Cedex  
FRANCE

**Prof. Dr. Carolina Araujo**

Instituto de Matematica Pura e  
Aplicada (IMPA)  
Estrada Dona Castorina 110  
Rio de Janeiro RJ 22460-320  
BRAZIL

**Dr. Ignacio Barros**

University of Antwerpen  
Middelheimlaan 1  
2020 Antwerpen  
BELGIUM

**Prof. Dr. Jérémy Blanc**

Institut de Mathématiques  
Université de Neuchâtel  
Unimail  
Rue Emile Argand 11  
2000 Neuchâtel  
SWITZERLAND

**Aurore Boitrel**

UFR Sciences  
Institut de Mathématiques d'Orsay  
Université Paris-Saclay  
Bâtiment 307, rue Michel Magat  
91405 Orsay Cedex  
FRANCE

**Anna Bot**

Departement Mathematik und  
Informatik der Universität Basel  
Fachbereich Mathematik  
Spiegelgasse 1  
4051 Basel  
SWITZERLAND

**Prof. Dr. Sebastien Boucksom**

IMJ-PRG  
Sorbonne Université – Campus Pierre et  
Marie Curie  
4 place Jussieu  
P.O. Box 247  
75252 Paris Cedex 05  
FRANCE

**Prof. Dr. Ana-Maria Castravet**

Laboratoire de Mathématiques  
Université Paris-Saclay, Versailles  
Bâtiment Fermat 3304  
45 Avenue des États Unis  
78035 Versailles Cedex  
FRANCE

**Prof. Dr. Ivan Cheltsov**

School of Mathematics  
University of Edinburgh  
King's Building Campus  
Peter Guthrie Tait Road  
Edinburgh EH9 3FD  
UNITED KINGDOM

**Dr. Thibaut Delcroix**

IMAG – UMR 5149  
Case courrier 051  
Université de Montpellier  
Place Eugène Bataillon  
34090 Montpellier  
FRANCE

**Elena Denisova**

School of Mathematics  
University of Edinburgh  
James Clerk Maxwell Bldg.  
King's Buildings, Mayfield Road  
Edinburgh EH9 3JZ  
UNITED KINGDOM

**Dr. Tiago Duarte Guerreiro**

Dept. of Mathematics  
University of Essex  
Wivenhoe Park  
Colchester CO4 3SQ  
UNITED KINGDOM

**Dr. Liana Heuberger**

Dept. of Mathematical Sciences  
University of Bath  
Claverton Down  
Bath BA2 7AY  
UNITED KINGDOM

**Prof. Dr. Alexander R. Duncan**

LeConte 448  
Department of Mathematics  
University of South Carolina  
1523 Greene Street  
Columbia SC, 29208  
UNITED STATES

**Dr. Anne-Sophie Kaloghiros**

Department of Mathematics and  
Statistics  
Brunel University  
Kingston Lane  
Uxbridge Middlesex UB8 3PH  
UNITED KINGDOM

**Erroxe Etxabarri Alberdi**

School of Mathematical Sciences  
The University of Nottingham  
University Park  
Nottingham NG7 2RD  
UNITED KINGDOM

**Dr. King-Leung Lee**

Département de Mathématiques  
Université Montpellier II  
Place Eugene Bataillon  
34095 Montpellier Cedex 5  
FRANCE

**Dr. Pascal Fong**

Université Paris-Saclay  
Faculté des sciences  
Laboratoire de Mathématique d'Orsay  
Equipe AGA  
rue Michel Magat  
91405 Orsay Cedex  
FRANCE

**Dr. Eveline Legendre**

Institut Camille Jordan  
Université Claude Bernard Lyon 1  
43 blvd. du 11 novembre 1918  
69622 Villeurbanne Cedex  
FRANCE

**Dr. Giulia Gugiatti**

Mathematics Section  
The Abdus Salam International Centre  
for Theoretical Physics (ICTP)  
Strada Costiera, 11  
34151 Trieste  
ITALY

**Dr. Yuchen Liu**

Department of Mathematics  
Lunt Hall  
Northwestern University  
Evanston, IL 60208-2730  
UNITED STATES

**Prof. Dr. Jürgen Hausen**

Fachbereich Mathematik  
Universität Tübingen  
Auf der Morgenstelle 10  
72076 Tübingen  
GERMANY

**Dr. Jesus Martinez-Garcia**

Dept. of Mathematics  
University of Essex  
Wivenhoe Park  
Colchester CO4 3SQ  
UNITED KINGDOM

**Dr. Enrica Mazzon**

Department of Mathematics  
University of Regensburg  
93040 Regensburg  
GERMANY

**Dr. Leonid Monin**

Institute of Mathematics  
EPFL  
1015 Lausanne  
SWITZERLAND

**Tran-Trung Nghiem**

Département de Mathématiques  
Université Montpellier II  
Place Eugene Bataillon  
34095 Montpellier Cedex 5  
FRANCE

**Dr. Andrea Petracci**

Dipartimento di Matematica  
Università di Bologna  
Piazza di Porta S. Donato, 5  
40126 Bologna  
ITALY

**Dr. Léonard Pille-Schneider**

Fakultät für Mathematik  
Universität Regensburg  
Universitätsstr. 31  
93053 Regensburg  
GERMANY

**Antoine Pinardin**

School of Mathematics  
University of Edinburgh  
James Clerk Maxwell Bldg.  
King's Buildings, Mayfield Road  
Edinburgh EH9 3JZ  
UNITED KINGDOM

**Dr. Roland Půček**

Mathematisches Institut  
Universität Jena  
Ernst-Abbe-Platz 2-4  
07743 Jena  
GERMANY

**Dr. Karin Schaller**

Fachbereich Mathematik und Informatik  
Freie Universität Berlin  
Arnimallee 3  
14195 Berlin  
GERMANY

**Dr. Julia Schneider**

Institut für Mathematik  
Universität Zürich  
Winterthurerstr. 190  
8057 Zürich  
SWITZERLAND

**Shreya Sharma**

Department of Mathematics  
University of South Carolina  
Columbia, SC 29208  
UNITED STATES

**Jonathan Smith**

Department of Mathematics  
University of South Carolina  
Columbia, SC 29208  
UNITED STATES

**Prof. Dr. Hendrik Süß**

Mathematisches Institut  
Universität Jena  
Ernst-Abbe-Platz 2-4  
07743 Jena  
GERMANY

**Prof. Dr. Carl Tipler**

Dept. de Mathématiques  
et Informatique  
Université de Bretagne Occidentale  
6, Avenue Victor Le Gorgeu  
29238 Brest Cedex 3  
FRANCE

**Dr. Nikolaos Tsakanikas**

Institut de Mathématiques  
Station 8, Bâtiment MA, MA C3 595  
École Polytechnique Fédérale de  
Lausanne (EPFL)  
1015 Lausanne  
SWITZERLAND

**Dr. Christian Urech**

Département Mathematik  
ETH-Zentrum  
Rämistr. 101  
8092 Zürich  
SWITZERLAND

**Ying Wang**

Department of Mathematics  
University of Michigan  
530 Church Street  
Ann Arbor, MI 48109-1043  
UNITED STATES

**Prof. Dr. Jaroslaw Wisniewski**

Instytut Matematyki  
Uniwersytet Warszawski  
ul. Banacha 2  
02-097 Warszawa  
POLAND

**Dr. Milena Wrobel**

Institut für Mathematik  
Carl-von-Ossietzky-Universität  
Oldenburg  
Ammerländer Heerstraße 114-118  
26129 Oldenburg  
GERMANY

**Dr. Zhixin Xie**

Institut Élie Cartan de Lorraine  
Université de Lorraine  
B.P. 70239, F-54506  
Vandœuvre-lès-Nancy  
FRANCE

**Dr. Egor Yasinsky**

Mathématiques et Informatique  
Université Bordeaux I  
351, cours de la Libération  
33405 Talence Cedex  
FRANCE

**Dr. Sokratis Zikas**

Université de Poitiers  
Département de Mathématiques  
Teleport 2 - BP 30179  
Blvd. Marie et Pierre Curie  
86962 Futuroscope Chasseneuil Cedex  
FRANCE

**Prof. Dr. Susanna Zimmermann**

Laboratoire de Mathématique d'Orsay  
Université Paris-Saclay  
Bâtiment 307  
rue Michel Magat  
91405 Orsay Cedex  
FRANCE