



Bendong Lou · Maolin Zhou

Convergence of solutions of the porous medium equation with reactions

Received 21 August 2022; revised 29 January 2024

Abstract. Consider the Cauchy problem of one-dimensional porous medium equation (PME) with reactions. We first prove a general convergence result, that is, any bounded global solution starting at a nonnegative compactly supported initial data converges as $t \rightarrow \infty$ to a nonnegative zero of the reaction term or a ground state stationary solution. Based on it, we give out a complete classification on the asymptotic behavior of the solutions for PME with monostable, bistable and combustion types of nonlinearities.

Keywords: porous medium equation with reactions, Cauchy problem, free boundary, asymptotic behavior, spreading and vanishing.

Contents

1. Introduction	2
2. Preliminaries	10
2.1. Well-posedness and a priori estimates	10
2.2. Waiting times of the free boundaries and compactly supported stationary solutions	11
2.3. Lipschitz continuity, Darcy law and strict monotonicity	18
3. Zero number argument and general convergence result	23
3.1. Zero number and intersection number properties	23
3.2. Monotonicity outside of the initial support	28
3.3. General convergence	30
4. Monostable RPME	39
4.1. Stationary solutions	39
4.2. Hair-trigger effect	41
5. Combustion RPME	42
5.1. Stationary solutions	42
5.2. Asymptotic behavior of the solutions	42
6. Bistable RPME	47

Bendong Lou: Mathematics and Science College, Shanghai Normal University, No. 100 Guilin Road, 200234 Shanghai, P. R. China; lou@shnu.edu.cn

Maolin Zhou: Chern Institute of Mathematics and The Key Laboratory of Pure Mathematics and Combinatorics (LPMC), Nankai University, No. 94 Weijin Road, 300071 Tianjin, P. R. China; zhouml123@nankai.edu.cn

Mathematics Subject Classification 2020: 35K65 (primary); 35K57, 35B40, 35K15 (secondary).

6.1. Stationary solutions	47
6.2. Asymptotic behavior	50
7. Complete vanishing phenomenon	52
A. Well-posedness and a priori estimates	56
A.1. Well-posedness	56
A.2. Persistence of positivity and finite propagation speed	56
A.3. A priori estimate for the pressure	58
References	63

1. Introduction

Consider the following reaction-diffusion equation (abbreviated as RDE in the following):

$$u_t = u_{xx} + f(u), \quad x \in \mathbb{R}, \quad t > 0. \quad (1.1)$$

In the last decades, many authors gave systemic qualitative study for this equation. The reaction terms include the following three typical types:

(f_m) monostable case, (f_b) bistable case, (f_c) combustion case.

In the monostable case, we assume $f = f(u) \in C^1([0, \infty))$ and

$$\begin{aligned} f(0) = f(1) = 0, \quad f'(0) > 0, \quad f'(1) < 0, \\ (1-u)f(u) > 0 \quad \text{for } u > 0, \quad u \neq 1. \end{aligned} \quad (1.2)$$

The well-known example is the logistic term, that is, the Verhulst law $f(u) = u(1-u)$. In the bistable case, we assume $f = f(u) \in C^1([0, \infty))$ and

$$f(0) = f(\theta) = f(1) = 0, \quad f(u) \begin{cases} < 0 & \text{in } (0, \theta), \\ > 0 & \text{in } (\theta, 1), \\ < 0 & \text{in } (1, \infty), \end{cases} \quad \int_0^1 f(s)ds > 0, \quad (1.3)$$

for some $\theta \in (0, 1)$, $f'(0) < 0$, $f'(1) < 0$. A typical example is $f(u) = u(u-\theta)(1-u)$ with $\theta \in (0, \frac{1}{2})$. In the combustion case, we assume there exists $\theta \in (0, 1)$ such that $f \in C^1([\theta, \infty))$ and

$$\begin{aligned} f(u) = 0 \quad \text{in } [0, \theta], \quad f(u) > 0 \quad \text{in } (\theta, 1), \quad f'(1) < 0, \quad f(u) < 0 \quad \text{in } (1, \infty), \\ \text{there exists a small } \delta > 0 \text{ such that } f'(u) > 0 \text{ for } u \in (\theta, \theta + \delta]. \end{aligned} \quad (1.4)$$

In 1937, Fisher [18] and Kolmogorov et al. [23] studied the RDE with logistic reaction in their pioneer works. In particular, they specified the *traveling wave solutions*. In 1975 and 1978, Aronson and Weinberger [6, 7] studied the asymptotic behavior for the solutions of the Cauchy problem of RDE. Among others, in the monostable case, they proved the *hair-trigger effect*, which says that *spreading happens* (that is, $u \rightarrow 1$ as $t \rightarrow \infty$, which is also called *persistence* or *propagation phenomenon*) for any positive solution of the monostable equation; in the bistable case, they gave sufficient conditions

for *spreading* and *vanishing* (that is, $u \rightarrow 0$ as $t \rightarrow \infty$, which is also called *extinction phenomenon*). In 1977, Fife and McLeod [17] also studied the bistable equation and proved the existence and stability of the traveling wave solution. In 2006, Zlatoš [32] gave further systemic study on the asymptotic behavior for the solutions of bistable and combustion equations. More precisely, he considered the Cauchy problem of (1.1) with initial data $u(x, 0) = \chi_{[-L, L]}(x)$ (which denotes the characteristic function over $[-L, L]$), and proved a trichotomy result: there exists a sharp value $L_* > 0$ such that when $L > L_*$ (resp. $L < L_*$), spreading (resp. vanishing) happens for the solution. In addition, the solution develops to a transition one if and only if $L = L_*$, which is a ground state solution in the bistable case and the largest zero $\theta \in (0, 1)$ of f in the combustion case. In 2010, Du and Matano [14] extended these results to the Cauchy problem with general initial data like $u(x, 0) = \sigma \phi(x)$ for positive number σ and continuous nonnegative compactly supported function ϕ . For bistable and combustion equations, they also proved the trichotomy and sharp transition results as in [32]. In 2015, Du and Lou [12] proved similar results for the nonlinear Stefan problem.

In this paper, we study the porous medium equation with reactions,

$$\begin{cases} u_t = (u^m)_{xx} + f(u), & x \in \mathbb{R}, t > 0, \\ u(x, 0) = u_0(x), & x \in \mathbb{R}, \end{cases} \quad (\text{CP})$$

where $m > 1$ is a constant. This problem can be used to model population dynamics with diffusion flux depending on the population density, the combustion, propagation of interfaces and nerve impulse propagation phenomena in porous media, as well as the propagation of intergalactic civilizations in the field of astronomy (cf. [21, 22, 25, 28] etc.). For simplicity, we use PME/RPME to refer to the porous media equation in (CP) without/with reaction.

The well-posedness of problem (CP) was studied in [2, 3, 27, 30] etc. For the qualitative property, the monostable equation was considered widely. For example, in 1979, Aronson [3] considered the problem with logistic reaction. He studied the existence of traveling wave solutions and compared the results with that in RDE. It was shown that (see also [9, 20, 29]) the monostable RPME has a traveling wave solution $u(x, t) = \tilde{U}(x - ct)$ if and only if $m \geq 1$ and only for the speed $c \geq c_*(m)$. For bistable and combustion RPMEs, however, there is only one traveling wave solution $\tilde{U}(x - c_*t)$ (cf. [20]). Recently, Du, Quirós and Zhou [15] considered the (N -dimensional) RPME with Fisher–KPP type of reaction. In particular, they presented a precise estimate for the spreading speed of the free boundary with a logarithmic correction. On the other hand, the qualitative properties for the RPMEs with bistable or combustion reaction were not as clear as in the monostable case. In 1982, Aronson, Crandall and Peletier [5] considered the bistable RPME in a bounded interval. Besides the well-posedness, they specified the stationary solutions which can be ω -limits of the solutions. Recently, Gárriz [19] considered problem (CP) with reactions of the monostable, bistable and combustion types. He gave sufficient conditions ensuring the spreading or vanishing, and used the traveling wave solution to characterize the spreading solutions starting at compactly supported initial data.




Fig. 1. An example of the initial data.

Our aim in this paper is to give out a complete classification for the asymptotic behavior of the solutions of (CP). We will specify its stationary solutions; prove a general convergence result, that is, any nonnegative bounded global solution converges as $t \rightarrow \infty$ to a stationary one. Then, based on the general result, we prove the hair-trigger effect for monostable PME and spreading-transition-vanishing trichotomy results on the asymptotic behavior for solutions of bistable or combustion RPMEs. In some sense, this paper can be regarded as an RPME version of the important works [12, 14, 32] for RDEs.

Our basic assumption on f is

$$\begin{aligned} f(u) &\in C^2([0, \infty)) \quad \text{with Lipschitz number } K, \\ f(0) &= 0 \quad \text{and} \quad f(u) < 0 \quad \text{for } u > 1. \end{aligned} \tag{F}$$

The C^2 smoothness is mainly used to give the a priori estimates, including the lower bound for v_{xx} (where v is the pressure, see appendix for details). In the study of asymptotic behavior, we actually only need $f \in C^1$. The assumption $f(u) < 0$ for $u > 1$ is natural in population dynamics, which means that the population has a finite capacity. If one only studies the well-posedness, this requirement can be weakened or omitted. The initial data are chosen from the following set (see Figure 1):

$$\begin{aligned} u_0 \in \mathfrak{X} := \{ \psi \in C(\mathbb{R}) \mid &\text{there exist } b > 0 \text{ and a finite number of open intervals} \\ &\text{in } [-b, b] \text{ such that } \psi(x) > 0 \text{ in these intervals,} \\ &\text{and } \psi(x) = 0 \text{ otherwise} \}. \end{aligned} \tag{I}$$

It is also possible to consider more general initial data, for example, $u_0 \in C(\mathbb{R})$ with compact support or $u_0 \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$. For simplicity and clarity of presentation, especially, for clarity of the free boundaries, we consider in this paper only the initial data in $\mathfrak{X}$. In this case, the solution has a left-most free boundary $l(t)$, a right-most free boundary $r(t)$, and finite number of interior free boundaries in $(l(t), r(t))$.

For any $T > 0$, denote $Q_T := \mathbb{R} \times (0, T)$. A function $u(x, t) \in C(Q_T) \cap L^\infty(Q_T)$ is called a *very weak solution* of (CP) if for any $\varphi \in C_c^\infty(Q_T)$, there holds

$$\begin{aligned} \int_{\mathbb{R}} u(x, T) \varphi(x, T) dx &= \int_{\mathbb{R}} u_0(x) \varphi(x, 0) dx + \iint_{Q_T} f(u) \varphi dx dt \\ &\quad + \iint_{Q_T} [u \varphi_t + u^m \varphi_{xx}] dx dt. \end{aligned} \tag{1.5}$$

As an extension of the definition, if u satisfies (1.5) with inequality “ $\geq$ ” (resp. “ $\leq$ ”), instead of equality, for every test function $\varphi \geq 0$, then u is called a *very weak supersolution/very weak upper solution* (resp. *very weak subsolution/very weak lower solution*) of (CP) (cf. [30, Chapter 5]).

From the references [5, 9, 27, 30] etc. we know that under assumptions (F) and (I), problem (CP) has a unique very weak solution $u(x, t) \in C(Q_T) \cap L^\infty(Q_T)$ for any $T > 0$, $u(x, t) \geq 0$ in Q_T , and the support of $u(\cdot, t)$, denoted by $\text{spt}[u(\cdot, t)]$, is contained in $[l(t), r(t)]$ for all $t > 0$. Moreover, we will show below that both $-l(t)$ and $r(t)$ are non-decreasing Lipschitz functions, as in PME, and so the following limits exist:

$$l^\infty := \lim_{t \rightarrow \infty} l(t), \quad r^\infty := \lim_{t \rightarrow \infty} r(t).$$

With the global existence in hand, it is possible to study the asymptotic behavior for the solutions. For RDEs, the equations are uniform parabolic and the strong maximum principle is applicable. So, any solution is a classical one and the convergence of u to its ω -limit is taken in the topology of $C_{\text{loc}}^{2,1}(\mathbb{R})$. For our RPME, however, we have very weak solutions with $C^{\alpha, \frac{\alpha}{2}}$ bounds in any compact domain. A nonnegative solution is not necessarily positive and classical. Hence, the convergence of a solution to its limit is first considered in the topology of $C_{\text{loc}}(\mathbb{R})$ or $L_{\text{loc}}^\infty(\mathbb{R})$. In addition, if $u(x, t_n) \rightarrow w(x) > 0$ as $t_n \rightarrow \infty$ in some domain J , then $u > 0$ in J by the positivity persistence and so it is a classical solution. Thus, the convergence $u(x, t_n) \rightarrow w(x)$ holds in C^2 topology in any compact subdomain of J . In summary, if no other specification, throughout this paper we use the following definition for ω -limits of u :

$$\lim_{n \rightarrow \infty} u(x, t_n) = w(x) \text{ means } \begin{cases} u(x, t_n) \rightarrow w(x) & \text{in } C_{\text{loc}}(\mathbb{R}) \text{ topology,} \\ u(x, t_n) \rightarrow w(x) & \text{in } C_{\text{loc}}^2(J) \text{ topology} \\ & \text{if } w(x) > 0 \text{ in } J. \end{cases} \quad (1.6)$$

In some cases, our problem may have a *ground state solution*, which means one of the following two types of nonnegative stationary solutions of (CP):

- *Type I ground state solution* (see Figure 2): $U_0(x) \in C^2(\mathbb{R})$ is an even stationary solution of (CP) with

$$U_0(x) > 0 > U'_0(x) \text{ for } x > 0, \quad U_0(+\infty) \in [0, 1);$$

- *Type II ground state solution* (see Figure 3): there exist a positive integer k , $z_i \in \mathbb{R}$, $i = 1, 2, \dots, k$, and $L > 0$ with $z_i + 2L \leq z_{i+1}$, $i = 1, 2, \dots, k-1$, such that

$$\mathcal{U}(x) = U(x - z_1) + U(x - z_2) + \dots + U(x - z_k), \quad x \in \mathbb{R}, \quad (1.7)$$

where $U(x) \in C(\mathbb{R}) \cap C^2(\mathbb{R} \setminus \{\pm L\})$ is an even stationary solution of problem (CP) with

$$\begin{aligned} U(x) &> 0 > U'(x) \quad \text{for } x \in (0, L), \\ U(x) &= 0 \quad \text{for } x \geq L, \\ (U^{m-1})'(L) &= 0. \end{aligned} \quad (1.8)$$




Fig. 2. Type I ground state solution.




Fig. 3. Type II ground state solution.

A Type I ground state solution is positive on the whole $\mathbb{R}$, which is the analog of that in bistable RDEs (cf. [14, 32]). For our RPME, this kind of solution exists if, in particular, f is a bistable nonlinearity with

$$f(u) = -\lambda u^\alpha(1 + o(1)) \quad \text{as } u \rightarrow 0 +, \quad (1.9)$$

for some $\lambda > 0$ and $\alpha \geq m$ (see Lemma 6.1). The component U in Type II ground state solution has a compact support, which is different from Type I. Such a solution exists when (1.9) holds for $0 < \alpha < m$ (see Lemma 6.1). The last equality in (1.8) is used to indicate that $U(x)$ is a stationary solution not only of the RPME but also of the Cauchy problem (CP). In fact, a necessary and sufficient condition ensuring that a stationary solution $\bar{u}$ of RPME with compact support $[-b, b]$ is also a stationary solution of the Cauchy problem (CP) is that the waiting times at its boundaries $\pm b$ are infinite, which requires that $(\bar{u}^{m-1})(\pm b) = 0$ (see Lemma 2.3 below).

Our first main theorem is a general convergence result, which says that any nonnegative bounded global solution of (CP) converges to a nonnegative zero of f or a ground state solution.

Theorem 1.1 (General convergence theorem). *Assume (F) and (I). Let $u(x, t)$ be a bounded, nonnegative, time-global solution of (CP). Then $u(\cdot, t)$ converges as $t \rightarrow \infty$, in the sense of (1.6), to a stationary solution of (CP), which is of one of the following types:*

- (i) a nonnegative zero of f ;
- (ii) $U_0(x - z_0)$ for some $z_0 \in [-b, b]$ and some Type I ground state solution U_0 ;
- (iii) $\mathcal{U}(x)$ as in (1.7) for $k \geq 1$ points $z_1, z_2, \dots, z_k \in [-b, b]$. In addition, l^∞, r^∞ are bounded in this case.

This theorem is a fundamental result and can be used conveniently to study various RPMEs. For example, when the reaction term in RPME is a typical monostable, bistable or combustion one, it is easy to clarify all of the nonnegative stationary solutions. Hence, using the general convergence result in the previous theorem, we can give a rather complete analysis on the asymptotic behavior for the solutions of (CP) with these reactions.

To obtain the above convergence result, our main tool is the so-called *zero number argument* in the version of porous medium equation, which can be used to describe the intersection points between two solutions. The main difficulty in establishing the argument is brought by the existence of free boundaries and the degeneracy of diffusion here, which is a significant difference compared with the classical heat equation. In particular, the intersection number may increase. For the convenience of the readers, we present the intersection number properties as a theorem below. Let u_1, u_2 be two solutions of (CP) with initial data in $\mathcal{X}$, then they are zero outside of their supports. It is a little confusing to define the intersection number between u_1 and u_2 outside of their supports. Instead, we denote by $Z_0(t)$ the number of their positive intersection points, that is,

$$Z_0(t) := \#\{x \in \mathbb{R} \mid u_1(x, t) = u_2(x, t) > 0\}, \quad t \geq 0, \quad (1.10)$$

where $\#S$ denotes the number of the elements in a set S .

Theorem 1.2 (Intersection number properties). *Assume (F). Let u_1, u_2 be bounded, non-negative, time-global solutions of (CP) with initial data in $\mathcal{X}$. If $Z_0(0) < \infty$, then $Z_0(t) < \infty$ for all $t > 0$, and there exists a time $T \geq 0$ such that*

$$Z_0(t) \text{ decreases for } t > T.$$

More specifically, there exist $\{t_j\}_{0 \leq j \leq k}$ with $1 \leq k \in \mathbb{N}$ and $0 = t_0 < t_1 < \dots < T := t_{k-1} < t_k = +\infty$ such that

- (i) $Z_0(t)$ decreases if $t_j < t < t_{j+1}$, $j = 0, 1, \dots, k-1$;
- (ii) $Z_0(t)$ strictly increases at $t = t_j$, $j = 1, \dots, k-1$, in the following sense:

$$\lim_{t \rightarrow t_j + 0} Z_0(t) > \lim_{t \rightarrow t_j - 0} Z_0(t).$$

The last property (that is, $Z_0(t)$ increases at t_j) happens in particular when a component where $u_1 > 0$ meets a component where $u_2 > 0$ while their boundaries approaching to each other (see more in Lemma 3.5). As we have mentioned above, this is significantly different from the classical zero number diminishing properties. Such a phenomenon happens only in PME where a solution has free boundaries but not in RDEs.

In order to describe the threshold of transition solutions, as in [12, 14, 32], we introduce a one-parameter family of initial data $u_0 = \phi_\sigma(x)$ satisfying the following conditions:

- (Φ_1) $\phi_\sigma \in \mathcal{X}$ for every $\sigma > 0$, and the map $\sigma \mapsto \phi_\sigma$ is continuous from $(0, \infty)$ to $L^\infty(\mathbb{R})$;
- (Φ_2) if $0 < \sigma_1 < \sigma_2$, then $\phi_{\sigma_1} \leq \phi_{\sigma_2}$, $\neq$;
- (Φ_3) $\lim_{\sigma \rightarrow 0} \|\phi_\sigma\|_{L^\infty(\mathbb{R})} = 0$.

Theorem 1.3. Assume $f \in C^2$ is a monostable reaction term satisfying (1.2). Let $u_\sigma(x, t)$ be the very weak solution of (CP) with $u_0 = \phi_\sigma(x)$ satisfying (Φ_1) – (Φ_3) . Then the hair-trigger effect holds: for all $\sigma > 0$, we have

$$u_\sigma(\cdot, t) \rightarrow 1 \quad \text{as } t \rightarrow \infty, \quad \text{in the topology of } L_{\text{loc}}^\infty(\mathbb{R}).$$

Remark 1.4. Theorem 1.3 is a direct application of Theorem 1.1 since the structure of the ω -limit set is rather simple in this case (see details in Section 4). Through a different proof, in [19] Gárriz obtained a deeper result on the hair-trigger effect for more general monostable RPME in N -dimensional space, where f satisfies $\liminf_{u \rightarrow 0+0} \frac{f(u)}{u^{\frac{m}{m+N}}} > 0$.

In the combustion case, we have the following result.

Theorem 1.5. Assume $f \in C^2$ is a combustion reaction term satisfying (1.4). Let $u_\sigma(x, t)$ be the very weak solution of (CP) with $u_0 = \phi_\sigma$ satisfying (Φ_1) – (Φ_3) . Then there exists $0 < \sigma_* \leq \infty$ such that the following trichotomy result holds:

(i) Spreading happens for $\sigma > \sigma_*$:

$$u_\sigma(\cdot, t) \rightarrow 1 \quad \text{as } t \rightarrow \infty, \quad \text{in the topology of } L_{\text{loc}}^\infty(\mathbb{R}).$$

(ii) Vanishing happens for $0 < \sigma < \sigma_*$:

$$u_\sigma(\cdot, t) \rightarrow 0 \quad \text{as } t \rightarrow \infty, \quad \text{in the topology of } L^\infty(\mathbb{R}).$$

(iii) Transition case for $\sigma = \sigma_*$:

$$u_\sigma(\cdot, t) \rightarrow \theta \quad \text{as } t \rightarrow \infty, \quad \text{in the topology of } L_{\text{loc}}^\infty(\mathbb{R}).$$

Moreover, for large t , u_σ has exactly two free boundaries $l(t) < r(t)$ satisfying

$$-l(t), r(t) = 2y_0\sqrt{t}[1 + o(1)] \quad \text{as } t \rightarrow \infty, \quad (1.11)$$

for some $y_0 \in (0, \theta^{\frac{m-1}{2}})$.

In the bistable case, if we further assume that $f'(0) < 0$, then problem (CP) has Type II ground state solutions, and so we have the following trichotomy result.

Theorem 1.6. Assume $f \in C^2$ is a bistable reaction term satisfying (1.3) and $f'(0) < 0$. Let $u_\sigma(x, t)$ be the very weak solution of (CP) with $u_0 = \phi_\sigma$ satisfying (Φ_1) – (Φ_3) . Then there exist $0 < \sigma_* \leq \sigma^* \leq \infty$ such that the following trichotomy holds:

(i) Spreading happens for $\sigma > \sigma^*$:

$$u_\sigma(\cdot, t) \rightarrow 1 \quad \text{as } t \rightarrow \infty, \quad \text{in the topology of } L_{\text{loc}}^\infty(\mathbb{R}).$$

(ii) Vanishing happens for $0 < \sigma < \sigma_*$:

$$u_\sigma(\cdot, t) \rightarrow 0 \quad \text{as } t \rightarrow \infty, \quad \text{in the topology of } L^\infty(\mathbb{R}).$$

(iii) *Transition case for $\sigma_* \leq \sigma \leq \sigma^*$:*

$$u_\sigma(\cdot, t) \rightarrow \mathcal{U}(x) \quad \text{as } t \rightarrow \infty, \quad \text{in the topology of } L^\infty(\mathbb{R}),$$

where $\mathcal{U}(x)$ is a Type II ground state solution defined as (1.7) for $k (\geq 1)$ points $z_1, z_2, \dots, z_k \in [-b, b]$.

Remark 1.7. (1) Under some additional conditions, we can show the sharpness of the transition solution: $\sigma_* = \sigma^*$ (see details in Lemma 6.3).

(2) The behavior of a transition solution in Theorem 1.6(iii) is quite different from that of the bistable RDE, where the transition solution converges to a Type I ground state solution which is positive on the whole $\mathbb{R}$.

(3) When spreading happens, the estimate on the propagation of the free boundaries has been shown in [19].

The main features of the paper are as follows:

- (1) We give out a complete classification for the asymptotic behavior of the solutions of (CP) with general nonlinear term.
- (2) We establish the zero number argument for RPME with solid proof.

This paper is arranged as follows. In Section 2, we first present some results on the well-posedness and the a priori estimates, which are not entirely new and some details are postponed to appendix. We also present sufficient conditions ensuring the waiting time of a free boundary to be 0, finite or infinite, specify the relationship between the stationary solutions of the RPME and those of the Cauchy problem, and derive the monotonicity and the Darcy law for the free boundaries. In Section 3, we study the monotonicity of the solution outside of the initial support, show the intersection number properties, and then use it, as well as the classical zero number argument, to prove the general convergence result. In Section 4, we consider the RPME with monostable reaction term and prove the hair-trigger effect. In Section 5, we consider the RPME with combustion reaction term: we present all possible nonnegative stationary solutions, prove a trichotomy result with a sharp transition for the asymptotic behavior. We also present a precise estimate for the propagation speed of the free boundaries of the transition solutions. In Section 6, we consider the RPME with bistable reaction term: we present all possible nonnegative stationary solutions, specify Types I and II ground state solutions, prove a trichotomy result with transition for the asymptotic behavior. In Section 7, we give a sufficient condition for the complete vanishing phenomena. Finally, for the convenience of the readers, we present in Appendix A the a priori estimates, well-posedness and other important properties for PME with general reaction terms. They are not entirely new but not well collected in the literature.

We collect some notations and symbols which will be used in this paper.

- For $\psi \in C(\mathbb{R})$, we use $\text{spt}[\psi(\cdot)]$ to denote its support.
- The expression $v(x) \rightarrow b$ as $x \rightarrow x_0 - 0$, or $\lim_{x \rightarrow x_0 - 0} v(x) = b$ means the left limit $\lim_{x < x_0, x \rightarrow x_0} v(x) = b$. The right limit is denoted similarly.

- $D_x^- v(x_0) = \lim_{x \rightarrow x_0-0} \frac{v(x)-v(x_0)}{x-x_0}$ denotes the left derivative of v at x_0 . The right derivative $D_x^+ v(x_0)$ is denoted similarly.
- $0 < 1 - x \ll 1$ means that $x < 1$ and $1 - x$ is sufficiently small.
- $a(x) \sim b(x)$ as $x \rightarrow x_0 \pm 0$ means that $a(x) = b(x)(1 + o(1))$ as $x \rightarrow x_0 \pm 0$.

2. Preliminaries

In this section, we first collect some basic results on the a priori estimates and the well-posedness. Then we specify the waiting time, stationary solutions, the regularity and the Darcy law of the free boundaries, which turn out to be not direct consequences of the corresponding results for PME.

2.1. Well-posedness and a priori estimates

As we have mentioned above, the existence and uniqueness of (CP) have been studied a lot by many authors (see, for example, [5, 9, 27, 30] etc.). In addition, by the assumption $f(u) < 0$ for $u > 1$ in (F) and by the comparison principle, it is easy to obtain

$$0 \leq u(x, t) \leq M'_0 := \max\{1, \|u_0\|_{L^\infty(\mathbb{R})}\}, \quad x \in \mathbb{R}, t > 0. \quad (2.1)$$

The persistence of positivity, the finite spreading speed and the a priori estimates were also studied widely, especially for the PME without reactions/sources or RPMEs with polynomial or logistic reactions. For our equation with a general reaction term, however, these properties seem not conveniently collected in literature, though they are not entirely new. We collect some of them here and present some detailed proof in the appendix.

1. *Positivity persistence.* If $u(x, t_0) \geq \varepsilon > 0$ in a neighborhood of $x_0 \in \mathbb{R}$, then $u(x_0, t) > 0$ for all $t \geq t_0$ (see Proposition A.2). Thus, u is classical near the line $\{(x_0, t) \mid t > t_0\}$.

2. *Finite propagation speed.* The solution has a left-most free boundary $l(t)$ and a right-most free boundary $r(t)$ (also called interfaces) which propagate in finite speed, that is, $\text{spt}[u(\cdot, t)] \subset [l(t), r(t)] \subset [-\bar{s}(t), \bar{s}(t)]$ with

$$\bar{s}(t) := O(1)(t+1)^{\frac{1}{2}} e^{\frac{K(m-1)(t+1)}{2}}, \quad t > 0$$

(see Proposition A.4).

In a gas flow problem through a porous medium, u denotes the density of the gas, and

$$v(x, t) := \frac{m}{m-1} [u(x, t)]^{m-1}$$

represents the *pressure* of the gas. In many cases, it is convenient to consider v rather than u . Using v , (CP) is converted into the following problem:

$$\begin{cases} v_t = (m-1)vv_{xx} + v_x^2 + g(v), & x \in \mathbb{R}, t > 0, \\ v(x, 0) = v_0(x) := \frac{m}{m-1} u_0^{m-1}(x), & x \in \mathbb{R}, \end{cases} \quad (\text{pCP})$$

with

$$g(v) := m \left(\frac{(m-1)v}{m} \right)^{\frac{m-2}{m-1}} f \left(\left(\frac{(m-1)v}{m} \right)^{\frac{1}{m-1}} \right).$$

For simplicity, we call the equation in (pCP) as *pRPME*. Since f is Lipschitz with number K , we have

$$|g(v)| \leq K(m-1)v, \quad v \geq 0.$$

Since u is bounded as in (2.1), we see that v is also bounded,

$$0 \leq v(x, t) \leq M_0 := \max \left\{ \frac{m}{m-1}, \|v_0\|_{L^\infty} \right\}, \quad x \in \mathbb{R}, t > 0.$$

3. *A priori estimates.* We have a priori estimates for v_x , v_{xx} and v_t as follows. If $v > 0$ in $R := (a, b) \times (0, T]$, then for any $0 < \delta < \frac{b-a}{2}$, $\tau < T$, there holds

$$|v_x(x, t)| \leq M_1(m, f, \delta, \tau), \quad (x, t) \in [a + \delta, b - \delta] \times [\tau, T].$$

There exist $\tau_0 = \tau_0(m, v_0)$ and $C = C(m, v_0)$ such that

$$v_{xx}(x, t) \geq -C(t + \tau_0), \quad x \in \mathbb{R}, t \in (0, T],$$

in the sense of distributions in $\mathbb{R} \times (0, T]$. For any $\tau > 0$, there holds

$$-C_3 t - C_4 \leq v_t(x, t) \leq C_1 t + C_2, \quad x \in \mathbb{R}, t \geq \tau,$$

where C_i depends on m , f and v_0 , while C_2 depends also on $v_t(x, \tau)$.

In addition, we have the following locally uniform Hölder estimates (cf. [10, 11] and [30, Theorem 7.18]): for any $\tau > 0$ and any compact domain $D \subset \mathbb{R} \times (\tau, \infty)$, there are positive constants C and α , both depending only on M'_0 , m and τ , such that

$$\|v\|_{C^{\alpha, \frac{\alpha}{2}}(D)} \leq C. \quad (2.2)$$

Note that the Hölder estimates in [10, 11, 30] are given for u , but it is easy to convert to that for v .

2.2. Waiting times of the free boundaries and compactly supported stationary solutions

In PME and RPMEs, the degeneracy of the equation not only causes finite propagation and free boundaries. It may even occur that a free boundary is stationary for a while if the mass distribution near the border of the support is very small.

Let $x_0 \in \mathbb{R}$ be a zero of $u_0(x)$. Define the *waiting time* at x_0 by

$$t^*(x_0) := \inf\{t > 0 \mid u(x_0, t) > 0\}.$$

For the one-dimensional PME, it is known that $t^*(x_0)$ can be 0 or a positive number, depending on the decay rate of u_0 near x_0 (cf. [30, §15.3], [31]). For our equation with a source, the situation is more complicated since positive reactions can promote the propagation while the negative reactions will restrain it.

We will show the following results:

- $t^*(x_0) = 0$ when the pressure v_0 near x_0 is bounded from below by a linear function.
- $t^*(x_0) > 0$ when the pressure v_0 lies below a quadratic function near x_0 . In addition, $t^*(x_0) < \infty$ if the reaction term $f \geq 0$ near 0 or if f is negative but v_0 lies above another quadratic function with big coefficient.
- $t^*(x_0) = \infty$ when the pressure v_0 is less than a quadratic function near x_0 and $f(u) \leq -K'u$ for some large $K' > 0$ and $0 \leq u \ll 1$.

For simplicity, we assume 0 is a zero of v_0 and consider its waiting time.

1. The case $t^*(0) = 0$.

Lemma 2.1. Assume (F) and (I). Assume $v_0(0) = 0$ and

$$v_0(x) \geq \rho x, \quad 0 \leq x \leq r_0,$$

for some $\rho > 0$, $r_0 > 0$, then $t^*(0) = 0$.

Proof. Set

$$a := \frac{\pi}{r_0}, \quad h := \frac{\rho r_0}{\pi}, \quad \mu := K(m-1) + mha^2, \quad b(t) := \frac{ha^2}{\mu}(e^{-\mu t} - 1),$$

and, for $t \geq 0$, define

$$\underline{v}(x, t) := \begin{cases} he^{-\mu t} \sin(ax - b(t)), & \frac{b(t)}{a} \leq x \leq \frac{b(t) + \pi}{a}, \\ 0, & \text{otherwise.} \end{cases}$$

By a direct calculation, we have

$$\underline{v}(x, 0) \leq v_0(x), \quad x \in \mathbb{R},$$

and, with $z := ax - b(t)$,

$$\begin{aligned} \mathcal{N}(\underline{v}) &:= \underline{v}_t - (m-1)\underline{v} \underline{v}_{xx} - \underline{v}_x^2 - g(\underline{v}) \\ &\leq \underline{v}_t - (m-1)\underline{v} \underline{v}_{xx} - \underline{v}_x^2 + K(m-1)\underline{v} \\ &= he^{-\mu t}[-\mu \sin z + (m-1)a^2 he^{-\mu t} \sin^2 z + K(m-1) \sin z \\ &\quad - b'(t) \cos z - a^2 he^{-\mu t} \cos^2 z] \\ &\leq a^2 h^2 e^{-\mu t}[-\sin z + e^{-\mu t} \cos z - e^{-\mu t} \cos^2 z] \\ &\leq a^2 h^2 e^{-2\mu t}[-\sin z + \cos z - \cos^2 z] \\ &= -\frac{1}{2}a^2 h^2 e^{-2\mu t}[(1 - \cos z)^2 + 2 \sin z - \sin^2 z] \\ &\leq 0, \quad \frac{b(t)}{a} \leq x \leq \frac{b(t) + \pi}{a}, \quad t \geq 0. \end{aligned}$$

Therefore, $\underline{v}(x, t)$ is a subsolution, and so by the comparison result (see, for example, [30, Theorem 6.5]), we have $v(0, t) \geq \underline{v}(0, t) > 0$ for small $t > 0$. Then $t^*(0) = 0$ since $b'(t) < 0$ for all $t \geq 0$. ■

2. The case $t^*(0) \in (0, \infty)$.

Lemma 2.2. Assume (F) and (I). Assume $v_0(0) = 0$,

$$v_0(x) \geq A_2 x^2, \quad 0 \leq x \leq r_0,$$

for some $r_0 > 0$ and some A_2 satisfying

$$A_2 > A_2(m, K) := \frac{K(m-1)}{2\sigma^{\frac{m}{m+1}} - 2\sigma}, \quad \sigma := \left(\frac{m}{m+1}\right)^{m+1},$$

and

$$v_0(x) \leq A_1 x^2, \quad x \geq 0,$$

for some $A_1 > A_2$. Then $t^*(0) \in (0, \infty)$.

Proof. First, we show that $t^*(0) > 0$. Set

$$a := \frac{(m-1)K}{2(m+1)A_1 + (m-1)K}, \quad T_1 := \frac{a}{(m-1)K}, \quad C_1 := \frac{1-a}{2(m+1)},$$

and define

$$\bar{v}(x, t) := \frac{C_1 x^2}{T_1 - t}, \quad x \in \mathbb{R}, \quad 0 \leq t < T_1.$$

Then $\bar{v}(x, 0) = A_1 x^2 \geq v_0(x)$, and

$$\begin{aligned} \mathcal{N}(\bar{v}) &:= \bar{v}_t - (m-1)\bar{v}\bar{v}_{xx} - \bar{v}_x^2 - g(\bar{v}) \\ &\geq \bar{v}_t - (m-1)\bar{v}\bar{v}_{xx} - \bar{v}_x^2 - K(m-1)\bar{v} \\ &= \frac{C_1 x^2}{(T_1 - t)^2} [1 - 2(m+1)C_1 - K(m-1)(T_1 - t)] \\ &\geq 0, \quad x \in \mathbb{R}, \quad t \in [0, T_1). \end{aligned}$$

By comparison, we have $v(x, t) \leq \bar{v}(x, t)$ in $t \in [0, T_1)$. Hence, the waiting time of v at 0 is not smaller than $T_1 = T_1(m, A_1, K)$.

Next, we prove $t^*(0) < \infty$. Set, for some $\delta \in (0, 1)$,

$$\tau := \frac{\sigma}{K(m^2 - 1)}, \quad C_2 := \frac{\delta r_0^2}{2(m+1)\tau^{\frac{1}{2(m+1)}}}, \quad x_1 := \left[\frac{C_2(1 + 2(m+1)A_2\tau)}{A_2\tau^{\frac{m}{m+1}}} \right]^{\frac{1}{2}}.$$

Then by the choice of A_2 , we have

$$\begin{aligned} \frac{x_1^2}{C_2} &= \frac{1}{A_2\tau^{\frac{m}{m+1}}} + \frac{2(m+1)\tau}{\tau^{\frac{m}{m+1}}} \\ &< 2 \frac{\sigma^{\frac{m}{m+1}} - \sigma}{K(m-1)\tau^{\frac{m}{m+1}}} + 2 \frac{(m+1)\tau}{\tau^{\frac{m}{m+1}}} \\ &= 2 \frac{[K(m^2 - 1)]^{\frac{m}{m+1}}}{K(m-1)}. \end{aligned}$$

Define

$$\underline{v}(x, t) := \begin{cases} C_2(t + \tau)^{-\frac{m}{m+1}} - \frac{(x - x_1)^2}{2(m+1)(t + \tau)}, & x \in [\underline{l}(t), \underline{r}(t)], \ 0 \leq t < T_2, \\ 0, & x < \underline{l}(t) \text{ or } x > \underline{r}(t), \ 0 \leq t < T_2, \end{cases}$$

where

$$\underline{l}(t) := x_1 - \sqrt{2(m+1)C_2}(t + \tau)^{\frac{1}{2(m+1)}}, \quad \underline{r}(t) := x_1 + \sqrt{2(m+1)C_2}(t + \tau)^{\frac{1}{2(m+1)}}$$

and

$$T_2 := \frac{1 - \sigma}{K(m^2 - 1)}.$$

We now show that $\underline{v}(x, t)$ is a subsolution in $0 \leq t < T_2$. In fact, by our construction it is easy to verify that

$$A_2 x^2 \geq \underline{v}(x, 0), \quad x \in [\underline{l}(0), \underline{r}(0)] \subset [0, r_0],$$

provided that $\delta \in (0, 1)$ is small, and the equality holds at

$$x = \hat{x} := \frac{x_1}{2(m+1)\tau A_2 + 1}.$$

In addition, for $x \in [\underline{l}(t), \underline{r}(t)]$, $0 \leq t \leq T_2$, by a direct calculation we have

$$\begin{aligned} \mathcal{N}(\underline{v}) &:= \underline{v}_t - (m-1)\underline{v}\underline{v}_{xx} - \underline{v}_x^2 - g(\underline{v}) \\ &\leq \underline{v}_t - (m-1)\underline{v}\underline{v}_{xx} - \underline{v}_x^2 + K(m-1)\underline{v} \\ &\leq \frac{C_2}{(t + \tau)^{\frac{2m+1}{m+1}}} \left[\frac{-1}{m+1} + K(m-1)(t + \tau) \right] \leq 0. \end{aligned}$$

This implies that $\underline{v}$ is a subsolution at least in the time interval $[0, T_2]$.

By the choice of the above parameters, we have

$$\frac{\underline{l}(0)}{\sqrt{2C_2}} = \left(\frac{1 + 2(m+1)A_2\tau}{A_2\tau^{\frac{m}{m+1}}} \right)^{\frac{1}{2}} - [2(m+1)\tau^{\frac{1}{m+1}}]^{\frac{1}{2}} > 0$$

and

$$\begin{aligned} \frac{\underline{l}(T_2)}{\sqrt{2C_2}} &< \left(\frac{[K(m^2 - 1)]^{\frac{m}{m+1}}}{K(m-1)} \right)^{\frac{1}{2}} - [(m+1)(T_2 + \tau)^{\frac{1}{m+1}}]^{\frac{1}{2}} \\ &= \left(\frac{[K(m^2 - 1)]^{\frac{m}{m+1}}}{K(m-1)} \right)^{\frac{1}{2}} - \left(\frac{m+1}{[K(m^2 - 1)]^{\frac{1}{m+1}}} \right)^{\frac{1}{2}} = 0. \end{aligned}$$

This means that before $t = T_2$, the left free boundary $\underline{l}(t)$ of $\underline{v}(x, t)$ moves leftward from a positive point to a negative one. Then, $\underline{v}(0, t)$ becomes positive at T_2 , and so does $v(0, t)$. This proves our lemma. $\blacksquare$

The case $t^*(0) > 0$ still includes two subcases: $t^*(0) \in (0, \infty)$ and $t^*(0) = \infty$. In the PME, it is known that only the former subcase is possible, which is called the *hole-filling* phenomenon (see, for example, [30, Theorem 15.15]). For our RPME with a reaction term, however, the latter is also possible.

3. *The case where $t^*(0) = \infty$ and compactly supported stationary solutions.* To discuss the possibility of infinite waiting time, we first specify the compactly supported stationary solutions, including sufficient and necessary conditions, examples and uniqueness. We remark that the following two concepts should be distinguished:

- compactly supported stationary solutions of the equation pRPME, and
- compactly supported stationary solutions of the Cauchy problem (pCP).

The latter means all compactly supported functions $\bar{v}$ such that the solution $v(x, t; \bar{v})$ of (pCP) with initial data $v_0 = \bar{v}$ satisfies $v(x, t; \bar{v}) \equiv \bar{v}$ for all $x \in \mathbb{R}$ and $t \geq 0$.

Lemma 2.3. *Assume $\bar{v}$ is a continuous nonnegative stationary solution of pRPME with support $[l, r]$ and $\bar{v}(x) > 0$ in (l, r) . Then*

- $\bar{v}$ is also a stationary solution of (pCP) if and only if $t^*(l), t^*(r) \geq \tau$ for some $\tau > 0$;
- a sufficient condition for (i) is $\bar{v}(x) \leq A(x-l)^2$ and $\bar{v}(x) \leq A(x-r)^2$ in (l, r) for some $A > 0$; a necessary condition for (i) is $\bar{v}'(l) = \bar{v}'(r) = 0$;
- $\bar{v}$ is not a stationary solution of (pCP) if $\bar{v}'(l) > 0$ or $\bar{v}'(r) < 0$;
- in any case, $\bar{v}$ is a time-independent (also called a stationary) very weak subsolution of (pCP), and $v(x, t; \bar{v}) \geq \bar{v}(x)$ for $x \in \mathbb{R}$, $t > 0$. The inequality is strict on $[l, r]$ when $\bar{v}$ is not a stationary solution of (pCP).

Proof. (i) If $t^*(l), t^*(r) \geq \tau$, then for all $t \in [0, \tau]$ we have

$$v(x, t; \bar{v}) = \bar{v}, \quad x \in \mathbb{R}, \quad l(t) = l, \quad r(t) = r.$$

Therefore, these equalities hold for all $t \geq 0$, and so $\bar{v}$ is a stationary solution of (pCP).

(ii) The sufficient condition follows from Lemma 2.2, and the necessary condition follows from Lemma 2.1.

(iii) The conclusion is a consequence of (ii).

(iv) Denote the corresponding density functions of $\bar{v}(x)$ and $v(x, t; \bar{v})$ by $\bar{u}(x)$ and $u(x, t; \bar{u})$, respectively. For any $T > 0$ and any test function $\varphi \in C_c^\infty(Q_T)$, $Q_T := \mathbb{R} \times (0, T)$, with $\varphi \geq 0$,

$$\begin{aligned} & \iint_{Q_T} [\bar{u}\varphi_t + \bar{u}^m\varphi_{xx}] dx dt \\ &= \int_{\mathbb{R}} \bar{u}(x)[\varphi(x, T) - \varphi(x, 0)] dx + \int_0^T \int_l^r (\bar{u}^m)_{xx}(x)\varphi(x, t) dx dt \\ & \quad + \int_0^T [(\bar{u}^m)_x(l)\varphi(l, t) - (\bar{u}^m)_x(r)\varphi(r, t)] dt \\ &\geq \int_{\mathbb{R}} \bar{u}(x)[\varphi(x, T) - \varphi(x, 0)] dx + \int_0^T \int_l^r (\bar{u}^m)_{xx}(x)\varphi(x, t) dx dt \\ &= \int_{\mathbb{R}} \bar{u}(x)[\varphi(x, T) - \varphi(x, 0)] dx - \iint_{Q_T} f(\bar{u}(x))\varphi(x, t) dx dt. \end{aligned}$$

So, $\bar{u}$ is a time-independent very weak supersolution of (CP), and then $u(x, t; \bar{u}) \geq \bar{u}(x)$ follows from the comparison result (cf. [30, Theorem 6.5]).

To show the strict inequality, we note that, in the domain (l, r) , both $\bar{v}$ and v are positive, and so the classical strong maximum principle is applied. On the other hand, by the conclusion in (i), the waiting times at l and r are zero since $\bar{v}$ is not a stationary solution of (pCP). So, $v(l, t; \bar{v}) > 0 = \bar{v}(l)$ and $v(r, t; \bar{v}) > 0 = \bar{v}(r)$. This proves the strict inequality on $[l, r]$. ■

Due to the strong maximum principle, the Cauchy problem of an RDE does not have compactly supported stationary solutions. For (pCP), however, such solutions are possible since the strong maximum principle is not true on the free boundaries. In fact, we will show in Section 5 (see details in Lemma 6.1) that if f is a bistable nonlinearity with

$$f(u) = -\lambda u^\alpha(1 + o(1)) \quad \text{as } u \rightarrow 0 + 0,$$

for some $\lambda > 0$ and $1 \leq \alpha < m$, then the bistable pRPME has nonnegative compactly supported stationary solution $\bar{v}$, which is a ground state solution satisfying

$$\bar{v}(x) > 0 \text{ in } (-L, L), \quad \bar{v}(x) = 0 \text{ for } |x| \geq L,$$

and

$$\bar{v}(x) \sim A(x \pm L)^{\frac{2(m-1)}{m-\alpha}} \quad \text{as } x \rightarrow \mp L \pm 0.$$

Since $\frac{2(m-1)}{m-\alpha} \geq 2$, by Lemma 2.3 (ii) we know that $\bar{v}$ is also a stationary solution of (pCP).

Comparing the ZKB solution of PME (named after Zel'dovich, Kompaneets and Barenblatt) with the solution of (pCP) for monostable and combustion reactions, it is easy to see that compactly supported stationary solutions do not exist in such problems. But they can exist for (pCP) with bistable reactions (see Lemma 6.1). We now show that it is unique when it exists.

Lemma 2.4. *Assume (F) and V is a compactly supported stationary solution of (pCP) with*

$$V(x) > 0 \text{ in } (-L, L), \quad V(x) = 0 \text{ for } |x| \geq L,$$

for some $L > 0$. Then

$$V(x) = V(-x) \text{ and } V'(x) < 0 \text{ in } (0, L), \quad V'(0) = V'(\pm L) = 0. \quad (2.3)$$

(So, V is a Type II ground state solution of (pCP).) Moreover, such compactly supported stationary solution is unique.

Proof. By the equation, we know that V is symmetric with respect to each positive critical point, so we have the symmetry and monotonicity in (2.3) and $V'(0) = 0$. The equality $V'(\pm L) = 0$ follows from Lemma 2.3.

We now prove the uniqueness of such compactly supported stationary solution. Set

$$w := \left(\frac{m-1}{m} V \right)^{\frac{1}{m-1}} V'. \quad (2.4)$$

Then $w(\pm L) = 0$. Hence, on the (V, w) -phase plane (cf. Sections 4.1 and 5.1 for more details), $V(x)$ corresponds to a homoclinic orbit starting and ending at $(0, 0)$. Using the equation in (pCP), we have

$$w' = -f\left(\left(\frac{m-1}{m}V\right)^{\frac{1}{m-1}}\right).$$

Multiplying this equation by (2.4) and integrating it over $[x, 0]$ for $x \in [-L, 0)$, we have

$$w^2(x) = F(V(x)) - F(V(0))$$

with

$$F(v) := -2 \int_0^v f\left(\left(\frac{m-1}{m}r\right)^{\frac{1}{m-1}}\right) \cdot \left(\frac{m-1}{m}r\right)^{\frac{1}{m-1}} dr.$$

Taking $x = -L$, we have $w^2(-L) = 0 = F(0) - F(V(0))$, and so $F(V(0)) = 0$. For $x \in (-L, 0)$, we have $V'(x) > 0$ and $w(x) > 0$. Thus,

$$F(v) > F(V(0)) = 0, \quad v \in (0, V(0)).$$

(This happens only if $f(u) < 0$ for $0 < u \ll 1$.) Therefore,

$$\Theta_1 := V(0)$$

is the smallest positive zero of $F(\cdot)$, and there exists $\Theta \in (0, \Theta_1)$ such that

$$\begin{aligned} f\left(\left(\frac{m-1}{m}\Theta\right)^{\frac{1}{m-1}}\right) &= g(\Theta) = 0, \\ f\left(\left(\frac{m-1}{m}v\right)^{\frac{1}{m-1}}\right) &> 0 \quad \text{for } v \in (\Theta, \Theta_1). \end{aligned} \tag{2.5}$$

Consequently, for $x < 0$, $V(x)$ is uniquely determined by the problem

$$\left(\frac{m-1}{m}V\right)^{\frac{1}{m-1}} V' = \sqrt{F(V(x))} \quad \text{for } x < 0, \quad V(0) = \Theta_1, \quad V'(0) = 0.$$

This proves the uniqueness of compactly supported stationary solution of (pCP), that is, the Type II ground state solution with one connected support. ■

We have given sufficient and necessary conditions, and uniqueness for compactly supported stationary solutions of the Cauchy problem (pCP). Clearly, they correspond to the analogs of the original problem (CP). Using them, we finally give some examples of initial data, which are not necessarily stationary solutions, such that the waiting times on their boundaries are infinite. For example, for a ground state solution V of the bistable PRPME in the previous lemma, we choose $v_0(x) \in C(\mathbb{R})$ such that

$$v_0(x) \leq V(x) \text{ in } \mathbb{R}, \quad v_0(x) > 0 \text{ in } (-L, L).$$

Then v_0 lies below V and has the same initial boundaries $\pm L$ as V . Since the boundaries $l(t)$ and $r(t)$ of $v(x, t; v_0)$ are monotone in time and since v always lies below V by comparison, we have $t^*(\pm L) = \infty$.

2.3. Lipschitz continuity, Darcy law and strict monotonicity

We now address the Lipschitz continuity, the Darcy law and the monotonicity for each free boundary. For definiteness, in this subsection, we work with the right free boundary $x = r(t)$, that is,

$$v(x, t) > 0 \text{ for } r(t) - 1 \ll x < r(t), \quad v(x, t) = 0 \text{ for } x \geq r(t).$$

We use one-sided partial derivatives in space and time,

$$D^\pm r(t) := r'(t \pm 0), \quad D_x^\pm v(x, t) = v_x(x \pm 0, t).$$

Also we use $r'(t)$ to denote the derivative of r when $D^+ r(t) = D^- r(t)$.

First, we recall the related results on PME.

Lemma 2.5 ([30, Chapter 15]). *Let v be the global solution of (pCP) with $g \equiv 0$, $r(t)$ be its right free boundary with waiting time $t^* \geq 0$. Then, for any $t_0 > t^*$, there exists a small $\varepsilon > 0$ such that*

(i) $r \in C^\infty((t_0 - \varepsilon, t_0 + \varepsilon))$, $v \in C^\infty(\mathcal{N}_\varepsilon)$ for

$$\mathcal{N}_\varepsilon = \{(x, t) \mid x \leq r(t) \text{ and } |(x - x_0, t - t_0)| \leq \varepsilon\}, \quad x_0 := r(t_0);$$

(ii) *there hold*

$$r'(t_0) = -D_x^- v(x_0, t_0) > 0, \quad r''(t_0) = m r'(t_0) D_{xx}^- v(x_0, t_0);$$

(iii) *for any positive integer j , there exists C_j depending on (x_0, t_0) , m , j , ε , v such that*

$$\left| \frac{\partial^j}{\partial x^j} v(x, t) \right| \leq C_j, \quad (x, t) \in \mathcal{N}_\varepsilon.$$

We will use this lemma to prove the important properties of the free boundaries, that is, the C^1 regularity and the Darcy law. Since we have no semiconvexity for the free boundary $r(t)$ of RPME as for the PME (cf. [30, §15.4.1]), we have to deal with the problem in a more complicated way.

We first show the left continuity of v_x .

Lemma 2.6. *Let v be the solution of (pCP), $r(t)$ be its right free boundary. Then*

$$D_x^- v(r(t), t) = \lim_{x \rightarrow r(t) - 0} v_x(x, t), \quad t > 0. \quad (2.6)$$

Proof. As we mentioned before, v_{xx} is bounded from below: $v_{xx} \geq -C_1 t - C_2$ for some $C_1, C_2 > 0$. Hence,

$$\tilde{v}(x, t) := v(x, t) + \frac{1}{2}(C_1 t + C_2)x^2$$

is a continuous and convex function of x for $t > 0$. So, for every $x_1 \in \mathbb{R}$, $t_1 > 0$, the function $\tilde{v}(\cdot, t_1)$ admits one-sided derivatives at x_1 : $D_x^+ \tilde{v}(x_1, t_1)$ and $D_x^- \tilde{v}(x_1, t_1)$ with $D_x^- \tilde{v}(x_1, t_1) \leq D_x^+ \tilde{v}(x_1, t_1)$, and the functions $D_x^\pm \tilde{v}(\cdot, t_1)$ are non-decreasing ones.

Consequently, $D_x^\pm v(x_1, t_1)$ exist with $D_x^- v(x_1, t_1) \leq D_x^+ v(x_1, t_1)$, and $D_x^\pm v(x, t_1) + (C_1 t_1 + C_2)x$ are non-decreasing in x . Note that, for small $\varepsilon > 0$, when $x \in I_1 := [r(t_1) - \varepsilon, r(t_1))$, $v(x, t_1)$ is smooth and so

$$v_x(x, t_1) + (C_1 t_1 + C_2)x \leq D_x^\pm v(r(t_1), t_1) + (C_1 t_1 + C_2)r(t_1).$$

The right-hand side is bounded since $D_x^- v(r(t_1), t_1) \leq 0 = D_x^+ v(r(t_1), t_1)$. So, $k_1 := \lim_{x \rightarrow r(t_1)-0} v_x(x, t_1)$ exists and $k_1 \leq D_x^- v(r(t_1), t_1)$. Conclusion (2.6) is proved if we can show the equality. By contradiction, we assume that $k_1 < D_x^- v(r(t_1), t_1) \leq 0$. Then there exists k_2 lying between them and

$$v_x(x, t_1) < k_2, \quad x \in I_1.$$

For any x_2, x_3 satisfying $r(t_1) - \varepsilon \ll x_3 < x_2 < r(t_1)$, we have

$$\frac{v(x_3, t_1) - v(x_2, t_1)}{x_3 - x_2} < k_2.$$

Taking the limit as $x_2 \rightarrow r(t_1) - 0$ and using the continuity of v , we obtain

$$\frac{v(x_3, t_1) - v(r(t_1), t_1)}{x_3 - r(t_1)} \leq k_2.$$

This implies that $D_x^- v(r(t_1), t_1) \leq k_2$, a contradiction. ■

Theorem 2.7. *Let v be the solution of (pCP), $r(t)$ be its right free boundary with waiting time $t^* \in [0, \infty)$. Then $r \in C^1((t^*, \infty))$ and the Darcy law holds:*

$$r'(t) = -D_x^- v(r(t), t) \geq 0, \quad t > t^*. \quad (2.7)$$

Moreover, $r'(t) \not\equiv 0$ in any interval in (t^*, ∞) .

Proof. We prove the Darcy law at any given time $t_0 > t^*$. Denote

$$x_0 := r(t_0) \quad \text{and} \quad \beta := -D_x^- v(x_0, t_0).$$

Step 1. We show that $D^+ r(t_0) = \beta$ when $\beta > 0$. Consider the following problem:

$$\begin{cases} \bar{v}_t = (m-1)\bar{v}\bar{v}_{xx} + \bar{v}_x^2 + K(m-1)\bar{v}, & x \in \mathbb{R}, t > 0, \\ \bar{v}(x, 0) = \bar{v}_0(x), & x \in \mathbb{R}, \end{cases}$$

where $\bar{v}_0(x)$ satisfies

$$\bar{v}'_0(x_0 - 0) = -\beta, \quad \bar{v}_0(x) \geq v(x, t_0) \quad \text{for } x \in \mathbb{R},$$

and $\bar{v}_0(x)$ is sufficiently smooth so that the right free boundary $\bar{r}(t)$ of $\bar{v}(\cdot, t)$ is smooth in $t \geq 0$. Then $\bar{v}$ is a supersolution of (pCP), and so

$$r(t + t_0) \leq \bar{r}(t), \quad t \geq 0.$$

To estimate $\bar{r}(t)$ precisely, we change the time variable from t to s by

$$s = S(t) := \frac{e^{K(m-1)t} - 1}{K(m-1)} \Leftrightarrow t = T(s) := \frac{\ln(1 + K(m-1)s)}{K(m-1)}$$

and define

$$V(x, s) := \frac{\bar{v}(x, T(s))}{1 + K(m-1)s},$$

then we have

$$\begin{cases} V_s = (m-1)V V_{xx} + V_x^2, & x \in \mathbb{R}, s > 0, \\ V(x, 0) = \bar{v}_0(x), & x \in \mathbb{R}. \end{cases}$$

By Lemma 2.5, for some small $\varepsilon_0 > 0$, the right free boundary $x = R(s) = \bar{r}(T(s))$ of $V(\cdot, s)$ satisfies

$$R'(s) = -D_x^- V(R(s), s), \quad |R''(s)| \leq \bar{C}_1, \quad 0 \leq s \leq S(\varepsilon_0),$$

or, equivalently,

$$\bar{r}'(t) = R'(s) \frac{ds}{dt} = -D_x^- \bar{v}(\bar{r}(t), t), \quad |\bar{r}''(t)| \leq \bar{C}, \quad 0 \leq t \leq \varepsilon_0.$$

Consequently, we have upper estimate for $r(t + t_0)$:

$$r(t + t_0) \leq \bar{r}(t) \leq \bar{r}(0) + \bar{r}'(0)t + \bar{C}t^2 \leq x_0 + \beta t + \bar{C}t^2, \quad 0 \leq t \leq \varepsilon_0. \quad (2.8)$$

Next, we consider the problem

$$\begin{cases} \underline{v}_t = (m-1)\underline{v}\underline{v}_{xx} + \underline{v}_x^2 - K(m-1)\underline{v}, & x \in \mathbb{R}, t > 0, \\ \underline{v}(x, 0) = \underline{v}_0(x), & x \in \mathbb{R}, \end{cases}$$

where $\underline{v}_0(x)$ satisfies

$$\underline{v}_0'(x_0 - 0) = -\beta, \quad \underline{v}_0(x) \leq v(x, t_0) \text{ for } x \in \mathbb{R},$$

and $\underline{v}_0(x)$ is sufficiently smooth so that the right free boundary $\underline{r}(t)$ of $\underline{v}(\cdot, t)$ is smooth in $t \geq 0$. Then $\bar{v}$ is a subsolution of (pCP) and, as above,

$$r(t + t_0) \geq \underline{r}(t) \geq x_0 + \beta t - \underline{C}t^2, \quad 0 \leq t \leq \varepsilon_0, \quad (2.9)$$

for some positive $\underline{C} > 0$. Combining with (2.8), we have $D^+r(t_0) = \beta$.

Step 2. We show that $D^+r(t) \not\equiv 0$ in any interval in (t^*, ∞) with positive measure. Note that this conclusion implies that $r(t)$ is strictly increasing in $t > t^*$. By the positivity persistence, $r(t)$ is increasing in $t > t^*$. Assume by contradiction that $D^+r(t) = 0$ for $t \in [t_1, t_2] \subset (t^*, \infty)$. We will derive

$$D^+r(t) = 0, \quad t \in [0, t_2], \quad (2.10)$$

which then contradicts the definition of t^* and the fact that $t_2 > t^*$.

To show (2.10), we suppose, on the contrary, that $t_3 := \inf\{\tilde{t} \mid D^+r(t) \equiv 0 \text{ in } [\tilde{t}, t_2]\} \in (0, t_1]$, and, by the continuity, there exists $\tau \in (t^*, t_3)$ such that

$$D^+r(\tau) > 0 \quad \text{and} \quad r(t_3) - r(\tau) < \Delta := \frac{P\pi}{\mu}(1 - e^{-\mu(t_2-t_1)}), \quad (2.11)$$

where $-P$ is the lower bound of v_{xx} : $v_{xx} > -P$ for $x \in \mathbb{R}$, $t \in [t^*, t_2]$, and $\mu := K(m-1) + mP\pi$. By Step 1, we have

$$0 < 2\rho := D^+r(\tau) = -D_x^-v(r(\tau), \tau).$$

Hence, for $\varepsilon := \frac{\rho}{P}$ we have

$$-D_x^-v(x, \tau) \geq \rho \quad \text{and} \quad v(x, \tau) \geq -\rho(x - r(\tau)), \quad r(\tau) - \varepsilon \leq x \leq r(\tau).$$

Denote

$$a := \frac{P\pi}{\rho}, \quad h := \frac{\rho^2}{P\pi}, \quad b(t) := \frac{P\pi}{\mu}(1 - e^{-\mu(t-\tau)}),$$

and using the subsolution $\underline{v}$ as in Lemma 2.1, we conclude that

$$r(t) \geq r(\tau) + b(t), \quad t > \tau. \quad (2.12)$$

In particular, at $t = t_3$ we have

$$r(t_3) = r(t_2) \geq r(\tau) + b(t_2) \geq r(\tau) + \Delta$$

that contradicts (2.11). This proves (2.10), and then the conclusion in this step is proved.

Step 3. We show that $D^-r(t_0) = \beta$ when $\beta > 0$. We assume that for some increasing sequence $\{t_n\}$ tending to t_0

$$\beta' := \lim_{n \rightarrow \infty} \frac{r(t_n) - r(t_0)}{t_n - t_0}, \quad D^+r(t_n) = \beta_n := -D_x^-v(r(t_n), t_n) > 0. \quad (2.13)$$

(The latter holds by Step 2.) If we can show that, for each of such time sequence $\{t_n\}$, the limit $\beta' = \beta$, then we obtain the conclusion $D^-r(t_0) = \beta$.

By contradiction, we assume $\beta' > \beta$. The reversed case is proved similarly. Then there exist a small $\delta > 0$ and a large n_0 such that

$$\frac{r(t_n) - r(t_0)}{t_n - t_0} \geq \beta + 2\delta, \quad n \geq n_0. \quad (2.14)$$

For any given $n \geq n_0$, using estimates (2.8) and (2.9) at the time t_n instead of t_0 , we see that, for some $\varepsilon_1 \in (0, \varepsilon_0)$ and some positive numbers $\underline{C}'$, $\bar{C}'$, there holds

$$r(t_n) + \beta_n t - \underline{C}'t^2 \leq r(t + t_n) \leq r(t_n) + \beta_n t + \bar{C}'t^2, \quad 0 \leq t \leq \varepsilon_1. \quad (2.15)$$

Taking $t = t_0 - t_n$ in the second inequality, we obtain

$$r(t_0) - r(t_n) \leq \beta_n(t_0 - t_n) + \bar{C}'(t_0 - t_n)^2.$$

Together with (2.14), we see that, when n is sufficiently large,

$$\beta_n \geq \beta + \delta.$$

Taking n large such that $\epsilon := t_0 - t_n \leq \frac{\epsilon_1}{2}$ and taking $t = 2\epsilon$ in (2.15), we have

$$\begin{aligned} r(t_0 + \epsilon) &\geq r(t_n) + 2\beta_n\epsilon + O(\epsilon^2) \\ &\geq r(t_n) + \beta_n\epsilon + \bar{C}'\epsilon^2 + (\beta + \delta)\epsilon + O(\epsilon^2) \\ &\geq r(t_0) + (\beta + \delta)\epsilon + O(\epsilon^2). \end{aligned}$$

Together with (2.8), we have

$$O(\epsilon^2) \geq \delta\epsilon + O(\epsilon^2).$$

This is a contradiction when ϵ is small or, equivalently, n is sufficiently large.

Step 4. We prove existence and continuity of $r'(t)$ at t_0 when $\beta = r'(t_0) > 0$. Combining Steps 1 and 3, we have

$$r'(t_0) = D^+r(t_0) = D^-r(t_0) = \beta \quad \text{when } \beta = -D_x^-v(x_0, t_0) > 0.$$

The proof in Step 3 also implies that, for any subsequence $\{n_i\}$ of $\{n\}$, $\beta_{n_i} \geq \beta + \delta$ is impossible, and so $\limsup_{n \rightarrow \infty} \beta_n \leq \beta$. In a similar way as in Step 3, one can show that $\beta_{n_i} \leq \beta - \delta$ is impossible, and so $\liminf_{n \rightarrow \infty} \beta_n \geq \beta$. Therefore, $r'(t)$ is left continuous at t_0 . On the other hand, if the time sequence $\{t_n\}$ in (2.13) is a decreasing one tending to t_0 from right side, then one can prove as above that $r'(t_n) \rightarrow \beta$, $n \rightarrow \infty$. This means that $r'(t)$ is right continuous at t_0 . Combining the above conclusions together, we get $r \in C^1$ at any time t_0 where $r'(t_0) > 0$.

Step 5. The case $\beta := -D_x^-v(x_0, t_0) = 0$. For any small $\varepsilon > 0$, we define

$$\tilde{v}_0(x) := v(x, t_0) + \varepsilon(x_0 - x), \quad x \in \mathbb{R},$$

and consider the equation of v for $t \geq t_0$, with initial data $v(x, t_0)$ replaced by $\tilde{v}_0(x)$. Then by Step 1, we have $D^+r(t_0) \leq -D_x^-\tilde{v}_0(x_0) = \varepsilon$. Since $\varepsilon > 0$ is arbitrary, we actually have $D^+r(t_0) = 0$. Now as the proof in Step 3, with $\beta > 0$ being replaced by $\beta = 0$, one can prove in a similar way that $D^-r(t_0) = 0$, and so $r'(t_0) = 0$. This implies that the Darcy law (2.7) continues to hold even if $\beta = 0$. Finally, the continuity of $r'(t)$ at t_0 where $r'(t_0) = 0$ can be shown as in Step 4.

This completes the proof of the theorem. ■

Remark 2.8. From this theorem, we know that for $t > t^*$, $r'(t) \geq 0$ and it is not identical zero in any time interval. Though we do not have the strong conclusion that $r'(t) > 0$ as in PME (cf. [30, Corollary 15.23]), the current properties are enough to continue the study for the asymptotic behavior of the solutions. For example, they are enough in the application of the zero number argument.

3. Zero number argument and general convergence result

In this section, we present the intersection number properties, prove that the solution is strictly monotone outside its initial support, and then prove the general convergence result.

3.1. Zero number and intersection number properties

A useful tool we will use in the qualitative study is the so-called zero number argument. For the convenience of the readers, we first prepare some basic results in this area. Consider

$$\eta_t = a(x, t)\eta_{xx} + b(x, t)\eta_x + c(x, t)\eta \quad (3.1)$$

in $E_0 := \{(x, t) \mid b_1(t) < x < b_2(t), t \in (t_1, t_2)\}$, where b_1 and b_2 are continuous functions in (t_1, t_2) . For each $t \in (t_1, t_2)$, denote by

$$\mathcal{Z}(t) := \#\{x \in \bar{J}(t) \mid \eta(\cdot, t) = 0\}$$

the number of zeros of $\eta(\cdot, t)$ in the interval $\bar{J}(t) := [b_1(t), b_2(t)]$. A point $x_0 \in \bar{J}(t)$ is called a *multiple zero* (or *degenerate zero*) of $\eta(\cdot, t)$ if $\eta(x_0, t) = \eta_x(x_0, t) = 0$. In 1988, Angenent [1] proved a zero number diminishing property, and in 1998, the conditions in [1] were weakened by Chen [8] for solutions with lower regularity. One of their results can be summarized as the following.

Proposition 3.1 (Zero number diminishing properties [1, 8]). *Assume the coefficients in (3.1) satisfies*

$$a, a^{-1}, a_t, a_x, b, c \in L^\infty. \quad (3.2)$$

Let η be a nontrivial $W_{p,\text{loc}}^{2,1}$ solution of (3.1). Further, assume that, for $i = 1, 2$, either

$$\eta(b_i(t), t) \neq 0, \quad t \in (t_1, t_2), \quad (3.3)$$

or

$$b_i(t) \in C^1 \quad \text{and} \quad \eta(b_i(t), t) \equiv 0, \quad t \in (t_1, t_2), \quad (3.4)$$

or

$$b_i(t) \in C^1 \quad \text{and} \quad \eta_x(b_i(t), t) \equiv 0, \quad t \in (t_1, t_2). \quad (3.5)$$

Then

- (i) $\mathcal{Z}(t)$ is finite and decreasing in $t \in (t_1, t_2)$;
- (ii) if $s \in (t_1, t_2)$ and $x_0 \in \bar{J}(s)$ is a multiple zero of $\eta(\cdot, s)$, then $\mathcal{Z}(s_1) > \mathcal{Z}(s_2)$ for all s_1, s_2 satisfying $t_1 < s_1 < s < s_2 < t_2$.

Note that, in the original results in [1, 8], the problem was considered in fixed intervals, that is, $b_i(t)$ are constants. Using our assumptions (3.3)–(3.5), for any time interval with small length, we can straighten the domain boundaries so that the zero number diminishing properties continue to hold. The boundedness assumption of a^{-1} in (3.2) is clearly a difficulty in using the zero number argument for RPMEs. Nevertheless, under some

additional conditions, we can prove a diminishing property for the number of intersection points between two solutions of (pCP), despite the degeneracy of the equation. The analogous results for RDEs were proved in [13, Lemma 2.4], but the proof for our current result is more complicated due to the degeneracy.

For $i = 1, 2$, let v_i be the solution of (pCP) with initial data in

$$\{\psi \in C(\mathbb{R}) \mid \psi(x) > 0 \text{ in } (l_i(0), r_i(0)), \psi(x) = 0 \text{ for } x \notin (l_i(0), r_i(0))\}.$$

Denote by $l_i(t)$ and $r_i(t)$ the left and right free boundaries of v_i , respectively. Write

$$l(t) := \max\{l_1(t), l_2(t)\}, \quad r(t) := \min\{r_1(t), r_2(t)\}, \quad t \geq 0.$$

Then in the case where $l(t) < r(t)$, the number of the positive intersection points between $v_1(\cdot, t)$ and $v_2(\cdot, t)$ in $\mathbb{R}$, which is denoted by $Z_0(t)$ as in (1.10), is the same as the number of intersection points in $J_0(t) := (l(t), r(t))$.

Theorem 3.2. *Assume (F), v_1, v_2 are given as above. Assume that $Z_0(0)$ is finite. Then there exists a time $\tau \in [0, +\infty]$ such that*

- (i) $Z_0(t)$ decreases for $t \in (0, \tau)$ and for $t \in (\tau, \infty)$;
- (ii) $\lim_{t \rightarrow \tau+0} Z_0(t) = 1$ and $\lim_{t \rightarrow \tau-0} Z_0(t) = 0$ if $\tau \in (0, \infty)$.

The proof of this theorem consists of the following two lemmas.

Lemma 3.3. *Assume (F), v_1, v_2 are given as above. Further, assume that $l(0) < r(0)$ and one of the following hypotheses holds:*

- (a) $l_1(0) \neq l_2(0), r_1(0) \neq r_2(0)$;
- (b) $Z_0(0) < \infty$.

Then

- (i) $Z_0(t)$ is decreasing in $t > 0$;
- (ii) $Z_0(t)$ is strictly decreasing when t passes through s if $v_1(\cdot, s) - v_2(\cdot, s)$ has a degenerate zero in $J_0(s)$ or if one interior zero moves to the boundary $l(t)$ or $r(t)$ as $t \rightarrow s - 0$.

Proof. We first prove the conclusions under the assumption (a). The main complexity is caused by the possibility that an interior intersection point moves up to the domain boundaries. Without loss of generality, we assume

$$l_1(t) < l_2(t) \text{ for small } t \geq 0, \quad r_1(t) < r_2(t) \text{ for all } t \geq 0.$$

Then the left boundaries $l_1(t)$ and $l_2(t)$ may meet after some time, but the right boundaries will not.

At the early stage, say $t \in [0, t_1]$ for some $t_1 > 0$, $l(t) = l_2(t) > l_1(t)$, and $\eta(x, t) := v_1(x, t) - v_2(x, t)$ satisfies a linear equation

$$\eta_t = a(x, t)\eta_{xx} + b(x, t)\eta_x + c(x, t)\eta, \quad x \in J_0(t) := (l(t), r(t)), \quad t \in (0, t_1], \quad (3.6)$$

with

$$a(x, t) := (m - 1)v_1(x, t), \quad b(x, t) := v_{1x}(x, t) + v_{2x}(x, t),$$

and

$$c(x, t) := (m - 1)v_{2xx}(x, t) + \begin{cases} \frac{g(v_1(x, t)) - g(v_2(x, t))}{v_1(x, t) - v_2(x, t)}, & v_1(x, t) \neq v_2(x, t), \\ 0, & v_1(x, t) = v_2(x, t). \end{cases}$$

By the continuity of v_1, v_2 ,

$$\eta(l(t), t) = v_1(l_2(t), t) \geq \rho, \quad \eta(r(t), t) = -v_2(r_1(t), t) \leq -\rho, \quad t \in [0, t_1],$$

for some small $\rho > 0$. Take two smooth curves $\hat{l}(t)$ and $\hat{r}(t)$ such that

$$\begin{aligned} l(t) < \hat{l}(t) < l(t) + 1, \quad r(t) - 1 < \hat{r}(t) < r(t), \\ \eta(x, t) \neq 0, \quad x \in [l(t), \hat{l}(t)] \cup [\hat{r}(t), r(t)], \quad t \in [0, t_1], \end{aligned} \quad (3.7)$$

and, for some $\rho_1 > 0$,

$$v_1(x, t), v_2(x, t) \geq \rho_1 \quad \text{in } E_1 := \{(x, t) \mid x \in \hat{J}(t) := [\hat{l}(t), \hat{r}(t)], t \in [0, t_1]\}.$$

Then both v_1 and v_2 are classical in E_1 , we can use the classical zero number diminishing property in E_1 to conclude that $\mathcal{Z}_{\hat{J}(t)}(t)$, which denotes the zero number of $\eta(\cdot, t)$ in the interval $\hat{J}(t)$, is finite, decreasing in t , and strictly decreasing when t passes through a moment s when $\eta(\cdot, s)$ has a degenerate zero. Note that the same conclusions hold true for $\mathcal{Z}_0(t)$ since it is equal to $\mathcal{Z}_{\hat{J}(t)}(t)$ by (3.7).

If $l_2(t)$ will never approach $l_1(t)$ for all $t > 0$, then the conclusions are proved.

We now consider the following case: there is a moment $T > t_1$ such that $l_2(t)$ moves leftward and meets $l_1(t)$ as $t \rightarrow T - 0$, and suppose that T is the first one of such moments. Due to the diminishing property for $\mathcal{Z}_0(t)$ in the time interval $(0, T)$, we see that there are at most finite moments when $\eta(\cdot, t)$ has degenerate zeros (only in $J_0(t)$). Thus, there exists $t_2 \in (t_1, T)$ such that $\eta(\cdot, t)$ has no degenerate zeros in $J_0(t)$ in the time interval (t_2, T) , that is, in this time interval $\eta(\cdot, t)$ has fixed finite number of non-degenerate zeros. For simplicity and without loss of generality, in the rest of the proof we assume that $\eta(\cdot, t)$ has no interior degenerate zeros in $J_0(t)$ for all $t \in (t_2, \infty)$. So, we can assume the null curves of η in $E_2 := \{(x, t) \mid x \in J_0(t), t_2 < t < T\}$ are

$$\gamma_1(t) < \gamma_2(t) < \cdots < \gamma_k(t), \quad t \in (t_2, T),$$

for some positive integer k . Using [16, Theorem 2] as in the proof of [13, Lemma 2.4], one can show that

$$x_j := \lim_{t \rightarrow T-0} \gamma_j(t), \quad j = 1, 2, \dots, k,$$

exist. We divide the situation into two cases.

Case 1. Assume that $l(T) < x_1$, that is, $l_2(t)$ moves leftward to catch up with $l_1(t)$ as $t \rightarrow T - 0$, while the left-most null curve $\gamma_1(t)$ remains on the right of $l(t)$ till $t = T$. In this case, by using the comparison principle for very weak solutions in the domain

$$E_3 := \{(x, t) \mid -\infty < x < \gamma_1(t), t \in (t_2, T)\},$$

we have

$$\eta(x, T) > 0 \text{ for } l(T) < x < x_1, \quad \eta(l(T), T) = \eta(x_1, T) = 0.$$

As a consequence, we have $Z_0(T) = k$. The boundary zero $l(T)$ of $\eta(\cdot, T)$ is not included in $Z_0(T)$. Recalling the Darcy law, we have the additional fact that $D_x^+ \eta(l(T), T) = 0$. We remark that, this will lead to a contradiction with the Hopf lemma immediately in the Stefan problems for RDEs (cf. [12, 13]). However, in the current problem, the fact that $D_x^+ \eta(l(T), T) = 0$ does not contradict the Hopf lemma, since the Hopf lemma is no longer necessarily true at $(l(T), T)$ where equation (3.6) is degenerate.

Next, we consider the time period from the time T to $t_3 \in (T, \infty]$, where t_3 is the smallest moment when $\gamma_1(t_3)$ meets $l(t_3)$. Using the comparison principle in the domain

$$E_4 := \{(x, t) \mid -\infty < x < \gamma_1(t), t \in [T, t_3)\},$$

we conclude that $\eta(x, t) \geq 0$ in this domain. This implies that, for $t \in [T, t_3)$, $l_1(t) \leq l_2(t)$, but the equality can hold from time to time. However, no matter if $l_1(t) < l_2(t)$ or $l_1(t) = l_2(t)$, we have $v_1, v_2 > 0$ for $x \in (l(t), \gamma_1(t)]$, and so the strong maximum principle holds in

$$E_5 := \{(x, t) \mid l(t) < x \leq \gamma_1(t), T \leq t < t_3\},$$

which implies that $\eta(x, t) > 0$ in E_5 . As a consequence, $Z_0(t) = k$ for $t \in [T, t_3)$.

At the moment t_3 , $\gamma_1(t_3)$ meets $l(t_3)$. The following analysis for $t \geq t_3$ is similar to what we will do in Case 2.

Case 2. Assume $\gamma_1(t)$ tends to $l(t)$ at $t \rightarrow T - 0$, and so $l_1(T) = l_2(T) = \gamma_1(T)$ (see Figure 4). Recall that we assumed before that $\eta(\cdot, t)$ no longer has interior degenerate zeros. However, this does not exclude the possibility that $\gamma_1(T) = \gamma_2(T) = \dots = \gamma_{j_0}(T) = l(T)$, that is, j_0 -null curves tend to the point $(l(T), T)$ at the same time as $t \rightarrow T - 0$. Since the discussion is similar, in what follows, we only consider the case

$$l_1(T) = l_2(T) = l(T) = \gamma_1(T) < \gamma_2(T) < \dots < \gamma_k(T).$$

Then we have $Z_0(T) = k - 1$, and

$$\eta(x, T) < 0, \quad l(T) < x < \gamma_2(T).$$

As above, we next consider the time period from T to $t_4 \in (T, \infty]$, where t_4 is the smallest moment when $\gamma_2(t_4)$ meets $l(t_4)$. Using the comparison principle for very weak solutions in the domain

$$E_6 := \{(x, t) \mid -\infty < x < \gamma_2(t), t \in [T, t_4)\},$$




Fig. 4. Vanishing of zero at $t = T$.

we conclude that $\eta(x, t) \leq 0$ in this domain. This implies that, for $t \in [T, t_4]$, $l_2(t) \leq l_1(t)$, but the equality can hold from time to time. However, no matter if $l_2(t) < l_1(t)$ or $l_2(t) = l_1(t)$, we have $v_1, v_2 > 0$ for $x \in (l(t), \gamma_2(t)]$, and so the strong maximum principle holds in

$$E_7 := \{(x, t) \mid l(t) < x \leq \gamma_2(t), T \leq t < t_4\},$$

which implies that $\eta(x, t) < 0$ in E_7 . As a consequence, $Z_0(t) = k - 1$ for $t \in [T, t_4]$.

In the case where $t_4 = \infty$, there is nothing left to prove. In the case where $t_4 < \infty$, we can analyze the situation for $t \geq t_4$ as in the current case.

In conclusion, we see that, in case when no interior degenerate zeros appear, the zero number of $\eta(\cdot, t)$ in the open interval $J_0(t)$ is decreasing. It is strictly decreasing when some interior null curves touch the boundaries.

Next, we consider the assumption (b). Without loss of generality, we assume that $l_1(0) = l_2(0)$, that is, the left free boundaries of v_1 and v_2 glue together at the beginning. Since $Z_0(0) < \infty$, there exists $\tilde{x} > l_1(0) = l_2(0)$, such that $v_1(x, 0)$ and $v_2(x, 0)$ has no intersection points in $(l_1(0), \tilde{x})$. This is a situation like that in Case 1 at time T . The rest of the discussion is then similar to the previous one.

This completes the proof of Lemma 3.3. ■

Remark 3.4. In the above proof, as well as in other related places of the paper, we use the maximum principle for η . One may worry about the unboundedness of the coefficient c in (3.6) at the points where $v_2 = 0$. We remark that, though we write $v_1 - v_2$ in the form of η for convenience, we actually compare the very weak solutions v_1 and v_2 . As in [30, Theorem 5.5] the comparison principle between v_1, v_2 holds safely. The main reason is that each of them can be approximated by a sequence of decreasing classical solutions.

Lemma 3.5. Assume (F), v_1, v_2 are given as above. Further assume that $l(0) \geq r(0)$. Then there exists a time $t_1 \in [0, \infty]$ such that

$$\begin{cases} Z_0(t) = 0, & t \leq t_1, \\ Z_0(t) = 1, & t_1 < t \ll t_1 + 1, \\ Z_0(t) \text{ decreases,} & t > t_1. \end{cases}$$

Proof. By the assumption $l(0) \geq r(0)$, we see that the supports of two initial data $v_1(x, 0)$ and $v_2(x, 0)$ do not have overlap. Without loss of generality, we assume v_1 lies on the left of v_2 in the beginning time, that is, $r_1(0) \leq l_2(0)$. Define $t_1 := \sup\{t \geq 0 \mid r_1(t) \leq l_2(t)\} \in [0, \infty]$, then $Z_0(t) = 0$ when $t \leq t_1$ by the definition of $Z_0(t)$. When $t_1 < \infty$, $r_1(t)$ will surpass $l_2(t)$ at $t = t_1$ (see Figure 5). According to the monotonicity of v_i near the free




Fig. 5. Generation of zero at $t = t_1$.

boundaries, we obtain $Z_0(t) = 1$ for $t_1 < t \ll t_1 + 1$. After t_1 , the problem can be treated like that in Lemma 3.3. Then the lemma follows. ■

Proof of Theorem 1.2. The proof is almost the same as that of Theorem 3.2. We only note that the support of initial data may not be singly connected, but consists of finite intervals. Similar to t_1 in Lemma 3.5, we define t_j as the j -th time for two components (in one of them $v_1 > 0$ and in the other $v_2 > 0$) to meet. It is easy to see that $\#\{t_j\}$ is at most finite up to the choice of initial data. However, more complex than the case in Lemma 3.5, $Z_0(t)$ may not increase strictly at $t = t_j$, since there may exist other interior degenerate zeros of $v_1 - v_2$ which vanishes at the same time. To exclude such cases, we select a subset of $\{t_j\}$ such that $Z_0(t)$ strictly increases at these times. This proves Theorem 1.2. ■

3.2. Monotonicity outside of the initial support

Assume the initial data u_0 of (CP) belongs to

$$\mathfrak{X}_1 := \{\psi \in C(\mathbb{R}) \mid \psi(x) > 0 \text{ for } x \in (-b, b), \text{ and } \psi(x) = 0 \text{ for } |x| \geq b\}.$$

Then the solution has exactly two free boundaries:

$$l(t) < r(t) \quad \text{with } l(0) = -b \text{ and } r(0) = b.$$

Moreover, $l(t)$ is strictly decreasing after the waiting time $t_1^*(-b)$, $r(t)$ is strictly increasing after the waiting time $t_2^*(b)$, and $\text{spt}[u(\cdot, t)] = [l(t), r(t)]$. We now prove a monotonicity property for $u(\cdot, t)$ outside of $[-b, b]$.

Lemma 3.6. Assume (F) and $u_0 \in \mathfrak{X}_1$. Then

- (i) for any given $\bar{x} > b$, if $r(\bar{t}) = \bar{x}$ for some $\bar{t} > t^*(b)$, then $v_x(\bar{x}, t) < 0$ for $t > \bar{t}$.
Similarly, for any given $\bar{x}' < -b$, if $l(\bar{t}') = \bar{x}'$ for some $\bar{t}' > t^*(-b)$, then $v_x(\bar{x}', t) > 0$ for $t > \bar{t}'$;
- (ii) $-2b \leq r(t) + l(t) \leq 2b$ for all $t > 0$.

Proof. (i) We only work on $\bar{x} > b$ since the case $\bar{x}' < -b$ is studied similarly. Given $\bar{x} > b$, either $r(t) \leq \bar{x}$ for all $t > 0$ or, by Theorem 2.7, there exists a unique $\bar{t} > t^*(b)$ such that $r(\bar{t}) = \bar{x} > b$ and $r(t) > \bar{x}$ for $t > \bar{t}$. We consider only the latter case and will use the so-called Aleksandrov's reflection principle. Set

$$\begin{aligned} \eta(x, t) &:= v(x, t) - v(2\bar{x} - x, t), \\ (x, t) \in E_1 &:= \{(x, t) \mid -\infty < x \leq \bar{x}, t \geq \bar{t}\}. \end{aligned}$$

Then, for some functions $c_1 = c_1(v_x)$ and $c_2 = c_2(m, g', v, v_{xx})$, η satisfies

$$\begin{cases} \eta_t = (m-1)v\eta_{xx} + c_1\eta_x + c_2\eta, & (x, t) \in E_1, \\ \eta(\bar{x}, t) = 0, & t \geq \bar{t}, \\ \eta(x, \bar{t}) \geq 0, & x \leq \bar{x}. \end{cases}$$

Using the maximum principle (see Remark 3.4 if one worries about the unboundedness of c_2), we have

$$\eta(x, t) \geq 0 \text{ in } E_1, \quad \eta_x(\bar{x}, t) \leq 0 \text{ for } t \geq \bar{t}. \quad (3.8)$$

Furthermore, $v(\bar{x}, t) > 0$ for $t > \bar{t}$ by the positivity persistence, and so both v and η are classical near the line $\{x = \bar{x}, t > \bar{t}\}$. This implies that the Hopf lemma is applicable and so

$$\eta_x(\bar{x}, t) = 2v_x(\bar{x}, t) < 0 \text{ for } t > \bar{t}.$$

(ii) The first inequality in (3.8) also implies that

$$l(t) \leq 2\bar{x} - r(t), \quad t > \bar{t},$$

that is,

$$l(t) + r(t) \leq 2\bar{x}, \quad t > \bar{t}.$$

Finally, since $\bar{x} - b > 0$ can be chosen as small as possible, so does $\bar{t} - t^*(b)$, we actually have $r(t) + l(t) \leq 2b$ for $t > 0$. In a similar way, one can show that $r(t) + l(t) \geq -2b$ for all $t > 0$. $\blacksquare$

In the above lemma, we consider initial data with exactly one connected support. Now we consider the case $u_0 \in \mathcal{X}$. More precisely, assume

$$\begin{aligned} u_0 \in \{ \psi \in C(\mathbb{R}) \mid & \text{there exist } -b \leq l_1 < r_1 \leq l_2 < r_2 \leq \dots \leq l_n < r_n \leq b \\ & \text{such that } \psi(x) > 0 \text{ in } (l_j, r_j) \text{ for } j = 1, \dots, n, \\ & \text{and } \psi(x) = 0 \text{ otherwise} \}. \end{aligned} \quad (3.9)$$

Note that this set is exactly the same as $\mathcal{X}$. Here we explicitly give the name of each boundary (see Figure 1). As we will see below, the purpose for choosing u_0 in this set is mainly for the clarity of the statement for the free boundaries. Our approach remains valid for general continuous and compactly supported initial data with infinitely many free boundaries.

For each $i = 1, 2, \dots, n$, denote by $l_i(t)$ (resp. $r_i(t)$) the free boundary of the solution starting at l_i (resp. r_i). Given $j \in \{1, 2, \dots, n-1\}$, the boundary point r_j either keeps stationary for all time (that is, $t^*(r_j) = \infty$) or it moves rightward after the finite waiting time. In the latter case, we define

$$T_j := \sup\{T \mid \text{there exists } x_j \in [r_j, l_{j+1}] \text{ such that } u(x_j, t) = 0 \text{ for } 0 \leq t \leq T\}.$$

In the case where $T_j = \infty$, the free boundaries $r_j(t)$ will never meet $l_{j+1}(t)$ or $r_j = l_{j+1}$ and both of them have infinite waiting times. In the case $T_j < \infty$, these two free boundaries meet in some finite time and then these free boundaries vanish after T_j . (Note that

the following case will not happen: they meet together in finite time, and this zero of u will never vanish. In fact, by Step 2 in the proof of Theorem 2.7, any free boundary will never stop once it begins to move.) Hence, for $t > T_j$, the intervals $(l_j(t), r_j(t))$ and $(l_{j+1}(t), r_{j+1}(t))$ merge into one, $(l_j(t), r_{j+1}(t))$. We remark that it may merge other intervals at the same time or at some time after T_j . Whatever happens, there exists a large T such that, for $t \geq T$, there exist fixed number of free boundaries:

$$l_1(t) \equiv l_{n_1}(t) < r_{n_2}(t) \leq l_{n_3}(t) < r_{n_4}(t) \leq \cdots \leq l_{n_{2k-1}}(t) < r_{n_{2k}}(t) \equiv r_n(t) \quad (3.10)$$

for $n_1, \dots, n_{2k} \in \{1, 2, \dots, n\}$ satisfying

$$n_1 \leq n_2 < n_3 \leq n_4 < \cdots < n_{2k-1} \leq n_{2k},$$

such that $v(x, t) > 0$ in $(l_{n_{2i-1}}(t), r_{n_{2i}}(t))$, $i = 1, 2, \dots, k$, and $v = 0$ otherwise. Moreover, by (3.9) and the monotonicity of $l_i(t)$, $r_i(t)$, we have

$$-b < r_{n_2}(t), \quad l_{n_{2k-1}}(t) < b, \quad t \geq 0. \quad (3.11)$$

Using Lemma 3.6 in each of these intervals, we have the following result.

Corollary 3.7. *Assume (F) and (I). Then there exists a large T such that for $t \geq T$, $u(\cdot, t)$ has $2k$ free boundaries as in (3.10) and (3.11). Moreover, for $1 \leq j \leq k$, $u(\cdot, t)$ is strictly increasing in $[l_{2j-1}(t), l_{2j-1}(0))$ and strictly decreasing in $(r_{2j}(0), r_{2j}(t)]$. In particular, $u(\cdot, t)$ is strictly increasing in $[l_1(t), l_1(0))$ and strictly decreasing in $(r_n(0), r_n(t)]$.*

3.3. General convergence

Proof of Theorem 1.1. We prove the results by using the pressure v instead of u .

Step 1. To show the ω -limit set is non-empty. By the locally uniform Hölder bound in (2.2), there exist $C > 0$, $\alpha_1 \in (0, 1)$, both depend on $\|v_0\|_C$ and m , such that for any $M > 0$ and any increasing time sequence $\{t_n\}$, there holds

$$\|v(x, t_n + t)\|_{C^{\alpha_1, \frac{\alpha_1}{2}}([-M, M] \times [-1, 1])} \leq C.$$

Hence, for any $\alpha \in (0, \alpha_1)$, there is a subsequence of $\{t_n\}$, denoted again by $\{t_n\}$, such that

$$\|v(x, t_n + t) - w_M(x, t)\|_{C^{\alpha, \frac{\alpha}{2}}([-M, M] \times [-1, 1])} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Using Cantor's diagonal argument, there exist a subsequence of $\{t_n\}$, denoted again by $\{t_n\}$, and a function $w(x, t) \in C^{\alpha, \frac{\alpha}{2}}(\mathbb{R} \times \mathbb{R})$ such that

$$v(x, t_n + t) \rightarrow w(x, t) \quad \text{as } n \rightarrow \infty, \quad \text{in the topology of } C_{\text{loc}}^{\alpha, \frac{\alpha}{2}}(\mathbb{R}^2).$$

In addition, if $w(x, t) > 0$ in a domain $E \subset \mathbb{R}^2$, then for any compact subset $D \subset E$, there exists small $\rho > 0$ such that

$$w(x, t), v(x, t_n + t) \geq \rho > 0, \quad (x, t) \in D, \quad n \gg 1.$$

Then, $v(x, t_n + t)$ is classical in D , and so, for any $\beta_1 \in (0, 1)$,

$$\|v(x, t_n + t)\|_{C^{2+\beta_1, 1+\frac{\beta_1}{2}}(D)} \leq C$$

for any large n and some C independent of n . This implies that a subsequence of $\{v(x, t_n + t)\}$ converges in $C^{2+\beta, 1+\frac{\beta}{2}}(D)$, $0 < \beta < \beta_1$, to the limit $w(x, t)$. Thus, $w(x, t)$ is a classical solution of (pCP) in the domain where $w(x, t) > 0$, and it is a very weak solution of (pCP) for $(x, t) \in \mathbb{R}^2$. Consequently, the ω -limit set of v in the topology of (1.6) is non-empty, compact, connected and invariant.

Step 2. Limits of the free boundaries. By Corollary 3.7, there exists a large T such that $v(\cdot, t)$ has exactly $2k$ free boundaries as in (3.10) and (3.11) for all $t \geq T$. For each $i = 1, 2, \dots, k$, since $l_{n_{2i-1}}(t)$ is decreasing and $r_{n_{2i}}(t)$ is increasing, the following limits exist:

$$\begin{aligned} l^\infty &:= \lim_{t \rightarrow \infty} l_{n_1}(t), & l_{n_{2i-1}}^\infty &:= \lim_{t \rightarrow \infty} l_{n_{2i-1}}(t), \\ r_{n_{2i}}^\infty &:= \lim_{t \rightarrow \infty} r_{n_{2i}}(t), & r^\infty &:= \lim_{t \rightarrow \infty} r_{n_{2k}}(t). \end{aligned} \quad (3.12)$$

We have the following possibilities.

Case 1. $k \geq 2$. In this case, all the free boundaries lie in $[-b, b]$ except for the left-most one $l_{n_1}(t) \equiv l_1(t)$ and the right-most one $r_{n_{2k}}(t) \equiv r_n(t)$. Together with Lemma 3.6 (ii), we have

$$\begin{aligned} l_1(t) &\geq -2b - r_{n_2}(t) \geq -3b, \\ r_n(t) &\leq 2b - l_{n_{2k-1}}(t) \leq 3b. \end{aligned}$$

Consequently, all the limits in (3.12) are in $[-3b, 3b]$.

Case 2. $k = 1$ and $-l^\infty, r^\infty < \infty$. In this case, $v(\cdot, t)$ has exactly one connected compact support $[l_{n_1}(t), r_{n_2}(t)] = [l_1(t), r_n(t)]$ which tends to the bounded interval $[l^\infty, r^\infty]$ as $t \rightarrow \infty$. Note that, even in this case, the support of the ω -limit $w(x, t_1)$ may have several separated intervals as in the ground state solution in (1.7), since the convergence $v \rightarrow w$ is taken in $C_{\text{loc}}^{\alpha, \frac{\alpha}{2}}(\mathbb{R})$ topology, the positivity of v does not guarantee the positivity of its limit w .

Case 3. $k = 1$ and $-l^\infty, r^\infty = \infty$. In this case, $v(\cdot, t)$ has exactly one connected compact support $[l_{n_1}(t), r_{n_2}(t)] = [l_1(t), r_n(t)]$ which tends to $\mathbb{R}$ as $t \rightarrow \infty$.

Step 3. Quasi-convergence and Type II ground state solutions in the case $-l^\infty, r^\infty < \infty$. This includes Cases 1 and 2 in Step 2. We first use the Lyapunov functional to prove the quasi-convergence result, that is, any ω -limit is a stationary solution of (pCP). Define

$$E[v(\cdot, t)] := \int_{l_1(t)}^{r_n(t)} \left[\frac{m-1}{2} v^{\frac{2}{m-1}} v_x^2 - G(v) \right] dx, \quad t \geq T, \quad (3.13)$$

with

$$G(v) := \int_0^v g(r) r^{\frac{3-m}{m-1}} dr,$$

which is convergent by $|g(v)| \leq K(m-1)|v|$. When v is the solution of (pCP), a direct calculation shows that

$$\frac{d}{dt} E[v(\cdot, t)] = - \int_{l_1(t)}^{r_n(t)} v^{\frac{3-m}{m-1}} v_t^2 dx \leq 0.$$

So, $E[v(\cdot, t)]$ is a Lyapunov functional. It is bounded from below by our assumption $l^\infty \leq l_1(t) < r_n(t) \leq r^\infty$. Hence, we can use the standard argument to show that any ω -limit of $v(\cdot, t)$ is a stationary solution of the equation pRPME.

We now explain that any ω -limit is also a stationary solution of the Cauchy problem (pCP). Assume, $w(x, t)$ is the limit function obtained in Step 1. For each $\tau_1 \in \mathbb{R}$, we claim that the solution $v(x, t; w(\cdot, \tau_1))$ of (pCP) with $v_0(x) = w(x, \tau_1)$ is a stationary one: $v(x, t; w(\cdot, \tau_1)) \equiv w(x, \tau_1)$. Since $w(x, \tau_1)$ has boundaries $l^\infty, l_{n_{2i-1}}^\infty, r_{n_{2i}}^\infty, r^\infty$ as in (3.12), by Lemma 2.3 we only need to show that the waiting time of each of these boundaries is not zero. Without loss of generality, we only prove $t^*(l^\infty) > 0$. By contradiction, assume the left boundary $l_w(t)$ of $v(x, t; w(\cdot, \tau_1))$ satisfies $l_w(\tau_2) < l^\infty$ for some $\tau_2 > 0$. Since $w(x, \tau_1 + \tau_2)$ is also an ω -limit of v ,

$$v(x, t_n + \tau_1 + \tau_2) \rightarrow w(x, \tau_1 + \tau_2) \equiv v(x, \tau_2; w(\cdot, \tau_1)), \quad n \rightarrow \infty.$$

(Note that $w(x, \tau_1 + t)$ solves (pCP) with initial data $w(x, \tau_1)$ at $t = 0$. It is nothing but $v(x, t; w(\cdot, \tau_1))$ by the uniqueness of the very weak solution of (pCP).) Thus, we have

$$l_1(t_n + \tau_1 + \tau_2) \rightarrow l_w(\tau_2) < l^\infty, \quad n \rightarrow \infty,$$

contradicts the definition of l^∞ . Therefore, $w(x, \tau_1)$ is a stationary solution of (pCP).

We continue to specify the possible shapes of the ω -limits of v . Denote by $w(x)$ one of them. By the monotonicity in Lemma 3.6 and Corollary 3.7, we see that either $w(x) \equiv 0$, or, it has maximums in $[-b, b]$. In the latter case, we claim that w does not have positive minimums. Otherwise, by the equation pRPME, w is symmetric with respect to each maximum point and each minimum point as long as it keeps positive, and so w must be a positive periodic function, which is impossible by the monotonicity outside of $[-b, b]$. Therefore, the only possible choice for w is that it has only maximum points in $[-b, b]$ but has no positive minimum points, and so it is a Type II ground state solution of (pCP). More precisely, for some $-b \leq z_1 < z_2 < \dots < z_k \leq b$ and $L > 0$ with $z_i + 2L \leq z_{i+1}$, $i = 1, 2, \dots, k-1$, there holds

$$w(x) = \mathcal{V}(x) := V(x - z_1) + V(x - z_2) + \dots + V(x - z_k), \quad x \in \mathbb{R}, \quad (3.14)$$

where $V(x)$ is the unique nonnegative compactly supported stationary solution of (pCP) as in Lemma 2.4.

Step 4. Convergence in case $-l^\infty, r^\infty < \infty$. In the previous step, we know that any ω -limit $w(x)$ of $v(x, t)$ is either 0 or a Type II ground state solution as $\mathcal{V}$. In the former case, we actually have the convergence result:

$$v(\cdot, t) \rightarrow 0 \quad \text{as } t \rightarrow \infty,$$

since the ω -limit set of v is connected. In the latter case, we now improve the conclusion in Step 3 to a convergence result: $v(x, t) \rightarrow \mathcal{V}(x)$ as $t \rightarrow \infty$, that is, the shifts $\{z_1, z_2, \dots, z_k\}$ in (3.14) is unique.

We first prove the following claim.

Claim 1. *There exists $T_1 > 0$ such that $v(\cdot, t)$ has exactly k maximums larger than Θ for $t \geq T_1$, where Θ is the largest positive zero of g in $(0, \Theta_1)$ as in (2.5).*

First, we show that $v(\cdot, t) - \Theta$ has exactly $2k$ non-degenerate zeros for large time. Using the standard zero number argument for $v(\cdot, t) - \Theta$, we know that, for some $T_2 > 0$, $v(\cdot, t) - \Theta$ has fixed even number of non-degenerate zeros when $t > T_2$. Since $\mathcal{V}$ defined by (3.14) is one ω -limit of v , we see that the fixed number must be $2k$. Denote zeros by

$$y_1(t) < y_2(t) < \dots < y_{2k-1}(t) < y_{2k}(t).$$

Next, we show that v has exactly one maximum $z_i(t)$ in $(y_{2i-1}(t), y_{2i}(t))$ for all large t and each $i = 1, 2, \dots, k$. We only prove the conclusion in $[y_1(t), y_2(t)]$. In this interval, $\zeta(x, t) := v_x(x, t)$ satisfies

$$\begin{cases} \zeta_t = (m-1)v\zeta_{xx} + (m+1)\zeta\zeta_x + g'(v)\zeta, & y_1(t) < x < y_2(t), \quad t > T_2, \\ \zeta(y_1(t), t) > 0 > \zeta(y_2(t), t), & t > T_2. \end{cases}$$

Since $v(x, t) > \Theta$ for $y_1(t) < x < y_2(t)$, $t > T_2$, the equation is a uniform parabolic one, and so we can use the classical zero number argument to conclude that, for some $T_3 > T_2$, $\zeta(\cdot, t)$ has fixed number of zeros in $(y_1(t), y_2(t))$. By our assumption, for some increasing time sequence $\{t_n\}$, we have

$$[y_1(t_n), y_2(t_n)] \subset (-L + z_1, L + z_1),$$

and $v(x, t_n) \rightarrow \mathcal{V}(x)$ as $n \rightarrow \infty$ in the topology $C_{\text{loc}}^{2,1}((-L + z_1, L + z_1))$. This implies that $v(x, t_n)$ has exactly one maximum point $z_1(t_n) \in (y_1(t_n), y_2(t_n))$ for large n . This prove Claim 1.

By $v(x, t_n) \rightarrow \mathcal{V}(x)$ again, for some large $T_4 > T_3$, $v(x, T_4) > \Theta$ near z_1 . For simplicity, when $t \geq T_4$, denote by $l(t)$, $r(t)$ the two free boundaries of the support of $v(\cdot, t)$ containing z_1 . They are nothing but $l_1(t) \equiv l_{n_1}(t)$ and $r_{n_2}(t)$ in (3.10). Then $-l(t)$ and $r(t)$ are non-decreasing continuous functions.

We now use Claim 1 to prove the convergence result, that is, the shift z_1 is unique. By contradiction, we assume that v has another ω -limit point

$$\check{\mathcal{V}}(x) := V(x - \check{z}_1) + V(x - \check{z}_2) + \dots + V(x - \check{z}_k),$$

and, without loss of generality, assume $z_1 < \check{z}_1$. Choose $x_0, x_1, x_2 \in [z_1, \check{z}_1]$ such that

$$x_0 \neq \frac{l(T_4) + r(T_4)}{2}, \quad x_0 - x_1 = x_2 - x_0 > 0,$$

that is, x_0 is the center of $[x_1, x_2] \subset [z_1, \check{z}_1]$ but not that of $[l(T_4), r(T_4)]$. Then, by the connectedness of the ω -limit set $\omega(v)$, each $V(x - x_i)$, $i = 0, 1, 2$, is a part of an ω -limit of v .

Define

$$\tilde{l}(t) := 2x_0 - r(t), \quad \tilde{r}(t) := 2x_0 - l(t), \quad \tilde{v}(x, t) := v(2x_0 - x, t)$$

for $x \in \mathbb{R}$, $t \geq T_4$. Then $\tilde{l}(t)$ and $\tilde{r}(t)$ are free boundaries of $\tilde{v}(\cdot, t)$, $\tilde{v}(x, t) > 0$ for $x \in (\tilde{l}(t), \tilde{r}(t))$, $t \geq T_4$, and

$$\tilde{v}_t = (m-1)\tilde{v}\tilde{v}_{xx} + \tilde{v}_x^2 + g(\tilde{v}), \quad x \in \mathbb{R}, \quad t \geq T_4.$$

It is easily seen that $\tilde{l}(t) < r(t)$, $\tilde{r}(t) > l(t)$, and so $v(\cdot, t)$ and $\tilde{v}(\cdot, t)$ have non-empty common domain $[\hat{l}(t), \hat{r}(t)]$ with

$$\hat{l}(t) := \max\{l(t), \tilde{l}(t)\} < x_0 < \hat{r}(t) := \min\{r(t), \tilde{r}(t)\}, \quad t \geq T_4.$$

By the choice of x_0 , we see that

$$l(T_4) \neq \tilde{l}(T_4) = 2x_0 - r(T_4), \quad r(T_4) \neq \tilde{r}(T_4) = 2x_0 - l(T_4).$$

Hence, we can use Lemma 3.3 which is valid in open intervals to conclude that the number of the zeros of

$$\eta(\cdot, t) := v(\cdot, t) - \tilde{v}(\cdot, t)$$

in the open interval $J_0(t) := (\hat{l}(t), \hat{r}(t))$ is finite, decreasing in $t > T_4$, and strictly decreasing when $\eta(\cdot, t)$ has degenerate zeros in $J_0(t)$. Consequently, for some large time $T_5 > T_4$,

$$x_0 \text{ is not a degenerate zero of } \eta(\cdot, t) := v(\cdot, t) - \tilde{v}(\cdot, t) \text{ for } t > T_5. \quad (3.15)$$

On the other hand, by the previous assumption, there exist two time sequences $\{s_n^{(1)}\}$ and $\{s_n^{(2)}\}$ with

$$\cdots < s_n^{(1)} < s_n^{(2)} < s_{n+1}^{(1)} < s_{n+1}^{(2)} < \cdots, \quad n = 1, 2, \dots,$$

such that, for $i = 1, 2$,

$$\|v(x, s_n^{(i)}) - V(x - x_i)\|_{L^\infty([-L+x_i, L+x_i])} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Therefore, for all large n we have

$$z_1(s_n^{(i)}) \approx x_i, \quad i = 1, 2.$$

By the continuity of $z_1(t)$, there exists $s_n \in (s_n^{(1)}, s_n^{(2)})$ such that $z_1(s_n) = x_0$. Thus, for large n we have

$$\begin{aligned} \eta(x_0, s_n) &= v(x_0, s_n) - \tilde{v}(x_0, s_n) = 0, \\ \eta_x(x_0, s_n) &= v_x(x_0, s_n) - \tilde{v}_x(x_0, s_n) = 0, \end{aligned}$$

contradicting (3.15). This proves the uniqueness of z_1 , and so the convergence in this step is proved.

Step 5. To show that any ω -limit is a stationary solution of pRPME in case $-l^\infty, r^\infty = \infty$. This is Case 3 in Step 2. For large t , say $t \geq T_6$, $v(\cdot, t)$ has exactly one connected support. Denote, for simplicity, its left and right free boundaries by $l(t)$ and $r(t)$, respectively. Due to the unboundedness of $l(t)$ and $r(t)$, the Lyapunov functional as in (3.13) is no longer a suitable tool to derive the quasi-convergence result. Instead, we will use the zero number argument to continue our reasoning.

Using the limit function $w(x, t)$ in Step 1, we want to show that, for each $t \in \mathbb{R}$, $w(x, t)$ is a stationary solution of (pCP). It is sufficient to prove that $w(x, 0)$ is so. When $w(x, 0) \equiv 0$ in $x \in \mathbb{R}$, there is nothing left to prove. So, we assume without loss of generality that $w(0, 0) > 0$. We now construct directly a positive stationary solution as the following:

$$(m-1)\bar{v}\bar{v}_{xx} + \bar{v}_x^2 + g(\bar{v}) = 0, \quad \bar{v}(0) = w(0, 0) > 0, \quad \bar{v}'(0) = w_x(0, 0). \quad (3.16)$$

As long as $\bar{v} > 0$, the equation is a non-degenerate second-order ordinary differential equation, and so the existence and uniqueness of the solution of (3.16) follow from the standard theory. Noticing from the equation we see that $\bar{v}$ is symmetric with respect to any positive critical point. Hence, we have the following possibilities:

- (a) $\bar{v}(x) > 0$ for all $x \in \mathbb{R}$;
- (b) $\bar{v}(x) > 0$ and $\bar{v}'(x) > 0$ in (l_0, ∞) for some $l_0 \in (-\infty, 0)$, and $\bar{v}(l_0) = 0$;
- (c) $\bar{v}(x) > 0$ and $\bar{v}'(x) < 0$ in $(-\infty, r_0)$ for some $r_0 \in (0, \infty)$, and $\bar{v}(r_0) = 0$;
- (d) $\bar{v}(x) > 0$ for $x \in (l_0, r_0)$ for some $l_0 < 0 < r_0$, $\bar{v}(l_0) = \bar{v}(r_0) = 0$ and it has a unique maximum point

$$z_0 := \frac{l_0 + r_0}{2}.$$

In this step, we will prove $w(x, 0) \equiv \bar{v}(x)$ in $\text{spt}[\bar{v}]$.

(a) Set

$$\eta(x, t) := v(x, t) - \bar{v}(x), \quad x \in \mathbb{R}, \quad t \geq T_6.$$

In the current case, we have

$$\eta(l(t), t) < 0, \quad \eta(r(t), t) < 0, \quad t \geq T_6, \quad (3.17)$$

and

$$\eta_t = a(x, t)\eta_{xx} + b(x, t)\eta_x + c(x, t)\eta, \quad x \in J_0(t) := (l(t), r(t)), \quad t \geq T_6, \quad (3.18)$$

with

$$a(x, t) := (m-1)v(x, t), \quad b(x, t) := v_x(x, t) + \bar{v}_x(x),$$

and

$$c(x, t) := (m-1)\bar{v}_{xx}(x) + \begin{cases} \frac{g(v(x, t)) - g(\bar{v}(x))}{v(x, t) - \bar{v}(x)}, & v \neq \bar{v}, \\ 0, & v = \bar{v}. \end{cases}$$

We will use Proposition 3.1 to study the zero number of $\eta(\cdot, t)$ in any bounded time interval $(t_1, t_2) \subset (T_6, \infty)$. For this purpose, we should require that a, b, c satisfy assumption (3.2). This is obviously not true in the domain

$$E_1 := \{(x, t) \mid x \in J_0(t), t_1 < t < t_2\},$$

since $a^{-1} = [(m-1)v]^{-1}$ is not bounded near the boundaries of E_1 . This difficulty will be solved below by cutting $J_0(t)$ a little bit near its boundaries. Since $l(t)$ and $r(t)$ are continuous and bounded functions in $[t_1, t_2]$, there exists $\rho_1 > 0$ small such that

$$\eta(l(t), t) = -\bar{v}(l(t)) \leq -\rho_1, \quad \eta(r(t), t) = -\bar{v}(r(t)) \leq -\rho_1, \quad t \in [t_1, t_2].$$

By the uniform continuity of v , there exists $\varepsilon_1 > 0$ small such that

$$\eta(x, t) < 0, \quad l(t) \leq x \leq l(t) + \varepsilon_1 \text{ or } r(t) - \varepsilon_1 \leq x \leq r(t), \quad t \in [t_1, t_2],$$

and, for some small $\rho_2 > 0$, there holds

$$v(x, t) \geq \rho_2 \quad \text{in } E_2 := \{(x, t) \mid l(t) + \varepsilon_1 \leq x \leq r(t) - \varepsilon_1, t_1 \leq t \leq t_2\}.$$

Now we see that a, b, c satisfy the conditions in (3.2) in the domain E_2 . Hence, the classical zero number diminishing properties as in Proposition 3.1 hold. Since $\varepsilon_1 > 0$, $t_1 - T_6 > 0$ can be chosen as small as possible and t_2 can be chosen as large as possible, we actually have the same zero number diminishing properties in E_1 with $(t_1, t_2) = (T_6, \infty)$. As a consequence, η has only degenerate zeros (not on the boundaries $l(t)$ and $r(t)$) in finite time, that is, there exists $T_7 > T_6$ such that $\eta(\cdot, t)$ no longer has degenerate zeros for $t \geq T_7$. Thus, for any $t \in [-1, 1]$ and all large n , $\eta(x, t_n + t) = v(x, t_n + t) - \bar{v}(x)$ has no degenerate zeros in $J_0(t_n + t)$. This implies that (see, for example, [14, Lemma 2.6]) its limit $w(x, t) - \bar{v}(x)$ either is 0 identically or has no degenerate zeros in the spatial domain where $w(\cdot, t) > 0$. The latter, however, contradicts the initial conditions in (3.16). This proves that $w(x, 0) \equiv \bar{v}(x)$ in $\text{spt}[\bar{v}]$.

(b) Now we assume $\bar{v}$ is a stationary solution as in the above listed case (b). Since $l(t) \rightarrow -\infty$ as $t \rightarrow \infty$, there exists $T_8 > T_6$ such that $l(t) < l_0$ for $t \geq T_8$. Hence, we should consider η in

$$E_3 := \{(x, t) \mid l_0 < x < r(t), t \geq T_8\}.$$

On its left boundary $\{x = l_0, t > T_8\}$, we have $\eta > 0$. By continuity, for any bounded time interval $(t_1, t_2) \subset (T_8, \infty)$, there exist $\varepsilon_2 > 0$ and $\rho_3 = \rho_3(t_1, t_2, l_0) > 0$ such that

$$\eta(x, t) > \rho_3, \quad l_0 \leq x \leq l_0 + \varepsilon_2, \quad t \in (t_1, t_2).$$

Furthermore, the term $\bar{v}_{xx}(l_0 + \varepsilon_2)$ in the coefficient c of (3.18) is also bounded, though $\bar{v}_{xx}(l_0)$ maybe not. Thus, we can use the zero number diminishing properties in the domain $\{(x, t) \mid l_0 + \varepsilon_2 < x < r(t), t_1 < t < t_2\}$. By the arbitrariness of ε_2 and (t_1, t_2) , we see that the zero number diminishing properties actually hold in the domain E_3 . The rest proof is similar to that in (a). This proves the quasi-convergence in case (b).

The proof for $w(x, 0) \equiv \bar{v}$, $x \in \text{spt}[\bar{v}]$, in cases (c) and (d) are similar to that in case (b).

Step 6. To show that any ω -limit of v is a positive stationary solution of (pCP) in the case $-l^\infty, r^\infty = \infty$. From the previous step we see that, in the case $-l^\infty, r^\infty = \infty$, any ω -limit of v must be a stationary solution of the equation pRPME as in (a)–(d). By the monotonicity in Corollary 3.7, cases (b) and (c) are impossible. We now prove that case (d) is also impossible.

(1) By the monotonicity of v and w outside of $[-b, b]$, we see that the maximum point $z_0 := \frac{l_0 + r_0}{2}$ of $\bar{v}(x) \equiv w(x, 0)|_{\text{spt}[\bar{v}]}$ lies in $[-b, b]$, and so, with $\ell := \frac{r_0 - l_0}{2}$, we have

$$-b - \ell \leq l_0 < z_0 < r_0 \leq b + \ell.$$

(2) We show that $v(-3b - 2\ell, t) < \bar{v}(z_0)$ for all $t > 0$. Assume by contradiction that the reversed inequality holds at some time $t_1 > 0$. Then by the monotonicity of v in $(l(t_1), -b)$, we have

$$v(x, t_1) \geq \bar{v}(x + z_0 + 3b + \ell), \quad x \in [-3b - 2\ell, -3b].$$

Since $\bar{v}(x + z_0 + 3b + \ell)$ is a time-independent weak subsolution of problem (pCP), by Lemma 2.3 we have

$$v(x, t) \geq \bar{v}(x + z_0 + 3b + \ell), \quad x \in \mathbb{R}, t \geq t_1,$$

and so

$$w(x, 0) \geq \bar{v}(x + z_0 + 3b + \ell), \quad x \in \mathbb{R}.$$

By the monotonicity of $w(x, 0)$ in $[-3b - 2\ell, -b]$, we have

$$w(x, 0) \geq \bar{v}(z_0), \quad x \in [-3b - \ell, -b]. \quad (3.19)$$

On the other hand, $w(x, 0) = 0$ at $l_0 \in [-b - \ell, b + \ell]$. So, $w(x, 0)$ has at least one maximum point $x_1 \in [-b, l_0]$. Without loss of generality, assume x_1 is the smallest one of such points. Then, as a stationary solution of pRPME, $w(x, 0)$ is symmetric with respect to x_1 . This implies by (3.19) that

$$w(x, 0) \geq \bar{v}(z_0), \quad x \in [-3b - \ell, 2x_1 + 3b + \ell] \supset [-b - \ell, b + \ell],$$

contradicts (1).

(3) If $\bar{v}(x) \equiv w(x, 0)|_{\text{spt}[\bar{v}]}$ is not only a stationary solution of the equation pRPME, but also a stationary solution of (pCP), then we can derive a contradiction as the following. Choose t' large such that $l(t') < -3b - 4\ell$, and choose L' such that

$$-L' + l_0 < l(t') < -L' + r_0$$

and $\bar{v}(x + L')$ has exactly one intersection point with $v(x, t')$. Then using the intersection number properties from Lemma 3.3 and using the fact in (2), we conclude that $l(t)$ will never pass across the left boundary $-L' + l_0$ of $\bar{v}(x + L')$. This contradicts the assumption $l^\infty = -\infty$.

(4) We then consider the case where $\bar{v}(x) \equiv w(x, 0)|_{\text{spt}[\bar{v}]}$ is just a stationary solution of the equation pRPME, but not a stationary one of (pCP). Denote by $v(x, t; \bar{v})$ the

solution of (pCP) with initial data $\bar{v}$, and denote its free boundaries by $\bar{l}(t)$ and $\bar{r}(t)$. By Lemma 2.3, the waiting times $t^*(l_0) = t^*(r_0) = 0$. Hence, $-\bar{l}(t)$ and $\bar{r}(t)$ are strictly increasing in $t > 0$, and

$$v(x, t; \bar{v})|_{[\bar{l}(t), \bar{r}(t)]} \equiv w(x, t)|_{[\bar{l}(t), \bar{r}(t)]}, \quad 0 < t \ll 1.$$

Since $\bar{v}$ is a (stationary) subsolution of (pCP), by Lemma 2.3 we have $v(x, t; \bar{v}) > \bar{v}(x)$ on the support of $\bar{v}$, and so $v(x, t; \bar{v})$ is strictly increasing in t . Therefore, for any small $\tau > 0$, we have

$$\bar{l}(2\tau) < \bar{l}(\tau) < \bar{l}(0) = l_0 < r_0 = \bar{r}(0) < \bar{r}(\tau) < \bar{r}(2\tau),$$

and

$$\begin{aligned} w(x, 0) &< w(x, \tau) \quad \text{for } x \in [l_0, r_0], \\ w(x, \tau) &< w(x, 2\tau) \quad \text{for } x \in [\bar{l}(\tau), \bar{r}(\tau)]. \end{aligned}$$

By Step 1, $v(x, t_n + 2\tau) \rightarrow w(x, 2\tau)$ as $n \rightarrow \infty$, and so for some large n_0 we have

$$v(x, t_{n_0} + 2\tau) \geq w(x, \tau), \quad x \in [\bar{l}(\tau), \bar{r}(\tau)].$$

From the previous step, we know that $w(x, \tau)$ is a stationary solution of pRPME in $[\bar{l}(\tau), \bar{r}(\tau)]$. Hence, $w(x, \tau)|_{[\bar{l}(\tau), \bar{r}(\tau)]}$ is a time-independent weak subsolution of (pCP). By comparison, we have

$$v(x, t) \geq w(x, \tau), \quad x \in [\bar{l}(\tau), \bar{r}(\tau)], \quad t > t_{n_0} + 2\tau.$$

This contradicts the fact that $w(x, 0)$ is an ω -limit of $v(x, t)$.

Consequently, the case (d) is also impossible. In other words, the only possible case is (a) in Step 5. We then have the following claim.

Claim 2. *When $-l^\infty, r^\infty = \infty$, any ω -limit $w(x)$ of $v(x, t; u_0)$ is a stationary solution of pRPME which is positive in $\mathbb{R}$, and so is a stationary solution of the Cauchy problem (pCP).*

Step 7. Convergence in case $-l^\infty, r^\infty = \infty$. By Claim 2, the convergence $v(x, t_n; u_0) \rightarrow w(x)$ in Step 1 holds actually in $C_{\text{loc}}^2(\mathbb{R})$ topology. Therefore, the solution $v(\cdot, t)$ is a classical one in $(l_{n_1}(t), r_{2n}(t))$. Then, using the classical zero number diminishing properties and using similar arguments as in proofs of [12, Theorem 1.1], [13, Theorem 1.1] (see also [14, Theorem 1.1]), we conclude that $v(\cdot, t)$ converges as $t \rightarrow \infty$ to either a nonnegative zero of g or

$$v(\cdot, t) \rightarrow V_0(\cdot - z_0) > 0,$$

where $z_0 \in [-b, b]$ and V_0 is an evenly decreasing positive stationary solution of pRPME, that is,

$$U_0(x) := \left(\frac{m-1}{m} V_0(x) \right)^{\frac{1}{m-1}} > 0, \quad x \in \mathbb{R}$$

is a Type I ground state solution of (CP) as given in Section 1.

This completes the proof of Theorem 1.1. ■

4. Monostable RPME

In this section, we study the hair-trigger effect for the solutions of (CP) or (pCP) with monostable type of reaction.

4.1. Stationary solutions

A stationary solution $u = q(x)$ of (CP) satisfies

$$(q^m)'' + f(q) = 0, \quad x \in J \subset \mathbb{R}. \quad (4.1)$$

We will use the phase portrait to present all the nonnegative stationary solutions that we are interested in. Equation (4.1) can be rewritten as a pair of first-order equations

$$\begin{cases} p = (q^m)' = mq^{m-1}q', \\ p' = -f(q), \end{cases} \quad (4.2)$$

or the first-order equation

$$pdp = -mq^{m-1}f(q)dq.$$

Its first integral is

$$p^2 = C - 2m \int_0^q r^{m-1} f(r) dr \quad (4.3)$$

for suitable choice of C . We consider only the trajectories in the region $\{q \geq 0\}$ which correspond to nonnegative solutions.

Case A. The points $(0, 0)$ and $(1, 0)$ are singular on the (q, p) -phase plane for system (4.2). Hence, they correspond to constant stationary solutions

$$U_q(x) \equiv q, \quad x \in \mathbb{R},$$

for $q = 0, 1$. We will show below that U_1 is the only possible choice for the ω -limit of the solution u of (CP).

Case B. When $C = C_0 := 2m \int_0^1 r^{m-1} f(r) dr$, the trajectories on the phase plane are two curves, whose functions are

$$p = P_0(q) := \left(2m \int_q^1 r^{m-1} f(r) dr \right)^{\frac{1}{2}} \quad \text{and} \quad p = -P_0(q), \quad 0 \leq q < 1.$$

They connect regular points $Q_{\pm} := (0, \pm\sqrt{C_0})$ to the singular point $(1, 0)$. Denote the corresponding stationary solution by $U^{\pm}(x)$. Then

$$\begin{aligned} U^+(l_0) &= 0, \quad U^+(x) > 0 \text{ for } x > l_0, \quad \text{and} \quad U^+(x) \nearrow 1 \text{ as } x \rightarrow +\infty; \\ U^-(r_0) &= 0, \quad U^-(x) > 0 \text{ for } x < r_0, \quad \text{and} \quad U^-(x) \nearrow 1 \text{ as } x \rightarrow -\infty, \end{aligned}$$

where l_0 is the left boundary of U^+ and r_0 is the right boundary of U^- .

Case C. When $C > C_0$, the trajectories start at $(0, \pm\sqrt{C})$ and go beyond the line $q = 1$, and so the corresponding stationary solutions are unbounded, which will not be used below.

Case D. When $0 < C < C_0$, the trajectory $p^2 + 2m \int_0^q r^{m-1} f(r) dr = C$ is a connected curve, symmetric with respect to the q -axis, and connecting the points $Q_+ := (0, \sqrt{C})$, $Q_0 := (q_0, 0)$ and $Q_- := (0, -\sqrt{C})$ for a unique $q_0 \in (0, 1)$ which is defined as the following:

$$C = 2m \int_0^{q_0} r^{m-1} f(r) dr.$$

This trajectory corresponds to a stationary solution $U_{q_0}(x)$, under the normalized condition $U'_{q_0}(0) = 0$, it satisfies

$$\begin{cases} U_{q_0}(0) = q_0, & U_{q_0}(\pm L_0) = 0 & \text{for some } L_0 > 0, \\ U'_{q_0}(x) < 0, & U_{q_0}(x) = U_{q_0}(-x) & \text{for } x \in (0, L_0). \end{cases}$$

We now study the order of the solution U in Cases B (in this case, it is U^+), C or D near its left boundary. Denote the left boundary by l_0 . By (4.3), we have

$$(U^m)' = \left(C - 2m \int_0^U r^{m-1} f(r) dr \right)^{\frac{1}{2}} \sim \sqrt{C}, \quad x \rightarrow l_0 + 0.$$

So,

$$U^m(x) \sim \sqrt{C}(x - l_0), \quad x \rightarrow l_0 + 0. \quad (4.4)$$

If we use $V := \frac{m}{m-1} U^{m-1}$ to denote the corresponding stationary solution of pRPME, we have

$$V(x) \sim \frac{m}{m-1} C^{\frac{m-1}{2m}} (x - l_0)^{\frac{m-1}{m}}, \quad x \rightarrow l_0 + 0. \quad (4.5)$$

Remark 4.1. We point out that the solutions $U^\pm$ in Case B and U_{q_0} in Case D are just stationary solutions of RPME but not stationary solutions of the Cauchy problem (CP). In fact, the expression in (4.5) and Lemma 2.1 imply that, when we choose $V(x)$ as an initial data of problem (pCP), the waiting time of its boundaries are 0. In particular, in the bistable RPME, U_{q_0} is not a Type II ground state solution.

We continue to specify the relationship between the support width $2L_0$ of U_{q_0} in Case D and its height q_0 .

Lemma 4.2. *When f is a monostable reaction term, there holds $L_0 \rightarrow 0$ as $q_0 \rightarrow 0 + 0$. In addition, if f satisfies $f(\rho u) > \rho^m f(u)$ for $\rho, u \in (0, 1)$, then L_0 is strictly increasing in $q_0 \in (0, 1)$.*

Proof. For any given small $\varepsilon > 0$, by $f'(0) > 0$, there exists small $\delta > 0$ such that

$$f(u) \geq \frac{\pi^2}{4\varepsilon^2} u^m, \quad u \in [0, \delta]. \quad (4.6)$$

For any $q_0 \in (0, \delta)$, by (4.3) we have

$$p^2 = 2m \int_q^{q_0} r^{m-1} f(r) dr \geq \frac{\pi^2}{4\varepsilon^2} (q_0^{2m} - q^{2m}).$$

Therefore, on $x \in [-L_0, 0]$ we have

$$p = (q^m)' \geq \frac{\pi}{2\varepsilon} q_0^m \left[1 - \left(\frac{q}{q_0} \right)^{2m} \right]^{\frac{1}{2}}.$$

Since $q(-L_0) = 0$, $q(0) = q_0$, by integrating the above inequality over $[-L_0, 0]$ we have

$$\frac{\pi}{2\varepsilon} L_0 \leq \int_0^1 \frac{dr}{\sqrt{1-r^2}} = \frac{\pi}{2},$$

that is, $L_0 \leq \varepsilon$.

Denote $\tilde{v} := u^m$ and $\tilde{f}(s) := f(s^{\frac{1}{m}})$, then the stationary problem

$$(u^m)'' + f(u) = 0, \quad u(x) > 0 \text{ in } (-L_0, L_0), \quad u(\pm L_0) = 0,$$

is equivalent to

$$\tilde{v}'' + \tilde{f}(\tilde{v}) = 0, \quad \tilde{v}(x) > 0 \text{ in } (-L_0, L_0), \quad \tilde{v}(\pm L_0) = 0.$$

By our assumption, the function $\tilde{f}$ is a Fisher–KPP type of nonlinearity: $\frac{\tilde{f}(s)}{s}$ is decreasing in $s \in (0, 1)$. Hence, $\tilde{v}(0)$ is strictly increasing in L_0 . This proves the lemma. ■

4.2. Hair-trigger effect

Now we can use the general convergence theorem to prove the hair-trigger effect easily.

Proof of Theorem 1.3. By Lemma 4.2, there exist sufficiently small $q_0 > 0$, small $L_0 > 0$ and some $x_0 \in \mathbb{R}$ such that

$$u_0(x) \geq U_{q_0}(x - x_0), \quad x \in [-L_0 + x_0, L_0 + x_0].$$

Since $U_{q_0}(x - x_0)$ is a time-independent very weak subsolution of (CP), by the comparison result in Lemma 2.3 (iv) we have

$$u(x, t) \geq U_{q_0}(x - x_0), \quad x \in \mathbb{R}, \quad t > 0.$$

In the monostable PME, the only stationary solution of (CP) larger than U_{q_0} is 1. So, by using the general convergence theorem we conclude that $u(x, t) \rightarrow 1$ in $L_{\text{loc}}^\infty(\mathbb{R})$ topology. This proves the hair-trigger effect. ■

Remark 4.3. From the above proofs we see that (4.6) holds if $\lim_{u \rightarrow 0+0} \frac{f(u)}{u^m} = \infty$. So, the hair-trigger effect for the monostable RPME actually holds under this condition rather than the stronger one $f'(0) > 0$.

5. Combustion RPME

In this section, we study the asymptotic behavior of the solutions of (CP) or (pCP) with combustion type of reaction: f satisfies (f_c) .

5.1. Stationary solutions

On the nonnegative stationary solutions of the equation in (CP), we first have Cases A–D types as in the monostable RPME. The difference is that in Case A the solutions are $U_q \equiv q$ for $q \in [0, \theta] \cup \{1\}$ in the combustion case (however, only U_0 , U_θ and U_1 are possible choices for the ω -limits of the solution u of (CP)); in Case D, the compactly supported stationary solutions U_{q_0} exist only for $q_0 \in (\theta, 1)$. The decay rates in (4.4) and (4.5) remain valid (in fact, they are true for $x \in [l_0, l_0 + C^{-\frac{1}{2}}\theta^m]$ due to $f(u) \equiv 0$ in $[0, \theta]$). As we mentioned in Remark 4.1, each U_{q_0} is just a stationary solution of the equation but not of the Cauchy problem (CP).

5.2. Asymptotic behavior of the solutions

Since the current problem has no Types I and II ground state solutions, by the general convergence results in Theorem 1.1 we know that $u(\cdot, t)$ converges as $t \rightarrow \infty$ to some constant solution in Case A.

Now we present a sufficient condition for the spreading phenomena: $u(\cdot, t) \rightarrow 1$ as $t \rightarrow \infty$.

Lemma 5.1. *Assume f is a combustion reaction, $u_0 \in \mathfrak{X}$ satisfies*

$$u_0(x) \geq U_{q_0}(x), \quad x \in \mathbb{R},$$

for some stationary solution U_{q_0} in Case D. Then spreading happens for the solution u of (CP).

Proof. By the comparison result, the solution $u(x, t)$ of (CP) remains above $U_{q_0}(x)$. Hence, any ω -limit of u is also above $U_{q_0}(x)$. The only possible choice for such stationary solutions in Case A is nothing but $U_1 \equiv 1$. Moreover, due to the finite propagation speed and the properties in Lemma 3.6, we know that $\text{spt}[u(\cdot, t)] = [l(t), r(t)]$ for large t , hence the convergence $u \rightarrow 1$ holds in the sense of (1.6). ■

Next, we present some sufficient conditions for the vanishing phenomenon $u(\cdot, t) \rightarrow 0$ as $t \rightarrow \infty$.

Lemma 5.2. *Assume f is a combustion reaction, $u_0 \in \mathfrak{X}$ satisfies $u_0(x) \leq \theta$. Then vanishing happens in the topologies of $L^\infty(\mathbb{R})$ for the solution u of (CP).*

Proof. The proof is simple since u actually solves the PME when $u(x, t) \leq \theta$, and so we can take a ZKB solution as a supersolution to suppress u to 0. ■

Remark 5.3. For any $q \in (0, \theta)$, $U_q \equiv q$ will not be an ω -limit. In fact, if $u(x, t_n) \rightarrow q$ ($n \rightarrow \infty$) in the $C_{\text{loc}}(\mathbb{R})$ topology, then, in the bounded interval $[-b, b]$, $u(x, T) < \theta$ for some large T . This inequality also holds in $[l(T), r(T)]$ by the monotonicity outside of $[-b, b]$. Hence, vanishing happens for u by the above lemma, a contradiction.

To prove the sharpness of the transition $u \rightarrow \theta$, we need the following lemmas.

Lemma 5.4. Assume (F). For $i = 1, 2$, assume $u_{i0} \in C(\mathbb{R})$ satisfies

$$u_{i0}(x) > 0 \text{ in } (l_i(0), r_i(0)), \quad u_{i0}(x) = 0 \text{ for } x \leq l_i(0) \text{ and } x \geq r_i(0),$$

and

$$u_{10}(x) \leq u_{20}(x), \quad x \in \mathbb{R}.$$

Let u_i be the solution of (CP) with initial data u_{i0} , and let $l_i(t)$, $r_i(t)$ be its left and right free boundaries. Then there exist $T \geq 0$, $\tau \geq 0$ and $\varepsilon_0 > 0$ such that

$$u_1(x, t) \leq u_2(x + \varepsilon, t + \tau), \quad x \in \mathbb{R}, \quad t \geq T, \quad \varepsilon \in [0, \varepsilon_0] \text{ or } \varepsilon \in [-\varepsilon_0, 0], \quad (5.1)$$

provided that one of the following conditions holds:

- (i) $r_2(t) - l_2(t) \not\equiv r_1(t) - l_1(t)$ for all $t \geq 0$;
- (ii) $t^*(l_2(0))$ or $t^*(r_2(0))$ is finite, and $r_2(t) - l_2(t) = r_1(t) - l_1(t)$ for all $t \geq 0$.

Proof. By comparison, we have

$$u_1(x, t) \leq u_2(x, t) \text{ for } x \in \mathbb{R} \text{ and } t \geq 0, \quad l_2(t) \leq l_1(t) < r_1(t) \leq r_2(t) \text{ for } t \geq 0.$$

(i) In the case $r_2(T_0) - l_2(T_0) > r_1(T_0) - l_1(T_0)$ for some $T_0 \geq 0$, we assume without loss of generality that $l_2(T_0) < l_1(T_0)$, and by continuity assume $T_0 > 0$. Then using the strong maximum principle in $\{(x, t) \mid l_1(t) \leq x < r_1(t), T_0 - 1 \ll t \leq T_0\}$, we have

$$u_1(x, T_0) < u_2(x, T_0) \quad \text{for } x \in [l_1(T_0), r_1(T_0)).$$

So, there exists a small $\varepsilon_0 > 0$ such that

$$u_1(x, T_0) < u_2(x + \varepsilon, T_0) \quad \text{for } x \in [l_1(T_0), r_1(T_0)], \quad \varepsilon \in [-\varepsilon_0, 0).$$

The conclusion then follows from the comparison principle for $T = T_0$ and $\tau = 0$.

(ii) Assume without loss of generality that $t^*(l_2(0)) < \infty$. Given $T_1 > t^*(l_2(0))$, then both $u_1(\cdot, t)$ and $u_2(\cdot, t)$ are strictly increasing in $J_1 := [x_1, l_1(0)]$, where $x_1 := l_2(T_1) = l_1(T_1)$. By the monotonicity of $l_2(t)$ (after the waiting time) we see that $l_2(t)$ is strictly decreasing in $t \geq T_1$. Hence, for any small $t > 0$ we have

$$l_2(T_1 + t) < x_1 = l_2(T_1) \quad \text{and} \quad 0 < u_2(x_1, T_1 + t) = u_1(x_2(t), T_1) < u_1(l_1(0), T_1)$$

for some $x_2(t) \in [x_1, l_1(0)]$. Choose $\tau > 0$ small and denote

$$E_1 := \{(x, t) \mid l_2(T_1 + t) \leq x \leq x_1, 0 \leq t \leq \tau\},$$

$$E_2 := \{(x, t) \mid x_1 \leq x \leq x_2(t), 0 \leq t \leq \tau\}.$$

By the monotonicity of $u_1(\cdot, T_1)$ and $u_2(\cdot, T_1 + \tau)$ in $[x_1, x_2(\tau)]$, we have

$$u_1(x, T_1) < u_1(x_2(\tau), T_1) = u_2(x_1, T_1 + \tau) < u_2(x, T_1 + \tau), \quad x \in [x_1, x_2(\tau)].$$

Thus, there exists $\varepsilon_0 > 0$ small such that

$$u_1(x, T_1) < u_2(x + \varepsilon, T_1 + \tau), \quad x \in [x_1, x_2(\tau)], \quad \varepsilon \in [-\varepsilon_0, 0]. \quad (5.2)$$

On the other hand, when $\tau > 0$ and ε_0 are sufficiently small we have

$$u_1(x, T_1) < u_2(x + \varepsilon, T_1 + \tau), \quad x \in [x_2(\tau), r_1(T_1)], \quad \varepsilon \in [-\varepsilon_0, 0].$$

Combining with (5.2), we have

$$u_1(x, T_1) < u_2(x + \varepsilon, T_1 + \tau), \quad x \in [l_1(T_1), r_1(T_1)], \quad \varepsilon \in [-\varepsilon_0, 0].$$

Then the conclusion follows from the comparison principle. ■

Lemma 5.5. *Under the assumptions of the previous lemma, if we further assume that $u_1(\cdot, t) \rightarrow \theta$ as $t \rightarrow \infty$, then $u_2(\cdot, t) \not\rightarrow \theta$ as $t \rightarrow \infty$.*

Proof. Since $u_1 \rightarrow \theta$, there exists $T' \geq T + 1$ for T in (5.1) such that

$$u_1(x, t) < \theta + \frac{\delta}{2}, \quad x \in \mathbb{R}, \quad t \geq T' - 1,$$

where δ is given in (1.4). Now, for $\lambda \in (0, 1)$, we define

$$w^\lambda(x, t) := \lambda^{-2} u_1(\lambda^m x, \lambda^2 t), \quad x \in \mathbb{R}, \quad t \geq T'.$$

We choose $\lambda_0 \in (0, 1)$ close enough to 1 so that, for every $\lambda \in [\lambda_0, 1)$,

$$w^\lambda(x, t) \leq \theta + \delta, \quad x \in \mathbb{R}, \quad t \geq T', \quad (5.3)$$

and, by (5.1),

$$w^\lambda(x, T') \leq u_2(x + \varepsilon_1, T'), \quad x \in [\lambda^{-m} l_1(T'), \lambda^{-m} r_1(T')] \quad (5.4)$$

for some ε_1 . Observe that w^λ satisfies the equation

$$w_t^\lambda = (w^\lambda)_{xx}^m + f(\lambda^2 w^\lambda), \quad x \in \mathbb{R}, \quad t > 0.$$

By (5.3) and the assumption in (f_c) , we have $f(\lambda^2 w^\lambda) \leq f(w^\lambda)$. (Here is the only place we use the assumption $f'(u) > 0$ in $(\theta, \theta + \delta]$.) Therefore, in view of (5.4), we find that w^λ is a subsolution of (CP) for $t \geq T'$. It follows that $u_2(x + \varepsilon_1, t) \geq w^\lambda(x, t)$ for $t \geq T'$. If $u_2 \rightarrow \theta$ ($t \rightarrow \infty$), then by taking limit as $t \rightarrow \infty$, we conclude that $\theta \geq \lambda^{-2}\theta$, a contradiction. ■

Proof of Theorem 1.5. From above lemmas, we know that $u(\cdot, t)$ converges as $t \rightarrow \infty$ to 0, or θ , or 1.

For the initial data $\phi_\sigma \in \mathcal{X}$, when $\sigma > 0$ is small, we have by (Φ_3) that $\phi_\sigma \leq \theta$. Hence, vanishing happens for u_σ by Lemma 5.2. Denote

$$\Sigma_0 := \{\sigma > 0 \mid \text{vanishing happens for } u_\sigma\}.$$

Then Σ_0 is not empty, and, by comparison, it is an interval. Since ϕ_σ depends on σ continuously, so does the solution u_σ (cf. [5]). Then, Σ_0 is an open interval $(0, \sigma_*)$ for some $\sigma_* \in (0, \infty]$.

In the case where $\sigma_* = \infty$ (this is the complete vanishing case, as stated in Theorem 7.1), there is nothing left to prove.

The left case is $\sigma_* \in (0, \infty)$. (This happens in particular when the condition in Lemma 5.1 holds.) In this case, the solution u_{σ_*} with critical initial data ϕ_{σ_*} converges to θ or 1. We will show that u_{σ_*} is a transition solution. Otherwise, spreading happens for $u_{\sigma_*} \rightarrow 1$, that is,

$$\sigma_* \in \Sigma_1 := \{\sigma \geq \sigma_* \mid \text{spreading happens for } u_\sigma\}.$$

Then, for any $q_0 \in (\theta, 1)$ and the corresponding stationary solution U_{q_0} in Case D in Section 5.1, there exists a sufficiently large T such that

$$u_{\sigma_*}(x, T) > U_{q_0}(x), \quad x \in \text{spt}[U_{q_0}].$$

By the continuous dependence of u on the initial data, for any σ satisfying $0 < \sigma_* - \sigma \ll 1$, we also have

$$u_\sigma(x, T) > U_{q_0}(x), \quad x \in \text{spt}[U_{q_0}].$$

Consequently, the ω -limits of u_σ are bigger than U_{q_0} , but these ω -limits are nothing but 1, that is, spreading happens for u_σ . This implies that Σ_1 is an open set and it contains σ_* , contradicts the definition of σ_* . Therefore, u_{σ_*} is a transition solution, that is, $u(x, t) \rightarrow \theta$ as $t \rightarrow \infty$, and so both of its left and right boundaries tend to infinity. It follows from Lemmas 5.4 and 5.5 that $u_\sigma \not\rightarrow \theta$ for any $\sigma > \sigma_*$. Consequently, $\Sigma_1 = (\sigma_*, \infty)$.

To finish the proof of Theorem 1.5, we only need to prove the asymptotic speeds of the free boundaries in (1.11) for the transition solution, which follows from Proposition 5.7 below.

This completes the proof of Theorem 1.5. ■

At the end of this section, we study the asymptotic speeds for the boundaries of the transition solution of (CP). In the current combustion case, the equation is PME $u_t = (u^m)_{xx}$ when $u \leq \theta$. So, it is convenient to compare the solutions of this equation, especially, the selfsimilar solutions, with the solution of (CP). For this purpose, we need a special selfsimilar solution which solves the following problem:

$$\begin{cases} v_t = (m-1)v v_{xx} + v_x^2, & 0 < x < \rho(t), \ t > 0, \\ v(0, t) = \Theta := \frac{m}{m-1} \theta^{m-1}, & t > 0, \\ v(\rho(t), t) = 0, & t > 0, \\ \rho'(t) = -v_x(\rho(t), t), & t > 0. \end{cases} \quad (5.5)$$

Lemma 5.6. *There exists y_0 satisfying*

$$0 < y_0 < \theta^{\frac{m-1}{2}}$$

such that, with $\rho(t) = 2y_0\sqrt{t}$, problem (5.5) has a selfsimilar solution $v(x, t) = V(\frac{x}{2\sqrt{t}})$.

Proof. The proof is divided into two steps.

Step 1. An auxiliary problem. We first consider the following initial value problem:

$$\begin{cases} \xi''(y) = -2y(\xi^{\frac{1}{m}})'(y) = -2\frac{y}{m}\xi^{\frac{1-m}{m}}\xi'(y), & y > 0, \\ \xi(0) = \theta^m, & \xi'(0) = -2\theta^{\frac{m+1}{2}}. \end{cases}$$

It is easily seen that, for any $h \in (0, \theta^m)$, the constant function $\xi \equiv h$ is a solution of the equation. Hence, it follows from $\xi'(0) < 0$ that $\xi'(y) < 0$ as long as $\xi > 0$. In other words, there exists $y_0 \in (0, \infty]$ such that

$$\xi'(y) < 0, \quad \xi(y) > 0, \quad 0 \leq y < y_0.$$

We now show that $y_0 < \infty$ and so $\xi(y_0) = 0$. In fact, for any $\tilde{y}, y \in (0, y_0)$ with $\tilde{y} < y$ we have

$$\xi''(s) < -2\frac{\tilde{y}}{m}\xi^{\frac{1-m}{m}}(s)\xi'(s), \quad 0 < s < \tilde{y}.$$

Integrating it over $[0, \tilde{y}]$, we have

$$\xi'(\tilde{y}) < -2\theta^{\frac{m+1}{2}} + 2\tilde{y}(\theta - \xi^{\frac{1}{m}}(\tilde{y})) < -2\theta^{\frac{m+1}{2}} + 2\theta\tilde{y}.$$

Integrating it again over $[0, y]$, we have

$$\xi(y) < \tilde{\xi}(y) := \theta^m - 2\theta^{\frac{m+1}{2}}y + \theta y^2.$$

Since $\tilde{\xi}(y)$ has a unique zero $y_* := \theta^{\frac{m-1}{2}}$, we conclude that

$$y_0 < y_*.$$

Step 2. Selfsimilar solution. A direct calculation shows that

$$u(x, t) = \xi^{\frac{1}{m}}\left(\frac{x}{2\sqrt{t}}\right)$$

is a selfsimilar solution of

$$\begin{cases} u_t = (u^m)_{xx}, & 0 < x < \rho(t), \quad t > 0, \\ u(0, t) = \theta, & u(\rho(t), t) = 0, \quad t > 0, \end{cases}$$

with $\rho(t) := 2y_0\sqrt{t}$. Therefore,

$$v(x, t) = V\left(\frac{x}{2\sqrt{t}}\right) := \frac{m}{m-1}\xi^{\frac{m-1}{m}}\left(\frac{x}{2\sqrt{t}}\right)$$

is a selfsimilar solution of the first equation in (5.5). It also satisfies the boundary conditions in (5.5) at $x = 0$ and $x = \rho(t)$. Moreover, the interface $x = \rho(t)$ satisfies the Darcy law (cf. [30, Theorem 15.19]):

$$\rho'(t) = -v_x(\rho(t), t).$$

In other words, we have $V'(y_0) = -2y_0$. This proves the lemma. $\blacksquare$

This lemma implies that $V(\frac{x}{2\sqrt{t}})$ is not only a solution of the equation in (5.5), but also a solution of the following Cauchy–Dirichlet problem:

$$\begin{cases} v_t = (m-1)vv_{xx} + v_x^2, & x > 0, t > 0, \\ v(0, t) = \Theta := \frac{m}{m-1}\theta^{m-1}, & t > 0, \end{cases}$$

with one free boundary $\rho(t)$.

Proposition 5.7. *Assume f satisfies (f_c) , u is a transition solution of (CP). Then its left and right free boundaries $l(t)$ and $r(t)$ satisfy the asymptotic speed as in (1.11).*

Proof. The conclusion can be proved in a similar way as the authors did for the reaction-diffusion equation in [13, Proposition 4.2]. Note that there are several key points in the proof:

- (1) using the zero number argument to study the number of intersection points between two solutions. Here we see that the argument remains valid for our problem by Lemma 3.3 and by the Darcy law;
- (2) using a selfsimilar solution satisfying the Darcy law (which is called the Stefan boundary condition in [13]) to construct lower and supersolutions. Here we only need to use the selfsimilar solution $V(\frac{x}{2\sqrt{t}})$ obtained in the previous lemma to replace the selfsimilar solution $\Phi(t, x)$ in [13];
- (3) using the limit $\frac{\theta(t)}{r(t)} \rightarrow 0$ as $t \rightarrow \infty$, as an extra condition, where $\theta(t)$ is the θ -level set of $u(\cdot, t)$. This limit remains true for our problem and the proof is the same as that in [13, Section 4.3]. $\blacksquare$

6. Bistable RPME

In this section, we study the asymptotic behavior for the solutions of (CP) or (pCP) with bistable type of reaction: f satisfies (f_b) .

6.1. Stationary solutions

On the nonnegative stationary solutions of the equation in (CP), we first have Cases A–D types as in the monostable RPME. The difference is that in Case A the solutions are U_q for $q = 0, \theta, 1$ in the current case; in Case D, the compactly supported stationary solutions U_{q_0} exists for $q_0 \in (\theta_1, 1)$, where θ_1 is defined below by (6.1). The decay rates

in (4.4) and (4.5) remain valid. As we mentioned in Remark 4.1, each U_{q_0} is just a stationary solution of the equation but not of the Cauchy problem (CP).

Besides these stationary solutions, for the bistable RPME, there are two other kinds of stationary solutions: periodic solutions and the *ground state solution*.

Case E. In the (q, p) -phase plane, besides $(0, 0)$ and $(1, 0)$, we have another singular point $(\theta, 0)$, which is a center. There are infinitely many closed trajectories surrounding $(\theta, 0)$. Each of them corresponds to a positive stationary periodic solution $U_{\text{per}}(x)$. By the monotonicity result in Corollary 3.7, none of them can be a candidate of the ω -limit for the solution of (CP) with $u_0 \in \mathcal{X}$.

Case F. In the phase plane, there is a homoclinic orbit which starts and ends at $(0, 0)$, crossing the point $(\theta_1, 0)$, with $\theta_1 \in (\theta, 1)$ given by

$$\int_0^{\theta_1} r^{m-1} f(r) dr = 0. \quad (6.1)$$

The function of this homoclinic orbit is

$$p^2 = -2m \int_0^q r^{m-1} f(r) dr, \quad 0 < q \leq \theta_1.$$

Denote the corresponding solution by $U(x)$, and assume it satisfies the normalized condition

$$U(0) = \theta_1, \quad U'(0) = 0.$$

Assume the largest interval where U remains positive is $(-L, L)$ for some $L \in (0, \infty]$. We can give a formula for L as the following. Denote $\eta := U^m$, then

$$\frac{d\eta}{dx} = p(\eta) := \left(-2m \int_0^{\eta^{\frac{1}{m}}} r^{m-1} f(r) dr \right)^{\frac{1}{2}}, \quad x \in (-L, 0]. \quad (6.2)$$

Therefore,

$$L := \int_0^{\theta_1^m} \frac{d\eta}{p(\eta)} = \int_0^{\theta_1^m} \left(-2m \int_0^{\eta^{\frac{1}{m}}} r^{m-1} f(r) dr \right)^{-\frac{1}{2}} d\eta.$$

Under the assumptions in (f_b) , we have $f'(0) < 0$ and so $f(q) \sim f'(0)q$ as $q \rightarrow 0+0$. We will see below that $L < \infty$ in this case. However, L might be ∞ if f is a bistable nonlinearity but without the assumption $f'(0) < 0$.

Lemma 6.1. *Assume $f \in C([0, \infty)) \cap C^1((0, \infty))$ is a bistable nonlinearity satisfying (1.3). Further, assume that*

$$f(q) = -\lambda q^\alpha (1 + o(1)) \quad \text{as } q \rightarrow 0+0, \quad (6.3)$$

for some $\lambda > 0$ and $\alpha > 0$. Then $L = \infty$ when $\alpha \geq m$; $L < \infty$ when $0 < \alpha < m$.

Proof. Substituting (6.3) into the formula $p(\eta)$ in (6.2), we obtain

$$p(\eta) \sim \left(\frac{2m\lambda}{m+\alpha} \right)^{\frac{1}{2}} \eta^{\frac{m+\alpha}{2m}} \quad \text{as } \eta \rightarrow 0+0. \quad (6.4)$$

Case 1. $\alpha \geq m$. When $0 < \delta \ll 1$, we have

$$L > \int_0^\delta \left(\frac{m+\alpha}{3m\lambda} \right)^{\frac{1}{2}} \eta^{-\frac{m+\alpha}{2m}} d\eta = \infty.$$

Thus, the solution $U(x) > 0$ in $\mathbb{R}$, it is a Type I ground state solution, as that in the bistable RDEs.

Case 2. $0 < \alpha < m$. When $0 < \delta \ll 1$, we have

$$L < \int_\delta^{\theta_1^m} \frac{d\eta}{p(\eta)} + \int_0^\delta \left(\frac{m+\alpha}{m\lambda} \right)^{\frac{1}{2}} \eta^{-\frac{m+\alpha}{2m}} d\eta < \infty.$$

By (6.4) and (6.2), we have

$$\frac{d\eta}{dx} \sim \left(\frac{2m\lambda}{m+\alpha} \right)^{\frac{1}{2}} \eta^{\frac{m+\alpha}{2m}} \quad \text{as } x \rightarrow -L + 0,$$

and so

$$\frac{2m}{m-\alpha} \left(\frac{m+\alpha}{2m\lambda} \right)^{\frac{1}{2}} \left(\eta^{\frac{m-\alpha}{2m}} \right)' \sim 1 \quad \text{as } x \rightarrow -L + 0.$$

Hence,

$$U(x) = \eta^{\frac{1}{m}}(x) \sim A_1(\alpha, \lambda)(x+L)^{\frac{2}{m-\alpha}} \quad \text{as } x \rightarrow -L + 0, \quad (6.5)$$

with

$$A_1(\alpha, \lambda) := \left(\frac{m-\alpha}{2m} \right)^{\frac{2}{m-\alpha}} \left(\frac{2m\lambda}{m+\alpha} \right)^{\frac{1}{m-\alpha}}.$$

From (6.5) we see that, at the boundaries $\pm L$, U is continuous for all $m > \alpha$, Lipschitz when $\alpha < m \leq 2 + \alpha$ and C^1 when $\alpha < m < 2 + \alpha$.

If we denote by $V(x) := \frac{m}{m-1} U^{m-1}$ the corresponding pressure of U , then we obtain

$$\begin{cases} V \in C([-L, L]), & V(0) = \Theta_1 := \frac{m}{m-1} \theta_1^{m-1}, & V(\pm L) = 0, \\ V(x) > 0, & V'(x) < 0, & V(x) = V(-x) \quad \text{for } x \in (0, L), \\ V(x) \sim A(x+L)^{\frac{2(m-1)}{m-\alpha}} & \text{as } x \rightarrow -L + 0 & \text{with } A := \frac{m}{m-1} A_1^{m-1}. \end{cases} \quad (6.6)$$

This proves the lemma. ■

Remark 6.2. In the special case when f is a bistable nonlinearity with $f'(0) < 0$ as in (f_b) , we have

$$\alpha = 1 \quad \text{and} \quad \lambda = -f'(0),$$

and so V in (6.6) satisfies:

$$V(x) \sim A(x+L)^2 \quad \text{as } x \rightarrow -L + 0 \quad \text{with } A := \frac{-(m-1)f'(0)}{2(m+1)}.$$

Thus, $V'(-L+0) = 0$, and so V or $U = (\frac{m-1}{m} V)^{\frac{1}{m-1}}$ is a Type II ground state solution. By Lemma 2.3, $V(x)$ is a stationary solution not only for the bistable RPME, but also for

the Cauchy problem (pCP). This is different from the compactly supported solutions in Case D, where they are only stationary solutions of the RPME but not of (pCP). Furthermore, the combinations of such solutions, like

$$\mathcal{V}(x) := V(x - z_1) + V(x - z_2) \quad \text{with } z_2 - z_1 \geq 2L, \quad (6.7)$$

are also Type II ground state stationary solutions of problem (pCP).

6.2. Asymptotic behavior

Proof of Theorem 1.6. We prove the trichotomy result in Theorem 1.6. By $f'(0) < 0$, it follows from Remark 6.2 that the problem has only Type II ground state solution as in (6.6) or (6.7). By the general convergence result from Theorem 1.1, we know that u converges as $t \rightarrow \infty$ to 0, θ , 1, or a Type II ground state solution.

The constant solution θ can be excluded easily from the candidates. In fact, for any positive stationary periodic solution $U_{\text{per}}(x)$ of the equation (see Case E in Section 6.1), by the classical zero number argument we see that the zero number of $u_\sigma(\cdot, t) - U_{\text{per}}(\cdot)$ is finite and decreasing in $t > 0$. However, if $u_\sigma \rightarrow \theta$ for some $\sigma > 0$, then the above zero number tends to that of $\theta - U_{\text{per}}(\cdot)$, which is infinite. So, we derive a contradiction.

For the initial data $\phi_\sigma \in \mathcal{X}$, when $\sigma > 0$ is small, we have by (Φ_3) that $\phi_\sigma \leq \theta$. Hence, an argument similar to that in Lemma 5.2 implies that vanishing happens for u_σ . Denote

$$\Sigma_0 := \{\sigma > 0 \mid \text{vanishing happens for } u_\sigma\}.$$

Then Σ_0 is an open interval $(0, \sigma_*)$ for some $\sigma_* \in (0, \infty]$, as in the combustion equation.

In the case $\sigma_* = \infty$ (this is the complete vanishing case, as stated in Theorem 7.1), there is nothing left to prove.

In the case σ_* is a positive number, we can show that u_{σ_*} is a transition solution, that is, it converges to some Type II ground state solution $\mathcal{U}$. Otherwise, it tends to 1. Then the set

$$\Sigma_1 := \{\sigma > 0 \mid u_\sigma \rightarrow 1 \text{ as } t \rightarrow \infty\}$$

is not empty. Since this set is open as proved in the combustion case and since $\sigma_* \in \Sigma_1$ by our assumption, we see that $\sigma \in \Sigma_1$ for any σ with $0 < \sigma_* - \sigma \ll 1$. This contradicts the definition of σ_* . In what follows, we assume $u(x, t)$ converges to a Type II ground state solution $\mathcal{U}(x)$ as in (1.7).

If Σ_1 is empty, then u_σ is a transition solution for all $\sigma \geq \sigma_*$ (this is the case $\sigma_* \in (0, \infty)$ and $\sigma^* = \infty$ in Theorem 1.6), and the proof is also completed. In the case where Σ_1 is a non-empty open set, by the comparison principle, it is actually an interval (σ^*, ∞) for some $\sigma^* \geq \sigma_*$. For each $\sigma \in [\sigma_*, \sigma^*]$, u_σ is a transition solution.

This completes the proof of Theorem 1.6. ■

On the sharpness of the transition solution: $\sigma_* = \sigma^*$, we give some sufficient conditions.

Lemma 6.3. *Let u_σ be the solution of (CP) with bistable f and with initial data ϕ_σ satisfying (Φ_1) – (Φ_3) . Assume u_{σ_*} is a transition solution, then, for any $\sigma > \sigma_*$, u_σ is no longer a transition one, if one of the following conditions holds:*

- (i) (Φ_2) is strengthened as $\phi_{\sigma_*}(x) \leq \phi_\sigma(x + \varepsilon)$ when $|\varepsilon|$ is small;
- (ii) *there is $i \in \{1, 2, \dots, k\}$ such that $r_i(t) - l_i(t) \neq r_{*i}(t) - l_{*i}(t)$, where $l_i(t)$, $r_i(t)$ are the left and right boundaries of the i -th connected component of the support $\text{spt}[u_\sigma(\cdot, t)]$, $l_{*i}(t)$, $r_{*i}(t)$ are the analogs of $\text{spt}[u_{\sigma_*}(\cdot, t)]$;*
- (iii) $l_i(t) \equiv l_{*i}(t)$, $r_i(t) \equiv r_{*i}(t)$ for all $i = \{1, 2, \dots, k\}$, and one of the waiting times of these boundaries is finite.

Proof. In a similar way as proving Lemma 5.4, we see that any of the three conditions (i)–(iii) implies that, for some $T \geq 0$, $\tau \geq 0$ and $\varepsilon_0 > 0$, there holds

$$u_{\sigma_*}(x, t) \leq u_\sigma(x + \varepsilon, t + \tau), \quad x \in [l_{*i}(t), r_{*i}(t)], \quad t \geq T, \quad \varepsilon \in [0, \varepsilon_0] \text{ or } \varepsilon \in [-\varepsilon_0, 0].$$

In the case where u_σ is also a transition solution, by taking limit as $t \rightarrow \infty$ in this inequality we conclude that

$$U(x - z_*) \leq U(x + \varepsilon - z_\sigma), \quad x \in [l_*^\infty, r_*^\infty], \quad \varepsilon \in [0, \varepsilon_0] \text{ or } \varepsilon \in [-\varepsilon_0, 0],$$

for some z_* , $z_\sigma \in [-b, b]$, where

$$l_*^\infty = \lim_{t \rightarrow \infty} l_{*i}(t), \quad r_*^\infty = \lim_{t \rightarrow \infty} r_{*i}(t).$$

This is impossible since U is a Type II ground state solution. ■

Remark 6.4. The only situation not included in (ii) and (iii) in the previous lemma is

$$l_i(t) \equiv l_{*i}(t) \equiv l_i(0), \quad r_i(t) \equiv r_{*i}(t) \equiv r_i(0) \quad \text{for all } i = \{1, 2, \dots, k\},$$

and all the waiting times of these boundaries are infinite.

We guess that, even in this case, the transition solution is also unique. The instability of Type II ground state solution should be the essential reason.

Remark 6.5. In the case where the initial data ϕ_σ has a unique maximum point, so does $u(\cdot, t)$. Then the limit of the transition solution is a Type II ground state solution $U(x - z)$ for some $z \in [-b, b]$. Even in this case, the transition solution may have a very wide support but just converges to a positive limit in a bounded interval. For example, assume $u(x, t; \sigma(b^2 - x^2))$ is a transition solution for some $\sigma > 0$, and $b > L$, then

$$\begin{aligned} u(x, t) &> 0 \quad \text{for } x \in (l(t), r(t)), \quad t > 0, \\ u(x, t) &\leq u(\pm L, t), \quad x \in (l(t), -L] \cup [L, r(t)), \quad t > 0, \end{aligned}$$

and

$$u(\pm L, t) \rightarrow 0, \quad \|u(\cdot, t) - U(\cdot)\|_{L^\infty([l(t), r(t)])} \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

7. Complete vanishing phenomenon

Note that Theorem 1.3 gives the hair-trigger effect for (CP) with monostable reactions (see also Remarks 1.4 and 4.3). In the bistable and combustion cases, however, Theorems 1.5 and 1.6 show that vanishing happens for the solutions u_σ when $\sigma < \sigma_*$, where $\sigma_* \in (0, \infty]$. A natural question is this: is it really possible that $\sigma_* = \infty$? In other words, is there $b > 0$ such that, for any initial data u_0 with support in $[-b, b]$, vanishing always happens for u , no matter how large $\|u_0\|_{L^\infty}$ is? We call such a result as a *complete vanishing phenomenon*. Of course, this phenomenon seems difficult in most cases. However, it does happen for some problems. Recently, Li and Lou [24] proved that such a phenomenon really happens in bistable and combustion RDEs provided the nonlinearity of $f(u)$ decreases sufficiently fast as $u \rightarrow \infty$. Also, it was shown in [12] that this phenomenon occurs in the free boundary problems for RDEs with bistable, combustion or even monostable reactions. We now show that this phenomenon happens for (CP), provided $f(u)$ decreases fast as $u \rightarrow \infty$.

Theorem 7.1. *Assume $f \in C^2$ is a bistable or combustion reaction term. Assume also that*

$$\liminf_{u \rightarrow +\infty} \frac{-f(u)}{u^{p_0}} > 0 \quad (\text{F})_\infty$$

for some $p_0 > m$. Then there exists a small $b > 0$ such that vanishing always happens for the solution of (CP) provided $\text{spt}[u_0] \subset [-b, b]$.

Proof. Recall that when f is a bistable or combustion nonlinearity, there exists $K > 0$ such that

$$f(u) \leq Ku, \quad u \geq 0.$$

Using the pressure notation v and

$$g(v) := m \left(\frac{(m-1)v}{m} \right)^{\frac{m-2}{m-1}} f \left(\left(\frac{(m-1)v}{m} \right)^{\frac{1}{m-1}} \right),$$

we see that

$$g(v) \leq K(m-1)v \quad \text{for } v \geq 0. \quad (7.1)$$

Denote

$$p_1 := \frac{m-2+p_0}{m-1}, \quad p := \frac{p_0+3m-4}{2(m-1)}, \quad \alpha := \frac{2}{p-2}.$$

Then $p_1 - p = p - 2 = \frac{p_0-m}{2(m-1)} = \frac{2}{\alpha}$. By (F) $_\infty$, we have

$$\liminf_{v \rightarrow +\infty} \frac{-g(v)}{v^{p_1}} > 0.$$

So, there exist $L_0 > 0$ and $M > 1$ such that

$$g(v) \leq -L_0 v^{p_1-p} v^p \leq -L v^p, \quad v \geq M, \quad (7.2)$$

where $L := L_0 M^{p_1-p}$.

Recall that θ is a positive zero of f in the bistable and combustion equations. Choose $M > 1$ larger if necessary (to be determined below) and denote $M_0 := (\frac{(m-1)M}{m})^{\frac{1}{m-1}}$, then we can choose $s_0 > 0$, with $0 < 1 - K(m-1)s_0 \ll 1$ when M_0 is sufficiently large, such that

$$\theta \frac{(\frac{1-K(m-1)s_0}{2K})^{\frac{1}{m+1}}}{(1 + \frac{(m-1)-K(m^2-1)s_0}{2})^{\frac{1}{m-1}}} = 2M_0 s_0^{\frac{1}{m+1}} > 1. \quad (7.3)$$

Define $b_1 = C_1(m)s_0^{\frac{1}{m+1}}$ by

$$\frac{(m-1)b_1^2}{2m(m+1)s_0^{\frac{2}{m+1}}} = 1 - \frac{1}{2^{m-1}}.$$

Then, we can choose $M > 1$ sufficiently large such that

$$b := M^{-\frac{1}{\alpha}} \leq \min\left\{\frac{b_1}{3}, 1\right\},$$

$$L := L_0 M^{p+1-p} \geq \frac{3c}{b_1 M} + 2^p[(m-1)\alpha(\alpha+1) + \alpha^2],$$

where $c > 0$ is defined by

$$c := 2^{\frac{1}{\alpha}} b^{-1-\alpha} \alpha^{-1} [2(m-1)\alpha(\alpha+1) + 2\alpha^2 + K(m-1)].$$

Denote $C := (2M_0)^{m-1} s_0^{\frac{m-1}{m+1}} > 1$, then

$$\left(C - \frac{(m-1)b_1^2}{2m(m+1)s_0^{\frac{2}{m+1}}}\right)^{\frac{1}{m-1}} \geq \frac{1}{2} C^{\frac{1}{m-1}} = M_0 s_0^{\frac{1}{m+1}}. \quad (7.4)$$

We will show that the complete vanishing phenomenon happens if

$$\text{spt}[u(x, 0)] \subset [-b, b].$$

Step 1. We first consider the stage when the solution is suppressed by a spatial homogeneous upper solution down to the range $v \leq M$. Consider the equation

$$v_{1t} = -Lv_1^2.$$

Then $v_1 = (Lt)^{-1}$ is monotonically decreasing from $+\infty$, and till it reaches M at time $t = t_1 := (LM)^{-1}$, it is a supersolution of (pCP) due to (7.2) and $p > 2$, no matter how large $\|u_0\|_{L^\infty}$ is. So,

$$v(x, t) \leq v_1(t_1) = M, \quad t \geq t_1.$$

This inequality gives an estimate for v from above. We also need estimates on the right and left boundaries. We state only the case on the right-hand side since the left-hand side is studied similarly. For c, b given above, we define

$$v_2(x, t) := (x - ct - b)^{-\alpha} - b^{-\alpha}, \quad b + ct < x \leq 2b + ct, \quad t \geq 0.$$

We now show that v_2 is also a supersolution of (pCP) in the domain

$$E := \{(x, t) \mid b + ct < x \leq 2b + ct, t \geq 0\}.$$

(1) First, on the interval $J_1 := [b + ct + 2^{-\frac{1}{\alpha}}b, 2b + ct]$, we have

$$2^{-\frac{1}{\alpha}}b \leq H := x - ct - b \leq b, \quad 0 \leq v_2 \leq b^{-\alpha},$$

and so by the first inequality in (7.1) and the choice of c , we have

$$\begin{aligned} \mathcal{N}(v_2) &:= v_{2t} - (m-1)v_2v_{2xx} - v_{2x}^2 - g(v_2) \\ &\geq \alpha c H^{-\alpha-1} - (m-1)\alpha(\alpha+1)v_2H^{-\alpha-2} - \alpha^2H^{-2\alpha-2} - K(m-1)v_2 \\ &\geq H^{-\alpha-2}[\alpha c H - (m-1)\alpha(\alpha+1)H^{-\alpha} - \alpha^2H^{-\alpha}] - K(m-1)b^{-\alpha} \\ &\geq b^{-\alpha-2}[\alpha c 2^{-\frac{1}{\alpha}}b - 2(m-1)\alpha(\alpha+1)b^{-\alpha} - 2\alpha^2b^{-\alpha}] - K(m-1)b^{-\alpha} \\ &\geq b^{-2\alpha-2}[\alpha c 2^{-\frac{1}{\alpha}}b^{1+\alpha} - 2(m-1)\alpha(\alpha+1) - 2\alpha^2 - K(m-1)] \geq 0. \end{aligned}$$

(2) Next, in the interval $J_2 := (b + ct, b + ct + 2^{-\frac{1}{\alpha}}b]$, we have

$$0 < H := x - ct - b \leq 2^{-\frac{1}{\alpha}}b, \quad v_2 \geq b^{-\alpha} \geq M,$$

and by the choice of L , we have

$$\begin{aligned} \mathcal{N}(v_2) &\geq H^{-\alpha-2}[\alpha c H - (m-1)\alpha(\alpha+1)H^{-\alpha} - \alpha^2H^{-\alpha}] + Lv_2^p \\ &\geq L(H^{-\alpha} - b^{-\alpha})^p - [(m-1)\alpha(\alpha+1) + \alpha^2]H^{-2\alpha-2} \\ &\geq 2^{-p}LH^{-\alpha p} - [(m-1)\alpha(\alpha+1) + \alpha^2]H^{-2\alpha-2} \\ &\geq H^{-2\alpha-2}[2^{-p}L - (m-1)\alpha(\alpha+1) - \alpha^2] \geq 0. \end{aligned}$$

(3) On the free boundary $r_2(t) := 2b + ct$ of v_2 , by the choice of c , we easily obtain

$$-v_{2x}(2b + ct, t) = \alpha b^{-\alpha-1} \leq c, \quad t \geq 0.$$

Therefore, v_2 is a supersolution of (pCP) in the domain E . As a consequence, at t_1 , the right free boundary $r(t)$ of v satisfies

$$r(t_1) \leq r_2(t_1) = 2b + ct_1 \leq 2b + \frac{c}{LM} \leq b_1$$

by the choice of b, c and L . Similarly, the left free boundary $l(t)$ of v satisfies $l(t_1) \geq -b_1$.

Using the notation of u , we conclude from above that

$$u(x, t_1) \leq \bar{u}(x) := M_0 \cdot \chi_{[-b_1, b_1]}.$$

Step 2. Next, we use the ZKB solution as another upper solution to suppress u further, so that it goes down below θ after some time. More precisely, consider

$$\begin{cases} u_{3t} = (u_3^m)_{xx} + Ku_3, & x \in \mathbb{R}, t > 0, \\ u_3(x, 0) = \bar{u}(x), & x \in \mathbb{R}. \end{cases}$$

Then u_3 is a upper solution of (CP) and

$$u(x, t + t_1) \leq u_3(x, t), \quad x \in \mathbb{R}, \quad t > 0.$$

Change the time variable from t to s by

$$s := \frac{e^{K(m-1)t} - 1}{K(m-1)} \Leftrightarrow t = T(s) := \frac{\ln(1 + K(m-1)s)}{K(m-1)}$$

and define

$$w(x, s) := u_3(x, T(s))[1 + K(m-1)s]^{-\frac{1}{m-1}},$$

then we have

$$\begin{cases} w_s = (w^m)_{xx}, & x \in \mathbb{R}, \quad s > 0, \\ w(x, 0) = \bar{u}(x), & x \in \mathbb{R}. \end{cases} \quad (7.5)$$

To estimate $w(x, s)$, we consider the ZKB solution

$$w_1(x, s) := s^{-\frac{1}{m+1}} \left(C - \frac{(m-1)x^2}{2m(m+1)s^{\frac{2}{m+1}}} \right)_+^{\frac{1}{m-1}}, \quad x \in \mathbb{R}, \quad s \geq s_0,$$

with s_0 and C given above. By the choice of b_1 , C and s_0 , for $x \in [-b_1, b_1]$ we have

$$\begin{aligned} w_1(x, s_0) &\geq w_1(\pm b_1, s_0) = s_0^{-\frac{1}{m+1}} \left(C - \frac{(m-1)b_1^2}{2m(m+1)s_0^{\frac{2}{m+1}}} \right)^{\frac{1}{m-1}} \\ &\geq \frac{1}{2} s_0^{-\frac{1}{m+1}} C^{\frac{1}{m-1}} > M_0 \geq \bar{u}(x). \end{aligned}$$

Therefore, w_1 is a upper solution of (7.5), and so

$$w(x, s) \leq w_1(x, s + s_0), \quad x \in \mathbb{R}, \quad s > 0.$$

In particular, at $x = 0$ and $s = s_1 := \frac{1-K(m+1)s_0}{2K}$ we have

$$w(0, s_1) \leq w_1(0, s_1 + s_0) = (s_1 + s_0)^{-\frac{1}{m+1}} C^{\frac{1}{m-1}}.$$

Using the notation u , we have

$$\begin{aligned} u(x, T(s_1) + t_1) &\leq u_3(x, T(s_1)) \\ &= w(x, s_1)[1 + K(m-1)s_1]^{-\frac{1}{m-1}} \\ &\leq (s_1 + s_0)^{-\frac{1}{m+1}} C^{\frac{1}{m-1}} [1 + K(m-1)s_1]^{-\frac{1}{m-1}} \\ &\leq \theta, \end{aligned}$$

by (7.3) and (7.4).

Step 3. By the sufficient condition for vanishing in bistable and combustion RPMEs (cf. Lemma 5.2), we conclude that $u \rightarrow 0$ as $t \rightarrow \infty$.

This proves the complete vanishing phenomenon in Theorem 7.1. ■

Appendix A. Well-posedness and a priori estimates

For our RPME with general reaction terms, it seems that the a priori estimates, well-posedness and other important properties as that in PME are not well collected in literature, though they are not entirely new. We present some details here for the convenience of the readers.

A.1. Well-posedness

The existence of very weak solution of (CP) has been well studied by many authors (cf. [5, 9, 27, 30, 31]).

Proposition A.1. *Assume (I) and f is Lipschitz continuous with $f(0) = 0$. Then problem (CP) has a unique very weak solution $u(x, t) \in C(Q_T) \cap L^\infty(Q_T)$ for any $T > 0$, and*

$$0 \leq u(x, t) \leq M'_0 := \max\{1, \|u_0\|_{L^\infty}\} \quad \text{in } Q_T = \mathbb{R} \times (0, T).$$

Moreover, there are two free boundaries $l(t) < r(t)$ such that $\text{spt}(u(\cdot, t)) \subset [l(t), r(t)]$ for all $t > 0$.

We cannot obtain $u(x, t) > 0$ as in RDEs due to the degeneracy of the equation at $u = 0$. Nevertheless, the solution u solves the equation in the classical sense by the standard parabolic theory in a neighborhood of every point (x, t) at which the solution is positive (see also [26]).

A.2. Persistence of positivity and finite propagation speed

We now show the following positivity persistence property as in PME.

Proposition A.2 (Persistence of positivity). *Assume $u(x_0, t_0) > 0$ for some $x_0 \in \mathbb{R}$, $t_0 > 0$. Then $u(x_0, t) > 0$ for all $t \geq t_0$. (Thus, u is classical near the line $\{(x_0, t) \mid t > t_0\}$.)*

Proof. Since $t_0 > 0$, we have $u(x, t_0) \in C(\mathbb{R})$, and so, by our assumption there exist $\varepsilon > 0$, $\delta > 0$ such that

$$u(x, t_0) \geq \varepsilon \quad \text{for } x \in J := [x_0 - \delta, x_0 + \delta].$$

We use K to denote the Lipschitz constant for f . Denote

$$\beta := \frac{1}{m-1}, \quad \mu := K + \frac{\pi^2}{4\delta^2} \varepsilon^{m-1},$$

and define

$$\underline{u}(x, t) := \begin{cases} \varepsilon e^{-\mu t} \left(\sin \frac{\pi(x - x_0 + \delta)}{2\delta} \right)^{\frac{1}{m}}, & x \in J, t \geq 0, \\ 0, & x \in \mathbb{R} \setminus J, t \geq 0. \end{cases}$$

It is easily seen that

$$\underline{u}(x, 0) \leq \varepsilon \leq u(x, t_0), \quad x \in J,$$

and

$$\underline{u}_t - (\underline{u}^m)_{xx} - f(\underline{u}) \leq \underline{u}_t - (\underline{u}^m)_{xx} + K\underline{u} \leq 0, \quad x \in \mathbb{R}, t \geq 0.$$

By the comparison principle, we have

$$u(x, t + t_0) \geq \underline{u}(x, t) > 0, \quad |x - x_0| < \delta, t > 0.$$

This proves the proposition. $\blacksquare$

Remark A.3. A positive lower bound can also be given by a subsolution in the form of ZKB solution

$$\underline{u}_1(x, t) := \frac{1}{(t+1)^\beta e^{K(t+1)}} \left(C - \frac{(x-x_0)^2 e^{(m-1)K(t+1)}}{2m} \right)_+^\beta, \quad x \in \mathbb{R}, t \geq 0, \quad (\text{A.1})$$

with

$$w_+ := \max\{w(x, t), 0\} \quad \text{and} \quad C > 0$$

satisfying

$$2mC e^{-(m-1)K} \leq \delta^2, \quad C^\beta \leq \varepsilon e^K.$$

Using this subsolution, we can conclude that

$$u(x, t) \geq \underline{u}_1(x, t) > 0 \quad \text{if } |x - x_0| < \underline{s}(t) := (2mC)^{\frac{1}{\beta}} e^{-\frac{(m-1)K(t+1)}{2}}, t \geq 0.$$

Note that, using this estimate we have $u > 0$ in a shrinking domain $(x_0 - \underline{s}(t), x_0 + \underline{s}(t))$, while using $\underline{u}$ we have $u > 0$ in a fixed domain $(x_0 - \delta, x_0 + \delta)$.

When $u_0 \geq 0$ is compactly supported, besides the upper bound in Proposition A.1, we also have the *finite propagation speed* as in PME.

Proposition A.4. Assume (I) and f is Lipschitz continuous with $f(0) = 0$. Then the support of $u(\cdot, t)$ lies in $[-\bar{s}(t), \bar{s}(t)]$ with

$$\bar{s}(t) := [C_1(t+1)]^{\frac{1}{2}} e^{\frac{(m-1)K(t+1)}{2}}, \quad t > 0, \quad (\text{A.2})$$

for some C_1 depending only on m, K and $\|u_0\|_{L^\infty}$, and so $-l(t), r(t) \leq \bar{s}(t)$.

Proof. We construct a supersolution of the ZKB form. Set $A := (4m\beta)^{-\beta}$, choose $C_1 > 0$ such that

$$Ae^K \left(C_1 - \frac{b^2}{e^{(m-1)K}} \right)^\beta \geq \|u_0\|_{L^\infty},$$

where $b > 0$ is the number in (I), and define

$$\bar{u}(x, t) := Ae^{K(t+1)} \left(C_1 - \frac{x^2}{(t+1)e^{(m-1)K(t+1)}} \right)_+^\beta, \quad x \in \mathbb{R}, t \geq 0.$$

By the choose of C_1 , we see that $\bar{u}(x, 0) \geq u_0(x)$, and by a direct calculation, we have

$$\begin{aligned} \bar{u}_t &\geq (\bar{u}^m)_{xx} + K\bar{u} \geq (\bar{u}^m)_{xx} + f(\bar{u}), \\ |x| \leq \bar{s}(t) &:= [C_1(t+1)]^{\frac{1}{2}} e^{\frac{(m-1)K(t+1)}{2}}, \quad t \geq 0. \end{aligned}$$

Hence, $\bar{u}$ is a supersolution of (CP), and the conclusion follows from the comparison. $\blacksquare$

Remark A.5. Our construction for the subsolution in (A.1) and the supersolution in this proof is inspired by the ZKB solution, but with obvious difference due to the presence of the linear term Ku .

A.3. A priori estimate for the pressure

As we have mentioned in Section 2, the pressure $v(x, t) := \frac{m}{m-1}[u(x, t)]^{m-1}$ solves

$$\begin{cases} v_t = (m-1)v v_{xx} + v_x^2 + g(v), & x \in \mathbb{R}, t > 0, \\ v(x, 0) = v_0(x) := \frac{m}{m-1}u_0^{m-1}(x), & x \in \mathbb{R}, \end{cases} \quad (\text{pCP})$$

with

$$g(v) := m \left(\frac{(m-1)v}{m} \right)^{\frac{m-2}{m-1}} f \left(\left(\frac{(m-1)v}{m} \right)^{\frac{1}{m-1}} \right)$$

satisfying

$$|g(v)| \leq K(m-1)v, \quad v \geq 0,$$

by (F). Since u is bounded as in Proposition A.1, we see that v is also bounded,

$$0 \leq v(x, t) \leq M_0 := \max \left\{ \frac{m}{m-1}, \|v_0\|_{L^\infty} \right\}, \quad x \in \mathbb{R}, t > 0.$$

Hence,

$$|g(v)| \leq G_0(m, K, v_0) := K(m-1)M_0, \quad 0 \leq v \leq M_0.$$

By the above proposition, we have the following result.

Proposition A.6. Assume (I) and f is Lipschitz continuous with $f(0) = 0$. Then problem (pCP) has a unique nonnegative solution $v(x, t) \in C(Q_T) \cap L^\infty(Q_T)$ for any $T > 0$.

Moreover, if $v(x_0, t_0) > 0$ for some $x_0 \in \mathbb{R}$ and $t_0 > 0$, then $v(x_0, t) > 0$ for all $t \geq t_0$, and so v is a classical solution near the line $\{(x_0, t) \mid t \geq t_0\}$.

The support of $v(\cdot, t)$ lies in $[-\bar{s}(t), \bar{s}(t)]$ with $\bar{s}(t)$ given by (A.2).

In what follows, we continue to present the a priori estimates for v_x , v_{xx} and v_t under further restrictions on $f: f \in C^2$. In this case, we have

$$|g'(v)| \leq G_1(m, K, v_0) \quad \text{and} \quad |g''(v)| \leq G_2(m, K, v_0), \quad 0 \leq v \leq M_0.$$

In a similar way as Aronson [2] (with obvious modification caused by the reaction term $g(v)$), we have the following.

Lemma A.7 (Uniform estimate for v_x [2]). Assume (I) and $f \in C^1([0, \infty))$ with $f(0) = 0$. Let v be a smooth positive classical solution of (pCP) in $R := (a, b) \times (0, T]$ for some $a, b, T \in \mathbb{R}$ with $b > a$, $T > 0$. Then for any $0 < \delta < \frac{b-a}{2}$, $\tau < T$ there holds

$$|v_x(x, t)| \leq M_1(m, M_0, G_0, G_1, \delta, \tau), \quad (x, t) \in [a + \delta, b - \delta] \times [\tau, T]. \quad (\text{A.3})$$

Moreover, if $\tau = 0$ and

$$M_1^0 := \max_{[a,b]} |v'_0(x)| < \infty,$$

then (A.3) holds in $(a + \delta, b - \delta) \times (0, T]$ for $M_1 = M_1(m, M_0, G_0, G_1, \delta, M_1^0)$.

Note that the a priori bound M_1 of v_x obtained in this lemma is uniform in T due to the fact that the bound of v is uniform. The result holds for classical solutions. For the very weak solution in $C(Q_T) \cap L^\infty(Q_T)$, however, one can show the following Lipschitz continuity for v and Hölder continuity for u (which can be regarded as the limit of a sequence of classical solutions).

Proposition A.8 (Lipschitz continuity for v and Hölder continuity for u [2]). *Assume (I) and $f \in C^1([0, \infty))$ with $f(0) = 0$. If we further assume that u_0^m is Lipschitz continuous, then for $T > \tau > 0$,*

$$|v(x, t) - v(y, t)| \leq C_1(m, \tau, \|u_0\|_{L^\infty}) \cdot |x - y|, \quad x, y \in \mathbb{R}, t \in [\tau, T],$$

$$|u(x, t) - u(y, t)| \leq C_2(m, \tau, \|u_0\|_{L^\infty}) \cdot |x - y|^v, \quad x, y \in \mathbb{R}, t \in [\tau, T],$$

$$v := \min\left\{1, \frac{1}{m-1}\right\}$$

and $\frac{\partial u^m}{\partial x}(x, t)$ exists and is continuous in $x \in \mathbb{R}$, with

$$\frac{\partial u^m}{\partial x}(x, t) = 0 \quad \text{if } u(x, t) = 0.$$

Proof. The proof follows from the previous lemma and the fact that the very weak solution u is the pointwise limit of a decreasing sequence of positive classical functions. (For the PME, this was shown in [26]. For the current problem, the proof is similar.) ■

Next, we consider the lower bound of v_{xx} . For the PME, Aronson and B enilan [4] gave a lower bound for v_{xx} in 1979,

$$v_{xx} \geq -\frac{1}{(m+1)t}, \quad t > 0. \quad (\text{A.4})$$

This inequality, usually known as the Aronson–B enilan estimate, is understood in the sense of distributions. It is optimal in the sense that equality is actually attained by the source-type or ZKB solutions. It is a significant novelty of the Cauchy problem, and is used so often in the theory of nonnegative solutions in the whole space. For our problem (pCP), we will also give a lower bound for v_{xx} , but with quite different order from (A.4).

Proposition A.9. *Assume (F), (I), and $v_0 \in C^2$ a.e. in $\mathbb{R}$, with*

$$0 \leq v_0(x) \leq M_0^0, \quad M_1^0 := \sup_{\mathbb{R}} |v'_0(x)| < \infty, \quad v''_0(x) \geq -M_2^0 > -\infty, \quad \text{a.e. } x \in \mathbb{R}.$$

Let $v \in C(Q_T) \cap L^\infty(Q_T)$ for any $T > 0$ be the solution of (pCP). Then

$$v_{xx}(x, t) \geq -C(t + \tau), \quad x \in \mathbb{R}, t \in (0, T],$$

for some $\tau = \tau(m, M_0^0, M_1^0, M_2^0, G_0, G_1, G_2)$ and $C = C(m, M_0^0, M_1^0, G_0, G_1, G_2)$, here the inequality holds in the sense of distributions in $\mathbb{R} \times (0, T]$.

Proof. We follow the idea in Aronson and B enilan [4] (see also [30,  9.3]).

(1) First, we assume $v > 0$ in $Q_T := \mathbb{R} \times (0, T]$. In this case, v is a classical solution of (pCP). We write the equation satisfied by $\eta := v_{xx}$ by differentiating the equation of v twice:

$$\eta_t = \mathcal{L}(\eta) + g''v_x^2$$

with

$$\mathcal{L}(\eta) := (m-1)v\eta_{xx} + 2mv_x\eta_x + (m+1)\eta^2 + g'(v)\eta.$$

Note that here we use the C^2 smoothness assumption for f . By Lemma A.7, we have

$$|v_x(x, t)| \leq M_1(m, M_0^0, M_1^0, G_0, G_1), \quad x \in \mathbb{R}, \quad t > 0.$$

Hence,

$$\eta_t \geq \mathcal{L}(\eta) - G_2(M_1 + 1)^2.$$

Now we construct a subsolution of this operator. Define

$$\tau_1 := \frac{G_1 + (m+1)M_2^0}{(m+1)G_2(M_1 + 1)^2}.$$

Then $\zeta(x, t) := -G_2(M_1 + 1)^2(t + \tau_1)$ satisfies

$$\begin{aligned} \zeta_t - \mathcal{L}(\zeta) + G_2(M_1 + 1)^2 \\ = G_2(M_1 + 1)^2(t + \tau_1)[G_1 - (m+1)G_2(M_1 + 1)^2(t + \tau_1)] \leq 0. \end{aligned}$$

In addition,

$$\zeta(x, 0) = -G_2(M_1 + 1)^2\tau_1 < -M_2^0 \leq v_0''(x) = \eta(x, 0+), \quad x \in \mathbb{R}.$$

By comparison, we have

$$v_{xx}(x, t) = \eta(x, t) \geq \zeta(x, t) = -G_2(M_1 + 1)^2(t + \tau_1), \quad x \in \mathbb{R}, \quad t > 0.$$

(2) We now construct a family of approximate positive solutions to approach the non-negative general solution. Set

$$u_{0\varepsilon}(x) := u_0(x) + \varepsilon, \quad x \in \mathbb{R}, \quad 0 < \varepsilon \ll 1.$$

Then the corresponding pressure is $v_{0\varepsilon} := \frac{m}{m-1}u_{0\varepsilon}^{m-1}$, which satisfies

$$0 < \varepsilon \leq v_{0\varepsilon}(x) \leq M_0^0 + 1, \quad |v_{0\varepsilon}'(x)| \leq M_1^0, \quad v_{0\varepsilon}''(x) \geq -M_2^0, \quad x \in \mathbb{R}.$$

According to the standard theory, problem (CP) with initial data $u_{0\varepsilon}$ has a unique solution $u_\varepsilon(x, t)$. In a similar way as in the previous step, we see that the corresponding pressure v_ε satisfies

$$0 < v_\varepsilon(x, t) \leq \max\left\{\frac{m}{m-1}, M_0^0 + 1\right\}, \quad x \in \mathbb{R}, \quad t > 0, \quad 0 < \varepsilon \ll 1,$$

$v_{\varepsilon x}$ satisfies

$$|v_{\varepsilon x}(x, t)| \leq M'_1(m, M_0^0, M_1^0, G_0, G_1), \quad x \in \mathbb{R}, t > 0, 0 < \varepsilon \ll 1,$$

and $v_{\varepsilon xx}$ satisfies

$$v_{\varepsilon xx}(x, t) \geq -C(t + \tau), \quad x \in \mathbb{R}, t \in (0, T], 0 < \varepsilon \ll 1, \quad (\text{A.5})$$

for

$$C = C(m, M_0^0, M_1^0, G_0, G_1, G_2) \quad \text{and} \quad \tau = \tau(m, M_0^0, M_1^0, M_2^0, G_0, G_1, G_2).$$

Since v_ε is strictly increasing in ε and since $v_\varepsilon \geq v$, there exists $\hat{v} \geq v \geq 0$ such that

$$\lim_{\varepsilon \rightarrow 0} v_\varepsilon(x, t) = \hat{v}(x, t) \quad \text{in the topology of } C_{\text{loc}}(Q_T).$$

(In particular, $\hat{v} \in C(Q_T) \cap L^\infty(Q_T)$.) Furthermore, in the area $D := \{(x, t) \in Q_T \mid \hat{v}(x, t) > 0\}$, by standard interior parabolic estimates we see that the convergence holds also in the topology $C_{\text{loc}}^{2,1}(D)$. Using the Lebesgue theorem to take limit as $\varepsilon \rightarrow 0$ in

$$\iint [v_\varepsilon \varphi_{xx} + C(t + \tau) \varphi] dx dt = \iint [v_{\varepsilon xx} + C(t + \tau) \varphi] dx dt \geq 0$$

for every $\varphi \in C_c^\infty(Q_T)$, $\varphi \geq 0$, we obtain

$$\iint [\hat{v} \varphi_{xx} + C(t + \tau) \varphi] dx dt \geq 0.$$

This means that (A.5) holds for $\hat{v}$ in the distribution sense.

To complete our proof, we only need to show that $\hat{v} \equiv v$. Denote by

$$\hat{u} = \left(\frac{m-1}{m} \hat{v} \right)^{\frac{1}{m-1}} \in C(Q_T) \cap L^\infty(Q_T)$$

the original density variable corresponding to the pressure $\hat{v}$. Recalling that u_ε is a classical solution of (CP) with initial data $u_{0\varepsilon}$, we have

$$\begin{aligned} \int_{\mathbb{R}} u_\varepsilon(x, T) \varphi(x, T) dx &= \int_{\mathbb{R}} u_{0\varepsilon}(x) \varphi(x, 0) dx + \iint_{Q_T} f(u_\varepsilon) \varphi dx dt \\ &\quad + \iint_{Q_T} [u_\varepsilon \varphi_t + u_\varepsilon^m \varphi_{xx}] dx dt. \end{aligned}$$

Taking limit as $\varepsilon \rightarrow 0$, we have

$$\begin{aligned} \int_{\mathbb{R}} \hat{u}(x, T) \varphi(x, T) dx &= \int_{\mathbb{R}} u_0(x) \varphi(x, 0) dx + \iint_{Q_T} f(\hat{u}) \varphi dx dt \\ &\quad + \iint_{Q_T} [\hat{u} \varphi_t + \hat{u}^m \varphi_{xx}] dx dt. \end{aligned}$$

This means that $\hat{u}$ is a very weak solution of (CP) with $\hat{u}(x, 0) = u_0$. By the uniqueness, $\hat{u}$ must be u , and so $\hat{v} \equiv v$. ■

Finally, we show that v is also Lipschitz in time.

Lemma A.10. *Assume the hypotheses in Proposition A.9 hold. Then for any $\tau > 0$ there holds*

$$v_t(x, t) \leq C_1 t + C_2, \quad x \in \mathbb{R}, \quad t \geq \tau, \quad (\text{A.6})$$

where C_1 depends on m, M_1^0, G_0 and G_1 , C_2 depends on $m, G_0, G_1, v_t(x, \tau)$, and

$$v_t \geq -C_3 t - C_4, \quad x \in \mathbb{R}, \quad t > 0,$$

where C_3 and C_4 depend on $m, M_0^0, M_1^0, G_0, G_1, G_2$.

Proof. By approximation, we may assume that v is positive and smooth. In this case, we have

$$0 < v \leq 1 + \|v_0\|_{L^\infty}, \quad |v_x(x, t)| \leq M_1(m, M_1^0, G_0, G_1), \quad x \in \mathbb{R}, \quad t > 0,$$

as above. We consider the function

$$P(x, t) := v_t + (m-1)v_x^2 = (m-1)vv_{xx} + mv_x^2 + g(v).$$

Then

$$\begin{aligned} P_x &= (m-1)vv_{xxx} + (3m-1)v_x v_{xx} + g'(v)v_x, \\ P_{xx} &= v_{txx} + 2(m-1)(v_{xx}^2 + v_x v_{xxx}), \\ P_t &= (m-1)vv_{txx} + (m-1)v_t v_{xx} + 2m(m+1)v_x^2 v_{xx} + 2m(m-1)vv_x v_{xx} \\ &\quad + 2mv_x g'(v)v_x + g'(v)v_t. \end{aligned}$$

Hence, we have

$$\begin{aligned} \mathcal{L}(P) &:= P_t - (m-1)vP_{xx} - 2v_x P_x = \tilde{g}(x, t, P) \\ &:= \frac{1}{v}[-(P-g)^2 + A_1(P-g) + A_2] + A_3, \end{aligned}$$

with

$$\begin{aligned} A_1 &:= (4m-1)v_x^2 + vg', \\ A_2 &:= -m(3m-1)v_x^4, \\ A_3 &:= 2mv_x g'(v)v_x + g'(v)(g - (m-1)v_x^2). \end{aligned}$$

By $f \in C^2$ and the bounds of v, v_x , we see that

$$|A_i(x, t)| \leq \bar{A}_i, \quad x \in \mathbb{R}, \quad t \geq \tau, \quad i = 1, 2, 3.$$

Choose

$$C_1 := \bar{A}_3, \quad C_2 := G_0 + \frac{\bar{A}_1 + \sqrt{\bar{A}_1^2 + 4\bar{A}_2}}{2} + (m-1)M_1^2 + \sup_{x \in \mathbb{R}} v_t(x, \tau),$$

then

$$-(C_1 t + C_2 - g)^2 + A_1(C_1 t + C_2 - g) + A_2 \leq 0,$$

and so

$$\mathcal{L}(C_1 t + C_2) - \tilde{g}(x, t, C_1 t + C_2) \geq 0$$

and

$$(C_1 t + C_2)|_{t=\tau} > C_2 > P(x, \tau).$$

Hence, $C_1 t + C_2$ is a supersolution of $\mathcal{L}(P) = \tilde{g}(x, t, P)$ in $t \geq \tau$, and so (A.6) follows from the maximum principle. Note that the auxiliary function P was used for PME in [30, Chapter 15], but the supersolution they used is $-\frac{C}{t}$, different from ours.

The lower bound for v_t is an immediate consequence of Proposition A.9:

$$v_t = (m-1)vv_{xx} + v_x^2 + g(v) \geq -C(m-1)(t+\tau)(1 + \|v_0\|_{L^\infty}) - G_0.$$

This proves the Lipschitz continuity for v in time. ■

Acknowledgments. We would like to thank the anonymous referees for their valuable suggestions and helpful comments.

Funding. Maolin Zhou is partially supported by National Key Research and Development Program of China (2021YFA1002400, 2020YFA0713300), NSF of China (12271437, 11971498), the Fundamental Research Funds for the Central Universities (Nankai University 63241642), Tianjin Natural Science Foundation Key Project (22JCZDJC00370), Tianjin Natural Science Foundation Outstanding Youth Project (23JCQJC00190) and Nankai Zhide Foundation. Bendong Lou is partially supported by the NSF of China (12071299, 12471199).

References

- [1] Angenent, S.: [The zero set of a solution of a parabolic equation](#). J. Reine Angew. Math. **390**, 79–96 (1988) Zbl [0644.35050](#) MR [953678](#)
- [2] Aronson, D. G.: [Regularity properties of flows through porous media](#). SIAM J. Appl. Math. **17**, 461–467 (1969) Zbl [0187.03401](#) MR [247303](#)
- [3] Aronson, D. G.: [Density-dependent interaction-diffusion systems](#). In: Dynamics and modelling of reactive systems (Proc. Adv. Sem., Math. Res. Center, Univ. Wisconsin, Madison, Wis., 1979), Publ. Math. Res. Center 44, Academic Press, New York, 161–176 (1980) MR [588018](#)
- [4] Aronson, D. G., Bénilan, P.: Régularité des solutions de l'équation des milieux poreux dans $\mathbf{R}^N$. C. R. Acad. Sci. Paris Sér. A-B **288**, 103–105 (1979) Zbl [0397.35034](#) MR [524760](#)
- [5] Aronson, D. G., Crandall, M. G., Peletier, L. A.: [Stabilization of solutions of a degenerate nonlinear diffusion problem](#). Nonlinear Anal. **6**, 1001–1022 (1982) Zbl [0518.35050](#) MR [678053](#)
- [6] Aronson, D. G., Weinberger, H. F.: [Nonlinear diffusion in population genetics, combustion, and nerve pulse propagation](#). In: Partial differential equations and related topics (Program, Tulane Univ., New Orleans, La., 1974), Lecture Notes in Math. 446, Springer, Berlin, 5–49 (1975) Zbl [0325.35050](#) MR [427837](#)
- [7] Aronson, D. G., Weinberger, H. F.: [Multidimensional nonlinear diffusion arising in population genetics](#). Adv. Math. **30**, 33–76 (1978) Zbl [0407.92014](#) MR [511740](#)

- [8] Chen, X.-Y.: [A strong unique continuation theorem for parabolic equations](#). Math. Ann. **311**, 603–630 (1998) Zbl [0990.35028](#) MR [1637972](#)
- [9] de Pablo, A., Vázquez, J. L.: [Travelling waves and finite propagation in a reaction-diffusion equation](#). J. Differential Equations **93**, 19–61 (1991) Zbl [0784.35045](#) MR [1122305](#)
- [10] DiBenedetto, E., Friedman, A.: [Addendum to “Hölder estimates for nonlinear degenerate parabolic systems”](#). J. Reine Angew. Math. **363**, 217–220 (1985) Zbl [0595.35065](#) MR [814022](#)
- [11] DiBenedetto, E., Friedman, A.: [Hölder estimates for nonlinear degenerate parabolic systems](#). J. Reine Angew. Math. **357**, 1–22 (1985) Zbl [0549.35061](#) MR [783531](#)
- [12] Du, Y., Lou, B.: [Spreading and vanishing in nonlinear diffusion problems with free boundaries](#). J. Eur. Math. Soc. (JEMS) **17**, 2673–2724 (2015) Zbl [1331.35399](#) MR [3420519](#)
- [13] Du, Y., Lou, B., Zhou, M.: [Nonlinear diffusion problems with free boundaries: Convergence, transition speed, and zero number arguments](#). SIAM J. Math. Anal. **47**, 3555–3584 (2015) Zbl [1321.35086](#) MR [3400577](#)
- [14] Du, Y., Matano, H.: [Convergence and sharp thresholds for propagation in nonlinear diffusion problems](#). J. Eur. Math. Soc. (JEMS) **12**, 279–312 (2010) Zbl [1207.35061](#) MR [2608941](#)
- [15] Du, Y., Quirós, F., Zhou, M.: [Logarithmic corrections in Fisher–KPP type porous medium equations](#). J. Math. Pures Appl. (9) **136**, 415–455 (2020) Zbl [1439.35238](#) MR [4076967](#)
- [16] Fernandez, F. J.: [Unique continuation for parabolic operators. II](#). Comm. Partial Differential Equations **28**, 1597–1604 (2003) Zbl [1029.35050](#) MR [2001174](#)
- [17] Fife, P. C., McLeod, J. B.: [The approach of solutions of nonlinear diffusion equations to travelling front solutions](#). Arch. Ration. Mech. Anal. **65**, 335–361 (1977) Zbl [0361.35035](#) MR [442480](#)
- [18] Fisher, R. A.: [The wave of advance of advantageous genes](#). Ann. Eugenics **7**, 335–369 (1937) Zbl [63.1111.04](#)
- [19] Gárriz, A.: [Propagation of solutions of the porous medium equation with reaction and their travelling wave behaviour](#). Nonlinear Anal. **195**, article no. 111736 (2020) Zbl [1442.35196](#) MR [4050784](#)
- [20] Gilding, B. H., Kersner, R.: [Travelling waves in nonlinear diffusion-convection reaction](#). Progr. Nonlinear Differential Equations Appl. 60, Birkhäuser, Basel (2004) Zbl [1073.35002](#) MR [2081104](#)
- [21] Gurney, W. S. C., Nisbet, R. M.: [A note on non-linear population transport](#). J. Theor. Biol. **56**, 249–251 (1976)
- [22] Gurtin, M. E., MacCamy, R. C.: [On the diffusion of biological populations](#). Math. Biosci. **33**, 35–49 (1977) Zbl [0362.92007](#) MR [682594](#)
- [23] Kolmogoroff, A., Petrowsky, I., Piscounoff, N.: [Étude de l’équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique](#). Bull. Univ. État Moscou, Sér. Int., Sect. A: Math. et Mécan. **1**, no. 6, 1–25 (1937) Zbl [0018.32106](#)
- [24] Li, Q., Lou, B.: [Vanishing phenomena in fast decreasing generalized bistable equations](#). J. Math. Anal. Appl. **500**, article no. 125096 (2021) Zbl [1460.35038](#) MR [4219962](#)
- [25] Newman, W. I., Sagan, C.: [Galactic civilizations: Population dynamics and interstellar diffusion](#). Icarus **46**, 293–327 (1981)
- [26] Oleĭnik, O. A., Kalashnikov, A. S., Zhou, Y.-L.: [The Cauchy problem and boundary problems for equations of the type of non-stationary filtration](#). Izv. Akad. Nauk SSSR Ser. Mat. **22**, 667–704 (1958) Zbl [0093.10302](#) MR [99834](#)
- [27] Sacks, P. E.: [The initial and boundary value problem for a class of degenerate parabolic equations](#). Comm. Partial Differential Equations **8**, 693–733 (1983) Zbl [0529.35038](#) MR [700733](#)
- [28] Sánchez-Garduño, F., Maini, P. K.: [Existence and uniqueness of a sharp travelling wave in degenerate non-linear diffusion Fisher–KPP equations](#). J. Math. Biol. **33**, 163–192 (1994) Zbl [0822.92021](#) MR [1314727](#)

- [29] Sherratt, J. A.: [On the form of smooth-front travelling waves in a reaction-diffusion equation with degenerate nonlinear diffusion](#). Math. Model. Nat. Phenom. **5**, 64–79 (2010) Zbl [1202.35050](#) MR [2681228](#)
- [30] Vázquez, J. L.: The porous medium equation. Mathematical theory. Oxford Math. Monogr., The Clarendon Press, Oxford (2007) Zbl [1107.35003](#) MR [2286292](#)
- [31] Wu, Z., Zhao, J., Yin, J., Li, H.: [Nonlinear diffusion equations](#). World Scientific, River Edge, NJ (2001) Zbl [0997.35001](#) MR [1881297](#)
- [32] Zlatoš, A.: [Sharp transition between extinction and propagation of reaction](#). J. Amer. Math. Soc. **19**, 251–263 (2006) Zbl [1081.35011](#) MR [2169048](#)