

Cluster algebras of finite mutation type with coefficients

Anna Felikson and Pavel Tumarkin

Abstract. We classify mutation-finite cluster algebras with arbitrary coefficients of geometric type. This completes the classification of all mutation-finite cluster algebras started in [Felikson, Shapiro, and Tumarkin, J. Eur. Math. Soc. 14 (2012)].

Contents

1. Introduction and main results	375
2. Background	379
3. Quivers from surfaces and peripheral laminations	383
4. Quivers of affine type	385
5. Extended affine quivers	389
6. Mutation classes of X_6 and X_7	392
7. Rank 2 quivers	396
8. Quivers from surfaces	396
9. Skew-symmetrizable mutation classes	407
10. Annulus property as criterion of mutation finiteness	414
References	417

1. Introduction and main results

Cluster algebras with coefficients were introduced in [13], the fourth paper in the series founding the theory of cluster algebras. Cluster algebras of geometric type are defined as those having their coefficients in tropical semifields. In particular, this includes the important case of cluster algebras with principal coefficients.

A cluster algebra of geometric type is completely defined by an integer $(m+n) \times n$ exchange matrix with skew-symmetrizable top $n \times n$ part (called *principal* or *mutable* part). Exchange matrices undergo involutive transformations called *mutations*,

Mathematics Subject Classification 2020: 13F60.

Keywords: cluster algebras, quiver, mutation.

all exchange matrices which can be obtained by iterative mutations form a *mutation class*. We say that a cluster algebra is *mutation-finite* if its mutation class is finite.

Coefficient-free mutation-finite cluster algebras were classified in [7, 8]. These algebras found various applications, including ones in quantum field theories (see e.g. [1, 2]).

In this paper, we classify all mutation-finite exchange matrices with arbitrary coefficients. We first restrict ourselves to matrices with skew-symmetric mutable part (this assumption will be dropped later). In this case the matrix can be represented by a quiver with vertices of two types: *mutable* (corresponding to the mutable part of the matrix) and *frozen* (such quivers are also called *ice quivers*). The quiver also undergoes mutations compatible with mutations of the matrix, we say that a quiver is mutation-finite if the corresponding exchange matrix is.

The first easy observation is that the mutable part of a mutation-finite quiver should be mutation-finite. Mutation-finite quivers without frozen vertices were classified in [8], the list consists of the following (overlapping) classes of quivers: rank 2 quivers, quivers originating from surfaces (see Section 2), quivers of finite type (i.e., with an orientation of a finite type Dynkin diagram in the mutation class), quivers of affine type (ones with an orientation of an affine type Dynkin diagram in the mutation class), quivers of extended affine types $E_6^{(1,1)}$, $E_7^{(1,1)}$ and $E_8^{(1,1)}$ (see Figure 2.2), exceptional quivers of types X_6 and X_7 (see also Figure 2.2).

Another easy observation is that it is enough to consider just one frozen vertex. Indeed, as there are no arrows between frozen vertices, the frozen vertices do not affect each other in the process of mutations.

Definition 1.1. Let Q be a quiver of finite mutation type (with vertices v_1, \dots, v_n all being mutable). Let q be an additional (frozen) vertex, denote by b_i the number of arrows connecting a vertex v_i of Q to q . We will say that the integer coefficient vector $\mathbf{b} = (b_1, \dots, b_n)$ is *admissible* if $\mathbf{b} \neq 0$ and the quiver spanned by Q and q with the unique frozen vertex q is of finite mutation type.

Therefore, the question of classification of mutation-finite exchange matrices is equivalent to finding all admissible vectors for every mutation-finite quiver without frozen vertices.

A distinguished class of cluster algebras consists of algebras of *finite type*: these were classified by Fomin and Zelevinsky in [12] by establishing a connection with Cartan–Killing classification of simple Lie algebras. They also proved in [13] that adding any coefficients to a cluster algebra of finite type results in a mutation-finite cluster algebra. Moreover, this characterizes cluster algebras of finite type: if every exchange matrix with given principal part is mutation-finite, then the principal part defines an algebra of finite type. A stronger conjecture was made in [13] stating that it

is sufficient to check the mutation-finiteness of the algebra with principal coefficients only, this was proved by Seven [20].

In particular, this provides the answer for the finite type.

Proposition 1.2 ([13]). *If Q is of finite type then any vector \mathbf{b} is admissible.*

The next large class of quivers consists of quivers from surfaces [10]. We first prove the following statement.

Proposition 1.3 (Theorem 3.2). *If Q is arising from a surface then \mathbf{b} is admissible if and only if it corresponds to a peripheral lamination.*

Due to results of Gu [15], Proposition 1.3 provides an algorithm which determines whether a given quiver from a surface with a frozen vertex is mutation-finite: using [15], one can reconstruct a triangulation, then one can reconstruct a lamination using a procedure from [11], and then it is straightforward to check whether a given lamination is peripheral. We will give a more explicit characterization of coefficient vectors corresponding to peripheral laminations in Section 8.

Next, we consider affine and exceptional mutation-finite classes. Every mutation class of quivers of affine type contains a representative with a double arrow, so the main tool in the considerations is the following necessary condition (which we call the *annulus property*).

Proposition 1.4 (Corollary 3.5). *Let Q be a quiver containing a double arrow from v_1 to v_2 . Then a vector \mathbf{b} is admissible only if $b_1 = -b_2 \leq 0$.*

In the affine case \tilde{A} we use Proposition 1.3 to show that the annulus property is also sufficient (see Lemma 4.1 and Remark 4.2).

The same result applies to other affine quivers, but here their treatment is based on their cluster modular groups studied in [19].

Proposition 1.5 (Theorem 4.3). *For the representatives of the mutation classes of affine types \tilde{D} and \tilde{E} shown in Figure 4.3, a vector is admissible if and only if it satisfies the annulus property.*

This result is then generalized to all quivers of affine type containing a double arrow.

Proposition 1.6 (Theorem 4.4). *If Q is a quiver of affine type containing a double arrow, then a vector \mathbf{b} is admissible if and only if \mathbf{b} satisfies the annulus property.*

For the extended affine quivers and quiver of type X_7 we take a specific representative Q from the mutation class (see Figures 5.1–6.1) and an element of the cluster modular group φ to show that the annulus property for Q is not compatible with the annulus property for $\varphi(Q)$, which results in the following statement.

Proposition 1.7 (Theorems 5.1–6.1). *Let Q be of type $E_6^{(1,1)}$, $E_7^{(1,1)}$, $E_8^{(1,1)}$ or X_7 . Then there is no admissible vector \mathbf{b} .*

The case of quiver Q of type X_6 is different: it admits an admissible vector such that the quiver spanned by Q and the frozen vertex q is isomorphic to X_7 . In fact, the quiver admits a series of admissible vectors as follows.

Proposition 1.8 (Theorem 6.2). *Let Q be a quiver of type X_6 with a single arrow from v_0 to v_5 , where v_5 is a leaf (see Figure 6.1). Then a vector \mathbf{b} is admissible if and only if $b_5 = -2b_0 \geq 0$ and all other b_i vanish. Admissible vectors for all representatives of the mutation class are shown in Figure 6.2.*

In the next proposition we consider the quivers of rank 2 (note that the first two parts have been already considered previously).

Proposition 1.9 (Theorem 7.1). *Let Q be a rank two quiver with the arrow from v_1 to v_2 of weight $a > 0$. Let $\mathbf{b} = (b_1, b_2)$ be an integer vector. Then*

- (1) *if $a = 1$ then \mathbf{b} is admissible for any b_1, b_2 ;*
- (2) *if $a = 2$ then \mathbf{b} is admissible if and only if $b_1 = -b_2 \leq 0$;*
- (3) *if $a > 2$ then there are no admissible vectors.*

Finally, we specify a particular triangulation for every surface and give the admissibility criterion for the corresponding quiver, see Theorem 8.2. The criterion is also based on the annulus property.

The results are extended to the general skew-symmetrizable case in Section 9 by using diagrams in place of quivers and orbifolds in place of surfaces. The modified annulus property for arrows of weight $(1, 4)$ is defined in Theorem 9.6.

We now combine the results in one theorem.

Theorem 1.10. *Let Q be a quiver/diagram and $\mathbf{b} = (b_1, \dots, b_n)$ be an integer vector. Then*

- (1) *if Q is of finite type then any vector \mathbf{b} is admissible;*
- (2) *if Q is of affine type and Q contains a double arrow or an arrow of weight $(1, 4)$, then a vector \mathbf{b} is admissible if and only if \mathbf{b} satisfies the annulus property;*
- (3) *if Q is arising from a surface/orbifold then \mathbf{b} is admissible if and only if it corresponds to some peripheral lamination; the criterion for admissibility is given for a specific representative of the mutation class in Theorems 8.2 and 9.8;*
- (4) *if Q is of type X_6 , all possible admissible vectors are shown in Figure 6.2;*
- (5) *otherwise, there is no admissible vector.*

A criterion for being mutation-finite can be also reformulated in terms of the annulus property applied to the whole mutation class, this was proposed by Sergey Fomin. The statement is similar to the analogous criterion for quivers/diagrams without frozen vertices (see e.g. [3, Corollary 8]).

Theorem 1.11 (Theorem 10.1). *Let Q be a quiver/diagram with a frozen vertex v . Suppose that the subquiver $Q \setminus v$ is mutation-finite. Then Q is mutation-finite if and only if the annulus property holds in every quiver/diagram Q' mutation-equivalent to Q for every double arrow contained in $Q' \setminus v$.*

From this one can conclude the following.

Corollary 1.12 (Corollary 10.2). *Let Q be a quiver/diagram with a frozen vertex. Then Q is mutation-finite if and only if for every quiver Q' in the mutation class of Q every rank 3 subquiver/subdiagram of Q' is mutation-finite.*

The paper is organized as follows. In Section 2 we recall necessary background concerning triangulated surfaces and laminations on them. Section 3 is devoted to quivers from surfaces and the connection between admissible vectors and peripheral laminations. In Section 4 we consider quivers of affine types, in Sections 5 and 6 we treat extended affine quivers and quivers of types X_6 and X_7 . In the short Section 7 we consider quivers of rank 2. Section 8 characterizes admissible vectors for a particular triangulation of a surface. In Section 9 all results are extended to the general context of skew-symmetrizable mutation classes. Finally, in Section 10 we discuss the criterion of mutation-finiteness in terms of the annulus property.

2. Background

2.1. Matrix mutation

We start by recalling the definition of matrix mutation (we adopt the notation used in [13]).

Given an integer skew-symmetric $n \times n$ matrix $B = (b_{ij})$, the mutation μ_k of B for $k \in \{1, \dots, n\}$ is defined by $\mu_k(B) = B' = (b'_{ij})$, where

$$b'_{ij} = \begin{cases} -b_{ij} & \text{if } i = k \text{ or } j = k, \\ b_{ij} + \operatorname{sgn}(b_{ik})[b_{ik}b_{kj}]_+ & \text{otherwise,} \end{cases}$$

where $\operatorname{sgn}(x)$ denotes the sign function, and $[x]_+ = \max\{x, 0\}$.

For an extended $m \times n$ matrix \bar{B} with $m > n$ and skew-symmetric principal part given by first n rows, the mutation is provided by the same formula.




Figure 2.1. Quiver mutation. Here p, q are positive, and the sign of r and $pq - r$ can be negative (which corresponds to opposite direction of the respective arrows).

Remark 2.1. A skew-symmetric $n \times n$ matrix $B = (b_{ij})$ can be represented by a quiver with n vertices v_1, \dots, v_n and b_{ij} arrows from v_i to v_j . Matrix mutation then can be reformulated in the quiver language, see Figure 2.1.

2.2. Construction of quivers from triangulations

We briefly recall the construction of quivers from triangulated surfaces [10].

Let S be a connected orientable surface with boundary and with a finite set M of marked points (such that every boundary component contains at least one marked point). Let T be a triangulation of S by the arcs having their endpoints in M . Suppose that T has no self-folded triangles (i.e. every triangle in T is bounded by three distinct arcs or boundary segments). We construct a quiver Q whose vertices v_1, \dots, v_n correspond to the arcs e_1, \dots, e_n of T . The number of arrows in Q from v_i to v_j is defined as

$$b_{ij} = \#\{\text{triangles with sides } e_i \text{ and } e_j, \text{ with } e_j \text{ following } e_i \text{ in clockwise order}\} - \#\{\text{triangles with sides } e_i \text{ and } e_j, \text{ with } e_j \text{ following } e_i \text{ in counterclockwise order}\}.$$

For more subtle rules for treating self-folded triangles see [10].

It is shown in [10] that mutations of the quiver Q correspond to flips of the triangulation T . It is easy to see from the definition above that combinatorially equivalent triangulations of S give rise to isomorphic quivers (we say that triangulations are combinatorially equivalent if one can be taken to the other by an orientation-preserving homeomorphism of the surface). As it is shown in [15], a triangulation of a surface can be uniquely reconstructed from the corresponding quiver (up to finitely many low rank examples).

2.3. Classification of mutation-finite quivers

We will heavily use the following theorem.




Figure 2.2. Eleven exceptional finite mutation classes.

Theorem 2.2 ([8]). *A connected mutation-finite quiver is either of rank 2, or a quiver arising from a triangulation of a surface, or a quiver mutation-equivalent to one of the eleven quivers $E_6, E_7, E_8, \tilde{E}_6, \tilde{E}_7, \tilde{E}_8, E_6^{(1,1)}, E_7^{(1,1)}, E_8^{(1,1)}, X_6, X_7$ shown in Figure 2.2.*

2.4. Laminations as coefficients for surface case

It is shown in [11] that in the case of a quiver from triangulated surface S , the coefficient vectors can be represented by *laminations* on the same surface S , and that the coefficient vectors can be computed from the triangulation and lamination using *shear coordinates*.

Definition 2.3 ([11, Definition 12.1]). *An integral unbounded measured lamination, or just a lamination for short, on a marked surface (S, M) is a finite collection of non-self-intersecting and pairwise non-intersecting curves in S , modulo isotopy relative to M , subject to the restrictions specified below. Each curve must be one of the following:*

- a closed curve (an embedded circle);
- a curve connecting two unmarked points on the boundary of S ;
- a curve starting at an unmarked point on the boundary and, at its other end, spiralling into a puncture (either clockwise or counterclockwise);
- a curve both of whose ends spiral into punctures (not necessarily distinct);

where the following types of curves are not allowed:

- a curve that bounds an unpunctured or once-punctured disk;
- a curve with two endpoints on the boundary of S which is isotopic to a piece of boundary containing no marked points, or a single marked point;
- a curve with two ends spiralling into the same puncture in the same direction without enclosing anything else.




Figure 2.3. Shear coordinates.

When speaking about two curves γ_1 and γ_2 (for example an arc of triangulation and a curve from a lamination) we always assume that the number of crossings is minimal possible for the curves in the homotopy classes of γ_1 and γ_2 , respectively.

Definition 2.4 ([11, Definition 12.2]). Let L be a lamination and let T be a triangulation without self-folded triangles of the same surface. For each arc γ in T , the corresponding *shear coordinate* of L with respect to the triangulation T , denoted by $b_\gamma(T, L)$, is defined as a sum of contributions from all intersections of curves in L with the arc γ . Specifically, such an intersection contributes $+1$ (resp., -1) to $b_\gamma(T, L)$ if the corresponding segment of a curve in L cuts through the quadrilateral surrounding γ cutting through edges as shown in Figure 2.3 on the left (resp., on the right). Note that at most one of these two types of intersection can occur. Note also that even though a spiralling curve can intersect an arc infinitely many times, the number of intersections that contribute to the computation of $b_\gamma(T, L)$ is always finite.

Shear coordinates can also be defined for arcs involved in self-folded triangles, see [11, Section 13].

Note that the vector $\mathbf{b} = (b_1, \dots, b_n)$ from Definition 1.1 consists of negative shear coordinates of the corresponding lamination.

It is known (see [11, Theorem 13.6], see also [9, Section 3]) that for a given triangulation T , the map

$$L \mapsto (b_\gamma(T, L))_{\gamma \in T}$$

provides a bijection between laminations and \mathbb{Z}^n .

In particular, for every triangulation T and every arc $\gamma_0 \in T$ there exists an *elementary lamination* L such that $b_{\gamma_0}(T, L) = 1$ and $b_{\gamma_i}(T, L) = 0$ for all $\gamma_i \in T$, $i \neq 0$. This elementary lamination consists of one curve which follows γ_0 but has its endpoints changed: for endpoints of γ_0 at a boundary marked point, the end of the elementary lamination is shifted to the left along the boundary, for endpoints of γ_0 at a puncture, the end of the elementary lamination is spiralling into the puncture counterclockwise if the end is untagged and clockwise otherwise. We will also use

negative elementary lamination defined by $b_{\gamma_0}(T, L) = -1$ and $b_{\gamma_i}(T, L) = 0$ for all $\gamma_i \in T, i \neq 0$. The negative elementary lamination also consists of one curve tracing the arc γ_0 , but having boundary endpoints shifted to the right and the puncture end points spiralling to the puncture in the clockwise direction.

3. Quivers from surfaces and peripheral laminations

Let Q be a quiver constructed by a triangulation T of a surface S . As it was mentioned in Section 2.4, choosing a coefficient vector \mathbf{b} is equivalent to a choice of a lamination L on S . Since mutations correspond to flips of triangulations (and we can reach every triangulation by a sequence of flips), the vector \mathbf{b} is admissible if and only if the shear coordinates of the lamination L on all triangulations of S take finitely many values only.

Definition 3.1. A curve on a marked surface S will be called *peripheral* if it belongs to some lamination on S and can be isotopically deformed to (a part of) a boundary component of S . By a *peripheral lamination* we understand a lamination consisting of peripheral curves.

In this section, we show that admissible vectors are in bijection with peripheral laminations (see Theorem 3.2). In Section 8 we will reformulate the result in terms of quivers.

Theorem 3.2. *Let Q be a quiver from a triangulated surface S . Then admissible vectors for Q are in bijection with peripheral laminations on S .*

Proof. We need to show that the vector of shear coordinates of a lamination L takes finitely many values if and only if L is peripheral.

First, consider a peripheral lamination L . It is preserved by any Dehn twist along any closed curve on the surface, and hence, it is preserved by the whole mapping class group of the surface (as the latter is generated by twists).

Observe that for a given surface S there is only a finite number of combinatorial types of triangulations (in particular, this is precisely the reason why quivers originating from surfaces are mutation-finite), and combinatorially equivalent triangulations can be taken to each other by elements of the mapping class group of S . This implies that given an initial triangulation T and the corresponding quiver Q , there is a finite number of mutation sequences applying which together with elements of the mapping class group we can reach any triangulation of S . Since shear coordinates of L are invariant under the action of the mapping class group, this implies that the vector of shear coordinates of L takes one of finitely many values.




Figure 3.1. Quiver \tilde{A}_1 , annulus and admissible coefficients.

Now, consider a lamination L which is not peripheral. Then there exists a closed curve C crossing L . Let T be a triangulation and $D_C^k(T)$, $k \in \mathbb{Z}$ be the images of T under iterative applications of Dehn twist D_C along C . We claim that shear coordinates of L with respect to $D_C^k(T)$ take infinitely many different values. Indeed, to apply D_C to T with keeping L intact is the same as applying D_C^{-1} to L and preserving T . As C intersects L , the Dehn twists $D_C^{-k}(L)$ will produce infinitely many different laminations. Due to the bijection between laminations and their shear coordinates, this implies that the shear coordinates of laminations $D_C^{-k}(L)$ with respect to triangulation T are different. Hence, the shear coordinates of L with respect to triangulations $D_C^k(T)$ are different, and thus take infinitely many values. This implies that non-peripheral laminations do not correspond to admissible vectors. ■

Notice that if a surface has no boundary, then it contains no peripheral curves. This gives rise to the following corollary of Theorem 3.2.

Corollary 3.3. *If a surface has no boundary, then the quiver of any of its triangulations has no admissible vectors.*

Example 3.4. Let Q be the affine quiver \tilde{A}_1 (two vertices v_1 and v_2 connected by a double arrow from v_1 to v_2). It corresponds to an annulus with one marked point on each boundary component, see Figure 3.1. The only peripheral curve on the annulus coincides with the unique closed curve inside this annulus (here we use the fact that every boundary component contains only one marked point). So, every peripheral lamination consists of an integer number of copies of this closed curve. As one can see from Figure 3.1, the corresponding coefficient vector satisfies

$$b_1 = -b_2 \leq 0$$

(recall that b_i denotes negative shear coordinate, i.e. the number of arrows from a vertex v_i to the frozen vertex).




Figure 4.1. Triangulated annulus $S_{p,q}$ with the corresponding quiver of type $\tilde{A}_{p,q}$.

The result of Example 3.4 can be reformulated as follows.

Corollary 3.5 (Annulus property). *Let Q be the rank 2 quiver with a double arrow from v_1 to v_2 and $\mathbf{b} = (b_1, b_2)$ be an admissible vector. Then $b_1 = -b_2 \leq 0$. This will be called the annulus property for $v_1 \Rightarrow v_2$.*

Corollary 3.5 leads to the following necessary condition for a coefficient vector \mathbf{b} to be admissible, which we will heavily use throughout the paper.

Definition 3.6 (Annulus property). For an arbitrary quiver Q , a coefficient vector $\mathbf{b} = (b_1, \dots, b_n)$ satisfies the *annulus property* if for every double arrow $v_i \Rightarrow v_j$ in Q we have $b_i = -b_j \leq 0$.

4. Quivers of affine type

Let L be a lamination on an annulus. A curve $C \in L$ is called *bridging* if it has endpoints on both boundary components of the annulus (in other words, if and only if it is not peripheral).

Lemma 4.1. *Let $S_{p,q}$ be the annulus with p and q boundary marked points triangulated as in Figure 4.1. Then a vector (b_1, \dots, b_n) is admissible if and only if it satisfies the annulus property.*

Proof. The annulus property $b_1 = -b_2 \leq 0$ is necessary by Corollary 3.5. We need to prove that it is also sufficient for admissibility of (b_1, \dots, b_n) . In view of Theorem 3.2, this is equivalent to proving that for every vector (b_1, \dots, b_n) satisfying

$$b_1 = -b_2 \leq 0,$$




Figure 4.2. Triangulated annulus $S_{1,1}$, its universal cover, and bridging curves with corresponding values of (b_1, b_2) (recall that we define b_i as negative shear coordinates).

there exists a peripheral lamination resulting in this vector. Since every vector is realisable by some lamination, we see that it is sufficient to show that a lamination satisfying the condition $b_1 = -b_2 \leq 0$ cannot contain bridging curves.

Suppose that L is a lamination on $S_{p,q}$ with $b_1 = -b_2 \leq 0$ and containing a bridging curve l . Consider the restriction \bar{L} of the lamination L to the shaded annulus $S_{1,1}$ with one marked point at each boundary component (see Figure 4.1). The restriction \bar{l} of the curve l to $S_{1,1}$ is a bridging curve for $S_{1,1}$. In Figure 4.2 we show a triangulated annulus (left) and its universal cover (right). For every bridging curve, we draw its lift (we normalize it by drawing the “lower” end in the same square of the universal cover) and compute its (negative) shear coordinates. Notice that every peripheral curve satisfies either $b_1 = b_2 = 0$ (if it is not the closed curve) or $b_1 = -b_2 = -1$ otherwise. The latter is not contained in L in presence of a bridging curve. Hence, peripheral curves in L do not affect b_1 and b_2 , and it is sufficient to check coordinates of all collections of mutually non-intersecting bridging curves.

Any bridging curve on $S_{1,1}$ can be obtained from any other bridging curve by application of a power of the Dehn twist along the unique closed curve, and if two curves differ by more than one twist then they intersect each other. One can easily see that no bridging curve on $S_{1,1}$ satisfies $|b_1| = |b_2|$, and coordinates (b_1, b_2) of a pair of bridging curves differing by one twist can take values

$$(-2k - 1, 2k + 3), \quad (1, 1), \quad (1, -1), \quad (-1, -1), \quad (-2k - 2, 2k + 1)$$

for $k \geq 0$ (see Figure 4.2). None of these satisfies $b_1 = -b_2 \leq 0$, so we obtain a contradiction. ■

Remark 4.2. Notice that the proof of Lemma 4.1 does not use any properties of the triangulation of $S_{p,q}$ outside of the shaded annulus. In other words, the same proof works for any triangulation of $S_{p,q}$ with the associated quiver containing a double arrow.

We will now use Lemma 4.1 to classify all admissible vectors for the remaining quivers of affine type.




Figure 4.3. Representatives of affine mutation classes $\tilde{A}_{p,q}$, \tilde{D}_n , \tilde{E}_n .

Take the representatives of the mutation classes \tilde{D}_n , \tilde{E}_6 , \tilde{E}_7 , \tilde{E}_8 shown in Figure 4.3. A necessary condition on an admissible vector follows from the annulus property. The following theorem shows that every coefficient not breaking the annulus property is admissible.

Theorem 4.3. *Let Q be the quiver of type \tilde{D}_n , \tilde{E}_6 , \tilde{E}_7 or \tilde{E}_8 shown in Figure 4.3. A coefficient vector \mathbf{b} is admissible if and only if it satisfies $b_0 = -b_1 \leq 0$.*

Before proving the theorem, we recall the notion of the *cluster modular group* as the group generated by sequences of mutations (followed by permutations of the vertices of a quiver if needed) preserving the initial quiver (see e.g. [5] for a detailed definition, where the term “mapping class group of a cluster algebra” is used instead, and [14, 19] for detailed descriptions of the cluster modular groups for affine and extended affine algebras).

Proof. The necessity of the assumption of the theorem follows from the annulus property. We now prove the sufficiency.

It is shown in [19] that the cluster modular group for Q is an abelian group generated by three mutation sequences (followed by certain permutations) described below. Define the sets of indices

$$I_{\text{odd}} = \begin{cases} \{i \in [5, k], i \text{ odd}\} & \text{for type } \tilde{E}_k, \\ \{i \in [3, n - 1], i \text{ odd}\} & \text{for type } \tilde{D}_n, \end{cases}$$

$$I_{\text{even}} = \begin{cases} \{i \in [5, k], i \text{ even}\} & \text{for type } \tilde{E}_k, \\ \{i \in [3, n - 1], i \text{ even}\} & \text{for type } \tilde{D}_n, \end{cases}$$

and define the composite mutations μ_{odd} and μ_{even} as compositions of commuting mutations in I_{odd} and I_{even} , respectively.

In these terms the generators of the cluster modular group can be written as follows:

$$\begin{aligned}
 \mu^{(1)} &= \mu_2 \circ \mu_1 \circ \mu_0 && \text{with cyclic permutation } (v_2 v_1 v_0), \\
 \mu^{(2)} &= \begin{cases} \mu_4 \circ \mu_3 \circ \mu_1 \circ \mu_0 & \text{with permutation } (v_3 v_1 v_0) \text{ for type } \tilde{E}_k, \\ \mu_n \circ \mu_1 \circ \mu_0 & \text{with permutation } (v_n v_1 v_0) \text{ for type } \tilde{D}_n, \end{cases} \\
 \mu^{(3)} &= \mu_{\text{even}} \circ \mu_{\text{odd}} \circ \mu_1 \circ \mu_0 && \text{with permutation } (v_5 v_1 v_0) \text{ or } (v_3 v_1 v_0) \\
 &&& \text{for } \tilde{E}_k \text{ and } \tilde{D}_n, \text{ respectively.}
 \end{aligned}$$

Inside Q consider the following subquivers which we will call *wings* (we list the vertices of the subquivers in the brackets):

$$Q_1 = \langle v_2 \rangle, \quad Q_2 = \langle v_3, v_4 \rangle \text{ or } \langle v_n \rangle \text{ for } \tilde{E}_k \text{ and } \tilde{D}_n \text{ resp.,} \quad Q_3 = \langle v_{I_{\text{odd}}}, v_{I_{\text{even}}} \rangle.$$

We claim that each of $\mu^{(k)}, k = 1, 2, 3$, only changes the value of b_i if $i \in Q_k$ and does not affect others. To see this, consider the subquivers $Q \setminus Q_i = \langle v_0, v_1, Q_j, Q_k \rangle, i, j, k$ distinct. Each of these corresponds to a triangulated annulus, with an annulus $S_{1,1}$ inside (which corresponds to the subquiver $\langle v_0, v_1 \rangle$) and polygons attached to each of its boundaries (which correspond to the wings). By Lemma 4.1, the assumption $b_0 = -b_1 \leq 0$ implies that the restriction of vector \mathbf{b} on $Q \setminus Q_i$ is defined by some peripheral lamination on the corresponding annulus. The mutation $\mu^{(k)}$ acts as a cyclic permutation of the boundary marked vertices of the triangulation corresponding to the k -th wing. Therefore, this element acts trivially on the wing Q_j and on the values of b_0 and b_1 , while the order of the action on Q_k is equal to the number of vertices in Q_k plus one. Since we could choose the quiver $Q \setminus Q_j$ instead, $\mu^{(k)}$ acts trivially on Q_i as well. Thus, the action of the whole cluster modular group on the vector \mathbf{b} has a finite orbit.

The rest of the proof is similar to the proof of Theorem 3.2. As Q is mutation-finite, the number of distinct mutation sequences modulo the action of the cluster modular group is finite. Together with the finiteness of the orbit of \mathbf{b} under the action of the cluster modular group this results in the admissibility of \mathbf{b} . ■

We now generalize the result of Theorem 4.3 to all quivers of affine type containing a double arrow.

Theorem 4.4. *If Q is of affine type and Q contains a double arrow, then a vector \mathbf{b} is admissible if and only if \mathbf{b} satisfies the annulus property.*

Proof. For the quivers of types \tilde{A} the statement follows from Remark 4.2. All quivers of type \tilde{D} are classified in [17], and it follows from the classification that any quiver with a double arrow can be obtained from the quiver Q of type \tilde{D}_n shown in Figure 4.3 by mutations in vertices v_i for $4 \leq i \leq n$. Therefore, we can mutate our quiver

to Q preserving the annulus property, so Theorem 4.3 implies that the vector \mathbf{b} is admissible.

The proof for types $\tilde{E}_6, \tilde{E}_7, \tilde{E}_8$ is similar: the inspection of the mutation classes shows that all quivers with double arrows are obtained from the quivers in Figure 4.3 by a sequence of mutations at vertices v_6, v_7, v_8 . As none of these vertices is connected to v_0 and v_1 , such a sequence of mutations cannot break the annulus property. ■

5. Extended affine quivers

In this section, we prove that there are no admissible vectors for extended affine types $E_{6,7,8}^{(1,1)}$. For every mutation class we choose a specific representative containing a double arrow and find an element μ from the cluster modular group such that the application of μ breaks the annulus property.

5.1. Mutation class of $E_6^{(1,1)}$

Theorem 5.1. *There is no admissible vector for a quiver in the mutation class of $E_6^{(1,1)}$.*

Proof. It is sufficient to prove the statement for one quiver from the mutation class. We consider the quiver Q shown in Figure 5.1 (left). Suppose that $\mathbf{b} = (b_1, \dots, b_8)$ is an admissible vector.

Plan of the proof and notation. The subquiver $\langle v_7, v_8 \rangle$ is of type \tilde{A}_1 , so from the annulus property for $v_8 \Rightarrow v_7$, we have

$$b_8 = -b_7 \leq 0.$$

We will find a sequence of mutations μ taking Q to the opposite quiver Q^{op} (where Q^{op} is obtained from Q by reversing all arrows) and check that after the application of the mutation sequence μ to \mathbf{b} the annulus property does not hold.

More precisely, let

$$\mu_* = \mu_3 \circ \mu_2 \circ \mu_1, \quad \mu_\diamond = \mu_7 \circ \mu_6 \circ \mu_5 \circ \mu_4$$

(notice that the components of each of these composite mutations commute) and consider the following sequence of three composite mutations:

$$\mu = \mu_* \circ \mu_\diamond \circ \mu_*.$$

We observe that $Q_1 := \mu_*(Q)$ is the quiver shown in Figure 5.1 (right), $Q_2 := \mu_\diamond \circ \mu_*(Q) = Q_1^{\text{op}}$ is the quiver opposite to Q_1 , and $Q_3 := \mu(Q) = Q^{\text{op}}$ is the quiver opposite to Q .




Figure 5.1. $Q = E_6^{(1,1)}$ (left) and $\mu_*(Q)$ (right).

We now compute how the vector \mathbf{b} changes under the sequence of mutations. Denote its components by $b_i^{(1)}$, $b_i^{(2)}$ and $b_i^{(3)}$ after applying μ_* , $\mu_\diamond \circ \mu_*$ and $\mu = \mu_* \circ \mu_\diamond \circ \mu_*$, respectively.

If \mathbf{b} is an admissible vector, then $(b_1^{(3)}, \dots, b_8^{(3)})$ satisfies the annulus property for $v_7 \Rightarrow v_8$ in Q_3 (as $Q_3 = Q^{\text{op}}$), i.e. one must have

$$b_7^{(3)} = -b_8^{(3)} \leq 0, \tag{5.1}$$

but the computation will show this implies $\mathbf{b} = 0$.

Computation of $b_8^{(3)}$. We start by computing $b_8^{(1)}$, $b_8^{(2)}$ and $b_8^{(3)}$:

$$\begin{aligned} b_8^{(1)} &= b_8 - [-b_1]_+ - [-b_2]_+ - [-b_3]_+ \leq b_8 \leq 0, \\ b_8^{(2)} &= b_8^{(1)} - [-b_4^{(1)}]_+ - [-b_5^{(1)}]_+ - [-b_6^{(1)}]_+ - [-b_7^{(1)}]_+ \leq b_8^{(1)} \leq 0, \\ b_8^{(3)} &= b_8^{(2)} - [-b_1^{(2)}]_+ - [-b_2^{(2)}]_+ - [-b_3^{(2)}]_+ \leq b_8^{(2)} \leq 0. \end{aligned}$$

If $b_8^{(3)} \neq 0$, then we obtain $b_8^{(3)} < 0$ which contradicts (5.1). Therefore, $b_8^{(3)} = 0$. Furthermore, since $b_8 \leq 0$ and all summands above are also non-positive, the condition $b_8^{(3)} = 0$ is satisfied if and only if $b_8 = 0$ and all entries in the computation above vanish. This results in the following constraints:

$$b_7 = b_8 = 0, \quad b_1, b_2, b_3 \geq 0, \quad b_4^{(1)}, b_5^{(1)}, b_6^{(1)}, b_7^{(1)} \geq 0, \quad b_1^{(2)}, b_2^{(2)}, b_3^{(2)} \geq 0. \tag{5.2}$$

Computation of $b_7^{(3)}$. Since $b_8^{(3)} = 0$, (5.1) implies that $b_7^{(3)} = 0$. Our goal is to express $b_7^{(3)}$ via the components of \mathbf{b} to find further constrains on b_i . We do this by first expressing $b_7^{(3)}$ via $b_i^{(2)}$, then computing required $b_i^{(2)}$ in terms of $b_j^{(1)}$, etc. While computing we will use the inequalities in (5.2):

$$\begin{aligned} b_7^{(1)} &= b_7 + [b_1]_+ + [b_2]_+ + [b_3]_+ = b_1 + b_2 + b_3, \\ b_7^{(2)} &= -b_7^{(1)} = -b_1 - b_2 - b_3, \\ b_7^{(3)} &= b_7^{(2)} + [b_1^{(2)}]_+ + [b_2^{(2)}]_+ + [b_3^{(2)}]_+ = b_7^{(2)} + b_1^{(2)} + b_2^{(2)} + b_3^{(2)}; \end{aligned}$$

$$\begin{aligned}
 b_i^{(1)} &= -b_i \leq 0 \quad \text{for } i = 1, 2, 3, \\
 b_4^{(1)} &= b_4 + b_1 \geq 0, \quad b_5^{(1)} = b_5 + b_2 \geq 0, \quad b_6^{(1)} = b_6 + b_3 \geq 0, \\
 b_7^{(1)} &= b_1 + b_2 + b_3 \geq 0; \\
 b_1^{(2)} &= b_1^{(1)} + b_4^{(1)} + b_7^{(1)} \\
 &= (-b_1) + (b_4 + b_1) + (b_1 + b_2 + b_3) = b_4 + b_1 + b_2 + b_3 \geq 0, \\
 b_2^{(2)} &= b_2^{(1)} + b_5^{(1)} + b_7^{(1)} = b_5 + b_1 + b_2 + b_3 \geq 0, \\
 b_3^{(2)} &= b_3^{(1)} + b_6^{(1)} + b_7^{(1)} = b_3 + b_1 + b_2 + b_3 \geq 0.
 \end{aligned}$$

Finally, we obtain

$$\begin{aligned}
 b_7^{(3)} &= b_7^{(2)} + b_1^{(2)} + b_2^{(2)} + b_3^{(2)} \\
 &= (b_4 + b_1) + (b_5 + b_2) + (b_6 + b_3) + b_1 + b_2 + b_3 = 0.
 \end{aligned}$$

Notice that every summand in the sum is non-negative. Therefore, every summand is zero, in particular, $b_1 = b_2 = b_3 = 0$, from which we have $b_4 = b_5 = b_6 = 0$. Since we also know $b_7 = b_8 = 0$, we conclude that $\mathbf{b} = 0$, which implies there are no non-zero admissible vectors. ■

5.2. Mutation class of $E_7^{(1,1)}$

Theorem 5.2. *There is no admissible vector for a quiver in the mutation class of $E_7^{(1,1)}$.*




Figure 5.2. $Q = E_7^{(1,1)}$.

Proof. The proof follows the same scheme as the one for the case of $E_6^{(1,1)}$, we omit explicit computations as they are very similar to the previous case but much longer.

We consider the quiver Q shown in Figure 5.2. Denote

$$\mu_* = \mu_5 \circ \mu_4 \circ \mu_3 \circ \mu_2 \circ \mu_1, \quad \mu_\diamond = \mu_8 \circ \mu_7 \circ \mu_6,$$

and consider

$$\mu = \mu_* \circ \mu_\diamond \circ \mu_* \circ \mu_\diamond \circ \mu_*.$$

Let $\mathbf{b} = (b_1, \dots, b_9)$ be an admissible vector. Denote by $\mathbf{b}' = (b'_1, \dots, b'_9)$ the result of mutation μ . One can check that $\mu(Q) = Q^{\text{op}}$. Then by the annulus property for Q we have

$$b_9 = -b_8 \leq 0,$$

and by the annulus property for $\mu(Q) = Q^{\text{op}}$ we need

$$b'_8 = -b'_9 \leq 0.$$

A computation similar to the one for $E_6^{(1,1)}$ shows that $b'_9 \leq b_9 \leq 0$, which implies $b'_9 = 0 = b'_8$ and similar constrains on the summands. Computing then b'_8 in exactly the same way as for $E_6^{(1,1)}$ we conclude that all $b_i = 0$ for $i = 1, \dots, 9$. ■

5.3. Mutation class of $E_8^{(1,1)}$

Theorem 5.3. *There is no admissible vector for a quiver in the mutation class of $E_8^{(1,1)}$.*




Figure 5.3. $Q = E_8^{(1,1)}$.

Proof. The proof is very similar to the one for $E_6^{(1,1)}$ and $E_7^{(1,1)}$. We consider the quiver Q shown in Figure 5.3. Denote

$$\mu_* = \mu_8 \circ \mu_6 \circ \mu_4 \circ \mu_3 \circ \mu_2, \quad \mu_\diamond = \mu_9 \circ \mu_7 \circ \mu_5 \circ \mu_1,$$

and consider

$$\mu = \mu_* \circ \mu_\diamond \circ \mu_* \circ \mu_\diamond \circ \mu_* \circ \mu_\diamond \circ \mu_* \circ \mu_\diamond \circ \mu_*.$$

As before, $\mu(Q) = Q^{\text{op}}$, and an explicit computation shows that the annulus property does not hold for $\mu(Q)$ unless $\mathbf{b} = 0$. ■

6. Mutation classes of X_6 and X_7

In this section, we show that there are no admissible vectors for mutation class X_7 . The proof is similar to the one for extended affine quivers. We also list all admissible vectors for quivers of type X_6 .




Figure 6.1. Quivers \$X_7\$ (left) and \$X_6\$ (right).

Theorem 6.1. *There is no admissible vector for a quiver in the mutation class of \$X_7\$.*

Proof. The idea is similar to the one we used for the case of \$E_6^{(1,1)}\$.

Consider the quiver \$Q\$ of type \$X_7\$ shown in Figure 6.1. From the annulus property for three double arrows, we get

$$b_1 = -b_2 \leq 0, \quad b_3 = -b_4 \leq 0, \quad b_5 = -b_6 \leq 0.$$

The composition of mutations

$$\mu_{012} = \mu_2 \circ \mu_1 \circ \mu_0$$

takes \$Q\$ to an isomorphic quiver with different location of double arrows (after \$\mu_{012}\$ the double arrows will be \$v_0v_2, v_3v_6\$ and \$v_4v_5\$). The annulus property for the mutated quiver \$\mu_{012}(Q)\$ after computing all entries would result in the following equations:

$$b_3 + b_6 + 2b_0 = 0, \quad b_4 + b_5 + 2b_0 = 0$$

(the third equation will be \$b_1 + b_2 = 0\$, which is satisfied automatically).

By symmetry, an application of another composition of mutations \$\mu_{034}\$ leads to the equations

$$b_1 + b_6 + 2b_0 = 0, \quad b_2 + b_5 + 2b_0 = 0,$$

and, similarly, one obtains from \$\mu_{056}\$ that

$$b_3 + b_2 + 2b_0 = 0, \quad b_4 + b_1 + 2b_0 = 0.$$

Adding all six equations together we get

$$2(b_1 + b_2) + 2(b_3 + b_4) + 2(b_5 + b_6) + 12b_0 = 0,$$

which implies \$b_0 = 0\$, and thus the six equations above result in

$$b_1 = b_3 = b_5 = -b_2 = -b_4 = -b_6 \leq 0.$$

So far, we have only used equalities arising from the annulus property but not the inequalities. Computing the value of b_3 after mutation μ_{012} (call it b'_3), one can find that if all assumptions from above hold then $b'_3 \leq 0$, while from the annulus property for $\mu_{012}(Q)$ one gets $b'_3 \geq 0$. This implies $b'_3 = 0$, which can hold in the only case of $b_3 = 0$ (similarly to E case), and hence $b_i = 0$ for all $i \in \{0, 1, 2, \dots, 6\}$. ■

Theorem 6.2. *Let Q be a quiver of type X_6 with a single arrow from v_0 to v_5 , where v_5 is a leaf (see Figure 6.1). Then a vector \mathbf{b} is admissible if and only if $b_5 = -2b_0 \geq 0$ and all other b_i vanish. Admissible vectors for all representatives of the mutation class are shown in Figure 6.2.*




Figure 6.2. Admissible vectors of all five quivers of type X_6 , $a \in \mathbb{N}$.

Proof. Let Q be as in Figure 6.1 (right). From the annulus property, we have

$$b_1 = -b_2 \leq 0, \quad b_3 = -b_4 \leq 0.$$

Two equations from μ_{012} and μ_{034} . Consider the sequences of mutations μ_{012}, μ_{034} . By the annulus properties in the resulting quivers, we get the following conditions:

$$b_4 + b_5 + 2b_0 = 0, \quad b_2 + b_5 + 2b_0 = 0. \tag{6.1}$$

Notice that we do not need to compute anything here: the equations follow from the computation for X_7 restricted to X_6 . From this we conclude that $b_2 = b_4$, i.e.

$$-b_1 = -b_3 = b_2 = b_4 = \beta$$

for some $\beta \geq 0$.

More equations from μ_5 . To obtain more equations, we will first apply mutation μ_5 to X_6 , then v_5 will be a source instead of a sink, so the resulting quiver will be isomorphic to the subquiver of X_7 where the vertex v_5 is removed.

More precisely, denote by b'_i the result of the application of μ_5 to \mathbf{b} , and call the image of b_5 by b'_6 (to use the restriction of X_7). Then the mutations μ_{012} and μ_{034} will lead to the following two equations:

$$b'_3 + b'_6 + 2b'_0 = 0, \quad b'_1 + b'_6 + 2b'_0 = 0. \tag{6.2}$$

The entries here are computed from mutation μ_5 as follows:

$$b'_1 = b'_3 = -\beta, \quad b'_2 = b'_4 = \beta, \quad b'_5 = -b_5, \quad b'_6 = b_0 + [b_5]_+,$$

and each of the equations in (6.2) leads to the following:

$$-\beta - b_5 + 2b_0 + 2[b_5]_+ = 0.$$

Since we also have $\beta + b_5 + 2b_0 = 0$ from equation (6.1), we obtain the following equations:

$$4b_0 + 2[b_5]_+ = 0, \quad 2\beta + 2b_5 - 2[b_5]_+ = 0,$$

which can be simplified to $2b_0 + [b_5]_+ = 0$ and $\beta = [-b_5]_+$.

We now have two cases to consider: either $b_5 \leq 0$ or $b_5 > 0$. If $b_5 \leq 0$, then $b_0 = 0$ and $\beta = -b_5$, so we obtain a vector

$$\mathbf{b} = (b_0, \dots, b_5) = (0, -\beta, \beta, -\beta, \beta, -\beta).$$

Applying mutation μ_{012} , we obtain a quiver with a double arrow v_4v_5 , and one can observe that the annulus property for this double arrow is not satisfied (unless $\beta = 0$ which implies $\mathbf{b} = 0$).

Therefore, we can assume $b_5 > 0$, so $\beta = 0$ and $b_5 = -2b_0$. We are left to show that all such vectors are admissible. The proof goes along the same lines as the proof of Theorem 4.3: we check that every generator of the cluster modular group leaves vector \mathbf{b} intact, where generators of the cluster modular group of the quiver of type X_6 shown in Figure 6.1 can be found in [18, Section 4.2].

■

Remark 6.3. Observe that the violation of the annulus property is the only argument used in the proof of Theorem 6.2 to show that a vector is not admissible. Therefore, we can conclude that for a quiver of type X_6 a vector is admissible if and only if the annulus property holds after every sequence of mutations. This observation will be generalized to all mutation-finite quivers in Section 10.

7. Rank 2 quivers

Theorem 7.1. *Let Q be a rank two quiver with the arrow from v_1 to v_2 of weight $a > 0$. Let $\mathbf{b} = (b_1, b_2)$ be an integer vector. Then*

- (1) *if $a = 1$ then \mathbf{b} is admissible for any b_1, b_2 ;*
- (2) *if $a = 2$ then \mathbf{b} is admissible if and only if $b_1 = -b_2 \leq 0$;*
- (3) *if $a > 2$ then there are no admissible vectors.*

Proof. The first and second parts concern finite and affine types.

To prove the third part, notice that after at most two mutations (and swapping the labels of v_1 and v_2 if needed) we may assume that $Q = v_1 \xrightarrow{a} v_2$, and $b_2 \geq 0 \geq b_1$. We may also assume that $|b_1| \geq |b_2|$ (otherwise replace μ_1 with μ_2 in the consideration below). Then after mutation μ_1 , we will get

$$b'_2 = b_2 - a(-b_1) = b_2 + ab_1 < b_2 + 2b_1 = (b_2 + b_1) + b_1 \leq b_1,$$

so, the absolute value of b_2 increases. Moreover, after swapping the labels of v_1 and v_2 the assumption above holds again, so we can mutate again to increase the components of the coefficient vector indefinitely. ■

8. Quivers from surfaces

In Section 3 we gave a general characterization of admissible vectors via peripheral laminations. We now want to make this more explicit by describing admissible vectors for a special triangulation from every mutation class. We exclude from our consideration disks with at most two punctures and unpunctured annuli as these correspond to quivers of finite or affine type and thus were considered either in [13] or in Section 4.

If a surface has no boundary, then, by Corollary 3.3, its quivers cannot have any admissible vector. Therefore, from now on we assume that a surface S has at least one boundary component.

A surface S contains the following *features*: boundary components (each with a number of boundary marked points), punctures and handles. To construct the triangulation we do the following:

- Choose any boundary component (we call it the *outer boundary component*) and a marked point p on it. All other boundary components will be called *inner* and the corresponding features will be called *holes*.




Figure 8.1. Standard triangulation of a surface with at least one boundary component.

- Place all features along a line from left to right, first all holes, then all handles, then all punctures, as in Figure 8.1, and enclose them by nested loops based at p so that every feature (except for the leftmost one in Figure 8.1) lies inside a digon with both vertices at p (recall that we excluded the case where S is a disk with one puncture).
- Triangulate the digons with features as follows:
 - each hole is enclosed by a loop x_i and the domain inside x_i triangulated as in Figure 8.2 (left);
 - each handle is enclosed by a loop y_i and the domain inside y_i triangulated as in Figure 8.2 (middle left);
 - each puncture inside a digon is connected by two arcs to two ends of the digon, see Figure 8.2 (middle right);
 - if there are no holes and handles, then the innermost monogon with two punctures is triangulated as in Figure 8.2 (right);
 - if the outer boundary contains other marked points than p , then the outermost loop at p separates a polygon (denote it S_{out}). S_{out} is triangulated as shown in Figure 8.1.

The quiver Q corresponding to the standard triangulation is shown in Figure 8.3. It consists of the following elements built into a chain (from right to left):

- quiver Q_{out} of triangulated outer polygon S_{out} ;
- quivers of digons with punctures;
- quivers of digons with handles;




Figure 8.2. Features (top row), their standard triangulations (middle row) and corresponding quivers (bottom). Columns from left to right: a digon with a hole, a digon with a handle, a digon with a puncture, a monogon with two punctures.




Figure 8.3. Quiver from standard triangulation.




Figure 8.4. Notation: subquivers of the quiver for standard triangulation.

- quivers of digons with holes;
- in case of absence of holes and handles, the leftmost element will be the quiver of a monogon with two punctures.

Notation 8.1. We will highlight the following subquivers of Q , as in Figure 8.4:

- Q_{out} : the subquiver of the outer polygon S_{out} (if the outer component contains other marked points than p);
- two vertices, v_1 and v_2 , connected to Q_{out} (see Figure 8.5 showing v_1 and v_2 depending on whether S_{out} is empty and whether the first feature from the right is a hole, a handle or a puncture), the arcs corresponding to v_1 and v_2 will be denoted by γ_1 and γ_2 ;
- Q_{in} : the subquiver corresponding to the inner boundary components, i.e. Q_{in} is spanned by all vertices corresponding to arcs of the triangulation with at least one endpoint on any of inner boundary components;
- the subquiver Q_I spanned by all other vertices of Q , where I is the index set of vertices not lying in Q_{out} , Q_{in} and different from v_1 and v_2 .

Theorem 8.2. *Let S be a surface with at least one boundary component distinct from a disk with at most two punctures and from an unpunctured annulus. Suppose that S is triangulated in the standard way. Then a coefficient vector $\mathbf{b} = (b_1, \dots, b_n)$ is admissible if and only if it satisfies the following conditions:*

- (a1) $b_i = 0$ for $i \in I$;
- (a2) the annulus property is satisfied;
- (a3) for the vertices v_1 and v_2 , one has $b_1 = -b_2 \leq 0$.




Figure 8.5. Vertices v_1 and v_2 for the cases when the rightmost feature is a hole (a), a handle (b), or a puncture (c), drawn for the case with $S_{\text{out}} \neq \emptyset$ (above) and for $S_{\text{out}} = \emptyset$ (below).

To prove the theorem we will use the following terminology.

Definition 8.3. Let L be a lamination and $C \in L$ be a curve. Let T be a triangulation. Then crossings of arcs of T with C cut C into *subsegments*, and by a *segment* we mean any connected union of subsegments of C (with respect to T). Two consecutive subsegments form a *crossing* with T . A crossing is *non-trivial* if its input into shear coordinates of L is non-zero, otherwise it is *trivial*. In the latter case, both subsegments can be isotopically deformed to be contained in a small neighbourhood of the same vertex q of the corresponding quadrilateral, and will be called *q-local*. The crossing formed by two *q-local* subsegments will be called *q-local*, as well as any segment formed of *q-local* subsegments.

We make the following elementary observation.

Proposition 8.4. Let T be a triangulation of a marked surface and L be a lamination. Choose $\gamma_i \in T$, and suppose that there exists a non-trivial crossing of γ_i with a curve $C \in L$. Then $b_i(L) \neq 0$ and $\text{sgn}(b_i(C)) = \text{sgn}(b_i(L))$.

The proof immediately follows from the definition of shear coordinates: segments inducing crossings of different signs inside a quadrilateral with diagonal γ_i intersect each other. The case of self-folded triangles is treated similarly.




Figure 8.6. Arcs γ_1 and γ_2 in case of no punctures in S (right), and otherwise (left). The grey circles indicate features (distinct from punctures on the right). If the outer boundary component contains a unique marked point, the arc γ_0 coincides with the outer boundary. Arcs γ_0, γ_1 and γ_2 form one triangle of the triangulation.

Proof of Theorem 8.2. In view of Theorem 3.2, we need to show that the conditions in the theorem hold if and only if the lamination is peripheral. The plan of the proof will be similar to the one of Lemma 4.1.

We will consider the arcs γ_1 and γ_2 corresponding to vertices v_1 and v_2 defined as shown in Figure 8.5. These arcs look as in Figure 8.6 depending on the presence of punctures in S .

Conditions (a1)–(a3) are necessary. We need to show that if L is peripheral, then (a1)–(a3) hold.

We start by proving (a3). Let L be a peripheral lamination. Notice that any peripheral curve homotopic to an inner boundary does not cross γ_1 and γ_2 . Consider peripheral curves homotopic to the outer boundary.

First, consider the closed curve C homotopic to the outer boundary, see Figure 8.7. It is easy to see that for this curve $b_1 = -b_2 = -1$. Furthermore, any non-closed peripheral curve has $b_1 = b_2 = 0$ (it either does not cross γ_1 and γ_2 at all, or consequently crosses p -locally all curves incident to p). So, no peripheral curve except for C can affect b_1 and b_2 , and hence condition (a3) is necessary.

Condition (a2) is necessary in view of Corollary 3.5. Condition (a1) is necessary since no peripheral curve crosses non-trivially any arc of the triangulation corresponding to any vertex of Q_I . Hence, conditions (a1)–(a3) are necessary.

Conditions (a1)–(a3) are sufficient. Next, we will prove that every lamination which is not peripheral contradicts some of conditions (a1)–(a3).




Figure 8.7. The closed curve C isotopic to the outer boundary component. The curves γ'_i play the role of γ_i in case of absence of punctures.

Suppose that L is a non-peripheral lamination and suppose that all conditions (a1)–(a3) are satisfied by L . Let us make several observations:

(O1) *No curve from L has any end on any inner boundary component except for the peripheral curves. No curve from L can spiral into a puncture. In particular, every curve $l \in L$ consists of finitely many subsegments.*

The first statement follows from condition (a2), the argument goes along the same lines as the part of the proof of Lemma 4.1 concerning bridging arcs. The second statement follows from the fact that a spiralling curve produces a non-zero shear coordinate on one of the two arcs incident to the puncture (see [11, Figure 36] and [6, Figure 6.3]) and from Proposition 8.4.

(O2) *Let x_i be an arc of T with both ends at p and enclosing exactly one inner boundary component, see Figure 8.2. Then for any curve $c \in L$ intersecting x_i the restriction of c onto the annulus cut out by x_i is a p -local segment of c .*

The statement follows immediately from (O1).

(O3) *Let $l \in L$, and let $\gamma \in T$ be incident to p and encircled by γ_1 , γ_2 , or $\gamma_1 \cup \gamma_2$. Then every intersection of l with γ is p -local.*

For arcs inside x_i this follows from (O2); all other arcs incident to p and encircled by γ_1 , γ_2 , or $\gamma_1 \cup \gamma_2$ correspond to vertices of Q belonging to Q_I , therefore the statement follows from Proposition 8.4 together with (O2).

(O4) *Let $l \in L$. Then every intersection of l with γ_i belongs to a p -local segment of l with two endpoints either on γ_i (if γ_i is a loop), or on $\gamma_1 \cup \gamma_2$ (otherwise).*




Figure 8.8. To the proof of (O5).

According to (O3), all subsegments of l inside a monogon bounded by γ_i (or the digon bounded by $\gamma_1 \cup \gamma_2$) are p -local, so they compose a p -local segment. Due to (O1), l is either closed or have both ends on the outer boundary component. Therefore, every maximal segment of l contained in γ_i (or in $\gamma_1 \cup \gamma_2$) has both ends on γ_i (or on $\gamma_1 \cup \gamma_2$, respectively).

(O5) Suppose that γ_1 and γ_2 are arcs with one endpoint in a puncture, as in Figure 8.6 (left). Let $l \in L$ and suppose that $b_1(l) \neq 0$ or $b_2(l) \neq 0$. Then l coincides with the closed curve C (see Figure 8.7).

Suppose that $b_1(l) \neq 0$ (the case of $b_2(l) \neq 0$ can be treated similarly). Let t_0 be an intersection point of l and γ_1 producing a non-trivial crossing. By (O4) there is a p -local segment t_0t_1 in l with $t_1 \in \gamma_1 \cup \gamma_2$, more precisely, $t_1 \in \gamma_2$, see Figure 8.8. Since the crossing at t_0 is non-trivial, the subsegment $t_{-1}t_0$ of l not lying on t_0t_1 should have its end t_{-1} on γ_2 . If $t_{-1} = t_1$ then l is the closed curve C .

Suppose that t_1 lies on γ_2 further from p than t_{-1} . Extending the segment $t_{-1}t_1 \in l$ past t_1 we will obtain a point t_2 on γ_1 lying further away from p than t_0 . By (O4), there is a p -local segment t_2t_3 of l with $t_3 \in \gamma_2$. Notice that we will get t_3 further away from p than t_1 . Continuing in the same way we will get infinitely many subsegments of l in contradiction to (O1). The case when t_1 lies on γ_2 closer to p than t_{-1} can be treated similarly (by extending the curve past t_{-1}).

(O6) Suppose that γ_1 and γ_2 are loops with both endpoints in p , as in Figure 8.6, right. Denote by l_1^\pm and l_2^\pm the positive and negative elementary laminations for γ_1 and γ_2 respectively, and by $D_C^r(l_i^\pm)$ twists along C applied to the curves above, $i = 1, 2, r \in \mathbb{Z}$. Denote by M the set of curves consisting of the closed curve C and the curves whose restriction onto $S \setminus S_{\text{out}}$ coincides with the restrictions of curves

$$l_1^+, \quad l_2^-, \quad D_C^r(l_1^+), \quad D_C^{-r}(l_2^-),$$

where $r > 0, i = 1, 2$. Then if $l \in L$ and $l \notin M$, we have $b_1(l) = b_2(l) = 0$.




Figure 8.9. To the proof of (O5): behaviour of curves on $S \setminus S_{out}$.

Let $l \in L$ be a curve, and suppose that at least one of $b_1(l)$ and $b_2(l)$ is not zero. This implies that l intersects at least one of γ_1 and γ_2 . Notice that $S \setminus S_{out}$ consists of one triangle bounded by $\gamma_0, \gamma_1, \gamma_2$ and two surfaces encircled by γ_1 and γ_2 respectively (here γ_0 may coincide with the outer boundary). In view of (O4), the segments of l contained inside the arcs γ_1 and γ_2 are p -local, and thus uniquely determined, see Figure 8.9 (left). We now want to list all possible subsegments of l inside the remaining triangle with two ends on γ_1 and γ_2 .

We say that a subsegment in the triangle joining γ_1 and γ_2 approaches γ_i from the right (left) if it is followed by a p -local segment whose other end can be reached by going around p counterclockwise (resp., clockwise). Since every subsegment joining γ_1 and γ_2 approaches them from one of the two sides, there are precisely four types of subsegments, they all are shown on Figure 8.9 (left).

Notice that two of the four subsegments intersect, which means that at most one of them can be a part of l ; we assume first that the one approaching both curves from the left does not appear. Gluing the p -local segments located inside γ_1 and γ_2 to all three remaining subsegments, we conclude that l can be assembled from the copies of the three curves shown in Figure 8.9 (middle); we will refer to these as to segments of types 1, 2, 3 respectively. These segments are attached to each other in l along p -local segments with both ends on the same curve γ_i .

Suppose that l does not contain any segment of type 3. It is easy to see that in this case none of the segments can be extended to an intersection with γ_0 (except for a p -local extension of a type 1 segment which has $b_1 = b_2 = 0$ and thus is excluded), which means that l is a closed curve. The only non-self-intersecting closed curve that can be composed out of segments of types 1 and 2 is the closed curve C .

Suppose now that l contains a segment of type 3. It can only be extended past its intersection with γ_1 by a type 1 segment, see Figure 8.9 (right). We obtain a segment with both ends on γ_2 . As it contains two p -local segments located inside γ_2 , we can determine which of the ends is closer to p along γ_2 , call it the lower end and the other one the upper end. Now, the upper end can be either joined to γ_0 or extended by a type 2 segment. Notice that if it is joined to γ_0 , then the lower end should also be

joined to γ_0 , and thus we obtain a restriction of l_1^+ onto $S \setminus S_{\text{out}}$. If the upper end is extended using a segment of type 2, then we obtain a new curve with both ends on γ_1 and well-defined upper and lower ends, so we can repeat the reasoning for the new upper end. We will need to connect the upper end to the boundary after finitely many steps (as l consists of finitely many of these segments). This will result in a restriction of $D_C^r(l_i^+)$, $i = 1, 2, r > 0$.

Finally, if while considering the four subsegments in the triangle we avoid the one approaching both curves from the right, then using precisely the same arguments we would obtain restrictions of curves $l_2^-, D_C^r(l_i^-)$ with $i = 1, 2, r < 0$.

(O7) *Let $l \in L$ and $b_1(l) \neq 0$ or $b_2(l) \neq 0$. Then l coincides with the closed curve C .*

If γ_1 and γ_2 are arcs incident to a puncture as in Figure 8.6 (left), then the statement follows immediately from (O5), so we may assume that γ_1 and γ_2 are loops as in Figure 8.6 (right).

Due to (O6), we need to consider the curves belonging to the set M only. Notice that any twist $D_C^k(l_i^+)$, $i = 1, 2, k \geq 0$ is not compatible with any twist $D_C^m(l_j^-)$, $j = 1, 2, m < 0$, since they contain intersecting subsegments (see Figure 8.9 (left)). Now, the negative shear coordinates (b_1, b_2) for $D_C^{k_1}(l_1^+)$ and $D_C^{k_2}(l_2^+)$ are equal to $(-(2k_1 + 1), 2k_1)$ and $(-2k_2, 2k_2 - 1)$, respectively. According to (a3) and Proposition 8.4, $k_1 \geq 0$ and $k_2 \geq 1$. It is easy to see that for any such curve the modulus of b_1 is strictly greater than the modulus of b_2 . For $D_C^m(l_j^-)$ the considerations are similar. For C ,

$$|b_1(C)| = |b_2(C)|.$$

Therefore, if L contains any curve from M except for C , then $|b_1(L)| \neq |b_2(L)|$ in contradiction to (a3).

Recall that L is a non-peripheral lamination. Let $C_{np} \in L$ be a non-peripheral curve and consider a lamination consisting of the single curve C_{np} (we will use the same notation for this lamination). We now show that there exists a non-peripheral curve which coincides with C_{np} inside $S \setminus S_{\text{out}}$ and has all shear coordinates equal to 0 in contradiction to [11].

Recall from Notation 8.1 that Q_{in} , Q_{out} and Q_I are subquivers of Q corresponding to inner boundary, outer boundary and the set defined in Figure 8.4. Denote by I_{in} and I_{out} the corresponding index sets. Denote also $Q_{12} = \langle v_1, v_2 \rangle$.

Observe:

- $b_i(C_{np}) = 0$ for $i \in I$ (by Proposition 8.4 and (a1));
- $b_i(C_{np}) = 0$ for $i = 1, 2$.

This follows immediately from Observation (O7) since C_{np} is non-peripheral, and hence does not coincide with C .




Figure 8.10. Shifting endpoints of the curve C_{np} on the outer boundary (to the segment on the left of p if the curve come to γ_0 from γ_2 , and to the segment on the right of p if it comes from γ_1).

- $b_i(C_{np}) = 0$ for $i \in I_{in}$.

This follows from applying observation (O2) to each inner boundary component. Therefore, we obtain that

- $b_i(C_{np}) = 0$ for $i \notin I_{out}$.

We are left to consider $b_i(C_{np})$ for $i \in I_{out}$ (notice that this only makes sense when S_{out} is non-empty). If ends of the curve C_{np} do not lie on the outer boundary component (i.e., C_{np} is closed), then C_{np} does not cross any arc corresponding to vertices of I_{out} and we have $b_i(C_{np}) = 0$ for $i \in I_{out}$. In this case all shear coordinates of C_{np} vanish, which contradicts [11]. Thus, we can assume that C_{np} has both ends on the outer boundary.

We will now modify the curve C_{np} by amending its intersection with the subsurface S_{out} only. The new curve C'_{np} is defined by shifting each endpoint of C_{np} to one of the boundary intervals containing the marked point p according to the following rules: the ends of segments crossing consequently γ_1 and γ_0 will be shifted clockwise along the outer boundary, and the ends of segments crossing consequently γ_2 and γ_0 will be shifted counterclockwise, see Figure 8.10. As a result, all crossings of C'_{np} with arcs in S_{out} are p -local (including the crossings with γ_0), and hence we get $b_i(C'_{np}) = 0$ for $i \in I_{out}$. As we also have $b_i(C'_{np}) = 0$ for $i \notin I_{out}$, we conclude that all shear coordinates of C'_{np} vanish, which leads to a contradiction.

This shows that non-peripheral lamination L satisfying (a1)–(a3) does not exist, which proves that the conditions (a1)–(a3) are sufficient. ■

9. Skew-symmetrizable mutation classes

In this section we consider the skew-symmetrizable case.

Let B be a skew-symmetrizable $n \times n$ matrix, i.e. there is an integer diagonal $n \times n$ matrix $D = (d_i)$ with positive entries such that BD is skew-symmetric. We suppose that B is mutation-finite and want to determine whether B can be complemented by one more row $(b_{n+1,1}, \dots, b_{n+1,n})$ so that the obtained $(n + 1) \times n$ matrix \tilde{B} will be still mutation-finite. As before, we call a vector $\mathbf{b} = (b_1, \dots, b_n)$ admissible if the matrix \tilde{B} composed of B and row $-\mathbf{b}$ is mutation-finite.

9.1. Diagrams and unfoldings

We recall basics on diagrams of skew-symmetrizable matrices.

Diagrams. According to [12], skew-symmetrizable matrices (b_{ij}) can be represented by diagrams with arrows from v_i to v_j of weight $-\text{sgn}(b_{ij}) b_{ij} b_{ji}$, which undergo mutations compatible with matrix mutations. A skew-symmetrizable matrix (b_{ij}) can be reconstructed by its diagram and the diagonal skew-symmetrizing matrix $D = (d_i)$. We will use a double arrow $i \rightrightarrows j$ to denote an arrow of weight 4 when $d_i = d_j$.

Notice that if B is skew-symmetrizable with the skew-symmetrizer $D = (d_i)$, then the $(n + 1) \times n$ matrix \tilde{B} can always be extended to a skew-symmetrizable $(n + 1) \times (n + 1)$ matrix by adding the $(n + 1)$ st column satisfying

$$b_{i,n+1} = -d_i b_{n+1,i},$$

and setting

$$d_{n+1} = 1.$$

This means that the matrix \tilde{B} can also be represented by a diagram (with arrows of weight $\text{sgn}(b_i) d_i b_i^2$ from v_i to the frozen vertex v_{n+1}).

One diagram with a frozen vertex may correspond to several skew-symmetrizable extended matrices, however, for any $k = 1, \dots, n$ mutations μ_k of such matrices always lead to the same extended diagram. We will call a diagram with a frozen vertex *mutation-finite* if it represents mutation-finite matrices (with respect to mutations in the first n vertices).

Mutation-finite diagrams without frozen vertices. It was shown in [6, 7] that mutation-finite diagrams either are skew-symmetric, or arise from triangulated orbifolds, or are of rank 2, or are mutation-equivalent to one of the seven types

$$F_4, \quad \tilde{G}_2, \quad \tilde{F}_4, \quad G_2^{(*,+)}, \quad G_2^{(*,*)}, \quad F_4^{(*,+)}, \quad F_4^{(*,*)}$$

shown in Figure 9.1.




Figure 9.1. Diagrams of exceptional skew-symmetrizable mutation-finite types.

We will consider the orbifolds, rank 2 diagrams, and each of the seven exceptional mutation-finite types separately, mostly based on the notion of *unfolding*.

Unfoldings. We briefly recall the definition of an unfolding of a skew-symmetrizable matrix introduced by A. Zelevinsky. For more details, see [7].

Let B be a skew-symmetrizable matrix with a skew-symmetrizer $D = (d_i)$. Suppose that we have chosen disjoint index sets E_1, \dots, E_k with $|E_i| = d_i$. Denote

$$m = \sum_{i=1}^k d_i.$$

Suppose also that we choose a skew-symmetric integer matrix C of size $m \times m$ with rows and columns indexed by the union of all E_i , such that

- (1) the sum of entries in each column of each $E_i \times E_j$ block of C equals b_{ij} ;
- (2) if $b_{ij} \geq 0$ then the $E_i \times E_j$ block of C has all entries non-negative.

Define a *composite mutation* $\hat{\mu}_i = \prod_{\hat{\tau} \in E_i} \mu_{\hat{\tau}}$ on C . This mutation is well defined, since all the mutations $\mu_{\hat{\tau}}, \hat{\tau} \in E_i$, for given i commute.

We say that C is an *unfolding* of B if C satisfies assertions (1) and (2) above, and for any sequence of iterated mutations $\mu_{k_1} \dots \mu_{k_m}(B)$, the matrix

$$C' = \hat{\mu}_{k_1} \dots \hat{\mu}_{k_m}(C)$$

satisfies assertions (1) and (2) with respect to $B' = \mu_{k_1} \dots \mu_{k_m}(B)$.

If C is an unfolding of a skew-symmetrizable integer matrix B , it is natural to define an *unfolding of a diagram* of B as a quiver of C . In general, we say that a quiver Q is an unfolding of a diagram Σ if there exist matrices B and C with diagram Σ and quiver Q respectively, and C is an unfolding of B . Note that a diagram may have many essentially different unfoldings.

We can also define an unfolding \tilde{C} of an extended skew-symmetrizable matrix \tilde{B} consisting of B and a row

$$(b_{n+1,1}, \dots, b_{n+1,n}) = -\mathbf{b} = -(b_1, \dots, b_n)$$

in the following way: it will consist of an unfolding C of B and a row vector $-\hat{\mathbf{b}}$ such that the block $E_{n+1} \times E_j$ consists of d_j equal entries $-b_{n+1,j}$. If we extend both matrices \tilde{B} and \tilde{C} with one additional column each to make them skew-symmetrizable and skew-symmetric respectively, then they will satisfy assertions (1) and (2) with respect to any sequence of mutations not including index $n + 1$.

This leads to a definition of an unfolding of a diagram with an additional frozen vertex. Such a diagram corresponds to an extended skew-symmetrizable matrix \tilde{B} , so we take an unfolding of it as defined above, add an additional column to make the obtained matrix skew-symmetric, and then take the corresponding quiver. Again, such a unfolding may not be unique.

Example 9.1. Consider the skew-symmetrizable exchange matrix B shown below and its diagram:


$$B = \begin{pmatrix} 0 & 1 \\ -4 & 0 \end{pmatrix}, \quad \bullet \xrightarrow{4} \bullet$$

We now can write the extended exchange matrix \tilde{B} with a coefficient vector (b_1, b_2) , add a column to make it skew-symmetrizable, and draw the corresponding diagram with a frozen vertex:

$$\tilde{B} = \begin{pmatrix} 0 & 1 \\ -4 & 0 \\ -b_1 & -b_2 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 0 & 1 & b_1 \\ -4 & 0 & 4b_2 \\ -b_1 & -b_2 & 0 \end{pmatrix} \rightsquigarrow \begin{array}{c} \square \\ \swarrow \text{sgn}(b_1)b_1^2 \quad \searrow 4\text{sgn}(b_2)b_2^2 \\ \bullet \xrightarrow{4} \bullet \end{array}$$

The results of unfoldings of both the matrix and the diagram are shown below:

$$\tilde{C} = \begin{pmatrix} 0 & 1 & 1 & 1 & 1 \\ -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ -b_1 & -b_2 & -b_2 & -b_2 & -b_2 \end{pmatrix} \rightsquigarrow \begin{pmatrix} 0 & 1 & 1 & 1 & 1 & b_1 \\ -1 & 0 & 0 & 0 & 0 & b_2 \\ -1 & 0 & 0 & 0 & 0 & b_2 \\ -1 & 0 & 0 & 0 & 0 & b_2 \\ -1 & 0 & 0 & 0 & 0 & b_2 \\ -b_1 & -b_2 & -b_2 & -b_2 & -b_2 & 0 \end{pmatrix}$$



In general, not every skew-symmetrizable matrix admits an unfolding. However, it is shown in [7] that every mutation-finite diagram without frozen vertices has a mutation-finite unfolding. This result provides us with a sufficient condition for a given coefficient vector to be admissible: we can always unfold a diagram with a frozen vertex to a quiver with a frozen vertex, and if the obtained quiver together with the unfolded coefficient vector is mutation-finite, then we immediately get the admissibility. A priori, this condition is not necessary for the admissibility: mutations of a diagram correspond to a very limited collection of mutations of the unfolded quiver, so the unfolded coefficient vector might not be admissible even when the initial vector is.

9.2. Diagrams from orbifolds and peripheral laminations

It is shown in [6] that the majority of skew-symmetrizable finite mutation classes originate from triangulated orbifolds. As in the surface case, coefficient vectors are in bijective correspondence with laminations, see [6] for details. Defining peripheral laminations in exactly the same way as for surfaces, and reasoning precisely as in Section 3, we obtain a similar result.

Theorem 9.2. *Let Σ be a diagram from a triangulated orbifold S . Then admissible vectors for Σ are in bijection with peripheral laminations on S .*

Similarly to the surface case, given a diagram from an orbifold, results of [16] allow one to reconstruct a triangulation, and results of [6, 11] allow one to reconstruct a lamination by a coefficient vector.

9.3. Rank 2 diagrams

Theorem 9.3. *Let Σ be a rank two diagram with the arrow from v_1 to v_2 of weight $a > 0$. Let $\mathbf{b} = (b_1, b_2)$ be an integer vector. Then*

- (1) *if $a < 4$, then \mathbf{b} is admissible for any b_1, b_2 ;*
- (2) *if $a = 4$, then \mathbf{b} is admissible if and only if $b_1 \leq 0 \leq b_2$ and $d_1 b_1^2 = d_2 b_2^2$;*
- (3) *if $a > 4$, then there are no admissible vectors.*

Proof. Part (1) concerns finite types, so it follows from [12].

For part (2) there are two cases: either $d_1 = d_2$, or we may assume that $d_1 = 1, d_2 = 4$. The former case is skew-symmetric and thus follows from Section 4. Let us now consider the latter case, the corresponding diagram with coefficient vector $\mathbf{b} = (b_1, b_2)$ is shown in Example 9.1. To prove the sufficiency, notice that in the assumptions (2) of the theorem the square roots of the weights of the diagram change

under mutations in the same way as the weights of arrows of the quiver $v_1 \Rightarrow v_2$ with coefficient vector $(-2b, b)$ with $b = b_2 \geq 0$, so the statement follows from Lemma 4.1 (alternatively, one can compute directly that both mutations act on the extended exchange matrix by multiplication by the negative identity matrix).

The proof of part (3) is similar to the skew-symmetric case. After at most two mutations (and swapping the labels of v_1 and v_2 if needed) we may assume that the diagram is $v_1 \xrightarrow{a} v_2$, and $b_1 \leq 0 \leq b_2$. We may also assume that $d_2 b_2^2 \leq d_1 b_1^2$ (otherwise replace μ_1 with μ_2 in the consideration below). Then after mutation μ_1 and swapping the labels of v_1 and v_2 we will obtain a diagram with coefficient vector (b'_1, b'_2) satisfying the same conditions and $|b'_i| > |b_i|, i = 1, 2$. Applying iterative mutations we can increase the components of the coefficient vector indefinitely. ■

9.4. Affine mutation classes

Every exceptional mutation class of diagrams of affine type contains a representative shown in Figure 9.2. Every mutation class of diagrams of affine type originating from an orbifold either contains a representative with a double arrow (we show one for every mutation class in Figure 9.2), or contains a representative with a subdiagram considered in Example 9.1 (see [6]). We treat these two cases separately, see Theorems 9.4 and 9.6.




Figure 9.2. Special representatives from non-skew-symmetric mutation classes of affine types.

Theorem 9.4. *Let Σ be a diagram of type $\tilde{G}_2, \tilde{F}_4, \tilde{B}_n$ or \tilde{C}_n shown in Figure 9.2. A coefficient vector \mathbf{b} is admissible if and only if it satisfies the annulus property.*

Proof. The annulus property is obviously a necessary condition for admissibility of a coefficient vector for given diagrams. To see that it is also sufficient, notice that these diagrams can be unfolded to quivers of type $\tilde{D}_4, \tilde{E}_6, \tilde{D}_n$ and $\tilde{A}_{n,n}$ shown in

Figure 4.3. The unfolded coefficient vectors still satisfy the annulus property, so every unfolded quiver with frozen vertex is mutation-finite by Theorem 4.3. This implies that the initial diagrams are mutation-finite as well, so the initial coefficient vectors are admissible. ■

Remark 9.5. By using unfoldings, we can extend the result of Theorem 4.4 to a general skew-symmetrizable case, i.e. for any diagram Σ of affine type containing a double arrow, a vector \mathbf{b} is admissible if and only if \mathbf{b} satisfies the annulus property.

In the case of a diagram containing a subdiagram from Example 9.1, we cannot use unfolding: the unfolded diagram is simply-laced, so the annulus property does not lead to any restrictions.

Theorem 9.6. *Let Σ be a diagram of affine type containing a subdiagram of type $v_1 \xrightarrow{4} v_2$ with $d_1 = 1, d_2 = 4$. A coefficient vector \mathbf{b} is admissible if and only if*

$$b_1 = -2b_2 \leq 0.$$

We will abuse notation by calling the condition in Theorem 9.6 the *annulus property* as well.

Proof of Theorem 9.6. The necessity follows from Theorem 9.3 (2).

To prove the sufficiency, notice that all diagrams in question correspond to an unpunctured disk with two orbifold points and several marked points at the boundary (see [6]). In particular, any triangulation corresponding to such a diagram consists of a monogon shown in Figure 9.3 (d) and a polygon S_{out} .

Now the proof is similar to part (2) of the rank 2 case. In the assumptions of the theorem, the square roots of the weights of the diagram change under mutations in the same way as the weights of arrows of the diagram of type \tilde{C}_n shown in Figure 9.2 with coefficient vector satisfying the annulus property, so the statement follows from Theorem 9.4. ■

9.5. Extended affine mutation classes

The result here is similar to the skew-symmetric case.

Theorem 9.7. *There are no admissible vectors for diagrams of any extended affine mutation class.*

Proof. Assume that Σ is one of the four diagrams of extended affine type (see Figure 9.1), and let \mathbf{b} be an admissible coefficient vector. It is clear that \mathbf{b} must satisfy the annulus property.




Figure 9.3. Standard triangulations of orbifolds: (a) a triangulated digon with an orbifold point, (b) corresponding quiver, (c) vertices v_1 and v_2 , (d) triangulated monogon with two orbifold points.

The diagrams of types $F_4^{(*,*)}$ and $G_2^{(*,+)}$ have unfoldings to quivers of type $E_6^{(1,1)}$, the diagram of type $F_4^{(*,+)}$ has an unfolding to a quiver of type $E_7^{(1,1)}$, and the diagram of type $G_2^{(*,*)}$ has an unfolding to a quiver of type $E_8^{(1,1)}$, where all the unfolded quivers are precisely those shown in Figures 5.1, 5.2, and 5.3, see [7]. Moreover, it is easy to see that the mutation sequences used in the proof of Theorems 5.1 and 5.3 are sequences of composite mutations (with respect to the unfoldings above) for certain sequences of mutations for the diagrams.

Therefore, if we take an unfolding Q of Σ with the unfolded coefficient vector $\hat{\mathbf{b}}$ and apply a mutation sequence $\hat{\mu}$ constructed in Section 5 such that $\hat{\mu}(\hat{\mathbf{b}})$ does not satisfy the annulus property, then there exists a mutation sequence μ of Σ such that $\mu(\mathbf{b})$ does not satisfy the annulus property either, which shows that \mathbf{b} cannot be admissible. ■

9.6. Diagrams from orbifolds

We now want to extend Theorem 8.2 to the orbifolds case. As in the surface case, we exclude finite and affine types, i.e. unpunctured disks with at most two orbifold points and once punctured disks with at most one orbifold point.

First, we can define a standard triangulation of an orbifold in a similar way. We add orbifold points to the list of features, and place them to the left of all other features. We then place the two leftmost orbifold points in a monogon (see Figure 9.3 (d)), and all the others in digons (Figure 9.3 (a)), the subdiagram corresponding to the digon is shown in Figure 9.3 (b).

Vertices v_1 and v_2 are defined in the same way as in the surface case. In the case of orbifold points being the only features (note that there should be at least three of them and thus there is at least one digon, otherwise the diagram is of finite or affine type), v_1 and v_2 are defined as in Figure 9.3 (c). The set I is also defined in the same way.

Theorem 9.8. *Let \mathcal{O} be an orbifold with at least one boundary component distinct from an unpunctured disk with at most two orbifold points and from once punctured disk with at most one orbifold point. Suppose that \mathcal{O} is triangulated in the standard way. Then a coefficient vector $\mathbf{b} = (b_1, \dots, b_n)$ is admissible if and only if it satisfies the following conditions:*

- (a1) $b_i = 0$ for $i \in I$;
- (a2) the annulus property is satisfied;
- (a3) for the vertices v_1, v_2 , one has $b_1 = -b_2 \leq 0$ if $d_1 \geq d_2$, and $b_1 = -2b_2 \leq 0$ if $d_1 < d_2$.

The proof is exactly the same as in the surface case. The only extra case is when the only features are orbifold points (otherwise, all the arcs incident to orbifold points belong to the set I), and, as in the surface case, the admissibility condition is prescribed by the shear coordinates $-b_1$ and $-b_2$ of the closed curve C . If $d_1 = 2d_2$, then one has $b_1 = -1 = -b_2$, and if $d_2 = 2d_1$, then one has $b_1 = -2$ and $b_2 = 1$, which gives precisely (a3).

10. Annulus property as criterion of mutation finiteness

We now prove a criterion of mutation finiteness in terms of the annulus property applied to the whole mutation class, this was proposed by Sergey Fomin. For simplicity, we consider the skew-symmetric case (i.e., quivers), which then can be easily generalized to the skew-symmetrizable case (see Remark 10.3).

Theorem 10.1. *Let Q be a quiver with a frozen vertex v . Suppose that the subquiver $Q \setminus v$ is mutation-finite. Then Q is mutation-finite if and only if the annulus property holds in every quiver Q' mutation-equivalent to Q for every double arrow contained in $Q' \setminus v$.*

Proof. The necessity follows from Corollary 3.5. Below we prove that the condition is also sufficient. As before, we denote by \mathbf{b} the coefficient vector associated with the vertex v .

Assume that the annulus property holds in any quiver of the mutation class. Let $Q_* = \langle Q \setminus v \rangle$ be the subquiver spanned by all mutable vertices. We will consider the following cases:

- if Q_* is of finite type, then every vector \mathbf{b} is admissible by [13];
- if Q_* is affine, then the statement follows from Theorem 4.4;
- if Q_* is mutation-equivalent to X_6 , then the statement follows from Remark 6.3;




Figure 10.1. Humphries generators of PMod in the case $g \geq 2$; small circles stay for punctures and boundary components (left). Construction of the annulus (right)

- if Q_* is mutation-equivalent to X_7 or $E_6^{(1,1)}$, $E_7^{(1,1)}$, $E_8^{(1,1)}$, then the proofs of Theorems 5.1–6.1 show that the annulus property cannot be satisfied in every quiver of the mutation class, and thus there are no admissible vectors;
- if Q_* is of rank 2, the statement follows immediately from Theorem 7.1;
- otherwise, Q_* is a quiver arising from a surface; the rest of the proof below is aimed at settling the question for this case.

From now on we will assume that Q_* is of surface type, and will assume that \mathbf{b} is *not* admissible. Our aim is to find a quiver in the mutation class for which the annulus property does not hold.

By Theorem 3.2, as \mathbf{b} is not admissible, it corresponds to a non-peripheral lamination L . Hence, there exists a closed curve intersecting the lamination L in a non-trivial way. We will pick such an intersecting curve in a particular way and will use it to construct a special triangulation, which will provide us with a quiver where annulus property does not hold.

Let g be the genus of the surface S corresponding to the quiver Q_* . We will consider separately the cases of $g \geq 2$, $g = 1$ and $g = 0$.

Case 1: $g \geq 2$. For a surface S of genus $g \geq 2$, the pure mapping class group $\text{PMod}(S)$ is generated by finitely many Dehn twists with respect to non-separating curves (see e.g. [4, Corollary 4.16]). One can choose these curves as in Figure 10.1 (left); these are called *Humphries generators*, see e.g. [4, Section 4.4.4]. Denote these curves C_1, \dots, C_m . As the lamination L is not peripheral, at least one of the curves C_1, \dots, C_m intersects L (otherwise, the pure mapping class group will act on L trivially). We will assume that C_1 intersects L .

Let M be a marked point (either a boundary marked point of a puncture). Let α_1 and α_2 be non-intersecting non-self-intersecting paths connecting M to each of the sides of C_1 , such paths exist since C_1 is a non-separating curve. Let γ_1 and γ_2 be loops

based at M and constructed by $\gamma_1 = \alpha_1\beta\alpha_1^{-1}$ and by $\gamma_2 = \alpha_2\beta\alpha_2^{-1}$, where β is the closed path along C_1 , see Figure 10.1 (right). Let T be any triangulation containing arcs γ_1 and γ_2 . We will show that in the triangulation T the annulus property breaks.

Indeed, as L crosses C_1 non-trivially, the restriction of L to the annulus bounded by γ_1 and γ_2 is a non-empty non-peripheral curve. Hence, the annulus property for this annulus does not hold in view of Lemma 4.1.

Case 2: $g = 1$. In this case, one cannot always find the generators of the mapping class group by twists along non-separating curves only, however there exists a set of generators by twists along finitely many non-separating curves and $k - 1$ boundary curves, where k is the number of boundary components on S (here, by a boundary curve we mean a closed peripheral curve along one of the boundary components), see [4].

If $k \leq 1$, then there still exists a set of generators by twists in non-separating curves, and we can use the same reasoning as before.

If $k > 1$, then for every generator we can find at least one marked point on each side with respect to the corresponding curve (i.e. for non-separating curves we proceed as before, and for a boundary curve we choose a marked point lying on the corresponding boundary component and a marked point lying on a different boundary component). So, we still are able to construct the annulus as in Figure 10.1 (right), but possibly using different marked points on different sides.

Case 3: $g = 0$. In this case, any generator of the pure mapping class group is a twist along a separating curve, but for any such curve one can find at least one marked point on each of its sides, and thus one can apply the same construction of an annulus as before. ■

Corollary 10.2. *Let Q be a quiver with a frozen vertex. Then Q is mutation-finite if and only if for every quiver Q' in the mutation class of Q every rank 3 subquiver of Q' is mutation-finite.*

Proof. The “only if” direction is evident. Suppose that Q is mutation-infinite. If the subquiver $Q_* = Q \setminus v$ (where v is the frozen vertex) is also mutation-infinite, then the mutation class of Q_* contains a quiver with an arrow of multiplicity higher than 2 and hence any connected rank 3 subquiver containing that arrow is mutation-infinite. If Q_* is mutation-finite, then, by Theorem 10.1, there is a quiver Q' in the mutation class of Q where the annulus property does not hold (for some vertices v_1, v_2 connected by a double arrow). Then the rank 3 subquiver of Q' spanned by $\langle v_1, v_2, v \rangle$ is mutation-infinite. ■

Remark 10.3. Both Theorem 10.1 and Corollary 10.2 can be easily generalized to the skew-symmetrizable case, where the annulus property is understood as in Theorem 9.6. The proofs are the same as in the skew-symmetric case.

Acknowledgements. We would like to thank Sergey Fomin for the question inspiring the current project and for helpful suggestions, Michael Shapiro for stimulating discussions, and Dani Kaufman for sharing then unpublished results of [19] with us. We are grateful to the anonymous referee for useful remarks. A substantial part of the paper was written at the Isaac Newton Institute for Mathematical Sciences, Cambridge; we are grateful to the organizers of the program “Cluster algebras and representation theory”, and to the Institute for support and hospitality during the program; this work was supported by EPSRC grant no EP/R014604/1.

Funding. Research of P.T. was supported in part by the Leverhulme Trust research grant RPG-2019-153.

References

- [1] M. Alim, S. Cecotti, C. Córdova, S. Espahbodi, A. Rastogi, and C. Vafa, [BPS quivers and spectra of complete \$\mathcal{N} = 2\$ quantum field theories](#). *Comm. Math. Phys.* **323** (2013), no. 3, 1185–1227 Zbl [1305.81118](#) MR [3106506](#)
- [2] S. Cecotti and C. Vafa, [Classification of complete \$\mathcal{N} = 2\$ supersymmetric theories in 4 dimensions](#). In *Surveys in differential geometry. Geometry and topology*, pp. 19–101, Surv. Differ. Geom. 18, International Press, Somerville, MA, 2013 Zbl [1320.81085](#) MR [3087917](#)
- [3] H. Derksen and T. Owen, [New graphs of finite mutation type](#). *Electron. J. Combin.* **15** (2008), no. 1, article no. 139 Zbl [1180.05052](#) MR [2465763](#)
- [4] B. Farb and D. Margalit, [A primer on mapping class groups](#). Princeton Math. Ser. 49, Princeton University Press, Princeton, NJ, 2012 Zbl [1245.57002](#) MR [2850125](#)
- [5] A. Felikson, M. Shapiro, H. Thomas, and P. Tumarkin, [Growth rate of cluster algebras](#). *Proc. Lond. Math. Soc. (3)* **109** (2014), no. 3, 653–675 Zbl [1386.13061](#) MR [3260289](#)
- [6] A. Felikson, M. Shapiro, and P. Tumarkin, [Cluster algebras and triangulated orbifolds](#). *Adv. Math.* **231** (2012), no. 5, 2953–3002 Zbl [1256.13014](#) MR [2970470](#)
- [7] A. Felikson, M. Shapiro, and P. Tumarkin, [Cluster algebras of finite mutation type via unfoldings](#). *Int. Math. Res. Not. IMRN* (2012), no. 8, 1768–1804 Zbl [1283.13020](#) MR [2920830](#)
- [8] A. Felikson, M. Shapiro, and P. Tumarkin, [Skew-symmetric cluster algebras of finite mutation type](#). *J. Eur. Math. Soc. (JEMS)* **14** (2012), no. 4, 1135–1180 Zbl [1262.13038](#) MR [2928847](#)

- [9] V. V. Fock and A. B. Goncharov, [Dual Teichmüller and lamination spaces](#). In *Handbook of Teichmüller theory. Vol. I*, pp. 647–684, IRMA Lect. Math. Theor. Phys. 11, European Mathematical Society, Zürich, 2007 Zbl [1162.32009](#) MR [2349682](#)
- [10] S. Fomin, M. Shapiro, and D. Thurston, [Cluster algebras and triangulated surfaces. I. Cluster complexes](#). *Acta Math.* **201** (2008), no. 1, 83–146 Zbl [1263.13023](#) MR [2448067](#)
- [11] S. Fomin and D. Thurston, [Cluster algebras and triangulated surfaces. II. Lambda lengths](#). *Mem. Amer. Math. Soc.* **255** (2018), no. 1223 Zbl [07000309](#) MR [3852257](#)
- [12] S. Fomin and A. Zelevinsky, [Cluster algebras. II. Finite type classification](#). *Invent. Math.* **154** (2003), no. 1, 63–121 Zbl [1054.17024](#) MR [2004457](#)
- [13] S. Fomin and A. Zelevinsky, [Cluster algebras. IV. Coefficients](#). *Compos. Math.* **143** (2007), no. 1, 112–164 Zbl [1127.16023](#) MR [2295199](#)
- [14] C. Fraser, [Braid group symmetries of Grassmannian cluster algebras](#). *Selecta Math. (N.S.)* **26** (2020), no. 2, article no. 17 Zbl [1436.13049](#) MR [4066538](#)
- [15] W. Gu, [A decomposition algorithm for the oriented adjacency graph of the triangulations of a bordered surface with marked points](#). *Electron. J. Combin.* **18** (2011), no. 1, article no. 91 Zbl [1217.05213](#) MR [2795772](#)
- [16] W. Gu, [The decomposition algorithm for skew-symmetrizable exchange matrices](#). *Electron. J. Combin.* **19** (2012), no. 2, article no. 54 Zbl [1256.65020](#) MR [2946112](#)
- [17] T. Henrich, [Mutation classes of diagrams via infinite graphs](#). *Math. Nachr.* **284** (2011), no. 17-18, 2184–2205 Zbl [1262.13039](#) MR [2859758](#)
- [18] T. Ishibashi, [Presentations of cluster modular groups and generation by cluster Dehn twists](#). *SIGMA Symmetry Integrability Geom. Methods Appl.* **16** (2020), article no. 025 Zbl [1439.13054](#) MR [4082289](#)
- [19] D. Kaufman and Z. Greenberg, [Cluster modular groups of affine and doubly extended cluster algebras](#). 2021, arXiv:[2107.10334v1](#)
- [20] A. I. Seven, [Mutation classes of finite type cluster algebras with principal coefficients](#). *Linear Algebra Appl.* **438** (2013), no. 12, 4584–4594 Zbl [1319.13013](#) MR [3039212](#)

Received 12 May 2022; revised 30 August 2023.

Anna Felikson

Department of Mathematical Sciences, Durham University, Mathematical Sciences and Computer Science Building, Upper Mountjoy Campus, Stockton Road, Durham DH1 3LE, UK; anna.felikson@durham.ac.uk

Pavel Tumarkin

Department of Mathematical Sciences, Durham University, Mathematical Sciences and Computer Science Building, Upper Mountjoy Campus, Stockton Road, Durham DH1 3LE, UK; pavel.tumarkin@durham.ac.uk