

On the interaction of the Coxeter transformation and the rowmotion bijection

René Marczinzik, Hugh Thomas, and Emine Yıldırım

Abstract. Let P be a finite poset and L the associated distributive lattice of order ideals of P . Let ρ denote the rowmotion bijection of the order ideals of P viewed as a permutation matrix and C the Coxeter matrix for the incidence algebra kL of L . Then, we show the identity $(\rho^{-1}C)^2 = \text{id}$, as was originally conjectured by Sam Hopkins. Recently, it was noted that the rowmotion bijection is a special case of the much more general grade bijection R that exists for any Auslander regular algebra. This motivates to study the interaction of the grade bijection and the Coxeter matrix for general Auslander regular algebras. For the class of higher Auslander algebras coming from n -representation finite algebras, we show that $(R^{-1}C)^2 = \text{id}$ if n is even and $(R^{-1}C + \text{id})^2 = 0$ when n is odd.

1. Introduction

Let A be a finite dimensional algebra over a field k with finite global dimension. We will always assume that k is a splitting field for the algebra A , which, for example, is true if k is algebraically closed or if A is a quiver algebra. We denote the indecomposable projective A -modules by P_i for $i = 1, \dots, n$. Then, the *Cartan matrix* M of A is defined as the $n \times n$ -matrix with entries $m_{i,j} := \dim_k \text{Hom}_A(P_i, P_j)$. The *Coxeter matrix* (a.k.a. Coxeter transformation) C of A is then defined as $C := -M^{-1}M^T$. Note that this is well defined as the Cartan matrix of an algebra of finite global dimension has determinant 1 or -1 , and thus, M is invertible over \mathbb{Z} , see, for example, [1, Proposition III.3.10]

The Coxeter matrix of a finite dimensional algebra with finite global dimension is the main object of study in the spectral theory of finite dimensional algebras, we refer, for example, to the survey article [15]. Of special interest in homological algebra are algebras with periodic Coxeter matrix, that is, $C^l = \text{id}$ for some $l \geq 1$. Algebras with periodic Coxeter matrix include, for example, fractionally Calabi–Yau algebras that arise in geometric and combinatorial contexts; we refer for example to [3, 21].

Mathematics Subject Classification 2020: 16G10 (primary); 16E10, 05E16, 05E18 (secondary).

Keywords: distributive lattices, grade bijection, Coxeter transformation.

When P is a finite poset, then the set of order ideals of P is a distributive lattice L and every finite distributive lattice arises this way and is uniquely determined by P . The Coxeter matrix of the incidence algebra kL can be directly read off from elementary combinatorial data involving the Möbius function of L . When S is an antichain of P , then set $I(S)$ to be the order ideal whose maximal elements are given by S and set $M(S)$ to be the order ideal whose minimal non-elements are given by S . The *rowmotion bijection* is given on the elements of L by sending $I(S)$ to $M(S)$, which defines a bijection. The rowmotion bijection is one of the main attractions in dynamical algebraic combinatorics, we refer, for example, to the articles [5, 19, 20]. When L has n elements, we can associate the rowmotion bijection to an $n \times n$ -permutation matrix in a natural way. While the Coxeter matrix and the rowmotion bijection are well studied objects in representation theory and combinatorics, respectively, it seems that no relation between them has been shown before.

The following theorem, which is our first main result, was first conjectured by Sam Hopkins, who noted the identity in several examples.

Theorem 1.1. *Let L be a finite distributive lattice with C the Coxeter matrix of the incidence algebra of kL and ρ the rowmotion bijection for L viewed as a permutation matrix. Then, $\rho^{-1}C$ has order two, that is $(\rho^{-1}C)^2 = \text{id}$.*

In the article [11] it was noted that the rowmotion bijection for a distributive lattice L is a special case of a much more general bijection that exists for any Auslander regular algebra. Recall here that a finite dimensional algebra A is called *Auslander regular* if A has finite global dimension and in the minimal injective coresolution

$$0 \rightarrow A \rightarrow I_0 \rightarrow I_1 \rightarrow \dots \rightarrow I_n \rightarrow 0$$

we have that the projective dimension of I_i is bounded by i for all $i \geq 0$. In [6], Iyama defined the so-called *grade bijection* of an Auslander regular algebra A . It is a permutation of the simple modules of A , whose precise definition we will recall in the next section. Again, we can associate a permutation matrix to the grade bijection that we will usually denote by R_A , or R for short, for an Auslander regular algebra A . One of the main results in [11] states that a finite lattice L is distributive if and only if the incidence algebra kL is Auslander regular. Moreover, when L is distributive, the grade bijection gives a homological realisation of the rowmotion bijection, when one identifies the elements of the lattice L with the simple kL -modules in a natural way. This leads to the natural question, whether for other Auslander regular algebras there is a simple relation between the Coxeter matrix and the grade bijection.

We give a positive answer for an important class of algebras that also appears in combinatorics. Namely, an algebra A is called *n -representation-finite* if A has global dimension at most n and there is an n -cluster tilting object M in $\text{mod } A$, which is

then uniquely given when we assume that M is basic. The notion of n -representation finite algebras was introduced in [10] and includes many important classes of algebras such as all path algebras of Dynkin type, higher Auslander algebras of Dynkin type A that have strong relations to the combinatorics of cyclic polytopes [16] and the 2-representation finite algebras that have strong relations to quivers with potential and Jacobian algebras [4]. For an n -representation finite A with n -cluster tilting module M , the endomorphism algebra $B := \text{End}_A(M)$ will be an higher Auslander algebra. Higher Auslander algebras are those algebras B with finite global dimension and a minimal injective coresolution

$$0 \rightarrow B \rightarrow I_0 \rightarrow I_1 \rightarrow \cdots \rightarrow I_n \rightarrow 0$$

such that the projective dimension of I_i is zero for $i = 0, 1, \dots, n - 1$, when n denotes the global dimension. Thus, by definition, higher Auslander algebras are always Auslander regular. Our second main result is as follows for such algebras B .

Theorem 1.2. *Let A be an n -representation finite algebra with n -cluster tilting module M . Let $B = \text{End}_A(M)$ with grade bijection R and Coxeter matrix C . If n is even, then $(R^{-1}C)^2 = \text{id}$ and if n is odd, then $(R^{-1}C + \text{id})^2 = 0$.*

2. Preliminaries

We assume that algebras are finite dimensional over a field k and that they are non-semisimple and connected unless stated otherwise. Additionally, we assume that k is a splitting field, which is, for example, automatic if k is algebraically closed or if A is a quiver algebra. Here, k being a splitting field for the k -algebra A means that every simple module A -module S has the property that $\text{End}_A(S) \cong k$; see, for example, [14, Chapter 7] for more equivalent characterisations and properties of splitting fields. We assume that modules are right modules unless otherwise stated. $D = \text{Hom}_k(-, k)$ denotes the duality and J denotes the Jacobson radical of an algebra. We assume that the reader is familiar with the basics of representation theory and homological algebra of finite dimensional algebras and we refer, for example, to the textbooks [2, 18]. Let $\nu_A := D \text{Hom}_A(-, A)$ denote the Nakayama functor of an algebra A and $\nu_A^{-1} = \text{Hom}_A(D(A), -)$ its inverse. It is well known that ν_A induces an equivalence between the category of projective A -modules and the category of injective A -modules with inverse ν_A^{-1} . The *global dimension* of an algebra A is defined as the supremum of all projective dimensions of A -modules. The *dominant dimension* of an algebra A with minimal injective coresolution

$$0 \rightarrow A \rightarrow I_0 \rightarrow I_1 \rightarrow \cdots$$

is defined as the smallest $n \geq 0$ such that I_n is not projective or as infinite if there is no such I_n . For an A -module M , $\text{add}(M)$ will denote the full subcategory of $\text{mod } A$ consisting of all direct summands of M^n for some natural number n . Let $K_0(A)$ denote the Grothendieck group of an algebra A with finite global dimension with basis given by the indecomposable projective modules $[P]$. For simplicity, we will often omit the usual brackets $[P]$ for an element in the Grothendieck group $K_0(A)$ when there is no danger of confusion. When A has finite global dimension, the *Coxeter transformation* of A is defined as $C_A([P]) := -[v_A(P)]$. When this linear transformation is expressed as a matrix with respect to the basis of $K_0(A)$ given by the classes of the indecomposable projective modules, we recover the matrix C from Section 1.

An algebra A is called *Auslander regular* when A has finite global dimension and there exists an injective coresolution

$$0 \rightarrow A \rightarrow I^0 \rightarrow I^1 \rightarrow \dots \rightarrow I^n \rightarrow 0$$

of the regular module such that $\text{pdim } I^i \leq i$ for all $i \geq 0$. Important classes of Auslander regular algebras are incidence algebras of distributive lattices (see [11]), higher Auslander algebras (see [7]) and blocks of category \mathcal{O} (see [13]). The *grade* of an A -module M is defined by

$$\text{grade } M := \inf \{i \geq 0 \mid \text{Ext}_A^i(M, A) \neq 0\}.$$

Dually, the *cograde* of a module M is defined as $\inf\{i \geq 0 \mid \text{Ext}_A^i(D(A), M) \neq 0\}$. For every Auslander regular algebra, there exists a bijection on the simple A -modules as follows.

Theorem 2.1 ([6, Theorem 2.10]). *Let A be an Auslander regular algebra. Then, there is a bijection Gr_A sending a simple A -module S to the simple A -module $\text{Gr}_A(S) = \text{top}(D \text{Ext}_A^{g_S}(S, A))$, where $g_S := \text{grade } S$. The grade of S is equal to the cograde of $\text{Gr}_A(S)$.*

We call the bijection Gr_A as in the previous theorem for Auslander regular algebras the *grade bijection* on simple A -modules. We refer to [6, Theorem 2.10] for a more general statement and proofs. In [11], it was shown that the grade bijection gives a categorification of the rowmotion map for incidence algebras of distributive lattices. We define the grade bijection on the Grothendieck group of an Auslander regular algebra A by

$$R_A([P_S]) = [P_{\text{Gr}_A(S)}],$$

where P_S denotes the projective cover of a simple module S . We define the *rowmotion Coxeter transformation* of an Auslander regular algebra A as $R_A^{-1}C_A$. Note that since R_A is a permutation matrix, we have $R_A^{-1} = R_A^T$. We will be mainly interested in the

minimal polynomial of $R_A^{-1}C_A$ and the relation $(R_A^{-1}C_A)^2 = \text{id}$. Note that since

$$C_A R_A^{-1} = R_A (R_A^{-1} C_A) R_A^{-1},$$

the operators $C_A R_A^{-1}$ and $R_A^{-1} C_A$ are similar and thus they have the same minimal polynomials and we have

$$(C_A R_A^{-1})^2 = \text{id}$$

if and only if $(R_A^{-1} C_A)^2 = \text{id}$.

3. The Coxeter transformation and the rowmotion bijection for distributive lattices

Let P denote a finite poset and L the distributive lattice of order ideals of P . Let S be an antichain of P and $I(S)$ the order ideal whose maximal elements are given by S and $M(S)$ the order ideal whose minimal non-elements are given by S . Formally,

$$I(S) = \{x \in P : x \leq s \text{ for some } s \in S\},$$

$$M(S) = P \setminus \{y \in P : y \geq s \text{ for some } s \in S\}.$$

The *Hasse quiver* of a poset P is the finite quiver with points $x \in P$ and an arrow $x \rightarrow y$ if x covers y (note that we use here the opposite convention compared to [11]). The *incidence algebra* kP of a poset P over a field k is defined as the quiver algebra kQ/\mathcal{I} with Q the Hasse quiver of P and the relations \mathcal{I} such that any two paths that start and end at the same points get identified.

The *rowmotion bijection* ρ for a distributive lattice L given as the set of order ideals of a poset P is defined as the permutation sending $I(S)$ to $M(S)$. As explained in the preliminaries, the rowmotion bijection is a special case of the more general grade bijection when viewing the incidence algebra of a distributive lattice as an Auslander regular algebra. We will denote by P_O the indecomposable projective module in the incidence algebra kL of a distributive lattice L corresponding to the order ideal O and $J_O := \nu_A(P_O)$ (we use J instead of I for the indecomposable injectives in this section to avoid confusion with $I(S)$). When $A = kL$ is the incidence algebra of a distributive lattice, we set


$$R_A([P_{I(S)}]) = [P_{M(S)}]$$

motivated by the fact that the grade bijection gives a homological realisation of the rowmotion bijection as explained after Theorem 2.1. We will use the following result, whose proof can be found in [11, Theorem 3.2] and where one can also find a description of the differentials. See also [21, Definition 2.3], where this construction was anticipated, though restricted to a particular case.


Theorem 3.1. *Let L be a distributive lattice given as the set of order ideals of a poset P . Then, a minimal projective resolution of the indecomposable injective module $J_{I(S)}$ for an antichain S is given as follows in kL :*

$$\begin{aligned}
 0 \rightarrow P_{M(S)} \rightarrow \bigoplus_{T \subseteq S, |T|=r-1} P_{M(T)} \rightarrow \cdots \rightarrow \bigoplus_{T \subseteq S, |T|=i} P_{M(T)} \rightarrow \cdots \\
 \cdots \rightarrow \bigoplus_{T \subseteq S, |T|=1} P_{M(T)} \rightarrow P_{M(\emptyset)} \rightarrow J_{I(S)} \rightarrow 0.
 \end{aligned}$$

Example 3.2. For an example of the previous theorem, consider the case where P is the poset shown on the left in the diagram below. The lattice L is the distributive lattice of order ideals of P , shown on the right.



Let $S = \{x_1, x_2\}$. The projective resolution of $J_{I(S)}$ given by Theorem 3.1 is as follows, where we represent kL modules by the corresponding representations of the Hasse quiver.



Corollary 3.3. *Let L be a distributive lattice given as the set of order ideals of a poset P . Let S denote an antichain of P . Then, the Coxeter transformation C_A for $A = kL$ is given by*

$$C_A([P_{I(S)}]) = - \sum_{T \subseteq S} (-1)^{|T|} [P_{M(T)}].$$

Theorem 3.4. *Let $A = kL$ be the incidence algebra of a distributive lattice L given as the set of order ideals of a poset P . Then, $(R_A^{-1}C_A)^2 = \text{id}$.*


Proof. We first look at $C_A R_A^{-1} C_A$ and determine the values on a basis:

$$\begin{aligned} C_A R_A^{-1} C_A([P_{I(S)}]) &= C_A R_A^{-1} \left(- \sum_{T \subseteq S} (-1)^{|T|} [P_{M(T)}] \right) \\ &= C_A \left(- \sum_{T \subseteq S} (-1)^{|T|} [P_{I(T)}] \right) \\ &= \sum_{T \subseteq S} \sum_{R \subseteq T} (-1)^{|T|} (-1)^{|R|} [P_{M(R)}] \\ &= \sum_{R \subseteq S} \sum_{R \subseteq T \subseteq S} (-1)^{|T| - |R|} [P_{M(R)}] \\ &= \sum_{R \subseteq S} [P_{M(R)}] \sum_{R \subseteq T \subseteq S} (-1)^{|T| - |R|}. \end{aligned}$$

If R is properly contained in S , the inner sum is over the $2^{|S|-|R|}$ choices for T , half with $|T| - |R|$ even and half with $|T| - |R|$ odd. The inner sum is therefore zero in this case. The only surviving term of the outer sum is $R = S$, in which case we also have $T = S$. The result is that $C_A R_A^{-1} C_A([P_{I(S)}]) = [P_{M(S)}]$ and if we now apply R_A^{-1} to both sides, we get that $R_A^{-1} C_A R_A^{-1} C_A([P_{I(S)}]) = [P_{I(S)}]$, as desired. ■

The next example shows that the identity $(R_A^{-1}C_A)^2 = \text{id}$, which is true when A is the incidence algebra of a distributive lattices, does not hold for Auslander regular incidence algebras of posets that are not lattices.

Example 3.5. The following poset with incidence algebra A is Auslander regular:



Let P_i denote the indecomposable projective A -modules and I_i the indecomposable injective A -modules. A computer program such as the GAP-package [17] can be used to show that A is Auslander regular and to find the grade bijection and Coxeter matrix. We just give a sketch for the calculations without all details in the following. One finds the injective resolution of each indecomposable projective, and then checks the

projective dimension of the injectives that appear. The minimal injective resolutions of indecomposable projectives are given by

$$\begin{aligned}
 0 &\rightarrow P_1 \rightarrow I_6 \rightarrow 0, \\
 0 &\rightarrow P_2 \rightarrow I_6 \rightarrow I_3 \rightarrow 0, \\
 0 &\rightarrow P_3 \rightarrow I_6 \rightarrow I_2 \rightarrow 0, \\
 0 &\rightarrow P_4 \rightarrow I_6 \rightarrow I_5 \rightarrow 0, \\
 0 &\rightarrow P_5 \rightarrow I_6 \rightarrow I_4 \rightarrow 0, \\
 0 &\rightarrow P_6 \rightarrow I_6 \rightarrow I_4 \oplus I_5 \rightarrow I_2 \oplus I_3 \rightarrow I_1 \rightarrow 0.
 \end{aligned}$$

Since the algebra is isomorphic to its opposite algebra, one can dually obtain the minimal projective resolution of the indecomposable injectives and from that the projective dimensions of the indecomposable injectives to see that A is indeed Auslander regular.

We emphasise that R_A here is given by the grade bijection. Currently, no purely combinatorial description of this bijection is available for Auslander regular incidence algebras of posets except in the case that the poset is a distributive lattice. Such a combinatorial description would be a generalisation of rowmotion on distributive lattices and would be very interesting to have. In forthcoming work [12], we show that the grade bijection R_A of a general Auslander regular algebra A is given by sending P_i to the last term in the minimal projective resolution of $v_A(P_i) = I_i$, which allows to calculate the grade bijection directly from the minimal projective resolutions of the I_i as above.

The matrix for R_A is given as follows:

$$\begin{pmatrix}
 0 & 0 & 0 & 0 & 0 & 1 \\
 0 & 0 & 1 & 0 & 0 & 0 \\
 0 & 1 & 0 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 1 & 0 \\
 0 & 0 & 0 & 1 & 0 & 0 \\
 1 & 0 & 0 & 0 & 0 & 0
 \end{pmatrix},$$

and the Coxeter matrix C_A is given as follows:

$$\begin{pmatrix}
 -1 & -1 & -1 & -1 & -1 & -1 \\
 1 & 0 & 1 & 0 & 0 & 0 \\
 1 & 1 & 0 & 0 & 0 & 0 \\
 -1 & 0 & 0 & 0 & 1 & 0 \\
 -1 & 0 & 0 & 1 & 0 & 0 \\
 1 & 0 & 0 & 0 & 0 & 0
 \end{pmatrix}.$$

The matrix $R_A^{-1}C_A$ has minimal polynomial $x^3 - x^2 - x + 1$, and thus, the identity $(R_A^{-1}C_A)^2 = \text{id}$ is not true for this poset.

4. n -representation finite algebras

We recall some basics on cluster-tilting modules and higher Auslander algebras. An A -module M is called n -cluster-tilting when M is a generator-cogenerator and $M^{\perp n} = \text{add}(M) = {}^{\perp n}M$, where $M^{\perp n} = \{N \in \text{mod } A \mid \text{Ext}_A^i(M, N) = 0 \text{ for all } 1 \leq i < n\}$ and ${}^{\perp n}M = \{N \in \text{mod } A \mid \text{Ext}_A^i(N, M) = 0 \text{ for all } 1 \leq i < n\}$. A 1-cluster tilting module M exists if and only if A is representation finite; in this case, $\text{add}(M) = \text{mod } A$. By the higher Auslander correspondence, see, for example, [7], M is n -cluster tilting if and only if the algebra $B := \text{End}_A(M)$ is a higher Auslander algebra of global dimension $n + 1$; that is, it has global dimension and dominant dimension equal to $n + 1$. We denote by

$$\tau_n := \tau \Omega^{n-1} \quad \text{for } n \geq 1$$

the n -Auslander-Reiten translate and by $\tau_n^{-1} := \Omega^{-(n-1)}\tau^{-1}$ the inverse n -Auslander-Reiten translate. Let A be an algebra, and let M be a generator-cogenerator of $\text{mod } A$ and $B := \text{End}_A(M)$. In the following, we will summarise several properties of B in this situation and refer, for example, to [2, Chapter VI.5] for more information. There is an equivalence of categories $\text{add}(M) \cong \text{proj } B$, given by $\text{Hom}_A(M, -)$. We denote the indecomposable projective B -module associated an indecomposable A -module $N \in \text{add}(M)$ by $L_N := \text{Hom}_A(M, N)$. Note that $v_B(L_N) = D \text{Hom}_B(L_N, B) = D \text{Hom}_B(\text{Hom}_A(M, N), \text{Hom}_A(M, M)) \cong D \text{Hom}_A(N, M)$, and thus, the indecomposable injective B -modules are given by $T_N := D \text{Hom}_A(N, M)$ for an indecomposable module $N \in \text{add}(M)$. We denote by S_N the simple B -module with projective cover L_N . In this section, the modules P_i will refer to terms in a minimal projective resolution of a module and not to the indecomposable projective modules of an algebra corresponding.

An algebra A is called n -representation-finite for some $n \geq 1$ if $\text{gldim } A \leq n$ and there is an n -cluster tilting module M in $\text{mod } A$, see, for example [10], where such algebras were studied systematically for the first time. Note that an n -cluster tilting module M in a n -representation-finite algebra is unique when we assume that M is basic. Note also that M necessarily contains the indecomposable projective and injective modules as direct summands. 1-representation-finite algebras are exactly the representation finite hereditary algebras. They are classified by Dynkin diagrams, see [2, Chapter VIII].

We collect several results that we will need. For a survey on n -cluster tilting categories and higher Auslander algebras, we refer to Section 2 of [9]. For the definition of n -almost split sequences and their basic properties, we refer to Section 2.3 in [9],

which also contains the next lemma. We recall that the length of an n -almost split sequence is equal to $n + 2$.

Lemma 4.1. *Let A be an algebra with n -cluster-tilting module M and $B := \text{End}_A(M)$.*

- (1) *Let N be an indecomposable non-projective summand of M . There is an n -almost split sequence*

$$0 \rightarrow \tau_n(N) \rightarrow \cdots \rightarrow N \rightarrow 0.$$

Applying $\text{Hom}_A(M, -)$ to it yields a minimal projective resolution of S_N :

$$0 \rightarrow L_{\tau_n(N)} \rightarrow \cdots \rightarrow L_N \rightarrow S_N \rightarrow 0. \tag{4.1}$$

- (2) *Let N be an indecomposable non-injective summand of M . There is an n -almost split sequence*

$$0 \rightarrow N \rightarrow \cdots \rightarrow \tau_n^{-1}(N) \rightarrow 0.$$

Applying $D \text{Hom}_A(-, M)$, we obtain a minimal injective coresolution of S_N :

$$0 \rightarrow S_N \rightarrow T_N \rightarrow \cdots \rightarrow T_{\tau_n^{-1}(N)} \rightarrow 0. \tag{4.2}$$

As an immediate consequence of this lemma, we deduce that if N is an indecomposable summand of M which is not projective, then the projective dimension of S_N is $n + 1$; similarly, if N is not injective, then the injective dimension of S_N is $n + 1$.

Lemma 4.2. *Let A be an n -representation-finite algebra with n -cluster tilting module M . Let $X, Y \in \text{add}(M)$.*

- (1) $\text{Hom}_A(\tau_n^{-1}(Y), X) = D \text{Ext}_A^n(X, Y)$.
- (2) $\text{Hom}_A(Y, \tau_n(X)) = D \text{Ext}_A^n(X, Y)$.

Proof. We show (1); the proof of (2) is dual. It holds in general (without the assumption that A has global dimension at most n) that $\underline{\text{Hom}}_A(\tau_n^{-1}(Y), X) = D \text{Ext}_A^n(X, Y)$ for any $X, Y \in \text{add}(M)$ for a cluster tilting module M , by [8, Theorem 2.3.1]. Now, using the assumption that A has global dimension at most n , we can additionally conclude that $\underline{\text{Hom}}_A(\tau_n^{-1}(Y), X) = \text{Hom}_A(\tau_n^{-1}(Y), X)$ by [10, Lemma 2.4(d)]. ■

Lemma 4.3. *Let N be an A -module of finite projective dimension. Then, $\text{pdim } N = \sup\{i \geq 0 \mid \text{Ext}_A^i(N, A) \neq 0\}$.*

Proof. See, for example, [2, Lemma VI.5.5]. ■

Given this lemma, it follows from the definition of n -representation finite algebra that the summands of M are either projective or of projective dimension n .

Lemma 4.4. *Let A be an n -representation finite algebra with n -cluster-tilting module M and $B := \text{End}_A(M)$.*

- (1) *Let N be an indecomposable projective summand of M . We have $T_N \cong L_{v_A(N)}$, and thus, T_N is projective as a B -module.*
- (2) *Let N be an indecomposable non-projective summand of M . Suppose that its minimal projective resolution is*

$$0 \rightarrow P_n \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow N \rightarrow 0.$$

Then, the projective resolution as a B -module of T_N is given by

$$0 \rightarrow L_{\tau_n(N)} \rightarrow L_{v_A(P_n)} \rightarrow \cdots \rightarrow L_{v_A(P_0)} \rightarrow T_N \rightarrow 0. \tag{4.3}$$

This then also gives an injective coresolution

$$0 \rightarrow L_{\tau_n(N)} \rightarrow T_{P_n} \rightarrow \cdots \rightarrow T_{P_0} \rightarrow T_N \rightarrow 0$$

of $L_{\tau_n(N)}$ with $T_{P_i} \cong L_{v_A(P_i)}$.

- (3) *Let N' be an injective indecomposable direct summand of M . Then, $L_{N'}$ is itself injective, so it is its own injective resolution.*
- (4) *There is a bijection between the indecomposable projective B -modules of injective dimension $n + 1$ and the indecomposable injective B -modules of projective dimension $n + 1$ given by $\Omega^{-(n+1)}$ with inverse Ω^{n+1} .*

Proof. (1) Since N is projective,

$$T_N = D \text{Hom}_A(N, M) \cong \text{Hom}_A(M, v_A(N)) = L_{v_A(N)}$$

by [18, Chapter III, Corollary 6.2], so T_N is itself projective.

(2) Since N is non-projective, we have already established that $\text{pdim } N = n$. In order to calculate $\text{Ext}^i(N, M)$, we would apply $\text{Hom}(-, M)$ to the projective resolution of N , and since $\text{Ext}^i(N, M) = 0$ except for $i = 0$ and $i = n$, we obtain an exact sequence

$$0 \rightarrow \text{Hom}_A(N, M) \rightarrow \text{Hom}_A(P_0, M) \rightarrow \text{Hom}_A(P_1, M) \rightarrow \cdots \tag{4.4}$$

$$\cdots \rightarrow \text{Hom}_A(P_n, M) \rightarrow \text{Ext}_A^n(N, M) \rightarrow 0.$$

Now, by the higher Auslander–Reiten formulas from Lemma 4.2, we have that

$$\text{Ext}_A^n(N, M) \cong D \text{Hom}_A(M, \tau_n(N)).$$

Note that $D \text{Hom}_A(N, M) =: T_N$. Thus, applying the duality D to equation (4.4), we obtain the exactness of equation (4.3). It is a projective resolution since the terms

$D \operatorname{Hom}_A(P_i, M) \cong L_{v_A(P_i)}$ are projective-injective by (1), which also shows that the second exact sequence is an injective coresolution of $L_{\tau_n(N)}$.

(3) This is dual to (1).

(4) The indecomposable projective B -modules which are not injective are the modules L_N for N an indecomposable direct summand of M which is not injective. By (4.3), we see that $\Omega^{-(n+1)}L_N = T_{\tau_n(N)}$. The modules of the form $T_{\tau_n(N)}$ are exactly the indecomposable injective B -modules which are not projective, and (4.3) shows that

$$\Omega^{n+1}T_{\tau_n(N)} = L_N. \quad \blacksquare$$

In the next theorem, we calculate the Coxeter transformation of higher Auslander algebras coming from n -cluster tilting modules in n -representation-finite algebras.

Theorem 4.5. *Let A be an n -representation-finite algebra with n -cluster-tilting module M , let $B = \operatorname{End}_A(M)$ and let N be an indecomposable A -module in $\operatorname{add}(M)$.*

- (1) *If N is projective, $C_B([L_N]) = -[L_{v_A(N)}]$.*
- (2) *If N is non-projective, let $0 \rightarrow P_n \rightarrow \dots \rightarrow P_1 \rightarrow P_0 \rightarrow N \rightarrow 0$ be a minimal projective resolution of N . Then,*

$$C_B([L_N]) = \sum_{i=0}^n (-1)^{i+1} [L_{v_A(P_i)}] + (-1)^n [L_{\tau_n(N)}].$$

Proof. By definition, $C_B([L_N]) = -[T_N]$. To express $[T_N]$ with respect to the basis of indecomposable projective B -modules, it suffices to calculate a projective resolution of T_N and take the alternating sum. We have found these projective resolutions in Lemma 4.4, and the claim follows. \blacksquare

Recall that we defined the grade bijection in the Grothendieck group of an Auslander regular algebra B on the indecomposable projective A -modules by $R_B([P_S]) = [P_{\operatorname{Gr}_B(S)}]$, where P_S denotes the projective cover of a simple module S and Gr_B the grade bijection on simple modules. We now calculate the grade bijection for higher Auslander algebras that are endomorphism rings of an n -cluster tilting module.

Theorem 4.6. *Let A be an algebra with n -cluster tilting module M having endomorphism algebra $B := \operatorname{End}_A(M)$ and let N be an indecomposable module in $\operatorname{add}(M)$. Then, the grade bijection R_B for B is given by*

$$R_B([L_N]) = [L_{v_A(N)}]$$

when N is projective and $R_B([L_N]) = [L_{\tau_n(N)}]$ else.

Proof. Assume first that N is projective. The simple B -module S_N injects into $T_N \cong L_{v_A(N)}$, which is a summand of B ; hence, $\operatorname{Hom}_B(S_N, B) \neq 0$ and thus $\operatorname{grade} S_N = 0$.

Applying $D \operatorname{Hom}_B(-, B)$ on $L_N \rightarrow S_N \rightarrow 0$, one gets $T_N \rightarrow D \operatorname{Hom}_B(S_N, B) \rightarrow 0$, and since T_N is projective and the middle term does not vanish, we have

$$\operatorname{top}(D \operatorname{Hom}_B(S_N, B)) = \operatorname{top}(T_N) = \operatorname{top}(L_{v_A(N)}) = S_{v_A(N)}.$$

Thus,

$$R_B([L_N]) = [L_{v_A(N)}].$$

Assume now that N is not projective. We will show that $D \operatorname{Ext}_B^i(S_N, B) = S_{\tau_n(N)}$, if $i = n + 1$ and $D \operatorname{Ext}_B^i(S_N, B) = 0$, else. This will show that $\operatorname{grade} S_N = n + 1$ and $R_B([L_N]) = [L_{\tau_n(N)}]$. By the projective resolution of S_N given in (4.1), we know that $D \operatorname{Ext}_B^i(S_N, B)$ is the i th homology of the complex obtained from

$$0 \rightarrow L_{\tau_n(N)} \rightarrow \cdots \rightarrow L_N \rightarrow 0$$

by applying $D \operatorname{Hom}_B(-, B)$. This complex equals

$$0 \rightarrow T_{\tau_n(N)} \rightarrow \cdots \rightarrow T_N \rightarrow 0,$$

which by (4.2) for $\tau_n(N)$ is exact except at the $(n + 1)$ th degree, where the homology equals $S_{\tau_n(N)}$. ■

The last theorem tells us how to calculate the grade bijection. In the next theorem, we will also use the inverse of the grade bijection, which is then given by $R_B^{-1}([L_N]) = [L_{v_A^{-1}(N)}]$ if N is injective and $R_B^{-1}([L_N]) = [L_{\tau_n^{-1}(N)}]$ otherwise.

Theorem 4.7. *Let A be an n -representation-finite algebra with n -cluster-tilting module M and $B := \operatorname{End}_A(M)$. Then, we have $(C_B R_B^{-1})^2 = \operatorname{id}$ if n is even and $(C_B R_B^{-1} + \operatorname{id})^2 = 0$ when n is odd.*

Proof. Supposing that N is an injective indecomposable summand of M , it follows that

$$C_B R_B^{-1}([L_N]) = C_B([L_{v_A^{-1}(N)}]) = -[L_N],$$

since $v_A^{-1}(N)$ is projective. If N is an indecomposable and non-injective summand of M , we have

$$C_B R_B^{-1}([L_N]) = C_B([L_{\tau_n^{-1}(N)}]) = \sum_{i=0}^n (-1)^{i+1} [L_{v_A(P_i)}] + (-1)^n [L_N],$$

where

$$0 \rightarrow P_n \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow \tau_n^{-1}(N) \rightarrow 0$$

denotes a minimal projective resolution of $\tau_n^{-1}(N)$.

It follows that with respect to the basis of the $[L_N]$, ordered such that the elements corresponding to the injective modules N precede all the others, the matrix of $C_B R_B^{-1}$ has the block form

$$\begin{pmatrix} -I_s & E \\ 0 & (-1)^n I_t \end{pmatrix}$$

for some matrix E , where I_r denotes the $r \times r$ identity matrix, s is the number of injective indecomposable modules N in $\text{add}(M)$, and t is the number of the non-injective such modules. If n is even, then the square of such a matrix is the identity I_{s+t} , whereas if n is odd, then by adding the identity I_{s+t} one gets the matrix

$$\begin{pmatrix} 0 & E \\ 0 & 0 \end{pmatrix}$$

whose square is zero. ■

Specialising to path algebras of Dynkin type gives the following special case.

Example 4.8. Let $A = KQ$ be a path algebra of Dynkin type and M the direct sum of all indecomposable A -modules, which is a 1-cluster tilting module. Let $B := \text{End}_A(M)$, which is the Auslander algebra of A . By Theorem 4.7, we have $(C_B R_B^{-1} + \text{id})^2 = 0$.

Acknowledgements. The authors would like to express their gratitude to Sam Hopkins for communicating his conjecture to us. This project began at a workshop on Dynamical Algebraic Combinatorics hosted (online) by Banff International Research Station in 2020. The authors are grateful to BIRS and to the organisers of the workshop, who provided the opportunity for the project to begin. We thank the referee for their detailed comments and suggestions which significantly improved the exposition of the paper. In particular, we followed the suggestion of the referee to include a shorter proof for Theorems 4.6 and 4.7.

Funding. R. Marczinzik is funded by the DFG with the project number 428999796. H. Thomas is partially supported by an NSERC Discover Grant RGPIN-2016-04872 and the Canada Research Chairs program, grant number CRC-2021-00120. Part of the work on this paper was undertaken during Emine Yıldırım’s tenure as INI-Simons Post Doctoral Research Fellow hosted by the Isaac Newton Institute for Mathematical Sciences (INI) participating in programme “Cluster algebras and representation theory” and by the Department of Pure Mathematics and Mathematical Statistics (DPMMS) at the University of Cambridge. This author would like to thank INI and DPMMS for support and hospitality during this fellowship, which was supported by Simons Foundation (Award ID 316017) and by EPSRC (grant no. EP/R014604/1).

References

- [1] I. Assem, D. Simson, and A. Skowroński, *Elements of the representation theory of associative algebras. Vol. 1*. London Math. Soc. Stud. Texts 65, Cambridge University Press, Cambridge, 2006 Zbl 1092.16001 MR 2197389
- [2] M. Auslander, I. Reiten, and S. O. Smalø, *Representation theory of Artin algebras*. Cambridge Stud. Adv. Math. 36, Cambridge University Press, Cambridge, 1997 Zbl 0834.16001 MR 1476671
- [3] A. Chan, E. Darpö, O. Iyama, and R. Marczinzik, Periodic trivial extension algebras and fractionally Calabi–Yau algebras. [v1] 2020, [v4] 2024, arXiv:2012.11927v4
- [4] M. Herschend and O. Iyama, Selfinjective quivers with potential and 2-representation-finite algebras. *Compos. Math.* **147** (2011), no. 6, 1885–1920 Zbl 1260.16016 MR 2862066
- [5] S. Hopkins, Order polynomial product formulas and poset dynamics. [v1] 2020, [v6] 2024, arXiv:2006.01568v6
- [6] O. Iyama, Symmetry and duality on n -Gorenstein rings. *J. Algebra* **269** (2003), no. 2, 528–535 Zbl 1034.16017 MR 2015291
- [7] O. Iyama, Auslander correspondence. *Adv. Math.* **210** (2007), no. 1, 51–82 Zbl 1115.16006 MR 2298820
- [8] O. Iyama, Higher-dimensional Auslander–Reiten theory on maximal orthogonal subcategories. *Adv. Math.* **210** (2007), no. 1, 22–50 Zbl 1115.16005 MR 2298819
- [9] O. Iyama, Auslander–Reiten theory revisited. In *Trends in representation theory of algebras and related topics*, pp. 349–397, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2008 Zbl 1206.16011 MR 2484730
- [10] O. Iyama, Cluster tilting for higher Auslander algebras. *Adv. Math.* **226** (2011), no. 1, 1–61 Zbl 1233.16014 MR 2735750
- [11] O. Iyama and R. Marczinzik, Distributive lattices and Auslander regular algebras. *Adv. Math.* **398** (2022), article no. 108233 Zbl 1495.16014 MR 4379204
- [12] V. Klasz, R. Marczinzik, and H. Thomas, On the grade bijection for Auslander regular algebras. Preprint
- [13] H. Ko, V. Mazorchuk, and R. Mrden, Some homological properties of category \mathcal{O} . VI. *Doc. Math.* **26** (2021), 1237–1269 Zbl 1486.16008 MR 4324465
- [14] T. Y. Lam, *A first course in noncommutative rings*. 2nd edn., Grad. Texts in Math. 131, Springer, New York, 2001 Zbl 0980.16001 MR 1838439
- [15] H. Lenzing and J. A. de la Peña, Spectral analysis of finite dimensional algebras and singularities. In *Trends in representation theory of algebras and related topics*, pp. 541–588, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2008 Zbl 1210.16013 MR 2484735
- [16] S. Oppermann and H. Thomas, Higher-dimensional cluster combinatorics and representation theory. *J. Eur. Math. Soc. (JEMS)* **14** (2012), no. 6, 1679–1737 Zbl 1254.05197 MR 2984586
- [17] The QPA Team, *QPA – Quivers, path algebras and representations – a GAP package, Version 1.33*. 2022, <https://folk.ntnu.no/oyvinso/QPA/>, visited on 11 September 2024

- [18] A. Skowroński and K. Yamagata, *Frobenius algebras. I. Basic representation theory*. EMS Textbk. Math., EMS Press, Zürich, 2011 Zbl 1260.16001 MR 2894798
- [19] J. Striker, *Dynamical algebraic combinatorics: promotion, rowmotion, and resonance*. *Notices Amer. Math. Soc.* **64** (2017), no. 6, 543–549 Zbl 1370.05219 MR 3585535
- [20] H. Thomas and N. Williams, *Rowmotion in slow motion*. *Proc. Lond. Math. Soc. (3)* **119** (2019), no. 5, 1149–1178 Zbl 1459.06010 MR 3968720
- [21] E. Yıldırım, *The Coxeter transformation on cominuscule posets*. *Algebr. Represent. Theory* **22** (2019), no. 3, 699–722 Zbl 1471.16028 MR 3951768

Received 25 January 2022; revised 8 April 2024.

René Marczinzik

Mathematical Institute of the University of Bonn, University of Bonn, Endenicher Allee 60, 53115 Bonn, Germany; marczire@math.uni-bonn.de

Hugh Thomas

Département de mathématiques, Université du Québec à Montréal, CP 8888, Succursale Centre-ville, Montréal QC, H3C 3P8, Canada; thomas.hugh_r@uqam.ca

Emine Yıldırım

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WB, UK; ey260@cam.ac.uk, emineyildirim@gmail.com