



Trace inequalities for Sobolev martingales

Dmitriy Stolyarov

Abstract. We study limiting trace inequalities in the style of Maz'ya and Meyers–Ziemer for Sobolev martingales. We develop the Bellman function approach to such estimates, which allows to provide sufficient and almost necessary conditions on the martingale space and the martingale transform under which the trace inequalities hold true.

1. Introduction

The space of summable functions and its subspaces play a special role in analysis. Many statements that are true for L_p spaces in the reflexive regime $1 < p < \infty$ break at the endpoints; one may recall the L_p boundedness of singular integrals as an example. Other remain true when interpreted properly, but the proofs might be more complicated. This is the instance for Sobolev embedding theorems. While the foundations of this theory in the reflexive regime were laid by Sobolev in [37], the endpoint case $p = 1$ was not covered until the work of Gagliardo [13] and Nirenberg [29]. Even then, many natural interesting questions were left open in the case $p = 1$. One of the directions of research was initiated by Bourgain and Brezis in [6]. Roughly speaking, we may describe Bourgain–Brezis inequalities as

$$(1.1) \quad \|f\|_{L_p(\mathbb{R}^d)} \lesssim \|Af\|_{L_1(\mathbb{R}^d)},$$

where A is a differential operator of order m or a Fourier multiplier with similar homogeneity properties, $p = d/(d - m)$, and f may satisfy additional constraints (such as $\operatorname{div} f = 0$). Usually, these inequalities are related to vectorial behavior of the function f or the operators in question (Af is usually a vector field or a differential form). We refer the reader to the papers [5, 7, 8, 11, 14, 17, 19, 20, 23, 24, 31, 32, 35, 39, 41, 42, 47–51], among many others, as well as the surveys [52] and [38] for many Bourgain–Brezis inequalities.

There is a classical trace inequality

$$(1.2) \quad \|f\|_{L_1(\mu)} \lesssim \left(\sup_{\substack{x \in \mathbb{R}^d \\ r > 0}} r^{1-d} \mu(B_r(x)) \right) \|\nabla f\|_{L_1(\mathbb{R}^d)}, \quad f \in C_0^\infty(\mathbb{R}^d),$$

proved by Maz'ya in [22] and Meyers and Ziemer in [26] (see [25] for generalizations, and Chapter 7 in [1] for more classical L_p -theory of trace inequalities with $p > 1$), which, seemingly does not have any relatives among Bourgain–Brezis inequalities. We call this inequality the trace inequality, because it allows to define traces of W_1^1 and even BV functions on sets of non-zero $(d - 1)$ -dimensional Hausdorff measure (we plug for μ the natural Frostman measure related to the set, see Chapter 8 in [21]). It is desirable to obtain similar inequalities in the generality of (1.1); first, because it is an $L_1 \rightarrow L_1$ inequality, which seems to be more difficult than the usual $L_1 \rightarrow L_p$ Bourgain–Brezis inequalities; second, such type inequalities often deliver information about geometric properties of corresponding BV-type measures. The depth of the desirable analog of (1.2) is emphasized by the fact that defining the trace on a hyperplane, the simplest $(d - 1)$ -dimensional set, in the generality of (1.1) was already a demanding problem. We refer the reader to [9, 12, 15, 16] for trace inequalities on hyperplanes for differential operators and L_1 -norms.

In this paper, we do not aim to say anything new about functions on \mathbb{R}^d . We wish to explore the aforementioned class of inequalities in the related martingale setting introduced in [4] (the discrete model might be traced to [18]). While there is a well-known analogy between singular integrals on \mathbb{R}^d and martingale transforms, the existence of an effect similar to that of Bourgain–Brezis inequalities in the martingale setting was noticed only in [4]. The consideration of these related discrete problems allows to guess the right way to prove statements in the Euclidean case. See [42] for the proof of several endpoint Bourgain–Brezis inequalities with the approach suggested by [4], [40] for a probabilistic model of weakly cancelling operators, and the implementation of the reasoning of that paper on the Euclidean setting in [41]. See [44] for Maz'ya's Φ -inequalities in the martingale setting, and [45] for the transference of that reasoning to the Euclidean setting.

In the forthcoming section, we introduce the martingale model. Section 3 contains the description of our methods. In [4], the authors relied upon common combinatorial tricks for proving martingale inequalities (mostly stopping times), here we will need to use a finer tool called the Bellman function or the Burkholder method. We refer the reader to the foundational papers [10, 28] and the books [30, 53] for the basics of the Bellman function method. Though the present paper is self-contained, it might be instructive to consult [44], where a simpler Bellman function was used to solve a related problem. For smoothness of exposition, we first prove the non-limiting L_1 -trace inequality from [4] using Bellman function in Section 4. This is already interesting, because the Bellman extremal problem arising here seems new. Section 5 introduces additional requirements on the spaces and functionals and states the main result, Theorem 5.11. The assumptions we impose on the martingale Sobolev space (see Definition 5.1) might be thought of as martingale versions of the requirements that the k -wave cone is empty introduced in [2] (see [43] as well). This is a simple condition on the operator A as in (1.1) sufficient to conclude that any vectorial measure $\mu = Af$ has the lower Hausdorff dimension at least k ; note that in some cases, this condition might be essentially sharpened, see [3]. Section 6 contains the proof of the main theorem, which has been reduced to a verification of a finite dimensional inequality by the Bellman function method in Section 4; the said finite dimensional inequalities are still non-trivial. The final section is devoted to two examples that emphasize the analogy between the martingale problems and similar questions in the Euclidean setting. We consider two particular cases of martingale Sobolev spaces that resemble the spaces W_1^1 and

the space of divergence-free measures, and apply our investigations to these two cases; the results are summarized in Theorems 7.17 and 7.18. We end the paper with formulating a conjecture about functions on the Euclidean space.

2. Sobolev martingales

Consider a probability space. If S is a finite algebra of measurable sets, a non-empty set $a \in S$ is called an atom provided there are no non-empty sets $b \in S$ such that $b \subset a$. Suppose that $m \geq 2$ is a fixed natural number and $\mathcal{F} = \{\mathcal{F}_n\}_{n \geq 0}$ is an m -uniform filtration (i.e., an increasing sequence of set algebras, $\mathcal{F}_n \subset \mathcal{F}_{n+1}$, such that each atom in \mathcal{F}_n is split into m atoms in \mathcal{F}_{n+1} having equal masses; we also assume that \mathcal{F}_0 is the trivial algebra) on the standard atomless probability space. The set of atoms of the algebra \mathcal{F}_n is denoted by \mathcal{AF}_n . There is a natural tree structure on the set of all atoms $\bigcup_n \mathcal{AF}_n$: the atom $\omega' \in \mathcal{AF}_{n+1}$ is a kid of $\omega \in \mathcal{AF}_n$ if $\omega' \subset \omega$. In such a case, ω is the parent of ω' , and we denote the parent of ω' by $(\omega')^\uparrow$. For each atom $a \in \mathcal{AF}_n$, we enumerate its kids with the numbers $1, 2, \dots, m$ and fix this enumeration. We will be using the notation $[1 \dots m]$ to denote the integer interval with the endpoints 1 and m , including the endpoints.

We may treat our probability space equipped with an m -uniform filtration as a metric space \mathbb{T} . The points of \mathbb{T} are infinite paths in the tree of atoms (we start from the whole space, then choose an atom in \mathcal{AF}_1 , then pass to one of its kids in \mathcal{AF}_2 , and so on). More formally, a path is a mapping $\gamma: \mathbb{N} \cup \{0\} \rightarrow [1 \dots m]$. A point of \mathbb{T} that corresponds to such a mapping γ may be thought of as the intersection of the atoms $\omega_n \in \mathcal{AF}_n$, $n = 0, 1, 2, \dots$, where ω_{n+1} is the $\gamma(n)$ -th kid of ω_n . This allows to interpret atoms as subsets of \mathbb{T} . The distance between the two paths γ_1 and γ_2 is defined by the formula

$$\text{dist}(\gamma_1, \gamma_2) = m^{-d}, \quad d = \max\{n \in \mathbb{N} \mid \gamma_1(j) = \gamma_2(j) \text{ for all } j < n\}.$$

With this metric, \mathbb{T} becomes a compact metric space. Another way to interpret \mathbb{T} is to identify a path $\gamma = (\gamma(0), \gamma(1), \gamma(2), \dots)$ with the formal series $\sum_{n=0}^{\infty} (\gamma(n) - 1)m^{-n-1}$, which induces the surjection $\mathbb{T} \rightarrow [0, 1]$. This suggests to call $\gamma(n)$ the n -th digit of γ . In this interpretation, the atoms become m -adic subintervals of $[0, 1]$. One may prove that \mathbb{T} is homeomorphic to a Cantor-type set.

Recall that a sequence of summable random variables $F = \{F_n\}_n$ is called a martingale adapted to \mathcal{F} , provided, first, each F_n is \mathcal{F}_n -measurable, and second, we have $F_n = \mathbb{E}(F_{n+1} \mid \mathcal{F}_n)$ for any $n \geq 0$. For a martingale F adapted to \mathcal{F} , let $\{dF_n\}_{n \geq 1}$ be the sequence of its martingale differences:

$$dF_{n+1} = F_{n+1} - F_n, \quad n \geq 0.$$

We refer the reader to Chapter VII in [36] for the general martingale theory. We will be working with a very narrow class of martingales adapted to uniform filtrations, and will not use anything other than terminology from the general theory.

Consider the linear space

$$V = \left\{ v \in \mathbb{R}^m \mid \sum_1^m v_j = 0 \right\}.$$

For each atom $\omega \in \mathcal{AF}_n$, the function dF_{n+1} attains at most m values on ω , and, thus, might be identified with an element of V . Namely, for any ω , we fix a bijective map

$$J_\omega: [1 \dots m] \rightarrow \{\omega' \in \mathcal{AF}_{n+1} \mid \omega = (\omega')^\uparrow\}$$

that maps the number j to the j -th kid of ω . This map may be extended linearly to the mapping between V and the space of restrictions of all possible martingale differences dF_{n+1} to ω . This extended map is also called J_ω .

Let us observe that each signed Borel measure μ with bounded variation on \mathbb{T} generates a martingale via the formula

$$(2.1) \quad F_n = m^n \sum_{\omega \in \mathcal{AF}_n} \mu(\omega) \chi_\omega, \quad n \in \mathbb{N} \cup \{0\}.$$

Note that the characteristic functions of atoms are continuous with respect to the metric d and form a total family in the space $C(\mathbb{T})$ (i.e., each measure on \mathbb{T} is uniquely defined by its values at the atoms). Using this fact, one may establish the one-to-one correspondence via (2.1) between finite signed measures on \mathbb{T} and L_1 -martingales adapted to \mathcal{F} . By an L_1 martingale we mean a martingale F such that $\sup_n \|F_n\|_{L_1}$ is finite, the latter expression being the L_1 norm of F .

The Hardy–Littlewood–Sobolev inequality for martingales (going back to [54]) reads as follows:

$$(2.2) \quad \left\| \sum_{n \geq 0} m^{-\alpha n} dF_n \right\|_{L_q} \lesssim \|F\|_{L_p},$$

where $1/p - 1/q = \alpha$ and $1 < p < q < \infty$. Here and in what follows, the notation $A \lesssim B$ means $A \leq CB$, where the constant C is uniform in a certain sense. For example, the constant in (2.2) should not depend on the particular choice of F , but it might depend on p and q . The parameter α should be interpreted as the order of the integration operator. See [27] and [46] for more general versions of the Hardy–Littlewood–Sobolev inequality for martingales.

Remark 2.1. Note that the inequality (2.2) is true when $p = 1$ and $1 - 1/q < \alpha$ since

$$\|dF_n\|_{L_q} \leq m^{(q-1)(n+1)/q} \|dF_n\|_{L_1} \leq 2m^{(q-1)(n+1)/q} \|F\|_{L_1}.$$

If the martingale F is \mathbb{R}^ℓ -valued, then $f_{n+1}|_\omega$ might be naturally identified with an element of $V \otimes \mathbb{R}^\ell$ (we apply J_ω to each of ℓ coordinates individually); here $\omega \in \mathcal{AF}_n$. In other words, we may think of the trace of f_{n+1} on $\omega \in \mathcal{AF}_n$ as of an $m \times \ell$ matrix such that the sum of the elements in each row equals zero. Let W be a linear subspace of $V \otimes \mathbb{R}^\ell$. Consider the subspace \mathbb{W} of the \mathbb{R}^ℓ -valued martingale space L_1 :

$$\mathbb{W} = \{F \text{ is an } L_1 \text{ martingale with values in } \mathbb{R}^\ell \mid \forall n, \omega \in \mathcal{AF}_n, dF_{n+1}|_\omega \in J_\omega[W]\}.$$

In view of (2.1), one should think of \mathbb{W} as of a BV-type space. The main common feature of \mathbb{W} and spaces of constrained vectorial measures such as $\nabla \text{BV} = \{\nabla f \mid f \in \text{BV}(\mathbb{R}^d)\}$ is that both spaces are invariant under certain group actions. The classical space ∇BV and

its generalizations are translation and dilation invariant, while \mathbb{W} is invariant with respect to shifts of the tree \mathbb{T} and, if $[1 \dots m]$ is equipped with a group structure and W is invariant, invariant with respect to other automorphisms of \mathbb{T} . The spaces \mathbb{W} were implicitly introduced in [18] and proved to be good discrete models for several problems in harmonic analysis on L_1 and the space of measures, see [4] and [40]; the former paper suggested the name ‘Sobolev martingales’ since the behavior of the spaces \mathbb{W} resembles that of \dot{W}_1^1 (it is more appropriate to compare \mathbb{W} with BV, since L_1 -martingales correspond to measures by (2.1); seemingly, this was overlooked by the authors of [4]).

3. Setting of the trace problem and the Bellman function

Let $\varphi: W \rightarrow V$ be a linear operator and let $\alpha \in (0, 1)$. Consider the operator

$$(3.1) \quad \mathbb{I}_\alpha[F] = \sum_n m^{-\alpha n} \sum_{\omega \in \mathcal{AF}_n} J_\omega [\varphi(J_\omega^{-1}[dF_{n+1}|\omega])], \quad F \in \mathbb{W}.$$

This operator maps martingales to functions; by the correspondence (2.1), we may interpret it as an operator that transforms measures into functions (the series converges in L_q with $q < 1/(1 - \alpha)$ by Remark 2.1). The operator J_ω is used to make the formula mathematically rigorous; informally, (3.1) may be written as

$$\mathbb{I}_\alpha[F] = \sum_n m^{-\alpha n} \sum_{\omega \in \mathcal{AF}_n} \varphi(dF_{n+1}|\omega), \quad F \in \mathbb{W}.$$

We note that the operator \mathbb{I}_α cannot be applied to an arbitrary \mathbb{R}^ℓ -valued martingale since φ is defined on W only and the condition $dF_{n+1}|\omega \in W$ is required. We will also use the notation

$$\left(\mathbb{I}_\alpha[F]\right)_N = \sum_{n=0}^{N-1} m^{-\alpha n} \sum_{\omega \in \mathcal{AF}_n} J_\omega [\varphi(J_\omega^{-1}[dF_{n+1}|\omega])], \quad F \in \mathbb{W}.$$

Note that $(\mathbb{I}_\alpha[F])_N$ is \mathcal{F}_N -measurable. Therefore, $\{(\mathbb{I}_\alpha[F])_n\}_n$ is a martingale.

Remark 3.1. Set $\ell = 1$. The choice of the identity operator for φ reduces \mathbb{I}_α to the martingale Riesz potential present in the Hardy–Littlewood–Sobolev inequality (2.2). This operator has positive kernel if we represent it as an integral operator. A different choice of the operator φ might make the operator \mathbb{I}_α more cancelling than the Riesz potential (usually, larger ℓ are needed for such a choice). This is similar to the difference between the classical Euclidean Riesz potential $|x|^{-\beta}$ and the kernel $x/|x|^{\beta+1}$. The most classical example arises in the regime $\beta = d$ (the dimension of the Euclidean space): the former operator does not act on L_1 at all, while the latter operator is a Calderón–Zygmund operator and maps L_1 to $L_{1,\infty}$. As we will see, these additional cancellations become important at the endpoints for the trace inequalities we consider (a similar effect appeared in [33]).

We will be studying the question: ‘for which measures ν on \mathbb{T} does the inequality

$$(3.2) \quad \|\mathbb{I}_\alpha[F]\|_{L_1(\nu)} \lesssim \|F\|_{L_1}, \quad F \in \mathbb{W},$$

hold true?’ This inequality allows to define L_1 -traces of $\mathbb{I}_\alpha[F]$ on lower-dimensional subsets of \mathbb{T} (this explains the name ‘trace inequality’). We will be working with simple martingales only, since usually the result may be extended to the class of all L_1 -martingales by a routine limiting argument.

Definition 3.2. Let $\alpha \in (0, 1)$ and let $p \in [1, \infty)$. We say that a measure ν with bounded variation on \mathbb{T} satisfies the (α, p) -Frostman condition provided

$$(3.3) \quad \forall n \geq 0 \quad \forall \omega \in \mathcal{AF}_n \quad (\nu(\omega))^{1/p} \lesssim m^{(\alpha-1)n}.$$

One may restate the (α, p) -Frostman condition in terms of Morrey space norms, see [33]. Sometimes, the (α, p) -Frostman condition is called the ball growth condition.

Remark 3.3. One may prove that the $(\alpha, 1)$ -Frostman condition is necessary for (3.2), provided φ is not degenerate. Namely, assume that for any $j \in [1 \dots m]$, there exists $v \in W$ such that the j -th coordinate of $\varphi[v]$, as an element of \mathbb{R}^m , is non-zero (we will later denote this number by $(\varphi[v])_j$, $j = 1, 2, \dots, m$). Then, the $(\alpha, 1)$ -Frostman condition is necessary for (3.2). One may prove this by plugging ‘elementary martingales’ into (3.2), i.e., martingales F for which $dF_{n+1}|_\omega$ is non-zero for only single n and single $\omega \in \mathcal{AF}_n$.

We will be working with a slightly more demanding ‘bilinear’ form of (3.2):

$$(3.4) \quad \|\mathbb{I}_\alpha[F]\|_{L_1(\nu)} \lesssim \left(\sup_{\substack{n \geq 0 \\ \omega \in \mathcal{AF}_n}} m^{(1-\alpha)n} \nu(\omega) \right) \|F\|_{L_1}, \quad F \in \mathbb{W}.$$

Definition 3.4. Define the Bellman function $\mathbb{B}: \mathbb{R}^\ell \times \mathbb{R} \times \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$ by the rule

$$(3.5) \quad \mathbb{B}(x, y, z, t, s) = \sup \left(\left\{ \int_{\mathbb{T}} |y + \mathbb{I}_\alpha[F]| d\nu \mid F_0 = x, \mathbb{E} |F_\infty| = z, \nu(\mathbb{T}) = t, \right. \right. \\ \left. \left. \sup_{\substack{n \geq 0 \\ \omega \in \mathcal{AF}_n}} m^{(1-\alpha)n} \nu(\omega) = s, F \in \mathbb{W} \right\} \right).$$

The supremum in the formula above is taken over the set of simple martingales F only; note that in such a case, we may assume that ν is generated (via (2.1)) by a simple martingale as well.

The Bellman domain (i.e., the set of points (x, y, z, t, s) for which the supremum in (3.5) is taken over a non-empty set) is

$$(3.6) \quad \Omega = \{(x, y, z, t, s) \in \mathbb{R}^\ell \times \mathbb{R} \times \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \mid |x| \leq z, t \leq s\}.$$

The Bellman function satisfies the boundary inequality (this follows from substitution of a constant martingale into (3.5))

$$(3.7) \quad \mathbb{B}(x, y, |x|, t, s) \geq |y|t.$$

We will use the symbol $\sum_{j=1}^m$ to denote the average, i.e.,

$$\sum_{j=1}^m a_j = \frac{1}{m} \sum_{j=1}^m a_j.$$

Proposition 3.5. *The Bellman function satisfies the main inequality*

$$(3.8) \quad \mathbb{B}(x, y, z, t, s) \geq m^{-\alpha} \sum_{j=1}^m \mathbb{B}(x_j, m^\alpha y + (\varphi[\vec{x}])_j, z_j, t_j, s_j),$$

where

$$(3.9) \quad x = \sum_{j=1}^m x_j, \quad z = \sum_{j=1}^m z_j, \quad t = \sum_{j=1}^m t_j, \\ \vec{x} = (x_1 - x, x_2 - x, \dots, x_m - x) \in W \quad \text{and} \quad s = \max(t, m^{1-\alpha} \max_{j=1, \dots, m} s_j).$$

Proposition 3.5 is similar to Lemma 3.4 in [44], so we omit its proof. Formally, we will not need it. A reverse statement is more important for us. We need to set up notation before formulating it.

Definition 3.6. The configuration space \mathfrak{W} is formed by strings $(\mathfrak{X}, y, \mathfrak{Z}, \mathfrak{T}, \mathfrak{S})$, where

$$\mathfrak{X} = \{x_j\}_{j=1}^m \in \mathbb{R}^{\ell m}, \quad y \in \mathbb{R}, \quad \mathfrak{Z} = \{z_j\}_{j=1}^m \in (\mathbb{R}_+)^m, \\ \mathfrak{T} = \{t_j\}_{j=1}^m \in (\mathbb{R}_+)^m, \quad \text{and} \quad \mathfrak{S} = \{s_j\}_{j=1}^m \in (\mathbb{R}_+)^m,$$

satisfy the splitting rules (3.9), i.e., \vec{x} constructed by the formulas listed in (3.9) belongs to W , and the natural domain requirements $|x_j| \leq z_j$ and $t_j \leq s_j$ for all $j = 1, 2, \dots, m$ are also fulfilled.

A sequence $\{H_n\}_n$ of summable random variables is called a supermartingale provided each H_n is \mathcal{F}_n -measurable (i.e., $\{H_n\}_n$ is adapted to \mathcal{F}) and for any $n \geq 0$,

$$\mathbb{E}(H_{n+1} | \mathcal{F}_n) \leq H_n.$$

If $\{H_n\}_n$ is a supermartingale, then, for any N ,

$$(3.10) \quad \mathbb{E} H_N \leq H_0.$$

Proposition 3.7. *Let $G: \Omega \rightarrow \mathbb{R}$ be a function that obeys the main inequality (3.8), i.e., such that*

$$(3.11) \quad G(x, y, z, t, s) \geq m^{-\alpha} \sum_{j=1}^m G(x_j, m^\alpha y + (\varphi[\vec{x}])_j, z_j, t_j, s_j)$$

for all $(\mathfrak{X}, y, \mathfrak{Z}, \mathfrak{T}, \mathfrak{S}) \in \mathfrak{W}$ and x, z, t, s generated from them by (3.9). Let v_n be the martingale generated by v via (2.1), and let

$$M_n(\omega) = \sup_{\substack{N \geq n, v \subset \omega, \\ v \in \mathcal{A}_{\mathcal{F}_N}}} m^{(1-\alpha)N} v(v).$$

Then, the process

$$(3.12) \quad m^{(1-\alpha)n} G(F_n, m^{\alpha n} (y_0 + (\mathbb{I}_\alpha[F])_n), \mathbb{E}(|F_\infty| | \mathcal{F}_n), m^{-n} v_n, m^{(\alpha-1)n} M_n)$$

is a supermartingale provided $F \in \mathbb{W}$ and $y_0 \in \mathbb{R}$.

Proposition 3.7 and Corollary 3.8 below are similar to Lemma 3.9 in [44].

Proof of Proposition 3.7. Let $w \in \mathcal{AF}_n$ and let w_1, w_2, \dots, w_m be its kids in \mathcal{AF}_{n+1} . Let

$$\begin{aligned} y &= m^{\alpha n} (y_0 + (\mathbb{I}_\alpha[F])_n(w)); & x_j &= F_{n+1}(w_j), & z_j &= \mathbb{E}(|F_\infty| \mid \mathcal{F}_{n+1})(w_j), \\ t_j &= v(w_j) = m^{-n-1} v_{n+1}(w_j), & s_j &= m^{(\alpha-1)(n+1)} M_{n+1}(w_j), & j &\in [1 \dots m]. \end{aligned}$$

By the martingale properties,

$$\begin{aligned} x &= F_n(w), & z &= \mathbb{E}(|F_\infty| \mid F_n)(w), & t &= v(w), & s &= m^{(\alpha-1)n} M_n(w), \\ \text{and } \bar{x} &= J_w^{-1}[dF_{n+1}|w], \end{aligned}$$

satisfy the splitting rules (3.9). In other words, the string $(\mathfrak{X}, y, \mathfrak{Z}, \mathfrak{T}, \mathfrak{S})$ formed by these numbers generated as above, belongs to the configuration space \mathfrak{W} . Therefore, the main inequality (3.11) applies, which means exactly the supermartingale property

$$\begin{aligned} &m^{(1-\alpha)n} G(F_n(w), m^{\alpha n} (y_0 + (\mathbb{I}_\alpha[F])_n(w)), \mathbb{E}(|F_\infty| \mid \mathcal{F}_n)(w), m^{-n} v_n(w), m^{(\alpha-1)n} M_n(w)) \\ &\geq m^{(1-\alpha)(n+1)} \mathbb{E} G(F_{n+1}, m^{\alpha(n+1)} (y_0 + (\mathbb{I}_\alpha[F])_{n+1}), \mathbb{E}(|F_\infty| \mid \mathcal{F}_{n+1}), m^{-n-1} v_{n+1}, \\ &\quad m^{(\alpha-1)(n+1)} M_{n+1}) \chi_w \end{aligned}$$

for the process (3.12); we have used the identity

$$m^{\alpha(n+1)} (y_0 + (\mathbb{I}_\alpha[F])_{n+1}(w_j)) = m^\alpha y + (\varphi[\bar{x}])_j \quad \text{on } w_j. \quad \blacksquare$$

Corollary 3.8. *Suppose there exists a finite function $G: \Omega \rightarrow \mathbb{R}$ that satisfies the main inequality (3.11), the boundary inequality $G(|x|, y, |x|, s, t) \geq |y|t$, and the estimate*

$$(3.13) \quad G(x, 0, z, t, s) \lesssim zs.$$

Then, the inequality (3.4) holds true for all measures ν that satisfy the $(\alpha, 1)$ -Frostman condition.

Proof. By Definition 3.4, it suffices to prove the estimate

$$(3.14) \quad \mathbb{B}(x, 0, z, t, s) \lesssim zs.$$

In fact, we will prove that $\mathbb{B} \leq G$ on Ω . Let $(x, y, z, t, s) \in \Omega$ and let (F, ν) be a pair of a simple martingale and a measure generated by a simple martingale that fulfills the requirements in (3.5) for these fixed x, z, t , and s . By Proposition 3.7, the corresponding process (3.12) is a supermartingale. Therefore,

$$\begin{aligned} G(x, y, z, t, s) &= G(F_0, y + (\mathbb{I}_\alpha[F])_0, \mathbb{E} |F_\infty|, v_0, M_0) \\ &\stackrel{(3.10)}{\geq} m^{(1-\alpha)N} \mathbb{E} G(F_N, m^{\alpha N} (y + (\mathbb{I}_\alpha[F])_N), \mathbb{E}(|F_\infty| \mid \mathcal{F}_N), m^{-N} v_N, m^{(\alpha-1)N} M_N), \end{aligned}$$

for any $N > 0$. Let us pick sufficiently large N (recall we are working with simple martingales). By the boundary inequality, the latter expression is bounded from below by

$$m^{(1-\alpha)N} \mathbb{E} m^{\alpha N} |y + (\mathbb{I}_\alpha[F])_N| m^{-N} v_N = \mathbb{E} |y + (\mathbb{I}_\alpha[F])_N| v_N = \int_{\mathbb{T}} |y + \mathbb{I}_\alpha[F]| d\nu.$$

Since F and v are arbitrary, we get

$$\mathbb{B}(x, y, z, t, s) \leq G(x, y, z, t, s),$$

which yields (3.14). ■

Remark 3.9. It is unclear whether the estimate (3.13) in the above corollary is necessary. Maybe, it is possible to derive (3.13) from the finiteness of G and the validity of the main inequality for it.

Definition 3.10. The functions G that satisfy the main inequality and the boundary inequality are usually called *supersolutions*.

4. Subcritical case

Let $v \in V$ be a vector. Consider the function $\kappa_v : (0, 1] \rightarrow \mathbb{R}$ given by the formula

$$\kappa_v(\theta) = \theta \log \left(\sum_{j=1}^m |1 + v_j|^{1/\theta} \right) = \log \|\mathbf{1} + v\|_{L_{1/\theta}}.$$

We may extend this function to 0 by continuity. Then,

$$(4.1) \quad \kappa_v(0) = \log \left(\max_{j=1, \dots, m} |1 + v_j| \right).$$

By Hölder's inequality, this function is convex and non-increasing. If, in addition, $v_j \geq -1$ for any j , then this function also satisfies $\kappa_v(1) = 0$. Consider yet another function $\kappa : [0, 1] \rightarrow \mathbb{R}$:

$$(4.2) \quad \kappa(\theta) = \sup \{ \kappa_v(\theta) \mid \exists a \in \mathbb{R}^\ell \setminus \{0\} \text{ such that } v \otimes a \in W \text{ and } \forall j, v_j \geq -1 \}.$$

The function κ is also convex, non-increasing, and satisfies $\kappa(1) = 0$. As an elementary computation shows,

$$(4.3) \quad \kappa'(1) = \inf \left\{ - \sum_{j=1}^m (1 + v_j) \log(1 + v_j) \mid \exists a \in \mathbb{R}^\ell \setminus \{0\} \text{ such that } v \otimes a \in W \text{ and } \forall j, v_j \geq -1 \right\}.$$

Heuristically, the function κ shows how rapidly can the L_p -norm of a rank-one martingale in \mathbb{W} grow. Theorem 1.11 in [4] shows that this function is linked to concentration of measures $\mu \in \mathbb{W}$: the lower Hausdorff dimension of $\mu \in \mathbb{W}$ is at least $1 + \kappa'(1)/\log m$ and this bound is sharp for every \mathbb{W} . We cite Theorem 6.2 from [4] (there was no operator φ in [4]; the theorem below is not sensitive to φ).

Theorem 4.1. *Assume v satisfies the $(\alpha, 1)$ -Frostman condition and*

$$(4.4) \quad \alpha > - \frac{\kappa'(1)}{\log m}.$$

Then, (3.4) holds true with any operator φ .

Let us give the Bellman function proof of this theorem (it relies upon the same principles as the original one in [4], but uses a different language).

Theorem 4.2. *Assume (4.4). Then, for any $\theta \in (0, 1)$ sufficiently close to one, the function*

$$(4.5) \quad G(x, y, z, t, s) = |y|t + M_1|x|t^{1-\theta}s^\theta + M_2(z - |x|)s$$

is a supersolution, provided $1 \ll M_1 \ll M_2$ (depending on θ).

The notation \ll means the statement holds true when M_1 is sufficiently large and M_2 is also sufficiently large (depending on the particular choice of M_1). Note that Theorem 4.2 implies Theorem 4.1 via Corollary 3.8.

Definition 4.3. If $G: \Omega \rightarrow \mathbb{R}$ is a function, then its discrepancy $\text{DISCR}[G]: \mathfrak{W} \rightarrow \mathbb{R}$ is

$$\begin{aligned} \text{DISCR}[G](\mathfrak{X}, y, \mathfrak{Z}, \mathfrak{T}, \mathfrak{S}) &= G(x, y, z, t, s) \\ &\quad - m^{-\alpha} \sum_{j=1}^m G(x_j, m^\alpha y + (\varphi[\vec{x}])_j, z_j, t_j, s_j). \end{aligned}$$

The main inequality says that $\text{DISCR}[G] \geq 0$ on \mathfrak{W} .

Proof of Theorem 4.2. We wish to prove

$$\text{DISCR}[|y|t + M_1|x|t^{1-\theta}s^\theta + M_2(z - |x|)s] \geq 0.$$

Let us first compute $\text{DISCR}[(z - |x|)s]$:

$$\begin{aligned} \text{DISCR}[(z - |x|)s] &= (z - |x|)s - m^{-\alpha} \sum_{j=1}^m (z_j - |x_j|)s_j \\ &= \left(\sum_{j=1}^m z_j - \sum_{j=1}^m |x_j| \right) s + \left(\sum_{j=1}^m |x_j| - |x| \right) s - m^{-\alpha} \sum_{j=1}^m (z_j - |x_j|)s_j \\ (4.6) \quad &= \left(\sum_{j=1}^m |x_j| - |x| \right) s + \sum_{j=1}^m (z_j - |x_j|)(m^{-1}s - m^{-\alpha}s_j). \end{aligned}$$

We see that this expression is always non-negative due to the splitting rules (3.9). What is more,

$$(4.7) \quad \text{DISCR}[(z - |x|)s] \gtrsim |\mathfrak{X}|s,$$

provided

$$(4.8) \quad \sum_{j=1}^m |x_j| \geq (1 + \varepsilon)|x|$$

for some fixed small ε (the notation $|\mathfrak{X}|$ means the Euclidean norm of $\mathfrak{X} = (x_1, x_2, \dots, x_m) \in \mathbb{R}^{\ell m}$). Now we turn to the function $|x|t^{1-\theta}s^\theta$ and estimate its discrepancy. Let p be such

that $(1 - \theta)p' = 1$, where $1/p + 1/p' = 1$. Then,

$$\begin{aligned}
 (4.9) \quad m^{-\alpha} \sum_{j=1}^m |x_j| t_j^{1-\theta} s_j^\theta &= m^{1-\alpha} \sum_{j=1}^m |x_j| t_j^{1-\theta} s_j^\theta \\
 &\leq m^{1-\alpha} \left(\sum_{j=1}^m |x_j|^p \right)^{1/p} \left(\sum_{j=1}^m t_j^{(1-\theta)p'} s_j^{\theta p'} \right)^{1/p'} \\
 &\stackrel{(1-\theta)p'=1}{\leq} m^{1-\alpha} \left(\sum_{j=1}^m |x_j|^p \right)^{1/p} \left(\frac{t}{m} (m^{(\alpha-1)s})^{\theta p'} \right)^{1/p'} \\
 &= m^{(1-\alpha)(1-\theta)-(1-\theta)} \left(\sum_{j=1}^m |x_j|^p \right)^{1/p} t^{1-\theta} s^\theta.
 \end{aligned}$$

Thus, if

$$(4.10) \quad m^{-\alpha(1-\theta)} \left(\sum_{j=1}^m |x_j|^p \right)^{1/p} \leq |x|,$$

then $\text{DISCR}[|x|t^{1-\theta}s^\theta] \geq 0$, i.e., the discrepancy of the second term in (4.5) is non-negative. It was proved in Lemma 2.1 of [4] that, given $p \in (1, \infty]$, for any $\tilde{\delta} > 0$ there exists $\varepsilon > 0$ such that

$$e^{-\kappa(p-1)} \left(\sum_{j=1}^m |x_j|^p \right)^{1/p} \leq (1 + \tilde{\delta})|x|, \quad \vec{x} \in W,$$

provided (4.8) is violated. Therefore, if $\alpha > -\kappa'(1)/\log m$, p is sufficiently close to one and ε is sufficiently small, then there exists a tiny δ such that

$$m^{-\alpha p/(p-1)} \left(\sum_{j=1}^m |x_j|^p \right)^{1/p} \leq (1 - \delta)|x|$$

whenever $\vec{x} \in W$ and $\sum_{j=1}^m |x_j| \leq (1 + \varepsilon)|x|$, which is slightly stronger than (4.10). Therefore,

$$(4.11) \quad \text{DISCR}[|x|t^{1-\theta}s^\theta] \gtrsim |\mathfrak{X}|t^{1-\theta}s^\theta,$$

provided (4.8) is violated with sufficiently small ε . We fix such an ε .

Combining (4.7), (4.11), and the simple estimate

$$(4.12) \quad |\text{DISCR}[|x|t^{1-\theta}s^\theta]| \lesssim |\mathfrak{X}|t^{1-\theta}s^\theta,$$

we obtain

$$(4.13) \quad \text{DISCR}[|x|t^{1-\theta}s^\theta + M(z - |x|)s] \gtrsim |\mathfrak{X}|t^{1-\theta}s^\theta,$$

provided M is sufficiently large (note that $t^{1-\theta}s^\theta \leq s$). Indeed, in the case (4.8) holds true, (4.13) follows from (4.7) and (4.12) by the choice of sufficiently large M ; in the case (4.8) is violated, we rely upon the positivity of $\text{DISCR}[(z - |x|)s]$ and (4.11)¹.

¹We will refer to this simple reasoning as to the flat/convex argument (see the explanation after the proof of the lemma).

Thus, it remains to prove

$$(4.14) \quad |\text{DISCR}[|y|t]| \lesssim |\mathfrak{X}|t.$$

This follows from direct computation,

$$(4.15) \quad \text{DISCR}[|y|t] = \sum_{j=1}^m (|y| - |y + m^{-\alpha}(\varphi[\vec{x}])_j|) t_j,$$

and the triangle inequality. ■

Remark 4.4. Lemma 6.2 in [4] says that if the function κ is affine and (3.2) holds true with any φ , then (4.4) is also true.

In [4], the condition (4.8) played the pivotal role (see Definition 3.1 in that paper). The atoms for which the corresponding quantities violated it were called ε -flat². The core of the reasoning is contained in Lemma 2.1 of [4], which manifests the principle that flat atoms (or flat configurations $\mathfrak{X} = (x_1, x_2, \dots, x_m)$ such that $\vec{x} \in W$) lie close to rank-one configurations. It is convenient to introduce the set of admissible 0-flat increments:

$$(4.16) \quad \mathbb{M}^W = \{v \in V \mid \forall j \in [1 \dots m], v_j \geq -1, \text{ and } \exists a \in \mathbb{R}^\ell \setminus \{0\} \text{ such that } v \otimes a \in W\}.$$

See [42] for ‘translation’ of these notions into the language of the Euclidean space (instead of \mathbb{T}). The following lemma justifies the heuristic principle that flat configurations are close to rank-one configurations.

Lemma 4.5. *Fix W . Let U be a neighborhood of the set*

$$(4.17) \quad \{\mathfrak{X} \in \mathbb{R}^m \otimes \mathbb{R}^\ell \mid \exists v \in \mathbb{M}^W \text{ and } a \in \mathbb{R}^\ell \setminus \{0\} \text{ such that } \vec{x} = v \otimes a, x = a\}.$$

Then, for sufficiently small ε , the violation of (4.8), together with the conditions $|x| = 1$ and $\vec{x} \in W$, imply $\mathfrak{X} \in U$.

Proof. Note that if $\mathfrak{X} = \{x_j\}_{j=1}^m$ violates (4.8) and $|x| = 1$, then $|x_j| \leq m + 1$ for every $j \in [1 \dots m]$ (we assume ε is sufficiently small). Therefore, the configurations \mathfrak{X} we consider are uniformly bounded as vectors in $\mathbb{R}^{m\ell}$. Assume the contrary: for any $n \in \mathbb{N}$, there exists $\mathfrak{X}^n = (x_1^n, x_2^n, \dots, x_m^n) \in \mathbb{R}^{m\ell}$ that does not belong to U , $|x^n| = 1$, $\vec{x}^n \in W$, and

$$\sum_{j=1}^m |x_j^n| \leq 1 + \frac{1}{n}.$$

Let \mathfrak{X} be a limit point of the sequence $\{\mathfrak{X}^n\}_n$. Then, $\sum_{j=1}^m |x_j| = 1$, which means that the triangle inequality $\sum_{j=1}^m |x_j| \geq 1$ turns into equality and there exists $v \in V$ with $v_j \geq -1$, $j \in [1 \dots m]$, such that $\vec{x} = v \otimes x$. Since $\vec{x} \in W$, we have $v \in \mathbb{M}^W$. Therefore, \mathfrak{X} belongs to (4.17), which contradicts the openness of U . ■

²The term ‘flat’ symbolizes that the martingale develops along the flat part of the boundary of the ball of L_1 .

5. Additional requirements on φ and W

Definition 5.1. We call the space $W \subset V \otimes \mathbb{R}^\ell$ *geometric* if the corresponding function κ is affine.

The condition that a space W is geometric is similar to the condition that the k -wave cone is empty (see also Lemma 7.4 below); the latter condition played an important role in [2] and [43].

Proposition 5.2. *The space W is geometric if there exists α such that*

- *the inequality*

$$(5.1) \quad \sum_{j=1}^m (1 + v_j) \log(1 + v_j) \leq \alpha \log m$$

holds true for any vector $v \in \mathbb{M}^W$;

- *there exists $v \in \mathbb{M}^W$ and $H \subset [1 \dots m]$ such that $|H| = m^{1-\alpha}$ and*

$$v_j = \begin{cases} m^\alpha - 1, & j \in H, \\ -1, & j \notin H; \end{cases}$$

in particular, (5.1) turns into equality on this vector.

Proof. Let W be a geometric space and let

$$\alpha = -\frac{\kappa'(1)}{\log m}.$$

Then, since κ is affine, we have that $\kappa(0) = \alpha \log m$. Thus, for arbitrary $v \in \mathbb{M}^W$, we have $(1 + v_j) \leq m^\alpha$ for all j , and, therefore,

$$(5.2) \quad \sum_{j=1}^m (1 + v_j) \log(1 + v_j) \leq \sum_{j=1}^m (1 + v_j) \alpha \log m = \alpha \log m.$$

Now let v be the vector at which the infimum in (4.3) is attained (by compactness, such a vector exists). The inequality (5.2) turns into equality by the choice of v . Thus, for any $j \in [1 \dots m]$, $1 + v_j$ equals either zero or m^α . The set where it equals m^α is the searched-for set H ; it is clear that $|H| = m^{1-\alpha}$ since v has mean zero. ■

Remark 5.3. If W is a geometric space, then $m^{1-\alpha}$ is an integer.

Remark 5.4. Proposition 5.2 may be reversed, the conditions described there are not only necessary, but also sufficient for W to be a geometric space.

Definition 5.5. The special vectors $v \otimes a$ from Proposition 5.2 for which (5.1) turns into equality are called *extremal*. The corresponding sets H are their supports³.

Remark 5.6. There might be several extremal vectors related to $a \in \mathbb{R}^\ell \setminus \{0\}$.

³This is a slight abuse of notation since H is the support of the function $j \mapsto 1 + v_j$, $j \in [1 \dots m]$.

Remark 5.7. The proof of Proposition 5.2 also says the following: if W is a geometric space and $\kappa_v(\theta) = \kappa(\theta)$ for some $v \in \mathbb{M}^W$, $\theta \in (0, 1)$, then $v \otimes a$ is an extremal vector for some $a \in \mathbb{R}^\ell \setminus \{0\}$. Indeed, by Jensen's inequality⁴,

$$\sum_{j=1}^m (1 + v_j) \log(1 + v_j) \leq \frac{1}{1 - \theta} \log \left(\sum_{j=1}^m (1 + v_j)^{1/\theta} \right)^\theta.$$

We know this inequality turns into equality. Thus, for each j , we have either $1 + v_j = 0$ or $1 + v_j = m^\alpha$.

Definition 5.8. Let $W \subset V \otimes \mathbb{R}^\ell$ be a geometric space. We say that an operator $\varphi: W \rightarrow V$ is *cancelling* if the identity

$$\forall j \in H, \quad (\varphi[w])_j = 0$$

holds true for any extremal vector $w = v \otimes a$ and its support $H \subset [1 \dots m]$.

The following proposition resembles Theorem 1.5 in [33].

Proposition 5.9. *Let W be a geometric space. If (3.4) holds true at the endpoint $\alpha = -\kappa'(1)/\log m$, then φ is a cancelling operator.*

Proof. Let $v \otimes a$ be an extremal vector. Consider the martingale F given by

$$(5.3) \quad F_n = \prod_{j=0}^n (1 + h_j) \otimes a, \quad \text{where} \quad h_j = \sum_{w \in \mathcal{AF}_j} J_w[v].$$

In such a case,

$$dF_{n+1} = F_n h_{n+1} \otimes a \quad \text{and} \quad F \in \mathbb{W}.$$

What is more,

$$\mathbb{I}_\alpha[F] = \sum_{n \geq 0} m^{-\alpha n} \sum_{w \in \mathcal{AF}_n} J_w[\varphi[v \otimes a]] F_n(w).$$

Let H be the support of v . By (5.3), we have $F_n(w) = 0$ provided there is a digit in the m -ary expansion of the 'number' w that does not belong to H . Let us compute the values of $F_n(w)$ at the other 'numbers' $w \in \mathcal{AF}_n$:

$$F_n(w) = m^{\alpha n} a.$$

Therefore, if we pick some large N and consider the stopped martingale

$$F^N = \{F_{\min(n, N)}\}_n,$$

then

$$\mathbb{I}_\alpha[F^N](w) = \sum_{j=0}^N (\varphi[v \otimes a])_{w(j)}$$

at any point w inside the set

$$\mathbb{H}_N = \{w \in \mathcal{AF}_n \mid \forall j \leq N, w(j) \in H\}.$$

⁴We use the weighted Jensen's inequality $\sum \alpha_j f(a_j) \leq f(\sum \alpha_j a_j)$ for a concave function f and substitute $\alpha_j := (1 + v_j)/m$, $a_j := (1 + v_j)^{1/\theta - 1}$, and $f := \log$.

The following two statements will contradict (3.4):

- we have

$$\lim_{N \rightarrow \infty} \sum_{w \in \mathbb{H}^N} \left| \sum_{j=0}^N (\varphi[v \otimes a])_{w(j)} \right| = \infty;$$

- the uniform probability measure on \mathbb{H}^N fulfills the $(\alpha, 1)$ -Frostman condition uniformly with respect to N .

The second statement follows from the fact that the said measure is represented via the correspondence (2.1) by the martingale $\{\prod_{j=0}^n (1 + h_j)\}_n$, which is the scalar version of F , and the bound

$$\left\| \prod_{j=0}^n (1 + h_j) \right\|_{L^\infty} \leq m^{\alpha n}.$$

To prove the first statement, we consider following the random variables ξ_j :

$$\xi_j(w) = (\varphi[v \otimes a])_{w(j)}, \quad w \in \mathbb{H}_N,$$

on the probability space \mathbb{H}_N equipped with the uniform probability measure. Note that these random variables are identically distributed and independent. We need to prove that $\mathbb{E} \left| \sum_{j=1}^N \xi_j \right| \rightarrow \infty$ as $N \rightarrow \infty$ (note that the mathematical expectation is computed with respect to the uniform measure on \mathbb{H}_N). In the case $\mathbb{E} \xi_j \neq 0$, this follows from the triangle inequality. In the remaining case $\mathbb{E} \xi_j = 0$, we use the central limit theorem

$$N^{-1/2} \sum_{j=0}^N \xi_j \rightarrow \mathcal{N}(0, \sigma) \quad \text{in distribution,}$$

where σ is the variance of ξ_j . Thus, $\mathbb{E} \left| \sum_{j=0}^N \xi_j \right| \sim c\sqrt{N} \rightarrow \infty$. ■

We will need yet another condition imposed on W . The necessity of this condition is doubtful.

Definition 5.10. Let $\alpha \in (0, 1)$ be fixed. We say that a geometric space $W \subset V \otimes \mathbb{R}^\ell$ is non-local (of order α) if for any extremal vector $v \otimes a \in \mathbb{R}^\ell$ supported on H , any $w \in W$, and any $b \in \mathbb{R}^\ell$ the identity

$$b + w_j = 0, \quad \forall j \notin H,$$

implies that w is proportional to $v \otimes a$.

In particular, the non-locality assumption implies that for any $H \subset [1 \dots m]$, all $a \in \mathbb{R}^\ell \setminus \{0\}$ such that there exists an extremal vector $v \otimes a$ supported on H , are proportional; this somehow resembles the strong cancellation condition in [15] (the similarity is emphasized by the Fourier-side sufficient condition of non-locality in Lemma 7.15 below). We are ready to formulate our main result – the endpoint trace theorem.

Theorem 5.11. *Let W be a geometric non-local space and let φ be a cancelling operator. Then, (3.4) holds true at the endpoint $\alpha = -\kappa'(1)/\log m$.*

This theorem has a Bellman function behind it.

Theorem 5.12. *Let W be a geometric non-local space and let φ be a cancelling operator. There exists a collection of constants $1 \ll C_1 \ll C_2 \ll C_3$ such that the function*

$$(5.4) \quad \Phi(x, y, z, t, s) = |y|t + C_1|x|t + C_2|x|\sqrt{st} + C_3(z - |x|)s$$

is a supersolution.

Theorem 5.12 implies Theorem 5.11 via Corollary 3.8, since Φ satisfies both the boundary inequality

$$\Phi(|x|, y, |x|, s, t) \geq |y|t$$

and (3.13).

Remark 5.13. Mysteriously, both terms $|x|t$ and $|x|\sqrt{st}$ are required in the sense that both functions

$$|y|t + C_2|x|\sqrt{st} + C_3(z - |x|)s \quad \text{and} \quad |y|t + C_1|x|t + C_3(z - |x|)s$$

are not supersolutions in general. We will not prove this claim. However, inspecting the proof of Theorem 5.12 at the end of the forthcoming section, one may see that the discrepancies of the two terms, namely, of $|x|t$ and of $|x|\sqrt{st}$, control the discrepancy of $|y|t$ at different places of the configuration space.

6. Proof of Theorem 5.12

In fact, the proof consists of an accurate computation and estimate of $\text{DISCR}[\Phi]$, which is done in several steps. We start with the function $|x|\sqrt{st} + M(z - |x|)s$ and prove that its discrepancy allows a bound from below better than simply zero (see Lemma 6.1 below). The general line is similar to the proof of Theorem 4.2: the discrepancy of the function in question is positive everywhere (by that theorem), except the extremal vectors and the corresponding extremal configurations, and we need to analyze its behavior close to those configurations. This is the most hard part of the proof. Then, we use these bounds to show that the function $|x|t + L(|x|\sqrt{st} + M(z - |x|)s)$ is a supersolution in Proposition 6.2. Finally, we use all these estimates to bound the discrepancy of $|y|t$ from below.

6.1. Estimates for $|x|\sqrt{st} + M(z - |x|)s$

One may verify that this function is a supersolution, provided M is sufficiently large, using the reasonings presented in the proof of Theorem 4.2 (see (4.13)). Now we need to prove that this function is a supersolution with certain excess. We derive from (4.6) and (4.9) that

$$\begin{aligned} \text{DISCR} [|x|\sqrt{st} + M(z - |x|)s] \\ \geq \sqrt{st} \left(|x| - m^{-\alpha/2} \left(\sum_{j=1}^m |x_j|^2 \right)^{1/2} + M \left(\sum_{j=1}^m |x_j| - |x| \right) \right) \end{aligned}$$

(recall $s \geq t$).

By homogeneity with respect to the variables x and z , we may assume $|x| = 1$. We may also assume $\sum_{j=1}^m |x_j| \leq (1 + \varepsilon)|x|$ for sufficiently small ε , the other case will be covered by the flat/convex argument and the choice of large M (or large C_3 for the initial function Φ). Thus, by Lemma 4.5, we need to study the behavior of the function

$$|x| - m^{-\alpha/2} \left(\sum_{j=1}^m |x_j|^2 \right)^{1/2} + M \left(\sum_{j=1}^m |x_j| - |x| \right)$$

when \vec{x} lies in a neighborhood of some $v \otimes a \in W$ and x lies in a neighborhood of a . Using the homogeneity with respect to the variables x and z , we may assume $|a| = 1$ and $\langle x, a \rangle = 1$, which does not change the things much since all our functions are homogeneous of order one with respect to x and z . Of course, $|x|$ is not necessarily equal to 1 anymore. Note that if $v \otimes a$ is not an extremal vector, then

$$(6.1) \quad \left(|x| - m^{-\alpha/2} \left(\sum_{j=1}^m |x_j|^2 \right)^{1/2} \right) > 0$$

provided \vec{x} is sufficiently close to $v \otimes a$ (since this function is positive at $x = v \otimes a$, see Remark 5.7). The case where $v \otimes a$ is an extremal vector is covered by the following lemma.

Lemma 6.1. *Assume $v \otimes a \in W$ is an extremal vector of the geometric non-local space W . If $\langle x, a \rangle = 1$, $\vec{x} \in W$ is sufficiently close to $v \otimes a$, and x is sufficiently close to a , then*

$$\text{DISCR} \left[|x| \sqrt{st} + M(z - |x|)s \right] \gtrsim \left(\sum_{j=1}^m |x_j - (1 + v_j)a| \right) \sqrt{st},$$

provided M is sufficiently large; here $v = (v_1, v_2, \dots, v_m)$.

Note that

$$\sum_{j=1}^m |x_j - (1 + v_j)a| = \sum_{j \in H} |x_j - m^\alpha a| + \sum_{j \notin H} |x_j|,$$

where H is the support of the extremal vector $v \otimes a$.

Proof. By the reasonings above, it suffices to prove that

$$(6.2) \quad \left(|x| - m^{-\alpha/2} \left(\sum_{j=1}^m |x_j|^2 \right)^{1/2} \right) + M \left(\sum_{j=1}^m |x_j| - |x| \right) \gtrsim \left(\sum_{j=1}^m |x_j - (1 + v_j)a| \right),$$

given the assumptions of the lemma. Let $\pi_x: \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ be the orthogonal projection onto the line spanned by x . The notation $\chi_{\mathbb{R}_-}[\pi_x[y]]$ means the negative part of the projection: it equals $|\pi_x[y]|$ if $\langle x, y \rangle \leq 0$ and zero otherwise. Let H be the support of $v \otimes a$. We expand the second term on the left-hand side of (6.2) (see the proof of Lemma 2.1 in [4])

for similar estimates):

$$\begin{aligned} \sum_{j=1}^m |x_j| - |x| &= \frac{1}{m} \left(\sum_{j=1}^m (|x_j| - |\pi_x[x_j]|) + \sum_{j=1}^m (|\pi_x[x_j]| - |x|) \right) \\ &\gtrsim \sum_{j \notin H} (|x_j| - |\pi_x[x_j]|) + \sum_{j \notin H} \chi_{\mathbb{R}_-}[\pi_x[x_j]]. \end{aligned}$$

Let

$$(6.3) \quad D = \left(\sum_{j=1}^m |x_j - (1 + v_j)a| \right),$$

which is equivalent to the distance between \mathfrak{X} and $(\mathbf{1}_{[1..m]} + v) \otimes a$. Then,

$$\begin{aligned} |x_j|^2 &= |m^\alpha a|^2 + 2 \langle x_j - m^\alpha a, m^\alpha a \rangle + O(D^2), & j \in H; \\ |x_j|^2 &= O(D^2), & j \notin H; \\ |x|^2 &= |a|^2 + 2 \langle x - a, a \rangle + O(D^2) = 1 + O(D^2), \end{aligned}$$

by our normalization $|a|^2 = \langle x, a \rangle = 1$. Therefore,

$$\begin{aligned} (6.4) \quad |x| - m^{-\alpha/2} \left(\sum_{j=1}^m |x_j|^2 \right)^{1/2} &= 1 + O(D^2) - m^{-\alpha/2} \left(m^{-1} \cdot m^{1-\alpha} \cdot m^{2\alpha} + \frac{2}{m} \sum_{j \in H} \langle x_j - m^\alpha a, m^\alpha a \rangle + O(D^2) \right)^{1/2} \\ &= 1 + O(D^2) - \left(1 + \frac{2}{m^{1+\alpha}} \sum_{j \in H} \langle x_j - m^\alpha a, m^\alpha a \rangle + O(D^2) \right)^{1/2} \\ &= -m^{-1-\alpha} \sum_{j \in H} \langle x_j - m^\alpha a, m^\alpha a \rangle + O(D^2). \end{aligned}$$

We use the identity

$$\left\langle \sum_{j \in H} x_j, a \right\rangle = m - \left\langle \sum_{j \notin H} x_j, a \right\rangle$$

(which follows from $\langle x, a \rangle = |a| = 1$) to rewrite the expression obtained in (6.4) as

$$-m^{-1-\alpha} \sum_{j \in H} \langle x_j - m^\alpha a, m^\alpha a \rangle + O(D^2) = \frac{1}{m} \left\langle \sum_{j \notin H} x_j, a \right\rangle + O(D^2).$$

Thus,

$$\begin{aligned} (6.5) \quad & \left(|x| - m^{-\alpha/2} \left(\sum_{j=1}^m |x_j|^2 \right)^{1/2} \right) + M \left(\sum_{j=1}^m |x_j| - |x| \right) \\ & \gtrsim \frac{1}{m} \left\langle \sum_{j \notin H} x_j, a \right\rangle + M \left(\sum_{j \notin H} (|x_j| - |\pi_x[x_j]|) + \sum_{j \notin H} \chi_{\mathbb{R}_-}[\pi_x[x_j]] \right) + O(D^2). \end{aligned}$$




Figure 1. The plane spanned by a and x .

Claim. *The assumption that W is a non-local space (together with the normalization $\langle a, x \rangle = |a|^2$) leads to the estimate*

$$(6.6) \quad \sum_{j \in H} |x_j - (1 + v_j)a| = \sum_{j \in H} |x_j - m^\alpha a| \lesssim \sum_{j \notin H} |x_j|.$$

To prove the claim, consider the linear subspace \mathfrak{L} of $\mathbb{R}^m \otimes \mathbb{R}^\ell$ formed by the strings $(\xi_1, \xi_2, \dots, \xi_m)$, where $\xi_j \in \mathbb{R}^\ell$, such that $(\xi_1 - \xi, \xi_2 - \xi, \dots, \xi_m - \xi) \in W$ and $\langle \xi, a \rangle = 0$; here, as usual, $\xi = \sum_{j=1}^m \xi_j$. Note that the string formed of vectors $x_j - m^\alpha a$ when $j \in H$ and x_j when $j \notin H$, belongs to this space. Therefore, it suffices to prove the inequality

$$(6.7) \quad \|\Xi\| \lesssim \sum_{j \notin H} |\xi_j|, \quad \Xi = (\xi_1, \xi_2, \dots, \xi_m) \in \mathfrak{L}.$$

Note that the function on the right hand-side of this inequality is one-homogeneous and even. It suffices to prove this function is positive outside the origin. In such a case, it is a norm on \mathfrak{L} , and the desired inequality (6.7) follows since any two norms on a finite dimensional space are equivalent. To prove

$$\sum_{j \notin H} |\xi_j| > 0, \quad \Xi \in \mathfrak{L} \setminus \{0\},$$

assume the contrary: let $\xi_j = 0$ when $j \notin H$ for some non-zero $\Xi \in \mathfrak{L}$. Since W is a non-local space (see Definition 5.10), this means $(\xi_1 - \xi, \xi_2 - \xi, \dots, \xi_m - \xi)$ is proportional to $v \otimes a$. In particular, since some of the ξ_j equals zero, ξ is parallel to a , which, by our assumption $\langle \xi, a \rangle = 0$, means $\xi = 0$. Therefore, $\Xi = 0$, a contradiction. Thus, we have proved the Claim. \blacksquare

In particular, $D \lesssim \sum_{j \notin H} |x_j|$. Therefore, by (6.5), the estimate

$$(6.8) \quad \sum_{j \notin H} |x_j| \lesssim \frac{1}{m} \left\langle \sum_{j \notin H} x_j, a \right\rangle + M \left(\sum_{j \notin H} (|x_j| - |\pi_x[x_j]|) + \sum_{j \notin H} \chi_{\mathbb{R}_-}[\pi_x[x_j]] \right)$$

yields (6.2). Assume $M > 1$, then, without loss of generality, we may replace the x_j with their projections onto the linear plane spanned by x and a (such a procedure will reduce the values $|x_j|$ in the formula above, while all the other quantities will remain the same). We will argue geometrically using Figure 1. We split the plane spanned by x and y by the lines $\langle y, a \rangle = \pm \frac{1}{3}|y|$ into four regions as it is drawn on Figure 1.

If x_j belongs to the second or the fourth domains, then $|x_j| \lesssim |x_j| - |\pi_x[x_j]|$ (recall that x is close to a and, thus, lies strictly inside the first domain). If x_j lies inside the first domain, then $|x_j| \lesssim \langle x_j, a \rangle$. Finally, in the third domain,

$$|x_j| \lesssim \chi_{\mathbb{R}_-}[\pi_x[x_j]],$$

and thus, the choice of large M implies

$$|x_j| + |\langle x_j, a \rangle| \leq M \chi_{\mathbb{R}_-}[\pi_x[x_j]]$$

in this case, which proves (6.8) (in fact, we have proved the individual estimate

$$|x_j| \lesssim \langle x_j, a \rangle + M(|x_j| - |\pi_x[x_j]|) + \chi_{\mathbb{R}_-}[\pi_x[x_j]]$$

provided M is sufficiently large). ■

6.2. Estimates for $|x|t + L(|x|\sqrt{st} + M(z - |x|)s)$

Proposition 6.2. *For sufficiently large L and M , the function*

$$|x|t + L(|x|\sqrt{st} + M(z - |x|)s)$$

is a supersolution.

Proof. We estimate

$$(6.9) \quad \text{DISCR}[|x|t + L(|x|\sqrt{st} + M(z - |x|)s)] \\ \geq \sum_{j=1}^m t_j (|x| - m^{-\alpha} |x_j|) + Lt \left(|x| - m^{-\alpha/2} \left(\sum_{j=1}^m |x_j|^2 \right)^{1/2} + M \left(\sum_{j=1}^m |x_j| - |x| \right) \right);$$

here we have used that $s_j \geq t_j$ for all j (and also (6.1)). Similar to the proof of Theorem 4.2, by Lemma 4.5, it suffices to consider the case where \vec{x} is close to some $v \otimes a$ and x is close to a (the other cases are covered by a combination of a compactness argument and the flat/convex argument). What is more, we may assume $v \otimes a$ is an extremal vector (say, by (6.1)). Let H be the support of v . We normalize x in such a way that $\langle x, a \rangle = 1$ and use Lemma 6.1 (or (6.2)):

$$\text{DISCR}[|x|t + L(|x|\sqrt{st} + M(z - |x|)s)] \\ \geq \sum_{j=1}^m t_j (|x| - m^{-\alpha} |x_j|) + Lt \sum_{i=1}^m |x_i - (1 + v_i)a|.$$

Note that for $j \notin H$, the expression in the parenthesis is always non-negative since x_j is close 0 in this case, while x is close to a . Fix some $j \in H$. Observe that the function $\mathfrak{X} \mapsto |x| - m^{-\alpha} |x_j|$ is Lipschitz on $\mathbb{R}^{m\ell}$ and vanishes at $(\mathbf{1}_{[1\dots m]} + v) \otimes a$, which means its absolute value is bounded by

$$L \sum_{i=1}^m |x_i - (1 + v_i)a|,$$

provided L is sufficiently large (because the latter expression is the distance between \mathfrak{X} and $(\mathbf{1}_{[1\dots m]} + v) \otimes a$).

Thus,

$$(6.10) \quad |x| - m^{-\alpha}|x_j| \geq 0, \quad j \notin H;$$

$$(6.11) \quad \left| |x| - m^{-\alpha}|x_j| \right| \lesssim \sum_{i=1}^m |x_i - (1 + v_i)a|, \quad j \in H,$$

and the proof is completed by choosing large L . \blacksquare

6.3. End of proof of Theorem 5.12

We rewrite the discrepancy of the first summand:

$$(6.12) \quad \text{DISCR}[|y|t] = \sum_{j=1}^m (|y| - |y + m^{-\alpha}(\varphi[\bar{x}])_j|)t_j.$$

We will treat this expression as a function of \mathfrak{X} and $\mathcal{T} = (t_1, t_2, \dots, t_m)$. Without loss of generality, we may assume $|x| = 1$ and $t = 1$. The estimate

$$\left| |y| - |y + m^{-\alpha}(\varphi[\bar{x}])_j| \right| \lesssim |\mathfrak{X}|$$

permits the application of the flat/convex argument, provided we have good estimates in neighborhoods of flat configurations, because if the configuration \mathfrak{X} satisfies (4.8), then the discrepancy of the fourth term in (5.4) majorizes the discrepancies of the other three terms. We also note that the set of all ε -flat configurations is compact, provided we require the normalization $|x| = 1$. In particular, given any flat configuration $((\mathbf{1}_{[1\dots m]} + v) \otimes a, \mathcal{T}_0)$, it suffices to find its neighborhood and a large constant K such that

$$\text{DISCR}[|y|t + K(|x|t + L(|x|\sqrt{st} + M(z - |x|)))] \geq 0$$

when $(\mathfrak{X}, \mathcal{T})$ lies inside the said neighborhood and $y, \mathfrak{Z} = (z_1, z_2, \dots, z_m)$, and $\mathfrak{S} = (s_1, s_2, \dots, s_m)$ are arbitrary. If $v \otimes a$ is not extremal, then the term

$$\text{DISCR}[|x|t + L(|x|\sqrt{st} + M(z - |x|))]$$

is bounded away from zero (uniformly with respect to \mathfrak{Z} and \mathfrak{S}) in a sufficiently small neighborhood of $((\mathbf{1}_{[1\dots m]} + v) \otimes a, \mathcal{T}_0)$ (say, by (6.1)), and the problem reduces to the choice of sufficiently large K . Assume $v \otimes a$ is extremal. As usual, we renormalize x to $\langle x, a \rangle = 1$. We estimate

$$\text{DISCR}[|y|t] \geq - \sum_{j \notin H} |\mathfrak{X}|t_j - \sum_{j \in H} \left(\sum_{i=1}^m |x_i - (1 + v_i)a| \right) t_j;$$

this bound comes from (6.12) and from the cancellation condition imposed on φ (for any $j \in H$, we have $(\varphi(v \otimes a))_j = 0$; recall also that D given in (6.3) is the distance between \mathfrak{X} and $(\mathbf{1}_{[1\dots m]} + v) \otimes a$). The second sum is compensated by the discrepancy of $|x|\sqrt{st} + M(z - |x|)s$, by Lemma 6.1. As for the first sum, we have

$$\sum_{j \notin H} |\mathfrak{X}|t_j \lesssim \sum_{j \notin H} (|x| - m^{-\alpha}|x_j|)t_j,$$

provided x is sufficiently close to $v \otimes a$. Thus, this part of the discrepancy may be compensated by $\text{DISCR}[|x|t]$ by choosing sufficiently large C_1 . \blacksquare

7. Two examples

Let now $m = \mu d$, where μ and d are natural numbers. We equip the set $[1 \dots m]$ with the group structure of $(\mathbb{Z}_\mu)^d$; this also identifies it with the cube $[0 \dots \mu - 1]^d$ in a natural way. Let

$$e_j = \underbrace{(0, 0, \dots, 0}_{j-1 \text{ times}}, 1, 0, \dots, 0) \in [0 \dots \mu - 1]^d, \quad j = 1, 2, \dots, d,$$

be the generators of $(\mathbb{Z}_\mu)^d$. It will be convenient to consider complex-valued functions and martingales, i.e., we set $\ell = 2l$ and identify \mathbb{C}^l and \mathbb{R}^ℓ in a natural way. In our two main examples, $l = d$. The first example will model the space $\text{BV}(\mathbb{R}^d)$,

$$(7.1) \quad W_\nabla = \left\{ g \in V \otimes \mathbb{C}^d \mid \exists f \in \mathbb{R}^m \text{ such that } \forall x \in [0 \dots \mu - 1]^d, j \in [1 \dots d], \right. \\ \left. g_j(x) = f(x + e_j) - f(x) \right\}.$$

The second will model the space of solenoidal charges,

$$(7.2) \quad W_{\text{div}} = \left\{ g \in V \otimes \mathbb{C}^d \mid \forall x \in [0 \dots \mu - 1]^d, \sum_{j=1}^d (g_j(x + e_j) - g_j(x)) = 0 \right\}.$$

Before we pass to the details, we survey simple facts about translation invariant spaces.

Definition 7.1. We say that a linear subspace $W \subset V \otimes \mathbb{C}^l$ is translation invariant if for any $f \in W$ and $y \in (\mathbb{Z}_\mu)^d$, the function $[0 \dots \mu - 1]^d \ni x \mapsto f(x + y)$ also belongs to W ; the addition sign means addition in $(\mathbb{Z}_\mu)^d$.

Note that the spaces W_∇ and W_{div} are translation invariant. Let us define the Fourier transform of a function $f: (\mathbb{Z}_\mu)^d \rightarrow \mathbb{C}^l$ by the formula

$$\hat{f}(\gamma) = \sum_{x \in (\mathbb{Z}_\mu)^d} e^{-2\pi i \gamma \cdot x / \mu} f(x), \quad \gamma \in (\mathbb{Z}_\mu)^d,$$

where $\gamma \cdot x$ is short for $\sum_{j=1}^d \gamma_j x_j \in \mathbb{Z}_\mu$, and we interpret this expression as a residue modulo μ (an integer number in $[0 \dots \mu - 1]$). The inverse Fourier transform is then given by the formula

$$\check{g}(x) = m^{-1} \sum_{\gamma \in (\mathbb{Z}_\mu)^d} e^{2\pi i \gamma \cdot x / \mu} g(\gamma), \quad x \in (\mathbb{Z}_\mu)^d.$$

We refer the reader to the book [34] for introduction to abstract Fourier analysis (though our case of finite groups is elementary, sometimes the abstract intuition seems useful). The following lemma is simple, so we leave it without proof.

Lemma 7.2. *The space $W \subset V \otimes \mathbb{C}^l$ is translation invariant if and only if there exist linear spaces $\Omega(\gamma) \subset \mathbb{C}^l$, $\gamma \in (\mathbb{Z}_\mu)^d \setminus \{0\}$, such that*

$$W = \left\{ f \in V \otimes \mathbb{C}^l \mid \forall \gamma \in (\mathbb{Z}_\mu)^d \setminus \{0\}, \hat{f}(\gamma) \in \Omega(\gamma) \right\}.$$

Remark 7.3. Since any element of W has mean zero, $\hat{f}(0) = 0$ for any $f \in W$. We may set $\Omega(0) = \{0\}$.

In the case $W = W_{\nabla}$, the corresponding spaces $\Omega(\gamma)$ are complex lines defined by the formula

$$(7.3) \quad \Omega_{\nabla}(\gamma) = (e^{2\pi i \gamma_1/\mu} - 1, e^{2\pi i \gamma_2/\mu} - 1, \dots, e^{2\pi i \gamma_d/\mu} - 1) \cdot \mathbb{C},$$

for $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_d) \in (\mathbb{Z}_{\mu})^d \setminus \{0\}$. In the case $W = W_{\text{div}}$, the spaces $\Omega(\gamma)$ are complex hyperplanes

$$\Omega_{\text{div}}(\gamma) = \left\{ \zeta \in \mathbb{C}^l \mid \sum_{j=1}^d \zeta_j (e^{2\pi i \gamma_j/\mu} - 1) = 0 \right\},$$

for $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_d) \in (\mathbb{Z}_{\mu})^d \setminus \{0\}$.

If W is a translation invariant space, the sets $\Omega^{-1}(a)$, where $a \in \mathbb{C}^l$, play an important role:

$$\Omega^{-1}(a) = \{\gamma \in (\mathbb{Z}_{\mu})^d \mid a \in \Omega(\gamma)\},$$

since these sets contain the spectra of rank-one vectors. By spectrum of a function we mean the support of its Fourier transform.

Lemma 7.4. *Let $k \in [0 \dots d]$. Assume W is a translation invariant subspace of $V \otimes \mathbb{C}^l$ such that*

$$(7.4) \quad \#(\Omega^{-1}(a) \cap (-\Omega^{-1}(a))) \leq \mu^k - 1$$

for any $a \in \mathbb{C}^l \setminus \{0\}$. Assume (7.4) turns into equality for at least one vector a , and that for this special a we have $\Omega^{-1}(a) = \Gamma \setminus \{0\}$, where Γ is a subgroup of index $d - k$ in $(\mathbb{Z}_{\mu})^d$. Then, W is a geometric space of order k/d .

Proof. Let $v \otimes a \in V \otimes \mathbb{C}^l$ be a rank-one vector. By definition,

$$\hat{v}(\gamma)a \in \Omega(\gamma), \quad \gamma \in (\mathbb{Z}_{\mu})^d.$$

Here we treat v as a real valued function on $[0 \dots \mu - 1]^d$. Therefore, $a \in \Omega(\gamma)$ whenever $\hat{v}(\gamma) \neq 0$. Since $\hat{v}(-\gamma) = \overline{\hat{v}(\gamma)}$, we have

$$\text{supp } \hat{v} \subset \Omega^{-1}(a) \cap (-\Omega^{-1}(a)).$$

Assume now that $v(x) \geq -1$ for any $x \in [0 \dots \mu - 1]^d$, and let $w(x) = 1 + v(x)$. Then, \hat{w} is supported in

$$(\Omega^{-1}(a) \cap (-\Omega^{-1}(a))) \cup \{0\}$$

and, since $\sum_x w(x) = m$ and the numbers $w(x)$ are non-negative, $|\hat{w}(\gamma)| \leq m$ for any γ . From this information and the bound (7.4), we conclude that

$$(7.5) \quad |w(x)| = m^{-1} \left| \sum_{\gamma \in (\mathbb{Z}_{\mu})^d} e^{2\pi i \gamma \cdot x/\mu} \hat{w}(\gamma) \right| \leq \mu^k, \quad x \in [0 \dots \mu - 1]^d.$$

Therefore, by (4.1), $\kappa_v(0) \leq k \log \mu$. By the convexity of the function κ_v and the identity $\kappa_v(1) = 0$,

$$\kappa_v(\theta) \leq k(\log \mu)(1 - \theta).$$

On the other hand, if the set

$$(\Omega^{-1}(a) \cap (-\Omega^{-1}(a))) \cup \{0\}$$

is a subgroup Γ , then we set $w = |\Gamma| \chi_{\Gamma^\perp}$ and see that $v \otimes a \in W$ for the corresponding v . For such a choice of v and w , the inequalities (7.5) turn into equalities when $x \in \Gamma^\perp$, which means

$$\kappa_v(\theta) = k \log \mu(1 - \theta) = \frac{k}{d} \log m(1 - \theta),$$

and Γ^\perp is the support of an extremal vector. Thus, the function κ is linear and W is a geometric space. ■

Remark 7.5. The proof above says that extremal vectors correspond to those $a \in \mathbb{C}^l \setminus \{0\}$, for which the set

$$(\Omega^{-1}(a) \cap (-\Omega^{-1}(a))) \cup \{0\}$$

is a subgroup Γ of index $d - k$. If $v \otimes a$ is an extremal vector, then its support is a coset of Γ^\perp .

Definition 7.6. A subset $\ell_D \subset (\mathbb{Z}_\mu)^d$ is called a combinatorial line generated by a set $D \subset [0 \dots \mu - 1]$ provided

$$\ell_D = \{x \in (\mathbb{Z}_\mu)^d \mid \exists b \in \mathbb{Z}_\mu \text{ such that } x_j = 0, j \notin D, x_j = b, j \in D\}.$$

Lemma 7.7. Let $W = W_{\nabla}$. Then, for any $a \in \mathbb{C}^d \setminus \{0\}$, either

$$(\Omega_{\nabla}^{-1}(a) \cap (-\Omega_{\nabla}^{-1}(a))) \cup \{0\} = \ell_D, \quad \text{for some } D \subset [1 \dots d],$$

or $\Omega_{\nabla}^{-1}(a) \cap (-\Omega_{\nabla}^{-1}(a)) = \emptyset$.

Proof. Let $a \in \mathbb{C}^l \setminus \{0\}$. Assume some points $\gamma^1, \gamma^2, \dots, \gamma^s$ belong to $\Omega_{\nabla}^{-1}(a)$. By (7.3), this means there exist non-zero complex numbers $\lambda_1, \lambda_2, \dots, \lambda_s$ such that

$$a = \lambda_k (e^{2\pi i \gamma_1^k / \mu} - 1, e^{2\pi i \gamma_2^k / \mu} - 1, \dots, e^{2\pi i \gamma_d^k / \mu} - 1), \quad k = 1, 2, \dots, s.$$

Define

$$(7.6) \quad D = \{j \in [1 \dots d] \mid a_j \neq 0\}.$$

Note that $\gamma_j^k = 0$ for any $k \in [1 \dots s]$ and $j \notin D$. If there are at least two distinct non-zero values a_{j_1} and a_{j_2} , then the ratio

$$1 \neq \frac{a_{j_1}}{a_{j_2}} = \frac{e^{2\pi i \gamma_{j_1}^k / \mu} - 1}{e^{2\pi i \gamma_{j_2}^k / \mu} - 1}$$

completely defines the elements $\gamma_{j_1}^k$ and $\gamma_{j_2}^k$ (i.e., these values do not depend on k), since the equation

$$e^{ix} - 1 = \alpha(e^{iy} - 1), \quad x, y \in (0, 2\pi),$$

has at most one solution when $\alpha \neq 1$. Thus, either all non-zero a_j are equal, or $s = 1$. In the latter case $a \notin \Omega_{\nabla}(-\gamma)$. In the former case, $\Omega_{\nabla}^{-1}(a) \cap (-\Omega_{\nabla}^{-1}(a)) = \ell_D$ with D given in (7.6). ■

Corollary 7.8. *The space W_{∇} is a geometric space of order $1/d$.*

Proof. We note that each combinatorial line is a subgroup of index $d - 1$, and then use Lemma 7.4. ■

Remark 7.9. The extremal vectors of W_{∇} are easy to describe: by Remark 7.5, each such vector is defined by its support, which is a coset of the subgroup ℓ_D^{\perp} , $D \subset [1 \dots d]$:

$$(7.7) \quad \ell_D^{\perp} = \left\{ x \in [0 \dots \mu - 1]^d \mid \sum_{j \in D} x_j = 0 \right\};$$

the corresponding vector a is parallel to $\sum_{j \in D} e_j$.

It is harder to describe the sets $\Omega_{\text{div}}^{-1}(a)$ in general. We will only prove that apart from several exclusions, these sets are small provided μ is sufficiently large.

Lemma 7.10. *Let $a \in \mathbb{C}^l \setminus \{0\}$. Either $\#\Omega_{\text{div}}^{-1}(a) \lesssim \mu^{d-2}$, or*

$$(7.8) \quad \exists j \in [1 \dots d] \text{ such that } \Omega_{\text{div}}^{-1}(a) = \{\gamma \in (\mathbb{Z}_{\mu})^d \mid \gamma_j = 0\}, \text{ or}$$

$$(7.9) \quad \exists j_1 \neq j_2 \in [1 \dots d] \text{ such that } \Omega_{\text{div}}^{-1}(a) = \{\gamma \in (\mathbb{Z}_{\mu})^d \mid \gamma_{j_1} = \gamma_{j_2}\}.$$

Remark 7.11. The set (7.8) corresponds to a being the j -th basic vector in \mathbb{C}^d (or a multiple of it), the set (7.9) corresponds to a being the difference of j_1 -th and j_2 -th basic vectors (or a multiple of it). Using Remark 7.5, we may describe the extremal vectors. In the case (7.8), each such vector is supported by a coset of the subgroup

$$\Gamma^{\perp} = \mathbb{Z}_{\mu} e_j;$$

in other words, for any extremal vector of this type, there exists $x \in (\mathbb{Z}_{\mu})^d$ such that it is supported by $x + \mathbb{Z}_{\mu} e_j$. In the case (7.9), the support of an extremal vector is a coset of the subgroup

$$\Gamma^{\perp} = \{x \in (\mathbb{Z}_{\mu})^d \mid x_{j_1} + x_{j_2} = 0\}.$$

The proof of Lemma 7.10 will take some time. We need an auxiliary lemma. Let $b \in \mathbb{C}$. Consider the set

$$S_{a,b} = \left\{ \zeta \in (\mathbb{Z}_{\mu})^d \mid \sum_{j=1}^d a_j (e^{2\pi i \zeta_j / \mu} - 1) = b \right\}.$$

Lemma 7.12. *Let all a_j , $j = 1, 2, \dots, d$, be non-zero. If $d \geq 3$, then $\#S_{a,b} \lesssim \mu^{d-2}$. In the case $d = 2$, either $\#S_{a,b} \leq 2$, or $a_1/a_2 = e^{2\pi i \zeta / \mu}$ for some $\zeta \in [0 \dots \mu - 1]$, $b = -a_1 - a_2$, and $\#S_{a,b} = \mu$ in this case.*

Proof. Let us study the structure of the sets $S_{a,b}$ in the case $d = 2$ first.

We rewrite the equation defining $S_{a,b}$ in the form

$$e^{2\pi i \xi_1/\mu} = -\frac{a_2}{a_1} e^{2\pi i \xi_2/\mu} + \frac{a_2 + b + a_1}{a_1}.$$

The left-hand and right-hand sides define two circles in the complex plane. In general, such two circles intersect by at most two points, which leads to the estimate $\#S_{a,b} \leq 2$. However, there is one exception: the two circles might coincide; in such a case, we cannot hope for anything better than $\#S_{a,b} \leq \mu$ (this estimate is definitely true). If the circles coincide, then $|a_1| = |a_2|$ and $b = -a_1 - a_2$. Since we search for solutions $\zeta_1, \zeta_2 \in [0 \dots \mu - 1]$, we also need $a_1/a_2 = e^{2\pi i \zeta/\mu}$, where $\zeta \in [0 \dots \mu - 1]$, to have $\#S_{a,b} = \mu$. We see that for any pair of non-zero numbers (a_1, a_2) , there exists at most one b such that $\#S_{a,b} > 2$.

Now let us pass to the case $d = 3$ and prove that in this case $\#S_{a,b} \leq 3\mu$. We consider the sets

$$\begin{aligned} S_{a,b,\xi_1} &= \{(\zeta_2, \zeta_3) \in (\mathbb{Z}_\mu)^2 \mid a_2(e^{2\pi i \zeta_2/\mu} - 1) + a_3(e^{2\pi i \zeta_3/\mu} - 1) \\ &= b - a_1(e^{2\pi i \xi_1/\mu} - 1)\}. \end{aligned}$$

We note that since $a_1 \neq 0$, $\#S_{a,b,\xi_1} \leq 2$ for all but at most one values of ξ_1 (for this special value, we have $\#S_{a,b,\xi_1} \leq \mu$). Summing over all ξ_1 , we get $\#S_{a,b} \leq 3\mu$.

Using induction and a similar ‘slicing’ trick, we prove that $\#S_{a,b} \leq 3\mu^{d-2}$ for all larger d . ■

Proof of Lemma 7.10. Now we allow some of the numbers a_j to be equal zero. If $d - 1$ of them vanish, we have (7.8). If $d - 2$ of them vanish, let a_{j_1} and a_{j_2} be non-zero. Then,

$$\#\Omega_{\text{div}}^{-1}(a) \leq \mu^{d-2} \#S_{(a_{j_1}, a_{j_2}), 0}.$$

If $a_{j_1} \neq -a_{j_2}$, then

$$\#S_{(a_{j_1}, a_{j_2}), 0} \leq 2,$$

by Lemma 7.12. If $a_{j_1} = -a_{j_2}$, then we arrive at (7.9). Finally, if $d - k$ of the coordinates of a vanish, $k > 2$, then

$$\#\Omega_{\text{div}}^{-1}(a) \lesssim \mu^{d-k} \mu^{k-2} \leq \mu^{d-2}$$

by Lemma 7.12. ■

Remark 7.13. We have proved that, apart from the exceptions, $\#\Omega_{\text{div}}^{-1}(a) \leq 3\mu^{d-2}$, which is less than μ^{d-1} if $\mu \geq 4$.

Corollary 7.14. For $\mu \geq 4$, the space W_{div} is geometric of order $(d - 1)/d$.

Proof. We note that the sets (7.8) and (7.9) are subgroups of index 1, and then we apply Lemma 7.4. ■

Now we pass to verification of the non-locality condition given in Definition 5.10.

Lemma 7.15. *Let $k \in [0 \dots d]$. Assume W is a translation invariant subspace of $V \otimes \mathbb{C}^l$ such that*

$$\#(\Omega^{-1}(a) \cap (-\Omega^{-1}(a))) \leq \mu^k - 1$$

for any $a \in \mathbb{C}^l \setminus \{0\}$. Assume also that if $\Omega^{-1}(a) = \Gamma \setminus \{0\}$, where Γ is a subgroup of index $d - k$ in $(\mathbb{Z}_\mu)^d$, then

$$(7.10) \quad \bigcap_{\gamma \in \Gamma} \Omega(\gamma) = \mathbb{C}a \quad \text{and}$$

$$(7.11) \quad \bigcap_{\gamma \in S} \Omega(\gamma) = \{0\}$$

for any coset $S \neq \Gamma$ of Γ . Then, W is a non-local space of order k/d .

Proof. By Remark 7.5 and translation invariance, it suffices to prove that if the function $c + f$, where c is a constant and $f \in W$, is supported in Γ^\perp for some subgroup Γ of index $d - k$, $\Gamma = \Omega^{-1}(a) \cup \{0\}$, then $c + f$ is proportional to $\chi_{\Gamma^\perp} \otimes a$. If $g = c + f$ is supported in Γ^\perp , then its Fourier transform is constant on each coset of Γ . In particular, for any such coset S and any $\zeta \in S$,

$$\hat{g}(\zeta) \in \bigcap_{\gamma \in S} \Omega(\gamma).$$

By (7.11), \hat{g} is supported in Γ and, by (7.10), \hat{g} is proportional to $\chi_\Gamma \otimes a$. Therefore, g is proportional to $\chi_{\Gamma^\perp} \otimes a$. ■

Corollary 7.16. *Let $\mu \geq 4$. The space W_∇ is non-local of order $1/d$ and the space W_{div} is non-local of order $(d - 1)/d$.*

Proof. We need to verify the assumptions of Lemma 7.15 for the cases W_∇ and W_{div} . The first case follows easily from Lemma 7.7, the second case follows from Lemma 7.10. ■

Let now $\varphi: W \rightarrow V$ be an operator that commutes with translations. We may extend it to an operator mapping $\mathbb{R}^m \otimes \mathbb{C}^l$ to V and then average over translations. The resulting operator will be translation invariant. In particular, there exists a function $M: (\mathbb{Z}_\mu)^d \rightarrow \mathbb{C}^l$ such that

$$(7.12) \quad \mathcal{F}[\varphi[f]](\gamma) = \langle M(\gamma), \hat{f}(\gamma) \rangle, \quad \gamma \in (\mathbb{Z}_\mu)^d, \quad f \in W,$$

the symbol \mathcal{F} means the Fourier transform. Here we use scalar product in \mathbb{C}^l . Without loss of generality, we may assume $M(\gamma) \in \Omega(\gamma)$ (in case this relation does not hold for some γ , we may redefine $M(\gamma)$ to be the orthogonal projection of $M(\gamma)$ onto $\Omega(\gamma)$; this will not change the formula (7.12)). In particular, $M(0) = 0$. Using the inverse Fourier transform, we have

$$(7.13) \quad \varphi[f](x) = \sum_{y \in (\mathbb{Z}_\mu)^d} \langle K(x - y), f(y) \rangle, \quad x \in (\mathbb{Z}_\mu)^d;$$

here $K = m\check{M}$. Assume now that W satisfies the assumptions of Lemma 7.4. Let $v \otimes a$ be an extremal vector. Then, by translation invariance and the condition $\sum_x K(x) = 0$, the cancellation condition from Definition 5.8 is equivalent to

$$0 = \varphi[v \otimes a](0) = \mu^\alpha \sum_{y \in H} \langle K(y), a \rangle.$$

Here we assume $v \otimes a$ is supported by a subgroup H (see Remark 7.5).

In the case $W = W_\nabla$, this reads as follows:

$$(7.14) \quad \forall D \subset [1 \dots d], \quad \sum_{\substack{x: \sum_{j \in D} x_j = 0}} \langle K(x), e_D \rangle = 0, \quad e_D = \sum_{j \in D} e_j.$$

Theorem 7.17. *Let W be given by (7.1). Let the operator φ be defined by (7.13). Then, the inequality (3.4) holds true if and only if the operator φ fulfills the cancellation condition (7.14).*

In the case $W = W_{\text{div}}$, the cancellation condition turns into:

$$(7.15) \quad \forall j \in [1 \dots d], \quad \sum_{y \in \mathbb{Z}_\mu} K_j(y e_j) = 0;$$

$$(7.16) \quad \forall i \neq j \in [1 \dots d], \quad \sum_{y \in \mathbb{Z}_\mu} K_j(y(e_j - e_i)) - K_i(y(e_j - e_i)) = 0.$$

Theorem 7.18. *Let W be given by (7.2). Let the operator φ be defined by (7.13). Then, the inequality (3.4) holds true if and only if the operator φ fulfills the cancellation conditions (7.15) and (7.16).*

Compare with condition (1.10) in [33] and Conjecture 1.6 of the same paper. It says that the inequality

$$\|K * f\|_{L_1(\nu)} \lesssim \|f\|_{L_1(\mathbb{R}^d)}, \quad \text{div } f = 0, \quad K * f = \sum_{j=1}^d K_j * f_j,$$

holds true for all measures ν satisfying the ball growth (or Frostman) condition $\nu(B_r(x)) \lesssim r$, if and only if the kernel $K: \mathbb{R}^d \rightarrow \mathbb{R}^d$ that is homogeneous of order -1 satisfies the condition

$$\sum_{j=1}^d (K_j(x) + K_j(-x)) x_j = 0$$

for any $x \in \mathbb{R}^d \setminus \{0\}$.

Note that since the kernel $K = (K_1, K_2, \dots, K_d)$ is homogeneous of order -1 , the latter condition may be informally interpreted that the (divergent) integral $\int_{\mathbb{R}} \langle K(tx), x \rangle dt$ vanishes for any x , which is somehow similar to (7.15) and (7.16). In a similar manner, Theorem 7.17 leads to another natural conjecture.

Conjecture 7.19. Assume the kernel $K: \mathbb{R}^d \rightarrow \mathbb{R}^d$ is homogeneous of order $1 - d$ and smooth outside the origin. The inequality

$$\|K * \nabla f\|_{L_1(\nu)} \lesssim \|\nabla f\|_{L_1(\mathbb{R}^d)}, \quad f \in C_0^\infty(\mathbb{R}^d),$$

holds true for all measures ν satisfying the condition $\nu(B_r(x)) \lesssim r^{d-1}$ if and only if for any $e \in S^{d-1}$,

$$\int_{e^\perp \cap S^{d-1}} \langle K(\zeta), e \rangle d\zeta = 0.$$

The integration is performed with respect to the natural Hausdorff measure on the $(d - 2)$ -dimensional sphere $e^\perp \cap S^{d-1}$.

Funding. Supported by the Russian Science Foundation grant no 19-71-30002.

References

- [1] Adams, D. R. and Hedberg, L. I.: *Function spaces and potential theory*. Grundlehren Math. Wiss. 314, Springer, Berlin, 1996. Zbl 0834.46021 MR 1411441
- [2] Arroyo-Rabasa, A., De Philippis, G., Hirsch, J. and Rindler, F.: [Dimensional estimates and rectifiability for measures satisfying linear PDE constraints](#). *Geom. Funct. Anal.* **29** (2019), no. 3, 639–658. Zbl 1420.49046 MR 3962875
- [3] Ayoush, R.: [On finite configurations in the spectra of singular measures](#). *Math. Z.* **304** (2023), no. 1, article no. 6, 17 pp. Zbl 07673779 MR 4569405
- [4] Ayoush, R., Stolyarov, D. and Wojciechowski, M.: [Sobolev martingales](#). *Rev. Mat. Iberoam.* **37** (2021), no. 4, 1225–1246. Zbl 1489.60071 MR 4269395
- [5] Bourgain, J. and Brezis, H.: [On the equation \$\operatorname{div} Y = f\$ and application to control of phases](#). *J. Amer. Math. Soc.* **16** (2003), no. 2, 393–426. Zbl 1075.35006 MR 1949165
- [6] Bourgain, J. and Brezis, H.: [New estimates for the Laplacian, the div-curl, and related Hodge systems](#). *C. R. Math. Acad. Sci. Paris* **338** (2004), no. 7, 539–543. Zbl 1101.35013 MR 2057026
- [7] Bourgain, J. and Brezis, H.: [New estimates for elliptic equations and Hodge type systems](#). *J. Eur. Math. Soc. (JEMS)* **9** (2007), no. 2, 277–315. Zbl 1176.35061 MR 2293957
- [8] Bousquet, P. and Van Schaftingen, J.: [Hardy–Sobolev inequalities for vector fields and canceling linear differential operators](#). *Indiana Univ. Math. J.* **63** (2014), no. 5, 1419–1445. Zbl 1325.46037 MR 3283556
- [9] Breit, D., Diening, L. and Gmeineder, F.: [On the trace operator for functions of bounded \$\mathbb{A}\$ -variation](#). *Anal. PDE* **13** (2020), no. 2, 559–594. Zbl 1450.46017 MR 4078236
- [10] Burkholder, D. L.: [Boundary value problems and sharp inequalities for martingale transforms](#). *Ann. Probab.* **12** (1984), no. 3, 647–702. Zbl 0556.60021 MR 0744226
- [11] Chanillo, S., Van Schaftingen, J. and Yung, P.-L.: [Bourgain–Brezis inequalities on symmetric spaces of non-compact type](#). *J. Funct. Anal.* **273** (2017), no. 4, 1504–1547. Zbl 1515.35021 MR 3661407
- [12] Diening, L. and Gmeineder, F.: [Sharp trace and Korn inequalities for differential operators](#). To appear in *Potential Anal.* (2024), DOI 10.1007/s11118-024-10165-1.

- [13] Gagliardo, E.: Ulteriori proprietà di alcune classi di funzioni in più variabili. *Ricerche Mat.* **8** (1959), 24–51. Zbl [0199.44701](#) MR [0109295](#)
- [14] Gmeineder, F. and Raiță, B.: [Embeddings for \$A\$ -weakly differentiable functions on domains](#). *J. Funct. Anal.* **277** (2019), no. 12, article no. 108278, 33 pp. Zbl [1440.46031](#) MR [4019087](#)
- [15] Gmeineder, F., Raiță, B. and Van Schaftingen, J.: [On limiting trace inequalities for vectorial differential operators](#). *Indiana Univ. Math. J.* **70** (2021), no. 5, 2133–2176. Zbl [1493.46054](#) MR [4340491](#)
- [16] Gmeineder, F., Raiță, B. and Van Schaftingen, J.: [Boundary ellipticity and limiting \$L^1\$ -estimates on halfspaces](#). *Adv. Math.* **439** (2024), article no. 109490, 25 pp. Zbl [07807342](#) MR [4690504](#)
- [17] Hernandez, F. and Spector, D.: [Fractional integration and optimal estimates for elliptic systems](#). *Calc. Var. Partial Differential Equations* **63** (2024), no. 5, article no. 117, 29 pp. Zbl [07845227](#) MR [4739434](#)
- [18] Janson, S.: [Characterizations of \$H^1\$ by singular integral transforms on martingales and \$\mathbb{R}^n\$](#) . *Math. Scand.* **41** (1977), no. 1, 140–152. Zbl [0369.42005](#) MR [0463789](#)
- [19] Kislyakov, S. V., Maksimov, D. V. and Stolyarov, D. M.: [Differential expressions with mixed homogeneity and spaces of smooth functions they generate in arbitrary dimension](#). *J. Funct. Anal.* **269** (2015), no. 10, 3220–3263. Zbl [1348.46037](#) MR [3401616](#)
- [20] Lanzani, L. and Stein, E. M.: [A note on div curl inequalities](#). *Math. Res. Lett.* **12** (2005), no. 1, 57–61. Zbl [1113.26015](#) MR [2122730](#)
- [21] Mattila, P.: *Geometry of sets and measures in Euclidean spaces*. Cambridge Stud. Adv. Math. 44, Cambridge University Press, Cambridge, 1995. Zbl [0819.28004](#) MR [1333890](#)
- [22] Maz’ya, V. G.: Strong capacity-estimates for “fractional” norms. *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **70** (1977), 161–168, 292. Zbl [0517.46028](#) MR [500118](#)
- [23] Maz’ya, V. G.: [Bourgain–Brezis type inequality with explicit constants](#). In *Interpolation theory and applications*, pp. 247–252. Contemp. Math. 445, American Mathematical Society, Providence, RI, 2007. Zbl [1143.46017](#) MR [2381898](#)
- [24] Maz’ya, V. G.: [Estimates for differential operators of vector analysis involving \$L^1\$ -norm](#). *J. Eur. Math. Soc. (JEMS)* **12** (2010), no. 1, 221–240. Zbl [1187.42011](#) MR [2578609](#)
- [25] Maz’ja, V. G.: *Sobolev spaces. With applications to elliptic partial differential equations*. Grundlehren Math. Wiss. 342, Springer, Berlin, 2011. Zbl [1217.46002](#) MR [0817985](#)
- [26] Meyers, N. G. and Ziemer, W. P.: [Integral inequalities of Poincaré and Wirtinger type for BV functions](#). *Amer. J. Math.* **99** (1977), no. 6, 1345–1360. Zbl [0416.46025](#) MR [0507433](#)
- [27] Nakai, E. and Sadasue, G.: [Martingale Morrey–Campanato spaces and fractional integrals](#). *J. Funct. Spaces Appl.* (2012), article no. 673929, 29 pp. Zbl [1254.46035](#) MR [2944703](#)
- [28] Nazarov, F. L. and Treil’, S. R.: The hunt for a Bellman function: applications to estimates for singular integral operators and to other classical problems of harmonic analysis. *Algebra i Analiz* **8** (1996), no. 5, 32–162; translation in *St. Petersburg Math. J.* **8** (1997), no. 5, 721–824. Zbl [0873.42011](#) MR [1428988](#)
- [29] Nirenberg, L.: On elliptic partial differential equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)* **13** (1959), 115–162. Zbl [0088.07601](#) MR [0109940](#)
- [30] Osękowski, A.: *Sharp martingale and semimartingale inequalities*. Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series) 72, Birkhäuser/Springer Basel AG, Basel, 2012. Zbl [1278.60005](#) MR [2964297](#)

- [31] Raiță, B.: L_1 -estimates for constant rank operators. Preprint 2018, arXiv:1811.10057v1.
- [32] Raiță, B.: [Critical \$L^p\$ -differentiability of \$BV^{\Delta}\$ -maps and canceling operators](#). *Trans. Amer. Math. Soc.* **372** (2019), no. 10, 7297–7326. Zbl 1429.26019 MR 4024554
- [33] Raiță, B., Spector, D. and Stolyarov, D.: [A trace inequality for solenoidal charges](#). *Potential Anal.* **59** (2023), no. 4, 2093–2104. Zbl 1541.46018 MR 4684387
- [34] Rudin, W.: *Fourier analysis on groups*. Interscience Tracts in Pure and Applied Mathematics 12, Interscience Publishers (a division of John Wiley & Sons), New York-London, 1962. Zbl 0107.09603 MR 0152834
- [35] Schikorra, A., Spector, D. and Van Schaftingen, J.: [An \$L^1\$ -type estimate for Riesz potentials](#). *Rev. Mat. Iberoam.* **33** (2017), no. 1, 291–303. Zbl 1375.47039 MR 3615452
- [36] Shiryaev, A. N.: *Probability-2*. Third edition. Grad. Texts in Math. 95, Springer, New York, 2019. Zbl 1472.60001 MR 3930599
- [37] Soboleff, S.: Sur un théorème d'analyse fonctionnelle. *Mat. Sbornik* **4(46)** (1938), no. 3, 471–497 (in Russian); translated in *Amer. Math. Soc. Transl.* **2** (1963) (34), 39–68. Zbl 64.1100.02
- [38] Spector, D.: [An optimal Sobolev embedding for \$L^1\$](#) . *J. Funct. Anal.* **279** (2020), no. 3, article no. 108559, 26 pp. Zbl 1455.46039 MR 4093790
- [39] Spector, D. and Van Schaftingen, J.: [Optimal embeddings into Lorentz spaces for some vector differential operators via Gagliardo's lemma](#). *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **30** (2019), no. 3, 413–436. Zbl 1442.46027 MR 4002205
- [40] Stolyarov, D.: [Martingale interpretation of weakly cancelling differential operators](#). *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **480** (2019), 191–198; translated in *J. Math. Sci., New York* **251** (2020), no. 2, 286–290. Zbl 1466.46030 MR 4058922
- [41] Stolyarov, D.: [Weakly canceling operators and singular integrals](#). *Tr. Mat. Inst. Steklova* **312** (2021), 259–271. Zbl 1462.42025 MR 4237403
- [42] Stolyarov, D.: [Hardy–Littlewood–Sobolev inequality for \$p = 1\$](#) . *Mat. Sb.* **213** (2022), no. 6, 125–174. Zbl 1531.46020 MR 4461456
- [43] Stolyarov, D.: [Dimension estimates for vectorial measures with restricted spectrum](#). *J. Funct. Anal.* **284** (2023), no. 1, article no. 109735, 16 pp. Zbl 07616885 MR 4500728
- [44] Stolyarov, D.: [On Maz'ya's \$\Phi\$ -inequalities for martingale fractional integration and their Bellman functions](#). *Michigan Math. J.* **74** (2024), no. 3, 509–526. Zbl 07899198 MR 4767503
- [45] Stolyarov, D.: [Fractional integration of summable functions: Maz'ya's \$\Phi\$ -inequalities](#). *Ann. Scuola Norm. Sup. Pisa* **25** (2024), 1727–1752. Zbl 07944264
- [46] Stolyarov, D. and Yarcev, D.: [Fractional integration for irregular martingales](#). *Tohoku Math. J. (2)* **74** (2022), no. 2, 253–261. Zbl 1503.60051 MR 4455867
- [47] Van Schaftingen, J.: [Estimates for \$L^1\$ -vector fields](#). *C. R. Math. Acad. Sci. Paris* **339** (2004), no. 3, 181–186. Zbl 1049.35069 MR 2078071
- [48] Van Schaftingen, J.: [A simple proof of an inequality of Bourgain, Brezis and Mironescu](#). *C. R. Math. Acad. Sci. Paris* **338** (2004), no. 1, 23–26. Zbl 1188.26015 MR 2038078
- [49] Van Schaftingen, J.: [Estimates for \$L^1\$ vector fields under higher-order differential conditions](#). *J. Eur. Math. Soc. (JEMS)* **10** (2008), no. 4, 867–882. Zbl 1228.46034 MR 2443922
- [50] Van Schaftingen, J.: [Limiting fractional and Lorentz space estimates of differential forms](#). *Proc. Amer. Math. Soc.* **138** (2010), no. 1, 235–240. Zbl 1184.35012 MR 2550188

- [51] Van Schaftingen, J.: [Limiting Sobolev inequalities for vector fields and canceling linear differential operators](#). *J. Eur. Math. Soc. (JEMS)* **15** (2013), no. 3, 877–921. Zbl [1284.46032](#) MR [3085095](#)
- [52] Van Schaftingen, J.: [Limiting Bourgain–Brezis estimates for systems of linear differential equations: theme and variations](#). *J. Fixed Point Theory Appl.* **15** (2014), no. 2, 273–297. Zbl [1311.35005](#) MR [3298002](#)
- [53] Vasyunin, V. and Volberg, A.: *The Bellman function technique in harmonic analysis*. Cambridge Stud. Adv. Math. 186, Cambridge University Press, Cambridge, 2020. Zbl [1498.42003](#) MR [4411371](#)
- [54] Watari, C.: [Multipliers for Walsh–Fourier series](#). *Tohoku Math. J. (2)* **16** (1964), 239–251. Zbl [0135.27502](#) MR [0206608](#)

Received June 21, 2023; revised October 31, 2024.

Dmitriy Stolyarov

Department of Mathematics and Computer Sciences, St. Petersburg State University
14th line 29b, Vasilyevski Island, 199178 St. Petersburg, Russia;
d.m.stolyarov@spbu.ru