

The category Θ_2 , derived modifications, and deformation theory of monoidal categories

Piergiorgio Panero and Boris Shoikhet

Abstract. A complex $C^\bullet(C, D)(F, G)(\eta, \theta)$, generalising the Davydov–Yetter complex of a monoidal category (Davydov (1998) and Yetter (1998)), is constructed. Here, C, D are \mathbb{k} -linear (corresp., dg) bicategories, $F, G: C \rightarrow D$ are \mathbb{k} -linear (corresp., dg) strong functors, and $\eta, \theta: F \Rightarrow G$ are strong natural transformations. Morally, it is a complex of “derived modifications” $\eta \Rightarrow \theta$; likewise for the case of dg categories, one has the complex of “derived natural transformations” $F \Rightarrow G$, given by the Hochschild cochain complex of C with coefficients in C -bimodule $D(F-, G-)$.

The complex $C^\bullet(C, D)(F, G)(\eta, \theta)$ naturally arises from a 2-cocellular dg vector space $A(C, D)(F, G)(\eta, \theta): \Theta_2 \rightarrow C^\bullet(\mathbb{k})$, as its Θ_2 -totalisation (here, Θ_2 is the category dual to the category of Joyal 2-disks (Joyal (1997))).

It is shown that for a \mathbb{k} -linear monoidal category C , the third cohomology vector space $H^3(C^\bullet(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id}))$ is isomorphic to the vector space of the outer (modulo twists) infinitesimal deformations of the \mathbb{k} -linear monoidal category which we call the *full* deformations. It means that the following data is to be deformed: (a) the underlying dg category structure, (b) the monoidal product on morphisms (the monoidal product on objects is a set-theoretical datum and is maintained under the deformation), and (c) the associator. The data (a), (b), (c) is subject to the (infinitesimal versions of) numerous monoidal compatibilities, which we interpret as the closeness of the corresponding degree 3 element. Similarly, $H^2(C^\bullet(C, D)(F, F)(\text{id}, \text{id}))$ is isomorphic to the vector space of the outer infinitesimal deformations of the strong monoidal functor F .

A relative totalisation $Rp_*A(C, D)(F, F)(\text{id}, \text{id})$ along the projection $p: \Theta_2 \rightarrow \Delta$ is defined, and it is shown to be a cosimplicial monoid, which fulfils the Batanin–Davydov 1-commutativity condition (Batanin and Davydov (2023)). Then it follows from loc. cit. that $C^\bullet(C, D)(F, F)(\text{id}, \text{id})$ is a $C_\bullet(E_2; \mathbb{k})$ -algebra. Conjecturally, $C^\bullet(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id})$ is a $C_\bullet(E_3; \mathbb{k})$ -algebra; however, the proof requires more sophisticated methods.

1. Introduction

1.1. In formal deformation theory, with a deformation functor (defined on a suitable category of commutative differential graded (dg) coalgebras), one associates a differential graded Lie algebra (or, more generally, an L_∞ algebra), whose completed chain Chevalley–Eilenberg complex pro-represents the deformation functor. In characteristic 0, such representability has been a proven statement [17, 24, 30]. Sometimes the underlying complex of the dg Lie algebra (corresp., of an L_∞ algebra) is called the deformation

complex. In many particular deformation problems, such complex is easier to find than the entire dg Lie algebra structure. The first cohomology of the deformation complex is isomorphic to the infinitesimal deformations mod out equivalences, whence the dg Lie algebra encodes, via the Maurer–Cartan equation, the formal (global) deformations mod out formal equivalences. When the “base of deformation” can be chosen as a E_n -coalgebra (in a suitable sense, e.g., a coalgebra over the dg operad $C_*(E_n; \mathbb{k})$, or a coalgebra over the Koszul resolution hoe_n of the operad $e_n = H_*(E_n; \mathbb{k})$, etc.), such that it gives rise to a deformation functor defined on a suitable category of E_n -coalgebras, the shifted by $[n]$ deformation complex has a structure of E_{n+1} -algebra, [24, 34, 36].

The higher structures on a deformation complex are important if we are interested in the formality phenomena. For instance, the Deligne conjecture asserts that the Gerstenhaber bracket on the Hochschild cohomological complex of a (dg) algebra A (which defines a dg Lie algebra structure) can be “extended” to a $C_*(E_2; \mathbb{k})$ -algebra structure (it has found several proofs; see [5, 22, 27, 28, 37]). It was used by Tamarkin [35] in his proof of (a stronger version of) the Kontsevich formality theorem [21]. The idea is roughly that the higher the structure on the deformation complex we consider, the more rigid the deformation complex with this structure becomes. Thus, the original formality theorem of Kontsevich was stated for dg Lie algebra structure on the Hochschild cohomological complex of a polynomial algebra and was proven by methods inspired by the Topological Quantum Field Theory. The idea of Tamarkin was to consider the entire higher structure of homotopy 2-algebra on the complex, and using the aforementioned rigidity, it can be proven by homotopy theoretical methods. The “transcendental part” of the proof becomes hidden in a solution to the Deligne conjecture.

1.2. In this paper, we are interested in the deformation theory of monoidal dg categories (and more generally of dg bicategories). Our interest originates in the (partly open) deformation theory of associative bialgebras. In this case, the deformation complex was constructed by Gerstenhaber and Schack [15]; its intrinsic interpretation in terms of abelian category of tetramodules over the bialgebra was given in [33]. This interpretation made it possible to compute the Gerstenhaber–Schack cohomology for the case of $B = S(V)$, the (co)free (co)commutative bialgebra. The answer was (as it had been conjectured by Kontsevich)

$$H_{\text{GS}}^k(S(V)) = \bigoplus_{a+b=k} \text{Hom}(V^{\otimes a}, V^{\otimes b}) = S^k(V^*[-1] \oplus V[-1]).$$

The symmetric algebra $H^*(B)\text{GS} = S(V^*[-1] \oplus V[-1])$ has a Poisson algebra structure of degree -2 , which comes from the degree -2 pairing $V^*[-1] \oplus V^*[-1] \rightarrow \mathbb{k}$. It gives, along with the graded commutative product, a structure of e_3 -algebra, where $e_3 = H_*(E_3; \mathbb{k})$. An interesting and important open question is how one can lift this structure to the Gerstenhaber–Schack complex for $B = S(V)$, or for general B .

Motivated by the problem of finding the higher structures on the (Gerstenhaber–Schack) deformation complex of a bialgebra B , we consider in this paper the deformation theory of a (dg or \mathbb{k} -linear) monoidal category C . In the example associated with the deformation theory of B , the monoidal category $C = \text{Mod}(B)$ is the category of left modules over

the underlying algebra B ; it is known to be a monoidal category: for $M, N \in \text{Mod}(B)$, the tensor product $M \otimes_{\mathbb{k}} N$ is a $B \otimes_{\mathbb{k}} B$ -module; then the precomposition along the coproduct map $\Delta: B \rightarrow B \otimes_{\mathbb{k}} B$ makes $M \otimes_{\mathbb{k}} N$ a B -module. The counit map $\varepsilon: B \rightarrow \mathbb{k}$ makes \mathbb{k} the unit for this monoidal structure.

The link between the deformations of $\text{Mod}(B)$ and the deformations of B is partially established by the Tannaka–Krein duality (though in what concerns the deformation complexes this link has to be understood better). An advantage of the monoidal category approach is that the higher structures on the deformation complex of a dg monoidal category are more manageable and can be explicitly found. A recent paper [6], where the authors deal with a truncation of our complexes, called the Davydov–Yetter complex of a monoidal category. In loc. cit. the authors constructed a $C.(E_3; \mathbb{k})$ -algebra structure on this truncated deformation complex. The Davydov–Yetter complex controls only the (infinitesimal) deformations of the associator, whence our complex controls the (infinitesimal) deformations of all linear data (the associator, the underlying dg category, the morphisms part of the product bifunctor, see (A1)–(A4) in Section 7).

The work in progress [32] aims to find a structure of $C.(E_3; \mathbb{k})$ -algebra on the (non-truncated) deformation complex, employing the technique of Batanin n -operads [2–4]. According to [3], an action of a contractible $(n - 1)$ -terminal (pruned and reduced) n -operad on a complex gives, via the symmetrisation functor and the cofibrant replacement, an action of the chain operad $C.(E_n; \mathbb{k})$ on it. In our opinion, the theory of n -operads provides, via the aforementioned result, a very flexible and powerful approach to higher generalisations of the Deligne conjecture. In [7], a version of Deligne conjecture for general n is stated, and it is proven in [8] for $n = 2$.

1.3. Let \mathbb{k} be a field. We consider \mathbb{k} -linear bicategories; see [20, 23]. We recall the basic definitions related to (enriched) bicategories in Section 4. A particular case of a (\mathbb{k} -linear) bicategory with a single object is a (\mathbb{k} -linear) monoidal category. For a \mathbb{k} -linear monoidal category C , we provide a complex $C^*(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id})$, whose 3rd cohomology is proven to parametrise the infinitesimal deformations of C mod out the infinitesimal equivalences (see Theorem 7.3). We also construct more general complexes. More precisely, for dg monoidal categories (resp., dg bicategories) C, D , two strong \mathbb{k} -linear monoidal (resp., strong bicategorical) functors $F, G: C \rightarrow D$, and two strong bicategorical natural transformations $\eta, \theta: F \Rightarrow G$, we provide a complex $C^*(C, D)(F, G)(\eta, \theta)$, whose 0-th cohomology is equal to the modifications $\eta \Rightarrow \theta$ (playing the role of 3-morphisms for the tricategory of bicategories, see Section 4). The entire complex $C^*(C, D)(F, G)(\eta, \theta)$ (or rather the closed elements of it) plays the role of *derived modifications*.

Thus, what we are dealing with here is a set-up for further theory one level (dimension) higher than the one developed in Tamarkin’s paper “What do dg categories form?” [37]. The work in progress [32] aims to construct a contractible Batanin 3-operad [3, 4] acting on the corresponding 3-quiver (whose underlying 2-quiver is a strict 2-category), which would provide a homotopy 3-algebra structure on $C^*(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id})$. Here, we construct the 3-quiver itself.

The complexes $C^\bullet(C, D)(F, G)(\eta, \theta)$ emerge as the totalisation¹ of 2-cocellular complexes, that is, of functors $A(C, D)(F, G)(\eta, \theta): \Theta_2 \rightarrow C^\bullet(\mathbb{k})$. Here, Θ_2 is the category dual to the category of Joyal 2-disks [10, 11, 19].

1.4. Our complex $C^\bullet(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id})$ can be thought of as a relaxed version of the Davydov–Yetter complex [13, 39] of a monoidal category, which fits better for aims of deformation theory. Recall that the Davydov–Yetter cochains in degree n are *natural* transformations from the functor M^n to itself, where

$$M^n(X_1, \dots, X_n) = X_1 \otimes (X_2 \otimes (\dots (X_{n-1} \otimes X_n) \dots)).$$

It gives rise to a cosimplicial (dg) vector space. Then, the Davydov–Yetter complex is defined as the totalisation of this cosimplicial vector space.

Often the Davydov–Yetter complex is said to compute the infinitesimal deformations of a monoidal category. In fact, it is not quite correct, where the Davydov–Yetter complex only encodes the deformations of the associator, leaving the underlying \mathbb{k} -linear category and the monoidal product *on morphisms* fixed.

In our set-up of \mathbb{k} -linear monoidal category, it is natural to deform all \mathbb{k} -linear data. More precisely, we assume that the monoidal product *on objects* remains fixed, while the underlying \mathbb{k} -linear category, the monoidal product *of morphisms*, and the associator are being deformed. We refer to such deformations as *full*.

Theorem 7.3 states that $H^3(C^\bullet(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id}))$ parametrises the infinitesimal *full* deformations of C mod out the infinitesimal equivalences.

The link between $C^\bullet(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id})$ and the Davydov–Yetter complex can be described as follows. The natural embedding $\Delta \times \Delta \rightarrow \Theta_2$ makes it possible to consider $C^\bullet(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id})$ as a bicomplex, with the horizontal differential d_0 and the vertical differential d_1 . One can show that $C_{DY}^\bullet(C)$ is the kernel of the vertical differential d_1 restricted to the 0-th row of this bicomplex (with the differential d_0).

The naturality of the Davydov–Yetter cochains is dropped in $C^\bullet(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id})$ and is replaced by the naturality with respect to monoidal structural maps. The reader is advised to look directly to Section 5.1 for more detail on the connection between the Davydov–Yetter and our complexes and on this restricted naturality for cochains.

One of our motivations here was the recent paper [6], where a $C.(E_3; \mathbb{k})$ algebra structure on the Davydov–Yetter complex $C_{DY}^\bullet(C)$ of a \mathbb{k} -linear monoidal category C was constructed. In [6], the authors consider more generally n -commutative cosimplicial monoids and prove that the totalisation of such cosimplicial monoid has a structure of homotopy $(n + 1)$ -algebra. On the other hand, it is shown in [6] that $C_{DY}^\bullet(C)$ is the totalisation of a 2-commutative cosimplicial monoid, which implies that $C_{DY}^\bullet(C)$ is a homotopy 3-algebra.

¹By Θ_2 -totalisation we mean here the corresponding (non-normalised) cochain Moore complex, as it is defined in (3.1), (3.2); see also discussion in Section 3.1.

Unfortunately, the 2-commutativity fails for our complex (even for the strict case). More precisely, there is a natural projection $p: \Theta_2 \rightarrow \Delta$, which defines a cosimplicial complex $R^* p_*(A(C, D)(F, G)(\eta, \theta))$.² For the case $F = G$, $\eta = \theta = \text{id}$, this cosimplicial complex is in turn a cosimplicial (dg) monoid. One easily shows that the Δ -totalisation of the latter monoid is equal to the Θ_2 -totalisation of $A(C, D)(F, G)(\eta, \theta)$. However, the cosimplicial monoid $Rp_*(A(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id}))$ fails to be 2-commutative. (It is, in a sense, a *homotopy 2-commutative monoid*, a cosimplicial monoid in which the 2-commutativity relation holds only up to homotopy, in a suitable sense, which conjecturally should be enough for its totalisation to be a homotopy 3-algebra.)³

On the other hand, for a strong bicategorical \mathbb{k} -linear functor $F: C \rightarrow D$, the cosimplicial monoid $R^* p_* A(C, D)(F, F)(\text{id}, \text{id})$ is 1-commutative (Proposition 6.4). Then it follows from [6, Cor. 2.46] that the complex $C^*(C, D)(F, F)(\text{id}, \text{id})$ is a homotopy 2-algebra (Theorem 6.6).

1.5. Organisation of the paper

The paper consists of 6 sections and 1 appendix.

In Section 2, we recall definitions and well-known results on the categories Θ_n , Joyal n -disks, and their interplay. None of the results of Section 2 is new; we basically follow [10, 11, 19].

In Section 3, we define the totalisation of a 2-cocellular complex (that is, of a functor $\Theta_2 \rightarrow C^*(\mathbb{k})$), as well as the relative totalisation along the projection $p: \Theta_2 \rightarrow \Delta$. For $X: \Theta_2 \rightarrow C^*(\mathbb{k})$, the relative totalisation $Rp_*(X)$ is a functor $\Delta \rightarrow C^*(\mathbb{k})$. In Proposition 3.7, we prove, for any X as above, the transitivity property for its totalisations:

$$\text{Tot}_{\Theta_2}(X) = \text{Tot}_{\Delta}(Rp_*(X)).$$

In Section 4, we recall the basic notions related to bicategories and introduce a (seemingly, new) concept of a 2-bimodule over a bicategory. We introduce a “bicategorical” version $\widehat{\Theta}_2$ of the category Θ_2 and study the left Kan extension along the projection $\widehat{\Theta}_2^{\text{op}} \rightarrow \Theta_2^{\text{op}}$.

In Section 5, we introduce our main new construction, the 2-cocellular complexes $A(C, D)(F, G)(\eta, \theta)$. The definition is more tricky than one could expect; namely, the components $A(C, D)(F, G)(\eta, \theta)_T$ are subcomplexes of the corresponding components of a “more natural” complex $\widehat{A}(C, D)(F, G)(\eta, \theta)_T$. The passage from \widehat{A} to A is performed by imposing the *bicategorical relations* (5.4)–(5.6) and taking the corresponding subspaces; without that, the assignment $T \rightsquigarrow \widehat{A}(C, D)(F, G)(\eta, \theta)_T$ itself fails to be 2-cocellular.

²One can not state that $R^* p_*(A(C, D)(F, G)(\eta, \theta))$ is the right homotopy Kan extension because $A(C, D)(F, G)(\eta, \theta)$ fails to be Reedy fibrant. One can alternatively define $R^* p_*(\dots)$ as “relative totalisation”; see Section 3.1.

³It would be interesting to define the concept of a cosimplicial homotopy n -commutative monoid; its totalisation should be an algebra over $C^*(E_{n+1}; \mathbb{k})$.

In Section 6, we employ the results of [6] for studying higher structures on the complexes $C^*(C, D)(F, G)(\eta, \theta)$. It is possible due to the transitivity property of Proposition 3.7. We prove that $C^*(C, D)(F, F)(\text{id}, \text{id})$ is a homotopy 2-algebra, for any strong \mathbb{k} -linear 2-functor, in Theorem 6.6.

Section 7 contains an identification of $H^3(C^*(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id}))$ with infinitesimal full deformations of C mod out infinitesimal equivalences. It justifies our complexes $C^*(C, D)(F, G)(\eta, \theta)$ as related to the deformation theory of monoidal \mathbb{k} -linear categories (and, more generally, of \mathbb{k} -linear bicategories).

In the appendix, we list the relations between (co)dimension 1 operators in Θ_2 , used throughout the paper.

2. The categories Θ_n

Here, we recall the definition of the categories $\Theta_n, n \geq 1$, and some related concepts.

2.1. n -ordinals and n -leveled trees

We denote by $[n]$ the ordinal $0 < 1 < \dots < n$ having $n + 1$ elements. Recall that the simplicial category Δ has objects $[n], n \geq 0$, that is, all non-empty finite ordinals. Its morphisms are the monotonous maps $f: [k] \rightarrow [\ell]$, that is, $f(i) \leq f(j)$ if $i \leq j$.

Recall the relations between the standard elementary face operators $\partial^i: [n - 1] \rightarrow [n]$ and the elementary degeneracy operators $\varepsilon^i: [n + 1] \rightarrow [n], i = 0, \dots, n$, in Δ :

$$\begin{aligned} \partial^j \partial^i &= \partial^i \partial^{j-1} && \text{if } i < j \\ \varepsilon^j \varepsilon^i &= \varepsilon^i \varepsilon^{j+1} && \text{if } i \leq j \\ \varepsilon^j \partial^i &= \begin{cases} \partial^i \varepsilon^{j-1} & \text{if } i < j \\ \text{id} & \text{if } i = j, j + 1 \\ \partial^{i-1} \varepsilon^j & \text{if } i > j + 1. \end{cases} \end{aligned} \tag{2.1}$$

An ordinal as above is also called a 1-ordinal. The following definition is due to M. Batanin.

Definition 2.1. An n -ordinal S is a sequence of surjective maps in Δ :

$$[k_n] \xrightarrow{\rho_{n-1}} [k_{n-1}] \xrightarrow{\rho_{n-2}} \dots \xrightarrow{\rho_1} [k_1] \xrightarrow{\rho_0} [0]. \tag{2.2}$$

The category Ord_n of n -ordinals has all n -ordinals as its objects, and the morphisms $S \rightarrow T$ are commutative diagrams

$$\begin{array}{ccccccc} [k_n] & \xrightarrow{\rho_{n-1}} & [k_{n-1}] & \xrightarrow{\rho_{n-2}} & \dots & \xrightarrow{\rho_1} & [k_1] & \xrightarrow{\rho_0} & [0] \\ \downarrow f_n & & \downarrow f_{n-1} & & & & \downarrow f_1 & & \downarrow \text{id} \\ [\ell_n] & \xrightarrow{\rho'_{n-1}} & [\ell_{n-1}] & \xrightarrow{\rho'_{n-2}} & \dots & \xrightarrow{\rho'_1} & [\ell_1] & \xrightarrow{\rho'_0} & [0] \end{array} \tag{2.3}$$

in which f_1, \dots, f_n are not necessarily maps in Δ , but a *weaker condition* holds: for any $a \in [k_j]$ the restriction of f_{j+1} to $\rho_j^{-1}(a)$ is order-preserving.

Remark: note that f_j 's are not necessarily morphisms in Δ ; the above condition is weaker than order-preserving.

An object of Ord_n is a non-empty n -ordinal.

When the assumption that the maps $\{\rho_i\}$ are surjective is dropped, the object (2.2) is called an n -level tree (or, shortly, an n -tree). A morphism of n -trees is defined as in (2.3).

The difference between the n -ordinals and the n -level trees is that the former have all input vertices at the top n -th level, whence an n -tree may have input vertices at all levels. Sometimes n -ordinals are called *pruned* n -trees.

Introduce some terminology related to leveled trees, which is used later in the paper.

We represent n -level trees as a collection of finite sets $\{T(i)\}_{0 \leq i \leq n}$ endowed with a map $i_T: T_{\geq 1} \rightarrow T$ which lowers the level by 1. The map i_T is defined as ρ_{i-1} at level i .

For $x \in T(i)$ we write $\text{ht}(x) = i$. By definition, $n = \text{ht}(T) = \max_{x \in T} \text{ht}(x)$. A vertex x of a leveled tree is called an *input*, or a *leaf*, if $i_T^{-1}(x) = \emptyset$. Note that for an n -leveled tree, the height of an input may be smaller than n (but there always exists an input of the height n).

An *edge* is a pair (x, y) with $x = i_T(y)$. The set of edges of T is denoted by $e(T)$. We define the *dimension* $d(T) = \#e(T)$. A leveled tree is called *linear* if $d(T) = \text{ht}(T)$.

For each vertex $x \in T$, the ordered set of incoming edges $e_x(T)$ is defined as $i_T^{-1}(x)$.

For a leveled tree T define a leveled tree \bar{T} as follows. For each $x \in T$, we set $e_x(\bar{T}) = e_x(T) \cup (x, x_-) \cup (x, x_+)$ with the order in which (x, x_-) is the minimal element and (x, x_+) is the maximal element. Thus, we add the leftmost and the rightmost element to each set $e_x(T)$. It results in $\bar{T}(i) = T(i) + 2T(i - 1)$, and $\text{ht}(\bar{T}) = \text{ht}(T) + 1$. A T -sector of height k is a triple $(x; y_L, y_R)$ where $x \in T(k)$, $y_L, y_R \in \bar{T}(k + 1)$, $i_{\bar{T}}(y_L) = i_{\bar{T}}(y_R) = x$, and y_L, y_R are consecutive elements of $\bar{T}(k + 1)$. We say that x supports a sector $(x; y_L, y_R)$. It follows that each input vertex x of T supports a unique sector (which is $(x; x_-, x_+)$).

2.2. The wreath product definition of Θ_n

The definition of the categories Θ_n , $n \geq 1$, is given inductively via the *wreath product* $\Delta \wr \mathcal{A}$; see below.

Definition 2.2. Let \mathcal{A} be a category. The objects of the category $\Delta \wr \mathcal{A}$ are tuples $([\ell]; A_1, \dots, A_\ell)$, where $A_1, \dots, A_\ell \in \mathcal{A}$. A morphism $\Phi: ([\ell]; A_1, \dots, A_\ell) \rightarrow ([m]; B_1, \dots, B_m)$ is a tuple $(\phi; \phi_1, \dots, \phi_\ell)$ where $\phi: [\ell] \rightarrow [m]$ is a morphism in Δ , and

$$\phi_i = (\phi_i^{\phi(i-1)+1}, \dots, \phi_i^{\phi(i)})$$

is a tuple of morphisms in \mathcal{A} , with $\phi_i^k: A_i \rightarrow B_k$. The composition is defined in the natural way.

The reader is advised to look at Lemma 2.9 which explains a natural framework in which the category $\Delta \wr \mathcal{A}$ emerges.

We set

$$\Theta_1 = \Delta \quad \text{and} \quad \Theta_n = \Delta \wr \Theta_{n-1}, \quad n \geq 2. \tag{2.4}$$

Remark 2.3. The case $\ell = 0$ is allowed for an object of Θ_n . In this case, the object $([0]; \emptyset)$ is final.

2.3. n -globular sets and strict n -categories

There is another category equivalent to the category Θ_n .

Recall that an n -globular set is the data one has on the underlying sets of objects, 1-morphisms, \dots , n -morphisms of a strict n -category. In this sense, it is a “pre- n -category”. For $n = 1$, it is a quiver.

The general definition is as follows.

Definition 2.4. An n -globular set is a collection of sets X_0, X_1, \dots, X_n and maps

$$X_n \begin{array}{c} \xrightarrow{s_{n-1}} \\ \xrightarrow{t_{n-1}} \end{array} X_{n-1} \begin{array}{c} \xrightarrow{s_{n-2}} \\ \xrightarrow{t_{n-2}} \end{array} \cdots X_1 \begin{array}{c} \xrightarrow{s_0} \\ \xrightarrow{t_0} \end{array} X_0$$

(here s_k are source maps and t_k are target maps), such that

$$s_k s_{k+1} = s_k t_{k+1}, \quad t_k s_{k+1} = t_k t_{k+1}, \quad 0 \leq k \leq n - 1.$$

For two n -globular sets X, Y , a morphism $f: X \rightarrow Y$ is defined as a sequence of maps $f_i: X_i \rightarrow Y_i, 0 \leq i \leq n$, which commute with the source and the target maps s and t .

The category of n -globular sets is denoted by Glob_n . The reader easily interprets the category Glob_n as some presheaf category.

The following question arises: *how can one define the free strict n -category generated by an n -globular set?* More precisely, the question is in defining the left adjoint functor ω_n to the forgetful functor $R: \text{Cat}_n \rightarrow \text{Glob}_n$. (The $n = 1$ case is the well-known construction of the free category generated by a quiver.)

The construction of M. Batanin [2, Sect. 4], which associates an n -globular set T^* with an n -ordinal T , is served to solve this problem.

We recall this construction, following a more explicit treatment given in [10, Lem. 1.2].

Lemma 2.5. *Let T be an n -leveled tree, and denote by T_k^* the set of all sectors of T of height $k, 0 \leq k \leq n$. Then, T^* is an n -globular set.*

Proof. Let $(x; y_L, y_R) \in T_k^*$. We have to define $s_{k-1}(x; y_L, y_R)$ and $t_{k-1}(x; y_L, y_R)$. Let x_L, x, x_R be the three consecutive elements in $\bar{T}(k)$. Define

$$s_{k-1}(x; y_L, y_R) = (i_T(x); x_L, x) \quad \text{and} \quad t_{k-1}(x; y_L, y_R) = (i_T(x); x, x_R).$$

One easily sees that the globular identities hold; see [10, Lem. 1.2] for more detail. ■

Example 2.6. Some examples for 2-level trees are shown in Figure 1.

Example 2.7. For the 2-level tree T having n vertices at level 1 with the preimages having ℓ_1, \dots, ℓ_n vertices at level 2, the 2-globular set T^* is schematically shown in Figure 2.




Figure 1. Examples of 2-level trees.




Figure 2. The 2-globular set T^* for T from Example 2.7.

With the help of the T^* construction, one can define the left adjoint $\omega_n: \text{Glob}_n \rightarrow \text{Cat}_n$ to the forgetful functor $R: \text{Cat}_n \rightarrow \text{Glob}_n$, as follows.

Let X be an n -globular set. We define an n -globular set $\omega_n(X)$ and prove that it is a strict n -category.

Set

$$(\omega_n(X))_k = \coprod_{T: \text{ht}(T) \leq k} \text{Hom}_{\text{Glob}_n}(T^*, X) \tag{2.5}$$

(one often uses the notation $\text{Hom}_{\text{Glob}_n}(T^*, X) = X^T$).

First of all, we show that $\omega_n(X)$ is an n -globular set.

Denote by $\partial_k T$ the $(k - 1)$ -leveled tree, obtained by removing all vertices of height higher than $k - 1$. There are two maps of n -globular sets $s_{k-1}^*, t_{k-1}^*: (\partial_k T)^* \rightarrow (\partial_{k+1} T)^*$. In general, a map of globular sets $S^* \rightarrow T^*$ is determined by its restriction to the input sectors of S^* ; see [10, Lem. 1.3]. The map s_{k-1}^* (corresp., t_{k-1}^*) is obtained by assigning to each input vertex x of $\partial_k T$ (which uniquely defined its input sector) the leftmost (resp., rightmost) input sector in $\partial_{k+1} T$ supported by x . One shows that the maps s_{k-1}^*, t_{k-1}^* satisfy the identities dual to the globular identities. Thus, for any n -globular set X , and for a k -leveled tree T , $k \leq n$, the precompositions with the maps s_{k-1}^*, t_{k-1}^* define maps

$$s_{k-1}, t_{k-1}: X^T \rightarrow X^{\partial_k T}.$$

It follows that these maps satisfy the globular identities. Thus, $\omega_n(X)$ is a globular set.

Next, prove that $\omega_n(X)$ is a strict n -category.

The following statement is proven in [11, Thm. 3.7].

Proposition 2.8. *For any n -ordinals S, T , one has*

$$\Theta_n(S, T) = \text{Cat}_n(\omega_n(\bar{S}^*), \omega_n(\bar{T}^*)). \quad \blacksquare$$

The proof is obtained, by induction, from the following nice interpretation of the wreath product, [11, Prop. 3.5].

Lemma 2.9. *Let a small category \mathcal{A} be a full subcategory of a cocomplete cartesian monoidal category \mathcal{V} . Then, $\Delta \wr \mathcal{A}$ is a full subcategory of $\mathcal{V}\text{-Cat}$.*

Proof. Any 1-ordinal $[n]$ can be considered a linear category \mathbf{n} with $n + 1$ objects $0, \dots, n$, with a single morphism in $\mathbf{n}(i, j)$ for $i \leq j$ and with empty set of morphisms otherwise. Having n objects A_1, \dots, A_n of \mathcal{A} , we regard them as objects of \mathcal{V} , and consider the linear \mathcal{V} -quiver:

$$0 \xrightarrow{A_1} 1 \xrightarrow{A_2} 2 \xrightarrow{A_3} \dots \xrightarrow{A_n} n.$$

Consider the \mathcal{V} -category generated by this quiver, and denote it by $F_{\mathcal{V}}(A_1, \dots, A_n)$ (here we use cocompleteness of \mathcal{V} to show that the forgetful functor from \mathcal{V} -categories to \mathcal{V} -quivers has a left adjoint).

For $B_1, \dots, B_m \in \mathcal{A}$, a \mathcal{V} -functor $\Phi: F_{\mathcal{V}}(A_1, \dots, A_n) \rightarrow F_{\mathcal{V}}(B_1, \dots, B_m)$ is defined by its restriction to “generators”, that is, by a map $\phi: [n] \rightarrow [m]$, and, for any $1 \leq i \leq n$, a morphism

$$A_i \rightarrow F_{\mathcal{V}}(\phi(i - 1), \phi(i)) = B_{\phi(i-1)+1} \times \dots \times B_{\phi(i)}.$$

We conclude that these \mathcal{V} -functors are the same as the morphisms

$$([n], A_1, \dots, A_n) \rightarrow ([m], B_1, \dots, B_m)$$

in $\Delta \wr \mathcal{A}$. \blacksquare

Example 2.10. For the case $\Theta_2 = \Delta \wr \Delta$, we set $\mathcal{V} = \text{Cat}$, using the imbedding $\Delta \rightarrow \text{Cat}$, $[n] \rightsquigarrow \mathbf{n}$. Thus, to the element $([n], [\ell_1], \dots, [\ell_n])$ is associated the 2-category generated by the following 2-globular set (see Figure 2).

Of course, this globular set is T^* , where T is the corresponding 2-ordinal $[\ell_1 + \dots + \ell_n + n - 1] \rightarrow [n - 1]$.

2.4. Disks

The category of disks was introduced in [19]. An *interval* is a finite ordinal, and a *map of intervals* is a map in Δ preserving the leftmost and the rightmost elements. The category of the non-empty intervals is denoted by Δ_f . Joyal (loc. cit.) showed that $\Delta_f^{\text{op}} \simeq \Delta_+$ where Δ_+ is the category of *all* finite ordinals (including the empty ordinal which is the initial object, we denote it by $[-1]$). The functor $F: \Delta_+^{\text{op}} \rightarrow \Delta_f$ is $[n] \mapsto \Delta_+([n], [1])$, $F([n]) = [n + 1]$. The dual functor $G: \Delta_f^{\text{op}} \rightarrow \Delta_+$ is $[n] \mapsto \Delta_f([n], [1])$, then the initial object $[-1]$ is $\Delta_f([0], [1])$, and in general $G([n]) = [n - 1]$.

Definition 2.11. A *disk of finite sets* D_\bullet is a sequence D_1, D_2, \dots of finite sets, equipped with the following data:

- (a) a map $p: D_k \rightarrow D_{k-1}$ such that for any $x \in D_{k-1}$ the pre-image $p^{-1}(x)$ has an interval structure, $k \geq 1$,
- (b) two maps $d_0, d_1: D_{k-1} \rightarrow D_k$ sending $x \in D_{k-1}$ to the leftmost and the rightmost elements of the interval $p^{-1}(x)$, $k \geq 1$,
- (c) for $k \geq 1$, the diagram

$$d_0(D_{k-1}) \cup d_1(D_{k-1}) \rightarrow D_k \begin{array}{c} \xrightarrow{d_0} \\ \xrightarrow{d_1} \end{array} D_{k+1}$$

is an equaliser, where the first arrow is the canonical embedding,

- (d) D_0 is a single point.

A map of two disks $F: D_\bullet \rightarrow D'_\bullet$ is a collection of maps $\{F_k: D_k \rightarrow D'_k\}_{k \geq 0}$ compatible with p, d_0, d_1 , such that for any $x \in D_k$ the map $p^{-1}(x) \rightarrow p^{-1}(F_k(x))$ is a map of intervals, $k \geq 0$.

The category of disks is denoted by Disk .

For a disk D_\bullet , the *interior* $i(D_k)$ is defined as $D_k \setminus \{d_0(D_{k-1}) \cup d_1(D_{k-1})\}$. It is an ordinal, and the sequence of maps of ordinals $p: i(D_k) \rightarrow i(D_{k-1})$, $k \geq 1$ makes $i(D_\bullet) = \{i(D_k)\}_{k \geq 0}$ a *leveled tree*. The height $\text{ht}(D_\bullet)$ is defined as the height of the level tree $i(D_\bullet)$. The category of disks of height $\leq n$ is denoted by Disk_n .

The functor i sends disks to leveled trees. The functor $T \mapsto \bar{T}$ is a left adjoint to it. For any leveled tree T , the leveled tree \bar{T} is a disk of finite sets. The elements of \bar{T} in the image of i are *internal*, and the elements in $\bar{T} \setminus T$ are *boundary*.

A map of disks $\bar{S} \rightarrow \bar{T}$ is “more general” than a map of leveled trees $S \rightarrow T$. The reason is that a map of disks $\bar{S} \rightarrow \bar{T}$ may map an internal point to a boundary point in \bar{T} . Thus, the category Ord_n is identified with a not full subcategory of Disk_n .

The following proposition is [10, Prop. 2.2].

Proposition 2.12. For any n -leveled trees S, T , one has

$$\text{Cat}_n(\omega_n(\bar{S}^*), \omega_n(\bar{T}^*)) = \text{Disk}_n(\bar{T}, \bar{S}).$$

Thus, the assignment $T \mapsto \bar{T}$ provides an equivalence of Θ_n^{op} and Disk_n . ■

Remark 2.13. We can restrict the assignment from the proof [10, Prop. 2.2] to the maps of disks $\bar{S} \rightarrow \bar{T}$ which come from maps of leveled trees $S \rightarrow T$ (that is, which map internal points to internal). The corresponding sub-category C of Cat_n has objects $\omega_n(\bar{T}^*)$, $T \in \text{Ord}_n$, and has the set of morphisms $C(\omega_n(\bar{S}^*), \omega_n(\bar{T}^*))$ which is the subset of $\text{Cat}_n(\omega_n(\bar{S}^*), \omega_n(\bar{T}^*))$ formed by maps of n -categories, preserving minima and maxima, in an appropriate sense. For $n = 2$, this equivalence is used by Tamarkin [37].

In fact, this equivalence (rather than the equivalence of Proposition 2.12) can be thought of as a proper analogue of the Joyal equivalence $\Delta_f \simeq \Delta_+^{\text{op}}$, for $n \geq 2$.

2.5. The categories Θ_n as higher analogues of the category Δ : inner and outer face maps

We have three equivalent descriptions of the category Θ_n which are

- (a) the definition via the wreath product (2.4),
- (b) the definition via morphisms of free strict n categories $\omega_n(\bar{T}^*)$, Proposition 2.8,
- (c) the dual of the category Disk_n , Proposition 2.12.

We will take advantage of all three equivalences. In particular, (c) is used to naturally define the realisation/totalisation, (b) is used to see that any strict n category C has a nerve which is a cocellular set $N(C): \Theta_n^{\text{op}} \rightarrow \text{Sets}$, and (a) is the most combinatorially explicit and manageable.

The existence of the nerve was the main motivation in [19], where the disk categories were defined. It also makes it possible to consider Θ_n as an analogue of Δ , for $n \geq 2$.

Note that the nerve $N(C)$ of the ordinary category C is a simplicial set, whose components can be defined as

$$N(C)_k = \text{Cat}(\mathbf{k}, C)$$

(where k is the linear category with $k + 1$ objects). We see directly that it gives rise to a simplicial set because a map $[k] \rightarrow [m]$ in Δ amounts to the same thing as a map of the linear categories $\mathbf{k} \rightarrow \mathbf{m}$.

Let now C be a strict n -category. Define its n -nerve as a cellular set

$$N(C): \Theta_n^{\text{op}} \rightarrow \text{Sets},$$

for which

$$N(C)_T = \text{Cat}_n(\omega_n(\bar{T}^*), C). \tag{2.6}$$

It gives rise to an n -cellular set because, by Proposition 2.8,

$$\Theta_n(S, T) = \text{Cat}_n(\omega_n(\bar{S}^*), \omega_n(\bar{T}^*)).$$

For any strict n -category C , the n -cellular set $N(T)$ has a property which is a higher n counterpart of the Boardman–Vogt inner horns filling property for $n = 1$, called the “weak Kan complexes”. The simplicial sets with inner horns filling condition were further studied by Joyal (under the name quasi-categories) and Lurie (under the name $(\infty, 1)$ -categories), as a model for weak analogues of ordinary categories. The aim in [19] was to define a model for weak analogues of strict n -categories, and it was the motivation for introducing the categories Θ_n .

In Θ_n , there are two classes of maps, *face maps* and *degeneracy maps*, and face maps are further subdivided to *outer face maps* and *inner face maps*.

The most direct way to define them is by using the category of disks Disk_n ; see Proposition 2.12.

Let S, T be two n -leveled trees.

A map $\bar{S} \rightarrow \bar{T}$ in Disk_n is called a *degeneracy* if it is an embedding on each interval $p^{-1}(x)$, and $|S| + 1 = |T|$.

A map $f: \bar{S} \rightarrow \bar{T}$ is called an *inner face map* if $|S| = |T| + 1$, f contracts two neighbour *inner* points a, b at some interval $p^{-1}(x)$ and is a *shuffling* map on the interiors of the intervals $p^{-1}(a)$ and $p^{-1}(b)$ (when such shuffling is fixed, the map of disks with these conditions is uniquely defined).

Let $x \in S$, $a \in p^{-1}(x)$, $a \in i(S)$ be an extreme (= leftmost or rightmost) vertex of $p^{-1}(x)$ in $i(S)$. We call a a *special extreme vertex* if a is also an input vertex of $i(S)$. Let $f: \bar{S} \rightarrow \bar{T}$ be a map in Disk_n , $|S| = |T| + 1$, a is a special extreme vertex in S , and b is a vertex in $\partial p^{-1}(x) \subset \bar{S}$ (left or right) neighbour to a (the vertex b is unique except for the case when $p^{-1}(x) \subset \bar{S}$ consists of 3 vertices, two of which are boundary). We call f the *outer face map* associated with (a, b) as above if $T = S \setminus \{a\}$ and f maps a to b .

It is clear that any surjective map of codimension 1 is either an inner or outer face map.

Remark 2.14. The inner (resp., all) face maps are used to define *inner horns* (resp., *all horns*) in [19, Def. 2] and to define *weak n -categories* (resp., *weak n -groupoids*) as cellular sets $X: \Theta_n \rightarrow \text{Sets}$ with inner (resp., all) horns filling property. This idea was further elaborated in [1, 10, 11].

2.6. The Reedy structure on Θ_2 , description of coface and codegeneracy maps

One can translate the above definition of coface and codegeneracy maps to the wreath product definition of Θ_2 . For $D = ([k]; [n_1], \dots, [n_k]) \in \Theta_2$, define *dimension* of D as

$$|D| = k + n_1 + \dots + n_k. \tag{2.7}$$

It was proven in [10, Lem. 2.4 (a)] that Θ_2 is a Reedy category, in which *degree* is equal to the dimension (2.7), and there are two classes of morphisms, coface maps and codegeneracy maps, which raise (corresp., lower) the degree. The construction was clarified in [12], using the wreath product definition. (In [10, 12], a Reedy structure on Θ_n for general $n \geq 1$ is defined.)

Recall following [12] the Reedy category structure on Θ_2 .

Recall that an object of Θ_2 is given by a tuple $([k]; [n_1], \dots, [n_k])$, a morphism $\Phi: ([n]; [\ell_1], \dots, [\ell_n]) \rightarrow ([m]; [k_1], \dots, [k_m])$ is $(\phi; \phi_1, \dots, \phi_n)$, where $\phi: [n] \rightarrow [m]$ is a morphism in Δ , and $\phi_i = (\phi_i^{\phi(i-1)+1}, \dots, \phi_i^{\phi(i)})$, $\phi_i^s: [\ell_i] \rightarrow [k_s]$ is a tuple of morphisms in Δ .

Define two subcategories $\Theta_2^-, \Theta_2^+ \subset \Theta_2$, such that $\text{Ob } \Theta_2^- = \text{Ob } \Theta_2^+ = \text{Ob } \Theta_2$. We say that $\Phi \in \Theta_2^-$ if ϕ is surjective, and for $\phi(i-1) < \phi(i)$ the map $\phi_i^{\phi(i)}: [k_i] \rightarrow [\ell_{\phi(i)}]$ is surjective. We say that a map $\Phi \in \Theta_2^+$ if ϕ is injective, and for any i the family of maps $\{\phi_i^j: [k_i] \rightarrow [\ell_j]\}_{\phi(i-1)+1 \leq j \leq \phi(i)}$ is *jointly* injective; that is, for any $a, a+1 \in [k_i]$ there is j such that $\phi_i^j(a) \neq \phi_i^j(a+1)$ (note that individual ϕ_i^j may not be injective for all j).

The following statement is a particular case of [12, Prop. 2.11].

Any $\Phi \in \Theta_2$ can be uniquely decomposed as $\Phi = \alpha^+ \circ \alpha^-$ with $\alpha_+ \in \Theta_2^+, \alpha_- \in \Theta_2^-$. One has $\Phi \in \Theta_2^+ \cap \Theta_2^-$ if $\Phi = \text{id}$, the morphisms in Θ_2^- decrease $|D|$, and the morphisms in Θ_2^+ raise $|D|$.

Below we list the $\text{codim} = \pm 1$ (with respect to $|-|$) coface and codegeneracy maps in the wreath product model of Θ_2 .

We denote by ∂^j the j -th coface maps $\partial^j: [n] \rightarrow [n + 1]$ in Δ , $0 \leq j \leq n + 1$.

Inner coface maps of codimension 1.

(F1) $n = m$, $\ell_i = k_i$ for $i \neq p$, $k_p = \ell_p + 1$, all $\phi_i^s = \text{id}$ except for $\phi_p^{\phi(p)}$ equal to the j -th coface map

$$\partial^j: [\ell_p] \rightarrow [\ell_p + 1], \quad j \neq 0, \ell_p + 1$$

(that is, ∂^j is an inner coface map in Δ). We denote this coface map ∂_p^j .

(F2) $m = n + 1$, the morphism $\phi: [m] = [n] \rightarrow [n + 1]$ is ∂^j , $j \neq 0, n + 1$. Next, $k_s = \ell_s$ except for $s = j, j + 1$, $k_j + k_{j+1} = \ell_j$, and all $\phi_s = \text{id}$ except for $s = j$. Let σ be a (k_j, k_{j+1}) -shuffle permutation in Σ_{ℓ_j} . The permutation σ defines two maps $p: [k_j - 1] \rightarrow [\ell_j - 1]$ and $q: [k_{j+1} - 1] \rightarrow [\ell_j - 1]$ in Δ . They define the Joyal dual maps $p^*: [\ell_j] \rightarrow [k_j]$ and $q^*: [\ell_j] \rightarrow [k_{j+1}]$ in Δ preserving the end-points. Then, $(p^*, q^*): [\ell_j] \rightarrow [k_j] \times [k_{j+1}]$, extended by the identity maps of the ordinals $[\ell_i]$, $i \neq j$, defines a map in Θ_2 . It is the $\text{codim} = 1$ coface map associated with a shuffle permutation σ .

We denote this coface map by $D_{j,\sigma}$.

Let us define p^*, q^* explicitly, unwinding the definition. We think of the sets $\{1, \dots, k_j\}$, $\{1, \dots, k_{j+1}\}$, $\{1, \dots, \ell_j\}$ as the elementary *arrows* in the interval categories I_{k_j} , $I_{k_{j+1}}$, and I_{ℓ_j} , correspondingly. Then, p^* and q^* are defined as follows. Both p^* and q^* preserve the end-points. If $\sigma^{-1}(\overrightarrow{i, i + 1}) = a, a + 1 \in I_{k_j}$, then

$$p^*(i) = a, \quad p^*(i + 1) = a + 1, \quad q^*(i) = q^*(i + 1).$$

If $\sigma^{-1}(\overrightarrow{i, i + 1}) = \overrightarrow{b, b + 1} \in I_{k_{j+1}}$, then

$$q^*(i) = b, \quad q^*(i + 1) = b + 1, \quad p^*(i) = p^*(i + 1).$$

Outer coface maps of codimension 1.

(F3) $n = m$, $\ell_i = k_i$ for $i \neq p$, $k_p = \ell_p + 1$, all $\phi_i^s = \text{id}$ except for $\phi_p^{\phi(p)}$ equal to the j -th coface map $\partial^j: [\ell_p] \rightarrow [\ell_p + 1]$, $j = 0, \ell_p + 1$ (that is, ∂^j is an outer coface map in Δ). We denote this coface map ∂_p^j .

(F4) The two remaining $\text{codim} 1$ coface maps are D_{\min} and D_{\max} . In both cases, $m = n + 1$. For the case of D_{\min} , $\phi = \partial^0$, and $k_1 = 0$, $k_s = \ell_{s-1}$ for $s \geq 1$, the maps $\phi_i = (\phi_i^{i+1}) = (\text{id})$. For the case of D_{\max} , $k_{m+1} = 0$, $\phi = \partial^{n+1}$, $\phi_i = (\phi_i^i) = (\text{id})$.

More generally, we call a map

$$\Phi: ([n]; [\ell_1], \dots, [\ell_n]) \rightarrow ([m]; [k_1], \dots, [k_m]), \quad \Phi = (\phi; \phi_1, \dots, \phi_n),$$

a *coface map* if $\phi: [n] \rightarrow [m]$ is injective, and each $\phi_i: [\ell_i] \rightarrow [k_{\phi(i-1)+1}] \times \dots \times [k_{\phi(i)}]$ is a (jointly) injective map (the latter means that for any $a, b \in [\ell_i]$, $a \neq b$, for at least one ϕ_i^s , $\phi(i - 1) + 1 \leq s \leq \phi(i)$, one has $\phi_i^s(a) \neq \phi_i^s(b)$).

Here is the list of elementary codegeneracy maps in Θ_2 :

(D1) $n = m$, $\ell_i = k_i$ for $i \neq p$, $k_p = \ell_p - 1$, all $\phi_i^s = \text{id}$ except for $\phi_p^{\phi(p)}$ equal to the j -th codegeneracy map $\varepsilon^j: [\ell_p] \rightarrow [\ell_p - 1]$.

We denote this codegeneracy map by ε_p^j .

(D2) $n - 1 = m$, the first component $p(\Phi)$ is $\varepsilon^p: [n] \rightarrow [n - 1]$. For any $[\ell_{p+1}]$, it extends uniquely to a morphism

$$\begin{aligned} \Phi: ([n]; [\ell_1], \dots, [\ell_p], [\ell_{p+1}], [\ell_{p+2}], \dots, [\ell_n]) \\ \rightarrow ([n - 1]; [\ell_1], \dots, [\ell_p], [\ell_{p+2}], \dots, [\ell_n]) \end{aligned}$$

for which $\phi_1, \dots, \phi_p, \phi_{p+2}, \dots, \phi_n$ are identity maps.

We denote this operator $\Upsilon_{\ell_{p+1}}^p$. Note that the morphism $\Upsilon_{\ell_{p+1}}^p$ is of codimension 1 iff $\ell_{p+1} = 0$. We define $\Upsilon_\ell^p = 0$ if $\ell \neq \ell_{p+1}$.

One can show that the morphisms in Θ_2 listed above form a set of generators for Θ_2 . For relations between these generators, see the appendix.

3. The totalisation of a 2-cocellular vector space

Here, we define the non-normalised Θ_2 -totalisation of a 2-cocellular complex. Also, we define a relative p -totalisation Rp_* , for the functor $p: \Theta_2 \rightarrow \Delta$, and prove the transitivity property saying that $\text{Tot}_\Delta \circ Rp_* = \text{Tot}_{\Theta_2}$.

3.1. Generalities on realisations and totalisations

Recall that for a general category Ξ and a functor $C: \Xi \rightarrow C^*(\mathbb{k})$, the corresponding *realisation* in $C^*(\mathbb{k})$ is a functor $\text{Sets}^{\Xi^{\text{op}}} \rightarrow C^*(\mathbb{k})$ defined as the left Kan extension of C along the Yoneda embedding $\Xi \rightarrow \text{Sets}^{\Xi^{\text{op}}}$. That is, the realisation depends on a functor $C: \Xi \rightarrow C^*(\mathbb{k})$. For $X \in \text{Sets}^{\Xi^{\text{op}}}$, we denote by $|X|_C$ or just $|C|$ its realisation with respect to the functor C . Dually, for the same Ξ and C , define the *totalisation* functor $\text{Sets}^\Xi \rightarrow C^*(\mathbb{k})$ as the right Kan extension of $C^{\text{op}}: \Xi^{\text{op}} \rightarrow C^*(\mathbb{k})^{\text{op}}$ by the dual Yoneda functor $\Xi^{\text{op}} \rightarrow (\text{Sets}^\Xi)^{\text{op}}$. The result is a functor $\text{Sets}^\Xi \rightarrow C^*(\mathbb{k})$. For $Y \in \text{Sets}^\Xi$, we denote by $\text{Tot}_C(Y)$ or $\text{Tot}(Y)$ its totalisation.

Remark 3.1. One similarly defines the realisation (resp., the totalisation) with values in any cocomplete (resp., complete) category E (replacing the category $C^*(\mathbb{k})$ in the above definition) out of a functor $C: \Xi \rightarrow E$. For example, for $\Xi = \Delta$ one can take $E = \mathcal{T}\text{op}$, $\Xi([n]) = \Delta^n$, or $E = \mathcal{C}\text{at}$, $\Xi([n]) = I_n$ (where I_n is the linearly ordered poset with $n + 1$ objects), etc. In the case $\Xi = \Delta$, $E = C^*(\mathbb{k})$, one can take $C([n]) = N(\mathbb{k}\Delta(=, [n]))$, or $C([n]) = C_{\text{Moore}}(\mathbb{k}\Delta(=, [n]))$ for a realisation/totalisation in $E = C^*(\mathbb{k})$ (here, C_{Moore} and N denote the Moore chain complex and the normalised Moore chain complex of a simplicial vector space).

It follows from the definition that the realisation commutes with all small colimits and the totalisation commutes with all small limits.

Using this observation, one proves that the realisation of a simplicial set X in $C^\bullet(\mathbb{k})$, for $C([n])$ equal to the non-normalised Moore complex of $\mathbb{k}\Delta(=, [n])$, is equal to the non-normalised Moore complex of X . This fact is fairly standard, but as we employ a similar argument for $\Xi = \Theta_2$ later in the paper, we recall the argument for completeness.

Represent $X = \text{colim}_{h_{[j]} \rightarrow X} h_{[j]}$, where $h_{[j]}([i]) = \Delta([i], [j])$. Then,

$$\begin{aligned} |X| &= |\text{colim}_{h_{[j]} \rightarrow X} h_{[j]}| = \text{colim}_{h_{[j]} \rightarrow X} |h_{[j]}| \stackrel{*}{=} \text{colim}_{h_{[j]} \rightarrow X} C([j]) \\ &= \text{colim}_{h_{[j]} \rightarrow X} C_{\text{Moore}}^\bullet(h_{[j]}) \end{aligned}$$

where in the equality marked by $*$ we use that the Yoneda functor is fully faithful and that the unit of the left Kan extension adjunction along a fully faithful functor is the identity [31, Cor. 1.4.5]. The rightmost complex has its i -th term equal to $\text{colim}_{h_{[j]} \rightarrow X} \mathbb{k}\Delta([i], [j])$. Then,

$$(\text{colim}_{h_{[j]} \rightarrow X} C^\bullet([j]))_i = \text{colim}_{h_{[j]} \rightarrow X} (\mathbb{k}h_{[j]}([i])) = (\text{colim}_{h_{[j]} \rightarrow X} \mathbb{k}h_{[j]})[i] = X([i]).$$

Similarly, we show, for $C([n]) = N(\mathbb{k}\Delta(=, [n]))$, the totalisation counterparts of these statements.

In this paper, we consider the realisation and the totalisation for

$$C([n]) = C_{\text{Moore}}^\bullet(\mathbb{k}\Delta(=, [n])),$$

for the case of Δ , and $C(T) = C^\bullet(\mathbb{k}\Theta_2(=, T))$, where $C^\bullet(=)$ is the complex defined in (3.1) and (3.2).

We also define “relative totalisation” with respect to the projection $p: \Theta_2 \rightarrow \Delta$. For the case of realisation, it can be defined as the following left Kan extension of Φ along the Yoneda embedding

$$\begin{array}{ccc} \Theta_2 & \longrightarrow & \text{Sets}^{\Theta_2^{\text{op}}} \\ \downarrow \Phi & & \downarrow \text{Lan} \\ C^\bullet(\mathbb{k})^{\Delta^{\text{op}}} & & \end{array}$$

where the functor Φ sends an object $T \in \Theta_2$ to the functor $\Delta^{\text{op}} \rightarrow C^\bullet(\mathbb{k})$, $[n] \mapsto \mathfrak{R}_{[n]}(T)$; see (3.3).

Note that the functor $\Delta \rightarrow C^\bullet(\mathbb{k})$, $[\ell] \mapsto C_{\text{Moore}}^\bullet(\mathbb{k}\Delta(=, [\ell]))$ is a resolution of the constant functor $[\ell] \mapsto \mathbb{k}$. The functor $T \mapsto C(T)$ is a projective resolution by Yonedas of the constant functor $T \mapsto \mathbb{k}$ (as is proven in Section 3.2), and the functor Φ above is a resolution of the functor $T \mapsto \{[n] \mapsto \mathbb{k}\Delta([n], p(T))\}$ (as it is proven in Lemma 3.3 below).

Morally, we wanted to talk about the homotopy right Kan extension along p , instead of the p -relative totalisation. We want to use the transitivity of the right homotopy Kan

extensions for the composition $\Theta_2 \xrightarrow{p} \Delta \rightarrow *$, which is clear from the derived functor of composition theorem, to prove Proposition 3.7. However, the totalisation agrees with the homotopy right Kan extension for the projection to $*$ only for *Reedy fibrant cosimplicial (resp., 2-cocellular) complexes of vector spaces* (see [18, Sect. 18.7]). Typically, the cosimplicial space whose Moore complex is the Hochschild complex is *not* Reedy fibrant (despite the fact that the corresponding bar-complex is Reedy cofibrant). To overcome this difficulty, we talk about the “relative totalisation” along $p: \Theta_2 \rightarrow \Delta$, choosing the functor Φ exactly as a resolution of $\mathbb{k}\Delta([n], p(T))$ by the (projective) Yoneda functors. Then, we prove the transitivity comparing the explicit formulas for the totalisations.

3.2. The (absolute) totalisation of a 2-cocellular vector space

Let $X: \Theta_2 \rightarrow C(\mathbb{k})$ be a cocellular complex. First of all, we define its non-normalised Moore complex explicitly, as the complex whose degree ℓ component is

$$\text{Tot}_{\Theta_2}(X)^\ell = \bigoplus_{\substack{T \in \Theta_2 \\ \dim T = \ell}} X_T \tag{3.1}$$

and the differential of degree $+1$ is equal to the sum of (taken with appropriate signs) all codimension 1 face maps:

$$\begin{aligned} d|_{X_T} = & \sum_{\substack{\text{coface maps } \partial_p^i \\ (F1), (F3)}} (-1)^{k_1 + \dots + k_{p-1} + p - 1 + i - 1} \partial_p^i \\ & + \sum_{\substack{\text{coface maps } D_{p,\sigma} \\ (F2)}} (-1)^{k_1 + \dots + k_{p-1} + p - 1 + \#(\sigma)} D_{p,\sigma} \\ & + D_{\min} + (-1)^{k_1 + \dots + k_n + n} D_{\max}, \end{aligned} \tag{3.2}$$

where $T = ([n]; [k_1], \dots, [k_n])$, $\dim T = k_1 + \dots + k_n + n$, and for the notations for the face maps, see Section 2.6.

Lemma 3.2. *One has $d^2 = 0$.*

Proof. It follows from relations (A.1)–(A.7) that the summands in d^2 come in pairs, in which the two operators are equal one to another. One checks by hand that for each pair the two terms have opposite signs, which makes them mutually cancelled. ■

This formula can be “explained” in the following way. We compute $\underline{\text{Hom}}_{\Theta_2}(K^\bullet, X)$ where K^\bullet is a (projective) resolution of the constant 2-cocellular vector space \mathbb{k} , formed by the (linearised) Yoneda modules $\mathbb{k}\Theta_2(-, T)$, for $T \in \Theta_2$ (here, $\underline{\text{Hom}}_{\Theta_2}$ stands for Hom taking values in $C^\bullet(\mathbb{k})$, and it can be equally defined as enriched natural transformations [31, Sect. 7.3]). In this way, it is the “semi-derived” functor of $\text{colim}_{\Theta_2} X$ because $\text{Hom}_{\Theta_2}(\mathbb{k}, X) = \text{colim}_{\Theta_2} X$, but it is not properly derived because X is not, in general, Reedy fibrant. We can not claim that for another choice of resolution of \mathbb{k} we get a quasi-isomorphic complex.

Here is an explicit resolution.

Fix $T \in \Theta_2$. Denote by $K^\bullet(T)$ a complex whose components are

$$K^{-\ell}(T) = \bigoplus_{\substack{T' \in \Theta_2 \\ |T'| = \ell}} \mathbb{k}\Theta_2(T', T).$$

The differential $d: K^{-\ell}(T) \rightarrow K^{-\ell+1}(T)$ is defined likewise the differential in (3.2), so that $K^\bullet(-, T)$ for a given T is the realisation of the cellular vector space $? \mapsto \mathbb{k}\Theta_2(? , T)$. We check $d^2 = 0$ as in Lemma 3.2.

Clearly the post-composition makes $K^\bullet(T)$ functorial in T . One remains to show that, for given T , $(K^\bullet(T), d)$ is quasi-isomorphic to \mathbb{k} .

Consider $(K^\bullet(T), d)$ as the total complex of a bicomplex, with “vertical” component of the differential

$$d_1 = \sum_{\text{coface maps } \partial_p^j(F1), (F3)} (-1)^{k_1 + \dots + k_{p-1} + p-1 + i-1} \partial_p^i$$

and the “horizontal” component

$$d_0 = \sum_{\substack{\text{coface maps} \\ D_{p,\sigma} (F2)}} (-1)^{k_1 + \dots + k_{p-1} + p-1 + \#(\sigma)} D_{p,\sigma} + D_{\min} + (-1)^{k_1 + \dots + k_n + n} D_{\max}.$$

We claim that $d_0 d_1 + d_1 d_0 = 0$; we check it later in the proof of Proposition 3.5.

3.3. The p -relative totalisation $Rp_*(X)$: an explicit description

3.3.1. The ordinary enriched right Kan extension of X along $p: \Theta_2 \rightarrow \Delta$ is taken equal to

$$p_*(X)([n]) = \underline{\text{Hom}}_{\Theta_2}(\mathbb{k}\Delta([n], p(-)), X(-)),$$

where, as above, $\underline{\text{Hom}}_{\Theta_2}$ taking values in $C^\bullet(\mathbb{k})$ is the enriched natural transformations. We replace $\mathbb{k}\Delta([n], p(T))$ by its resolution $\mathfrak{R}_{[n]}$ by Yoneda modules. Note that it is not the homotopy right Kan extension along p , for the case X is not Reedy fibrant. Therefore, we can not say that for any other choice of resolution we get a quasi-isomorphic complex. The resulting Δ -complex $\text{Hom}_{\Theta_2}(\mathfrak{R}, X)$ is “closely related” to our relative realisation.

We have to resolve the functor $T \mapsto \mathbb{k}\Delta([n], p(T))$ (for a given $[n] \in \Delta$) by the Yoneda functors $h_{T'}(T) = \mathbb{k}\Theta_2(T', T)$. Below we provide an explicit resolution $\mathfrak{R}_{[n]}^\bullet$ (which is a complex of vector spaces over \mathbb{k}).

The degree ℓ component is

$$\mathfrak{R}_{[n]}^\ell(T) = \bigoplus_{\substack{T' \in \Theta_2, p(T') = [n] \\ \dim T' = n - \ell}} \mathbb{k}\Theta_2(T', T). \tag{3.3}$$

Thus, the complex $\mathfrak{R}_{[n]}^\bullet$ has non-zero components in degrees ≤ 0 .

The differential $d: \mathfrak{R}_{[n]}^\ell \rightarrow \mathfrak{R}_{[n]}^{\ell+1}$ is defined as the alternated sum of the “vertical” face operators (acting on the first argument T'), that is, of face operators (F1) and (F3) from the list in Section 2.6. More precisely, for $T = ([q]; [t_1], \dots, [t_q])$, $T' = ([n]; [k_1], \dots, [k_n])$, $\Phi = (\phi; \phi_1, \dots, \phi_n): T' \rightarrow T$, one has

$$d(\Phi) = \sum_{s=1}^n \sum_{i=0}^{k_s} (-1)^{k_1 + \dots + k_{s-1} + s - 1 + i} \Phi_{s,i}, \tag{3.4}$$

where $\Phi_{s,i}: T'_{s,i} \rightarrow T$ is defined as the pre-composition $\Phi \circ \partial_s^i$ (see Section 2.6, (F1) and (F3)), and $T'_{s,i} = ([n]; [m_1], \dots, [m_n])$, where $m_j = k_j$ for $j \neq s$, $m_s = k_s - 1$, and $\partial_s^i: T'_{s,i} \rightarrow T'$ is the corresponding “vertical” face operator. Note that this pre-composition does not affect the “horizontal” map ϕ . It is clear that $d^2 = 0$.

Lemma 3.3. *The following statements are true:*

- (1) *degree 0 cohomology of $\mathfrak{R}_{[n]}^\bullet$ is isomorphic to $\mathbb{k}\Delta([n], p(T))$,*
- (2) *the higher cohomology (in the negative degrees ≤ -1) vanishes.*

Proof. (1) The degree 0 component $\mathfrak{R}_{[n]}^0$ is a direct sum $\oplus \mathbb{k}\Phi$, where

$$\Phi: ([n]; [0], \dots, [0]) \rightarrow T,$$

which is the same as $\Phi = (\phi: [n] \rightarrow [q]; T_1, \dots, T_n)$ where $T_i \in [t_{\phi(i-1)+1}] \times \dots \times [t_{\phi(i)}]$, an element (recall that $[q] = p(T)$). Degree 0 cohomology is equal to the quotient-space by the image of $\oplus \mathbb{k}\Phi'$, with $\Phi': ([n]; [0], \dots, [0], [1], [0], \dots, [0]) \rightarrow T$. For a given $\phi: [n] \rightarrow [q]$, all choices of T_i become equal in the quotient-space $H^0(\mathfrak{R}_{[n]}^\bullet) = \mathfrak{R}_{[n]}^0/d(\mathfrak{R}_{[n]}^{-1})$. It shows that $H^0(\mathfrak{R}_{[n]}^\bullet) \simeq \mathbb{k}\Delta([n], [q]) = \mathbb{k}\Delta([n], p(T))$.

(2) We construct a contracting homotopy operator H of degree -1 , that is, an operator such that $(dH + Hd)|_{\mathfrak{R}_{[n]}^\ell} = c_\ell$, where c_ℓ is the multiplication by an integer c_ℓ , non-zero for $\ell \neq 0$. This H is constructed in a standard way as the alternated sum of the “vertical” degeneracy operators. ■

Remark 3.4. It is clear that the complex $\mathfrak{R}_{[n]}^\bullet$ is a direct sum $\oplus_\phi \mathfrak{R}_{[n],\phi}^\bullet$ over $\phi: [n] \rightarrow p(T)$ because the differential does not affect ϕ . Each complex $\mathfrak{R}_{[n],\phi}^\bullet$ is a resolution of \mathbb{k} (where \mathbb{k} denotes the complex-object \mathbb{k} in degree 0).

One checks that $\mathfrak{R}_{[n]}^\bullet$ is a functor $\Theta_2 \rightarrow C^\bullet(\mathbb{k})$, where the action of Θ_2 is given by the *post-composition*. It commutes with the differential as the general post-composition and pre-composition do.

3.3.2. The action of Δ . Our next task is to endow our resolution $\mathfrak{R}_{[n]}^\bullet$ with a structure of a functor $\Delta^{\text{op}} \rightarrow C^\bullet(\mathbb{k})$, when $[n]$ varies. Note that unlike for the cohomology $\mathbb{k}\Delta([n], p(T))$ of $\mathfrak{R}_{[n]}^\bullet$, the “lifted” action of Δ on $\mathfrak{R}_{[n]}^\bullet$ does not come automatically.

We need to define the actions of the elementary face operators ∂^i and the elementary degeneracy operators ε^j in Δ , which we denote, in this context, by Ω_Δ^i and Υ_Δ^j , correspondingly.

Here are the definitions.

Let $\Phi: T' \rightarrow T$ be an element in $\mathfrak{R}_{[n]}^\ell$, $p(T') = [n]$.

$$\Omega_\Delta^i(\Phi) = \sum_\sigma \pm \Phi \circ D_{i,\sigma} \pm \Phi \circ D_{\min} \pm \Phi \circ D_{\max}, \quad \Upsilon_\Delta^j(\Phi) = \pm \Phi \circ \Upsilon_0^j,$$

where $\Upsilon_0^j: ([n]; [\ell_1], \dots, [\ell_j], [0], [\ell_{j+2}], \dots, [\ell_n]) \rightarrow T$. Note that we take only T' with $[\ell_{j+1}] = [0]$.

(See Section 2.6 for the notations $D_{i,\sigma}$ and Υ^j .)

Proposition 3.5. *The following statements are true:*

- (1) *the operators Ω_Δ^i and Υ_Δ^j define maps of complexes $\Omega_\Delta^i: \mathfrak{R}_{[n]}^\bullet \rightarrow \mathfrak{R}_{[n-1]}^\bullet$ and $\Upsilon_\Delta^j: \mathfrak{R}_{[n]}^\bullet \rightarrow \mathfrak{R}_{[n+1]}^\bullet$, preserving the cohomological degree,*
- (2) *the operators Ω_Δ^i and Υ_Δ^j fulfil the simplicial relations, defining a simplicial object $\mathfrak{R}_?^\bullet: \Delta^{\text{op}} \rightarrow C^\bullet(\mathbb{k})$, functorial in T . The cohomology $H^\bullet(\mathfrak{R}_{[n]}^\bullet)$ with respect to the differential (3.4), with its simplicial action, is isomorphic to $\Delta([n], p(T))$ with its natural simplicial action.*

The proof of Proposition 3.5 is a straightforward computation using relations between the generators of Θ_2 listed in the appendix. It is reproduced in [29, Appx. B].

3.3.3. Finally, here is $Rp_*(X)$. For $X \in \text{Fun}(\Theta_2, C^\bullet(\mathbb{k}))$, define $Rp_*(X)[n] \in C^\bullet(\mathbb{k})$ as the following complex:

$$0 \rightarrow Rp_*(X)[n]^0 \xrightarrow{d} Rp_*(X)[n]^1 \xrightarrow{d} Rp_*(X)[n]^2 \xrightarrow{d} \dots$$

where

$$Rp_*(X)[n]^\ell = \bigoplus_{\substack{T \in \Theta_2, p(T)=[n] \\ \dim T = n + \ell}} X(T)$$

and the differential d is the alternated sum of “vertical” face operators (of type (F1) and (F3) in Section 2.6):

$$d|_{X(T)} = \sum_{i=1}^{p(T)} \sum_{j=0}^{T_i} (-1)^{T_1 + \dots + T_{i-1} + j} \partial_i^j$$

where we write $T = (p(T); [T_1], \dots, [T_{p(T)}])$.

According to Proposition 3.5, when $[n] \in \Delta$ varies, it gives rise to a functor $\Delta \rightarrow C^\bullet(\mathbb{k})$.

More precisely, we have formulas for the coface maps

$$\Omega_\Delta^i: Rp_*(X)[n] \rightarrow Rp_*(X)[n + 1], \quad i = 0, \dots, n + 2$$

similar to the ones stated in Proposition 3.5 and the codegeneracy maps $\Upsilon_\Delta^j: Rp_*(X)[n] \rightarrow Rp_*(X)[n - 1]$, $j = 0, \dots, n$:

$$\Omega_\Delta^i = \sum_\sigma \pm D_{i,\sigma} \pm D_{\min} \pm D_{\max} \tag{3.5}$$

and

$$\Upsilon_{\Delta}^j = \pm \Upsilon_0^j \tag{3.6}$$

in the notations of Section 2.6. The signs are the same as in Proposition 3.5.

Proposition 3.6. *The coface and codegeneracy operators (3.5) and (3.6) commute with the differentials on $Rp_*(X)[n]$ for a given n and satisfy the standard cosimplicial identities.*

For a proof, we have two options. We can either note that the proof of Proposition 3.5 is literally applied to a proof of Proposition 3.6, with the same computations, or, alternatively, we can deduce it from Proposition 3.5 and the definition of $Rp_*(X)[=]$ as the right Kan extension

$$\begin{array}{ccc} \Theta_2^{\text{op}} & \longrightarrow & (\text{Sets}^{\Theta_2^{\text{op}}})^{\text{op}} \\ \downarrow \Phi & \nearrow \text{Ran} = Rp_*(-) & \\ (C^\bullet(\mathbb{k})^{\Delta^{\text{op}}})^{\text{op}} & & \end{array}$$

Here, Φ is defined as above, $T \mapsto \{[n] \mapsto \mathfrak{R}_{[n]}(T)\}$. In this way, the Δ -action on $\mathfrak{R}_{[?]}(T)$ given by Proposition 3.5 is translated to the action of Δ on $Rp_*(X)[?]$. ■

3.3.4. $\text{Tot}_{\Delta}(Rp_*(X)) \simeq \text{Tot}_{\Theta_2}(X)$. Here we prove the following.

Proposition 3.7. *Let $X: \Theta_2 \rightarrow C(\mathbb{k})$ be a 2-cocellular vector space. Then, the Δ -totalisation $\text{Tot}_{\Delta}(Rp_*(X))$ of $Rp_*(X)$ is a complex isomorphic to the Θ_2 -totalisation $\text{Tot}_{\Theta_2}(X)$.*

Proof. When one applies the usual non-normalised cochain complex functor to the cosimplicial vector space $Rp_*(X)$, we get exactly the formula (3.1) for the (non-normalised) Θ_2 -totalisation. ■

4. Bicategories and 2-bimodules over bicategories

4.1. Reminder on bicategories

Here, we recall the basic definitions related to bicategories, basically aiming to fix our terminology and notations. The reference are [9, 20, 23, 25].

Shortly, a bicategory is a lax category enriched in $\mathcal{C}\text{at}$. Here, “lax” indicates that the associativity of composition holds up to a 2-arrow which is an isomorphism. Below is a detailed definition (of a small bicategory).

Definition 4.1. A small bicategory C consists of the following data:

- (1) a set C_0 whose elements are called the objects of C ,
- (2) for $x, y \in C_0$, a set $C_1(x, y)$ whose elements are called 1-morphisms or 1-arrows,
- (3) for any $x, y \in C_0$ and $f, g \in C_1(x, y)$, a set $C_2(x, y)(f, g)$ called 2-morphisms or 2-arrows,

- (4) for any $x, y \in C_0$, one has a small category $C(x, y)$ whose objects are $C_1(x, y)$ and whose arrows $f \rightarrow g$ are $C_2(x, y)(f, g)$,
- (5) for any $x, y, z \in C_0$, there is a composition $m_{x,y,z}: C(y, z) \times C(x, y) \rightarrow C(x, z)$, which is a functor of categories,
- (6) for any $x, y, z, w \in C_0$, there is a natural transformation

$$\begin{array}{ccc}
 C(z, w) \times C(y, z) \times C(x, y) & \xrightarrow{m_{y,z,w} \times \text{id}} & C(y, w) \times C(x, y) \\
 \downarrow \text{id} \times m_{x,y,z} & \Uparrow \alpha_{x,y,z,w} & \downarrow m_{x,y,w} \\
 C(z, w) \times C(x, z) & \xrightarrow{m_{x,z,w}} & C(x, w)
 \end{array}$$

whose components are given by isomorphisms (for which we often use the notation $\alpha_{h,g,f}$ assuming $\alpha_{x,y,z,w}(h \times g \times f)$),

- (7) for any $x, y \in C_0$, there are 2-arrows $\rho_{x,y}$ and $\lambda_{x,y}$ defined as

$$\begin{array}{ccc}
 C(x, y) \times \mathbf{1} & & \mathbf{1} \times C(x, y) \\
 \text{id} \times \text{id}_x \downarrow & \searrow \cong & \downarrow \text{id}_y \times \text{id} \\
 C(x, y) \times C(x, x) & \xrightarrow{m_{x,x,y}} & C(x, y) & \quad & C(y, y) \times C(x, y) & \xrightarrow{m_{x,y,y}} & C(x, y) \\
 \Uparrow \rho_{x,y} & & & & \Uparrow \lambda_{x,y} & & \\
 & & & & & &
 \end{array}$$

whose components are isomorphisms (we often denote $\rho_{x,y}(f \times \text{id})$ and $\lambda_{x,y}(\text{id} \times g)$ by $\rho(f)$ and $\lambda(g)$, correspondingly)

which are subject to the following identities:

- (i) The associator $\alpha_{x,y,z,w}$ is subject to the usual pentagon diagram:

$$\begin{array}{ccc}
 & (t \circ h) \circ (g \circ f) & \\
 \alpha_{t,h,g \circ f} \nearrow & & \searrow \alpha_{t \circ h,g,f} \\
 t \circ (h \circ (g \circ f)) & & ((t \circ h) \circ g) \circ f \\
 m_{t,\alpha_{h,g,f}} \searrow & & \nearrow m_{\alpha_{t,h,g},f} \\
 t \circ ((h \circ g) \circ f) & \xrightarrow{\alpha_{t,h \circ g,f}} & (t \circ (h \circ g)) \circ f
 \end{array}$$

- (ii) for any $x, y \in C_0$, the unit 2-arrows

$$\lambda(f): m_{x,y,y}(\text{id}_y, f) \Rightarrow f$$

and

$$\rho(f): m_{x,x,y}(f, \text{id}_x) \Rightarrow f$$

are subject to the following unit identity, for $f \in C_1(x, y), g \in C_1(y, z)$:

$$\begin{array}{ccc}
 g \circ (\text{id}_y \circ f) & \xrightarrow{\alpha_{x,y,z,w}} & (g \circ \text{id}_y) \circ f \\
 \searrow \text{id}(g) \circ \lambda(f) & & \swarrow \rho(g) \circ \text{id}(f) \\
 & g \circ f &
 \end{array}$$

The composition of 2-arrows coming from the composition in $C(x, y)$ is called *vertical* (notation: $f \circ_v g$, where $f, g \in C(x, y)$), while the composition of 2-arrows coming from $m_{x,y,z}: C(y, z) \times C(x, z) \rightarrow C(x, z)$ is called *horizontal* (notation: $g \circ_h f$, where $f \in C(x, y), g \in C(y, z)$).

Examples. (i) When $\alpha_{x,y,z,w}, \lambda(f), \rho(f)$ are the identity 2-arrows, one gets the concept of a 2-category (which is then a “strict” bicategory).

(ii) When $C_0 = \{*\}$, the bicategory is a monoidal category.

Next, we recall lax-functors of bicategories and lax-natural transformations thereof. Also, we recall the concept of a *modification* between two lax natural transformations, playing the role of 3-arrows in the (weak) 3-category of bicategories (they form a tricategory; see [16]).

Definition 4.2. Let C, D be bicategories. A *lax-functor* $F: C \rightarrow D$ is given by the following data:

- (1) an assignment $F: C_0 \rightarrow D_0$ of objects,
- (2) for any $x, y \in C_0$, a functor $F_{x,y}: C(x, y) \rightarrow D(Fx, Fy)$,
- (3) for any $x, y, z \in C_0$, there is a natural transformation $\phi_{x,y,z}$ making the diagram below commutative:

$$\begin{array}{ccc}
 C(y, z) \times C(x, y) & \xrightarrow{m_{x,y,z}^C} & C(x, z) \\
 \downarrow F_{y,z} \times F_{x,y} & \Rightarrow \phi_{x,y,z} & \downarrow F_{x,y} \\
 D(Fy, Fz) \times D(Fx, Fy) & \xrightarrow{m_{Fx,Fy,Fz}^D} & D(Fx, Fz)
 \end{array}$$

(we use notations $\phi_{g,f}: Fg \circ Ff \Rightarrow F(gf)$),

- (4) for any $x \in C_0$, there is a natural transformation ϕ_x making the diagram below commutative:

$$\begin{array}{ccc}
 \mathbf{1} & \xrightarrow{I_x} & C(x, x) \\
 \downarrow = & \Rightarrow \phi_x & \downarrow F_{x,x} \\
 \mathbf{1} & \xrightarrow{I_{Fx}} & D(Fx, Fx)
 \end{array}$$

which are subject to the following properties:

(i) the hexagon diagram

$$\begin{array}{ccccc}
 (Fh \circ Fg) \circ Ff & \xrightarrow{\phi \circ \text{id}} & F(h \circ g) \circ Ff & \xrightarrow{\phi} & F((h \circ g) \circ f) \\
 \downarrow \alpha^D & & & & \downarrow F(\alpha^C) \\
 Fh \circ (Fg \circ Ff) & \xrightarrow{\text{id} \circ \phi} & Fh \circ F(g \circ f) & \xrightarrow{\phi} & F(h \circ (g \circ f))
 \end{array}$$

commutes,

(ii) the unit commutative diagrams for ϕ_x (see, e.g., [23, Sect. 1.1]).

When $\{\phi_{x,y,z}\}$ and $\{\phi_x\}$ are isomorphisms, the lax functor F is called *strong*; when they are identity natural transformations, the lax functor is called *strict*.

Definition 4.3. Let C, D be bicategories, $F = (F, \phi), G = (G, \psi): C \rightarrow D$ lax-functors. A *lax natural transformation* $\eta: F \Rightarrow G$ is given by the following data:

- (1) a collection of 1-arrows $\eta_x: Fx \rightarrow Gx, x \in C_0$,
- (2) a collection of natural transformations

$$\begin{array}{ccc}
 C(x, y) & \xrightarrow{F_{x,y}} & D(Fx, Fy) \\
 \downarrow G_{x,y} & \Rightarrow_{\eta_{x,y}} & \downarrow (\eta_y)_* \\
 D(Gx, Gy) & \xrightarrow{(\eta_x)_*} & D(Fx, Gy)
 \end{array}$$

which are subject to commutative diagrams expressing the compatibility of $\{\eta_f\}$ with the composition $f_2 \circ f_1$ and of $\{\eta_x\}$ with unit morphisms; see, e.g., [23, Sect. 1.2]. The 2-arrows $\eta_{x,y}$ are given in components by 2-arrows

$$\eta_f: G(f) \circ \eta_x \Rightarrow \eta_y \circ F(f): F(x) \rightarrow G(y), \quad \text{for any } f \in C_1(x, y).$$

A lax-monoidal transformation of bicategories is called *strong* if the 2-arrows $\{\eta_f\}$ are isomorphisms, and it is called *strict* if they are identity 2-arrows.

Definition 4.4. Let C, D be bicategories, $F, G: C \rightarrow D$ lax-functors, $\eta, \theta: F \Rightarrow G$ lax natural transformations. A *modification* $\kappa: \eta \Rightarrow \theta$ is given by a collection of 2-arrows $\{\Gamma_x\}$ for $x \in C_0$:

$$\begin{array}{ccc}
 & \theta_x & \\
 & \curvearrowright & \\
 Fx & & Gx \\
 & \Downarrow \Gamma_x & \\
 & \curvearrowleft & \\
 & \eta_x &
 \end{array}$$

(denoted as $\Gamma: \eta \Rightarrow \theta$) which are subject to the following axiom: for any $f: x \rightarrow y \in C_1$,

the diagram below commutes:

$$\begin{array}{ccc}
 Gf \circ \eta_x & \xrightarrow{\text{id} \circ \Gamma_x} & Gf \circ \theta_x \\
 \eta_f \downarrow & & \downarrow \theta_f \\
 \eta_y \circ Ff & \xrightarrow{\Gamma_y \circ \text{id}} & \theta_y \circ Ff
 \end{array}$$

Definition 4.5. For a symmetric monoidal category \mathcal{V} , by \mathcal{V} -enriched bicategory we mean a modification of the previous definition, in which $C_2(x, y)(f, g) \in \mathcal{V}$ (while $C_0, C_1(x, y) \in \text{Sets}$), so that each category $C(x, y)$ is a \mathcal{V} -enriched category, for any $x, y \in C_0$, and the compositions $m_{x,y,z}$ are given by \mathcal{V} -functors. Similarly, one defines \mathcal{V} -lax functors, \mathcal{V} -lax natural transformations, and \mathcal{V} -modifications. Namely, one adjusts axioms by requiring that the 2-arrows $\phi_x, \phi_{x,y,z}, \eta_x, \eta_{x,y}, \Gamma_x$ are objects of \mathcal{V} . The reader will find details in [20].

A particular case we deal with in this paper is $\mathcal{V} = C^*(\mathbb{k})$, the category of complexes over a field \mathbb{k} . In this case, we call a \mathcal{V} -enriched bicategory a dg bicategory.

4.1.1. Coherence for bicategories. We will need the coherence theorem for bicategories [9, 23, 25].

Let C, D be two bicategories. They are said to be *biequivalent* if there are *strong* functors $F: C \rightarrow D$ and $G: D \rightarrow C$ and *strong* natural transformations $\text{id}_C \rightarrow G \circ F$ and $F \circ G \rightarrow \text{Id}_D$ (that is, we use only the underlying 1-category structure on the bicategory $[C, D]$; the modifications are irrelevant for this definition). The coherence theorem for bicategories is the following result.

Theorem 4.6. *Every bicategory is biequivalent to a 2-category.*

See, e.g., [23, Sect. 2.3] for a short proof, based on the Yoneda embedding. ■

In a 2-category, the associator and the identity maps are equal to identity. The following statement, which we primarily will use in the paper, is a direct consequence of Theorem 4.6.

Corollary 4.7. *Let C be a bicategory, $x, y \in C_0$, $f, g: C_1(x, y)$. Let $\eta, \theta: f \Rightarrow g$ be two natural transformations each of which is a composition of the associators and the unit maps. Then, $\eta = \theta$.* ■

A similar statement holds for enriched bicategories, in particular, for dg bicategories.

4.2. 2-bimodules over a bicategory

Let \mathcal{V} be a symmetric monoidal category, C a \mathcal{V} -enriched bicategory (see Definition 4.5). A C -2-bimodule consists of the following data:

- (i) a 2-globular set M whose second component M_2 is \mathcal{V} -enriched, and $M_{\leq 1} = C_{\leq 1}$,

(ii) the upper and the lower vertical compositions:

$$\bar{\circ}_-^v: M_2(g, h) \otimes C_2(f, g) \rightarrow M_2(f, h), \quad \bar{\circ}_+^v: C_2(g, h) \otimes M_2(f, g) \rightarrow M_2(f, h)$$

where $f, g, h \in C_1(x, y)$,

(iii) the left and the right horizontal compositions

$$\bar{\circ}_-^h: M_2(g_1, g_2) \otimes C_2(f_1, f_2) \rightarrow M_2(g_1 f_1, g_2 f_2),$$

$$\bar{\circ}_+^h: C_2(g_1, g_2) \otimes M_2(f_1, f_2) \rightarrow M_2(g_1 f_1, g_2 f_2),$$

where $f_1, f_2 \in C_1(x, y)$, $g_1, g_2 \in C_1(y, z)$,

which are subject to the following properties:

(1) the vertical compositions are strictly associative, in the sense that for $f_1, f_2, f_3, f_4 \in C_1(x, y)$, the following three identities hold:

$$m \bar{\circ}_-^v(\psi \circ^v \phi) = (m \bar{\circ}_-^v \psi) \bar{\circ}_-^v \phi,$$

where $m \in M_2(f_3, f_4)$, $\phi \in C_2(f_1, f_2)$, $\psi \in C_2(f_2, f_3)$,

$$\psi \bar{\circ}_+^v(m \bar{\circ}_-^v \phi) = (\psi \bar{\circ}_+^v m) \bar{\circ}_-^v \phi,$$

where $\psi \in C_2(f_3, f_4)$, $m \in M_2(f_2, f_3)$, $\phi \in C_2(f_1, f_2)$,

$$\psi \bar{\circ}_+^v(\phi \bar{\circ}_+^v m) = (\psi \circ \phi) \bar{\circ}_+^v m,$$

where $\psi \in C_2(f_3, f_4)$, $\phi \in C_2(f_2, f_3)$, $m \in M_2(f_1, f_2)$,

(2) the horizontal compositions are associative up to the associator α in C , in the sense that for $x, y, z, w \in C_0$, $f_1, f_2 \in C_1(x, y)$, $g_1, g_2 \in C_1(y, z)$, $h_1, h_2 \in C_1(z, w)$, one has

$$\alpha_{h_2, g_2, f_2} \bar{\circ}_+^v(m \bar{\circ}_-^h(\psi \circ^h \phi)) = ((m \bar{\circ}_-^h \psi) \bar{\circ}_-^h \phi) \bar{\circ}_-^v \alpha_{f_1, g_1, h_1}, \quad (4.1)$$

where $\phi \in C_2(f_1, f_2)$, $\psi \in C_2(g_1, g_2)$, $m \in M_2(h_1, h_2)$,

$$\alpha_{f_2, g_2, h_2} \bar{\circ}_+^v(\psi \bar{\circ}_+^h(m \bar{\circ}_-^h \phi)) = ((\psi \bar{\circ}_+^h m) \bar{\circ}_-^h \phi) \bar{\circ}_-^v \alpha_{f_1, g_1, h_1}, \quad (4.2)$$

where $\phi \in C_2(f_1, f_2)$, $m \in M_2(g_1, g_2)$, $\psi \in C_2(h_1, h_2)$,

$$\alpha_{f_2, g_2, h_2} \bar{\circ}_+^v(\psi \bar{\circ}_+^h(\phi \bar{\circ}_+^h m)) = ((\psi \circ \phi) \bar{\circ}_+^h m) \bar{\circ}_-^v \alpha_{f_1, g_1, h_1}, \quad (4.3)$$

where $\phi \in C_2(g_1, g_2)$, $\psi \in C_2(h_1, h_2)$, $m \in C_2(f_1, f_2)$,

(3) the four Eckmann–Hilton identities (depending on the place among the four arguments at which an element of M_2 is, then there are elements of C_2 on the three others). One among these 4 identities reads:

$$(\phi_3 \bar{\circ}_+^v m) \bar{\circ}_-^h(\phi_2 \circ^v \phi_1) = (\phi_3 \circ^h \phi_2) \bar{\circ}_+^v(m \bar{\circ}_-^v \phi_1),$$

where $f_1, f_2, f_3 \in C_1(x, y)$, $g_1, g_2, g_3 \in C_1(y, z)$, $\phi_1 \in C_2(f_1, f_2)$, $\phi_2 \in C_2(f_2, f_3)$, $\phi_3 \in C_2(g_1, g_2)$, $m \in M_2(g_2, g_3)$; the three other identities are similar and are left to the reader,

(4) let $x, y \in C_0, f, g \in C_1(x, y), m \in M_2(f, g)$; then,

$$\rho(g)\bar{\sigma}_+^v(m\bar{\sigma}_-^h\text{id}(x))\bar{\sigma}_-^v\rho(f)^{-1}=m, \quad \lambda(g)\bar{\sigma}_+^v(\text{id}(y)\bar{\sigma}_+^hm)\bar{\sigma}_-^v\lambda(f)^{-1}=m. \quad (4.4)$$

The category of 2-bimodules over a bicategory C is denoted by $\text{Bimod}_2(C)$.

As a trivial example, $M = C$ satisfies the axioms; it is called the *tautological 2-bimodule* over C .


4.3. Examples

4.3.1. The free 2-bimodule. There is a forgetful functor U from C -2-bimodules to 2-globular sets whose second component is enriched over \mathcal{V} . Assume for simplicity that $\mathcal{V} = \text{Vect}(\mathbb{k})$ or $C^*(\mathbb{k})$. The functor U has a left adjoint L ; the corresponding C -2-bimodule $L(M_0)$ is called the *free C -2-bimodule generated by the \mathcal{V} -enriched 2-globular set M_0* . We set $L(M_0)_{\leq 1} = C_{\leq 1}$, let $x, y \in C_0$, and $f, g \in C_1(x, y)$. Define

$$M(f, g) = \bigoplus_{\substack{z_0, z_1 \in C_0 \\ \alpha \in C_1(x, z_0), \beta \in C_1(z_1, y) \\ f_0, g_0 \in C_1(z_0, z_1)}} C(\beta \circ g_0 \circ \alpha, g) \otimes M_0(f_0, g_0) \otimes C(f, \beta \circ f_0 \circ \alpha) / \sim$$

where the equivalence relation identifies the following elements, in notations of diagram (4.6) below:

$$(\phi_2 \circ^v (\sigma_2 \circ^h \text{id}_{g_0} \circ^h \sigma_1)) \otimes m \otimes \phi_1 \sim \phi_2 \otimes m \otimes ((\sigma_2 \circ^h \text{id}_{f_0} \circ^h \sigma_1) \circ^v \phi_1), \quad (4.5)$$



One easily checks that indeed

$$\text{Hom}_{\text{Bimod}_2}(L(M_0), N) \simeq \text{Hom}_{\text{Glob}_2}(M_0, U(N)).$$

4.3.2. The 2-bimodule $M(C, D)(F, G)(\eta, \theta)$. Let C, D be \mathbb{k} -linear bicategories,

$$F, G: C \rightarrow D$$

strong lax functors, $\eta, \theta: F \Rightarrow G$ two strong lax natural transformations.

We associate with this data the following 2- C -bimodule $M(C, D)(F, G)(\eta, \theta)$:

$$M(C, D)(F, G)(\eta, \theta)(f, g) = D(\eta(y) \circ F(f), G(g) \circ \theta(x)), \tag{4.7}$$

where $f, g \in C(x, y)$.


Lemma 4.8. *In notations as above, $M(C, D)(F, G)(\eta, \theta)$ is a 2-bimodule over C .*

Proof. Let $m \in M(C, D)(F, G)(\eta, \theta)(f, g)$.

For a 2-morphism $\alpha: f' \Rightarrow f$, the vertical composition $m \circ^v \alpha$ is defined as the vertical composition $m \circ^v (\eta(y) \circ F(\alpha))$ in D . Similarly, for $\beta: g \Rightarrow g'$, the vertical composition $\beta \circ^v m$ is defined as the vertical composition $(G(\beta) \circ \theta(x)) \circ^v m$.

For $f_0, g_0 \in C(w, x), \alpha \in C(f_0, g_0)$, define the 2-morphism $(m \circ^h \alpha)_0$ in D as the horizontal composition in $m \circ^h F(\alpha)$ in D post-composed vertically with the 2-morphism $\theta(x) \circ F(g_0) \Rightarrow G(g_0) \circ \theta(x_0)$ (whiskered by $G(g)$). Define the horizontal composition $m \circ^h \alpha = \psi \circ^v (m \circ^h \alpha)_0 \circ^v \phi$ where ϕ is the vertical composition of 2-arrows $\eta(y) \circ F(f \circ f_0) \xrightarrow{\sim} \eta(y) \circ (F(f) \circ F(f_0)) \xrightarrow{\sim} (\eta(y) \circ F(f)) \circ F(f_0)$, and ψ is the composition $G(g) \circ (G(g_0) \circ \eta(w)) \xrightarrow{\sim} (G(g) \circ G(g_0)) \circ \eta(w) \xrightarrow{\sim} G(g \circ g_0) \circ \eta(w)$.

For $f_1, g_1 \in C(y, z), \beta \in C(f_1, g_1)$, define the 2-morphism $(\beta \circ^v m)_0$ in D as the horizontal composition $m \circ^h G(\beta)$ in D pre-composed vertically with the 2-morphism $\eta(z) \circ F(f_1) \Rightarrow G(f_1) \circ \eta(y)$ (whiskered by $F(f)$). Define the horizontal composition $\beta \circ^v m = \psi \circ^v (\beta \circ^v m)_0 \circ^v \phi$, where ϕ is the vertical composition of 2-arrows $\eta(z) \circ F(f_1 \circ f) \xrightarrow{\sim} \eta(z) \circ (F(f_1) \circ F(f)) \xrightarrow{\sim} (\eta(z) \circ F(f_1)) \circ F(f)$, and ψ is the vertical composition $G(g_1) \circ (G(g) \circ \theta(x)) \xrightarrow{\sim} (G(g_1) \circ G(g)) \circ \theta(x) \xrightarrow{\sim} G(g_1 \circ g) \circ \theta(x)$.



One has to check the compatibilities with the associator (4.1)–(4.3) and the compatibility with the unit (4.4). It is left to the reader. ■

Note that the particular case $C = D, F = G = \text{Id}, \eta = \theta = \text{id}$ recovers the tautological 2-bimodule over C (for which $M_2(x, y) = C_2(x, y)$).

4.4. The category $\widehat{\Theta}_2$

In analogy with the category Θ_2 , we define here its “bicategorical extension” $\widehat{\Theta}_2$, playing an important role in definition of the complex $A(C, D)(F, G)(\eta, \theta)$ in the next section.

The objects of $\widehat{\Theta}_2$ are the same as the objects of Θ_2 , that is, the 2-level trees T . The morphisms are defined in terms of 2-globular sets T^* ; namely,

$$\widehat{\Theta}_2(S, T) = \text{Hom}_{\text{Bicat}}(\widehat{\omega}_2(S^*), \widehat{\omega}_2(T^*)),$$

where $\widehat{\omega}_2(D)$ denotes the free *bicategory* generated by the 2-globular set D , and the set $\text{Hom}_{\text{Bicat}}(-, -)$ stands for the set of *strict* functors of bicategories.

In sequel, we need somewhat a more explicit description of morphisms of $\widehat{\Theta}_2$.

First of all, we have the following explicit description of the bicategory $\widehat{\omega}_2(T^*)$ (which follows from the Coherence Theorem 4.6). The vertices are the 0-cells of T^* which we denote by $\{0, 1, \dots, k\}$. Consider an ordered set S with n elements. We consider an *extended* set $S(m)$, adding some m marked new elements to the n elements of S , along with a total order on $S(m)$ compatible with the total order on S . Define an *extended parenthesising* of S as a full parenthesising of $S(m)$, $m \geq 0$. The 1-morphism set $(\widehat{\omega}_2)_1(i, j)$ is empty if $i < j$ and is the set of *extended parenthesising*s of the ordered set of elementary 1-morphisms in the paths $\lambda(i, j)$ from i to j in $(T^*)_1$ (the marked elements indicate the positions of identity 1-morphisms). The two such paths are considered different also if they differ only by parenthesising or number and position of marked elements. The composition of 1-morphisms is defined naturally.

The set of 2-morphisms $\widehat{\omega}_2(\lambda(i, j), \lambda'(i, j))$ consists of a single element if $\lambda'(i, j)$ dominates $\lambda(i, j)$ (that is, in each column between i and j of T^* , the element of $\lambda'(i, j)$ in this column is \geq than the element of $\lambda(i, j)$ in this column, and the domination relation does not depend on the new marked elements) and is an empty set otherwise. The vertical and the horizontal compositions of 2-arrows are defined by the unique possible way.

It gives rise to the following wreath-product-like description of morphisms in $\widehat{\Theta}_2$.

A morphism Φ from $([k]; [n_1], \dots, [n_k])$ to $([\ell]; [m_1], \dots, [m_\ell])$ in $\widehat{\Theta}_2$ is given by $(\phi, \{\phi_i\})$

- (i) a morphism $\phi: [k] \rightarrow [\ell]$ in Δ ,
- (ii) for each $1 \leq i \leq k$ maps $\phi_i^j: [n_i] \rightarrow [n_j]$, $\phi(i-1) + 1 \leq j \leq \phi(i)$,
- (iii) for each $a \in [n_i]$ an extended parenthesising of the $(\phi(i) - \phi(i-1))$ -element set (considered as the set $\{\phi_i^j(a)\}_j$) (for given i , the number of added marked elements and the extended parenthesising may depend on $a \in [n_i]$).

The composition of morphism is defined naturally. The only new feature compared with the case of Θ_2 is that here we also compose parenthesising, in the operadic way.

We are interested in the explicit form of the left Kan extension $\text{Lan}_{p^{\text{op}}}(F)$ of a functor $F: \widehat{\Theta}_2^{\text{op}} \rightarrow \mathcal{E}$ along the functor

$$p^{\text{op}}: \widehat{\Theta}_2^{\text{op}} \rightarrow \Theta_2^{\text{op}},$$

where in our main examples \mathcal{E} is an abelian category.

Let $\phi: D \rightarrow D'$ be a map in Θ_2 ; we are interested in a lift $\hat{\phi}: D \rightarrow D'$ in $\hat{\Theta}_2$, so that $p(\hat{\phi}) = \phi$. It is clear that each map ϕ admits a non-empty (in fact, a finite) set of lifts. They depend on (multiple) insertions of the associator 2-arrows and the unit 2-arrows.

Lemma 4.9. *Let $F: \hat{\Theta}_2^{\text{op}} \rightarrow \mathcal{E}$ be a functor to an abelian category \mathcal{E} . Then, the left Kan extension $\text{Lan}_{p^{\text{op}}}(F)$ along the projection $p^{\text{op}}: \hat{\Theta}_2^{\text{op}} \rightarrow \Theta_2^{\text{op}}$ is given by*

$$\text{Lan}_{p^{\text{op}}}(F)(D) = F(D)/V_D,$$

where $V_D \subset F(D)$ is generated by the elements of the form $\hat{\phi}^*(F)(\xi) - \hat{\phi}^*(F)(\xi)$, where $\phi: D \rightarrow D_1$ is a map in Θ_2 , and $\hat{\phi}, \hat{\phi}^*$ are its two lifts to a morphism in $\hat{\Theta}_2$, $\xi \in F(D_1)$.

Proof. By the classical formula for a point-wise Kan extension, one has

$$\text{Lan}_{p^{\text{op}}}(F)(D) = \text{colim}_{D_1 \rightarrow D \in p^{\text{op}}/D} F(D_1).$$

The colimit is taken over the comma-category whose objects are $D_1 \in \hat{\Theta}_2^{\text{op}}$ and a map $p(D_1) = D_1 \rightarrow D$ in Θ_2^{op} ; the morphisms $\phi: (i_1: D_1 \rightarrow D) \rightarrow (i_2: D_2 \rightarrow D)$ are maps $\hat{\phi}: D_1 \rightarrow D_2$ in $\hat{\Theta}_2^{\text{op}}$ such that $i_1 = i_2 \circ p^{\text{op}}(\hat{\phi})$. Replacing the opposite categories by the non-opposite ones, the colimit is taken over the category I_D whose objects are morphisms $i_1: D \rightarrow D_1$ in $\hat{\Theta}_2$, and a morphism $\phi: (i_1: D \rightarrow D_1) \rightarrow (i_2: D \rightarrow D_2)$ is given by a map $\hat{\phi}: D_2 \rightarrow D_1$ in $\hat{\Theta}_2$ such that $i_1 = p(\hat{\phi}) \circ i_2$.

Clearly, any object $\phi: D \rightarrow D_1$ of I_D admits a morphism to $\text{id}: D \rightarrow D$ (one has to take any lift $\hat{\phi}: D \rightarrow D_1$ of ϕ). That is, $\text{id}: D \rightarrow D$ is a generalised final object: any other object admits a (non-unique) morphism to it. It follows that the colimit is a quotient of $F(D)$. One easily gets the description of the quotient given in the statement. ■

Now we want to get a more explicit description of the quotient in the statement of Lemma 4.9. It turns out that we easily can restrict ourselves with some “minimal” maps $\phi: D \rightarrow D_1$ in Θ_2 . We call them “standard maps”. There are two types of them: the associator morphism type and the unit morphism type.

The associator type maps. Let $D = ([k]; [n_1], \dots, [n_k])$,

$$D_1 = ([k + 2]; [n_1], [n_2], \dots, [n_a], [n_a], [n_a], [n_{a+1}], \dots, [n_k]).$$

Consider the map

$$\Phi = (\phi, \{\phi_i^j\}): D \rightarrow D_1,$$

where $\phi(0) = 0, \dots, \phi(a - 1) = a - 1, \phi(a) = a + 2, \dots, \phi(k) = k + 2, \phi_i^j = \text{id}$ for all i and j (clearly one has a unique j for $i \neq a$, and there are $j = a, a + 1, a + 2$ for $i = a$). Consider a lift $\hat{\Phi}$ for which $\phi_a(\ell)$ is either $(\ell_a \ell_{a+1}) \ell_{a+2}$ or $\ell_a(\ell_{a+1} \ell_{a+2})$ (the choice depends on $\ell \in [n_a]$) (we denote by ℓ_j the element $\ell \in [n_j]$). One has to specify the image of the two-morphisms. Clearly, the two-morphisms corresponding to 1-morphisms in $[n_i]$ (considered as the interval category I_{n_i}) are defined as the corresponding two-morphisms,

for $i \neq a$. When $i = a$, the prescription is as follows: the image of the 2-morphism corresponding to $m_\ell := \ell \rightarrow \ell + 1$ in $[n_a]$ is

$$\left\{ \begin{array}{l} ((m_\ell)_a \circ_h (m_\ell)_{a+1}) \circ_h (m_\ell)_{a+2} \\ \quad \text{if both } \phi_a(\ell) \text{ and } \phi_a(\ell + 1) \text{ are of type } (**)* \\ (m_\ell)_a \circ_h ((m_\ell)_{a+1} \circ_h (m_\ell)_{a+2}) \\ \quad \text{if both } \phi_a(\ell) \text{ and } \phi_a(\ell + 1) \text{ are of type } *(**) \\ \alpha \circ_v (((m_\ell)_a \circ_h (m_\ell)_{a+1}) \circ_h (m_\ell)_{a+2}) \\ \quad \text{if } \phi_a(\ell) = (\ell_a \ell_{a+1}) \ell_{a+2}, \phi_a(\ell + 1) = (\ell + 1)_a ((\ell + 1)_{a+1} (\ell + 1)_{a+2}) \\ (((m_\ell)_a \circ_h (m_\ell)_{a+1}) \circ_h (m_\ell)_{a+2}) \circ_v \alpha^{-1} \\ \quad \text{if } \phi_a(\ell) = \ell_a (\ell_{a+1} \ell_{a+2}), \phi_a(\ell + 1) = ((\ell + 1)_a (\ell + 1)_{a+1}) (\ell + 1)_{a+2} \end{array} \right.$$

(note that we use here non-conventional writing from the left to the right for the composition of 1-morphisms and for the horizontal composition of 2-morphisms).

We refer to such morphisms as Type A ones.

The unit type maps. Here, $D = ([k]; [n_1], \dots, [n_k])$, $\Phi = \text{Id}: D \rightarrow D$ in Θ_2 , but $\widehat{\Phi}: D \rightarrow D$ in $\widehat{\Theta}_2$ is defined by insertion of several marked points to $[k]$ (corresponding to the identity 1-morphisms). Accordingly, each ϕ_i^j is the identity modulo the marked points, but with their presence it is given by some parenthesisising (of words only 1 element of which is not marked). The images of elementary 2-morphisms are defined accordingly, pre- or post-composing vertically with the unit 2-morphisms or their inverse ones.

We refer to such morphisms as Type U ones.

Proposition 4.10. *In the notations of Lemma 4.9,*

$$\text{Lan}_{p^{\text{op}}}(F)(D) = F(D)/W_D,$$

where $W_D \subset F(D)$ is generated by the elements of the form $\widehat{\phi}^*(F)(\xi) - \widehat{\phi}^*(F)(\xi)$, where $\phi: D \rightarrow D_1$ is a map in Θ_2 listed in Type A or in Type U, and $\widehat{\phi}, \widehat{\phi}$ are its two lifts to a morphism in $\widehat{\Theta}_2$, listed in Type A or Type U, $\xi \in F(D_1)$.

Proof. It follows from Lemma 4.9 and (a rather evident) observation that any morphism $\widehat{\phi}: D \rightarrow D_1$ factors as $D \xrightarrow{\widehat{\psi}} D' \rightarrow D_1$, where $\widehat{\psi}$ is of Type A or of Type U. ■

5. The 2-cocellular vector space $A(C, D)(F, G)(\eta, \theta)$

5.1. The Davydov–Yetter complex and an attempt of generalisation

Let C, D be \mathbb{k} -linear monoidal categories (that is, \mathbb{k} -linear bicategories with a single object), $F: C \rightarrow D$ a strong monoidal functor. For $n \geq 1$, denote

$$F^{\otimes n}: C^{\otimes n} \rightarrow D$$

defined on objects as

$$F^{\otimes n}(X_1, \dots, X_n) = F(X_1) \otimes_D F(X_2) \otimes_D \cdots \otimes_D F(X_n)$$

(one has to fix any parenthesising of the right-hand side, for example, from the left to the right).

Define

$$\begin{aligned} A^n(F) &= \text{Nat}(F^{\otimes n}, F^{\otimes n}) \\ &= \prod_{X_1, \dots, X_n \in C} D(F(X_1) \otimes \cdots \otimes F(X_n), F(X_1) \otimes \cdots \otimes F(X_n))_{\text{Nat}}, \end{aligned}$$

where Nat stands for (linear) natural transformations.

The assignment $[n] \rightsquigarrow A^n(F)$ gives rise to a cosimplicial object. Its non-normalised dg totalisation is called *the Davydov–Yetter complex* of $F: C \rightarrow D$. Let us recall this definition, with notations for simplicial coface and codegeneracy maps from the beginning of Section 2.1. Let $\Psi \in \widehat{A}_n(F)$. The elementary coface maps $\partial^i: [n-1] \rightarrow [n]$, $1 \leq i \leq n-1$, act by plugging $X_i \otimes X_{i+1}$ in place of the i -th argument X_i of Ψ , followed by the application of the colax-map $F(X_i \otimes X_{i+1}) \rightarrow F(X_i) \otimes F(X_{i+1})$ and rearranging the parentheses (note that by the MacLane coherence theorem one need not specify the way by which the parentheses are rearranged, as any two such maps are equal). The extreme coface map ∂^0 acts by $\Psi \mapsto \text{id}_{F(X_0)} \otimes \Psi(X_1 \otimes \cdots \otimes X_k)$, and the other extreme coface map ∂^k acts by $\Psi \mapsto \Psi \otimes \text{id}_{F(X_k)}$, followed by the necessary reparenthesising. The codegeneracy map ε^i acts on k -cochain Ψ by plugging the monoidal unit e to the i -th position of Ψ , thus decreasing the number of arguments by 1, followed by the necessary rearrangements. The reader is referred to [14, Chap. 7] or [6] for a more detailed description.

Now the question is: does the construction still give rise to a cosimplicial object in \mathcal{V} when the polynaturality condition of Davydov–Yetter is dropped? The answer is negative, unless the monoidal category is strict (the associator and the unit maps are the identity maps), because the relations in Δ are no longer respected. Denote the action of elementary cosimplicial operators as above by \mathcal{O} . Then, for instance, the actions of $\mathcal{O}(\partial^{i+1}) \circ \mathcal{O}(\partial^i)$ differ from $\mathcal{O}(\partial^i) \circ \mathcal{O}(\partial^i)$ only by the i -th argument, which is $(X_i \otimes X_{i+1}) \otimes X_{i+2}$ for the first composition, and $X_i \otimes (X_{i+1} \otimes X_{i+2})$ for the second one. These two expressions are mapped one to another by the associator map α . Therefore, in order for the relation $\partial^{i+1}\partial^i = \partial^i\partial^i$ to be respected under the action \mathcal{O} , one has to *require the naturality with respect to α* on the i -th argument. Similar considerations are applied to the unit map and degeneracy operators.

It is clear from this reasoning that the naturality under all monoidal *structure morphisms* maps (that is, compositions of products, the associator, the unit maps, and its inverse, with the identity maps on some factors) acting in each argument is the *minimal* naturality condition one has to impose on the cochains to get a cosimplicial object in \mathcal{V} .

A drawback of the Davydov–Yetter complex, from the point of view of the deformation theory, is that it controls only the deformation of the associator. The full deformation

theory of a linear monoidal category should also control deformations of the underlying linear category (which are controlled by the Hochschild cochain complex of this underlying category) and the action of *morphisms* on the monoidal product (the monoidal product on objects is non-linear data and thus is assumed to remain unchanged under the deformation). So our goal is to define a “bigger” bicomplex of Davydov–Yetter type, whose “vertical” differential is of the Hochschild type, and whose “horizontal” differential is of Davydov–Yetter type. The classical Davydov–Yetter complex is obtained by a sort of truncation. This truncation is the kernel of the vertical (Hochschild type) differential of the 0-th row of the bicomplex (so the truncated complex is a subcomplex of the 0-th row of our bicomplex). Taking this kernel affects imposing the Davydov–Yetter naturality condition.

5.2. A functor $\text{Bar}: \widehat{\Theta}_2 \rightarrow \text{Bimod}_2$

Let C be small dg bicategory. In this subsection, we define a functor

$$\text{Bar}(C): \widehat{\Theta}_2^{\text{op}} \rightarrow \text{Bimod}_2(C)$$

(playing the role of a bicategorical bar-complex), as follows.

Let $T = ([k]; [n_1], \dots, [n_k])$ be an object of Θ_2 . Let $x', y' \in C_0$, $f_0, g_0 \in C_1(x', y')$. Define

$$(M_T)_2(f_0, g_0) = \bigoplus_{\substack{x_0, \dots, x_k \in C_0, x_0 = x, x_k = y \\ f_{ij} \in C_1(x_{i-1}, x_i), j = 0 \dots n_i \\ f_{\circ \min} = f_0, f_{\circ \max} = g_0}} \bigotimes_{i=1 \dots k} C(f_{i,j-1}, f_{i,j}), \tag{5.1}$$

where

$$\begin{aligned} f_{\circ \min} &:= f_{k0} \circ (f_{k-1,0} \circ (\dots (f_{20} \circ f_{10}) \dots)), \\ f_{\circ \max} &= f_{kn_k} \circ (f_{k-1,n_{k-1}} \circ (\dots (f_{2n_2} \circ f_{1n_1}) \dots)). \end{aligned} \tag{5.2}$$

It is considered the second component of the enriched 2-globular set M_T such that $(M_T)_{\leq 1} = C_{\leq 1}$.

Next, define a dg 2-bimodule $\text{Bar}(C)_T$ over C as

$$\text{Bar}(C)_T = L(M_T), \tag{5.3}$$

where $L: \text{Glob}_2 \rightarrow \text{Bimod}_2$ is the left adjoint functor to the forgetful functor, which was discussed in Section 4.3.1.

Explicitly, for $x, y \in C_0$, $f, g \in C_1(x, y)$, one has

$$\begin{aligned} &\text{Bar}(C)_T(f, g) \\ &= \bigoplus_{\alpha \in C_1(x, x'), \beta \in C_1(y', y)} C(f, \beta \circ (f_0 \circ \alpha)) \otimes_{\mathbb{k}} (M_T)_2(f_0, g_0) \otimes_{\mathbb{k}} C(\beta \circ (g_0 \circ \alpha), g) / \sim, \end{aligned}$$

where \sim is the relation in (4.5).

Proposition 5.1. *In the notations as above, the assignment $T \rightsquigarrow \text{Bar}(C)_T$ is an object part of a functor*

$$\text{Bar}(C): \widehat{\Theta}_2 \rightarrow \text{Bimod}_2(C).$$

Remark 5.2. It will be clear from the proof that the assignment $T \rightsquigarrow M_T, \text{Ob}(\widehat{\Theta}_2) \rightarrow \text{Glob}_2$ cannot be extended to a functor $\widehat{\Theta}_2 \rightarrow \text{Glob}_2$. So it becomes a functor only after the application of $L(-)$.

Proof. If we considered an ordinary (set-enriched) bicategory C , we would use the fact that the set of strict functors $\text{Hom}_{\text{Bicat}}^{\text{str}}(\widehat{\omega}_2(T^*), C)$ is functorial with respect to strict bicategory maps $\widehat{\Theta}_2(S, T) = \text{Hom}_{\text{Bicat}}^{\text{str}}(\widehat{\omega}_2(S^*), \widehat{\omega}_2(T^*))$. On the other hand, a strict functor $\widehat{\omega}_2(T) \rightarrow C$ is the same as a map of 2-globular sets $T^* \rightarrow U(C)$, where $U(C)$ denotes the underlying 2-globular object. A map of 2-globular sets $T^* \rightarrow U(C)$ is defined by its values on the sets of i -cells, $i = 0, 1, 2$. The set of these maps is “very closed” to our M_T . More precisely, the set of maps of 2-globular sets $T^* \rightarrow U(C)$ is

$$M'_T = \coprod_{\substack{x_0, \dots, x_k \in C_0 \\ f_{ij} \in C_1(x_{i-1}, x_i), j=0 \dots n_i}} \prod_{\substack{i=1 \dots k \\ j=1 \dots n_i}} C(f_{i,j-1}, f_{i,j})$$

(with dropped condition $f_{\circ \min} = f_0, f_{\circ \max} = g_0$). ■

It follows from the discussion just above that the assignment $T \rightsquigarrow M'_T$ gives rise to a functor $M'(C): \widehat{\Theta}_2^{\text{op}} \rightarrow \mathcal{V}$.

Note that if a map $p(\Phi) \in \Theta_2$ is not dominant (that is, does not necessarily preserve all minima and maxima), the projection along several factors is used to define an action of Φ .

We are mostly interested in the case when the bicategory C is dg \mathbb{k} -linear, and in this case, the projections of $V \otimes W$ to V and to W are not defined. Moreover, it is not true that, for a \mathbb{k} -linear bicategory C ,

$$M''_T = \bigoplus_{\substack{x_0, \dots, x_k \in C_0 \\ f_{ij} \in C_1(x_{i-1}, x_i), j=0 \dots n_i}} \bigotimes_{\substack{i=1 \dots k \\ j=1 \dots n_i}} C(f_{i,j-1}, f_{i,j})$$

is $\text{Hom}_{\text{Bicat}(\mathbb{k})}^{\text{str}}(\mathbb{k}\widehat{\omega}_2(T^*), C)$, where $\mathbb{k}\widehat{\omega}_2(T^*)$ denotes the free \mathbb{k} -linear bicategory generated by T^* made \mathbb{k} -linear in dimension 2. Indeed, a homogeneous element of a complex of vector space V is not the same as $C^*(\mathbb{k})(\mathbb{k}, V)$.

On the other hand, the same formulas with the direct product of sets replaced by the tensor product of complexes of \mathbb{k} -vector space define an action of $\widehat{\Theta}_2^{\text{dom}}$ on M'' , where the upper script dom stands for the subcategory of morphisms Φ such that $p(\Phi)$ is dominant.

Now instead of non-existing projections we use the “from inside” action in the free 2-bimodule case (it can be considered a 2-dimensional version of the two-sided bar-construction). It gives an action of general (possibly non-dominant) morphisms. Also, the parenthesis in (5.2) is fixed, so after an application of a morphism in $\widehat{\Theta}_2$, one may get a wrong parenthesis in $f_{\circ \min}$ and $f_{\circ \max}$. The same holds when extra identity 1-morphisms are present after an application of a morphism in $\widehat{\Theta}_2$. In these cases, we have

to reduce $f_{\circ \min}$ and $f_{\circ \max}$ to the standard parenthesisating and without extra identity morphisms, by applying the associator and the unit 2-morphisms “from inside” on the (upper and lower) 2-bimodule arguments. The reader easily checks that this description indeed gives rise to a functor $\text{Bar}(C): \widehat{\Theta}_2 \rightarrow \text{Bimod}_2(C)$.

5.3. The 2-cocellular complex $A(C, D)(F, G)(\eta, \theta)$

Let C, D be dg bicategories, $F, G: C \rightarrow D$ strong functors, $\eta, \theta: F \Rightarrow G$ strong natural transformations. We associate with this data a complex $A(C, D)(F, G)(\eta, \theta) \in C^\bullet(\mathbb{k})$.

In Section 5.2, we associated with a dg bicategory C a functor

$$\text{Bar}(C): \widehat{\Theta}_2^{\text{op}} \rightarrow \text{Bimod}_2(C).$$

Let $p: \widehat{\Theta}_2 \rightarrow \Theta_2$ be the projection which is the identity on objects. The left Kan extension $\text{Lan}_{p^{\text{op}}}(\text{Bar}(C))$ is a functor $\Theta_2^{\text{op}} \rightarrow \text{Bimod}(C)$.

Define

$$\begin{aligned} A^\Theta(C, D)(F, G)(\eta, \theta)_T \\ = \underline{\text{Bimod}}_2(C)((\text{Lan}_{p^{\text{op}}} \text{Bar}(C))_T, M(C, D)(F, G)(\eta, \theta)) \in C^\bullet(\mathbb{k}). \end{aligned}$$

Here, the Hom is taken internally with respect to complexes of vector spaces, so it takes values in $C^\bullet(\mathbb{k})$, and the 2- C -bimodule $M(C, D)(F, G)(\eta, \theta)$ was defined in Section 4.3.2.

For the final object $T_* = ([0]; \emptyset)$, we set

$$A^\Theta(C, D)(F, G)(\eta, \theta)_{T_*} = \prod_{X \in \text{Ob}(C)} D_2(\eta_X, \theta_X).$$

Finally, define

$$A(C, D)(F, G)(\eta, \theta) = \text{Tot}_{T \in \Theta_2} A^\Theta(C, D)(F, G)(\eta, \theta)_T.$$

In Sections 5.5 and 5.6 below, we unwind this definition making it more explicit.

We consider this complex as “derived modifications” $\eta \Rightarrow \theta$ (see Definition 4.4 for the definition of a classical modification).

Remark 5.3. Likewise, for the dg 1-categories C, D , and two dg functors $F, G: C \rightarrow D$, the Hochschild complex

$$\text{Hoch}^\bullet(C, D(F(-), G(=))) = \underline{\text{Bimod}}(C)(\text{Bar}(C), D(F(-), G(=)))$$

is interpreted as “derived natural transformations” $F \Rightarrow G$, in the sense that closed degree 0 0-cochains correspond to classical dg natural transformations. Here, $\text{Bar}(C)$ is the classical bar-complex of the dg category C , taking values in C -bimodules.

More precisely, one has the following proposition.

Proposition 5.4. *Assume that C, D are dg bicategories, $F, G: C \rightarrow D$ strong functors, $\eta, \theta: F \Rightarrow G$ strong dg transformations. Then, the vector space $V \subset A^\Theta(C, D)(F, G)(\eta, \theta)_{T_*}$ of degree 0 closed elements in $A(C, D)(F, G)(\eta, \theta)$ is isomorphic to the vector space of degree 0 modifications $\eta \Rightarrow \theta$.*

Proof. A general element Ψ of $A^\Theta(C, D)(F, G)(\eta, \theta)_{T_*}$ belongs to $\prod_{X \in C} D(\eta_X, \theta_X)$. Its boundary belongs to the arity $T_1 = ([1]; [0])$. A general cochain in $A^\Theta(C, D)(F, G)(\eta, \theta)_{T_1}$ is an element of

$$\prod_{f: X \rightarrow Y \in C_1} D_2(\eta_Y \circ F(f), G(f) \circ \theta_X).$$

Let Ψ have the components $\Psi_X \in D_2(\eta_X, \theta_X)$, $X \in C_0$. Then, $d\Psi$ has the components $(d\Psi)_f$, $f \in C_1(X, Y)$, which are

$$(d\Psi)_f = \Psi_Y \circ F(f) - G(f) \circ \Psi_X.$$

(The two summands come from the two morphisms $T_\emptyset \rightarrow T_1$ in Θ_2 corresponding to the two morphisms $[0] \rightarrow [1]$ in Δ ; they are D_{\min} and D_{\max} in the notations of Section 2.6, (F4).)

Then, $(d\Psi)_f = 0$ for any $f \in C(X, Y)$ is precisely the condition for Ψ being a modification. ■

5.4. Some properties of the category of 2-bimodules

Lemma 5.5. *Let C be a \mathbb{k} -linear bicategory, \mathbb{k} a field. The category $\text{Bimod}_2(C)$ of 2-bimodules over C is abelian \mathbb{k} -linear. The 2-bimodules $L(M)$ are projective, where L is the left adjoint to the forgetful functor $U: \text{Bimod}_2(C) \rightarrow \text{Glob}_2(\mathbb{k})$; see Section 4.2.*

Proof. Consider the truncation functor

$$\xi: \text{Glob}_2(\mathbb{k}) \rightarrow \text{Glob}_1,$$

where $\text{Glob}_2(\mathbb{k})$ denotes the category of 2-globular objects enriched in $\text{Vect}(\mathbb{k})$ in degree 2 (so that the degree 0 and the degree 1 components are sets). It is clear that the comma-category $\xi \setminus Y$ is an abelian \mathbb{k} -linear category for any $Y \in \text{Glob}_1$. One can consider the forgetful functor U as a functor $U_\xi: \text{Bimod}_2(C) \rightarrow \xi \setminus C_{\leq 1}$.

Let $f: M \rightarrow N$ be a morphism of 2-bimodules over C . The kernel and the cokernel of $U_\xi(f)$ have natural structures of C -2-bimodules. Then, the first statement follows from the fact that the comma-category $\xi \setminus C_{\leq 1}$ is abelian.

For the second statement, there is an adjunction

$$L_\xi: \xi \setminus C_{\leq 1} \xrightleftharpoons{\quad} \text{Bimod}_2(C): U_\xi,$$

where the free 2-bimodule functor L_ξ is the same as the functor L defined in Section 4.2. The equality

$$\text{Bimod}_2(C)(L(M), N) = (\xi \setminus C_{\leq 1})(M, U_\xi(N))$$

and the projectivity of $M \in \xi \setminus C_{\leq 1}$ proves the second statement. ■

Remark 5.6. Consider $B(C) = \text{Real}_{T \in \Theta_2} \text{Lan}_{p^{\text{op}}} \text{Bar}(C)_T$, where *Real* stands for the Θ_2 -realisation. It is *not* true that $B(C)$ is a resolution of the tautological 2-bimodule C over C . Consequently, Lemma 5.5 does not imply that

$$A(C, D)(F, G)(\eta, \theta) = \text{RHom}_{\text{Bimod}_2(C)}(C, M(C, D)(F, G)(\eta, \theta)). \quad (*)$$

As we show now, $(*)$ cannot be true.

Indeed, consider the most limit case of a \mathbb{k} -linear bicategory, namely, when it has a single object and a single 1-morphism (the identity morphism *id* of this object). Such a category is just a commutative algebra $X = C_2(\text{id}, \text{id})$ over \mathbb{k} . Our results of Section 7 show that $H^3(A(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id}))$ computes infinitesimal deformations of X as a commutative \mathbb{k} -algebra. When X is singular, such cohomology may not vanish. On the other hand, the category of 2-bimodules over such C is just the category of left X -modules. If $(*)$ was true, we would have that this cohomology is $H^3(\text{RHom}_{\text{Mod}(X)}(X, X))$, but the latter cohomology vanishes for any X .

It would be interesting to compare, for such a 1-terminal bicategory C , the cohomology of $A(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id})$ with the Andre–Quillen cohomology of X .

5.5. The complex $A(C, D)(F, G)(\eta, \theta)$: an explicit description

In this subsection, we provide a more direct description of $A(C, D)(F, G)(\eta, \theta)$ and of the differential on it, defined in Section 5.3, where C, D are \mathbb{k} -linear bicategories, F, G strong functors, η, θ strong natural transformations. The reader easily checks that the description given below agrees with the one given in Section 5.3.

We define $A(C, D)(F, G)(\eta, \theta)_T$, for $T = ([k]; [n_1], \dots, [n_k])$, as a graded subspace (subcomplex) of $\hat{A}(C, D)(F, G)(\eta, \theta)_T$; the latter is defined as follows.

For $x, y \in C_0, f, g \in C_1(x, y), n \geq 0$, denote

$$C_{x,y}^n(f, g) = \bigoplus_{\substack{f_0, \dots, f_n \in C_1(x,y) \\ f_0=f, f_n=g}} C_1(f_{n-1}, f_n) \otimes_{\mathbb{k}} C_1(f_{n-2}, f_{n-1}) \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} C_1(f_0, f_1)$$

(when $n = 0$, we must have $f = g$). We use the notation θ for an element in $C_{x,y}^n(f, g)$, and then $\theta(i) = f_i$. Also, we use the notation $\theta(\sigma_1, \dots, \sigma_n)$ for such an element, where $\sigma_i \in C_2(f_{i-1}, f_i)$.

We will need one more notation. Let $x, y, z, w \in C_0, f_i \in C_1(x, y), f'_i \in C_1(y, z), f''_i \in C_1(z, w), \sigma_i \in C_2(f_{i-1}, f_i), \sigma'_i \in C_1(f'_{i-1}, f'_i), \sigma''_i \in C_1(f''_{i-1}, f''_i)$. Denote $\Sigma_i^- = \sigma''_i \circ^h (\sigma'_i \circ^h \sigma_i), \Sigma_i^+ = (\sigma'_i \circ^h \sigma_i) \circ^h \sigma_i$. Then, we denote by $\theta^{(3)}(\Sigma_1, \dots, \Sigma_k)$ the corresponding element in $C_{x,w}^k$, where each Σ_i is either Σ_i^+ or Σ_i^- . In particular, we may consider $\theta^{(3)}(\Sigma_1^-, \dots, \alpha \circ^h \Sigma_i^-, \Sigma_{i+1}^+, \dots, \Sigma_k^+)$ where α is the associator.

Then, for $T = ([k]; [n_1], \dots, [n_k])$, one has

$$\begin{aligned} & \hat{A}(C, D)(F, G)(\eta, \theta)_T \\ &= \prod_{\substack{x_0, \dots, x_k \in C_0 \\ f_i, g_i \in C_1(x_{i-1}, x_i)}} \underline{\text{Hom}}_{\mathbb{k}}(C_{x_0, x_1}^{n_1}(f_1, g_1) \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} C_{x_{k-1}, x_k}^{n_k}(f_k, g_k), \\ & \quad D(\eta_{x_k} \circ f_{\text{tot}}^F, g_{\text{tot}}^G \circ \theta_{x_0})), \end{aligned}$$

where

$$\begin{aligned} f_{\text{tot}}^F &= F(f_k) \circ (F(f_{k-1}) \circ (\dots (F(f_2) \circ F(f_1)) \dots)), \\ g_{\text{tot}}^G &= G(g_k) \circ (G(g_{k-1}) \circ (\dots (G(g_2) \circ G(g_1)) \dots)). \end{aligned}$$

For $T = ([0]; \emptyset)$, we set

$$\widehat{A}(C, D)(F, G)_{([0]; \emptyset)} = \prod_{x \in C_0} D_2(\eta_x, \theta_x).$$

Now, $A(C, D)(F, G)(\eta, \theta)_T$ is a subcomplex of $\widehat{A}(C, D)(F, G)(\eta, \theta)_T$ formed by the cochains $\Psi \in \widehat{A}(C, D)(F, G)(\eta, \theta)$ depicted by the following conditions.

The conditions are divided into two groups: the associator and the unit map conditions. They are direct consequences of Proposition 4.10.

The associator conditions read:

$$\begin{aligned} &\Psi(\theta_1 \otimes \dots \otimes \theta_{j-1} \otimes \theta_j^{(3)}(\Sigma_1^-, \dots, \alpha \circ^v \Sigma_i^-, \Sigma_{i+1}^+, \dots, \Sigma_{n_j}^+) \otimes \theta_{j+1} \otimes \dots \otimes \theta_k) \\ &= \Psi(\theta_1 \otimes \dots \otimes \theta_{j-1} \otimes \theta_j^{(3)}(\Sigma_1^-, \dots, \Sigma_i^-, \Sigma_{i+1}^+ \circ^v \alpha, \Sigma_{i+2}^+, \dots, \Sigma_{n_j}^+) \\ &\quad \otimes \theta_{j+1} \otimes \dots \otimes \theta_k), \quad 1 \leq j \leq k, \quad 1 \leq i \leq n_j - 1, \\ &\Psi(\theta_1 \otimes \dots \otimes \theta_{j-1} \otimes \theta_j^{(3)}(\Sigma_1^+, \dots, \alpha^{-1} \circ^v \Sigma_i^+, \Sigma_{i+1}^-, \dots, \Sigma_k^-) \otimes \theta_{j+1} \otimes \dots \otimes \theta_k) \\ &= \Psi(\theta_1 \otimes \dots \otimes \theta_{j-1} \otimes \theta_j^{(3)}(\Sigma_1^+, \dots, \Sigma_i^+, \Sigma_{i+1}^- \circ^v \alpha^{-1}, \Sigma_{i+2}^-, \dots, \Sigma_k^-) \\ &\quad \otimes \theta_{j+1} \otimes \dots \otimes \theta_k), \quad 1 \leq j \leq k, \quad 1 \leq i \leq n_j - 1, \\ &\Psi(\theta_1 \otimes \dots \otimes \theta_j^{(3)}(\Sigma_1^+ \circ^v \alpha, \Sigma_2^+, \dots, \Sigma_{n_j}^+) \otimes \dots \otimes \theta_k) \\ &= \Psi(\theta_1 \otimes \dots \otimes \theta_j^{(3)}(\Sigma_1^+, \dots, \Sigma_{n_j}^+) \otimes \dots \otimes \theta_k) \circ^v \tilde{\alpha}_j, \quad 1 \leq j \leq k, \tag{5.4} \\ &\Psi(\theta_1 \otimes \dots \otimes \theta_j^{(3)}(\Sigma_1^-, \dots, \Sigma_{n_j-1}^-, \alpha \circ^v \Sigma_{n_j}^-) \otimes \dots \otimes \theta_k) \\ &= \tilde{\alpha}_j \circ^v \Psi(\theta_1 \otimes \dots \otimes \theta_j^{(3)}(\Sigma_1^-, \dots, \Sigma_{n_j}^-) \otimes \dots \otimes \theta_k), \quad 1 \leq j \leq k, \\ &\Psi(\theta_1 \otimes \dots \otimes \theta_j^{(3)}(\Sigma_1^- \circ^v \alpha^{-1}, \Sigma_2^-, \dots, \Sigma_{n_j}^-) \otimes \dots \otimes \theta_k) \\ &= \Psi(\theta_1 \otimes \dots \otimes \theta_j^{(3)}(\Sigma_1^-, \Sigma_2^-, \dots, \Sigma_{n_j}^-) \otimes \dots \otimes \theta_k) \circ^v \tilde{\alpha}_j^{-1}, \quad 1 \leq j \leq k, \\ &\Psi(\theta_1 \otimes \dots \otimes \theta_j^{(3)}(\Sigma_1^+, \dots, \alpha^{-1} \circ^v \Sigma_{n_j}^+) \otimes \dots \otimes \theta_k) \\ &= \tilde{\alpha}_j^{-1} \circ^v \Psi(\theta_1 \otimes \dots \otimes \theta_j^{(3)}(\Sigma_1^+, \dots, \Sigma_{n_j}^+) \otimes \dots \otimes \theta_k), \quad 1 \leq j \leq k, \end{aligned}$$

where α is the associator 2-morphism, and $\tilde{\alpha}_j$ (resp., $\tilde{\alpha}_j^{-1}$) stand for the whiskering of α (resp., of α^{-1}) acting on j -th output with suitable identity 2-morphisms.

When $n_j = 0$ for some j , the list above for this j gives (and is reduced to) the following equations:

$$\begin{aligned} &\tilde{\alpha}_j \circ^v \Psi(\theta_1 \otimes \dots \otimes \theta^{(3)}(\text{id}_{f'' \circ (f' \circ f)}) \otimes \dots \otimes \theta_k) \\ &= \Psi(\theta_1 \otimes \dots \otimes \theta^{(3)}(\text{id}_{(f'' \circ f) \circ f}) \otimes \dots \otimes \theta_k) \circ^v \tilde{\alpha}_j, \end{aligned}$$

$$\begin{aligned} & \tilde{\alpha}_j^{-1} \circ^v \Psi(\theta_1 \otimes \cdots \otimes \theta_j^{(3)}(\text{id}_{(f'' \circ f') \circ f}) \otimes \cdots \otimes \theta_k) \\ & = \Psi(\theta_1 \otimes \cdots \otimes \theta_j^{(3)}(\text{id}_{f'' \circ (f' \circ f)}) \otimes \cdots \otimes \theta_k) \circ^v \tilde{\alpha}_j^{-1} \end{aligned} \quad (5.5)$$

(compare with the discussion in Section 5.1).

The unit map conditions read:

$$\begin{aligned} & \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\sigma_1, \dots, \lambda^{-1} \circ^v \sigma_i, \text{id} \circ^h \sigma_{i+1}, \dots, \text{id} \circ^h \sigma_{n_j}) \otimes \cdots \otimes \theta_k) \\ & = \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\sigma_1, \dots, \sigma_i, (\text{id} \circ^h \sigma_{i+1}) \circ^v \lambda^{-1}, \text{id} \circ \sigma_{i+1}, \dots, \text{id} \circ \sigma_{n_j}) \\ & \quad \otimes \cdots \otimes \theta_k), 1 \leq j \leq k \\ & \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\text{id} \circ^h \sigma_1, \dots, \text{id} \circ^h \sigma_{i-1}, \sigma_i \circ^v \lambda, \sigma_{i+1}, \dots, \sigma_{n_j})) \\ & = \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\text{id} \circ^h \sigma_1, \dots, \lambda \circ^v (\text{id} \circ^h \sigma_{i-1}), \sigma_i, \dots, \sigma_{n_j}) \\ & \quad \otimes \theta_{j+1} \otimes \cdots \otimes \theta_k), 1 \leq j \leq k \\ & \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\sigma_1 \circ^v \lambda, \sigma_2, \dots, \sigma_{n_j}) \otimes \cdots \otimes \theta_k) \\ & = \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\sigma_1, \dots, \sigma_{n_j}) \otimes \theta_{j+1} \otimes \cdots \otimes \theta_k) \circ^v \tilde{\lambda}_j, 1 \leq j \leq k \\ & \Psi(\theta_1 \otimes \cdots \otimes \theta_j((\text{id} \circ^h \sigma_1) \circ^v \lambda^{-1}, \text{id} \circ^h \sigma_2, \dots, \text{id} \circ^h \sigma_{n_j}) \otimes \cdots \otimes \theta_k) \\ & = \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\text{id} \circ^h \sigma_1, \dots, \text{id} \circ^h \sigma_{n_j}) \otimes \theta_{j+1} \otimes \cdots \otimes \theta_k) \circ^v \tilde{\lambda}_j^{-1}, 1 \leq j \leq k \\ & \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\text{id} \circ^h \sigma_1, \dots, \text{id} \circ^h \sigma_{n_j-1}, \lambda \circ^v (\text{id} \circ^h \sigma_{n_j}) \otimes \cdots \otimes \theta_k)) \\ & = \tilde{\lambda}_j \circ^v \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\text{id} \circ^h \sigma_1, \dots, \text{id} \circ^h \sigma_{n_j}) \otimes \theta_{j+1} \otimes \cdots \otimes \theta_k), 1 \leq j \leq k \\ & \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\sigma_1, \dots, \sigma_{n_j-1}, \lambda^{-1} \circ^v \sigma_{n_j}) \otimes \cdots \otimes \theta_k) \\ & = \tilde{\lambda}_j^{-1} \circ^v \Psi(\theta_1 \otimes \cdots \otimes \theta_j(\sigma_1, \dots, \sigma_{n_j}) \otimes \cdots \otimes \theta_k), 1 \leq j \leq k, \end{aligned} \quad (5.6)$$

where $\lambda_f: \text{id} \circ^h f \rightarrow f$ is the left unit map, and $\tilde{\lambda}_j$ (resp., $\tilde{\lambda}_j^{-1}$) stands for whiskering of λ acting on the j -th factor of the output with the suitable identity maps.

As well, one has similar relations for the right unit map ρ , which we do not write down here.

The next point is to extend the assignment

$$T \rightsquigarrow A(C, D)(F, G)(\eta, \theta)$$

to a functor

$$A(C, D)(F, G)(\eta, \theta): \Theta_2 \rightarrow C^*(\mathbb{k}).$$

Once again, the existence of such functor follows from the construction given in Section 5.3, and our task here is to provide explicit formulas for the action of morphisms of Θ_2 .

We restrict ourselves to the case when C and D are *strict* 2-categories, F, G *strict* functors, and η, θ *strict* natural transformations. The reason is that even this case shows the nature of the aforementioned action, but essentially simplifies the formulas. The formulas for the Θ_2 -action in the general case differ by numerous conjunctions with the structure 2-isomorphisms.

5.6. An explicit description of the complex $A(C, D)(F, G)(\eta, \theta)$, II

5.6.1. Let C be a dg bicategory, $x, y \in C_0$, and let $\phi: [m] \rightarrow [n]$ be a morphism in Δ . We associate with ϕ a map of complexes

$$C_{x,y}(\phi): C_{x,y}^n(f_0, f_n) \rightarrow C_2 \underset{A_L(\phi)}{(f_{\phi(m)}, f_n)} \otimes_{\mathbb{k}} C_{x,y}^m \underset{M(\phi)}{(f_{\phi(0)}, f_{\phi(m)})} \otimes_{\mathbb{k}} C_2 \underset{A_R(\phi)}{(f_0, f_{\phi(0)})} \quad (5.7)$$

as follows.

We use the notation $\sigma_n \otimes \dots \otimes \sigma_1$ for an element in $C_{x,y}^n(f_0, f_n)$, where $\sigma_i \in C_2(f_{i-1}, f_i)$ (a general element in $C_{x,y}^n(f_0, f_n)$ is a linear combination of such elements).

The two ‘‘extreme’’ factors $A_L(\phi)$ and $A_R(\phi)$ are defined as the compositions

$$A_L(\phi) = \sigma_n \circ^v \dots \circ^v \sigma_{\phi(m)+1}, \quad A_R(\phi) = \sigma_{\phi(0)} \circ^v \dots \circ^v \sigma_1$$

($A_L(\phi)$ is by definition equal to $\text{id}(f_n)$ if $\phi(m) = n$, and $A_R(\phi)$ is $\text{id}(f_0)$ if $\phi(0) = 0$).

The middle factor

$$M(\phi)(\sigma_{\phi(m)} \otimes \dots \otimes \sigma_{\phi(0)+1}) = \omega_m \otimes \dots \otimes \omega_1 \in C_{x,y}^m(f_{\phi(0)}, f_{\phi(m)})$$

is defined by

$$\omega_a = \omega_a(\phi) = \begin{cases} \sigma_{c-1} \circ^v \dots \circ^v \sigma_b: f_b \rightarrow f_c & \text{if } \phi(a-1) = b, \phi(a) = c, c > b, \\ \text{id}(f_b) & \text{if } \phi(a-1) = \phi(a) = b. \end{cases}$$

It completes the definition of $C_{x,y}(\phi)$.

Remark 5.7. The reader certainly realises that this construction just mimics the classical nerve construction in a non-cartesian monoidal (\mathbb{k} -linear) case. In the classical case of enrichment in Sets, one just projects along the two extreme factors $A_L(\phi)$ and $A_R(\phi)$. Our ‘‘bimodule’’ version is a way to phrase out the same construction in the situation when the projections (with respect to the monoidal product) do not exist.

5.6.2. Let C be as above, and let $x_0, \dots, x_k \in C_0$. Assume we are given maps in Δ $\phi_1: [m] \rightarrow [n_1], \dots, \phi_k: [m] \rightarrow [n_k]$. For $f_{i_0}, \dots, f_{i_{n_i}} \in C_1(x_{i-1}, x_i)$, $1 \leq i \leq k$, define

$$f_{\min} = f_{\otimes 0} = f_{k0} \circ f_{k-1,0} \circ \dots \circ f_{10}, \quad f_{\max} = f_{\otimes n} = f_{kn_k} \circ \dots \circ f_{1n_1}$$

and, for $0 \leq s \leq m$,

$$f_{\otimes \phi(s)} = f_{k\phi_k(s)} \circ \dots \circ f_{1\phi_1(s)}.$$

We define a map generalising the map defined above when we had $k = 1$:

$$\begin{aligned} C_{x_0, \dots, x_k}(\phi_1, \dots, \phi_k): C_{x_0, x_1}^{n_1}(f_{10}, f_{1n_1}) \otimes_{\mathbb{k}} \dots \otimes_k C_{x_{k-1}, x_k}^{n_k}(f_{k0}, f_{kn_k}) \\ \rightarrow C_2 \underset{A_L(\phi_1, \dots, \phi_k)}{(f_{\otimes \phi(m)}, f_{\otimes n})} \otimes_{\mathbb{k}} C_{x_0, x_k}^m \underset{B(\phi_1, \dots, \phi_k)}{(f_{\otimes \phi(0)}, f_{\otimes \phi(m)})} \otimes_{\mathbb{k}} C_2 \underset{A_R(\phi_1, \dots, \phi_k)}{(f_{\otimes 0}, f_{\otimes \phi(0)})}, \quad (5.8) \end{aligned}$$

where

$$\begin{aligned} A_L(\phi_1, \dots, \phi_k) &= A_L(\phi_k) \circ^h A_L(\phi_{k-1}) \circ^h \dots \circ^h A_L(\phi_1) \in C_2(f_{\otimes\phi(m)}, f_{\otimes}), \\ A_R(\phi_1, \dots, \phi_k) &= A_R(\phi_k) \circ^h \dots \circ^h A_R(\phi_1) \in C_2(f_{\otimes 0}, f_{\otimes\phi(0)}), \\ B(\phi_1, \dots, \phi_k) &= \Omega_1 \otimes_{\mathbb{k}} \dots \otimes_{\mathbb{k}} \Omega_m \in C_2(f_{\otimes\phi(0)}, f_{\otimes\phi(m)}), \end{aligned} \quad (5.9)$$

where

$$\Omega_i = \omega_i(\phi_k) \circ^h \omega_i(\phi_{k-1}) \circ^h \dots \circ^h \omega_i(\phi_1) \in C_2(f_{\otimes\phi(i-1)}, f_{\otimes\phi(i)}).$$

5.6.3. Let $T = ([k]; [m_1], \dots, [m_k])$, $S = ([\ell]; [n_1], \dots, [n_\ell])$, and let $\Phi = (\phi; \{\phi^{i,\ell}\}): T \rightarrow S$, a morphism in Θ_2 . Here, we construct a map of complexes

$$\Phi_*: A(C, D)(F, G)(\eta, \theta)_T \rightarrow A(C, D)(F, G)(\eta, \theta)_S.$$

Let $\Psi \in A(C, D)(F, G)(\eta, \theta)_T$, and let $x_0, \dots, x_\ell \in C_0$, $\{f_{ij} \in C_1(x_{i-1}, x_i)\}_{i=1 \dots \ell, j=0 \dots n_i}$, and

$$\{\sigma_{ij} \in C_2(f_{i,j-1}, f_{i,j})\}_{i=1 \dots \ell, j=1 \dots n_i}$$

a datum “of shape S ”.

We have to define

$$\Phi_*(\Psi)(\{\sigma_{ij}\}) \in D_2(\eta_{x_\ell} \circ F(f_{\ell 0}) \circ \dots \circ F(f_{10}), G(f_{\ell n_\ell}) \circ \dots \circ G(f_{1n_1}) \circ \theta_{x_0}).$$

Let $\min = \phi(0)$, $\max = \phi(k)$.

For each $0 \leq i \leq k-1$, one gets a sequence of maps $\{\phi^{i,j}: [m_i] \rightarrow [n_j]\}_{\phi(i-1)+1 \leq j \leq \phi(i)}$. Assume $\phi(i-1) < \phi(i)$, and then the construction of Section 5.6.2 ((5.8) and (5.9)) gives a map

$$\begin{aligned} C_{x_{\phi(i-1)}, x_{\phi(i-1)+1}}^{n_{\phi(i-1)+1}}(f_{\phi(i-1)+1,0}, f_{\phi(i-1)+1, n_{\phi(i-1)+1}}) \otimes_{\mathbb{k}} \dots \otimes_{\mathbb{k}} C_{x_{\phi(i)-1}, x_{\phi(i)}}^{n_{\phi(i)}}(f_{\phi(i),0}, f_{\phi(i), n_{\phi(i)}}) \\ \rightarrow C_2(f_{\otimes\phi_i(m_i)}, f_{\otimes i, \max}) \otimes_{\mathbb{k}} C_{x_{\phi(i-1)}, x_{\phi(i)}}^{m_i}(f_{\otimes\phi_i(0)}, f_{\otimes\phi_i(m_i)}) \otimes_{\mathbb{k}} C_2(f_{\otimes i, 0}, f_{\otimes\phi_i(0)}), \end{aligned}$$

$A_L^i \qquad \qquad \qquad B^i \qquad \qquad \qquad A_R^i$

where

$$f_{\otimes\phi_i(s)} = f_{\phi(i), \phi_i^{\phi(i)}(s)} \circ \dots \circ f_{\phi(i-1)+1, \phi_i^{\phi(i-1)+1}(s)},$$

where $0 \leq s \leq m_i$, and

$$\begin{aligned} f_{\otimes i, 0} &= f_{\phi(i), 0} \circ f_{\phi(i)-1, 0} \circ \dots \circ f_{\phi(i-1)+1, 0} \\ f_{\otimes i, \max} &= f_{\phi(i), n_{\phi(i)}} \circ f_{\phi(i)-1, n_{\phi(i)-1}} \circ \dots \circ f_{\phi(i-1)+1, n_{\phi(i-1)+1}}. \end{aligned}$$

For the case when $\phi(i-1) = \phi(i)$, we set

$$\begin{aligned} A_L^i &= \text{id} \in C_2(\text{id}_{x_{\phi(i)}}, \text{id}_{x_{\phi(i)}}), \\ A_R^i &= \text{id} \in C_2(\text{id}_{x_{\phi(i)}}, \text{id}_{x_{\phi(i)}}), \\ B^i &= \text{id}_{x_{\phi(i)}} \xrightarrow{\text{id}} \text{id}_{x_{\phi(i)}} \xrightarrow{\text{id}} \dots \xrightarrow{\text{id}} \text{id}_{x_{\phi(i)}}, \end{aligned}$$

where B^i is the chain with m_i arrows.

Define

$$G(A_L^\otimes) = G(A_L^k) \circ^h G(A_L^{k-1}) \circ^h \dots \circ^h G(A_L^1),$$

$$F(A_R^\otimes) = F(A_R^k) \circ^h F(A_R^{k-1}) \circ^h \dots \circ^h F(A_R^1).$$

For a string

$$f_{j0} \xrightarrow{\sigma_{j1}} f_{j1} \xrightarrow{\sigma_{j2}} f_{j2} \dots \xrightarrow{\sigma_{jn_j}} f_{jn_j}$$

($f_{js} \in C_1(x_{j-1}, x_j)$ for $0 \leq s \leq n_j$), denote

$$\sigma_{j,\text{tot}} = \sigma_{jn_j} \circ^v \dots \circ^v \sigma_{j0}.$$

Let $\min = \phi(0)$, $\max = \phi(k)$, and denote

$$\sigma_{\otimes \min} = \sigma_{\min,\text{tot}} \circ^h \sigma_{\min-1,\text{tot}} \circ^h \dots \circ^h \sigma_{1,\text{tot}} \circ^h \sigma_{0,\text{tot}}$$

$$\sigma_{\otimes (\max+1)} = \sigma_{\ell,\text{tot}} \circ^h \sigma_{\ell-1,\text{tot}} \circ^h \dots \circ^h \sigma_{\max+1,\text{tot}}.$$

Finally, we have

$$\Phi_*(\Psi) = G(\sigma_{\otimes (\max+1)}) \circ^h ((G(A_L^\otimes)) \circ^v \Psi(B^1 \otimes_{\mathbb{k}} \dots \otimes_{\mathbb{k}} B^k) \circ^v (F(A_R^\otimes))) \circ^h F(\sigma_{\otimes \min}). \tag{5.10}$$

Formula (5.10) is an explicit expression for the definition of $A(C, D)(F, G)(\eta, \theta)$, in particular when C, D, F, G, η, θ are strict. The general case differs by numerous insertions of structure 2-isomorphisms, which makes them more complicated and tedious, but can be written down in a similar way.

Note that it follows from the discussion in Sections 4.4, 5.2, and 5.3 that the prescription (5.10) gives rise to a functor $\Theta_2 \rightarrow C^*(\mathbb{k})$.

5.7. An example: a shuffle permutation

Consider in more detail the action of an inner face map $D_{j,\sigma}$ (F2) (see Section 2.6) on $A(C, D)(F, G)(\eta, \theta)$, corresponding to an (m_j, m_{j+1}) -shuffle permutation σ .

Let $t \in \Sigma_{\ell_j}$ be an (m_j, m_{j+1}) -shuffle, $\ell_j = m_j + m_{j+1}$. Let $p^*: [\ell_j] \rightarrow [m_j]$ and $q^*: [\ell_j] \rightarrow [m_{j+1}]$ be the two maps Joyal dual to the natural embeddings $[m_j - 1] \rightarrow [\ell_j - 1]$ and $[m_{j+1} - 1] \rightarrow [\ell_j - 1]$ (see Section 2.6, (F2)). Let

$$T = ([n]; [\ell_1], \dots, [\ell_n]), \quad S = ([n + 1]; [\ell_1], \dots, [\ell_{j-1}], [m_j], [m_{j+1}], [\ell_{j+1}], \dots, [\ell_{n+1}]).$$

Consider the morphism $\Phi = D_{j,t}: T \rightarrow S$ in Θ_2 , corresponding to the shuffle t .

Then, the morphism $D_{j,t}$ acts on $A(C, D)(F, G)(\eta, \theta)$ as follows.

Use the notation \underline{f}_S for a chain of 2-morphisms $\{\sigma_{si}: f_{s,i-1} \rightarrow f_{si}\}$ in C :

$$f_{s0} \xrightarrow{\sigma_{s1}} f_{s2} \xrightarrow{\sigma_{s2}} \dots \xrightarrow{\sigma_{s,m_s}} f_{s,m_s}.$$

For a cochain $\Psi \in A(C, D)(F, G)(\eta, \theta)_T$, one has

$$(D_{j,t_*}(\Psi))(f_{\underline{1}}, \dots, f_{\underline{n+1}}) = \Psi_T(f_{\underline{1}}, \dots, f_{\underline{j-1}}, \underline{g}_j, f_{\underline{j+2}}, \dots, f_{\underline{n+1}}),$$

where \underline{g}_j is the chain

$$f_{j+1,0} \circ f_{j0} \xrightarrow{\omega_1} \dots \xrightarrow{\omega_{m_j+m_{j+1}}} f_{j+1,m_{j+1}} \circ f_{j,m_j}$$

and

$$\omega_i = \begin{cases} \text{id} \circ^h \sigma_{ja} & \text{if } t^{-1}(i) = a, 0 \leq a \leq m_j \\ \sigma_{j+1,b} \circ^h \text{id} & \text{if } t^{-1}(i) = b, m_j + 1 \leq b \leq m_j + m_{j+1}. \end{cases}$$

5.8. Normalised vs non-normalised chain complexes of a 2-cellular object in $C^*(\mathbb{k})$

In the proof of Theorem 7.3, we use that the Θ_2 -cochain complex of $A(C, D)(F, G)(\eta, \theta)$ is quasi-isomorphic to its *normalised* subcomplex

$$A(C, D)(F, G)(\eta, \theta)_{\text{norm}}(C, D)(F, G)(\eta, \theta).$$

The latter is, by definition, the sub-complex which consists of all cochains Ψ which are equal to 0 if some of its 2-morphism arguments $\sigma_{i,j}$ are the identity morphism of a 1-morphism.

In Proposition 5.9, we prove that the complexes

$$A(C, D)(F, G)(\eta, \theta) \quad \text{and} \quad A_{\text{norm}}(C, D)(F, G)(\eta, \theta)$$

are quasi-isomorphic. Theorem 7.3 is a statement about cohomology. Therefore, due to Proposition 5.9, one can assume in its proof that we work with the normalised complex.

Recall that for a simplicial object in an abelian category \mathcal{A} , its normalised Moore complex $N(X)$ is defined as the quotient-complex of the ordinary Moore complex $C(X)$ by the subcomplex $DC(X)$ spanned by elements of the form $s_i y$ (here, s_i stands for the simplicial version of the degeneracy morphisms $\varepsilon_i \in \Delta$; see Section 2.1).

Recall the following classical result, in a slightly more general version.

Proposition 5.8. *Let $X: \Delta^{\text{op}} \rightarrow C^*(\mathbb{k})$ be a simplicial object in $C^*(\mathbb{k})$. Then, the total sum complex $\text{Tot}^{\oplus}(C(X))$ of the Moore complex of X is quasi-isomorphic to the total sum complex $\text{Tot}^{\oplus}(N(X))$ of the normalised Moore complex.*

Proof. The proof given in [26, Sect. VIII. 6] can be easily adopted to this case. Indeed, MacLane constructs a map $g: C(Y)/DC(Y) \rightarrow C(Y)$, for Y a simplicial object in an abelian category, such that g is a “quasi-inverse” to the natural projection $\pi: C(Y) \rightarrow C(Y)/DC(Y)$ in the sense that $\pi \circ g = \text{id}$, and $g \circ \pi$ is chain homotopic to the identity. The chain homotopy constructed in loc. cit. clearly commutes with “inner” differentials on X_i s. Consequently, if one defines $\pi': \text{Tot}^{\oplus}(C(X)) \rightarrow \text{Tot}^{\oplus}(N(X))$ and $g': \text{Tot}^{\oplus}(N(X)) \rightarrow \text{Tot}^{\oplus}(C(X))$, one still has $\pi'g' = \text{id}$ and $g'\pi'$ chain homotopic to the identity. ■

The next step is to generalise Proposition 5.8 to the case of 2-cellular objects in $C^\bullet(\mathbb{k})$, that is, to the case of functors $X: \Theta_2^{\text{op}} \rightarrow C^\bullet(\mathbb{k})$.

For $Y: \Theta_2^{\text{op}} \rightarrow \text{Vect}(\mathbb{k})$, its chain complex is defined as the complex $C(Y)$, with

$$C_{-\ell}(Y) = \bigoplus_{T, \dim T = \ell} Y_T$$

with the differential dual to (3.2), and its normalised complex is defined as the quotient-complex of $C(Y)$ by the subcomplex $DC(Y)$ generated by the elements $\varepsilon_p^j(y)$ of type (D1) (see Section 2.6), $y \in Y_D$:

$$N(Y) = C(Y)/DC(Y).$$

That is, we use only “vertical” degeneracy morphisms of type (D1), *not* “horizontal” degeneracy morphisms of type (D2), in the definition of $DC(Y)$.

For the case of a functor $X: \Theta_2^{\text{op}} \rightarrow C^\bullet(\mathbb{k})$ as above, $C(X), DC(X), N(X)$ are defined as $\text{Tot}^\oplus(C(X)), \text{Tot}^\oplus(DC(X)), \text{Tot}^\oplus(N(X))$, correspondingly.

Proposition 5.9. *Let $X: \Theta_2^{\text{op}} \rightarrow C^\bullet(\mathbb{k})$ be a 2-cellular complex. Then, the natural projection $\pi: \text{Tot}^\oplus(C(X)) \rightarrow \text{Tot}^\oplus(N(X))$ is a quasi-isomorphism of complexes.*

Proof. One can not follow directly the same line as in the proof of [26, Chap. VIII, Thm. 6.1], for the following reason. The subspaces $D_i C(X)$, $i \geq 0$ (or rather their direct analogues) are *not* subcomplexes of $C(X)$ because the components $D_{j,\sigma}$ of type (F2) (see Section 2.6) in the differential (3.2) may *increase* i . Indeed, these components act as “deshuffling” of two neighbour columns, resulting in a column of a greater length, so this operation may send $\varepsilon_p^i y$ to $\varepsilon_q^{i'}(y')$ with $i' > i$ (here, $q = p$ or $p - 1$).

To overcome this obstacle, we employ the following spectral sequence argument.

Denote by $F_N \subset C(X)$ the subspace spanned by X_T , $T = ([n]; [\ell_1], \dots, [\ell_n])$ with $n \leq N$. Then, F_N is a subcomplex: the boundary operators of type (F1) and (F3) preserve n , and the boundary operators of types (F2) and (F4) decrease n by 1; see Section 2.6.

We get an exhausting ascending filtration of $C(X)$ by subcomplexes:

$$F_0 \subset F_1 \subset F_2 \subset \dots$$

A similar filtration exists for $N(X)$ as well; denote the corresponding subspaces by F'_N . The natural projection $\pi: C(X) \rightarrow N(X)$ sends F_N to F'_N ; hence, π induces a map of the corresponding spectral sequences. Denote these spectral sequences by $\{E_n^{pq}\}$ and $\{E'_n{}^{pq}\}$, so that π induces a map $\pi_*: (E_n^{pq}, d_n) \rightarrow (E'_n{}^{pq}, d'_n)$.

The spectral sequences at the term E_0 (resp., E'_0) are non-zero at the lower half plane $y \leq 0$, and the differential d_0 is horizontal. So the spectral sequences converge by dimensional reasons.

Lemma 5.10. *The map $\pi_*: (E_0^{*\ell}, d_0) \rightarrow (E'_0{}^{*\ell}, d'_0)$ is a quasi-isomorphism, for any $\ell \leq 0$. In particular, π_* defines an isomorphism $\pi_*: E_1^{pq} \rightarrow E'_1{}^{pq}$, for all p, q .*

Proof. For any fixed ℓ , the complex $(E_0^{*\ell}, d_0)$ is $C^{(\ell)}(X)$, whose degree $-n$ component is equal to the direct sum $\bigoplus_T X_T$ over $T = ([\ell]; [n_1], \dots, [n_\ell])$ with $\dim T = n$, and with

the differential components given only by (F1) and (F3) types; see Section 2.6. That is, the contribution of types (F2) and (F4) components in (3.2) becomes 0 in the associated graded complex $C^{(\ell)}(X) = F_\ell/F_{\ell-1}$. The complex $(E_0^{*,\ell}, d'_0)$ has a similar description.

It makes it possible for us to employ the construction of the proof of [26, Chap. VIII, Thm. 6.1]. Namely, we define *subcomplexes* $D_i C^{(\ell)}(X)$, for any $i \geq 0$, such that

$$D_{i+1}C^{(\ell)}(X) \supset D_i C^{(\ell)}(X) \quad \text{and} \quad DC^{(\ell)}(X) = \bigcup_{i \geq 0} D_i C^{(\ell)}(X).$$

As in loc. cit., we construct a map $h_i: C^{(\ell)}(X) \rightarrow C^{(\ell)}(X)$ chain homotopic to id and mapping D_i to D_{i-1} . The composition of these maps is well defined, is chain homotopic to id, and sends $DC^{(\ell)}(X)$ to 0. It gives a map $g: N^{(\ell)}(X) \rightarrow C^{(\ell)}(X)$ such that $\pi_*g = \text{id}$ and $g\pi_*$ is chain homotopic to id, which completes the proof. ■

It follows from this lemma that π_* defines an isomorphism at E_∞ sheet; hence, π is a quasi-isomorphism. ■

We can prove the following theorem.

Theorem 5.11. *The natural embedding*

$$i: A_{\text{norm}}(C, D)(F, G)(\eta, \theta) \rightarrow A(C, D)(F, G)(\eta, \theta)$$

is a quasi-isomorphism of complexes.

Proof. One can apply the arguments “dual” to the ones provided above, for the case of a 2-cocellular complex. Define $\Phi^p \subset A(C, D)(F, G)(\eta, \theta) = A$ as the subcomplex formed by cochains which vanish on F_p defined above. Then, $\{\Phi^p\}$ form a descending filtration of A . This filtration is complete in the sense that

$$A = \lim_{\leftarrow} A / \Phi^p A.$$

The term E_0 lives in $x \geq 0$ half of the plane; thus, it is bounded below. It follows from the complete convergence theorem [38, Thm. 5.5.10] (in its version when d has degree +1) that the sequence converges. A similar convergent spectral sequence exists on A_{norm} , given by the filtration $\Phi^{p'} A_{\text{norm}} = (\Phi^p A) \cap A_{\text{norm}}$. The argument given in the 2-cellular case gives a quasi-isomorphism of complexes $\Phi^{p'} A_{\text{norm}} / \Phi^{(p+1)'} A_{\text{norm}} \rightarrow \Phi^p A / \Phi^{p+1} A$ induced by i . Then, the result follows from the complete comparison theorem [38, Thm. 5.5.11]. ■

6. The p -relative totalisation $Rp_*(A(C, D)(F, G)(\eta, \theta))$ and higher structures via Davydov–Batatin

We know from Propositions 3.5 and 3.7 that the relative totalisation of the cosimplicial vector space $Rp_*(A(C, D)(F, G)(\eta, \theta))$ is a cosimplicial vector space, and its Δ -Moore complex is equal to the (absolute) Θ_2 -totalisation such that its non-normalised Moore complex is isomorphic to the (absolute) Θ_2 -totalisation of $A(C, D)(F, G)(\eta, \theta)$.

In this section, we apply some results of [6] for studying the higher structures on the complexes $C^\bullet(C, D)(F, G)(\eta, \theta)$. We show that $(Rp_*)(X)$ enjoys, for the case

$$X = A(C, D)(F, F)(\text{id}, \text{id}),$$

the property of being a *1-commutative* cosimplicial monoid, in the sense of [6]. Consequently, $C^\bullet(C, D)(F, F)(\text{id}, \text{id})$ is a homotopy 2-algebra, for any \mathbb{k} -linear strict 2-functor $F: C \rightarrow D$.

At the same time, for the case

$$X = A(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id}),$$

the cosimplicial monoid $(Rp_*)(X)$ is *not* 2-commutative (unlike the case of the Davydov–Yetter complex). At the moment, we do not know the correct homotopy refinement of 2-commutativity, which would imply that $C^\bullet(C, D)(\text{Id}, \text{Id})(\text{id}, \text{id})$ is a homotopy 3-algebra.

6.1. The totalisation $\text{Tot}_{\Theta_2} A(C, D)(F, F)(\text{id}, \text{id})$ is a homotopy 2-algebra

Recall a *cosimplicial monoid* X (in a symmetric monoidal category \mathcal{C}) is a cosimplicial object in the category of monoids $\text{Mon}(\mathcal{C})$. The question raised in [6] is the following:

Which condition on X implies that the totalisation $\text{Tot}(X)$ admits an action of an operad (homotopy equivalent to) E_n ?

It follows immediately that the condition that X is a cosimplicial monoid implies that X^\bullet is a monoid with respect to the Batanin \square -product [4]. Thus, it follows from loc. cit. that for a cosimplicial monoid X^\bullet , the totalisation $\text{Tot}(X)$ is an A_∞ monoid, that is, a E_1 -algebra.

In [6, Sect. 2.2], the following definition is given.

Definition 6.1. Let $\tau: [p] \rightarrow [m]$ and $\pi: [q] \rightarrow [m]$ be two maps in Δ . A *shuffling of length* n of τ, π is a decomposition of the images of τ and π into disjoint union of *connected intervals*

$$\begin{aligned} \text{Im}(\tau) &= A_1 \cup A_2 \cup \dots \cup A_s, & A_1 < A_2 < \dots < A_s \\ \text{Im}(\pi) &= B_1 \cup B_2 \cup \dots \cup B_t, & B_1 < B_2 < \dots < B_t \\ s + t &= n + 1 \end{aligned}$$

which satisfy either

$$A_1 \leq B_1 \leq A_2 \leq B_2 \leq \dots$$

or

$$B_1 \leq A_1 \leq B_2 \leq A_2 \leq \dots$$

(that is, the rightmost end-point of A_i may coincide with the leftmost end-point of the sequel B).

The linking number $\mathbf{lk}(\tau, \pi)$ is defined as the minimal possible shuffling of τ, π has length n .

See [6, Sect. 2.2], for examples.

Definition 6.2. Let X be a cosimplicial monoid, $n \geq 0$. X is called n -commutative if for any $\tau: [p] \rightarrow [m]$, $\pi: [q] \rightarrow [m]$ in Δ with $\mathbf{lk}(\tau, \pi) \leq n$, the diagram below commutes:

$$\begin{array}{ccc} X(p) \otimes X(q) & \xrightarrow{X(\tau) \otimes X(\pi)} & X(m) \otimes X(m) \\ \downarrow & & \downarrow \mu \\ X(q) \otimes X(p) & \xrightarrow{X(\pi) \otimes X(\tau)} & X(m) \otimes X(m) \xrightarrow{\mu} X(m). \end{array}$$

The following result is proven in [6, Thm. 2.45, Cor. 2.46].

Theorem 6.3. Let X be an n -commutative cosimplicial monoid in $C(\mathbb{k})$. Then, there is an action of the operad homotopy equivalent to $C.(E_{n+1}, \mathbb{k})$ on the totalisation $\text{Tot}(X) \in C(\mathbb{k})$.

In [6], some explicit formulas for the degree $-n$ Lie bracket are provided; see [6, Sects. 2.9 and 2.10].

We easily prove the following proposition.

Proposition 6.4. Let C, D be \mathbb{k} -linear bicategories, $F: C \rightarrow D$ a strong bicategorical functor. Then, the cosimplicial vector space $Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))$ is a 1-commutative cosimplicial monoid.

Proof. Let $\tau_{m,n}: [n] \rightarrow [m+n]$ and $\pi_{m,n}: [m] \rightarrow [m+n]$ be defined as $\tau_{m,n}(i) = i$ and $\pi_{m,n}(j) = n + j$. It is clear that $\mathbf{lk}(\tau_{m,n}, \pi_{m,n}) = 1$. Moreover, the general case of the linking number 1 is reduced to this particular case, due to the following simple observation [6, Lem. 2.1].

Let $\tau: [p] \rightarrow [m]$, $\pi: [q] \rightarrow [m]$ be morphisms in Δ and

$$[p] \rightarrow [p'] \xrightarrow{\tau'} [m], \quad [q] \rightarrow [q'] \xrightarrow{\pi'} [m]$$

their epi-mono factorisations. Then, $\mathbf{lk}(\tau, \pi) = \mathbf{lk}(\tau', \pi')$.

We check the 1-commutativity of $Rp_*(A(F, F))$. We firstly assume that C is strict and F is strict. Let $\Phi \in Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))^n$, $\Psi \in Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))^m$ be represented by cochains $\Phi \in A(C, D)(F, F)(\text{id}, \text{id})_D$, $\Psi \in A(C, D)(F, F)(\text{id}, \text{id})_{D'}$, with $p(D) = [n]$, $p(D') = [m]$. Assume $D = ([n]; [k_1], \dots, [k_n])$ and $D' = ([m]; [\ell_1], \dots, [\ell_m])$. Then, $\tau_{m,n}(\Phi)$ takes a non-zero value on the object

$$\widehat{D} = ([n+m]; [k_1], \dots, [k_n], [0], \dots, [0])$$

and is equal to

$$\tau_{m,n}(\Phi)(-, X_{n+1}, \dots, X_{m+n}) = \Phi(-) \circ^h (\text{id}_{F(X_{n+1} \circ \dots \circ X_{m+n})}).$$

Analogously, $\pi_{m,n}(\Psi)$ takes a non-zero value on $\widehat{D}' = ([m+n]; [0], \dots, [0], [\ell_1], \dots, [\ell_m])$, and

$$\pi_{m,n}(\Psi)(Y_1, \dots, Y_m, -) = \text{id}_{F(Y_1 \circ \dots \circ Y_m)} \circ^h \Psi(-).$$

Finally, for their product in the monoid $Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))^{m+n}$, one has

$$\begin{aligned} \tau_{m,n}(\Phi) * \pi_{m,n}(\Psi)(T_1, \dots, T_{m+n}) &= (\Phi(T_1, \dots, T_n) \circ^h \text{id}_{F(X_{n+1} \circ \dots \circ X_{m+n})}) \circ^v (\text{id}_{F(Y_1 \circ \dots \circ Y_n)} \circ^h \Psi(T_{n+1}, \dots, T_{m+n})) \\ &= \Phi(T_1, \dots, T_n) \circ^h \Psi(T_{n+1}, \dots, T_{m+n}), \end{aligned} \tag{6.1}$$

where T_i is a string of morphisms of length k_i for $1 \leq i \leq n$ and ℓ_{j-n} for $j = n + 1, \dots, n + m$, starting at X_i and ending at Y_i .

We clearly get the same expression when computing

$$\pi_{m,n}(\Psi) * \tau_{m,n}(\Phi)(T_1, \dots, T_{m+n}),$$

and the 1-commutativity for $Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))$ follows. It completes the proof for F strict.

Now, when C is a bicategory and F is a strong bicategorical functor, we argue as follows.

There are two sorts of the structure isomorphisms which figure in (the strong counterpart of) (6.1). These two sorts are (a) the structure constraints of the bicategory and (b) the structure constraints of the functor F . It follows from Definition 4.2 that the structure constraints of type (a) commute with the structure constraints of type (b); on the other hand, the elements $Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))$ commute with the constraints of type (a), in the sense of (5.4)–(5.6). These two properties imply that the presence of these constraints does not affect the previous speculation in the strict case. ■

Remark 6.5. The fulfilment of the Batanin–Davydov 1-commutativity condition [6] for $Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))$ is a lucky situation, which is easily generalised from the corresponding proof for the classical Davydov–Yetter complex in [6, Thm. 3.4]. Namely, (although our cochains are not natural transformations) one does not use the naturality of cochains for general morphisms in this proof. One does use the naturality with respect to the identity morphisms, which automatically holds.

The case of the 2-commutativity of $Rp_*(\text{Id}, \text{Id})$ is not that lucky because the corresponding proof for the classical counterpart given in [6, Thm. 3.8] essentially uses the naturality for non-identity morphisms. Our cochains are not natural transformations, which results in the failure of 2-commutativity for $Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))$. However, a sort of “homotopy 2-commutativity” still holds.

Theorem 6.6. *Let C, D be \mathbb{k} -linear bicategories, $F: C \rightarrow D$ a strong bicategorical functor. Then, the 2-cocellular totalisation $\text{Tot}_{\Theta_2}(A(C, D)(F, F)(\text{id}, \text{id}))$ has a structure of an algebra over an operad homotopically equivalent to $C_*(E_2; \mathbb{k})$.*

Proof. By Proposition 3.7,

$$\text{Tot}_{\Theta_2}(A(C, D)(F, F)(\text{id}, \text{id})) \sim \text{Tot}_{\Delta}(Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))).$$

By Proposition 6.4, $Rp_*(A(C, D)(F, F)(\text{id}, \text{id}))$ is a 1-commutative cosimplicial monoid. Then, the result follows from Theorem 6.3. ■

7. The totalisations $\text{Tot}_{\Theta_2}(A(C, D)(F, F)(\text{id}, \text{id}))$ and $\text{Tot}_{\Theta_2}(A(C, C)(\text{Id}, \text{Id})(\text{id}, \text{id}))$ as deformation complexes

7.1. Infinitesimal deformation theory of a monoidal dg category C

We start with the deformations of a monoidal category C .

Let C be a \mathbb{k} -linear monoidal category (or a monoidal dg category over \mathbb{k}). The deformations we consider are *formal* deformations; that is, C_t may not make sense unless $t = 0$. That is, the category C_t is a monoidal category over the formal power series $\mathbb{k}[[t]]$.

We consider *flat* deformations C_t of C in the following sense: the set of objects, the vector spaces (complexes) $C_t(x, y)$ of morphisms, and the monoidal product *on objects* remain undeformed.

Then, the data which is being deformed is as follows:

- (A1) the composition of morphisms $m_{X,Y,Z}: C(Y, Z) \otimes C(X, Y) \rightarrow C(X, Z)$,
 $X, Y, Z \in C$,
- (A2) for $f \in C(X, X')$, $g \in C(Y, Y')$, the monoidal products of *morphisms* $m_{X,g} = \text{id}_X \otimes g: C(X, Y) \rightarrow C(X, Y')$ and $m_{f,Y} = f \otimes \text{id}_Y: C(X, Y) \rightarrow C(X', Y)$,
- (A3) the associator $\alpha_{X,Y,Z}: X \otimes (Y \otimes Z) \rightarrow (X \otimes Y) \otimes Z$, $X, Y, Z \in C$,
- (A4) the left and right unit maps $\lambda_X: e \otimes X \rightarrow X$ and $\rho_X: X \otimes e \rightarrow X$.

It is assumed that (a) the identity morphism id_X , $X \in C$, (b) the monoidal unit e , (c) the maps λ_Y , ρ_X , and (d) $m_{f,e}$ and $m_{e,g}$ are stable under the deformations, and (e) $m_{X,\text{id}_Y} = m_{\text{id}_X,Y} = \text{id}_{X \otimes Y}$.

The following example shows that this set-up is realistic.

Example 7.1. Let A be a bialgebra over \mathbb{k} , $C = \text{Mod}(A)$ the category of left A -modules over the underlying algebra. It is a monoidal category in a standard way: for two modules M, N , the tensor product of the underlying vector spaces $M \otimes_{\mathbb{k}} N$ is naturally an $A \otimes A$ -module, and the precomposition with $\Delta: A \rightarrow A \otimes A$ makes it an A -module.

Assume that A is a Hopf algebra. Then, the monoidal product $A \otimes A$ of two *free* modules of rank 1 is a free module again, whose underlying vector space is *canonically* isomorphic to $A \otimes A_u$, where A_u is the underlying vector space of A .

Indeed, define the maps $\alpha: A \otimes A \rightarrow A \otimes A_u$ and $\beta: A \otimes A_u \rightarrow A \otimes A$ as

$$\begin{aligned} \alpha(a \otimes b) &= \sum a_1 \otimes S(a_2)b, \\ \beta(a \otimes b) &= \sum a_1 \otimes a_2b, \end{aligned}$$

where $S: A \rightarrow A$ is the antipode, and we use the Swindler notations $\Delta(a) = \sum a_1 \otimes a_2$.

Assume that α and β are maps of A -modules, and $\alpha \circ \beta = \text{id}$, $\beta \circ \alpha = \text{id}$. It proves the claim.

Thus, if we consider a deformation A_t of a Hopf algebra A , the \mathbb{k} -linear subcategory $C_{\text{free}}(A_t)$ is a deformation of a monoidal \mathbb{k} -linear category $C_{\text{free}}(A)$, for which the conditions (A1)–(A3) are fulfilled.

The data listed in (A1)–(A4) is subject to the following axioms:

- (R1) the composition $m_{X,Y,Z}$ in (A1) is associative,
- (R2) for maps in (A2), one has $(f \otimes \text{id}_y) \circ (\text{id}_x \otimes g) = (\text{id}_x \otimes g) \circ (f \otimes \text{id}_y)$ (both sides are equal to $f \otimes g: X \otimes Y \rightarrow X' \otimes Y'$; therefore, the deformation of $f \otimes g$ is determined by deformations of $f \otimes \text{id}_y$ and $\text{id}_x \otimes g$),
- (R3) for any two composable morphisms $X \xrightarrow{f} X' \xrightarrow{f'} X''$, and any $Y \in C$, one has $m_{f',Y} \circ m_{f,Y} = m_{f' \circ f, Y}$; similarly, for any two composable morphisms

$$Y \xrightarrow{g} Y' \xrightarrow{g'} Y'',$$

and any $X \in C$, one has $m_{X,g'} \circ m_{X,g} = m_{X,g' \circ g}$,

- (R4) this and the next two axioms express naturality of the associator. Let $f: X \rightarrow X'$ be a morphism in C , and Y, Z objects. The following diagram commutes:

$$\begin{array}{ccc} X \otimes (Y \otimes Z) & \xrightarrow{\alpha_{X,Y,Z}} & (X \otimes Y) \otimes Z \\ m_{f,Y \otimes Z} \downarrow & & \downarrow m_{m_{f,Y},Z} \\ X' \otimes (Y \otimes Z) & \xrightarrow{\alpha_{X',Y,Z}} & (X' \otimes Y) \otimes Z \end{array}$$

- (R5) let $g: Y \rightarrow Y'$ be a morphism in C , X, Z objects. The following diagram commutes:

$$\begin{array}{ccc} X \otimes (Y \otimes Z) & \xrightarrow{\alpha_{X,Y,Z}} & (X \otimes Y) \otimes Z \\ m_{X,m_{g,Z}} \downarrow & & \downarrow m_{m_{X,g},Z} \\ X \otimes (Y' \otimes Z) & \xrightarrow{\alpha_{X,Y',Z}} & (X \otimes Y') \otimes Z \end{array}$$

- (R6) let $h: Z \rightarrow Z'$ be a morphism in C , X, Y objects. Then, the following diagram commutes:

$$\begin{array}{ccc} X \otimes (Y \otimes Z) & \xrightarrow{\alpha_{X,Y,Z}} & (X \otimes Y) \otimes Z \\ m_{X,m_{Y,h}} \downarrow & & \downarrow m_{X \otimes Y,h} \\ X \otimes (Y \otimes Z') & \xrightarrow{\alpha_{X,Y,Z'}} & (X \otimes Y) \otimes Z' \end{array}$$

- (R7) the pentagon equation for the associator is

$$\begin{array}{ccc} & (X \otimes Y) \otimes (Z \otimes T) & \\ & \nearrow \alpha_{X,Y,Z \otimes T} & \searrow \alpha_{X \otimes Y,Z,T} \\ X \otimes (Y \otimes (Z \otimes T)) & & ((X \otimes Y) \otimes Z) \otimes T \\ & \searrow m_{X,\alpha_{Y,Z,T}} & \nearrow m_{\alpha_{X,Y,Z},T} \\ X \otimes ((Y \otimes Z) \otimes T) & \xrightarrow{\alpha_{X,Y \otimes Z,T}} & (X \otimes (Y \otimes Z)) \otimes T \end{array} \tag{7.1}$$

(R8) left unit functionality: for any map $f: X \rightarrow X'$, the diagram

$$\begin{array}{ccc} X \otimes e & \xrightarrow{\rho_X} & X \\ m_{f,e} \downarrow & & \downarrow f \\ X' \otimes e & \xrightarrow{\rho_{X'}} & X' \end{array}$$

commutes,

(R9) right unit functionality: for any $g: Y \rightarrow Y'$, the diagram

$$\begin{array}{ccc} e \otimes Y & \xrightarrow{\lambda_Y} & Y \\ m_{e,g} \downarrow & & \downarrow g \\ e \otimes Y' & \xrightarrow{\lambda_{Y'}} & Y' \end{array}$$

commutes,

(R10) left right unit compatibility: the two possible maps $\lambda_e, \rho_e: e \otimes e \rightarrow e$ are equal.

Among the deformations C_t , there are ones which we consider “trivial”. This appears in the literature under the name “twist”; however, here we consider “upgraded” twists acting not only on the associator, but also on the underlying category structure and on the action of morphisms on the monoidal product.

In the deformation theory, one identifies two deformations if one is obtained from another by a twist and interests in the “quotient-space”.

Lemma 7.2. *Let C be a \mathbb{k} -linear (or dg over \mathbb{k}) monoidal category, and denote by C_u the underlying \mathbb{k} -linear quiver of C . Assume that, for any $X, Y \in C$, we are given an isomorphism $\varphi_{X,Y}: C(X, Y) \rightarrow C(X, Y)$, and an isomorphism $\psi_{X,Y} \in C(X \otimes Y, X \otimes Y)$. Then, these data give rise to a monoidal equivalence functor F from C to another monoidal \mathbb{k} -linear (resp., dg over \mathbb{k}) category \tilde{C} on the quiver C_u , such that F is the identity map on any object of C .*

Proof. It is standard. We define F on morphisms by $F(f) = \varphi_{X,Y}(f)$ if $f \in C(X, Y)$, and define monoidal constraints $F(X \otimes Y) \rightarrow F(X) \otimes F(Y)$ as the isomorphisms $\psi_{X,Y}$. Then, the monoidal category structure on \tilde{C} is uniquely determined by the requirement that F is a monoidal functor.

We assume that $\varphi_{X,X}(\text{id}_X) = \text{id}_X$, $\psi_{e,Y} = \text{id}_{e \otimes Y}$, $\psi_{X,e} = \text{id}_{X \otimes e}$. As well, we assume that the constraint $F(e) \rightarrow e$ is the identity map.

For the convenience of the reader, we provide explicit formulas for the new composition of morphisms, for the action of morphisms on the tensor product, and for the associator. We use the same notations decorated by \sim for the corresponding data (A1)–(A3) of the new category on \tilde{C} . We use the same notations as in (A1)–(A3). One has

$$\tilde{m}_{X,Y,Z}(f, g) = \varphi_{X,Z}(m_{X,Y,Z}(\varphi_{Y,Z}^{-1}(g), \varphi_{X,Y}^{-1}(f))) \tag{7.2}$$

$$\begin{aligned} \tilde{m}_{X,g} \tilde{\circ} \psi_{X,Y} &= \psi_{X,Y'} \tilde{\circ} \varphi_{X \otimes Y, X \otimes Y'}(m_{X, \varphi_{Y',g}^{-1}}) \\ \tilde{m}_{f,Y} \tilde{\circ} \psi_{X,Y} &= \psi_{X',Y} \tilde{\circ} \varphi_{X \otimes Y, X' \otimes Y}(m_{\varphi_{X',f,Y}^{-1}}), \end{aligned} \tag{7.3}$$

where $\tilde{\circ}$ denotes the composition in \tilde{C} (given by (7.2)).

The two last equations follow from the commutative diagram:

$$\begin{array}{ccc} F(X \otimes Y) & \longrightarrow & F(X) \otimes F(Y) \\ F(f \otimes g) \downarrow & & \downarrow F(f) \otimes F(g) \\ F(X' \otimes Y') & \longrightarrow & F(X') \otimes F(Y'), \end{array}$$

where $\tilde{\circ}$ is the composition in \tilde{C} , and $F(?) = ?$ for any object $? \in C$:

$$\tilde{\alpha}_{X,Y,Z} = \tilde{m}_{\psi_{X,Y}, Z} \tilde{\circ} \psi_{X \otimes Y, Z} \tilde{\circ} \varphi(\alpha_{X,Y,Z}) \tilde{\circ} \psi_{X',Y \otimes Z}^{-1} \tilde{\circ} \tilde{m}_{X, \psi_{Y,Z}^{-1}}. \tag{7.4}$$

It comes from the commutative diagram:

$$\begin{array}{ccc} F(X \otimes (Y \otimes Z)) & \xrightarrow{F(\alpha)} & F((X \otimes Y) \otimes Z) \\ \psi_{X,Y \otimes Z} \downarrow & & \downarrow \psi_{X \otimes Y, Z} \\ F(X) \otimes F(Y \otimes Z) & & F(X \otimes Y) \otimes F(Z) \\ \text{id}_X \otimes \psi_{Y,Z} \downarrow & & \downarrow \psi_{X,Y} \otimes \text{id}_Z \\ F(X) \otimes (F(Y) \otimes F(Z)) & \xrightarrow{\tilde{\alpha}} & (F(X) \otimes F(Y)) \otimes F(Z), \end{array}$$

where $\tilde{\circ}$ is the composition in \tilde{C} , and $F(?) = ?$ for any object $? \in C$.

One can check directly that $\tilde{m}_{X,Y,Z}, \tilde{m}_{X,g}, \tilde{m}_{f,Y}, \tilde{\alpha}_{X,Y,Z}$ satisfy (R1)–(R7) and thus define a monoidal category \tilde{C} , such that the functor

$$F: C \rightarrow \tilde{C}$$

is a monoidal equivalence. ■

Note that in the assumption of the lemma, $\varphi_{X,Y}$ and $\psi_{X,Y}$ are arbitrary isomorphisms. Now we switch back to the formal deformation theory.

By definition, a *trivial deformation* depends on the following data:

(T1) a formal power series $\varphi_{X,Y}: C(X, Y) \rightarrow C(X, Y)$, for any $X, Y \in C$ of the form

$$\varphi_{X,Y}(t) = \text{Id}_{C(X,Y)} + t \cdot \varphi_{X,Y}^1 + t^2 \cdot \varphi_{X,Y}^2 + \dots, \tag{7.5}$$

where $\varphi_{X,Y}^i \in \text{Hom}_{\mathbb{k}}(C(X, Y), C(X, Y)), i \geq 1$,

(T2) a formal power series $\psi_{X,Y}: C(X \otimes Y, X \otimes Y)$, for any $X, Y \in C$, of the form

$$\psi_{X,Y} = \text{Id}_{X \otimes Y} + t \cdot \psi_{X,Y}^1 + t^2 \cdot \psi_{X,Y}^2 + \dots \tag{7.6}$$

where $\psi_{X,Y}^i \in C(X \otimes Y, X \otimes Y), i \geq 1$.

Out of this data, a formal deformation of C is constructed as in (7.2)–(7.4).

One defines the concepts of an *infinitesimal deformation* and of a *trivial infinitesimal deformation* of a monoidal (linear or dg) category by replacing in the previous definitions the ring of formal power series $\mathbb{k}[[t]]$ by the dual numbers $\mathbb{k}[t]/(t^2)$. We say that two infinitesimal deformations belong to the same equivalence class if the corresponding monoidal categories are equivalent by an (extended) twist, as in Lemma 7.2 but over $\mathbb{k}[t]/(t^2)$.

One has the following theorem.

Theorem 7.3. *Let C be a \mathbb{k} -linear (or a dg over \mathbb{k}) monoidal category. The third cohomology $H^3(\text{Tot}_{\Theta_2} A((C, C)(\text{Id}, \text{Id})(\text{id}, \text{id})))$ is isomorphic to the equivalence classes of infinitesimal deformations (in the sense specified above) of the monoidal (dg) category C .*

Proof. The proof is a rather long but standard computation, for which we refer the reader to [29, Sect. 5]. ■

7.2. Infinitesimal deformation theory of a strict monoidal functor

The following theorem is proven analogously but is easier than Theorem 7.3, and we leave the details to the reader.

Theorem 7.4. *Let C, D be \mathbb{k} -linear (or dg over \mathbb{k}) monoidal categories, $F: C \rightarrow D$ a monoidal functor. The second cohomology $H^2(\text{Tot}_{\Theta_2} A(C, D)(F, F)(\text{id}, \text{id}))$ is isomorphic to the equivalence classes of infinitesimal deformations of the functor F .*

By Theorem 6.6, $\text{Tot}_{\Theta_2} A(C, D)(F, F)$ is a homotopy 2-algebra. In fact, one can construct a dg Lie algebra on $\text{Tot}_{\Theta_2} A(C, D)(F, F)[1]$ directly (without any use of loc. cit.), and develop, via the Maurer–Cartan equation and the deformation functor associated with dg Lie algebra formalism, the “global” deformation theory for $F: C \rightarrow D$ over $\mathbb{k}[[t]]$.

Appendix: Relations in Θ_2

One has the following relations between the elementary face and degeneracy maps in Θ_2 , which are checked straightforwardly:

$$D_{q,\sigma'} D_{p,\sigma} = D_{p,\sigma} D_{q-1,\sigma'} \quad \text{if } p < q - 1, \tag{A.1}$$

$$D_{q,\sigma_2} D_{q-1,\sigma_1} = D_{q-1,\eta_2} D_{q-1,\eta_1}. \tag{A.2}$$

Here is an explanation of the notations: any (a, b) -shuffle σ_1 and $(a + b, c)$ -shuffle σ_2 define uniquely a (b, c) -shuffle η_1 and an $(a, b + c)$ -shuffle η_2 such that $\sigma_2 \circ (\sigma_1, \text{id}_c) = \eta_2 \circ (\text{id}_a, \eta_1)$ (the latter is an (a, b, c) -shuffle). Consider

$$\partial_p^j \partial_q^i = \partial_q^i \partial_p^j \quad \text{if } p \neq q, \tag{A.3}$$

$$\partial_p^j \partial_p^i = \partial_p^i \partial_p^{j-1} \quad \text{if } i < j, \tag{A.4}$$

$$D_{q,\sigma} \partial_p^j = \begin{cases} \partial_{p+1}^j D_{q,\sigma} & \text{if } p > q, \\ \partial_p^j D_{q,\sigma} & \text{if } p < q, \end{cases} \tag{A.5}$$

$$D_{p,\sigma} \partial_p^i = \begin{cases} \partial_p^a D_{p,\bar{\sigma}} & \text{if } \sigma^{-1}(\overrightarrow{i, i+1}) = \overrightarrow{a, a+1} \in [0, k_p], \\ \partial_{p+1}^b D_{p,\bar{\sigma}} & \text{if } \sigma^{-1}(\overrightarrow{i, i+1}) = \overrightarrow{b, b+1} \in [k_p, k_p + k_{p+1}], \end{cases} \quad (\text{A.6})$$

where $\bar{\sigma}$ is the shuffle obtained from σ by collapsing $\sigma^{-1}(\overrightarrow{i, i+1})$, and $\{k_s\}$ is used as in Section 2.6, (F2). Consider

$$\begin{aligned} \partial_p^i D_{\min} &= D_{\min} \partial_{p-1}^i & \text{if } p \geq 1, \\ D_{p,\sigma} D_{\min} &= D_{\min} D_{p-1,\sigma} & \text{if } p \geq 1 \end{aligned} \quad (\text{A.7})$$

and similarly for D_{\max} . Moreover,

$$\begin{aligned} \varepsilon_p^j \circ \varepsilon_q^i &= \varepsilon_q^i \circ \varepsilon_p^j & \text{if } p \neq q, \\ \varepsilon_p^j \circ \varepsilon_p^i &= \varepsilon_p^i \circ \varepsilon_p^{j-1} & \text{if } i \leq j, \\ \Upsilon_0^q \circ \Upsilon_0^p &= \Upsilon_0^p \circ \Upsilon_0^{q+1} & \text{if } p \leq q, \\ \Upsilon_\ell^q \circ \varepsilon_p^j &= \begin{cases} \varepsilon_{p-1}^j \circ \Upsilon_\ell^q & \text{if } p > q + 1, \\ \varepsilon_p^j \circ \Upsilon_\ell^q & \text{if } p \leq q, \\ \Upsilon_{\ell+1}^q & \text{if } p = q + 1, \end{cases} \\ \partial_p^i \circ \varepsilon_q^j &= \varepsilon_q^j \circ \partial_p^i & \text{if } p \neq q, \\ \varepsilon_p^j \circ \partial_p^i &= \begin{cases} \partial_p^i \circ \varepsilon_p^{j-1} & \text{if } i < j, \\ \text{id} & \text{if } i = j, j + 1, \\ \partial_p^{i-1} \circ \varepsilon_p^j & \text{if } i > j + 1, \end{cases} \\ \Upsilon_\ell^q \circ \partial_p^j &= \begin{cases} \partial_{p-1}^j \circ \Upsilon_\ell^q & \text{if } p > q + 1, \\ \partial_p^j \circ \Upsilon_\ell^q & \text{if } p \leq q, \\ \Upsilon_{\ell-1}^q & \text{if } p = q + 1, \end{cases} \\ D_{q,\sigma} \circ \varepsilon_p^i &= \begin{cases} \varepsilon_{p+1}^i \circ D_{q,\sigma} & \text{if } q < p, \\ \varepsilon_p^i \circ D_{q,\sigma} & \text{if } q > p, \\ \varepsilon_p^a \circ D_{q,\sigma'} & \text{if } q = p, \sigma^{-1}(\overrightarrow{i, i+1}) = \overrightarrow{a, a+1} \in [0, k_p], \\ \varepsilon_{p+1}^b \circ D_{q,\sigma'} & \text{if } q = p, \sigma^{-1}(\overrightarrow{i, i+1}) = \overrightarrow{b, b+1} \in [k_p, k_p + k_{p+1}], \end{cases} \end{aligned}$$

where σ' is obtained from σ by adding a new element (blowing up) at $\sigma^{-1}(\overrightarrow{i, i+1})$. Then,

$$\begin{aligned} \Upsilon_0^q \circ D_{p,\sigma} &= \begin{cases} D_{p,\sigma} \circ \Upsilon_0^{q-1} & \text{if } p < q, \\ D_{p-1,\sigma} \circ \Upsilon_0^q & \text{if } p > q + 1, \\ \text{id} & \text{if } p = q, \sigma = (0, k_p + k_{p+1}), \\ \text{id} & \text{if } p = q + 1, \sigma = (k_p + k_{p+1}, 0), \end{cases} \\ D_{\min} \circ \varepsilon_p^i &= \varepsilon_{p+1}^i \circ D_{\min}, \end{aligned}$$

$$\begin{aligned}
 D_{\max} \circ \varepsilon_p^i &= \varepsilon_p^i \circ D_{\max}, \\
 \Upsilon_0^q \circ D_{\min} &= \begin{cases} D_{\min} \circ \Upsilon_0^{q-1} & \text{if } q > 0, \\ \text{id} & \text{if } q = 0, \end{cases} \\
 \Upsilon_0^q \circ D_{\max} &= \begin{cases} D_{\max} \circ \Upsilon_0^q & \text{if } q < n + 1, \\ \text{id} & \text{if } q = n + 1. \end{cases}
 \end{aligned}$$

Acknowledgements. The authors are thankful to Michael Batanin for his interest and suggestions.

Funding. The work of Piergiorgio Panero was supported by the FWO Research Project G060118N. The work of Boris Shoikhet was supported by Support Grant for International Mathematical Centres Creation and Development, by the Ministry of Higher Education and Science of Russian Federation and PDMI RAS agreement No. 075-15-2022-289 issued on April 6, 2022.

References

- [1] D. Ara, [Higher quasi-categories vs higher Rezk spaces](#). *J. K-Theory* **14** (2014), no. 3, 701–749 Zbl [1322.18002](#) MR [3350089](#)
- [2] M. Batanin, [Monoidal globular categories as a natural environment for the theory of weak \$n\$ -categories](#). *Adv. Math.* **136** (1998), no. 1, 39–103 Zbl [0912.18006](#) MR [1623672](#)
- [3] M. Batanin, [Symmetrisation of \$n\$ -operads and compactification of real configuration spaces](#). *Adv. Math.* **211** (2007), no. 2, 684–725 Zbl [1146.18003](#) MR [2323542](#)
- [4] M. Batanin, [The Eckmann-Hilton argument and higher operads](#). *Adv. Math.* **217** (2008), no. 1, 334–385 Zbl [1138.18003](#) MR [2365200](#)
- [5] M. Batanin and C. Berger, [The lattice path operad and Hochschild cochains](#). In *Alpine perspectives on algebraic topology*, pp. 23–52, Contemp. Math. 504, American Mathematical Society, Providence, RI, 2009 Zbl [1221.18006](#) MR [2581904](#)
- [6] M. Batanin and A. Davydov, [Cosimplicial monoids and deformation theory of tensor categories](#). *J. Noncommut. Geom.* **17** (2023), no. 4, 1167–1229 Zbl [1524.18038](#) MR [4653784](#)
- [7] M. Batanin and M. Markl, [Centers and homotopy centers in enriched monoidal categories](#). *Adv. Math.* **230** (2012), no. 4–6, 1811–1858 Zbl [1279.18004](#) MR [2927355](#)
- [8] M. Batanin and M. Markl, [Operadic categories and duoidal Deligne’s conjecture](#). *Adv. Math.* **285** (2015), 1630–1687 Zbl [1360.18009](#) MR [3406537](#)
- [9] J. Bénabou, [Introduction to bicategories](#). In *Reports of the Midwest Category Seminar*, pp. 1–77, Lecture Notes in Math. 47, Springer, Berlin, 1967 MR [0220789](#)
- [10] C. Berger, [A cellular nerve for higher categories](#). *Adv. Math.* **169** (2002), no. 1, 118–175 Zbl [1024.18004](#) MR [1916373](#)
- [11] C. Berger, [Iterated wreath product of the simplex category and iterated loop spaces](#). *Adv. Math.* **213** (2007), no. 1, 230–270 Zbl [1127.18008](#) MR [2331244](#)
- [12] J. E. Bergner and C. Rezk, [Reedy categories and the \$\Theta\$ -construction](#). *Math. Z.* **274** (2013), no. 1–2, 499–514 Zbl [1270.55014](#) MR [3054341](#)

- [13] A. A. Davydov, [Monoidal categories](#). *J. Math. Sci. (New York)* **88** (1998), no. 4, 457–519
Zbl [0907.18002](#) MR [1613071](#)
- [14] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, [Tensor categories](#). Math. Surveys Monogr. 205, American Mathematical Society, Providence, RI, 2015 Zbl [1365.18001](#) MR [3242743](#)
- [15] M. Gerstenhaber and S. D. Schack, [Bialgebra cohomology, deformations, and quantum groups](#). *Proc. Nat. Acad. Sci. U.S.A.* **87** (1990), no. 1, 478–481 Zbl [0695.16005](#) MR [1031952](#)
- [16] R. Gordon, A. J. Power, and R. Street, [Coherence for tricategories](#). *Mem. Amer. Math. Soc.* **117** (1995), no. 558, vi+81 Zbl [0836.18001](#) MR [1261589](#)
- [17] A. Guan, A. Lazarev, Y. Sheng, and R. Tang, Review of deformation theory II: a homotopical approach. *Adv. Math. (China)* **49** (2020), no. 3, 278–298 Zbl [1463.18001](#) MR [4140875](#)
- [18] P. S. Hirschhorn, [Model categories and their localizations](#). Math. Surveys Monogr. 99, American Mathematical Society, Providence, RI, 2003 MR [1944041](#)
- [19] A. Joyal, Disks, duality, and Θ -categories. 1997, preprint
- [20] G. M. Kelly, [Basic concepts of enriched category theory](#). London Math. Soc. Lecture Note Ser. 64, Cambridge University Press, Cambridge, 1982 Zbl [0478.18005](#) MR [0651714](#)
- [21] M. Kontsevich, [Deformation quantization of Poisson manifolds](#). *Lett. Math. Phys.* **66** (2003), no. 3, 157–216 Zbl [1058.53065](#) MR [2062626](#)
- [22] M. Kontsevich and Y. Soibelman, Deformations of algebras over operads and the Deligne conjecture. In *Conférence Moshé Flato 1999, Vol. I (Dijon)*, pp. 255–307, Math. Phys. Stud. 21, Kluwer Academic Publishers, Dordrecht, 2000 Zbl [0972.18005](#) MR [1805894](#)
- [23] T. Leinster, Basic bicategories. 1998, arXiv:[arXiv:math/9810017v1](#)
- [24] J. Lurie, Formal moduli problems. [www.math.ias.edu/~lurie/](#) visited on 4 September 2024
- [25] S. MacLane, [Categories for the working mathematician](#). Grad. Texts in Math. 5, Springer, New York, 1971 Zbl [0705.18001](#) MR [0354798](#)
- [26] S. MacLane, [Homology. Reprint of the 1975 edition](#). Classics Math., Springer, Berlin, 1995 Zbl [0818.18001](#) MR [1344215](#)
- [27] J. E. McClure and J. H. Smith, [A solution of Deligne’s Hochschild cohomology conjecture](#). In *Recent progress in homotopy theory (Baltimore, MD, 2000)*, pp. 153–193, Contemp. Math. 293, American Mathematical Society, Providence, RI, 2002 Zbl [1009.18009](#) MR [1890736](#)
- [28] J. E. McClure and J. H. Smith, [Multivariable cochain operations and little \$n\$ -cubes](#). *J. Amer. Math. Soc.* **16** (2003), no. 3, 681–704 Zbl [1014.18005](#) MR [1969208](#)
- [29] P. Panero and B. Shoikhet, The category Θ_2 , derived modifications, and deformation theory of monoidal categories. [v1] 2022, [v2] 2024, arXiv:[2210.01664v2](#)
- [30] J. P. Pridham, [Unifying derived deformation theories](#). *Adv. Math.* **224** (2010), no. 3, 772–826 Zbl [1195.14012](#) MR [2628795](#)
- [31] E. Riehl, [Categorical homotopy theory](#). New Math. Monogr. 24, Cambridge University Press, Cambridge, 2014 Zbl [1317.18001](#) MR [3221774](#)
- [32] B. Shoikhet, 2-shuffles, 3-operads, and Deligne conjecture in dimension 3. In progress
- [33] B. Shoikhet, [Tetramodules over a bialgebra form a 2-fold monoidal category](#). *Appl. Categ. Structures* **21** (2013), no. 3, 291–309 Zbl [1288.18006](#) MR [3047010](#)
- [34] B. Shoikhet, [Differential graded categories and Deligne conjecture](#). *Adv. Math.* **289** (2016), 797–843 Zbl [1333.18010](#) MR [3439700](#)
- [35] D. Tamarkin, Another proof of M. Kontsevich formality theorem. 1998, arXiv:[math/9803025v4](#)
- [36] D. Tamarkin, Deformation complex of a d -algebra is a $(d + 1)$ -algebra. 2000, arXiv:[math/0010072v1](#)

- [37] D. Tamarkin, [What do dg-categories form?](#) *Compos. Math.* **143** (2007), no. 5, 1335–1358
Zbl [1138.18004](#) MR [2360318](#)
- [38] C. A. Weibel, [An introduction to homological algebra](#). Cambridge Stud. Adv. Math. 38, Cambridge University Press, Cambridge, 1994 Zbl [0797.18001](#) MR [1269324](#)
- [39] D. N. Yetter, [Braided deformations of monoidal categories and Vassiliev invariants](#). In *Higher category theory (Evanston, IL, 1997)*, pp. 117–134, Contemp. Math. 230, American Mathematical Society, Providence, RI, 1998 Zbl [0927.18003](#) MR [1664995](#)

Received 3 March 2023; revised 15 June 2024.

Piergiorgio Panero

Universiteit Antwerpen, Campus Middelheim, Wiskunde en Informatica, Gebouw G,
Middelheimlaan 1, 2020 Antwerpen, Belgium; panero1994@gmail.com

Boris Shoikhet

Euler International Mathematical Institute (EIMI), 10 Pesochnaya Embankment, 197376 Saint
Petersburg, Russia; shoikhet@pdmi.ras.ru