

Finite Ramsey theory through category theory

Sławomir SOLECKI

Abstract. We present a new, category-theoretic point of view on finite Ramsey theory. Our aims are as follows:

- to define the category-theoretic notions needed for the development of finite Ramsey theory;
- to state, in terms of these notions, the general fundamental Ramsey results (of which various concrete Ramsey results are special cases); and
- to give self-contained proofs within the category-theoretic framework of these general results.

We also provide some concrete illustrations of the general method.

Mathematics Subject Classification 2020: 05D10 (primary); 05C55, 18A22 (secondary).

Keywords: Ramsey theory, category theory, frank functors.

1. Introduction

Ramsey theory is a subfield of combinatorics whose roots lie in logic, see [15]; but see also [21] for another early source of Ramsey-theoretic ideas. It is an area that aims to find implementations of the slogan “total disorder is unavoidable”—its theorems search, in a space of elements under discussion, for subspaces whose elements are not distinguishable from each other. Such theorems have a common form but they differ deeply in the type of elements and spaces they concern and in the methods employed in their proofs. At this point of its development, Ramsey theory strongly connects not only with logic but also with topological dynamics [10, 22] and Banach space theory [1], and, of course, it continues to be an important branch of combinatorics [14].

The present paper treats finite Ramsey theory, the part of Ramsey theory in which the elements are finite, the spaces of elements are finite, while the proofs rely on iterative inductive arguments. The paper can be seen as a contribution to the efforts aimed at unifying Ramsey-theoretic results, as in [8] for finite structural Ramsey theory, in [2] and [20] for infinite-dimensional Ramsey theory, and in [18] for the ultrafilter methods in Ramsey theory.

We give now examples of some classical Ramsey statements, namely Ramsey's original theorem [15], van der Waerden's theorem [21], and the dual Ramsey theorem of Graham–Rothschild [5]. Our goal in reminding the reader about these results is partly to clarify what type of theorems form the starting point of the category theory approach presented in this paper. For a positive integer r , by an r -coloring of a set A we understand a function on A with at most r values. Ramsey's original theorem proved in [15] is the following statement.

For positive integers r and k, l , there exists a positive integer m such that, for each r -coloring of all k -element subsets of an m -element set X , there exists an l -element set $Y \subseteq X$ such that all k -element subsets of Y get the same color.

We cover this theorem as part of our framework in Section 4.4. The following statement is van der Waerden's theorem [21].

For positive integers r, k , there is a positive integer l such that, for each r -coloring of the set $\{1, 2, \dots, l\}$, there exists an arithmetic progression of length k , whose elements get the same color.

A purely combinatorial phrasing of the arithmetic statement above was found by Hales and Jewett in [7]. We treat this combinatorial statement as part of our approach in Section 3.5. Finally, we state the dual Ramsey theorem of Graham–Rothschild [5]. By a k -partition of a set X we understand a family of k non-empty sets whose union is X .

For positive integers r and k, l , there exists a positive integer m such that, for each r -coloring of all k -partitions of an m -element set X , there exists an l -partition P of X such that all k -partitions of X that are coarser than P get the same color.

Again, the above statement can be seen as a particular case of the general theorems of this paper. We do not present this derivation in detail but we invite the reader to do it on their own, perhaps following the lead of [16, Section 8.2].

We move to discussing the category-theoretic set-up. If C is a category, by $\text{ob}(C)$ we denote the class of all objects of C ; for $a, b \in \text{ob}(C)$, $\text{hom}(a, b)$ stands for the class of all morphisms in C from a to b . It has been known for some time, at least since the early 1970s, that Ramsey-theoretic statements are naturally expressed in the language of category theory. Given a category C , an object $a \in \text{ob}(C)$ is said to have the *Ramsey property* if for each object $b \in \text{ob}(C)$ and a positive integer r , there is $c \in \text{ob}(C)$ such that for each r -coloring of $\text{hom}(a, c)$, there is $g \in \text{hom}(b, c)$ that makes the set

$$g \cdot \text{hom}(a, b) = \{g \cdot f \mid f \in \text{hom}(a, b)\}$$

monochromatic. One then calls C a *Ramsey category* if each of its objects has the Ramsey property. It is also known that the following notion is a useful refinement of the Ramsey property. For $a, b \in \text{ob}(C)$, the *Ramsey degree* of the pair a, b is the smallest positive integer k , if such a number exists, such that for each positive integer r , there exists $c \in \text{ob}(C)$ with the property that, for each r -coloring χ of $\text{hom}(a, c)$, there exists $g \in \text{hom}(b, c)$ with χ attaining at most k values on $g \cdot \text{hom}(a, b)$. If such a number k does not exist, we say that the Ramsey degree of the pair a, b is ∞ . We write

$$(1) \quad \text{rd}(a, b)$$

for the Ramsey degree of a, b . More studied, and often more useful, notion is that of Ramsey degree of a single object. *Ramsey degree of a* is defined by

$$(2) \quad \text{rd}(a) = \sup_{b \in \text{ob}(C)} \text{rd}(a, b).$$

Having the Ramsey property is expressible in terms of the Ramsey degree— a has the Ramsey property precisely when $\text{rd}(a) = 1$. This observation leads to an important easing of the condition of the category C being Ramsey to the condition of C having *finite Ramsey degrees*, which asserts that, for each $a \in \text{ob}(C)$, $\text{rd}(a) < \infty$. The importance of finiteness of Ramsey degree stems partly from it being a refinement of the Ramsey property and partly, and more significantly, from its relevance to topological dynamics as indicated in [10] and, especially, in [22].

Going beyond just formulating Ramsey-theoretic notions, several papers used the language of category theory to carry out proofs of finite Ramsey-theoretic statements; see, for example, [4, 9, 11–13]. This was usually done by identifying a category A that was known to be Ramsey or to have a related property, and then transferring Ramseyness or the related property from A to another category B by finding an, often subtle, connection between A and B . Another distinct approach was presented by Leeb in [11]. In a number of specific categories, he verified certain identities, called by him Pascal identities, and then working separately in each of these categories, but using similarly structured arguments, he showed that the categories are Ramsey, from which various classical Ramsey theorems followed. In [9], certain Ramsey-theoretic constructions were revealed to be canonical category theory constructions. In his paper [6], Gromov advocated for a broad use of category theory in Ramsey theory.

In the current paper, expanding and simplifying the author's approach of [16] and following the spirit of [6], we present a way of seeing finite Ramsey theory in the category-theoretic terms that is global, in the sense that the whole theory is developed from scratch in the category-theoretic framework as opposed to transferring specific Ramsey results between categories or running separate proofs in different categories. We change the usual perspective and consider functors, rather than categories, as

fundamental to the development. More precisely, we formulate a general pigeonhole principle (P) for functors between categories. On our view, this is the main notion of finite Ramsey theory, and the theory is concerned with proving (P) for various functors. To this end, we show that (P) persists under several natural operations applied to functors. Further, we formulate a localized version (FP) of (P) and show that, for essentially all relevant functors, (FP) implies (P). The advantage of having this implication resides in a relative ease of proving the localized pigeonhole principle (FP) for many functors in comparison with proving (P) itself. These general theorems allow us to establish results giving an upper bound on Ramsey degrees. As explained above, such upper estimates are generalizations of the statement that the category is Ramsey thereby making it possible to deduce various concrete Ramsey theorems.

It may be worth emphasizing that the point we are making is not so much that the concrete Ramsey theorems can be derived using our methods, it is more that they are particular cases (for concrete functors) of our results. We give several examples of theorems that can be seen as such—Ramsey’s theorem [15] and the product Ramsey theorem, both in Section 4.4; the Hales–Jewett theorem [7] in Section 3.5; Fouché’s Ramsey theorem for trees [3] in Section 4.5. These examples should be viewed merely as illustrations since many other theorems, for example, the Ramsey theorems for trees due to various mathematicians that are surveyed in [17] and treated there with the methods of [16], the Ramsey theorems considered in [16], for example, the dual Ramsey theorem of Graham–Rothschild [5], and the dual Ramsey theorem for trees from [19], can all be seen as, in essence, particular cases of the general theorems of this paper.

As already mentioned, our approach here builds on [16]. The main advances with respect to that paper consist of the use of categories and functors instead of various types of ad hoc structures (in particular, eliminating partial or linear orders from the general structures), weakening of the localized pigeonhole principle (from (LP) in [16] to (FP) here), obtaining upper estimates on the Ramsey degree rather than just Ramsey statements, and an overall substantial simplification of the presentation.

The following conventions will be used throughout. By \mathbb{N} we understand the set of all natural numbers including 0. For $m, n \in \mathbb{Z}$, we write

$$[m, n] = \{i \in \mathbb{Z} \mid m \leq i \leq n\}.$$

For $n \in \mathbb{N}$ and $m = 1$, we shorten the above piece of notation to

$$[n] = [1, n].$$

In particular, $[0] = \emptyset$. The cardinality of a set x will be denoted by $|x|$. So, if x is finite, then $|x| \in \mathbb{N}$. For a functor γ defined on a category C and for an object a of C and a

morphism f of C , we often write

$$\gamma a \text{ and } \gamma f$$

for $\gamma(a)$ and $\gamma(f)$.

2. Condition (P) and frank functors

2.1. Formulation of condition (P). We regard the following statement as the fundamental pigeonhole principle for a functor $\delta: C \rightarrow D$ between the categories C and D . In a nutshell, it says that, in a suitable sense, δ controls colorings.

Definition 2.1. Let δ be a functor defined on a category C , and let $a, b \in \text{ob}(C)$. We declare δ to *fulfill* (P) at a, b if for each $r \in \mathbb{N}$, there exists $c \in \text{ob}(C)$ such that, for each r -coloring χ of $\text{hom}(a, c)$, there exists $g \in \text{hom}(b, c)$ with

$$\delta f_1 = \delta f_2 \implies \chi(g \cdot f_1) = \chi(g \cdot f_2),$$

for all $f_1, f_2 \in \text{hom}(a, b)$.

For ease of phrasing, we adopt the following conventions. We say that δ *fulfills* (P) at $a \in \text{ob}(C)$ if it fulfills (P) at a, b for all $b \in \text{ob}(C)$. We say simply that δ *fulfills* (P) if it fulfills (P) at all $a, b \in \text{ob}(C)$. Note that the condition of fulfilling (P) at a is stronger than fulfilling (P) at a, b . We will be mostly interested in this stronger condition, but using the weaker condition is somewhat easier and permits us to make more precise statements in some situations.

The fundamental connection of property (P) with Ramsey-theoretic notions of Ramsey degrees (recall (1) and (2) here) goes through the following proposition, which improves the trivial bound $\text{rd}(a, b) \leq |\text{hom}(a, b)|$.

Proposition 2.2. *Let Δ be a family of functors whose domains are all equal to C .*

(i) *If each $\delta \in \Delta$ fulfills (P) at $a, b \in \text{ob}(C)$, then*

$$\text{rd}(a, b) \leq \min_{\delta \in \Delta} |\delta(\text{hom}(a, b))|.$$

(ii) *If each $\delta \in \Delta$ fulfills (P) at $a \in \text{ob}(C)$, then*

$$\text{rd}(a) \leq \sup_{b \in \text{ob}(C)} \min_{\delta \in \Delta} |\delta(\text{hom}(a, b))|.$$

Proof. By the definition of $\text{rd}(a)$ in (2), it suffices to show (i). This amounts to proving that given $r \in \mathbb{N}$, there exists c such that for each r -coloring of $\text{hom}(a, c)$, there exists $g \in \text{hom}(b, c)$ with the number of colors attained on $g \cdot \text{hom}(a, b)$ bounded by $|\delta(\text{hom}(a, b))|$. This statement is an immediate consequence of δ fulfilling (P) at a . ■

2.2. Definition of frank functors. When dealing with property (P) for a functor δ , sometimes we will need to make an additional surjectivity assumption on δ . This surjectivity assumption is critical and seems interesting enough to isolate it here.

Definition 2.3. A functor $\delta: C \rightarrow D$ between two categories C and D is called *frank* if, for all $a \in \text{ob}(C)$ and $b' \in \text{ob}(D)$, there exists $b \in \text{ob}(C)$ with

$$\delta(b) = b' \quad \text{and} \quad \delta(\text{hom}(a, b)) = \text{hom}(\delta a, \delta b).$$

In the second equality in the definition above, the inclusion $\delta(\text{hom}(a, b)) \subseteq \text{hom}(\delta a, \delta b)$ follows just from δ being a functor. Therefore, the point of the equality is that δ is surjective as a function from $\text{hom}(a, b)$ to $\text{hom}(\delta a, \delta b)$.

Lemma 2.4. (i) *The identity functor is frank.*

(ii) *The composition of frank functors is a frank functor.*

(iii) *If $\delta: C \rightarrow D$ is a frank functor, then, for all $d_1, d_2 \in \text{ob}(D)$, there are $c_1, c_2 \in \text{ob}(C)$ such that*

$$\delta(c_1) = d_1, \quad \delta(c_2) = d_2, \quad \text{and} \quad \delta(\text{hom}(c_1, c_2)) = \text{hom}(d_1, d_2).$$

Proof. Points (i) and (ii) are almost immediate, and we leave checking them to the reader. To see (iii), given d_1 , use frankness of δ to find c_1 with $\delta(c_1) = d_1$. Now use frankness of δ again to find c_2 with $\delta(c_2) = d_2$ and $\delta(\text{hom}(c_1, c_2)) = \text{hom}(d_1, d_2)$. ■

2.3. Unifying assumptions. Without harming applicability in finite Ramsey theory of the general theorems presented below, one may always make the following unifying assumptions:

- *functors are frank;*
- *categories are such that $\text{hom}(a, b)$ is finite for all objects a, b .*

All the general theorems proved below hold under these assumptions. Of course, when stating the theorems, we make assumptions that are appropriate (minimal) for each theorem.

2.4. Examples—frank functors fulfilling (P). We describe here some examples of frank functors with property (P).

1. Let C be a category. The identity functor $C \rightarrow C$ is frank and fulfills (P) at each pair of objects of C .

2. We define here a category P and a frank functor $\partial_P: P \rightarrow P$ that will play an important auxiliary role in proving the Hales–Jewett theorem. Fulfilling of condition (P) by ∂_P is the usual pigeonhole principle.

Objects of P are

- pairs (k, i) , where $k \in \mathbb{N}$, $k \geq 1$ if $i \in \{0, 2\}$, and $k \geq 2$ if $i = 1$.

Morphisms of P from an object (k, i) to an object (k', i') will be certain functions from $[k']$ to $[k]$, whose nature will depend on the second coordinates i and i' . We define the morphisms as follows:

- $p \in \text{hom}((l, 2), (m, 2))$, if $p: [m] \rightarrow [l]$ is a surjection with $p(i) \leq p(i + 1) \leq p(i) + 1$ for $i \in [m - 1]$;
- $x \in \text{hom}((k_1, 1), (l, 2))$, if $x: [l] \rightarrow [k_1]$ and there are $1 \leq a < a + 1 < b \leq l$ such that x is constant on the intervals $[1, a]$, $[a + 1, b - 1]$, and $[b, l]$ and

$$x(1) = k_1 \quad \text{and} \quad x(l) = k_1 - 1;$$

- $x \in \text{hom}((k_0, 0), (l, 2))$, if $x: [l] \rightarrow [k_0]$ and there are $1 \leq a < a + 1 < b \leq l$ such that x is constant on the intervals $[1, a]$, $[a + 1, b - 1]$, and $[b, l]$ and

$$x(1) = x(l) = k_0;$$

- there are no other morphisms except for identities.

Composition of morphisms in P will be the composition of functions taken with reverse order:

- for $p \in \text{hom}((l, 2), (m, 2))$ and $x \in \text{hom}((k, i), (l, 2))$, with $i \in \{0, 1\}$, let

$$p \cdot x = x \circ p;$$

- for morphisms $p \in \text{hom}((l, 2), (m, 2))$ and $q \in \text{hom}((m, 2), (n, 2))$, let

$$q \cdot p = p \circ q.$$

The functor $\partial_P: P \rightarrow P$ acts non-trivially only on objects of the form $(k, 1)$ and on morphisms in $\text{hom}((k, 1), (l, 2))$, on which it lowers the top value by 1. So the functor ∂_P is defined by:

- $\partial_P(k, i) = (k, i)$, for $i \neq 1$, and $\partial_P(k, 1) = (k - 1, 0)$;
- $\partial_P p = p$, for $p \in \text{hom}((l, 2), (m, 2))$;
- $\partial_P x = x$, for $x \in \text{hom}((k_0, 0), (l, 2))$;
- $\partial_P x = \min(x, k_1 - 1)$, for $x \in \text{hom}((k_1, 1), (l, 2))$.

Lemma 2.5. *The following statements hold:*

- (i) $\partial_P: P \rightarrow P$ is a functor and it is frank.
- (ii) ∂_P fulfills (P).

Proof. Point (i) is straightforward, and we leave checking it to the reader.

To see (ii), we check (P) at $(k, 1)$, $(l, 2)$, which is the only not entirely trivial case. After fixing $r \in \mathbb{N}$, we need to find an object $(m, 2)$ for which the conclusion of (P) holds. It is good to keep in mind that all we are doing is proving a version of the standard pigeonhole principle.

Put $m = (l - 1)r + 2$. Let χ be an r -coloring of $\text{hom}((k_1, 1), (m, 2))$, that is, an r -coloring of the set of all $x: [m] \rightarrow [k_1]$, for which there exist $1 \leq a_x < a_x + 1 < b_x \leq m$ such that x is constant on the intervals $[1, a_x]$, $[a_x + 1, b_x - 1]$, and $[b_x, m]$, and $x(1) = k_1 - 1$ and $x(m) = k_1$. Consider the r -coloring χ' of $[m - 1]$ given by

$$\chi'(j) = \chi(x_j),$$

where $x_j: [m] \rightarrow [k_1]$, with $j \in [m - 1]$, is such that

$$x_j \upharpoonright [j] = k_1 - 1 \quad \text{and} \quad x_j \upharpoonright [j + 1, m] = k_1.$$

Note that $x_j \in \text{hom}((k_1, 1), (m, 2))$, so $\chi(x_j)$ is defined. By our choice of m , χ' is constant on a subset of $[m - 1]$ of size l . So there exists $p: [m] \rightarrow [l]$, a surjection with $p(j) \leq p(j + 1) \leq p(j) + 1$ for $j \in [m - 1]$, that is, $p \in \text{hom}((l, 2), (m, 2))$, and such that χ' is constant on the set

$$J = \{j \in [m - 1] \mid p(j) < p(j + 1)\}.$$

The above condition on χ' means that $\chi(x_j)$ is constant as j varies over J .

We claim that this p works. Indeed, let $x, x' \in \text{hom}((k_1, 1), (l, 2))$ be such that $\partial_P x = \partial_P x'$. Then either $x = x'$ or there are $i, i' \in [l - 1]$ such that

$$\begin{aligned} x \upharpoonright [i] &= k_1 - 1, & x \upharpoonright [i + 1, m] &= k_1, \\ x' \upharpoonright [i'] &= k_1 - 1, & x' \upharpoonright [i' + 1, m] &= k_1. \end{aligned}$$

In the first case, clearly $\chi(x \circ p) = \chi(x' \circ p)$. In the second case, for $j, j' \in J$ specified by

$$i = p(j) < p(j + 1) \quad \text{and} \quad i' = p(j') < p(j' + 1),$$

we have $x \circ p = x_j$ and $x' \circ p = x_{j'}$. Thus, we get

$$\chi(x \circ p) = \chi(x_j) = \chi(x_{j'}) = \chi(x' \circ p),$$

as required. ■

3. The following example does fulfill condition (P), but we postpone its verification till we have a general result from which it will follow readily. The example will serve to prove the standard Ramsey theorem.

We define a category R and a functor $\partial_R: R \rightarrow R$. The objects of the underlying category will be natural numbers n and morphisms will be, essentially, subsets x of $[n]$. For technical reasons, for such a set x , we will need to remember which $[n]$ it is designated to be a subset of; so the morphisms will actually be pairs (x, n) . For $n \in \mathbb{N}$, we write

$$(3) \quad n \dot{-} 1 = \max(n - 1, 0).$$

Objects of R are:

- n , for $n \in \mathbb{N}$.

Morphisms of R are described as follows:

- $(x, n) \in \text{hom}(m, n)$, for $x \subseteq [n]$ and $|x| = m$.

Composition in R is defined by the following rule:

- for morphisms $(x, m) \in \text{hom}(l, m)$ and $(y, n) \in \text{hom}(m, n)$, let

$$(y, n) \cdot (x, m) = (f_y(x), n),$$

where $f_y: [m] \rightarrow y$ is the unique increasing bijection.

The functor $\partial_R: R \rightarrow R$ is defined using the notation set up by (3):

- $\partial_R n = n \dot{-} 1$;
- $\partial_R(x, n) = (x \setminus \{\max x\}, n \dot{-} 1)$, for $(x, n) \in \text{hom}(m, n)$ with $x \neq \emptyset$;
- $\partial_R(\emptyset, n) = (\emptyset, n \dot{-} 1)$, for $(\emptyset, n) \in \text{hom}(0, n)$.

3. Propagating condition (P)

In this section, we prove three theorems that let us transfer property (P) from one functor to another.

3.1. Composition. The following theorem asserts that, under appropriate assumptions, property (P) is preserved under composition of functors.

Theorem 3.1. *Let $\gamma: C \rightarrow D$ and $\delta: D \rightarrow E$ be functors with γ being frank. Let $a, b \in \text{ob}(C)$. If γ fulfills (P) at a and δ fulfills (P) at $\gamma(a), \gamma(b)$, then $\delta \circ \gamma$ fulfills (P) at a, b .*

Proof. We write $\delta\gamma$ for $\delta \circ \gamma$.

Fix $a, b \in \text{ob}(C)$ with the aim to show that $\delta\gamma$ has (P) at a, b . In order to do this, let $r \in \mathbb{N}$. The objects a, b and the natural number r will remain fixed for the rest of this proof.

Claim. *There is $c \in \text{ob}(C)$ such that for every r -coloring χ of $\text{hom}(\gamma a, \gamma c)$, there is $g \in \text{hom}(b, c)$ such that for $f_1, f_2 \in \text{hom}(a, b)$*

$$(4) \quad \delta\gamma f_1 = \delta\gamma f_2 \implies \chi(\gamma(g \cdot f_1)) = \chi(\gamma(g \cdot f_2)).$$

Proof of Claim. Since δ fulfills (P) at γa and γb , we can find $d \in \text{ob}(D)$ such that for every r -coloring χ of $\text{hom}(\gamma a, d)$, there is $g' \in \text{hom}(\gamma b, d)$ such that, for $h_1, h_2 \in \text{hom}(\gamma a, \gamma b)$, we have

$$(5) \quad \delta h_1 = \delta h_2 \implies \chi(g' \cdot h_1) = \chi(g' \cdot h_2).$$

By frankness of the functor γ , there is $c \in \text{ob}(C)$ such that $d = \gamma c$ and, for every $g' \in \text{hom}(\gamma b, d)$, there is $g \in \text{hom}(b, c)$ with $g' = \gamma g$. We claim that this c makes the conclusion of the Claim true.

Let χ be an r -coloring of $\text{hom}(\gamma a, \gamma c) = \text{hom}(\gamma a, d)$. By our choice of d , there is $g' \in \text{hom}(\gamma b, d)$ such that, for $h_1, h_2 \in \text{hom}(\gamma a, \gamma b)$, condition (5) holds. Let $g \in \text{hom}(b, c)$ be such that $g' = \gamma g$. In order to check condition (4), fix $f_1, f_2 \in \text{hom}(a, b)$ with

$$(6) \quad \delta\gamma f_1 = \delta\gamma f_2.$$

Note that $\gamma f_1, \gamma f_2 \in \text{hom}(\gamma a, \gamma b)$, and therefore by (5) and (6), we have

$$\chi(g' \cdot (\gamma f_1)) = \chi(g' \cdot (\gamma f_2)).$$

Since

$$g' \cdot (\gamma f_1) = (\gamma g) \cdot (\gamma f_1) = \gamma(g \cdot f_1) \quad \text{and} \quad g' \cdot (\gamma f_2) = (\gamma g) \cdot (\gamma f_2) = \gamma(g \cdot f_2),$$

it follows that $\chi(\gamma(g \cdot f_1)) = \chi(\gamma(g \cdot f_2))$, which gives (4) and the Claim. \square

Now, we prove the conclusion of the theorem from the claim. We are seeking $c \in \text{ob}(C)$ with the following property: for each r -coloring χ of $\text{hom}(a, c)$ there exists $g \in \text{hom}(b, c)$ such that, for $f_1, f_2 \in \text{hom}(a, b)$,

$$(7) \quad \delta\gamma f_1 = \delta\gamma f_2 \implies \chi(g \cdot f_1) = \chi(g \cdot f_2).$$

We apply the claim to obtaining $c' \in \text{ob}(C)$. Next, recall that we assume that γ fulfills (P) at a , so it fulfills (P) at a, c' , which, for the given r , yields $c \in \text{ob}(C)$. We claim that this c works.

Let χ be an r -coloring of $\text{hom}(a, c)$. By the choice of c there exists $g' \in \text{hom}(c', c)$ such that for $f_1, f_2 \in \text{hom}(a, b)$ and $h_1, h_2 \in \text{hom}(b, c')$,

$$(8) \quad \gamma(h_1 \cdot f_1) = \gamma(h_2 \cdot f_2) \implies \chi(g' \cdot (h_1 \cdot f_1)) = \chi(g' \cdot (h_2 \cdot f_2)).$$

We define an r -coloring $\bar{\chi}$ on $\text{hom}(\gamma a, \gamma c')$. First we specify $\bar{\chi}$ of the subset

$$\{\gamma(h \cdot f) \mid f \in \text{hom}(a, b), h \in \text{hom}(b, c')\}$$

of $\text{hom}(\gamma a, \gamma c')$. So, for $f \in \text{hom}(a, b)$ and $h \in \text{hom}(b, c')$, let

$$(9) \quad \bar{\chi}(\gamma(h \cdot f)) = \chi(g' \cdot h \cdot f).$$

The function $\bar{\chi}$ is well-defined by (8). Now we extend $\bar{\chi}$ to an r -coloring of the whole set $\text{hom}(\gamma a, \gamma c')$ in an arbitrary way. We denote this extension again by $\bar{\chi}$. By our choice of c' from Claim, there exists $g'' \in \text{hom}(b, c')$ such that, for $f_1, f_2 \in \text{hom}(a, b)$,

$$(10) \quad \delta \gamma f_1 = \delta \gamma f_2 \implies \bar{\chi}(\gamma(g'' \cdot f_1)) = \bar{\chi}(\gamma(g'' \cdot f_2)).$$

Combining (10) with (9), we see that, for $f_1, f_2 \in \text{hom}(a, b)$,

$$\delta \gamma f_1 = \delta \gamma f_2 \implies \chi((g' \cdot g'') \cdot f_1) = \chi((g' \cdot g'') \cdot f_2).$$

Thus, $g = g' \cdot g'' \in \text{hom}(b, c)$ is as required by (7). ■

The following corollary improves the estimate from Proposition 2.2. For a family Δ of endofunctors of a category C , by $\langle \Delta \rangle$ we denote the semigroup generated by Δ using composition, that is,

$$\langle \Delta \rangle = \{\bar{\delta} \mid \bar{\delta} = \delta_1 \circ \dots \circ \delta_n, \text{ for } \delta_1, \dots, \delta_n \in \Delta\}.$$

Corollary 3.2. *Let Δ be a family of endofunctors of C with each endofunctor in Δ being frank and fulfilling (P).*

(i) *For each $a, b \in \text{ob}(C)$, we have*

$$\text{rd}(a, b) \leq \min\{|\bar{\delta}(\text{hom}(a, b))| \mid \bar{\delta} \in \langle \Delta \rangle\}.$$

(ii) *For each $a \in \text{ob}(C)$,*

$$\text{rd}(a) \leq \sup_{b \in \text{ob}(C)} \min\{|\bar{\delta}(\text{hom}(a, b))| \mid \bar{\delta} \in \langle \Delta \rangle\}.$$

Proof. It is enough to check (i) as (ii) follows from (i) immediately. By Lemma 2.4 (ii), every endofunctor in $\langle \Delta \rangle$ is frank. By Theorem 3.1, each endofunctor in $\langle \Delta \rangle$ fulfills (P) at a, b . The conclusion follows from Proposition 2.2 (i). ■

3.2. Products. We define the finitely supported product of categories in a natural way. Let $C_i, i \in I$, be a family of categories. Define

$$\bigotimes_I C_i$$

as follows. Objects of this category are of the form

$$(c_i)_{i \in K},$$

where $K \subseteq I$ is finite and $c_i \in \text{ob}(C_i)$. Morphisms are of the form

$$(f_i)_{i \in K},$$

where $K \subseteq I$ is finite and $f_i \in \text{hom}(c_i, d_i)$, for $c_i, d_i \in \text{ob}(C_i)$, and we declare the above morphism to be a morphism from $(c_i)_{i \in K}$ and $(d_i)_{i \in K}$. To relax the notation, we will write

$$(c_i)_K \quad \text{and} \quad (f_i)_K$$

for the object $(c_i)_{i \in K}$ and the morphism $(f_i)_{i \in K}$, respectively.

Assume now we have two families of categories C_i and D_i with $i \in I$. Let $\delta_i: C_i \rightarrow D_i$ be a functor. Define $\otimes_I \delta_i: \otimes_I C_i \rightarrow \otimes_I D_i$ by letting

$$(\otimes_I \delta_i)((c_i)_K) = (\delta_i(c_i))_K \quad \text{and} \quad (\otimes_I \delta_i)((f_i)_K) = (\delta_i(f_i))_K.$$

It is immediate that $\otimes_{i \in I} \delta_i$ is a functor. If $C_i = C$ and $\delta_i = \delta$ for all $i \in I$, we write $\otimes_I C$ and $\otimes_I \delta$ for $\otimes_I C_i$ and $\otimes_I \delta_i$, respectively.

The following lemma is easy to check and we leave doing it to the reader.

Lemma 3.3. *If each functor $\delta_i: C_i \rightarrow D_i, i \in I$ is frank, then $\otimes_I \delta_i: \otimes_I C_i \rightarrow \otimes_I D_i$ is frank.*

The following theorem gives the transfer of property (P) from the factors to the product of categories.

Theorem 3.4. *Let $\delta_i: C_i \rightarrow D_i, i \in I$ be frank functors. Let $K \subseteq I$ be finite, and assume that $\text{hom}(a, b)$ is finite for all $a, b \in \text{ob}(C_i)$ with $i \in K$. If δ_i fulfills (P) at $a_i \in \text{ob}(C_i)$, for each $i \in K$, then $\otimes_I \delta_i$ fulfills (P) at $(a_i)_K$.*

Proof. Fix an enumeration of K , that is,

$$K = \{i_j \mid j = 1, \dots, k\}.$$

We define the following families of categories

$$E_i^0 = C_i, \quad \text{for all } i \in I,$$

and, for $p = 1, \dots, k + 1$,

$$E_i^p = \begin{cases} D_i, & i \in I \setminus K; \\ D_i, & i = i_j, \text{ for some } j < p; \\ C_i, & i = i_j, \text{ for some } j \geq p. \end{cases}$$

In particular, $E_i^1 = D_i$ for all $i \in I \setminus K$, and $E_i^1 = C_i$ for all $i \in K$, and $E_i^{k+1} = D_i$ for all $i \in I$. Similarly, define $\delta_i^p: E_i^p \rightarrow E_i^{p+1}$, for $i \in I$ and $p = 0, \dots, k$, by letting

$$\delta_i^0 = \begin{cases} \delta_i, & i \in I \setminus K; \\ \text{id}_{C_i}, & i \in K; \end{cases}$$

and, for $p \geq 1$,

$$\delta_i^p = \begin{cases} \delta_i, & i = i_p; \\ \text{id}_{D_i}, & i \in I \setminus K \text{ or } i = i_j \text{ for some } j < p; \\ \text{id}_{C_i}, & i = i_j \text{ for some } j > p. \end{cases}$$

For $p = 0, \dots, k$, consider the functor

$$\otimes_I \delta_i^p: \bigotimes_I E_i^p \rightarrow \bigotimes_I E_i^{p+1}.$$

Set $\widehat{\delta}^p = \otimes_I \delta_i^p$, and note that, by Lemma 3.3, each $\widehat{\delta}^p$ is frank and that

$$\otimes_I \delta_i = \widehat{\delta}^k \circ \dots \circ \widehat{\delta}^0.$$

Fix $a_i \in \text{ob}(C_i)$, for $i \in K$. By Theorem 3.1, it suffices to show that, for each $p \leq k$, $\widehat{\delta}^p$ fulfills (P) at

$$(11) \quad (\widehat{\delta}^{p-1} \circ \dots \circ \widehat{\delta}^0)((a_i)_K) \quad \text{and} \quad (b'_i)_{K'},$$

where $(b'_i)_{K'}$ is an arbitrary object of $\text{ob}(\bigotimes_I E_i^p)$. This is clear for $p = 0$ and, for arbitrary p , when $K' \neq K$ as in this case there are no morphisms between the two objects in (11). Now consider the case $p \geq 1$ and $K = K'$. Set

$$(a'_i)_K = (\widehat{\delta}^{p-1} \circ \dots \circ \widehat{\delta}^0)((a_i)_K),$$

and note that

$$a'_i = \begin{cases} a_i, & \text{if } i = i_j \text{ for } j \geq p; \\ \delta_i(a_i), & \text{if } i = i_j \text{ for } j < p. \end{cases}$$

Let

$$R = r^M, \quad \text{where } M = \prod_{i \in K, i \neq i_p} |\text{hom}(a'_i, b'_i)|.$$

By our assumption of finiteness of $\text{hom}(a, b)$ for $a, b \in \text{ob}(C_i)$ and frankness of δ_i in conjunction with Lemma 2.4 (iii), for all $i \in K$, we see that M , and so also R , is finite. Find $c \in C_{i_p}$ that witnesses property (P) for the functor δ_{i_p} at a_{i_p}, b_{i_p} with R colors.

We claim that the object $(c_i)_K$ of $\otimes_I C_i$ with

$$c_i = \begin{cases} b'_i, & \text{if } i \neq i_p; \\ c, & \text{if } i = i_p \end{cases}$$

witnesses property (P) for the functor $\widehat{\delta}^p$ at $(a'_i)_K$ and $(b'_i)_K$ with r colors. Indeed, let χ be an r -coloring of $\text{hom}((a'_i)_K, (c_i)_K)$. Define an R -coloring χ' of $\text{hom}(a_{i_p}, c)$ by letting $\chi'(h)$ be the sequence

$$(12) \quad (\chi((f_i)_K) \mid f_{i_p} = h, f_i \in \text{hom}(a'_i, b'_i) \text{ for } i \neq i_p).$$

By our choice of c , there is $g \in \text{hom}(b_{i_p}, c)$ such that for $f, f' \in \text{hom}(a_{i_p}, b_{i_p})$ we have

$$(13) \quad \delta_{i_p}(f) = \delta_{i_p}(f') \implies \chi'(g \cdot f) = \chi'(g \cdot f').$$

Let $(g_i)_K \in \text{hom}((b'_i)_K, (c_i)_K)$ be such that $g_{i_p} = g$ and, for $i \in K, i \neq i_p$, g_i is the identity in $\text{hom}(b'_i, c_i) = \text{hom}(b'_i, b'_i)$. It is now easy to check, from (12) and (13), that for $(f_i)_K, (f'_i)_K \in \text{hom}((a'_i)_K, (b'_i)_K)$,

$$\widehat{\delta}^p((f_i)_K) = \widehat{\delta}^p((f'_i)_K) \implies \chi((g_i)_K \cdot (f_i)_K) = \chi((g_i)_K \cdot (f'_i)_K),$$

as required. ■

3.3. Example—a frank functor for the Hales–Jewett theorem. We give now one concrete application of Theorem 3.4 that will be relevant for the proof of the Hales–Jewett theorem in Section 3.5. Recall the category P and the functor ∂_P from Section 2.4. From Lemma 2.5 and Theorem 3.4, we immediately get:

Corollary 3.5. *The functor $\otimes_{\mathbb{N}} \partial_P: \otimes_{\mathbb{N}} P \rightarrow \otimes_{\mathbb{N}} P$ is frank and fulfills (P).*

3.4. Modeling. We present a notion that allows property (P) to be transferred from one functor to another. We aim to give this notion its natural generality. Two related notions have already been proposed in the literature. One was the notion of interpretability that was defined by the author in [16, Section 6.2], the other one was the notion of

pre-adjunction defined by Mašulović in [13]. Our notion of modeling generalizes both these definitions.

Let C and D be categories. We define now the notion of cross-relatedness that will play an auxiliary, but important, role in the definition of modeling.

Definition 3.6. Let $c_1, c_2, c_3 \in \text{ob}(C)$ and $d_1, d_2, d_3 \in \text{ob}(D)$. We say that c_1, c_2, c_3 and d_1, d_2, d_3 are *cross-related* if there are functions

$$\begin{aligned}\phi &: \text{hom}(c_1, c_2) \times \text{hom}(d_2, d_3) \rightarrow \text{hom}(d_1, d_2), \\ \psi &: \text{hom}(d_2, d_3) \rightarrow \text{hom}(c_2, c_3), \\ \zeta &: \text{hom}(d_1, d_3) \rightarrow \text{hom}(c_1, c_3)\end{aligned}$$

such that, for $(f, g) \in \text{hom}(c_1, c_2) \times \text{hom}(d_2, d_3)$,

$$\zeta(g \cdot \phi(f, g)) = \psi(g) \cdot f.$$

Intuitively, one can see the notion of cross-relatedness as a way to define composition “ $g \cdot f$ ” of morphisms $f \in \text{hom}(c_1, c_2)$ by morphisms $g \in \text{hom}(d_2, d_3)$. Of course, literally, such composition does not exist as $\text{hom}(c_1, c_2)$ is computed in C while $\text{hom}(d_2, d_3)$ in D . But it can be defined in a generalized sense, in fact, in two ways. In order to do it, one stipulates that there exist functions

$$\phi: \text{hom}(c_1, c_2) \rightarrow \text{hom}(d_1, d_2) \quad \text{and} \quad \psi: \text{hom}(d_2, d_3) \rightarrow \text{hom}(c_2, c_3)$$

that allow one to compute the composition “ $g \cdot f$,” for $f \in \text{hom}(c_1, c_2)$ and $g \in \text{hom}(d_2, d_3)$, in two ways:

$$g \cdot \phi(f) \in \text{hom}(d_1, d_3) \quad \text{and} \quad \psi(g) \cdot f \in \text{hom}(c_1, c_3).$$

To relate these two results one stipulates further that there is a function

$$\zeta: \text{hom}(d_1, d_3) \rightarrow \text{hom}(c_1, c_3)$$

such that

$$\zeta(g \cdot \phi(f)) = \psi(g) \cdot f.$$

Being cross-related asserts that the above procedure can be implemented additionally allowing ϕ to depend on g .

Observe that one can formulate the definition of cross-relatedness without invoking ζ , as, to ensure that such a function ζ exists, it suffices to assume that $g \cdot \phi(g, f)$ determines $\psi(g, f) \cdot f$, that is, for all $(f, g), (f', g') \in \text{hom}(c_1, c_2) \times \text{hom}(d_2, d_3)$,

$$g \cdot \phi(f, g) = g' \cdot \phi(f', g') \implies \psi(g) \cdot f = \psi(g') \cdot f'.$$

When it is important to remember how cross-relatedness of the triples c_1, c_2, c_3 and d_1, d_2, d_3 is witnessed, we say that c_1, c_2, c_3 and d_1, d_2, d_3 are *cross-related* by (ϕ, ψ) omitting ζ from the notation for the reasons explained above.

The notion of cross-relatedness defined here is new; but in [16], a version of it with $\phi(f, g)$ depending only on f is present implicitly; in [13] another version of it is implicit, where ϕ and ψ are defined globally as functions from C to D and from D to C , respectively, and ζ is assumed to be equal to ψ .

Let $\gamma: C \rightarrow E$ and $\delta: D \rightarrow F$ be functors, and let $d_1, d_2 \in \text{ob}(D)$ and $c_1, c_2 \in \text{ob}(C)$.

Definition 3.7. We say that γ at c_1, c_2 is *modeled* by δ at d_1, d_2 if, for each $d_3 \in \text{ob}(D)$, there is $c_3 \in \text{ob}(C)$ so that c_1, c_2, c_3 and d_1, d_2, d_3 are cross-related by some (ϕ, ψ) such that

$$(14) \quad \gamma(f) = \gamma(f') \implies \delta(\phi(f, g)) = \delta(\phi(f', g)),$$

for all $f, f' \in \text{hom}(c_1, c_2)$ and $g \in \text{hom}(d_2, d_3)$.

Now, we have the main result of this section on transferring property (P) through modeling.

Theorem 3.8. *Let γ be a functor with domain C . Let $a, b \in \text{ob}(C)$. If γ at a, b is modeled by δ at d_1, d_2 with δ fulfilling (P) at d_1, d_2 , then γ fulfills (P) at a, b .*

Proof. Fix the number of colors $r \in \mathbb{N}$. Let δ be a functor with domain D such that γ at a, b is modeled by δ at $d_1, d_2 \in \text{ob}(D)$ with δ fulfilling (P) at d_1, d_2 . Let $d_3 \in \text{ob}(D)$ witness property (P) for δ at d_1, d_2 with r colors. Find $c \in \text{ob}(C)$ given for d_3 by the definition of modeling. So a, b, c and d_1, d_2, d_3 are cross-related by a pair of functions (ϕ, ψ) as in the definition of modeling.

We claim that c witnesses property (P) for γ at a, b with r colors. Let χ be an r -coloring of $\text{hom}(a, c)$. For $f \in \text{hom}(a, b)$ and $g \in \text{hom}(d_2, d_3)$, define

$$(15) \quad \chi'(g \cdot \phi(f, g)) = \chi(\psi(g) \cdot f).$$

Note that χ' is well defined since a, b, c and d_1, d_2, d_3 are cross-related by (ϕ, ψ) . The function χ' is defined on a subset of $\text{hom}(d_1, d_3)$. We extend it to $\text{hom}(d_1, d_3)$ in an arbitrary way to get an r -coloring χ' of $\text{hom}(d_1, d_3)$. Now, by our choice of d_3 , there exists $g \in \text{hom}(d_2, d_3)$ such that, for each $h, h' \in \text{hom}(d_1, d_2)$,

$$\delta h = \delta h' \implies \chi'(g \cdot h) = \chi'(g \cdot h');$$

in particular, for so chosen g , for all $f, f' \in \text{hom}(a, b)$, we have

$$\delta(\phi(f, g)) = \delta(\phi(f', g)) \implies \chi'(g \cdot \phi(f, g)) = \chi'(g \cdot \phi(f', g)).$$

By our choice of δ , whose relationship with γ is given by (14), and the definition (15) of χ' , the implication above yields, for all $f, f' \in \text{hom}(a, b)$,

$$\gamma f = \gamma f' \implies \chi(\psi(g) \cdot f) = \chi(\psi(g) \cdot f').$$

Thus, condition (P) for γ at a, b is proved. ■

To transfer bounds on Ramsey degree, only the following version of modeling is needed.

Definition 3.9. Let $c_1, c_2 \in \text{ob}(C)$ and let $d_1, d_2 \in \text{ob}(D)$. We say that c_1, c_2 is *R-modeled* by d_1, d_2 if for each $d_3 \in \text{ob}(D)$ there exists $c_3 \in \text{ob}(C)$ such that c_1, c_2, c_3 and d_1, d_2, d_3 are cross-related.

With the above definition, one proves the following proposition, which strengthens [13].

Proposition 3.10. *Let C, D be categories, and let $a, b \in \text{ob}(C)$ and $d_1, d_2 \in \text{ob}(D)$. If a, b is R-modeled by d_1, d_2 , then*

$$\text{rd}(a, b) \leq \text{rd}(d_1, d_2).$$

Proof. This proof is parallel to the proof of Theorem 3.8. Let $k = \text{rd}(d_1, d_2)$. We can assume that $k < \infty$. We show that $\text{rd}(a, b) \leq k$. Fix $r \in \mathbb{N}$. Let now $d_3 \in \text{ob}(D)$ witness $\text{rd}(d_1, d_2) \leq k$ for r -colorings. Let $c \in \text{ob}(C)$ be provided from the definition of R-modeling so that a, b, c and d_1, d_2, d_3 are cross-related by (ϕ, ψ) .

We claim that c chosen above witnesses $\text{rd}(a, b) \leq k$ for r -colorings. To check this claim, let now χ be an r -coloring of $\text{hom}(a, c)$. Define a coloring χ' of $\text{hom}(d_1, d_3)$ exactly as is done around formula (15) in the proof of Theorem 3.8. Now by our choice of k , there is $g \in \text{hom}(d_2, d_3)$ such that χ' attains at most k colors on the set

$$\{g \cdot f \mid f \in \text{hom}(d_1, d_2)\}.$$

By the definition of χ' , we see that for so chosen g , χ attains at most k colors on the set

$$\{\psi(g) \cdot f \mid f \in \text{hom}(a, b)\},$$

as required. ■

3.5. Example—the Hales–Jewett theorem. We fix $k_0 \in \mathbb{N}$. Below, for a function f , $\text{im}(f)$ will stand for the set of all values of f .

We define a category HJ_{k_0} and its endofunctor ∂_{k_0} , which will be used to prove the Hales–Jewett theorem.

Objects of HJ_{k_0} are as follows:

- natural numbers $l \in \mathbb{N}$;
- surjections $v: [-k_0, 0] \rightarrow [k]$, for some $k \in \mathbb{N}$.

In the remainder of this section, l , possibly with subscripts, will stand for objects of HJ_{k_0} of the first kind above and v , possibly with subscripts, will stand for objects of the second kind.

Morphisms of HJ_{k_0} will be appropriate functions. We describe them as follows:

- $f \in \text{hom}(v, l)$ is a function $f: [l] \rightarrow \text{im}(v)$;
- $g \in \text{hom}(l_1, l_2)$ is a function $g: [l_2] \rightarrow [-k_0, l_1]$ such that $\text{im}(g) \supseteq [l_1]$;
- there are no other morphisms except for the identities.

In order to define composition of morphisms in HJ_{k_0} , we need to introduce a new piece of notation. For two functions h and h' domains are disjoint intervals I and I' of \mathbb{Z} , respectively, let

$$h \frown h'$$

stand for the function whose domain is $I \cup I'$ and whose restrictions to I and I' are h and h' , respectively. Now composition in HJ_{k_0} is defined as follows:

- for $f \in \text{hom}(v, l_1)$ and $g \in \text{hom}(l_1, l_2)$, let

$$g \cdot f = (v \frown f) \circ g;$$

- for $g_1 \in \text{hom}(l_1, l_2)$ and $g_2 \in \text{hom}(l_2, l_3)$, let

$$g_2 \cdot g_1 = ((\text{id}_{[-k_0, 0]}) \frown g_1) \circ g_2.$$

Note that $v \frown f$ and $(\text{id}_{[-k_0, 0]}) \frown g_1$, whose domains are $[-k_0, l_1]$ and $[-k_0, l_2]$, respectively.

Before we define the functor $\partial_{k_0}: \text{HJ}_{k_0} \rightarrow \text{HJ}_{k_0}$, we need to modify (3). For $n \in \mathbb{N}$, $n \geq 1$, let

$$n \dot{-} 1 = \max(n - 1, 1).$$

The functor $\partial_{k_0}: \text{HJ}_{k_0} \rightarrow \text{HJ}_{k_0}$ is now defined on objects by:

- $\partial_{k_0}(l) = l$;
- $\partial_{k_0}(v) = \min(v, \max(\text{im}(v)) \dot{-} 1)$;

and on morphisms by:

- $\partial_{k_0}(g) = g$, for $g \in \text{hom}(l_1, l_2)$;
- $\partial_{k_0}(f) = \min(f, \max(\text{im}(v)) \dot{-} 1)$, for $f \in \text{hom}(v, l)$.

Lemma 3.11. *The following statements hold:*

- (i) ∂_{k_0} is a functor and it is frank.
(ii) The functor $\partial_{k_0}: \text{HJ}_{k_0} \rightarrow \text{HJ}_{k_0}$ fulfills (P).

Proof. (i) is done by an easy check that we leave to the reader.

We now prove (ii). Note that it is clear that ∂_{k_0} fulfills (P) at pairs of objects of the form (l_1, l_2) , (v_1, v_2) , and (l, v) . Indeed, the morphism sets between objects in each of these pairs are empty with the exception of l_1, l_2 with $l_1 \leq l_2$, in which case, δ_{k_0} is equal to the identity map on $\text{hom}(l_1, l_2)$.

It remains to check that ∂_{k_0} fulfills (P) at pairs of the form $v, l \in \text{ob}(\text{HJ}_{k_0})$. This goal will be achieved by showing that ∂_{k_0} at v, l is modeled by $\otimes_{\mathbb{N}} \partial$ at a pair of objects of $\otimes_{\mathbb{N}} P$ that we will choose below, and using Theorem 3.8 and Corollary 3.5. Let $k_1 \in \mathbb{N}$ be such that $\text{im}(v) = [k_1]$. So, at this point, we have fixed v, l, k_1 , and, of course, k_0 .

We define the two objects of $\otimes_{\mathbb{N}} P$ that will be used to model ∂_{k_0} at v, l . Let $a = (a_i)_{i=1}^l$ and $b = (b_i)_{i=1}^l$, where $a_i, b_i \in \text{ob}(P)$, for $1 \leq i \leq l$, be defined by

$$a_i = (k_1, 1) \quad \text{and} \quad b_i = (3, 2).$$

So a, b are objects in $\otimes_{\mathbb{N}} P$. To see that ∂_{k_0} at v, l is modeled by $\otimes_{\mathbb{N}} \partial_P$ at a, b , we fix an arbitrary object $c \in \text{ob}(\otimes_{\mathbb{N}} P)$. We can assume that $c = (c_i)_{i=1}^l$ for some $c_i \in \text{ob}(P)$ with $c_i = (m_i, 2)$ for some $m_i \in \mathbb{N}$, $1 \leq i \leq l$. We need to find an object l' in HJ_{k_0} and functions

$$\phi: \text{hom}(v, l) \rightarrow \text{hom}(a, b) \quad \text{and} \quad \psi: \text{hom}(b, c) \rightarrow \text{hom}(l, l')$$

such that

$$(16) \quad p \cdot \phi(f) = p' \cdot \phi(f') \implies \psi(p) \cdot f = \psi(p') \cdot f',$$

for all $p, p' \in \text{hom}(b, c)$, $f, f' \in \text{hom}(v, l)$.

Translating (FP) to the situation dealt with here, the function ϕ should be defined on $\text{hom}(v, l) \times \text{hom}(b, c)$; but our ϕ will not depend on the second coordinate.

We can define ϕ right away. For $f \in \text{hom}(v, l)$, set

$$\phi(f) = (\phi_1(f), \dots, \phi_l(f)),$$

where, for each $1 \leq i \leq l$, $\phi_i(f): [3] \rightarrow [k_1]$ is defined by letting

$$(\phi_i(f))(j) = \begin{cases} k_1, & \text{if } j = 1; \\ f(i), & \text{if } j = 2; \\ \max(1, k_1 - 1), & \text{if } j = 3. \end{cases}$$

Note that $\phi_i(f) \in \text{hom}(a_i, b_i)$, and, therefore, $\phi(f) \in \text{hom}(a, b)$.

With the definition of ϕ in hand, we state (16) in more basic terms. Note that each $p \in \text{hom}(b, c)$ is of the form

$$(17) \quad p = (p_1, \dots, p_l),$$

where p_i is in $\text{hom}(a_i, b_i)$, that is, it is a non-decreasing surjections such that $\text{im}(p_i) = [3]$ and $\text{dom}(p_i) = [m_i]$, for each $1 \leq i \leq l$. Similarly, we represent $p' \in \text{hom}(b, c)$ as (p'_1, \dots, p'_l) . Now, (16) becomes

$$(18) \quad (\phi_i(f) \circ p_i = \phi_i(f') \circ p'_i, \text{ for } 1 \leq i \leq l) \implies f \circ \psi(p) = f' \circ \psi(p')$$

for all $p, p' \in \text{hom}(b, c)$, $f, f' \in \text{hom}(v, l)$.

It remains to define l' and ψ for which (18) holds. Set

$$l' = m_1 + \dots + m_l.$$

For p as in (17), define $g = \psi(p) \in \text{hom}(l, l')$ as follows. If $j \in [m_1 + \dots + m_l]$, let i be the unique natural number such that

$$m_1 + \dots + m_{i-1} < j \leq m_1 + \dots + m_i.$$

Then if $p_i(j - (m_1 + \dots + m_{i-1})) = 2$, let

$$g(j) = i,$$

otherwise, let $g(j)$ be a number in $[-k_0, 0]$ such that

$$v(g(j)) = k_1, \text{ if } p_i(j - (m_1 + \dots + m_{i-1})) = 1,$$

and

$$v(g(j)) = \max(1, k_1 - 1), \text{ if } p_i(j - (m_1 + \dots + m_{i-1})) = 3.$$

With the definitions above, it is easy to check that, for $p \in \text{hom}(b, c)$ represented as in (17) and for $f \in \text{hom}(v, l)$, we have

$$v \wedge (\phi_1(f) \circ p_1) \wedge \dots \wedge (\phi_l(f) \circ p_l) = f \circ \psi(p),$$

from which (18) follows immediately.

To finish the proof of (FP), it remains to show that for $f, f' \in \text{hom}(v, l)$,

$$\partial_{k_0} f = \partial_{k_0} f' \implies \left(\bigotimes_{\mathbb{N}} \partial_P \right) (\phi(f)) = \left(\bigotimes_{\mathbb{N}} \partial_P \right) (\phi(f')),$$

which amount to proving

$$\partial_{k_0} f = \partial_{k_0} f' \implies \partial_P(\phi_i(f)) = \partial_P(\phi_i(f')), \text{ for } i = 1, \dots, l. \quad \blacksquare$$

The following is the Hales–Jewett theorem, see [7] and [14].

Corollary 3.12. *For each $k, l \in \mathbb{N}$ and $r \in \mathbb{N}$, there exists $m \in \mathbb{N}$ such that for each r -coloring of all functions from $[m]$ to $[-k, 0]$, there is $g: [m] \rightarrow [-k, l]$, with $\text{im}(g) \supseteq [l]$, such that the set*

$$\{f \circ g \mid f: [-k, l] \rightarrow [-k, 0], f \upharpoonright [-k, 0] = \text{id}_{[-k, 0]}\}$$

is monochromatic.

Proof. Fix $k, l \in \mathbb{N}$. Consider the category HJ_k and the objects id_k and l in it. Note that the conclusion of the corollary follows from $\text{rd}(\text{id}_k, l) = 1$. To prove this equality, observe that the set

$$(\partial_k \circ \cdots \circ \partial_k)(\text{hom}(\text{id}_k, l)),$$

where ∂_k is composed $k - 1$ times, has one element. By Lemma 3.11 combined with Corollary 3.2 (i) for $\Delta = \{\partial_k\}$, the equality $\text{rd}(\text{id}_k, l) = 1$ follows immediately. ■

With some additional routine work, the methods used to prove Corollary 3.12 can be adapted to proving more general versions of the Hales–Jewett theorem as in [16, Section 8.1] or [19, Lemma 3.3]. In these generalizations, one obtains concrete Ramsey statements, in which $\text{rd}(a, b)$ may be strictly bigger than 1.

4. Proving condition (P)

The goal of this section is to formulate a local version of condition (P) and prove that, in most circumstances, it implies (P).

4.1. Condition (FP). We state the local version, we call (FP), of the pigeonhole principle (P). In applications, it is often easier to check directly (FP) than (P). Let $\delta: C \rightarrow D$ be a functor. Recall the statement of (P) for a functor δ from Definition 2.1. Note that the property of $g \in \text{hom}(b, c)$ in condition (P) can be rephrased as follows:

For each $f' \in \delta(\text{hom}(a, b))$, $\chi(g \cdot f)$ is constant for $f \in \text{hom}(a, b)$ with $\delta f = f'$.

Above, g is chosen first and independently of f' . This feature is relaxed when passing to (FP) from (P), namely, in (FP) it suffices to find g that depends on f' . The price of this relaxation is included in the second point of condition (FP). It has to do with controlling the behavior of δg in a suitable way.

It will be convenient to introduce the following piece of notation. Let C be a category C and δ a functor defined on C . For $a, b \in \text{ob}(C)$ and $h \in \delta(\text{hom}(a, b))$, let

$$\text{hom}(a, b)_h = \{f \in \text{hom}(a, b) \mid \delta f = h\}.$$

Definition 4.1. We say that δ fulfills (FP) at $a, b \in \text{ob}(C)$, if for $r \in \mathbb{N}$ and a finite non-empty set $s \subseteq \delta(\text{hom}(a, b))$ the following condition holds:

There exist $c \in \text{ob}(C)$, $f' \in s$, and $g' \in \delta(\text{hom}(b, c))$ such that for each r -coloring χ of $\text{hom}(a, c)$, there exists $g \in \text{hom}(b, c)$ with

- $g \cdot \text{hom}(a, b)_{f'}$ χ -monochromatic; and
- $(\delta g) \cdot e = g' \cdot e$, for each $e \in s$.

As with condition (P), we say that δ fulfills (FP) at $a \in \text{ob}(C)$, if it fulfills (FP) at a, b for all $b \in \text{ob}(C)$; and we simply say that δ fulfills (FP), if it fulfills (FP) at a, b for all $a, b \in \text{ob}(C)$.

4.2. Example—a frank functor with (FP) for Ramsey’s theorem. Recall Example 3 from Section 2.4. We check condition (FP) for ∂_R from this example.

Lemma 4.2. $\partial_R: R \rightarrow R$ fulfills (FP).

Proof. This proof amounts to an application of the standard pigeonhole principle. Fix $r \in \mathbb{N}$. Let k, l be two objects in R , and let $\emptyset \neq s \subseteq \partial_R(\text{hom}(k, l))$. To avoid trivial cases, we can assume that $\text{hom}(k, l)$ has at least two elements and that $k \geq 1$, therefore, $1 \leq k < l$.

For the two objects k and l and the number of colors r , we need to find, in the notation of (FP), and object c and two morphisms f' and g' . First, we define the object by letting

$$m = (r + 1)l \in \text{ob}(R).$$

Next, we define the two objects. Pick $(x', l - 1) \in \partial_R(\text{hom}(k, l))$ so that

$$(19) \quad (x', l - 1) \in s \quad \text{and} \quad \max x' = \max \{ \max x'' \mid (x'', l - 1) \in s \},$$

and let

$$(20) \quad (y', m - 1) = ([l - 1], m - 1) \in \partial_R(\text{hom}(l, m)).$$

By convention, if $k = 1$, we interpret the above definition to give $x' = \emptyset$. We claim that this choice of the object m and the morphisms $(x', l - 1)$ and $(y', m - 1)$ ensures that (FP) are satisfied.

To prove this claim, let χ be an r -coloring of $\text{hom}(k, m)$. For $i \in [l, m]$, set

$$\mathbf{x}_i = (x' \cup \{i\}, m) \in \text{hom}(k, m),$$

and consider the r -coloring of $[l, m]$ given by

$$(21) \quad [l, m] \ni i \rightarrow \chi(\mathbf{x}_i).$$

Set

$$p = \max x' \leq l - 1,$$

with $p = 0$, if $x' = \emptyset$, by convention. Note that, by the choice of m , there is a subset I of $[l, m]$ of size $l - p$ on which the r -coloring (21) is constant, which means that χ is constant on \mathbf{x}_i as i varies over I . Define

$$\mathbf{y} = ([p] \cup I, m) \in \text{hom}(l, m).$$

This is the morphism g in the notation from (FP). We need to check that \mathbf{y} satisfies the two points displayed in (FP).

For $(x, l) \in \text{hom}(k, l)$ with $\partial_R(x, l) = (x', l - 1)$, we have

$$\mathbf{y} \cdot (x, l) = \mathbf{x}_i, \quad \text{for some } i \in I.$$

Therefore, $\chi(\mathbf{y} \cdot (x, l))$ is constant for $(x, l) \in \text{hom}(k, l)$ with $\partial_R(x, l) = (x', l - 1)$, and the first point in (FP) is checked for \mathbf{y} . To see the second point, note that, for each $(x'', l - 1) \in s$, we have

$$\partial_{R\mathbf{y}} \cdot (x'', l - 1) = (x'', m - 1) = (y', m - 1) \cdot (x'', l - 1),$$

using (19) to get the first equality and (19) and (20) to get the second one. Thus, (FP) follows. \blacksquare

4.3. Condition (FP) implies (P). The following theorem is the main result of Section 4. It shows that under mild assumptions the local condition (FP) implies (P). In concrete situations, (FP) is usually much easier to check than (P).

Theorem 4.3. *Let $\delta: C \rightarrow D$ be a functor, and let $a \in \text{ob}(C)$. If δ fulfills (FP) at a , then δ fulfills (P) at a, b for each $b \in \text{ob}(C)$ with $\delta(\text{hom}(a, b))$ finite.*

Proof. Fix a functor δ and an object a . In order to prove that δ fulfills (P) at a, b , we fix $r > 0$ and $b \in \text{ob}(C)$ with finite $\delta(\text{hom}(a, b))$. Set $n = |\delta(\text{hom}(a, b))|$. By recursion, we construct

- $c_k \in \text{ob}(C)$, for $0 \leq k \leq n$;
- $g'_k \in \delta(\text{hom}(c_{k-1}, c_k))$, for $1 \leq k \leq n$;
- $f'_k \in \delta(\text{hom}(a, b))$, for $1 \leq k \leq n$.

Note that we enumerate the c_k -s starting with $k = 0$ and the g'_k -s and f'_k -s starting with $k = 1$. These objects will have the following properties for $0 \leq k \leq n$, where we note that the first point in (c) makes sense as, by the conditions above, we have that $g'_{k-1} \cdots g'_1 \cdot f'_k \in \delta(\text{hom}(a, c_{k-1}))$.

- (a) $c_0 = b$;
- (b) $f'_k \neq f'_i$ for all $i < k$;
- (c) for each r -coloring χ of $\text{hom}(a, c_k)$, there exists $g \in \text{hom}(c_{k-1}, c_k)$ such that
- χ is constant on $g \cdot (\text{hom}(a, c_{k-1})_{g'_{k-1} \cdots g'_1 \cdot f'_k})$;
 - $(\delta g) \cdot g'_{k-1} \cdots g'_1 \cdot f' = g'_k \cdot g'_{k-1} \cdots g'_1 \cdot f'$ for all $f' \in \delta(\text{hom}(a, b))$ with $f' \notin \{f_i \mid i \leq k\}$.

To start the construction, we set $c_0 = b$. The conditions above for $k = 0$ hold with (b) and (c) being vacuously true. Assume that $0 < k \leq n$ and the construction has been carried out up to stage $k - 1$. Consider the finite set

$$s = \{g'_{k-1} \cdots g'_1 \cdot f' \mid f' \in \delta(\text{hom}(a, b)) \setminus \{f'_i \mid i < k\}\}.$$

By our choice of g'_i for $1 \leq i < k$ and the assumption that $k \leq n$, we have that

$$\emptyset \neq s \subseteq \delta(\text{hom}(a, c_{k-1})).$$

Now condition (FP) applied to a, c_{k-1} , and the set s above allows us to pick $c_k \in \text{ob}(C)$, $f'_k \in \delta(\text{hom}(a, b))$, and $g'_k \in \delta(\text{hom}(c_{k-1}, c_k))$ so that conditions (b) and (c) hold. The construction has been carried out.

Observe that by the choice of n and condition (b), we have

$$(22) \quad \delta(\text{hom}(a, b)) = \{f'_k \mid 1 \leq k \leq n\}.$$

We claim that $c = c_n$ witnesses that δ fulfills (P) at a, b with r colors; that is, for each r -coloring χ of $\text{hom}(a, c)$, there is $g \in \text{hom}(b, c)$ such that, for $h_1, h_2 \in \text{hom}(a, b)$, we have

$$(23) \quad \delta h_1 = \delta h_2 \implies \chi(g \cdot h_1) = \chi(g \cdot h_2).$$

In order to prove the statement above, fix an r -coloring χ of $\text{hom}(a, c)$. We recursively produce

$$g_n \in \text{hom}(c_{n-1}, c_n), \dots, g_1 \in \text{hom}(c_0, c_1)$$

starting with g_n and ending with g_1 as follows. Having produced g_n, \dots, g_{k+1} , we consider the r -coloring of $\text{hom}(a, c_k)$ given by

$$\text{hom}(a, c_k) \ni f \rightarrow \chi(g_n \cdots g_{k+1} \cdot f).$$

By (c), we get $g_k \in \text{hom}(c_{k-1}, c_k)$ such that

$$(24) \quad \chi(g_n \cdots g_{k+1} \cdot g_k \cdot f) \text{ is constant for } f \in \text{hom}(a, c_{k-1})_{g'_{k-1} \cdots g'_1 \cdot f'_k}$$

and

$$(25) \quad (\delta g_k) \cdot g'_{k-1} \cdots g'_1 \cdot f'_j = g'_k \cdot g'_{k-1} \cdots g'_1 \cdot f'_j, \quad \text{for } j > k.$$

Now we show that

$$g = g_n \cdots g_1 \in \text{hom}(b, c)$$

witnesses that the implication in (23) holds. Let $h_1, h_2 \in \text{hom}(a, b)$ be such that $\delta h_1 = \delta h_2$. This common value can be taken to be f'_k for some $1 \leq k \leq n$ by (22). For $i = 1, 2$, an iterative application of condition (25) gives

$$\begin{aligned} g'_{k-1} \cdot g'_{k-2} \cdots g'_1 \cdot f'_k &= \delta g_{k-1} \cdot \delta g_{k-2} \cdots \delta g_1 \cdot \delta h_i \\ &= \delta(g_{k-1} \cdot g_{k-2} \cdots g_1 \cdot h_i), \end{aligned}$$

and so

$$g_{k-1} \cdot g_{k-2} \cdots g_1 \cdot h_i \in (\text{hom}(a, c_{k-1}))_{g'_{k-1} \cdots g'_1 \cdot f'_k},$$

which in light of (24) implies that

$$\chi(g_n \cdots g_k \cdot (g_{k-1} \cdots g_1 \cdot h_1)) = \chi(g_n \cdots g_k \cdot (g_{k-1} \cdots g_1 \cdot h_2)).$$

Thus, (23) is proved. ■

The following corollary follows immediately from Corollary 3.2 (i) and Theorem 4.3.

Corollary 4.4. *Let Δ be a family of frank endofunctors of C . Let $a \in \text{ob}(C)$. Assume that each $\delta \in \Delta$ fulfills (FP) at a and $\text{hom}(a, b)$ is finite for all $b \in \text{ob}(C)$. Then, for all $b \in \text{ob}(C)$,*

$$\text{rd}(a, b) \leq \min\{|\bar{\delta}(\text{hom}(a, b))| \mid \bar{\delta} \in \langle \Delta \rangle\}.$$

4.4. Example—Ramsey's theorem and its product. As an illustration, we derive now the classical Ramsey theorem and the product Ramsey theorem from the general results established earlier.

Corollary 4.5. *Given $k, l \in \mathbb{N}$, for each $r \in \mathbb{N}$, there exists $m \in \mathbb{N}$ such that, for each r -coloring of all k -element subsets of $[m]$, there exists $b \subseteq [m]$ of size l such that all k -element subsets of b get the same color.*

Proof. Since, by Lemma 4.2, ∂_R fulfills (FP), and $\text{hom}(k, l)$ is finite, for all $k, l \in \text{ob}(R)$, it follows from that $\Delta = \{\partial_R\}$ satisfies all the assumptions of Corollary 4.4. Thus, we get the conclusion after noticing that the set

$$(\partial_R \circ \cdots \circ \partial_R)(\text{hom}(k, l))$$

has only one element (its only element is the empty set), where ∂_R is composed k times. ■

Corollary 4.6. *Let $r \in \mathbb{N}$ and let k_1, \dots, k_l and p_1, \dots, p_l be natural numbers. There exist natural numbers q_1, \dots, q_l such that for each r -coloring of the set*

$$\{(a_1, \dots, a_l) \mid a_i \subseteq [q_i], |a_i| = k_i, \text{ for } i \leq l\}$$

there exist $b_1 \subseteq [q_1], \dots, b_l \subseteq [q_l]$ with $|b_i| = p_i$, for each $i \leq l$, and such that the set

$$\{(a_1, \dots, a_l) \mid a_i \subseteq b_i, |a_i| = k_i, \text{ for } i \leq l\}$$

is monochromatic.

Proof. By Lemma 4.2, ∂_R fulfills (FR). Since $\text{hom}(k, l)$ is finite, for all $k, l \in \text{ob}(R)$, Theorem 4.3 implies that ∂_R fulfills (P). Thus, by Theorem 3.4, $\otimes_{\mathbb{N}} \partial_R: \otimes_{\mathbb{N}} R \rightarrow \otimes_{\mathbb{N}} R$ fulfills (P). Now, the conclusion follows from Corollary 3.2 (i). ■

4.5. Example—Fouché’s Ramsey theorem for trees. We present here one more elaborate example of using the general theory. We derive from it Fouché’s Ramsey theorem for trees as proved in [3].

First, we collect basic definitions concerning trees and a type of morphism between them. By a *tree* we understand a *finite, non-empty* partial order such that each two elements have a common predecessor and the set of predecessors of each element is linearly ordered. A *leaf* is a maximal element of a tree. By convention, we regard every node of a tree as one of its own predecessors and as one of its own successors.

Each tree T carries a binary function \wedge_T that assigns to each $v, w \in T$ the largest element $v \wedge_T w$ of T that is a predecessor of both v and w . For a tree T and $v \in T$, let

$$\text{im}_T(v)$$

be the set of all *immediate successors* of v , and we do not regard v as one of them. Let

$$\text{ht}_T(v)$$

be the cardinality of the set of all predecessors of v (including v), and let

$$\text{ht}(T) = \max\{\text{ht}_T(v) : v \in T\}.$$

For a tree T , let

$$\text{br}(T)$$

be the maximum of cardinalities of $\text{im}_T(v)$ for $v \in T$.

A tree T is called *ordered* if for each $v \in T$ there is a fixed linear order of $\text{im}_T(v)$. Such an assignment allows us to define the lexicographic linear order \leq_T on all the nodes of T by specifying that $v \leq_T w$ precisely when

- v is a predecessor of w ; or
- v is not a predecessor of w , and w is not a predecessor of v , and the predecessor of v in $\text{im}_{\mathcal{T}}(v \wedge w)$ is less than or equal to the predecessor of w in $\text{im}_{\mathcal{T}}(v \wedge w)$ in the given order on $\text{im}_{\mathcal{T}}(v \wedge w)$.

A *height-preserving embedding* f from an ordered tree S to an ordered tree T is an injective function $f: S \rightarrow T$ such that

- f is order preserving between \leq_S and \leq_T ;
- $f(v \wedge_S w) = f(v) \wedge_T f(w)$, for $v, w \in S$; and
- $\text{ht}_S(v) = \text{ht}_T(f(v))$, for $v \in S$.

Note that preservation of order by f is equivalent to saying that for every $v \in S$ and all $w_1, w_2 \in \text{im}_S(v)$ with $w_1 \leq_S w_2$, we have $f(w_1) \leq_T f(w_2)$ in $\text{im}_T(f(v))$.

Theorem 4.7 (Fouché [3]). *Let $r \in \mathbb{N}$ and let S and T be ordered trees. There is an ordered tree V such that $\text{ht}(V) = \text{ht}(T)$ and for each r -coloring of all height-preserving embeddings from S to V there is a height-preserving embedding $g: T \rightarrow V$ such that the set*

$$\{g \circ f: f \text{ a height-preserving embedding of } S \text{ to } T\}$$

is monochromatic.

We define a category and an endofunctor on it that are appropriate for the theorem above. Consider the category \mathcal{T} whose objects are ordered trees. Given $S, T \in \text{ob}(\mathcal{T})$, with $\text{ht}(S) = \text{ht}(T)$, $\text{hom}(S, T)$ consists of all height-preserving embeddings from S to T . There are no other morphisms in \mathcal{T} ; in particular, if $\text{ht}(S) \neq \text{ht}(T)$, then $\text{hom}(S, T) = \emptyset$.

We now define a functor $\partial^*: \mathcal{T} \rightarrow \mathcal{T}$. Given $T \in \text{ob}(\mathcal{T})$, put

$$\partial^*T = \begin{cases} \{v \in T: \text{ht}(v) < \text{ht}(T)\}, & \text{if } \text{ht}(T) > 1; \\ T, & \text{if } \text{ht}(T) = 1. \end{cases}$$

We will write T^* for ∂^*T . Now, define the functor ∂^* on morphisms of \mathcal{T} by letting, for $f: S \rightarrow T$,

$$\partial^*f = f \upharpoonright S^*.$$

Lemma 4.8. *∂^* is a frank functor.*

Proof. It is clear that ∂^* is a functor as for morphisms $f: S \rightarrow T$ and $g: T \rightarrow V$ we have $f(S^*) \subseteq T^*$ and hence

$$\partial^*(g \circ f) = (g \circ f) \upharpoonright S^* = g \circ (f \upharpoonright S^*) = (g \upharpoonright T^*) \circ (f \upharpoonright S^*) = (\partial^*g) \circ (\partial^*f).$$

Now, to check that ∂^* is frank, we fix two objects of \mathcal{T} , that is, two ordered trees S and T' . We need to find $T \in \text{ob}(\mathcal{T})$ such that $T^* = T'$ and $\partial^*(\text{hom}(S, T)) = \text{hom}(S^*, T^*)$. If $\text{ht}(S) \neq \text{ht}(T') + 1$, then any $T \in \mathcal{T}$ with $T^* = T'$ works, since then $\text{hom}(S^*, T^*) = \emptyset$ and $\text{hom}(S, T) = \emptyset$. So we assume that $\text{ht}(S) = \text{ht}(T') + 1$. We need to find $T \in \text{ob}(\mathcal{T})$ such that

- $T^* = T'$; and
- for each height-preserving embedding $f': S^* \rightarrow T^*$, there is a height-preserving embedding $f: S \rightarrow T$ such that $f' = f \upharpoonright S^*$.

One defines T so that $\text{ht}(T) = \text{ht}(S)$, $T^* = T'$, and, for each leaf w of T' ,

$$|\text{im}_T(w)| = \text{br}(S).$$

One then linearly orders T by extending the linear order on T' in an arbitrary way as long as the resulting order makes T into an ordered tree. It is then clear that each height-preserving embedding $f': S^* \rightarrow T'$ extends to a height-preserving embedding $f: S \rightarrow T$ by mapping elements of $\text{im}_S(v)$, for each leaf v of S^* , to $\text{im}_T(f'(v))$ in an injective and order-preserving fashion. ■

Proof of Theorem 4.7. To obtain the conclusion of the theorem, one needs to check that $\text{rd}(S, T) \leq 1$ for all $S, T \in \text{ob}(\mathcal{T})$. Note that the set

$$(\partial^* \circ \dots \circ \partial^*)(\text{hom}(S, T))$$

has at most one element, where ∂^* is composed $\text{ht}(S) - 1$ many times. Indeed, if $\text{ht}(S) \neq \text{ht}(T)$, then $\text{hom}(S, T) = \emptyset$; if $\text{ht}(S) = \text{ht}(T)$, then this set contains only the unique function from a one-node tree to a one-node tree. Thus, by Lemma 4.8 and Corollary 4.4, it will suffice to check that ∂^* fulfills (FP) at each pair of objects of \mathcal{T} . So fix $r \in \mathbb{N}$, $S, T \in \text{ob}(\mathcal{T})$, and $\emptyset \neq s \subseteq \text{hom}(S^*, T^*)$. Non-emptiness of s implies that $\text{ht}(S) = \text{ht}(T)$; we call this common height h . We need to produce $V \in \text{ob}(\mathcal{T})$ with $\text{ht}(V) = h$, and $f' \in s$ and $g' \in \text{hom}(T^*, V^*)$ so that the conclusion of (FP) holds for these choices.

We let $f' \in s$ be arbitrary and g' be equal to the identity map on T^* . The tree V will be chosen so that $V^* = T^*$. It suffices to specify, for each leaf w of T^* with $\text{ht}_{T^*}(w) = h - 1$, the number of elements in $\text{im}_V(w)$. If w is not of the form $f'(v)$ for a leaf v of S^* , let $\text{im}_V(w)$ be empty. Now, let v_1, \dots, v_l list all the leaves of S^* of height $h - 1$; since $\text{ht}(S) = h$ such leaves exist. Put

$$K_i = \text{im}_S(v_i) \quad \text{and} \quad k_i = |K_i|, \quad \text{for } 1 \leq i \leq l,$$

and also

$$P_i = \text{im}_T(f'(v_i)) \quad \text{and} \quad p_i = |P_i|, \quad \text{for } 1 \leq i \leq l.$$

Let the natural numbers q_i , for $1 \leq i \leq l$, be gotten from Corollary 4.6 for the sequences (k_i) and (p_i) , and the number of colors r . Let $Q_i = \text{im}_V(f'(v_i))$ have size q_i . This procedure defines V .

Now, it is enough to do the following: for an r -coloring χ of $\text{hom}(S, V)$, find a height-preserving embedding $g: T \rightarrow V$ such that

- (i) $\chi(g \circ f)$ is constant on the set of height-preserving embeddings $f: S \rightarrow T$ with $f \upharpoonright S^* = f'$;
- (ii) $g \upharpoonright T^* = \text{id}_{T^*}$.

We fix χ as above. For a tuple (a_i) such that $a_i \subseteq Q_i$ and $|a_i| = k_i$, for each $1 \leq i \leq l$, define $f_{(a_i)}: S \rightarrow V$ by letting

$$\begin{aligned} f_{(a_i)} \upharpoonright S^* &= f'; \\ f_{(a_i)} \upharpoonright K_i &: K_i \rightarrow a_i \text{ the unique order-preserving function.} \end{aligned}$$

Note that $f_{(a_i)}$ is a height-preserving embedding. Further note that

$$\begin{aligned} (26) \quad (f \in \text{hom}(S, T), f \upharpoonright S^* = f' \text{ and } g \in \text{hom}(T, V), g \upharpoonright T^* = \text{id}_{T^*}) \\ \implies (g \circ f = f_{(a_i)}, \text{ where } a_i = g(f(K_i))). \end{aligned}$$

Color tuples (a_i) such that $a_i \subseteq Q_i$ and $|a_i| = k_i$ by letting

$$\chi'((a_i)) = \chi(f_{(a_i)}).$$

By our choice of (q_i) , there are sets $b_i \subseteq Q_i$ with $|b_i| = p_i$ and such that all tuples (a_i) with $a_i \subseteq b_i$ get the same color with respect to χ' . Let $g: T \rightarrow V$ be defined by letting

$$\begin{aligned} g \upharpoonright T^* &= \text{id}_{T^*}; \\ g \upharpoonright Q_i &: Q_i \rightarrow b_i \text{ the unique order-preserving function.} \end{aligned}$$

By (26), for each $f: S \rightarrow T$ with $f \upharpoonright S^* = f'$, we have that

$$g \circ f = f_{(a_i)},$$

for some (a_i) with $a_i \subseteq b_i$ and $|a_i| = k_i$. Thus, $\chi(g \circ f)$ is constant, as required by (i). Obviously, g fulfills (ii). ■

Acknowledgments. I would like to thank Sebastian Junge, whose remarks improved the presentation of the material in this paper.

Funding. Research supported by NSF grant DMS-1954069.

References

- [1] S. A. ARGYROS and S. TODORCEVIC, *Ramsey methods in analysis*. Adv. Courses Math. CRM Barcelona, Birkhäuser, Basel, 2005. MR 2145246
- [2] T. J. CARLSON, [Some unifying principles in Ramsey theory](#). *Discrete Math.* **68** (1988), no. 2–3, 117–169. Zbl 0817.04002 MR 926120
- [3] W. L. FOUCHÉ, [Symmetries and Ramsey properties of trees](#). *Discrete Math.* **197–198** (1999), 325–330. Zbl 0945.05045 MR 1674871
- [4] R. L. GRAHAM, K. LEEB and B. L. ROTHSCHILD, [Ramsey’s theorem for a class of categories](#). *Adv. Math.* **8** (1972), no. 3, 417–433. Zbl 0243.18011 MR 306010
- [5] R. L. GRAHAM and B. L. ROTHSCHILD, [Ramsey’s theorem for \$n\$ -parameter sets](#). *Trans. Amer. Math. Soc.* **159** (1971), 257–292. Zbl 0233.05003 MR 284352
- [6] M. L. GROMOV, [Colorful categories](#). (in Russian) *Uspekhi Mat. Nauk* **70** (2015), no. 4 (424), 3–76; [English translation: Russian Math. Surveys](#) **70** (2015), no. 4, 591–655. Zbl 1329.05206 MR 3400569
- [7] A. W. HALES and R. I. JEWETT, [Regularity and positional games](#). *Trans. Amer. Math. Soc.* **106** (1963), no. 2, 222–229. Zbl 0113.14802 MR 143712
- [8] J. HUBIČKA and J. NEŠETŘIL, [All those Ramsey classes \(Ramsey classes with closures and forbidden homomorphisms\)](#). *Adv. Math.* **356** (2019), article no. 106791. Zbl 1421.05091 MR 4001036
- [9] S. JUNGE, [Categorical view of the Partite lemma in structural Ramsey theory](#). *Appl. Categ. Structures* **31** (2023), no. 4, article no. 30. Zbl 07725392 MR 4629482
- [10] A. S. KECHRIS, V. G. PESTOV and S. TODORCEVIC, [Fraïssé limits, Ramsey theory, and topological dynamics of automorphism groups](#). *Geom. Funct. Anal.* **15** (2005), no. 1, 106–189. Zbl 1084.54014 MR 2140630
- [11] K. LEEB, *Vorlesungen über Pascaltheorie*. Arbeitsber. Inst. Math. Masch. Datenverarb. 6, Friedrich-Alexander-Universität, Erlangen–Nürnberg, 1973.
- [12] D. MAŠULOVIĆ, [A dual Ramsey theorem for permutations](#). *Electron. J. Combin.* **24** (2017), no. 3, article no. P3.39. Zbl 1369.05200 MR 3691556
- [13] D. MAŠULOVIĆ, [Pre-adjunctions and the Ramsey property](#). *European J. Combin.* **70** (2018), 268–283. Zbl 1384.05153 MR 3779618
- [14] J. NEŠETŘIL, Ramsey theory. In *Handbook of combinatorics, Vol. 2*, pp. 1331–1403, Elsevier, Amsterdam, 1995. Zbl 0848.05065 MR 1373681
- [15] F. P. RAMSEY, [On a problem of formal logic](#). *Proc. London Math. Soc. (2)* **30** (1929), no. 4, 264–286. Zbl 55.0032.04 MR 1576401
- [16] S. SOLECKI, [Abstract approach to finite Ramsey theory and a self-dual Ramsey theorem](#). *Adv. Math.* **248** (2013), 1156–1198. Zbl 1283.05176 MR 3107538

- [17] S. SOLECKI, [Abstract approach to Ramsey theory and Ramsey theorems for finite trees](#). In *Asymptotic Geometric Analysis*, pp. 313–340, Fields Inst. Commun. 68, Springer, New York, 2013. Zbl [1267.05287](#) MR [3075998](#)
- [18] S. SOLECKI, [Monoid actions and ultrafilter methods in Ramsey theory](#). *Forum Math. Sigma* **7** (2019), article no. e2. Zbl [1404.05211](#) MR [3904213](#)
- [19] S. SOLECKI, [Dual Ramsey theorem for trees](#). *Combinatorica* **43** (2023), no. 1, 91–128. Zbl [07745869](#) MR [4622590](#)
- [20] S. TODORCEVIC, [Introduction to Ramsey spaces](#). Ann. of Math. Stud. 174, Princeton University Press, Princeton, NJ, 2010. Zbl [1205.05001](#) MR [2603812](#)
- [21] B. L. VAN DER WAERDEN, Beweis einer Baudetschen Vermutung. *Nieuw Arch. Wisk. (2)* **15** (1927), 212–216. Zbl [53.0073.12](#)
- [22] A. ZUCKER, [Topological dynamics of automorphism groups, ultrafilter combinatorics, and the generic point problem](#). *Trans. Amer. Math. Soc.* **368** (2016), no. 9, 6715–6740. Zbl [1359.37024](#) MR [3461049](#)

(Reçu le 24 mai 2022)

Sławomir SOLECKI, Department of Mathematics, Cornell University, Ithaca, NY 14853, USA;
e-mail: ssolecki@cornell.edu