

Patterson–Sullivan densities in convex projective geometry

Pierre-Louis Blayac

Abstract. For any rank-one convex projective manifold with a compact convex core, we prove that there exists a unique probability measure of maximal entropy on the set of unit tangent vectors whose geodesic is contained in the convex core, and that it is mixing. We use this to establish asymptotics for the number of closed geodesics. In order to construct the measure of maximal entropy, we develop a theory of Patterson–Sullivan densities for general rank-one convex projective manifolds. In particular, we establish a Hopf–Tsuji–Sullivan–Roblin dichotomy, and prove that, when it is finite, the measure on the unit tangent bundle induced by a Patterson–Sullivan density is mixing under the action of the geodesic flow.

Contents

1. Introduction	1
2. Reminders	11
3. Construction of the Sullivan measures	29
4. The Shadow lemma	32
5. The convergent case of the HTSR dichotomy	38
6. The divergent case of the HTSR dichotomy	39
7. Preparatory results on convex cocompact projective manifolds	53
8. The measure of maximal entropy	62
9. Counting closed geodesics	71
References	78

1. Introduction

Compact real hyperbolic manifolds are fundamental objects in geometry and dynamical systems; their geodesic flows are among the prime examples of chaotic systems

Mathematics Subject Classification 2020: 22D40 (primary); 20F67, 20F69, 22E40, 37D40 (secondary).

Keywords: ergodic theory, geodesic flow, convex projective geometry, Patterson–Sullivan densities, measure of maximal entropy.

which led to standard notions such as ergodicity or entropy. One classical way to deform the geometry of a compact hyperbolic manifold M_0 is to deform its Riemannian metric to allow variable negative curvature. It was one breakthrough of the theory of Anosov flows to establish that many dynamical features of the induced geodesic flow remain after such deformations (e.g. existence and uniqueness of the measure of maximal entropy, proved independently by Bowen [19] and Margulis [50]).

There is another interesting way to deform the geometry of M_0 . Consider the holonomy representation of the fundamental group $\rho_0: \pi_1(M_0) \rightarrow \text{PO}(d, 1)$, where d is the dimension of M_0 . Under certain conditions, one can deform it continuously (using e.g. bending, see [42]) to get a representation ρ valued in $\text{PGL}_{d+1}(\mathbb{R})$ which is not conjugate to a representation into $\text{PO}(d, 1)$. A theorem of Koszul [47, Cor. p. 103], combined with a theorem of Benoist [9, Thm. 1.1] (due to Choi–Goldman [25] for $d = 2$), ensures that the representation remains faithful and discrete, and $\rho(\pi_1(M_0))$ preserves and acts cocompactly on a properly convex open subset Ω of the real projective space $\text{P}(V)$, where $V = \mathbb{R}^{d+1}$. The quotient $\Omega/\rho(\pi_1(M_0))$ is a compact *convex projective manifold* which, like Riemannian manifolds, admits a geodesic flow, as we now recall.

In general, a convex projective manifold is a quotient $M = \Omega/\Gamma$ of a properly convex open set $\Omega \subset \text{P}(V)$ by a discrete group $\Gamma \subset \text{PGL}(V)$ of projective transformations preserving Ω . If M is compact, then we say that Γ divides Ω and that Ω is a divisible convex set. The set Ω admits a Finsler metric, called the Hilbert metric, which is proper and Γ -invariant (hence Γ acts properly discontinuously on Ω). Moreover, the intersection of any projective line with Ω is a geodesic for the Hilbert metric, which we call a *straight line* of Ω . Thus, there is a natural geodesic flow $(\phi_t)_{t \in \mathbb{R}}$ on the unit tangent bundle $T^1M := T^1\Omega/\Gamma$, which parametrises straight lines. Benoist [8] initiated the study of $(\phi_t)_{t \in \mathbb{R}}$ in the divisible case. He proved that if M is obtained by deforming a compact hyperbolic manifold, then Ω is *strictly convex*, i.e. its boundary $\partial\Omega \subset \text{P}(V)$ does not contain any non-trivial projective segment. He also proved that if M is compact, then Ω is strictly convex if and only if the geodesic flow $(\phi_t)_{t \in \mathbb{R}}$ is Anosov.

In this paper, we are particularly interested in divisible convex sets that are *not* strictly convex. Classical examples are the *higher-rank symmetric* divisible convex sets, namely the projective models of the symmetric spaces of $\text{PGL}_n(\mathbb{K})$, where $n \geq 3$ and \mathbb{K} is \mathbb{R} or \mathbb{C} or the classical quaternionic (or octonionic for $n = 3$) division algebra (for more details see [11, §2.4]). Other interesting, irreducible examples were constructed by Benoist [10], followed by Marquis [51], Ballas–Danciger–Lee [4] and Choi–Lee–Marquis [26], in dimensions 3 to 7.

Bray [21, 22] studied the geodesic flow of 3-dimensional irreducible compact convex projective manifolds $M = \Omega/\Gamma$ for which Ω is not necessarily strictly convex. Although the theory of Anosov flows does not apply in this setting, he managed to

construct, on the unit tangent bundle, a flow-invariant ergodic measure with maximal entropy [21, Thm. 1.1]. For this he used Benoist’s precise and beautiful geometric description of such 3-manifolds [10, Thm. 1.1]. He also adopted methods that had been elaborated for non-positively curved Riemannian manifolds, whose geodesic flows are not Anosov in general. More precisely, he drew on the work of Knieper [45] and Roblin [56], whose main tool are *Patterson–Sullivan densities*.

In the present article we study the dynamics of the geodesic flow of convex projective manifolds that have arbitrary dimension and are not necessarily compact. Like Bray, we use methods from the non-positively curved Riemannian world, in particular inspired by Knieper and Roblin. We generalise and improve Bray’s results [21], and develop more systematically the theory of Patterson–Sullivan densities in this setting.

1.1. Rank-one convex projective manifolds

Knieper [45] considered non-positively curved compact Riemannian manifolds which satisfy a property called *rank-one*. These generalise negatively curved compact manifolds, whose geodesic flow is uniformly hyperbolic, in only requiring that the geodesic flow have a hyperbolic behaviour along at least one geodesic, which is said to be *rank-one* (see [46, Def. 5.1.1]). Similarly, we will consider *rank-one* convex projective manifolds, which generalise convex projective manifolds $M = \Omega/\Gamma$ where Ω is strictly convex and $\partial\Omega$ is smooth (by which we mean \mathcal{C}^1 , see Section 2.2). We will use the following definition, recently introduced by M. Islam [40].

We say that a point of the boundary $\partial\Omega$ is *strongly extremal* if it does not belong to any non-trivial segment contained in $\partial\Omega$. If Ω is strictly convex, then all points of $\partial\Omega$ are strongly extremal. We say that a point is *smooth* if it admits a unique supporting hyperplane; such points are also commonly called \mathcal{C}^1 . We denote by $\partial_{\text{sse}}\Omega$ the set of smooth, strongly extremal points of $\partial\Omega$. Given any vector $v \in T^1\Omega$, we denote by $\pi v \in \Omega$ its footpoint and by $\phi_{\pm\infty}v = \lim_{t \rightarrow \pm\infty} \pi\phi_t v$ the intersection points of the projective line generated by v with $\partial\Omega$. We denote by $\text{Aut}(\Omega)$ the subgroup of $\text{PGL}(V)$ consisting of elements that preserve Ω , called automorphisms of Ω .

Definition 1.1 ([40, Def. 6.2 and Prop. 6.3]). Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set. A vector $v \in T^1\Omega$, and the geodesic of Ω spanned by v , are called *rank-one* if $\phi_{\infty}v$ and $\phi_{-\infty}v$ belong to $\partial_{\text{sse}}\Omega$. An infinite-order automorphism of Ω is said to be *rank-one* if it preserves a rank-one geodesic of Ω .

Let $\Gamma \subset \text{Aut}(\Omega)$ be a discrete subgroup, and $M := \Omega/\Gamma$. A vector v of T^1M , and the geodesic of M spanned by v , are said to be *rank-one* if any lift of v to $T^1\Omega$ is rank-one. The convex projective manifold (or orbifold) M is *rank-one* if it contains a rank-one periodic vector, i.e. if Γ contains a rank-one element.

We will see (Fact 2.11) that a rank-one automorphism g of Ω is a *proximal* element of $\mathrm{PGL}(V)$, i.e. it has an attracting fixed point in $\mathrm{P}(V)$; its inverse being also rank-one and hence proximal, g is said to be *biproximal*. We will also see that a rank-one automorphism preserves a unique rank-one geodesic in Ω .

If Ω is strictly convex and $\partial\Omega$ is smooth, then $\partial_{\mathrm{sse}}\Omega$ is the whole projective boundary $\partial\Omega$, all geodesics of Ω are rank-one and any biproximal automorphism of Ω is rank-one. In fact, if Ω is strictly convex *or* if $\partial\Omega$ is smooth, then $M = \Omega/\Gamma$ is rank-one as soon as Γ contains a biproximal element, by Facts 2.13 and 2.14 below.

When Ω is not strictly convex, an example of a geodesic in Ω that is *not* rank-one is a geodesic that is contained in a *properly embedded simplex* (PES), i.e. a projective simplex S of dimension $k \geq 2$ whose relative interior (see Section 2.2) is equal to $S \cap \Omega$. Such a simplex S can be interpreted as a flat of Ω since it is isometric to \mathbb{R}^k endowed with some norm. In many examples of convex projective manifolds $M = \Omega/\Gamma$, for instance when M is 3-dimensional, compact and irreducible, the maximal (for inclusion) PES's of Ω satisfy good properties (such as being isolated, see [10, 18, 41]) which imply that M is rank-one. More precisely, Islam used [41] to establish [40, Prop. A.2] that if $\Gamma \subset \mathrm{Aut}(\Omega)$ is a non-virtually abelian, discrete subgroup which is relatively hyperbolic with respect to a collection of virtually abelian subgroups of rank at least two, and which acts convex cocompactly on Ω in the sense of Danciger–Guéritaud–Kassel [35] (this notion is introduced in the next Section 1.2), then Ω/Γ is rank-one. Islam's argument is explained in the particular case of 3-dimensional compact convex projective manifolds in Example 2.17.

Rank-one manifolds are interesting because they include a diversity of examples, and are in some sense generic: in both the Riemannian and the convex projective settings, there exist higher-rank rigidity theorems which classify compact higher-rank (i.e. not rank-one) manifolds. See the work of Ballmann [5, Cor. 1] and Burns–Spatzier [24, Thm. 5.1] in the Riemannian case and the recent work of A. Zimmer [64, Thm. 1.4] in the convex projective case.

1.2. The Bowen–Margulis measure on quotients of convex cocompact actions

The generalisation of Bray's results [21] that we are about to state is analogous to [45, Thm. 1.1.i], which says that any non-positively curved rank-one compact Riemannian manifold has a unique measure of maximal entropy. However, our result does not restrict to compact manifolds: it concerns the more general class of manifolds that are quotients of *convex cocompact actions*.

Let $M = \Omega/\Gamma$ be a convex projective manifold. Following Danciger–Guéritaud–Kassel, the action of Γ on Ω is said to be *naively convex cocompact* if there exists a non-empty Γ -invariant convex subset of Ω on which Γ acts cocompactly; when Ω is not strictly convex, this notion is not quite satisfactory, because a small deformation

of Γ in $\mathrm{PGL}(V)$ may not preserve any properly convex open set (see [35, §4.1]). Therefore, we will consider another stronger definition, introduced by Danciger–Guéritaud–Kassel.

The *full orbital limit set* of Γ is the union over all $x \in \Omega$ of the set of accumulation points of the orbit $\Gamma \cdot x$, and is denoted by $\Lambda_{\Omega}^{\mathrm{orb}}(\Gamma)$, or simply Λ^{orb} when the context is clear. When Ω is not strictly convex, the set of accumulation of points of an orbit $\Gamma \cdot x$ may depend on the choice of $x \in \Omega$ (e.g. if Ω is a triangle in $\mathrm{P}(\mathbb{R}^3)$, and Γ is generated by an infinite-order non-proximal element). The convex hull in Ω of the full orbital limit set is denoted by $\mathcal{C}_{\Omega}^{\mathrm{cor}}(\Gamma)$; it is Γ -invariant.

Definition 1.2 ([35, Def. 1.11]). Let $\Omega \subset \mathrm{P}(V)$ be a properly convex open set, and $\Gamma \subset \mathrm{Aut}(\Omega)$ a discrete subgroup. The action of Γ on Ω is said to be *convex cocompact* if $\mathcal{C}_{\Omega}^{\mathrm{cor}}(\Gamma)$ is non-empty and has compact quotient by Γ ; the projection in M of $\mathcal{C}_{\Omega}^{\mathrm{cor}}(\Gamma)$ is called the *convex core* of M .

We refer to [35, §§1.4–1.7, 4.1–10.7] and [36] for more details and examples on convex cocompactness. Note that if Γ divides a properly convex open set Ω , then the convex hull of any Γ -orbit in Ω is equal to Ω (this is due to Vey [61, Prop. 3]), hence $\mathcal{C}_{\Omega}^{\mathrm{cor}}(\Gamma) = \Omega$ and Γ acts convex cocompactly on Ω .

Danciger–Guéritaud–Kassel [35, Cor. 4.8] proved that if the action of Γ on Ω is convex cocompact, then Λ^{orb} is closed. We denote by $T^1 M_{\mathrm{cor}} \subset T^1 M$ the $(\phi_t)_{t \in \mathbb{R}}$ -invariant (and compact if Γ is convex cocompact) subset consisting of those vectors whose orbit under the geodesic flow is contained in the convex core, i.e. the endpoints of any lift to Ω of the geodesic are in Λ^{orb} .

A rank-one convex projective manifold is said to be *non-elementary* if its fundamental group does not contain \mathbb{Z} as a finite-index subgroup.

Theorem 1.3. *Let $\Omega \subset \mathrm{P}(V)$ be a properly convex open set, and $\Gamma \subset \mathrm{Aut}(\Omega)$ a discrete subgroup. Assume that Γ acts convex cocompactly on Ω and that $M = \Omega/\Gamma$ is rank-one and non-elementary. Then there exists a unique $(\phi_t)_{t \in \mathbb{R}}$ -invariant probability measure m on $T^1 M_{\mathrm{cor}}$ with maximal entropy (Bowen–Margulis measure). Moreover, m is mixing.*

Recall that if (X, m) is a measured space with m finite and invariant under a measurable flow $(\phi_t)_{t \in \mathbb{R}}$, then m is said to be *mixing* if

$$\lim_{t \rightarrow \infty} m(\phi_t(A) \cap B) m(X) = m(A) m(B)$$

for all measurable subsets $A, B \subset X$. Reminders on the notion of entropy are given in Sections 2.13 and 2.14.

Remark 1.4. In a first version of this paper, we had proved Theorem 1.3 under the stronger assumption that Γ acts strongly irreducibly on $\mathrm{P}(V)$.

Replacing this assumption by non-elementary was made possible by Facts 2.18 and 6.13 below, which are proved in the paper [17, Prop. 2.4 and Lem. 2.8] in collaboration with Feng Zhu.

Note that some results in the present paper (in particular, Theorem 1.6) are used in [17]. However, *there is no circular reasoning between the two papers*, since Section 2 of [17], where is proved what we need here, does not use the present paper.

Moreover, Facts 2.18 and 6.13 can also be seen as consequences of more general results established in the author's PhD thesis [15, Lem. 3.1.3.2, Prop. 3.2.2, Lem. 3.2.4, Cor. 5.2.7, Prop. 5.3.1].

Following Knieper [45], we will use Theorem 1.3 to establish asymptotic estimates on the number of closed geodesics on M , and equidistribution results on the Lebesgue measures on closed geodesics. (Note that some counting results for the Riemannian case were already present in another earlier work of Knieper [44].)

Recall that in our convex projective setting, the *critical exponent* of Γ is defined as

$$\delta_\Gamma := \limsup_{r \rightarrow \infty} \frac{1}{r} \log(\#\{\gamma \in \Gamma : d_\Omega(o, \gamma o) \leq r\}),$$

where o is any point of Ω , and d_Ω is the Hilbert metric on Ω ; it does not depend on the Γ -invariant properly convex open set Ω , see Section 2.5. For any element $g \in \text{PGL}(V)$, we set

$$\ell(g) := \frac{1}{2} \log\left(\frac{\lambda_1(\tilde{g})}{\lambda_{d+1}(\tilde{g})}\right), \quad (1.1)$$

where $\tilde{g} \in \text{GL}(V)$ is any lift of g , the integer $d + 1$ is the dimension of V , and $\lambda_1(\tilde{g}) \geq \dots \geq \lambda_{d+1}(\tilde{g})$ are the moduli of the (complex) eigenvalues of \tilde{g} . Given a convex projective manifold $M = \Omega/\Gamma$ and a positive number $T > 0$, we denote by $[\Gamma]_T$ (resp. $[\Gamma]_T^{\text{sing}}$, resp. $[\Gamma]_T^{\text{r1}}$) the set of conjugacy classes of elements (resp. of non-rank-one elements, resp. of rank-one elements) $\gamma \in \Gamma$ such that $\ell(\gamma) \leq T$. When $\gamma \in \Gamma$ is rank-one, we denote by $\mathcal{L}[\gamma]$ the Lebesgue measure on the rank-one closed geodesic associated to γ , normalised to be a probability measure. The link between closed geodesics on M and conjugacy classes of Γ is recalled in Section 2.7.

Proposition 1.5. *In the setting of Theorem 1.3, one can find constants $0 < \delta < \delta_\Gamma$ and $C > 0$ such that for any $T > C$,*

- (1) $(1/CT)e^{\delta_\Gamma T} \leq \#[\Gamma]_T \leq (C/T)e^{\delta_\Gamma T}$;
- (2) $\#[\Gamma]_T^{\text{sing}} \leq e^{\delta T}$;
- (3) $(1/\#[\Gamma]_T^{\text{r1}}) \sum_{[\gamma] \in [\Gamma]_T^{\text{r1}}} \mathcal{L}[\gamma] \xrightarrow{T \rightarrow \infty} m$ for the weak* topology.

In the case that M is compact, Proposition 1.5(1) was previously established by Islam [40, Thm. 1.12] using different techniques. One may compare Proposition 1.5(2) with another result of Islam [40, Thm. 1.11] concerning random walks

on Γ . We prove a slightly more general version of Proposition 1.5 (3) in Proposition 9.5 below.

1.3. The machinery of Patterson–Sullivan densities

The main idea to prove Theorem 1.3 is to use *conformal densities* (Definition 2.20), also called *Patterson–Sullivan densities*. They are, for an arbitrary proper metric space (X, d) acted on by a discrete subgroup $\Gamma \subset \text{Isom}(X, d)$, families of finite measures (Patterson–Sullivan measures) on the *horoboundary* $\partial_h X$ of (X, d) . More precisely, Patterson–Sullivan measures are quasi-invariant under the action of Γ , and there is an explicit formula for the Radon–Nikodym cocycle (see Definition 2.21). They were originally introduced by Patterson [53] and Sullivan [58] for discrete groups of isometries of real hyperbolic spaces, whose horoboundary is the boundary at infinity. They were later used in much more general geometric settings, such as non-positively curved Riemannian manifolds by Knieper [45], CAT(−1)-spaces by Roblin [56], and even CAT(0)-spaces by Picaud–Link [48] and Ricks [55]. In all these settings, the horoboundary is equal to the visual boundary.

Conformal densities were also brought to convex projective geometry, in the strictly convex and smooth case by Crampon in his PhD thesis [32] and more recently by F. Zhu [63], and in the non-strictly convex case in dimension 3 by Bray [21]. While Zhu does not adapt Knieper’s work (hence does not prove the existence and uniqueness of the measure of maximal entropy), his adaptation of Roblin’s results goes further than in the present paper, enabling him for instance to obtain more precise estimates for various counting problems. In the paper [17], the author and Zhu generalise the results of [63] to the non-strictly convex case.

Let $M = \Omega / \Gamma$ be a convex projective manifold. If $\partial\Omega$ is smooth, then the horoboundary $\partial_h \Omega$ is the projective boundary $\partial\Omega$ (see Fact 2.26). When $\partial\Omega$ is not smooth, the situation is more delicate since $\partial_h \Omega$ is different from $\partial\Omega$. To handle this difficulty, Bray’s strategy was to weaken the definition of conformal densities so that it has a meaning on $\partial\Omega$. We will do something slightly different: we will use a result of Walsh [62, Thm. 1.3], who proved in general that the horoboundary $\partial_h \Omega$ *dominates* $\partial\Omega$, in the sense that the identity on Ω extends to a continuous map $\partial_h \Omega \rightarrow \partial\Omega$, which is onto by density of Ω . We will define conformal densities on $\partial\Omega$ simply as the push-forwards of conformal densities on $\partial_h \Omega$ by the natural projection $\partial_h \Omega \rightarrow \partial\Omega$.

Let us detail the steps that we will follow below, adapted from the general theory of conformal densities. Recall that conformal densities depend on a parameter $\delta \geq 0$. Fix $o \in \Omega$.

- (1) Construct a δ_Γ -conformal density (this is actually very general, see Fact 2.22).

(2) Given $\delta \geq 0$ and a δ -conformal density $(\mu_x)_{x \in \Omega}$ on $\partial_{\text{h}}\Omega$, construct a measure on $\partial_{\text{h}}\Omega^2 \times \mathbb{R}$ which is invariant under the actions of Γ and \mathbb{R} , and equivalent to $\mu_o \times \mu_o$ times the Lebesgue measure. Derive from it a $(\phi_t)_{t \in \mathbb{R}}$ -invariant measure on T^1M , called the induced *Sullivan measure*.

(3) Given $\delta \geq 0$ and a δ -conformal density $(\mu_x)_{x \in \Omega}$ on $\partial_{\text{h}}\Omega$, prove a *Hopf–Tsuji–Sullivan–Roblin (HTSR) dichotomy*. It states in particular that the sum

$$\sum_{\gamma \in \Gamma} e^{-\delta d_{\Omega}(o, \gamma o)}$$

is infinite if and only if the induced Sullivan measure is ergodic under the geodesic flow; in this case $\delta = \delta_{\Gamma}$, the δ_{Γ} -conformal density is unique and its induced Sullivan measure is called the *Bowen–Margulis measure*. Recall that ergodic means that any $(\phi_t)_{t \in \mathbb{R}}$ -invariant measurable set has null or full measure. (A set has full measure if its complement has null measure.)

(4) Assuming that

$$\sum_{\gamma \in \Gamma} e^{-\delta_{\Gamma} d_{\Omega}(o, \gamma o)}$$

is infinite, prove that if the Bowen–Margulis measure is finite, then it is mixing under the geodesic flow.

(5) Use mixing to solve counting problems. There are many counting problems, such as estimating, as R tends to infinity, the number of points in a fixed orbit $\Gamma \cdot o$ that lie in the ball of radius R centred at o , or the number of closed geodesics with length less than R .

As we are particularly interested in the compact case, we decided to follow, instead of Roblin’s, Knieper’s approach. It gives estimates for the number of closed geodesics with length less than R as R grows, but also establishes that the Bowen–Margulis measure is the unique measure with maximal entropy.

Recall that if

$$\sum_{\gamma \in \Gamma} e^{-\delta_{\Gamma} d_{\Omega}(o, \gamma o)} = \infty,$$

then Γ is said to be *divergent*; this notion does not depend on the Γ -invariant properly convex open set $\Omega \subset P(V)$, see Section 2.5.

One strength of the theory of conformal densities in the various geometric settings we have mentioned is that many properties of compact manifolds still hold under the following much weaker assumptions: asking Γ to be divergent and the Bowen–Margulis measure to be finite. For example, these assumptions hold if Γ acts convex cocompactly on Ω and $M = \Omega / \Gamma$ is rank-one (see Proposition 1.7).

Another broad class of manifolds which are generally well understood are the geometrically finite manifolds. In some settings, such as Riemannian geometry with

variable negative curvature, there is no implication between geometric finiteness and finiteness of the Bowen–Margulis measure (see [56, §1.F]). Crampon–Marquis [33, Def. 1.4] defined geometrically finite convex projective manifolds $M = \Omega/\Gamma$ in the case where Ω is strictly convex and $\partial\Omega$ is smooth. They carefully investigated the dynamics of the geodesic flow on these manifolds, without using conformal densities. In particular, they extended Benoist’s result that the flow is uniformly hyperbolic [34, Thm. 1.2] (under an additional assumption of asymptotically hyperbolic cusps). Zhu and the author [17, Thm. C] proved (generalising earlier results by Zhu [63, Thm. 10 and Prop. 14]) that for these manifolds Γ is divergent and the Bowen–Margulis measure is finite.

1.4. The Hopf–Tsuji–Sullivan–Roblin dichotomy

Let us explain more precisely Steps (1), (2) and (3) of the previous section. Let $M = \Omega/\Gamma$ be a non-elementary rank-one convex projective manifold. On the one hand, Steps (1) and (2), i.e. constructing a δ_Γ -conformal density on $\partial_{\mathfrak{h}}\Omega$ and the induced Sullivan measure on T^1M , do not require major changes compared to the case where M is a real hyperbolic manifold; these steps are done in Sections 2.9 and 3.3. On the other hand, Step (3), namely the HTSR dichotomy, is more delicate; let us state it formally. For this, we need to recall the definition of two limit sets, both contained in the full orbital limit set.

The *conical limit set* of the action of Γ on Ω is denoted by $\Lambda_\Omega^{\text{con}}(\Gamma)$ (or simply Λ^{con} when the context is clear), and defined as follows. Fix $o \in \Omega$, then $\xi \in \partial\Omega$ belongs to Λ^{con} if there exists a sequence $(\gamma_n)_{n \in \mathbb{N}} \in \Gamma^{\mathbb{N}}$ going to infinity such that the sequence $(d_\Omega(\gamma_n o, [o, \xi]))_{n \in \mathbb{N}}$ is bounded; this does not depend on the choice of o . The *proximal limit set* of Γ , denoted by $\Lambda^{\text{prox}}(\Gamma)$, or simply Λ^{prox} when the context is clear, is the closure of the set of attracting fixed points of proximal elements of Γ .

In general, $\Lambda^{\text{con}} \subset \Lambda^{\text{orb}}$ and $\Lambda^{\text{prox}} \subset \overline{\Lambda^{\text{orb}}}$. Furthermore, Danciger–Guéritaud–Kassel [35, Cor. 4.8 and Lem. 4.18] established that Γ acts convex cocompactly on Ω if and only if $\mathcal{C}_\Omega^{\text{cor}}(\Gamma)$ is non-empty and $\Lambda_\Omega^{\text{con}}(\Gamma)$ and $\Lambda_\Omega^{\text{orb}}(\Gamma)$ are equal and closed.

In [14, Def. 1.1] we introduced a $(\phi_t)_{t \in \mathbb{R}}$ -invariant closed subset of T^1M , called the *biproximal unit tangent bundle* and denoted by T^1M_{bip} ; it consists of those vectors $v \in T^1M$ such that $\phi_{\pm\infty} \tilde{v} \in \Lambda^{\text{prox}}$ for any lift $\tilde{v} \in T^1\Omega$. When M is rank-one and Γ is divergent, the following result gives a new interpretation of T^1M_{bip} , as the support of the Bowen–Margulis measure on T^1M .

Theorem 1.6. *Let $o \in \Omega \subset \mathbb{P}(V)$ be a pointed properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Let $\delta \geq 0$, let $(v_x)_{x \in \Omega}$ be a δ -conformal density on $\partial\Omega$ and let m be the induced Sullivan measure. Then there are two possibilities:*

- (1) either $\sum_{\gamma} e^{-\delta d_{\Omega}(o, \gamma o)} < \infty$, and then $\nu_o(\Lambda^{\text{con}}) = 0$ and the dynamical systems $(T^1 M, \phi_t, m)$, $(\text{Geod}^{\infty}(\Omega), \Gamma, \nu_o^2)$ and $(T^1 \Omega, \Gamma \times \mathbb{R}, \tilde{m})$ are dissipative and non-ergodic;
- (2) or $\sum_{\gamma} e^{-\delta d_{\Omega}(o, \gamma o)} = \infty$, in which case $\delta = \delta_{\Gamma}$ (and Γ is divergent), and
- $(\nu_x)_{x \in \Omega}$ is the only δ_{Γ} -conformal density (up to a scalar multiple), and m is called the Bowen–Margulis measure on $T^1 M$;
 - $\nu_o(\partial_{\text{sse}} \Omega \cap \Lambda^{\text{prox}} \cap \Lambda^{\text{con}}) = \nu_o(\partial \Omega)$ and ν_o is non-atomic, in particular $\text{supp}(m) = T^1 M_{\text{bip}}$;
 - $(T^1 M, \phi_t, m)$, $(\partial \Omega^2, \Gamma, \nu_o^2)$ and $(T^1 \Omega, \Gamma \times \mathbb{R}, \tilde{m})$ are conservative and ergodic;
 - if m is finite then it is mixing.

In (1) we denote by $\text{Geod}^{\infty}(\Omega)$ the space of straight geodesics of Ω , which consists of pairs $(\xi, \eta) \in \partial \Omega^2$ such that $[\xi, \eta] \cap \Omega$ is non-empty. Reminders on the dynamical notions of dissipativity and conservativity are given in Section 2.11.

To establish the mixing property in Theorem 1.6(2) when m is finite, we use Babillot’s strategy of proof [3], and also some general results of Coudène [30] in ergodic theory that are inspired by [3]. In particular, cross-ratios of quadruples of points on the boundary of a properly convex open set are a crucial component of the proof of the mixing property. Zhu proved the mixing property [63, Thm. 18] in the case where Ω is strictly convex with \mathcal{C}^1 boundary by using the same strategy.

As we explained in Section 1.3, in order to prove Theorem 1.3 we will use Theorem 1.6(2); hence we need to check that the latter can be applied to the case where Γ acts convex cocompactly on Ω and $M = \Omega / \Gamma$ is rank-one. This is the goal of the next proposition.

Proposition 1.7. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup. Suppose $M = \Omega / \Gamma$ is rank-one and non-elementary. Then Γ is divergent, and the Bowen–Margulis measure is finite.*

In another article [16], we prove that for any non-elementary rank-one compact convex projective manifold $M = \Omega / \Gamma$, we have $\Lambda^{\text{prox}} = \partial \Omega$. This implies that

$$T^1 M_{\text{bip}} = T^1 M,$$

and from Theorem 1.6 and Proposition 1.7 we derive the following.

Corollary 1.8. *The measure of maximal entropy of any non-elementary rank-one compact convex projective manifold has full support.*

Benoist [8, Prop. 6.7] proved that, in the setting of Corollary 1.8, if M is not hyperbolic, then the measure of maximal entropy is singular with respect to the Lebesgue measure.

Organisation of the paper. In Section 2 we collect definitions and basic properties in convex projective geometry, on conformal densities and on entropy.

Sections 3 to 6 are based on Roblin’s work [56]. In Section 3 we define the Hopf coordinates and the Gromov product in the setting of convex projective geometry, in order to make sense of Sullivan’s formula [58, Prop. 11], which defines Sullivan measures. In Section 4 we state and prove a convex projective version of the Shadow lemma, a fundamental result in the study of conformal densities. In Section 5 we establish the convergent case of the HTSR dichotomy. In Section 6 we assume that Γ is divergent, and follow closely Roblin’s proof of HTSR dichotomy in order to establish the convex projective version of Theorem 1.6(2). The proof is divided into several steps: proving that the conical limit set has full measure (Section 6.1); proving that $\partial_{\text{sse}}\Omega$ has full measure (Section 6.3); proving that the Bowen–Margulis measure is ergodic, and moreover mixing when finite (Section 6.6); and finally proving that Λ^{prox} has full measure (Section 6.7).

Sections 7 to 9 are based on Knieper’s work [45]. In Section 7 we collect some properties of convex cocompact projective actions; in particular we prove Proposition 1.7, and give two refined versions of the Shadow lemma (Lemma 4.2) which will be used in Section 8. In Section 8 we prove Theorem 1.3. The main three steps are: estimating the measure of small dynamical balls (Section 8.1); bounding from below the entropy of the Bowen–Margulis measure (Section 8.2); proving the uniqueness of the measure of maximal entropy (Section 8.4). In Section 9 we establish Proposition 1.5. The main steps are: bounding the number of rank-one closed geodesics from below (Section 9.1) and from above (Section 9.2); bounding from above the number of non-rank-one closed geodesics (Section 9.3); and finally proving the equidistribution of closed geodesics (Section 9.4).

2. Reminders

2.1. Properly convex open subsets of $\mathbf{P}(\mathbb{R}^{d+1})$ and their geodesic flow

In the whole paper we fix a real vector space $V = \mathbb{R}^{d+1}$, where $d \geq 1$. Let $\Omega \subset \mathbf{P}(V)$ be a properly convex open set. Recall that Ω admits an $\text{Aut}(\Omega)$ -invariant proper metric called the *Hilbert metric* and defined by the following formula: for $(a, x, y, b) \in \partial\Omega \times \Omega \times \Omega \times \partial\Omega$ aligned in this order (see Figure 1),

$$d_{\Omega}(x, y) = \frac{1}{2} \log([a, x, y, b]), \quad (2.1)$$

where $[a, x, y, b]$ is the cross-ratio of the four points, normalised so that $[0, 1, t, \infty] = t$.




Figure 1. The Hilbert metric and the geodesic flow ($t = d_\Omega(x, y)$).

Recall that if Ω is an ellipsoid, then (Ω, d_Ω) is the Klein model of the real hyperbolic space of dimension d ; if Ω is a d -simplex, then (Ω, d_Ω) is isometric to \mathbb{R}^d endowed with a hexagonal norm.

Any discrete subgroup $\Gamma \subset \text{PGL}(V)$ of automorphisms of Ω preserves d_Ω , hence must act properly discontinuously on Ω ; therefore the quotient $M = \Omega/\Gamma$ is an orbifold. Furthermore, M is a manifold if the action is free (i.e. if Γ is torsion-free, by Brouwer's fixed point theorem, applied to the convex hull of a finite orbit of a torsion element). Note that by Selberg's lemma [57], if Γ is finitely generated, then it has a torsion-free finite-index subgroup. We will work in general with Γ not necessarily torsion-free, so we set the notation $T^1M = T^1\Omega/\Gamma$.

The straight lines of Ω can be parametrised to be geodesics, which are said to be *straight*. However, an interesting feature in the non-strictly convex case is that when there are two coplanar non-trivial segments in $\partial\Omega$, one can construct geodesics which are not straight, see the broken green segment in Figure 1. In order to define the geodesic flow we only take into account straight geodesics: for v in $T^1\Omega$, let $t \mapsto c(t)$ be the parametrisation of the projective line tangent to v such that c is an isometric embedding from \mathbb{R} to Ω and $c'(0) = v$. For $t \in \mathbb{R}$ we set $\phi_t(v) = c'(t) \in T^1\Omega$. See Figure 1.

The geodesic flow on $T^1\Omega/\Gamma$ is well defined because the two actions of $\text{Aut}(\Omega)$ and $(\phi_t)_{t \in \mathbb{R}}$ on $T^1\Omega$ commute. We denote by $\pi: T^1M \rightarrow M$ and $\pi: T^1\Omega \rightarrow \Omega$ the projections, we define the following metrics:

$$\begin{aligned} \forall x, y \in M, \quad d_M(x, y) &= \min\{d_\Omega(\tilde{x}, \tilde{y}) : \tilde{x}, \tilde{y} \in \Omega \text{ lifts of } x, y\}, \\ \forall v, w \in T^1\Omega, \quad d_{T^1\Omega}(v, w) &= \max_{0 \leq t \leq 1} d_\Omega(\pi\phi_t v, \pi\phi_t w), \\ \forall v, w \in T^1M, \quad d_{T^1M}(v, w) &= \max_{0 \leq t \leq 1} d_M(\pi\phi_t v, \pi\phi_t w) \\ &\leq \min\{d_{T^1\Omega}(\tilde{v}, \tilde{w}) : \tilde{v}, \tilde{w} \in T^1\Omega \text{ lifts of } v, w\}. \end{aligned}$$

A very useful inequality is provided by the following lemma.

Lemma 2.1 ([31, Lem. 8.3]). *Let Ω be a properly convex open subset of $\mathbb{P}(V)$. Let c_1 and c_2 be two straight geodesics parametrised with constant speed, but not necessarily with the same speed. Then for all $0 \leq t \leq T$,*

$$d_\Omega(c_1(t), c_2(t)) \leq d_\Omega(c_1(0), c_2(0)) + d_\Omega(c_1(T), c_2(T)).$$

See [14, §A] for a proof of the above lemma that fill in a missing detail in the original proof.

2.2. Terminology on convex sets and duality

We recall here some terminology on convex sets.

Notation 1. For any subset X of the projective space $\mathbb{P}(V)$, the closure (resp. interior, resp. boundary) of X , denoted by \bar{X} (resp. $\text{int}(X)$, resp. ∂X), will always be considered *with respect to* $\mathbb{P}(V)$.

Let $K \subset \mathbb{P}(V)$ be properly convex.

- The *relative interior* (resp. *relative boundary*) of K , denoted by $\text{int}_{\text{rel}}(K)$ (resp. $\partial_{\text{rel}}K$) is its topological interior (resp. boundary) with respect to the projective subspace it spans.
- For $x \in \bar{K}$, the *open face* of x in \bar{K} , denoted by $F_K(x)$, consists of the points $y \in \bar{K}$ such that $[x, y]$ is contained in the relative interior of a segment contained in \bar{K} . The *closed face* of x is $\bar{F}_K(x) = \overline{F_K(x)}$.
- A point $x \in \partial_{\text{rel}}K$ is said to be *extremal* (resp. *strongly extremal*) if $F_K(x) = \{x\}$ (resp. $x \notin \bar{F}_K(y)$ for $y \in \partial_{\text{rel}}K \setminus \{x\}$); one says that K is *strictly convex* if all the points in the relative boundary are extremal (and hence strongly extremal).
- Assume that K spans $\mathbb{P}(V)$ and let $\xi \in \partial K$. A *supporting hyperplane* of K at ξ is a hyperplane which contains ξ but does not intersect $\text{int}(K)$. Note that there always exists such a hyperplane. The point ξ is said to be a *smooth point* of ∂K if there is only one supporting hyperplane of K at ξ , denoted by $T_\xi \partial K$.

Let us recall the notion of duality for properly convex open sets. We identify the dual projective space $\mathbb{P}(V^*)$ with the set of projective hyperplanes of $\mathbb{P}(V)$. Let Ω be a properly convex open subset of $\mathbb{P}(V)$. The dual of Ω , denoted by Ω^* , is the properly convex open subset of $\mathbb{P}(V^*)$ defined as the set of projective hyperplanes which do not intersect $\bar{\Omega}$. We naturally identify $\text{PGL}(V)$ and $\text{PGL}(V^*)$, then $\text{Aut}(\Omega)$ identifies with $\text{Aut}(\Omega^*)$, and the attracting (resp. repelling) fixed point of the action on $\mathbb{P}(V^*)$ of any biproximal element $g \in \text{PGL}(V)$ is $x_g^+ \oplus x_g^0$ (resp. $x_g^- \oplus x_g^0$).

Observe that a hyperplane H is a smooth point of $\partial\Omega^*$ if and only if its intersection with $\bar{\Omega}$ is reduced to a singleton.

2.3. More metrics and upper semi-continuity of balls

In this section we define two metrics, one on the usual boundary $\partial\Omega$ of any properly convex open set $\Omega \subset \mathbb{P}(V)$, and one on the closure $\bar{\Omega}$. These notions are used in the proofs of Lemma 4.4 and Proposition 6.2.

Definition 2.2. Let $x \in \Omega \subset \mathbb{P}(V)$ be a pointed properly convex open set, and $t > 0$ be a positive parameter.

- Let $\xi, \eta \in \partial\Omega$. The *simplicial distance* between ξ and η is defined as follows (and it is possibly infinite):

$$d_{\text{spl}}(\xi, \eta) := \inf\{k : \exists a_0, \dots, a_k \in \partial\Omega : a_0 = \xi, a_k = \eta, \\ \forall 0 \leq i < k, [a_i, a_{i+1}] \subset \partial\Omega\}.$$

- Let $\xi, \eta \in \bar{\Omega}$. When ξ and η are on the same open face F of $\bar{\Omega}$, we set $d_{\bar{\Omega}}(\xi, \eta) := d_F(\xi, \eta)$, and otherwise we set $d_{\bar{\Omega}}(\xi, \eta)$ to be infinite.

The open balls of radius R and centred at ξ are respectively denoted by $B_{\text{spl}}(\xi, R)$ and $B_{\bar{\Omega}}(\xi, R)$.

Observe that d_{spl} and $d_{\bar{\Omega}}$ are lower semi-continuous. In particular, closed balls for the metric $d_{\bar{\Omega}}$ are upper semi-continuous with respect to the Hausdorff topology (in the sense of Fact 2.3 below). Given a metrisable locally compact topological space X , recall that the *Hausdorff topology* on the set of compact subsets of X is a metrisable topology such that a sequence of compact subsets $(K_n)_n$ converges to $K \subset X$ compact if and only if any converging sequence $(x_n)_n \in \prod_n K_n$ has its limit in K , and any point of K is the limit of a sequence $(x_n)_n \in \prod_n K_n$.

Fact 2.3. Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set. For any $R > 0$, the map

$$\begin{aligned} \bar{B}_{\bar{\Omega}}(\cdot, R) : \bar{\Omega} &\rightarrow \{\text{compact subsets of } \bar{\Omega}\}, \\ \xi &\mapsto \bar{B}_{\bar{\Omega}}(\xi, R) \end{aligned}$$

is upper semi-continuous in the following sense: all accumulation points of $\bar{B}_{\bar{\Omega}}(\eta, R)$ when $\eta \rightarrow \xi$ must be contained in $\bar{B}_{\bar{\Omega}}(\xi, R)$.

2.4. Benzécri's compactness theorem and proper densities

In this section we recall Benzécri's famous compactness theorem, and we see a first consequence for $\text{PGL}(V)$ -equivariant volume form on properly convex open sets. We

denote by \mathcal{E}_V (resp. \mathcal{E}_V^\bullet) the set of (resp. pointed) properly convex open set of $\mathbb{P}(V)$ (resp. endowed with the pointed Hausdorff topology).

Fact 2.4 ([12, Ch. 5, §2, Thm. 2]). The action of $\mathrm{PGL}(V)$ on \mathcal{E}_V^\bullet is continuous, proper and cocompact.

We recall the notion of a proper density on the set of properly convex open sets. They prescribe the way to choose a volume form on each properly convex open set in a $\mathrm{PGL}(V)$ -equivariant manner. For the whole paper we fix a density $\mathrm{Vol}_{\mathbb{P}(V)}$ on $\mathbb{P}(V)$, seen as a measure.

Definition 2.5. A *proper density* on \mathcal{E}_V is a map of the form $\Omega \mapsto \mathrm{Vol}_\Omega$, where $\Omega \subset \mathbb{P}(V)$ is a properly convex open set and Vol_Ω is a density on Ω with Radon–Nikodym derivative $f(x, \Omega) > 0$ with respect to $\mathrm{Vol}_{\mathbb{P}(V)}$, satisfying the following three conditions.

Continuity. The function $f: \mathcal{E}_V^\bullet \rightarrow \mathbb{R}_{>0}$ is continuous.

Monotone decreasing. Let (x, Ω) and $(y, \Omega') \in \mathcal{E}_V$. If $x = y \in \Omega \subset \Omega'$, then

$$f(x, \Omega') \leq f(x, \Omega).$$

$\mathrm{PGL}(V)$ -equivariance. For any $T \in \mathrm{PGL}(V)$,

$$T_* \mathrm{Vol}_\Omega = \mathrm{Vol}_{T(\Omega)}.$$

See [60] for more details and examples. We fix for the whole paper a proper density $\Omega \mapsto \mathrm{Vol}_\Omega$ on \mathcal{E}_V . One of the key observations that we will need on proper densities is that for any $R > 0$, the following quantities are positive and finite (this is a direct consequence of Fact 2.4):

$$\begin{aligned} 0 < \chi_-(R) &:= \min_{(x, \Omega) \in \mathcal{E}_V^\bullet} \mathrm{Vol}_\Omega(\bar{B}_\Omega(x, R)) \leq \chi_+(R) \\ &:= \max_{(x, \Omega) \in \mathcal{E}_V^\bullet} \mathrm{Vol}_\Omega(\bar{B}_\Omega(x, R)) < \infty. \end{aligned} \quad (2.2)$$

2.5. The critical exponent

Let $\Gamma \subset \mathrm{PGL}(V)$ be a discrete subgroup which preserves a properly convex open set $\Omega \subset \mathbb{P}(V)$. We defined in the introduction (Sections 2.5 and 1.3) the critical exponent of Γ and when Γ is divergent. In this section we explain why these definitions do not depend on Ω .

Recall that $V = \mathbb{R}^{d+1}$ has a canonical Euclidean structure. For any $g \in \mathrm{GL}(V) = \mathrm{GL}_{d+1}(\mathbb{R})$, we denote by $\mu_1(g) \geq \dots \geq \mu_{d+1}(g)$ the singular values of g . For any

$g \in \mathrm{PGL}(V)$, we set

$$\kappa(g) := \frac{1}{2} \log \left(\frac{\mu_1(\tilde{g})}{\mu_{d+1}(\tilde{g})} \right),$$

where $\tilde{g} \in \mathrm{GL}(V)$ is any lift of g .

Fact 2.6 ([35, Prop. 10.1]). Let $\Omega \subset \mathrm{P}(V)$ be a properly convex open subset and let $x \in \Omega$. Then there exists a constant $C > 0$ such that for any $g \in \mathrm{Aut}(\Omega)$, one has

$$|d_\Omega(x, gx) - \kappa(g)| \leq C.$$

As a consequence, if $\Gamma \subset \mathrm{PGL}(V)$ is a discrete subgroup which preserves a properly convex open set $\Omega \subset \mathrm{P}(V)$, then

$$\delta_\Gamma = \limsup_{r \rightarrow \infty} \frac{1}{r} \log(\#\{\gamma \in \Gamma : \kappa(\gamma) \leq r\}), \quad (2.3)$$

and

$$\Gamma \text{ is divergent if and only if } \sum_{\gamma \in \Gamma} e^{-\delta_\Gamma \kappa(\gamma)} = \infty. \quad (2.4)$$

If Γ does not preserve any properly convex open set, then we take (2.3) and (2.4) as definitions.

We end this section with the following classical fact, which is a consequence for instance of the Tits alternative and of the sub-additivity of κ . It applies to all strongly irreducible discrete subgroups of $\mathrm{PGL}(V)$, and non-elementary rank-one discrete groups of properly convex open sets.

Fact 2.7. Let $\Gamma \subset \mathrm{PGL}(V)$ be a discrete subgroup that is not virtually solvable. Then $\delta_\Gamma > 0$.

2.6. Proximal linear transformations

In this section we recall the notion of a proximal linear transformation, which was used in the definition of the proximal limit set Λ^{prox} and the biproximal unit tangent bundle $T^1 M_{\mathrm{bip}}$.

Notation 2. If W_1 and W_2 are two subspaces of V such that $W_1 \cap W_2 = \{0\}$, we write $W_1 \oplus W_2 \subset V$ for their direct sum and $\mathrm{P}(W_1) \oplus \mathrm{P}(W_2) = \mathrm{P}(W_1 \oplus W_2)$ for its projectivisation. In particular, if $x, y \in \mathrm{P}(V)$ are two distinct points, we write $x \oplus y$ for the projective line through x and y .

Definition 2.8. A linear transformation $g \in \mathrm{End}(V)$ is *proximal* if it has exactly one complex eigenvalue with maximal modulus among all eigenvalues, and if this eigenvalue has multiplicity 1. The associated eigenline in $\mathrm{P}(V)$ is the attracting fixed point of g and is denoted by x_g^+ .

An invertible linear transformation $g \in \mathrm{GL}(V)$ is said to be *biproximal* if g and g^{-1} are proximal. The attracting fixed point of g^{-1} is the repelling fixed point of g and is denoted by x_g^- . The projective line $x_g^+ \oplus x_g^-$ (see Notation 2) is the *axis* of g and is denoted by $\mathrm{Axis}(g)$. The g -invariant complementary subspace to the axis of g is denoted by x_g^0 . Note that the notions of biproximality, attracting/repelling fixed point, and axis, are well defined for the image of g in $\mathrm{PGL}(V)$.

More generally for any projective transformation $g \in \mathrm{PGL}(V)$, one can define the attracting subspace x_g^+ of g as the span of all generalised eigenspaces associated to eigenvalues with maximal norm, and x_g^0 and x_g^- can be defined similarly.

Remark 2.9. The set of proximal linear transformations is open in $\mathrm{End}(V)$, and the map sending a proximal linear transformation to the pair (attracting fixed point, maximal eigenvalue) is continuous.

Remark 2.10. As observed by Guivarc’h [39, Thm. 1] (see also [6, Lem. 3.6.ii]), for any irreducible subgroup $\Gamma \subset \mathrm{PGL}(V)$ which contains a proximal element, the proximal limit set is the smallest closed Γ -invariant non-empty subset of $\mathrm{P}(V)$; in particular, the action of Γ on Λ^{prox} is minimal (i.e. any orbit is dense). Indeed, consider any proximal element $\gamma \in \Gamma$, and let $\mathrm{P}(W) \subset \mathrm{P}(V)$ be the γ -invariant complementary subspace to x_γ^+ . By irreducibility, any closed Γ -invariant non-empty subset $X \subset \mathrm{P}(V)$ contains a point x outside $\mathrm{P}(W)$, and then x_γ^+ , which is the limit of the sequence $(\gamma^n x)_{n \in \mathbb{N}}$, belongs to X .

2.7. Periodic straight geodesics and elements of Γ

In this section we recall the link between periodic geodesics in $T^1\Omega/\Gamma$ and conjugacy classes of Γ . Let $\Omega \subset \mathrm{P}(V)$ be a properly convex open set. Let $g \in \mathrm{Aut}(\Omega)$. Then

$$\ell(g) = \inf\{d_\Omega(x, g \cdot x) : x \in \Omega\} \geq 0, \quad (2.5)$$

where $\ell(g)$ was defined in (1.1). The right-hand side of (2.5) is called the *translation length* of g . See [27, Prop. 2.1] for a proof.

Combined with an elementary computation, (2.5) yields:

Fact 2.11. Let $\Omega \subset \mathrm{P}(V)$ be a properly convex open set and $\Gamma \subset \mathrm{Aut}(\Omega)$ a discrete subgroup; denote Ω/Γ by M . Then for any lift in Ω of any periodic straight geodesic of M , there is an automorphism $\gamma \in \Gamma$ which preserves it and acts by positive translation on it. Let $\tilde{\gamma} \in \mathrm{GL}(V)$ be a lift of γ . The endpoints in $\partial\Omega$ of the geodesic are fixed by γ , the associated eigenvalues of $\tilde{\gamma}$ are $\lambda_1(\tilde{\gamma})$ and $\lambda_{d+1}(\tilde{\gamma})$, and the length of the geodesic in M is the translation length of γ . If furthermore these endpoints are extremal (for example if γ is rank-one, see Definition 1.1), then γ is biproximal.

Definition 2.12. Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Let $\gamma \in \Gamma$ be a biproximal element whose axis meets Ω . Then the periodic geodesic associated to γ is said to be *biproximal*, and the vectors along this geodesic are said to be *biproximal periodic*.

There are cases where $\gamma \in \Gamma$ is biproximal but its axis does not intersect Ω (e.g. when Ω is a triangle, or is higher-rank symmetric). Then we cannot make sense of a straight periodic geodesic associated to γ .

2.8. Biproximal vs rank-one periodic geodesics

In this section we explain when a biproximal periodic geodesic is rank-one. We saw in Fact 2.11 that a rank-one periodic geodesic is always biproximal. Conversely, the end-points of a biproximal periodic geodesic are smooth, but not always strongly extremal.

Fact 2.13 ([14, Lem. 3.1]). Let $\Omega \subset P(V)$ be a properly convex open set. Let g be a biproximal automorphism of Ω . Then $\text{Axis}(g) \cap \Omega$ is non-empty if and only if x_g^+ is smooth; in this case $T_{x_g^+} \partial\Omega = x_g^+ \oplus x_g^0$.

The following result is later completed by Corollary 6.9.

Fact 2.14 ([14, Lem. 3.2], [40, Prop. 6.3]). Let $\Omega \subset P(V)$ be a properly convex open set. Let $g \in \text{PGL}(V)$ be a biproximal automorphism of Ω . Then the following are equivalent:

- (a) g is rank-one;
- (b) $x_g^+, x_g^- \in \partial\Omega$ are smooth and strongly extremal points;
- (c) x_g^+ is strongly extremal;
- (d) the element g seen as an automorphism of Ω^* is rank-one;
- (e) the axis of g in $P(V)$ intersects Ω , and the axis of g in $P(V^*)$ intersects Ω^* ;
- (f) $d_{\text{spl}}(x_g^+, x_g^-) \geq 3$.

Proof. In [14], the point (g) was missing, so we briefly check that it is equivalent to the other points.

- (c) implies (g) because it implies that $d_{\text{spl}}(x_g^+, x_g^-) = \infty$.
- If $d_{\text{spl}}(x_g^+, x_g^-) \geq 3$, then the axis of g in $P(V)$ intersects Ω so x_g^+ is smooth. Furthermore, x_g^0 does intersect $\partial\Omega$, because otherwise $d_{\text{spl}}(x_g^+, x_g^-) \leq 2$, since $x_g^+ \oplus x_g^0$ and $x_g^- \oplus x_g^0$ are supporting hyperplanes of Ω . ■

These characterisations of rank-one automorphisms yield a characterisation of rank-one manifolds. To see this we need the following fact.

Fact 2.15 ([7, Prop. 1.1] and [6, Lem. 3.6.iv]). Let $\Gamma \subset \mathrm{PGL}(V)$ be a strongly irreducible subgroup.

- (1) If Γ preserves a properly convex open set $\Omega \subset \mathbb{P}(V)$, then it contains a proximal element.
- (2) If Γ contains a proximal element, then $\{(x_\gamma^+, x_\gamma^-) : \gamma \in \Gamma \text{ biproximal}\}$ is dense in $\Lambda^{\mathrm{prox}} \times \Lambda^{\mathrm{prox}}$.

Corollary 2.16. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \mathrm{Aut}(\Omega)$ a strongly irreducible discrete subgroup. Then $M = \Omega/\Gamma$ is rank-one if and only if Λ^{prox} contains two points at simplicial distance at least 3.*

Proof. If M is rank-one, then Λ^{prox} contains two strongly extremal points by definition. Conversely, if there exists $\xi, \eta \in \Lambda^{\mathrm{prox}}$ at simplicial distance at least 3. By lower semi-continuity of d_{spl} and by Fact 2.15 (2), there exists $\gamma \in \Gamma$ such that

$$d_{\mathrm{spl}}(x_\gamma^+, x_\gamma^-) \geq 3.$$

By Fact 2.14 (f), γ is rank-one and M as well. ■

Example 2.17 below is an elementary application of the previous result. It can also be seen as a consequence of Zimmer’s higher-rank rigidity [64, Thm. 1.4].

Example 2.17. Any 3-dimensional irreducible compact convex projective manifold is rank-one.

Proof. Let $M = \Omega/\Gamma$ be an irreducible compact convex projective manifold of dimension 3. By [10, Prop. 3.10.a], $\Lambda^{\mathrm{prox}} = \partial\Omega$, and by [10, Thm. 1.1.d and Prop. 3.8], $\partial\Omega \setminus \partial_{\mathrm{sse}}\Omega$ is contained in the union of countably many properly embedded triangles (PET) (namely PES’s of dimension 2) of Ω . These cannot cover the whole boundary $\partial\Omega$ (for instance, because they have Lebesgue measure zero), hence there is a point in $\partial_{\mathrm{sse}}\Omega = \partial_{\mathrm{sse}}\Omega \cap \Lambda^{\mathrm{prox}}$. By irreducibility, there is another one, and we can conclude using Corollary 2.16. ■

Remark 2.10 and Fact 2.15 are useful results, and we will need them to hold for certain discrete groups $\Gamma \subset \mathrm{PGL}(V)$ that are not necessarily irreducible. The following fact is exactly what will be needed. Its proof is quite similar to the proof of Fact 2.15, yet somehow easier, because rank-one elements are easier to manipulate than general biproximal elements (for instance for rank-one groups the existence of an invariant convex set make things easier to visualise, while Fact 2.15 holds for groups that do not necessarily preserve a convex set).

Fact 2.18 ([17, Prop. 2.4]). Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a non-elementary rank-one discrete subgroup. Then $\Lambda^{\text{prox}} \subset \bar{\Omega}$ is the smallest Γ -invariant closed subset, and it has no isolated point. Furthermore,

$$\{(x_\gamma^+, x_\gamma^-) \in \Lambda^{\text{prox}} \times \Lambda^{\text{prox}} : \gamma \in \Gamma \text{ rank-one}\}$$

is dense in $\Lambda^{\text{prox}} \times \Lambda^{\text{prox}}$.

Fact 2.18 can also be seen as a consequence of [15, Lem. 3.1.3.2, Prop. 3.2.2, Lem. 3.2.4, Prop. 5.3.1].

2.9. Horoboundaries and Patterson–Sullivan densities

In this section we recall the classical definitions of horofunctions and horoboundary, and how they can be used to define Patterson–Sullivan densities.

Definition 2.19. Let (X, d) be a proper metric space. The *horofunction* at points $x, y, z \in X$ is defined as follows:

$$b_z(x, y) = d(x, z) - d(y, z).$$

We now recall the definition of the horocompactification. By a *compactification* of a topological space X we mean a compact topological space Y together with an embedding $X \hookrightarrow Y$ with open and dense image; then the subset $Y \setminus X = \partial_Y X$ is called the boundary of the compactification. Another compactification Z *dominates* Y if there is a continuous map from Z to Y which is compatible with the embeddings of X . Using the Arzelà–Ascoli theorem, one can show that the following is well defined.

Definition 2.20 ([38, §3]). Let (X, d) be a proper metric space. The *horocompactification* of (X, d) is the smallest compactification \bar{X}^{h} of X such that $z \mapsto b_z(x, y)$ extends continuously to \bar{X}^{h} for every $x, y \in X$. The *horoboundary* $\partial_{\text{h}} X$ of (X, d) is the boundary of \bar{X}^{h} .

Note that \bar{X}^{h} is metrisable and the function $(\xi, x, y) \in \bar{X}^{\text{h}} \times X \times X \mapsto b_\xi(x, y)$ is continuous. Any isometry of X extends continuously to a bi-Lipschitz homeomorphism of \bar{X}^{h} .

We now recall the definition of Patterson–Sullivan measures on the horoboundary of a proper metric space.

Definition 2.21. Let (X, d) be a proper metric space. Let $\Gamma < \text{Isom}(X, d)$ be a discrete subgroup. Given $\delta \in \mathbb{R}$, a $(\Gamma$ -equivariant) δ -*conformal density* on $\partial_{\text{h}} X$ is a family of finite measures $(\mu_x)_{x \in X}$ on $\partial_{\text{h}} X$ such that

- μ_y is absolutely continuous with respect to μ_x for all $x, y \in X$, and the Radon–Nikodym derivative is:

$$\frac{d\mu_y}{d\mu_x}(\xi) = e^{-\delta b_\xi(y,x)};$$

(This implies that the family is entirely determined by μ_o for any $o \in X$.)

- for every $\gamma \in \Gamma$ and $x \in X$, the push-forward by γ of μ_x is:

$$\gamma_*\mu_x = \mu_{\gamma x}.$$

Let us recall the classical example of a conformal density, which we will need in this paper. For any measured metric space (X, d, μ) with infinite mass, the *volume entropy* is, for any $o \in X$,

$$\delta_\mu := \limsup_{r \rightarrow \infty} \frac{\log \mu(B_X(o, r))}{r} \in \mathbb{R}_{\geq 0} \cup \{\infty\}.$$

Fact 2.22 ([53, §3]). Let (X, d) be a proper metric space, let $o \in X$ be a basepoint, let Γ be a non-compact closed subgroup of $\text{Isom}(X, d)$ and let Vol be a Γ -invariant Radon measure on X . We assume that the volume entropy δ_{Vol} is finite. Then there exists a continuous non-decreasing function $h: \mathbb{R}_+ \rightarrow \mathbb{R}_{>0}$ such that

- $\int_{x \in X} h(d(o, x)) e^{-\delta_{\text{Vol}} d(o, x)} d\text{Vol}(x) = \infty$;
- for every $\varepsilon > 0$, there exists $R > 0$ such that $h(r + t) \leq e^{\varepsilon t} h(r)$ for any $r \geq R$ and $t \geq 0$.

Furthermore, if we define, for $s > \delta_{\text{Vol}}$, the probability measure $\mu_{o,s}$ on X such that

$$\mu_{o,s}(A) = \frac{\int_{x \in A} h(d(o, x)) e^{-sd(o, x)} d\text{Vol}(x)}{\int_{x \in X} h(d(o, x)) e^{-sd(o, x)} d\text{Vol}(x)},$$

for any Borel subset $A \subset X$, then any accumulation point of $(\mu_{o,s})_{s \rightarrow \delta_{\text{Vol}}}$ in the space $\mathcal{P}(\bar{X}^h)$ of probability measures on \bar{X}^h is supported on $\partial_h \Omega$ and is a δ_{Vol} -conformal density.

A typical example of Γ -invariant Radon measure on X is the push-forward by an orbital map of the Haar measure on Γ .

Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set. One can check that the critical exponent δ_Γ of any discrete group $\Gamma \subset \text{Aut}(\Omega)$ is equal to the volume entropy of the push-forward by any orbital map of the counting measure on Γ , and also that $\delta_\Gamma \leq \delta_{\text{Vol}\Omega}$.

Fact 2.23 ([59, Thm. 2]). Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set. Then $\delta_{\text{Vol}\Omega} \leq \dim(V) - 2$. In particular, for any discrete subgroup $\Gamma \subset \text{Aut}(\Omega)$, the numbers δ_Γ and $\delta_{\text{Vol}\Omega}$ are finite.

This will allow us to apply Fact 2.22 to our convex projective setting.

2.10. The horoboundary of a properly convex open set

In this section we recall results on the horoboundary of a properly convex open set that were mentioned in the introduction. A direct and elementary proof of the following result may be found in [15, Fact 6.1.2].

Fact 2.24 ([62, Thm. 1.3]). Let $\Omega \subset P(V)$ be a properly convex open set. Then the horocompactification $\bar{\Omega}^h$ dominates the projective compactification $\bar{\Omega}$.

Notation 3. Given a properly convex open set $\Omega \subset P(V)$, we denote by π_h the map $\partial_h \Omega \rightarrow \partial \Omega$.

Definition 2.25. Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. For $\delta \geq 0$, a δ -conformal density on $\partial \Omega$ is the push-forward by π_h of a δ -conformal density on $\partial_h \Omega$. Note that such a family is made of Γ -quasi-invariant measures.

If it was not true that $\partial_h \Omega$ dominates $\partial \Omega$, we could have considered a common refinement of $\partial_h \Omega$ and $\partial \Omega$, where the conformal densities are also well defined.

Thanks to the following fact, we will from now on abusively identify any smooth point on the projective boundary with its preimage in the horoboundary.

Fact 2.26 ([21, Lem. 3.2]). Let $\Omega \subset P(V)$ be a properly convex open set. Let ξ be a smooth point of $\partial \Omega$. Then it has only one preimage by $\pi_h: \partial_h \Omega \rightarrow \partial \Omega$.

2.11. The Hopf decomposition and quotients of measures

Let us fix for the whole section a locally compact, second countable and unimodular group G acting measurably on a standard Borel space X , and a G -invariant and σ -finite measure \tilde{m} on X ; fix a Haar measure on G , denoted by dg , and an integrable positive function σ on X . For any non-negative measurable function f on X , we denote by $\int_G f$ the G -invariant measurable function defined by

$$\int_G f(x) = \int_G f(gx) dg;$$

this also denotes the induced function on $G \backslash X$.

A measurable subset $A \subset X$ is said to *wandering* under the action of G if for \tilde{m} -almost any $x \in A$, the *transporter* $T(x, A) := \{g \in G : gx \in A\}$ is relatively compact. The following fact is classical, and serves as a definition of the Hopf decomposition.

Fact 2.27. Let $\mathcal{C} := \{\int_G \sigma = \infty\}$ and $\mathcal{D} := \{\int_G \sigma < \infty\} \subset X$. The decomposition $X = \mathcal{C} \sqcup \mathcal{D}$ is a *Hopf decomposition*, in the sense that every wandering subset of \mathcal{C}

has \tilde{m} -measure zero, and \mathcal{D} is a countable union of wandering subsets of X . Any two Hopf decompositions agree on some \tilde{m} -full subset of X . The dynamical system (X, G, \tilde{m}) is said to be *conservative* (resp. *dissipative*) if $\tilde{m}(\mathcal{D}) = 0$ (resp. $\tilde{m}(\mathcal{C}) = 0$).

Proof. Let $A \subset \mathcal{C}$ be a measurable subset, and let us prove that $\int_G 1_A$ is infinite on \tilde{m} -almost every point of A . On the one hand, if for $R > 0$, we denote

$$A_R := \left\{ \int_G 1_A \leq R \right\} \cap A,$$

then

$$\int_{X \times G} \sigma(gx) 1_{A_R}(x) \, dg \, d\tilde{m}(x) = \infty \cdot \tilde{m}(A_R),$$

while on the other hand, since G is unimodular and \tilde{m} is G -invariant,

$$\begin{aligned} \int_{X \times G} \sigma(gx) 1_{A_R}(x) \, dg \, d\tilde{m}(x) &= \int_{X \times G} \sigma(x) 1_{A_R}(gx) \, dg \, d\tilde{m}(x) \\ &= \int_X \sigma \left(\int_G 1_{A_R} \right) d\tilde{m} \leq R \int_X \sigma \, d\tilde{m} < \infty. \end{aligned}$$

Therefore, $\tilde{m}(A_R) = 0$ for any $R > 0$, hence $\tilde{m}(\cup_R A_R) = 0$. For any compact subset $K \subset G$, we consider

$$B_K := \left\{ \int_K \sigma > \frac{1}{2} \int_G \sigma \right\} \subset \mathcal{D},$$

and observe that it is wandering. Indeed if $x \in B_K$ and $g \in G$ are such that $gx \in B_K$, then

$$\int_K \sigma(x) + \int_{Kg} \sigma(x) > \int_G \sigma(x),$$

thus $K \cap Kg \neq \emptyset$ and $g \in K^{-1} \cdot K$, which is compact. Furthermore, $\mathcal{D} = \cup_n B_{K_n}$ if $G = \cup_n K_n$. \blacksquare

Note that if the action of G on X is *smooth* (namely $G \backslash X$ is a standard Borel space) and has compact stabilisers, then (X, G, \tilde{m}) is dissipative. In particular, this observation applies when X is a locally compact second countable topological space and the action of G is continuous and proper.

Definition 2.28. If (X, G, \tilde{m}) is dissipative, the *quotient* of \tilde{m} on $G \backslash X$ is defined as

$$m := \left(\int_G \sigma \right)^{-1} \pi_*(\sigma \tilde{m}),$$

where π denotes the projection $X \rightarrow G \backslash X$. For any non-negative (or integrable) function f on X ,

$$\int_X f \, d\tilde{m} = \int_{G \backslash X} \left(\int_G f \right) dm,$$

and m is independent of the choice of σ (but it depends on the choice of dg).

Fact 2.29. Suppose G contains a unimodular, normal and closed subgroup H whose action on X is dissipative and smooth; fix a Haar measure on H . Set $\pi: X \rightarrow H \backslash X$ and m to be the quotient of \tilde{m} on $H \backslash X$; this measure is G/H -invariant. Then the Hopf decomposition of X projects under π onto the Hopf decomposition of $H \backslash X$. In particular, (X, G, \tilde{m}) is conservative (resp. dissipative) if and only if $(H \backslash X, G/H, m)$ is conservative (resp. dissipative).

Proof. Observe that the Haar measure on G/H is the quotient of the Haar measure on G by the action of H . Therefore, if σ is a positive integrable function on X , then $\int_H \sigma$ is a positive integrable function on $H \backslash X$, and

$$\int_G \sigma = \int_{G/H} \int_H \sigma.$$

Moreover, for any G -invariant measurable subset $\tilde{A} \subset X$ whose projection A in $H \backslash X$ is measurable,

$$\tilde{m}(\tilde{A}) = \int_{H \backslash X} \left(\int_H 1_{\tilde{A}} \right) dm = \int_{H \backslash X} \| \text{Haar}_H \| 1_A dm$$

is zero if and only if $m(A) = 0$ (with the convention $\infty \cdot 0 = 0$). ■

Recall that if the action of G is continuous, then a point $x \in X$ is said to be *recurrent* if for any neighbourhood U of x , the set $\{g \in G : gx \in U\}$ is not relatively compact; if $G = \mathbb{R}$, then x is call *forward recurrent* (resp. *backward recurrent*) if $\{t > 0 : \phi_t y \in U\} \subset \mathbb{R}$ (resp. $\{t < 0 : \phi_t y \in U\} \subset \mathbb{R}$) is unbounded. The following fact is classical and will be used in Section 6.3.

Fact 2.30. Assume that X is a locally compact topological space with countable basis, and that the action of G is continuous.

- (1) If \tilde{m} is conservative, then \tilde{m} -almost all points are recurrent.
- (2) If \tilde{m} is conservative and $G = \mathbb{R}$, then m -almost all points are forward and backward recurrent.
- (3) If \tilde{m} is ergodic, then \tilde{m} -almost all points have a dense G -orbit in $\text{supp}(\tilde{m})$.
- (4) If \tilde{m} is ergodic and conservative, and $G = \mathbb{R}$, then \tilde{m} -almost all points have a dense forward orbit and a dense backward orbit in $\text{supp}(\tilde{m})$.

2.12. Criteria for ergodicity and for mixing

Inspired by Babillot's proof of mixing [3], Coudène stated and proved a criterion for the ergodicity property, and another for the mixing property, both based on the following definition.

Definition 2.31. Consider a Borel flow $(\phi_t)_{t \in \mathbb{R}}$ on a metric space (X, d) . For each point $x \in X$, we define the *strong stable manifold*

$$W^{ss}(x) = \{y \in X : d(\phi_t x, \phi_t y) \xrightarrow{t \rightarrow \infty} 0\}.$$

We define the *strong unstable manifold* $W^{us}(x)$ to be the strong stable manifold of the time-reversed flow. Consider a $(\phi_t)_{t \in \mathbb{R}}$ -invariant σ -finite measure on X . A measurable function $f: X \rightarrow \mathbb{R}$ is said to be *W^{ss} -invariant* when there exists a measurable subset $E \subset X$ with full measure such that for all x and y in E , if they are on the same strong stable manifold then $f(x) = f(y)$. The notion of *W^{su} -invariance* is similarly defined.

Coudène's criteria are the following.

Fact 2.32 ([29]). Let (X, d) be a metric space, $(\phi_t)_t$ a measurable flow on it, m a conservative $(\phi_t)_t$ -invariant measure, and assume that some m -full subset of X is covered by a countable family of open sets with finite m -measure. If every W^{ss} , W^{su} and $(\phi_t)_t$ -invariant measurable function is essentially constant then the flow is ergodic.

Fact 2.33 ([28]). Consider a Borel flow preserving a finite measure on a metric space. If every W^{ss} - and W^{su} -invariant Borel function is essentially constant then the flow is mixing.

2.13. Topological entropy

In this section we recall the definition of topological entropy.

Definition 2.34. Let $\phi = (\phi_t)_{t \in \mathbb{R}}$ be a continuous flow on a compact metric space (X, d) .

- Let $\varepsilon > 0$. A subset $S \subset X$ is *(d, ε) -separated* if $d(s, s') \geq \varepsilon$ for all $s \neq s'$ in S . We denote by $N(d, \varepsilon)$ the maximal cardinality of such a set S .
- Let $\varepsilon > 0$. A subset $S \subset X$ is *(d, ε) -spanning* if for any $x \in X$, there exists $s \in S$ with $d(x, s) < \varepsilon$. We denote by $S(d, \varepsilon)$ the minimal cardinality of such a set S .
- We take the classical notation $d^{(t)}(x, y) := \max_{0 \leq s \leq t} d(\phi_s x, \phi_s y)$ for $t \geq 0$ and $x, y \in X$; this defines a family of metrics on X .
- The *topological entropy* of ϕ on X is:

$$h_{\text{top}}(\phi) := \lim_{\varepsilon \rightarrow 0} \limsup_{t \rightarrow \infty} \frac{\log N(d^{(t)}, \varepsilon)}{t} = \lim_{\varepsilon \rightarrow 0} \limsup_{t \rightarrow \infty} \frac{\log S(d^{(t)}, \varepsilon)}{t}.$$

One defines similarly the topological entropy of a homeomorphism by replacing $t \in \mathbb{R}_{\geq 0}$ by $n \in \mathbb{Z}_{\geq 0}$ in Definition 2.34. Note that if $\phi = (\phi_t)_{t \in \mathbb{R}}$ is a continuous

flow on a compact metric space (X, d) , then the topological entropies of the flow and of the underlying time-one map are the same. Also note that the reparametrised flow $(\phi_{kt})_{t \in \mathbb{R}}$, where $k \in \mathbb{R}_{\neq 0}$, has topological entropy equal to $|k|$ times the topological entropy of $(\phi_t)_{t \in \mathbb{R}}$.

Remark 2.35. Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup; set $M = \Omega / \Gamma$. Then by definition, for any $t \geq 0$, for all $v, w \in T^1M$ and all possible lifts $\tilde{v}, \tilde{w} \in T^1\Omega$,

$$\begin{aligned} d_{T^1M}^{(t)}(v, w) &= \max_{0 \leq s \leq t+1} d_M(\pi\phi_s v, \pi\phi_s w) \\ &\leq d_{T^1\Omega}^{(t)}(\tilde{v}, \tilde{w}) = \max_{0 \leq s \leq t+1} d_\Omega(\pi\phi_s \tilde{v}, \pi\phi_s \tilde{w}). \end{aligned}$$

Yet, we will sometimes prefer to work with dynamical balls in the universal cover, and this is why we introduce another notation (which is a bit sloppy unfortunately). For $t \geq 0$ and $v, w \in T^1M$,

$$\tilde{d}_{T^1M}^{(t)}(v, w) := \min\{d_{T^1\Omega}^{(t)}(\tilde{v}, \tilde{w}) : \tilde{v}, \tilde{w} \in T^1\Omega \text{ lifts of } v, w\},$$

and we denote by $\tilde{B}_{T^1M}^{(t)}(v, R)$ the open ball of radius $R \geq 0$ and centred at $v \in T^1M$ for the metric $\tilde{d}_{T^1M}^{(t)}$. Observe that $\tilde{d}_{T^1M}^{(t)} \geq d_{T^1M}^{(t)}$ and that we do not have equality in general; however, we have equality on

$$\left\{d_{T^1M}^{(t)} < \frac{\text{inj}(M)}{2}\right\},$$

where $\text{inj}(M)$ is the injectivity radius of M , namely

$$\text{inj}(M) := \inf\{d_\Omega(x, \gamma x) : x \in \Omega, \gamma \in \Gamma \setminus \{\text{id}\}\},$$

To conclude this section, we give a relation between the critical exponent, the volume entropy and the topological entropy, which is originally due to Manning [49]. Our setting is not exactly the same as his, but the proof works perfectly thanks to Fact 2.1 and Selberg's lemma.

Fact 2.36. Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact, strongly irreducible and discrete subgroup. Let Vol be a Γ -invariant Radon measure on $\mathcal{C}_\Omega^{\text{cor}}(\Gamma)$. Then

$$\delta_\Gamma = \delta_{\text{Vol}} \geq h_{\text{top}}(T^1M_{\text{cor}}, (\phi_t)_{t \in \mathbb{R}}) \geq h_{\text{top}}(T^1M_{\text{bip}}, (\phi_t)_{t \in \mathbb{R}}).$$

Proof. The fact that $h_{\text{top}}(T^1M_{\text{cor}}, (\phi_t)_{t \in \mathbb{R}}) \geq h_{\text{top}}(T^1M_{\text{bip}}, (\phi_t)_{t \in \mathbb{R}})$ is immediate from the definition of topological entropy.

Let us prove that $\delta_\Gamma = \delta_{\text{Vol}}$. Consider $o \in \mathcal{C}^{\text{cor}}$ and $R > 0$ large enough such that $\Gamma \cdot B_\Omega(o, R)$ contains \mathcal{C}^{cor} (which is acted on cocompactly by Γ). Then, on the one hand,

$$\begin{aligned} \text{Vol}(B_\Omega(o, r)) &\leq \sum_{\gamma: d_\Omega(o, \gamma o) \leq r+R} \text{Vol}(B_\Omega(\gamma o, R)) \\ &\leq \text{Vol}(B_\Omega(o, R)) \#\{\gamma : d_\Omega(o, \gamma o) \leq r + R\} \end{aligned}$$

for any $r > 0$, hence $\delta_{\text{Vol}} \leq \delta_\Gamma$, and $\text{Vol}(B_\Omega(o, R)) > 0$.

On the other hand, consider

$$C = \#\{\gamma : d_\Omega(o, \gamma o) \leq 2R\},$$

so that for any $r > 0$, one can find a subset A of $\{\gamma : d_\Omega(o, \gamma o) \leq r\}$ with size greater than $C^{-1}\#\{\gamma : d_\Omega(o, \gamma o) \leq r\} - 1$ and whose image by the orbital map $\gamma \mapsto \gamma o$ is $(2R, d_\Omega)$ -separated; then

$$\begin{aligned} C^{-1}\#\{\gamma : d_\Omega(o, \gamma o) \leq r\} - 1 &\leq \text{Vol}(B_\Omega(o, R))^{-1} \sum_{\gamma \in A} \text{Vol}(B_\Omega(\gamma o, R)) \\ &\leq \text{Vol}(B_\Omega(o, R))^{-1} \text{Vol}(B_\Omega(o, r + R)), \end{aligned}$$

therefore $\delta_\Gamma \leq \delta_{\text{Vol}}$.

Let us prove that $\delta_\Gamma \geq h_{\text{top}}(T^1 M_{\text{cor}}, (\phi_t)_{t \in \mathbb{R}})$. Fix $\varepsilon > 0$, and fix a maximal $(\varepsilon/4, d_\Omega)$ -spanning set A of $B_\Omega(o, R)$. Let B_r be the set of vectors based at a point of A and pointing at a point of

$$\{\gamma a : a \in A, d_\Omega(o, \gamma o) \leq r + 2R\};$$

for each $\tilde{v} \in B_r$ choose $v' \in T^1 M_{\text{cor}}$ at distance less than $\varepsilon/2$ from the projection in $T^1 M$ of \tilde{v} (when such a vector exists), and denote by B'_r the collection of v' . By Fact 2.1, B'_r is a $(\varepsilon, d_{T^1 M}^{(r)})$ -spanning set of $T^1 M_{\text{cor}}$ for any $r > 0$. Furthermore,

$$\#B'_r \leq \#B_r \leq \#A^2 \cdot \#\{\gamma : d_\Omega(o, \gamma o) \leq r + 2R\},$$

hence $\delta_\Gamma \geq h_{\text{top}}(T^1 M_{\text{cor}}, (\phi_t)_{t \in \mathbb{R}})$. ■

We will see in Proposition 8.4 that $h_{\text{top}}(T^1 M_{\text{cor}}, (\phi_t)_{t \in \mathbb{R}}) = \delta_\Gamma$.

2.14. Measure-theoretic entropy

Definition 2.37. Let f be an invertible measurable map of a measurable space X , which preserves a probability measure μ . Let \mathcal{P} be a finite measurable partition of X .

- The *entropy* of \mathcal{P} is

$$H_\mu(\mathcal{P}) := - \sum_{P \in \mathcal{P}} \mu(P) \log \mu(P).$$

- For each integer $n \geq 1$, we set

$$\mathcal{P}^{(n)} := \{P_0 \cap f^{-1}P_1 \cap \dots \cap f^{-n+1}P_{n-1} : P_k \in \mathcal{P}, 0 \leq k \leq n-1\}.$$

- The entropy of f with respect to μ and \mathcal{P} is

$$H_\mu(f, \mathcal{P}) := \lim_{n \rightarrow \infty} \frac{H_\mu(\mathcal{P}^{(n)})}{n}.$$

- The entropy of f with respect to μ is

$$h_\mu(f) := \sup\{H_\mu(f, \mathcal{Q}) : \mathcal{Q} \text{ measurable partition}\}.$$

Let us now consider a measurable flow $(\phi_t)_{t \in \mathbb{R}}$ on a measurable space X preserving a probability measure μ . The entropy of the flow is defined to be the entropy of the time-one map ϕ_1 .

Remark 2.38. The reparametrised flow $(\phi_{kt})_{t \in \mathbb{R}}$ has measure-theoretic entropy equal to $|k|$ times the measure-theoretic entropy of $(\phi_t)_{t \in \mathbb{R}}$.

The relation between topological and measure-theoretic entropies is given by the following famous principle; for more details, see [43, Thm. 4.5.3].

Fact 2.39 (Variational principle). Let $\phi = (\phi_t)_{t \in \mathbb{R}}$ a continuous flow on a compact metric space (X, d) . Denote by $\mathcal{P}^\phi(X)$ the set of ϕ -invariant probability measures on X . Then

$$h_{\text{top}}(\phi) = \sup_{\mu \in \mathcal{P}^\phi(X)} h_\mu(\phi).$$

We recall the definition of entropy-expansive maps from Bowen [20].

Definition 2.40. Consider a homeomorphism f of a metric space (X, d) and $\varepsilon > 0$. The map f is said to be (d, ε) -entropy-expansive if for each $x \in X$, the action of f on the *Bowen ball*

$$Z_\varepsilon(x) := \{y \in X : \forall n \in \mathbb{Z}, d(f^n x, f^n y) \leq \varepsilon\}$$

has zero entropy.

Remark 2.41. We note that if f is (d, ε) -entropy-expansive, then f^n is $(d^{(n)}, \varepsilon)$ -entropy-expansive.

The following is one of the main properties of entropy-expansive maps.

Fact 2.42 ([20, Thm. 3.5]). Let ε be a positive number; let f be a ε -entropy-expansive homeomorphism of a metric space (X, d) ; let μ be a f -invariant probability measure on X ; let \mathcal{P} be a finite measurable partition all of whose elements have diameter less than ε . Then

$$h_\mu(f) = H_\mu(f, \mathcal{P}).$$

Remark 2.43. Let $\varepsilon > 0$; let f be an ε -entropy-expansive homeomorphism of a metric space (X, d) . Fact 2.42 tells us that the measure-theoretic entropy depends upper semi-continuously on the measure (with respect to the weak* topology), and that there exists a measure of maximal entropy if X is compact.

The geodesic flow on a convex projective manifold is entropy-expansive if the injectivity radius is non-zero.

Fact 2.44 ([22, Thm. 6.2]). Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Let us assume that the injectivity radius ε_0 of $M = \Omega/\Gamma$ is non-zero. Then the time-one map of the geodesic flow on T^1M is $(d_{T^1M}, \varepsilon_0/3)$ -entropy-expansive.

It applies in particular if Γ acts convex cocompactly on Ω and if Γ is torsion-free. However, we will consider cases where Γ is not torsion-free, and the geodesic flow on T^1M is *not* entropy-expansive: for instance if Ω is the Poincaré disk and Γ is a cocompact triangle group. To overcome this issue, we will use Selberg’s lemma (if Γ acts convex cocompactly on Ω , then it is finitely generated; see [23, Thm. 8.10]) and the following elementary observation.

Observation 2.45. Let $(\phi_t)_t$ be a measurable flow on a measurable space X that preserves a probability measure μ , and G a finite group that acts measurably on X and commutes with $(\phi_t)_t$. We denote by $\pi: X \rightarrow X/G$ the natural projection, and $(\phi_t)_t$ the induced flow on the quotient. Then $h_\mu(X, (\phi_t)_t) = h_{\pi_*\mu}(X/G, (\phi_t)_t)$. In particular, if X is a compact topological space and the actions of $(\phi_t)_t$ and G are continuous, then $h_{\text{top}}(X, (\phi_t)_t) = h_{\text{top}}(X/G, (\phi_t)_t)$.

3. Construction of the Sullivan measures

3.1. The Gromov product

Let us define the Gromov product. It will be used to define the Sullivan measures. Recall that given a properly convex open set $\Omega \subset \mathbb{P}(V)$ with three points $x, \xi, \eta \in \Omega$, the Gromov product is defined by

$$2\langle \xi, \eta \rangle_x := d_\Omega(x, \xi) + d_\Omega(x, \eta) - d_\Omega(\xi, \eta) \geq 0.$$

The following are three immediate properties of the Gromov product. Let $y \in \Omega$:

$$x \in [\xi, \eta] \Rightarrow \langle \xi, \eta \rangle_x = 0; \quad (3.1)$$

$$2\langle \xi, \eta \rangle_x = 2\langle \xi, \eta \rangle_y + b_\xi(x, y) + b_\eta(x, y); \quad (3.2)$$

$$|\langle \xi, \eta \rangle_x - \langle \xi, \eta \rangle_y| \leq d_\Omega(x, y). \quad (3.3)$$

We denote by $\text{Geod}_h(\Omega)$ (resp. $\text{Geod}_h^\infty(\Omega)$) the set of pairs (ξ, η) in $(\bar{\Omega}^h)^2$ (resp. $\partial_h \Omega^2$) such that $[\xi, \eta] \cap \Omega \neq \emptyset$, where $[\xi, \eta] := [\pi_h(\xi), \pi_h(\eta)]$.

Proposition 3.1. *Let $\Omega \subset P(V)$ be a properly convex open set. The map*

$$(\xi, \eta, x) \mapsto \langle \xi, \eta \rangle_x$$

defined on Ω^3 extends continuously to $\text{Geod}_h(\Omega) \times \Omega$, as well as to (3.1), (3.2) and (3.3).

Proof. Let $A := \{(\xi, \eta, x, y) \in \text{Geod}_h(\Omega) \times \Omega^2 : y \in [\xi, \eta]\}$. The projection

$$(\xi, \eta, x, y) \mapsto (\xi, \eta, x)$$

from A to $\text{Geod}_h(\Omega) \times \Omega$ is continuous, surjective and open. Therefore, it is enough to prove that the function $(\xi, \eta, x, y) \mapsto \langle \xi, \eta \rangle_x$ defined on $A \cap \Omega^4$ extends continuously to A . This is a consequence of (3.1) and (3.2) combined, which yield, for any $(\xi, \eta, x, y) \in A \cap \Omega^4$,

$$\langle \xi, \eta \rangle_x = b_\xi(x, y) + b_\eta(x, y).$$

Note that (3.1) extends to $(\xi, \eta) \in \text{Geod}_h(\Omega)$ because the set $\{(\xi, \eta, x) \in \Omega^3 : x \in [\xi, \eta]\}$ is dense $\{(\xi, \eta, x) \in \text{Geod}_h(\Omega) \times \Omega : x \in [\xi, \eta]\}$. ■

3.2. The Hopf coordinates

Let us define the Hopf parametrisation of the unit tangent bundle $T^1\Omega$ of a properly convex open set Ω , which depends on the choice of a fixed basepoint $o \in \Omega$.

Definition 3.2. Let $\Omega \subset P(V)$ be a properly convex open set and fix a basepoint $o \in \Omega$. The Hopf parametrisation based at o is the $(\phi_t)_t$ -equivariant continuous surjective map

$$\text{Hopf}_o: \text{Geod}_h^\infty(\Omega) \times \mathbb{R} \rightarrow T^1\Omega,$$

that sends $(\xi, \eta, t) \in \text{Geod}_h^\infty(\Omega) \times \mathbb{R}$ to the vector $\text{Hopf}_o(\xi, \eta, t)$ which is tangent to the projective line $\pi_h \xi \oplus \pi_h \eta$ and such that $b_\eta(o, \pi \text{Hopf}_o(\xi, \eta, t)) = t$.

When the context is clear, we will simply write Hopf instead of Hopf_o . Note that if $x \in \Omega$ then for any $(\xi, \eta, t) \in \text{Geod}_h^\infty(\Omega) \times \mathbb{R}$,

$$\text{Hopf}_x(\xi, \eta, t) = \text{Hopf}_o(\xi, \eta, t + b_\eta(o, x)).$$

Therefore, if we consider the following action of $\text{Aut}(\Omega)$:

$$g \cdot (\xi, \eta, t) = (g\xi, g\eta, t + b_\eta(g^{-1}o, o))$$

for $g \in \text{Aut}(\Omega)$ and $(\xi, \eta, t) \in \text{Geod}_h^\infty(\Omega)$, then Hopf_o is $\text{Aut}(\Omega)$ -equivariant. Note also that for any $(\xi, \eta, t) \in \text{Geod}_h^\infty(\Omega)$,

$$-\text{Hopf}_o(\xi, \eta, t) = \text{Hopf}_o(\eta, \xi, \langle \xi, \eta \rangle_o - t), \quad (3.4)$$

where for any $v \in T^1\Omega$, the vector $-v$ satisfies $\phi_{\pm\infty}(-v) = \phi_{\mp\infty}v$ and $\pi(-v) = \pi v$.

3.3. The Sullivan measures

In this section, given a conformal density, we construct the associated $(\phi_t)_{t \in \mathbb{R}}$ -invariant Sullivan measures on $T^1\Omega$ and on the quotient $T^1M = T^1\Omega/\Gamma$.

Definition 3.3. Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a strongly irreducible discrete subgroup; denote $M = \Omega/\Gamma$. Let $\delta \geq 0$ and let $(\mu_x)_x$ be a δ -conformal density on $\partial_h\Omega$. The *Sullivan measure* \tilde{m}_h on $\partial_h\Omega^2 \times \mathbb{R}$ induced by $(\mu_x)_x$ is defined by the following:

$$d\tilde{m}_h(\xi, \eta, t) = e^{2\delta\langle \xi, \eta \rangle_o} 1_{\text{Geod}_h^\infty(\Omega)}(\xi, \eta) d\mu_o(\xi) d\mu_o(\eta) dt,$$

where $o \in \Omega$; the measure \tilde{m}_h does not depend on o and is $(\phi_t)_t$ -invariant. Then we push it forward via the Hopf coordinates to get the induced *Sullivan measure* on $T^1\Omega$:

$$\tilde{m} := (\text{Hopf}_o)_* \tilde{m}_h.$$

It does not depend on o , and it is Γ and $(\phi_t)_t$ -invariant. Hence, it yields an induced $(\phi_t)_t$ -invariant *Sullivan measure* on T^1M , denoted by m .

Observe that the formula $e^{2\delta\langle \xi, \eta \rangle_o} 1_{\text{Geod}_h^\infty(\Omega)}(\xi, \eta) d\mu_o(\xi) d\mu_o(\eta)$ yields a Γ -invariant measure on $\text{Geod}_h^\infty(\Omega)$, which is in fact the quotient of the Sullivan measure under the action of \mathbb{R} .

Proposition 3.4. *Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Let $\delta \geq 0$ and let $(\mu_x)_x$ be a δ -conformal density on $\partial_h\Omega$. Then the Sullivan measures on $\text{Geod}_h(\Omega) \times \mathbb{R}$, $T^1\Omega$ and T^1M are Radon. If Γ is strongly irreducible and $T^1M_{\text{bip}} \neq \emptyset$, or if M is rank-one and non-elementary, then these measures are non-zero.*

Proof. The fact that the Sullivan measures are Radon is a direct consequence of the continuity of the Gromov product (Proposition 3.1).

Suppose that Γ is strongly irreducible and $T^1M_{\text{bip}} \neq \emptyset$, or that M is rank-one and non-elementary. Then there exists $(\xi, \eta) \in \text{Geod}^\infty(\Omega) \cap (\Lambda^{\text{prox}})^2$. Let U and V be neighbourhoods of ξ and η in $\partial\Omega$ such that

$$U \times V \subset \text{Geod}^\infty(\Omega).$$

Since $\nu_o = \pi_{\text{hor}*}\mu_o$ is Γ -quasi-invariant, its support is Γ -invariant, and hence contains Λ^{prox} (see Remark 2.10 and Fact 2.18). Therefore, $\nu_o^2(U \times V) > 0$, and

$$\tilde{m}_h(\pi_h^{-1}(U) \times \pi_h^{-1}(V) \times \mathbb{R}) > 0. \quad \blacksquare$$

4. The Shadow lemma

In this section we establish the Shadow lemma (Lemma 4.2) which consists of estimates on the measures of shadows. The measure is a conformal density on $\partial\Omega$, and shadows are subsets of the projective boundary, defined as follows. Recall that the Shadow lemma is a classical result in the theory of conformal densities, and we adapt here its classical proof to the convex projective setting.

Definition 4.1. Let $\Omega \subset P(V)$ be a properly convex open set. Take $x, y \in \Omega$ and $r > 0$. Set

$$\begin{aligned} \mathcal{O}_r(x, y) &:= \{\xi \in \partial\Omega : [x, \xi] \cap B_\Omega(y, r) \neq \emptyset\}; \\ \mathcal{O}_r^+(x, y) &:= \{\xi \in \partial\Omega : \exists z \in B_\Omega(x, r) \text{ such that } [z, \xi] \cap B_\Omega(y, r) \neq \emptyset\}; \\ \mathcal{O}_r^-(x, y) &:= \{\xi \in \partial\Omega : \forall z \in B_\Omega(x, r), [z, \xi] \cap B_\Omega(y, r) \neq \emptyset\}. \end{aligned}$$

See in Figure 2 an example of a shadow.

Lemma 4.2. Let $o \in \Omega \subset P(V)$ be a pointed properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup; set $M = \Omega/\Gamma$. Suppose that Γ is strongly irreducible and T^1M_{bip} is non-empty, or that M is rank-one and non-elementary. Consider $\delta \geq 0$ and a δ -conformal density $(\nu_x)_{x \in \Omega}$ on $\partial\Omega$. Then there exists $R_0 > 0$ such that for any $R \geq R_0$, one can find $C = C(R) > 0$ such that for each $\gamma \in \Gamma$,

$$C^{-1}e^{-\delta d_\Omega(o, \gamma o)} \leq \nu_o(\mathcal{O}_R(o, \gamma o)) \leq \nu_o(\mathcal{O}_R^+(o, \gamma o)) \leq Ce^{-\delta d_\Omega(o, \gamma o)}.$$

We will actually need two more Shadow lemmas: Lemma 4.6 and Corollary 7.3. They both make stronger assumptions on the convex projective manifold $M = \Omega/\Gamma$, and either are consequences of Lemma 4.2, or have a very similar proof.

4.1. Preliminaries

In this section we prove two classical intermediate lemmas, used in the proof of the Shadow lemma.

Lemma 4.3. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set. Let $\xi \in \bar{\Omega}^{\text{h}}$ and $x, y \in \Omega$. If $y \in [x, \xi]$, then $b_\xi(x, y) = d_\Omega(x, y)$. Let $r > 0$. If $\pi_{\text{h}}(\xi) \in \mathcal{O}_r^+(x, y)$ (see Definition 4.1), then*

$$d_\Omega(x, y) - 4r \leq b_\xi(x, y) \leq d_\Omega(x, y).$$

Proof. One easily sees that $b_\xi(x, y) - d_\Omega(x, y) = -2\langle \xi, x \rangle_y$ if $\xi \in \Omega$, and this extends to $\xi \in \bar{\Omega}^{\text{h}}$ by continuity. This, by (3.1), implies that $b_\xi(x, y) = d_\Omega(x, y)$ if $y \in [x, \xi]$. Assume that $\xi \in \mathcal{O}_r^+(x, y)$. The triangular inequality gives $b_\xi(x, y) \leq d_\Omega(x, y)$; let us establish $b_\xi(x, y) \geq d_\Omega(x, y) - 4r$. By definition of $\mathcal{O}_r^+(x, y)$, we can find $x' \in B_\Omega(x, r)$ and $y' \in B_\Omega(y, r) \cap [x', \xi]$. Then

$$\begin{aligned} b_\xi(x, y) &= b_\xi(x, x') + b_\xi(x', y') + b_\xi(y', y) \\ &\geq -d_\Omega(x, x') + d_\Omega(x', y') - d_\Omega(y', y) \\ &\geq d_\Omega(x, y) - 2d_\Omega(x, x') - 2d_\Omega(y, y') \\ &\geq d_\Omega(x, y) - 4r. \end{aligned} \quad \blacksquare$$

Lemma 4.4. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup; set $M = \Omega/\Gamma$. Suppose that Γ is strongly irreducible and T^1M_{bip} is non-empty, or that M is rank-one and non-elementary. Consider a Γ -quasi-invariant finite Borel measure ν on $\partial\Omega$. Then there is $\varepsilon > 0$ and $R > 0$ such that for all $x \in \Omega$,*

$$\nu(\mathcal{O}_R(x, o)) \geq \varepsilon.$$

Proof. By contradiction suppose that there are sequences $(R_n)_n \in \mathbb{R}_{>0}^{\mathbb{N}}$ and $(x_n)_n \in \Omega$ such that

$$R_n \xrightarrow[n \rightarrow \infty]{} \infty \quad \text{and} \quad \nu(\mathcal{O}_{R_n}(x_n, o)) \xrightarrow[n \rightarrow \infty]{} 0.$$

We can assume that $(x_n)_n$ converges to some $\xi \in \bar{\Omega}$. If $\xi \in \Omega$ then for n such that $R_n \geq d_\Omega(o, \xi) + 1$ and $d_\Omega(x_n, \xi) < 1$, we have $\mathcal{O}_{R_n}(x_n, o) = \partial\Omega$ which is absurd; hence, $\xi \in \partial\Omega$. We claim that $\nu(\bar{B}_{\text{spl}}(\xi, 1)) = 1$. It is enough to prove that

$$\partial\Omega \setminus \bar{B}_{\text{spl}}(\xi, 1) \subset \bigcup_n \bigcap_{k \geq n} \mathcal{O}_{R_k}(x_k, o).$$

See Figure 2. Let $\eta \in \partial\Omega \setminus \bar{B}_{\text{spl}}(\xi, 1)$. Fixing an affine chart containing $\bar{\Omega}$, we can consider for each n the following points of Ω :

$$y := \frac{1}{2}(\xi - \eta) + \eta \quad \text{and} \quad y_n := \frac{1}{2}(x_n - \eta) + \eta,$$




Figure 2. The sequence of increasing shadows in the proof of Lemma 4.4.

where the map $x \mapsto (x - \eta)/2 + \eta$ is defined on the affine chart as the homothety centred at η and with ratio one half. We can then find n large enough so that $B_\Omega(o, R_n)$ contains a neighbourhood U of y , and so that y_n is contained in U . This implies that $\eta \in \mathcal{O}_{R_n}(x_n, o)$, which concludes the proof of the claim that $\nu(\bar{B}_{\text{spl}}(\xi, 1)) = 1$.

Since Λ^{prox} is the smallest Γ -invariant closed subset of $\bar{\Omega}$ (Remark 2.10 and Fact 2.18), and ν is Γ -quasi-invariant, we deduce that Λ^{prox} is contained in the support of ν . In order to get a contradiction, let us prove that Λ^{prox} is not contained in $\bar{B}_{\text{spl}}(\xi, 1)$. By assumption we can find $\eta, \eta' \in \Lambda^{\text{prox}}$ such that $(\eta \oplus \eta') \cap \Omega \neq \emptyset$. Using again that $\Lambda^{\text{prox}} \subset \bar{\Omega}$ is minimal for the action of Γ , we deduce the existence of a sequence $(\gamma_n)_n \in \Gamma^{\mathbb{N}}$ such that $(\gamma_n \xi)_n$ converges to η . But then $(\bar{B}_{\text{spl}}(\gamma_n \xi, 1))_n$ sub-converges to $\bar{B}_{\text{spl}}(\eta, 1)$ (i.e. any accumulation point for the Hausdorff topology is contained in $\bar{B}_{\text{spl}}(\eta, 1)$); hence, $\bar{B}_{\text{spl}}(\gamma_n \xi, 1)$ does not contain η' for n large enough. ■

We will need exactly twice a refined version of the previous lemma, which bounds from below the measure of *scarce shadows*. It will be used to prove the refined version of the Shadow lemma for scarce shadows (Lemma 4.6), and to prove Proposition 6.2. This version needs M to be rank-one, and the statement is a bit more technical. The proof is almost the same.

Lemma 4.5. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Suppose $M = \Omega/\Gamma$ is rank-one and non-elementary. Consider a Γ -quasi-invariant Borel finite measure ν on $\partial\Omega$. Then there exist $\varepsilon > 0$ and $R_0 > 0$ such that $\nu(\mathcal{O}_{\bar{R}}(x, o)) \geq \varepsilon$ for all $R \geq R_0$ and $x \in \Omega \setminus B_\Omega(o, 2R)$.*

Proof. By contradiction we suppose the existence of sequences $(R_n)_n \in \mathbb{R}_{>0}^{\mathbb{N}}$ and $(x_n)_n \in \Omega^{\mathbb{N}}$ such that for all $n \in \mathbb{N}$,

$$R_n \xrightarrow[n \rightarrow \infty]{} \infty,$$

while

$$d_\Omega(x_n, o) - R_n \xrightarrow[n \rightarrow \infty]{} \infty \quad \text{and} \quad \nu(\mathcal{O}_{R_n}^-(x_n, o)) \xrightarrow[n \rightarrow \infty]{} 0.$$

We can assume, up to extracting, that $(x_n)_n$ converges to some point $\xi \in \partial\Omega$. We claim that $\nu(\bar{B}_{\text{spl}}(\xi, 2)) = 1$ (see Definition 2.2). It is enough to prove that

$$\partial\Omega \setminus \bar{B}_{\text{spl}}(\xi, 2) \subset \bigcup_n \bigcap_{k \geq n} \mathcal{O}_{R_k}^-(x_k, o).$$

Let $\eta \in \partial\Omega \setminus \bar{B}_{\text{spl}}(\xi, 2)$. Fixing an affine chart containing $\bar{\Omega}$, we can consider for each n the following compact subsets of Ω :

$$K := \frac{1}{2}(\bar{B}_{\text{spl}}(\xi, 1) - \eta) + \eta \quad \text{and} \quad K_n := \frac{1}{2}(\bar{B}_{\Omega}(x_n, R_n) - \eta) + \eta,$$

where the map $x \mapsto (x - \eta)/2 + \eta$ is defined on the affine chart as the homothety centred at η and with ratio one half. We observe that all accumulation points of the sequence $(K_n)_n$ are contained in K . Indeed, any accumulation point of $(\bar{B}_{\Omega}(x_n, R_n))_n$ is convex, contains ξ and is moreover contained in $\partial\Omega$ since x_n goes faster to infinity than R_n ; hence, it is contained in $\bar{B}_{\text{spl}}(\xi, 1)$. Therefore we can find n large enough so that $B_{\Omega}(o, R_n)$ contains a neighbourhood U of K , and so that K_n is contained in U .

Since Λ^{prox} is the smallest Γ -invariant closed subset of $\bar{\Omega}$ (Fact 2.18), and ν is Γ -quasi-invariant, we deduce that Λ^{prox} is contained in the support of ν . In order to get a contradiction, let us prove that Λ^{prox} is not contained in $\bar{B}_{\text{spl}}(\xi, 2)$. By assumption we can find two distinct points η, η' in $\Lambda^{\text{prox}} \cap \partial_{\text{sse}}\Omega$. Using again that $\Lambda^{\text{prox}} \subset \bar{\Omega}$ is minimal for the action of Γ , we deduce the existence of a sequence $(\gamma_n)_n \in \Gamma^{\mathbb{N}}$ such that $(\gamma_n \xi)_n$ converges to η . But then $(\bar{B}_{\text{spl}}(\gamma_n \xi, 2))_n$ sub-converges to $\bar{B}_{\text{spl}}(\eta, 2) = \{\eta\}$; hence, $\bar{B}_{\text{spl}}(\gamma_n \xi, 2)$ does not contain η' for n large enough. ■

4.2. Proof of Lemma 4.2 and another Shadow lemma

Proof of Lemma 4.2. We compute for $\alpha \in \{\emptyset, +, -\}$:

$$\begin{aligned} \nu_o(\mathcal{O}_R^\alpha(o, \gamma o)) &= \mu_o(\pi_h^{-1}(\mathcal{O}_R^\alpha(o, \gamma o))) \\ &= \mu_{\gamma^{-1}o}(\pi_h^{-1}(\mathcal{O}_R^\alpha(\gamma^{-1}o, o))) \\ &= \int_{\pi_h^{-1}(\mathcal{O}_R^\alpha(\gamma^{-1}o, o))} e^{-\delta b_{\tilde{\xi}}(\gamma^{-1}o, o)} d\mu_o(\tilde{\xi}). \end{aligned}$$

Hence, on one hand $b_{\tilde{\xi}}(\gamma^{-1}o, o) \geq d_{\Omega}(o, \gamma^{-1}o) - 4R$ because of Lemma 4.3, so

$$\begin{aligned} \nu_o(\mathcal{O}_R^+(o, \gamma o)) &\leq \int_{\pi_h^{-1}(\mathcal{O}_R^+(\gamma^{-1}o, o))} e^{-\delta d_{\Omega}(o, \gamma o) + 4\delta R} d\mu_o(\tilde{\xi}) \\ &\leq e^{4\delta R} e^{-\delta d_{\Omega}(o, \gamma o)} \nu_o(\mathcal{O}_R^+(\gamma^{-1}o, o)) \\ &\leq e^{4\delta R} e^{-\delta d_{\Omega}(o, \gamma o)} \nu_o(\partial\Omega). \end{aligned}$$

On the other hand, we can use Lemma 4.4, to obtain $\varepsilon > 0$ and R_0 such that for $R \geq R_0$ and for $\gamma \in \Gamma$ such that $d_\Omega(o, \gamma o) \geq R_0$,

$$\begin{aligned} v_o(\mathcal{O}_R(o, \gamma o)) &\geq \int_{\pi_h^{-1}(\mathcal{O}_R(\gamma^{-1}o, o))} e^{-\delta d_\Omega(o, \gamma o)} d\mu_o(\tilde{\xi}) \\ &\geq v_o(\mathcal{O}_R(\gamma^{-1}o, o)) e^{-\delta d_\Omega(o, \gamma o)} \\ &\geq \varepsilon e^{-\delta d_\Omega(o, \gamma o)}. \end{aligned} \quad \blacksquare$$

As for Lemma 4.4, there exists a refined version of the Shadow lemma (Lemma 4.2) for scarce shadows. We will only need it once, for the proof of Proposition 6.2. Its proof is exactly the same as that of Lemma 4.2, except that we use instead Lemma 4.4.

Lemma 4.6. *Let $o \in \Omega \subset \mathbb{P}(V)$ be a pointed properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Suppose $M = \Omega/\Gamma$ is rank-one and non-elementary. Consider $\delta \geq 0$ and a δ -conformal density $(v_x)_{x \in \Omega}$ on $\partial\Omega$. Then there exists $R_0 > 0$ such that for any $R \geq R_0$, one can find $C = C(R) > 0$ such that for each $\gamma \in \Gamma$ satisfying $d_\Omega(o, \gamma o) \geq 2R$,*

$$C^{-1} e^{-\delta d_\Omega(o, \gamma o)} \leq v_o(\mathcal{O}_R^-(o, \gamma o)).$$

Proof. We make the same computation as in the proof of Lemma 4.2:

$$v_o(\mathcal{O}_R^-(o, \gamma o)) = \int_{\pi_h^{-1}(\mathcal{O}_R^-(\gamma^{-1}o, o))} e^{-\delta b_{\tilde{\xi}}(\gamma^{-1}o, o)} d\mu_o(\tilde{\xi}).$$

We can use Lemma 4.5, to obtain $\varepsilon > 0$ and R_0 such that for $R \geq R_0$ and for $\gamma \in \Gamma$ such that $d_\Omega(o, \gamma o) \geq 2R$,

$$\begin{aligned} v_o(\mathcal{O}_R^-(o, \gamma o)) &\geq \int_{\pi_h^{-1}(\mathcal{O}_R^-(\gamma^{-1}o, o))} e^{-\delta d_\Omega(o, \gamma o)} d\mu_o(\tilde{\xi}) \\ &\geq v_o(\mathcal{O}_R^-(\gamma^{-1}o, o)) e^{-\delta d_\Omega(o, \gamma o)} \\ &\geq \varepsilon e^{-\delta d_\Omega(o, \gamma o)}. \end{aligned} \quad \blacksquare$$

4.3. First consequences

In this section we deduce from the Shadow lemma that there is no conformal density with parameter $0 \leq \delta < \delta_\Gamma$. We also prove that open faces of the conical limit set are neglected by conformal densities.

Proposition 4.7. *Let $o \in \Omega \subset \mathbb{P}(V)$ be a pointed properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup; denote $M = \Omega/\Gamma$. Suppose that Γ is strongly irreducible and $T^1 M_{\text{bip}}$ is non-empty, or that M is rank-one and non-elementary. Consider*

$\delta \geq 0$ such that there exists a δ -conformal density on $\partial\Omega$. Then $\delta \geq \delta_\Gamma$, and there is some constant $C > 0$ such that

$$\#\{\gamma \in \Gamma : d_\Omega(o, \gamma o) \leq r\} \leq C e^{\delta_\Gamma r}.$$

Proof. Let $(v_x)_{x \in \Omega}$ be a δ -conformal density on $\partial\Omega$. We consider R and $C > 0$ from the Shadow lemma (Lemma 4.2), such that for each automorphism $\gamma \in \Gamma$,

$$v_o(\mathcal{O}_R(o, \gamma o)) \geq C^{-1} e^{-\delta d_\Omega(o, \gamma o)}.$$

For each $r > 0$, we give ourselves a maximal $(1 + 4R)$ -separated subset F_r of

$$\Gamma \cdot o \cap B_\Omega(o, r + 1) \setminus B_\Omega(o, r).$$

One can easily see that the shadows $(\mathcal{O}_R(o, x))_{x \in F_r}$ are pairwise disjoint, therefore

$$\begin{aligned} 1 &\geq \sum_{x \in F_r} v_o(\mathcal{O}_R(o, x)) \\ &\geq C^{-1} \sum_{x \in F_r} e^{-\delta d_\Omega(o, x)} \\ &\geq C^{-1} e^{-\delta} e^{-\delta r} \#F_r \\ &\geq C^{-1} e^{-\delta} \#\{\gamma : d_\Omega(o, \gamma o) \leq 1 + 4R\}^{-1} \\ &\quad \times e^{-\delta r} \#\{\gamma : \gamma o \in B_\Omega(o, r + 1) \setminus B_\Omega(o, r)\}. \end{aligned}$$

This implies that $\#\{\gamma : d_\Omega(o, \gamma o) \leq r\} \leq C' e^{\delta r}$ for any $r > 0$, for some $C' > 0$ independent of r . By definition, this implies that $\delta \geq \delta_\Gamma$, and since by Fact 2.22 there exists a δ_Γ -conformal density, $\#\{\gamma : d_\Omega(o, \gamma o) \leq r\} \leq C'' e^{\delta_\Gamma r}$ for any $r > 0$, for some $C'' > 0$ independent of r . ■

Proposition 4.8. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup; set $M = \Omega/\Gamma$. Suppose that Γ is strongly irreducible and $T^1 M_{\text{bip}}$ is non-empty, or that M is rank-one and non-elementary. Consider $\delta \geq 0$ and a δ -conformal density $(v_x)_{x \in \Omega}$ on $\partial\Omega$. Then $v_x(F_\Omega(\xi)) = 0$ for all $x \in \Omega$ and $\xi \in \Lambda^{\text{con}}$.*

Proof. Let $o \in \Omega$. Observe that the open face $F_\Omega(\xi)$ is contained in Λ^{con} by [35, Cor. 4.10]. It is enough to prove that $v_o(B_{\bar{\Omega}}(\xi, R)) = 0$ for any $R > 0$. By definition, there are sequences $(\gamma_n)_n \in \Gamma^{\mathbb{N}}$ and $(x_n)_n \in [o, \xi]^{\mathbb{N}}$ going to infinity such that $(d_\Omega(\gamma_n o, x_n))_n$ is bounded; denote $R' = \sup_n d_\Omega(\gamma_n o, x_n)$. Using the Shadow lemma (Lemma 4.2), we can find $C > 0$ such that for any n ,

$$v_o(B_{\bar{\Omega}}(\xi, R)) \leq v_o(\mathcal{O}_R(o, x_n)) \leq v_o(\mathcal{O}_{R+R'}(o, \gamma_n o)) \leq C e^{-\delta d_\Omega(o, \gamma_n o)}.$$

This last term goes to zero as n tends to infinity since $\delta \geq \delta_\Gamma > 0$ (Fact 2.7 and Proposition 4.7). ■

5. The convergent case of the HTSR dichotomy

Next we establish the convergent case of the HTSR dichotomy (Theorem 1.6(1)).

5.1. The conical limit set has zero measure

We prove that, in the convergent case of the HTSR dichotomy, any δ -conformal density gives zero measure to the conical limit set.

Proposition 5.1. *Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Suppose that Γ is strongly irreducible and T^1M_{bip} is non-empty, or that M is rank-one and non-elementary. Let $\delta \geq \delta_\Gamma$ with $\sum_{\gamma \in \Gamma} e^{-\delta d_\Omega(o, \gamma o)}$ finite, and consider a δ -conformal density $(v_x)_{x \in \Omega}$ on $\partial\Omega$. Then $v_x(\Lambda^{\text{con}}) = 0$ for any $x \in \Omega$.*

Proof. We consider $R > 0$ and prove that $v_o(\Lambda_R^{\text{con}}) = 0$. By definition, for any $r > 0$, the set Λ_R^{con} is contained in the union, over all $\gamma \in \Gamma$ such that $d_\Omega(o, \gamma o) \geq r$, of the shadows $\mathcal{O}_R(o, \gamma o)$. As a consequence, by the Shadow lemma (Lemma 4.2), we can find a constant $C > 0$ such that

$$v_o(\Lambda_R^{\text{con}}) \leq C \sum_{\gamma: d_\Omega(o, \gamma o) \geq r} e^{-\delta d_\Omega(o, \gamma o)}$$

for any $r > 0$, and this last quantity converges to zero as r goes to infinity. ■

5.2. Proof of Theorem 1.6(1)

Let μ_o be a δ_Γ -conformal density on $\partial_h\Omega$ such that $(\pi_h)_*\mu_o = v_o$, and let \tilde{m} be the Sullivan measure on $T^1\Omega$ induced by μ_o . According to Proposition 4.7, we have $\delta \geq \delta_\Gamma$. Let us assume that $\sum_\gamma e^{-\delta d_\Omega(o, \gamma o)}$ is convergent.

By Fact 2.29, in order to prove that (T^1M, ϕ_t, m) and $(\text{Geod}^\infty(\Omega), \Gamma, v_o^2)$ are dissipative, it is enough to prove that $(T^1\Omega, \Gamma \times \mathbb{R}, \tilde{m})$ is dissipative. If by contradiction it is not the case, then the conservative part contains a compact subset K of positive measure. This means that for almost any vector $v \in K$, using notations from Section 2.11,

$$\infty = \int_{\Gamma \times \mathbb{R}} 1_K(v) = \sum_{\gamma \in \Gamma} \int_{-\infty}^{\infty} 1_K(\gamma \phi_t v) dt,$$

and hence there exist diverging sequences $(\gamma_n)_n \in \Gamma^{\mathbb{N}}$ and $(t_n)_n \in \mathbb{R}^{\mathbb{N}}$ such that $\gamma_n \phi_{t_n} v \in K$; if $(t_n)_n$ tends to ∞ (resp. $-\infty$), then $\phi_\infty v$ (resp. $\phi_{-\infty} v$) is in Λ^{con} , which contradicts Proposition 5.1 and the definition of \tilde{m} since for almost every vector v in $T^1\Omega$, the endpoints $\phi_{\pm\infty} v$ are not in Λ^{con} .

Suppose by contradiction that (T^1M, ϕ_t, m) is ergodic. Let $K \subset T^1\Omega$ be a compact neighbourhood of a vector v in $T^1\Omega_{\text{bip}} := \pi_\Gamma^{-1}T^1M_{\text{bip}}$. By Fact 2.30(3), almost

any $(\phi_t)_t$ orbit is dense in the support of m , which contains T^1M_{bip} (this was basically proved in Proposition 3.4). By dissipativity (and because m is Radon), almost any orbit spends a finite time in $\pi_\Gamma K$. Hence, there exists a vector $w \in T^1\Omega$ and a time $T > 0$ such that

$$\pi_\Gamma(K) \cap T^1M_{\text{bip}} \subset \phi_{[-T, T]}\pi_\Gamma w;$$

in other words, as the action of Γ on $T^1\Omega$ is properly discontinuous, we may find a finite subset $A \subset \Gamma$ such that

$$K \cap T^1\Omega_{\text{bip}} \subset A\phi_{[-T, T]}w.$$

Thus, $\phi_\infty(K \cap T^1\Omega_{\text{bip}})$, which is a neighbourhood in Λ^{prox} of $\phi_\infty v$, is contained in $\phi_\infty A\phi_{[-T, T]}w$, which is equal to $A\phi_\infty w$ and hence is finite. This contradicts the fact that Λ^{prox} has no isolated point (Remark 2.10 and Fact 2.18).

6. The divergent case of the HTSR dichotomy

In this section, we adapt some proofs of Roblin [56, pp. 19–23] to our convex projective setting, in order to establish Theorem 1.6. We will fix a Patterson–Sullivan density, and need several time to prove that some Γ -invariant subset A of $\partial\Omega$ is given full measure by the Patterson–Sullivan density. Thanks to the following elementary observation, it is often enough to show that A has non-zero measure.

Observation 6.1. Let $\Omega \subset P(V)$ be a properly convex open set, $\Gamma \subset \text{Aut}(\Omega)$ a closed subgroup, $A \subset \partial\Omega$ a Γ -invariant measurable subset, and $\delta \geq 0$. Then for any δ -conformal density $(\nu_x)_{x \in \Omega}$ on $\partial\Omega$, the restrictions $(\nu_x|_A)_{x \in \Omega}$ also form a δ -conformal density. This implies the following. Suppose that for any non-zero δ -conformal density $(\nu_x)_{x \in \Omega}$, we have $\nu_x(A) > 0$ (for any $x \in \Omega$). Then for any δ -conformal density $(\nu_x)_{x \in \Omega}$, we have $\nu_x(\partial\Omega \setminus A) = 0$ (for any $x \in \Omega$).

Proof. That restrictions to A of δ -conformal densities are still δ -conformal densities follows easily from the definition. Let us prove the second statement.

We make a proof by contrapositive: let $B = \partial\Omega \setminus A$ suppose that there exists a δ -conformal density $(\nu_x)_x$ such that $\nu_x(B) > 0$ for some (hence for any) $x \in \Omega$. Then by the first part of the observation $(\nu_x|_B)_x$ is a non-zero δ -conformal density that gives zero measure to A , concluding the proof. ■

6.1. The conical limit set has full measure

Let $o \in \Omega \subset P(V)$ be a pointed properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Recall that the conical limit set, denoted by Λ^{con} , is the union, over

$R > 0$, of the sets $\Lambda_R^{\text{con}}(\Gamma, \Omega, o) = \Lambda_R^{\text{con}}$, consisting of points $\xi \in \partial\Omega$ for which there exists a sequence $(\gamma_n)_{n \in \mathbb{N}} \in \Gamma^{\mathbb{N}}$ going to infinity such that $d_\Omega([o, \xi], \gamma_n o) < R$ for each $n \in \mathbb{N}$.

Thanks to Observation 6.1, we only need to prove that the conical limit set has non-zero measure. Roblin’s proof of this [56, item (f), p. 19] in CAT(−1) geometry actually works verbatim in the present context; we rewrite it here for the convenience of the (non-French-speaking) reader.

Proposition 6.2. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Suppose Γ is divergent, and $M = \Omega/\Gamma$ is rank-one and non-elementary. Consider a δ_Γ -conformal density $(\nu_x)_{x \in \Omega}$ on $\partial\Omega$. Then $\nu_x(\partial\Omega \setminus \Lambda^{\text{con}}) = 0$ for any $x \in \Omega$.*

The idea of the proof of Proposition 6.2 is to find a compact subset $K \subset T^1M$ such that the set of vectors $v \in K$ whose geodesic come back infinitely often to K has positive m -measure where m is a Sullivan measure induced by $(\nu_x)_{x \in \Omega}$. There are two main ingredients. The first one is a generalisation of the Borel–Cantelli lemma, due to Rényi. One can find a proof of it in [1, Lem. 2]; it is the consequence of the following estimate: for any non-negative square-integrable function g on a probability space, for any $0 < a < E(g)$, where $E(g)$ is the expectation of g , one has

$$P(g > a) \geq \frac{E(g)^2}{E(g^2)} \left(1 + \frac{a}{E(g) - a}\right)^{-2}.$$

Fact 6.3 ([54, p. 391]). Let (X, μ) be a measurable space equipped with a finite positive measure. Let $(A_t)_{t \geq 0}$ be a family of subsets of X such that the function $(x, t) \in X \times \mathbb{R}_{\geq 0} \mapsto 1_{A_t}(x)$ is measurable. Let us assume that $\int_0^\infty \mu(A_t) dt = \infty$, and that, for some constant $C > 0$,

$$\int_0^T \int_0^T \mu(A_t \cap A_s) dt ds \leq C \left(\int_0^T \mu(A_t) dt \right)^2, \tag{6.1}$$

for T large enough. Then the set of $x \in X$ such that

$$\int_0^\infty 1_{A_t}(x) dt = \infty$$

has μ -measure greater than or equal to $1/C$.

This lemma will be applied to $A_t = K \cap \phi_{-t}K$. The second ingredient consists of estimates that will allow us to check that the assumptions of Fact 6.3 are satisfied.

Lemma 6.4. *Let $o \in \Omega \subset \mathbb{P}(V)$ be a pointed properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Suppose $M = \Omega/\Gamma$ is rank-one and non-elementary. Consider a δ_Γ -conformal density $(\nu_x)_{x \in \Omega}$ on $\partial\Omega$, with induced Sullivan measure m on T^1M .*

Let $\nu_o \in T_o^1\Omega$. For $R > 0$ large enough, if we denote by K the projection in T^1M of $\bar{B}_{T^1\Omega}(\nu_o, R)$, then there exist constants $C > 0$ and T_0 such that for any $T > T_0$, we have the estimates

$$\int_0^T m(K \cap \phi_{-t}K) dt \geq C^{-1} \sum_{\substack{g \in \Gamma \\ d_\Omega(o, go) \leq T}} e^{-\delta_\Gamma d_\Omega(o, go)}, \quad (6.2)$$

$$\int_0^T \int_0^T m(K \cap \phi_{-t}K \cap \phi_{-t-s}K) ds, dt \leq C \left(\sum_{\substack{g \in \Gamma \\ d_\Omega(o, go) \leq T}} e^{-\delta_\Gamma d_\Omega(o, go)} \right)^2. \quad (6.3)$$

Proof. We assume that ν_o is a probability measure. Let $(\mu_x)_{x \in \Omega}$ be a δ_Γ -conformal density on $\partial_h\Omega$ that induces $(\nu_x)_{x \in \Omega}$ and m ; we denote by \tilde{m} the induced Sullivan measure on $T^1\Omega$. We fix $R > 0$ large enough so that we can apply the Shadow lemma (Lemmas 4.2 and 4.6) and Lemmas 4.4 and 4.5 to it and to $R' := (R - 2)/6$. One can find a constant $C_1 > 0$ such that

$$\begin{aligned} m(K \cap \phi_{-t}K) &\geq C_1^{-1} \sum_{g \in \Gamma} \tilde{m}(\tilde{K} \cap \phi_{-t}g\tilde{K}), \\ m(K \cap \phi_{-t}K \cap \phi_{-t-s}K) &\leq \sum_{g, h \in \Gamma} \tilde{m}(\tilde{K} \cap \phi_{-t}g\tilde{K} \cap \phi_{-t-s}h\tilde{K}) \end{aligned}$$

for all $t, s \geq 0$. Indeed, we just have to recall the definition of m , which is the quotient of \tilde{m} under the action of Γ (Definition 2.28), then

$$\begin{aligned} \sum_{g \in \Gamma} \tilde{m}(\tilde{K} \cap \phi_{-t}g\tilde{K}) &= \int_{T^1\Omega} \sum_{g \in \Gamma} 1_{\tilde{K}} 1_{g\phi_{-t}\tilde{K}} d\tilde{m} \\ &= \int_{T^1M} \sum_{h \in \Gamma} \sum_{g \in \Gamma} (1_{\tilde{K}} \circ h) \cdot (1_{g\phi_{-t}\tilde{K}} \circ h) dm \\ &= \int_{T^1M} \underbrace{\left(\sum_{h \in \Gamma} 1_{h\tilde{K}} \right)}_{\leq C_1 1_K} \cdot \underbrace{\left(\sum_{g \in \Gamma} 1_{g\phi_{-t}\tilde{K}} \right)}_{\leq C_1 1_{\phi_{-t}K}} dm, \end{aligned}$$

where $C_1 > 0$ is a constant which is independent of t . Similarly,

$$\begin{aligned} \sum_{g, h \in \Gamma} \tilde{m}(\tilde{K} \cap \phi_{-t}g\tilde{K} \cap \phi_{-t-s}h\tilde{K}) \\ &= \int_{T^1M} \sum_{k \in \Gamma} \sum_{g, h \in \Gamma} (1_{\tilde{K}} \circ k) \cdot (1_{g\phi_{-t}\tilde{K}} \circ k) \cdot (1_{h\phi_{-t-s}\tilde{K}} \circ k) dm \\ &= \int_{T^1M} \underbrace{\left(\sum_{k \in \Gamma} 1_{k\tilde{K}} \right)}_{\geq 1_K} \cdot \underbrace{\left(\sum_{g \in \Gamma} 1_{g\phi_{-t}\tilde{K}} \right)}_{\geq 1_{\phi_{-t}K}} \cdot \underbrace{\left(\sum_{h \in \Gamma} 1_{h\phi_{-t-s}\tilde{K}} \right)}_{\geq 1_{\phi_{-t-s}K}} dm. \end{aligned}$$

Therefore, in order to prove Lemma 6.4, it is enough to find a constant $C_2 > 0$ such that

$$\begin{aligned} a &:= \int_0^T \sum_{g \in \Gamma} \tilde{m}(\tilde{K} \cap \phi_{-t} g \tilde{K}) dt \geq C_2^{-1} \sum_{g: d_\Omega(o, go) \leq T} e^{-\delta_\Gamma d_\Omega(o, go)}, \\ b &:= \int_0^T \int_0^T \sum_{g, h \in \Gamma} \tilde{m}(\tilde{K} \cap \phi_{-t} g \tilde{K} \cap \phi_{-s} h \tilde{K}) ds dt \\ &\leq C_2 \left(\sum_{g: d_\Omega(o, go) \leq T} e^{-\delta_\Gamma d_\Omega(o, go)} \right)^2. \end{aligned}$$

We first establish the estimate of b . For all $0 \leq t, s \leq T$ and $g, h \in \Gamma$, for any triple (ξ, η, τ) in $\text{Hopf}^{-1}(\tilde{K} \cap \phi_{-t-s} h \tilde{K})$, one observes that $\eta \in \pi_h^{-1} \mathcal{O}_R^+(o, ho)$ and

$$d_\Omega(o, \pi \text{Hopf}(\xi, \eta, \tau)) \leq R;$$

this last inequality implies that $|\tau| \leq R$ and $\langle \xi, \eta \rangle_o \leq R$; then by the Shadow lemma (Lemma 4.2), and by definition of \tilde{m} ,

$$\begin{aligned} \tilde{m}(\tilde{K} \cap \phi_{-t} g \tilde{K} \cap \phi_{-s} h \tilde{K}) &\leq 2R e^{2\delta_\Gamma R} \nu_0(\mathcal{O}_R^+(o, ho)) \\ &\leq 2R e^{2\delta_\Gamma R} C e^{-\delta_\Gamma d_\Omega(o, ho)}. \end{aligned}$$

Furthermore, by triangular inequality, if $\tilde{K} \cap \phi_{-t} g \tilde{K} \cap \phi_{-s} h \tilde{K}$ is non-empty, then

$$|d_\Omega(o, go) - t|, |d_\Omega(go, ho) - s|, |d_\Omega(o, ho) - t - s| \leq 2R$$

and

$$d_\Omega(o, go) + d_\Omega(go, ho) \leq d_\Omega(o, ho) + 6R.$$

Combining these estimates, we obtain

$$\begin{aligned} b &\leq \int_0^T \int_0^T \sum_{g, h \in \Gamma} 2R e^{2\delta_\Gamma R} C e^{-\delta_\Gamma d_\Omega(o, ho)} 1_{\left\{ \begin{array}{l} |d_\Omega(o, go) - t|, |d_\Omega(go, ho) - s| \leq 2R \\ d_\Omega(o, go) + d_\Omega(go, ho) \leq d_\Omega(o, ho) + 6R \end{array} \right\}} ds dt \\ &\leq 2R^2 R^2 R e^{2\delta_\Gamma R} C \sum_{\substack{d_\Omega(o, go) \leq T+2R \\ d_\Omega(o, g^{-1}ho) \leq T+2R}} e^{-\delta_\Gamma (d_\Omega(o, go) + d_\Omega(o, g^{-1}ho) - 6R)} \\ &\leq 8R^3 C e^{8\delta_\Gamma R} \left(\sum_{d_\Omega(o, go) \leq T+2R} e^{-\delta_\Gamma d_\Omega(o, go)} \right)^2. \end{aligned}$$

We end the estimation of b by noting that for all $T \geq 0$ and $A \geq 0$, using again the Shadow lemma (Lemma 4.2),

$$\begin{aligned} \sum_{T \leq d_{\Omega}(o, go) \leq T+A} e^{-\delta_{\Gamma} d_{\Omega}(o, go)} &\leq C \sum_{T \leq d_{\Omega}(o, go) \leq T+A} \nu_o(\mathcal{O}_R(o, go)) \\ &\leq C \int_{\partial\Omega} \sum_{T \leq d_{\Omega}(o, go) \leq T+A} 1_{\mathcal{O}_R(o, go)}(\xi) d\nu_o(\xi) \\ &\leq C \#\{g : d_{\Omega}(o, go) \leq 4R + 2A\}. \end{aligned}$$

We now proceed to the minoration of a . Take $g \in \Gamma$, take $t \geq 0$ at distance less than R' from $d_{\Omega}(o, go)$; let us prove that

$$\text{Hopf}(\pi_h^{-1}(\mathcal{O}_{R'}^-(go, o)) \times \pi_h^{-1}(\mathcal{O}_{R'}^-(o, go)) \times [0, R']) \subset \tilde{K} \cap \phi_{-t} g \tilde{K}.$$

Take $(\xi, \eta, \tau) \in \pi_h^{-1}(\mathcal{O}_{R'}^-(go, o)) \times \pi_h^{-1}(\mathcal{O}_{R'}^-(o, go)) \times [0, R']$. According to Observation 6.5 below, there exists $s_1 < s_2$ such that

$$d_{\Omega}(\pi \text{Hopf}(\xi, \eta, s_1), o) \leq R' \quad \text{and} \quad d_{\Omega}(\pi \text{Hopf}(\xi, \eta, s_2), go) \leq R'.$$

This implies that $|s_1| = |b_{\eta}(\pi \text{Hopf}(\xi, \eta, s_1), o)| \leq R'$, hence

$$d_{\Omega}(\pi \text{Hopf}(\xi, \eta, \tau), o) \leq |\tau| + |s_1| + d_{\Omega}(\pi \text{Hopf}(\xi, \eta, s_1), o) \leq 3R'.$$

Finally, $d_{T^1\Omega}(\text{Hopf}(\xi, \eta, \tau), v_o) \leq 3R' + 2 \leq R$, which means that $\text{Hopf}(\xi, \eta, \tau) \in \tilde{K}$.

In order to prove the inclusion in $\phi_{-t} g \tilde{K}$, we note that since $s_2 - s_1$ is the distance between $\pi \text{Hopf}(\xi, \eta, s_2)$ and $\pi \text{Hopf}(\xi, \eta, s_1)$, then by triangular inequality $|s_2 - d_{\Omega}(o, go)| \leq 3R'$; therefore,

$$\begin{aligned} d_{\Omega}(\pi \text{Hopf}(\xi, \eta, t + \tau), go) &\leq |\tau| + |t - d_{\Omega}(o, go)| \\ &\quad + |d_{\Omega}(o, go) - s_2| + d_{\Omega}(\pi \text{Hopf}(\xi, \eta, s_2), go) \\ &\leq 6R'. \end{aligned}$$

Finally,

$$d_{T^1\Omega}(\text{Hopf}(\xi, \eta, t + \tau), gv_o) \leq 6R' + 2 = R,$$

which means that $\text{Hopf}(\xi, \eta, \tau) \in \phi_{-t} g \tilde{K}$.

As a consequence, if $d_{\Omega}(o, go) \geq 2R'$, then by the Shadow lemma (Lemma 4.6) and Lemma 4.5,

$$\begin{aligned} \tilde{m}(\tilde{K} \cap \phi_{-t} g \tilde{K}) &\geq \nu_o(\mathcal{O}_{R'}^-(o, go)) \nu_o(\mathcal{O}_{R'}^-(go, o)) R' \\ &\geq R' C^{-2} e^{-\delta_{\Gamma} d_{\Omega}(o, go)}. \end{aligned}$$

We conclude that for $T \geq R'$,

$$\begin{aligned} & \int_0^T \sum_{g \in \Gamma} \tilde{m}(\tilde{K} \cap \phi_{-t} g \tilde{K}) dt \\ & \geq R'^2 C^{-2} \left(\sum_{g: d_\Omega(o, go) \leq T} e^{-\delta_\Gamma d_\Omega(o, go)} - \sum_{g: d_\Omega(o, go) \leq 2R'} e^{-\delta_\Gamma d_\Omega(o, go)} \right). \quad \blacksquare \end{aligned}$$

Observation 6.5. Let $n \geq 1$ be an integer, A and B be two non-empty disjoint compact subsets of \mathbb{R}^n , and $\xi, \eta \in \mathbb{R}^n$ be two points. If for all $a \in A$ and $b \in B$, the intersections $[a, \eta] \cap B$ and $[b, \xi] \cap A$ are non-empty, then one can find $a \in A$ and $b \in B$ such that ξ, a, b, η are aligned in this order.

Proof of Proposition 6.2. Let m be a Sullivan measure associated to $(v_x)_{x \in \Omega}$ on T^1M . By definition of m and of the conical limit set, it is enough to find a compact set $K \subset T^1M$ large enough so that the set of vectors $v \in K$ such that

$$\int_0^\infty 1_K(\phi_t v) dt = \infty$$

has non-zero m -measure.

Let $R > 0$ be large enough so that we can apply Lemma 6.4, and let K be the projection in T^1M of $\bar{B}_{T^1\Omega}(v_o, R)$, where $v_o \in T_o^1\Omega$ and $o \in \Omega$. We want to apply Fact 6.3 for $(X, \mu) = (T^1M, m|_K)$ and $A_t := K \cap \phi_{-t}K$ for all $t \geq 0$. The measure μ is finite because m is Radon and K is compact. Since Γ is divergent (this is important since it is the only place where we need this assumption), and by the estimate (6.2) of Lemma 6.4, the integral $\int_0^\infty \mu(A_t) dt$ diverges. It remains to check that (6.1) is satisfied, but this is a direct consequence of the estimates (6.2) and (6.3) of Lemma 6.4, and of the fact that

$$\int_0^T \int_0^T \mu(A_t \cap A_s) dt ds \leq 2 \int_0^T \int_0^T \mu(A_t \cap A_{t+s}) dt ds. \quad \blacksquare$$

6.2. The geodesic flow is conservative

In this section, we prove that in the setting of Theorem 1.6(2) the geodesic flow is conservative.

Proposition 6.6. *Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a divergent discrete subgroup. Suppose $M = \Omega/\Gamma$ is rank-one and non-elementary. Let $(v_x)_{x \in \Omega}$ be a δ_Γ -conformal density on $\partial\Omega$ and m an induced Sullivan measure on T^1M . Then m is conservative under the action of $(\phi_t)_t$.*

Proof. Let $(\mu_x)_{x \in \Omega}$ be a δ_Γ -conformal density on $\partial_{\text{h}}\Omega$ such that $(\pi_{\text{h}})_*\mu_x = \nu_x$ for any $x \in \Omega$, and let \tilde{m} be the induced Sullivan measure on $T^1\Omega$.

According to Fact 2.29, it is enough to prove that $(T^1\Omega, \Gamma \times \mathbb{R}, \tilde{m})$ is conservative. Let σ be an integrable, positive and continuous function on $T^1\Omega$, and let us prove that $\int_{\Gamma \times \mathbb{R}} \sigma = \infty$ almost surely (recall the notation from Section 2.11). For almost any vector $v \in T^1\Omega$, we have $\phi_\infty v \in \Lambda^{\text{con}}$ by Proposition 6.2, and by definition of \tilde{m} ; let v be such a vector. Then we can find $R > 0$ such that $\phi_\infty v \in \Lambda_R^{\text{con}}$; in other words, there exist $(\gamma_n)_n \in \Gamma^{\mathbb{N}}$ and $(t_n)_n \in \mathbb{R}^{\mathbb{N}}$ such that $\gamma_n \neq \gamma_k$ and $d_\Omega(\pi\phi_{t_n}\gamma_n v, o) \leq R$ for $n \neq k$. Let $\varepsilon := \min\{\sigma(w) : d_\Omega(\pi w, o) \leq R + 1\} > 0$. Then

$$\int_{\Gamma \times \mathbb{R}} \sigma(v) \geq \sum_{n \geq 0} \int_{t=t_n-1}^{t_n+1} \sigma(\phi_t \gamma v) dt \geq \sum_{n \geq 0} 2\varepsilon = \infty. \quad \blacksquare$$

6.3. The smooth and strongly extremal points have full measure

In this section we combine the conservativity of Sullivan’s measures with a result of Benzécri to establish that in the divergent case, the smooth and strongly extremal points have full measure. The following result is a particular case of a more general result of Benzécri. Recall that \mathcal{E}_V^\bullet is the space of pointed properly convex open subsets of $\mathbb{P}(V)$, endowed with the Hausdorff topology, on which $\text{PGL}(V)$ acts properly cocompactly (Fact 2.4).

Fact 6.7 ([12, Prop. 5.3.9]). Suppose that $\dim(V) = 3$. Let $(x, \Omega) \in \mathcal{E}_V^\bullet$ and $\xi \in \partial_{\text{sing}}\Omega$. Let $T \in \mathcal{E}_V$ be a triangle and $o \in T$. Then we have the following convergence in $\mathcal{E}_V^\bullet/\text{PGL}(V)$:

$$[y, \Omega] \xrightarrow[\substack{y \rightarrow \xi \\ y \in [x, \xi)}}{]} [o, T].$$

Proof. We briefly recall the proof of this fact, which is very easy in this particular case. Consider a projective line $\mathbb{P}(W)$ that does not intersect $[x, \xi]$, and for each $y \in [x, \xi)$, denote by $g_y \in \text{PGL}(V)$ the unique element that fixes ξ and $\mathbb{P}(W)$, and such that $g_y y = x$. Let $p: \mathbb{P}(V) \setminus \{\xi\} \rightarrow \mathbb{P}(W)$ be the stereographic projection. Since ξ is a singular point of $\partial\Omega$, the image $p(\Omega)$ is a properly convex open subset of $\mathbb{P}(W)$, i.e. the interior of a segment. As y tends to ξ , the properly convex open set $g_y\Omega$ converges to the convex hull of $p(\Omega)$ and ξ , which is a triangle containing x . \blacksquare

Lemma 6.8. Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup; set $M = \Omega/\Gamma$. Let $v \in T^1\Omega$ with a forward recurrent projection in T^1M . If $\phi_\infty v \notin \partial_{\text{sse}}\Omega$, then

$$d_{\text{spl}}(\phi_{-\infty} v, \phi_\infty v) = 2.$$

A similar statement holds for backward recurrent vectors.

In particular, the above lemma implies that within set of vectors $v \in T^1\Omega$ with forward and backward recurrent projection, the subset made of non-rank-one vectors is closed.

Proof. Observe that the inequality $d_{\text{spl}}(\phi_{-\infty}w, \phi_{\infty}w) \geq 2$ holds for any vector w . Since the projection of v in T^1M is forward recurrent, there exist diverging sequences $(t_n)_n \in [0, \infty)^{\mathbb{N}}$ and $(\gamma_n)_n \in \Gamma^{\mathbb{N}}$ such that $(\gamma_n\phi_{t_n}v)_n$ tends to v .

Suppose that $\phi_{\infty}v$ is singular. Then there exists a projective plane $P(W) \subset P(V)$ which contains v and such that ξ is a singular point of $\partial\Omega \cap P(W)$. Up to extraction, we can assume that $(\gamma_n P(W))_n$ converges to some $P(W')$. By construction, $(\gamma_n\pi\phi_{t_n}v, \Omega \cap \gamma_n P(W))_n$ converges to $(\pi v, \Omega \cap P(W'))$. Since $\phi_{\infty}v$ is singular, we can apply Fact 6.7, and we obtain that $\Omega \cap P(W')$ is a triangle that contains $\phi_{\pm\infty}v$, hence

$$d_{\text{spl}}(\phi_{-\infty}v, \phi_{\infty}v) \leq 2.$$

Suppose there exists $\xi \in \partial\Omega \setminus \{\phi_{\infty}v\}$ such that $[\xi, \phi_{\infty}v] \subset \partial\Omega$. We can take ξ extremal. Up to extraction, we can assume that $(\gamma_n\xi)_n$ converges to some $\xi' \in \partial\Omega$. Observe that

$$[\phi_{\infty}v, \xi'] \subset \partial\Omega,$$

since it is the limit of the sequence of segments $([\gamma_n\phi_{\infty}v, \gamma_n\xi])_n$ that are contained in the boundary (see Figure 3). Since ξ is extremal, the Hilbert distance of $\pi\phi_{t_n}v$ to $[\phi_{-\infty}v, \xi] \cap \Omega$ tends to infinity with n , and hence the Hilbert distance of $\gamma_n\pi\phi_{t_n}v$ to $[\gamma_n\phi_{-\infty}v, \gamma_n\xi]$ tends to infinity. This last segment tends to the segment $[\phi_{-\infty}v, \xi']$, while $\gamma_n\pi\phi_{t_n}v \rightarrow \pi v$. This implies that $[\phi_{-\infty}v, \xi'] \subset \partial\Omega$, since otherwise the Hilbert distance from $\gamma_n\pi\phi_{t_n}v$ to $[\gamma_n\phi_{-\infty}v, \gamma_n\xi]$ would converge to the (finite) Hilbert distance from πv to $[\phi_{-\infty}v, \xi']$. Thus, $d_{\text{spl}}(\phi_{-\infty}v, \phi_{\infty}v) \leq 2$. ■

Observe that the following corollary of Lemma 6.8 generalises with a shorter proof a result of Islam [40, Prop. 6.3]. It can also be deduced from Lemma 7.7 (6).

Corollary 6.9. *Let $\Omega \subset P(V)$ be a properly convex open set and $g \in \text{Aut}(\Omega)$ with $\ell(g) > 0$ and such that g fixes two points $\xi, \eta \in \partial\Omega$ with $d_{\text{spl}}(\xi, \eta) > 2$. Then g is rank-one and $\{\xi, \eta\} = \{x_g^+, x_g^-\}$.*

Proof. We apply Lemma 6.8 to any vector tangent to $[\xi, \eta]$, whose projection in the quotient of $T^1\Omega$ by the group generated by g is periodic, and hence it is forward recurrent. ■

We now combine Lemma 6.8 with Poincaré's recurrence theorem (Fact 2.30 (2)) to establish the following.




Figure 3. Illustration of the second part of the proof of Lemma 6.8.

Proposition 6.10. *Let $\Omega \subset P(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a divergent discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Then any δ_Γ -conformal density on $\partial\Omega$ gives full measure to $\partial_{\text{sse}}\Omega$.*

Proof. Let $o \in \Omega$ and $(\nu_x)_{x \in \Omega}$ a δ_Γ -conformal density on $\partial\Omega$. Observe that $\partial_{\text{sse}}\Omega$ is Γ -invariant and measurable, hence, by Observation 6.1, in order to show that $\partial_{\text{sse}}\Omega$ has full ν_o -measure, it is enough to prove that $\nu_o(\partial_{\text{sse}}\Omega) > 0$. Let us assume by contradiction that

$$\nu_o(\partial_{\text{sse}}\Omega) = 0.$$

Let m be an induced Sullivan measure on T^1M . By assumption, m gives zero measure to the set of vectors in T^1M who have a lift $v \in T^1\Omega$ such that $\phi_\infty v$ is smooth and strongly extremal. By Proposition 6.6 and Fact 2.30 (2), m gives full measure to the set of forward recurrent vectors. Let $K \subset T^1\Omega$ denote the closed set of vectors $v \in T^1\Omega$ such that $d_{\text{spl}}(\phi_{-\infty}v, \phi_\infty v) = 2$. By Lemma 6.8, \tilde{m} gives full measure to the set of vectors $v \in K$ with a forward recurrent projection in T^1M . Let $v \in T^1\Omega$ be a periodic rank-one vector; it is in the support of \tilde{m} and in the open set $T^1\Omega \setminus K$, so $\tilde{m}(T^1\Omega \setminus K) > 0$. This is a contradiction. ■

6.4. Smooth points and strong stable manifolds

Let $\Omega \subset P(V)$ be a properly convex open set. The following fact describes the strong stable manifolds (Definition 2.31) of the action of the geodesic flow on $T^1\Omega$.

Fact 6.11 ([14, Cor. 4.5]). Let $\Omega \subset P(V)$ be a properly convex open set. Let $v, w \in T^1\Omega$ be such that $\phi_\infty v = \phi_\infty w$ is a smooth point of $\partial\Omega$ and $b_{\phi_\infty v}(\pi v, \pi w) = 0$ (see Figure 4). Then $(d_{T^1\Omega}(\phi_t v, \phi_t w))_t$ converges non-increasingly to zero as t goes to infinity, i.e. $w \in W^{ss}(v)$.




Figure 4. v and w in the same strong stable manifold.

Let us examine how Fact 6.11 can be rephrased using the Hopf parametrisation (Definition 3.2). Fix $o \in \Omega$. Let $(\xi, \eta, t) \in \text{Geod}_h^\infty(\Omega) \times \mathbb{R}$. If $\pi_h(\eta)$ is a smooth point of $\partial\Omega$, then by Fact 6.11,

$$\text{Hopf}_o(\xi', \eta, t) \in W^{ss}(\text{Hopf}_o(\xi, \eta, t)) \quad (6.4)$$

for any $\xi' \in \partial_h\Omega$ such that $(\xi', \eta) \in \text{Geod}_h^\infty(\Omega)$. The parametrisation of the unstable manifolds is more subtle. Using (3.4) and the fact that the unstable manifolds are the stable manifolds of the reversed flow, one can see that if $\pi_h\xi$ is smooth, then

$$\text{Hopf}_o(\xi, \eta', t + \rho_{\xi, \eta}^o(\eta')) \in W^{su}(\text{Hopf}_o(\xi, \eta, t)) \quad (6.5)$$

for any $\eta' \in \partial_h\Omega$ such that $(\xi, \eta') \in \text{Geod}_h(\Omega)$, where

$$\rho_{\xi, \eta}^o(\eta') := 2\langle \xi, \eta' \rangle_o - 2\langle \xi, \eta \rangle_o.$$

6.5. A cross-ratio on the boundary

Following the article [52], whose setting is that of negatively curved manifolds, let us define a cross-ratio, denoted by B , for four points on the boundary of a properly convex open set $\Omega \subset \mathbb{P}(V)$. It should not be confused with the cross-ratio of four aligned points of the projective space, denoted with brackets, and used to defined the Hilbert metric d_Ω (see (2.1)).

Recall that the cross-ratio of four points $\xi, \xi', \eta, \eta' \in \Omega$ is defined as

$$B(\xi, \xi', \eta, \eta') := d_\Omega(\xi, \eta) + d_\Omega(\xi', \eta') - d_\Omega(\xi, \eta') - d_\Omega(\xi', \eta). \quad (6.6)$$

Fix $o \in \Omega$. One can check that

$$B(\xi, \xi', \eta, \eta') = \rho_{\xi, \eta}^o(\eta') + \rho_{\xi', \eta'}^o(\eta), \quad (6.7)$$

and this implies that B extends continuously to the set of quadruples $(\xi, \xi', \eta, \eta') \in (\overline{\Omega}^h)^4$ such that (ξ, η) , (ξ', η) , (ξ, η') and (ξ', η') belong to $\text{Geod}_h \Omega$. See Figure 5 for a geometric interpretation using horospheres (which also works with spheres when ξ, ξ', η, η' belong to Ω).

The next lemma is classical and relates special values of B , called *periods*, with equation (2.5).

Lemma 6.12. *Let $\Omega \subset P(V)$ be a properly convex open set. Let $g \in \text{Aut}(\Omega)$ be a biproximal automorphism whose axis intersect Ω . Let $\xi \in \overline{\Omega}^h$ be such that (ξ, x_g^+) and (ξ, x_g^-) are in $\text{Geod}_h \Omega$ (recall that x_g^\pm is a smooth point of $\partial \Omega$ by Fact 2.13, so it identifies with its preimage in $\partial_h \Omega$). Then $B(x_g^+, x_g^-, \xi, g\xi) = 2\ell(g)$.*

Proof. By continuity of the cross-ratio, we can assume that $\xi \in \Omega$. Then using (6.6) or (6.7), one can show that

$$B(x_g^+, x_g^-, \xi, g\xi) = b_{x_g^+}(\xi, g\xi) + b_{x_g^-}(g\xi, \xi).$$

Now consider $x \in \Omega$ on the axis of g . We compute

$$\begin{aligned} b_{x_g^+}(\xi, g\xi) &= b_{x_g^+}(\xi, x) + b_{x_g^+}(x, gx) + b_{x_g^+}(gx, g\xi) \\ &= d_\Omega(x, gx) + b_{x_g^+}(\xi, x) + b_{g^{-1}x_g^+}(x, \xi) \quad (\text{by Lemma 4.3}) \\ &= \ell(g). \end{aligned}$$

By taking the inverse, we get

$$b_{x_g^-}(g\xi, \xi) = b_{x_{g^{-1}}^+}(g\xi, g^{-1}g\xi) = \ell(g^{-1}) = \ell(g). \quad \blacksquare$$

6.6. W^{ss} and W^{su} -invariant functions are essentially constant

In this section we prove that the W^{ss} and W^{su} -invariant functions on T^1M are essentially constant, and we derive as a corollary that the flow is ergodic and mixing. We will need the following result, about the local non-arithmeticity of the length spectrum; it is similar to [14, Prop. 4.1] but does not need the assumption of strong irreducibility.

Fact 6.13 ([17, Lem. 2.8]). Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Then the set of translation lengths $\ell(\gamma)$ of rank-one elements $\gamma \in \Gamma$ generates a dense subgroup of \mathbb{R} .

The previous fact can also be seen as a consequence of the more general result [15, Cor. 5.2.7].




Figure 5. Illustration of computations (6.8), where $d_\Omega(x, y) = B(\xi, \xi', \eta, \eta')$.

Proposition 6.14. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup with $M = \Omega/\Gamma$ divergent rank-one and non-elementary. Let m be the Sullivan measure on T^1M induced by a δ_Γ -conformal density $(\nu_x)_{x \in \Omega}$. Then any W^{ss} and W^{su} -invariant function on T^1M is essentially constant with respect to m .*

Proof. We fix $o \in \Omega$, and may assume that $\nu_o(\partial\Omega) = 1$. Let f be a W^{ss} and W^{su} -invariant function on T^1M . By Proposition 6.10, the measure m gives full measure to the (measurable) set $T^1M_{\text{sse}} \subset T^1M$ of vectors v with $\phi_\infty v, \phi_{-\infty} v \in \partial_{\text{sse}}\Omega$, hence it suffices to show that the restriction $f|_{T^1M_{\text{sse}}}$ is essentially constant. We lift $f|_{T^1M_{\text{sse}}}$, via the Hopf parametrisation, to a function \tilde{f} on $\{(\xi, \eta) \in \partial_{\text{sse}}\Omega : \xi \neq \eta\} \times \mathbb{R}$ that we extend to $\partial_{\text{sse}}\Omega^2 \times \mathbb{R}$ by setting $\tilde{f}(\xi, \xi) = 0$ for any ξ (recall that we abusively identify $\partial_{\text{sse}}\Omega$ with its preimage in $\partial_{\text{h}}\Omega$). Let us show that \tilde{f} is ν_o^2 times Lebesgue-essentially constant. Recall that ν_o is non-atomic by Propositions 4.8 and 6.2, thus for ν_o -almost all $\xi, \eta \in \partial_{\text{sse}}\Omega$ we have $\xi \neq \eta$.

The function f is W^{ss} and W^{su} -invariant, so by (6.4) and (6.5), for ν_o -almost $\xi, \xi', \eta, \eta' \in \partial_{\text{sse}}\Omega$ and Lebesgue-almost any $t \in \mathbb{R}$, the quantity $\rho_{\xi, \eta}(\eta')$ is well defined and

$$\tilde{f}(\xi, \eta, t) = \tilde{f}(\xi', \eta, t) = \tilde{f}(\xi, \eta', t + \rho_{\xi, \eta}(\eta')).$$

This implies in particular that there exist $\xi_0 \neq \eta_0 \in \partial_{\text{sse}}\Omega$ such that, if we denote $g(t) := \tilde{f}(\xi_0, \eta_0, t)$ for any $t \in \mathbb{R}$, then for ν_o -almost all $\xi, \eta \in \partial_{\text{sse}}\Omega$ and Lebesgue-almost any $t \in \mathbb{R}$, the quantity $\rho_{\xi_0, \eta_0}(\eta)$ is well defined and

$$\tilde{f}(\xi, \eta, t + \rho_{\xi_0, \eta_0}(\eta)) = g(t).$$

It is enough to establish that the measurable function g is essentially constant with respect to the Lebesgue measure. We denote by H the additive real subgroup consisting of numbers τ such that $g(t + \tau) = g(t)$ for Lebesgue-almost every $t \in \mathbb{R}$.

A classical result says that H is a closed subgroup of \mathbb{R} . (To see this first reduce to the case where g is bounded and then note that H is the stabiliser of $g(t)dt$ for the continuous action of \mathbb{R} on the space of Radon measures on \mathbb{R} .) To finish the proof of Proposition 6.14 it is enough, according to Fact 6.13, to prove that $2\ell(\gamma) \in H$ for any rank-one element $\gamma \in \Gamma$. To this end we observe that many cross-ratios belong to H : for ν_o -almost all $\xi, \eta, \xi', \eta' \in \partial_{\text{sse}}\Omega$ and Lebesgue-almost every $t \in \mathbb{R}$, the quantities $\rho_{\xi_0, \eta_0}(\eta)$, $\rho_{\xi, \eta}(\eta')$, $\rho_{\xi', \eta'}(\eta)$ and $B(\xi, \xi', \eta, \eta')$ are well defined and we have the following series of equalities (see Figure 5 for a geometric interpretation):

$$\begin{aligned}
 g(t) &= \tilde{f}(\xi, \eta, t + \rho_{\xi_0, \eta_0}(\eta)) \\
 &= \tilde{f}(\xi, \eta', t + \rho_{\xi_0, \eta_0}(\eta) + \rho_{\xi, \eta}(\eta')) \\
 &= \tilde{f}(\xi', \eta', t + \rho_{\xi_0, \eta_0}(\eta) + \rho_{\xi, \eta}(\eta')) \\
 &= \tilde{f}(\xi', \eta, t + \rho_{\xi_0, \eta_0}(\eta) + \rho_{\xi, \eta}(\eta') + \rho_{\xi', \eta'}(\eta)) \\
 &= \tilde{f}(\xi, \eta, t + \rho_{\xi_0, \eta_0}(\eta) + B(\xi, \xi', \eta, \eta')) \\
 &= g(t + B(\xi, \xi', \eta, \eta')).
 \end{aligned} \tag{6.8}$$

Consider a rank-one element $\gamma \in \Gamma$ and a point $\xi \in \partial_{\text{sse}}\Omega \cap \text{supp}(\nu_o) \setminus \{x_g^-, x_g^+\}$. Then $2\ell(\gamma)$ is equal to $B(x_\gamma^+, x_\gamma^-, \xi, \gamma\xi)$ by Lemma 6.12, and furthermore this quantity belongs to H because H is closed, B is continuous and $x_\gamma^+, x_\gamma^-, \xi, \gamma\xi$ are in $\text{supp}(\mu_o)$. ■

Corollary 6.15. *In the setting of Proposition 6.14, if Γ is divergent, then m is ergodic under the action of the geodesic flow. Furthermore, the Γ -quasi-invariant measures ν_o^2 and ν_o on $\partial\Omega^2$ and $\partial\Omega$ are ergodic under the action of Γ . In particular, the δ_Γ -conformal density is unique up to a scalar multiple. If moreover m is finite, then it is mixing.*

Proof. The ergodicity of m is a direct consequence of Fact 2.32, Proposition 6.6 and Proposition 6.14, which can be applied since m is Radon.

The ergodicity of ν_o^2 is a direct consequence of that of m , since there is correspondence between Γ -invariant measurable subsets of $\text{Geod}^\infty\Omega$ (which has full ν_o^2 -measure) and $(\phi_t)_t$ -invariant measurable subsets of T^1M , sending sets with full (resp. null) measure to sets with full (resp. null) measure. For similar reasons, the ergodicity of ν_o is a direct consequence of that of ν_o^2 .

Let $(\nu'_x)_{x \in \Omega}$ be another δ_Γ -conformal density such that $\nu'_o(\partial\Omega) = \nu_o(\partial\Omega)$. Then the family $(\nu_x + \nu'_x)_{x \in \Omega}$ is also a conformal density. The measures ν_o and ν'_o are absolutely continuous with respect to $\nu_o + \nu'_o$; the Radon–Nikodym derivatives are Γ -invariant, by definition of conformal densities, and hence constant $(\nu_o + \nu'_o)$ -almost surely by ergodicity. Thus $\nu_o = \nu'_o$.

If m is finite, then it is mixing by Fact 2.33 and Proposition 6.14. ■

6.7. The support of conformal densities

Let us establish that the support of the δ_Γ -conformal density, for Γ divergent and $M = \Omega/\Gamma$ rank-one, is exactly the proximal limit set. This is a consequence of conservativity and ergodicity of the Bowen–Margulis measure, Fact 2.30 (4), and the fact that $T^1 M_{\text{bip}}$ is maximal among the flow-invariant closed subsets of $T^1 M$ on which the geodesic flow is forward topologically transitive (i.e. has a dense forward orbit); this last result was proved in [14].

Proposition 6.16. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a divergent discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Then Λ^{prox} is the support of any δ_Γ -conformal density and $T^1 M_{\text{bip}}$ is the support of any Bowen–Margulis measure.*

Proof. Let $(v_x)_{x \in \Omega}$ be the δ_Γ -conformal density on $\partial\Omega$, and $o \in \Omega$. Since Λ^{prox} is the smallest Γ -invariant closed subset of $\bar{\Omega}$ (Fact 2.18) and v_o is Γ -quasi-invariant, we have

$$\Lambda^{\text{prox}} \subset \text{supp}(v_o) \quad \text{and} \quad T^1 M_{\text{bip}} \subset \text{supp}(m).$$

Since m is conservative and ergodic by Proposition 6.6 and Corollary 6.15, it is implied by Fact 2.30 (4) that the geodesic flow is forward topologically transitive on $\text{supp}(m)$ (i.e. has a dense forward orbit). According to [14, Thm. 1.2], this implies that

$$\text{supp}(m) = T^1 M_{\text{bip}}.$$

Suppose by contradiction there is a point $\xi \in \text{supp}(v_o) \setminus \Lambda^{\text{prox}}$. Since M is rank-one, we can find a strongly extremal point $\eta \in \Lambda^{\text{prox}}$. Then we can find a neighbourhood $U \times V \subset \text{Geod}^\infty(\Omega)$ of (ξ, η) such that $U \cap \Lambda^{\text{prox}}$ is empty. Note that $v_o(U)v_o(V) > 0$ since $\xi, \eta \in \text{supp}(v_o)$. Hence, m gives non-zero measure to the set of vectors v with lifts $\tilde{v} \in T^1\Omega$ such that $(\phi_{-\infty}v, \phi_\infty v) \in U \times V$; but this set is not contained in $T^1 M_{\text{bip}}$: this is a contradiction. ■

6.8. Proof of Theorem 1.6 (2)

Theorem 1.6 (2) is a direct consequence of Propositions 4.7, 4.8, 6.6, 6.10 and 6.16, Corollary 6.15 and Fact 2.29. Observe that $(\partial\Omega^2, \nu_o^2, \Gamma)$ being conservative and ergodic is due to the fact that $\text{Geod}^\infty(\Omega) \subset \partial\Omega^2$ has full ν_o^2 -measure, since ν_o is nonatomic and gives full measure to the set of strongly extremal points.

7. Preparatory results on convex cocompact projective manifolds

In real hyperbolic geometry, many properties of compact manifolds are actually also true for a broader class of manifolds: the convex cocompact ones, and the proofs of these properties generally work verbatim. This observation remains valid in convex projective geometry, thanks to the recent notion of convex cocompactness developed by Danciger, Guéritaud and Kassel in this setting (see Definition 1.2). Note that most of the results of the remainder of the paper concern convex cocompact actions. The present section gathers results on convex cocompact actions, and in particular more precise Shadow lemmas. An important result of Danciger–Guéritaud–Kassel that we will need is the following.

Fact 7.1 ([35, Prop. 5.10]). Let $\Omega \in \mathcal{P}(V)$ be a properly convex open set and $\Gamma \subset \mathrm{PGL}(V)$ a discrete subgroup. Suppose that Γ acts convex cocompactly on Ω and Ω^* . Then $\bar{\Omega} \setminus \Lambda_{\Omega}^{\mathrm{orb}}(\Gamma)$ has bisaturated boundary, in the sense that $[\xi, \eta] \cap \Omega$ is non-empty for all $\xi \in \Lambda^{\mathrm{orb}}$ and $\eta \in \bar{\Omega} \setminus \Lambda^{\mathrm{orb}}$.

Observe that, given a strongly irreducible discrete subgroup $\Gamma \subset \mathrm{PGL}(V)$ that acts convex cocompactly on a properly convex open $\Omega \subset \mathcal{P}(V)$, if we want to understand the dynamics of the geodesic flow on $(T^1\Omega/\Gamma)_{\mathrm{cor}}$, then we can assume that Γ also acts convex cocompactly on Ω^* , since otherwise we may take another Γ -invariant properly convex open set Ω' such that Γ acts convex cocompactly on both Ω' and $(\Omega')^*$, and we have $(T^1\Omega/\Gamma)_{\mathrm{cor}} = (T^1\Omega'/\Gamma)_{\mathrm{cor}}$ (see [35, Prop. 5.10 and Cor. 4.17]).

7.1. Proof of Proposition 1.7

Consider $o \in \mathcal{C}_{\Omega}^{\mathrm{cor}}(\Gamma)$. We set $\mathrm{Vol} := \sum_{\gamma \in \Gamma} \mathcal{D}_{\gamma o}$, where \mathcal{D}_x denotes the Dirac mass at $x \in \Omega$; this defines a Γ -invariant Radon measure on Ω , which is supported on $\mathcal{C}_{\Omega}^{\mathrm{cor}}(\Gamma)$. We apply Fact 2.22 to Vol , in order to obtain a δ_{Γ} -conformal density ν_o on $\partial\Omega$ which is supported on Λ^{orb} . Recall that $\Lambda^{\mathrm{orb}} = \Lambda^{\mathrm{con}}$ (see Section 1.2), hence $\nu_o(\Lambda^{\mathrm{con}}) = 1$ and therefore Γ is divergent by Theorem 1.6. By the same theorem, the Bowen–Margulis measure m is supported on T^1M_{cor} . Since m is Radon and T^1M_{cor} is compact, m must be finite.

7.2. A Shadow lemma for convex cocompact actions

In this section we prove a Shadow lemma for when Γ satisfies the stronger assumption that it acts convex cocompactly on Ω . The main interest of the new Shadow lemma is that it estimates the measure of *small* shadows.

We denote $\mathrm{ray}_x(A) = \bigcup_{\xi \in A} [x, \xi)$ for $x \in \Omega$ and $A \subset \partial\Omega$.

Lemma 7.2. *Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup. Suppose that Γ is strongly irreducible and $T^1 M_{\text{bip}}$ is non-empty, or that M is rank-one and non-elementary. Consider $\delta \geq 0$ and a δ -conformal density $(\nu_x)_{x \in \Omega}$ on $\partial\Omega$. Let $K \subset \Omega$ be a compact subset and $\mathcal{C} = \Gamma \cdot K$. Then there exists $R_0 > 0$ such that for any $R > 0$, we can find a constant $C = C(R) > 0$ such that for all $x, y \in \mathcal{C}$,*

$$C^{-1} e^{-\delta d_\Omega(x,y)} \leq \nu_x(\mathcal{O}_{R_0+R}(x, y)) \leq \nu_x(\mathcal{O}_{R_0+R}^+(x, y)) \leq C e^{-\delta d_\Omega(x,y)},$$

and if furthermore $y \in \text{ray}_x(\text{supp}(\nu_o))$, then

$$\nu_y(\mathcal{O}_R(x, y)) \geq C^{-1} \quad \text{and} \quad \nu_x(\mathcal{O}_R(x, y)) \geq C^{-1} e^{-\delta d_\Omega(x,y)}.$$

Proof. Let $R > 0$. By definition of the conformal density, $\nu_x \leq e^{\delta d_\Omega(x,y)} \nu_y$ for any $x, y \in \Omega$. Using the triangular inequality and Fact 2.1, it is elementary to see that for all $x, y, x', y' \in \Omega$,

$$\mathcal{O}_R^\alpha(x, y) \subset \mathcal{O}_{R+d_\Omega(x,x')+d_\Omega(y,y')}^\alpha(x', y') \quad \text{for } \alpha = \emptyset \text{ or } +.$$

By Γ -equivariance it is enough to prove the lemma for $x \in K$. Let D be the diameter of K for the Hilbert metric, and fix $o \in K$. For any $y \in \mathcal{C}$, there exists $\gamma \in \Gamma$ such that $d_\Omega(y, \gamma o) \leq D$, and then for any $x \in K$,

$$\begin{aligned} e^{-\delta \Gamma D} \nu_o(\mathcal{O}_R(o, \gamma o)) &\leq \nu_x(\mathcal{O}_{R+2D}(x, y)) \\ &\leq \nu_x(\mathcal{O}_{R+2D}^+(x, y)) \leq e^{\delta \Gamma D} \nu_o(\mathcal{O}_{R+4D}^+(o, \gamma o)). \end{aligned}$$

Now we can use the Shadow lemma (Lemma 4.2) to bound the right-most and left-most terms when R is greater than some R_0 , and we obtain the first estimate.

Let us show the second estimate. We set

$$\varepsilon := \inf\{\nu_x(\mathcal{O}_R(y, x')) : x \in K, y \in \Omega, x' \in K \cap \text{ray}_y(\text{supp } \nu_o)\} > 0.$$

We then make the same computations as in the Shadow lemma (Lemma 4.2). We take a δ -conformal density $(\mu_x)_{x \in \Omega}$ on $\partial_{\text{h}}\Omega$ such that $(\pi_{\text{hor}})_* \mu_x = \nu_x$ for any $x \in \Omega$. Let $x \in K$ and $y \in \text{ray}_x(\text{supp}(\nu_o)) \cap \mathcal{C}$; by definition of K there exists $\gamma \in \Gamma$ such that $\gamma^{-1}y \in K$. Then

$$\begin{aligned} \nu_x(\mathcal{O}_R(x, y)) &= \mu_x(\pi_{\text{hor}}^{-1}(\mathcal{O}_R(x, y))) \\ &= \mu_{\gamma^{-1}x}(\pi_{\text{hor}}^{-1}(\mathcal{O}_R(\gamma^{-1}x, \gamma^{-1}y))) \\ &= \int_{\pi_{\text{hor}}^{-1}(\mathcal{O}_R(\gamma^{-1}x, \gamma^{-1}y))} e^{-\delta b_{\tilde{\xi}}(\gamma^{-1}x, \gamma^{-1}y)} d\mu_x(\tilde{\xi}) \\ &\geq \int_{\pi_{\text{hor}}^{-1}(\mathcal{O}_R(\gamma^{-1}x, \gamma^{-1}y))} e^{-\delta d_\Omega(x,y)} d\mu_x(\tilde{\xi}) \\ &\geq \nu_x(\mathcal{O}_R(\gamma^{-1}x, \gamma^{-1}y)) e^{-\delta d_\Omega(x,y)} \\ &\geq \varepsilon e^{-\delta d_\Omega(x,y)}. \end{aligned}$$

■

As an immediate corollary, we obtain the following.

Corollary 7.3. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact and discrete subgroup. Suppose that Γ is strongly irreducible and T^1M_{bip} is non-empty, or that M is rank-one and non-elementary. Consider $\delta \geq 0$ and a δ -conformal density $(\nu_x)_{x \in \Omega}$ on $\partial\Omega$. Then there exists $R_0 > 0$ such that for any $R > 0$, we can find a constant $C = C(R) > 0$ such that for all $x, y \in \mathcal{E}^{\text{cor}}$,*

$$C^{-1}e^{-\delta d_\Omega(x,y)} \leq \nu_x(\mathcal{O}_{R_0+R}(x,y)) \leq Ce^{-\delta d_\Omega(x,y)},$$

and if furthermore $y \in \text{ray}_x(\text{supp}(\nu_o))$, then

$$\nu_y(\mathcal{O}_R(x,y)) \geq C^{-1} \quad \text{and} \quad \nu_x(\mathcal{O}_R(x,y)) \geq C^{-1}e^{-\delta d_\Omega(x,y)}.$$

7.3. Non-straight closed geodesics

In this section we investigate non-rank-one automorphisms of properly convex open sets which realise their translation length.

7.3.1. Some standard facts. The following fact says that the only case where a non-straight geodesic can appear is the one shown in Figure 1.

Fact 7.4. Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set. For $x, y \in \Omega$ distinct, consider (a_{xy}, b_{xy}) in $\partial\Omega^2$ such that a_{xy}, x, y, b_{xy} are aligned in this order. Let $x, y, z \in \Omega$ be pairwise distinct. Then $d_\Omega(x, z) = d_\Omega(x, y) + d_\Omega(y, z)$ if and only if $a_{xz} \in [a_{xy}, a_{yz}]$ and $b_{xz} \in [b_{xy}, b_{yz}]$.

This property can be used to prove the following.

Fact 7.5 ([37]). Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, $I \subset \mathbb{R}$ a non-trivial interval and $c: I \rightarrow \Omega$ an isometric embedding. For all $t < s \in I$, consider (a_{ts}, b_{ts}) in $\partial\Omega^2$ such that $a_{ts}, c(t), c(s), b_{ts}$ are aligned in this order. Let F_+ (resp. F_-) be the smallest closed face of Ω that contains $\{b_{ts} : t < s \in I\}$ (resp. $\{a_{ts} : t < s \in I\}$). Then F_+ and F_- are proper faces of Ω , whose dimension is the dimension of the convex hull of $c(I)$ minus 1.

Moreover, if $\sup I = \infty$ (resp. $\inf I = -\infty$), then $(c(t))_t$ converges to a point of F_+ (resp. F_-) when t goes to $+\infty$ (resp. $-\infty$).

The following generalises the fact that $x_g^+ \in \partial\Omega$ and $x_g^+ \oplus x_g^0 \cap \Omega = \emptyset$ for any biproximal automorphism g of a properly convex open set Ω .

Fact 7.6. Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set. Let $g \in \mathbb{P}(\text{End}(V))$ be in the closure of $\text{Aut}(\Omega)$, then its kernel and its image intersect $\bar{\Omega}$ but not Ω .

7.3.2. Analysis of an automorphism that attains its translation length.

Lemma 7.7. *Let Ω be a properly convex open set. Let $g \in \text{Aut}(\Omega)$ with $\ell(g) > 0$ and suppose there exists $x \in \Omega$ such that $d_\Omega(x, gx) = \ell(g)$. Consider $a, b \in \partial\Omega$ such that a, x, gx, b are aligned in this order. Then*

- (1) *the path $c: \mathbb{R} \rightarrow \Omega$, defined by $c(t) \in [g^n x, g^{n+1} x]$ and $d_\Omega(c(t), g^n x) = t - n$ for all $n \in \mathbb{Z}$ and $t \in [n, n + 1]$, is a geodesic;*
- (2) *the restriction of g to $x_g^+ \oplus x_g^-$ is diagonalisable over \mathbb{C} ;*
- (3) *the restriction of g to the span $\mathbb{P}(W)$ of $\{g^n x\}_n$ is biproximal;*
- (4) *$x_g^+ \oplus x_g^0 \cap \mathbb{P}(W) \cap \bar{\Omega}$ (resp. $x_g^- \oplus x_g^0 \cap \mathbb{P}(W) \cap \bar{\Omega}$) is the smallest g -invariant closed face of Ω that contains b (resp. a);*
- (5) *if $x, gx, g^2 x$ are not aligned, then $x_g^0 \cap \bar{\Omega}$ is non-empty;*
- (6) *if g is not rank-one, then $d_{\text{spl}}(a, b) = 2$.*

Proof. Let us check that (1) holds. Consider three real numbers $r \leq s \leq t$, pick three integers $k \leq n \leq m$ such that $k \leq r \leq k + 1$ and $n \leq s \leq n + 1$ and $m \leq t \leq m + 1$. By (2.5), we have

$$\begin{aligned}
 d_\Omega(c(r), c(t)) &\geq d_\Omega(g^k x, g^{m+1} x) - d_\Omega(g^k x, c(r)) - d_\Omega(c(t), g^{m+1} x) \\
 &\geq \ell(g^{m+1-k}) - d_\Omega(g^k x, c(r)) - d_\Omega(c(t), g^{m+1} x) \\
 &= \sum_{i=k}^m d_\Omega(g^i x, g^{i+1} x) - d_\Omega(g^k x, c(r)) - d_\Omega(c(t), g^{m+1} x) \\
 &= d_\Omega(c(r), g^{k+1} x) + \sum_{i=k+1}^{m-1} d_\Omega(g^i x, g^{i+1} x) + d_\Omega(g^m x, c(t)) \\
 &\geq d_\Omega(c(r), c(s)) + d_\Omega(c(s), c(t)).
 \end{aligned}$$

For all distinct pair of points $(y, z) \in \bar{\Omega}^2$, let us denote by $s_y^z \in \partial\Omega$ the point such that y, z, s_y^z are aligned in this order. Let $\mathbb{P}(W) \subset \mathbb{P}(V)$ be the span of $\{g^n x\}_{n \in \mathbb{Z}}$, with dimension $k \geq 1$; we set $\Omega' = \Omega \cap \mathbb{P}(W)$. By Fact 7.5, the smallest closed face of Ω that contains $\{g^n b\}_{n \in \mathbb{Z}}$ (resp. $\{g^n a\}_{n \in \mathbb{Z}}$), denoted by F_+ (resp. F_-), is proper, and therefore its span $\mathbb{P}(W_+)$ (resp. $\mathbb{P}(W_-)$) has dimension $k - 1$. Moreover, $s_{g^n x}^{g^m x} \in F_+$ (resp. F_-) for all $m > n$ (resp. $m < n$), and $(g^n x)_n$ converges to some point $\xi_+ \in F_+ \cap x_g^+$ (resp. $\xi_- \in F_- \cap x_g^-$), which is fixed by g , when n goes to $+\infty$ (resp. $-\infty$).

Consider a lift $\tilde{g} \in \text{GL}(V)$ of g that preserves one connected component $C \subset V \setminus \{0\}$ of the preimage of Ω , and such that $\lambda_1(\tilde{g}) = 1$. Let us examine the Jordan normal form of \tilde{g} : there exists $\ell \geq 0$ and a decomposition

$$\tilde{g} = u_1 + \cdots + u_\alpha - (u'_1 + \cdots + u'_{\alpha'}) + r_1^{\theta_1} v_1 + \cdots + r_\beta^{\theta_\beta} v_\beta + h, \quad (7.1)$$

which satisfies the following. The product of any two matrices in this sum is zero. The integers $\alpha, \beta \geq 0$ are not both zero. The sequence $((1/n^\ell)h^n)_n$ tends to zero. The matrices $u_1, \dots, u_\alpha, u'_1, \dots, u'_{\alpha'}$ are all conjugate to the matrix with zeros everywhere except on the upper-left block of size $\ell + 1$, which is the exponential of the upper-triangular matrix whose (i, j) -entry is 1 if $j = i + 1$ and zero otherwise. For $\theta \in \mathbb{R}$ and $1 \leq i \leq \beta$, the matrices r_i^θ and v_i are simultaneously conjugate to the matrices with zeros everywhere except on the upper-left block of size $2\ell + 2$, where they are respectively of the form

$$\begin{pmatrix} \text{rot}^\theta & & & \\ & \ddots & & \\ & & & \text{rot}^\theta \end{pmatrix} \quad \text{and} \quad \exp \begin{pmatrix} 0 & I_2 & & \\ & \ddots & \ddots & \\ & & \ddots & I_2 \\ & & & 0 \end{pmatrix},$$

with

$$\text{rot}^\theta = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \quad \text{and} \quad I_2 = \text{rot}^0.$$

Finally, $\theta_1, \dots, \theta_\beta \in \mathbb{R} \setminus \pi\mathbb{Z}$. Let

$$\begin{aligned} \bar{u}_i &= \lim_{n \rightarrow \infty} \frac{\ell!}{n^\ell} u_i^n, & \bar{u}'_i &= \lim_{n \rightarrow \infty} \frac{\ell!}{n^\ell} u_i'^n \quad \text{for } 1 \leq i \leq \alpha, \\ \bar{v}_i &= \lim_{n \rightarrow \infty} \frac{\ell!}{n^\ell} v_i^n & & \text{for } 1 \leq i \leq \beta; \end{aligned}$$

the set of accumulation points of $(r_i^{n\theta_i})_n$ is $\{r_i^\theta : \theta \in \overline{\theta_i\mathbb{Z} + 2\pi\mathbb{Z}}\}$ for $1 \leq i \leq \beta$. Let $\tilde{x} \in C$ be a lift of $x \in \Omega$. The accumulation points of $((\ell!/n^\ell)\tilde{g}^n \tilde{x})_n$ are

$$\left\{ \sum_i \bar{u}_i \tilde{x} + (-1)^m \sum_i \bar{u}'_i \tilde{x} + \sum_i r_i^{\theta'_i} \bar{v}_i \tilde{x} : m \in \{0, 1\}, \theta'_i \in \overline{\theta_i(2\mathbb{Z} + m) + 2\pi\mathbb{Z}} \right\},$$

which are non-zero by Fact 7.6. Since $(g^n x)_n$ converges to ξ^+ , these accumulation points are all in $\xi^+ \cap C$, and this implies that $\bar{u}'_i \tilde{x} = 0$ for $1 \leq i \leq \alpha'$ and $\bar{v}_i \tilde{x} = 0$ for $1 \leq i \leq \beta$. Up to considering another basis, we can assume that $\bar{u}_i \tilde{x} = 0$ for any $2 \leq i \leq \alpha$.

The element \tilde{g} commutes with every matrix of its decomposition (7.1), and hence also with $\{\bar{u}_i\}_{1 \leq i \leq \alpha}$, $\{\bar{u}'_i\}_{1 \leq i \leq \alpha'}$, $\{r_i^\theta\}_{1 \leq i \leq \beta, \theta \in \mathbb{R}}$, and $\{\bar{v}_i\}_{1 \leq i \leq \beta}$. Thus $\{\bar{u}_i \tilde{g}^n \tilde{x}\}_{2 \leq i \leq \alpha}$, $\{\bar{u}'_i \tilde{g}^n \tilde{x}\}_{1 \leq i \leq \alpha'}$, and $\{\bar{v}_i \tilde{g}^n \tilde{x}\}_{1 \leq i \leq \beta}$ are zero for any $n \geq 0$. By construction of W , all elements $\{\bar{u}_i\}_{2 \leq i \leq \alpha}$, $\{\bar{u}'_i\}_{1 \leq i \leq \alpha'}$, and $\{\bar{v}_i\}_{1 \leq i \leq \beta}$ are zero on W .

Suppose by contradiction that the restriction of g to x_g^+ is not diagonalisable over \mathbb{C} , i.e. that $\ell > 0$. Then $\bar{u}_1 \tilde{\xi}_+ = 0$ for any lift $\tilde{\xi}_+ \in W$ of ξ_+ . This, and the fact that $\bar{u}_1 \tilde{x} \neq 0$, imply that $\bar{u}_1 \tilde{y} \neq 0$ for any lift $\tilde{y} \in C$ of $y := s_{\xi_+}^x \in F_-$. As

a consequence, $(g^n y)_n$ converges to ξ_+ . Since F_- is closed, it contains ξ_+ , as well as x : a contradiction. For the same reasons, the restriction of g to x_g^- is diagonalisable over \mathbb{C} , and this concludes the proof of (2).

The sequence of restrictions $(\tilde{g}_{|W}^n)_n$ converges to $\bar{u}_{1|W}$ which is proximal because it is a projector with rank 1. Therefore, $\tilde{g}_{|W}^n$ is proximal for n large enough, and so is $\tilde{g}_{|W}$. For the same reasons, $\tilde{g}_{|W}^{-1}$ is proximal, and this concludes the proof of (3).

In order to establish (4), it is enough to prove that F_+ (resp. F_-) is contained in $x_g^+ \oplus x_g^0$ (resp. $x_g^- \oplus x_g^0$), since its dimension is $k - 1$. Pick $\xi \in F_+$; the sequence $(g^n \xi)_{n \geq 0}$ cannot converge to ξ_- since this point does not belong to F_+ . That $g_{|P(W)}$ is biproximal implies that $\xi \in x_g^+ \oplus x_g^0$. The proof of $F_- \subset x_g^- \oplus x_g^0$ is similar.

Suppose that x, gx, g^2x are not aligned. Then $b \in F_+ \setminus \{\xi_+\} \subset x_g^+ \oplus x_g^0 \setminus x_g^+$, hence $(g^{-n}b)_n$ accumulates in x_g^0 . This proves (5).

Suppose that g is not rank-one. Since $x \in [a, b] \cap \Omega$, the simplicial distance between a and b is at least 2. If x, gx, g^2x are aligned, then $a = \xi_-$ and $b = \xi_+$ are fixed by g , and $d_{\text{spl}}(a, b) \leq 2$ by Corollary 6.9. Otherwise, there is $\xi \in x_g^0 \cap \partial\Omega'$, and $d_{\text{spl}}(a, b) \leq d_{\text{spl}}(a, \xi) + d_{\text{spl}}(\xi, b) \leq 2$ since $\xi \in F_+ \cap F_-$. This proves (6). ■

7.4. Non-straight closed geodesics on convex cocompact manifolds

Let $\Gamma \subset \text{PGL}(V)$ be a discrete subgroup that preserves a properly convex open set $\Omega \subset P(V)$; set $M := \Omega/\Gamma$. In Section 2.7, we have associated to each periodic geodesic of M a conjugacy class of Γ . When Γ acts convex cocompactly on Ω , one can produce a weak converse of this; to each automorphism of $\gamma \in \Gamma$, we are able to associate a geodesic segment of length $\ell(\gamma)$, which is closed on M and is freely homotopic to γ , but which is not necessarily closed in T^1M .

Fact 7.8 ([35, Prop. 10.4]). Consider a discrete subgroup $\Gamma \subset \text{PGL}(V)$ that acts convex cocompactly on a properly convex open sets $\Omega \subset P(V)$. Then for any $\gamma \in \Gamma$, there exists $x \in \mathcal{C}_{\Omega}^{\text{cor}}(\Gamma)$ such that $d_{\Omega}(x, \gamma x) = \ell(\gamma)$.

The following corollary ensures the closed curve can be taken in T^1M_{cor} .

Corollary 7.9. *Let $\Gamma \subset \text{PGL}(V)$ be a convex cocompact discrete subgroup. Consider a Γ -invariant properly convex open set $\Omega \subset P(V)$ such that Γ acts convex cocompactly on Ω and Ω^* ; set $M = \Omega/\Gamma$. Let $\gamma \in \Gamma$ and $v \in T^1\Omega$ be such that $\ell(\gamma) > 0$ and $\gamma\pi v = \pi\phi_{\ell(\gamma)}v$. Then $\phi_{\pm\infty}v \in \Lambda^{\text{orb}}$ and $\pi_{\Gamma}v \in T^1M_{\text{cor}}$.*

Proof. According to Fact 7.1, the properly convex set $\bar{\Omega} \setminus \Lambda^{\text{orb}}$ has bisaturated boundary (i.e. $[\xi, \eta] \cap \Omega \neq \emptyset$ for all $\xi \in \Lambda^{\text{orb}}$ and $\eta \in \bar{\Omega} \setminus \Lambda^{\text{orb}}$). By Lemma 7.7, $\phi_{\pm\infty}v \in x_{\gamma}^{\pm} \oplus x_{\gamma}^0 \cap \partial\Omega$, hence the segment between $\phi_{\pm\infty}v$ and the limit of $(\gamma^{\pm n}\pi v)_n$ is contained in $\partial\Omega$. This implies that $\phi_{\pm\infty}v \in \Lambda^{\text{orb}}$ since $\bar{\Omega} \setminus \Lambda^{\text{orb}}$ has bisaturated boundary and since $\lim_{n \rightarrow \infty} \gamma^n \pi v \in \Lambda^{\text{orb}}$. Thus v belongs to T^1M_{cor} . ■

The next corollary ensures that when Γ is convex cocompact, if $(T^1\Omega_{\max}/\Gamma)_{\text{bip}}$ is non-empty then we can find $\Omega \subset \Omega_{\max}$ such that Ω/Γ is rank-one. This corollary generalises results of Islam [40, Lem. 6.4] and Zimmer [64, Thm. 7.1] for compact convex projective manifolds; one may also compare it to a remark of Islam [40, Rem. A.1.C] which concerns not necessarily compact convex projective manifolds with a compact convex core.

Corollary 7.10. *Let $\Gamma \subset \text{PGL}(V)$ be a discrete subgroup that acts convex cocompactly on a properly convex open set $\Omega \subset \text{P}(V)$.*

- *Any biproximal element of Γ whose dual axis in $\text{P}(V^*)$ intersects Ω^* is rank-one; in particular, if $T^1(\Omega^*/\Gamma)_{\text{bip}} \neq \emptyset$, then Ω/Γ is rank-one.*
- *Suppose that Γ acts convex cocompactly on Ω^* , then any biproximal element of Γ whose axis intersects Ω is rank-one; in particular, if $(T^1\Omega/\Gamma)_{\text{bip}} \neq \emptyset$, then Ω/Γ is rank-one.*

Proof. The first point is an immediate consequence of Lemma 7.7 (5) and Fact 7.8. Let us establish the second point. Let $\gamma \in \Gamma$ be biproximal with $\text{Axis}(\gamma) \cap \Omega \neq \emptyset$. Since Γ acts convex cocompactly on Ω^* , the first point implies that γ , seen as a rank-one automorphism of Ω^* , is rank-one. Thus, γ is a rank-one automorphism of Ω by Fact 2.14. ■

7.5. A closing lemma

In this section we state a closing lemma, generalising [22, Thm. 4.4] and weaker than the classical one from Anosov [2, Lem. 13.1]. We briefly recall the idea: whenever a geodesic segment comes back sufficiently close to its starting point (no matter how long it is), we can find closed geodesic which tracks it. The following version is a more geometrical formulation of the closing lemma. We state the dynamical version below.

Lemma 7.11. *Let $\Omega \subset \text{P}(V)$ be a properly convex open set, $x \in \Omega$ and $(\xi_-, \xi_+) \in \partial\Omega^2$. Assume that $d_{\text{spl}}(\xi_+, \bar{F}_\Omega(\xi_-)) \geq 2$ and $d_{\text{spl}}(\xi_-, \bar{F}_\Omega(\xi_+)) \geq 2$. Then there exists $R > 0$ such that for any neighbourhood W of $\bar{B}_\Omega(\xi_-, R) \times \bar{B}_\Omega(\xi_+, R)$ in $\bar{\Omega}^2$, there exists a neighbourhood U of (ξ_-, ξ_+) such that for any $g \in \text{Aut}(\Omega)$, if $(g^{-1}x, gx) \in U$, then g is biproximal and $(x_g^-, x_g^+) \in W$.*

To prove the previous result, we need the following generalisation of the fact that projective transformations with an attracting fixed point are proximal.

Fact 7.12 ([13]). *Let $g \in \text{PGL}(V)$ contract a properly convex open set, in the sense that there exists a properly convex open set $\Omega \subset \text{P}(V)$ such that $g(\bar{\Omega}) \subset \Omega$. Then g is proximal.*

Proof of Lemma 7.11. The case where $\dim(V) = 2$ is trivial, and we assume that $\dim(V) \geq 3$.

(1) By assumption, we can find $R > 0$ large enough so that $\mathcal{O}_R(\xi_{\pm}, x)$ contains $\bar{F}_{\Omega}(\xi_{\mp})$.

(2) By lower semi-continuity of $d_{\bar{\Omega}}$, we may find a neighbourhood

$$U_{\pm} \subset \bar{\Omega} \setminus \bar{B}_{\Omega}(x, R)$$

of ξ_{\pm} such that $\mathcal{O}_R(\xi, x)$ contains $\bar{\mathcal{O}}_R(x, y)$ for all $\xi \in U_{\pm}$ and $y \in \Omega \cap U_{\mp}$.

(3) Our assumption ensures that $d_{\text{spl}}(\xi, \bar{F}_{\Omega}(\xi_{\mp})) \geq 2$ for any $\xi \in \bar{B}_{\bar{\Omega}}(\xi_{\pm}, R)$, so we can find $R' \geq R$ large enough so that $\mathcal{O}_{R'}(\xi, x)$ contains $\bar{F}_{\Omega}(\xi_{\mp})$ for any $\xi \in \bar{B}_{\bar{\Omega}}(\xi_{\pm}, R)$.

(4) By lower semi-continuity of $d_{\bar{\Omega}}$, we may find a neighbourhood

$$W_{\pm} \subset \bar{\Omega} \setminus \bar{B}_{\Omega}(x, R')$$

of $\bar{B}_{\bar{\Omega}}(\xi_{\pm}, R)$ such that $\mathcal{O}_{R'}(\xi, x)$ contains $\bar{\mathcal{O}}_{R'}(\xi, y)$ for all $\xi \in W_{\pm}$ and $y \in \Omega \cap W_{\mp}$, and such that $W_- \times W_+ \subset W$.

(5) Take a neighbourhood $U'_{\pm} \subset U_{\pm}$ of ξ_{\pm} such that $\bar{\mathcal{O}}_R(x, y)$ is contained in W_{\pm} for any $y \in \Omega \cap U'_{\pm}$.

Consider $g \in \text{Aut}(\Omega)$ such that $g^{\pm 1}x \in U'_{\pm}$, and let us show that g is biproximal with (x_g^-, x_g^+) in W . By (2) and since $gx \in U_+$ and $g^{-1}x \in U_-$, we have

$$g\bar{\mathcal{O}}_R(g^{-1}x, x) = \bar{\mathcal{O}}_R(x, gx) \subset \mathcal{O}_R(g^{-1}x, x).$$

Hence, g fixes some point $\eta_+ \in \bar{\mathcal{O}}_R(x, gx)$ by the Brouwer fixed point theorem. Symmetrically, g fixes some point $\eta_- \in \bar{\mathcal{O}}_R(x, g^{-1}x)$.

By (5), the point η_+ lies in W_+ , and η_- lies in W_- . By (4), this implies that

$$g\bar{\mathcal{O}}_{R'}(\eta_-, x) = \bar{\mathcal{O}}_{R'}(\eta_-, gx) \subset \mathcal{O}_{R'}(\eta_-, x).$$

Therefore, according to Fact 7.12, the projection $g' \in \text{PGL}(V/\eta_-)$ of g is proximal, and its attracting fixed point corresponds in $\text{P}(V)$ to a line of the form $\eta_- \oplus \zeta_+$, where $\zeta_+ \in \bar{\mathcal{O}}_{R'}(\eta_-, gx)$ is fixed by g .

By Fact 7.6, since $\ell(g) \geq \ell(g') > 0$ and since $\eta_- \oplus \zeta_+$ intersects Ω , we either have $(\eta_-, \zeta_+) \in x_g^- \times x_g^+$ or $(\eta_-, \zeta_+) \in x_g^+ \times x_g^-$. The latter case contradicts the fact that $\dim(V) \geq 3$ and g' is proximal. Hence $\eta_- \in x_g^-$ and $\zeta_+ \in x_g^+$, and g is proximal with $\zeta_+ = x_g^+$. Symmetrically, g^{-1} is also proximal and $\eta_+ \in x_g^+$. We have proved that g is biproximal with $(x_g^-, x_g^+) = (\eta_-, \eta_+) \in W$. ■

Corollary 7.13. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set. Let $x \in \Omega$, $(\xi_-, \xi_+) \in \partial\Omega^2$ and W be a neighbourhood of (ξ_-, ξ_+) . Then there exists a neighbourhood U of (ξ_-, ξ_+) such that for any $g \in \text{Aut}(\Omega)$ with $(g^{-1}x, gx) \in U$,*

- *if $d_{\text{spl}}(\xi_-, \xi_+) \geq 3$, then g is rank-one;*
- *if ξ_- and ξ_+ are extremal and $d_{\text{spl}}(\xi_-, \xi_+) \geq 2$, then g is biproximal and $(x_g^-, x_g^+) \in W$;*
- *if ξ_- and ξ_+ are distinct and strongly extremal, then g is rank-one and $(x_g^-, x_g^+) \in W$.*

Given a convex projective orbifold M , we use $\tilde{B}_{T^1M}^{(t)}$ to denote the open balls for the metric $\tilde{d}_{T^1\Omega}^{(t)}$ (see Section 2.13).

Lemma 7.14. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup; denote $M = \Omega/\Gamma$. Consider $\alpha > 0$ and $v_0 \in T^1M$ such that the endpoints $\phi_{-\infty}\tilde{v}_0$ and $\phi_{\infty}\tilde{v}_0$ of any lift \tilde{v}_0 are extremal (resp. strongly extremal). Then there exists $\varepsilon > 0$ satisfying the following. For any $v \in B_{T^1M}(v_0, \varepsilon)$, and for any time $t > \alpha$, if $\phi_t v \in B_{T^1M}(v_0, \varepsilon)$, then one can find a biproximal (resp. rank-one) periodic vector $w \in \tilde{B}_{T^1M}^{(t)}(v, \alpha)$ with period in $[t - \alpha, t + \alpha]$.*

Proof. Let W be a neighbourhood of $(\phi_{-\infty}\tilde{v}_0, \phi_{\infty}\tilde{v}_0)$ such that

$$[\xi_-, \xi_+] \cap B_{T^1\Omega}\left(\phi_t\tilde{v}_0, \frac{\alpha}{8}\right) \neq \emptyset$$

for any $0 \leq t \leq 1$. By Corollary 7.13, we can find a neighbourhood $U = U_- \times U_+ \subset W$ of $\phi_{\pm\infty}\tilde{v}_0$ such that for any $\gamma \in \Gamma$, if $(\gamma^{-1}\pi\tilde{v}_0, \gamma\pi\tilde{v}_0) \in U$ then γ is biproximal (resp. rank-one) and $(x_\gamma^-, x_\gamma^+) \in W$. Let $t_0 > 0$ and $\varepsilon_1 < \alpha/8$ be such that

$$\bar{B}_\Omega(\pi\phi_{\pm t}\tilde{v}, \varepsilon_1) \subset U_\pm$$

for any $\tilde{v} \in \bar{B}_{T^1\Omega}(\tilde{v}_0, \varepsilon_1)$ and $t \geq t_0$.

Now consider $t \geq t_0$ and $v \in B_{T^1M}(v_0, \varepsilon_1)$ such that $\phi_t v \in B_{T^1M}(v_0, \varepsilon_1)$. We can find a lift $\tilde{v} \in B_{T^1\Omega}(\tilde{v}_0, \varepsilon_1)$ and an element $\gamma \in \Gamma$ such that $\phi_t\tilde{v} \in B_{T^1\Omega}(\gamma\tilde{v}_0, \varepsilon_1)$. Then $(\gamma^{-1}\pi\tilde{v}_0, \gamma\pi\tilde{v}_0) \in U$, hence γ is biproximal (resp. rank-one) and $(x_\gamma^-, x_\gamma^+) \in W$. Be definition of W , we can find $\tilde{w} \in B_{T^1\Omega}(\tilde{v}_0, \alpha/8)$ tangent to the axis of γ . Then

$$d_{T^1\Omega}(\tilde{w}, \tilde{v}) \leq \frac{\alpha}{4} \quad \text{and} \quad d_{T^1\Omega}(\gamma\tilde{w}, \phi_t\tilde{v}) \leq \frac{\alpha}{4};$$

since $\gamma\tilde{w} = \phi_{\ell(\gamma)}\tilde{w}$, by the triangular inequality we have

$$|\ell(\gamma) - t| \leq \frac{\alpha}{2},$$

and $d_{T^1\Omega}^{(t)}(\tilde{w}, \tilde{v}) \leq \alpha$.

To finish the proof, it remains to find $\varepsilon < \varepsilon_1$ such that for all $\alpha \leq t \leq t_0$ and $v \in B_{T^1M}(v_0, \varepsilon)$, if $\phi_t v \in B_{T^1M}(v_0, \varepsilon)$, then one can find a biproximal (resp. rank-one) periodic vector $w \in \tilde{B}_{T^1M}^{(t)}(v, \alpha)$ with period in $[t - \alpha, t + \alpha]$. If v_0 is not periodic, then we take $\varepsilon < \varepsilon_1$ small enough so that $B_{T^1M}(v_0, \varepsilon) \subset B_{T^1M}^{(t_0)}(v_0, \varepsilon_2/2)$, where

$$\varepsilon_2 := \min_{\alpha \leq t \leq t_0} d_{T^1M}(\phi_t v_0, v_0).$$

If v_0 is periodic, then it is biproximal (resp. rank-one) since $\phi_{\pm\infty} \tilde{v}_0$ are extremal (resp. strongly extremal). We then take $\varepsilon < \varepsilon_1$ small enough so that $B_{T^1M}(v_0, \varepsilon) \subset B_{T^1M}^{(t_0)}(v_0, \varepsilon_2/2)$, where

$$\varepsilon_2 := \min_{t \in \mathcal{T}} d_{T^1M}(\phi_t v_0, v_0)$$

and \mathcal{T} is the set of times $\alpha \leq t \leq t_0$ such that $\phi_s v_0 \neq v_0$ for any $t - \alpha < s < t + \alpha$, concluding the proof. ■

8. The measure of maximal entropy

In this section we prove that on a non-elementary rank-one convex projective manifold $M = \Omega/\Gamma$, if Γ acts convex cocompactly on Ω , then the *Bowen–Margulis probability measure* (i.e. the unique Bowen–Margulis measure with total mass 1) is the unique measure with maximal entropy.

8.1. The measure of dynamical balls

In this section we derive from the Shadow lemma (Corollary 7.3) an estimate for the measure of dynamical balls (namely balls for the metrics $(d^{(t)})_{t \geq 0}$ defined in Definition 2.34). The idea is very similar to the computations in the proof of Proposition 6.2, which are actually Roblin’s computations; the difference here is that our shadows are a priori *small*, and this is why we need the stronger Shadow lemma (Corollary 7.3), which works for small shadows. However, Corollary 7.3 only works for usual shadows $\mathcal{O}_R(x, y)$, and not for the scarce shadows of the form $\mathcal{O}_R^-(x, y)$; to overcome this issue we use the following lemma, which is a consequence of Benzécri’s compactness theorem (Fact 2.4).

Lemma 8.1. *For any $\varepsilon > 0$, there exists $\varepsilon' > 0$ such that for any properly convex open set $\Omega \subset \mathbb{P}(V)$, for any $x, y \in \Omega$ at distance at least 1, if we have two points on the boundary $\xi \in \mathcal{O}_{\varepsilon'}(y, x)$ and $\eta \in \mathcal{O}_{\varepsilon'}(x, y)$, then the line $\xi \oplus \eta$ intersects both balls $B_\Omega(x, \varepsilon)$ and $B_\Omega(y, \varepsilon)$.*

Proof. By symmetry it is enough to prove that $\xi \oplus \eta$ meets $B_\Omega(x, \varepsilon)$. We assume by contradiction that there exist sequence $(\varepsilon'_n)_{n \in \mathbb{N}} \in \mathbb{R}_{>0}^{\mathbb{N}}$ converging to 0 and a sequence $(\Omega_n)_{n \in \mathbb{N}} \in \mathcal{E}_V^{\mathbb{N}}$ such that for each $n \in \mathbb{N}$, we can find $x_n, y_n \in \Omega_n$ at distance at least 1 and $\xi_n \in \mathcal{O}_{\varepsilon'_n}(y_n, x_n)$ and $\eta_n \in \mathcal{O}_{\varepsilon'_n}(x_n, y_n)$ such that

$$(\xi_n \oplus \eta_n) \cap B_\Omega(x_n, \varepsilon) = \emptyset.$$

By Benzécri's compactness theorem (Fact 2.4), we can assume, up to extraction, that $((\Omega_n, x_n))_n$ converges to some pointed properly convex open set $(\Omega, x) \in \mathcal{E}_V^\bullet$, bounded in some affine chart that we fix. Up to extraction we assume that Ω_n is also bounded in the affine chart for any $n \in \mathbb{N}$, and that $(y_n)_{n \in \mathbb{N}}$ (resp. $(\xi_n)_{n \in \mathbb{N}}$ and $(\eta_n)_{n \in \mathbb{N}}$) converges to $y \in \bar{\Omega} \setminus B_\Omega(x, 1)$ (resp. ξ and $\eta \in \partial\Omega$) such that

$$(\xi \oplus \eta) \cap B_\Omega(x, \varepsilon) = \emptyset.$$

By definition of the Hilbert metric one checks that

$$B_{\Omega_n}(x_n, \varepsilon'_n) \subset (1 - e^{-2\varepsilon'_n})(\Omega_n - x_n) + x_n,$$

hence $(B_{\Omega_n}(x_n, \varepsilon'_n))_{n \in \mathbb{N}}$ converges to the singleton $\{x\}$ for the Hausdorff topology, and similarly $(B_{\Omega_n}(y_n, \varepsilon'_n))_{n \in \mathbb{N}}$ converges to the singleton $\{y\}$. This implies that $x \in [\xi, \eta]$, which contradicts $(\xi \oplus \eta) \cap B_\Omega(x, \varepsilon) = \emptyset$. ■

Lemma 8.2. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup; set $M = \Omega/\Gamma$. Suppose that Γ is strongly irreducible and $T^1M_{\text{bip}} \neq \emptyset$, or that M is rank-one and non-elementary. Let m be a Sullivan measure on T^1M induced by a δ_Γ -conformal density. Then for any compact subset $K \subset T^1M$, for any $r > 0$, there exists a constant $C > 0$ such that given any time $t > 0$, for any $v \in \Gamma \cdot K$ such that $\phi_t v \in \Gamma \cdot K$,*

$$m(\tilde{B}_{T^1M}^{(t)}(v, r)) \leq C e^{-\delta_\Gamma t},$$

and if $v \in T^1M_{\text{bip}}$, then

$$C^{-1} e^{-\delta_\Gamma t} \leq m(\tilde{B}_{T^1M}^{(t)}(v, r)).$$

Proof. Let $(\mu_x)_{x \in \Omega}$ be a δ_Γ -conformal density on $\partial_h \Omega$ which induces a δ_Γ -conformal density $(\nu_x)_{x \in \Omega}$ on $\partial\Omega$ and the Sullivan measures \tilde{m} on $T^1\Omega$ and m on T^1M . Let $\tilde{v} \in T^1\Omega$ be a lift of v . We have

$$C_0^{-1} \tilde{m}(B_{T^1\Omega}^{(t)}(\tilde{v}, r)) \leq m(\tilde{B}_{T^1M}^{(t)}(v, r)) \leq \tilde{m}(B_{T^1\Omega}^{(t)}(\tilde{v}, r))$$

for any $t \geq 0$, where

$$C_0 = \max_{\tilde{w} \in \pi^{-1}K} \#\{\gamma : d_\Omega(\pi \tilde{w}, \gamma \pi \tilde{w}) \leq 4r\}.$$

Let us prove the upper bound in the lemma. Consider $\tilde{w} \in B_{T^1\Omega}^{(t)}(\tilde{v}, r)$. We make the following observations.

- The Lebesgue measure of the set of times $s \in \mathbb{R}$ such that $\phi_s \tilde{w} \in B_{T^1\Omega}^{(t)}(\tilde{v}, r)$ is less than $2r$.
- $\phi_\infty \tilde{w} \in \mathcal{O}_r^+(\pi \tilde{v}, \pi \phi_t \tilde{v})$.
- $\langle \xi, \eta \rangle_{\pi \tilde{v}} \leq r$ for all $\xi \in \pi_h^{-1}(\phi_{-\infty} \tilde{w})$ and $\eta \in \pi_h^{-1}(\phi_\infty \tilde{w})$.

Combined with the definition of \tilde{m} (see Section 3.3), they yield:

$$\tilde{m}(B_{T^1\Omega}^{(t)}(\tilde{v}, r)) \leq e^{2\delta_\Gamma r} \cdot \nu_{\pi \tilde{v}}(\partial\Omega) \cdot \nu_{\pi \tilde{v}}(\mathcal{O}_r^+(\pi \tilde{v}, \pi \phi_t \tilde{v})) \cdot 2r.$$

We deduce from this and the Shadow lemma (Lemma 7.2) the desired upper bound.

Let us prove the lower bound. We apply Lemma 8.1 to $\varepsilon := r/16$ to obtain $\varepsilon' > 0$. Then for all $\eta \in \mathcal{O}_{\varepsilon'}(\pi \tilde{v}, \pi \phi_{t+1} \tilde{v})$ and $\xi \in \mathcal{O}_{\varepsilon'}(\pi \phi_{t+1} \tilde{v}, \pi \tilde{v})$, we are able to find $\tilde{w} \in B_{T^1\Omega}^{(t)}(\tilde{v}, r/2)$ tangent to $\xi \oplus \eta$. Observe that the Lebesgue measure of the set of times $s \in \mathbb{R}$ such that $\phi_s \tilde{w} \in B_{T^1\Omega}^{(t)}(\tilde{v}, r)$ is greater than r . This means (remembering that the Gromov product is always non-negative) that

$$\tilde{m}(B_{T^1\Omega}^{(t)}(\tilde{v}, r)) \geq \nu_{\pi v}(\mathcal{O}_{\varepsilon'}(\pi \tilde{v}, \pi \phi_{t+1} \tilde{v})) \cdot \nu_{\pi v}(\mathcal{O}_{\varepsilon'}(\pi \phi_{t+1} \tilde{v}, \pi \tilde{v})) \cdot r.$$

We conclude the proof thanks to the Shadow lemma (Lemma 7.2), and the fact that $\Lambda^{\text{prox}} \subset \text{supp}(\nu_o)$. ■

In the convex cocompact case, this yields the following.

Corollary 8.3. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup; set $M = \Omega/\Gamma$. Suppose that Γ is strongly irreducible and $T^1M_{\text{bip}} \neq \emptyset$, or that M is rank-one and non-elementary. Let m be a Sullivan measure on T^1M induced by a δ_Γ -conformal density. Then for any $r > 0$, there exists a constant $C > 0$ such that given any time $t > 0$, for any $v \in T^1M_{\text{cor}}$,*

$$m(\tilde{B}_{T^1M}^{(t)}(v, r)) \leq C e^{-\delta_\Gamma t},$$

and if $v \in T^1M_{\text{bip}}$, then

$$C^{-1} e^{-\delta_\Gamma t} \leq m(\tilde{B}_{T^1M}^{(t)}(v, r)).$$

8.2. The Bowen–Margulis measure has maximal entropy

In this section we prove that any Sullivan measure induced by a δ_Γ -conformal density has maximal entropy, which is equal to δ_Γ . For this we do not need Theorem 1.6.

Proposition 8.4. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Let m be the Bowen–Margulis probability measure on T^1M . Then m has maximal entropy on T^1M_{cor} ; in other words,*

$$h_m(T^1M_{\text{bip}}, (\phi_t)_t) = h_{\text{top}}(T^1M_{\text{bip}}, (\phi_t)_t) = h_{\text{top}}(T^1M_{\text{cor}}, (\phi_t)_t) = \delta_\Gamma.$$

Proof. We can assume without loss of generality that Γ is torsion-free by Observation 2.45 and since for any finite-index subgroup Γ' of Γ , the measure m is the push-forward of the Bowen–Margulis probability measure on $T^1\Omega/\Gamma'$. Let ε_0 be the injectivity radius of M . Let \mathcal{P} be a finite measurable partition of T^1M_{cor} whose elements have diameter less than $\varepsilon_0/3$. According to the definition of the measure-theoretic entropy, to the variational principle (Fact 2.39), and to Fact 2.36, it is enough to prove that

$$H_m(\phi_1, \mathcal{P}) \geq \delta_\Gamma.$$

For $n \geq 1$, we observe that by definition (and by Lemma 2.1), any element of $\mathcal{P}^{(n)}$ has diameter less than $\varepsilon_0/3$ with respect to the metric $d_{T^1M}^{(n)}$. Therefore, by Corollary 8.3, there exists a constant $C > 0$ such that for any $n \geq 1$, any element of $\mathcal{P}^{(n)}$ has an m -measure less than $Ce^{-\delta_\Gamma n}$. We now conclude the proof by the following computation:

$$\begin{aligned} H_m(\mathcal{P}^{(n)}) &= - \sum_{P \in \mathcal{P}^{(n)}} m(P) \log(m(P)) \\ &\geq - \sum_{P \in \mathcal{P}^{(n)}} m(P) \log(Ce^{-\delta_\Gamma n}) \\ &\geq \delta_\Gamma n - \log(C). \end{aligned}$$

The inequality $H_m(\phi_1, \mathcal{P}) \geq \delta_\Gamma$ then follows immediately from Definition 2.37:

$$H_m(\phi_1, \mathcal{P}) := \lim_{n \rightarrow \infty} \frac{H_m(\mathcal{P}^{(n)})}{n}. \quad \blacksquare$$

8.3. Separated sets in dynamical balls

In this section we prove a technical lemma which bound from above the size a separated set in a dynamical ball. To perform this estimate we will use the notion of proper densities (Definition 2.5).

Corollary 8.5. *Consider $0 < r < R$. Let $\Omega \in \mathcal{E}_V$, let $\Gamma \subset \text{Aut}(\Omega)$ be a discrete subgroup, let $M = \Omega/\Gamma$ and let $t \geq 0$.*

- (1) For any vector $v \in T^1M$, the cardinality of any $(\tilde{d}_{T^1M}^{(t)}, r)$ -separated set of $\tilde{B}_{T^1M}^{(t)}(v, R)$ is less than

$$\chi_+ \left(R + \frac{r}{4} \right)^2 \cdot \chi_- \left(\frac{r}{4} \right)^{-2}$$

(see (2.2)).

- (2) Consider a $(\tilde{d}_{T^1M}^{(t)}, r)$ -separated subset $\{v_1, \dots, v_k\}$ of T^1M (and of size k), take a vector $w_i \in \tilde{B}_{T^1M}^{(t)}(v_i, R)$ for each $i = 1, \dots, k$. Then one can find a subset I of $\{1, \dots, k\}$ of size greater than

$$k \cdot \chi_+ \left(2R + \frac{r}{2} \right)^{-2} \cdot \chi_- \left(\frac{r}{4} \right)^2$$

such that $\{w_i : i \in I\}$ is $(\tilde{d}_{T^1M}^{(t)}, r)$ -separated.

Proof. Let us first prove (1) when Γ is trivial. Let $A \subset B_{T^1\Omega}^{(t)}(v, R)$ be a $(d_{T^1\Omega}^{(t)}, r)$ -separated set. We set

$$B := \{(\pi w, \pi \phi_t w) : w \in A\} \subset B_\Omega(\pi v, R) \times B_\Omega(\pi \phi_t v, R).$$

By Lemma 2.1, and since A is $(d_{T^1\Omega}^{(t)}, r)$ -separated, we see that B is $(d, r/2)$ -separated for the metric d on Ω^2 , defined by

$$d((x, y), (x', y')) = \max(d_\Omega(x, x'), d_\Omega(y, y'))$$

for $(x, y), (x', y') \in \Omega^2$; this exactly means that for all $(x, y) \neq (x', y') \in B$,

$$\left(B_\Omega\left(x, \frac{r}{4}\right) \times B_\Omega\left(y, \frac{r}{4}\right) \right) \cap \left(B_\Omega\left(x', \frac{r}{4}\right) \times B_\Omega\left(y', \frac{r}{4}\right) \right) = \emptyset.$$

As a consequence,

$$\begin{aligned} \#A = \#B &\leq \chi_- \left(\frac{r}{4} \right)^{-2} \sum_{(x,y) \in B} \text{Vol}_\Omega \left(B_\Omega \left(x, \frac{r}{4} \right) \right) \text{Vol}_\Omega \left(B_\Omega \left(y, \frac{r}{4} \right) \right) \\ &\leq \chi_- \left(\frac{r}{4} \right)^{-2} \text{Vol}_\Omega^2 \left(\bigsqcup_{(x,y) \in B} B_\Omega \left(x, \frac{r}{4} \right) \times B_\Omega \left(y, \frac{r}{4} \right) \right) \\ &\leq \chi_- \left(\frac{r}{4} \right)^{-2} \text{Vol}_\Omega^2 \left(B_\Omega \left(\pi v, R + \frac{r}{4} \right) \times B_\Omega \left(\pi \phi_t v, R + \frac{r}{4} \right) \right) \\ &\leq \chi_+ \left(R + \frac{r}{4} \right)^2 \chi_- \left(\frac{r}{4} \right)^{-2}. \end{aligned}$$

Next, let us prove (1) when Γ is not necessarily trivial. Let $A \subset \tilde{B}_{T^1M}^{(t)}(v, R)$ be a $(\tilde{d}_{T^1M}^{(t)}, r)$ -separated set. Consider a lift $\tilde{v} \in T^1\Omega$ of v , and a lift $\tilde{A} \subset \tilde{B}_{T^1\Omega}^{(t)}(\tilde{v}, R)$

of A . Then \tilde{A} is $(d_{T^1\Omega}^{(t)}, r)$ -separated, and therefore it has cardinality less than

$$\chi_+ \left(R + \frac{r}{4} \right)^2 \chi_- \left(\frac{r}{4} \right)^{-2},$$

and so does A .

Let us establish (2). We construct I by induction. We set

$$i_0 = 0 \quad \text{and} \quad I_0 := \{1, \dots, k\}.$$

For $j \geq 1$, if (I_0, \dots, I_{j-1}) and (i_0, \dots, i_{j-1}) have been constructed, we set

$$i_j := \min I_{j-1} > i_{j-1} \quad \text{and} \quad I_j := \{i \in I_{j-1} : \tilde{d}_{T^1M}^{(t)}(w_{i_j}, w_i) \geq r\} \subsetneq I_{j-1}.$$

This process eventually stops, at the n -th step for some $n \in \{1, \dots, k\}$ such that $I_n = \emptyset$. The set $\{w_{i_j} : 1 \leq j \leq n\}$ is $(\tilde{d}_{T^1M}^{(t)}, r)$ -separated by construction. In order to prove that k is bounded above by $n \cdot \chi_+(2R + r/2)^2 \cdot \chi_-(r/4)^{-2}$, it is enough to see that for each $0 \leq j \leq n-1$,

$$\#I_{j+1} \geq \#I_j - \chi_+ \left(2R + \frac{r}{2} \right)^2 \cdot \chi_- \left(\frac{r}{4} \right)^{-2}.$$

This is a consequence of (1) and of the fact that $I_j \setminus I_{j+1}$ is contained in the set of indices i such that $v_i \in \tilde{B}_{T^1\Omega}^{(t)}(v_{j+1}, r + 2R)$. \blacksquare

8.4. The measure of maximal entropy is unique

In order to prove the uniqueness of the measure of maximal entropy, we will use our estimates on the size of the dynamical balls Corollary 8.3. However, one of these estimates only holds for balls centred at vectors in T^1M_{bip} , whereas we would like the uniqueness of the measure of maximal entropy on T^1M_{cor} . This is why we will need Corollary 8.5 and the following lemma.

For the rest of this section, given a convex projective manifold $M = \Omega/\Gamma$, we denote by $T^1\Omega_{\text{cor}}$ (resp. $T^1\Omega_{\text{bip}}$) the set (depending on Γ) of vectors $v \in T^1\Omega$ such that $\phi_{\pm\infty}v \in \Lambda_{\Omega}^{\text{orb}}(\Gamma)$ (resp. $\Lambda^{\text{prox}}(\Gamma)$).

Lemma 8.6. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup. Suppose $M = \Omega/\Gamma$ is rank-one and non-elementary. Then there exists $R > 0$ such that for every point $\xi \in \Lambda^{\text{orb}}$ we can find $\eta \in \Lambda^{\text{prox}}$ such that $d_{\bar{\Omega}}(\xi, \eta) \leq R$. In particular, according to Lemma 2.1, for any $v \in T^1\Omega_{\text{cor}}$, we can find a vector $w \in T^1\Omega_{\text{bip}}$ such that $d_{T^1\Omega}(\phi_t v, \phi_t w) \leq 2R$ for any $t \in \mathbb{R}$.*

Proof. Consider o in the convex hull of Λ^{orb} in Ω and the δ_{Γ} -conformal density ν_o on $\partial\Omega$. Let $R > 0$ be given by the Shadow lemma (Corollary 7.3). According to

Fact 2.3, it is enough to show that for any $x \in [o, \xi)$, the shadow $\mathcal{O}_R(o, x)$ intersect Λ^{prox} ; this is implied by the fact that $\nu_o(\Lambda^{\text{prox}}) = 1$ (by Theorem 1.6 and Proposition 1.7) and $\nu_o(\mathcal{O}_R(o, x)) > 0$. ■

In the case where M is compact, the proof of Theorem 1.3 below can be shorten by using the following result; for instance, we do not need Section 8.3 and Lemma 8.6 in this case.

Fact 8.7 ([16, Thm. 1.3]). Let $\Omega \subset P(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a non-elementary rank-one discrete subgroup that acts cocompactly on Ω . Then $\Lambda^{\text{prox}} = \partial\Omega$.

Proof of Theorem 1.3. It is enough to prove that the Bowen–Margulis probability measure m on T^1M is the unique measure of maximal entropy, since m is mixing by Theorem 1.6.

We can assume that Γ is torsion-free by Observation 2.45 and since for any finite-index subgroup Γ' of Γ , the measure m is the push-forward of the Bowen–Margulis probability measure on $T^1\Omega/\Gamma'$; let ε_0 be the injectivity radius of Ω/Γ . Consider a $(\phi_t)_{t \in \mathbb{R}}$ -invariant probability measure m' on T^1M_{cor} which is different from m . Since m is ergodic (Theorem 1.6 and Proposition 1.7), m' cannot be absolutely continuous with respect to m . By Radon–Nikodym Theorem we can decompose m' into a sum $tm'' + (1-t)m$, where $0 < t \leq 1$ and m'' is $(\phi_t)_{t \in \mathbb{R}}$ -invariant and singular with respect to m . Then

$$h_{m'}(\phi) = th_{m''}(\phi) + (1-t)\delta_\Gamma$$

(see [43, Cor. 4.3.17]), and we only need to prove that $h_{m''}(\phi) < \delta_\Gamma$. Note that since $\Lambda^{\text{orb}} \setminus \Lambda^{\text{prox}}$ does not intersect $\partial_{\text{sse}}\Omega$. Without loss of generality, we assume that $m'' = m'$ is singular with respect to m . Let $A \subset T^1M_{\text{sse}}$ be a flow-invariant measurable subset such that $m(A) = 1$, while $m'(A) = 0$.

Fix $\varepsilon > 0$ and let $K_1 \subset A \cap T^1M_{\text{cor}}$ and $K_2 \subset T^1M_{\text{cor}} \setminus A$ be compact subsets such that $m(K_1) \geq 1 - \varepsilon$ and $m'(K_2) \geq 1 - \varepsilon$. Observe that

$$\min\{\tilde{d}_{T^1M}^{(2t)}(\phi_{-t}v, \phi_{-t}w) : v \in K_1, w \in K_2\} \xrightarrow{t \rightarrow \infty} \infty. \tag{8.1}$$

Indeed, otherwise there would exist $v \in K_1$ and $w \in K_2$ such that

$$\sup_{t > 0} \tilde{d}_{T^1M}^{(2t)}(\phi_{-t}v, \phi_{-t}w) < \infty.$$

Then we could find lifts $\tilde{v}, \tilde{w} \in T^1\Omega$ such that

$$\sup_{t > 0} d_{T^1\Omega}^{(2t)}(\phi_{-t}\tilde{v}, \phi_{-t}\tilde{w}) < \infty,$$

which implies, since $\phi_\infty \tilde{v}$ and $\phi_{-\infty} \tilde{v}$ are extremal, that $\phi_{\pm\infty} \tilde{v} = \phi_{\pm\infty} \tilde{w}$, and hence $w \in \phi_{\mathbb{R}} v \subset A$, which is a contradiction.

Let $n \geq 1$ and consider a maximal $(d_{T^1 M}^{(2n)}, \varepsilon_0/8)$ -separated set

$$\{v_1, \dots, v_k\} \subset T^1 M_{\text{cor}},$$

which is ordered so that for any $i = 1, \dots, k$, the ball $B_{T^1 M_{\text{cor}}}^{(2n)}(v_i, \varepsilon_0/8)$ intersects $\phi_{-n} K_2$ if and only if $i \leq l$, where $1 \leq l \leq k$ is some integer.

Construct by induction the finite measurable partition $\mathcal{P} = \{P_1, \dots, P_k\}$ of $T^1 M_{\text{cor}}$ so that

$$\begin{aligned} B_{T^1 M_{\text{cor}}}^{(2n)}\left(v_{i+1}, \frac{\varepsilon_0}{16}\right) &\subset P_{i+1} := B_{T^1 M_{\text{cor}}}^{(2n)}\left(v_{i+1}, \frac{\varepsilon_0}{8}\right) \setminus (P_1 \cup \dots \cup P_i) \\ &\subset B_{T^1 M_{\text{cor}}}^{(2n)}\left(v_{i+1}, \frac{\varepsilon_0}{8}\right). \end{aligned}$$

Note that P_i has diameter less than $\varepsilon_0/3$ with respect to $d_{T^1 M}^{(2n)}$ for each $1 \leq i \leq k$, therefore we can combine Remarks 2.38 and 2.41 with Facts 2.44 and 2.42 to obtain

$$h_{m'}(\phi_1) = \frac{1}{2n} h_{m'}(\phi_{2n}) = \frac{1}{2n} H_{m'}(\phi_{2n}, \mathcal{P}) \leq \frac{1}{2n} H_{m'}(\mathcal{P}).$$

Now we use the classical fact that for all $q \in \mathbb{N}_{>0}$ and $a_1, \dots, a_q > 0$, if $s := a_1 + \dots + a_q \leq 1$ then

$$-\sum_i a_i \log(a_i) \leq -s \log s + s \log q \leq s \log(q) + \frac{1}{e},$$

and we compute

$$\begin{aligned} H_{m'}(\mathcal{P}) &= -\sum_{i=1}^k m'(P_i) \log(m'(P_i)) \\ &\leq m'\left(\bigcup_{i=1}^l P_i\right) \log(l) + m'\left(\bigcup_{i=l+1}^k P_i\right) \log(k) + \frac{2}{e}. \end{aligned}$$

Note that on one hand, $\phi_{-n} K_2$ is contained in $\bigcup_{i=1}^l P_i$, hence

$$m'\left(\bigcup_{i=1}^l P_i\right) \geq 1 - \varepsilon.$$

On the other hand $\phi_{-n} K_1$ does not intersect $\bigcup_{i=1}^l P_i$ for n large enough. Indeed if there exist $i \in \{1, \dots, l\}$ and $v \in \phi_{-n} K_1 \cap P_i$, then we take a vector

$$w \in \phi_{-n} K_2 \cap B_{T^1 M}^{(2n)}\left(v_i, \frac{\varepsilon_0}{8}\right)$$

and use the triangular inequality to obtain that $\tilde{d}_{T^1M}^{(2n)}(\phi_{-n}\phi_n v, \phi_{-n}\phi_n w) \leq \varepsilon_0/4$, which is not compatible with (8.1) for n large. As a consequence,

$$m\left(\bigcup_{i=1}^l P_i\right) \leq \varepsilon.$$

We now need to bound from above k and l . If M is compact then it is easier to conclude the proof: we can use Fact 8.7, which implies that

$$T^1M = T^1M_{\text{cor}} = T^1M_{\text{bip}},$$

and apply directly Corollary 8.3 to get

$$l \leq \frac{m(\bigcup_{i=1}^l P_i)}{\min\{m(P_i) : 1 \leq i \leq l\}} \leq \varepsilon C e^{2n\delta_\Gamma},$$

and similarly $k \leq C e^{2n\delta_\Gamma}$, where C only depends on ε_0 . Thus, we obtain

$$2nh_{m'}(\phi_1) \leq (1 - \varepsilon) \log(\varepsilon) + \log(C) + 2n\delta_\Gamma + \frac{2}{e},$$

which is strictly less than $2n\delta_\Gamma$ for ε small enough.

In the general case, we need to take into account that T^1M_{cor} and T^1M_{bip} might be different and that Corollary 8.3 only holds on T^1M_{bip} ; this is where we need Corollary 8.5 and Lemma 8.6. Thanks to the latter, there exists $R > 0$, which only depends on Γ and Ω , and such that for each $1 \leq i \leq k$, we can find

$$w_i \in \tilde{B}_{T^1M}^{(2n)}(v_i, R) \cap T^1M_{\text{bip}}.$$

Then we can use Corollary 8.5 (2) to find $I \subset \{1, \dots, k\}$ such that $\{w_i : i \in I\}$ is $(d_{T^1M}^{(2n)}, \varepsilon_0/8)$ -separated and

$$\begin{aligned} k &\leq \#I \cdot \chi_+\left(2R + \frac{\varepsilon_0}{16}\right)^2 \chi_-\left(\frac{\varepsilon_0}{32}\right)^{-2} \\ &\leq \frac{m(\bigcup_{i \in I} B_{T^1M}^{(2n)}(w_i, \varepsilon_0/16))}{\min\{m(B_{T^1M}^{(2n)}(w_i, \varepsilon_0/16)) : i \in I\}} \chi_+\left(2R + \frac{\varepsilon_0}{16}\right)^2 \chi_-\left(\frac{\varepsilon_0}{32}\right)^{-2} \\ &\leq C' e^{2n\delta_\Gamma}, \end{aligned}$$

where C' only depends on Γ and Ω . Similarly, we can find

$$w'_i \in \phi_{-n}K_2 \cap B_{T^1M}^{(2n)}\left(v_i, \frac{\varepsilon_0}{8}\right) \quad \text{and} \quad w''_i \in \tilde{B}_{T^1M}^{(2n)}(w'_i, R) \cap T^1M_{\text{bip}}$$

for each $i = 1, \dots, l$. Corollary 8.5 (2) gives us $I' \subset \{1, \dots, l\}$ such that $\{w_i'' : i \in I'\}$ is $(d_{T^1M}^{(2n)}, \varepsilon_0/8)$ -separated and

$$\begin{aligned} l &\leq \#I' \cdot \chi_+ \left(2R + \frac{\varepsilon_0}{8}\right)^2 \chi_- \left(\frac{\varepsilon_0}{32}\right)^{-2} \\ &\leq \frac{m(\bigcup_{i \in I'} B_{T^1M}^{(2n)}(w_i'', \varepsilon_0/16))}{\min\{m(B_{T^1M}^{(2n)}(w_i'', \varepsilon_0/16)) : i \in I'\}} \chi_+ \left(2R + \frac{\varepsilon_0}{8}\right)^2 \chi_- \left(\frac{\varepsilon_0}{32}\right)^{-2} \\ &\leq \varepsilon C'' e^{2n\delta_\Gamma}, \end{aligned}$$

where C'' only depends on Γ and Ω , and we have used the fact that $B_{T^1M}^{(2n)}(w_i'', \varepsilon_0/16)$ does not intersect $\phi_{-n}K_1$ for any $i = 1, \dots, l$ and for n large enough (again because of (8.1)). As we explained in the compact case, this implies that

$$2nh_{m'}(\phi_1) \leq (1 - \varepsilon) \log(\varepsilon) + \log \max(C', C'') + 2n\delta_\Gamma + \frac{2}{e},$$

which is strictly less than $2n\delta_\Gamma$ for ε small enough. ■

9. Counting closed geodesics

In this section we keep on adapting Knieper’s article [45] in order to prove Proposition 1.5, which gives asymptotic estimates for the number of closed geodesics of length less than t , when t goes to infinity.

These estimates do not all need the results of the previous sections. More precisely, to prove the upper bound on the number of rank-one close geodesics in (1) we do not need Theorem 1.6. To prove the lower bound in (1) we need the mixing property of the Bowen–Margulis measure, but not the uniqueness of the measure of maximal entropy. To establish the upper bound on the number of non-rank-one closed geodesics in (2) and the equidistribution of closed geodesics we need uniqueness of the measure of maximal entropy.

Recall that $[\Gamma]$ is the set of conjugacy classes of Γ , and $[\Gamma]^{r1} \subset [\Gamma]$ (resp. $[\Gamma]^{\text{sing}}$) consists of conjugacy classes of rank-one (resp. non-rank-one) elements of Γ . For each subset $A \subset [\Gamma]$ and interval $I \subset \mathbb{R}$, we set $A_I = \{c \in A : \ell(c) \in I\}$, and we write $A_T = A_{[0, T]}$ for any $T \geq 0$.

9.1. The lower bound

In this section we use the mixing of the Bowen–Margulis measure to obtain a lower bound on the number of closed geodesics.

Proposition 9.1. *Let $\Omega \subset P(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Then there exists a constant $C > 0$ such that for any $T > C$,*

$$\#[\Gamma]_T^{\text{r1}} \geq \frac{C}{T} e^{\delta_\Gamma T}.$$

Proof. Without loss of generality, we may assume that Γ is torsion-free, since for any finite-index subgroup $\Gamma' \subset \Gamma$, for any element $\gamma \in \Gamma'$, there at most $[\Gamma : \Gamma']$ conjugacy classes of Γ' inside the conjugacy class of γ in Γ . Let $\varepsilon_0 > 0$ be the injectivity radius of M and m the Bowen–Margulis probability measure. Fix a compact subset $K \subset T^1 M_{\text{sse}}$ whose measure $m(K)$ is positive. Using Lemma 7.14, we can find

$$0 < \varepsilon < \frac{\varepsilon_0}{3}$$

such that for any vector $v \in K$, for any time $t \geq 1$, if $d_{T^1 M}(v, \phi_t v) \leq \varepsilon$ then there exists a rank-one periodic vector $w \in B_{T^1 M}^{(t)}(v, \varepsilon_0/6)$ with period in $[t - 1, t + 1]$. Let us denote by $R_t \subset K$ the subset of vectors $v \in K$ such that $d_{T^1 M}(v, \phi_t v) \leq \varepsilon$; we are going to bound from below its measure. To that end take \mathcal{P} a finite measurable partition of $T^1 M_{\text{bip}}$ with diameter less than ε , and compute the limit, using the mixing property, established in Theorem 1.6:

$$m(R_t) \geq \sum_{P \in \mathcal{P}} m(P \cap K \cap \phi_t(P \cap K)) \xrightarrow{t \rightarrow \infty} \sum_{P \in \mathcal{P}} m(P \cap K)^2 \geq \frac{m(K)^2}{\#\mathcal{P}} > 0.$$

On the other hand we can bound from above this measure thanks to the closing lemma and Corollary 8.3. For any conjugacy class $c \in [\Gamma]^{\text{r1}}$, fix a vector $v_c \in T^1 M$ tangent to the projection in $T^1 M$ of the axis of any element of c :

$$\begin{aligned} m(R_t) &\leq \sum_{c \in [\Gamma]_{t+1}^{\text{r1}}} \sum_{k=0}^{\lfloor 6(t+1)/\varepsilon_0 \rfloor} m\left(B_{T^1 M}^{(t)}\left(\phi_{k(\varepsilon_0/6)} v_c, \frac{\varepsilon_0}{3}\right)\right) \\ &\leq C e^{-\delta_\Gamma t} \left(\frac{6(t+1)}{\varepsilon_0} + 1\right) \#[\Gamma]_{t+1}. \end{aligned}$$

This ends the proof. ■

9.2. The upper bound on the number of rank-one closed geodesics

To bound from above the number of rank-one closed geodesics, we do not need Theorem 1.6.

Proposition 9.2. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Then there exists a constant $C > 0$ such that for any $T > 0$,*

$$\#[\Gamma]_T^{r_1} \leq \frac{C}{T} e^{\delta_\Gamma T}.$$

Proof. Let $\Gamma' \subset \Gamma$ be a finite-index torsion-free subgroup; we set $M' = \Omega/\Gamma'$ and take $\varepsilon_0 < 1$ smaller than the injectivity radius of M' . For each rank-one conjugacy class c of Γ , choose $\gamma_c \in c$ and $v_c \in \text{Axis}(\gamma_c)$. Consider an integer $n \geq 1$. One easily checks using the triangular inequality that any vector of $T^1\Omega$ belongs to at most

$$C_1 := \max_{x \in \Omega} \#\left\{ \gamma \in \Gamma : d_\Omega(x, \gamma x) \leq 1 + \frac{2\varepsilon_0}{3} \right\}$$

balls of the family

$$\left\{ B_{T^1\Omega}^{(n+1)}\left(\phi_k v_c, \frac{\varepsilon_0}{6}\right) : c \in [\Gamma]_{[n, n+1]}^{r_1}, 0 \leq k < n \right\}.$$

By Corollary 8.3, we can bound from below the m' -measure of the projection in T^1M' of these balls by $C^{-1}e^{-\delta_\Gamma n}$ for some constant $C > 0$, where m' is the Bowen–Margulis probability measure on T^1M' . Since m' is a probability measure, we obtain

$$n\#[\Gamma]_{[n, n+1]}^{r_1} \leq C_1 C e^{\delta_\Gamma n \varepsilon_0/3}. \quad \blacksquare$$

9.3. The upper bound on the number of non-rank-one closed geodesics

Let us bound from above the number of non-rank-one conjugacy classes of Γ . The idea is that to each non-rank-one conjugacy class we can associate a closed geodesic which is contained in a flow-invariant closed subset of T^1M_{cor} to which the Bowen–Margulis measure gives zero measure; this implies that the topological entropy of the geodesic flow on this subset is smaller than δ_Γ .

Proposition 9.3. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Let $K \subset T^1M_{\text{cor}}$ be the $(\phi_t)_t$ -invariant closed subset that consists of the vectors whose lifts $v \in T^1\Omega$ are such that $d_{\text{spl}}(\phi_{-\infty}v, \phi_\infty v) \leq 2$. Then the topological entropy of $(\phi_t)_t$ on K is strictly smaller than δ_Γ .*

Proof. According to Remark 2.43 and Observation 2.45, there exists a probability measure m' on K whose entropy is the topological entropy on K . Observe that K is disjoint from the set T^1M_{sse} of vectors whose lifts $v \in T^1\Omega$ satisfy $\phi_{-\infty}v, \phi_\infty v \in \partial_{\text{sse}}\Omega$, while the Bowen–Margulis probability measure m_Γ is concentrated on T^1M_{sse} by Theorem 1.6. Thus, m' and m_Γ are different. By Theorem 1.3, the entropy of m' must be strictly smaller than δ_Γ . \blacksquare

Corollary 9.4. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Then the exponential growth rate of the number of non-rank-one conjugacy classes of translation length shorter than t , when t grows, is strictly less than δ_Γ .*

Proof. Let $\Gamma' \subset \Gamma$ be a finite-index torsion-free subgroup; we set $M' = \Omega/\Gamma'$ and ε_0 to be the injectivity radius of M' ; we denote by ε_0 the injectivity radius of M' . It is easy to show that there are only finitely many conjugacy class c of Γ such that $\ell(c) \leq \varepsilon_0$. Using Corollary 7.9, for each conjugacy class c with $\ell(c) > 0$ we can find an element $\gamma_c \in c$ and a vector $v_c \in T^1\Omega$ such that $\phi_{\pm\infty} v_c \in \Lambda^{\text{orb}}$ and $\gamma_c \pi v_c = \pi \phi_{\ell(c)} v_c$. Consider $t > 0$. Using the triangular inequality, one can check that any vector of $T^1\Omega$ belongs to at most

$$C_1 := \max_{x \in \mathcal{C}_\Omega^{\text{con}}(\Gamma)} \#\left\{ \gamma \in \Gamma : d_\Omega(x, \gamma x) \leq 1 + \frac{2\varepsilon_0}{3} \right\}$$

balls of the family

$$\left\{ B_{T^1\Omega}^{(t+1)}\left(v_c, \frac{\varepsilon_0}{6}\right) : c \in [\Gamma]_{[t, t+1]}^{\text{sing}} \right\}.$$

Therefore, we can extract from $\{v_c : c \in [\Gamma]_{[t, t+1]}^{\text{sing}}\}$ a $(d_{T^1\Omega}^{(t+1)}, \varepsilon_0/6)$ -separated family of size at least $C_1^{-1} \#[\Gamma]_{[t, t+1]}^{\text{sing}}$. The projection in T^1M' of this family belongs to the set K in Proposition 9.3 by Lemma 7.7 (6). By definition of the topological entropy, the exponential growth rate of the size of such a family, when t goes to infinity, is bounded above by the topological entropy on K , which is strictly less than δ_Γ by Proposition 9.3 above. ■

9.4. Sums of uniform measures on closed geodesics

Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set and $\Gamma \subset \text{Aut}(\Omega)$ a discrete subgroup; denote $M = \Omega/\Gamma$. We introduce a few notations. For any conjugacy class $c \in [\Gamma]^{\text{r1}}$, we denote by $\mathcal{L}c$ the unique $(\phi_t)_{t \in \mathbb{R}}$ -invariant probability measure supported on the projection in T^1M of the axis of any element of c . For each finite subset $A \subset [\Gamma]^{\text{r1}}$, we consider

$$\mathcal{L}A = \frac{1}{\#A} \sum_{c \in A} \mathcal{L}c.$$

Proposition 9.5. *Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Let $A \subset [\Gamma]^{\text{r1}}$ be such that $\log(\#A_T)/T$ converges to δ_Γ when T tends to infinity. Then $\mathcal{L}A_T$ converges to the Bowen–Margulis probability measure when T goes to infinity.*

Proof. Let $\Gamma' \subset \Gamma$ be a torsion-free finite-index subgroup; let ε_0 be the injectivity radius of $M' = \Omega/\Gamma'$. For each conjugacy class $c \in [\Gamma]^{11}$, we choose a representative $\gamma_c \in \Gamma$, and we call $\mathcal{L}'c$ the unique $(\phi_t)_t$ -invariant probability measure on the projection in T^1M' of the axis of γ_c . For any finite subset $B \subset [\Gamma]^{11}$, we set

$$\mathcal{L}'B = (\#B)^{-1} \sum_{c \in B} \mathcal{L}'c.$$

By Theorem 1.3, it is enough to show that any accumulation point $m' \lim_{k \rightarrow \infty} \mathcal{L}'A_T$ on T^1M' has entropy bounded below by δ_Γ .

Let us give ourselves a finite measurable partition \mathcal{P} of T^1M' of diameter less than $\varepsilon_0/3$ and such that for any element $P \in \mathcal{P}$, we have $m'(\partial P) = 0$. Then

$$h_{m'}(\phi) \geq H_{m'}(\phi, \mathcal{P}) = \lim_{n \rightarrow \infty} \frac{1}{n} H_{m'}(\mathcal{P}^{(n)}).$$

Fix $n \geq 1$ and note that for each $P \in \mathcal{P}^{(n)}$, we have $m'(\partial P) = 0$. As a consequence

$$\begin{aligned} H_{m'}(\mathcal{P}^{(n)}) &= \lim_{k \rightarrow \infty} H_{\mathcal{L}'A_{T_k}}(\mathcal{P}^{(n)}) \\ &\geq \liminf_{T \rightarrow \infty} H_{\mathcal{L}'A_T}(\mathcal{P}^{(n)}). \end{aligned}$$

Consider $\alpha > 0$ and let us show that

$$\liminf_{T \rightarrow \infty} H_{\mathcal{L}'A_T}(\mathcal{P}^{(n)}) \geq n(\delta_\Gamma - \alpha).$$

Let $T_0 > 0$ be large enough so that $\#A_T \geq e^{(\delta_\Gamma - \alpha)T}$ for any $T \geq T_0$. Take $T > T_0$ and decompose $[0, T]$ into disjoint intervals:

$$[0, T] = [0, T_0] \sqcup \bigsqcup_{I \in \mathcal{I}_T} I,$$

such that each $I \in \mathcal{I}_T$ has diameter less than 1, and $\#\mathcal{I}_T = \lceil T - T_0 \rceil$. Then by [43, Prop. 4.3.3.6], we have

$$\begin{aligned} \liminf_{T \rightarrow \infty} H_{\mathcal{L}'A_T}(\mathcal{P}^{(n)}) &\geq \liminf_{T \rightarrow \infty} \frac{\#A_{T_0}}{\#A_T} H_{\mathcal{L}'A_{T_0}}(\mathcal{P}^{(n)}) + \sum_{I \in \mathcal{I}_T} \frac{\#A_I}{\#A_T} H_{\mathcal{L}'A_I}(\mathcal{P}^{(n)}) \\ &\geq \liminf_{T \rightarrow \infty} \sum_{I \in \mathcal{I}_T} \frac{\#A_I}{\#A_{]T_0, T]}} H_{\mathcal{L}'A_I}(\mathcal{P}^{(n)}) \\ &\geq \liminf_{T \rightarrow \infty} \frac{n}{T} \sum_{I \in \mathcal{I}_T} \frac{\#A_I}{\#A_{]T_0, T]}} H_{\mathcal{L}'A_I}(\mathcal{P}^{(\lceil T \rceil)}), \end{aligned}$$

where we have used the Euclidean division $\lceil T \rceil = q_T n + r_T$ with the following classical inequality (see e.g. [43, Prop. 4.3.3.1–4]):

$$\begin{aligned} H_{\lambda_I}(\mathcal{P}^{(n)}) &\geq \frac{1}{q_T} H_{\lambda_I}(\mathcal{P}^{(\lceil T \rceil)}) - \frac{1}{q_T} H_{\lambda_I}(\mathcal{P}^{(r_T)}) \\ &\geq \frac{n}{T+1} H_{\lambda_I}(\mathcal{P}^{(\lceil T \rceil)}) - \frac{n}{T-n} \log(\#\mathcal{P}^n). \end{aligned}$$

Using the triangular inequality, one checks that for any $P \in \mathcal{P}^{(\lceil T \rceil)}$ and any $I \in \mathcal{I}_T$, there are at most $C_1 = \max_{x \in \mathcal{C}_{\Omega}^{\text{cor}}(\Gamma)} \#\{\gamma \in \Gamma : d_{\Omega}(x, \gamma x) \leq 1 + 2\varepsilon_0/3\}$ conjugacy classes $c \in A_I$ such that $\mathcal{L}'c(P) > 0$; this implies that $\mathcal{L}'A_I(P) \leq C_1 \#A_I^{-1}$. Hence,

$$H_{\mathcal{L}'A_I}(\mathcal{P}^{(\lceil T \rceil)}) \geq \log(\#A_I) - \log(C_1),$$

and we resume our computation using the concavity of the logarithm and Cauchy–Schwarz inequality:

$$\begin{aligned} \frac{n}{T} \sum_{I \in \mathcal{I}_T} \frac{\#A_I}{\#A_I]_{T_0, T]} H_{\mathcal{L}'A_I}(\mathcal{P}^{(\lceil T \rceil)}) &\geq \frac{n}{T} \sum_{I \in \mathcal{I}_T} \frac{\#A_I}{\#A_I]_{T_0, T]} \log(\#A_I) - \frac{n}{T} \log(C_1) \\ &\geq \frac{n}{T} \log\left(\frac{1}{\#A_I]_{T_0, T]} \sum_{I \in \mathcal{I}_T} \#A_I^2\right) - \frac{n}{T} \log(C_1) \\ &\geq \frac{n}{T} \log\left(\frac{\#A_I]_{T_0, T]}{\#\mathcal{I}_T}\right) - \frac{n}{T} \log(C_1) \\ &\geq \frac{n}{T} \log\left(\frac{\#A_T - \#A_{T_0}}{\lceil T - T_0 \rceil}\right) - \frac{n}{T} \log(C_1) \\ &\geq \frac{n}{T} \log\left(\frac{e^{(\delta_{\Gamma} - \alpha)T} - \#A_{T_0}}{\lceil T - T_0 \rceil}\right) - \frac{n}{T} \log(C_1). \end{aligned}$$

This last term converges to $n(\delta_{\Gamma} - \alpha)$ as T goes to infinity, concluding the proof. ■

We can now end the proof of Proposition 1.5.

Proof of Proposition 1.5. It is the immediate combination of Facts 2.22 and 2.23, Theorem 1.3, Propositions 8.4, 9.1, 9.2, 9.5, and Corollary 9.4. ■

9.5. Periodic geodesics and conjugacy classes

In this section, we estimate, in some cases, the asymptotic of the ratio of the number of rank-one closed geodesics by the number of rank-one conjugacy classes. The same discussion is carried in more details in [17, §6.1]; here the discussion is kept at its strict minimum: we only give the necessary definitions and observations.

Definition 9.6. Let $M = \Omega/\Gamma$ be a non-elementary rank-one convex projective orbifold. The *core-fixing subgroup* of Γ is the kernel of the restriction of Γ to the span of Λ_Γ .

Observation 9.7 ([17, Obs. 6.14]). Let $M = \Omega/\Gamma$ be a rank-one non-elementary convex projective orbifold, and let $F \subset \Gamma$ be the core-fixing subgroup. If $T^1\Omega_{\text{bip}}$ is the preimage by $\pi_\Gamma: T^1\Omega \rightarrow T^1M$ of T^1M_{bip} , then the set of vectors $v \in T^1\Omega_{\text{bip}}$ with $\text{Stab}_\Gamma(v) = F$ is open and dense in $T^1\Omega_{\text{bip}}$, and Γ -invariant.

Proposition 9.8. Let $\Omega \subset \mathbb{P}(V)$ be a properly convex open set, and $\Gamma \subset \text{Aut}(\Omega)$ a convex cocompact discrete subgroup with $M = \Omega/\Gamma$ rank-one and non-elementary. Let $F \subset \Gamma$ be the core-fixing subgroup. Let $K \subset T^1M_{\text{bip}}$ be the set vectors of whose lifts $v \in T^1\Omega$ satisfy $\text{Stab}_\Gamma(v) \neq F$. Then the topological entropy of the geodesic flow on K is strictly smaller than δ_Γ , as well as the exponential growth rate of the number of rank-one conjugacy classes of Γ with axis in K .

Proof. The subset $K \subset T^1M_{\text{bip}}$ is closed and $(\phi_t)_t$ -invariant, and has empty-interior by Observation 9.7; thus, it is given zero measure by the Bozen–Margulis measure, by Theorem 1.6. By Remark 2.43 and Observation 2.45, we can find a probability measure on K whose entropy δ is the topological entropy of $(\phi_t)_t$; this measure is different from the Bowen–Margulis probability measure since the latter gives zero measure to K . Hence, $\delta < \delta_\Gamma$ by Theorem 1.3.

Let $\Gamma' \subset \Gamma$ be a finite-index torsion-free subgroup; we set $M' = \Omega/\Gamma'$ and ε_0 to be the injectivity radius of M' ; we denote by ε_0 the injectivity radius of M' . Recall that there are only finitely many conjugacy class c of Γ such that $\ell(c) \leq \varepsilon_0$. For each rank-one conjugacy class $c \in [\Gamma]^{r_1}$ with $\ell(c) > 0$ we can find an element $\gamma_c \in c$ and a vector $v_c \in T^1\Omega_{\text{bip}} := \pi_\Gamma^{-1}T^1M_{\text{bip}}$ such that $\gamma_c \pi v_c = \pi \phi_{\ell(c)} v_c$. Let $A \subset [\Gamma]^{r_1}$ be the subset made of conjugacy classes c such that $\text{Stab}_\Gamma(v_c) \neq F$. Consider $t > 0$. Using the triangular inequality, one can check that any vector of $T^1\Omega$ belongs to at most

$$C_1 := \max_{x \in \mathcal{C}_\Omega^{\text{cor}}(\Gamma)} \#\left\{ \gamma \in \Gamma : d_\Omega(x, \gamma x) \leq 1 + \frac{2\varepsilon_0}{3} \right\}$$

balls of the family

$$\left\{ B_{T^1\Omega}^{(t+1)}\left(v_c, \frac{\varepsilon_0}{6}\right) : c \in A_{[t, t+1]} \right\}.$$

Therefore, we can extract from $\{v_c : c \in A_{[t, t+1]}\}$ a $(d_{T^1\Omega}^{(t+1)}, \varepsilon_0/6)$ -separated family of size at least $C_1^{-1} \#A_{[t, t+1]}$. The projection in T^1M' of this family is $(d_{T^1M'}^{(t+1)}, \varepsilon_0/6)$ -separated, and belongs to the preimage by $T^1M' \rightarrow T^1M$ of K . By definition of the topological entropy, the exponential growth rate of the size of such a family, when t goes to infinity, is bounded above by δ , which is strictly less than δ_Γ . ■

Remark 9.9. Let $M = \Omega/\Gamma$ be a non-elementary rank-one convex projective orbifold with compact convex core. As in [17, Obs. 6.15 and Prop. 6.16], one may use the notion of strongly primitive rank-one elements (i.e. rank-one elements $\gamma \in \Gamma$ such that $\ell(\gamma) \leq \ell(\gamma')$ for any rank-one element $\gamma' \in \Gamma$ with the same axis as γ) to prove that, if the core-fixing subgroup $F \subset \Gamma$ is the centraliser, then

$$\#[\Gamma]_T^{r1} \underset{T \rightarrow \infty}{\sim} \#[\Gamma]_T^{\text{pr1}} \underset{T \rightarrow \infty}{\sim} \#F \cdot \#\mathcal{G}_T^{r1},$$

where $[\Gamma]^{\text{pr1}}$ is the set of strongly primitive rank-one conjugacy classes in Γ and \mathcal{G}_T^{r1} is the set of rank-one periodic orbits in T^1M with period at most T .

The proof in [17] uses equidistribution results which may be replaced for the present purpose by Proposition 1.5 (one would also need Proposition 9.8). This way, one actually obtains that $\#[\Gamma]_T^{\text{pr1}}/\#\mathcal{G}_T^{r1}$ goes *exponentially fast* to $\#F$, while in [17] no rate of convergence is given.

Acknowledgements. I am grateful to Yves Benoist, Harrison Bray, Olivier Glorieux, Ludovic Marquis, Frédéric Paulin and Samuel Tapie for helpful discussions. I thank my adviser Fanny Kassel for her time, help, advice and encouragements.

I warmly thank Feng Zhu for many fruitful discussions which led to the paper [17], and made me realise that the strongly irreducible assumption in the main results of the present paper could be replaced by a weaker non-elementary assumption.

Funding. This project received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC starting grant DiGGeS, grant agreement No. 715982).

References

- [1] J. Aaronson and D. Sullivan, [Rational ergodicity of geodesic flows](#). *Ergodic Theory Dynam. Systems* **4** (1984), no. 2, 165–178 Zbl [0599.58029](#) MR [0766098](#)
- [2] D. V. Anosov, Geodesic flows on closed Riemannian manifolds of negative curvature. *Trudy Mat. Inst. Steklov.* **90** (1967) Zbl [0176.19101](#) MR [0224110](#)
- [3] M. Babillot, [On the mixing property for hyperbolic systems](#). *Israel J. Math.* **129** (2002), 61–76 Zbl [1053.37001](#) MR [1910932](#)
- [4] S. A. Ballas, J. Danciger, and G.-S. Lee, [Convex projective structures on nonhyperbolic three-manifolds](#). *Geom. Topol.* **22** (2018), no. 3, 1593–1646 Zbl [1388.57017](#) MR [3780442](#)
- [5] W. Ballmann, [Nonpositively curved manifolds of higher rank](#). *Ann. of Math. (2)* **122** (1985), no. 3, 597–609 Zbl [0585.53031](#) MR [0819559](#)
- [6] Y. Benoist, [Propriétés asymptotiques des groupes linéaires](#). *Geom. Funct. Anal.* **7** (1997), no. 1, 1–47 Zbl [0947.22003](#) MR [1437472](#)

- [7] Y. Benoist, [Automorphismes des cônes convexes](#). *Invent. Math.* **141** (2000), no. 1, 149–193 Zbl [0957.22008](#) MR [1767272](#)
- [8] Y. Benoist, Convexes divisibles. I. In *Algebraic groups and arithmetic*, pp. 339–374, Tata Institute of Fundamental Research, Mumbai, 2004 Zbl [1084.37026](#) MR [2094116](#)
- [9] Y. Benoist, [Convexes divisibles. III](#). *Ann. Sci. École Norm. Sup. (4)* **38** (2005), no. 5, 793–832 Zbl [1085.22006](#) MR [2195260](#)
- [10] Y. Benoist, [Convexes divisibles. IV](#). *Invent. Math.* **2006**, no. 2, 249–278 Zbl [1107.22006](#) MR [2218481](#)
- [11] Y. Benoist, A survey on divisible convex sets. In *Geometry, analysis and topology of discrete groups*, pp. 1–18, Adv. Lect. Math. (ALM) 6, International Press, Somerville, MA, 2008 Zbl [1154.22016](#) MR [2464391](#)
- [12] J.-P. Benzécri, [Sur les variétés localement affines et localement projectives](#). *Bull. Soc. Math. France* **88** (1960), 229–332 Zbl [0098.35204](#) MR [0124005](#)
- [13] G. Birkhoff, [Extensions of Jentzsch’s theorem](#). *Trans. Amer. Math. Soc.* **85** (1957), 219–227 Zbl [0079.13502](#) MR [0087058](#)
- [14] P.-L. Blayac, Topological mixing of the geodesic flow on convex projective manifolds. [v1] 2020, [v2] 2021, arXiv:[2009.05035v2](#), to appear in *Ann. Inst. Fourier*.
- [15] P.-L. Blayac, Dynamical aspects of convex projective structures. PhD thesis, Laboratoire Alexander Grothendieck, Université Paris-Saclay, 2021
- [16] P.-L. Blayac, [The boundary of rank-one divisible convex sets](#). *Bull. Soc. Math. France* **152** (2024), no. 1, 1–17 Zbl [1540.22021](#) MR [4762175](#)
- [17] P.-L. Blayac and F. Zhu, [Ergodicity and equidistribution in Hilbert geometry](#). *J. Mod. Dyn.* **19** (2023), 879–945 Zbl [1535.22018](#) MR [4671566](#)
- [18] M. D. Bobb, [Codimension-1 simplices in divisible convex domains](#). *Geom. Topol.* **25** (2021), no. 7, 3725–3753 Zbl [1537.57038](#) MR [4372640](#)
- [19] R. Bowen, [Periodic points and measures for Axiom A diffeomorphisms](#). *Trans. Amer. Math. Soc.* **154** (1971), 377–397 Zbl [0212.29103](#) MR [0282372](#)
- [20] R. Bowen, [Entropy-expansive maps](#). *Trans. Amer. Math. Soc.* **164** (1972), 323–331 Zbl [0229.28011](#) MR [0285689](#)
- [21] H. Bray, [Ergodicity of Bowen–Margulis measure for the Benoist 3-manifolds](#). *J. Mod. Dyn.* **16** (2020), 305–329 Zbl [1467.37036](#) MR [4182399](#)
- [22] H. Bray, [Geodesic flow of nonstrictly convex Hilbert geometries](#). *Ann. Inst. Fourier (Grenoble)* **70** (2020), no. 4, 1563–1593 Zbl [1489.37040](#) MR [4245581](#)
- [23] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*. Grundlehren Math. Wiss. 319, Springer, Berlin, 1999 Zbl [0988.53001](#) MR [1744486](#)
- [24] K. Burns and R. Spatzier, [Manifolds of nonpositive curvature and their buildings](#). *Inst. Hautes Études Sci. Publ. Math.* (1987), no. 65, 35–59 Zbl [0643.53037](#) MR [0908215](#)
- [25] S. Choi and W. M. Goldman, [Convex real projective structures on closed surfaces are closed](#). *Proc. Amer. Math. Soc.* **118** (1993), no. 2, 657–661 Zbl [0810.57005](#) MR [1145415](#)
- [26] S. Choi, G.-S. Lee, and L. Marquis, [Convex projective generalized Dehn filling](#). *Ann. Sci. Éc. Norm. Supér. (4)* **53** (2020), no. 1, 217–266 Zbl [1477.57022](#) MR [4093439](#)

- [27] D. Cooper, D. D. Long, and S. Tillmann, [On convex projective manifolds and cusps](#). *Adv. Math.* **277** (2015), 181–251 Zbl [1419.57039](#) MR [3336086](#)
- [28] Y. Coudène, [On invariant distributions and mixing](#). *Ergodic Theory Dynam. Systems* **27** (2007), no. 1, 109–112 Zbl [1129.37005](#) MR [2297089](#)
- [29] Y. Coudène, [The Hopf argument](#). *J. Mod. Dyn.* **1** (2007), no. 1, 147–153 Zbl [1109.37024](#) MR [2261076](#)
- [30] Y. Coudène, [Sur le mélange du flot géodésique](#). In *Géométrie ergodique*, pp. 13–24, Monogr. Enseign. Math. 43, L’Enseignement Mathématique, Geneva, 2013 Zbl [1316.37004](#) MR [3220549](#)
- [31] M. Crampon, [Entropies of strictly convex projective manifolds](#). *J. Mod. Dyn.* **3** (2009), no. 4, 511–547 Zbl [1189.37034](#) MR [2587084](#)
- [32] M. Crampon, [Dynamics and entropies of Hilbert metrics](#). PhD thesis, Institut de Recherche Mathématique Avancée, Université de Strasbourg, Strasbourg, 2011
- [33] M. Crampon and L. Marquis, [Finitude géométrique en géométrie de Hilbert](#). *Ann. Inst. Fourier (Grenoble)* **64** (2014), no. 6, 2299–2377 Zbl [1306.22005](#) MR [3331168](#)
- [34] M. Crampon and L. Marquis, [Le flot géodésique des quotients géométriquement finis des géométries de Hilbert](#). *Pacific J. Math.* **268** (2014), no. 2, 313–369 Zbl [1321.37026](#) MR [3227438](#)
- [35] J. Danciger, F. Guéritaud, and F. Kassel, [Convex cocompact actions in real projective geometry](#). *Ann. Sci. Éc. Norm. Supér. (4)*, **57** (2024), no. 6, pp. 1753–1843
- [36] J. Danciger, F. Guéritaud, F. Kassel, G.-S. Lee, and L. Marquis, [Convex cocompactness for Coxeter groups](#). *J. Eur. Math. Soc. (JEMS)* **27** (2025), no. 1, 119–181 Zbl [7981648](#) MR [4859567](#)
- [37] T. Foertsch and A. Karlsson, [Hilbert metrics and Minkowski norms](#). *J. Geom.* **83** (2005), no. 1-2, 22–31 Zbl [1084.52008](#) MR [2193224](#)
- [38] M. Gromov, [Hyperbolic manifolds, groups and actions](#). In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, NY, 1978)*, pp. 183–213, Ann. of Math. Stud 97, Princeton University Press, Princeton, NJ, 1981 Zbl [0467.53035](#) MR [0624814](#)
- [39] Y. Guivarc’h, [Produits de matrices aléatoires et applications aux propriétés géométriques des sous-groupes du groupe linéaire](#). *Ergodic Theory Dynam. Systems* **10** (1990), no. 3, 483–512 Zbl [0715.60008](#) MR [1074315](#)
- [40] M. Islam, [Rank-one Hilbert geometries](#). [v1] 2019, [v3] 2024, arXiv:1912.13013v3
- [41] M. Islam and A. Zimmer, [Convex cocompact actions of relatively hyperbolic groups](#). *Geom. Topol.* **27** (2023), no. 2, 417–511 Zbl [1515.20241](#) MR [4589559](#)
- [42] D. Johnson and J. J. Millson, [Deformation spaces associated to compact hyperbolic manifolds](#). In *Discrete groups in geometry and analysis (New Haven, Conn., 1984)*, pp. 48–106, Progr. Math. 67, Birkhäuser Boston, Boston, MA, 1987 Zbl [0664.53023](#) MR [0900823](#)
- [43] A. Katok and B. Hasselblatt, [Introduction to the modern theory of dynamical systems](#). Encyclopedia Math. Appl. 54, Cambridge University Press, Cambridge, 1995 Zbl [0878.58020](#) MR [1326374](#)
- [44] G. Knieper, [On the asymptotic geometry of nonpositively curved manifolds](#). *Geom. Funct. Anal.* **7** (1997), no. 4, 755–782 Zbl [0896.53033](#) MR [1465601](#)

- [45] G. Knieper, [The uniqueness of the measure of maximal entropy for geodesic flows on rank 1 manifolds](#). *Ann. of Math. (2)* **148** (1998), no. 1, 291–314 Zbl [0946.53045](#) MR [1652924](#)
- [46] G. Knieper, [Hyperbolic dynamics and Riemannian geometry](#). In *Handbook of dynamical systems, Vol. 1A*, pp. 453–545, North-Holland, Amsterdam, 2002 Zbl [1049.37020](#) MR [1928523](#)
- [47] J.-L. Koszul, [Déformations de connexions localement plates](#). *Ann. Inst. Fourier (Grenoble)* **18** (1968), no. 1, 103–114 Zbl [0167.50103](#) MR [0239529](#)
- [48] G. Link and J.-C. Picaud, [Ergodic geometry for non-elementary rank one manifolds](#). *Discrete Contin. Dyn. Syst.* **36** (2016), no. 11, 6257–6284 Zbl [1366.37090](#) MR [3543588](#)
- [49] A. Manning, [Topological entropy for geodesic flows](#). *Ann. of Math. (2)* **110** (1979), no. 3, 567–573 Zbl [0426.58016](#) MR [0554385](#)
- [50] G. A. Margulis, [Certain measures that are connected with U-flows on compact manifolds](#). *Funct. Anal. Appl.* **4** (1970), 55–67 Zbl [0245.58003](#) MR [0272984](#)
- [51] L. Marquis, [Espace des modules de certains polyèdres projectifs miroirs](#). *Geom. Dedicata* **147** (2010), 47–86 Zbl [1243.52015](#) MR [2660566](#)
- [52] J.-P. Otal, [Sur la géométrie symplectique de l'espace des géodésiques d'une variété à courbure négative](#). *Rev. Mat. Iberoamericana* **8** (1992), no. 3, 441–456 Zbl [0777.53042](#) MR [1202417](#)
- [53] S. J. Patterson, [The limit set of a Fuchsian group](#). *Acta Math.* **136** (1976), no. 3-4, 241–273 Zbl [0336.30005](#) MR [0450547](#)
- [54] A. Rényi, *Probability theory*. North-Holland Ser. Appl. Math. Mech. 10, North-Holland, Amsterdam-London; Elsevier, New York, 1970 Zbl [0233.60001](#) MR [0315747](#)
- [55] R. Ricks, [The unique measure of maximal entropy for a compact rank one locally CAT\(0\) space](#). *Discrete Contin. Dyn. Syst.* **41** (2021), no. 2, 507–523 Zbl [1460.37034](#) MR [4191516](#)
- [56] T. Roblin, [Ergodicité et équidistribution en courbure négative](#). *Mém. Soc. Math. Fr. (N.S.)* (2003), no. 95 Zbl [1056.37034](#) MR [2057305](#)
- [57] A. Selberg, [On discontinuous groups in higher-dimensional symmetric spaces](#). In *Contributions to function theory (Internat. Colloq. Function Theory, Bombay, 1960)*, pp. 147–164, Tata Institute of Fundamental Research, Bombay, 1960 Zbl [0201.36603](#) MR [0130324](#)
- [58] D. Sullivan, [The density at infinity of a discrete group of hyperbolic motions](#). *Inst. Hautes Études Sci. Publ. Math.* (1979), no. 50, 171–202 Zbl [0439.30034](#) MR [0556586](#)
- [59] N. Tholozan, [Volume entropy of Hilbert metrics and length spectrum of Hitchin representations into \$\mathrm{PSL}\(3, \mathbb{R}\)\$](#) . *Duke Math. J.* **166** (2017), no. 7, 1377–1403 Zbl [1375.53022](#) MR [3649358](#)
- [60] C. Vernicos, [Approximability of convex bodies and volume entropy in Hilbert geometry](#). *Pacific J. Math.* **287** (2017), no. 1, 223–256 Zbl [1364.53068](#) MR [3613440](#)
- [61] J. Vey, [Sur les automorphismes affines des ouverts convexes saillants](#). *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)* **24** (1970), 641–665 Zbl [0206.51302](#) MR [0283720](#)
- [62] C. Walsh, [The horofunction boundary of the Hilbert geometry](#). *Adv. Geom.* **8** (2008), no. 4, 503–529 Zbl [1155.53335](#) MR [2456635](#)

- [63] F. Zhu, Ergodicity and equidistribution in strictly convex Hilbert geometry. [v1] 2020, [v3] 2021, [arXiv:2008.00328v3](https://arxiv.org/abs/2008.00328v3)
- [64] A. Zimmer, [A higher-rank rigidity theorem for convex real projective manifolds](#). *Geom. Topol.* **27** (2023), no. 7, 2899–2936 Zbl [1534.53050](#) MR [4645488](#)

Received 13 April 2023.

Pierre-Louis Blayac

Laboratoire Alexander Grothendieck, Institut des Hautes Études Scientifiques,
Université Paris-Saclay, 35 route de Chartres, 91440 Bures-sur-Yvette, France;
blayac@unistra.fr, plblayac@hotmail.fr