

Local complete intersections and Weierstrass points

André Contiero and Sarah Mazzini

Abstract. This work presents a simple proof that the moduli space of complete integral Gorenstein curves with a prescribed symmetric Weierstrass semigroup becomes a weighted projective space, even for fields of positive characteristic, when the associated monomial curve is a local complete intersection.

1. Introduction

Given a numerical semigroup $S \subset \mathbb{N}$ of genus $g \geq 1$, minimally generated by a_1, \dots, a_r , let $\mathcal{M}_{g,1}^S$ be the moduli space parameterizing smooth pointed curves defined over an algebraically closed field \mathbf{k} (or compact Riemann surfaces when $\mathbf{k} = \mathbb{C}$), whose Weierstrass semigroup at the marked point is S . It is well known that $\mathcal{M}_{g,1}^S$ can be empty depending on S , but when it is non-empty, a major and very classical problem is to describe the moduli space $\mathcal{M}_{g,1}^S$ and its compactification.

By allowing singularities, any numerical semigroup S can be realized as the Weierstrass semigroup of a projectivization of the affine monomial curve

$$\mathcal{C}_S := \{(t^{a_1}, \dots, t^{a_r}); t \in \mathbf{k}\} \subset \mathbb{A}^r.$$

Herzog [10] showed that the ideal of \mathcal{C}_S can be generated by suitable polynomials in $\mathbf{k}[X_{a_1}, \dots, X_{a_r}]$ which are differences of two monomials with the same weighted degree, namely

$$G_{d_j}^{(0)} := X_{a_1}^{\alpha_{1,j}} \cdots X_{a_r}^{\alpha_{r,j}} - X_{a_1}^{\beta_{1,j}} \cdots X_{a_r}^{\beta_{r,j}},$$

where $\alpha_{i,j} \cdot \beta_{i,j} = 0$ and $\sum a_i \alpha_{i,j} = \sum a_i \beta_{i,j}$ for $1 \leq i \leq r$ and $1 \leq j \leq m$.

The purpose of this paper is to establish the following result.

Mathematics Subject Classification 2020: 14H10 (primary); 14H55, 14B07, 14M10 (secondary).

Keywords: Weierstrass points, deformation of singularities, local complete intersections, moduli of curves.

Main Theorem. *If S is such that the affine monomial curve $\mathcal{C}_S = \text{Spec } \mathbf{k}[S]$ is a local complete intersection and $\text{char}(\mathbf{k}) = 0$ or a prime not dividing any exponent $\alpha_{i,j}$ and $\beta_{i,j}$ of the defining equations of \mathcal{C}_S , then a compactification*

$$\overline{\mathcal{M}_{g,1}^S} = \mathbb{P}(\mathbf{T}^{1,-}),$$

is constructed and the closure is compounded by integral Gorenstein curves with a smooth point whose Weierstrass semigroup is S .

The vector space $\mathbf{T}^{1,-}$ stands for the negatively graded part of the first module of the cotangent complex associated to the semigroup algebra $\mathbf{k}[S] = \bigoplus_{n \in S} \mathbf{k}t^n$,

$$\mathbf{T}^1(\mathbf{k}[S]) = \mathbf{T}^{1,-}(\mathbf{k}[S]) \oplus \mathbf{T}^{1,+}(\mathbf{k}[S]).$$

We recall that a numerical semigroup S is a complete intersection if the affine monomial curve \mathcal{C}_S is a complete intersection in \mathbb{A}^r , where r is the embedding dimension of \mathcal{C}_S , i.e., the smallest number of elements required to generate S . Equivalently, the semigroup algebra $\mathbf{k}[S]$ is a complete intersection when we consider it as the quotient of $\mathbf{k}[X_{a_1}, \dots, X_{a_r}]$ by the kernel \mathbf{I} of the surjective map

$$\begin{aligned} \mathbf{k}[X_{a_1}, \dots, X_{a_r}] &\longrightarrow \mathbf{k}[S], \\ X_{a_i} &\longmapsto t^{a_i}, \end{aligned}$$

and \mathbf{I} is the defining ideal of $\mathcal{C}_S \subset \mathbb{A}^r$.

The affine monomial curve \mathcal{C}_S has a unique unibranch singular point at the origin $\mathbf{0}$, with singularity degree $g = g(S)$. Therefore \mathcal{C}_S , or even its closure in a suitable (weighted) projective space, is a local complete intersection if and only if the local ring of its unique singularity is a complete intersection. Since \mathcal{C}_S is affine and a locally complete intersection, a minimal free resolution of the local ring singularity lifts to a minimal free resolution of the semigroup algebra $\mathbf{k}[S]$, and hence \mathcal{C}_S is a global complete intersection in \mathbb{A}^r .

If $\mathbf{k}[S]$ is a complete intersection, then there are no obstructions to formally deform \mathcal{C}_S in characteristic zero, the second cohomology module of the cotangent complex associated to \mathcal{C}_S is null, $\mathbf{T}_{\mathcal{C}_S}^2 = 0$, as shown in [11]. Hence, we can conclude that $\mathcal{M}_{g,1}^S$ is smooth. Furthermore, the base space \mathcal{T}^- of the miniversal deformation in negative degrees is an affine space \mathbb{A}^N . Therefore, we can deduce that a closure of $\mathcal{M}_{g,1}^S$ is also a projective space, whenever we apply Pinkham's construction of $\mathcal{M}_{g,1}^S$ for smooth fibers $\mathcal{X}^- \rightarrow \mathcal{T}^-$ of the miniversal deformation (see Section 3.4 for more details). The advantage of our techniques is that the proof of the main theorem presented here is rather explicit and simple, and it also works for fields of positive characteristic and describes the curves that compound the boundary.

A rather simple proof of the main theorem in characteristic zero can be obtained as follows.

Proof. The dimension of $\mathcal{M}_{g,1}^S$ is at least $2g - 1 - \dim \mathbf{T}^{1,+}$, cf. [5, Theorem 2.4]. Since the monomial curve in \mathbb{P}^{g-1} associated to \mathcal{C}_S is a local complete intersection and $\text{char}(\mathbf{k}) = 0$, we are able to show that $\dim \mathbf{T}^{1,-} = 2g - \dim \mathbf{T}^{1,+}$, meaning that the Tjurina number of a complete intersection singularity is $2g$. At this point we just have to apply the results due to Stöhr and Contiero–Stöhr [6, 20] assuring that $\overline{\mathcal{M}_{g,1}^S}$ is a closed subset of $\mathbb{P}(\mathbf{T}^{1,-})$. ■

The way we prove the main theorem for also fields of positive characteristic is to apply a variant of Hauser’s algorithm (see [8, 9] and [18]) by deforming the affine monomial curve $\mathcal{C}_S \subset \mathbb{A}^r$ instead of the associated canonical Gorenstein monomial curve in \mathbb{P}^{g-1} , as required by Stöhr’s original construction ([20]). The first step is to take the unfold of the $r - 1$ defining equations of the ideal of \mathcal{C}_S . Next, since \mathcal{C}_S is a complete intersection, we can show that no relations between the unfolded coefficients arise from syzygies, with the exception of $\frac{1}{2}r(r + 1)$ normalizations to zero. This is where the condition on the characteristic of the ground field appears. Hence, the closure of the moduli space $\mathcal{M}_{g,1}^S$ is $\mathbb{P}(V)$, where V is the \mathbf{k} -vector space spanned by the normalized unfolded coefficients. Finally, we just need to note that V is in bijection with $\mathbf{T}^{1,-}$, cf. [20, Appendix].

We obtain the following two naive and immediate consequences of the above main theorem, provided that $\text{char}(\mathbf{k}) = 0$ or a prime not dividing any exponent $\alpha_{i,j}$ and $\beta_{i,j}$ of the defining equations of \mathcal{C}_S .

Corollary 1.1 (Schlessinger [17] and Pinkham [16]). *A complete intersection numerical semigroup is realized as a Weierstrass semigroup of a smooth curve.*

Corollary 1.2. *If \mathcal{C}_S is a local complete intersection, then the associated affine monomial curve can be negatively smoothed without any obstruction.*

In general, it is very difficult to describe a compactification of $\mathcal{M}_{g,1}^S$ and the curves that make up its boundary. The authors are aware of two main approaches to considering geometric features of a closure of $\mathcal{M}_{g,1}^S$ and properties of curves on its boundary. In the following two subsections, we cite some results concerning these two approaches. There are many high-standing works that are not cited here, most of which are referenced in the works cited below.

1.1. $\mathcal{M}_{g,1}^S$ coming from versal deformation

The general theory of versal deformations of singularities dates back to the 1960s and 1970s, with the remarkable works of Schlessinger [17] and Artin [1]. The connection between the spaces $\mathcal{M}_{g,1}^S$ and the miniversal deformation in negative degrees was made by Pinkham in his Ph.D. thesis [16], using an affine monomial curve associated

with the semigroup S . We shall briefly describe this connection in Section 3.4 below, as it is one of the main techniques used in this paper.

Several works have investigated the application of versal deformations to the study of $\mathcal{M}_{g,1}^S$. As demonstrated by Pinkham in [16], the miniversal deformation provides a method for constructing a compactification of $\mathcal{M}_{g,1}^S$. The resulting closure of $\mathcal{M}_{g,1}^S$ is totally described just for a few families of semigroups, as we note below.

In [20], Stöhr presents a rather explicit way to construct a compactification of $\mathcal{M}_{g,1}^S$ as a variant of Hauser's algorithm, when S is assumed to be a suitable symmetric semigroup. Stöhr's construction relies on the unfold of the defining equations of the canonically embedded projective monomial curve associated to \mathcal{C}_S , extending Petri's analysis of the canonical ideal and then exploring appropriate syzygies coming from the defining equations of \mathcal{C}_S . It is obtained a compactification of $\mathcal{M}_{g,1}^S$ as a closed subset of a weighted projective space by allowing irreducible Gorenstein curves at the boundary. Later on, Contiero–Stöhr [6] and Contiero–Fontes [4] extend Stöhr's construction to all symmetric semigroups, making it totally implementable as well. In Section 3.4 below, we briefly recall this construction. We also refer to [13], where the second author presents some algorithms to compute the defining equations of \mathcal{C}_S and their unfolding, the defining equations of $\overline{\mathcal{M}_{g,1}^S}$, and the equivariant tangent space $\mathbf{T}_{\mathcal{C}_S}^1$ of the versal deformation space, whenever S is symmetric.

Nakano [14] computed $\mathcal{M}_{g,1}^S$ using Pinkham's approach by computationally determining the base space of the miniversal deformation of the monomial curve \mathcal{C}_S in negative degrees, for $g \leq 5$. He shows that for $g \leq 5$ the base space is an irreducible rational variety, except in one case: the semigroup $\langle 4, 6, 11, 13 \rangle$ when it has the structure of a projective quasi-cone over $\mathbb{P}^1 \times \mathbb{P}^3$. In this case, the base space is also irreducible, but in negative degrees it contains two components, one smooth and the other containing a curve with a double point (see [5, Remark 2.9]).

In a recent paper [19], Stevens extends the results of Nakano [14] and explicitly computes the defining equations of the moduli space $\mathcal{M}_{g,1}^S$ for many cases of genus at most seven and determines the dimension for all semigroups of genus not greater than seven. Stevens uses Hauser's algorithm in most cases, but in one case, he uses the projection method developed by De Jong and Van Straten [7].

2. Reviewing Weierstrass points

We recall that a numerical semigroup S is a subset of the nonnegative integers \mathbb{N} containing 0, closed under addition such that only a finite number of elements are missing from S . The genus of S is the number of its gaps, i.e., the number of positive

integers that are not in S ,

$$g(S) := \#(\mathbb{N} \setminus S) = \#\{1 = \ell_1 < \dots < \ell_g\},$$

and we easily see that the largest gap ℓ_g is not bigger than $2g - 1$.

Given an irreducible smooth pointed curve $(\mathcal{C}, P) \in \mathcal{M}_{g,1}$ of genus g , its associated Weierstrass semigroup S_P is the subset of all nonnegative integers n such that

$$H^0(\mathcal{C}, \mathcal{O}_{\mathcal{C}}((n-1)P)) \subsetneq H^0(\mathcal{C}, \mathcal{O}_{\mathcal{C}}(nP)), \tag{2.1}$$

i.e., $n \in S_P$ if and only if there is a rational function on \mathcal{C} whose pole divisor is exactly nP . The point $P \in \mathcal{C}$ is called a Weierstrass point if S_P is different from the ordinary semigroup $\{0, g + 1, g + 2, \dots\}$. The Riemann–Roch theorem implies that the genus of the Weierstrass semigroup S_P is equal to the genus of the curve \mathcal{C} . It is well known that only a finite number of Weierstrass points exist on a curve.

Since the i -th gap of S defines an upper semicontinuous function on $\mathcal{M}_{g,1}$, it follows that $\mathcal{M}_{g,1}^S$ is a locally closed subset of $\mathcal{M}_{g,1}$. However, it is also well known that the moduli space $\mathcal{M}_{g,1}^S$ can be empty, meaning that there are numerical semigroups that cannot be realized as Weierstrass semigroups of a smooth pointed curve. There is no purely arithmetical criterion for determining when a numerical semigroup is realizable, but one necessary numerical condition is given by Buchweitz in [3].

On the other hand, one can see that any numerical semigroup can be realized as a Weierstrass semigroup of a monomial curve. Taking $S := \langle a_1, \dots, a_r \rangle$, a numerical semigroup, let $\mathbf{k}[S] := \bigoplus_{n \in S} \mathbf{k}t^n$ be the associated semigroup algebra. The affine monomial curve attached to S is

$$\mathcal{C}_S = \text{Spec } \mathbf{k}[S] \subset \mathbb{A}^r,$$

which in parametric terms is just

$$\mathcal{C}_S = \{(t^{a_1}, \dots, t^{a_r}) \in \mathbb{A}^r; t \in \mathbb{A}^1\}.$$

It is easy to produce the closure of \mathcal{C}_S in a weighted projective space \mathbb{P}^r by adding just a smooth point P at infinity, and so the Weierstrass semigroup at P is S . Here, a Weierstrass point on an integral curve \mathcal{C} at a smooth point P is defined in the same way that when \mathcal{C} is smooth, i.e., for a smooth point P on \mathcal{C} , a positive integer n is a nongap if and only if equation (2.1) holds.

A criterion for determine if a numerical semigroup S is realizable was given by Pinkham in his Ph.D. thesis. Namely, a numerical semigroup S is realizable if and only if the affine monomial curve \mathcal{C}_S admits a negative smoothing, cf. [16, p. 108]. Dealing with this criterion is unfortunately far from easy.

3. Gorenstein curves and subcanonical points

In this section, \mathcal{C} stands for a non-hyperelliptic Gorenstein curve with a smooth subcanonical point P , i.e., the associated Weierstrass semigroup S at P is symmetric. Recall that a numerical semigroup is symmetric if the Frobenius number ℓ_g of S is the biggest possible, $\ell_g = 2g - 1$. Equivalently,

$$\ell_{g-i} = 2g - 1 - n_i \quad (0 \leq i \leq g - 1),$$

where $0 = n_0 < n_1 < n_2 < \dots$ are the nongaps of the semigroup. Since it is assumed S to be non-hyperelliptic, we may impose that $\ell_2 = 2$, equivalently, $n_{g-1} = 2g - 2$.

We also fix at once a system of generators, $S := \langle a_1, \dots, a_r \rangle$. We are interested in two suitable systems of generators: the minimal system, where r is the embedding dimension of S , and the canonical system of generators, i.e., $r = g - 1$. In the former case, S is generated by its first g nongaps, $S = \langle n_0, n_1, \dots, n_{g-1} \rangle$.

As a general and important comment, if a curve \mathcal{C} is a local complete intersection, then \mathcal{C} is also Gorenstein and non-hyperelliptic, because the dualizing sheaf of a local complete intersection always induces an embedding.

3.1. On P -hermitian bases

By virtue of the Max Noether theorem for non-hyperelliptic Gorenstein curves ([6]), the maps

$$\text{Sym}^n H^0(\mathcal{C}, \omega) \longrightarrow H^0(\mathcal{C}, \omega^n)$$

are surjective for all $n \geq 1$, where $\omega \cong \mathcal{O}_{\mathcal{C}}((2g - 2)P)$ is the dualizing sheaf of \mathcal{C} . Hence, each vector space $H^0(\mathcal{C}, \omega^n)$ admits a so-called P -hermitian basis, i.e., given $n \geq 1$, for each nongap $S \leq n(2g - 2)$ we can choose a meromorphic function on \mathcal{C} of the form $\mathbf{x}_S^\alpha := x_{a_0}^{\alpha_0} \dots x_{a_r}^{\alpha_r}$ satisfying

$$\text{ord}_{\infty, P} \mathbf{x}_S^\alpha = \sum \alpha_i a_i = s,$$

where each x_{a_i} is a regular function on $\mathcal{C} \setminus \{P\}$ whose pole order at P is $\text{ord}_{\infty, P}(x_{a_i}) = a_i$. We also may declare $a_0 = 0$, so one can assume that $x_{a_0} = 1$. Hence, each $H^0(\mathcal{C}, \omega^n)$ admits a base formed by meromorphic functions on \mathcal{C} whose pole orders at P are pairwise distinct.

In order to have a uniqueness between the chosen basis elements \mathbf{x}_S^α , one can take them in a way that $\alpha := (\alpha_0, \dots, \alpha_r) \in \mathbb{N}^{r+1}$ is a minimal element according to the lexicographical order

$$\left(\sum_{i=0}^r \alpha_i, \sum_{i=0}^r a_i \alpha_i, -\alpha_0, -\alpha_{r-1}, \dots, -\alpha_1 \right). \tag{3.1}$$

Hence,

$$H^0(\mathcal{C}, \omega^n) = \text{Spam} \bigcup_{s \leq n(2g-2)} \{\mathbf{x}_s^\alpha; \alpha \text{ is minimal}\}. \quad (3.2)$$

For each $n \geq 1$, Δ_n stands for the vector subspace of $\mathbf{k}[X_{a_0}, \dots, X_{a_r}]$ spanned by the lifting of the above monomial basis of $H^0(\mathcal{C}, \omega^n)$, namely

$$\Delta_n := \text{Spam} \left(\bigcup_s \{\mathbf{X}_s^\alpha; s \leq n(2g-2) \text{ and } \alpha \text{ minimal}\} \right), \quad (3.3)$$

where $\mathbf{X}_s^\alpha := X_{a_0}^{\alpha_0} \dots X_{a_r}^{\alpha_r}$, with $\sum_{i=1}^r a_i \alpha_i = s$.

We define $\deg(X_{a_i}) = a_i$. It follows from the Riemann–Roch theorem for singular curves that $\dim_{\mathbf{k}} \Delta_n = (2n-1)(g-1)$ and so

$$\dim \mathbf{k}[X_{a_0}, \dots, X_{a_r}]_{\leq n} = \binom{n+g-1}{n} - (2n-1)(g-1),$$

where $\mathbf{k}[X_{a_0}, \dots, X_{a_r}]_{\leq n}$ stands for the vector space over \mathbf{k} given by the isobaric polynomials of (weighted) degree not bigger than $n(2g-2)$.

Remark 3.1. Considering the canonical system of generators for $S = \langle n_0, \dots, n_{g-1} \rangle$, the above process produces a basis for Δ_n (respectively for $H^0(\mathcal{C}, \omega^n)$) that is formed just by monomials on X_{n_i} (respectively on x_{n_i}) all of the same degree n , which does not happen when we consider the minimal system of generators. For instance, the base elements of Δ_2 , respectively Δ_3 , are given by quadratic forms $X_{a_s} X_{b_s}$ with $S \leq 4g-4$, respectively by cubic forms $X_{u_\sigma} X_{v_\sigma} X_{w_\sigma}$ with $\sigma \leq 6g-6$, according to the order fixed in (3.1). We may also conclude that

$$\dim \mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}]_n = \binom{n+g-1}{n} - (2n-1)(g-1),$$

where $\mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}]_n$ stands for the usual vector space given by the forms of degree n .

3.2. The ideal of the canonically embedded \mathcal{C}

We start by identifying \mathcal{C} with its image under the canonical embedding given by its dualizing sheaf ω . In this way, \mathcal{C} can be viewed as a curve of genus g and degree $2g-2$ in \mathbb{P}^{g-1} . Let $I(\mathcal{C}) = \bigoplus_{j=2}^{\infty} I_j(\mathcal{C})$ be the homogeneous ideal of \mathcal{C} . By Riemann's theorem, for each $j \geq 2$, the codimension of $I_j(\mathcal{C})$ in the vector space $\mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}]_j$ is $(2j-1)(g-1) = \dim_{\mathbf{k}} \Delta_j$. Then we obtain

$$\mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}]_j = \Delta_j \oplus I_j(\mathcal{C}), \quad \text{for each } j \geq 2.$$

Recall, see [15, Theorems 1.7 and 1.9], that each nongap $s \leq 4g - 4$ can be written in ν_s different ways as a sum of two nongaps not bigger than $2g - 2$, namely

$$s = a_{s_1} + b_{s_1} = \cdots = a_{s\nu_s} + b_{s\nu_s},$$

where $a_{s_1} < \cdots < a_{s\nu_s}$ and $a_{s_i} \leq b_{s_i}$, for all $i = 1, \dots, \nu_s$. Analogously, there are ν_σ different ways to write each nongap $\sigma \leq 6g - 6$ as a sum of three nongaps,

$$\sigma = u_{\sigma 1} + v_{\sigma 1} + w_{\sigma 1} = \cdots = u_{\sigma\nu_\sigma} + v_{\sigma\nu_\sigma} + w_{\sigma\nu_\sigma}.$$

Since $x_{a_{s_i}}x_{b_{s_i}}$ and $x_{u_{\sigma j}}x_{v_{\sigma j}}x_{w_{\sigma j}}$ are elements in $H^0(\mathcal{C}, \omega^2)$ and $H^0(\mathcal{C}, \omega^3)$, respectively, we may assume that $x_{a_{s_1}}x_{b_{s_1}} := x_{a_s}x_{b_s}$ and $x_{u_{\sigma 1}}x_{v_{\sigma 1}}x_{w_{\sigma 1}} := x_{u_\sigma}x_{v_\sigma}x_{w_\sigma}$ are base elements of Δ_2 and Δ_3 , respectively, cf. Remark 3.1. Hence, for each $i = 2, \dots, \nu_s$ and each $j = 2, \dots, \nu_\sigma$ the elements $x_{a_{s_i}}x_{b_{s_i}}$ and $x_{u_{\sigma j}}x_{v_{\sigma j}}x_{w_{\sigma j}}$ can be written as a linear combination of base elements, preserving the pole order at P , namely

$$\begin{aligned} x_{a_{s_i}}x_{b_{s_i}} &= c_{s_i s}x_{a_s}x_{b_s} + \sum_{n < s} c_{s i n}x_{a_n}x_{b_n}, \\ x_{u_{\sigma j}}x_{v_{\sigma j}}x_{w_{\sigma j}} &= d_{\sigma j \sigma}x_{u_\sigma}x_{v_\sigma}x_{w_\sigma} + \sum_{m < \sigma} d_{\sigma j m}x_{u_m}x_{v_m}x_{w_m} \end{aligned}$$

where n and m run over the nongaps and $c_{s i n}, d_{\sigma j m} \in \mathbf{k}$ are constants. We also may assume that $c_{s i s} = d_{\sigma j \sigma} = 1$, because they must be different from zero and so we can multiply them by suitable constants. By construction, the $\frac{1}{2}(g-2)(g-3)$ quadratic forms

$$F_{s_i} := X_{a_{s_i}}X_{b_{s_i}} - X_{a_s}X_{b_s} - \sum_{n=0}^{s-1} c_{s i n}X_{a_n}X_{b_n} \in \mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}] \quad (3.4)$$

and the $\binom{g+2}{3} - (5g-5)$ cubic forms

$$\begin{aligned} G_{\sigma j} &= X_{a_{\sigma j}}X_{b_{\sigma j}}X_{c_{\sigma j}} - X_{a_\sigma}X_{b_\sigma}X_{c_\sigma} \\ &\quad - \sum_{n=0}^{\sigma-1} d_{\sigma j n}X_{a_n}X_{b_n}X_{c_n} \in \mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}], \end{aligned} \quad (3.5)$$

vanish identically on the canonical curve \mathcal{C} , are linearly independent and because of their number, form a basis of the vector spaces $I_2(\mathcal{C})$ and $I_3(\mathcal{C})$, respectively.

Petri's analysis remains true for canonical Gorenstein curves and assures that the ideal $I(\mathcal{C})$ is generated by quadratic relations, provided \mathcal{C} is non-hyperelliptic, non-trigonal and not isomorphic to a quintic plane curve. When \mathcal{C} is trigonal or isomorphic to a quintic plane curve, it assures that $I(\mathcal{C})$ is generated by quadratic and cubic forms. It turns out that if S is such that $3 < n_1 < g$ and $S \neq \langle 4, 5 \rangle$, then \mathcal{C} is nontrigonal

and not isomorphic to a quintic plane curve. Hence the ideal $I(\mathcal{C})$ is generated by the quadratic forms in equation (3.4), see [20] or Theorem 3.4 below. Therefore, if S is such that $n_1 = 3$, $n_1 = g$ or $S = \langle 4, 5 \rangle$, then the ideal $I(\mathcal{C})$ is generated by the quadratic forms in (3.4) and suitable cubic forms picked up from (3.5), cf. [4, Theorem 3.7].

It is worth to note that each non-hyperelliptic numerical semigroup S can be realized as the Weierstrass semigroup of a Gorenstein (canonical) curve. Namely, taking the canonical monomial curve

$$\mathcal{C}^{(0)} := \{(a^{n_0} b^{\ell_{g-1}} : a^{n_1} b^{\ell_{g-1}-1} : \dots : a^{n_{g-1}} b^{\ell_1-1}; (a : b) \in \mathbb{P}^1)\} \subset \mathbb{P}^{g-1}, \quad (3.6)$$

the Weierstrass semigroup at $P = (0 : \dots : 0 : 1)$ is equal to S , cf. [20, p. 190]. Moreover, the ideal of $\mathcal{C}^{(0)}$ is generated by the following $\frac{1}{2}(g-2)(g-3)$ *folded quadratic forms* (see [6, Lemma 2.3]):

$$F_{si}^{(0)} = X_{a_{si}} X_{b_{si}} - X_{a_s} X_{b_s}, \quad (3.7)$$

provided that $3 < n_1 < g$ and $S \neq \langle 4, 5 \rangle$. In addition, if $n_1 = 3$, $n_1 = g$ or $S = \langle 4, 5 \rangle$, then the ideal of $I(\mathcal{C}^{(0)})$ is generated by the above $\frac{1}{2}(g-2)(g-3)$ *folded quadratic forms* and suitable *folded cubic forms*

$$G_{\sigma j}^{(0)} = X_{a_{\sigma j}} X_{b_{\sigma j}} X_{c_{\sigma j}} - X_{a_{\sigma}} X_{b_{\sigma}} X_{c_{\sigma}}, \quad (3.8)$$

cf. [4, Lemma 3.3].

3.3. Unfolding the defining equations

Given the monomial curve $\mathcal{C}_S \subset \mathbb{A}^r$ associated to any non-ordinary numerical semigroup $S = \langle a_1, \dots, a_r \rangle$, a result due to Herzog, [10], assures that the generators of the ideal $I(\mathcal{C}_S)$ can be chosen to be *isobaric forms* which are given by the difference of two monomials in the variables X_{a_1}, \dots, X_{a_r} , namely

$$\mathbf{X}_s^\alpha - \mathbf{X}_s^\beta,$$

such that $\alpha_i \beta_i = 0$ for $i = 1, \dots, r$ and $\sum a_i \alpha_i = \sum a_i \beta_i = s$ is its weight.

When we assume non-hyperelliptic symmetric semigroups, we can consider two systems of generators for S , namely the minimal and the canonical ones. By considering the minimal system of generators, a_1, \dots, a_r , we can choose a basis of $\Delta_i \subset \mathbf{k}[X_{a_0}, X_{a_1}, \dots, X_{a_r}]$, for $i \geq 2$, that is given by the lifting of the P -hermitian basis of $H^0(C, \omega^i)$, where ω is the dualizing sheaf of \mathcal{C}_S , see equation (3.3) of Section 3.1. If $H^{(0)}$ is a generating form of $I(\mathcal{C}_S)$, say $H^{(0)} = \mathbf{X}_s^\alpha - \mathbf{X}_s^\beta \in I(\mathcal{C}_S)$ of

weight s , let n be the smallest positive integer such that $s \leq n(2g - 2)$. Thus the *unfold* of $H^{(0)}$ is the polynomial

$$H_s = H_s^{(0)} + \sum_{j < s} c_{sj} \mathbf{X}_j^{\mathcal{Y}}|_{X_{a_0}=1} \in \mathbf{k}[\{c_{sj}\}] \otimes \mathbf{k}[X_{a_1}, \dots, X_{a_r}], \quad (3.9)$$

where each $\mathbf{X}_j^{\mathcal{Y}}$ is the unique basis element of Δ_n of weight j , $\mathbf{X}_j^{\mathcal{Y}}|_{X_{a_0}=1}$ is the monomial obtained from $\mathbf{X}_j^{\mathcal{Y}}$ making $X_{a_0} = 1$ and c_{sj} are variables over the ground field \mathbf{k} . We attach a weight $s - j$ to each c_{sj} . Since the weight of X_{a_0} is zero, the *unfold* of $H^{(0)}$ is also an isobaric form of degree s .

By considering the canonical system of generators for \mathbf{S} , namely n_0, n_1, \dots, n_{g-1} , the canonical ideal of $\mathcal{C}^{(0)}$ is generated by isobaric forms that are also homogeneous polynomials (quadratic and cubic) in the usual sense, cf. equation (3.7) and equation (3.8). Then the *unfold* of a defining quadratic form $F_{si}^{(0)}$ is

$$F_{si} = F_{si}^{(0)} - \sum_{n=0}^{s-1} c_{sin} X_{a_n} X_{b_n} \in \mathbf{k}[\{c_{sij}\}] \otimes \mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}], \quad (3.10)$$

while the *unfold* of a cubic defining form $G_{sj}^{(0)}$ is

$$G_{sj} = G_{sj}^{(0)} - \sum_{n=0}^{\sigma-1} d_{sjn} X_{a_n} X_{b_n} X_{c_n} \in \mathbf{k}[\{d_{sij}\}] \otimes \mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}]. \quad (3.11)$$

Note that the *unfold* of the quadratic and cubic defining equations of the canonical curve $\mathcal{C}^{(0)}$ are again quadratic and cubic forms in $\mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}]$ and isobaric as well, provided the weight of c_{sin} and d_{sjn} are $s - n$ and $\sigma - n$, respectively.

It is evident that the *unfold* of the defining equations of a monomial curve is a perturbation of its defining ideal. To get a deformation preserving at least the dimension and the arithmetical genus over the fibers, these perturbations cannot be chosen independently. Generally, they are related by *syzygetic* relations. This is precisely the subject of the next subsection.

3.4. A variant of Hauser’s algorithm

In his Ph.D. thesis [16], Pinkham constructs the moduli space $\mathcal{M}_{g,1}^{\mathbf{S}}$ using equivariant (versal) deformation theory. In short, Pinkham starts by considering the versal deformation space of the affine monomial curve $\mathcal{C}_{\mathbf{S}}$, say

$$\begin{array}{ccc} \mathcal{X}_{t_0} \cong \mathcal{C}_{\mathbf{S}} & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \{t_0\} = \text{Spec } \mathbf{k} & \longrightarrow & \tilde{\mathcal{T}} \end{array}$$

where $\mathcal{T} = \text{Spec } A$ and A is a local, complete noetherian \mathbf{k} -algebra, cf. [1]. The \mathbb{G}_m -action on \mathcal{C}_S , given by $(\zeta, X_{a_i}) \mapsto \zeta^{a_i} X_{a_i}$, can be extended to the total and parameter spaces, \mathcal{X} and \mathcal{T} , inducing a grading on the tangent space $\mathbf{T}_{\mathcal{C}_S}^1 \cong \mathbf{T}^1(\mathbf{k}[S])$ to \mathcal{T} , that is the cotangent complex associated to \mathcal{C}_S .

We declare that a deformation has negative weight $-e$ if it decreases the weights of the defining equations of the curve and the corresponding deformation variable has then (positive) weight e . It is more than convenient to note that the unfolds of the defining forms of \mathcal{C}_S and $\mathcal{C}^{(0)}$ in Equations (3.9), (3.10) and (3.11) of the preceding subsection occur in negative degrees, once provided that they define a deformation of \mathcal{C}_S and $\mathcal{C}^{(0)}$, respectively.

Let \mathbf{I} be the ideal of A generated by the elements corresponding to the positive graded part $\mathbf{T}^{1,+}(\mathbf{k}[S])$. The space $\mathcal{T}^- := \text{Spec } A/\mathbf{I}$ is the subspace of \mathcal{T} in negative degrees and the restriction $\mathcal{X}^- \rightarrow \mathcal{T}^-$ is the versal deformation in negative degrees,

$$\begin{array}{ccc} \mathcal{X}_{t_0} \cong \mathcal{C}_S & \longrightarrow & \mathcal{X}^- \\ \downarrow & & \downarrow \\ \{t_0\} = \text{Spec } \mathbf{k} & \longrightarrow & \mathcal{T}^- = \text{Spec}(A/\mathbf{I}). \end{array}$$

In addition, the total space \mathcal{X}^- and the parameter space \mathcal{T}^- are both defined by polynomials. In general, the total and parameter spaces associated to an analytic singularity cannot be defined by polynomial equations alone, and sometimes do not have a finite dimension. However, this does not happen when deforming quasi-homogeneous singularities.

Next Pinkham produces a fiberwise compactification $\overline{\mathcal{X}}^- \rightarrow \mathcal{T}^-$ of the versal deformation in negative degrees $\mathcal{X}^- \rightarrow \mathcal{T}^-$ without compactifying the parameter and the total space, avoiding technical problems coming from inverse limits. Doing this, Pinkham shows that each fiber of $\overline{\mathcal{X}}^- \rightarrow \mathcal{T}^-$ is an integral curve in a weighted projective space with one point P at infinity whose associated Weierstrass semigroup is exactly S . All the fibers over a given \mathbb{G}_m orbit of \mathcal{T}^- are isomorphic, and two fibers are isomorphic if and only if they lie in the same orbit. This is proved in [16] for smooth fibers and in general in the Appendix of [12].

Now, let us invert the above considerations starting with a possible singular integral curve \mathcal{C} of arithmetic genus $g > 1$ defined over \mathbf{k} . Given a smooth point P of \mathcal{C} , let S be the Weierstrass semigroup of \mathcal{C} at P . Consider the line bundle $L = \mathcal{O}_{\mathcal{C}}(P)$ and form the ring of sections $\mathcal{R} = \bigoplus_{i=0}^{\infty} H^0(\mathcal{C}, L^i)$. This leads to an embedding of $\mathcal{C} = \mathbb{P}(\mathcal{R})$ in a weighted projective space, with coordinates X_{a_0}, \dots, X_{a_r} with $\text{deg}(X_{a_0}) = 1$. The space $\text{Spec } \mathcal{R}$ is the corresponding quasi-cone in an affine space. Setting $X_{a_0} = 0$ defines the monomial curve \mathcal{C}_S , all other fibers are isomorphic to $\mathcal{C} \setminus P$. In particular, if \mathcal{C} is smooth, this construction defines a smoothing of \mathcal{C}_S . Then Pinkham establishes the following result.

Theorem 3.2 ([16, Theorem 13.9]). *Let $\mathcal{X}^- \rightarrow \mathcal{T}^-$ be the equivariant miniversal deformation in negative degrees of the monomial curve \mathcal{C}_S for a given semigroup S and denote by \mathcal{U}^- the open subset of \mathcal{T}^- given by the points with smooth fibers. Then the moduli space $\mathcal{M}_{g,1}^S$ is isomorphic to the quotient $\mathcal{M}_{g,1}^S = (\mathcal{U}^-)/\mathbb{G}_m$ of \mathcal{U}^- by the \mathbb{G}_m -action.*

The remainder of this subsection is devoted to explicitly describing Pinkham’s Theorem [16, Theorem 13.9] in the case where S is assumed to be a non-hyperelliptic symmetric semigroup. We present the construction initiated by Stöhr [20], and subsequently developed by Contiero–Stöhr [6] and Contiero–Fontes [4]. This construction can be viewed as a variant of Hauser’s algorithm for computing the versal deformation space of a singularity; see [8, 9] and [18].

We start by fixing a numerical symmetric semigroup $S = \langle n_0, n_1, \dots, n_{g-1} \rangle$ of genus $g > 3$ satisfying

$$3 < n_1 < g \quad \text{and} \quad S \neq \langle 4, 5 \rangle.$$

These restrictions are also imposed to avoid simple and well-known cases. If $n_1 = 3$ or $S = \langle 3, 4 \rangle$, then \mathcal{C}_S is a plane curve. Additionally, if $n_1 = g$, then S is not a complete intersection, see [2], and the associated moduli is studied in [4].

So, if \mathcal{C} is a Gorenstein curve with a smooth point whose associated Weierstrass semigroup is S , then \mathcal{C} can be identified with this image under the canonical embedding in such a way that the Weierstrass point P that realizes S is the point $P = (0 : \dots : 0 : 1)$. Hence, by Section 3.2, the canonical ideal $I(\mathcal{C}) \subset \mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}]$ is generated by the $\frac{1}{2}(g - 2)(g - 3)$ quadratic forms

$$F_{si} = X_{a_{si}} X_{b_{si}} - X_{a_s} X_{b_s} - \sum_{n=0}^{s-1} c_{sin} X_{a_n} X_{b_n} \in \mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}],$$

where c_{sij} are suitable constants in \mathbf{k} , the forms $F_{si}^{(0)} = X_{a_{si}} X_{b_{si}} - X_{a_s} X_{b_s}$ generate the ideal of the canonical monomial curve $\mathcal{C}^{(0)} \subset \mathbb{P}^{g-1}$ defined in (3.6), and each $X_{a_j} X_{b_j}$ belongs to the fixed base Δ_2 in equation (3.3).

Now let us invert the considerations on the previous paragraph. Let

$$F_{si}^{(0)} = X_{a_{si}} X_{b_{si}} - X_{a_s} X_{b_s}$$

be the defining polynomials of the canonical curve $\mathcal{C}^{(0)}$ as in (3.7). Now let us take their unfolding

$$F_{si} = F_{si}^{(0)} - \sum_{n=0}^{s-1} c_{sin} X_{a_n} X_{b_n} \in \mathbf{k}[\{c_{sij}\}] \otimes \mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}],$$

defined in Section 3.3, equation (3.10). We want to determine the constants c_{sin} in order that the intersection of the $V(F_{si})$ in \mathbb{P}^{g-1} is a canonical Gorenstein curve of genus g whose Weierstrass semigroup at the smooth point P is S .

Since the coordinates functions x_n introduced in Section 3.1, where $n \in S$ and $n \leq 2g - 2$, are not uniquely determined by their pole divisor nP , we may transform

$$X_{n_i} \mapsto X_{n_i} + \sum_{j=0}^{i-1} \alpha_{ij} X_{n_{i-j}},$$

for each $i = 1, \dots, g - 1$, and so we can normalize $\frac{1}{2}g(g - 1)$ of the coefficients c_{sin} to be zero, see [20, Proposition 3.1]. Due to these normalizations and the normalizations of the coefficients $c_{sin} = 1$ with $n = s$, the only freedom left to us is to transform $x_{n_i} \mapsto \alpha^{n_i} x_{n_i}$ for $i = 1, \dots, g - 1$.

The first step to the explicit construction of a compactification of $\mathcal{M}_{g,1}^S$ due to Contiero–Stöhr is the following lemma.

Lemma 3.3 ([6, Syzygy lemma]). *For each of the $\frac{1}{2}(g - 2)(g - 5)$ quadratic binomials $F_{s'i'}^{(0)}$ different from $F_{n_i+2g-2,1}^{(0)}$, $i = 0, \dots, g - 3$, there is a syzygy of the form*

$$X_{2g-2} F_{s'i'}^{(0)} + \sum_{nsi} \varepsilon_{nsi}^{(s'i')} X_n F_{si}^{(0)} = 0 \tag{3.12}$$

where the coefficients $\varepsilon_{nsi}^{s'i'}$ are integers equal to 1, -1 or 0 and where the sum is taken over the nongaps $n \leq 2g - 2$ and the double indices si with $n + s = 2g - 2 + s$.

The algorithmic construction of the closure of $\mathcal{M}_{g,1}^S$ starts by replacing the initial binomials $F_{s'i'}^{(0)}$ and $F_{si}^{(0)}$ in equation (3.12) by the corresponding unfolded forms $F_{s'i'}$ and F_{si} displayed in equation (3.10) of Section 3.3, obtaining a linear combination of cubic monomials of weight smaller than $s' + 2g - 2$. By virtue of [6, Lemma 2.4] and its proof, this linear combination of cubic monomials admits the following decomposition:

$$X_{2g-2} F_{s'i'} + \sum_{nsi} \varepsilon_{nsi}^{(s'i')} X_n F_{si} = \sum_{nsi} \eta_{nsi}^{(s'i')} X_n F_{si} + R_{s'i'},$$

where the sum on the right hand side is taken over the nongaps $n \leq 2g - 2$ and the double indexes si with $n + s < s' + 2g - 2$, the coefficients $\eta_{nsi}^{(s'i')}$ are constants, and where $R_{s'i'}$ is a linear combination of cubic monomials of pairwise different weights smaller than $s' + 2g - 2$.

For each nongap $m < s' + 2g - 2$, let $\varrho_{s'i'm}$ be the unique coefficient of $R_{s'i'}$ of weight m . It is a quasi-homogeneous polynomial expression of weight $s' + 2g - 2 - m$ in the coefficients c_{sin} .

All the objects that are required to construct the compactification of $\mathcal{M}_{g,1}^S$ were introduced above. The main results due to Stöhr and Contiero–Stöhr are the following.

Theorem 3.4 (Cf. [6, Theorem 2.6]). *Let $S \subset \mathbb{N}$ be a numerical symmetric semigroup of genus g satisfying $3 < n_1 < g$ and $S \neq \langle 4, 5 \rangle$. Then the $\frac{1}{2}(g - 2)(g - 3)$ quadratic forms $F_{si} = F_{si}^{(0)} - \sum_{n=0}^{s-1} c_{sin} X_{a_n} X_{b_n}$ cut out a canonical integral Gorenstein curve in \mathbb{P}^{g-1} if and only if the coefficients c_{sin} satisfy the quasi-homogeneous equations $\varrho_{s'i'm} = 0$. In this case, the point $P = (0 : 0 : \dots : 1)$ is a smooth point of the canonical curve with Weierstrass semigroup S .*

Theorem 3.5 (Cf. [6, Theorem 2.7]). *Let $S \subset \mathbb{N}$ a symmetric numerical semigroup of genus $g := \#(\mathbb{N} \setminus S)$ satisfying $3 < n_1 < g$ and $S \neq \langle 4, 5 \rangle$. The isomorphism classes of the pointed complete integral Gorenstein curves with Weierstrass semigroup S correspond bijectively to the orbits of the $\mathbb{G}_m(\mathbf{k})$ -action*

$$(c, \dots, c_{sin}, \dots) \mapsto (\dots, c^{s-n} c_{sin}, \dots)$$

on the affine quasi-cone of the vectors whose coordinates are the coefficients c_{sin} of the $\frac{1}{2}(g - 2)(g - 3)$ normalized quadratic forms F_{si} that satisfy the quasi-homogeneous equations $\varrho_{s'i'm} = 0$.

4. The main theorem

Let S be a non-hyperelliptic symmetric semigroup and $0 = n_0 < n_1 < \dots < n_{g-1}$ its canonical system of generators. Let us also take $a_1 < \dots < a_r$ a minimal system of generators of S . Considering the polynomial rings $\mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}]$ and $\mathbf{k}[X_{a_1}, \dots, X_{a_r}]$, we attach to the variable X_k the degree $\deg(X_k) = k$ and then $\deg(X_k^\alpha) = \alpha \cdot \deg(X_k)$. The map

$$\begin{aligned} \Pi : \mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}] &\longrightarrow \mathbf{k}[X_{a_1}, \dots, X_{a_r}], \\ X_{n_0} &\longmapsto 1, \\ X_{n_i} &\longmapsto \mathbf{X}_{n_i}^\alpha \end{aligned}$$

where $\mathbf{X}_{n_i}^\alpha$ is the monomial in the variables X_{a_i} introduced in the above Section 3.1, is a graded homomorphism between $\mathbf{k}[X_{n_0}, \dots, X_{n_{g-1}}]$ and $\mathbf{k}[X_{a_1}, \dots, X_{a_r}]$. Henceforward, the *shrinking map* stands to this homomorphism Π .

4.1. Proof of the main theorem

Let S be a numerical symmetric non-hyperelliptic semigroup of genus $g > 1$. Then S is realized as the Weierstrass semigroup of the canonical monomial curve $\mathcal{C}^{(0)}$ at the point $P = (0 : \dots : 0 : 1)$. Considering the affine open chart $X_0 = 1$, the parametrization of $\mathcal{C}^{(0)}|_{X_0=1}$ is given by

$$\mathcal{C}^{(0)}|_{X_0=1} = \{(t^{n_1}, t^{n_2}, \dots, t^{n_{g-1}}); t \in \mathbb{A}^1\} \subset \mathbb{A}^{g-1}.$$

On the other hand, let a_1, \dots, a_r be the minimal system of generators of S and consider the affine monomial curve $\mathcal{C}_S = \text{Spec } \mathbf{k}[S]$.

Then $\mathcal{C}_S \simeq \mathcal{C}^{(0)}|_{X_0=1} = \mathcal{C}^{(0)} \setminus P$ because their coordinate rings are both $\mathbf{k}[S]$. Since the projectivization of $\mathcal{C}^{(0)} \setminus P$ and \mathcal{C}_S are obtained by adding a single point $P = (0 : \dots : 0 : 1) \in \mathbb{P}^{g-1}$ and $Q = (0 : \dots : 0 : 1) \in \mathbb{P}^r$, respectively, at infinity, we conclude that the projectivization of this two curves are also isomorphic. Since we do not lose information on the coefficients of the unfolded quadratic forms that generate the monomial curve $\mathcal{C}^{(0)}$, we can adapt Stöhr’s construction to provide a compactification of $\mathcal{M}_{g,1}^S$, using the monomial affine curve \mathcal{C}_S instead of the canonical one $\mathcal{C}^{(0)}$. To do this we shrink all the forms that are involved in Stöhr’s construction, in particular the P -hermitian basis and the unfolded quadratic forms.

Let us fix an algebraic closed field \mathbf{k} of arbitrary characteristic. In order to prove the main theorem, we first shall prove the following theorem.

Theorem 4.1. *Let \mathcal{C} a non-hyperelliptic Gorenstein curve defined over \mathbf{k} and S a complete intersection numerical semigroup. Then \mathcal{C} realizes S at a smooth point $P \in \mathcal{C}$ if and only if there is an embedding of \mathcal{C} into \mathbb{P}^r such that the defining equations of $\mathcal{C} \setminus P$ are given by the unfolding of the $r - 1$ defining equations of $\mathcal{C}_S \subseteq \mathbb{A}^r$.*

Proof. Let \mathcal{C} be a complete integral Gorenstein curve and P be a smooth point on \mathcal{C} whose Weierstrass semigroup is equal to S . Let us take the line bundle $\mathcal{L} = \mathcal{O}_{\mathcal{C}}(P)$ and its associated ring of sections $\mathcal{R} = \bigoplus_{i=0}^{\infty} H^0(\mathcal{C}, \mathcal{L}^i)$. Since we are fixing a minimal system of generators for S , the ring \mathcal{R} induces an embedding of $\mathcal{C} = \mathbb{P}(\mathcal{R})$ in a weighted projective space, with coordinates Y_0, \dots, Y_r with $\deg Y_0 = 0$. The space $\text{Spec } \mathcal{R}$ is the corresponding quasi-cone in an affine space. Setting $Y_0 = 0$ defines the monomial curve \mathcal{C}_S and all other fibers are isomorphic to $\mathcal{C} \setminus P$. In particular, $\mathcal{C} \setminus P$ is obtained by a deformation of \mathcal{C}_S , as predicted by Pinkham’s construction. Since every deformation of \mathcal{C}_S that realizes S at an added point at the infinity is obtained by unfolding the defining equations of \mathcal{C}_S , cf. Theorem 3.4, we are done.

Conversely, let \mathcal{D} be an affine curve in \mathbb{A}^r that is given by the unfold of the regular sequence $G_{kj}^{(0)} \in \mathbf{k}[X_{a_1}, \dots, X_{a_r}]$ that generates the ideal of \mathcal{C}_S , where each $G_{kj}^{(0)}$ is an isobaric polynomial of degree k . So the ideal of \mathcal{D} is given by $G_{kj} = G_{kj}^{(0)} + \sum_i e_i \beta_i$, with $\beta_i \in \Gamma_2$, where Γ_2 is the shrink P -hermitian basis of $H^0(\mathcal{C}^{(0)}, \omega^2)$ fixed in (3.2), and $e_i \in \mathbf{k}$.

Following Stöhr’s construction, a curve is in $\overline{\mathcal{M}_{g,1}^S}$ if and only if it satisfies some quasi-homogeneous equations $Q_{s'i'n} = 0$ that come from suitable syzygies of the generators $F_{si}^{(0)}$ of the affine monomial curve $\mathcal{C}^{(0)}$, cf. Theorem 3.4. Now, the Syzygy lemma 3.12 assures the existence of $\frac{1}{2}(g - 2)(g - 5)$ syzygies of the form

$$S_{s'i'} := X_{2g-2} F_{s'i'}^{(0)} + \sum_{nsi} \varepsilon_{nsi}^{(s'i')} X_n F_{si}^{(0)} = 0.$$

Taking the image of these syzygies under the shrink map Π , we get

$$\Pi(S_{s'i'}) = \sum_{j=1}^{r-1} M_{s'i'j} G_{kj}^{(0)} = 0,$$

where $M_{s'i'j} \in \mathbf{k}[X_{a_1}, \dots, X_{a_r}]$ are isobaric polynomials of weight $s' - k$. Using the Koszul complex, we are able to show that the relations between the generators $G_{ki}^{(0)}$, for $i = 1, \dots, r - 1$, must be trivial, because \mathcal{C}_S is a complete intersection. Thus, when we exchange $G_{kj}^{(0)}$ by the unfold G_{kj} , the relations $Q_{s'i'm} = 0$ between the coefficients given by Theorem 3.4 are trivially satisfied. Then the projectivized unfolds of the forms that generate \mathcal{D} cut out an integral curve in \mathbb{P}^r with Weierstrass semigroup S in $Q(0 : \dots : 0 : 1)$. ■

Proof of the main theorem. By virtue of the above Theorem 4.1, an integral curve is in $\overline{\mathcal{M}}_{g,1}^S$ if and only if it is given by the unfolding of the regular sequence of the complete intersection monomial curve \mathcal{C}_S . Then the space $\overline{\mathcal{M}}_{g,1}^S$ is just determined by the coefficients of the unfolded forms. The coordinate functions x_n , for $n \in S$, were chosen as functions with pole divisors nP , so they are not uniquely determined. Hence we are able to do the following changes of variables:

$$X_n \mapsto X_n + \sum_{m=0}^{n-1} d_{nm} X_m,$$

where the coefficients d_{nm} are constant. As there are r minimal generators in S we can normalize $\frac{1}{2}r(r + 1)$ coefficients with weights determined in the unfolds of the generator polynomials of the complete intersection affine curve, provided the characteristic of the ground field \mathbf{k} is zero or not a prime divisor of any exponent of the defining equations of \mathcal{C}_S . After these normalizations, the only change we can make is to transform $x_{a_i} \mapsto c^{a_i} x_{a_i}$, $i = 1, \dots, r - 1$, for some $c \in \mathbb{G}_m(\mathbf{k}) = \mathbf{k}^*$. According to [20, Appendix] the coefficients of the normalized unfolded polynomials form a basis for the negatively-graded part of the first cohomology module of the cotangent complex $\mathbf{T}^{1,-}(\mathbf{k}[S])$. Hence we conclude that $\overline{\mathcal{M}}_{g,1}^S = \mathbb{P}(\mathbf{T}^{1,-}(\mathbf{k}[S]))$. ■

4.2. Examples

Example 4.1. We start with a simple example in codimension 2. Given a positive integer τ , consider the semigroup

$$S = \langle 4, 3 + 4\tau, 6 + 4\tau \rangle = 4\mathbb{N} \sqcup (3 + 4\tau + 4\mathbb{N}) \sqcup (6 + 4\tau + 4\mathbb{N}) \sqcup (9 + 8\tau + 4\mathbb{N}),$$

of genus $g = 3 + 4\tau$ and whose Frobenius number is $\ell_g = 5 + 8\tau = 2g - 1$, so S is symmetric. Consider the affine monomial curve

$$\mathcal{C}_S := \{(t^4, t^{3+4\tau}, t^{6+4\tau}); t \in \mathbf{k}\} \subset \mathbb{A}^3$$

and $\mathcal{C}^{(0)} \subset \mathbb{P}^{g-1}$ the canonical monomial curve where $P = (0 : \cdots : 0 : 1)$ realizes S . Let $\{x_0, x_{n_1}, \dots, x_{n_{g-1}}\}$ be a basis for $H^0(\mathcal{C}^{(0)}, \mathcal{O}(P))$. For short we use

$$x := x_4, \quad y_3 := x_{3+4\tau} \quad \text{and} \quad y_6 := x_{6+4\tau}.$$

Thus a P -hermitian base of $H^0(\mathcal{C}^{(0)}, \omega) = H^0(\mathcal{C}^{(0)}, \mathcal{O}(4 + 8\tau)P)$ is given by

$$\begin{cases} x^0, x, \dots, x^{2\tau+1}, \\ x^0 y_3, x y_3, \dots, x^\tau y_3, \\ x^0 y_6, x y_6, \dots, x^{\tau-1} y_6, \end{cases} \quad \tau \geq 1.$$

We can consider $y_9 := x_{9+8\tau}$ as the product $y_3 y_6$. Hence the P -hermitian basis for the bicanonical divisor $H^0(\mathcal{C}^{(0)}, \mathcal{O}(8 + 16\tau))$ is given by the $3g - 3$ elements

$$\begin{cases} x^0, \dots, x^{2+4\tau}, \\ x^0 y_3, x y_3, \dots, x^{1+3\tau} y_3, \\ x^0 y_6, x y_6, \dots, x^{3\tau} y_6, \\ x^0 y_3 y_6, x y_3 y_6, \dots, x^{2\tau-1} y_3 y_6. \end{cases}$$

Lifting the P -hermitian basis elements, we attach the variables x, y_3 and y_6 to X, Y_3 and Y_6 of weights $4, 3 + 4\tau$ and $6 + \tau$, respectively. For short we use

$$Z_{4i} := X^i, \quad Z_{j+4\tau+4i} := X^i Y_j, \quad Z_{9+8\tau+4i} := X^i Y_3 Y_6.$$

As the curve \mathcal{D} is a complete intersection, there are two polynomials in $\mathbf{k}[X, Y_3, Y_6]$ that vanish in \mathcal{D} and generate its ideal. They are

$$G_1 = Y_3^2 - Y_6 X^\tau \quad \text{and} \quad G_2 := Y_6^2 - X^{3+2\tau}.$$

The unfolds of the above polynomials are

$$\tilde{G}_1 = Y_3^2 - Y_6 X^\tau - \sum_{j=1}^{6+8\tau} a_j Z_{6+8\tau-j} \quad \text{and} \quad \tilde{G}_2 := Y_6^2 - X^{3+2\tau} - \sum_{j=k}^{12+8\tau} b_k Z_{12+8\tau-k},$$

where the sums vary between the positive integers j and k such that $6 + 8\tau - j \in S$ and $12 + 8\tau - k \in S$. Doing the variable changes of the form

$$\begin{aligned} X &\mapsto X + \alpha_4, \\ Y_3 &\mapsto Y_3 + \beta_{3+4(\tau-1)} X + \beta_{3+4\tau}, \\ Y_6 &\mapsto Y_6 + \gamma_3 Y_3 + \gamma_{2+4\tau} X + \gamma_{6+4\tau} \end{aligned}$$

we can normalize 6 coefficients of the unfolded forms to zero, provided that the characteristic of \mathbf{k} is zero or an odd prime that not divides τ . The unfold of the polynomials G_1 and G_2 have $3 + 4\tau$ and $9 + 3\tau$ coefficients, respectively. Then the parameter space depends on $6 + 7\tau$ coefficients, i.e.,

$$\overline{\mathcal{M}_{g,1}^S} \simeq \mathbb{P}^{5+7\tau}.$$

In the particular case $\tau = 1$, we have $S = \langle 4, 7, 10 \rangle$ and $g = 7$. The canonical ideal of the monomial curve $\mathcal{C}^{(0)}$ is generated by 10 quadratic forms, namely

$$\begin{aligned} F_{8,1}^{(0)} &= X_4^2 - X_0X_8, & F_{11,1}^{(0)} &= X_4X_7 - X_0X_{11}, & F_{12,1}^{(0)} &= X_4X_8 - X_0X_{12}, \\ F_{14,1}^{(0)} &= X_7^2 - X_4X_{10}, & F_{15,1}^{(0)} &= X_7X_8 - X_4X_{11}, & F_{16,1}^{(0)} &= X_8^2 - X_4X_{12}, \\ F_{18,1}^{(0)} &= X_8X_{10} - X_7X_{11}, & F_{19,1}^{(0)} &= X_8X_{11} - X_7X_{12}, & F_{20,1}^{(0)} &= X_{10}^2 - X_8X_{12}, \\ & & F_{22,1}^{(0)} &= X_{11}^2 - X_{10}X_{12}. \end{aligned}$$

The only syzygy coming from the Syzygy lemma 3.3 is

$$X_{12}F_{14,1}^{(0)} - X_{10}F_{16,1}^{(0)} + X_7F_{19,1}^{(0)} - X_8F_{18,1}^{(0)} = 0.$$

Applying the shrinking map, i.e., considering this syzygy in $\mathbf{k}[X_4, X_7, X_{10}]$, we have the trivial syzygy

$$X_4^3(X_7^2 - X_4X_{10}) - X_4^3(X_7^2 - X_4X_{10}) = X_4^3F_{14,1}^{(0)} - X_4^3F_{14,1}^{(0)} = 0.$$

Thus, as there are no non-trivial syzygies, the space of parameters depends on 13 coefficients, and

$$\overline{\mathcal{M}_{g,1}^S} \simeq \mathbb{P}^{12}.$$

Example 4.2. Let us now consider an example in codimension 4. For each $\tau > 0$, consider the semigroup $S = \langle 16, 1 + 16\tau, 2 + 16\tau, 4 + 16\tau, 8 + 16\tau \rangle$, whose genus is 32τ and Frobenius number $\ell_g = 64\tau - 1$. The affine monomial curve

$$\mathcal{D} = \{(t^{16}, t^{1+16\tau}, t^{2+16\tau}, t^{4+16\tau}, t^{8+16\tau}); t \in \mathbb{k}\} \subset \mathbb{A}^5$$

is a complete intersection in \mathbb{A}^4 , its ideal is generated by

$$G_1 = Y_1^2 - Y_2X^\tau, \quad G_2 = Y_2^2 - Y_4X^\tau, \quad G_3 = Y_4^2 - Y_8X^\tau, \quad G_4 = Y_8^2 - X^{3\tau},$$

where $X := X_{16}$, $Y_1 := Y_{1+16\tau}$, $Y_2 := Y_{2+16\tau}$, $Y_4 := Y_{4+16\tau}$, and $Y_8 := Y_{8+16\tau}$. Unfolding the defining polynomials of \mathcal{D} we get

$$\begin{aligned} \tilde{G}_1 &= Y_1^2 - Y_2X^\tau - \sum_{j=1}^{2+32\tau} a_j Z_{2+32\tau-j}, & \tilde{G}_2 &= Y_2^2 - Y_4X^\tau - \sum_{j=k}^{4+32\tau} b_k Z_{4+32\tau-k}, \\ \tilde{G}_3 &= Y_4^2 - Y_8X^\tau - \sum_{j=u}^{8+32\tau} c_u Z_{8+32\tau-u}, & \tilde{G}_4 &= Y_8^2 - X^{3\tau} - \sum_{j=v}^{16+32\tau} d_v Z_{16+32\tau-v}, \end{aligned}$$

with

$$\begin{aligned}
 Z_{16i} &:= X^i, & Z_{1+16\tau+8i} &= X^i Y_1, \\
 Z_{2+16\tau+8i} &= X^i Y_2, & Z_{3+32\tau+8i} &= X^i Y_1 Y_2, \\
 Z_{4+16\tau+8i} &= X^i Y_4, & Z_{5+32\tau+8i} &= X^i Y_1 Y_4, \\
 Z_{6+32\tau+8i} &= X^i Y_2 Y_4, & Z_{7+48\tau+8i} &= X^i Y_1 Y_2 Y_4, \\
 Z_{8+16\tau+8i} &= X^i Y_8, & Z_{9+32\tau+8i} &= X^i Y_1 Y_8, \\
 Z_{10+32\tau+8i} &= X^i Y_2 Y_8, & Z_{11+48\tau+8i} &= X^i Y_1 Y_2 Y_8, \\
 Z_{12+32\tau+8i} &= X^i Y_4 Y_8, & Z_{13+48\tau+8i} &= X^i Y_1 Y_4 Y_8, \\
 Z_{14+48\tau+8i} &= X^i Y_2 Y_4 Y_8, & Z_{15+64\tau+8i} &= X^i Y_1 Y_2 Y_4 Y_8.
 \end{aligned}$$

We can normalize 15 coefficients from the unfolding polynomials using

$$\begin{aligned}
 X &\longmapsto X + \alpha_{16}, \\
 Y_1 &\longmapsto Y_1 + \beta_{-15+16\tau} X + \beta_{1+16\tau}, \\
 Y_2 &\longmapsto Y_2 + \gamma_1 Y_1 + \gamma_{-14+16\tau} X + \gamma_{2+16\tau}, \\
 Y_4 &\longmapsto Y_4 + \theta_2 Y_2 + d_3 Y_1 + \theta_{-12+16\tau} X + \theta_{4+16\tau}, \\
 Y_8 &\longmapsto Y_8 + \mu_4 Y_4 + \mu_6 Y_2 + \mu_7 Y_1 + \mu_{-8+16\tau} X + \mu_{8+16\tau}.
 \end{aligned}$$

Hence, counting coefficients we can conclude that $\overline{\mathcal{M}_{g,1}^S} \simeq \mathbb{P}^{8+24\tau}$.

The GAP System’s semigroup package simplifies finding complete intersection numerical semigroups, like $S = \langle 32, 33, 34, 36, 40, 48 \rangle$ of genus $g = 80$. Following the procedure presented here, verifying $\overline{\mathcal{M}_{g,1}^S} \simeq \mathbb{P}^{53}$ becomes straightforward. For any such semigroup S , a family like $S = \langle 32, 1 + 32\tau, 2 + 32\tau, 4 + 32\tau, 8 + 32\tau, 16 + 32\tau \rangle$ ($\tau \geq 1$) can be considered. Our procedure readily adapts to any family member, as shown in Examples 4.1 and 4.2.

Acknowledgments. We would like to thank the anonymous reviewer for valuable comments that improved the quality of the paper.

Funding. During the preparation of this paper the first author was partially supported by ICTP-INdAM Research in Pairs Programme, Trieste, Italy and by Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG), Brazil, grant no. APQ-00798-18. The second author was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001.

References

- [1] M. Artin, *Lectures on deformations of singularities, Notes by C. S. Seshadri*, Allen Tannenbaum., Lect. Math. Phys. 54, Tata Institute of Fundamental Research, Bombay, 1976 Zbl [0395.14003](#)
- [2] I. Bermejo, I. García-Marco, and J. J. Salazar-González, [An algorithm for checking whether the toric ideal of an affine monomial curve is a complete intersection](#). *J. Symbolic Comput.* **42** (2007), no. 10, 971–991 Zbl [1147.14026](#) MR [2361674](#)
- [3] R.-O. Buchweitz, *Über Deformationen monomialer Kurvensingularitäten und Weierstrasspunkte auf Riemannschen flächen*, Ph.D. thesis, Hannover, 1976
- [4] A. Contiero and A. Fontes, [The locus of curves with an odd subcanonical point](#). *Doc. Math.* **28** (2023), no. 5, 1027–1052 Zbl [07787210](#) MR [4705605](#)
- [5] A. Contiero, A. L. Fontes, J. Stevens, and J. Q. Vargas, [On nonnegatively graded Weierstrass points](#). 2021, arXiv:[2111.07721v1](#)
- [6] A. Contiero and K.-O. Stöhr, [Upper bounds for the dimension of moduli spaces of curves with symmetric Weierstrass semigroups](#). *J. Lond. Math. Soc. (2)* **88** (2013), no. 2, 580–598 Zbl [1288.14016](#) MR [3106737](#)
- [7] T. de Jong and D. van Straten, [Deformations of the normalization of hypersurfaces](#). *Math. Ann.* **288** (1990), no. 3, 527–547 Zbl [0715.32013](#) MR [1079877](#)
- [8] H. Hauser, [An algorithm of construction of the semiuniversal deformation of an isolated singularity](#). In *Singularities, Part 1 (Arcata, Calif., 1981)*, pp. 567–573, Proc. Sympos. Pure Math. 40, American Mathematical Society, Providence, RI, 1983 Zbl [0523.58012](#) MR [0713093](#)
- [9] H. Hauser, [La construction de la déformation semi-universelle d'un germe de variété analytique complexe](#). *Ann. Sci. École Norm. Sup. (4)* **18** (1985), no. 1, 1–56 Zbl [0583.32052](#) MR [0803194](#)
- [10] J. Herzog, [Generators and relations of abelian semigroups and semigroup rings](#). *Manuscripta Math.* **3** (1970), 175–193 Zbl [0211.33801](#) MR [0269762](#)
- [11] S. Lichtenbaum and M. Schlessinger, [The cotangent complex of a morphism](#). *Trans. Amer. Math. Soc.* **128** (1967), 41–70 Zbl [0156.27201](#) MR [0209339](#)
- [12] E. Looijenga, [The smoothing components of a triangle singularity. II](#). *Math. Ann.* **269** (1984), no. 3, 357–387 Zbl [0568.14003](#) MR [0761312](#)
- [13] S. Mazzini, *Intersecções completas e pontos de Weierstrass*. Ph.D. thesis, Universidade Federal de Minas Gerais, Belo Horizonte, 2022
- [14] T. Nakano, [On the moduli space of pointed algebraic curves of low genus. II. Rationality](#). *Tokyo J. Math.* **31** (2008), no. 1, 147–160 Zbl [1145.14025](#) MR [2426799](#)
- [15] G. Oliveira, [Weierstrass semigroups and the canonical ideal of nontrigonal curves](#). *Manuscripta Math.* **71** (1991), no. 4, 431–450 Zbl [0742.14029](#) MR [1104994](#)
- [16] H. C. Pinkham, *Deformations of algebraic varieties with G_m action*. Astérisque 20, Société Mathématique de France, Paris, 1974 Zbl [0304.14006](#) MR [0376672](#)
- [17] M. Schlessinger, *Infinitesimal deformations of singularities*. Ph.D. Thesis, Harvard Univ., Cambridge, Mass., 1964

- [18] J. Stevens, [Computing versal deformations of singularities with Hauser's algorithm](#). In *Deformations of surface singularities*, pp. 203–228, Bolyai Soc. Math. Stud. 23, János Bolyai Math. Soc., Budapest, 2013 Zbl [1315.32001](#) MR [3203579](#)
- [19] J. Stevens, [The dimension of the moduli space of pointed algebraic curves of low genus](#). *Comm. Algebra* **51** (2023), no. 7, 2963–2977 Zbl [1520.14068](#) MR [4571079](#)
- [20] K.-O. Stöhr, [On the moduli spaces of Gorenstein curves with symmetric Weierstrass semi-groups](#). *J. Reine Angew. Math.* **441** (1993), 189–213 Zbl [0771.14009](#) MR [1228616](#)

Received 4 February 2024.

André Contiero

Departamento de Matemática, Universidade Federal de Minas Gerais, Av. Antônio Carlos, 6627, 31270-901 Belo Horizonte, Brazil; contiero@ufmg.br

Sarah Mazzini

Instituto de Matemática e Estatística, Universidade Federal de Uberlândia, Avenida João Naves de Ávila, 2121, 38408-100 Uberlândia, Brazil; sarahmazzini@ufu.br