

Universal behavior of the BCS energy gap

Joscha Henheik and Asbjørn Bækgaard Lauritsen

Abstract. We consider the BCS energy gap $\Xi(T)$ (essentially given by $\Xi(T) \approx \Delta(T, \sqrt{\mu})$, the BCS order parameter) at all temperatures $0 \leq T \leq T_c$ up to the critical one, T_c , and show that, in the limit of weak coupling, the ratio $\Xi(T)/T_c$ is given by a universal function of the relative temperature T/T_c . On the one hand, this recovers a recent result by Langmann and Triola [Phys. Rev. B 108 (2023), no. 10, article no. 104503] on three-dimensional s -wave superconductors for temperatures bounded uniformly away from T_c . On the other hand, our result lifts these restrictions, as we consider arbitrary spatial dimensions $d \in \{1, 2, 3\}$, discuss superconductors with non-zero angular momentum (primarily in two dimensions), and treat the perhaps physically most interesting (due to the occurrence of the superconducting phase transition) regime of temperatures close to T_c .

1. Introduction

The Bardeen–Cooper–Schrieffer (BCS) [1] theory of superconductivity exhibits many interesting features. Notably, it predicts, for s -wave superconductors (i.e., those for which the gap function has angular momentum $\ell = 0$, i.e., is radially symmetric), that the superconducting energy gap Ξ (essentially given by $\Xi \approx \Delta(\sqrt{\mu})$, the BCS order parameter) is proportional to the critical temperature T_c with a universal proportionality constant independent of the microscopic details of the electronic interactions, i.e., the specific superconductor. At zero temperature, the claimed universality is the (approximate) formula $\Xi/T_c \approx \pi e^{-\gamma} \approx 1.76$ with $\gamma \approx 0.57$ the Euler–Mascheroni constant, a property which is well known in the physics literature [1, 37]. More recently, based on the variational formulation of BCS theory, first introduced by Leggett [33], and later developed on mostly by Hainzl and Seiringer with others [13, 20, 23, 25], it has been put on rigorous grounds in various (physically quite different) limiting regimes [13, 21, 22, 26–28, 32] (see Section 1.2.1 for details). The general picture in all these works is that the universal behavior appears in a limit where “superconductivity is weak,” meaning that T_c is small.

Mathematics Subject Classification 2020: 81Q10 (primary); 46N50, 82D55 (secondary).

Keywords: BCS theory, Ginzburg–Landau theory, energy gap, universality.

The predicted universality at *positive* temperature is notably less studied. It is expected that the ratio $\Xi(T)/T_c$ is given by some universal function of the relative temperature T/T_c , see [30,34] and Figure 1. For three-dimensional superconductors,¹ this has recently been shown in [30] (building on ideas of [31]) for temperatures uniformly in an interval $[0, (1 - \varepsilon)T_c]$ for any $\varepsilon > 0$ in an appropriate limit where T_c is small. Perhaps, the most interesting regime of temperatures, however, are those close to the critical temperature, due to the phase transition occurring there. For such temperatures, one expects² the behavior [39, eq. (3.54)]

$$\frac{\Xi(T)}{T_c} \approx C_{\text{univ}} \sqrt{\frac{1 - T}{T_c}}, \quad C_{\text{univ}} = \sqrt{\frac{8\pi^2}{7\zeta(3)}} \approx 3.06. \tag{1.1}$$

Notably, the critical exponent $1/2$ (i.e., the order parameter $\Delta(\sqrt{\mu}) \approx \Xi$ vanishing as a square root) agrees with the prediction from the phenomenological Landau theory [29] for second order phase transitions (not to be confused with the *Ginzburg–Landau* theory of superconductivity [8, 17, 18]) in mean-field systems.

In this paper, we extend the previously shown universality in three important directions. Firstly, we consider all spatial dimensions $d \in \{1, 2, 3\}$. Secondly, we treat the full range of temperatures $0 \leq T \leq T_c$. Thirdly, we extend the result to the case of non-zero angular momentum in two dimensions, in particular proving the formula in (1.1). Interestingly, the case of non-zero angular momentum in two dimensions has the exact same universal behavior as *s*-wave superconductors in any dimensions: independently of the angular momentum, we find the same universal function describing the ratio $\Xi(T)/T_c$. This is substantially different from the three-dimensional case, where one still expects some sort of universal behavior to occur, only the universal function strongly depends on the angular momentum, see, e.g., [38] and Remark 2.15 below.

One of the central ideas in the analysis of temperatures close to the critical temperature is the use of *Ginzburg–Landau* (GL) theory. In the physics literature, it is well known that, for temperatures close to the critical one, BCS theory is well approximated by GL theory [18]. This correspondence has been studied, and put on rigorous grounds, quite recently in the mathematical physics literature [10, 11, 14, 16]. See Section 1.2.2 for more details.

¹In [30], only the three-dimensional case is considered explicitly. However, their arguments seem to be easily extendable to handle also the cases of one- and two-dimensional superconductors.

²Historically, the first article suggesting the square root behavior near T_c is by Buckingham [5]. In [1, eq. (3.31)], Bardeen, Cooper, and Schrieffer verified this in their original model, however, with the numerical constant given by 3.2 instead of $C_{\text{univ}} \approx 3.06$.




Figure 1. The ratio of the BCS energy gap and the critical temperature, Ξ/T_c , is well approximated by a *universal function* of the relative temperature T/T_c , which is given by $f_{\text{BCS}}(\sqrt{1 - T/T_c})$ with f_{BCS} defined in (2.9) below. At $T = 0$, it approaches the well-known constant $\pi e^{-\gamma} \approx 1.76$, with $\gamma \approx 0.57$ being the Euler–Mascheroni constant.

1.1. Mathematical formulation of BCS theory

We consider a gas of fermions in \mathbb{R}^d for $d = 1, 2, 3$ at temperature $T > 0$ and chemical potential $\mu > 0$. The interaction is described by a two-body, real-valued and reflection-symmetric potential $V \in L^1(\mathbb{R}^d)$, for which we assume the following.

Assumption 1.1. We have that $V \in L^{p_V}(\mathbb{R}^d)$ for $p_V = 1$ if $d = 1$, $p_V \in (1, \infty)$ if $d = 2$, or $p_V = 3/2$ if $d = 3$.

A BCS state Γ is given by a pair of functions (γ, α) and can be conveniently represented as a 2×2 matrix valued Fourier multiplier on $L^2(\mathbb{R}^d) \oplus L^2(\mathbb{R}^d)$ of the form

$$\hat{\Gamma}(p) = \begin{pmatrix} \hat{\gamma}(p) & \hat{\alpha}(p) \\ \overline{\hat{\alpha}(p)} & 1 - \hat{\gamma}(p) \end{pmatrix} \quad (1.2)$$

for all $p \in \mathbb{R}^d$. Here, $\hat{\gamma}(p)$ denotes the Fourier transform of the one particle density matrix and $\hat{\alpha}(p)$ is the Fourier transform of the Cooper pair wave function. We require reflection symmetry of $\hat{\alpha}$, i.e., $\hat{\alpha}(-p) = \hat{\alpha}(p)$, as well as $0 \leq \hat{\Gamma}(p) \leq 1$ as a matrix. Recall the definition of the BCS free energy functional [20, 33], which is given by

$$\mathcal{F}_T[\Gamma] := \int_{\mathbb{R}^d} (p^2 - \mu) \hat{\gamma}(p) \, dp - TS[\Gamma] + \int_{\mathbb{R}^d} V(x) |\alpha(x)|^2 \, dx, \quad (1.3)$$

where the entropy per unit volume is defined as

$$S[\Gamma] = - \int_{\mathbb{R}^d} \text{Tr}_{\mathbb{C}^2} [\hat{\Gamma}(p) \log \hat{\Gamma}(p)] \, dp.$$

The variational problem associated with the BCS functional is studied on

$$\mathcal{D} := \{ \Gamma \text{ as in (1.2)} : 0 \leq \hat{\Gamma} \leq 1, \hat{\gamma} \in L^1(\mathbb{R}^d, (1 + p^2) \, dp), \alpha \in H^1_{\text{sym}}(\mathbb{R}^d) \}.$$

The following proposition provides the foundation for studying this problem.

Proposition 1.2 ([20], see also [25]). *Under Assumption 1.1 on V , the BCS free energy is bounded below on \mathcal{D} and attains its minimum.*

However, in general, the minimizer is not necessarily unique. This potential non-uniqueness shall not bother us at this stage, but will be of importance later on (see Sections 1.1.1 and 2.3). The Euler–Lagrange equation for α associated with the minimization problem is the celebrated *BCS gap equation*

$$\Delta(p) = - \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} \hat{V}(p - q) \frac{\Delta(q)}{K_T^\Delta(q)} \, dq, \tag{1.4}$$

satisfied by $\Delta(p) = -2(2\pi)^{-d/2} (\hat{V} \star \hat{\alpha})(p)$, where α is the off-diagonal entry of a minimizing $\Gamma \in \mathcal{D}$ of (1.3), see [20, 25]. Here, $\hat{V}(p) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} V(x) e^{-ipx} \, dx$ denotes the Fourier transform of V , and we have introduced the notation

$$K_T^\Delta(p) = \frac{E^\Delta(p)}{\tanh\left(\frac{E_\Delta(p)}{2T}\right)} \quad \text{with } E_\Delta(p) = \sqrt{(p^2 - \mu)^2 + |\Delta(p)|^2}.$$

The gap equation can equivalently be written as

$$(K_T^\Delta + V)\alpha = 0, \tag{1.5}$$

where $K_T^\Delta(p)$ is understood as a multiplication operator in momentum space and $V(x)$ is understood as a multiplication operator in position space. The Euler–Lagrange equation for γ (see [20, 25]) is given by

$$\hat{\gamma}(p) = \frac{1}{2} - \frac{p^2 - \mu}{2K_T^\Delta(p)}. \tag{1.6}$$

Remark 1.3 (Log-divergence). For $T = \Delta = 0$ we have $K_{T=0}^{\Delta=0}(p) = |p^2 - \mu|$. This gives rise to a logarithmic divergence in (1.4). Understanding how to treat this log-divergence was one of the key insights of Langmann and Triola [30].

1.1.1. Critical temperature and energy gap. The system described by \mathcal{F}_T is *superconducting* if and only if any minimizer Γ of \mathcal{F}_T has off-diagonal entry $\alpha = \Gamma_{12} \neq 0$ (or, equivalently, (1.4) has a solution $\Delta \neq 0$). The question, whether a system is superconducting or not can be reduced to a linear criterion involving the pseudo-differential operator with symbol

$$K_T(p) \equiv K_T^0(p) = \frac{p^2 - \mu}{\tanh\left(\frac{p^2 - \mu}{2T}\right)}.$$

In fact, as shown in [20], the system is superconducting if and only if the operator $K_T + V$ has at least one negative eigenvalue. Moreover, there exists a unique *critical temperature* $T_c \geq 0$ being defined as

$$T_c := \inf\{T > 0 : K_T + V \geq 0\} \quad (1.7)$$

for which $K_{T_c} + V \geq 0$ and $\inf \text{spec}(K_T + V) < 0$ for all $T < T_c$. By Assumption 1.1 and the asymptotic behavior $K_{T_c}(p) \sim p^2$ for $|p| \rightarrow \infty$, the critical temperature is well defined by Sobolev's inequality [35, Theorem 8.3]. Note that, for $T \geq T_c$, the BCS functional (1.3) is uniquely minimized by the normal state $\Gamma_{\text{FD}} \equiv (\gamma_{\text{FD}}, 0)$, where

$$\hat{\gamma}_{\text{FD}}(p) = \frac{1}{1 + e^{\frac{1}{T}(p^2 - \mu)}} \quad (1.8)$$

is the usual Fermi–Dirac distribution. In contrast, for temperatures $0 \leq T < T_c$ strictly below the critical temperature, the normal state Γ_{FD} is not a minimizer of (1.3) and it is a priori not clear whether or not the minimizer of (1.3) is unique.

In this paper we deal, with two different cases: in the case of *s*-wave superconductivity, we will assume properties of V such that the minimizer is unique; and, in the case of 2-dimensional non-zero angular momentum, we will assume properties of V such that there are at most 2 minimizers, see Section 2.3.

For the *s*-wave case, we assume the following.

Assumption 1.4. Let the (real valued) interaction potential $V \in L^1(\mathbb{R}^d)$ be *radially symmetric* and assume that V is of *negative type*, i.e., $\hat{V} \leq 0$ and $\hat{V}(0) < 0$.

As shown in [21], Assumption 1.4 implies that, in particular, the critical temperature is non-zero, i.e., $T_c > 0$.³ Moreover, as already indicated above, it ensures that the minimizer of (1.3) is unique. While this fact is already known at zero temperature [21, Lemma 2], we are not aware of any place in the literature where the extension to positive temperature is given. As we will need this extension, we formulate it in the following proposition and give a proof in Section A.1.

³To be precise, the arguments in [21] cover only the case $d = 3$, but, as already noted in [14], they are immediately transferable to the cases $d = 1, 2$.

Proposition 1.5 (Uniqueness of minimizers for potentials of negative type). *Let V satisfy Assumptions 1.1 and 1.4, and consider the BCS functional (1.3). Then, we have the following.*

- (i) *For $0 \leq T < T_c$, let $\Gamma \equiv (\gamma, \alpha)$ be a minimizer of the BCS functional (1.3) (which exists by means of Proposition 1.2). Then the operator $K_T^\Delta + V$ from (1.5) is non-negative and α is its unique ground state with eigenvalue zero 0.*
- (ii) *The minimizer $\Gamma =: \Gamma_* \equiv (\gamma_*, \alpha_*)$ of (1.3) is unique up to a phase of α_* and can be chosen to have strictly positive Fourier transform $\hat{\alpha}_*$. Moreover, both γ_* and α_* are radial functions.*

In particular, under Assumption 1.4, we have that the energy gap

$$\boxed{\Xi(T) := \inf_{p \in \mathbb{R}^d} \sqrt{(p^2 - \mu)^2 + |\Delta(p)|^2}} \tag{1.9}$$

(Δ being the (up to multiplication by a constant phase) unique non-zero solution of (1.4) and temperatures $0 \leq T < T_c$) is well defined.

In case there is more than one solution Δ of the BCS gap equation (1.4) (i.e., more than one minimizer of the BCS functional), we may for each such Δ define the energy gap Ξ as in (1.9). In the case of two dimensions with (definite) non-zero angular momentum, we shall prove that there exist exactly two (up to multiplication of either by a constant phase) such functions, Δ_\pm . They however satisfy $|\Delta_+| = |\Delta_-|$, and so the energy gap Ξ is also here uniquely defined. For the details see Section 2.3.

Remark 1.6. The energy gap is essentially the same as the order parameter $|\Delta(\sqrt{\mu})|$, as we show in (3.17) and (3.28) below. In particular, one may replace Ξ with $|\Delta(\sqrt{\mu})|$ in our main results, Theorem 2.1 and Theorems 2.4 and 2.11.

1.1.2. Weak coupling. We consider here the weak-coupling limit, where the interaction is of the form λV for a $\lambda > 0$, and we consider the limit $\lambda \rightarrow 0$. In the weak-coupling limit, an important role is played by the (rescaled) operator $\mathcal{V}_\mu: L^2(\mathbb{S}^{d-1}) \rightarrow L^2(\mathbb{S}^{d-1})$ [7, 21, 24, 28]. This operator, which is defined as

$$(\mathcal{V}_\mu u)(p) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{S}^{d-1}} \hat{V}(\sqrt{\mu}(p - q))u(q) \, d\omega(q), \tag{1.10}$$

where $d\omega$ denotes the uniform (Lebesgue) measure on the unit sphere \mathbb{S}^{d-1} , measures the strength of the interaction potential \hat{V} on the Fermi surface. The pointwise evaluation of \hat{V} (and in particular on a codim-1 submanifold) is well defined, since we assume that $V \in L^1(\mathbb{R}^d)$.

The lowest eigenvalue $e_\mu = \inf \text{spec } \mathcal{V}_\mu$ is of particular importance. Note that \mathcal{V}_μ is a trace-class operator (see the argument above [13, eq. (3.2)]) with

$$\text{tr}(\mathcal{V}_\mu) = \frac{|\mathbb{S}^{d-1}|}{(2\pi)^d} \int_{\mathbb{R}^d} V(x) dx = \frac{|\mathbb{S}^{d-1}|}{(2\pi)^{d/2}} \widehat{V}(0).$$

For radial potentials, one sees that the eigenfunctions of \mathcal{V}_μ are the spherical harmonics.

For potentials of negative type, we have $\widehat{V}(0) < 0$, and so $e_\mu < 0$. This corresponds to an attractive interaction between (some) electrons on the Fermi sphere. Further, one easily sees that the constant function $u(p) = (|\mathbb{S}^{d-1}|)^{-1/2}$ is an eigenfunction of \mathcal{V}_μ , which, since $\widehat{V} \leq 0$ by Assumption 1.4, is in fact the ground state by the Perron–Frobenius theorem, i.e.,

$$e_\mu = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{S}^{d-1}} \widehat{V}(\sqrt{\mu} - q\sqrt{\mu}) d\omega(q). \quad (1.11)$$

In two dimensions, the spherical harmonics take the form

$$u_{\pm\ell}(p) = (2\pi)^{-1/2} e^{\pm i\ell\varphi},$$

with φ denoting the angle of $p \in \mathbb{R}^2$ in polar coordinates. In this case, the ground state space of \mathcal{V}_μ is spanned by $\{u_{\pm\ell}\}_{\ell \in \mathcal{L}}$ for some set of angular momenta \mathcal{L} . If $\ell_0 \neq 0$ for some $\ell_0 \in \mathcal{L}$, then the ground state is at least twice degenerate, since then both $u_{\pm\ell_0}$ are eigenfunctions with this lowest eigenvalue.

1.2. Previous mathematical results

So far, all mathematical results on solutions of the BCS gap equation (1.4) focused either on zero temperature, $T = 0$, or the regime close to the critical one, $T \approx T_c$, where the transition from superconducting to normal behavior is described by Ginzburg–Landau theory.

1.2.1. BCS theory in limiting regimes: Universality at $T = 0$. At zero temperature it is expected that the ratio of the energy gap and the critical temperature is given by a universal constant,

$$\frac{\Xi(T = 0)}{T_c} \approx \pi e^{-\gamma}, \quad (1.12)$$

with $\gamma \approx 0.577$ the Euler–Mascheroni constant in a limiting regime where “superconductivity is weak,” meaning that T_c is small.

In the literature, three such limits have been studied. Historically, the first regime which has been considered is the *weak coupling limit* in three spatial dimensions [13, 21], which we recently extended to one and two dimensions in [28]. The critical temperature in the *low density limit* in three dimensions was studied in [22], and later complemented by a study of the energy gap by one of us in [32], thus, in combination, yielding the above-mentioned universal behavior. Finally, we considered the *high density limit*, again in three dimension, in [26, 27] and proved (1.12) in this regime.

1.2.2. Superconductors close to T_c : Ginzburg–Landau theory. For temperatures close to the critical one, BCS theory is well approximated by Ginzburg–Landau (GL) theory. In contrast to the microscopic BCS model, GL theory is a phenomenological model, which describes the superconductor on a macroscopic scale. Moreover, as suggested by (1.1), a natural parameter measuring “closeness to T_c ” is the parameter $h = \sqrt{1 - T/T_c}$. A rigorous analysis of various aspects of BCS theory in the limit $h \rightarrow 0$ was then studied in [14–16], very recently also allowing for general external fields [10, 11]. Of particular interest to us it is the fact that any minimizer of the BCS functional (γ, α) has $\alpha \approx h\psi\alpha_0$, with $\alpha_0 \in \ker(K_{T_c} + \lambda V)$ fixed and $\psi \in \mathbb{C}$ a minimizer of the corresponding GL functional, see [16, Theorem 2.10].

1.3. Outline of the paper

The rest of this paper is structured as follows. In Section 2 we present our main result, starting with the prototypical universality in the original BCS model (Section 2.1). Afterwards, in Sections 2.2 and 2.3 we describe our results on universality for *s*-wave superconductors in arbitrary dimension $d \in \{1, 2, 3\}$, and for two-dimensional superconductors having pure angular momentum, respectively. The proofs of these results are given in Section 3, while several additional proofs are deferred to Appendix A.

2. Main Result

We next describe the main results of the paper. We first consider the example of an interaction as considered by BCS [1]. The reason for doing this is twofold:

- (1) it highlights how the universal function $\Xi(h)/T_c \approx f_{\text{BCS}}(h)$ appears;
- (2) a central idea in the proof of removing the log-divergence is already present in the BCS gap equation (1.4). (Recall Theorem 1.3.)

2.1. Energy gap in the original BCS approximation

In their seminal work [1], Bardeen, Cooper, and Schrieffer modeled the interaction by a so-called *separable potential* $V(x, y)$ (i.e., factorizing and depending not only on

the relative coordinate $x - y$), whose Fourier transform $\widehat{V}(p, q)$ is a product of two *radial single variable* functions, that are compactly supported in the shell

$$\mathcal{S}_\mu(T_D) := \{p \in \mathbb{R}^d : |p^2 - \mu| \leq T_D\} \quad (2.1)$$

around the Fermi surface $\{p \in \mathbb{R}^d : p^2 = \mu\}$, the only (material dependent) parameter being the so-called *Debye temperature* $0 < T_D < \mu$. Switching from momentum p to energy $\epsilon = p^2 - \mu$, the just-mentioned single variable functions are chosen in such a way that⁴

$$\widehat{V}(\epsilon, \epsilon')N(\epsilon') = -\lambda_{\text{BCS}}\theta\left(1 - \left|\frac{\epsilon}{T_D}\right|\right)\theta\left(1 - \left|\frac{\epsilon'}{T_D}\right|\right), \quad \lambda_{\text{BCS}} > 0, \quad (2.2)$$

where the electronic density of states (DOS) is denoted by $N(\epsilon) \sim (\epsilon + \mu)^{(d-2)/2}$ and θ is the Heaviside function ($\theta(t) = 1$ for $t > 0$ and $\theta(t) = 0$ otherwise).

In this case, the (unique non-negative) solution to the BCS gap equation (1.4) is given by

$$\Delta(\epsilon) = \Delta \cdot \theta\left(1 - \left|\frac{\epsilon}{T_D}\right|\right), \quad (2.3)$$

for some temperature dependent constant $\Delta \geq 0$, which is determined by the scalar gap equation (cf. [1, eq. (3.27)])

$$\frac{1}{\lambda_{\text{BCS}}} = \int_0^{T_D} \frac{\tanh\left(\frac{\sqrt{\epsilon^2 + \Delta^2}}{2T}\right)}{\sqrt{\epsilon^2 + \Delta^2}} d\epsilon, \quad (2.4)$$

for any temperature $0 \leq T < T_c$. In turn, the critical temperature $T_c > 0$ is determined by (2.4) with $\Delta = 0$, i.e.,

$$\frac{1}{\lambda_{\text{BCS}}} = \int_0^{T_D} \frac{\tanh\left(\frac{\epsilon}{2T_c}\right)}{\epsilon} d\epsilon. \quad (2.5)$$

In case of a small BCS coupling parameter, $\lambda_{\text{BCS}} \ll 1$,⁵ it holds that T_c is exponentially small in λ_{BCS} , i.e., $T_c \sim e^{-1/\lambda_{\text{BCS}}}$ (see [1, eq. (3.29)]). Moreover, it is easily checked that Δ as a function of temperature is monotonically decreasing in the interval $[0, T_c]$ and satisfies $\Delta(T = 0) \sim e^{-1/\lambda_{\text{BCS}}}$, similarly to the critical temperature.

⁴Assuming that \widehat{V} is constant throughout the energy shell (2.1) (as done in [1]), the BCS coupling parameter emerges as $\lambda_{\text{BCS}} = -\widehat{V}(0, 0)N(0)$.

⁵This can happen for various reasons. One example is that V itself is scaled by a coupling parameter $\lambda > 0$, i.e., $V \rightarrow \lambda V$, and one considers the limit $\lambda \rightarrow 0$, as done in Sections 2.2–2.3.

Next, changing variables as $x := \epsilon/T_c$ and setting $\delta := \Delta/T_c$ as well as⁶

$$h := \sqrt{1 - \frac{T}{T_c}} \quad \text{for } 0 \leq T \leq T_c, \tag{2.6}$$

we can subtract (2.4) and (2.5) to find

$$\int_0^{T_D/T_c} \left\{ \frac{\tanh\left(\frac{\sqrt{x^2 + \delta^2}}{2(1-h^2)}\right)}{\sqrt{x^2 + \delta^2}} - \frac{\tanh\left(\frac{x}{2}\right)}{x} \right\} dx = 0. \tag{2.7}$$

Note that this difference formula (2.7) removes the divergences of (2.4)–(2.5) as $\lambda_{\text{BCS}} \rightarrow 0$.

The proof of the following proposition is given in Section 3.1. (In the statement of Theorem 2.1, one may replace Ξ by the order parameter $\Delta(\sqrt{\mu})$, see Theorem 1.6 above.)

Proposition 2.1 (Energy gap in the original BCS model [1]). *Let $\mu > 0$, fix a Debye temperature $0 < T_D < \mu$, and let $\lambda_{\text{BCS}} > 0$ be the BCS coupling parameter as above. Let the critical temperature T_c and the gap function $\Delta(p)$ be defined via (2.3)–(2.5).*

Then the energy gap Ξ (defined in (1.9)) as a function of $h = \sqrt{1 - T/T_c}$ for $0 \leq T \leq T_c$ (recall (2.6)) is given by

$$\Xi(h) = T_c f_{\text{BCS}}(h) (1 + O(e^{-1/\lambda_{\text{BCS}}})) \tag{2.8}$$

uniformly in $h \in [0, 1]$, where the function $f_{\text{BCS}}: [0, 1] \rightarrow [0, \infty)$ is implicitly defined via

$$\int_{\mathbb{R}} \left\{ \frac{\tanh\left(\frac{\sqrt{s^2 + f_{\text{BCS}}(h)^2}}{2(1-h^2)}\right)}{\sqrt{s^2 + f_{\text{BCS}}(h)^2}} - \frac{\tanh\left(\frac{s}{2}\right)}{s} \right\} ds = 0 \tag{2.9}$$

and plotted in Figure 2.

This means that, independent of the material dependent Debye temperature $T_D > 0$ and the chemical potential $\mu > 0$, the energy gap Ξ within the original BCS approximation [1], follows a universal curve, described by (2.8), in the limit of weak BCS coupling. A similar formula for f_{BCS} like (2.9) (but as a function of $x := 1 - h^2$) also appeared in the monograph of Leggett [34, eq. (5.5.21)]. We now list a few basic properties of f_{BCS} , whose proofs we omit, as they can be obtained by means of the

⁶As mentioned above, the parameter h is commonly used (see, e.g., [14, 16]) in the context of Ginzburg–Landau theory, where it served as a ‘semiclassical’ small parameter in the derivation this theory.




Figure 2. Sketch of the function f_{BCS} obtained via the implicit relation (2.9).

implicit function theorem and further elementary tools (see also [30, Lemma 1] as well as Lemmas 3.1 and 3.13 below). Almost all of these properties become apparent from Figure 2.

Lemma 2.2 (Properties of f_{BCS}). *There exists a unique implicitly defined solution function $f_{\text{BCS}}: [0, 1] \rightarrow [0, \infty)$ of (2.9). Moreover, f_{BCS} has the following properties:*

- (i) *it is strictly monotonically increasing in $[0, 1]$;*
- (ii) *it is C^1 in $(0, 1)$ and has continuous one-sided derivatives at the boundaries 0 and 1;*
- (iii) *it has the boundary values $f_{\text{BCS}}(0) = 0$, $f'_{\text{BCS}}(0) = C_{\text{univ}}$ and $f_{\text{BCS}}(1) = \pi e^{-\gamma} \approx 1.76$, $f'_{\text{BCS}}(1) = 0$. Here, $\gamma \approx 0.57$ is the Euler–Mascheroni constant and*

$$C_{\text{univ}} := \sqrt{\frac{8\pi^2}{7\zeta(3)}} \approx 3.06, \quad (2.10)$$

where $\zeta(s)$ denotes Riemann's ζ -function.

Remark 2.3 (Contact interactions). Our proof of Proposition 2.1 can easily be generalized to all BCS models, in which the energy gap is constant (at least near the Fermi surface).

(a) In case of a delta potential, $V(x) = -\delta(x)$ in one spatial dimension, $d = 1$, the gap function solving (1.4) is given by a constant (simply because here \hat{V} is constant).

This setting can be analyzed similarly (in a weak coupling limit, i.e., replacing $V \rightarrow \lambda V$ and taking $\lambda \rightarrow 0$) as done in Proposition 2.1 for the original BCS model [1].

(b) Also, for contact interactions in three spatial dimensions, $d = 3$, the situation is similar. This setting is studied in [3,4], where it is shown that for a suitable sequence of potentials V_ℓ converging to a point interaction with scattering length $a < 0$, the gap function Δ_ℓ converges (uniformly on compact sets, see [4, eq. (14)]) to a constant Δ solving the gap equation

$$-\frac{1}{4\pi a} = \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} \left(\frac{1}{K_T^\Delta(p)} - \frac{1}{p^2} \right) dp.$$

Replacing the limit of weak coupling by a small scattering length limit, $a \rightarrow 0$, one can obtain a result similar to Proposition 2.1.

2.2. Universal behavior of the s -wave BCS energy gap

After having discussed the prototypical universality in the seminal BCS paper [1], we can now formulate our main result on general s -wave superconductors with local interactions. The proof of Theorem 2.4 is presented in Sections 3.2–3.4, while the main ideas are briefly described in Remark 2.7 below. (We remark that, in the statement of Theorem 2.4, one may replace Ξ by the order parameter $\Delta(\sqrt{\mu})$, see Remark 1.6 above.)

Theorem 2.4 (BCS energy gap for s -wave superconductors). *Let $d \in \{1, 2, 3\}$, $\mu > 0$, $\lambda > 0$ and let V satisfy Assumptions 1.1 and 1.4. Let T_c and Ξ be as in (1.7) and (1.9) with interaction λV and Δ the unique non-zero solution the BCS gap equation (1.4) with interaction λV .*

Then, with $f_{\text{BCS}}(h)$, $h = \sqrt{1 - T/T_c}$ being the function defined via (2.9), we have the following.

(a) *Assuming additionally that $|\cdot|^2 V \in L^1(\mathbb{R}^d)$, it holds that*

$$\Xi(h) = T_c f_{\text{BCS}}(h) (1 + O(h^{-1} e^{-c/\lambda})) \tag{2.11}$$

for some constant $c > 0$ independent of λ and h .

(b) *Assuming additionally that $(1 + |\cdot|) V \in L^2(\mathbb{R}^d)$, it holds that*

$$\Xi(h) = T_c f_{\text{BCS}}(h) (1 + O(h e^{c'/\lambda}) + o_{\lambda \rightarrow 0}(1)) \tag{2.12}$$

for some constant $c' > 0$ independent of λ and h and where $o_{\lambda \rightarrow 0}(1)$ vanishes as $\lambda \rightarrow 0$ uniformly in h .

For the special case $h = 1$, i.e., $T = 0$, (2.11) reproduces the results from [21] (for $d = 3$) and [28] (for $d = 1, 2$), which state the universality

$$\lim_{\lambda \rightarrow 0} \frac{\Xi(T = 0)}{T_c} = \frac{\pi}{e^\nu}$$

at $T = 0$. Moreover, by (2.11) again, we find that, uniformly in temperatures bounded away from T_c , i.e., $h \in [\varepsilon, 1]$ for some fixed $\varepsilon > 0$,

$$\lim_{\lambda \rightarrow 0} \frac{\Xi(h)}{T_c} = f_{\text{BCS}}(h),$$

recovering the universality result in [30] (for $d = 3$), with an exponential speed $O(e^{-c/\lambda})$ of convergence. In the complementary case, for temperatures very close to the critical temperature, $T \approx T_c$, the question of universality is (i) physically more interesting due to the phase transition from superconducting to normal behavior and (ii) mathematically more delicate than in the previous scenarios. This is because now there are *two* small parameters λ and h , instead of λ only, and the error term in (2.11) might actually be large compared to one. However, now involving both, (2.11) and (2.12), we find that

$$\lim_{\substack{\lambda, h \rightarrow 0 \\ e^{-c/\lambda} \ll h}} \frac{\Xi(h)}{T_c h} = C_{\text{univ}} \quad \text{and} \quad \lim_{\substack{\lambda, h \rightarrow 0 \\ h \ll e^{-c'/\lambda}}} \frac{\Xi(h)}{T_c h} = C_{\text{univ}} \quad (2.13)$$

with the aid of Lemma 2.2 (iii). In particular, the ratio $\Xi(h)/(T_c h)$ converges to the *same* universal constant C_{univ} (recall (2.10)) in both orders of limits, $\lim_{\lambda \rightarrow 0} \lim_{h \rightarrow 0}$ and $\lim_{h \rightarrow 0} \lim_{\lambda \rightarrow 0}$.

Remark 2.5 (Joint limit). A careful inspection of the proof reveals that the constants c and c' satisfy $c < c'$. In particular, the proof does *not* allow the two regimes considered in (2.13) to be overlapping and we cannot prove that $\lim_{\lambda, h \rightarrow 0} \Xi(h)/T_c h = C_{\text{univ}}$ in any joint limit. We expect this to hold in any joint limit, however, as we saw for the particular example from [1] in Theorem 2.1

Remark 2.6 (Comparison of assumptions with [30]). Compared to the similar result in [30] our assumptions hold for a slightly different class of potentials. The assumptions of [30] are essentially on the smoothness of the interaction V (formulated via some regularity/decay assumption on the Fourier transform \hat{V}). Our assumptions, on the other hand, are on the regularity/decay of V . In particular, our assumptions cover the examples of [30, Table I] which are *not* covered by the assumptions of [30]. These are (in three dimensions)

$$V_{\text{Yukawa}}(x) = \frac{e^{-|x|}}{4\pi|x|}, \quad V_{aY+bE}(x) = \frac{(2a + b|x|)e^{-|x|}}{8\pi|x|}, \quad V_{x\text{-box}}(x) = \frac{3\theta(1 - |x|)}{4\pi}.$$

Remark 2.7 (On the proof). The main ideas in the proof of Theorem 2.4 are the following.

(a) For part (a), we crucially use that both $K_T^\Delta + \lambda V$ and $K_{T_c} + \lambda V$ have lowest eigenvalue zero. We then consider their corresponding Birman–Schwinger (BS) operators and use that, for λ small enough, two naturally associated operators on the Fermi sphere both have the *same* ground state. Evaluating the difference of these two associated operators in this common ground state, we find that a *difference* of two logarithmically divergent integrals, similarly to (2.9), vanishes up to exponentially small errors $O(e^{-c/\lambda})$.

The removal of the log-divergence in this way (which – in a similar fashion – was the major insight in [30]) is the key idea to (i) access also non-zero temperatures and (ii) obtain extremely precise error estimates (compared to all the previous results mentioned in Section 1.2.1).

(b) For part (b), we employ Ginzburg–Landau (GL). The principal realm of GL theory is to describe superconductors and superfluids close to their critical temperature T_c . In this regime, when superconductivity is weak, the main idea is that the prime competitor for developing a small off-diagonal component $\hat{\alpha}$ for a BCS minimizer, is the *normal state* $\Gamma_{\text{FD}} = (\gamma_{\text{FD}}, 0)$, with γ_{FD} given by the usual Fermi–Dirac distribution (recall (1.8)). Moreover, to leading order, the off-diagonal component $\hat{\alpha}$ lies in the kernel (which agrees with the ground state space) of the operator $K_{T_c} + \lambda V$.

The main input, which we use, is that every minimizer (γ, α) of the BCS functional has $\alpha \approx h\psi\alpha_0$ with $\alpha_0 \in \ker(K_{T_c} + \lambda V)$ fixed and $\psi \in \mathbb{C}$ minimizing the corresponding GL functional [16, Theorem 2.10]. Taking the convolution of $\hat{\alpha}_0$ with \hat{V} , we find the universal constant (2.10) appearing in $\Xi/(T_c h) \approx C_{\text{univ}}$.

Moreover, the “additional assumptions” in Theorem 2.4 are not quite rigid, meaning that they can be weakened in the following sense.

(a) In case that $|\cdot|^{2\alpha} V \in L^1$ for $0 < \alpha \leq 1$, the error term in (2.11) should instead be $O(h^{-1}e^{-c\alpha/\lambda})$ with the constant c then being independent also of α .

(b) In case that $(1 + |\cdot|)V \in L^p(\mathbb{R}^d)$ for $p < 2$, the factor h in the first error term in (2.12) would not appear raised to the first power but with exponent

$$\begin{aligned} & \frac{3p - 4}{p} \quad \text{for } d = 1, \\ & \frac{3p - 4}{p} - \epsilon \quad \text{for } d = 2, \\ & \frac{4p - 6}{p} \quad \text{for } d = 3. \end{aligned}$$

Remark 2.8 (Other limits). Although, in this paper, we considered only the weak coupling limit, we expect the relation $\Xi(h) \approx T_c f_{\text{BCS}}(h)$ to hold also in other limiting regimes in which “superconductivity is weak,” that is, e.g., the low-⁷ and high-density limit, that were studied in [22, 32] and [26, 27], respectively. This idea is already contained in [30], where the authors considered a “universal” parameter λ in [30, eq. (7)], which can be small for various physical situations.

Remark 2.9 (Non-universality). We recover also the formula [30, eq. (16)]

$$\frac{\Delta(p)}{\Delta(\sqrt{\mu})} = F(p) + O(e^{-c/\lambda}) \quad (2.14)$$

for some function F not depending on the temperature and some constant $c > 0$. The function F depends on the interaction V however. For this reason, (2.14) is called a “non-universal” feature in [30]. The proof of (2.14) is given in Section 3.3.

2.3. The case of pure angular momentum for $d = 2$

We generalize Theorem 2.4 from s -wave superconductors to two-dimensional systems which have a definite (or *pure*) angular momentum $\ell_0 \in \mathbb{N}_0$, which can differ from 0.

Assumption 2.10 (Pure angular momentum). Let $V \in L^1(\mathbb{R}^2)$ be radially symmetric and *attractive on the Fermi sphere*, i.e., the lowest eigenvalue e_μ of \mathcal{V}_μ is strictly negative (recall (1.10)–(1.11)). Moreover, suppose that for all $\lambda > 0$ small enough the lowest eigenvalue of $K_{T_c} + \lambda V$ is at most twice degenerate, i.e., $\dim \ker(K_{T_c} + \lambda V) \in \{1, 2\}$.

Since K_{T_c} commutes with the Laplacian, Assumption 2.10 ensures the ground state of $K_{T_c} + \lambda V$ to have definite angular momentum. More precisely, it holds that

$$\ker(K_{T_c} + \lambda V) = \text{span}\{\rho\} \otimes \mathcal{S}_{\ell_0} \quad (2.15)$$

with

$$\mathcal{S}_{\ell_0} := \text{span}\{e^{\pm i\ell_0\varphi}\} \subset L^2(\mathbb{S}^1)$$

for some $\ell_0 \in \mathbb{N}_0$, where $\rho \in L^2((0, \infty); r dr)$ is a (λ -dependent) radial function.⁸

We can now formulate our main result in the case of pure angular momentum for $d = 2$. (We again remark that, in the statement of Theorem 2.11, one may replace Ξ by the order parameter $|\Delta(\sqrt{\mu})|$, see Theorem 1.6 above.)

⁷For dimensions $d = 1, 2$, the same caveats mentioned in [28, Remark 2.8] apply.

⁸In fact, the angular momentum of the kernel of $K_{T_c} + \lambda V$ must be *even*, i.e., $\ell_0 \in 2\mathbb{N}_0$. This is because BCS theory is formulated for reflection symmetric α , whence $K_{T_c} + \lambda V$ is naturally defined on the space of reflection symmetric functions only.

Theorem 2.11 (BCS energy gap for $2d$ pure angular momentum). *Let $d = 2$, $\mu > 0$, and let V satisfy Assumptions 1.1 and 2.10. Define the critical temperature T_c and the energy gap Ξ as in (1.7) and (1.9), with interaction λV for a $\lambda > 0$ and Δ being any (arbitrary!) non-zero solution of the BCS gap equation (1.4) with interaction λV .*

Then, with $f_{\text{BCS}}(h)$, $h = \sqrt{1 - T/T_c}$ being the function defined via (2.9), we have the following.

- (a) *Assume additionally that $V \in L^2(\mathbb{R}^2)$, $\widehat{V} \in L^r(\mathbb{R}^2)$ for some $1 \leq r < 2$ and that $|\cdot|^2 V \in L^1(\mathbb{R}^d)$. Then there exists $0 \leq \widetilde{T} < T_c$ with $\widetilde{T}/T_c \leq e^{-c/\lambda}$ for some $c > 0$, such that for all temperatures $T \in (\widetilde{T}, T_c)$ it holds that*

$$\Xi(h) = T_c f_{\text{BCS}}(h)(1 + O(h^{-1}e^{-c/\lambda})) \tag{2.16}$$

for some constant $c > 0$ independent of λ and h .

- (b) *Assuming additionally that $(1 + |\cdot|)V \in L^2(\mathbb{R}^d)$, it holds that*

$$\Xi(h) = T_c f_{\text{BCS}}(h)(1 + O(he^{c'/\lambda}) + o_{\lambda \rightarrow 0}(1))$$

for some constant $c' > 0$ independent of λ and h and where $o_{\lambda \rightarrow 0}(1)$ vanishes as $\lambda \rightarrow 0$ uniformly in h .

The proof of Theorem 2.11 is given in Section 3.5.

Remark 2.12 (On the assumptions). The additional assumptions in part (a) here compared to Theorem 2.4 (namely $V \in L^2$ and $\widehat{V} \in L^r$) are those of [9, Theorem 2.1]. The proof of (2.16) centrally uses this result. As discussed in [9, Remark 2.3], these additional assumptions are expected to be of a technical nature.

Remark 2.13 (The temperature \widetilde{T}). The presence of the temperature \widetilde{T} in Theorem 2.11 (a) arises from the first excited eigenvalue of $K_{T_c} + \lambda V$, see [9, Remark 2.2]. As discussed in the proof, the temperatures T_c and \widetilde{T} are given by $T_c = T_c(\ell_0)$ and $\widetilde{T} = T_c(\ell_1)$, the critical temperatures restricted to angular momenta ℓ_0 and ℓ_1 , for some angular momenta $\ell_0 \neq \ell_1$, see also [9, Remark 2.2]. For temperatures $T \in (\widetilde{T}, T_c)$, the BCS minimizer(s) then have angular momentum ℓ_0 , see [9, Theorem 2.1]. For temperatures $T < \widetilde{T}$, however, we do not in general know whether the BCS minimizer(s) have angular momentum ℓ_0 . The proof crucially uses that the minimizer(s) have a definite angular momentum. If we however know, a priori, that the BCS minimizer(s) have angular momentum ℓ_0 for some larger ranger of temperatures (T_1, T_c) , then the formula in (2.16) holds in this larger range of temperatures.

Remark 2.14 (Nodes of the gap function). As already mentioned in Section 1.1.1, we establish during the proof that any solution Δ of the BCS gap equation (1.4) has a radially symmetric absolute value, $|\Delta(p)|$, which is, moreover, independent of the particular solution Δ . In particular, every solution Δ of the BCS gap equation (1.4)

does *not* have nodes on the Fermi surface. This contrasts many examples of d -wave superconductors in the physics literature, where a (necessarily) non-radial interaction V leads to a gap function Δ with nodes on the Fermi surface, see, e.g., [2, 12, 36, 40].

Remark 2.15 (Non-extension to three dimensions). The formula $\Xi(h) \approx T_c f_{\text{BCS}}(h)$ is *not* expected to hold in three dimensions for non-zero angular momentum, see for instance [38, Figure 14.6]. More precisely, we have the following.

(i) For non-zero angular momentum in three dimensions, our method of proving Theorem 2.11 (a) breaks down. In fact, we crucially use that $K_T^\Delta + \lambda V \geq 0$ for $\Delta = -2\lambda \widehat{V}\alpha$ with α a minimizer of the BCS functional. However, as shown in [9, Proposition 2.11], this implies that $|\hat{\alpha}|$ is a radial function. In particular, in three dimensions, α (and therefore also Δ) cannot have a definite non-zero angular momentum.

(ii) Assume that we know *a priori* that a solution of the BCS gap equation (1.4) (in spherical coordinates) satisfies $\Delta(p, \omega) = \Delta_0(p)Y_\ell^m(\omega)$ – at least to leading order.⁹ Here, Y_ℓ^m is the usual L^2 -normalized (complex) spherical harmonic with $\ell \in \mathbb{N}_0$ and $m \in \{-\ell, \dots, \ell\}$. Then, by application of [16, Theorem 2.10], following very similar arguments to Sections 3.4 and 3.5.2, we find that the radial part of the gap function is given by

$$|\Delta_0(\sqrt{\mu})| \approx c_{\ell,m} C_{\text{univ}} h T_c \quad (2.17)$$

on the Fermi sphere $\{p^2 = \mu\}$. Here C_{univ} was defined in (2.10) and we denoted

$$\begin{aligned} c_{\ell,m} &:= \left(\int_{\mathbb{S}^2} |Y_\ell^m(\omega)|^4 d\omega \right)^{-1/2} \\ &= \left(\sum_{L=0}^{2\ell} \frac{(2\ell+1)^2}{4\pi(2L+1)} |\langle \ell, \ell; 0, 0 | L; 0 \rangle|^2 |\langle \ell, \ell; m, m | L; 2m \rangle|^2 \right)^{-1/2} \end{aligned} \quad (2.18)$$

with $\langle \ell_1, \ell_2; m_1, m_2 | L; M \rangle$ being the well-tabulated Clebsch–Gordan coefficients (see, e.g., [6, p. 1046]). Relation (2.18) shows that, in particular, even in the subspace of fixed angular momentum $\ell \neq 0$, the behavior (2.17) is non-universal due to a non-trivial dependence on $m \in \{-\ell, \dots, \ell\}$, as, for example (see [16, eq. (6.8)]),

$$c_{2,0} = \sqrt{\frac{28\pi}{15}} \quad \text{and} \quad c_{2,\pm 1} = c_{2,\pm 2} = \sqrt{\frac{14\pi}{5}}.$$

⁹If one models the interaction V by a rank one projection $V = |\chi\rangle \langle \chi|$ (similarly to (2.2)), instead of a multiplication operator, such a form of Δ can easily be enforced by taking $\chi(p, \omega) = \chi_0(p)Y_\ell^m(\omega)$.

For temperatures $0 \leq T \leq T_c$ and $h := \sqrt{1 - T/T_c}$, we expect (2.17) to generalize to

$$|\Delta_0(\sqrt{\mu})| \approx T_c f_{\text{BCS}}^{(\ell,m)}(h)$$

with $f_{\text{BCS}}^{(\ell,m)}: [0, 1] \rightarrow [0, \infty)$ being implicitly defined via

$$\int_0^\infty ds \int_{\mathbb{S}^2} d\omega \left\{ \frac{\tanh\left(\frac{\sqrt{s^2 + (f_{\text{BCS}}^{(\ell,m)}(h))^2 |Y_\ell^m(\omega)|^2}}{2(1-h^2)}\right)}{\sqrt{s^2 + (f_{\text{BCS}}^{(\ell,m)}(h))^2 |Y_\ell^m(\omega)|^2}} - \frac{\tanh\left(\frac{s}{2}\right)}{s} \right\} |Y_\ell^m(\omega)|^2 = 0, \quad (2.19)$$

similarly to [38, eq. (14.33)]. For $\ell = m = 0$, (2.19) yields that $f_{\text{BCS}}^{(0,0)} = (4\pi)^{1/2} f_{\text{BCS}}$ with f_{BCS} from (2.9) due to the L^2 -normalization of the spherical harmonics (recall $\Delta(p, \omega) = \Delta_0(p) Y_\ell^m(\omega)$).

A detailed analysis of the three-dimensional case with non-zero angular momentum is deferred to future work.

3. Proofs of the main results

This section contains the proofs of our main results formulated in Section 2.

3.1. Proof of Proposition 2.1

For ease of notation, we shall henceforth write λ instead of λ_{BCS} . From the explicit form (2.3), it is clear that $\Xi = \Delta$ and $\delta(h) \equiv \delta = \Delta/T_c$ is determined through (2.7). Hence, the goal is to show that $\delta(h)/f_{\text{BCS}}(h) = 1 + O(e^{-1/\lambda})$ uniformly in $h \in [0, 1]$. The proof of this is conducted in three steps.

3.1.1. A priori bound on δ . We shall prove the following lemma.

Lemma 3.1. *For $\delta = \delta(h)$ defined through (2.7) and $\lambda > 0$ small enough, it holds that*

$$\delta(h) \leq Ch. \quad (3.1)$$

Proof. First, we note that $\delta(h) \leq C$ uniformly for $h \in [0, 1]$. This easily follows from observing that $\delta(h)$ is strictly monotonically increasing (as follows from elementary monotonicity properties of the integrand in (2.7)) and $\delta(1)$ is necessarily bounded.

In order to show (3.1), we employ the implicit function theorem to derive an asymptotic ODE for $\delta(h)$. For this purpose, we now introduce the function (recall-

ing $T_c \sim e^{-1/\lambda}$)

$$G_\lambda: [0, 1] \times [0, \infty) \rightarrow \mathbb{R}, \quad (h, \delta) \mapsto \int_0^{T_D/T_c} \left\{ \frac{\tanh\left(\frac{\sqrt{x^2+\delta^2}}{2(1-h^2)}\right)}{\sqrt{x^2+\delta^2}} - \frac{\tanh\left(\frac{x}{2}\right)}{x} \right\} dx$$

and trivially note that (2.7) is equivalent to $G_\lambda(h, \delta(h)) = 0$. Since G_λ is C^1 (away from the boundary) in δ and h (this easily follows from dominated convergence), we can apply the implicit function theorem to obtain the differential equation

$$\frac{\partial\delta(h)}{\partial h} = \frac{(1-h^2)h}{\delta(h)} \left(\int_0^{T_D/T_c} \frac{1}{\cosh^2\left(\frac{\sqrt{x^2+\delta^2}}{2(1-h^2)}\right)} dx \Big/ \int_0^{T_D/T_c} \frac{g_1\left(\frac{\sqrt{x^2+\delta^2}}{1-h^2}\right)}{\frac{\sqrt{x^2+\delta^2}}{1-h^2}} dx \right), \quad (3.2)$$

where we introduced the auxiliary functions

$$g_0(z) := \frac{\tanh\left(\frac{z}{2}\right)}{z}, \quad g_1(z) := -g'_0(z) = z^{-1}g_0(z) - \frac{1}{2}z^{-1}\frac{1}{\cosh^2\left(\frac{z}{2}\right)}. \quad (3.3)$$

It is elementary to check that the even function $z \mapsto g_1(z)/z$ is (strictly) positive and (strictly) decreasing for $z \in [0, \infty)$. In combination with $\delta(h) \leq C$ and $T_c \sim e^{-1/\lambda}$, one can thus bound the denominator on the right-hand side of (3.2) from below. Together with an upper bound on the integral in the numerator (obtained by using elementary monotonicity properties of the hyperbolic cosine), we find that

$$\begin{aligned} \frac{\partial\delta(h)}{\partial h} &\leq C' \frac{h}{\delta(h)} \left(\int_0^\infty \frac{1}{\cosh^2(x)} dx \Big/ \int_0^C \frac{g_1(\sqrt{x^2+C^2})}{\sqrt{x^2+C^2}} dx \right) \\ &\leq C'' \frac{h}{\delta(h)} \end{aligned} \quad (3.4)$$

for $h > 0$ and $\lambda > 0$ small enough (to ensure $T_D/T_c \geq C$).

Finally, the differential inequality (3.4) can be integrated using the boundary condition $\delta(0) = 0$ to conclude the desired.¹⁰ ■

¹⁰Strictly speaking, this requires to extend the function $\delta(h)$ in $(0, 1)$, obtained via the implicit function theorem for G_λ , to the boundary points 0. In order to do so, note that, for $h \in (0, 1/2)$, (3.2) yields

$$\frac{\partial\delta(h)}{\partial h} \sim \frac{h}{\delta(h)},$$

from which we immediately conclude that $|\partial_h\delta(h)| \leq C$, uniformly in $(0, 1/2)$. Hence, $\delta(h)$ continuously extends to 0. The same is true for its derivative by means of (3.2) again. We remark that by a similar argument, $\delta(h)$ can be extended to 1 as well.

3.1.2. Uniform error estimate. Having Lemma 3.1 as an input, we shall now prove the following.

Lemma 3.2. *For $\delta = \delta(h)$ defined through (2.7), it holds that*

$$\int_{T_D/T_c}^{\infty} \left| \frac{\tanh\left(\frac{\sqrt{x^2+\delta^2}}{2(1-h^2)}\right)}{\sqrt{x^2+\delta^2}} - \frac{\tanh\left(\frac{x}{2}\right)}{x} \right| dx \leq Ch^2e^{-2/\lambda}. \tag{3.5}$$

Proof. First, we add and subtract $\tanh(x/2)/\sqrt{x^2+\delta^2}$ in (3.5). Then, we employ $T_c \sim e^{-1/\lambda}$ and Lemma 3.1 to estimate

$$\int_{T_D/T_c}^{\infty} \left| \frac{\tanh\left(\frac{\sqrt{x^2+\delta^2}}{2(1-h^2)}\right)}{\sqrt{x^2+\delta^2}} - \frac{\tanh\left(\frac{x}{2}\right)}{\sqrt{x^2+\delta^2}} \right| dx \leq Ch^2 \int_{T_D/T_c}^{\infty} \frac{1}{\cosh^2\left(\frac{x}{2}\right)} dx \leq Ch^2e^{-2/\lambda}$$

and

$$\int_{T_D/T_c}^{\infty} \left| \frac{\tanh\left(\frac{x}{2}\right)}{\sqrt{x^2+\delta^2}} - \frac{\tanh\left(\frac{x}{2}\right)}{x} \right| dx \leq Ch^2 \int_{T_D/T_c}^{\infty} \frac{1}{x^3} dx \leq Ch^2e^{-2/\lambda}.$$

Combining these bounds yields the claim by means of the triangle inequality. ■

From Lemma 3.2 and (2.7), we immediately conclude that

$$\int_{\mathbb{R}} \left\{ \frac{\tanh\left(\frac{\sqrt{x^2+\delta^2}}{2(1-h^2)}\right)}{\sqrt{x^2+\delta^2}} - \frac{\tanh\left(\frac{x}{2}\right)}{x} \right\} dx = O(h^2e^{-2/\lambda}). \tag{3.6}$$

3.1.3. Comparison with f_{BCS} . Given (3.6), the remaining task is to show that, because δ approximately solves the defining equation of f_{BCS} , it is *actually* close to f_{BCS} . This is the content of the following lemma.

Lemma 3.3. *Fix $h \in [0, 1]$. If $\phi \in [0, \infty)$ satisfies¹¹*

$$\int_{\mathbb{R}} \left\{ \frac{\tanh\left(\frac{\sqrt{x^2+\phi^2}}{2(1-h^2)}\right)}{\sqrt{x^2+\phi^2}} - \frac{\tanh\left(\frac{x}{2}\right)}{x} \right\} dx = R \tag{3.7}$$

for some $|R| \leq C$, then

$$\phi = f_{\text{BCS}}(h) + O(|R|^{1/2}) \tag{3.8}$$

with f_{BCS} defined in (2.9).

¹¹We follow the convention that, if $h = 1$, we replace $\tanh(\dots)$ by the constant function 1.

Hence, combining (3.6) with (3.8) and invoking Lemma 2.2 (iii), we find that

$$\delta(h) = f_{\text{BCS}}(h) + O(he^{-1/\lambda}) = f_{\text{BCS}}(h)(1 + O(e^{-1/\lambda})).$$

This concludes the proof of Proposition 2.1.

Proof of Lemma 3.3. First, we note that, given $h \in [0, 1]$, (3.7) has a solution $\phi \in [0, \infty)$ if and only if $R \leq \log(1/(1 - h^2))$. Then, as in the proof of [30, Lemma 6] (see [30, eq. (C50)]), we find that

$$\phi = e^{-R} f_{\text{BCS}}(\sqrt{h^2 + (1 - h^2)(1 - e^R)}). \tag{3.9}$$

Taylor expanding in R around 0, using regularity of f_{BCS} from Lemma 2.2, we get

$$e^R \phi = f_{\text{BCS}}(h) + \int_0^1 f'_{\text{BCS}}(\sqrt{h^2 + (1 - h^2)(1 - e^{tR})}) \frac{-(1 - h^2)Re^{tR}}{2\sqrt{h^2 + (1 - h^2)(1 - e^{tR})}} dt.$$

To bound the integral, we change variables to $s = h^2 + (1 - h^2)(1 - e^{tR})$ and bound $|f'_{\text{BCS}}(h)| \leq C$ by Theorem 2.2. We split into cases depending on the sign of R .

Case $R > 0$. For $R > 0$, the integral is bounded by

$$\begin{aligned} \int_{h^2 - (1 - h^2)(e^R - 1)}^{h^2} \frac{ds}{2\sqrt{s}} &= h - \sqrt{h^2 - (1 - h^2)(e^R - 1)} \\ &= \frac{(1 - h^2)(e^R - 1)}{\sqrt{h^2 - (1 - h^2)(e^R - 1)} + h}. \end{aligned}$$

Noting that $\delta/(\sqrt{\varepsilon - \delta} + \sqrt{\varepsilon}) \leq \sqrt{\delta}$ and that $(1 - h^2)(e^R - 1) \leq CR$, we find that the integral is bounded by \sqrt{R} .

Case $R < 0$. For $R < 0$, the integral is similarly bounded by

$$\int_{h^2}^{h^2 + (1 - h^2)(1 - e^R)} \frac{ds}{2\sqrt{s}} = \sqrt{h^2 + (1 - h^2)(1 - e^R)} - h \leq C \sqrt{|R|}.$$

Plugging these bounds into (3.9), we conclude the desired. ■

3.2. Proof of Theorem 2.4 (a)

We give here the proof of Theorem 2.4 (a). The argument is divided into several steps.

3.2.1. A priori spectral information. For any temperature T , we have by Theorem 1.5 that there exists a unique (up to a constant global phase) minimizer (γ_*, α_*) of the BCS functional. The function α_* is radial and has $\hat{\alpha}_* > 0$. Moreover, the operator $K_T^\Delta + \lambda V$ has lowest eigenvalue 0 and α_* is the unique eigenfunction with this eigenvalue.

3.2.2. Weak a priori bound on Δ . From the proof of Theorem 1.2 in [25], we have the following bound on the minimising (γ_*, α_*) of the BCS functional [25, eq. 3.12]

$$\begin{aligned} & \int (1 + p^2)(|\hat{\alpha}_*(p)|^2 + \hat{\gamma}_*(p))dp \\ & \leq 8T \int \log(1 + e^{-(p^2-\mu)/4T})dp + 8 \int \left[\frac{p^2}{4} - 1 + C_2(\lambda) \right]_- dp \leq C\lambda, \end{aligned}$$

with $C_2(\lambda) = \inf \text{spec}(p^2/4 + \lambda V) \leq 0$, and, thus, C_λ uniformly bounded for λ small enough. In particular, $\|\alpha_*\|_{H^1} \leq C$ uniformly for λ small enough. By Sobolev’s inequality, [35, Theorem 8.3] we then have

$$\begin{aligned} \|\alpha\|_{L^\infty}^2 & \leq C \|\nabla\alpha\|_{L^2} \|\alpha\|_{L^2} \leq C & (d = 1) \\ \|\alpha\|_{L^q}^2 & \leq C \|\nabla\alpha\|_{L^2} \|\alpha\|_{L^2} \leq C & (d = 2) \\ \|\alpha\|_{L^6}^2 & \leq C \|\nabla\alpha\|_{L^2}^2 \leq C & (d = 3) \end{aligned}$$

for any $2 \leq q < \infty$. Thus,

$$\begin{aligned} \|\Delta\|_{L^\infty} & \leq 2\lambda \|V\alpha\|_{L^1} \\ & \leq \begin{cases} 2\lambda \|V\|_{L^1} \|\alpha\|_{L^\infty} & \text{for } d = 1, \\ 2\lambda \|V\|_{L^p} \|\alpha\|_{L^2}^{(2pq-2q-2p)/(pq-2p)} \|\alpha\|_{L^q}^{(2q-pq)/(pq-2p)} & \text{for } d = 2, \\ 2\lambda \|V\|_{L^{3/2}} \|\alpha\|_{L^2}^{1/2} \|\alpha\|_{L^6}^{1/2} & \text{for } d = 3 \end{cases} \\ & \leq C\lambda \end{aligned} \tag{3.10}$$

uniformly for λ small enough (for any $1 < p < \min\{2, p_V\}$ by choosing q large enough in dimension $d = 2$). In particular, we see that $\Delta(p) \rightarrow 0$ as $\lambda \rightarrow 0$.

3.2.3. Birman–Schwinger principle. Next, by the Birman–Schwinger principle [13, 20, 25], the fact that $K_T^\Delta + \lambda V$ has lowest eigenvalue 0 with α_* being the unique eigenvector is equivalent to the Birman–Schwinger operator

$$B_{T,\Delta} := \lambda V^{1/2} \frac{1}{K_T^\Delta} |V|^{1/2}$$

having -1 as its lowest eigenvalue and $\phi = V^{1/2}\alpha_*$ being the unique eigenfunction corresponding to this eigenvalue. Here we use the convention

$$V^{1/2}(x) = \operatorname{sgn}(V(x))|V(x)|^{1/2}.$$

We decompose the Birman–Schwinger operator into a dominant singular term and an error term. For this purpose, we define the (rescaled) Fourier transform restricted to the sphere $\mathfrak{F}_\mu: L^1(\mathbb{R}^d) \rightarrow L^2(\mathbb{S}^{d-1})$ by

$$(\mathfrak{F}_\mu \psi)(p) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} \psi(x) e^{-i\sqrt{\mu}p \cdot x} dx,$$

which is well defined by the Riemann–Lebesgue lemma. Define then

$$m(T, \Delta) = \frac{1}{|\mathbb{S}^{d-1}|} \int_{|p| \leq \sqrt{2\mu}} \frac{1}{K_T^\Delta(p)} dp \tag{3.11}$$

and decompose

$$B_{T,\Delta} = \lambda m(T, \Delta) V^{1/2} \mathfrak{F}_\mu^\dagger \mathfrak{F}_\mu |V|^{1/2} + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2},$$

with $M_{T,\Delta}$ defined such that this holds. Analogously to [13, Lemma 2] and [28, Lemma 3.4], we have the following lemma, the proof of which (it is analogous to the one of [28, Lemma 3.4]) is given in Section A.2.1.

Lemma 3.4. *We have $\|V^{1/2} M_{T,\Delta} |V|^{1/2}\|_{\text{HS}} \leq C$ for small λ uniformly in T and Δ , where $\|\cdot\|_{\text{HS}}$ denotes the Hilbert–Schmidt norm of an operator.*

We conclude that $1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}$ is invertible for sufficiently small λ and thus, analogously to [21, Lemma 4] and [27, Lemma 13 and Proposition 15] the fact that $B_{T,\Delta}$ has lowest eigenvalue -1 is equivalent to the fact that the operator

$$S_{T,\Delta} := \lambda m(T, \Delta) \mathfrak{F}_\mu |V|^{1/2} \frac{1}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger \tag{3.12}$$

acting on $L^2(\mathbb{S}^{d-1})$ has lowest eigenvalue -1 . Moreover, the function $\mathfrak{F}_\mu |V|^{1/2} \phi = \mathfrak{F}_\mu V \alpha_*$ is the unique eigenfunction of $S_{T,\Delta}$ corresponding to the eigenvalue -1 .

3.2.4. A priori bounds on Δ . For the analysis of the integral $m(T, \Delta)$ we need some a priori bounds on Δ . Analogously as in [21, 28] we need some control of $\Delta(p)$ in terms of $\Delta(\sqrt{\mu})$ and some type of Lipschitz-continuity of Δ . These are collected in the following lemma.

Lemma 3.5. *The function Δ satisfies the bounds*

$$\Delta(p) \leq C\Delta(\sqrt{\mu}), \tag{3.13}$$

$$|\Delta(p) - \Delta(q)| \leq C\Delta(\sqrt{\mu})|p - q| \tag{3.14}$$

for sufficiently small λ .

Proof. As noted above, the function $\mathfrak{F}_\mu|V|^{1/2}\phi = \mathfrak{F}_\mu V\alpha$ is the eigenfunction of $S_{T,\Delta}$ corresponding to the lowest eigenvalue -1 .

Further, to leading order, $S_{T,\Delta}$ is proportional to $\mathcal{V}_\mu = \mathfrak{F}_\mu V \mathfrak{F}_\mu^\dagger$. Since the constant function $u = |\mathbb{S}^{d-1}|^{-1/2} \in L^2(\mathbb{S}^{d-1})$ is the ground state of \mathcal{V}_μ (see the argument around (1.11)), the same is also true for $S_{T,\Delta}$ whenever λ is small enough. Hence, one can easily see that

$$\frac{1}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger u$$

is an eigenvector of $B_{T,\Delta}$ corresponding to the eigenvalue -1 and thus proportional to $\phi = V^{1/2}\alpha_*$. Thus, with \mathfrak{F} denoting the usual Fourier transform, by expanding $1/(1+x) = 1-x/(1+x)$, we have

$$\begin{aligned} \Delta &= -2\lambda \mathfrak{F}|V|^{1/2}\phi = f(\lambda)\mathfrak{F}|V|^{1/2} \frac{1}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger u \\ &= f(\lambda) \left(\int_{\mathbb{S}^{d-1}} \widehat{V}(p - \sqrt{\mu}q) d\omega(q) + \lambda \eta_\lambda(p) \right) \end{aligned}$$

for some constant $f(\lambda)$ (where we absorbed the factor $|\mathbb{S}^{d-1}|^{-1/2} (2\pi)^{-d/2}$ into $f(\lambda)$). One easily verifies that

$$\eta_\lambda = -(2\pi)^{d/2} \mathfrak{F}|V|^{1/2} \frac{V^{1/2} M_{T,\Delta} |V|^{1/2}}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger u$$

has $\|\eta_\lambda\|_\infty \leq C$ uniformly in $\lambda < \lambda_0$ for some λ_0 by Theorem 3.4. By evaluating at $p = \sqrt{\mu}$, we find $|f(\lambda)| \leq C\Delta(\sqrt{\mu})$ for small λ , and thus the global bound (3.13). Moreover, we have the Lipschitz-bound

$$\begin{aligned} &|\Delta(p) - \Delta(q)| \\ &\leq |f(\lambda)| |p - q| \left\| |x| |V|^{1/2} \frac{1}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger u \right\|_{L^1} \\ &\leq C\Delta(\sqrt{\mu}) |p - q| \| |x|^2 V \|_{L^1}^{1/2} \frac{1}{1 - \lambda \|V^{1/2} M_{T,\Delta} |V|^{1/2}\|} \|V^{1/2} \mathfrak{F}_\mu^\dagger u\|_{L^2} \\ &\leq C\Delta(\sqrt{\mu}) |p - q| \end{aligned}$$

for sufficiently small λ . ■

3.2.5. First order. Expanding the resolvent in (3.12) to first order in a geometric series, we see that $S_{T,\Delta}$ to leading order is proportional to the operator \mathcal{V}_μ (defined in (1.10) above). Moreover, we have

$$\begin{aligned} -1 &= \lambda m(T, \Delta) \inf \operatorname{spec} \left(\mathfrak{F}_\mu |V|^{1/2} \frac{1}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger \right) \\ &= \lambda m(T, \Delta) \inf \operatorname{spec} \mathcal{V}_\mu (1 + O(\lambda)) \\ &= \lambda e_\mu m(T, \Delta) (1 + O(\lambda)). \end{aligned}$$

In particular,

$$m(T, \Delta) = -\frac{1}{\lambda e_\mu} (1 + O(\lambda)) \rightarrow \infty, \quad \text{as } \lambda \rightarrow 0.$$

3.2.6. Exponential vanishing of Δ . Pointwise, we may bound $K_T^\Delta \geq E_\Delta$. Thus, by the first-order analysis above, we have

$$\begin{aligned} \frac{-1}{\lambda e_\mu} (1 + O(\lambda)) = m(T, \Delta) &= \frac{1}{|\mathbb{S}^{d-1}|} \int_{|p| < \sqrt{2\mu}} \frac{1}{K_T^\Delta} dp \\ &\leq \frac{1}{|\mathbb{S}^{d-1}|} \int_{|p| < \sqrt{2\mu}} \frac{1}{\sqrt{|p^2 - \mu|^2 + |\Delta(p)|^2}} dp. \end{aligned}$$

The latter integral is calculated in [21, 28]. The same calculation is valid here by the bounds (3.13) and (3.14) and the fact that $\Delta(p) \rightarrow 0$ by (3.10). That is,

$$\frac{-1}{\lambda e_\mu} \leq \mu^{d/2-1} \left(\log \frac{\mu}{\Delta(\sqrt{\mu})} + O(1) \right)$$

in the limit $\lambda \rightarrow 0$. We conclude

$$\Delta(\sqrt{\mu}) \lesssim e^{-c/\lambda} \quad \text{as } \lambda \rightarrow 0$$

(with $c = -(1/e_\mu)\mu^{d/2-1}$).

The constant c shall henceforth be used generically and its precise value might change from line to line.

3.2.7. Infinite order. Recall that, for small λ , the unique eigenfunction of $S_{T,\Delta}$ corresponding to the eigenvalue -1 is given by the constant function u . Thus, for small λ , we have

$$-1 = \lambda m(T, \Delta) \left\langle u \left| \mathfrak{F}_\mu |V|^{1/2} \frac{1}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger \right| u \right\rangle.$$

Combining this for the temperatures T and T_c , we find

$$\begin{aligned} & m(T, \Delta) - m(T_c, 0) \\ &= \frac{-1}{\lambda} \left[\frac{1}{\langle u \mid \mathfrak{F}_\mu |V|^{1/2} \frac{1}{1+\lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger \mid u \rangle} \right. \\ &\quad \left. - \frac{1}{\langle u \mid \mathfrak{F}_\mu |V|^{1/2} \frac{1}{1+\lambda V^{1/2} M_{T_c,0} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger \mid u \rangle} \right] \\ &= \frac{-1}{\lambda e_\mu^2} (1 + O(\lambda)) \langle u \mid \mathfrak{F}_\mu |V|^{1/2} A_{T,\Delta,T_c} V^{1/2} \mathfrak{F}_\mu^\dagger \mid u \rangle, \end{aligned}$$

for small enough λ by expanding to first order in the denominator and noting that $\text{inf spec } \mathcal{V}_\mu = e_\mu$, where

$$A_{T,\Delta,T_c} := \frac{1}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} - \frac{1}{1 + \lambda V^{1/2} M_{T_c,0} |V|^{1/2}}.$$

The proof of the following lemma (which is somewhat analogous to the proof of Theorem 3.4) is given in Section A.2.2.

Lemma 3.6. *There exists $\lambda_0 > 0$ such that, for $0 < \lambda < \lambda_0$,*

$$\|V^{1/2} (M_{T,\Delta} - M_{T_c,0}) |V|^{1/2}\|_{\text{HS}} \leq C e^{-c/\lambda},$$

for some constants $C, c > 0$ uniformly in λ .

Having Theorem 3.6 at hand, we write the difference as a telescoping sum

$$\begin{aligned} A_{T,\Delta,T_c} &= \sum_{k=1}^{\infty} (-\lambda)^k [(V^{1/2} M_{T,\Delta} |V|^{1/2})^k - (V^{1/2} M_{T_c,0} |V|^{1/2})^k] \\ &= \sum_{k=1}^{\infty} (-\lambda)^k \sum_{\ell=0}^{k-1} (V^{1/2} M_{T,\Delta} |V|^{1/2})^{k-1-\ell} V^{1/2} (M_{T,\Delta} - M_{T_c,0}) |V|^{1/2} \\ &\quad \times (V^{1/2} M_{T_c,0} |V|^{1/2})^\ell. \end{aligned}$$

Thus, by Theorems 3.4 and 3.6, we have

$$\begin{aligned} \|A_{T,\Delta,T_c}\|_{\text{HS}} &\leq \sum_{k=1}^{\infty} \lambda^k \sum_{\ell=0}^{k-1} C^{k-1-\ell} \times C e^{-c/\lambda} \times C^\ell \\ &\leq \sum_{k=1}^{\infty} k \lambda^k C^k e^{-c/\lambda} \leq C \lambda e^{-c/\lambda}. \end{aligned}$$

We conclude that

$$|m(T, \Delta) - m(T_c, 0)| \leq C e^{-c/\lambda}. \tag{3.15}$$

3.2.8. Calculation of the integral $m(T, \Delta) - m(T_c, 0)$. To extract the asymptotics in (2.11) from the bound in (3.15), we calculate the difference $m(T, \Delta) - m(T_c, 0)$ and show that it is essentially the left-hand side of (2.9). The argument is essentially given in [30, Appendix C.4]. For completeness, we give the argument here.

By changing variables to $s = (p^2 - \mu)/\mu$ and defining

$$x(s) = \frac{\Delta(\sqrt{\mu(1+s)})}{\mu},$$

we get

$$\begin{aligned} & m(T, \Delta) - m(T_c, 0) \\ &= \int_0^{\sqrt{2\mu}} \left(\frac{1}{K_T^\Delta} - \frac{1}{K_{T_c}} \right) p^{d-1} dp \\ &= \frac{\mu^{d/2-1}}{2} \int_{-1}^1 \left(\frac{\tanh\left(\frac{\sqrt{s^2+x(s)^2}}{2T/\mu}\right)}{\sqrt{s^2+x(s)^2}} - \frac{\tanh\frac{s}{2T_c/\mu}}{s} \right) (1+s)^{d/2-1} ds. \end{aligned}$$

This is of the form where we can use [30, Lemma 5].

Lemma 3.7 ([30, Lemma 5]). *Let g, G be functions with $g(0) = G(0) = 1$ and $g \in L^\infty$ and let $\tau, \tau_c, \delta > 0$. Assume that $\tilde{g}(s) := (g(s) - 1)/s$ and $\tilde{G}(s) := (G(s) - 1)/s$ satisfy $\tilde{g}, \tilde{G} \in L^\infty(\mathbb{R})$. Let $s_1 > 0$ such that $g(s) > 1/2$ for $|s| < s_1$ and define*

$$\begin{aligned} J_{\tau,\delta,\tau_c}(g, G) &= \int_{\mathbb{R}} \left\{ \frac{\tanh\frac{\sqrt{s^2+g(s)^2\delta^2}}{2\tau}}{\sqrt{s^2+g(s)^2\delta^2}} - \frac{\tanh\frac{s}{2\tau_c}}{s} \right\} G(s) ds, \\ J_{\tau,\delta,\tau_c}^{(0)} &= J_{\tau,\delta,\tau_c}(1, 1) = \int_{\mathbb{R}} \left\{ \frac{\tanh\frac{\sqrt{s^2+\delta^2}}{2\tau}}{\sqrt{s^2+\delta^2}} - \frac{\tanh\frac{s}{2\tau_c}}{s} \right\} ds. \end{aligned}$$

Then

$$\begin{aligned} & |J_{\tau,\delta,\tau_c}(g, G) - J_{\tau,\delta,\tau_c}^{(0)}| \\ & \leq \|\tilde{G}\|_{L^\infty} (4\tau + 4\tau_c + \pi\delta\|g\|_{L^\infty}) + 4\delta\|\tilde{g}\|_{L^\infty} \left(1 + \|g\|_{L^\infty}\right) \left(1 + \frac{\delta}{2s_1}\right). \end{aligned}$$

To apply this lemma we write

$$x(s) = \frac{\Delta(\sqrt{\mu(1+s)})}{\Delta(\sqrt{\mu})} \frac{\Delta(\sqrt{\mu})}{\mu} = g(s)\delta.$$

Then $g \in L^\infty$ uniformly in λ by (3.13) and $\tilde{g} \in L^\infty$ uniformly in λ by (3.14). Finally, clearly $G(s) = (1 + s)^{d/2-1} \chi_{|s| \leq 1}$ has $\tilde{G} \in L^\infty$. We conclude that

$$m(T, \Delta) - m(T_c, 0) = \frac{\mu^{d/2-1}}{2} \int_{\mathbb{R}} \left\{ \frac{\tanh \frac{\sqrt{s^2 + (\Delta(\sqrt{\mu})/\mu)^2}}{2T/\mu}}{\sqrt{s^2 + (\frac{\Delta(\sqrt{\mu})}{\mu})^2}} - \frac{\tanh \frac{s}{2T_c/\mu}}{s} \right\} ds + O(e^{-c/\lambda}).$$

Writing $T = T_c(1 - h^2)$, recalling the bound in (3.15) and changing variables, we find

$$\int_{\mathbb{R}} \left\{ \frac{\tanh \frac{\sqrt{s^2 + (\Delta(\sqrt{\mu})/T_c)^2}}{2(1-h^2)}}{\sqrt{s^2 + (\frac{\Delta(\sqrt{\mu})}{T_c})^2}} - \frac{\tanh \frac{s}{2}}{s} \right\} ds = R,$$

with $R = O(e^{-c/\lambda})$. Hence, by Lemma 3.3, we find that

$$\frac{\Delta(\sqrt{\mu})}{T_c} = f_{\text{BCS}}(h) + O(|R|^{1/2}) = f_{\text{BCS}}(h)(1 + O(h^{-1}e^{-c/\lambda})) \tag{3.16}$$

since $f_{\text{BCS}}(h) \sim h$ for small h by Lemma 2.2.

3.2.9. Comparing $\Delta(\sqrt{\mu})$ and Ξ . We finally prove that Ξ is essentially given by $\Delta(\sqrt{\mu})$.

Clearly, $\Xi = \inf_p E_\Delta(p) \leq E_\Delta(\sqrt{\mu}) = \Delta(\sqrt{\mu})$. To show a corresponding lower bound, consider p with $|p^2 - \mu| \leq \Xi \leq \Delta(\sqrt{\mu})$. Then, by (3.14) and the bound $\Delta(\sqrt{\mu}) = O(e^{-c/\lambda})$, we have

$$\begin{aligned} \Delta(p) &\geq \Delta(\sqrt{\mu}) - C\Delta(\sqrt{\mu})|p - \sqrt{\mu}| \\ &\geq \Delta(\sqrt{\mu})(1 - C\Delta(\sqrt{\mu})) \\ &\geq \Delta(\sqrt{\mu})(1 + O(e^{-c/\lambda})). \end{aligned}$$

We conclude that

$$\Xi = \Delta(\sqrt{\mu})(1 + O(e^{-c/\lambda})). \tag{3.17}$$

Together with (3.16), this concludes the proof of Theorem 2.4 (a).

3.3. Non-universal property of Δ : Proof of equation (2.14)

From the Birman–Schwinger argument (Section 3.2.3), we have that $\phi = V^{1/2}\alpha_*$ is the (unique) eigenvector of

$$\frac{\lambda m(T, \Delta)}{1 + \lambda V^{1/2} M_{T, \Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger \mathfrak{F}_\mu |V|^{1/2}$$

corresponding to the eigenvalue -1 . Recalling that $\Delta = -2\lambda\mathfrak{F}V\alpha_*$, we thus get the equation

$$\Delta = -\mathfrak{F}|V|^{1/2} \frac{\lambda m(T, \Delta)}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger \Delta(\sqrt{\mu})$$

with $\Delta(\sqrt{\mu})$ being the constant function on the unit sphere of value $\Delta(\sqrt{\mu})$. Recall that

$$-1 = \lambda m(T, \Delta) \left\langle u \left| \mathfrak{F}_\mu |V|^{1/2} \frac{1}{1 + \lambda V^{1/2} M_{T,\Delta} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger \right| u \right\rangle$$

for small enough λ . By the same argument as in Section 3.2.7, we may replace $M_{T,\Delta}$ by $M_{0,0}$, its corresponding value at $T = \Delta = 0$, up to errors of order $e^{-c/\lambda}$. (Concretely, one can define $M_{0,0}$ via the representation of its kernel as given in (A.3)–(A.5), (A.8), and (A.9), setting $T = \Delta = 0$.) Hence, for sufficiently small λ ,

$$\begin{aligned} \frac{\Delta}{\Delta(\sqrt{\mu})} &= \frac{|\mathbb{S}^{d-1}|^{1/2} \mathfrak{F}|V|^{1/2} \frac{1}{1 + \lambda V^{1/2} M_{0,0} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger u}{\langle u \left| \mathfrak{F}_\mu |V|^{1/2} \frac{1}{1 + \lambda V^{1/2} M_{0,0} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger \right| u \rangle} + O(e^{-c/\lambda}) \\ &= F + O(e^{-c/\lambda}). \end{aligned}$$

Clearly, the function F does not depend on the temperature T .

3.4. Ginzburg–Landau theory: Proof of Theorem 2.4 (b)

As mentioned above, the proof of Theorem 2.4 (b) builds on Ginzburg–Landau (GL) theory. For convenience of the reader, we recall the main input from GL theory for the purpose of the present paper in Proposition 3.9 below. More general and detailed statements can be found in the original papers [10, 11, 14, 16]. In particular, these works allow for external fields or ground state degeneracy (cf. Lemma 3.8 below), respectively.

As a preparation for Proposition 3.9, we have the following lemma.

Lemma 3.8 (Ground state of $K_{T_c} + \lambda V$). *Let V satisfy Assumptions 1.1 and 1.4. Then, $K_{T_c} + \lambda V$ has 0 as a non-degenerate ground state eigenvalue and its $L^2(\mathbb{R}^d)$ -normalized ground state α_0 can be chosen to have strictly positive Fourier transform. Moreover, it holds that $\hat{\alpha}_0 \in L^\infty(\mathbb{R}^d)$.*

Proof. Since $T_c > 0$ (recall the discussion below Assumption 1.4), we first note that the Fourier multiplier K_{T_c} is strictly positive. Then, using $\hat{V} \leq 0$, the claim follows from a Perron–Frobenius type argument (see also [21] and [14, Assumption 2]). The fact that $\hat{\alpha}_0 \in L^\infty(\mathbb{R}^d)$ follows from [14, Proposition 2] by invoking Assumption 1.1. ■

We can now formulate the main results from GL theory, needed for the present paper.

Proposition 3.9 (Ginzburg–Landau theory, see [16, Theorem 2.10]). *Let V be a function satisfying Assumptions 1.1 and 1.4 and suppose that $0 \leq T < T_c$. Then, using the notations from Proposition 1.5 and Lemma 3.8, we have that*

$$\mathcal{F}_T[\Gamma_*] - \mathcal{F}_T[\Gamma_{FD}] = h^4 \mathcal{E}_{GL}(\psi_{GL}) + \mathcal{O}(h^6) \quad \text{as } h \rightarrow 0,$$

where $\psi_{GL} \neq 0$ minimizes the Ginzburg–Landau “functional” $\mathcal{E}_{GL}: \mathbb{C} \rightarrow \mathbb{R}$,

$$\begin{aligned} \mathcal{E}_{GL}(\psi) = |\psi|^4 & \left[\frac{1}{T_c^3} \int_{\mathbb{R}^d} \frac{g_1\left(\frac{p^2 - \mu}{T_c}\right)}{\left(\frac{p^2 - \mu}{T_c}\right)} |K_{T_c}(p)|^4 |\hat{\alpha}_0(p)|^4 \, dp \right] \\ & - |\psi|^2 \left[\frac{1}{2T_c} \int_{\mathbb{R}^d} \frac{1}{\cosh^2\left(\frac{p^2 - \mu}{2T_c}\right)} |K_{T_c}(p)|^2 |\hat{\alpha}_0(p)|^2 \, dp \right]. \end{aligned} \tag{3.18}$$

Here, we used the auxiliary function g_1 from (3.3). Moreover, we can decompose the off-diagonal element $\hat{\alpha}_*$ of Γ_* as

$$\hat{\alpha}_* = h|\psi_0|\hat{\alpha}_0 + \hat{\xi} \tag{3.19}$$

where $\|\hat{\xi}\|_{L^2} = \mathcal{O}(h^2)$ and $\psi_0 \neq 0$ approximately minimizes (3.18), i.e.,

$$\mathcal{E}_{GL}(\psi_0) \leq \mathcal{E}_{GL}(\psi_{GL}) + \mathcal{O}(h^2). \tag{3.20}$$

Remark 3.10. We emphasize that all error terms in the above proposition (and also the implicit constants hidden in $\hat{\alpha}_0 \in L^\infty(\mathbb{R}^d)$) are *not* uniform in λ . This crucially limits the applicability of our GL theory-based method for temperatures slightly away from the critical one with $e^{-c/\lambda} \ll h \ll 1$ (cf. the error bound in (2.12)). Indeed, a careful examination of the proofs in [14, 16] reveals that the hidden dependencies on the critical temperature T_c are at most inverse polynomially and hence exponential in λ , i.e., $e^{c/\lambda}$ for some $c > 0$ (independent of λ and h).

3.4.1. Minimizing the Ginzburg–Landau functional. Given the inputs from GL theory, Theorem 2.4 (b) is based on the following Proposition 3.11, the proof of which we postpone after finishing the proof of Theorem 2.4 (b).

Proposition 3.11. *The (up to a phase unique) minimizer ψ_{GL} of the GL functional (3.18) satisfies*

$$|\psi_{GL}| = C_{\text{univ}} \frac{T_c}{\Delta_0(\sqrt{\mu})} (1 + o_{\lambda \rightarrow 0}(1)), \tag{3.21}$$

where $\Delta_0 := -2(2\pi)^{-d/2}\lambda\widehat{V} \star \widehat{\alpha}_0$,¹² the constant C_{univ} is given in (2.10), and the error is uniform in h .

The fact that $|\psi_0| > 0$ approximately minimizes (3.18) (see (3.20)) implies that (recalling Remark 3.10)

$$|\psi_0| = |\psi_{\text{GL}}| + O(h e^{c/\lambda}).$$

Therefore, by means of (3.19) in combination with Proposition 3.11, we infer

$$\widehat{\alpha}_* = C_{\text{univ}} T_c h \frac{\widehat{\alpha}_0}{\Delta_0(\sqrt{\mu})} (1 + o_{\lambda \rightarrow 0}(1) + O(h e^{c/\lambda})) + \widehat{\xi}.$$

Thus, after taking the convolution with $\lambda\widehat{V}$,

$$\Delta = C_{\text{univ}} T_c h \frac{\Delta_0}{\Delta_0(\sqrt{\mu})} (1 + o_{\lambda \rightarrow 0}(1) + O(h e^{c/\lambda})) + \Delta_\xi, \tag{3.22}$$

where $\Delta > 0$ is the unique solution of the BCS gap equation (1.4) (see Proposition 1.5) and we denoted $\Delta_\xi := -2(2\pi)^{-d/2}\lambda\widehat{V} \star \widehat{\xi}$.

3.4.2. A priori bounds on Δ_0 . For the proof of Theorem 2.4 (b), we need some a priori bounds on Δ_0 analogously to those of Section 3.2.4. The bounds follow from the following lemma, the proof of which is analogous to the argument of Sections 3.2.3 and 3.2.4 and given in Section A.2.3.

Lemma 3.12 (cf. [21, Lemma 4] and [27, Lemma 13]). *Let $\alpha_0 \in H^1(\mathbb{R}^d)$, with $\widehat{\alpha}_0 > 0$, be the unique $H^1(\mathbb{R}^d)$ -normalized ground state of $K_{T_c} + \lambda V$ from Lemma 3.8. Moreover, let $u(p) = (|\mathbb{S}^{d-1}|)^{-1/2}$ be the constant function on the sphere \mathbb{S}^{d-1} and let*

$$\widehat{\phi}(p) = -\frac{1}{(2\pi)^{d/2}} \int_{\mathbb{S}^{d-1}} \widehat{V}(p - \sqrt{\mu}q) \, d\omega(q). \tag{3.23}$$

Then, $\Delta_0 = -2(2\pi)^{-d/2}\lambda\widehat{V} \star \widehat{\alpha}_0$ can be expanded as

$$\Delta_0(p) = f(\lambda)[\widehat{\phi}(p) + \lambda\eta_\lambda(p)] \tag{3.24}$$

for some positive function $f(\lambda)$ and $\|\eta_\lambda\|_{L^\infty(\mathbb{R}^d)}$ bounded uniformly in $\lambda > 0$.

After realizing $\widehat{\phi}(\sqrt{\mu}) = -e_\mu$ by (1.11), we conclude for small enough $\lambda > 0$ that

$$\Delta_0(p) - \Delta_0(\sqrt{\mu}) = \frac{[\widehat{\phi}(p) - \widehat{\phi}(\sqrt{\mu})] + \lambda[\eta_\lambda(p) - \eta_\lambda(\sqrt{\mu})]}{-e_\mu + \lambda\eta_\lambda(\sqrt{\mu})} \Delta_0(\sqrt{\mu}).$$

¹²Note that Δ_0 was denoted by t in [14, 16]. We also remark that $\Delta_0(\sqrt{\mu}) \neq 0$ as follows from $\widehat{\alpha}_0 > 0$ (by Lemma 3.8) and $\widehat{V} \leq 0$ (by Assumption 1.4).

Now, it is an easy computation to see that $|\hat{\varphi}(p) - \hat{\varphi}(q)| \leq C \min\{|p| - |q|, 1\}$ for all $p, q \in \mathbb{R}^d$. Thus,

$$|\Delta_0(p) - \Delta_0(\sqrt{\mu})| \leq C(\min\{|p| - \sqrt{\mu}, 1\} + \lambda)\Delta_0(\sqrt{\mu}). \tag{3.25}$$

3.4.3. A priori bounds on Δ_ξ . For the following arguments, we need two estimates on $\Delta_\xi = -2(2\pi)^{-d/2}\lambda\hat{V} \star \hat{\xi}$.

- First, it is a simple consequence of Young’s inequality and $\|\hat{\xi}\|_{L^2} = \|\xi\|_{L^2} = O(h^2e^{c/\lambda})$ that

$$\|\Delta_\xi\|_{L^\infty} = \|V\|_{L^2} O(h^2e^{c/\lambda}). \tag{3.26}$$

- Second, we note that $\Delta_\xi(p) - \Delta_\xi(q)$ is (proportional to) the Fourier transform of $V(x)(1 - e^{i(p-q)\cdot x})\xi(x)$, and

$$\begin{aligned} \|V(x)(1 - e^{i(p-q)\cdot x})\|_{L^2}^2 &= \int_{\mathbb{R}^d} |V(x)|^2 |1 - e^{i(p-q)\cdot x}|^2 dx \\ &\leq C|p - q|^2 \int_{\mathbb{R}^d} |V(x)|^2 |x|^2 dx. \end{aligned}$$

Using radiality of Δ and Δ_0 , we conclude the radiality of Δ_ξ and therefore

$$|\Delta_\xi(p) - \Delta_\xi(q)| \leq \|p| - |q|\| \cdot \|V\|_{L^2} O(h^2e^{c/\lambda}). \tag{3.27}$$

Recall that $\|(1 + |\cdot|)V\|_{L^2} < \infty$ by assumption.

3.4.4. Comparing $\Delta(\sqrt{\mu})$ and Ξ . We aim at proving

$$\Xi = \Delta(\sqrt{\mu})(1 + O(\lambda + h^2e^{c/\lambda})). \tag{3.28}$$

In order to see this, we note that clearly $\Xi = \sqrt{(p^2 - \mu)^2 + |\Delta(p)|^2} \leq \Delta(\sqrt{\mu})$. For the reverse inequality, let $p \in \mathbb{R}^d$ with $|p^2 - \mu| \leq \Xi \leq \Delta(\sqrt{\mu})$. Then

$$\begin{aligned} &|\Delta(p) - \Delta(\sqrt{\mu})| \\ &\leq CT_c h(1 + o_{\lambda \rightarrow 0}(1) + O(he^{c/\lambda})) \cdot (|p| - \sqrt{\mu} + \lambda) + C|p| - \sqrt{\mu}|h^2e^{c/\lambda} \end{aligned}$$

by application of (3.25) and (3.27). Using that $\Delta(\sqrt{\mu}) \sim T_c h$, as a consequence of (3.22) for h small enough (meaning $he^{c/\lambda} \ll 1$), we then conclude

$$|\Delta(p) - \Delta(\sqrt{\mu})| \leq C(\lambda + h^2e^{c/\lambda})\Delta(\sqrt{\mu}).$$

In combination with the upper bound, this proves (3.28).

3.4.5. Conclusion: Proof of Theorem 2.4 (b). We evaluate (3.22) at $p = \sqrt{\mu}$, so that we find

$$\Xi = [C_{\text{univ}} T_c h(1 + o_{\lambda \rightarrow 0}(1) + O(h e^{c/\lambda})) + O(h^2 e^{c/\lambda})] \cdot (1 + O(\lambda + h^2 e^{c/\lambda}))$$

with the aid of (3.26) and (3.28). Collecting all the error terms leaves us with

$$\Xi = C_{\text{univ}} T_c h(1 + o_{\lambda \rightarrow 0}(1) + O(h e^{c/\lambda})).$$

Hence, using $f_{\text{BCS}}(h) = C_{\text{univ}} h + O(h^2)$, by Lemma 2.2, we arrive at Theorem 2.4 (b).

3.4.6. Proof of Proposition 3.11. In the following estimates, we use the shorthand notations (recall the definition of the auxiliary function g_1 from (3.3))

$$f_4(p) := \frac{g_1\left(\frac{p^2 - \mu}{T_c}\right)}{\left(\frac{p^2 - \mu}{T_c}\right)} \quad f_2(p) := \frac{1}{\cosh^2\left(\frac{p^2 - \mu}{2T_c}\right)},$$

such that the absolute value of the minimizer ψ_{GL} of (3.18) is given by

$$\begin{aligned} |\psi_{\text{GL}}| &= T_c \left(\frac{\int_{\mathbb{R}^d} f_2(p) |K_{T_c}(p)|^2 |\hat{\alpha}_0(p)|^2 dp}{4 \int_{\mathbb{R}^d} f_4(p) |K_{T_c}(p)|^4 |\hat{\alpha}_0(p)|^4 dp} \right)^{1/2} \\ &= T_c \left(\frac{\int_{\mathbb{R}^d} f_2(p) |\Delta_0(p)|^2 dp}{\int_{\mathbb{R}^d} f_4(p) |\Delta_0(p)|^4 dp} \right)^{1/2}. \end{aligned}$$

We denoted $\Delta_0 = -2(2\pi)^{-d/2} \lambda \hat{V} \star \hat{\alpha}_0$ (as in Proposition 3.11) and used that $\alpha_0 \in \ker(K_{T_c} + \lambda V)$. Note that $\Delta_0 = |\Delta_0|$, by means of Proposition 1.5 and $\hat{V} \leq 0$ from Assumption 1.4.

Next, we add and subtract $|\Delta_0(\sqrt{\mu})|^2$ (resp. $|\Delta_0(\sqrt{\mu})|^4$) in the integral in the numerator (resp. denominator). The terms involving $|\Delta_0(\sqrt{\mu})|^j$ are evaluated as follows.

Lemma 3.13 (Emergence of C_{univ} in GL theory). *Let $\mu > 0$. In the limit $T_c/\mu \rightarrow 0$, we have that (recall C_{univ} from (2.10))*

$$\left(\int_{\mathbb{R}^d} \frac{1}{\cosh^2\left(\frac{p^2 - \mu}{2T_c}\right)} dp \right) / \left(\int_{\mathbb{R}^d} \frac{g_1\left(\frac{p^2 - \mu}{T_c}\right)}{\left(\frac{p^2 - \mu}{T_c}\right)} dp \right)^{1/2} \rightarrow C_{\text{univ}} \quad (3.29)$$

for all $d = 1, 2, 3$.

Proof. Since the integrands are both radial, we switch to spherical coordinates and neglect the common $|\mathbb{S}^{d-1}|$ -factor in the numerator and the denominator. By splitting the remaining radial integration according to $p^2 \leq \mu$ and $p^2 \geq \mu$ and changing the

integration variables from $(p^2 - \mu)/2T_c$ to $-t$ (resp. t), we find that the numerator of (3.29) is equal to

$$2T_c\mu^{(d-2)/2} \left[\int_0^{\mu/2T_c} \left(1 - 2\frac{T_c}{\mu}t\right)^{(d-2)/2} + \int_0^\infty \left(1 + 2\frac{T_c}{\mu}t\right)^{(d-2)/2} \right] \left(\frac{1}{\cosh^2(t)}\right) dt. \tag{3.30}$$

Similarly, we find the that denominator of (3.29) is equal to

$$\frac{2T_c\mu^{(d-2)/2}}{8} \left[\int_0^{\mu/2T_c} \left(1 - 2\frac{T_c}{\mu}t\right)^{(d-2)/2} + \int_0^\infty \left(1 + 2\frac{T_c}{\mu}t\right)^{(d-2)/2} \right] \times \left(\frac{\tanh(t)}{t^3} - \frac{1}{t^2 \cosh^2(t)}\right) dt. \tag{3.31}$$

We now take the ratio of (3.30) and (3.31) and send $T_c/\mu \rightarrow 0$. By means of the dominated convergence theorem (note that the integrand in (3.31) behaves as t^{-3} for large t), we thus find the limit being given as the ratio of

$$\int_0^\infty \frac{1}{\cosh^2(t)} dt \quad \text{and} \quad \frac{1}{8} \int_0^\infty \left(\frac{\tanh(t)}{t^3} - \frac{1}{t^2 \cosh^2(t)}\right) dt.$$

While the former is easily evaluated as being equal to one, the latter is given by $7\zeta(3)/8\pi^2$ (see, e.g., [19, (3.333.3)]). This proves the claim. ■

With the aid of (3.25) and noting $f_j > 0$, the resulting differences (from adding and subtracting $|\Delta_0(\sqrt{\mu})|^j$) can be estimated as

$$\left| \int_{\mathbb{R}^d} f_j(p)(|\Delta_0(p)|^j - |\Delta_0(\sqrt{\mu})|^j) dp \right| \leq C |\Delta_0(\sqrt{\mu})|^j \int_{\mathbb{R}^d} f_j(p)(\min(|p| - \sqrt{\mu}, 1) + \lambda) dp$$

for $j = 2, 4$. These integrals can be treated analogously to (3.30) and (3.31) in Lemma 3.13 (for the $|p| - \sqrt{\mu}$ -term, note that f_j essentially concentrates around $|p| \approx \sqrt{\mu}$) and we find them to be smaller than the corresponding leading term

$$\int_{\mathbb{R}^d} f_j(p)|\Delta_0(\sqrt{\mu})|^j dp$$

in the limit $\lambda \rightarrow 0$ (and hence $T_c \rightarrow 0$). Therefore,

$$\begin{aligned} |\psi_{\text{GL}}| &= \frac{T_c}{\Delta_0(\sqrt{\mu})} \left(\frac{\int_{\mathbb{R}^d} f_2(p) \, dp}{\int_{\mathbb{R}^d} f_4(p) \, dp} \right)^{1/2} \cdot (1 + o_{\lambda \rightarrow 0}(1)) \\ &= C_{\text{univ}} \frac{T_c}{\Delta_0(\sqrt{\mu})} \cdot (1 + o_{\lambda \rightarrow 0}(1)), \end{aligned}$$

where we used Lemma 3.13 in the last step. As the GL functional (3.18) is entirely independent of the relative difference to the critical temperature $(T_c - T)/T_c = h^2$, it is clear that all the errors here hold uniform in the parameter h . This finishes the proof of Proposition 3.11.

3.5. Pure angular momentum for $d = 2$: Proof of Theorem 2.11

3.5.1. Part (a). The proof of Theorem 2.11 (a) is mostly the same as that of Theorem 2.4 (a). We sketch the argument here, highlighting the few differences.

The operator \mathcal{V}_μ . Using the Birman–Schwinger principle on the operator $K_{T_c} + \lambda V$ (as is done in [13, 21]), we find that, for sufficiently small λ , the lowest eigenvalue e_μ of \mathcal{V}_μ (recall (1.10)) is an eigenvalue for angular momentum ℓ_0 , since this is the angular momentum of the ground state(s) of $K_{T_c} + \lambda V$ by assumption. Further, since V is radial, the eigenfunctions of \mathcal{V}_μ all have a definite angular momentum. In particular, the first excited state has some angular momentum $\ell_1 \neq \ell_0$:

$$e_\mu^{(1)} = \inf_{u \perp u_{\pm \ell_0}} \langle u | \mathcal{V}_\mu | u \rangle = \langle u_{\pm \ell_1} | \mathcal{V}_\mu | u_{\pm \ell_1} \rangle,$$

with $u_{\pm \ell}(p) = (2\pi)^{-1/2} e^{\pm i \ell \varphi}$ the eigenfunctions of angular momentum ℓ . Here, φ denotes the angle of $p \in \mathbb{R}^2$ in polar coordinates. Note that $e_\mu^{(1)} \leq 0$, since \mathcal{V}_μ is a compact operator on an infinite-dimensional space.

A priori spectral information. It is proved in [9, Theorem 2.1] that there exists a temperature \tilde{T} such that for temperatures $\tilde{T} < T < T_c$ the minimizers of the BCS functional are given by

$$\hat{\alpha}_\pm(p) = e^{\pm i \ell_0 \varphi} \hat{\alpha}_0(p)$$

where φ denotes the angle of $p \in \mathbb{R}^2$ in polar coordinates, $\hat{\alpha}_0$ is a radial function, and ℓ_0 is the angular momentum given by (2.15). The BCS gap functions are then $\Delta_\pm(p) = \Delta_0(p) e^{\pm i \ell_0 \varphi}$, with Δ_0 a radial function.¹³ Further, $K_T^{\Delta_0} + \lambda V \geq 0$ for temperatures $T \in (\tilde{T}, T_c)$ [9, Proposition 4.3] and $\ker(K_T^{\Delta_0} + \lambda V) = \text{span}\{\alpha_+, \alpha_-\}$.

¹³This should not be confused with the function Δ_0 used in Section 3.4.

The temperature \tilde{T} . As discussed in [9, Remark 2.6] the temperature \tilde{T} is given by $\tilde{T} = T_c(\ell_1)$, the critical temperature when restricted to angular momentum ℓ_1 . Following the argument in [13] (see also [24, Theorem 1]), we find

$$\tilde{T} \leq \begin{cases} Ce^{1/\lambda e_\mu^{(1)}}, & e_\mu^{(1)} < 0, \\ Ce^{-c/\lambda^2}, & e_\mu^{(1)} = 0. \end{cases}$$

Recalling that $T_c \sim e^{1/\lambda e_\mu}$ and that $e_\mu < e_\mu^{(1)} \leq 0$, then clearly $\tilde{T}/T_c \leq Ce^{-c/\lambda}$ for some $c > 0$.

Weak a priori bound on Δ_\pm . Exactly as in (3.10), we have that $\|\Delta_\pm\|_{L^\infty} \leq C\lambda$.

Birman–Schwinger principle. As in Section 3.2.3, by the Birman–Schwinger principle, we have that

$$B_{T,\Delta_0} = V^{1/2} \frac{1}{K_T^{\Delta_0}} |V|^{1/2}$$

has -1 as its lowest eigenvalue; only the eigenspace is spanned by the two vectors $\phi_\pm = V^{1/2}\alpha_\pm$. By a completely analogous argument is in Section 3.2.3, we find that

$$S_{T,\Delta_0} = \lambda m(T, \Delta_0) \mathfrak{F}_\mu |V|^{1/2} \frac{1}{1 + \lambda V^{1/2} M_{T,\Delta_0} |V|^{1/2}} V^{1/2} \mathfrak{F}_\mu^\dagger$$

has -1 as its lowest eigenvalue with corresponding eigenspace spanned by $\mathfrak{F}_\mu V\alpha_\pm$.

A priori bounds on Δ_\pm . Analogously to Theorem 3.5 we claim the following.

Lemma 3.14. *The functions Δ_\pm satisfy the bounds (with slight abuse of notation, recall that Δ_0 is a radial function)*

$$|\Delta_\pm(p)| \leq C|\Delta_0(\sqrt{\mu})|, \quad |\Delta_\pm(p) - \Delta_\pm(q)| \leq C|\Delta_0(\sqrt{\mu})||p - q|.$$

Proof. The proof is analogous to that of Theorem 3.5. First, we note that $\mathcal{V}_\mu = \mathfrak{F}_\mu V \mathfrak{F}_\mu^\dagger$ has eigenfunctions of lowest eigenvalue $u_{\pm\ell_0}(p) = (2\pi)^{-1/2} e^{\pm i\ell_0\varphi}$ and that the operator S_{T,Δ_0} preserves the angular momentum. Analogously to the proof of Theorem 3.5, we find

$$\Delta_\pm(p) = f_\pm(\lambda) \left(\int_{\mathbb{S}^1} \hat{V}(p - \sqrt{\mu}q) u_{\pm\ell_0}(q) d\omega(q) + \lambda \eta_{\lambda\pm}(p) \right)$$

with $\|\eta_{\lambda\pm}\|_{L^\infty} \leq C$ uniformly in λ . Evaluating on the Fermi surface $\{p^2 = \mu\}$, we get (recall that $\inf \text{spec } \mathcal{V}_\mu = e_\mu$)

$$\Delta_\pm(\sqrt{\mu}p/|p|) = f_\pm(\lambda) (e_\mu u_{\pm\ell_0}(p/|p|) + \lambda \eta_{\lambda\pm}(\sqrt{\mu}p/|p|)).$$

In particular, we conclude that $|\Delta_0(\sqrt{\mu})| = |\Delta_{\pm}(\sqrt{\mu}p/|p|)| > 0$ for λ small enough and that $|f_{\pm}(\lambda)| \leq C|\Delta_0(\sqrt{\mu})|$. We conclude the rest of the proof exactly as for Theorem 3.5. ■

The remaining parts of the argument (first order analysis of m , the exponential vanishing of Δ_{\pm} , infinite order analysis of m , calculation of the integral $m(T, \Delta_0) - m(T_c, 0)$, and comparing Δ_{\pm} on the Fermi surface with Ξ) are exactly as in Sections 3.2.5–3.2.9, only replacing Δ and u by Δ_{\pm} and $u_{\pm\ell_0}$, respectively. This concludes the proof of Theorem 2.11 (a).

3.5.2. Part (b). Again, we highlight only the main differences compared to the proof of Theorem 2.4 (b).

Ginzburg–Landau functional. Since every function $\hat{\alpha}_0$ in kernel of $K_{T_c} + \lambda V$ can be written (in polar coordinates) as

$$\hat{\alpha}_0(p, \varphi) = \hat{\rho}(p)[\psi_+ e^{i\ell_0\varphi} + \psi_- e^{-i\ell_0\varphi}]$$

for an appropriate normalized $\hat{\rho} \in L^2((0, \infty); p dp)$ and $\psi_{\pm} \in \mathbb{C}$ by Assumption 2.10 (cf. (2.15)), the analog of the Ginzburg–Landau functional (3.18) becomes [16, Theorems 2.10 and 3.5]

$$\begin{aligned} \mathcal{E}_{\text{GL}}(\psi_+, \psi_-) &= [|\psi_+|^4 + |\psi_-|^4 + 4|\psi_+|^2|\psi_-|^2] \\ &\quad \times \left[\frac{2\pi}{T_c^3} \int_0^{\infty} \frac{g_1((p^2 - \mu)/T_c)}{(p^2 - \mu)/T_c} |K_{T_c}(p)|^4 |\hat{\rho}(p)|^4 p dp \right] \\ &\quad - [|\psi_+|^2 + |\psi_-|^2] \\ &\quad \times \left[\frac{\pi}{T_c} \int_0^{\infty} \frac{1}{\cosh^2((p^2 - \mu)/(2T_c))} |K_{T_c}(p)|^2 |\hat{\rho}(p)|^2 p dp \right]. \end{aligned} \tag{3.32}$$

Minimizers of the GL functional. In contrast to (3.18), the functional (3.32) now has *two* (up to a phase unique) minimizers. This follows from observing that

$$\mathcal{E}_{\text{GL}}(\psi_+, \psi_-) = \mathcal{E}_{\text{GL}}(\psi_-, \psi_+)$$

and that one of the ψ_{\pm} is necessarily zero for any minimizer of (3.32). In fact, these minimizers are

$$(|\psi_{\text{GL}}|, 0) \quad \text{and} \quad (0, |\psi_{\text{GL}}|),$$

with $|\psi_{\text{GL}}|$ given in (3.21), but with $\Delta_0 := -2(2\pi)^{-d/2} \lambda \hat{V} \star \hat{\rho}$, where $\hat{\rho}$ is understood as a radial function in $L^2(\mathbb{R}^2)$.¹⁴ Hence, using the notation from Proposition 3.9 (see

¹⁴The fact that $\Delta_0(\sqrt{\mu}) \neq 0$ can be seen in a similar way as in the proof of Lemma 3.14.

also [16, Theorem 2.10], which provides a general analog of (3.19)–(3.20), valid also for the concrete functional (3.32) and (3.22), we find that (up to a constant phase) every non-zero solution of the BCS gap equation (1.4) can be written as

$$\Delta_{\pm}(p, \varphi) = C_{\text{univ}} T_c h \frac{\Delta_0(p)}{\Delta_0(\sqrt{\mu})} e^{\pm i \ell_0 \varphi} (1 + o_{\lambda \rightarrow 0}(1) + O(h e^{c/\lambda})) + \Delta_{\xi}(p, \varphi).$$

The rest of the argument (a priori bounds on $\Delta_0(p) e^{\pm i \ell_0 \varphi}$ and Δ_{ξ} , comparison of $|\Delta_{\pm}|$ on the Fermi surface with Ξ) works completely analogously to Section 3.4, with similar adjustments as explained in Section 3.5.1. This concludes the proof of Theorem 2.11 (b).

A. Additional proofs

A.1. Uniqueness of the minimizer: Proof of Proposition 1.5

Finally, we present the proof of Proposition 1.5.

Proof of Proposition 1.5. We remark that the argument of the proof has already partly been sketched in [9, 25]. The key observation for our proof is that, if $\widehat{V} \leq 0$, then

$$\langle \widehat{\alpha} | \widehat{V} \star \widehat{\alpha} \rangle \geq \langle |\widehat{\alpha}| | \widehat{V} \star |\widehat{\alpha}| \rangle. \tag{A.1}$$

Let $(\widehat{\gamma}, \widehat{\alpha})$ minimize the BCS functional (1.3). Then, by means of (A.1), we have $\mathcal{F}_T[(\widehat{\gamma}, \widehat{\alpha})] \geq \mathcal{F}_T[(\widehat{\gamma}, |\widehat{\alpha}|)]$, hence also $(\widehat{\gamma}, |\widehat{\alpha}|)$ is a minimizer. Consequently, the (inverse) Fourier transform of $|\widehat{\alpha}|$ is an eigenvector of $K_T^{\Delta} + V$ with

$$\Delta = -2(2\pi)^{-d/2} \widehat{V} \star |\widehat{\alpha}|$$

with the eigenvalue zero. Note that, using continuity of Δ and the BCS gap equation (1.4), we not only have $|\widehat{\alpha}| \geq 0$ but also $|\widehat{\alpha}| > 0$ everywhere (see [21, Lemma 2.1]). By the observation (A.1) again, we find that, for any ground state $\widehat{\alpha}_{\text{GS}}$ of $K_T^{\Delta} + V$, also $|\widehat{\alpha}_{\text{GS}}|$ is a ground state. But $|\widehat{\alpha}_{\text{GS}}|$ is non-orthogonal to $|\widehat{\alpha}|$, which implies that zero has to be the lowest eigenvalue of $K_T^{\Delta} + V$, i.e.,

$$K_T^{\Delta} + V \geq 0. \tag{A.2}$$

By writing out (A.1), we see that the inequality is actually an application of Cauchy–Schwarz, and thus becomes strict, unless $\widehat{\alpha}(p) = e^{i\phi} |\widehat{\alpha}(p)|$ for some fixed $\phi \in \mathbb{R}$. Therefore, by repeating the above arguments, we find that the ground state of (A.2) is non-degenerate, and we have proven (i).

In order to prove (ii), let $\Gamma_i \equiv (\gamma_i, \alpha_i)$, $i = 1, 2$, be two (non-trivial) minimizers of the BCS functional (1.3), and denote the corresponding gap functions by Δ_1 resp. Δ_2 .

We now apply the relative entropy identity (see [14] and [16, Proposition 5.2]) and a simple trace inequality (see [14, Lemma 3] and [16, Lemma 5.7]) to find that

$$\mathcal{F}_T[\Gamma_1] - \mathcal{F}_T[\Gamma_2] \geq \langle (\alpha_1 - \alpha_2) | K_T^{\Delta_1} + V | (\alpha_1 - \alpha_2) \rangle \geq 0$$

(and the same inequality with indices 1 and 2 interchanged) by means of (A.2). Since $\mathcal{F}_T[\Gamma_1] = \mathcal{F}_T[\Gamma_2]$, this implies $(\alpha_1 - \alpha_2) \in \ker(K_T^{\Delta_1} + V)$, and thus $\alpha_2 = \psi_{21}\alpha_1$ for some $\psi_{21} \in \mathbb{C} \setminus \{0\}$ (recall from (i) that $\ker(K_T^{\Delta_1} + V)$ is one-dimensional). From this we conclude

$$(K_T^{\psi_{21}\Delta_1} + V)\alpha_1 = 0.$$

Now, the pointwise strict monotonicity of $|\psi_{21}| \mapsto K_T^{\psi_{21}\Delta_1}(p)$ together with the fact that one can choose $|\hat{\alpha}_1|$ to be strictly positive, implies that $|\psi_{21}| = 1$ and we have shown uniqueness of minimizers up to a constant phase, which can be chosen in such a way that it ensures strict positivity of $\hat{\alpha}$. Finally, it is shown in [9, Proposition 2.9] that if α is not radial, then (A.2) is violated. Radiality of the corresponding γ follows from (1.6). This finishes the proof. ■

A.2. Proofs of technical lemmas within the proof of Theorem 2.4

This section contains the proofs of Lemmas 3.4, 3.6, and 3.12.

A.2.1. Proof of Lemma 3.4. The argument is slightly different in dimensions $d = 1, 2, 3$. The case $d = 3$ is similar to [13, 21] and the case $d = 1, 2$ is similar to [28].

The case $d = 3$. We write

$$\begin{aligned} V^{1/2}M_{T,\Delta}|V|^{1/2} &= V^{1/2}\frac{1}{p^2}|V|^{1/2} + V^{1/2}\left(M_{T,\Delta} - \frac{1}{p^2}\chi_{|p|>\sqrt{2\mu}}\right)|V|^{1/2} \\ &\quad - V^{1/2}\frac{1}{p^2}\chi_{|p|\leq\sqrt{2\mu}}|V|^{1/2}. \end{aligned} \tag{A.3}$$

The first term in (A.3) has kernel (proportional to)

$$V(x)^{1/2}\frac{1}{|x-y|}|V(y)|^{1/2} \in L^2(\mathbb{R}^3 \times \mathbb{R}^3) \tag{A.4}$$

by the Hardy–Littlewood–Sobolev inequality [35, Theorem 4.3]. The kernel of the second term in (A.3) is given by

$$\begin{aligned} V(x)^{1/2}|V(y)|^{1/2} &\frac{1}{(2\pi)^3} \left[\int_{|p|<\sqrt{2\mu}} \frac{1}{K_T^\Delta(p)} (e^{ip(x-y)} - e^{i\sqrt{\mu}p/|p|(x-y)}) dp \right. \\ &\quad \left. + \int_{|p|>\sqrt{2\mu}} \left(\frac{1}{K_T^\Delta(p)} - \frac{1}{p^2} \right) e^{ip(x-y)} dp \right]. \end{aligned} \tag{A.5}$$

We compute the angular integral first. In the first term, the integral is

$$4\pi \int_0^{\sqrt{2\mu}} \frac{1}{K_T^\Delta(p)} \left[\frac{\sin |p||x-y|}{|p||x-y|} - \frac{\sin \sqrt{\mu}|x-y|}{\sqrt{\mu}|x-y|} \right] |p|^2 dp. \tag{A.6}$$

Here we bound $\left| \frac{\sin a}{a} - \frac{\sin b}{b} \right| \leq C \frac{|a-b|}{a+b}$ for $a, b > 0$. Thus, we get the bound

$$|(A.6)| \leq C \int_0^{\sqrt{2\mu}} \frac{1}{K_T^\Delta(p)} \frac{|p - \sqrt{\mu}|}{p + \sqrt{\mu}} p^2 dp \leq C \int_0^{\sqrt{2\mu}} \frac{1}{|p^2 - \mu|} \frac{|p - \sqrt{\mu}|}{p + \sqrt{\mu}} p^2 dp \leq C.$$

In the second term, the integral is (bounded by)

$$\begin{aligned} & 4\pi \int_{\sqrt{2\mu}}^\infty \left| \frac{1}{K_T^\Delta(p)} - \frac{1}{p^2} \right| \frac{|\sin |p||x-y||}{|p||x-y|} |p|^2 dp \\ & \leq \frac{4\pi}{|x-y|} \int_{\sqrt{2\mu}}^\infty \left| \frac{1}{K_T^\Delta(p)} - \frac{1}{p^2} \right| p dp. \end{aligned}$$

To bound the remaining integral, we bound

$$\begin{aligned} \left| \frac{1}{K_T^\Delta(p)} - \frac{1}{p^2} \right| & \leq \frac{\left| \tanh \frac{\sqrt{|p^2 - \mu|^2 + \Delta(p)^2}}{2T} - 1 \right|}{\sqrt{|p^2 - \mu|^2 + \Delta(p)^2}} \\ & \quad + \left| \frac{1}{\sqrt{|p^2 - \mu|^2 + \Delta(p)^2}} - \frac{1}{|p^2 - \mu|} \right| \\ & \quad + \left| \frac{1}{|p^2 - \mu|} - \frac{1}{p^2} \right|. \end{aligned}$$

Note first that $|\tanh x - 1| \leq 2e^{-2x}$. Thus, we have

$$\begin{aligned} & \int_{\sqrt{2\mu}}^\infty \frac{\left| \tanh \frac{\sqrt{|p^2 - \mu|^2 + \Delta(p)^2}}{2T} - 1 \right|}{\sqrt{|p^2 - \mu|^2 + \Delta(p)^2}} p dp \\ & \leq 2 \int_{\sqrt{2\mu}}^\infty e^{-\sqrt{|p^2 - \mu|^2 + \Delta(p)^2}/T} \frac{1}{\sqrt{|p^2 - \mu|^2 + \Delta(p)^2}} p dp \\ & \leq 2 \int_{\sqrt{2\mu}}^\infty e^{-|p^2 - \mu|/T} \frac{p}{|p^2 - \mu|} dp \leq CT. \end{aligned}$$

Next, we estimate

$$\begin{aligned}
 & \left| \frac{1}{\sqrt{|p^2 - \mu|^2 + \Delta(p)^2}} - \frac{1}{|p^2 - \mu|} \right| \\
 &= \frac{1}{|p^2 - \mu|} \frac{\Delta(p)^2}{\sqrt{|p^2 - \mu|^2 + \Delta(p)^2} (|p^2 - \mu| + \sqrt{|p^2 - \mu|^2 + \Delta(p)^2})} \\
 &\leq \frac{1}{|p^2 - \mu|} \frac{\|\Delta\|_{L^\infty}^2}{\sqrt{|p^2 - \mu|^2 + \|\Delta\|_{L^\infty}^2} (|p^2 - \mu| + \sqrt{|p^2 - \mu|^2 + \|\Delta\|_{L^\infty}^2})} \\
 &\leq \frac{\|\Delta\|_{L^\infty}^2}{2|p^2 - \mu|^3}, \tag{A.7}
 \end{aligned}$$

using pointwise monotonicity in $\Delta(p)$. Thus, changing variables to $u = p^2 - \mu$ we have

$$\int_{\sqrt{2\mu}}^{\infty} \left| \frac{1}{\sqrt{|p^2 - \mu|^2 + \Delta(p)^2}} - \frac{1}{|p^2 - \mu|} \right| p dp \leq \frac{1}{4} \|\Delta\|_{L^\infty}^2 \int_{\mu}^{\infty} \frac{1}{u^3} du \leq C \|\Delta\|_{L^\infty}^2.$$

Finally,

$$\int_{\sqrt{2\mu}}^{\infty} \left| \frac{1}{|p^2 - \mu|} - \frac{1}{p^2} \right| p dp \leq C.$$

We conclude that the kernel of the second term in (A.3) is bounded by

$$|V(x)|^{1/2} \left(\frac{1 + T + \|\Delta\|_{L^\infty}^2}{|x - y|} + 1 \right) |V(y)|^{1/2} \in L^2(\mathbb{R}^3 \times \mathbb{R}^3).$$

Finally, the last term of (A.3) has kernel

$$4\pi V(x)^{1/2} |V(y)|^{1/2} \int_0^{\sqrt{2\mu}} \frac{\sin p|x - y|}{p|x - y|} dp \in L^2(\mathbb{R}^3 \times \mathbb{R}^3) \tag{A.8}$$

since the integral is bounded by $\sqrt{2\mu}$.

The cases $d = 1$ and $d = 2$. The kernel of $M_{T,\Delta}$ is given by

$$\begin{aligned}
 & M_{T,\Delta}(x, y) \\
 &= \frac{1}{(2\pi)^d} \left[\int_{|p| < \sqrt{2\mu}} \frac{1}{K_T^\Delta} (e^{ip(x-y)} - e^{i\sqrt{\mu}p/|p|(x-y)}) + \int_{|p| > \sqrt{2\mu}} \frac{1}{K_T^\Delta} e^{ip(x-y)} \right]. \tag{A.9}
 \end{aligned}$$

Now, one may bound $K_T^\Delta \geq |p^2 - \mu|$ uniformly in T, Δ . Then, we may bound $M_{T,\Delta}$ exactly as in [28, Lemma 3.4]. That is, we have the bounds

$$\begin{aligned} & \|V^{1/2} M_{T,\Delta} |V|^{1/2}\|_{\text{HS}}^2 \\ & \lesssim \begin{cases} \|V\|_{L^1}^2 + \|V\|_{L^1} \int_{\mathbb{R}} |V(x)| [1 + \log(1 + \sqrt{\mu}|x|)]^2 dx, & d = 1, \\ \|V\|_{L^1}^2 + \|V\|_{L^p}^2, & d = 2, \end{cases} \end{aligned}$$

for any $1 < p \leq 4/3$. This concludes the proof of Theorem 3.4.

A.2.2. Proof of Lemma 3.6. To bound the difference, first note that, by computing the angular integrals, we have

$$\begin{aligned} & [M_{T,\Delta} - M_{T_c,0}](x, y) \\ & = \frac{|\mathbb{S}^{d-1}|}{(2\pi)^d} \left[\int_0^{\sqrt{2\mu}} \left(\frac{1}{K_T^\Delta} - \frac{1}{K_{T_c}} \right) (j_d(p|x-y|) - j_d(\sqrt{\mu}|x-y|)) p^{d-1} dp \right. \\ & \quad \left. + \int_{\sqrt{2\mu}}^\infty \left(\frac{1}{K_T^\Delta} - \frac{1}{K_{T_c}} \right) j_d(p|x-y|) p^{d-1} dp \right] \tag{A.10} \end{aligned}$$

where

$$j_d(x) = \frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} e^{ix\omega} d\omega = \begin{cases} \cos x, & d = 1, \\ J_0(|x|), & d = 2, \\ \frac{\sin|x|}{|x|}, & d = 3, \end{cases}$$

with J_0 the zero'th Bessel function.

Bounding (A.10) is similar in spirit to the proof of Theorem 3.4 above. We bound

$$\left| \frac{1}{K_T^\Delta} - \frac{1}{K_{T_c}} \right| \leq \frac{|\tanh(\frac{E_\Delta}{2T}) - 1|}{E_\Delta} + \left| \frac{1}{E_\Delta} - \frac{1}{|p^2 - \mu|} \right| + \frac{|1 - \tanh(\frac{|p^2 - \mu|}{2T_c})|}{|p^2 - \mu|}.$$

We bound the first term as follows:

$$\begin{aligned} \frac{|\tanh(\frac{E_\Delta}{2T}) - 1|}{E_\Delta} & \leq 2e^{-E_\Delta/T} \frac{1}{E_\Delta} \\ & \leq 2e^{-|p^2 - \mu|/T} \frac{1}{|p^2 - \mu|} \\ & \leq 2e^{-|p^2 - \mu|/T_c} \frac{1}{|p^2 - \mu|}. \end{aligned}$$

Similarly,

$$\frac{|1 - \tanh(\frac{|p^2 - \mu|}{2T_c})|}{|p^2 - \mu|} \leq 2e^{-|p^2 - \mu|/T_c} \frac{1}{|p^2 - \mu|}.$$

Finally, we estimate, exactly as in (A.7) in the course of proving Theorem 3.4,

$$\begin{aligned} & \left| \frac{1}{E_\Delta} - \frac{1}{|p^2 - \mu|} \right| \\ & \leq \frac{1}{|p^2 - \mu|} \frac{\|\Delta\|_{L^\infty}^2}{\sqrt{|p^2 - \mu|^2 + \|\Delta\|_{L^\infty}^2} (|p^2 - \mu| + \sqrt{|p^2 - \mu|^2 + \|\Delta\|_{L^\infty}^2})} \\ & \leq \frac{\|\Delta\|_{L^\infty}^2}{2|p^2 - \mu|^3}. \end{aligned}$$

We will use the first bound for the first integral in (3.13) and the second bound for the second integral in (3.13). Note further that

$$\frac{1}{\sqrt{x^2 + A^2}(x + \sqrt{x^2 + A^2})}$$

is decreasing in x and $|p^2 - \mu| \geq \sqrt{\mu}|p - \sqrt{\mu}|$. That is, we have the bound

$$\begin{aligned} & \left| \frac{1}{K_T^\Delta} - \frac{1}{K_{T_c}} \right| \\ & \leq 4e^{-|p^2 - \mu|/T_c} \frac{1}{|p^2 - \mu|} + \chi_{|p| > \sqrt{2\mu}} \frac{\|\Delta\|_{L^\infty}^2}{2|p^2 - \mu|^3} \\ & \quad + \chi_{|p| < \sqrt{2\mu}} \frac{1}{\sqrt{\mu}|p - \sqrt{\mu}|} \\ & \quad \times \frac{\|\Delta\|_{L^\infty}^2}{\sqrt{|\mu|p - \sqrt{\mu}|^2 + \|\Delta\|_{L^\infty}^2} (\sqrt{\mu}|p - \sqrt{\mu}| + \sqrt{|\mu|p - \sqrt{\mu}|^2 + \|\Delta\|_{L^\infty}^2})}. \end{aligned} \tag{A.11}$$

In the first integral, in (A.10) we bound $|j_a(a) - j_a(b)| \leq C|a - b|$. Then, the contribution of the first term of (A.11) to the first integral in (A.10) is bounded by (changing variables to $s = \sqrt{\mu}|p - \sqrt{\mu}|/T_c$)

$$\begin{aligned} & \int_0^{\sqrt{2\mu}} e^{-|p^2 - \mu|/T_c} \frac{1}{|p^2 - \mu|} |p - \sqrt{\mu}| |x - y| p^{d-1} dp \\ & \leq C|x - y| \int_0^{\sqrt{2\mu}} e^{-\sqrt{\mu}|p - \sqrt{\mu}|/T_c} dp \\ & \leq T_c|x - y| \int_0^{\mu/T_c} e^{-s} ds \\ & \leq CT_c|x - y|. \end{aligned}$$

Next, the contribution of the last term of (A.11) is bounded by (changing variables to $s = \sqrt{\mu}|p - \sqrt{\mu}|/\|\Delta\|_{L^\infty}$)

$$\begin{aligned} & \int_0^{\sqrt{2\mu}} \frac{\|\Delta\|_{L^\infty}^2}{\sqrt{\mu}|p - \sqrt{\mu}|^2 + \|\Delta\|_{L^\infty}^2} (|p - \sqrt{\mu}|\sqrt{\mu} + \sqrt{\mu}|p - \sqrt{\mu}|^2 + \|\Delta\|_{L^\infty}^2) \\ & \qquad \qquad \qquad \times \frac{|p - \sqrt{\mu}||x - y|p^{d-1}}{|p - \sqrt{\mu}|\sqrt{\mu}} dp \\ & \leq C\|\Delta\|_{L^\infty}|x - y| \int_0^{\mu/\|\Delta\|_{L^\infty}} \frac{1}{\sqrt{s^2 + 1}(s + \sqrt{s^2 - 1})} ds \\ & \leq C\|\Delta\|_{L^\infty}|x - y|. \end{aligned}$$

Next, we estimate the last integral of (A.10). Here we note that $|j_d(a)| \leq C$. Then, the contributions of the first and second term in (A.11) to (A.10) is bounded by

$$\begin{aligned} \int_{\sqrt{2\mu}}^\infty e^{-|p^2 - \mu|/T_c} \frac{1}{|p^2 - \mu|} p^{d-1} dp & \leq C \int_{\sqrt{2\mu}}^\infty e^{-|p^2 - \mu|/T_c} |p^2 - \mu|^{d/2-2} p dp \\ & \leq CT_c e^{-\mu/T_c} \leq CT_c \end{aligned}$$

and

$$\int_{\sqrt{2\mu}}^\infty \frac{\|\Delta\|_{L^\infty}^2}{|p^2 - \mu|^3} p^{d-1} dp \leq C\|\Delta\|_{L^\infty}^2.$$

We conclude that (using $\|\Delta\|_{L^\infty} \leq CT_c$)

$$\begin{aligned} \|V^{1/2}(M_{T,\Delta} - M_{T_c,0})|V|^{1/2}\|_{\text{HS}}^2 & \leq CT_c^2 \iint |V(x)||V(y)|(\mu|x - y|^2 + 1) dx dy \\ & \leq CT_c^2(\mu\| |\cdot|^2 V \|_1 \|V\|_1 + \|V\|_1^2) \leq Ce^{-c/\lambda} \end{aligned}$$

by assumption on V . This finishes the proof of Theorem 3.6.

A.2.3. Proof of Lemma 3.12. The proof is very similar to the ones of [21, Lemma 4] and [27, Lemma 13], and follows from a Birman–Schwinger argument analogously to Sections 3.2.3 and 3.2.4.

First of all, recall from Proposition 1.5 (ii), that $K_{T_c} + \lambda V$ has 0 as a (non-degenerate) ground state eigenvalue, which, by the Birman–Schwinger principle, is equivalent to the fact that the Birman–Schwinger operator $B_{T_c} := \lambda V^{1/2} K_{T_c}^{-1} |V|^{1/2}$ has -1 as its (non-degenerate) ground state eigenvalue. As in Section 3.2.3, defining $m(T_c) :=$

$m(T_c, 0)$ (recall (3.11)), we decompose B_{T_c} as

$$B_{T_c} = \lambda m(T_c) V^{1/2} \mathfrak{F}_\mu^\dagger \mathfrak{F}_\mu |V|^{1/2} + \lambda V^{1/2} M_{T_c} |V|^{1/2},$$

where M_{T_c} is such that this holds. It has been shown in [13, Lemma 2] (for $d = 3$) and [28, Lemma 3.4] (for $d = 1, 2$) that the Hilbert–Schmidt norm $\|V^{1/2} M_{T_c} |V|^{1/2}\|_{\text{HS}}$ of the second term is uniformly bounded for small T_c (i.e., small λ).

Then, by an argument completely analogous to the one in the proof of Lemma 3.5 in Section 3.2.4, we find that $\Delta_0 = f(\lambda)[\hat{\varphi} + \lambda\eta_\lambda]$, with $\hat{\varphi}$ defined in (3.23) and η_λ has $\|\eta_\lambda\|_{L^\infty} \leq C$ uniformly in small λ (cf. (3.24)). This concludes the proof of Theorem 3.12.

Acknowledgments. We thank Andreas Deuchert, Christian Hainzl, Edwin Langmann, Marius Lemm, Robert Seiringer, and Jan Philip Solovej for helpful discussions, and Edwin Langmann and Robert Seiringer for valuable comments on an earlier version of the manuscript.

Funding. Joscha Henheik gratefully acknowledges partial financial support by the ERC Advanced Grant “RMTBeyond” No. 101020331. Asbjørn Bækgaard Lauritsen gratefully acknowledges partial financial support by the Austrian Science Fund (FWF) through grant DOI [10.55776/I6427](https://doi.org/10.55776/I6427) (as part of the SFB/TRR 352).

References

- [1] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, [Theory of superconductivity](#). *Phys. Rev.* (2) **108** (1957), 1175–1204 [Zbl 0090.45401](#) [MR 0095694](#)
- [2] L.S. Borkowski and P.J. Hirschfeld, [Distinguishing \$d\$ -wave superconductors from highly anisotropic \$s\$ -wave superconductors](#). *Phys. Rev. B* **49** (1994), no. 21, 15404–15407
- [3] G. Bräunlich, C. Hainzl, and R. Seiringer, [Translation-invariant quasi-free states for fermionic systems and the BCS approximation](#). *Rev. Math. Phys.* **26** (2014), no. 7, article no. 1450012 [Zbl 1327.82099](#) [MR 3246962](#)
- [4] G. Bräunlich, C. Hainzl, and R. Seiringer, [On the BCS gap equation for superfluid fermionic gases](#). In *Mathematical results in quantum mechanics*, pp. 127–137, World Sci. Publ., Hackensack, NJ, 2015 [Zbl 1343.82059](#) [MR 3381143](#)
- [5] M. J. Buckingham, [Very high frequency absorption in superconductors](#). *Phys. Rev.* **101** (1956), 1431–1432
- [6] C. Cohen-Tannoudji, B. Diu, and F. Laloe, *Quantum mechanics*. Vol. 2. Second edn., Wiley-VCH, Weinheim, 2020.
- [7] J.-C. Cuenin and K. Merz, [Weak coupling limit for Schrödinger-type operators with degenerate kinetic energy for a large class of potentials](#). *Lett. Math. Phys.* **111** (2021), no. 2, article no. 46 [Zbl 1464.58002](#) [MR 4241156](#)

- [8] P.-G. De Gennes, *Superconductivity of metals and alloys*. W. A. Benjamin, New York and Amsterdam, 1966 Zbl [0138.22801](#)
- [9] A. Deuchert, A. Geisinger, C. Hainzl, and M. Loss, [Persistence of translational symmetry in the BCS model with radial pair interaction](#). *Ann. Henri Poincaré* **19** (2018), no. 5, 1507–1527 Zbl [1393.82028](#) MR [3784920](#)
- [10] A. Deuchert, C. Hainzl, and M. O. Maier, [Microscopic derivation of Ginzburg-Landau theory and the BCS critical temperature shift in a weak homogeneous magnetic field](#). *Probab. Math. Phys.* **4** (2023), no. 1, 1–89 Zbl [1520.35147](#) MR [4567398](#)
- [11] A. Deuchert, C. Hainzl, and M. O. Maier, [Microscopic derivation of Ginzburg-Landau theory and the BCS critical temperature shift in general external fields](#). *Calc. Var. Partial Differential Equations* **62** (2023), no. 7, article no. 203 Zbl [1522.35483](#) MR [4621520](#)
- [12] J. D. Fletcher, A. Serafin, L. Malone, J. G. Analytis, J.-H. Chu, A. S. Erickson, I. R. Fisher, and A. Carrington, [Evidence for a nodal-line superconducting state in LaFePO](#). *Phys. Rev. Lett.* **102** (2009), no. 14, article no. 147001
- [13] R. L. Frank, C. Hainzl, S. Naboko, and R. Seiringer, [The critical temperature for the BCS equation at weak coupling](#). *J. Geom. Anal.* **17** (2007), no. 4, 559–567 Zbl [1137.82025](#) MR [2365659](#)
- [14] R. L. Frank, C. Hainzl, R. Seiringer, and J. P. Solovej, [Microscopic derivation of Ginzburg–Landau theory](#). *J. Amer. Math. Soc.* **25** (2012), no. 3, 667–713 Zbl [1251.35156](#) MR [2904570](#)
- [15] R. L. Frank, C. Hainzl, R. Seiringer, and J. P. Solovej, [The external field dependence of the BCS critical temperature](#). *Comm. Math. Phys.* **342** (2016), no. 1, 189–216 Zbl [1334.82080](#) MR [3455149](#)
- [16] R. L. Frank and M. Lemm, [Multi-component Ginzburg-Landau theory: microscopic derivation and examples](#). *Ann. Henri Poincaré* **17** (2016), no. 9, 2285–2340 Zbl [1351.82117](#) MR [3535864](#)
- [17] V. L. Ginzburg and L. D. Landau, On the theory of superconductivity. *Zh. Eksp. Teor. Fiz.* **20** (1950), 1064–1082. English translation in V. L. Ginzburg, *On superconductivity and superfluidity*, pp. 113–137, Springer, Berlin etc., 2009 Zbl [1541.82017](#)
- [18] L. P. Gor’kov, Microscopic derivation of the Ginzburg–Landau equations in the theory of superconductivity (in Russian). *Zhurn. Eksper. Teor. Fiz.* **36** (1959), 1918–1923. English translation: *Sov. Phys. JETP* **9** (1959), no. 6, 1364–1367 Zbl [0088.45602](#)
- [19] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*. Seventh edn., Academic Press, Boston, 2007 Zbl [1208.65001](#) MR [2360010](#)
- [20] C. Hainzl, E. Hamza, R. Seiringer, and J. P. Solovej, [The BCS functional for general pair interactions](#). *Comm. Math. Phys.* **281** (2008), no. 2, 349–367 Zbl [1161.82027](#) MR [2410898](#)
- [21] C. Hainzl and R. Seiringer, [Critical temperature and energy gap for the BCS equation](#). *Phys. Rev. B* **77** (2008), no. 18, article no. 184517 Zbl [0801.4159](#)
- [22] C. Hainzl and R. Seiringer, [The BCS critical temperature for potentials with negative scattering length](#). *Lett. Math. Phys.* **84** (2008), no. 2-3, 99–107 Zbl [1164.81006](#) MR [2415542](#)

- [23] C. Hainzl and R. Seiringer, [Spectral properties of the BCS gap equation of superfluidity](#). In *Mathematical results in quantum mechanics*, pp. 117–136, World Scientific, Hackensack, NJ, 2008 Zbl [1156.82413](#) MR [2466682](#)
- [24] C. Hainzl and R. Seiringer, [Asymptotic behavior of eigenvalues of Schrödinger type operators with degenerate kinetic energy](#). *Math. Nachr.* **283** (2010), no. 3, 489–499 Zbl [1193.58003](#) MR [2643024](#)
- [25] C. Hainzl and R. Seiringer, [The Bardeen–Cooper–Schrieffer functional of superconductivity and its mathematical properties](#). *J. Math. Phys.* **57** (2016), no. 2, article no. 021101 Zbl [1360.81315](#) MR [3463277](#)
- [26] J. Henheik, [The BCS critical temperature at high density](#). *Math. Phys. Anal. Geom.* **25** (2022), no. 1, article no. 3 Zbl [1484.81035](#) MR [4366075](#)
- [27] J. Henheik and A. B. Lauritsen, [The BCS energy gap at high density](#). *J. Stat. Phys.* **189** (2022), no. 1, article no. 5 Zbl [1492.81055](#) MR [4458584](#)
- [28] J. Henheik, A. B. Lauritsen, and B. Roos, [Universality in low-dimensional BCS theory](#). *Rev. Math. Phys.* **36** (2024), no. 9, article no. 2360005 Zbl [07938498](#) MR [4802022](#)
- [29] L. D. Landau, On the theory of phase transitions. *Zh. Eksp. Teor. Fiz.* **7** (1937), 19–32
- [30] E. Langmann and C. Triola, [. Phys. Rev. B](#) **108** (2023), no. 10, article no. 104503
- [31] E. Langmann, C. Triola, and A. V. Balatsky, [Ubiquity of superconducting domes in the Bardeen–Cooper–Schrieffer theory with finite-range potentials](#). *Phys. Rev. Lett.* **122** (2019), no. 15, article no. 157001
- [32] A. B. Lauritsen, [The BCS energy gap at low density](#). *Lett. Math. Phys.* **111** (2021), no. 1, article no. 20 Zbl [1467.82103](#) MR [4218360](#)
- [33] A. J. Leggett, [Diatomic molecules and Cooper pairs](#). In *Modern trends in the theory of condensed matter*, pp. 13–27 Lecture Notes in Phys. 115, Springer, Berlin etc., 1980 MR [0583566](#)
- [34] A. J. Leggett, *Quantum liquids: Bose condensation and Cooper pairing in condensed-matter systems*. Oxford university press, Oxford, 2006 Zbl [1509.82002](#)
- [35] E. H. Lieb and M. Loss, *Analysis*. Second edn., Grad. Stud. Math. 14, American Mathematical Society, Providence, RI, 2001 Zbl [0966.26002](#) MR [1817225](#)
- [36] K. Maki and H. Won, Why *d*-wave superconductivity? *J. Physique I France* **6** (1996), no. 12, 2317–2326
- [37] P. Nozières and S. Schmitt-Rink, [Bose condensation in an attractive fermion gas: From weak to strong coupling superconductivity](#). *J. Low Temp. Phys.* **59** (1985), 195–211
- [38] C. P. Poole, H. A. Farach, R. J. Creswick, and R. Prozorov, *Superconductivity*. Second edn., Elsevier, Amsterdam, 2007
- [39] M. Tinkham, *Introduction to superconductivity*. Dover Publications, Mineola, NY, 2004
- [40] Y Zhang, ZR Ye, QQ Ge, F Chen, Juan Jiang, M Xu, BP Xie, and DL Feng, [Nodal superconducting-gap structure in ferropnictide superconductor BaFe₂\(As_{0.7}P_{0.3}\)₂](#). *Nature Physics* **8** (2012), no. 5, 371–375

Received 2 August 2024.

Joscha Henheik

Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria;
joscha.henheik@ist.ac.at

Asbjørn Bækgaard Lauritsen

Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria;
current address: CNRS & CEREMADE, Université Paris–Dauphine, PSL University,
Place du Maréchal de Lattre de Tassigny, 75775 Paris Cedex 16, France;
asbjorn.lauritsen@dauphine.psl.eu