

Noncommutative scheme theory and the Serre–Artin–Zhang–Verevkin theorem for semi-graded rings

Andrés Chacón and Armando Reyes

Abstract. In this paper, we present a noncommutative scheme theory for the semi-graded rings generated in degree one defined by Lezama and Latorre [Internat. J. Algebra Comput. 27 (2017), 361–389] following the ideas about schematicity introduced by Van Oystaeyen and Willaert [J. Pure Appl. Algebra 104 (1995), 109–122] for \mathbb{N} -graded algebras. With this theory, we prove the Serre–Artin–Zhang–Verevkin theorem for several families of non- \mathbb{N} -graded algebras and finitely non- \mathbb{N} -graded algebras appearing in ring theory and noncommutative algebraic geometry. Our treatment contributes to the research on this theorem presented by Lezama from a different point of view.

Dedicated to Professor Lorenzo Acosta

1. Introduction

In his beautiful paper [42], Serre proved a theorem that describes the coherent sheaves on a projective scheme in terms of graded modules. Briefly, a commutative graded \mathbb{k} -algebra is associated to a projective scheme $\text{Proj} A$, and the geometry of this scheme can be described in terms of the quotient category $\text{qgr} A = \text{gr} A / \text{tors}$, where $\text{gr} A$ denotes the category of graded modules and tors denotes its subcategory of torsion modules. For A a finitely generated commutative graded \mathbb{k} -algebra and X its associated projective scheme, if $\text{coh} X$ denotes the category of coherent sheaves on X and $\mathcal{O}_X(n)$ is the n th power of the twisting sheaf on X [20, page 117], then we have a functor $\Gamma_* : \text{coh} X \rightarrow \text{qgr} A$ given by

$$\Gamma_*(\mathcal{F}) = \bigoplus_{d=-\infty}^{\infty} H^0(X, \mathcal{F} \otimes \mathcal{O}_X(d)).$$

Serre’s theorem [42, Section 59, Proposition 7.8, page 252], [19, 3.3.5], and [20, Proposition II. 5.15], asserts that if A is generated over \mathbb{k} by elements of degree one, then Γ_* defines an equivalence of categories $\text{coh} X \rightarrow \text{qgr} A$.

Artin and Zhang [5] extended Serre’s theorem to the noncommutative setting in the following way: let A be an \mathbb{N} -graded algebra over a commutative Noetherian ring. They defined the associated projective scheme to be the pair $\text{Proj} A = (\text{qgr} A, \mathcal{A})$, where $\text{qgr} A$ is the quotient category above, \mathcal{A} is the image of A in $\text{qgr} A$ and plays the role of the structure sheaf of $\text{Proj} A$, and s , called the *polarization* defined by the projective embedding (this definition is the same as is given by Verevkin [56, 57]), is given by the shifting of the degrees in $\text{gr} A$. Since Serre’s theorem does not hold for all commutative graded algebras, i.e., the functor defined by Γ_* need not be an equivalence, Artin and Zhang’s definition of $\text{Proj} A$ is compatible with the classical definition for commutative graded rings only under some additional hypotheses such that A is generated in degree one. In the literature, the noncommutative version of Serre’s theorem is known as *Serre–Artin–Zhang–Verevkin theorem* [5, 56, 57]. Several authors have investigated the results of commutative algebraic geometry, but now in the noncommutative setting following Artin, Zhang, and Verevkin’s ideas (e.g., [12, 26–28, 41, 43, 53–55, 59] and references therein).

On the other hand, Manin [30] commented on the failure of attempts to obtain a non-commutative scheme theory à la Grothendieck for quantized algebras. Nevertheless, Van Oystaeyen and Willaert [33] studied this Proj by developing a kind of scheme theory similar to the commutative theory. They noticed that this theory is possible only if the connected and \mathbb{N} -graded algebra considered contains “enough” Ore sets. Algebras satisfying this condition are called *schematic*. They constructed a *generalized Grothendieck topology* for the free monoid on all Ore sets of a schematic algebra R , and defined a *non-commutative site* (cf. [51]) as a category with coverings on which sheaves can be defined, and formulated the Serre’s theorem. As a consequence of their treatment, an equivalence between the category of all coherent sheaves and the category $\text{Proj} R$ was obtained in the sense of Artin [3]. Some years later, Van Oystaeyen and Willaert [50–52] presented a sequel of [33] in which they studied the cohomology of these algebras and proved a lifting property for Ore sets. This allowed to present many examples of schematic algebras like homogenizations of almost commutative algebras, Rees rings of universal enveloping algebras of Lie algebras, and three-dimensional Sklyanin algebras. A detailed treatment of schematic algebras can be found in Van Oystaeyen’s book [49].

A few years ago, Lezama and Latorre [29] introduced the *semi-graded rings* with the aim of generalizing the \mathbb{N} -graded rings, the finitely \mathbb{N} -graded algebras and several algebras appearing in ring theory and noncommutative geometry that are not \mathbb{N} -graded algebras in a non-trivial sense. In that paper, they investigated some geometrical properties of semi-graded rings, within which is the Serre–Artin–Zhang–Verevkin theorem following Artin, Zhang, and Verevkin’s ideas (see also [27]). In this way, having in mind Van Oystaeyen and Willaert’s ideas developed in [33] about a scheme theory for Proj in the setting of \mathbb{N} -graded algebras, it is natural to ask by a noncommutative scheme theory for semi-graded rings, and hence, to investigate the *schematicness* of these objects in a more general context than \mathbb{N} -graded rings. This is the purpose of the paper. As expected, we generalize the results established in [33] for \mathbb{N} -graded algebras to the semi-graded setting (as a matter of fact, we do not impose the condition of connectedness of the algebra),

and present another approach (Examples 5.27 and 5.28 show that the theory presented by Lezama [27, 29] and the one developed in this paper are independent) to the Serre–Artin–Zhang–Verevkin theorem for semi-graded rings which is a fundamental problem proposed for these objects [27, Section 1.4, Problem 1].

The article is organized as follows. In Section 2, we recall some key facts about torsion theory, Serre’s theorem in the commutative case and the noncommutative setting of \mathbb{N} -graded rings following the ideas presented by Artin and Zhang [5] and Van Oystaeyen and Willaert [33, 52]. Next, in Section 3, we consider some preliminaries about semi-graded rings and semi-graded modules, and present some facts about the localization of these objects. In Section 4, we formulate the notion of *schematicness* (Definition 4.1) for semi-graded rings without the assumption of connectedness established by Van Oystaeyen and Willaert for \mathbb{N} -graded rings. Section 5 contains the definition of *noncommutative site* with the aim of establishing the Serre–Artin–Zhang–Verevkin theorem in the semi-graded setting (Theorem 5.23). Our results generalize those corresponding in the case of \mathbb{N} -graded rings (Remark 5.24) and allow to guarantee that other non- \mathbb{N} -graded algebras (even not connected) to be schematic (Example 5.25). As we said above, Examples 5.27 and 5.28 show that the theory presented by Lezama about Serre–Artin–Zhang–Verevkin theorem and the one developed in this paper are independent. Finally, in Section 6, we present some ideas for a future work that are motivated by different topics concerning schematic algebras [49, 50, 52, 58].

Throughout the paper, the term ring means an associative ring with identity not necessarily commutative. The letter \mathbb{k} denotes an arbitrary field, and all algebras are \mathbb{k} -algebras. The symbols \mathbb{N} and \mathbb{Z} denote the set of natural numbers including zero, and the ring of integer numbers, respectively. For a ring R and a subset I of R , $I \triangleleft_l R$ means that I is a left ideal of R . $Z(R)$ denotes the center of R , while the category of left R -modules is written as $R - \text{Mod}$.

2. Serre’s theorem and schematic algebras

We recall briefly some notions of algebraic geometry which are key in the proof of *Serre’s theorem*.

Following Hartshorne [20], if $C = \mathbb{k} \oplus C_1 \oplus C_2 \oplus \cdots$ is a positively graded commutative Noetherian ring generated in degree one, consider $Y = \text{Proj} C$ and $Y(f) = \{\mathfrak{p} \in Y \mid f \notin \mathfrak{p}\}$, the Zariski open set corresponding to a homogeneous element $f \in C$. It is well-known that there is a finite subset $\{f_i \mid f_i \in C_1\}$ such that $Y = \bigcup_i Y(f_i)$. Equivalently, for every choice of $d_i \in \mathbb{N}$, there exists $n \in \mathbb{N}$ with $(C_+)^n \subseteq \sum_i C f_i^{d_i}$. In this way, for any finitely generated graded C -module M , we have that

$$\begin{aligned} \Gamma_*(M) &:= \bigoplus_{n \in \mathbb{Z}} \Gamma(Y, \widetilde{M(n)}) \\ &= Q_{\kappa_+}(M) = \varprojlim_i Q_{f_i}(M), \end{aligned}$$

where $\widetilde{M(n)}$ denotes the sheaf of modules associated to the shifted module $M(n)$, $Q_{\kappa+}$ is the module of quotients of M with respect to the classical torsion of modules in the \mathbb{N} -graded case, and $Q_{f_i}(M)$ is the localization of M at $\{1, f_i, f_i^2, \dots\}$. Of course,

$$Q_f(M) = \varprojlim_i Q_{f f_i}(M),$$

where the inverse systems are defined as $g \leq h$ if and only if $Y(g) \subseteq Y(h)$. This is precisely the key fact to prove *Serre's theorem*: the category of coherent \mathcal{O}_Y -modules is equivalent with a certain quotient category.

In the noncommutative setting, for a *noncommutative positively graded Noetherian \mathbb{k} -algebra* $R = \mathbb{k} \oplus R_1 \oplus R_2 \oplus \dots$ with $R = \mathbb{k}[R_1]$ (notice that R is connected, that is, $R_0 = \mathbb{k}$), Van Oystaeyen and Willaert [33] presented their interpretation of Serre's theorem for algebras with enough Ore sets called *schematic algebras*. With the aim of presenting the key ideas developed by them, we start by recalling some notions of torsion theory that we will use freely throughout the paper. For more details, we refer to Goldman [16], Stenstrom [44] or Van Oystaeyen [48].

Definition 2.1 ([33, Section 2]). Let \mathcal{L} be a set of left ideals of an arbitrary ring R . \mathcal{L} is said to be a *filter* if it satisfies the following conditions:

- (T₁) if $I \in \mathcal{L}$ and $I \subseteq J$, then $J \in \mathcal{L}$,
- (T₂) if $I, J \in \mathcal{L}$, then $I \cap J \in \mathcal{L}$,
- (T₃) if $I \in \mathcal{L}$ and $a \in R$, then $(I : a) := \{r \in R \mid ra \in I\} \in \mathcal{L}$.

The functor $\kappa : R - \text{Mod} \rightarrow R - \text{Mod}$ defined by

$$\kappa(M) = \{m \in M \mid \text{there exists } I \in \mathcal{L} \text{ with } \text{Im} = 0\}$$

is a *left exact preradical*, that is, a left exact subfunctor of the identity functor on the category $R - \text{Mod}$. A module M satisfying $\kappa(M) = M$ is called a *κ -torsion module*, and if $\kappa(M) = 0$, then M is said to be a *κ -torsion-free module*. It is straightforward to see that the family of torsion modules are closed under quotient objects and coproducts, while the torsion-free modules are closed under subobjects and products.

The filter \mathcal{L} is called *idempotent* (also called a *Gabriel topology*) when it satisfies the following condition.

- (T₄) If $I \triangleleft_l R$ and there exists $J \in \mathcal{L}$ such that for all $a \in J$ the relation $(I : a) \in \mathcal{L}$ holds, then $I \in \mathcal{L}$.

Condition (T₄) implies that \mathcal{L} is closed under products and that the functor κ is *radical*, that is, $\kappa(M/\kappa(M)) = 0$, for all $M \in R - \text{Mod}$.

Proposition 2.2. *If R is a left Noetherian ring and $J_1 \supseteq J_2 \supseteq \dots$ is a descending chain of two-sided ideals of R , then the set*

$$\mathcal{A} = \{I \triangleleft_l R \mid \text{there exist elements } n, m \in \mathbb{N} \text{ with } (J_m)^n \subseteq I\}$$

is an idempotent filter.

Proof. (T₁) If $(J_m)^n \subseteq I$ and $I \subseteq I'$, then it is clear that $(J_m)^n \subseteq I'$.

(T₂) If $I, I' \in \mathcal{A}$, then there exist elements $n_1, n_2, m_1, m_2 \in \mathbb{N}$ such that $(J_{m_1})^{n_1} \subseteq I$ and $(J_{m_2})^{n_2} \subseteq I'$. If we consider $n := \max\{n_1, n_2\}$ and $m := \max\{m_1, m_2\}$, it follows that $(J_m)^n \subseteq I \cap I'$.

(T₃) If $I \in \mathcal{A}$, then there exist elements $n, m \in \mathbb{N}$ such that $(J_m)^n \subseteq I$. Fix $a \in R$ and let $r \in (J_m)^n$. Since $(J_m)^n$ is an ideal of R , then $ra \in (J_m)^n \subseteq I$, and so, $r \in (I : a)$, that is, $(J_m)^n \subseteq (I : a)$.

(T₄) Let $I \triangleleft_l R$ and $J \in \mathcal{A}$ such that for all $a \in J$, we have that $(I : a) \in \mathcal{A}$. Since $J \in \mathcal{A}$ there exist elements $n, m \in \mathbb{N}$ with $(J_m)^n \subseteq J$. By assumption, R is left Noetherian, so $(J_m)^n$ is finitely generated by some elements a_1, \dots, a_l . Notice that $a_i \in J$ for $1 \leq i \leq l$, whence $(I : a_i) \in \mathcal{A}$. In this way, there exist elements $k_i, j_i \in \mathbb{N}$ such that $(J_{j_i})^{k_i} \subseteq (I : a_i)$. If $k := \max\{k_i\}_{1 \leq i \leq l}$ and $j := \max\{j_i\}_{1 \leq i \leq l}$, then $(J_j)^k \subseteq (I : a_i)$ for every $1 \leq i \leq l$.

Let $r \in (J_j)^k$ and $s \in (J_m)^n$. There exist elements $r_1, \dots, r_l \in R$ such that $s = \sum_{i=1}^l r_i a_i$, and so, $rs = \sum_{i=1}^l r r_i a_i$. Since $(J_j)^k$ is an ideal of R , we have that $r r_i \in (J_j)^k$ for all i . Thus, $r r_i a_i \in I$ whence $rs \in I$. It follows that $(J_{j+m})^{k+n} \subseteq (J_j)^k (J_m)^n \subseteq I$. ■

Consider a ring R , \mathcal{L} an idempotent filter of left ideals of R and its associated radical κ . For an R -module M , we recall the quotient module $Q_\kappa(M)$ of M (Definition 2.4). With this aim, we introduce Definition 2.3.

Definition 2.3. Let $M \in R - \text{Mod}$. Consider the family Ω_M of pairs (I, f) with $I \in \mathcal{L}$ and $f : I \rightarrow M$ an R -homomorphism. We define the relation \sim on Ω_M as $(I_1, f_1) \sim (I_2, f_2)$ if and only if there exists an element $J \in \mathcal{L}$ such that $J \subseteq I_1 \cap I_2$ and $f|_J = g|_J$.

It is straightforward to see that \sim is an equivalence relation. The equivalence class of the element (I, f) is denoted as $[I, f]$, and the set of equivalence classes will be written as $M_\mathcal{L}$. For two elements $[I, f], [J, g] \in M_\mathcal{L}$, we define their sum as $[I, f] + [J, g] = [I \cap J, f + g]$. It is easy to see that this sum is well defined and that $(M_\mathcal{L}, +)$ is an Abelian group.

It is also easy to see that if $I, J \in \mathcal{L}$ and $f \in \text{Hom}(I, R)$, then $f^{-1}(J) \in \mathcal{L}$. In this way, for elements $[I, f] \in R_\mathcal{L}$ and $[J, g] \in M_\mathcal{L}$, we can define their product as $[I, f] \cdot [J, g] = [f^{-1}(J), g \circ f]$. Notice that this product is well defined, and so, $R_\mathcal{L}$ is actually a ring with identity $[R, \text{id}_R]$. Thus, $M_\mathcal{L}$ is a left $R_\mathcal{L}$ -module.

Let $m \in M$. We define the application $\beta(m) : R \rightarrow M$ given by $\beta(m)(r) = rm$. It is well known that $\beta : M \rightarrow \text{Hom}(R, M)$ is an isomorphism of R -modules. If we consider $\varphi_M : M \rightarrow M_\mathcal{L}$ defined by $\varphi_M(m) = [R, \beta(m)]$, it follows that φ_R is a ring homomorphism, and so, we can consider $R_\mathcal{L}$ and $M_\mathcal{L}$ as R -modules with the action given by $r[I, f] := [R, \beta(r)][I, f]$. Note that φ_M is actually a homomorphism of R -modules. Since $\text{Ker}(\varphi_M) = \kappa(M)$, the fundamental isomorphism theorem guarantees that $\varphi_M(M) \cong M/\kappa(M)$. In this way, if $\kappa(M) = 0$, then we can embed M into $M_\mathcal{L}$.

For an element $\xi \in M_\mathcal{L}$ given by $\xi = [I, f]$ and an element $a \in I$, notice that $a\xi = [R, \beta(f(a))] = \varphi_M(f(a))$ which shows that $I\xi \subseteq \varphi_M(M)$, that is, $\text{Coker}(\varphi_M)$ is a κ -torsion module.

Considering the notation and terminology above, we present the definition of the quotient module of an object M in $R - \text{Mod}$.

Definition 2.4. The *quotient module* of M with respect to κ is defined as $Q_\kappa(M) = (M/\kappa(M))_{\mathcal{L}}$. Since \mathcal{L} is idempotent, it follows that $\kappa(M/\kappa(M)) = 0$. Hence, we can embed $M/\kappa(M)$ into $Q_\kappa(M)$.

Equivalently, the *quotient module* of M with respect to κ is given by

$$Q_\kappa(M) = \varinjlim_{I \in \mathcal{L}} \text{Hom}_R(I, M/\kappa(M)),$$

where $Q_\kappa(M)$ turns out to be a module over the ring $Q_\kappa(R)$.

Following [44], recall that an R -module E is κ -*injective* if for every R -module M and each submodule N such that $\kappa(M/N) = M/N$, every R -homomorphism $N \rightarrow E$ can be extended to an R -homomorphism $M \rightarrow E$. We say that E is κ -*closed* (also known as *faithfully κ -injective*) if the extension of the homomorphism is unique. It is straightforward to see that E is κ -closed if and only if E is κ -injective and κ -torsion-free. By using these notions, we can characterize $Q_\kappa(M)$ in the following way: $Q_\kappa(M)$ is the unique κ -closed module containing $N = M/\kappa(M)$ such that $Q_\kappa(M)/N$ is κ -torsion.

Example 2.5 ([33, page 111]). (i) Consider S a left Ore set in an arbitrary ring R . The set

$$\mathcal{L}(S) = \{I \triangleleft_l R \mid I \cap S \neq \emptyset\}$$

is an idempotent filter. If κ_S denotes its corresponding radical and $Q_S(M)$ is the module of quotients of M , then it can be seen that $Q_S(M)$ is isomorphic to $S^{-1}M$, i.e., the classical Ore localization of M at S .

(ii) If $R = \bigoplus_{k \geq 0} R_k$ is a positively graded Noetherian ring and R_+ denotes the two-sided ideal $\bigoplus_{k \geq 1} R_k$, by Proposition 2.2 the set

$$\mathcal{L}(\kappa_+) = \{I \triangleleft_l R \mid \text{there exists } n \in \mathbb{N} \text{ with } (R_+)^n \subseteq I\}$$

is an idempotent filter. The corresponding radical is denoted by κ_+ .

From the treatment above and having in mind that the filter $\mathcal{L}(\kappa_+)$ is idempotent, Van Oystaeyen and Willaert [33] formed the *quotient category* $(R, \kappa_+)\text{-gr}$, that is, the full subcategory of $Q_{\kappa_+}(R)\text{-gr}$ consisting of modules of the form $Q_{\kappa_+}(M)$ for some graded R -module M . Notice that $(R, \kappa_+)\text{-gr}$ is equivalent to the full subcategory of $R\text{-gr}$ consisting of the κ_+ -closed modules. Define $\text{Proj } R$ as the *Noetherian objects* in $(R, \kappa_+)\text{-gr}$. Since they wanted to describe the objects of $\text{Proj } R$ by means of objects of usual module categories in the same way as for commutative algebras, they needed modules determined by Ore localizations. This is the content of the following definition.

Definition 2.6 ([33, Definition 1]). The noncommutative positively graded Noetherian \mathbb{k} -algebra

$$R = \mathbb{k} \oplus R_1 \oplus R_2 \oplus \cdots \quad \text{with } R = \mathbb{k}[R_1]$$

is *schematic* if there is a finite set I of homogeneous left Ore sets of R such that for every $S \in I$ we have that $S \cap R_+ \neq \emptyset$, and such that one of the following equivalent properties is satisfied:

- (i) for each element $(r_S)_{S \in I} \in \prod_{S \in I} S$ there exists $m \in \mathbb{N}$ such that $(R_+)^m \subseteq \sum_{S \in I} Rr_S$,
- (ii) $\bigcap_{S \in I} \mathcal{L}(S) = \mathcal{L}(\kappa_+)$,
- (iii) $\bigcap_{S \in I} \kappa_S(M) = \kappa_+(M)$ for all $M \in R\text{-Mod}$,
- (iv) $\bigwedge_{S \in I} \kappa_S = \kappa_+$ where \bigwedge denotes the infimum of torsion theories.

In [33, 51], Van Oystaeyen and Willaert constructed the *noncommutative site*, a category with coverings on which sheaves can be defined, and formulated the Serre’s theorem. Examples 2.7 and 2.9 contain remarkable examples of schematic algebras.

Example 2.7. Recall that if R is a *positively filtered* \mathbb{k} -algebra by the family $(F_n R)_{n \geq 0}$ (i.e., $F_0 R = \mathbb{k}$), $\sigma : R \rightarrow G(R)$ is the principal symbol map, and \hat{R} is the Rees-ring of R , it is well known that $G(R)$ and \hat{R} are positively graded and there is a canonical central element X in \hat{R} of degree 1 such that $\hat{R}/\langle X \rangle \cong G(R)$. If \hat{R} is Noetherian, this is equivalent to $G(R)$ being Noetherian or the filtration of R being Zariskian. For R positively filtered by $(F_n R)_{n \geq 0}$, if $G(R)$ is schematic then \hat{R} is schematic [52, Theorem 1]. In this way, since for an almost commutative ring R there exists a filtration on R such that $G(R)$ is commutative, it follows that its Rees-ring is schematic. For example, the algebra R generated by three elements x, y and z of degree 1 with relations $xy - yx = z^2$, $xz - zx = 0$, and $yz - zy = 0$, is schematic since it is the Rees-ring of the first Weyl algebra $A_1(\mathbb{k})$ with respect to the Bernstein-filtration (this algebra is known as the *homogenized Weyl algebra*).

Van Oystaeyen and Willaert [52, page 199] said that “it is probably not true that the class of schematic algebras is closed under iterated Ore extensions since Ore sets in a ring R need not be Ore in an Ore extension $R[x; \sigma, \delta]$ ”. Nevertheless, the following proposition shows that under suitable conditions, these extensions are schematic.

Proposition 2.8 ([52, Theorem 3]). *Given a positively graded ring R which is generated by R_1 and which is schematic by means of Ore sets S_i , given σ a graded automorphism of R and δ a σ -derivation of degree 1, then for all $s_i \in \prod S_i$ and for all $m \in \mathbb{N}$, there exists $p \in \mathbb{N}$ such that*

$$(R[x; \sigma, \delta]_+)^p \subseteq \sum_i R[x; \sigma, \delta]s_i + R[x; \sigma, \delta]x^m,$$

where $R[x; \sigma, \delta]$ denotes the Ore extension considered with graduation $(R[x; \sigma, \delta])_n = \bigoplus_{k=0}^n R_k x^{n-k}$.

Proposition 2.8 is one of the results that Van Oystaeyen and Willaert [52] used to show that the algebras in Example 2.9 are schematic.

Example 2.9 ([52, Examples 2–5]). (i) The *coordinated ring of quantum 2×2 -matrices* $\mathcal{O}_q(M_2(\mathbb{C}))$ with $q \in \mathbb{C}$ is generated by elements a, b, c and d subject to the relations

$$\begin{aligned} ba &= q^{-2}ab, & ca &= q^{-2}ac, & bc &= cb, \\ db &= q^{-2}bd, & dc &= q^{-2}cd, & ad - da &= (q^2 - q^{-2})bc. \end{aligned}$$

(ii) *Quantum Weyl algebras* $A_n^{\bar{q}, \Lambda}$ defined by Alev and Dumas [1] are given by an $n \times n$ matrix $\Lambda = (\lambda_{ij})$ with $\lambda_{ij} \in \mathbb{k}^*$ and a row vector $\bar{q} = (q_1, \dots, q_n)$, where $q_i \neq 0$ for every i , the algebra is generated by elements $x_1, \dots, x_n, y_1, \dots, y_n$ subject to relations ($i < j$) given by

$$\begin{aligned} x_i x_j &= \mu_{ij} x_j x_i, \\ x_i y_j &= \lambda_{ji} y_j x_i, \\ y_j y_i &= \lambda_{ji} y_i y_j, \\ x_j y_i &= \mu_{ij} y_i x_j, \\ x_j y_j &= 1 + q_j y_j x_j + \sum_{i < j} (q_i - 1) y_i x_i, \quad \text{where } \mu_{ij} = \lambda_{ij} q_i. \end{aligned}$$

(iii) *Three-dimensional Sklyanin algebras* $A_{\mathbb{k}}$ over a field \mathbb{k} according to Artin et al. [4] are graded \mathbb{k} -algebras generated by three homogeneous elements x, y and z of degree 1 satisfying the relations

$$axy + byx + cz^2 = 0, \quad ayz + bzy + cx^2 = 0, \quad azx + bxz + cy^2 = 0,$$

where $a, b, c \in \mathbb{k}$.

(iv) Color Lie super algebras are defined by Rittenberg and Wyler [40].

Remark 2.10. Of course, there are examples of non-schematic algebras. If we take the graded algebra $\mathbb{k}\{x, y\}/\langle yx - xy - x^2 \rangle$ and suppose that $\text{char}(\mathbb{k}) = 0$, then its subalgebra generated by y and xy is not left schematic [52, page 203].

3. Semi-graded rings

Lezama and Latorre [29] presented a first approach to the noncommutative algebraic geometry for non- \mathbb{N} -graded algebras and finitely non-graded algebras by defining a new class of rings, the *semi-graded rings*. These rings extend several kinds of noncommutative rings of polynomial type such as Ore extensions [31, 32], families of differential operators generalizing Weyl algebras and universal enveloping algebras of finite dimensional Lie algebras [6, 7, 43], algebras appearing in mathematical physics [22, 37, 60], down-up algebras [9, 10, 25], ambiskew polynomial rings [23, 24], 3-dimensional skew polynomial rings [8, 34, 37, 41], PBW extensions [7], and skew PBW extensions [15], among others. A detailed list of examples of semi-graded rings and their relationships with other

algebras can be found in Fajardo et al. [14]. Ring-theoretical, algebraic and geometric properties of semi-graded rings have been investigated in the literature by several authors (e.g., [2, 11, 21, 35, 36, 38, 45–47] and references therein).

Definition 3.1 ([29, Definition 2.1]). Let R be a ring. R is said to be *semi-graded* (SG) if there exists a collection $\{R_n\}_{n \in \mathbb{Z}}$ of subgroups R_n of the additive group R^+ such that the following conditions hold:

- (i) $R = \bigoplus_{n \in \mathbb{Z}} R_n$,
- (ii) for every $m, n \in \mathbb{Z}$, we have that $R_m R_n \subseteq \bigoplus_{k \leq m+n} R_k$,
- (iii) $1 \in R_0$.

The collection $\{R_n\}_{n \in \mathbb{Z}}$ is called a *semi-graduation* of R , and we say that the elements of R_n are *homogeneous of degree n* .

We say that R is *positively semi-graded* if $R_n = 0$ for every $n < 0$. If R and S are semi-graded rings and $f : R \rightarrow S$ is a ring homomorphism, then we say that f is *homogeneous* if $f(R_n) \subseteq S_n$ for every $n \in \mathbb{Z}$.

Definitions 3.2 and 3.3 recall the notion of finitely semi-graded ring and finitely semi-graded algebra, respectively.

Definition 3.2 ([29, Definition 2.4]). A ring R is called *finitely semi-graded* (FSG) if it satisfies the following conditions:

- (i) R is SG,
- (ii) there exist finitely many elements $x_1, \dots, x_n \in R$ such that the subring generated by R_0 and x_1, \dots, x_n coincides with R ,
- (iii) for every $n \geq 0$, we have that R_n is a free R_0 -module of finite dimension.

Definition 3.3 ([28, Definition 10]). A \mathbb{k} -algebra R is said to be *finitely semi-graded* (FSG) if the following conditions hold:

- (i) R is an FSG ring with semi-graduation given by $R = \bigoplus_{n \geq 0} R_n$,
- (ii) for every $m, n \geq 1$, we have that $R_m R_n \subseteq R_1 \oplus \dots \oplus R_{m+n}$,
- (iii) R is connected, i.e., $R_0 = \mathbb{k}$,
- (iv) R is generated in degree 1.

From Definition 3.3, it is straightforward to see that if R is a \mathbb{k} -algebra, then

$$R_+ := \bigoplus_{n \geq 1} R_n$$

is a maximal ideal of R .

\mathbb{N} -graded rings are SG. Finitely graded \mathbb{k} -algebras, PBW extensions [7], 3-dimensional skew polynomial rings [8], down-up algebras [9, 10], diffusion algebras [22], and skew PBW extensions [15] are examples of FSG rings.

Definition 3.4 ([29, Definition 2.2]). Let R be an SG ring and let M be an R -module. We say that M is *semi-graded* if there exists a collection $\{M_n\}_{n \in \mathbb{Z}}$ of subgroups M_n of the additive group M^+ such that the following conditions hold:

- (i) $M = \bigoplus_{n \in \mathbb{Z}} M_n$,
- (ii) for every $m \geq 0$ and $n \in \mathbb{Z}$, we have that $R_m M_n \subseteq \bigoplus_{k \leq m+n} M_k$.

The collection $\{M_n\}_{n \in \mathbb{Z}}$ is called a *semi-graduation* of M , and we say that the elements of M_n are *homogeneous of degree n* .

M is said to be *positively semi-graded* if $M_n = 0$ for every $n < 0$. Let $f : M \rightarrow N$ be a homomorphism of R -modules, where M and N are semi-graded R -modules. We say that f is *homogeneous* if $f(M_n) \subseteq N_n$ for every $n \in \mathbb{Z}$.

Definition 3.5 ([29, Definition 2.3]). Let R be an SG ring, M an SG R -module and N a submodule of M . We say that N is a *semi-graded (SG) submodule* of M if $N = \bigoplus_{n \in \mathbb{Z}} N_n$, where $N_n = M_n \cap N$. In this case, N is an SG R -module.

Proposition 3.6 ([29, Proposition 2.6]). *If R is an SG ring, M is an SG R -module and N is a submodule of M , then the following conditions are equivalent:*

- (1) N is a semi-graded submodule of M ,
- (2) for every $z \in N$, the homogeneous components of z are in N ,
- (3) M/N is an SG R -module with semi-graduation given by

$$(M/N)_n = (M_n + N)/N \quad \text{for every } n \in \mathbb{Z}.$$

If M is an SG R -module and $\{N_i\}_{i \in I}$ is a family of SG submodules of M , then it is clear that $\bigcap_{i \in I} N_i$ is an SG submodule of M .

Let X be a subset of M . We define the SG submodule generated by X as the intersection of all SG submodules containing X , and we will denote it as $\langle X \rangle^{\text{SG}}$. If $X = \{x_1, \dots, x_n\}$, then we write $\langle X \rangle^{\text{SG}} = \langle x_1, \dots, x_n \rangle^{\text{SG}}$. We will say that M is a *finitely generated* SG R -module if there exist finitely many elements m_1, \dots, m_n such that $M = \langle m_1, \dots, m_n \rangle^{\text{SG}}$. If M is simultaneously a module over different kinds of rings and there is a risk of confusion, we write $\langle - \rangle_R^{\text{SG}}$ to indicate the ring R we are considering.

In a similar way, if R is a positively SG ring, for $t \in \mathbb{N}$ we define $R_{\geq t}$ as the intersection of all two-sided ideals that are SG submodules containing $\bigoplus_{k \geq t} R_k$.

Remark 3.7. If R is a positively SG left Noetherian ring, then Proposition 2.2 shows that

$$\mathcal{L}(\kappa_+) = \{I \triangleleft_l R \mid \text{there exist } n, m \in \mathbb{N} \text{ with } (R_{\geq m})^n \subseteq I\}$$

is an idempotent filter. The corresponding left exact radical is denoted by κ_+ and $\mathcal{Q}_{\kappa_+}(M)$ is the module of quotients of M .

Next, we want to formalize several constructions concerning semi-graded rings which are necessary to formulate the Serre's theorem.

3.1. Localization of semi-graded rings

With the aim of defining *good* Ore sets (Definition 3.8), for R an SG ring and an element $n \in \mathbb{Z}$, we consider the following sets:

$$\begin{aligned} R'_n &= \{r \in R_n \mid \text{for all } m \in \mathbb{Z}, \text{ and for all } h \in R_m, rh \in R_{n+m}\}, \\ R''_n &= \{r \in R'_n \mid \text{for all } m \in \mathbb{Z}, \text{ and for all } h \in R_m, hr \in R_{n+m}\}, \\ R' &= \bigcup_{n \in \mathbb{Z}} R'_n, \\ R'' &= \bigcup_{n \in \mathbb{Z}} R''_n. \end{aligned}$$

Definition 3.8. Let R be an SG ring and consider a left Ore set S of R . We say that S is *good* if the following conditions hold:

- (i) $S \subseteq R''$,
- (ii) if $s \in S$ and $r \in R'$, then there exist elements $u \in R'$ and $v \in S$ such that $us = vr$.

From Definition 3.8, it follows that for any elements $s_1, \dots, s_k \in S$, there exist $r_1, \dots, r_k \in R'$ such that $r_i s_i = r_j s_j \in S$ for every i, j .

Definition 3.9. Let R be an SG ring and M an SG R -module. We say that M is *localizable semi-graded* (LSG) if for every element $(n, m) \in \mathbb{Z}^2$ the inclusion $R'_n M_m \subseteq M_{n+m}$ holds.

Proposition 3.10. Let R be an SG ring, S a good left Ore set and M an LSG R -module. Then, $S^{-1}M$ is an LSG R -module with semi-graduation given by

$$(S^{-1}M)_n = \left\{ \frac{f}{s} \mid f \in \bigcup_{k \in \mathbb{Z}} M_k, \deg(f) - \deg(s) = n \right\}.$$

Proof. First of all, let us show that $(S^{-1}M)_n$ is a subgroup of $S^{-1}M$. It is clear that $0 = \frac{0}{1} \in (S^{-1}M)_n$ and that $(S^{-1}M)_n$ has additive inverses. Consider elements $\frac{p}{s}, \frac{q}{t} \in (S^{-1}M)_n$. Then, $\deg(p) - \deg(s) = \deg(q) - \deg(t) = n$. There exist elements $u \in R'$ and $v \in S$ such that $us = vt \in S$. Note that $\deg(u) + \deg(s) = \deg(v) + \deg(t)$. Since $u, v \in R'$, it follows that up and vq are homogeneous elements satisfying $\deg(up) = \deg(u) + \deg(p) = \deg(v) + \deg(q) = \deg(vq)$, whence $\frac{p}{s} + \frac{q}{t} = \frac{up+vt}{vt}$ is a homogeneous elements of degree $\deg(v) + \deg(q) - (\deg(v) + \deg(t)) = n$.

It is clear that $S^{-1}M$ is the sum of the subgroups $(S^{-1}M)_n$, so let us show that the sum is direct. Consider the sum

$$\sum_{i=1}^k \frac{m_i}{s_i} = 0$$

of homogeneous elements of $S^{-1}M$ with different degrees, that is, $\deg(m_i) - \deg(s_i) \neq \deg(m_j) - \deg(s_j)$ for $i \neq j$. There exist elements $r_1, \dots, r_k \in R'$ such that $r_i s_i = r_j s_j$

for all i, j , which implies that

$$0 = \sum_{i=1}^k \frac{m_i}{s_i} = \frac{\sum_{i=1}^k r_i m_i}{r_1 s_1}.$$

Hence, there exists an element $s \in S$ such that $0 = s \sum_{i=1}^k r_i m_i = \sum_{i=1}^k s r_i m_i$. Since $s, r_i \in R'$ and m_i is homogeneous for $0 \leq i \leq k$, then every one of the terms above is homogeneous. By using that $r_i s_i = r_j s_j$, we have the equality $\deg(r_i) + \deg(s_i) = \deg(r_j) + \deg(s_j)$, whence $\deg(s) + \deg(r_i) + \deg(m_i) \neq \deg(s) + \deg(r_j) + \deg(m_j)$, which shows that $s r_i m_i = 0$. Thus, $0 = \frac{r_i m_i}{r_i s_i} = \frac{m_i}{s_i}$.

Now, let us see that $R_a(S^{-1}M)_b \subseteq \bigoplus_{k \leq a+b} (S^{-1}M)_k$. Let $r \in R_a$ and $\frac{m}{s} \in (S^{-1}M)_b$. There exist elements $r' \in R$ and $s' \in S$ such that $r's = s'r$. Since $s, s' \in R''$ and r is homogeneous, we can take the element r' being homogeneous. Then, $\deg(r') = \deg(s') + \deg(r) - \deg(s)$, and using that $r \frac{m}{s} = \frac{r'm}{s'}$ and $r'm \in \bigoplus_{k \leq \deg(r') + \deg(m)} M_k$, it follows that $\frac{r'm}{s'} \in \bigoplus_{k \leq \deg(r') + \deg(m) - \deg(s')} (S^{-1}M)_k$. Since $\deg(r') + \deg(m) - \deg(s') = \deg(r) + \deg(m) - \deg(s) = a + b$, then $r \frac{m}{s} \in \bigoplus_{k \leq a+b} (S^{-1}M)_k$. This fact proves that $S^{-1}M$ is an SG R -module.

If we consider above the element $r \in R'_a$, then we can take $r' \in R'$ to obtain that $r'm \in M_{\deg(r') + \deg(m)}$, and so, $r \frac{m}{s} \in (S^{-1}M)_{a+b}$. This shows that $S^{-1}M$ is an LSG R -module. ■

The next result shows that the localization of an SG ring by considering a good Ore set is also an SG ring.

Proposition 3.11. *Let R be an SG ring and S a good left Ore set. Then, $S^{-1}R$ is an SG ring with semigraduation given by*

$$(S^{-1}R)_n = \left\{ \frac{f}{s} \mid f \in \bigcup_{k \in \mathbb{Z}} R_k, \deg(f) - \deg(s) = n \right\}.$$

Proof. It is clear that R is an LSG R -module so $S^{-1}R$ is an SG R -module with the semigraduation above, and hence, $S^{-1}R = \bigoplus_{k \in \mathbb{Z}} (S^{-1}R)_k$. It is easy to see that $1 = \frac{1}{1} \in (S^{-1}R)_0$. We only have to show that

$$(S^{-1}R)_n (S^{-1}R)_m \subseteq \bigoplus_{k \leq n+m} (S^{-1}R)_k.$$

Let $\frac{r_1}{s_1} \in (S^{-1}R)_n$ and $\frac{r_2}{s_2} \in (S^{-1}R)_m$. There exist elements $u \in R$ and $v \in S$ such that $vr_1 = us_2$ which implies that $\frac{r_1}{s_1} \frac{r_2}{s_2} = \frac{ur_2}{vs_1}$. Again, since $s_2, v \in R''$ and r_1 is homogeneous we can take u as a homogeneous element. Hence, $ur_2 \in \bigoplus_{k \leq \deg(u) + \deg(r_2)} R_k$, and so, $\frac{ur_2}{vs_1} \in \bigoplus_{k \leq \deg(u) + \deg(r_2) - \deg(v) - \deg(s_1)} (S^{-1}R)_k$, i.e., $\frac{ur_2}{vs_1} \in \bigoplus_{k \leq n+m} (S^{-1}R)_k$. ■

Proposition 3.12. *Let R be an SG ring, S a good left Ore set and M an LSG R -module. Then, $S^{-1}M$ is an SG $S^{-1}R$ -module.*

Proof. We know that $S^{-1}M$ is an $S^{-1}R$ -module and an SG R -module which guarantees the direct sum $S^{-1}M = \bigoplus_{k \in \mathbb{Z}} (S^{-1}M)_k$. In this way, we have to prove that $(S^{-1}R)_n (S^{-1}M)_m \subseteq \bigoplus_{k \leq n+m} (S^{-1}M)_k$. Consider elements $\frac{r}{s_1} \in (S^{-1}R)_n$ and $\frac{a}{s_2} \in (S^{-1}M)_m$. There exist elements $u \in R$ and $v \in S$ such that $vr = us_2$ which implies that $\frac{r}{s_1} \frac{a}{s_2} = \frac{ua}{vs_1}$. Since $s_2, v \in R''$ and r is homogeneous, we can take u as an homogeneous element. Thus, $ua \in \bigoplus_{k \leq \deg(u) + \deg(a)} R_k$, and so, $\frac{ua}{vs_1} \in \bigoplus_{k \leq \deg(u) + \deg(a) - \deg(v) - \deg(s_1)} (S^{-1}R)_k$, i.e., $\frac{ua}{vs_1} \in \bigoplus_{k \leq n+m} (S^{-1}R)_k$. ■

3.2. Category of semi-graded rings

We define the *category SGR of semi-graded rings* whose objects are the semi-graded rings and morphisms are the homogeneous ring homomorphisms. For a semi-graded ring R , $\text{SGR} - R$ will denote the *category of semi-graded modules over R* where the morphisms are the homogeneous R -homomorphisms. It is straightforward to see that $\text{SGR} - R$ is preadditive, and that the zero object of the category is the trivial module.

Let $f : M \rightarrow N$ be a morphism in $\text{SGR} - R$. Since $\text{Ker}(f)$ and $\text{Im}(f)$ are semi-graded submodules, it follows that $N/\text{Im}(f)$ is a semi-graded module. This fact guarantees that the category $\text{SGR} - R$ has kernels and cokernels. If f is a monomorphism of $\text{SGR} - R$, then f is the kernel of the canonical homomorphism $j : N \rightarrow N/\text{Im}(f)$. If f is an epimorphism, then f is the cokernel of the inclusion $i : \text{Ker}(f) \rightarrow M$. In this way, the category $\text{SGR} - R$ is normal and conormal.

If $\{M_i\}_{i \in I}$ is a family of objects of $\text{SGR} - R$, then their direct sum $\bigoplus_{i \in I} M_i$ is a semi-graded ring with semi-graduation given by

$$\left(\bigoplus_{i \in I} M_i \right)_p := \bigoplus_{i \in I} (M_i)_p \quad \text{for each } p \in \mathbb{Z}.$$

It is easy to see that this object with the natural inclusions coincides with the coproduct of the family of objects $\{M_i\}_{i \in I}$ in $\text{SGR} - R$. Therefore, $\text{SGR} - R$ is an Abelian category.

We define $\text{LSG} - R$ as the full subcategory of $\text{SGR} - R$ whose objects are the LSG modules. This subcategory is closed for subobjects, quotients and coproducts, which shows that it is Abelian.

4. Schematicness of semi-graded rings

Following Van Oystaeyen and Willaert's ideas developed in [33], in this section, we define the notion of *schematicness* in the setting of semi-graded rings. For a positively SG ring R , we define $R_+ = \bigoplus_{k \geq 1} R_k$ and say that a left Ore set S is *non-trivial* if $S \cap R_+ \neq \emptyset$.

Definition 4.1. Let R be a positively SG left Noetherian ring. R is called (*left*) *schematic* if there is a finite set I of non-trivial good left Ore sets of R such that for each $(x_S)_{S \in I} \in \prod_{S \in I} S$ there exist $t, m \in \mathbb{N}$ such that $(R_{\geq t})^m \subseteq \sum_{S \in I} R x_S$.

The following result illustrates some characterizations of being schematic (cf. Definition 2.6).

Proposition 4.2. *Let R be a positively SG left Noetherian algebra and S_1, \dots, S_n a finite set of non-trivial good left Ore sets of R . The following conditions are equivalent:*

- (1) *for each $(x_1, \dots, x_n) \in \prod_{i=1}^n S_i$ there exist elements $t, m \in \mathbb{N}$ such that $(R_{\geq t})^m \subseteq \sum_{s \in I} Rx_s$,*
- (2) *let $I \triangleleft_l R$. If I has no trivial intersection with every S_i , then I contains a power of $R_{\geq t}$ for some $t \in \mathbb{N}$,*
- (3) $\bigcap_{i=1}^n \mathcal{L}(S_i) = \mathcal{L}(\kappa_+)$.

Proof. The equivalence (1) \Leftrightarrow (2) and the implication (3) \Rightarrow (1) are straightforward.

(1) \Rightarrow (3) Let $I \in \bigcap_{i=1}^n \mathcal{L}(S_i)$. There exist elements x_1, \dots, x_n such that $x_i \in I \cap S_i$ for every i . Thus, $\sum_{i=1}^n Rx_i \subseteq I$ and there exist t, m with $(R_{\geq t})^m \subseteq I$, which shows that $I \in \mathcal{L}(\kappa_+)$.

Now, let $I \in \mathcal{L}(\kappa_+)$. There exist t, m such that $(R_{\geq t})^m \subseteq I$. By using that $S_i \cap R_+ \neq \emptyset$, there exist elements $s_i \in S_i$ such that $\deg(s_i) \geq 1$ for all i . Then, $s_i^t \in R_{\geq t}$, $s_i^{tm} \in (R_{\geq t})^m \subseteq I$, and therefore, $I \cap S_i \neq \emptyset$. This shows that $I \in \bigcap_{i=1}^n \mathcal{L}(S_i)$. ■

If R is schematic by considering the good left Ore sets S_i , then $\bigcap_{i=1}^n \kappa_{S_i}(M) = \kappa_+(M)$ for every R -module M . If M is an LSG R -module, then for each $i = 1, \dots, n$ we have that $\kappa_{S_i}(M)$ is an SG submodule, and so, $\kappa_+(M)$ is also an SG submodule. These facts imply that $M/\kappa_+(M)$ is an SG R -module, and so, it is a submodule of $Q_{\kappa_+}(M)$. The idea is to show that $Q_{\kappa_+}(M)$ is semi-graded. For the remainder of the section, we will take $\mathcal{L} := \mathcal{L}(\kappa_+)$.

Let us start by taking an LSG R -module M such that $\kappa_+(M) = 0$. It is clear that $Q_{\kappa_+}(M) = M_{\mathcal{L}}$ and $\varphi_M(M) \cong M$. Thus, $\varphi_M(M)$ is a submodule of $M_{\mathcal{L}}$ which is an SG R -module where $\varphi_M(m)$ is homogeneous of degree k if and only if m is homogeneous of degree k . If we want $M_{\mathcal{L}}$ to be an LSG R -module, it must satisfy that if ξ is homogeneous of degree k , then for every $s \in R'$ the element $s\xi \in (\varphi_M(M))_{\deg(s)+k}$. Since there exists $I \in \mathcal{L}$ with $I\xi \subseteq \varphi_M(M)$ the following definition makes sense.

Definition 4.3. Let $\xi \in M_{\mathcal{L}}$. We say that the element ξ is *homogeneous of degree k* if there exists $I \in \mathcal{L}$ such that $I\xi \subseteq \varphi_M(M)$, and for every element $s \in I \cap R'$ we have that $s\xi \in (\varphi_M(M))_{\deg(s)+k}$.

Notice that if the condition above is satisfied for I , then it also holds for every $J \subseteq I$. Lemma 4.5 shows that this condition is true for ideals containing I .

Remark 4.4. Since the good Ore sets S_i are non-trivial there exist elements $s'_i \in S_i \cap R_+$ for $i = 1, \dots, n$, whence $\alpha_i = \deg(s'_i) > 0$. If we define $m := \text{lcm}\{\alpha_i\}_{1 \leq i \leq n}$ and $s''_i := (s'_i)^{m/\alpha_i}$, we obtain that $s''_i \in S_i \cap R_+$, and all of them have the same degree. If we consider an element $I \in \mathcal{L}$, there exist $t, n \in \mathbb{Z}$ such that $(R_{\geq t})^n \subseteq I$. Thus, $s_i = (s''_i)^{tn} \in I \cap S_i$

which implies that $\sum_{i=1}^n R s_i \subseteq I$. In this way, for each $I \in \mathcal{L}$, there exist elements $s_i \in S_i$, all with the same positive degree, satisfying the relation $\sum_{i=1}^n R s_i \subseteq I$.

Lemma 4.5. *Let $I, J \in \mathcal{L}$ be ideals such that $I \subseteq J$ and $I\xi, J\xi \subseteq \varphi_M(M)$. If for every $s \in I \cap R'$, the element $s\xi$ belongs to $(\varphi_M(M))_{\deg(s)+k}$, then the same property holds for each $s \in J \cap R'$.*

Proof. Let $s \in J \cap R'$. Then, $s\xi \in \varphi_M(M)$, and so, there exist homogeneous elements ξ_j , $j = 1, \dots, l_r$ of $\varphi_M(M)$ with $\xi_j \in (\varphi_M(M))_j$ and $s\xi = \sum \xi_j$. As we said before, if the property holds for I , then it is true for any ideal contained in I , so we can take $I = (R_{\geq t})^m$ for some $t, m \in \mathbb{N}$. From above, there exist $s_i \in S_i$ for $i = 1, \dots, n$ such that $\sum R s_i \subseteq I$ and $\deg(s_i) = \beta$ for each $1 \leq i \leq n$. In particular, every element $s_i \in I$ whence $s_i s \in I$ (recall that I is a two-sided ideal). By assumption, $s_i s\xi \in (\varphi_M(M))_{\deg(s)+\beta+k}$ for each i .

On the other hand, if we consider the expression $s_i s\xi = \sum s_i \xi_j$ in terms of homogeneous elements of $\varphi_M(M)$, then for each $j \neq k + \deg(s)$ the equality $s_i \xi_j = 0$ holds. Since this is true for every i it follows that $\xi_j \in \bigcap_{i=1}^n \kappa_{S_i}(\varphi_M(M)) = \kappa_+(\varphi_M(M)) = 0$ (recall that $\kappa_+(M) = 0$). Therefore, $s\xi = \xi_{\deg(s)+k}$. ■

From Lemma 4.5, it is sufficient to guarantee the property by considering any ideal I such that $I\xi \subseteq \varphi_M(M)$. Our next purpose is to give a more simple method to verify that the element ξ is homogeneous. Let $\xi = [I, f]$. Since $I\xi \subseteq \varphi_M(M)$ the element ξ is homogeneous of degree k if and only if for each $s \in I \cap R'$ the element $s\xi = [R, \beta(f(s))] = \varphi_M(f(s)) \in (\varphi_M(M))_{\deg(s)+k}$, or equivalently, for all $s \in I \cap R'$ the element $f(s)$ belongs to $M_{\deg(s)+k}$.

For a morphism $f : I \rightarrow M$, we will say that f is *homogeneous of degree k* if for each $s \in I \cap R'$ the element $f(s)$ is homogeneous of degree $\deg(s) + k$. Hence, $[I, f]$ is homogeneous of degree k (in $M_{\mathcal{L}}$) if and only if f is homogeneous of degree k . Let $(M_{\mathcal{L}})_k$ be the family of homogeneous elements of degree k . It is clear that $(M_{\mathcal{L}})_k$ is a subgroup and $\varphi_M(M_k) \subseteq (M_{\mathcal{L}})_k$.

Remark 4.6. We will say that the morphism $f : I \rightarrow M$ is *strongly homogeneous of degree k* if for every homogeneous element $s \in I$ the element $f(s)$ is homogeneous of degree $\deg(s) + k$. It is clear that in the setting of graded rings, the notions of homogeneous morphism and strongly homogeneous morphism coincide.

On the other hand, $[I, f]$ it will be called *strongly homogeneous of degree k* if some of its representative elements is strongly homogeneous of degree k . Let $(\overline{M_{\mathcal{L}}})_k$ be the family of strongly homogeneous elements of degree k . It is straightforward to see that $\overline{R_{\mathcal{L}}} = \bigoplus (R_{\mathcal{L}})_k$ is a graded ring and $\overline{M_{\mathcal{L}}} = \bigoplus (\overline{M_{\mathcal{L}}})_k$ is an $\overline{R_{\mathcal{L}}}$ -graded module. Note also that if $s \in R''$ then $\varphi_R(s) \in \overline{R_{\mathcal{L}}}$; in particular, $\overline{R_{\mathcal{L}}}$ is an extension of the graded ring $\varphi_R(\bigoplus R_k'')$. As it is clear, $\overline{R_{\mathcal{L}}}$ is an R -submodule of $R_{\mathcal{L}}$ if and only if R is graded. This last remark shows that in the setting of non-graded rings is not appropriate to consider strongly homogeneous morphisms.

Proposition 4.7. *The sum $\sum (M_{\mathcal{L}})_k$ is direct.*

Proof. Let $[I_i, f_i] \in (M_{\mathcal{L}})_{k_i}$ for $i = 1, \dots, m$ with $k_i \neq k_j$ if $i \neq j$. Notice that if $\sum [I_i, f_i] = 0$ then there exists $J \subseteq \bigcap I_i$, $J \in \mathcal{L}$, such that $(\sum f_i)|_J = \sum f_i|_J = 0$. We can take $J = \sum R s_j$ for some $s_j \in S_j$. Let $s \in J \cap R'$ with $\deg(s) = l$. Then, $0 = (\sum f_i)(s) = \sum f_i(s)$ and since $f_i(s)$ is homogeneous of degree $l + k_i$ and all elements k_i are different, then we have a sum of homogeneous elements of different degrees equal to zero, whence $f_i(s) = 0$ for each i . In particular, $f_i(s_j) = 0$ for all i, j . Therefore, $f_i(x) = 0$ for all $x \in J$, and so, $[J, f_i|_J] = [I_i, f_i] = 0$. ■

Let $[I, f] \in M_{\mathcal{L}}$ with $I = \sum_{i=1}^n R s_i$ for some elements $s_i \in S_i \cap R''_k$. Since there are finitely s_i 's we may assume that the homogeneous decompositions of the elements $f(s_i)$ have the same length, say $f(s_i) = \sum_{t=\alpha}^{\beta} (f(s_i))_{t+k}$, where $(f(s_i))_j$ is the j th homogeneous component of $f(s_i)$. By taking $f_t(s_i) = (f(s_i))_{t+k}$, we have that $f(s_i) = \sum_{t=\alpha}^{\beta} f_t(s_i)$. For elements $t = \alpha, \dots, \beta$, we define the maps $f_t : I \rightarrow M$ in the natural way as $f_t(\sum a_i s_i) = \sum a_i f_t(s_i)$. However, we have to show that these maps are well defined. This is the content of the following proposition.

Proposition 4.8. *f_t is well defined for every element $t = \alpha, \dots, \beta$.*

Proof. We divide the proof into three parts.

- Suppose that $0 = \sum a_i s_i$, with $a_i \in R_{k_1}$ for every i (recall that $s_i \in R''_k$). Fix i . Since $s_j \in R''$ for each $1 \leq j \leq n$, there exist elements $u_j \in R'$ and $v_j \in S_i$ such that $u_j s_i = v_j s_j$. In particular, $\deg(u_j) = \deg(v_j)$ and since $u_j, v_j \in R'$, $u_j f(s_i) = v_j f(s_j)$ and M is LSG, if we compare the homogeneous components of the same degree then we obtain that $u_j f_t(s_i) = v_j f_t(s_j)$ for each $\alpha \leq t \leq \beta$.

Now, by using that $v_1 \in S_i$ and $a_1 \in R$ there exist elements $b_1 \in R$ and $c_1 \in S_i$ such that $b_1 v_1 = c_1 a_1$. Repeating this argument with the elements v_2 and $c_1 a_2$ we find that $b_2 \in R$ and $c_2 \in S_i$ satisfy the equality $b_2 v_2 = c_2 c_1 a_2$. Continuing in this way, for every $1 \leq j \leq n$ we will find elements $b_j \in R$ and $c_j \in S_i$ such that $b_j v_j = \prod_{i=1}^j c_i a_i$ (notice that the elements b_i 's can be taken homogeneous). If we define $c := \prod_{i=1}^n c_i \in S_i$ and $d_j = \prod_{i=j+1}^n c_i b_j$, then we have that $d_j v_j = c a_j$ for every $1 \leq j \leq n$. Hence, $0 = c \sum_{j=1}^n a_j s_j = \sum_{j=1}^n d_j v_j s_j = \sum_{j=1}^n d_j u_j s_i = r s_i$, where $r = \sum_{j=1}^n d_j u_j$. Note that the elements $d_j u_j$ are homogeneous of the same degree, which implies that r is also homogeneous. Since $0 = r s_i$, by the first condition of the noncommutative localization there exists an element $s \in S_i$ such that $s r = 0$.

Considering the equalities

$$s c \sum_{j=1}^n a_j f_t(s_j) = s \sum_{j=1}^n d_j v_j f_t(s_j) = s \sum_{j=1}^n d_j u_j f_t(s_i) = s r f_t(s_i) = 0,$$

it follows that $\sum_{j=1}^n a_j f_t(s_j) \in \kappa_{S_i}(M)$. Since this holds for every element i we have that $\sum_{j=1}^n a_j f_t(s_j) \in \bigcap \kappa_{S_i}(M) = \kappa_+(M) = 0$, whence $\sum_{j=1}^n a_j f_t(s_j) = 0$.

- Suppose that $0 = \sum_{i=1}^n a_i s_i$ (the elements a_i 's are not necessarily homogeneous). Since there are only finitely elements a_i 's we consider the sum $a_i = \sum_{j=l_1}^{l_2} b_{i,j}$ with $b_{i,j} \in R_j$. In this way, $0 = \sum_{i=1}^n \sum_{j=l_1}^{l_2} b_{i,j} s_i = \sum_{j=l_1}^{l_2} \sum_{i=1}^n b_{i,j} s_i$. Now, using that $\sum_{i=1}^n b_{i,j} s_i \in R_{j+k}$ is the homogeneous component of degree $j+k$, it follows that $0 = \sum_{i=1}^n b_{i,j} s_i$. By the first part above, we can assert that $\sum_{i=1}^n b_{i,j} f_t(s_i) = 0$, whence

$$\begin{aligned} 0 &= \sum_{j=l_1}^{l_2} \sum_{i=1}^n b_{i,j} f_t(s_i) \\ &= \sum_{i=1}^n \sum_{j=l_1}^{l_2} b_{i,j} f_t(s_i) \\ &= \sum_{i=1}^n a_i f_t(s_i) \quad \text{for } \alpha \leq t \leq \beta. \end{aligned}$$

- Let r be an element of $\sum R s_i$. Suppose that we have two expressions for r given by $r = \sum a_i s_i = \sum b_i s_i$. Then, $0 = \sum (a_i - b_i) s_i$. By the second part above, we obtain that $\sum (a_i - b_i) f_t(s_i) = 0$, and so, $\sum a_i f_t(s_i) = \sum b_i f_t(s_i)$ for $\alpha \leq t \leq \beta$. This means that the expression for $f_t(r)$ does not depend on the decomposition of r . ■

From the proof of Proposition 4.8, it follows that the maps f_t 's are R –homomorphisms. The next proposition establishes that these are homogeneous of degree t .

Proposition 4.9. *The map f_t is homogeneous of degree t .*

Proof. Consider $s \in I \cap R'$ with $\deg(s) = l$. Let $(f_t(s))_m$ be the homogeneous component of degree m in the expression of $f_t(s)$. For a fixed i , there exist elements $v_i \in S_i$ and $u_i \in R'$ such that $u_i s_i = v_i s$, which implies that $v_i f_t(s) = u_i f_t(s_i)$. Since $f_t(s_i) \in M_{t+k}$, $u_i, v_i \in R''$ and M is LSG, when we compare the homogeneous components of these elements we obtain that if $m \neq t+l$, then $v_i (f_t(s))_m = 0$, whence $(f_t(s))_m \in \kappa_{S_i}(\varphi_M(M))$. Note that this fact holds for every i , hence, $(f_t(s))_m \in \bigcap \kappa_{S_i}(\varphi_M(M)) = \kappa_+(\varphi_M(M)) = 0$. Therefore, $f_t(s) = (f_t(s))_{t+l}$ which guarantees that f_t is homogeneous of degree t . ■

Propositions 4.7, 4.8, and 4.9 imply the following important result.

Proposition 4.10. *If M is an LSG R –module with $\kappa_+(M) = 0$, then $Q_{\kappa_+}(M) = M_{\mathcal{L}}$ is an LSG R –module with semigraduation given by*

$$M_{\mathcal{L}} = \bigoplus_k (M_{\mathcal{L}})_k.$$

Theorem 4.11. *If M is an LSG R –module, then $Q_{\kappa_+}(M)$ is an LSG R –module.*

Proof. It follows from Proposition 4.10 and the equalities $\kappa_+(M/\kappa_+(M)) = 0$, $Q_{\kappa_+}(M) = Q_{\kappa_+}(M/\kappa_+(M))$. ■

5. Serre–Artin–Zhang–Verevkin theorem

In this section, we prove the Serre–Artin–Zhang–Verevkin theorem for semi-graded rings (Theorem 5.23) using a different approach than the one presented by Lezama [27, 29].

Briefly, this theorem was partially formulated by Lezama and Latorre [29, Theorem 6.12], where it was assumed that the semi-graded left Noetherian ring is a domain. Nevertheless, as is well known, the Serre–Artin–Zhang–Verevkin theorem for finitely graded algebras does not include this restriction, so that this assumption was eliminated by Lezama [27, Theorem 1.24] (see also [14, Section 18.4, Theorem 18.5.13]). More exactly, he proved the theorem for an SG ring $R = \bigoplus_{n \geq 0} R_n$ satisfying the following conditions:

- (C1) R is left Noetherian,
- (C2) R_0 is left Noetherian,
- (C3) for every n , we have that R_n is a finitely generated left R_0 -module,
- (C4) $R_0 \subset Z(R)$.

Notice that condition (C4) implies that R_0 is a commutative Noetherian ring.

Universal enveloping algebras of finite-dimensional Lie algebras, some quantum algebras with three generators and some examples of 3-dimensional skew polynomial algebras [8, 34, 37] illustrate the Serre–Artin–Zhang–Verevkin theorem [27, Example 1.26] and [14, Example 18.5.15].

We start with the following preliminary result.

Lemma 5.1. *Let R be a positively SG left Noetherian ring and S a non-trivial left Ore set of R . Then, $\mathcal{L}(\kappa_+) \subseteq \mathcal{L}(S)$.*

Proof. Let $I \in \mathcal{L}(\kappa_+)$. There exist elements $t, n \in \mathbb{N}$ such that $R_{\geq t}^n \subseteq I$. Since S is non-trivial there exists $s \in S$ with $\deg(s) \geq 1$ whence $s^{tn} \in R_{\geq t}^n$. This fact shows that $S \cap I \neq \emptyset$. ■

Lemma 5.1 says that if M is an R -module and S is a non-trivial left Ore set of R then $\kappa_+(M) \subseteq \kappa_S(M)$.

Lemma 5.2. *Let R be a positively SG left Noetherian ring and S a non-trivial good left Ore set. If M is an LSG R -module then $S^{-1}(M) \cong S^{-1}(Q_{\kappa_+}(M))$.*

Proof. Let

$$f: S^{-1}M \longrightarrow S^{-1}(M/\kappa_+(M)),$$

$$\frac{m}{s} \longmapsto \frac{\bar{m}}{s}.$$

It is clear that f is surjective. Let $\frac{m}{s} \in \text{Ker}(f)$. Then, $\frac{\bar{m}}{s} = 0$, and so, there exists $s' \in S$ such that $s'\bar{m} = 0$, i.e., $s'm \in \kappa_+(M) \subseteq \kappa_S(M)$. There exists $s'' \in S$ with $s''s'm = 0$ and since $s''s' \in S$ it follows that $\frac{m}{s} = 0$. Therefore, $S^{-1}(M) \cong S^{-1}(M/\kappa_+(M))$.

Now, let

$$g: S^{-1}(M/\kappa_+(M)) \longrightarrow S^{-1}(Q_{\kappa_+}(M)),$$

$$\frac{\bar{a}}{s} \longmapsto \frac{h(\bar{a})}{s},$$

where h is the isomorphism between $M/\kappa_+(M)$ and $\varphi_M(M)$. Since h is injective so g also is. Let $\frac{\xi}{s} \in S^{-1}(Q_{\kappa_+}(M))$. Then, there exist elements $t, n \in \mathbb{N}$ such that $(R_{\geq t}^n)\xi \subseteq \varphi_M(M)$. Since S is non-trivial repeating the argument above in the proof of Lemma 5.1 we can assert that there exists $s' \in S$ such that $s'' = (s')^{tn} \in R_{\geq t}^n$. In this way $s''\xi \in \varphi_M(M)$, and so, there exist $m \in M$ such that $s''\xi = \varphi_M(m)$, whence $g(\frac{\bar{m}}{s''s}) = \frac{h(\bar{m})}{s''s} = \frac{\varphi_M(m)}{s''s} = \frac{s''\xi}{s''s} = \frac{\xi}{s}$. We conclude that $S^{-1}(Q_{\kappa_+}(M)) \cong S^{-1}(M/\kappa_+(M)) \cong S^{-1}(M)$. ■

For the rest of this section, R denotes a schematic ring (recall that by Definition 4.1 R is left Noetherian). Consider the full subcategory $(R, \kappa_+) - \text{LSG}$ of $\text{LSG} - R$ whose objects are the κ_+ -closed modules. If M is an R -module κ_+ -closed and N is a submodule of M , then N is κ_+ -closed if and only if M/N is κ_+ -torsion-free [44, Proposition 4.2, Chapter IX]. Hence, it is clear that the intersection of κ_+ -closed modules is κ_+ -closed. This fact allows us to consider the submodule κ_+ -closed generated by a subset of M . If we define

$$N^c = \{x \in M \mid (N : x) \in \mathcal{L}(\kappa_+)\},$$

then it is clear that N^c is the submodule κ_+ -closed generated by N , and in fact $N^c = M$ if and only if M/N is κ_+ -torsion.

Notice that in the category $(R, \kappa_+) - \text{LSG}$ the subobjects are the submodules $\text{LSG-}\kappa_+$ -closed that are closed under arbitrary intersections. The submodule $\text{LSG-}\kappa_+$ -closed generated by $X \subseteq M$ will be denoted as $\langle X \rangle^{\text{SG-}\kappa}$. We will say that M is $\text{LSG-}\kappa_+$ -finitely generated if there exists a finite set $X \subseteq M$ with $\langle X \rangle^{\text{SG-}\kappa} = M$. Let $\text{Proj}R$ be the full subcategory of $(R, \kappa_+) - \text{LSG}$ consisting of $\text{LSG-}\kappa_+$ -finitely generated modules.

Proposition 5.3. *If N is an SG submodule of M , then N^c also is.*

Proof. Let $m = m_1 + \cdots + m_k \in N^c$ with $m_i \in M_{I_i}$. There exists $I \in \mathcal{L}(\kappa_+)$ such that $I \subseteq (N : m)$. Since R is schematic by the good left Ore set S_i for $i = 1, \dots, n$, say, then there exist elements $s_i \in S_i$ with $\sum R s_i \subseteq I$, whence $s_i m \in N$ for all i . Since N is SG and $s_i \in R''$ then $s_i m_j \in N$ for each i, j . Thus, $\sum_{i=1}^n R s_i \subseteq (N : m_j)$ which shows that $m_j \in N^c$. ■

From these facts, we have the equality $\langle X \rangle^{\text{SG-}\kappa} = (\langle X \rangle^{\text{SG}})^c$ for each $X \subseteq M$. In this way, M is $\text{LSG-}\kappa_+$ -finitely generated if and only if there exists a finite set $X \subseteq M$ such that $(\langle X \rangle^{\text{SG}})^c = M$, or equivalently, M/M_1 is κ_+ -torsion with $M_1 = \langle X \rangle^{\text{SG}}$.

We define the notion of *noncommutative site*.

Definition 5.4. Let \mathcal{O} be the set of non-trivial good left Ore sets of R and \mathcal{W} the free monoid on \mathcal{O} . We define the category $\underline{\mathcal{W}}$ as follows: the objects of $\underline{\mathcal{W}}$ are the elements

of \mathcal{W} , while for two words W and W' we define the morphisms of $\underline{\mathcal{W}}$, denoted by $\text{Hom}(W', W)$, as a singleton $\{W' \rightarrow W\}$ if there exists an increasing injection from the letters of W to the letters of W' , i.e., $W = S_1 \dots S_n$ and $W' = V_0 S_1 V_1 S_2 V_2 \dots S_n V_n$ for some letters S_i and some (possibly empty) words V_i . In other cases, $\text{Hom}(W', W)$ is defined to be empty.

It is easy to see that $\underline{\mathcal{W}}$ is a *thin category* (i.e., a category where between two objects there is at most one morphism). We denote the empty word as 1, which is the final object of the category.

If $W = S_1 \dots S_n \in \mathcal{W} \setminus \{1\}$ and M is an LSG R -module, we define

$$Q_W(M) = S_n^{-1} R \otimes_R \dots \otimes_R S_1^{-1} R \otimes_R M.$$

Lemma 5.2 asserts that if $W \neq 1$ then $Q_W(M) \cong Q_W(Q_{\kappa_+}(M))$.

If $W = S_1 \dots S_n \in \mathcal{W} \setminus \{1\}$, we say that $w \in W$ if $w = s_1 \dots s_n$ with $s_i \in S_i$. We associate a set of left ideals to W , namely,

$$\mathcal{L}(W) = \{I \triangleleft_l R \mid \text{there exists } w \in W \text{ such that } w \in I\}.$$

Let $\mathcal{L}(1) := \mathcal{L}(\kappa_+)$.

Lemma 5.5. *Let $W \in \mathcal{W} \setminus \{1\}$ and $w, w' \in W$. Then, there exists $w'' \in W$ such that $w'' = aw$ and $w'' = bw'$ for some elements $a, b \in R$.*

Proof. We prove the assertion by induction on the length of elements of W . If $W = S_1$, then by Ore's condition, there exist elements $a \in R$ and $b \in S_1$ such that $aw = bw' \in S_1$.

Let the assertion hold for every element of length k . Let $W = S_1 \dots S_{k+1}$, $\tilde{W} = S_2 \dots S_{k+1}$, $w = s_1 \dots s_{k+1}$, $w' = s'_1 \dots s'_{k+1} \in W$, $x = s_2 \dots s_{k+1}$, and $x' = s'_2 \dots s'_{k+1}$. By the inductive step, there exist elements $a, b \in R$ such that $ax = bx' \in \tilde{W}$. Since S_1 is a left Ore set then there exist $s''_1 \in S_1$ and $a_1 \in R$ such that $a_1 s_1 = s''_1 a$. Hence, $a_1 w = a_1 s_1 x = s''_1 a x = s''_1 b x' \in W$. Again, by the Ore's condition, there exist $s^*_1 \in S_1$ and $b_1 \in R$ such that $b_1 s'_1 = s^*_1 s''_1 b$, and so, $b_1 w' = b_1 s'_1 x' = s^*_1 s''_1 b x' = s^*_1 a_1 w \in W$. ■

Remark 5.6. Lemma 5.5 can be extended to a finite collection of words, i.e. if $w_1, \dots, w_i \in W$ then there exist $a_1, \dots, a_n \in R$ such that $a_1 w_1 = a_2 w_2 = \dots = a_n w_n \in W$.

Lemma 5.7. *Let $W \in \mathcal{W} \setminus \{1\}$, $w \in W$ and $a \in R$. There exist elements $w' \in W$ and $b \in R$ with $w'a = bw$.*

Proof. We prove by induction on the length of words of W . If $W = S_1$, then the assertion is precisely the Ore's condition.

Suppose that the lemma holds for each element of length k . Let $W = S_1 \dots S_{k+1}$, $\tilde{W} = S_2 \dots S_{k+1}$, $w = s_1 \dots s_{k+1}$, and $x = s_2 \dots s_{k+1}$. By the inductive step, there exist elements $x' \in \tilde{W}$ and $b \in R$ such that $x'a = bx$. Since S_1 is an Ore set there exist $s'_1 \in S_1$ and $b' \in R$ such that $s'_1 b = b' s_1$, whence $s'_1 x'a = s'_1 b x = b' s_1 x = b' w$. ■

Lemmas 5.5 and 5.7 allow us to conclude that $\mathcal{L}(W)$ is a filter. In the case $W \neq 1$ we will call κ_W the *pre-radical* associated to $\mathcal{L}(W)$. It can be seen that for every LSG R -module M the following equality holds:

$$\kappa_W(M) = \{m \in M \mid \text{there exists } w \in W \text{ such that } wm = 0\} = \text{Ker}(M \rightarrow Q_W(M)).$$

Following [33, page 113] a *global cover* is a finite subset $\{W_i \mid i \in I\}$ of \mathcal{W} such that $\bigcap_{i \in I} \mathcal{L}(W_i) = \mathcal{L}(\kappa_+)$. For $W \in \mathcal{W}$, $\text{Cov}(W)$ is defined as the set of all sets of the morphisms of $\underline{\mathcal{W}}$ of the form $\{W_i \rightarrow W \mid i \in I\}$, where $\{W_i \mid i \in I\}$ is a global cover. It is clear that $\{1\}$ is a global cover that will be called trivial. Notice that the schematic condition guarantees the existence of at least one non-trivial global cover. This collection of coverings is not a Grothendieck topology of $\underline{\mathcal{W}}$ but satisfies similar conditions (Proposition 5.9) that allow us to talk about sheaves on $\underline{\mathcal{W}}$. For this reason, Van Oystaeyen and Willaert called the category $\underline{\mathcal{W}}$ with this coverings the *noncommutative site* (cf. [51]).

The proof of the following lemma is analogous to the setting of graded rings [33, Lemma 1]. We include it for the completeness of the paper.

Lemma 5.8. *If $\{W_i \mid i \in I\}$ is a global cover, then for all $V \in \mathcal{W}$ we have that*

$$\bigcap_{i \in I} \mathcal{L}(W_i V) = \mathcal{L}(V).$$

Proof. If $I \in \mathcal{L}(V)$, there exists $v \in V$ such that $v \in I$. For $w_i \in W_i$, we have that $w_i v \in W_i V$ and $w_i v \in I$, and so, $I \in \mathcal{L}(W_i V)$. It follows that $\mathcal{L}(V) \subseteq \bigcap_{i \in I} \mathcal{L}(W_i V)$.

Let $I \in \bigcap_{i \in I} \mathcal{L}(W_i V)$. For each i , there exist $v_i \in V$ and $w_i \in W_i$ such that $w_i v_i \in I$. By Remark 5.6, there exist $a_1, \dots, a_n \in R$ and $v \in V$ such that $v = a_i v_i$ for every i . Lemma 5.7 guarantees that there exist $w'_i \in W_i$ and $b_i \in R$ with $w'_i a_i = b_i w_i$. Since $w'_i \in \sum R w'_i$ and $\{W_i \mid i \in I\}$ is a global cover there exist elements $n, t \in \mathbb{N}$ such that $R_{\geq t}^n \subseteq \sum R w'_i$. Multiplying by v we obtain $(R_{\geq t}^n) v \subseteq I$. If S is the first letter of V , there exists $s \in S \cap R_{\geq t}^n$. Finally, $sv \in I$ and $sv \in V$, and thus, $I \in \mathcal{L}(V)$. ■

Proposition 5.9. *The category $\underline{\mathcal{W}}$ together with the sets $\text{Cov}(W)$ for any element $W \in \underline{\mathcal{W}}$ satisfies the following properties:*

- (G₁) $\{W \rightarrow W\} \in \text{Cov}(W)$,
- (G₂) $\{W_i \rightarrow W \mid i \in I\} \in \text{Cov}(W)$ and for all $i \in I : \{W_{ij} \rightarrow W_i \mid j \in I_i\} \in \text{Cov}(W_i)$ we have that $\{W_{ij} \rightarrow W_i \rightarrow W \mid i \in I, j \in I_i\} \in \text{Cov}(W)$,
- (G₃) if $\{W_i \rightarrow W \mid i \in I\} \in \text{Cov}(W)$, $W' \rightarrow W \in \underline{\mathcal{W}}$, and if we define $W_i W \times_W W' = W_i W'$, then $\{W_i W \times_W W' \rightarrow W' \mid i \in I\} \in \text{Cov}(W')$.

Proof. (G₁) holds since $\{1\}$ is a global cover. (G₂) is a direct consequence of Lemma 5.8. (G₃) is clear. ■

Definitions 5.10 and 5.11 introduce the notion of presheaf and sheaf, respectively, in the setting of the category $\underline{\mathcal{W}}$.

Definition 5.10. A presheaf \mathcal{F} on $\underline{\mathcal{W}}$ is a contravariant functor from $\underline{\mathcal{W}}$ to the category $\text{LSG} - R$ such that for all $W \in \underline{\mathcal{W}} \setminus \{1\}$ the sections $\mathcal{F}(W)$ of \mathcal{F} on W are SG $S^{-1}R$ -modules, where S denotes the last letter of W and $\mathcal{F}(1)$ is an SG $Q_{\kappa_+}(R)$ -module.

Since $\mathcal{F}(1)$ denotes the global sections, we will denote it as $\Gamma_*(\mathcal{F})$. We abbreviate $\mathcal{F}(V \rightarrow W)$ as $\rho_V^W : \mathcal{F}(W) \rightarrow \mathcal{F}(V)$. If $W = 1$, then we will write ρ_V instead of ρ_V^1 .

Definition 5.11. A presheaf \mathcal{F} on $\underline{\mathcal{W}}$ is a *sheaf* if it satisfies the following two properties:

- (i) *Separatedness*: for all elements $W \in \mathcal{W}$ and each global cover $\{W_i \mid i \in I\}$, if $m \in \mathcal{F}(W)$ satisfies that for every $i \in I$ we have that $\rho_{W_i W}^W(m) = 0$ in $\mathcal{F}(W_i W)$, then $m = 0$,
- (ii) *Gluing*: $\forall W \in \mathcal{W}$ and each global cover $\{W_i \mid i \in I\}$, given $(m_i) \in \prod_i \mathcal{F}(W_i W)$ satisfying

$$\rho_{W_i W_j W}^{W_i W}(m_i) = \rho_{W_i W_j W}^{W_j W}(m_j) \quad \text{for all } (i, j) \in I \times I,$$

there exists an element $m \in \mathcal{F}(W)$ such that

$$\rho_{W_i W}^W(m) = m_i \quad \text{for all } i \in I.$$

Proposition 5.12. A presheaf \mathcal{F} is a sheaf if and only if for every word W and each global cover $\{W_i \mid i \in I\}$, $\mathcal{F}(W)$ (with the arrows given by \mathcal{F}) is the limit of the diagram

$$\begin{array}{ccc} \mathcal{F}(W_i W) & \longrightarrow & \mathcal{F}(W_i W_j W) \\ & \searrow & \nearrow \\ \mathcal{F}(W_j W) & \longrightarrow & \mathcal{F}(W_j W_i W) \end{array} \quad (5.1)$$

Proof. Suppose that \mathcal{F} is a sheaf. Let M be an SG R -module with morphisms $f_i : M \rightarrow \mathcal{F}(W_i W)$ which are compatibles with the morphisms $\rho_{W_i W_j W}^{W_i W}$ and $\rho_{W_j W_i W}^{W_j W}$. Consider an element $m \in M$. By using this compatibility, we have that the element $(f_i(m))$ of $\prod_i \mathcal{F}(W_i W)$ satisfies the equality

$$\rho_{W_i W_j W}^{W_i W} f_i(m) = \rho_{W_i W_j W}^{W_j W} f_j(m) \quad \text{for all } (i, j) \in I \times I.$$

In this way, there is a unique element $m' \in \mathcal{F}(W)$ such that $\rho_{W_i W}^W(m') = f_i(m)$, for each $i \in I$. If we define the map $\beta : M \rightarrow \mathcal{F}(W)$ as $\beta(m) = m'$, then it is clear that β is a homogeneous R -homomorphism and it is the only one that satisfies the equality $\rho_{W_i W}^W \circ \beta = f_i$ for all $i \in I$. Hence, $\mathcal{F}(W)$ is the limit of the diagram (5.1).

On the other hand, suppose that $\mathcal{F}(W)$ is the limit of the diagram (5.1). Let $m \in \mathcal{F}(W)$ such that $\rho_{W_i W}^W(m) = 0$ for each $i \in I$. This means that $m \in \bigcap \text{Ker}(\rho_{W_i W}^W)$, and by assumption on $\mathcal{F}(W)$ we have that $\bigcap \text{Ker}(\rho_{W_i W}^W) = 0$, whence $m = 0$. Let

$$A := \left\{ (m_i) \in \prod \mathcal{F}(W_i W) \mid \rho_{W_i W_j W}^{W_i W}(m_i) = \rho_{W_i W_j W}^{W_j W}(m_j) \text{ for all } (i, j) \in I \times I \right\}.$$

It is clear that A is an SG R -submodule of $\prod \mathcal{F}(W_i W)$, which guarantees the existence of only one homogeneous R -homomorphism $\beta : A \rightarrow \mathcal{F}(W)$ such that $\rho_{W_i W}^W \circ \beta = \pi_i$ for all $i \in I$, where π_i denotes the usual projection. In this way, $\beta(m_i)$ is the element that satisfies the gluing condition. ■

Definition 5.13. Let M be an LSG R -module. We define the presheaf \hat{M} in the following way: for objects, we have that $\hat{M}(1) = Q_{\kappa_+}(M)$, and for $W \in \mathcal{W} \setminus \{1\}$, we have that $\hat{M}(W) = Q_W(M)$. Now, for morphisms, if $W \neq 1$ then to the map $V \rightarrow W$ we assign it the canonical morphism such that the following diagram commutes:

$$\begin{array}{ccc} Q_W(M) & \xrightarrow{\rho_V^W} & Q_V(M) \\ \uparrow & \nearrow & \\ M & & \end{array}$$

while for the morphism $W \rightarrow 1$, we assign the composition map

$$Q_{\kappa_+}(M) \rightarrow Q_W(Q_{\kappa_+}(M)) \rightarrow Q_W(M),$$

where the first arrow is the natural map and the second arrow is precisely the isomorphism obtained in Lemma 5.2.

If $W = S_1 \dots S_n$ and $w \in W$, say $w = s_1 \dots s_n$, then the element $\frac{1}{s_n} \otimes \dots \otimes \frac{1}{s_1} \otimes m \in Q_W(M)$ will be denoted as $\frac{m}{w}$. In particular, $\frac{m}{1}$ stands for $1 \otimes m$ in $Q_S(M)$, for $1 \otimes 1 \otimes m$ in $Q_{ST}(M)$, and so on, which element is meant depends on the module it belongs to.

The proofs of the following two lemmas follow the same ideas as those presented in the setting of \mathbb{N} -graded rings [33, Lemmas 2 and 3].

Lemma 5.14. *Given elements $\frac{m}{w} \in Q_W(M)$ and $a \in R$, there exist $w' \in W$ and $b \in R$ such that $w'a = bw$ and $a\frac{m}{w} = \frac{bm}{w'} \in Q_W(M)$.*

Proof. Let $W = S_1 \dots S_n$ and $w = s_1 \dots s_n$. We consider $a_n = a$ and define a_i recursively. More exactly, for an element a_i the Ore's condition guarantees the existence of elements $s'_i \in S_i$ and $a_{i-1} \in R$ such that $s'_i a_i = a_{i-1} s_i$. Hence, $a\frac{m}{w} = \frac{1}{s_n} \otimes \dots \otimes \frac{1}{s_1} \otimes a_0 m$. If we define $b = a_0$ and $w' = s'_1 \dots s'_n$, then the assertion follows. ■

Lemma 5.15. *If $\frac{m}{w} = \frac{m'}{1}$ in $Q_W(M)$ for some element $m' \in M$, then there exist $\tilde{w} \in W$ and $r \in R$ such that $\tilde{w} = rw$ and $\tilde{w}m' = rm$.*

Proof. Induction on the length n of the word $w = s_1 \dots s_n$. The case $n = 1$ is clear. Suppose that the assertion holds for any word of length less than n . Let $W' = S_1 \dots S_{n-1}$ and $w' = s_1 \dots s_{n-1}$. Then, $\frac{1}{s_n} \otimes \frac{m}{w'} = 1 \otimes \frac{m'}{1} \in S_n^{-1}(Q_{W'})$, so that there exist elements $a, b \in R$ such that $a\frac{m}{w'} = b\frac{m'}{1} = \frac{bm'}{1} \in Q_{W'}(M)$ and $as_n = b \in S_n$. By Lemma 5.14, there exist elements $w'' \in W'$ and $c \in R$ with $w''a = cw'$ and $\frac{cm}{w''} = a\frac{m}{w'} = \frac{bm'}{1}$. By hypothesis, there exist $w''' \in W'$ and $d \in R$ such that $w''' = dw''$ and $w'''bm' = dcm$, so that if we consider $\tilde{w} = w'''b$ and $x = dc$ the assertion follows. ■

Let $\{W_i \mid i \in I\}$ be a global cover. The limit of the diagram

$$\begin{array}{ccc}
 Q_W(Q_{W_i}(M)) & \longrightarrow & Q_W(Q_{W_j}(Q_{W_i}(M))) \\
 & \searrow & \nearrow \\
 Q_W(Q_{W_j}(M)) & \longrightarrow & Q_W(Q_{W_i}(Q_{W_j}(M)))
 \end{array} \quad (5.2)$$

will be denoted by $\Gamma_W(\hat{M})$. Notice that due to the universal property of the limit, for the family $\{M \rightarrow Q_{W_i}(M) \mid i \in I\}$ there is a unique morphism $\varphi : M \rightarrow \Gamma_1(\hat{M})$. This morphism is of great importance in the following lemma.

Lemma 5.16. *Let $\varphi : M \rightarrow \Gamma_1(\hat{M})$ the morphism described above. Then, $\text{Coker}(\varphi)$ is κ_+ -torsion.*

Proof. Let $\xi = (\frac{m_i}{w_i})_i \in \Gamma_1(\hat{M})$ with $w_i \in W_i$ and $\frac{1}{w_i} \otimes 1 \otimes m_i = 1 \otimes \frac{1}{w_j} \otimes m_j \in Q_{W_i}(Q_{W_j}(M)) \forall i, j$. Fix j . Then, $\frac{1 \otimes m_i}{w_i} = \frac{\frac{1}{w_j} \otimes m_j}{1} \in Q_{W_i}(M)$ whence by Lemma 5.15, there exist elements $w'_i \in W_i$ and $a_i \in R$ such that $w'_i = a_i w_i$ and $w'_i(\frac{1}{w_j} \otimes m_j) = a_i(1 \otimes m_i)$, that is, $w'_i(\frac{m_j}{w_j}) = \frac{a_i m_i}{1} \in Q_{W_j}(M)$. By taking $I := \sum_{i \in I} R w'_i$, it is clear that $I \in \bigcap_i \mathcal{L}(W_i) = \mathcal{L}(\kappa_+)$. There exist elements $t_j, n_j \in \mathbb{N}$ such that $R_{\geq t_j}^{n_j} \subseteq I$, which shows that $(R_{\geq t_j}^{n_j}) \frac{m_j}{w_j}$ is contained in the direct image of the map $M \rightarrow Q_{W_j}(M)$, that is, $(R_{\geq t_j}^{n_j}) \frac{m_j}{w_j} \subseteq \text{Im}(M \rightarrow Q_{W_j}(M))$. If $n := \max\{n_i \mid i \in I\}$ and $t := \max\{t_i \mid i \in I\}$, then it is straightforward to see that this reasoning is true for every element j .

Let $a \in R_{\geq t}^n$. Then, $a\xi = (\frac{n_i}{1})_i$ for some elements $n_i \in M$ with $1 \otimes 1 \otimes n_i = 1 \otimes 1 \otimes n_j$ in $Q_{W_i}(Q_{W_j}(M))$ for every i, j . Fix i . Lemma 5.15 guarantees that for each j there exist elements $\tilde{w}_j \in W_i$ and $x_j \in R$ such that $\tilde{w}_j = x_j$ and $\tilde{w}_j \frac{n_i}{1} = x_j \frac{n_j}{1}$. Now, by Remark 5.6, we can find elements $w_i^* \in W_i$ with $w_i^* \frac{n_i}{1} = w_i^* \frac{n_j}{1}$ for all j . Hence, $w_i^* a\xi = \varphi(w_i^* n_i)$. As above, by defining $J = \sum_{i \in I} R w_i^*$, there exist elements $t'(a)$ and $n'(a)$ (notice that all elements depend on a) such that $(R_{\geq t'(a)}^{n'(a)}) a\xi \subseteq \varphi(M)$. Since R is a left Noetherian ring, $R_{\geq t}^n$ is finitely generated, say by the elements a_1, \dots, a_r . By defining, $n' = \max\{n(a_k) \mid 1 \leq k \leq r\}$ and $t' = \max\{t'(a_k) \mid 1 \leq k \leq r\}$ we have that $(R_{\geq t+t'}^{n+n'}) \xi \subseteq \varphi(M)$, i.e., $\text{Coker}(\varphi)$ is κ_+ -torsion, which concludes the proof. ■

Proposition 5.17. *The presheaf \hat{M} is a sheaf.*

Proof. Fix a global cover $\{W_i \mid i \in I\}$ and let $\varphi : M \rightarrow \Gamma_1(\hat{M})$ be the map established in Lemma 5.16. Let us see that $\Gamma_1(\hat{M}) \cong Q_{\kappa_+}(M) = \hat{M}(1)$. Since for the family $\{Q_{\kappa_+}(M) \rightarrow Q_{W_i} \mid i \in I\}$ the universal property of the limit guarantees the existence of a unique morphism $\phi : Q_{\kappa_+}(M) \rightarrow \Gamma_1(\hat{M})$, we obtain the following commutative diagram:

$$\begin{array}{ccccc}
 & & \Gamma_1(\hat{M}) & \longrightarrow & Q_{W_i}(M) \longrightarrow Q_{W_j}(Q_{W_i}(M)) \\
 & \nearrow \varphi & \uparrow \phi & \nearrow \phi_i & \\
 M & \longrightarrow & Q_{\kappa_+}(M) & &
 \end{array} \quad (5.3)$$

It is clear that $\text{Ker}(\phi) \subseteq \bigcap \text{Ker}(\phi_i) = \bigcap \kappa_{W_i}(Q_{\kappa_+}(M)) = \kappa_+(Q_{\kappa_+}(M)) = 0$. On the other hand, since $\text{Im}(\varphi) \subseteq \text{Im}(\phi)$ and $\Gamma_1(\widehat{M})/\text{Im}(\varphi)$ is κ_+ -torsion (Lemma 5.16) it follows that $\Gamma_1(\widehat{M})/\text{Im}(\phi)$ is κ_+ -torsion also. Besides, if S_i is the last letter of W_i then $Q_{W_i}(M)$ is κ_{S_i} -torsion-free, and so, it is κ_+ -torsion-free. In this way, $\Gamma_1(\widehat{M})$ is the limit of objects that are κ_+ -torsion-free, and it is clear that $\Gamma_1(\widehat{M})$ is κ_+ -torsion-free also. Since we have the short exact sequence

$$0 \rightarrow Q_{\kappa_+}(M) \rightarrow \Gamma_1(\widehat{M}) \rightarrow \Gamma_1(\widehat{M})/\text{Im}(\phi) \rightarrow 0,$$

it follows that $Q_{\kappa_+}(M) \cong \Gamma_1(\widehat{M})$ [16, Proposition 3.4].

Finally, by recalling that Q_W is an exact functor that commutes with finite limits, if $W \neq 1$, then we have that

$$\Gamma_W(\widehat{M}) \cong Q_W(\Gamma_1(\widehat{M})) \cong Q_W(Q_{\kappa_+}(M)) \cong Q_W(M) = \widehat{M}(W).$$

By Remark 5.12, it follows that \widehat{M} is a sheaf. ■

Next, we define the notion of affine cover and quasi-coherent sheaf.

Definition 5.18. An *affine cover* is a finite subset $\{T_i \mid i \in I\}$ of \mathcal{O} such that $\bigcap_{i \in I} \mathcal{L}(T_i) = \mathcal{L}(\kappa_+)$.

Definition 5.19. A sheaf \mathcal{F} is *quasi-coherent* if there exists an affine cover $\{T_i \mid i \in I\}$, and for each $i \in I$ there exists an SG $T_i^{-1}R$ -module M_i such that for all morphisms $V \rightarrow W$ in the category $\underline{\mathcal{W}}$, we have a commutative diagram given by

$$\begin{array}{ccc} \mathcal{F}(T_i W) & \xrightarrow{\rho_{T_i V}^{T_i W}} & \mathcal{F}(T_i V) \\ \downarrow & & \downarrow \\ Q_W(M_i) & \longrightarrow & Q_V(M_i) \end{array} \quad (5.4)$$

where the vertical maps are isomorphisms in $\text{LSG} - R$ and $Q_1(*) := Q_{\kappa_+}(*)$. \mathcal{F} is called *coherent* if moreover all M_i are finitely generated SG $T_i^{-1}R$ -modules.

Remark 5.20. Note that the sheaf \widehat{M} is quasi-coherent for each object in the category $\text{LSG} - R$. If M is finitely generated SG module, then \widehat{M} is coherent.

The proof of the following proposition is analogous to the proof of [33, Theorem 1]. For the completeness of the paper, we include it here.

Proposition 5.21. If \mathcal{F} is a quasi-coherent sheaf on \mathcal{W} and $\Gamma_*(\mathcal{F})$ denotes its global sections $\mathcal{F}(1)$, then \mathcal{F} is isomorphic to $\widehat{\Gamma_*(\mathcal{F})}$, the sheaf associated to $\Gamma_*(\mathcal{F})$.

Proof. First of all, notice that we can suppose that M_i is κ_+ -closed because if this is not the case then we can replace it by $Q_{\kappa_+}(M_i)$ and the commutative diagram 5.4 holds. We want to see that $\mathcal{F}(W) \cong Q_W(\Gamma_*(\mathcal{F}))$ for all $W \in \mathcal{W}$. If $W \neq 1$ then by Remark 5.12

and the fact that Q_W commutes with finite limits (recall that Q_W is an exact functor), it follows that $\mathcal{F}(W)$ is the limit of the diagram (5.1), while $Q_W(\Gamma_*(\mathcal{F}))$ is the limit of the diagram

$$\begin{array}{ccc} Q_W(\mathcal{F}(W_i)) & \longrightarrow & Q_W(\mathcal{F}(W_i W_j)) \\ & \searrow & \nearrow \\ Q_W(\mathcal{F}(W_j)) & \longrightarrow & Q_W(\mathcal{F}(W_j W_i)) \end{array} \quad (5.5)$$

Notice that we have the isomorphism $\mathcal{F}(T_i) \cong Q_{\kappa_+}(M_i) = M_i$, and by the diagram 5.4, for every W there exists an isomorphism $\psi_i^W : Q_W(\mathcal{F}(T_i)) \rightarrow \mathcal{F}(T_i W)$. If $W = S_1 \dots S_n$ and $W_t := S_1 \dots S_t$, then we obtain the following commutative diagram:

$$\begin{array}{ccccccc} \mathcal{F}(T_i) & \xrightarrow{\rho_{T_i W_1}^{T_i}} & \mathcal{F}(T_i W_1) & \xrightarrow{\rho_{T_i W_2}^{T_i W_1}} & \mathcal{F}(T_i W_2) & \longrightarrow \dots \longrightarrow & \mathcal{F}(T_i W_{n-1}) & \xrightarrow{\rho_{T_i W}^{T_i W_{n-1}}} & \mathcal{F}(T_i W) \\ \downarrow & & \downarrow \psi_i^{W_1} & & \downarrow \psi_i^{W_2} & & \downarrow \psi_i^{W_{n-1}} & & \downarrow \psi_i^W \\ \mathcal{F}(T_i) & \longrightarrow & Q_{W_1}(\mathcal{F}(T_i)) & \longrightarrow & Q_{W_2}(\mathcal{F}(T_i)) & \longrightarrow \dots \longrightarrow & Q_{W_{n-1}}(\mathcal{F}(T_i)) & \longrightarrow & Q_W(\mathcal{F}(T_i)) \end{array}$$

Since $Q_{W_t}(\mathcal{F}(T_i))$ is an $S_t^{-1}R$ -module, and so, $\mathcal{F}(T_i W_t)$ also is, for an element $s_t \in S_t$, we can multiply by s_t^{-1} , whence the commutativity of the diagram above guarantees that

$$\begin{aligned} & \psi_i^W \left(\frac{1}{s_n} \otimes \dots \otimes \frac{1}{s_1} \otimes m \right) \\ &= s_n^{-1} \rho_{T_i W}^{T_i W_{n-1}} \left(s_{n-1}^{-1} \rho_{T_i W_{n-1}}^{T_i W_{n-2}} \left(\dots s_2^{-1} \rho_{T_i W_2}^{T_i W_1} (s_1^{-1} \rho_{T_i W_1}^{T_i} (m)) \dots \right) \right). \end{aligned} \quad (5.6)$$

On the other hand, we have that

$$\mathcal{F}(T_i T_j W) \cong Q_{T_j W}(\mathcal{F}(T_i)) = Q_W(Q_{T_j}(\mathcal{F}(T_i))) \cong Q_W(\mathcal{F}(T_i T_j)).$$

If we write $\psi_{ij}^W : Q_W(\mathcal{F}(T_i T_j)) \rightarrow \mathcal{F}(T_i T_j W)$ as the isomorphism above, by using a similar diagram to the above, it can be seen that

$$\begin{aligned} & \psi_{ij}^W \left(\frac{1}{s_n} \otimes \dots \otimes \frac{1}{s_1} \otimes m \right) \\ &= s_n^{-1} \rho_{T_i T_j W}^{T_i T_j W_{n-1}} \left(s_{n-1}^{-1} \rho_{T_i T_j W_{n-1}}^{T_i T_j W_{n-2}} \left(\dots s_2^{-1} \rho_{T_i T_j W_2}^{T_i T_j W_1} (s_1^{-1} \rho_{T_i T_j W_1}^{T_i} (m)) \dots \right) \right). \end{aligned} \quad (5.7)$$

Notice that $\rho_{T_i T_j W_t}^{T_i W_t}$ and $\rho_{T_j T_i W_t}^{T_i W_t}$ are $S_t^{-1}R$ -linear for $t = 1, \dots, n$, since both are R -homomorphisms between $S_t^{-1}R$ -modules. In this way, the expressions (5.6) and (5.7) imply the commutativity of the following two diagrams:

$$\begin{array}{ccc} Q_W(\mathcal{F}(T_i)) & \xrightarrow{Q_W(\rho_{T_i T_j}^{T_i})} & Q_W(\mathcal{F}(T_i T_j)) \\ \downarrow \psi_i & & \downarrow \psi_{ij} \\ \mathcal{F}(T_i W) & \xrightarrow{\rho_{T_i T_j W}^{T_i W}} & \mathcal{F}(T_i T_j W) \end{array} \quad \begin{array}{ccc} Q_W(\mathcal{F}(T_i)) & \xrightarrow{Q_W(\rho_{T_j T_i}^{T_i})} & Q_W(\mathcal{F}(T_j T_i)) \\ \downarrow \psi_i & & \downarrow \psi_{ji} \\ \mathcal{F}(T_i W) & \xrightarrow{\rho_{T_j T_i W}^{T_i W}} & \mathcal{F}(T_j T_i W) \end{array}$$

Hence, it is clear that diagrams (5.1) and (5.5) have isomorphic limits, that is, $\mathcal{F} \cong Q_W(\mathcal{F}(\Gamma_*))$. Besides, for a morphism $V \rightarrow W$ the map ρ_V^W is determined by the maps $\rho_{T_i V}^{T_i W}$ and $\rho_{T_i T_j V}^{T_i T_j W}$, which shows that the diagram

$$\begin{array}{ccc} Q_W(\Gamma_*(\mathcal{F})) & \longrightarrow & Q_V(\Gamma_*(\mathcal{F})) \\ \downarrow & & \downarrow \\ \mathcal{F}(W) & \xrightarrow{\rho_V^W} & \mathcal{F}(V) \end{array}$$

is commutative.

For $W = 1$, we have to show that $\Gamma_*(\mathcal{F}) \cong Q_{\kappa_+}(\Gamma_*(\mathcal{F}))$. Since $\mathcal{F}(T_i)$ and $\mathcal{F}(T_i T_j)$ are κ_+ -torsion-free ($\mathcal{F}(T_i) \cong M_i$ and $\mathcal{F}(T_i T_j) \cong T_j^{-1}(\mathcal{F}(T_i))$), then $\Gamma_*(\mathcal{F})$ is κ_+ -torsion-free because it is the limit of objects κ_+ -torsion-free. Let us see that $\Gamma_*(\mathcal{F})$ is κ_+ -injective. By [16, Proposition 3.2], it is sufficient to show that for all $I \in \mathcal{L}(\kappa_+)$ every R -homomorphism $f : I \rightarrow \Gamma_*(\mathcal{F})$ can be extended to a R -homomorphism $g : R \rightarrow \Gamma_*(\mathcal{F})$.

Since $\mathcal{F}(T_i)$ is κ_+ -injective the map $\rho_{T_i} \circ f$ can be extended to a map $g_i : R \rightarrow \mathcal{F}(T_i)$. If $x_i = g_i(1)$, then $g_i(r) = r x_i$ for every $r \in R$. In particular, for each $a \in I$ we have that $a \rho_{T_i T_j}^{T_i}(x_i) = \rho_{T_i T_j}^{T_i}(\rho_{T_i}(f(a))) = \rho_{T_i T_j}^{T_j}(\rho_{T_j}(f(a))) = a \rho_{T_i T_j}^{T_j}(x_j)$, which shows that there exists an element $x \in \Gamma_*(\mathcal{F})$ such that $\rho_{T_i}(x) = x_i$ for every i . Notice that the map $g : I \rightarrow \Gamma_*(\mathcal{F})$ defined by $g(r) = r x$ extends f , so we conclude $\Gamma_*(\mathcal{F}) = Q_{\kappa_+}(\Gamma_*(\mathcal{F}))$. ■

Theorem 5.22. *The category of quasi-coherent sheaves is equivalent to the category $(R, \kappa_+) - \text{LSG}$.*

Proof. Let \mathcal{F} be a quasi-coherent sheaf. From the last part of the proof of the Proposition 5.21 we have that $\Gamma_*(\mathcal{F})$ is an object of $(R, \kappa_+) - \text{LSG}$. Moreover, if M belongs to $(R, \kappa_+) - \text{LSG}$ then $M \cong Q_{\kappa_+}(M)$. In this way, $\widehat{\square}$ and $\Gamma_* \square$ are functors between the category $(R, \kappa_+) - \text{LSG}$ and the category of quasi-coherent sheaves, which are equivalent by Propositions 5.17 and 5.21. ■

By taking the *category of coherent sheaves* $\text{coh}R$ as the full subcategory of quasi-coherent sheaves that consists of coherent sheaves, and having in mind that $\text{Proj}R$ is the full subcategory of $(R, \kappa_+) - \text{LSG}$ consisting of $\text{LSG-}\kappa_+$ -finitely generated modules, we arrive at the most important result of the paper: the *Serre–Artin–Zhang–Verevkin theorem* for semi-graded rings.

Theorem 5.23 (Serre–Artin–Zhang–Verevkin theorem). *The category of coherent sheaves is equivalent to $\text{Proj}R$.*

Proof. From Theorem 5.22, it is sufficient to show that M belongs to $\text{Proj}R$ if and only if \widehat{M} is coherent.

We fix a cover $\{T_i \mid i \in I\}$. If M is an element of $\text{Proj}R$, then there exist $m_1, \dots, m_k \in M$ such that M/N is κ_+ -torsion with $N = \langle m_1, \dots, m_k \rangle^{\text{SG}}$. Let $f_i : M \rightarrow T_i^{-1}M$ be the canonical map. It is straightforward to see that $\langle \frac{m_1}{1}, \dots, \frac{m_k}{1} \rangle_R^{\text{SG}} = f_i(N)$, and that the

$T_i^{-1}M$ -submodule $J_i = \langle \frac{m_1}{1}, \dots, \frac{m_k}{1} \rangle_{T_i^{-1}R}^{\text{SG}}$ satisfies the relation $f_i(N) \subseteq J_i$. Let $m \in M$. Since M/N is κ_+ -torsion there exists $I \in \mathcal{L}(\kappa_+)$ such that $Im \subseteq N$. By using that T_i is non-trivial, there exists $t_i \in I \cap T_i$, whence $t_i m \in N$, which shows that $\frac{t_i m}{1} \in f_i(N)$. Since J_i is a $T_i^{-1}R$ -module and $\frac{t_i m}{1} \in J_i$ then $J_i = T_i^{-1}M$, that is, $T_i^{-1}M$ is finitely generated as an SG $T_i^{-1}R$ -module.

Suppose that every one of the $T_i^{-1}M$ is a finitely generated SG $T_i^{-1}R$ -module. Note that if $T_i^{-1}M = \langle \frac{m_{1,i}}{s_{1,i}}, \dots, \frac{m_{t_i,i}}{s_{t_i,i}} \rangle_{T_i^{-1}R}^{\text{SG}}$ then $T_i^{-1}M = \langle \frac{m_{1,i}}{1}, \dots, \frac{m_{t_i,i}}{1} \rangle_{T_i^{-1}R}^{\text{SG}}$. Since there are finite elements i 's, the union set $\bigcup_{i \in I} \{m_{1,i}, \dots, m_{t_i,i}\}$ is also finite, $\{m_1, \dots, m_k\}$ say. If we define $N = \langle m_1, \dots, m_k \rangle^{\text{SG}}$, it is straightforward to see that $T_i^{-1}N$ is an SG $T_i^{-1}R$ -submodule of $T_i^{-1}M$, whence $T_i^{-1}N = T_i^{-1}M$.

For an element $m \in M$, since $\frac{m}{1} \in T_i^{-1}N$ there exist elements $n_i \in N$ and $t_i \in T_i$ such that $\frac{m}{1} = \frac{n_i}{t_i}$. There exist $c_i, d_i \in R$ such that $c_i m = d_i n_i$ and $c_i = d_i t_i \in T_i$, whence $c_i m \in N$. By using that $c_i \in T_i$ for each $i \in I$ and that $\{T_i \mid i \in I\}$ is an affine cover it follows that $I = \sum Rc_i \in \mathcal{L}(\kappa_+)$. We conclude that $Im \subseteq N$, and therefore, M/N is κ_+ -torsion. ■

Next, we show that the notion of schematicness in the semi-graded setting generalizes the corresponding concept in the case of connected and \mathbb{N} -graded algebras introduced and studied by Van Oystaeyen and Willaert [33, 49, 50, 52, 58].

Remark 5.24. Consider a positively graded left Noetherian ring R . It is clear that $R_+ = R_{\geq 1}$. Note that if R is generated in degree one then $R_{\geq t} = (R_+)^t$, which shows that $\mathcal{L}(\kappa_+) = \{I \triangleleft_l R \mid \text{there exists } n \in \mathbb{N} \text{ with } (R_+)^n \subseteq I\}$. On the other hand, the LSG modules are exactly the same \mathbb{N} -graded modules and the good left Ore sets coincide with the homogeneous left Ore sets. In this way, the notion of schematic ring presented in this paper generalizes the corresponding notion introduced by Van Oystaeyen and Willaert [33]. Last, but not least, notice that in the \mathbb{N} -graded setting the left Noetherianity of R implies that the finitely generated objects of $(R, \kappa_+) - \text{LSG}$ are the Noetherian objects, which shows that Theorem 5.23 generalizes [33, Theorem 3].

We present some examples that illustrate our Theorem 5.23 in the case of non- \mathbb{N} -graded rings where [33, Theorem 3] cannot be applied.

Example 5.25. (i) Consider the first Weyl algebra $A_1(\mathbb{k}) = \mathbb{k}\{x, y\}/\langle yx - xy - 1 \rangle$ over a field \mathbb{k} of $\text{char}(\mathbb{k}) = p > 0$. It is well known that $A_1(\mathbb{k})$ is a non- \mathbb{N} -graded ring, the set $\{x^n y^m \mid n, m \in \mathbb{N}\}$ is \mathbb{k} -basis of $A_1(\mathbb{k})$, and that $A_1(\mathbb{k})$ is a Noetherian ring. Since $x^p, y^p \in Z(A_1(\mathbb{k}))$ it is clear that $\{x^{pk} \mid k \in \mathbb{N}\}$ and $\{y^{pk} \mid k \in \mathbb{N}\}$ are good left Ore sets. Besides, if $k_1, k_2 \in \mathbb{N}$ then $A_1(\mathbb{k})x^{pk_1} + A_1(\mathbb{k})y^{pk_2}$ is a two-sided ideal of $A_1(\mathbb{k})$ which is a left SG submodule, whence $A_1(\mathbb{k})_{\geq pk_1 + pk_2} \subseteq A_1(\mathbb{k})x^{pk_1} + A_1(\mathbb{k})y^{pk_2}$. Therefore, $A_1(\mathbb{k})$ is a schematic algebra and Theorem 5.23 holds.

(ii) In a similar way, it can be shown that the n th Weyl algebra $A_n(\mathbb{k})$ is schematicness when $\text{char}(\mathbb{k}) = p > 0$.

(iii) The well-known Jordan plane $\mathbb{k}\{x, y\}/\langle yx - xy - y^2 \rangle$ is schematic when $\text{char}(\mathbb{k}) = p > 0$ since the sets $\{x^{pk} \mid k \in \mathbb{N}\}$ and $\{y^{pk} \mid k \in \mathbb{N}\}$ are good left Ore sets.

For the skew PBW extensions introduced by Gallego and Lezama [15] (cf. [36, 39]), which are examples of non- \mathbb{N} -graded rings, Proposition 5.26 establishes sufficient conditions to guarantee their schematicness.

Proposition 5.26. *Let $A = \sigma(R)\langle x_1, \dots, x_n \rangle$ be a bijective skew PBW extension over a left Noetherian ring R with the usual semi-graduation, that is, $\deg(x_i) = 1$ and $\deg(r) = 0$, for every i and each $r \in R$. If for every i , there exists $m_i \geq 1$ such that $x_i^{m_i} \in Z(A)$ then A is schematic.*

Proof. From [14, Theorem 3.1.5] we know that A is left Noetherian. Since $x_i^{m_i} \in Z(A)$ it follows that $\{x_i^{m_i m} \mid m \in \mathbb{N}\}$ is a non-trivial good left Ore set for every i . Let us see that these sets satisfy the schematicness condition. Let $t_i \in \mathbb{N}$. Then, $\sum_{i=1}^n R x_i^{m_i t_i}$ is a two-sided ideal and an SG submodule of A . If $t := \sum m_i t_i$, then $\bigoplus_{m \geq t} R_m \subseteq \sum_{i=1}^n R x_i^{m_i t_i}$, and thus, $R_{\geq t} \subseteq \sum_{i=1}^n R x_i^{m_i t_i}$. ■

Examples 5.27 and 5.28 show that the theory presented by Lezama about Serre–Artin–Zhang–Verevkin theorem and the one developed in this paper are independent.

Example 5.27. Proposition 5.26 guarantees that if R is a left Noetherian noncommutative ring then $A = R[x]$ is schematic, and so, Theorem 5.23 holds for A . Notice that this result cannot be obtained from the theory developed by Lezama [27, 29] because it does not satisfy Lezama’s assumption (C4) that says that $A_0 = R$ is a commutative ring. In the particular case of the \mathbb{k} -algebra $R = M_n(\mathbb{k})$, since R is not connected it does not satisfy the definition of schematicness given by Van Oystaeyen and Willaert (Definition 2.6), and it is not a *finitely semi-graded algebra* in the sense of Lezama [29, Definition 2.4]. However, from our point of view, the algebra is schematic and Theorem 5.23 holds.

Example 5.28. Consider A as the 3-dimensional skew polynomial algebra subject to the relations

$$yz = zy, \quad xz = zx, \quad yx = xy - z.$$

Following the ideas presented by Lezama [27, 29], it can be seen that this algebra satisfies the Serre–Artin–Zhang–Verevkin theorem [14, Example 18.5.15 (v)].

It is straightforward to see that the following relations hold:

$$y^n x = xy^n - ny^{n-1}z \quad \text{and} \quad yx^n = x^n y - nx^{n-1}z \quad \text{for } n > 0.$$

If $\text{char}(\mathbb{k}) = p > 0$ then $x^p, y^p, z \in Z(A)$, and so, Proposition 5.26 implies that A is schematic.

Consider the case $\text{char}(\mathbb{k}) = 0$. Let us see that $A''_n = \{az^n \mid a \in \mathbb{k}, n \in \mathbb{N}\}$. With this aim, consider $\alpha \in A''_n$. Then, α is a homogeneous element of degree n , and we can write

$$\alpha = \sum_{i+j \leq n} a_{i,j} x^i y^j z^{n-i-j}.$$

Since that

$$\begin{aligned}
 \alpha x &= \sum_{i+j \leq n} a_{i,j} x^i y^j x z^{n-i-j} \\
 &= \sum_{i+j \leq n} a_{i,j} x^i (x y^j - j y^{j-1} z) z^{n-i-j} \\
 &= \sum_{i+j \leq n} a_{i,j} x^{i+1} y^j z^{n-i-j} - \sum_{i+j \leq n} j a_{i,j} x^i y^{j-1} z^{n-i-j+1},
 \end{aligned}$$

the element αx is homogeneous of degree $n + 1$, and so, $j a_{i,j} = 0$ for each i, j . In a similar way, for the element $y \alpha$ we obtain that $i a_{i,j} = 0$ for each i, j . These facts imply that the only non-zero coefficient is precisely $a_{0,0}$, that is, $\alpha = a_{0,0} z^n$. This shows that

$$A''_n = \{a z^n \mid a \in \mathbb{k}, n \in \mathbb{N}\}.$$

Now, let us prove that A is not schematic. Since $z \in Z(A)$ then $S = \{a z^k \mid a \in \mathbb{k}^*, k \in \mathbb{N}\}$ is a good left Ore set. Note that for all $m \in \mathbb{N}$ we have that $x^m \in A_{\geq m} \setminus Rz$, whence S does not satisfy the schematicness condition. Besides, due to the reasoning above it is clear that S contains any other good left Ore of A , and so, if S does not satisfy the schematicness condition, then no other set will.

Finally, Proposition 5.29 presents necessary conditions to assert the schematicness of skew PBW extensions with two indeterminates.

Proposition 5.29. *Let $A = \sigma(\mathbb{k})\langle x, y \rangle$ be a skew PBW extension over \mathbb{k} defined by the relation*

$$yx = dxy + ex + fy + g, \quad \text{where } d \in \mathbb{k}^* \text{ and } e, f, g \in \mathbb{k}. \quad (5.8)$$

A is schematic if and only if one of the following cases holds:

- (1) $yx = dxy$ (quantum plane, Manin's plane),
- (2) $yx = xy + g$ with $\text{char}(\mathbb{k}) = p > 0$,
- (3) $yx = dxy + g$ with $d \neq 1$ and $d^p = 1$ for some $p \in \mathbb{N}$.

Proof. We divide the proof into four parts.

- (a) Let $P := dx + f$, $Q := ex + g$, $\bar{P} := dy + e$ and $\bar{Q} := fy + g$. Notice that the binomial theorem holds for P and \bar{P} , that is,

$$P^i = \sum_{k=0}^i \binom{i}{k} d^{i-k} f^k x^{i-k} \quad \bar{P}^i = \sum_{k=0}^i \binom{i}{k} d^{i-k} e^k y^{i-k} \quad \text{for all } i > 0,$$

and

$$yx = Py + Q = x\bar{P} + \bar{Q}.$$

Let us see some relations of commutativity between x and y .

For $n \geq 1$, the following identities

$$\begin{aligned} yx^n &= P^n y + \sum_{i=0}^{n-1} P^{n-1-i} x^i Q, \\ y^n x &= x \bar{P}^n + \sum_{i=0}^{n-1} \bar{P}^{n-1-i} y^i \bar{Q} \end{aligned}$$

hold.

The case $n = 1$ is clear. Suppose that the assertion holds for n . Then,

$$\begin{aligned} yx^{n+1} &= \left(P^n y + \sum_{i=0}^{n-1} P^{n-1-i} x^i Q \right) x \\ &= P^n (Py + Q) + \sum_{i=0}^{n-1} P^{n-1-i} x^{i+1} Q \\ &= P^{n+1} y + \left(P^n + \sum_{i=0}^{n-1} P^{n-1-i} x^{i+1} \right) Q \\ &= P^{n+1} y + \sum_{i=0}^n P^{n-i} x^i Q, \end{aligned}$$

which concludes the proof. In a similar way, we can prove the other equality.

- (b) For $n > 0$, we write $\Delta_n := \sum_{i=0}^{n-1} d^i$.

Let us see that if $\xi = ax^n$ (resp., $\xi = ay^n$) belongs to R_n'' with $a \neq 0$, then $f = 0$ (resp., $e = 0$). In the case $Q \neq 0$ (resp., $\bar{Q} \neq 0$), it follows that $\Delta_n = 0$.

The equalities

$$\begin{aligned} y\xi &= ayx^n \\ &= a \left(P^n y + \sum_{i=0}^{n-1} P^{n-1-i} x^i Q \right) \\ &= a \left(\sum_{k=0}^n \binom{n}{k} d^{n-k} f^k x^{n-k} y + \sum_{i=0}^{n-1} P^{n-1-i} x^i Q \right) \end{aligned}$$

show that the element $y\xi$ is homogeneous of degree $n + 1$, and that the monomials having the indeterminate y satisfy that if $k \neq 0$ then $a \binom{n}{k} d^{n-k} f^k = 0$. In particular, if $k = n$ then $af^n = 0$ whence $f = 0$.

Now, with respect to the other monomials, it is clear that these form a polynomial element of degree less than $n + 1$, which shows that

$$a \sum_{i=0}^{n-1} P^{n-1-i} x^i Q = 0.$$

Since $f = 0$, we get that $P = dx$, and so,

$$0 = a \sum_{i=0}^{n-1} P^{n-1-i} x^i Q = a \left(\sum_{i=0}^{n-1} d^{n-1-i} \right) x^{n-1} Q.$$

Thus, if $Q \neq 0$, then

$$0 = \sum_{i=0}^{n-1} d^{n-1-i} = \sum_{i=0}^{n-1} d^i = \Delta_n.$$

The proof for the case ay^n is analogous.

(c) Let $n \geq 1$ and consider the expression

$$\xi = \sum_{i=0}^n a_i x^i y^{n-i} \in R_n''.$$

Let us show that if $\bar{Q} \neq 0$ (resp., $Q \neq 0$) and $\xi \neq a_n x^n$ (resp., $\xi \neq a_0 y^n$), then $e = 0$ (resp., $f = 0$) and $\Delta_k = 0$ for some $0 \leq k \leq n$.

Note that ξx is a homogeneous element of degree $n + 1$. We have the following equalities:

$$\begin{aligned} \xi x &= a_n x^{n+1} + \sum_{i=0}^{n-1} a_i x^i y^{n-i} x \\ &= a_n x^{n+1} + \sum_{i=0}^{n-1} a_i x^i \left(x \bar{P}^{n-i} + \sum_{j=0}^{n-1-i} \bar{P}^{n-1-i-j} y^j \bar{Q} \right) \\ &= a_n x^{n+1} + \sum_{i=0}^{n-1} \left(a_i x^{i+1} \bar{P}^{n-i} + a_i x^i \sum_{j=0}^{n-1-i} \bar{P}^{n-1-i-j} y^j \bar{Q} \right). \end{aligned} \quad (5.9)$$

Suppose that there exists $0 \leq i \leq n - 1$ such that $a_i \neq 0$ and let $t := \min\{0 \leq i \leq n \mid a_i \neq 0\}$. Then,

$$\xi x = a_n x^{n+1} + \sum_{i=t}^{n-1} \left(a_i x^{i+1} \bar{P}^{n-i} + a_i x^i \sum_{j=0}^{n-1-i} \bar{P}^{n-1-i-j} y^j \bar{Q} \right).$$

Notice that the lower exponent of x appears when $i = t$ and we have that

$$a_t x^t \sum_{j=0}^{n-1-t} \bar{P}^{n-1-t-j} y^j \bar{Q}$$

is a polynomial element of degree less than $n + 1$ that has no other terms of ξx , whence necessarily this polynomial has to be the zero element. Since $a_t \neq 0$, it follows that

$$\left(\sum_{j=0}^{n-1-t} \bar{P}^{n-1-t-j} y^j \right) \bar{Q} = 0.$$

By using that $\bar{Q} \neq 0$, we have that $\sum_{j=0}^{n-1-t} \bar{P}^{n-1-t-j} y^j = 0$. Hence,

$$\begin{aligned}
 0 &= \sum_{j=0}^{n-1-t} \bar{P}^{n-1-t-j} y^j \\
 &= \sum_{j=0}^{n-1-t} \left(\sum_{k=0}^{n-1-t-j} \binom{n-1-t-j}{k} d^{n-1-t-j-k} e^k y^{n-1-t-j-k} \right) y^j \\
 &= \sum_{j=0}^{n-1-t} \sum_{k=0}^{n-1-t-j} \binom{n-1-t-j}{k} d^{n-1-t-j-k} e^k y^{n-1-t-k}. \tag{5.10}
 \end{aligned}$$

The coefficient of the monomial y^0 is obtained when $j = 0$ and $k = n - 1 - t$, which implies that

$$\binom{n-1-t}{n-1-t} d^{n-1-t-0-(n-1-t)} e^{n-t-1} = e^{n-t-1} = 0,$$

whence $n - 1 \neq t$ and $e = 0$. By replacing in the expression (5.10), it follows that

$$0 = \sum_{j=0}^{n-1-t} d^{n-1-t-j} y^{n-1-t},$$

and so,

$$0 = \sum_{j=0}^{n-1-t} d^{n-1-t-j} = \sum_{j=0}^{n-1-t} d^j = \Delta_{n-t}.$$

The condition that there exists $n > 0$ such that $\Delta_n = 0$ is recursive, so we will call it *Condition U*. It can be seen that this condition is satisfied if and only if one of the following conditions hold:

- $d = 1$ and $\text{char}(\mathbb{k}) = p > 0$,
- $d \neq 1$ and there exists $p > 0$ such that $d^p = 1$.

- (d) With the analysis above, we can determine the schematicness of the skew PBW extensions defined by relation (5.8).

First of all, note that if $d = 0$, Part (c) implies that $A'' = \mathbb{k}$ since *Condition U* does not hold. Thus, the skew PBW extension A is not schematic. From now on, consider $d \neq 0$. It is clear that the case $e = f = g = 0$ shows that A is schematic. Let us see what happens if one of these three elements is non-zero and *Condition U* does not hold.

Let $\xi \in A''_n$ with $n > 1$. If $g \neq 0$, then $Q \neq 0 \neq \bar{Q}$, and by Part (c) we have that $\xi = a_n x^n = a_0 y^n$, whence $\xi = 0$, and so, $A'' = \mathbb{k}$, which shows that A is not schematic. If $e \neq 0$ then $Q \neq 0$, and Part (c) implies that $\xi = a_0 y^n$, that is $\xi = 0$ (Part (b) above). In this case A is not schematic. Similarly, if $f \neq 0$ then A is not schematic.

Let us see the case where *Condition U* holds (with the less value of p satisfying this condition), and two of the three elements e, f, g being non-zero. If $e \neq 0 \neq f$, then Part (c) implies that $\xi = 0$, whence A is not schematic. If $e \neq 0 \neq g$ and $f = 0$, it follows that $x^p \in Z(R)$. On the other hand, $\bar{Q} \neq 0 \neq \bar{Q}$, and so, Part (c) shows that $\xi = a_n x^n$ with $p \mid n$. In this way, $A'' = \{ax^{pk} \mid a \in \mathbb{k}, k \in \mathbb{N}\}$ and hence, $S = \{ax^{pk} \mid a \in \mathbb{k}^*, k \in \mathbb{N}\}$ is the greatest left Ore set, and since S does not satisfy the condition of schematicness (due to the powers of y), it is clear that A is not schematic. Analogously, one can check that if $f \neq 0 \neq g$ and $e = 0$ then A is not schematic.

Now, let us see the situation where *Condition U* is satisfied and only one element is non-zero. If $e \neq 0$ and $f = 0 = g$, then $\bar{Q} = 0$, and so, equation 5.9 can be written as

$$\xi x = a_n x^{n+1} + \sum_{i=0}^{n-1} \sum_{k=0}^{n-i} \binom{n-i}{k} a_i d^{n-i-k} e^k x^{i+1} y^{n-i-k}.$$

Note that every value of i corresponds to only one power of x , and when $k \neq 0$ the degree of $x^{i+1} y^{n-i-k}$ is less than $n + 1$. These facts show $\binom{n-i}{k} a_i d^{n-i-k} e^k = 0$ for each $0 \leq i \leq n - 1$ and all $0 < k \leq n - i$. In particular, if $k = 1$ then $a_i = 0$, and so, $\xi = a_n x^n$. Part (b) above implies that $p \mid n$ whence A is not schematic by the same reason as above in the case $e \neq 0 \neq g$ and $f = 0$. Analogously, if $f \neq 0$ and $e = 0 = g$ it follows that A is not schematic.

Finally, if *Condition U* holds and $g \neq 0$ with $e = 0 = f$ then it is straightforward to see that $x^p, y^p \in Z(R)$, whence A is schematic. ■

Remark 5.30. Proposition 5.29 shows that there are Ore extensions over schematic rings that are not schematic. This is consistent with Proposition 2.8.

6. Conclusions and future work

In this paper, we have defined the notion of schematic ring in the context of semi-graded objects and illustrated our Theorem 5.23 with some non- \mathbb{N} -graded algebras. With the aim of obtaining new examples of schematic algebras in this more general setting, it is of interest to generalize the criterion formulated by Van Oystayen and Willaert [52, Lemma 2] that says that if R is an \mathbb{N} -graded \mathbb{k} -algebra such that its center $Z(R)$ is Noetherian and such that R is a finitely generated $Z(R)$ -module, then R is schematic. The importance of this criterion can be appreciated in [13] where the authors investigated the schematicness of *skew Ore polynomials of higher order generated by homogeneous quadratic relations* defined by Golovashkin and Maksimov [17, 18]. Since these algebras are non- \mathbb{N} -graded, the research on its schematicness will be crucial for another families of noncommutative rings.

Now, having in mind that Willaert [58] studied the least possible number of Ore sets satisfying the condition of schematicness for \mathbb{N} -graded algebras and called it the *schematic dimension*, a natural task is to investigate this notion in the setting of semi-graded rings. Also, an important topic of future research for these rings is the Čech cohomology developed by Van Oystayen and Willaert [50, 52].

Funding. The second author was supported by the research fund of Faculty of Science, Code HERMES 53880, Universidad Nacional de Colombia - Sede Bogotá, Colombia.

References

- [1] J. Alev and F. Dumas, [Sur le corps des fractions de certaines algèbres quantiques](#). *J. Algebra* **170** (1994), no. 1, 229–265 Zbl [0820.17015](#) MR [1302839](#)
- [2] V. A. Artamonov, [Derivations of skew PBW-extensions](#). *Commun. Math. Stat.* **3** (2015), no. 4, 449–457 Zbl [1338.16030](#) MR [3432214](#)
- [3] M. Artin, [Geometry of quantum planes](#). In *Azumaya algebras, actions, and modules (Bloomington, IN, 1990)*, pp. 1–15, Contemp. Math. 124, American Mathematical Society, Providence, RI, 1992 Zbl [0749.14002](#) MR [1144023](#)
- [4] M. Artin, J. Tate, and M. Van den Bergh, [Some algebras associated to automorphisms of elliptic curves](#). In *The Grothendieck Festschrift, Vol. I*, pp. 33–85, Progr. Math. 86, Birkhäuser, Boston, MA, 1990 Zbl [0744.14024](#) MR [1086882](#)
- [5] M. Artin and J. J. Zhang, [Noncommutative projective schemes](#). *Adv. Math.* **109** (1994), no. 2, 228–287 Zbl [0833.14002](#) MR [1304753](#)
- [6] V. V. Bavula, Generalized Weyl algebras and their representations. *St. Petersburg Math. J.* **4** (1992), no. 1, 71–92 Zbl [0807.16027](#)
- [7] A. D. Bell and K. R. Goodearl, Uniform rank over differential operator rings and Poincaré–Birkhoff–Witt extensions. *Pacific J. Math.* **131** (1988), no. 1, 13–37 Zbl [0598.16002](#) MR [0917863](#)
- [8] A. D. Bell and S. P. Smith, PBW bases for some 3-dimensional skew polynomial algebras. 2018, arXiv:[1805.03489v1](#)
- [9] G. Benkart, [Down-up algebras and Witten’s deformations of the universal enveloping algebra of \$\mathfrak{sl}_2\$](#) . In *Recent progress in algebra (Taejon/Seoul, 1997)*, pp. 29–45, Contemp. Math. 224, American Mathematical Society, Providence, RI, 1999 Zbl [0922.17007](#) MR [1653061](#)
- [10] G. Benkart and T. Roby, [Down-up algebras](#). *J. Algebra* **209** (1998), no. 1, 305–344 Zbl [0922.17006](#) MR [1652138](#)
- [11] F. Calderón and A. Reyes, Some interactions between Hopf Galois extensions and noncommutative rings. *Univ. Sci.* **27** (2022), no. 2, 58–161
- [12] T. Cassidy and M. Vancliff, [Corrigendum: Generalizations of graded Clifford algebras and of complete intersections](#). *J. Lond. Math. Soc. (2)* **90** (2014), no. 2, 631–636 Zbl [1303.16029](#) MR [3263968](#)
- [13] A. Chacón and A. Reyes, On the schematicness of some ore polynomials of higher order generated by homogeneous quadratic relations. *J. Algebra Appl.* (2024), DOI [10.1142/S021949882550207X](#)

- [14] W. Fajardo, C. Gallego, O. Lezama, A. Reyes, H. Suárez, and H. Venegas, *Skew PBW extensions—ring and module-theoretic properties, matrix and Gröbner methods, and applications*. Algebr. Appl. 28, Springer, Cham, 2020 Zbl 1489.16002 MR 4241412
- [15] C. Gallego and O. Lezama, *Gröbner bases for ideals of σ -PBW extensions*. Comm. Algebra **39** (2011), no. 1, 50–75 Zbl 1259.16053 MR 2770878
- [16] O. Goldman, *Rings and modules of quotients*. J. Algebra **13** (1969), 10–47 Zbl 0201.04002 MR 0245608
- [17] A. V. Golovashkin and V. M. Maksimov, *Skew Ore polynomials of higher orders generated by homogeneous quadratic relations*. Russian Math. Surveys **53** (1998), no. 2, 384–386 Zbl 0949.16024
- [18] A. V. Golovashkin and V. M. Maksimov, *Algebras of skew polynomials generated by quadratic homogeneous relations*. J. Math. Sci. (N. Y.) **129** (2005), no. 2, 3757–3771 Zbl 1144.16311
- [19] A. Grothendieck, *Éléments de géométrie algébrique. II. Étude globale élémentaire de quelques classes de morphismes*. Inst. Hautes Études Sci. Publ. Math. (1961), no. 8, 1–222 MR 217084
- [20] R. Hartshorne, *Algebraic geometry*. Grad. Texts in Math. 52, Springer, New York-Heidelberg, 1977 Zbl 0367.14001 MR 0463157
- [21] E. Hashemi, K. Khalilnezhad, and A. Alhevaz, *(Σ, Δ) -compatible skew PBW extension ring*. Kyungpook Math. J. **57** (2017), no. 3, 401–417 Zbl 1380.16023 MR 3722698
- [22] A. P. Isaev, P. N. Pyatov, and V. Rittenberg, *Diffusion algebras*. J. Phys. A **34** (2001), no. 29, 5815–5834 Zbl 1053.16019 MR 1854981
- [23] D. A. Jordan, *Down-up algebras and ambiskew polynomial rings*. J. Algebra **228** (2000), no. 1, 311–346 Zbl 0958.16030 MR 1760967
- [24] D. A. Jordan and I. E. Wells, *Invariants for automorphisms of certain iterated skew polynomial rings*. Proc. Edinburgh Math. Soc. (2) **39** (1996), no. 3, 461–472 Zbl 0864.16027 MR 1417689
- [25] E. Kirkman, I. M. Musson, and D. S. Passman, *Noetherian down-up algebras*. Proc. Amer. Math. Soc. **127** (1999), no. 11, 3161–3167 Zbl 0940.16012 MR 1610796
- [26] J. O. Lezama, *Computation of point modules of finitely semi-graded rings*. Comm. Algebra **48** (2020), no. 2, 866–878 Zbl 1436.16030 MR 4068919
- [27] O. Lezama, *Some open problems in the context of skew PBW extensions and semi-graded rings*. Commun. Math. Stat. **9** (2021), no. 3, 347–378 Zbl 1490.16060 MR 4299903
- [28] O. Lezama and J. Gómez, *Koszulity and point modules of finitely semi-graded rings and algebras*. Symmetry **11** (2019), no. 7, 1–22
- [29] O. Lezama and E. Latorre, *Non-commutative algebraic geometry of semi-graded rings*. Internat. J. Algebra Comput. **27** (2017), no. 4, 361–389 Zbl 1382.16024 MR 3668100
- [30] Y. I. Manin, *Topics in noncommutative geometry*. M. B. Porter Lectures, Princeton University Press, Princeton, NJ, 1991 MR 1095783
- [31] O. Ore, *Linear equations in non-commutative fields*. Ann. of Math. (2) **32** (1931), no. 3, 463–477 Zbl 57.0166.01 MR 1503010
- [32] O. Ore, *Theory of non-commutative polynomials*. Ann. of Math. (2) **34** (1933), no. 3, 480–508 Zbl 0007.15101 MR 1503119
- [33] F. V. Oystaeyen and L. Willaert, *Grothendieck topology, coherent sheaves and Serre’s theorem for schematic algebras*. J. Pure Appl. Algebra **104** (1995), no. 1, 109–122 Zbl 0854.16027 MR 1359695
- [34] I. T. Redman, *The homogenization of the three-dimensional skew polynomial algebras of type I*. Comm. Algebra **27** (1999), no. 11, 5587–5602 Zbl 0942.16030 MR 1713054

- [35] A. Reyes, [Armendariz modules over skew PBW extensions](#). *Comm. Algebra* **47** (2019), no. 3, 1248–1270 Zbl [1444.16035](#) MR [3938553](#)
- [36] A. Reyes and C. Rodríguez, [The McCoy condition on skew Poincaré–Birkhoff–Witt extensions](#). *Commun. Math. Stat.* **9** (2021), no. 1, 1–21 Zbl [1479.16035](#) MR [4222493](#)
- [37] A. Reyes and C. Sarmiento, [On the differential smoothness of 3-dimensional skew polynomial algebras and diffusion algebras](#). *Internat. J. Algebra Comput.* **32** (2022), no. 3, 529–559 Zbl [1503.16032](#) MR [4417484](#)
- [38] A. Reyes and H. Suárez, [Skew Poincaré–Birkhoff–Witt extensions over weak compatible rings](#). *J. Algebra Appl.* **19** (2020), no. 12, article no. 2050225 Zbl [1464.16019](#) MR [4160976](#)
- [39] A. Reyes and H. Suárez, [Radicals and Köthe’s conjecture for skew PBW extensions](#). *Commun. Math. Stat.* **9** (2021), no. 2, 119–138 Zbl [1470.16050](#) MR [4258034](#)
- [40] V. Rittenberg and D. Wyler, [Generalized superalgebras](#). *Nuclear Phys. B* **139** (1978), no. 3, 189–202 MR [0491859](#)
- [41] A. L. Rosenberg, [Noncommutative algebraic geometry and representations of quantized algebras](#). Math. Appl. 330, Kluwer, Dordrecht, 1995 MR [1347919](#)
- [42] J.-P. Serre, [Faisceaux algébriques cohérents](#). *Ann. of Math. (2)* **61** (1955), 197–278 Zbl [0067.16201](#) MR [0068874](#)
- [43] S. P. Smith, [A class of algebras similar to the enveloping algebra of \$\mathfrak{sl}\(2\)\$](#) . *Trans. Amer. Math. Soc.* **322** (1990), no. 1, 285–314 Zbl [0732.16019](#) MR [0972706](#)
- [44] B. Stenström, [Rings of quotients. An introduction to methods of ring theory](#). Die Grundlehren der mathematischen Wissenschaften 217, Springer, New York–Heidelberg, 1975 Zbl [0296.16001](#) MR [0389953](#)
- [45] H. Suárez, A. Chacón, and A. Reyes, [On NI and NJ skew PBW extensions](#). *Comm. Algebra* **50** (2022), no. 8, 3261–3275 Zbl [1497.16019](#) MR [4429459](#)
- [46] H. Suárez, A. Reyes, and Y. Suárez, [Homogenized skew PBW extensions](#). *Arab. J. Math. (Springer)* **12** (2023), no. 1, 247–263 Zbl [1514.16024](#) MR [4552851](#)
- [47] A. B. Tumwesigye, J. Richter, and S. Silvestrov, [Centralizers in PBW extensions](#). In *Algebraic structures and applications*, pp. 469–490, Springer Proc. Math. Stat. 317, Springer, Cham, 2020 MR [4113917](#)
- [48] F. Van Oystaeyen, [On graded rings and modules of quotients](#). *Comm. Algebra* **6** (1978), no. 18, 1923–1959 Zbl [0388.16001](#) MR [0511169](#)
- [49] F. Van Oystaeyen, [Algebraic geometry for associative algebras](#). Monogr. Textbooks Pure Appl. Math. 232, Marcel Dekker, New York, 2000 Zbl [0986.16014](#) MR [1807463](#)
- [50] F. Van Oystaeyen and L. Willaert, [Cohomology of schematic algebras](#). *J. Algebra* **185** (1996), no. 1, 74–84 Zbl [0864.16038](#) MR [1409975](#)
- [51] F. Van Oystaeyen and L. Willaert, [The quantum site of a schematic algebra](#). *Comm. Algebra* **24** (1996), no. 1, 209–222 Zbl [0847.16025](#) MR [1370531](#)
- [52] F. Van Oystaeyen and L. Willaert, [Examples and quantum sections of schematic algebras](#). *J. Pure Appl. Algebra* **120** (1997), no. 2, 195–211 Zbl [0892.16020](#) MR [1466620](#)
- [53] M. Vanclicff, [The interplay of algebra and geometry in the setting of regular algebras](#). In *Commutative algebra and noncommutative algebraic geometry. Vol. I*, pp. 371–390, Math. Sci. Res. Inst. Publ. 67, Cambridge University Press, New York, 2015 Zbl [1359.16025](#) MR [3525477](#)
- [54] M. Vanclicff and K. Van Rompay, [Four-dimensional regular algebras with point scheme, a non-singular quadric in \$\mathbf{P}^3\$](#) . *Comm. Algebra* **28** (2000), no. 5, 2211–2242 Zbl [0961.16019](#) MR [1757458](#)

- [55] P. Veerapen, [Graded Clifford algebras and graded skew Clifford algebras and their role in the classification of Artin–Schelter regular algebras](#). *Adv. Appl. Clifford Algebr.* **27** (2017), no. 3, 2855–2871 Zbl [1380.16042](#) MR [3688849](#)
- [56] A. B. Verëvkin, [Injective Serre sheaves](#). *Mat. Zametki* **52** (1992), no. 4, 35–41 Zbl [0812.16007](#) MR [1203950](#)
- [57] A. B. Verëvkin, [On a noncommutative analogue of the category of coherent sheaves on a projective scheme](#). In *Algebra and analysis (Tomsk, 1989)*, pp. 41–53, Amer. Math. Soc. Transl. Ser. 2 151, American Mathematical Society, Providence, RI, 1992 Zbl [0920.14001](#) MR [1191171](#)
- [58] L. Willaert, A new dimension for schematic algebras. In *Rings, Hopf algebras, and Brauer groups (Antwerp/Brussels, 1996)*, pp. 325–332, Lecture Notes in Pure and Appl. Math. 197, Dekker, New York, 1998 Zbl [0920.16019](#) MR [1615908](#)
- [59] J. J. Zhang and J. Zhang, [Double Ore extensions](#). *J. Pure Appl. Algebra* **212** (2008), no. 12, 2668–2690 Zbl [1157.16009](#) MR [2452318](#)
- [60] A. S. Zhedanov, [“Hidden symmetry” of Askey–Wilson polynomials](#). *Theor. Math. Phys.* **89** (1991), no. 2, 1146–1157 Zbl [0782.33012](#)

Received 20 April 2023; revised 30 August 2024.

Andrés Chacón

Escuela de Matemáticas, Universidad Industrial de Santander, Carrera 27 Calle 9, Bucaramanga, Colombia; andchaca@uis.edu.co

Armando Reyes

Departamento de Matemáticas, Universidad Nacional de Colombia, Carrera 45, no. 26-85, Bogotá, D.C., Colombia; mareyesv@unal.edu.co