

On the Cahn–Hilliard equation with kinetic rate dependent dynamic boundary conditions and non-smooth potentials: Well-posedness and asymptotic limits

Maoyin Lv and Hao Wu

Abstract. We analyze a class of Cahn–Hilliard equations with kinetic rate dependent dynamic boundary conditions that describe possible short-range interactions between the binary mixture and the solid boundary. In the presence of surface diffusion on the boundary, the initial boundary value problem can be viewed as a transmission problem consisting of Cahn–Hilliard-type equations both in the bulk and on the boundary. We first establish the existence, uniqueness, and continuous dependence of global weak solutions. In the construction of weak solutions, an explicit convergence rate in terms of the parameter for the Yosida approximation is obtained. Under some additional assumptions, we further prove the existence and uniqueness of global strong solutions. Next, we study the asymptotic limit as the coefficient of the boundary diffusion goes to zero and show that the limit problem with a forward-backward dynamic boundary condition is well posed in a suitable weak formulation. At last, we investigate the asymptotic limits as the kinetic rate tends to zero and infinity, respectively. Our results are valid for a general class of bulk and boundary potentials with double-well structure, including the physically relevant logarithmic potential and the non-smooth double obstacle potential.

Contents

1. Introduction	177
2. Main results	184
3. Well-posedness in the presence of surface diffusion	199
4. Asymptotic limit as $\delta \rightarrow 0$ and well-posedness without surface diffusion	216
5. Asymptotic limits with respect to kinetic rate: $L \rightarrow 0$ or $L \rightarrow +\infty$	222
A. Appendix	230
References	231

1. Introduction

The Cahn–Hilliard equation was proposed in [8] as a phenomenological model to describe spinodal decomposition in binary alloys. It characterizes the fundamental process of phase

Mathematics Subject Classification 2020: 35K35 (primary); 35K61, 35B20, 35B40, 80A22 (secondary).
Keywords: Cahn–Hilliard equation, dynamic boundary condition, bulk-boundary interaction, non-smooth potential, well-posedness, asymptotic limit.

separation due to certain non-Fickian diffusion driven by the gradient of chemical potential. The Cahn–Hilliard equation belongs to the so-called diffuse interface models, in which the free interface between two components of the mixture is represented by a thin layer with finite thickness. The diffuse interface methodology avoids tracking free interfaces explicitly as in the classical free boundary problems, and it provides a thermodynamically consistent description for the evolution of complex geometries. In particular, topological changes in free interfaces can be captured in a natural and efficient way. For further information, we refer to, e.g., [1, 2, 25] and the references therein. As a representative of the diffuse interface models, the Cahn–Hilliard equation has become a useful tool for the study of a wide variety of segregation-like phenomena arising, for instance, in material science, image inpainting, biology, and fluid mechanics. In recent years, the study of boundary effects in the phase separation process of binary mixtures has attracted a lot of attention. To describe short-range interactions of the binary mixture with the solid wall, several types of dynamic boundary conditions for the Cahn–Hilliard equation have been proposed and investigated in the literature; see, for instance, [40, 53, 63] and the references therein.

1.1. Model description

Let $T \in (0, +\infty)$ be an arbitrary but fixed final time and $\Omega \subset \mathbb{R}^d$ ($d \in \{2, 3\}$) a smooth bounded domain with boundary $\Gamma := \partial\Omega$. In this study, we consider the following initial boundary value problem of the Cahn–Hilliard equation subject to a class of dynamic boundary conditions [38, 42, 49]:

$$\partial_t \varphi = \Delta \mu, \quad \text{in } \Omega \times (0, T), \tag{1.1}$$

$$\mu = -\Delta \varphi + F'(\varphi), \quad \text{in } \Omega \times (0, T), \tag{1.2}$$

$$\begin{cases} L \partial_{\mathbf{n}} \mu = \theta - \mu & \text{if } L \in [0, +\infty), \\ \partial_{\mathbf{n}} \mu = 0 & \text{if } L = +\infty, \end{cases} \quad \text{on } \Gamma \times (0, T), \tag{1.3}$$

$$\partial_t \varphi = \Delta_{\Gamma} \theta - \partial_{\mathbf{n}} \mu, \quad \text{on } \Gamma \times (0, T), \tag{1.4}$$

$$\theta = \partial_{\mathbf{n}} \varphi - \delta \Delta_{\Gamma} \varphi + G'(\varphi), \quad \text{on } \Gamma \times (0, T), \tag{1.5}$$

$$\varphi|_{t=0} = \varphi_0, \quad \text{in } \bar{\Omega}. \tag{1.6}$$

Here, the phase function $\varphi : \bar{\Omega} \times (0, T) \rightarrow \mathbb{R}$ is related to local concentrations of the two components of a binary mixture. The total free energy functional associated with the system (1.1)–(1.5) is given by

$$E(\varphi) := \underbrace{\int_{\Omega} \left(\frac{1}{2} |\nabla \varphi|^2 + F(\varphi) \right) dx}_{\text{bulk free energy}} + \underbrace{\int_{\Gamma} \left(\frac{\delta}{2} |\nabla_{\Gamma} \varphi|^2 + G(\varphi) \right) dS}_{\text{surface free energy}}. \tag{1.7}$$

In particular, the surface free energy on the boundary is introduced to describe possible short-range interactions between the solid wall and components of the mixture [27]. The

nonlinear potential functions F and G denote free energy densities in the bulk and on the boundary, respectively. To describe the phase separation process, they usually present a double-well structure, that is, with two minima and a local unstable maximum in between. Typical and physically significant examples of such potentials include the logarithmic potential [8] and the double obstacle potential [5]:

$$\begin{aligned}
 F_{\log}(r) &:= (1+r)\ln(1+r) + (1-r)\ln(1-r) - c_1 r^2, \quad r \in (-1, 1), \quad (1.8) \\
 F_{2\text{obs}}(r) &:= \begin{cases} c_2(1-r^2) & \text{if } |r| \leq 1, \\ +\infty & \text{if } |r| > 1, \end{cases} \quad (1.9)
 \end{aligned}$$

where the constants satisfy $c_1 > 1$ and $c_2 > 0$ so that $F_{\log}, F_{2\text{obs}}$ are nonconvex. In practice, regular double-well potential of polynomial type such as

$$F_{\text{reg}}(r) := \frac{1}{4}(r^2 - 1)^2, \quad r \in \mathbb{R}, \quad (1.10)$$

and its generalizations are widely used (see [53]). In (1.2), $\mu : \Omega \times (0, T) \rightarrow \mathbb{R}$ stands for the chemical potential in the bulk, while in (1.5), $\theta : \Gamma \times (0, T) \rightarrow \mathbb{R}$ stands for the chemical potential on the boundary. They can be expressed as Fréchet derivatives of the bulk and surface free energies in (1.7), respectively. The symbol Δ denotes the Laplace operator in Ω , Δ_Γ denotes the Laplace–Beltrami operator on Γ , ∇ denotes the gradient operator in the bulk, and ∇_Γ denotes the tangential (surface) gradient operator. In the boundary conditions (1.3)–(1.5), the symbol \mathbf{n} denotes the outward normal vector on Γ and $\partial_{\mathbf{n}}$ means the outward normal derivative on Γ . The nonlinearities F' and G' in (1.2) and (1.5) simply denote derivatives of the related potentials. Nevertheless, when non-smooth potentials are taken into account, F' and G' correspond to the subdifferential of the convex part (may be multivalued graphs) plus the derivative of the smooth concave contribution. For example, we have $F'_{2\text{obs}}(r) = \partial I_{[-1,1]}(r) - 2c_2 r$, where $\partial I_{[-1,1]}(r)$ is the subdifferential of the indicator function of $[-1, 1]$. In this case, one should replace the equality in (1.2) and (1.5) by inclusion. The boundary conditions (1.3)–(1.5) allow descriptions of the physically realistic scenario with possible mass transfer between bulk and boundary as well as a dynamic angle between the free interface (separating components of the binary mixture) and the solid boundary at the contact line. In this aspect, we refer to [37], where a general thermodynamically consistent Navier–Stokes–Cahn–Hilliard system with dynamic boundary conditions for incompressible two-phase flows with non-matched densities was introduced and analyzed (see [30] for the existence of global weak solutions in a more general setting with singular potentials). Besides, we refer to [23] for a related Cahn–Hilliard–Brinkman model on incompressible creeping two-phase flows through a porous medium.

In our problem (1.1)–(1.6), we maintain two parameters $L \in [0, +\infty]$ and $\delta \in [0, +\infty)$ that are important in the subsequent analysis. Other coefficients are set to be one for the sake of simplicity.

We first explain the role of $L \in [0, +\infty]$. The bulk and boundary chemical potentials μ, θ are coupled through the boundary condition (1.3), which accounts for possible adsorption or desorption processes between the materials in the bulk and on the boundary [42].

The mass flux $\partial_{\mathbf{n}}\mu$, which describes the motion of materials towards and away from the boundary, is driven by the difference in the chemical potentials. In this sense, the coefficient $1/L$ can be interpreted as a kinetic rate. The value of L distinguishes different types of bulk-boundary interactions. The case $L = 0$ was introduced by Goldstein, Miranville, and Schimperna [38] (GMS in short) for the phase separation of a binary mixture confined to a bounded domain with porous walls. The GMS model extends the Cahn–Hilliard equation with Wentzell-type boundary conditions proposed by Gal [29] (see [31, 62] for related mathematical analysis). Taking $L = 0$ in (1.3), we obtain the Dirichlet boundary condition $\mu = \theta$ on $\Gamma \times (0, T)$, which implies that the chemical potentials μ and θ are always in the chemical equilibrium (see [38] for more general situations with a factor that can be a uniformly bounded positive function). The case $L = +\infty$ was introduced by Liu and Wu [49] (LW in short) based on an energetic variational approach that combines the least action principle and Onsager’s principle of maximum dissipation. Then, the homogeneous Neumann boundary condition for μ implies that there is no mass transfer between the bulk and boundary. In such a situation, the chemical potentials μ and θ are not directly coupled. Nevertheless, interactions between the bulk and the surface materials take place through the phase function φ . Indeed, let us introduce a new unknown

$$\psi = \varphi|_{\Gamma}, \quad \text{on } \Gamma \times (0, T), \quad (1.11)$$

where $\varphi|_{\Gamma}$ denotes the trace of φ (cf., [53, 55]). Then, we find that the system (1.1)–(1.5) yields a sort of transmission problem between the dynamics in the bulk Ω and the one on the boundary Γ . Finally, the case $L \in (0, +\infty)$ was recently introduced by Knopf, Lam, Liu, and Metzger [42] (KLLM in short). The corresponding Robin-type boundary condition (1.3) describes the situation that the chemical potentials μ and θ are not in equilibrium and they are related through the mass flux. Formally speaking, the KLLM model (with $0 < L < +\infty$) can be regarded as an interpolation between the GMS model ($L = 0$, instantaneous mass transfer) and the LW model ($L = +\infty$, no mass transfer) via a finite, positive relaxation parameter L (see [42] for a rigorous verification).

Next, let us comment on the parameter $\delta \in [0, +\infty)$, which acts as a weight for surface diffusion effects on the boundary Γ . When $\delta > 0$, (1.4) together with (1.5) leads to a Cahn–Hilliard-type dynamic boundary condition. The case $\delta = 0$ is closely related to the evolution of a free interface in contact with the solid boundary, that is, the moving contact line problem [11, 57]. From the mathematical point of view, without surface diffusion, the boundary conditions (1.4)–(1.5) (formally) reduce to

$$\partial_t \varphi - G''(\varphi) \Delta_{\Gamma} \varphi = G^{(3)}(\varphi) |\nabla_{\Gamma} \varphi|^2 + \Delta_{\Gamma} (\partial_{\mathbf{n}} \varphi) - \partial_{\mathbf{n}} \mu, \quad \text{on } \Gamma \times (0, T). \quad (1.12)$$

In the regime that the potential G is non-convex, in particular, $G''(\varphi) \leq 0$, we obtain a backward heat equation on Γ , whose well-posedness is usually a delicate issue. As pointed out in [18, 19], (1.12) yields a forward-backward dynamic boundary condition, complemented with a Cahn–Hilliard equation (1.1)–(1.2) in the bulk.

The values of L and δ also lead to differences in some basic properties for the thermodynamically consistent problem (1.1)–(1.6), such as mass conservation and energy

dissipation (see [38, 42, 49, 63]). For a sufficiently regular solution, we have the energy dissipation law

$$\frac{d}{dt} E(\varphi(t)) + \int_{\Omega} |\nabla \mu(t)|^2 dx + \int_{\Gamma} |\nabla_{\Gamma} \theta(t)|^2 dS + \chi(L) \int_{\Gamma} |\mu(t) - \theta(t)|^2 dS = 0$$

for all $t \in (0, T)$, where $\chi(L) = 1/L$ if $L \in (0, +\infty)$ and $\chi(L) = 0$ if $L = 0$ or $+\infty$. This implies that, for all $L \in [0, +\infty]$, the total free energy $E(\varphi)$ is decreasing as time evolves. Nevertheless, the form of the free energy E depends on δ (recall (1.7)) and the energy dissipation varies according to L . On the other hand, we obtain the conservation of total mass for $L \in [0, +\infty)$,

$$\int_{\Omega} \varphi(t) dx + \int_{\Gamma} \varphi(t) dS = \int_{\Omega} \varphi_0 dx + \int_{\Gamma} \varphi_0 dS \quad \forall t \in [0, T],$$

while for $L = +\infty$, mass conservation laws hold in the bulk and on the boundary separately:

$$\int_{\Omega} \varphi(t) dx = \int_{\Omega} \varphi_0 dx, \quad \int_{\Gamma} \varphi(t) dS = \int_{\Gamma} \varphi_0 dS \quad \forall t \in [0, T].$$

1.2. A brief overview of related literature

The Cahn–Hilliard equation with different types of dynamic boundary conditions has been extensively studied from various viewpoints [53, 63]. For the case with a dynamic boundary condition of Allen–Cahn type, that is, $\partial_t \varphi - \delta \Delta_{\Gamma} \varphi + G'(\varphi) + \partial_{\mathbf{n}} \varphi = 0$ on $\Gamma \times (0, T)$, we quote [11, 13, 17, 21, 34, 35, 56, 58, 64] among the vast literature. We note that the dynamic boundary condition of Allen–Cahn type corresponds to an L^2 -relaxation of the surface energy (see [57]), while in our problem (1.1)–(1.6), the dynamic boundary condition of Cahn–Hilliard type accounts for the mass transport on Γ with a $(H^1)'$ -relaxation dynamics.

For mathematical analysis of the GMS model ($L = 0$), we refer to [12, 14, 16, 18, 22, 28, 38] and the references therein. In [38], existence, uniqueness, regularity, and long-time behavior of global weak solutions were established under general assumptions on the nonlinearities, with G being regular. On the other hand, well-posedness of the GMS model with non-smooth bulk and boundary potentials was proved in [16], where the boundary potential was assumed to dominate the bulk one. See also [28] for the strict separation property and long-time behavior, and [22] when convection effects were taken into account. In [18], the asymptotic analysis as $\delta \rightarrow 0$, i.e., the surface diffusion term on the dynamic boundary condition tends to 0, was carried out in a very general setting with nonlinear terms admitting maximal monotone graphs both in the bulk and on the boundary.

Concerning the LW model ($L = +\infty$), well-posedness and long-time behavior were first established in [49] when F and G are suitable regular potentials. By introducing a slightly weaker notion of the solution, the authors of [32] proved existence of global

weak solutions via a gradient flow approach, removing the additional geometric assumption imposed in [49] for the case $\delta = 0$. Well-posedness for the LW model with general non-smooth potentials has been obtained in [20]. In [19], the asymptotic analysis as $\delta \rightarrow 0$ was investigated with non-smooth potentials in the bulk and on the boundary. A recent work [51] established the strict separation property and showed that every global weak solution will converge to a single equilibrium. We also mention [54] for the existence of global attractors and [52] for the numerical analysis.

For the KLLM model ($0 < L < +\infty$), weak and strong well-posedness results were established in [42] with suitable regular potentials F and G . Under a similar setting, long-time behaviors, such as existence of a global attractor/exponential attractors and convergence to a single equilibrium, were obtained in [33]. Asymptotic limits as $L \rightarrow 0$ and $L \rightarrow +\infty$ were rigorously verified in [42] with regular potentials; see also [33] for further properties in the limit $L \rightarrow 0$. A nonlocal variant of the KLLM model (including a non-local dynamic boundary condition) was proposed and investigated in [44]. Besides, for simulations and numerical analysis, we refer to [3, 42]. In the recent contribution [46], a class of more general bulk-surface convective Cahn–Hilliard systems with dynamic boundary conditions and regular potentials F, G were investigated. Therein, the trace relation (1.11) for the phase function is further relaxed as follows (cf., (1.3) for the chemical potentials):

$$\begin{cases} J \partial_n \varphi = \psi - \varphi|_\Gamma & \text{if } J \in [0, +\infty), \\ \partial_n \varphi = 0 & \text{if } J = +\infty, \end{cases} \quad \text{on } \Gamma \times (0, T).$$

The Robin approximation with $J \in (0, +\infty)$ describes a scenario, where the boundary phase variable and the trace of the bulk phase variable are not proportional (see [41] for a similar consideration for the LW model). Existence of weak solutions for $J, L \in (0, +\infty)$ was proved [46] by means of a Faedo–Galerkin approach. For other cases, existence results were obtained by studying the asymptotic limits as sending J, L to 0 and $+\infty$, respectively. It is worth mentioning that all the related results mentioned above for the KLLM model were achieved for regular potentials F, G , singular potentials, such as the logarithmic potential (1.8) and the double-obstacle potential (1.9), are unfortunately not admissible. The only known result on the existence of weak solutions to the KLLM model with singular potentials including (1.8) can be obtained as a consequence of [30] on a generalized Navier–Stokes–Cahn–Hilliard system with dynamic boundary conditions. However, uniqueness and regularity properties were not available therein due to the presence of fluid interaction.

1.3. Goal of the paper

In this paper, we aim to study well-posedness and asymptotic limits as $\delta \rightarrow 0, L \rightarrow 0$, or $L \rightarrow +\infty$ of the initial boundary value problem (1.1)–(1.6) with a wide class of bulk/boundary potentials F, G that have a double-well structure, in particular, including the non-smooth logarithmic potential (1.8) and the double-obstacle potential (1.9).

(1) Well-posedness of the KLLM model ($0 < L < +\infty$) with surface diffusion ($\delta > 0$). We prove the existence of global weak solutions (see Theorem 2.7) and their continuous dependence on the data that yields the uniqueness (see Theorem 2.9). Thanks to the solvability of a second-order elliptic problem with bulk-surface coupling [43], we are allowed to apply the approach in [16] for the GMS model to conclude the existence result. To this end, we consider a regularized problem by adding viscous terms in the Cahn–Hilliard equation as well as the dynamic boundary condition and substituting the maximal monotone graphs with their Yosida regularizations. The resulting regularized problem can be solved by the abstract theory of doubly nonlinear evolution inclusions (see Proposition 3.1). After that, we derive suitable estimates for the approximating solutions, which are uniform with respect to the parameter ε for the Yosida regularization. Passing to the limit as $\varepsilon \rightarrow 0$, we can construct a global weak solution by the compactness argument. The continuous dependence estimate can be proved by the energy method. Moreover, we establish an $O(\varepsilon^{1/2})$ -estimate for the convergence of approximating phase functions in $L^\infty(0, T; (\mathcal{H}^1)') \cap L^2(0, T; \mathcal{V}^1)$ (see Proposition 3.7), which seems to be the first result of this kind for problem (1.1)–(1.6). Finally, after deriving some higher-order (in time) uniform estimates for the approximating solutions, we obtain the existence of a unique global strong solution (see Theorem 2.11). Our contribution extends previous works on well-posedness of the KLLM model with regular potentials [42, 46].

(2) Asymptotic limit as $\delta \rightarrow 0$ for the KLLM model with fixed $L \in (0, +\infty)$. We show that weak solutions to problem (1.1)–(1.6) obtained in Theorems 2.7, 2.9 converge as $\delta \rightarrow 0$ (in the sense of a subsequence) and thus prove the existence of weak solutions to the limit problem in which the boundary condition (1.5) is replaced by the one with $\delta = 0$ (see Theorem 2.15). Besides, we establish a continuous dependence result for the phase function with respect to the data (see Theorem 2.17). This implies that the coupling of a forward-backward-type boundary condition with the Cahn–Hilliard equation in the bulk can be well posed in a suitable sense. Analogous conclusions have been obtained for the GMS model and the LW model with general bulk/boundary potentials, respectively, in [18] and [19]. Hence, our results fill the gap left by [18, 19]. Like in [18, 19], the solution of the limit problem with vanishing surface diffusion loses some spatial regularity, so several terms on the boundary including $\partial_n \varphi$ should be understood in a weaker sense.

(3) Asymptotic limit as $L \rightarrow 0$ or $L \rightarrow +\infty$ for the KLLM model with fixed $\delta \in (0, +\infty)$. In presence of the surface diffusion in the KLLM model, we rigorously justify the limit case of instantaneous reaction as $L \rightarrow 0$ (see Theorem 2.20), where the chemical potentials are in equilibrium, and a vanishing reaction rate as $L \rightarrow +\infty$ (see Theorem 2.23), where the chemical potentials are not directly coupled. Comparing with the previous results in [42] that are only valid for suitable regular potentials, the main novelty in our analysis is the treatment of non-smooth bulk and boundary potentials that include the logarithmic potential (1.8) and the double obstacle potential (1.9). In order to derive uniform estimates with respect to the parameter L , different approaches have to be applied in the regime of vanishing (or large) kinetic rate. We remark that the asymptotic

limit with respect to L turns out to be more involved in the case of vanishing surface diffusion and general potentials. This issue will be studied in a future work.

Plan of the paper. The remaining part of this paper is organized as follows. In Section 2, we introduce our notation and assumptions and then state the main results. In Section 3, we consider the KLLM model with surface diffusion, proving the existence of global weak solutions, the continuous dependence, and the existence of global strong solutions. Besides, we obtain a convergence rate for the approximating phase functions. In Section 4, we investigate the asymptotic limit as $\delta \rightarrow 0$, which yields the existence of global weak solutions to the limit problem with vanishing surface diffusion. A continuous dependence estimate is also derived. In Section 5, we study the asymptotic limits with respect to the kinetic rate as $L \rightarrow 0$, $L \rightarrow +\infty$, in presence of the surface diffusion. In the Appendix, we list some useful tools that are frequently used in this paper.

2. Main results

In this section, we first recall some notation for the functional settings; then, we describe our problem and state the main results.

2.1. Preliminaries

For any real Banach space X , we denote its norm by $\|\cdot\|_X$, its dual space by X' , and the duality pairing between X' and X by $\langle \cdot, \cdot \rangle_{X', X}$. If X is a Hilbert space, its inner product will be denoted by $(\cdot, \cdot)_X$. The space $L^q(0, T; X)$ ($1 \leq q \leq +\infty$) denotes the set of all strongly measurable q -integrable functions with values in X , or, if $q = +\infty$, essentially bounded functions. The space $C([0, T]; X)$ denotes the Banach space of all bounded and continuous functions $u : [0, T] \rightarrow X$ equipped with the supremum norm, while $C_w([0, T]; X)$ denotes the topological vector space of all bounded and weakly continuous functions.

Let Ω be a bounded domain in \mathbb{R}^d ($d \in \{2, 3\}$) with sufficiently smooth boundary $\Gamma := \partial\Omega$ (at least Lipschitz). The associated outward unit normal vector field on Γ is denoted by \mathbf{n} . We use $|\Omega|$ and $|\Gamma|$ to denote the Lebesgue measure of Ω and the Hausdorff measure of Γ , respectively. For any $1 \leq q \leq +\infty$, $k \in \mathbb{N}$, the standard Lebesgue and Sobolev spaces on Ω are denoted by $L^q(\Omega)$ and $W^{k,q}(\Omega)$. Here, we use \mathbb{N} for the set of natural numbers including zero. For $s \geq 0$ and $q \in [1, +\infty)$, we denote by $H^{s,q}(\Omega)$ the Bessel-potential spaces and by $W^{s,q}(\Omega)$ the Slobodeckij spaces. If $q = 2$, it holds $H^{s,2}(\Omega) = W^{s,2}(\Omega)$ for all s and these spaces are Hilbert spaces. We will use the notation $H^s(\Omega) = H^{s,2}(\Omega) = W^{s,2}(\Omega)$ and $H^0(\Omega)$ can be identified with $L^2(\Omega)$. The Lebesgue spaces, Sobolev spaces, and Slobodeckij spaces on the boundary Γ can be defined analogously, provided that Γ is sufficiently regular. We write

$$H^s(\Gamma) = H^{s,2}(\Gamma) = W^{s,2}(\Gamma)$$

and identify $H^0(\Gamma)$ with $L^2(\Gamma)$. Hereafter, the following shortcuts will be applied:

$$H := L^2(\Omega), \quad H_\Gamma := L^2(\Gamma), \quad V := H^1(\Omega), \quad V_\Gamma := H^1(\Gamma).$$

For every $y \in (H^1(\Omega))'$, we denote by $\langle y \rangle_\Omega = |\Omega|^{-1} \langle y, 1 \rangle_{(H^1(\Omega))', H^1(\Omega)}$ its generalized mean value over Ω . If $y \in L^1(\Omega)$, then its spatial mean is given by

$$\langle y \rangle_\Omega = |\Omega|^{-1} \int_\Omega y \, dx.$$

The spatial mean for a function y_Γ on Γ , denoted by $\langle y_\Gamma \rangle_\Gamma$, can be defined in a similar manner. Then, we introduce the spaces for functions with zero mean:

$$\begin{aligned} V_0 &:= \{y \in V : \langle y \rangle_\Omega = 0\}, & V_0^* &:= \{y^* \in V' : \langle y^* \rangle_\Omega = 0\}, \\ V_{\Gamma,0} &:= \{y_\Gamma \in V_\Gamma : \langle y_\Gamma \rangle_\Gamma = 0\}, & V_{\Gamma,0}^* &:= \{y_\Gamma^* \in V_\Gamma' : \langle y_\Gamma^* \rangle_\Gamma = 0\}. \end{aligned}$$

The following Poincaré–Wirtinger inequalities in Ω and on Γ hold (see, e.g., [26, Theorem 2.12] for the case on Γ):

$$\|u - \langle u \rangle_\Omega\|_H \leq C_\Omega \|\nabla u\|_H \quad \forall u \in V, \tag{2.1}$$

$$\|u_\Gamma - \langle u_\Gamma \rangle_\Gamma\|_{H_\Gamma} \leq C_\Gamma \|\nabla_\Gamma u_\Gamma\|_{H_\Gamma} \quad \forall u_\Gamma \in V_\Gamma, \tag{2.2}$$

where C_Ω (resp., C_Γ) is a positive constant depending only on Ω (resp., Γ). For basic facts of calculus on surfaces, see, e.g., [26, Section 2].

Consider the Neumann problem

$$\begin{cases} -\Delta u = y, & \text{in } \Omega, \\ \partial_{\mathbf{n}} u = 0, & \text{on } \Gamma. \end{cases} \tag{2.3}$$

Owing to the Lax–Milgram theorem, for every $y \in V_0^*$, problem (2.3) admits a unique weak solution $u \in V_0$ satisfying

$$\int_\Omega \nabla u \cdot \nabla \zeta \, dx = \langle y, \zeta \rangle_{V', V} \quad \forall \zeta \in V.$$

Thus, we can define the solution operator $\mathcal{N}_\Omega : V_0^* \rightarrow V_0$ such that $u = \mathcal{N}_\Omega y$. Analogously, let us consider the surface Poisson equation

$$-\Delta_\Gamma u_\Gamma = y_\Gamma, \quad \text{on } \Gamma, \tag{2.4}$$

where $\Delta_\Gamma = \operatorname{div}_\Gamma \nabla_\Gamma$ denotes the Laplace–Beltrami operator on Γ . For every $y_\Gamma \in V_{\Gamma,0}^*$, problem (2.4) admits a unique weak solution $u_\Gamma \in V_{\Gamma,0}$ satisfying

$$\int_\Gamma \nabla_\Gamma u_\Gamma \cdot \nabla_\Gamma \zeta_\Gamma \, dS = \langle y_\Gamma, \zeta_\Gamma \rangle_{V_\Gamma', V_\Gamma} \quad \forall \zeta_\Gamma \in V_\Gamma.$$

Then, we can define the solution operator $\mathcal{N}_\Gamma : V_{\Gamma,0}^* \rightarrow V_{\Gamma,0}$ such that $u_\Gamma = \mathcal{N}_\Gamma y_\Gamma$. By virtue of these definitions, we can introduce the following equivalent norms:

$$\begin{aligned} \|y\|_{V_0^*} &:= \left(\int_\Omega |\nabla \mathcal{N}_\Omega y|^2 dx \right)^{1/2} & \forall y \in V_0^*, \\ \|y\|_{V'} &:= \left(\|y - \langle y \rangle_\Omega\|_{V_0^*}^2 + |\langle y \rangle_\Omega|^2 \right)^{1/2} & \forall y \in V', \\ \|y_\Gamma\|_{V_{\Gamma,0}^*} &:= \left(\int_\Gamma |\nabla_\Gamma \mathcal{N}_\Gamma y_\Gamma|^2 dS \right)^{1/2} & \forall y_\Gamma \in V_{\Gamma,0}^*, \\ \|y_\Gamma\|_{V'_\Gamma} &:= \left(\|y_\Gamma - \langle y_\Gamma \rangle_\Gamma\|_{V_{\Gamma,0}^*}^2 + |\langle y_\Gamma \rangle_\Gamma|^2 \right)^{1/2} & \forall y_\Gamma \in V'_\Gamma. \end{aligned}$$

Next, we introduce the product spaces

$$\mathcal{L}^q := L^q(\Omega) \times L^q(\Gamma) \quad \text{and} \quad \mathcal{W}^{k,q} := W^{k,q}(\Omega) \times W^{k,q}(\Gamma)$$

for $q \in [1, +\infty]$ and $k \in \mathbb{N}$. In particular, we use the notation

$$\mathcal{H}^k := H^k(\Omega) \times H^k(\Gamma) = \mathcal{W}^{k,2}.$$

Like before, we can identify \mathcal{H}^0 with \mathcal{L}^2 . For any $k \in \mathbb{N}$, \mathcal{H}^k is a Hilbert space endowed with the standard inner product

$$((y, y_\Gamma), (z, z_\Gamma))_{\mathcal{H}^k} := (y, z)_{H^k(\Omega)} + (y_\Gamma, z_\Gamma)_{H^k(\Gamma)} \quad \forall (y, y_\Gamma), (z, z_\Gamma) \in \mathcal{H}^k$$

and the induced norm $\|\cdot\|_{\mathcal{H}^k} := (\cdot, \cdot)_{\mathcal{H}^k}^{1/2}$. We introduce the duality pairing

$$\langle (y, y_\Gamma), (\zeta, \zeta_\Gamma) \rangle_{(\mathcal{H}^1)', \mathcal{H}^1} = (y, \zeta)_{L^2(\Omega)} + (y_\Gamma, \zeta_\Gamma)_{L^2(\Gamma)} \quad \forall (y, y_\Gamma) \in \mathcal{L}^2, (\zeta, \zeta_\Gamma) \in \mathcal{H}^1.$$

By the Riesz representation theorem, this product can be extended to a duality pairing on $(\mathcal{H}^1)' \times \mathcal{H}^1$.

For any $k \in \mathbb{Z}^+$, we introduce the Hilbert space

$$\mathcal{V}^k := \{(y, y_\Gamma) \in \mathcal{H}^k : y|_\Gamma = y_\Gamma \text{ a.e. on } \Gamma\},$$

endowed with the inner product $(\cdot, \cdot)_{\mathcal{V}^k} := (\cdot, \cdot)_{\mathcal{H}^k}$ and the associated norm $\|\cdot\|_{\mathcal{V}^k} := \|\cdot\|_{\mathcal{H}^k}$. Here, $y|_\Gamma$ stands for the trace of $y \in H^k(\Omega)$ on the boundary Γ , which makes sense for $k \in \mathbb{Z}^+$. The duality pairing on $(\mathcal{V}^1)' \times \mathcal{V}^1$ can be defined in a similar manner. For convenience, we also use the notation

$$\tilde{\mathcal{V}}^k := \{(y, y_\Gamma) \in H^k(\Omega) \times H^{k-1/2}(\Gamma) : y|_\Gamma = y_\Gamma \text{ a.e. on } \Gamma\}, \quad k \in \mathbb{Z}^+.$$

Thanks to the trace theorem, for every $y \in H^k(\Omega)$, $k \in \mathbb{Z}^+$, it holds $(y, y|_\Gamma) \in \tilde{\mathcal{V}}^k$.

For any given $m \in \mathbb{R}$, we set

$$\mathcal{L}_{(m)}^2 := \{(y, y_\Gamma) \in \mathcal{L}^2 : |\Omega| \langle y \rangle_\Omega + |\Gamma| \langle y_\Gamma \rangle_\Gamma = m\}.$$

The closed linear subspaces

$$\mathcal{H}_{(0)}^k = \mathcal{H}^k \cap \mathcal{L}_{(0)}^2, \quad \mathcal{V}_{(0)}^k = \mathcal{V}^k \cap \mathcal{L}_{(0)}^2, \quad k \in \mathbb{Z}^+,$$

are Hilbert spaces endowed with the inner products $(\cdot, \cdot)_{\mathcal{H}^k}$ and the associated norms $\|\cdot\|_{\mathcal{H}^k}$, respectively. For $L \in [0, +\infty)$ and $k \in \mathbb{Z}^+$, we introduce the notation

$$\mathcal{H}_L^k := \begin{cases} \mathcal{H}^k & \text{if } L \in (0, +\infty), \\ \mathcal{V}^k & \text{if } L = 0, \end{cases} \quad \mathcal{H}_{L,0}^k := \begin{cases} \mathcal{H}_{(0)}^k & \text{if } L \in (0, +\infty), \\ \mathcal{V}_{(0)}^k & \text{if } L = 0. \end{cases}$$

Consider the bilinear form

$$\begin{aligned} & a_L((y, y_\Gamma), (z, z_\Gamma)) \\ & := \int_{\Omega} \nabla y \cdot \nabla z \, dx + \int_{\Gamma} \nabla_{\Gamma} y_{\Gamma} \cdot \nabla_{\Gamma} z_{\Gamma} \, dS + \chi(L) \int_{\Gamma} (y - y_{\Gamma})(z - z_{\Gamma}) \, dS \end{aligned}$$

for all $(y, y_{\Gamma}), (z, z_{\Gamma}) \in \mathcal{H}^1$, where

$$\chi(L) = \begin{cases} 1/L & \text{if } L \in (0, +\infty), \\ 0 & \text{if } L = 0. \end{cases}$$

For any $(y, y_{\Gamma}) \in \mathcal{H}_{L,0}^1$, $L \in [0, +\infty)$, we define

$$\|(y, y_{\Gamma})\|_{\mathcal{H}_{L,0}^1} := ((y, y_{\Gamma}), (y, y_{\Gamma}))_{\mathcal{H}_{L,0}^1}^{1/2} = [a_L((y, y_{\Gamma}), (y, y_{\Gamma}))]^{1/2}. \quad (2.5)$$

We note that, for $(y, y_{\Gamma}) \in \mathcal{V}_{(0)}^1 \subseteq \mathcal{H}_{L,0}^1$, $\|(y, y_{\Gamma})\|_{\mathcal{H}_{L,0}^1}$ does not depend on L , since the third term in a_L simply vanishes. The following Poincaré type inequality has been proved in [43, Lemma A.1].

Lemma 2.1. *There exists a constant $c_P > 0$ depending only on $L \in [0, +\infty)$ and Ω such that*

$$\|(y, y_{\Gamma})\|_{\mathcal{L}^2} \leq c_P \|(y, y_{\Gamma})\|_{\mathcal{H}_{L,0}^1} \quad \forall (y, y_{\Gamma}) \in \mathcal{H}_{L,0}^1. \quad (2.6)$$

Hence, for every $L \in [0, +\infty)$, $\mathcal{H}_{L,0}^1$ is a Hilbert space with the inner product $(\cdot, \cdot)_{\mathcal{H}_{L,0}^1}$. The induced norm $\|\cdot\|_{\mathcal{H}_{L,0}^1}$ prescribed in (2.5) is equivalent to the standard one $\|\cdot\|_{\mathcal{H}^1}$ on $\mathcal{H}_{L,0}^1$. Besides, for any fixed $\delta > 0$, we define the bilinear form

$$b_{\delta}((y, y_{\Gamma}), (z, z_{\Gamma})) := \int_{\Omega} \nabla y \cdot \nabla z \, dx + \delta \int_{\Gamma} \nabla_{\Gamma} y_{\Gamma} \cdot \nabla_{\Gamma} z_{\Gamma} \, dS \quad \forall (y, y_{\Gamma}), (z, z_{\Gamma}) \in \mathcal{H}^1.$$

It is easy to check that

$$\|(y, y_{\Gamma})\|_{\mathcal{V}_{(0)}^1} := [b_{\delta}((y, y_{\Gamma}), (y, y_{\Gamma}))]^{1/2} \quad \forall (y, y_{\Gamma}) \in \mathcal{V}_{(0)}^1 \quad (2.7)$$

yields a norm equivalent to $\|\cdot\|_{\mathcal{H}^1}$ on $\mathcal{V}_{(0)}^1$.

For $L \in [0, +\infty)$, let us consider the following elliptic boundary value problem:

$$\begin{cases} -\Delta u = y & \text{in } \Omega, \\ -\Delta_\Gamma u_\Gamma + \partial_{\mathbf{n}} u = y_\Gamma & \text{on } \Gamma, \\ L \partial_{\mathbf{n}} u = u_\Gamma - u|_\Gamma & \text{on } \Gamma. \end{cases} \tag{2.8}$$

Define the space

$$\mathcal{H}_{(0)}^{-1} := \{(y, y_\Gamma) \in (\mathcal{H}^1)'\} : |\Omega|\langle y \rangle_\Omega + |\Gamma|\langle y_\Gamma \rangle_\Gamma = 0\}.$$

The chain of inclusions holds

$$\mathcal{H}_{L,0}^1 \subseteq \mathcal{H}_{(0)}^1 \subset \mathcal{L}_{(0)}^2 \subset \mathcal{H}_{(0)}^{-1} \subset (\mathcal{H}^1)' \subseteq (\mathcal{H}_L^1)' \subset (\mathcal{H}_{L,0}^1)'.$$

It has been shown in [43, Theorem 3.3] that, for every $(y, y_\Gamma) \in \mathcal{H}_{(0)}^{-1}$, problem (2.8) admits a unique weak solution $(u, u_\Gamma) \in \mathcal{H}_{L,0}^1$ satisfying the weak formulation

$$a_L((u, u_\Gamma), (\zeta, \zeta_\Gamma)) = \langle (y, y_\Gamma), (\zeta, \zeta_\Gamma) \rangle_{(\mathcal{H}_L^1)', \mathcal{H}_L^1} \quad \forall (\zeta, \zeta_\Gamma) \in \mathcal{H}_L^1$$

and the estimate

$$\|(u, u_\Gamma)\|_{\mathcal{H}^1} \leq C \|(y, y_\Gamma)\|_{(\mathcal{H}_L^1)'} \tag{2.9}$$

for some constant $C > 0$ depending only on L and Ω . Furthermore, if the domain Ω is of class C^{k+2} and $(y, y_\Gamma) \in \mathcal{H}_{L,0}^k$, $k \in \mathbb{N}$, the following regularity estimate holds:

$$\|(u, u_\Gamma)\|_{\mathcal{H}^{k+2}} \leq C \|(y, y_\Gamma)\|_{\mathcal{H}^k}. \tag{2.10}$$

The above facts enable us to define the solution operator

$$\mathcal{S}^L : \mathcal{H}_{(0)}^{-1} \rightarrow \mathcal{H}_{L,0}^1, \quad (y, y_\Gamma) \mapsto (u, u_\Gamma) = \mathcal{S}^L(y, y_\Gamma) = (\mathcal{S}_\Omega^L(y, y_\Gamma), \mathcal{S}_\Gamma^L(y, y_\Gamma)).$$

We mention that similar results for the special case $L = 0$ have also been presented in [16]. A direct calculation yields that

$$((u, u_\Gamma), (z, z_\Gamma))_{\mathcal{H}^2} = ((u, u_\Gamma), \mathcal{S}^L(z, z_\Gamma))_{\mathcal{H}_{L,0}^1} \quad \forall (u, u_\Gamma) \in \mathcal{H}_{(0)}^1, (z, z_\Gamma) \in \mathcal{L}_{(0)}^2.$$

Thanks to [43, Corollary 3.5], we can introduce the inner product on $\mathcal{H}_{(0)}^{-1}$ as

$$((y, y_\Gamma), (z, z_\Gamma))_{0,*} := (\mathcal{S}^L(y, y_\Gamma), \mathcal{S}^L(z, z_\Gamma))_{\mathcal{H}_{L,0}^1} \quad \forall (y, y_\Gamma), (z, z_\Gamma) \in \mathcal{H}_{(0)}^{-1}.$$

The associated norm $\|(y, y_\Gamma)\|_{0,*} := ((y, y_\Gamma), (y, y_\Gamma))_{0,*}^{1/2}$ is equivalent to the standard dual norm $\|\cdot\|_{(\mathcal{H}_L^1)'}$ on $\mathcal{H}_{(0)}^{-1}$. For any $(y, y_\Gamma) \in (\mathcal{H}^1)'$, we define the generalized mean

$$\bar{m}(y, y_\Gamma) := \frac{|\Omega|\langle y \rangle_\Omega + |\Gamma|\langle y_\Gamma \rangle_\Gamma}{|\Omega| + |\Gamma|}. \tag{2.11}$$

Then, it follows that

$$\|(y, y_\Gamma)\|_* := (\|(y, y_\Gamma) - \bar{m}(y, y_\Gamma)\mathbf{1}\|_{0,*}^2 + |\bar{m}(y, y_\Gamma)|^2)^{1/2} \quad \forall (y, y_\Gamma) \in (\mathcal{H}^1)'$$

is equivalent to the usual dual norm $\|\cdot\|_{(\mathcal{H}_L^1)'}$ on $(\mathcal{H}^1)'$.

Define the closed linear subspaces $\tilde{\mathcal{V}}_{(0)}^k := \tilde{\mathcal{V}}^k \cap \mathcal{L}_{(0)}^2$. The following Poincaré-type inequality has been proved in [18, Lemma A.1].

Lemma 2.2. *There exists a constant $\tilde{c}_P > 0$ depending only on Ω such that*

$$\|y\|_H \leq \tilde{c}_P \|\nabla y\|_H \quad \forall (y, y_\Gamma) \in \tilde{\mathcal{V}}_{(0)}^1. \tag{2.12}$$

Moreover, we have the following interpolation inequality (see [42, Lemma 2.3]).

Lemma 2.3. *For any $\gamma > 0$, there exists a constant C_γ depending only on $L \in [0, +\infty)$, γ , and Ω such that*

$$\|y\|_H + \|y_\Gamma\|_{H_\Gamma} \leq \gamma \|\nabla y\|_H + C_\gamma \|(y, y_\Gamma)\|_{0,*} \quad \forall (y, y_\Gamma) \in \tilde{\mathcal{V}}_{(0)}^1. \tag{2.13}$$

Throughout the paper, the symbol C stands for generic positive constants that may depend on Ω , the final time T , and the coefficients and the norms of functions involved in the assumptions of either our statements or our approximation. Specific dependence will be pointed out if necessary. Besides, we use different symbols to denote precise constants that we would refer to.

2.2. Problem setting

In what follows, we assume that $\Omega \subset \mathbb{R}^d$ ($d \in \{2, 3\}$) is a bounded domain of class C^2 , $T \in (0, +\infty)$ is an arbitrary but fixed final time, and we set

$$Q := \Omega \times (0, T), \quad \Sigma := \Gamma \times (0, T).$$

For $L \in [0, +\infty]$, $\delta \in [0, +\infty)$, using (1.11), we can rewrite our target problem (1.1)–(1.6) as follows:

$$\partial_t \varphi = \Delta \mu \quad \text{in } Q, \tag{2.14}$$

$$\mu = -\Delta \varphi + \xi + \pi(\varphi) - f, \quad \xi \in \beta(\varphi) \quad \text{in } Q, \tag{2.15}$$

$$\partial_t \psi = \Delta_\Gamma \theta - \partial_{\mathbf{n}} \mu \quad \text{on } \Sigma, \tag{2.16}$$

$$\theta = \partial_{\mathbf{n}} \varphi - \delta \Delta_\Gamma \psi + \xi_\Gamma + \pi_\Gamma(\psi) - f_\Gamma, \quad \xi_\Gamma \in \beta_\Gamma(\psi) \quad \text{on } \Sigma, \tag{2.17}$$

$$\varphi = \psi \quad \text{on } \Sigma, \tag{2.18}$$

$$\begin{cases} L \partial_{\mathbf{n}} \mu = \theta - \mu & \text{if } L \in [0, +\infty), \\ \partial_{\mathbf{n}} \mu = 0 & \text{if } L = +\infty, \end{cases} \quad \text{on } \Sigma, \tag{2.19}$$

$$\varphi|_{t=0} = \varphi_0 \quad \text{in } \Omega, \tag{2.20}$$

$$\psi|_{t=0} = \psi_0 \quad \text{on } \Gamma, \tag{2.21}$$

where $f : Q \rightarrow \mathbb{R}$, $f_\Gamma : \Sigma \rightarrow \mathbb{R}$, $\varphi_0 : \Omega \rightarrow \mathbb{R}$, $\psi_0 : \Gamma \rightarrow \mathbb{R}$ are given data. For convenience, we denote (2.14)–(2.21) by problem $(S_{L,\delta})$.

Hereafter, we consider the nonlinear bulk and boundary potentials F and G that can be decomposed as

$$F = \hat{\beta} + \hat{\pi}, \quad G = \hat{\beta}_\Gamma + \hat{\pi}_\Gamma.$$

Besides, we make the following assumptions:

- (A1) $\hat{\beta}, \hat{\beta}_\Gamma : \mathbb{R} \rightarrow [0, +\infty]$ are lower semicontinuous and convex functions with $\hat{\beta}(0) = \hat{\beta}_\Gamma(0) = 0$. Their subdifferentials

$$\beta := \partial \hat{\beta}, \quad \beta_\Gamma := \partial \hat{\beta}_\Gamma$$

are maximal monotone graphs in $\mathbb{R} \times \mathbb{R}$, with effective domains $D(\beta)$ and $D(\beta_\Gamma)$, respectively. Since 0 is a minimum point of both $\hat{\beta}, \hat{\beta}_\Gamma$, it follows that $0 \in \beta(0)$ and $0 \in \beta_\Gamma(0)$.

- (A2) $D(\beta_\Gamma) \subseteq D(\beta)$ and there exist positive constants $\varrho, c_0 > 0$ such that

$$|\beta^\circ(r)| \leq \varrho |\beta_\Gamma^\circ(r)| + c_0 \quad \forall r \in D(\beta_\Gamma), \tag{2.22}$$

where β° denotes the minimal section of the graph β , i.e., $\beta^\circ(r) := \{r^* \in \beta(r) : |r^*| = \inf_{s \in \beta(r)} |s|\}$ and the same definition applies to β_Γ° .

- (A3) $\hat{\pi}, \hat{\pi}_\Gamma \in C^1(\mathbb{R})$ and their derivatives $\pi := \hat{\pi}'$, $\pi_\Gamma := \hat{\pi}'_\Gamma$ are globally Lipschitz continuous with Lipschitz constants denoted by K and K_Γ , respectively.

- (A4) $(f, f_\Gamma) \in L^2(0, T; \mathcal{V}^1)$.

- (A5) $(\varphi_0, \psi_0) \in \mathcal{V}^1$ satisfying $\hat{\beta}(\varphi_0) \in L^1(\Omega)$, $\hat{\beta}_\Gamma(\psi_0) \in L^1(\Gamma)$ as well as

$$\langle \varphi_0 \rangle_\Omega, \langle \psi_0 \rangle_\Gamma \in \text{Int } D(\beta_\Gamma).$$

In order to obtain some further regularity properties of the weak solution, that is, (2.29), we need the following additional assumption on the singular potentials F and G .

- (A1)' The nonlinear convex functions satisfy $\hat{\beta}, \hat{\beta}_\Gamma \in C([-1, 1]) \cap C^2(-1, 1)$. Their derivatives $\beta = \hat{\beta}'$, $\beta_\Gamma = \hat{\beta}'_\Gamma$ satisfy $\beta, \beta_\Gamma \in C^1(-1, 1)$ and are monotone increasing functions such that

$$\begin{aligned} \lim_{r \rightarrow -1} \beta(r) &= -\infty, & \lim_{r \rightarrow 1} \beta(r) &= +\infty, \\ \lim_{r \rightarrow -1} \beta_\Gamma(r) &= -\infty, & \lim_{r \rightarrow 1} \beta_\Gamma(r) &= +\infty. \end{aligned}$$

We also extend $\hat{\beta}(r) = \hat{\beta}_\Gamma(r) = +\infty$ for any $r \notin [-1, 1]$. Without loss of generality, we assume $\hat{\beta}(0) = \hat{\beta}_\Gamma(0) = \beta(0) = \beta_\Gamma(0) = 0$. This also entails that $\hat{\beta}(r), \hat{\beta}_\Gamma(r) \geq 0$ for all $r \in [-1, 1]$.

Remark 2.4. All the typical examples of potentials given in (1.8)–(1.10) fulfill the assumptions (A1)–(A3), provided that the boundary potential dominates the one in the bulk

as demanded in (2.22). The condition (2.22) follows the setting in [9, 15, 16, 20, 21, 49, 59]. An alternative choice is that the bulk potential dominates the boundary one (see, e.g., [34, 35]). These compatibility conditions are useful when we deal with the bulk-surface interaction and derive necessary *a priori* estimates. Here, we take the classical choice of a dominating boundary potential in order to avoid further technicalities. From (A2) and (A5), we find $\bar{m}(\varphi_0, \psi_0) \in \text{Int } D(\beta_\Gamma)$. Besides, we note that, for non-smooth potentials (1.8) and (1.9), only the logarithmic potential (1.8) satisfies (A1)', but the double obstacle potential (1.9) does not.

For convenience, below we will use the bold notation

$$\boldsymbol{\varphi} = (\varphi, \psi), \quad \boldsymbol{\mu} = (\mu, \theta), \quad \boldsymbol{\xi} = (\xi, \xi_\Gamma), \quad \boldsymbol{\pi} = (\pi, \pi_\Gamma), \quad \boldsymbol{f} = (f, f_\Gamma), \quad \boldsymbol{\varphi}_0 = (\varphi_0, \psi_0),$$

and also for generic elements $\mathbf{y} = (y, y_\Gamma)$ in the product spaces $\mathcal{L}^2, \mathcal{H}^1, (\mathcal{H}^1)'$, etc. Owing to the solvability of the elliptic problem (2.8) for $L \in (0, +\infty)$, inspired by [15, 16, 47], we can formally write our problem $(S_{L,\delta})$ into a suitable abstract formulation. To this end, we define the projection operator

$$\mathbf{P} : \mathcal{L}^2 \rightarrow \mathcal{L}^2_{(0)}, \quad (y, y_\Gamma) \mapsto (y - \bar{m}(y, y_\Gamma), y_\Gamma - \bar{m}(y, y_\Gamma)),$$

where \bar{m} is the generalized mean given by (2.11). Then, for any functions $\mathbf{z} \in \mathcal{L}^2$ and $\mathbf{y} \in \mathcal{L}^2_{(0)}$, it holds

$$(\mathbf{P}\mathbf{z}, \mathbf{y})_{\mathcal{L}^2} = \int_\Omega (z - \bar{m}(z))y \, dx + \int_\Gamma (z_\Gamma - \bar{m}(z))y_\Gamma \, dS = (\mathbf{z}, \mathbf{y})_{\mathcal{L}^2}. \quad (2.23)$$

Concerning the equations (2.14), (2.16), (2.19), using (2.8), we can (formally) write

$$\mathbf{P}\boldsymbol{\mu} = \mathfrak{G}^L(-\partial_t \boldsymbol{\varphi}) \quad \text{in } \mathcal{H}^1_{(0)}.$$

On the other hand, consider the lower semicontinuous and convex functional $\Phi_\delta : \mathcal{L}^2_{(0)} \rightarrow [0, +\infty]$ given by

$$\Phi_\delta(\mathbf{z}) := \begin{cases} \frac{1}{2} \int_\Omega |\nabla z|^2 \, dx + \frac{\delta}{2} \int_\Gamma |\nabla_\Gamma z_\Gamma|^2 \, dS & \text{if } \mathbf{z} \in \mathcal{V}^1_{(0)}, \\ +\infty & \text{otherwise.} \end{cases}$$

For any $\mathbf{z} \in \mathcal{V}^1_{(0)}$, we have $2\Phi_\delta(\mathbf{z}) = b_\delta(\mathbf{z}, \mathbf{z})$. Thanks to [16, Lemma C], the subdifferential $\partial\Phi_\delta$ on $\mathcal{L}^2_{(0)}$ fulfills

$$\partial\Phi_\delta(\mathbf{z}) = (-\Delta z, \partial_n z - \delta \Delta_\Gamma z_\Gamma) \quad \forall \mathbf{z} \in D(\partial\Phi_\delta) = \mathcal{H}^2 \cap \mathcal{V}^1_{(0)}. \quad (2.24)$$

Set

$$\boldsymbol{\omega} = (\omega, \omega_\Gamma) := \boldsymbol{\varphi} - \bar{m}_0 \mathbf{1} \quad \text{with } \bar{m}_0 = \bar{m}(\boldsymbol{\varphi}_0) = \frac{|\Omega| \langle \varphi_0 \rangle_\Omega + |\Gamma| \langle \psi_0 \rangle_\Gamma}{|\Omega| + |\Gamma|}.$$

Then, our target problem $(S_{L,\delta})$ with $L, \delta \in (0, +\infty)$ is equivalent to the Cauchy problem for a suitable evolution equation:

$$\begin{cases} \mathfrak{S}^L(\omega'(t)) + \mathbf{P}\mu(t) = \mathbf{0} & \text{in } \mathcal{H}_{(0)}^1 & \text{for a.a. } t \in (0, T), \\ \mu(t) = \partial\Phi_\delta(\omega(t)) + \xi(t) \\ \quad + \pi(\omega(t) + \bar{m}_0\mathbf{1}) - f(t) & \text{in } \mathcal{L}^2 & \text{for a.a. } t \in (0, T), \\ \xi(t) \in \beta(\omega(t) + \bar{m}_0\mathbf{1}) & \text{in } \mathcal{L}^2 & \text{for a.a. } t \in (0, T), \\ \omega|_{t=0} = \omega_0 = (\omega_0, \omega_{\Gamma_0}) & \text{in } \mathcal{L}_{(0)}^2, \end{cases} \quad (2.25)$$

where $\beta(z) := (\beta(z), \beta_\Gamma(z_\Gamma))$ and $\omega_0 := \varphi_0 - \bar{m}_0\mathbf{1}$.

2.3. Statement of results

In this subsection, we summarize the main results of this paper.

2.3.1. Well-posedness for $L, \delta \in (0, +\infty)$. Let us start with the case $L, \delta \in (0, +\infty)$. First, we give the definition of weak solutions.

Definition 2.5. Let $L, \delta \in (0, +\infty)$ be fixed parameters. Suppose that the assumptions (A1)–(A5) are satisfied. The triplet (φ, μ, ξ) is called a weak solution to problem $(S_{L,\delta})$ on $[0, T]$, if the following conditions are fulfilled.

- (1) The functions φ, μ, ξ have the regularity properties

$$\begin{aligned} \omega &= \varphi - \bar{m}_0\mathbf{1} \in H^1(0, T; \mathcal{H}_{(0)}^{-1}) \cap L^\infty(0, T; \mathcal{V}_{(0)}^1) \cap L^2(0, T; \mathcal{V}^2), \\ \mu &\in L^2(0, T; \mathcal{H}^1), \quad \xi \in L^2(0, T; \mathcal{L}^2). \end{aligned}$$

- (2) The variational formulation

$$\langle \partial_t \varphi(t), \mathbf{y} \rangle_{(\mathcal{H}^1)', \mathcal{H}^1} + a_L(\mu(t), \mathbf{y}) = 0 \quad \forall \mathbf{y} \in \mathcal{H}^1 \quad (2.26)$$

holds for almost all $t \in (0, T)$ and we have

$$\mu = -\Delta\varphi + \xi + \pi(\varphi) - f \quad \text{a.e. in } Q, \quad (2.27)$$

$$\theta = \partial_n\varphi - \delta\Delta_\Gamma\psi + \xi_\Gamma + \pi_\Gamma(\psi) - f_\Gamma \quad \text{a.e. on } \Sigma, \quad (2.28)$$

with

$$\xi \in \beta(\varphi) \quad \text{a.e. in } Q, \quad \xi_\Gamma \in \beta_\Gamma(\psi) \quad \text{a.e. on } \Sigma.$$

- (3) The initial conditions are satisfied:

$$\varphi|_{t=0} = \varphi_0 \quad \text{a.e. in } \Omega, \quad \psi|_{t=0} = \psi_0 \quad \text{a.e. on } \Gamma.$$

Remark 2.6. The regularity of φ implies that $\varphi \in C_w([0, T]; \mathcal{V}^1) \cap C([0, T]; \mathcal{L}^2)$. The variational equality (2.26) provides a representation of the time derivative $\partial_t\varphi$ as an element of the dual space $L^2(0, T; (\mathcal{H}^1)')$.

Theorem 2.7 (Existence of weak solution for $L, \delta \in (0, +\infty)$). *Suppose that the assumptions (A1)–(A5) are satisfied. For any given $L, \delta \in (0, +\infty)$, problem $(S_{L,\delta})$ admits a weak solution (φ, μ, ξ) on $[0, T]$ in the sense of Definition 2.5. Moreover, if the assumption (A1)' is fulfilled, we have the following additional regularity,*

$$\varphi \in L^2(0, T; \mathcal{W}^{2,p}), \quad (\beta(\varphi), \beta_\Gamma(\psi)) \in L^2(0, T; \mathcal{L}^p), \tag{2.29}$$

where $p \in [1, 6]$ if $d = 3$ and $p \in [1, +\infty)$ if $d = 2$.

Remark 2.8. When $(f, f_\Gamma) \equiv (0, 0)$, we can conclude the energy equality

$$\begin{aligned} E(\varphi(t), \psi(t)) &+ \int_0^t \int_\Omega |\nabla \mu(s)|^2 \, dx \, ds + \int_0^t \int_\Gamma |\nabla_\Gamma \theta(s)|^2 \, dS \, ds \\ &+ \chi(L) \int_0^t \int_\Gamma |\mu(s) - \theta(s)|^2 \, dS \, ds = E(\varphi_0, \psi_0) \quad \forall t \in [0, T], \end{aligned}$$

where $E(\varphi, \psi)$ is given by (1.7) (recalling that $\psi = \varphi|_\Gamma$). The energy equality presents the dissipative nature of the bulk-surface coupled system (2.14)–(2.21). Besides, the mapping $t \mapsto E(\varphi(t))$ is absolutely continuous for all $t \in [0, T]$. This fact allows us to further conclude $\varphi \in C([0, T]; \mathcal{V}^1)$. For the derivation of the energy equality, we refer to [50, Proposition 3.2] for the case that F, G satisfy (A1)'. For the general case under the assumption (A1), including double obstacle potentials, the proof can be carried out in the same way.

Uniqueness of the phase function φ associated with problem $(S_{L,\delta})$ is guaranteed by the following continuous dependence estimate.

Theorem 2.9 (Continuous dependence for $L, \delta \in (0, +\infty)$). *Suppose that the assumptions (A1)–(A5) are satisfied. Let $(\varphi_i, \mu_i, \xi_i)$, $i \in \{1, 2\}$, be two weak solutions to problem $(S_{L,\delta})$ corresponding to the data $(\varphi_{0,i}, f_i)$ obtained in Theorem 2.7, with*

$$\bar{m}(\varphi_{0,1}) = \bar{m}(\varphi_{0,2}) = \bar{m}_0.$$

Then, there exists a constant $C > 0$, depending only on K, K_Γ, Ω , and T , such that

$$\begin{aligned} \|\varphi_1(t) - \varphi_2(t)\|_{0,*}^2 &+ \int_0^t \|\varphi_1(s) - \varphi_2(s)\|_{\mathcal{V}_1^1}^2 \, ds \\ &\leq C \left(\|\varphi_{0,1} - \varphi_{0,2}\|_{0,*}^2 + \int_0^t \|f_1(s) - f_2(s)\|_{(\mathcal{V}^1)'}^2 \, ds \right) \quad \forall t \in [0, T]. \end{aligned} \tag{2.30}$$

In order to prove the existence of strong solutions, some additional regularity assumptions on φ_0 and f are needed:

(A6) $f := (f, f_\Gamma) \in H^1(0, T; \mathcal{L}^2)$.

(A7) $\varphi_0 \in \mathcal{V}^2$ and the family

$$\{\partial \Phi_\delta(\varphi_0 - \bar{m}_0 \mathbf{1}) + \beta_\varepsilon(\varphi_0) + \pi(\varphi_0) - f(0) : \varepsilon \in (0, \varepsilon_0)\}$$

is bounded in \mathcal{H}^1 for some $\varepsilon_0 \in (0, 1)$. Here, $\beta_\varepsilon = (\beta_\varepsilon, \beta_{\Gamma,\varepsilon})$ denotes the Yosida regularization for maximal monotone operators $\beta = (\beta, \beta_\Gamma)$; see (3.1)–(3.2) in Section 3.

Definition 2.10. Let $L, \delta \in (0, +\infty)$ be fixed parameters. Suppose that the assumptions (A1)–(A7) are satisfied. The triplet (φ, μ, ξ) is called a strong solution to problem $(S_{L,\delta})$ on $[0, T]$, if the following additional regularity properties

$$\begin{aligned} \varphi &\in W^{1,\infty}(0, T; (\mathcal{H}^1)') \cap H^1(0, T; \mathcal{V}^1) \cap L^\infty(0, T; \mathcal{H}^2), \\ \mu &\in L^\infty(0, T; \mathcal{H}^1) \cap L^2(0, T; \mathcal{H}^2), \\ \xi &\in L^\infty(0, T; \mathcal{L}^2) \end{aligned}$$

are satisfied, and it holds

$$\begin{aligned} \partial_t \varphi &= \Delta \mu, && \text{a.e. in } Q, \\ \mu &= -\Delta u + \xi + \pi(\varphi) - f, \quad \xi \in \beta(\varphi), && \text{a.e. in } Q, \\ \varphi &= \psi, \quad L \partial_{\mathbf{n}} \mu = \theta - \mu, && \text{a.e. on } \Sigma, \\ \partial_t \psi &= \Delta_\Gamma \theta - \partial_{\mathbf{n}} \mu, && \text{a.e. on } \Sigma, \\ \theta &= \partial_{\mathbf{n}} \varphi - \delta \Delta_\Gamma \psi + \xi_\Gamma + \pi_\Gamma(\psi) - f_\Gamma, \quad \xi_\Gamma \in \beta_\Gamma(\psi), && \text{a.e. on } \Sigma, \\ \varphi|_{t=0} &= \varphi_0, && \text{a.e. in } \Omega, \\ \psi|_{t=0} &= \psi_0, && \text{a.e. on } \Gamma. \end{aligned}$$

Theorem 2.11 (Existence and uniqueness of strong solution for $L, \delta \in (0, +\infty)$). *Suppose that the assumptions (A1)–(A7) are satisfied. For any given $L, \delta \in (0, +\infty)$, problem $(S_{L,\delta})$ admits a unique strong solution on $[0, T]$ in the sense of Definition 2.10.*

Remark 2.12. With minor modifications in the proof (cf., [16, 18]), we can still obtain the conclusions of Theorems 2.7, 2.9, and 2.11, if (A4) is replaced by the following alternative assumption:

$$(A4)' \quad \mathbf{f} = (f, f_\Gamma) \in W^{1,1}(0, T; \mathcal{L}^2).$$

Note that (A6) implies (A4)'.

2.3.2. Asymptotic limit: $L \in (0, +\infty)$ fixed, $\delta \rightarrow 0$. In the case of $L \in (0, +\infty)$ and $\delta = 0$ (i.e., without surface diffusion), well-posedness of problem $(S_{L,0})$ can be established via the asymptotic limit of problem $(S_{L,\delta})$ as $\delta \rightarrow 0$. To this end, we first introduce the notion of weak solutions to problem $(S_{L,0})$.

Definition 2.13. Let $L \in (0, +\infty)$ be a fixed parameter, $\delta = 0$. Suppose that the assumptions (A1)–(A4) are satisfied. Besides, we assume that

$$(A5)' \quad (\varphi_0, \psi_0) \in \tilde{\mathcal{V}}^1 \text{ satisfying } \hat{\beta}(\varphi_0) \in L^1(\Omega), \hat{\beta}_\Gamma(\psi_0) \in L^1(\Gamma) \text{ as well as}$$

$$\langle \varphi_0 \rangle_\Omega, \langle \psi_0 \rangle_\Gamma \in \text{Int } D(\beta_\Gamma).$$

The triplet (φ, μ, ξ) is called a weak solution to problem $(S_{L,0})$ on $[0, T]$, if the following conditions are fulfilled.

(1) The functions φ, μ, ξ have the regularity properties

$$\begin{aligned} \omega &= \varphi - \bar{m}_0 \mathbf{1} \in H^1(0, T; \mathcal{H}_{(0)}^{-1}) \cap L^\infty(0, T; \tilde{\mathcal{V}}_{(0)}^1) \cap C_w([0, T]; \tilde{\mathcal{V}}^1), \\ \Delta\varphi &\in L^2(0, T; H), \quad \mu \in L^2(0, T; \mathcal{H}^1), \\ \xi &\in L^2(0, T; H \times (H^{1/2}(\Gamma))') \end{aligned}$$

such that $\varphi|_{t=0} = \varphi_0$ in $\tilde{\mathcal{V}}^1$ (by weak continuity).

(2) The variational formulations

$$\langle \partial_t \varphi(t), y \rangle_{(\mathcal{H}^1)', \mathcal{H}^1} + a_L(\mu(t), y) = 0 \quad \forall y \in \mathcal{H}^1 \tag{2.31}$$

and

$$\begin{aligned} (\mu(t), z)_{\mathcal{X}^2} &= \int_{\Omega} \nabla \varphi(t) \cdot \nabla z \, dx + (\xi(t), z)_H + \langle \xi_\Gamma(t), z_\Gamma \rangle_{(H^{1/2}(\Gamma))', H^{1/2}(\Gamma)} \\ &\quad + (\pi(\varphi(t)) - f(t), z)_{\mathcal{X}^2} \quad \forall z \in \tilde{\mathcal{V}}^1 \end{aligned} \tag{2.32}$$

are satisfied almost everywhere in $(0, T)$, and

$$\xi \in \beta(\varphi), \quad \text{a.e. in } Q, \tag{2.33}$$

$$\int_{\Sigma} \hat{\beta}_\Gamma(\psi) \, dS \, dt + \int_0^T \langle \xi_\Gamma, \zeta_\Gamma - \psi \rangle_{(H^{1/2}(\Gamma))', H^{1/2}(\Gamma)} \, dt \leq \int_{\Sigma} \hat{\beta}_\Gamma(\zeta_\Gamma) \, dS \, dt \tag{2.34}$$

for all $\zeta_\Gamma \in L^2(0, T; H^{1/2}(\Gamma))$, where the last integral is intended to be $+\infty$ whenever $\hat{\beta}_\Gamma(\zeta_\Gamma) \notin L^1(\Sigma)$.

Remark 2.14. We can deduce that

$$\mu = -\Delta\varphi + \xi + \pi(\varphi) - f \quad \text{a.e. in } Q, \tag{2.35}$$

$$\theta = \partial_n \varphi + \xi_\Gamma + \pi_\Gamma(\psi) - f_\Gamma \quad \text{in } (H^{1/2}(\Gamma))' \quad \text{a.e. in } (0, T). \tag{2.36}$$

Indeed, (2.35) follows from the fact $\Delta\varphi \in L^2(0, T; H)$ and the weak formulation (2.32). Concerning (2.36), since for almost all $t \in (0, T)$ it holds that $\varphi(t) \in V$ and $\Delta\varphi(t) \in H$, we deduce from [7, Theorem 2.27] that $\partial_n \varphi(t)$ is well defined in $(H^{1/2}(\Gamma))'$. Hence, (2.36) can be deduced from (2.32) by using (2.35).

Theorem 2.15 (Asymptotic limit: $L \in (0, +\infty)$ fixed, $\delta \rightarrow 0$). *Let $L \in (0, +\infty)$ be given. Suppose that the assumptions (A1)–(A4) together with (A5)' are satisfied. We consider a family of data $\{\varphi_0^\delta, f^\delta\}_{\delta \in (0,1)}$ that satisfy the assumptions (A4) and (A5). Assume in addition that there exists a constant $M > 0$ such that, for all $\delta \in (0, 1)$,*

$$\delta \|\nabla_\Gamma \psi_0^\delta\|_{H_\Gamma}^2 + \|\hat{\beta}(\varphi_0^\delta)\|_{L^1(\Omega)} + \|\hat{\beta}_\Gamma(\psi_0^\delta)\|_{L^1(\Gamma)} \leq M, \tag{2.37}$$

$$\|f^\delta\|_{L^2(0,T;\mathcal{H}^1)} \leq M, \tag{2.38}$$

and as $\delta \rightarrow 0$,

$$\varphi_0^\delta \rightharpoonup \varphi_0 \text{ weakly in } \tilde{\mathcal{V}}^1, \quad \mathbf{f}^\delta \rightharpoonup \mathbf{f} \text{ weakly in } L^2(0, T; \mathcal{L}^2). \quad (2.39)$$

Let $\{(\varphi^\delta, \boldsymbol{\mu}^\delta, \boldsymbol{\xi}^\delta)\}_{\delta \in (0,1)}$ be weak solutions to problem $(S_{L,\delta})$ corresponding to data $\{\varphi_0^\delta, \mathbf{f}^\delta\}_{\delta \in (0,1)}$ that are determined by Theorem 2.7. Then, problem $(S_{L,0})$ admits a weak solution $(\varphi, \boldsymbol{\mu}, \boldsymbol{\xi})$ in the sense of Definition 2.13 such that as $\delta \rightarrow 0$ (in the sense of a subsequence),

$$\begin{aligned} \varphi^\delta &\rightharpoonup \varphi && \text{weakly in } H^1(0, T; (\mathcal{H}^1)'), \\ &&& \text{weakly star in } L^\infty(0, T; \tilde{\mathcal{V}}^1), \\ &&& \text{strongly in } C([0, T]; \mathcal{L}^2), \\ \Delta \varphi^\delta &\rightharpoonup \Delta \varphi && \text{weakly in } L^2(0, T; H), \\ \boldsymbol{\mu}^\delta &\rightharpoonup \boldsymbol{\mu} && \text{weakly in } L^2(0, T; \mathcal{H}^1), \\ \boldsymbol{\xi}^\delta &\rightharpoonup \boldsymbol{\xi} && \text{weakly in } L^2(0, T; H \times (H^1(\Gamma))'), \\ -\delta \Delta_\Gamma \psi^\delta + \xi_\Gamma^\delta &\rightharpoonup \xi_\Gamma && \text{weakly in } L^2(0, T; (H^{1/2}(\Gamma))'), \\ \delta \varphi^\delta &\rightarrow \mathbf{0} && \text{strongly in } L^\infty(0, T; \mathcal{V}^1). \end{aligned}$$

Remark 2.16. For any initial data φ_0 satisfying $(A5)'$, the existence of an approximating sequence $\{\varphi_0^\delta\}_{\delta \in (0,1)}$ as in Theorem 2.15 is guaranteed by [18, Proposition A.2].

Uniqueness of the phase function φ associated with problem $(S_{L,0})$ is ensured by the following continuous dependence estimate.

Theorem 2.17 (Continuous dependence, $L \in (0, +\infty)$, $\delta = 0$). *Let $L \in (0, +\infty)$ be given. Suppose that the assumptions (A1)–(A3) are satisfied. Let $(\varphi_i, \boldsymbol{\mu}_i, \boldsymbol{\xi}_i)$, $i \in \{1, 2\}$, be two weak solutions to problem $(S_{L,0})$ corresponding to the data $\{(\varphi_{0,i}, \mathbf{f}_i)\}_{i=1,2}$ satisfying (A4) and $(A5)'$ with $\bar{m}(\varphi_{0,1}) = \bar{m}(\varphi_{0,2}) = \bar{m}_0$. Then, there exists a constant $C > 0$, depending only on K, K_Γ, Ω , and T , such that*

$$\begin{aligned} &\|\varphi_1(t) - \varphi_2(t)\|_{0,*}^2 + \int_0^t \|\varphi_1(s) - \varphi_2(s)\|_{\tilde{\mathcal{V}}_1^1}^2 \, ds \\ &\leq C \left(\|\varphi_{0,1} - \varphi_{0,2}\|_{0,*}^2 + \int_0^t \|\mathbf{f}_1(s) - \mathbf{f}_2(s)\|_{(\tilde{\mathcal{V}}_1^1)'}^2 \, ds \right) \quad \forall t \in [0, T]. \quad (2.40) \end{aligned}$$

Remark 2.18. Thanks to Theorem 2.17, every convergent subsequence $\{\varphi^{\delta_k}\}$ in Theorem 2.15 converges to the same limit φ . In the case when the two graphs β, β_Γ exhibit the same growth, we can obtain further results on refined convergence and error estimate; see Corollary 4.7.

2.3.3. Asymptotic limits: $\delta \in (0, +\infty)$ fixed, $L \rightarrow 0$ or $L \rightarrow +\infty$. Problem $(S_{L,\delta})$ with $L \in (0, +\infty)$ can be viewed as an interpolation of the GMS model (with $L = 0$) and the

LW model (with $L = +\infty$). Assuming $\delta > 0$, below we provide a rigorous investigation of the asymptotic limits as $L \rightarrow 0$ and $L \rightarrow +\infty$, respectively.

First, we introduce the definition of weak solutions to the GMS model (cf., [16, Definition 2.1]).

Definition 2.19. Let $\delta \in (0, +\infty)$ be a fixed parameter, $L = 0$. Suppose that (A1)–(A5) are satisfied. The triplet (φ, μ, ξ) is called a weak solution of problem $(S_{0,\delta})$ on $[0, T]$, if the following conditions are fulfilled.

- (1) The functions φ, μ, ξ have the regularity properties

$$\begin{aligned} \varphi &= \omega + \bar{m}_0 \mathbf{1} \in H^1(0, T; (\mathcal{V}^1)') \cap L^\infty(0, T; \mathcal{V}^1) \cap L^2(0, T; \mathcal{V}^2), \\ \mu &\in L^2(0, T; \mathcal{V}^1), \quad \xi \in L^2(0, T; \mathcal{L}^2). \end{aligned}$$

- (2) The variational formulation

$$\langle \partial_t \varphi(t), y \rangle_{(\mathcal{V}^1)', \mathcal{V}^1} + a_0(\mu(t), y) = 0 \quad \forall y \in \mathcal{V}^1 \quad (2.41)$$

holds for almost all $t \in (0, T)$ and we have

$$\mu = -\Delta \varphi + \xi + \pi(\varphi) - f \quad \text{a.e. in } Q, \quad (2.42)$$

$$\theta = \partial_n \varphi - \delta \Delta_\Gamma \psi + \xi_\Gamma + \pi_\Gamma(\psi) - f_\Gamma \quad \text{a.e. on } \Sigma \quad (2.43)$$

with

$$\xi \in \beta(\varphi) \quad \text{a.e. in } Q, \quad \xi_\Gamma \in \beta_\Gamma(\psi) \quad \text{a.e. on } \Sigma.$$

- (3) The initial conditions are satisfied

$$\varphi|_{t=0} = \varphi_0 \quad \text{a.e. in } \Omega, \quad \psi|_{t=0} = \psi_0 \quad \text{a.e. on } \Gamma.$$

Theorem 2.20 (Asymptotic limit: $\delta \in (0, +\infty)$ fixed, $L \rightarrow 0$). *Let $\delta \in (0, +\infty)$ be given. Suppose that the assumptions (A1)–(A5) are satisfied. For every $L \in (0, 1)$, let $(\varphi^L, \mu^L, \xi^L)$ be a weak solution to problem $(S_{L,\delta})$ corresponding to the data (φ_0, f) . Then, there exists a triplet $(\varphi^0, \mu^0, \xi^0)$ such that as $L \rightarrow 0$ (in the sense of a subsequence),*

$$\varphi^L \rightarrow \varphi^0 \quad \text{weakly in } H^1(0, T; (\mathcal{V}^1)') \cap L^2(0, T; \mathcal{V}^2), \quad (2.44)$$

$$\text{weakly star in } L^\infty(0, T; \mathcal{V}^1), \quad (2.45)$$

$$\text{strongly in } C([0, T]; \mathcal{L}^2), \quad (2.46)$$

$$\mu^L \rightarrow \mu^0 \quad \text{weakly in } L^2(0, T; \mathcal{H}^1), \quad (2.47)$$

$$\xi^L \rightarrow \xi^0 \quad \text{weakly in } L^2(0, T; \mathcal{L}^2) \quad (2.48)$$

with

$$\|\theta^L - \mu^L|_\Gamma\|_{L^2(0,T;H_\Gamma)} \leq C\sqrt{L}. \quad (2.49)$$

The limit triplet $(\varphi^0, \mu^0, \xi^0)$ is a weak solution of the GMS model $(S_{0,\delta})$ corresponding to the data (φ_0, f) in the sense of Definition 2.19.

Remark 2.21. Thanks to [16, Theorem 2.1], the limit function φ^0 , as a solution to problem $(S_{0,\delta})$, is unique. Hence, every convergent subsequence $\{\varphi^{L_k}\}$ in Theorem 2.20 converges to the same limit φ^0 .

Next, we introduce the definition of weak solutions to the LW model (cf., [20, Definition 2.1]).

Definition 2.22. Let $\delta \in (0, +\infty)$ be a fixed parameter, $L = +\infty$. Suppose that the assumptions (A1)–(A5) are satisfied. The triplet (φ, μ, ξ) is called a weak solution to problem $(S_{\infty,\delta})$ on $[0, T]$, if the functions φ, μ, ξ have the regularity properties

$$\begin{aligned} \varphi &\in H^1(0, T; V') \cap L^\infty(0, T; V) \cap L^2(0, T; H^2(\Omega)), \\ \psi &\in H^1(0, T; V'_\Gamma) \cap L^\infty(0, T; V_\Gamma) \cap L^2(0, T; H^2(\Gamma)), \\ \mu &\in L^2(0, T; V), \quad \xi \in L^2(0, T; H), \\ \theta &\in L^2(0, T; V_\Gamma), \quad \xi_\Gamma \in L^2(0, T; H_\Gamma), \end{aligned}$$

and they satisfy

$$\begin{aligned} (\partial_t \varphi, z)_{V',V} + \int_\Omega \nabla \mu \cdot \nabla z \, dx &= 0 \quad \forall z \in V, && \text{a.e. in } (0, T), \\ \mu &= -\Delta \varphi + \xi + \pi(\varphi) - f, \quad \xi \in \beta(\varphi), && \text{a.e. in } Q, \\ \varphi &= \psi, && \text{a.e. on } \Sigma, \\ (\partial_t \psi, z_\Gamma)_{V'_\Gamma, V_\Gamma} + \int_\Gamma \nabla_\Gamma \theta \cdot \nabla_\Gamma z_\Gamma \, dS &= 0 \quad \forall z_\Gamma \in V_\Gamma, && \text{a.e. in } (0, T), \\ \theta &= \partial_n \varphi - \delta \Delta_\Gamma \psi + \xi_\Gamma + \pi_\Gamma(\psi) - f_\Gamma, \quad \xi_\Gamma \in \beta_\Gamma(\psi), && \text{a.e. on } \Sigma, \\ \varphi|_{t=0} &= \varphi_0, \quad \text{a.e. in } \Omega, \quad \psi|_{t=0} = \psi_0, && \text{a.e. on } \Gamma. \end{aligned}$$

Theorem 2.23 (Asymptotic limit: $\delta \in (0, +\infty)$ fixed, $L \rightarrow +\infty$). *Let $\delta \in (0, +\infty)$ be given. Suppose that the assumptions (A1)–(A5) are satisfied with π, π_Γ being differentiable on \mathbb{R} . For every $L \in (1, +\infty)$, let $(\varphi^L, \mu^L, \xi^L)$ be a weak solution to problem $(S_{L,\delta})$ corresponding to the data (φ_0, \mathbf{f}) . Then, there exists a triplet $(\varphi^\infty, \mu^\infty, \xi^\infty)$ such that as $L \rightarrow +\infty$ (in the sense of a subsequence),*

$$\begin{aligned} \varphi^L &\rightarrow \varphi^\infty \quad \text{weakly in } H^1(0, T; V') \cap L^2(0, T; H^2(\Omega)), \\ &\quad \text{weakly star in } L^\infty(0, T; V), \\ &\quad \text{strongly in } C([0, T]; H), \\ \psi^L &\rightarrow \psi^\infty \quad \text{weakly in } H^1(0, T; V'_\Gamma) \cap L^2(0, T; H^2(\Gamma)), \\ &\quad \text{weakly star in } L^\infty(0, T; V_\Gamma), \\ &\quad \text{strongly in } C([0, T]; H_\Gamma), \\ \mu^L &\rightarrow \mu^\infty \quad \text{weakly in } L^2(0, T; \mathcal{H}^1), \\ \xi^L &\rightarrow \xi^\infty \quad \text{weakly in } L^2(0, T; \mathcal{L}^2), \end{aligned}$$

and

$$\frac{1}{L} \|\theta^L - \mu^L|_\Gamma\|_{L^2(0,T;H_\Gamma)} \leq \frac{C}{\sqrt{L}}.$$

The limit triplet $(\varphi^\infty, \mu^\infty, \xi^\infty)$ is a weak solution of the LW model corresponding to the data (φ_0, f) in the sense of Definition 2.22.

Remark 2.24. Thanks to [20, Theorem 2.4], the limit function φ^∞ , as a solution to problem $(S_{\infty,\delta})$, is unique. Hence, every convergent subsequence $\{\varphi^{L_k}\}$ in Theorem 2.23 converges to the same limit φ^∞ .

3. Well-posedness in the presence of surface diffusion

In this section, we focus on the case when the surface diffusion on Γ is present, that is, $\delta > 0$. Then, we prove Theorems 2.7, 2.9, and 2.11 on the well-posedness of problem $(S_{L,\delta})$ for any given $L, \delta \in (0, +\infty)$.

3.1. Approximation via Yosida regularization

To prove the existence of weak solutions to problem $(S_{L,\delta})$ with general potentials, we approximate the maximal monotone operators β and β_Γ by means of suitable Yosida regularizations; see, e.g., [16, 20] for the case of Cahn–Hilliard equations with dynamic boundary conditions. For each $\varepsilon \in (0, 1)$, we define $\beta_\varepsilon, \beta_{\Gamma,\varepsilon} : \mathbb{R} \rightarrow \mathbb{R}$, along with the associated resolvent operators $J_\varepsilon, J_{\Gamma,\varepsilon} : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\beta_\varepsilon(r) := \frac{1}{\varepsilon}(r - J_\varepsilon(r)) := \frac{1}{\varepsilon}(r - (I + \varepsilon\beta)^{-1}(r)), \tag{3.1}$$

$$\beta_{\Gamma,\varepsilon}(r) := \frac{1}{\varepsilon\varrho}(r - J_{\Gamma,\varepsilon}(r)) := \frac{1}{\varepsilon\varrho}(r - (I + \varepsilon\varrho\beta_\Gamma)^{-1}(r)) \tag{3.2}$$

for all $r \in \mathbb{R}$, where $\varrho > 0$ is the constant given in the condition (2.22). The related Moreau–Yosida regularizations $\hat{\beta}_\varepsilon, \hat{\beta}_{\Gamma,\varepsilon}$ of $\hat{\beta}, \hat{\beta}_\Gamma : \mathbb{R} \rightarrow \mathbb{R}$ are given by (cf., e.g., [60])

$$\begin{aligned} \hat{\beta}_\varepsilon(r) &:= \inf_{s \in \mathbb{R}} \left\{ \frac{1}{2\varepsilon}|r - s|^2 + \hat{\beta}(s) \right\} = \frac{1}{2\varepsilon}|r - J_\varepsilon(r)|^2 + \hat{\beta}(J_\varepsilon(r)) = \int_0^r \beta_\varepsilon(s) \, ds, \\ \hat{\beta}_{\Gamma,\varepsilon}(r) &:= \inf_{s \in \mathbb{R}} \left\{ \frac{1}{2\varepsilon\varrho}|r - s|^2 + \hat{\beta}_\Gamma(s) \right\} = \int_0^r \beta_{\Gamma,\varepsilon}(s) \, ds \quad \forall r \in \mathbb{R}. \end{aligned}$$

From (A1), (A2), we find $\beta_\varepsilon(0) = \beta_{\Gamma,\varepsilon}(0) = 0$. Besides, $\beta_\varepsilon, \beta_{\Gamma,\varepsilon}$ are Lipschitz continuous with Lipschitz constants $1/\varepsilon$ and $1/\varepsilon\varrho$, respectively (see [6, Propositions 2.6 and 2.7]). Thus, it follows that $\hat{\beta}_\varepsilon$ and $\hat{\beta}_{\Gamma,\varepsilon}$ are nonnegative convex functions with (at most) quadratic growth. Moreover, it holds (cf., e.g., [4, 60])

$$|\beta_\varepsilon(r)| \leq |\beta^\circ(r)| \quad \text{for } r \in D(\beta), \quad |\beta_{\Gamma,\varepsilon}(r)| \leq |\beta_\Gamma^\circ(r)| \quad \text{for } r \in D(\beta_\Gamma), \tag{3.3}$$

$$0 \leq \hat{\beta}_\varepsilon(r) \leq \hat{\beta}(r), \quad 0 \leq \hat{\beta}_{\Gamma,\varepsilon}(r) \leq \hat{\beta}_\Gamma(r) \quad \forall r \in \mathbb{R}. \tag{3.4}$$

Thanks to [9, Lemma 4.4], we keep the compatibility condition

$$|\beta_\varepsilon(r)| \leq \varrho|\beta_{\Gamma,\varepsilon}(r)| + c_0 \quad \forall r \in \mathbb{R}, \tag{3.5}$$

with the same constants ϱ and c_0 as in (2.22). Additionally, the following inequalities hold for $\varepsilon \in (0, 1)$ (see [34, Section 5]): for any given $r_0 \in \text{Int } D(\beta_\Gamma)$,

$$\beta_\varepsilon(r)(r - r_0) \geq \delta_0|\beta_\varepsilon(r)| - c_1, \quad \beta_{\Gamma,\varepsilon}(r)(r - r_0) \geq \delta_0|\beta_{\Gamma,\varepsilon}(r)| - c_1 \tag{3.6}$$

for all $r \in \mathbb{R}$, where δ_0, c_1 are positive constants depending on r_0 but independent of ε .

Recalling the abstract formulation (2.25) for problem $(S_{L,\delta})$, it is straightforward to check that the first and second equations in (2.25) yield the single evolution equation:

$$\mathfrak{C}^L(\omega'(t)) + \partial\Phi_\delta(\omega(t)) = \mathbf{P}(-\xi(t) - \pi(\omega(t) + \bar{m}_0\mathbf{1}) + \mathbf{f}(t)) \quad \text{in } \mathcal{L}^2_{(0)} \tag{3.7}$$

for almost all $t \in (0, T)$, where

$$\xi(t) \in \beta(\omega(t) + \bar{m}_0\mathbf{1}) \quad \text{in } \mathcal{L}^2 \quad \text{for a.a. } t \in (0, T).$$

For each $\varepsilon \in (0, 1)$, let us consider the following approximating problem for (3.7): find $\omega_\varepsilon := (\omega_\varepsilon, \omega_{\Gamma,\varepsilon})$ satisfying

$$\begin{aligned} \varepsilon\omega'_\varepsilon(t) + \mathfrak{C}^L(\omega'_\varepsilon(t)) + \partial\Phi_\delta(\omega_\varepsilon(t)) \\ = \mathbf{P}(-\beta_\varepsilon(\omega_\varepsilon(t) + \bar{m}_0\mathbf{1})) \\ + \mathbf{P}(-\pi(\omega_\varepsilon(t) + \bar{m}_0\mathbf{1})) + \mathbf{P}(\mathbf{f}(t)) \quad \text{in } \mathcal{L}^2_{(0)} \quad \text{for a.a. } t \in (0, T), \end{aligned} \tag{3.8}$$

$$\omega_\varepsilon|_{t=0} = \omega_0 := \varphi_0 - \bar{m}_0\mathbf{1}, \quad \text{in } \mathcal{L}^2_{(0)}. \tag{3.9}$$

Proposition 3.1. *Assume that the assumptions (A1)–(A5) are satisfied. For each $\varepsilon \in (0, 1)$, problem (3.8)–(3.9) admits a unique solution*

$$\omega_\varepsilon \in H^1(0, T; \mathcal{L}^2_{(0)}) \cap C([0, T]; \mathcal{V}^1_{(0)}) \cap L^2(0, T; \mathcal{V}^2_{(0)}).$$

Proof. The proof mainly follows the argument for [16, Proposition 4.1], which is based on the abstract theory of doubly nonlinear evolution inclusions [24] combined with the contraction mapping principle.

By the definition of \mathfrak{C}^L , it is straightforward to check that

$$\begin{aligned} ((\varepsilon I + \mathfrak{C}^L)\mathbf{z}, \mathbf{z})_{\mathcal{X}^2} &= \varepsilon\|\mathbf{z}\|_{\mathcal{X}^2}^2 + (\mathfrak{C}^L\mathbf{z}, \mathbf{z})_{\mathcal{X}^2} \\ &= \varepsilon\|\mathbf{z}\|_{\mathcal{X}^2}^2 + (\mathfrak{C}^L\mathbf{z}, \mathfrak{C}^L\mathbf{z})_{\mathcal{H}^1_{L,0}} \\ &\geq \varepsilon\|\mathbf{z}\|_{\mathcal{X}^2}^2 \quad \forall \mathbf{z} \in \mathcal{L}^2_{(0)}. \end{aligned}$$

On the other hand, thanks to (2.9), we find

$$\|(\varepsilon I + \mathfrak{C}^L)\mathbf{z}\|_{\mathcal{X}^2} \leq \varepsilon\|\mathbf{z}\|_{\mathcal{X}^2} + \|\mathfrak{C}^L\mathbf{z}\|_{\mathcal{X}^2} \leq (\varepsilon + C)\|\mathbf{z}\|_{\mathcal{X}^2} \quad \forall \mathbf{z} \in \mathcal{L}^2_{(0)}.$$

As a consequence, for each $\varepsilon \in (0, 1)$, the operator $\varepsilon I + \mathfrak{G}^L$ is coercive and with linear growth in $\mathcal{X}_{(0)}^2$. These facts enable us to apply the abstract theory [24, Theorem 2.1] with the particular choices $A = \varepsilon I + \mathfrak{G}^L$, $B = \partial\Phi_\delta$, $H = \mathcal{X}_{(0)}^2$, and $V = \mathcal{V}_{(0)}^1$ therein. Hence, for any given function $\tilde{\omega} \in C([0, T]; \mathcal{X}_{(0)}^2)$, we can conclude that there exists a unique

$$\omega \in H^1(0, T; \mathcal{X}_{(0)}^2) \cap L^\infty(0, T; \mathcal{V}_{(0)}^1) \subset C([0, T]; \mathcal{X}_{(0)}^2)$$

satisfying $\partial\Phi_\delta(\omega) \in L^2(0, T; \mathcal{X}_{(0)}^2)$ and

$$\begin{aligned} & (\varepsilon I + \mathfrak{G}^L)\omega'(t) + \partial\Phi_\delta(\omega(t)) \\ &= \mathbf{P}(-\beta_\varepsilon(\tilde{\omega}(t) + \bar{m}_0\mathbf{1}) - \pi(\tilde{\omega}(t) + \bar{m}_0\mathbf{1}) + f(t)) \quad \text{in } \mathcal{X}_{(0)}^2 \quad \text{for a.a. } t \in (0, T), \\ & \omega|_{t=0} = \omega_0 \quad \text{in } \mathcal{X}_{(0)}^2. \end{aligned}$$

In this way, we can define the map $\Psi : \tilde{\omega} \mapsto \omega$ from $C([0, T]; \mathcal{X}_{(0)}^2)$ into itself. Next, for given $\tilde{\omega}^{(i)} \in C([0, T]; \mathcal{X}_{(0)}^2)$, $i \in \{1, 2\}$, we set $\omega^{(i)} := \Psi(\tilde{\omega}^{(i)})$. Consider the equation of their differences

$$\begin{aligned} & \varepsilon((\omega^{(1)})' - (\omega^{(2)})') + \mathfrak{G}^L((\omega^{(1)})' - (\omega^{(2)})') + \partial\Phi_\delta(\omega^{(1)} - \omega^{(2)}) \\ &= \mathbf{P}(-\beta_\varepsilon(\tilde{\omega}^{(1)} + \bar{m}_0\mathbf{1}) + \beta_\varepsilon(\tilde{\omega}^{(2)} + \bar{m}_0\mathbf{1}) - \pi(\tilde{\omega}^{(1)} + \bar{m}_0\mathbf{1}) + \pi(\tilde{\omega}^{(2)} + \bar{m}_0\mathbf{1})). \end{aligned}$$

Taking the \mathcal{X}^2 inner product with $\omega^{(1)} - \omega^{(2)}$ and using the monotonicity of $\partial\Phi_\delta$, we obtain

$$\begin{aligned} & \frac{\varepsilon}{2} \frac{d}{dt} \|\omega^{(1)} - \omega^{(2)}\|_{\mathcal{X}_{(0)}^2}^2 + \frac{1}{2} \frac{d}{dt} \|\omega^{(1)} - \omega^{(2)}\|_{0,*}^2 \\ & \leq (-\beta_\varepsilon(\tilde{\omega}^{(1)} + \bar{m}_0\mathbf{1}) + \beta_\varepsilon(\tilde{\omega}^{(2)} + \bar{m}_0\mathbf{1}), \omega^{(1)} - \omega^{(2)})_{\mathcal{X}^2} \\ & \quad + (-\pi(\tilde{\omega}^{(1)} + \bar{m}_0\mathbf{1}) + \pi(\tilde{\omega}^{(2)} + \bar{m}_0\mathbf{1}), \omega^{(1)} - \omega^{(2)})_{\mathcal{X}^2} \\ & \leq C_\varepsilon \|\tilde{\omega}^{(1)} - \tilde{\omega}^{(2)}\|_{\mathcal{X}_{(0)}^2} \|\omega^{(1)} - \omega^{(2)}\|_{\mathcal{X}_{(0)}^2}, \end{aligned} \tag{3.10}$$

where the last inequality follows from the fact that $\beta_\varepsilon, \beta_{\Gamma,\varepsilon}, \pi, \pi_\Gamma$ are Lipschitz continuous, and $C_\varepsilon > 0$ depends on K, K_Γ, ε . Using the Cauchy–Schwarz inequality and Grönwall’s lemma, we deduce from (3.10) that

$$\|\omega^{(1)}(t) - \omega^{(2)}(t)\|_{\mathcal{X}_{(0)}^2}^2 \leq C_\varepsilon \int_0^t \|\tilde{\omega}^{(1)}(s) - \tilde{\omega}^{(2)}(s)\|_{\mathcal{X}_{(0)}^2}^2 \, ds \quad \forall t \in [0, T],$$

which implies

$$\|\omega^{(1)}(t) - \omega^{(2)}(t)\|_{\mathcal{X}_{(0)}^2}^2 \leq C_\varepsilon t \|\tilde{\omega}^{(1)} - \tilde{\omega}^{(2)}\|_{C([0,T]; \mathcal{X}_{(0)}^2)}^2 \quad \forall t \in [0, T].$$

For $k \in \mathbb{Z}^+$, we set $\omega_k^{(i)} := \Psi^k(\tilde{\omega}^{(i)})$. By iteration, we infer from the above estimates that

$$\|\omega_k^{(1)}(t) - \omega_k^{(2)}(t)\|_{\mathcal{X}_{(0)}^2}^2 \leq C_\varepsilon \left(\frac{t^k}{k!}\right) \|\tilde{\omega}^{(1)} - \tilde{\omega}^{(2)}\|_{C([0,T]; \mathcal{X}_{(0)}^2)}^2 \quad \forall t \in [0, T].$$

Hence, there exists some $k_0 \in \mathbb{Z}^+$ large enough such that

$$\|\omega_{k_0}^{(1)} - \omega_{k_0}^{(2)}\|_{C([0,T]; \mathcal{L}_{(0)}^2)}^2 \leq \frac{1}{2} \|\tilde{\omega}^{(1)} - \tilde{\omega}^{(2)}\|_{C([0,T]; \mathcal{L}_{(0)}^2)}^2.$$

This yields that Ψ^{k_0} is a contraction mapping from $C([0, T]; \mathcal{L}_{(0)}^2)$ into itself.

Therefore, thanks to the contraction mapping principle, Ψ^{k_0} admits a unique fixed point $\omega_\varepsilon \in C([0, T]; \mathcal{L}_{(0)}^2)$. It follows that

$$\Psi(\omega_\varepsilon) = \Psi(\Psi^{k_0}(\omega_\varepsilon)) = \Psi^{k_0}(\Psi(\omega_\varepsilon)).$$

Hence, we find $\Psi(\omega_\varepsilon) = \omega_\varepsilon$ because of the uniqueness of ω_ε . By the definition of Ψ , ω_ε is indeed a solution to problem (3.8)–(3.9). Its uniqueness easily follows from an estimate similar to (3.10) and Grönwall’s lemma.

For every $\varepsilon \in (0, 1)$, we have shown that problem (3.8)–(3.9) admits a unique solution such that

$$\omega_\varepsilon \in H^1(0, T; \mathcal{L}_{(0)}^2) \cap L^\infty(0, T; \mathcal{V}_{(0)}^1) \quad \text{with } \partial\Phi_\delta(\omega_\varepsilon) \in L^2(0, T; \mathcal{L}_{(0)}^2).$$

In view of (2.24), we can apply the elliptic estimate [43, Theorem 3.3] to conclude that $\omega_\varepsilon \in L^2(0, T; \mathcal{V}_{(0)}^2)$. Finally, observing that

$$H^1(0, T; \mathcal{L}_{(0)}^2) \cap L^2(0, T; \mathcal{V}_{(0)}^2) \subset C([0, T]; \mathcal{V}_{(0)}^1),$$

we arrive at the conclusion of Proposition 3.1. ■

3.2. Uniform estimates

For every $\varepsilon \in (0, 1)$, in view of Proposition 3.1, we set $\varphi_\varepsilon := \omega_\varepsilon + \bar{m}_0 \mathbf{1}$ and

$$\mu_\varepsilon(t) := \varepsilon \omega'_\varepsilon(t) + \partial\Phi_\delta(\omega_\varepsilon(t)) + \beta_\varepsilon(\varphi_\varepsilon(t)) + \pi(\varphi_\varepsilon(t)) - f(t) \quad \text{for a.a. } t \in (0, T).$$

Then, the evolution equation (3.8) can be written as

$$\mathcal{G}^L(\omega'_\varepsilon(t)) + \mu_\varepsilon(t) - m_\varepsilon(t) \mathbf{1} = \mathbf{0} \quad \text{in } \mathcal{L}_{(0)}^2 \quad \text{for a.a. } t \in (0, T) \tag{3.11}$$

with

$$m_\varepsilon(t) := \bar{m}(\beta_\varepsilon(\varphi_\varepsilon(t)) + \pi(\varphi_\varepsilon(t)) - f(t)) \quad \text{for a.a. } t \in (0, T). \tag{3.12}$$

It follows that

$$\mathbf{P}\mu_\varepsilon = \mu_\varepsilon - m_\varepsilon \mathbf{1} \in L^2(0, T; \mathcal{H}_{(0)}^1) \quad \text{and} \quad m_\varepsilon \in L^2(0, T).$$

As a consequence, $\mu_\varepsilon = (\mu_\varepsilon, \theta_\varepsilon) \in L^2(0, T; \mathcal{H}^1)$ and the following weak formulations hold:

$$\begin{aligned} & \int_\Omega \partial_t \omega_\varepsilon(t) y \, dx + \int_\Gamma \partial_t \omega_{\Gamma, \varepsilon}(t) y_\Gamma \, dS + \int_\Omega \nabla \mu_\varepsilon(t) \cdot \nabla y \, dx + \int_\Gamma \nabla_\Gamma \theta_\varepsilon(t) \cdot \nabla_\Gamma y_\Gamma \, dS \\ & + \frac{1}{L} \int_\Gamma (\mu_\varepsilon(t) - \theta_\varepsilon(t))(y - y_\Gamma) \, dS = 0 \quad \forall y \in \mathcal{H}_{(0)}^1 \end{aligned} \tag{3.13}$$

and

$$\begin{aligned}
 & \int_{\Omega} \mu_{\varepsilon}(t) z \, dx + \int_{\Gamma} \theta_{\varepsilon}(t) z_{\Gamma} \, dS \\
 &= \varepsilon \int_{\Omega} \partial_t \omega_{\varepsilon}(t) z \, dx + \varepsilon \int_{\Gamma} \partial_t \omega_{\Gamma, \varepsilon}(t) z_{\Gamma} \, dS + \int_{\Omega} \nabla \omega_{\varepsilon}(t) \cdot \nabla z \, dx \\
 &+ \delta \int_{\Gamma} \nabla_{\Gamma} \omega_{\Gamma, \varepsilon}(t) \cdot \nabla_{\Gamma} z_{\Gamma} \, dS + \int_{\Omega} [\beta_{\varepsilon}(\varphi_{\varepsilon}(t)) + \pi(\varphi_{\varepsilon}(t)) - f(t)] z \, dx \\
 &+ \int_{\Gamma} [\beta_{\Gamma, \varepsilon}(\psi_{\varepsilon}(t)) + \pi_{\Gamma}(\psi_{\varepsilon}(t)) - f_{\Gamma}(t)] z_{\Gamma} \, dS \quad \forall z \in \mathcal{V}^1
 \end{aligned} \tag{3.14}$$

for almost all $t \in (0, T)$. In particular, it follows from (3.14) that

$$\mu_{\varepsilon} = \varepsilon \partial_t \omega_{\varepsilon} - \Delta \omega_{\varepsilon} + \beta_{\varepsilon}(\varphi_{\varepsilon}) + \pi(\varphi_{\varepsilon}) - f, \quad \text{a.e. in } Q, \tag{3.15}$$

$$\theta_{\varepsilon} = \varepsilon \partial_t \omega_{\Gamma, \varepsilon} + \partial_{\mathbf{n}} \omega_{\varepsilon} - \delta \Delta_{\Gamma} \omega_{\Gamma, \varepsilon} + \beta_{\Gamma, \varepsilon}(\psi_{\varepsilon}) + \pi_{\Gamma}(\psi_{\varepsilon}) - f_{\Gamma}, \quad \text{a.e. on } \Sigma. \tag{3.16}$$

Besides, since $\omega'_{\varepsilon} \in L^2(0, T; \mathcal{L}^2_{(0)})$, we infer from (3.11) and the elliptic estimate (2.10) that $\mu_{\varepsilon} \in L^2(0, T; \mathcal{H}^2)$. This allows us to deduce from (3.13) the following pointwise relations:

$$\partial_t \omega_{\varepsilon} = \Delta \mu_{\varepsilon}, \quad \text{a.e. in } Q, \tag{3.17}$$

$$\partial_t \omega_{\Gamma, \varepsilon} = \Delta_{\Gamma} \theta_{\varepsilon} - \partial_{\mathbf{n}} \mu_{\varepsilon}, \quad \text{a.e. on } \Sigma, \tag{3.18}$$

$$L \partial_{\mathbf{n}} \mu_{\varepsilon} = \theta_{\varepsilon} - \mu_{\varepsilon}, \quad \text{a.e. on } \Sigma. \tag{3.19}$$

Now, we proceed to derive uniform estimates with respect to the approximating parameter $\varepsilon \in (0, 1)$.

Lemma 3.2. *The mass is conserved in time, that is,*

$$\bar{m}(\varphi_{\varepsilon}(t)) = \bar{m}(\varphi_0) = \bar{m}_0 \quad \forall t \in [0, T]. \tag{3.20}$$

Proof. The conclusion (3.20) easily follows from the definition of φ_{ε} and the fact that $\omega_{\varepsilon} \in C([0, T]; \mathcal{L}^2_{(0)})$. ■

Lemma 3.3. *There exists a positive constant M_1 , independent of $\varepsilon, \delta \in (0, 1)$, such that*

$$\begin{aligned}
 & \varepsilon^{1/2} \|\omega_{\varepsilon}\|_{L^{\infty}(0, T; \mathcal{L}^2_{(0)})} + \|\omega_{\varepsilon}\|_{L^{\infty}(0, T; \mathcal{H}^{-1}_{(0)})} + \|\omega_{\varepsilon}\|_{L^2(0, T; \mathcal{V}^1_{(0)})} \\
 &+ \|\beta_{\varepsilon}(\varphi_{\varepsilon})\|_{L^1(0, T; L^1(\Omega))} + \|\beta_{\Gamma, \varepsilon}(\psi_{\varepsilon})\|_{L^1(0, T; L^1(\Gamma))} \leq M_1,
 \end{aligned} \tag{3.21}$$

where the norm $\|\cdot\|_{\mathcal{V}^1_{(0)}}$ is defined as in (2.7).

Proof. We adopt the same strategy as in [9, 15, 16]. Testing (3.8) at time $s \in (0, T)$ by ω_{ε} , using (2.23) and the definition of the subdifferential $\partial \Phi_{\delta}$ with the fact $\Phi_{\delta}(\mathbf{0}) = 0$, we obtain

$$\begin{aligned}
 & \varepsilon (\omega'_{\varepsilon}(s), \omega_{\varepsilon}(s))_{\mathcal{L}^2} + (\mathfrak{C}^L(\omega'_{\varepsilon}(s)), \omega_{\varepsilon}(s))_{\mathcal{L}^2} + \Phi_{\delta}(\omega_{\varepsilon}(s)) \\
 &+ (\beta_{\varepsilon}(\varphi_{\varepsilon}(s)), \omega_{\varepsilon}(s))_{\mathcal{L}^2} \leq (f(s) - \pi(\varphi_{\varepsilon}(s)), \omega_{\varepsilon}(s))_{\mathcal{L}^2}
 \end{aligned} \tag{3.22}$$

for almost all $s \in (0, T)$. Recalling the assumptions (A2) and (A5), we find $\bar{m}_0 \in \text{Int } D(\beta_\Gamma)$. Then, we can apply the inequalities in (3.6) with $r_0 = \bar{m}_0$ to get

$$\begin{aligned} & (\beta_\varepsilon(\varphi_\varepsilon(s)), \omega_\varepsilon(s))_{\mathcal{X}^2} \\ &= \int_\Omega \beta_\varepsilon(\varphi_\varepsilon(s))(\varphi_\varepsilon(s) - \bar{m}_0) \, dx + \int_\Gamma \beta_{\Gamma,\varepsilon}(\psi_\varepsilon(s))(\psi_\varepsilon(s) - \bar{m}_0) \, dS \\ &\geq \delta_0 \int_\Omega |\beta_\varepsilon(\varphi_\varepsilon(s))| \, dx - c_1|\Omega| + \delta_0 \int_\Gamma |\beta_{\Gamma,\varepsilon}(\psi_\varepsilon(s))| \, dS - c_1|\Gamma| \end{aligned} \tag{3.23}$$

for almost all $s \in (0, T)$. Furthermore, owing to the assumption (A3) and the interpolation inequality (2.13), there exists a positive constant \tilde{M}_1 , depending only on $K, K_\Gamma, \pi(\bar{m}_0), \pi_\Gamma(\bar{m}_0), |\Omega|$, and $|\Gamma|$, such that

$$\begin{aligned} & (\mathbf{f}(s) - \boldsymbol{\pi}(\varphi_\varepsilon(s)), \omega_\varepsilon(s))_{\mathcal{X}^2} \\ &\leq \frac{1}{2} \int_\Omega |\mathbf{f}(s)|^2 \, dx + \frac{1}{2} \int_\Omega |\omega_\varepsilon(s)|^2 \, dx + K \int_\Omega |\omega_\varepsilon(s)|^2 \, dx \\ &\quad + \frac{1}{2} \int_\Gamma |\mathbf{f}_\Gamma(s)|^2 \, dS + \frac{1}{2} \int_\Gamma |\omega_{\Gamma,\varepsilon}(s)|^2 \, dS + K_\Gamma \int_\Gamma |\omega_{\Gamma,\varepsilon}(s)|^2 \, dS \\ &\quad + |\pi(\bar{m}_0)| \int_\Omega |\omega_\varepsilon(s)| \, dx + |\pi_\Gamma(\bar{m}_0)| \int_\Gamma |\omega_{\Gamma,\varepsilon}(s)| \, dS \\ &\leq \tilde{M}_1(1 + \|\mathbf{f}(s)\|_{\mathcal{X}^2}^2 + \|\omega_\varepsilon(s)\|_{0,*}^2) + \frac{1}{4} \|\nabla \omega_\varepsilon(s)\|_H^2 \quad \text{for a.a. } s \in (0, T). \end{aligned} \tag{3.24}$$

From (3.22)–(3.24), we obtain

$$\begin{aligned} & \varepsilon \frac{d}{ds} \|\omega_\varepsilon(s)\|_{\mathcal{X}^2}^2 + \frac{d}{ds} \|\omega_\varepsilon(s)\|_{0,*}^2 + \frac{1}{2} \|\nabla \omega_\varepsilon(s)\|_H^2 + \delta \|\nabla_\Gamma \omega_{\Gamma,\varepsilon}(s)\|_{H_\Gamma}^2 \\ &\quad + 2\delta_0 \int_\Omega |\beta_\varepsilon(\varphi_\varepsilon(s))| \, dx + 2\delta_0 \int_\Gamma |\beta_{\Gamma,\varepsilon}(\psi_\varepsilon(s))| \, dS \\ &\leq 2c_1(|\Omega| + |\Gamma|) + 2\tilde{M}_1(1 + \|\mathbf{f}(s)\|_{\mathcal{X}^2}^2 + \|\omega_\varepsilon(s)\|_{0,*}^2) \quad \text{for a.a. } s \in (0, T), \end{aligned}$$

which combined with Grönwall’s lemma leads to the conclusion (3.21). ■

Lemma 3.4. *There exists a positive constant M_2 , independent of $\varepsilon, \delta \in (0, 1)$, such that*

$$\begin{aligned} & \|\omega_\varepsilon\|_{L^\infty(0,T;V_{(0)}^1)} + \varepsilon^{1/2} \|\omega'_\varepsilon\|_{L^2(0,T;\mathcal{X}_{(0)}^2)} + \|\omega'_\varepsilon\|_{L^2(0,T;\mathcal{H}_{(0)}^{-1})} \\ &\quad + \|\hat{\beta}_\varepsilon(\varphi_\varepsilon)\|_{L^\infty(0,T;L^1(\Omega))} + \|\hat{\beta}_{\Gamma,\varepsilon}(\psi_\varepsilon)\|_{L^\infty(0,T;L^1(\Gamma))} \leq M_2, \end{aligned} \tag{3.25}$$

where the norm $\|\cdot\|_{V_{(0)}^1}$ is defined as in (2.7).

Proof. Testing (3.8) at time $s \in (0, T)$ by $\omega'_\varepsilon = \varphi'_\varepsilon$, using (2.23) and the chain rule for subdifferentials (see, e.g., [60, Lemma 4.3], and also [23, Proposition A.1]), we find

$$\begin{aligned} & \frac{d}{ds} \left(\Phi_\delta(\omega_\varepsilon(s)) + \int_\Omega (\hat{\beta}_\varepsilon(\varphi_\varepsilon(s)) + \hat{\pi}(\varphi_\varepsilon(s))) \, dx + \int_\Gamma (\hat{\beta}_{\Gamma,\varepsilon}(\psi_\varepsilon(s)) + \hat{\pi}_\Gamma(\psi_\varepsilon(s))) \, dS \right) \\ &\quad + \varepsilon \|\omega'_\varepsilon(s)\|_{\mathcal{X}^2}^2 + \|\omega'_\varepsilon(s)\|_{0,*}^2 \\ &= (\mathbf{P}\mathbf{f}(s), \omega'_\varepsilon(s))_{\mathcal{X}^2} \quad \text{for a.a. } s \in (0, T). \end{aligned} \tag{3.26}$$

Since $\mathfrak{S}^L(\omega'_\varepsilon(s)) \in \mathcal{V}_{(0)}^1$ for almost all $s \in (0, T)$, we observe that

$$(\mathbf{P}\mathbf{f}(s), \omega'_\varepsilon(s))_{\mathcal{X}^2} = (\mathbf{P}\mathbf{f}(s), \mathfrak{S}^L(\omega'_\varepsilon(s)))_{\mathcal{H}_{L,0}^1} \leq \frac{1}{2} \|\omega'_\varepsilon(s)\|_{0,*}^2 + \frac{1}{2} \|\mathbf{f}(s)\|_{\mathcal{V}^1}^2. \quad (3.27)$$

Integrating (3.26) over $(0, t)$ with respect to s , we infer from (A3) (i.e., $\hat{\pi}$, $\hat{\pi}_\Gamma$ are at most with quadratic growth), Lemma 3.2, (3.27), and (3.4) that

$$\begin{aligned} & \frac{1}{2} \|\nabla \omega_\varepsilon(t)\|_H^2 + \frac{\delta}{2} \|\nabla \omega_{\Gamma,\varepsilon}(t)\|_{H_\Gamma}^2 + \int_\Omega \hat{\beta}_\varepsilon(\varphi_\varepsilon(t)) \, dx + \int_\Gamma \hat{\beta}_{\Gamma,\varepsilon}(\psi_\varepsilon(t)) \, dS \\ & \quad + \varepsilon \int_0^t \|\omega'_\varepsilon(s)\|_{\mathcal{X}^2}^2 \, ds + \frac{1}{2} \int_0^t \|\omega'_\varepsilon(s)\|_{0,*}^2 \, ds \\ & \leq \frac{1}{2} \|\omega_0\|_{\mathcal{V}_{(0)}^1}^2 + \int_\Omega \hat{\beta}(\varphi_0) \, dx + \int_\Gamma \hat{\beta}_\Gamma(\psi_0) \, dS + \int_\Omega \hat{\pi}(\varphi_0) \, dx - \int_\Omega \hat{\pi}(\varphi_\varepsilon(t)) \, dx \\ & \quad + \int_\Gamma \hat{\pi}_\Gamma(\psi_0) \, dS - \int_\Gamma \hat{\pi}_\Gamma(\psi_\varepsilon(t)) \, dS + \frac{1}{2} \int_0^t \|\mathbf{f}(s)\|_{\mathcal{V}^1}^2 \, ds \\ & \leq \frac{1}{2} \|\omega_0\|_{\mathcal{V}_{(0)}^1}^2 + \int_\Omega \hat{\beta}(\varphi_0) \, dx + \int_\Gamma \hat{\beta}_\Gamma(\psi_0) \, dS \\ & \quad + C(1 + \|\varphi_0\|_{\mathcal{X}^2}^2 + \|\omega_\varepsilon(t) + \bar{m}_0 \mathbf{1}\|_{\mathcal{X}^2}^2) \\ & \quad + \frac{1}{2} \int_0^t \|\mathbf{f}(s)\|_{\mathcal{V}^1}^2 \, ds \quad \forall t \in [0, T]. \end{aligned} \quad (3.28)$$

The interpolation inequality (2.13) implies

$$\|\omega_\varepsilon(t)\|_{\mathcal{X}^2}^2 \leq \gamma \|\nabla \omega_\varepsilon(t)\|_H^2 + C_\gamma \|\omega_\varepsilon(t)\|_{0,*}^2 \quad \forall \gamma > 0.$$

Then, taking γ sufficiently small (independent of δ), we infer from (3.21), (3.28) and (A5) that

$$\begin{aligned} & \frac{1}{4} \|\nabla \omega_\varepsilon(t)\|_H^2 + \frac{\delta}{2} \|\nabla \omega_{\Gamma,\varepsilon}(t)\|_{H_\Gamma}^2 + \int_\Omega \hat{\beta}_\varepsilon(\varphi_\varepsilon(t)) \, dx + \int_\Gamma \hat{\beta}_{\Gamma,\varepsilon}(\psi_\varepsilon(t)) \, dS \\ & \quad + \varepsilon \int_0^t \|\omega'_\varepsilon(s)\|_{\mathcal{X}^2}^2 \, ds + \frac{1}{2} \int_0^t \|\omega'_\varepsilon(s)\|_{0,*}^2 \, ds \\ & \leq C + \frac{1}{2} \int_0^t \|\mathbf{f}(s)\|_{\mathcal{V}^1}^2 \, ds \quad \forall t \in [0, T], \end{aligned} \quad (3.29)$$

which gives the estimate (3.25). ■

Lemma 3.5. *There exist positive constants M_3 and M_4 , independent of $\varepsilon, \delta \in (0, 1)$, such that*

$$\|\beta_\varepsilon(\varphi_\varepsilon)\|_{L^2(0,T;L^1(\Omega))} + \|\beta_{\Gamma,\varepsilon}(\psi_\varepsilon)\|_{L^2(0,T;L^1(\Gamma))} \leq M_3, \quad (3.30)$$

$$\|m_\varepsilon\|_{L^2(0,T)} + \|\mu_\varepsilon\|_{L^2(0,T;\mathcal{H}^1)} \leq M_4. \quad (3.31)$$

Proof. We can deduce from (3.22), (3.23) that

$$\begin{aligned}
 & \delta_0 \int_{\Omega} |\beta_{\varepsilon}(\varphi_{\varepsilon}(s))| \, dx + \delta_0 \int_{\Gamma} |\beta_{\Gamma,\varepsilon}(\psi_{\varepsilon}(s))| \, dS \\
 & \leq c_1(|\Omega| + |\Gamma|) + (\mathbf{f}(s) - \boldsymbol{\pi}(\varphi_{\varepsilon}(s)) - \varepsilon \boldsymbol{\omega}'_{\varepsilon}(s), \boldsymbol{\omega}_{\varepsilon}(s))_{\mathcal{X}^2} \\
 & \quad - (\mathfrak{G}^L(\boldsymbol{\omega}'_{\varepsilon}(s)), \boldsymbol{\omega}_{\varepsilon}(s))_{\mathcal{X}^2} \\
 & \leq c_1(|\Omega| + |\Gamma|) + (\|\mathbf{f}(s)\|_{\mathcal{X}^2} + \|\boldsymbol{\pi}(\varphi_{\varepsilon}(s))\|_{\mathcal{X}^2} + \varepsilon \|\boldsymbol{\omega}'_{\varepsilon}(s)\|_{\mathcal{X}^2}) \|\boldsymbol{\omega}_{\varepsilon}(s)\|_{\mathcal{X}^2} \\
 & \quad + \|\boldsymbol{\omega}'_{\varepsilon}(s)\|_{0,*} \|\boldsymbol{\omega}_{\varepsilon}(s)\|_{0,*} \quad \text{for a.a. } s \in (0, T).
 \end{aligned} \tag{3.32}$$

Due to Lemma 3.4 and (A4), the right-hand side of (3.32) is uniformly bounded in $L^2(0, T)$. This yields the estimate (3.30). Next, by the definition of $m_{\varepsilon}(s)$ (recall (3.12)), we have

$$\begin{aligned}
 |m_{\varepsilon}(s)|^2 & \leq \frac{6}{(|\Omega| + |\Gamma|)^2} (\|\beta_{\varepsilon}(\varphi_{\varepsilon}(s))\|_{L^1(\Omega)}^2 + \|\pi(\varphi_{\varepsilon}(s))\|_{L^1(\Omega)}^2 + |\Omega| \|f(s)\|_H^2 \\
 & \quad + \|\beta_{\Gamma,\varepsilon}(\psi_{\varepsilon}(s))\|_{L^1(\Gamma)}^2 + \|\pi_{\Gamma}(\psi_{\varepsilon}(s))\|_{L^1(\Gamma)}^2 + |\Gamma| \|f_{\Gamma}(s)\|_{H_{\Gamma}}^2)
 \end{aligned}$$

for almost all $s \in (0, T)$. Integrating over $(0, T)$, using Lemma 3.4, (3.30) and (A3), (A4), we obtain the first estimate in (3.31). On the other hand, it follows from (3.11) that

$$\|\mathbf{P}\boldsymbol{\mu}_{\varepsilon}(s)\|_{\mathcal{H}_{L,0}^1} = \|\boldsymbol{\omega}'_{\varepsilon}(s)\|_{0,*}. \tag{3.33}$$

This fact combined with Poincaré’s inequality (2.6) yields

$$\begin{aligned}
 \|\boldsymbol{\mu}_{\varepsilon}(s)\|_{\mathcal{H}^1} & \leq \|\mathbf{P}\boldsymbol{\mu}_{\varepsilon}(s)\|_{\mathcal{H}^1} + \|m_{\varepsilon}(s)\mathbf{1}\|_{\mathcal{H}^1} \leq C \|\mathbf{P}\boldsymbol{\mu}_{\varepsilon}(s)\|_{\mathcal{H}_{L,0}^1} + \|m_{\varepsilon}(s)\mathbf{1}\|_{\mathcal{X}^2} \\
 & \leq C \|\boldsymbol{\omega}'_{\varepsilon}(s)\|_{0,*} + (|\Omega| + |\Gamma|)^{1/2} |m_{\varepsilon}(s)| \quad \text{for a.a. } s \in (0, T).
 \end{aligned}$$

Recalling (3.25), we obtain the second estimate in (3.31). ■

Lemma 3.6. *There exist positive constants M_5 , M_6 , and M_7 , independent of $\varepsilon \in (0, 1)$, such that*

$$\|\beta_{\varepsilon}(\varphi_{\varepsilon})\|_{L^2(0,T;H)} + \|\beta_{\varepsilon}(\psi_{\varepsilon})\|_{L^2(0,T;H_{\Gamma})} \leq M_5, \tag{3.34}$$

$$\|\beta_{\Gamma,\varepsilon}(\psi_{\varepsilon})\|_{L^2(0,T;H_{\Gamma})} \leq M_6, \tag{3.35}$$

$$\|\boldsymbol{\omega}_{\varepsilon}\|_{L^2(0,T;\mathcal{X}^2)} \leq M_7. \tag{3.36}$$

Proof. The proof of (3.34) follows the argument for [16, Lemma 4.4]; we sketch it here for completeness. Testing (3.15) by $\beta_{\varepsilon}(\varphi_{\varepsilon}) \in L^2(0, T; V)$, using (3.16) and noting that $(\beta_{\varepsilon}(\varphi_{\varepsilon}))|_{\Gamma} = \beta_{\varepsilon}(\psi_{\varepsilon}) \in L^2(0, T; V_{\Gamma})$, we find

$$\begin{aligned}
 & \int_{\Omega} \beta'_{\varepsilon}(\varphi_{\varepsilon}(s)) |\nabla \omega_{\varepsilon}(s)|^2 \, dx + \|\beta_{\varepsilon}(\varphi_{\varepsilon}(s))\|_H^2 \\
 & \quad + \delta \int_{\Gamma} \beta'_{\varepsilon}(\psi_{\varepsilon}(s)) |\nabla_{\Gamma} \omega_{\Gamma,\varepsilon}(s)|^2 \, dS + \int_{\Gamma} \beta_{\Gamma,\varepsilon}(\psi_{\varepsilon}(s)) \beta_{\varepsilon}(\psi_{\varepsilon}(s)) \, dS \\
 & = (f(s) + \mu_{\varepsilon}(s) - \varepsilon \partial_t \omega_{\varepsilon}(s) - \pi(\varphi_{\varepsilon}(s)), \beta_{\varepsilon}(\varphi_{\varepsilon}(s)))_H \\
 & \quad + (f_{\Gamma}(s) + \theta_{\varepsilon}(s) - \varepsilon \partial_t \omega_{\Gamma,\varepsilon}(s) - \pi_{\Gamma}(\psi_{\varepsilon}(s)), \beta_{\varepsilon}(\psi_{\varepsilon}(s)))_{H_{\Gamma}}
 \end{aligned} \tag{3.37}$$

for almost all $s \in (0, T)$. Recalling the compatibility condition (3.5) and noticing that $\beta_\varepsilon(r)$, $\beta_{\Gamma,\varepsilon}(r)$ have the same sign for all $r \in \mathbb{R}$, we get

$$\begin{aligned} \int_{\Gamma} \beta_{\Gamma,\varepsilon}(\psi_\varepsilon(s)) \beta_\varepsilon(\psi_\varepsilon(s)) \, dS &= \int_{\Gamma} |\beta_{\Gamma,\varepsilon}(\psi_\varepsilon(s))| |\beta_\varepsilon(\psi_\varepsilon(s))| \, dS \\ &\geq \frac{1}{\varrho} \int_{\Gamma} |\beta_\varepsilon(\psi_\varepsilon(s))|^2 \, dS - \frac{c_0}{\varrho} \int_{\Gamma} |\beta_\varepsilon(\psi_\varepsilon(s))| \, dS \\ &\geq \frac{1}{2\varrho} \int_{\Gamma} |\beta_\varepsilon(\psi_\varepsilon(s))|^2 \, dS - \frac{c_0^2}{2\varrho} |\Gamma|. \end{aligned}$$

On the other hand, we observe that

$$\int_{\Omega} \beta'_\varepsilon(\varphi_\varepsilon(s)) |\nabla \omega_\varepsilon(s)|^2 \, dx \geq 0, \quad \int_{\Gamma} \beta'_\varepsilon(\psi_\varepsilon(s)) |\nabla_{\Gamma} \omega_{\Gamma,\varepsilon}(s)|^2 \, dS \geq 0.$$

By Young's inequality and the Lipschitz continuity of π and π_{Γ} , we can find a positive constant \tilde{M}_5 , independent of $\varepsilon \in (0, 1)$, such that

$$\begin{aligned} &(f(s) + \mu_\varepsilon(s) - \varepsilon \partial_t \omega_\varepsilon(s) - \pi(\varphi_\varepsilon(s)), \beta_\varepsilon(\varphi_\varepsilon(s)))_H \\ &\leq \frac{1}{2} \|\beta_\varepsilon(\varphi_\varepsilon(s))\|_H^2 + \tilde{M}_5 (1 + \|f(s)\|_H^2 + \|\mu_\varepsilon(s)\|_H^2 + \|\omega_\varepsilon(s)\|_H^2 + \varepsilon \|\partial_t \omega_\varepsilon(s)\|_H^2) \end{aligned}$$

and

$$\begin{aligned} &(f_{\Gamma}(s) + \theta_\varepsilon(s) - \varepsilon \partial_t \omega_{\Gamma,\varepsilon}(s) - \pi_{\Gamma}(\psi_\varepsilon(s)), \beta_\varepsilon(\psi_\varepsilon(s)))_{H_{\Gamma}} \\ &\leq \frac{1}{4\varrho} \|\beta_\varepsilon(\psi_\varepsilon(s))\|_{H_{\Gamma}}^2 \\ &\quad + \varrho \tilde{M}_5 (1 + \|f_{\Gamma}(s)\|_{H_{\Gamma}}^2 + \|\theta_\varepsilon(s)\|_{H_{\Gamma}}^2 + \|\omega_{\Gamma,\varepsilon}(s)\|_{H_{\Gamma}}^2 + \varepsilon \|\partial_t \omega_{\Gamma,\varepsilon}(s)\|_{H_{\Gamma}}^2). \end{aligned}$$

Combining the above inequalities, integrating (3.37) in $(0, T)$ with respect to s , we infer from Lemmas 3.4, 3.5 that

$$\frac{1}{2} \int_0^T \|\beta_\varepsilon(\varphi_\varepsilon(s))\|_H^2 \, ds + \frac{1}{4\varrho} \int_0^T \|\beta_\varepsilon(\psi_\varepsilon(s))\|_{H_{\Gamma}}^2 \, ds \leq C, \quad (3.38)$$

where $C > 0$ is independent of $\varepsilon \in (0, 1)$. This yields the estimate (3.34).

By comparison in (3.15) and the estimates just proved, we have

$$\|\Delta \omega_\varepsilon\|_{L^2(0,T;H)} \leq C, \quad (3.39)$$

where $C > 0$ is independent of $\varepsilon \in (0, 1)$. Next, testing (3.16) by $\beta_{\Gamma,\varepsilon}(\psi_\varepsilon) \in L^2(0, T; V_{\Gamma})$, we can deduce that

$$\begin{aligned} &\delta \int_{\Gamma} \beta'_{\Gamma,\varepsilon}(\psi_\varepsilon(s)) |\nabla_{\Gamma} \omega_{\Gamma,\varepsilon}(s)|^2 \, dS + \|\beta_{\Gamma,\varepsilon}(\psi_\varepsilon(s))\|_{H_{\Gamma}}^2 \\ &= (f_{\Gamma}(s) + \theta_\varepsilon(s) - \varepsilon \partial_t \omega_{\Gamma,\varepsilon}(s) - \partial_{\mathbf{n}} \omega_\varepsilon(s) - \pi_{\Gamma}(\psi_\varepsilon(s)), \beta_{\Gamma,\varepsilon}(\psi_\varepsilon(s)))_{H_{\Gamma}} \\ &\leq \frac{1}{2} \|\beta_{\Gamma,\varepsilon}(\psi_\varepsilon(s))\|_{H_{\Gamma}}^2 + 3(\|f_{\Gamma}(s)\|_{H_{\Gamma}}^2 + \|\theta_\varepsilon(s)\|_{H_{\Gamma}}^2 \\ &\quad + \varepsilon \|\partial_t \omega_{\Gamma,\varepsilon}(s)\|_{H_{\Gamma}}^2 + \|\partial_{\mathbf{n}} \omega_\varepsilon(s)\|_{H_{\Gamma}}^2 + \|\pi_{\Gamma}(\psi_\varepsilon(s))\|_{H_{\Gamma}}^2) \end{aligned} \quad (3.40)$$

for almost all $s \in (0, T)$. Integrating (3.40) over $(0, T)$, we infer from Lemmas 3.4, 3.5 that

$$\|\beta_{\Gamma,\varepsilon}(\psi_\varepsilon)\|_{L^2(0,T;H_\Gamma)} \leq C(1 + \|\partial_n \omega_\varepsilon\|_{L^2(0,T;H_\Gamma)}). \tag{3.41}$$

By comparing the terms in (3.16), we infer that

$$\|\partial_n \omega_\varepsilon - \Delta_\Gamma \omega_{\Gamma,\varepsilon}\|_{L^2(0,T;H_\Gamma)} \leq C \left(1 + \frac{1}{\delta}\right) (1 + \|\partial_n \omega_\varepsilon\|_{L^2(0,T;H_\Gamma)}), \tag{3.42}$$

where the constant $C > 0$ is independent of $\varepsilon \in (0, 1)$. Combining (3.39) and (3.42), we can apply the elliptic estimate (2.10) (with $L = 0, k = 0$) to get

$$\|\omega_\varepsilon\|_{L^2(0,T;\mathcal{H}^2)} \leq C(1 + \|\partial_n \omega_\varepsilon\|_{L^2(0,T;H_\Gamma)}), \tag{3.43}$$

where the constant $C > 0$ is independent of ε but depends on δ . Thanks to the trace theorem and the Ehrling lemma (see Lemma A.2), we have

$$\begin{aligned} \|\partial_n \omega_\varepsilon\|_{L^2(0,T;H_\Gamma)} &\leq C \|\omega_\varepsilon\|_{L^2(0,T;H^{2-r}(\Omega))} \quad \text{for some } r \in (0, 1/2) \\ &\leq \epsilon \|\omega_\varepsilon\|_{L^2(0,T;H^2(\Omega))} + C_\epsilon \|\omega_\varepsilon\|_{L^2(0,T;V)}. \end{aligned}$$

Hence, taking $\epsilon > 0$ sufficiently small, we can conclude (3.36) from (3.43), with some constant $M_7 > 0$ independent of ε (but depends on δ). Finally, this estimate together with (3.41) easily yields the estimate (3.35). ■

3.3. Existence and uniqueness of weak solutions

Proof of Theorem 2.7. We are in a position to prove the existence of weak solutions to problem $(S_{L,\delta})$. This can be done by passing to the limit in the approximating problem as $\varepsilon \rightarrow 0$, applying a standard compactness argument as in [16]. Owing to the uniform estimates derived in Lemmas 3.2–3.6, there exist a subsequence of ε (not relabeled) and some limit functions $\omega = (\omega, \omega_\Gamma), \mu = (\mu, \theta), \xi = (\xi, \xi_\Gamma)$ and \tilde{m} such that

$$\omega_\varepsilon \rightharpoonup \omega \quad \text{weakly star in } L^\infty(0, T; \mathcal{V}_{(0)}^1), \tag{3.44}$$

$$\omega_\varepsilon \rightharpoonup \omega \quad \text{weakly in } H^1(0, T; \mathcal{H}_{(0)}^{-1}) \cap L^2(0, T; \mathcal{V}^2), \tag{3.45}$$

$$\varepsilon \omega_\varepsilon \rightarrow \mathbf{0} \quad \text{strongly in } H^1(0, T; \mathcal{L}^2), \tag{3.46}$$

$$\mu_\varepsilon \rightharpoonup \mu \quad \text{weakly in } L^2(0, T; \mathcal{H}^1), \tag{3.47}$$

$$\beta_\varepsilon(\varphi_\varepsilon) \rightarrow \xi \quad \text{weakly in } L^2(0, T; H), \tag{3.48}$$

$$\beta_{\Gamma,\varepsilon}(\psi_\varepsilon) \rightarrow \xi_\Gamma \quad \text{weakly in } L^2(0, T; H_\Gamma), \tag{3.49}$$

$$m_\varepsilon \rightharpoonup \tilde{m} \quad \text{weakly in } L^2(0, T). \tag{3.50}$$

From (3.45), due to the Aubin–Lions–Simon lemma (see Lemma A.1), we find

$$\omega_\varepsilon \rightarrow \omega \quad \text{strongly in } C([0, T]; \mathcal{L}_{(0)}^2) \cap L^2(0, T; \mathcal{V}_{(0)}^1), \tag{3.51}$$

which implies

$$\varphi_\varepsilon \rightarrow \varphi = \omega + \bar{m}_0 \mathbf{1} \quad \text{strongly in } C([0, T]; \mathcal{L}^2) \cap L^2(0, T; \mathcal{V}^1). \quad (3.52)$$

The above facts also imply that

$$\varphi|_{t=0} = \varphi_0 \quad \text{a.e. in } \Omega, \quad \psi|_{t=0} = \psi_0 \quad \text{a.e. on } \Gamma.$$

Moreover, (3.52) and the Lipschitz continuity of π , π_Γ ensure that

$$\pi(\varphi_\varepsilon) \rightarrow \pi(\varphi) \quad \text{strongly in } C([0, T]; \mathcal{L}^2),$$

and $\tilde{m} = \bar{m}(\xi + \pi(\varphi) - f)$ with $\xi = (\xi, \xi_\Gamma)$. Due to the maximal monotonicity of β and β_Γ , by applying [4, Proposition 2.2] and (3.48), (3.49), (3.52), we obtain

$$\xi \in \beta(\varphi) \quad \text{a.e. in } Q, \quad \xi_\Gamma \in \beta_\Gamma(\psi) \quad \text{a.e. on } \Sigma. \quad (3.53)$$

Now, we are able to pass to the limit in the weak formulations (3.13), (3.14) to recover (2.26) and

$$(\mu(t), z)_{\mathcal{L}^2} = b_\delta(\varphi(t), z) + (\xi(t) + \pi(\varphi(t)) - f(t), z)_{\mathcal{L}^2} \quad \forall z \in \mathcal{V}^1 \quad (3.54)$$

for almost all $t \in (0, T)$. Since $z \in \mathcal{V}^1$ is arbitrary, we can conclude that the limit triplet (φ, μ, ξ) is indeed a weak solution to problem $(S_{L,\delta})$.

Finally, we establish the additional regularity (2.29) under the assumption (A1)'. The estimate for nonlinearities (β, β_Γ) is similar to [28] for the case $L = 0$ and we just sketch it. As in [36], for each $k \in \mathbb{N} \setminus \{1\}$, we introduce the Lipschitz continuous function $h_k : \mathbb{R} \rightarrow \mathbb{R}$ by

$$h_k(r) := \begin{cases} -1 + \frac{1}{k} & \text{if } r < -1 + \frac{1}{k}, \\ r & \text{if } -1 + \frac{1}{k} \leq r \leq 1 - \frac{1}{k}, \\ 1 - \frac{1}{k} & \text{if } r > 1 - \frac{1}{k}. \end{cases}$$

We define $\varphi_k := h_k \circ \varphi$, $\psi_k := h_k \circ \psi$ and denote $\varphi_k := (\varphi_k, \psi_k)$ for all $k \in \mathbb{N} \setminus \{1\}$. It follows that $\varphi_k \in L^\infty(0, T; \mathcal{V}^1)$ and

$$\nabla \varphi_k = \nabla \varphi \chi_{[-1+\frac{1}{k}, 1-\frac{1}{k}]}, \quad \nabla_\Gamma \psi_k = \nabla_\Gamma \psi \chi_{[-1+\frac{1}{k}, 1-\frac{1}{k}]},$$

where the function $\chi_{[-1+\frac{1}{k}, 1-\frac{1}{k}]}$ is the characteristic function of interval $[-1 + \frac{1}{k}, 1 - \frac{1}{k}]$. For $k \in \mathbb{N} \setminus \{1\}$ and $p \geq 2$, we see that

$$\beta_k := |\beta(\varphi_k)|^{p-2} \beta(\varphi_k) \in L^\infty(0, T; V), \quad \beta_{\Gamma,k} := |\beta_\Gamma(\psi_k)|^{p-2} \beta_\Gamma(\psi_k) \in L^\infty(0, T; V_\Gamma)$$

are well defined, and

$$\nabla \beta_k = (p-1)|\beta(\varphi_k)|^{p-2} \beta'(\varphi_k) \nabla \varphi_k, \quad \nabla_\Gamma \beta_{\Gamma,k} = (p-1)|\beta_\Gamma(\psi_k)|^{p-2} \beta'_\Gamma(\psi_k) \nabla_\Gamma \psi_k.$$

Besides, $\beta_k|_\Gamma = |\beta(\psi_k)|^{p-2}\beta(\psi_k) \in L^\infty(0, T; V_\Gamma)$. Define

$$\tilde{\mu} := \mu - \pi(\varphi) - f \quad \text{and} \quad \tilde{\theta} := \theta - \pi_\Gamma(\psi) - f_\Gamma.$$

Then, it holds $(\tilde{\mu}, \tilde{\theta}) \in L^2(0, T; \mathcal{L}^p)$, where $p \in [2, 6]$ if $d = 3$ and $p \in [2, +\infty)$ if $d = 2$. By a similar process as in [28], we can deduce that

$$\begin{aligned} & \|\beta(\varphi_k)\|_{L^2(0,T;L^p(\Omega))} + \|\beta_\Gamma(\psi_k)\|_{L^2(0,T;L^p(\Gamma))} \\ & \leq C_p(\|\tilde{\mu}\|_{L^2(0,T;L^p(\Omega))} + \|\tilde{\theta}\|_{L^2(0,T;L^p(\Gamma))} + 1). \end{aligned}$$

Passing to the limit as $k \rightarrow +\infty$, owing to Fatou’s lemma, we conclude that

$$(\beta(\varphi), \beta_\Gamma(\psi)) \in L^2(0, T; \mathcal{L}^p), \quad \begin{cases} p \in [2, 6] & \text{if } d = 3, \\ p \in [2, +\infty) & \text{if } d = 2. \end{cases} \quad (3.55)$$

Next, since φ satisfies the following bulk-surface elliptic problem

$$\begin{cases} -\Delta\varphi = \tilde{\mu} - \beta(\varphi), & \text{a.e. in } \Omega, \\ \varphi = \psi, & \text{a.e. on } \Gamma, \\ -\Delta_\Gamma\psi + \partial_n\varphi = \tilde{\theta} - \beta_\Gamma(\psi), & \text{a.e. on } \Gamma, \end{cases}$$

applying (3.55) and the L^p -regularity theory for the bulk-surface elliptic system (see [45, Proposition A.1]), we find

$$\varphi \in L^2(0, T; \mathcal{W}^{2,p}), \quad \begin{cases} p \in [2, 6] & \text{if } d = 3, \\ p \in [2, +\infty) & \text{if } d = 2. \end{cases} \quad (3.56)$$

The case $p \in [1, 2)$ follows from (3.55), (3.56) and Hölder’s inequality.

Hence, we finish the proof of Theorem 2.7. ■

Proof of Theorem 2.9. In what follows, we prove the continuous dependence on the initial data and the force terms. This can be done by the standard energy method. For $i \in \{1, 2\}$, let $(\varphi_i, \mu_i, \xi_i)$ be a weak solution to problem $(S_{L,\delta})$ corresponding to the data $(\varphi_{0,i}, f_i)$ with

$$\bar{m}(\varphi_{0,1}) = \bar{m}(\varphi_{0,2}) = \bar{m}_0.$$

Set $\omega_i := \varphi_i - \bar{m}_0\mathbf{1}$. We consider the difference between (3.7), at the time $s \in (0, T)$, for $\omega_1(s) = (\omega_1(s), \omega_{\Gamma,1}(s))$ and $\omega_2(s) = (\omega_2(s), \omega_{\Gamma,2}(s))$, that is,

$$\begin{aligned} & \mathfrak{E}^L(\omega'_1(s) - \omega'_2(s)) + \partial\Phi_\delta(\omega_1(s) - \omega_2(s)) \\ & = \mathbf{P}(-\xi_1(s) + \xi_2(s)) \\ & \quad + \mathbf{P}(-\pi(\omega_1(s) + \bar{m}_0\mathbf{1}) + \pi(\omega_2(s) + \bar{m}_0\mathbf{1})) + \mathbf{P}(f_1(s) - f_2(s)). \end{aligned}$$

Taking \mathcal{L}^2 inner product with $\omega_1(s) - \omega_2(s)$, by the Lipschitz continuity of π , π_Γ , we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{ds} \|\omega_1(s) - \omega_2(s)\|_{0,*}^2 + \|\omega_1(s) - \omega_2(s)\|_{\mathcal{V}_{(0)}^1}^2 \\ & \quad + (\xi_1(s) - \xi_2(s), \omega_1(s) - \omega_2(s))_{\mathcal{X}^2} \\ & = -(\pi(\omega_1(s) + \bar{m}_0 \mathbf{1}) - \pi(\omega_2(s) + \bar{m}_0 \mathbf{1}), \omega_1(s) - \omega_2(s))_{\mathcal{X}^2} \\ & \quad + (f_1(s) - f_2(s), \omega_1(s) - \omega_2(s))_{\mathcal{X}^2} \\ & \leq (K + K_\Gamma) \|\omega_1(s) - \omega_2(s)\|_{\mathcal{X}^2}^2 \\ & \quad + \|f_1(s) - f_2(s)\|_{(\mathcal{V}^1)'} \|\omega_1(s) - \omega_2(s)\|_{\mathcal{V}^1} \end{aligned} \tag{3.57}$$

for almost all $s \in (0, T)$. Since $\omega_1(s) - \omega_2(s) \in \mathcal{V}_{(0)}^1$ for almost all $s \in (0, T)$, by the Ehrling lemma (see Lemma A.2) and Poincaré’s inequality (2.6), we get

$$\|\omega_1(s) - \omega_2(s)\|_{\mathcal{X}^2}^2 \leq \epsilon \|\omega_1(s) - \omega_2(s)\|_{\mathcal{V}_{(0)}^1}^2 + C_\epsilon \|\omega_1(s) - \omega_2(s)\|_{0,*}^2$$

for any $\epsilon > 0$. Besides, it follows from (3.53) and the monotonicity of β , β_Γ that

$$\begin{aligned} & (\xi_1(s) - \xi_2(s), \omega_1(s) - \omega_2(s))_{\mathcal{X}^2} \\ & = (\xi_1(s) - \xi_2(s), \varphi_1(s) - \varphi_2(s))_{\mathcal{X}^2} \geq 0 \quad \text{for a.a. } s \in (0, T). \end{aligned}$$

Inserting the above inequalities into (3.57), taking $\epsilon > 0$ sufficiently small and applying Young’s inequality, we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{ds} \|\omega_1(s) - \omega_2(s)\|_{0,*}^2 + \|\omega_1(s) - \omega_2(s)\|_{\mathcal{V}_{(0)}^1}^2 \\ & \leq \frac{1}{2} \|\omega_1(s) - \omega_2(s)\|_{\mathcal{V}_{(0)}^1}^2 + C \|\omega_1(s) - \omega_2(s)\|_{0,*}^2 + C \|f_1(s) - f_2(s)\|_{(\mathcal{V}^1)'}^2 \end{aligned}$$

for almost all $s \in (0, T)$. Then, by Grönwall’s lemma, we arrive at the conclusion (2.30). ■

Thanks to the uniqueness of the phase function φ associated with problem $(S_{L,\delta})$, we are able to provide an estimate on the convergence rate for $\varphi_\epsilon \rightarrow \varphi$ in terms of the parameter ϵ of the Yosida approximation.

Proposition 3.7 (Convergence rate). *Suppose that the assumptions of Theorem 2.7 are satisfied. Let φ_ϵ be the unique solution to the approximating problem (3.8)–(3.9) and φ the unique weak solution to problem $(S_{L,\delta})$. Then, we have*

$$\|\varphi_\epsilon - \varphi\|_{L^\infty(0,T;(\mathcal{X}^1)')} + \|\varphi_\epsilon - \varphi\|_{L^2(0,T;\mathcal{V}^1)} \leq C \sqrt{\epsilon} \quad \forall \epsilon \in (0, 1),$$

where the constant $C > 0$ depends on $\hat{\pi}$, $\hat{\pi}_\Gamma$, $\hat{\beta}$, $\hat{\beta}_\Gamma$, and the \mathcal{V}^1 -norm of the initial data φ_0 , but not on ϵ .

Proof. For any $\varepsilon_1, \varepsilon_2 \in (0, 1)$, let ω_{ε_1} and ω_{ε_2} be the corresponding solutions to the approximating problem (3.8)–(3.9), respectively. Recalling the relation

$$\varphi_{\varepsilon_i} = \omega_{\varepsilon_i} + \bar{m}_0 \mathbf{1}, \quad i \in \{1, 2\},$$

we consider their difference $\varphi_* = \varphi_{\varepsilon_1} - \varphi_{\varepsilon_2}$. It follows from (3.8) that

$$\begin{aligned} & \mathfrak{C}^L(\varphi'_*) + \partial\Phi_\delta(\varphi_*) \\ &= -\varepsilon_1 \varphi'_{\varepsilon_1} + \varepsilon_2 \varphi'_{\varepsilon_2} + \mathbf{P}(-\beta_{\varepsilon_1}(\varphi_{\varepsilon_1}) + \beta_{\varepsilon_2}(\varphi_{\varepsilon_2})) \\ & \quad + \mathbf{P}(-\pi(\varphi_{\varepsilon_1}) + \pi(\varphi_{\varepsilon_2})) \quad \text{in } \mathcal{L}^2_{(0)}. \end{aligned} \tag{3.58}$$

Testing (3.58) by φ_* and integrating in $[0, t] \subset [0, T]$, we obtain

$$\begin{aligned} & \frac{1}{2} \|\varphi_*(t)\|_{0,*}^2 + \int_0^t \|\varphi_*(s)\|_{\mathcal{V}^1_{(0)}}^2 \, ds \\ &= \int_0^t (-\varepsilon_1 \varphi'_{\varepsilon_1}(s) + \varepsilon_2 \varphi'_{\varepsilon_2}(s), \varphi_*(s))_{\mathcal{X}^2} \, ds \\ & \quad + \int_0^t (-\beta_{\varepsilon_1}(\varphi_{\varepsilon_1}(s)) + \beta_{\varepsilon_2}(\varphi_{\varepsilon_2}(s)), \varphi_*(s))_{\mathcal{X}^2} \, ds \\ & \quad + \int_0^t (-\pi(\varphi_{\varepsilon_1}(s)) + \pi(\varphi_{\varepsilon_2}(s)), \varphi_*(s))_{\mathcal{X}^2} \, ds \\ &=: J_1 + J_2 + J_3. \end{aligned} \tag{3.59}$$

Using Lemma 3.4, we obtain that

$$\begin{aligned} J_1 &= \int_0^t (-\varepsilon_1 \mathfrak{C}^L(\varphi'_{\varepsilon_1}(s)) + \varepsilon_2 \mathfrak{C}^L(\varphi'_{\varepsilon_2}(s)), \varphi_*(s))_{\mathcal{H}^1_{L,0}} \, ds \\ &\leq C \int_0^t (\varepsilon_1 \|\varphi'_{\varepsilon_1}(s)\|_{0,*} + \varepsilon_2 \|\varphi'_{\varepsilon_2}(s)\|_{0,*}) \|\varphi_*(s)\|_{\mathcal{V}^1} \, ds \\ &\leq C(\varepsilon_1 + \varepsilon_2). \end{aligned} \tag{3.60}$$

Concerning J_2 , we apply the argument in [10]. It follows from the definition of the Yosida approximation that $J_\varepsilon \varphi_\varepsilon = \varphi_\varepsilon - \varepsilon \beta_\varepsilon(\varphi_\varepsilon)$. From this, we find

$$\begin{aligned} & - \int_0^t \int_\Omega (\beta_{\varepsilon_1}(\varphi_{\varepsilon_1}) - \beta_{\varepsilon_2}(\varphi_{\varepsilon_2})) \varphi_* \, dx \, ds \\ &= - \int_0^t \int_\Omega (\beta_{\varepsilon_1}(\varphi_{\varepsilon_1}) - \beta_{\varepsilon_2}(\varphi_{\varepsilon_2})) (J_{\varepsilon_1}(\varphi_{\varepsilon_1}) - J_{\varepsilon_2}(\varphi_{\varepsilon_2})) \, dx \, ds \\ & \quad - \int_0^t \int_\Omega (\beta_{\varepsilon_1}(\varphi_{\varepsilon_1}) - \beta_{\varepsilon_2}(\varphi_{\varepsilon_2})) (\varepsilon_1 \beta_{\varepsilon_1}(\varphi_{\varepsilon_1}) - \varepsilon_2 \beta_{\varepsilon_2}(\varphi_{\varepsilon_2})) \, dx \, ds. \end{aligned} \tag{3.61}$$

The first term on the right-hand side of (3.61) is nonpositive by the monotonicity of β and the fact that $\beta_\varepsilon(\varphi_\varepsilon) \in \beta(J_\varepsilon(\varphi_\varepsilon))$ (cf., [39, (2.7)]). On the other hand, the second term on

the right-hand side of (3.61) can be estimated by Hölder’s inequality and (3.34), that is,

$$\begin{aligned}
 & - \int_0^t \int_{\Omega} (\beta_{\varepsilon_1}(\varphi_{\varepsilon_1}) - \beta_{\varepsilon_2}(\varphi_{\varepsilon_2}))(\varepsilon_1 \beta_{\varepsilon_1}(\varphi_{\varepsilon_1}) - \varepsilon_2 \beta_{\varepsilon_2}(\varphi_{\varepsilon_2})) \, dx \, ds \\
 & = -\varepsilon_1 \|\beta_{\varepsilon_1}(\varphi_{\varepsilon_1})\|_{L^2(0,T;H)}^2 - \varepsilon_2 \|\beta_{\varepsilon_2}(\varphi_{\varepsilon_2})\|_{L^2(0,T;H)}^2 \\
 & \quad + (\varepsilon_1 + \varepsilon_2) \int_0^t \int_{\Omega} \beta_{\varepsilon_1}(\varphi_{\varepsilon_1}) \beta_{\varepsilon_2}(\varphi_{\varepsilon_2}) \, dx \, ds \\
 & \leq (\varepsilon_1 + \varepsilon_2) \|\beta_{\varepsilon_1}(\varphi_{\varepsilon_1})\|_{L^2(0,T;H)} \|\beta_{\varepsilon_2}(\varphi_{\varepsilon_2})\|_{L^2(0,T;H)} \\
 & \leq C(\varepsilon_1 + \varepsilon_2).
 \end{aligned}$$

Similarly, by the monotonicity of β_{Γ} and (3.35), we can obtain the following estimate for boundary potentials:

$$- \int_0^t \int_{\Gamma} (\beta_{\Gamma,\varepsilon_1}(\psi_{\varepsilon_1}) - \beta_{\Gamma,\varepsilon_2}(\psi_{\varepsilon_2})) \psi_* \, dS \, ds \leq C(\varepsilon_1 + \varepsilon_2).$$

The above estimates provide the control of J_2 :

$$J_2 \leq C(\varepsilon_1 + \varepsilon_2). \tag{3.62}$$

To handle J_3 , using (A3) and the Ehrling lemma (see Lemma A.2), we find

$$\begin{aligned}
 J_3 & \leq \max\{K, K_{\Gamma}\} \int_0^t \|\varphi_*(s)\|_{\mathcal{L}^2}^2 \, ds \\
 & \leq \max\{K, K_{\Gamma}\} \epsilon \int_0^t \|\varphi_*(s)\|_{\mathcal{V}_1^1}^2 \, ds + \max\{K, K_{\Gamma}\} C_{\epsilon} \int_0^t \|\varphi_*(s)\|_{0,*}^2 \, ds. \tag{3.63}
 \end{aligned}$$

Combining (3.59), (3.60), (3.62), (3.63), and taking $\epsilon > 0$ sufficiently small, we obtain

$$\|\varphi_*(t)\|_{0,*}^2 + \int_0^t \|\varphi_*(s)\|_{\mathcal{V}_1^1}^2 \, ds \leq C(\varepsilon_1 + \varepsilon_2) + C \int_0^t \|\varphi_*(s)\|_{0,*}^2 \, ds \quad \forall t \in [0, T].$$

Then, we deduce from Grönwall’s lemma that

$$\|\varphi_*\|_{L^{\infty}(0,T;\mathcal{H}_0^{-1})}^2 + \|\varphi_*\|_{L^2(0,T;\mathcal{V}_1^1)}^2 \leq C(\varepsilon_1 + \varepsilon_2), \tag{3.64}$$

where the constant $C > 0$ is independent of ε_1 and ε_2 .

The estimate (3.64) implies that

$$\{\varphi_{\varepsilon}\} \text{ is a Cauchy sequence in } L^{\infty}(0, T; \mathcal{H}_0^{-1}) \cap L^2(0, T; \mathcal{V}^1)$$

as $\varepsilon \rightarrow 0$. Denote its limit by φ . Recalling Theorems 2.7, 2.9, we find $\varphi_{\varepsilon} \rightarrow \varphi$ strongly in $L^2(0, T; \mathcal{V}^1) \cap C([0, T]; \mathcal{L}^2)$ and φ is the unique weak solution to problem $(S_{L,\delta})$. Hence, we can pass to the limit $\varepsilon_2 \rightarrow 0$ in (3.64) and obtain

$$\|\varphi_{\varepsilon_1} - \varphi\|_{L^{\infty}(0,T;(\mathcal{H}^1)')}^2 + \|\varphi_{\varepsilon_1} - \varphi\|_{L^2(0,T;\mathcal{V}^1)}^2 \leq C \varepsilon_1,$$

which completes the proof. ■

Remark 3.8. In the recent work [39], the authors studied the nonlocal Cahn–Hilliard equation

$$\begin{aligned} \partial_t \varphi &= \Delta \mu && \text{in } Q, \\ \mu &= B(\varphi) + \beta(\varphi) + \pi(\varphi) && \text{in } Q, \\ \partial_{\mathbf{n}} \mu &= 0 && \text{on } \Sigma, \end{aligned}$$

with $B\varphi(x) := \int_{\Omega} J(x - y)(\varphi(x) - \varphi(y)) \, dy$ and $J \in L^2_{\text{loc}}(\mathbb{R}^d)$. Assuming some additional assumptions on the nonlinearity π and applying the theory of Hilbert–Schmidt operator for the nonlocal term, they proved the convergence rate

$$\|\varphi_{\varepsilon} - \varphi\|_{L^2(0,T;H)} \leq C \sqrt{\varepsilon},$$

where ε is the parameter of the Yosida approximation.

3.4. Existence and uniqueness of strong solutions

Proof of Theorem 2.11. Now, let us consider strong solutions to problem $(S_{L,\delta})$. The uniqueness is guaranteed by Theorem 2.9. As in [16], we can prove the existence by deriving some uniform estimates for higher-order norms of the approximating solutions. For any $h \in (0, 1)$ sufficiently small, we consider the difference between (3.8), at time s and $s + h$, that is,

$$\begin{aligned} &\varepsilon(\omega'_\varepsilon(s + h) - \omega'_\varepsilon(s)) + \mathfrak{S}^L(\omega'_\varepsilon(s + h) - \omega'_\varepsilon(s)) + \partial \Phi_\delta(\omega_\varepsilon(s + h) - \omega_\varepsilon(s)) \\ &= \mathbf{P}(-\beta_\varepsilon(\omega_\varepsilon(s + h) + \bar{m}_0 \mathbf{1}) + \beta_\varepsilon(\omega_\varepsilon(s) + \bar{m}_0 \mathbf{1})) \\ &\quad + \mathbf{P}(-\pi(\omega_\varepsilon(s + h) + \bar{m}_0 \mathbf{1}) + \pi(\omega_\varepsilon(s) + \bar{m}_0 \mathbf{1})) \\ &\quad + \mathbf{P}(f(s + h) - f(s)) \end{aligned}$$

for almost all $s \in (0, T - h]$. Taking \mathcal{L}^2 inner product with $\omega_\varepsilon(s + h) - \omega_\varepsilon(s)$, then integrating from 0 to t with respect to s , and dividing the resultant by h^2 , using the monotonicity of β_ε and $\beta_{\Gamma,\varepsilon}$, we obtain

$$\begin{aligned} &\frac{\varepsilon}{2} \left\| \frac{\omega_\varepsilon(t + h) - \omega_\varepsilon(t)}{h} \right\|_{\mathcal{X}^2}^2 + \frac{1}{2} \left\| \frac{\omega_\varepsilon(t + h) - \omega_\varepsilon(t)}{h} \right\|_{0,*}^2 \\ &\quad + \frac{1}{2} \int_0^t \left\| \frac{\omega_\varepsilon(s + h) - \omega_\varepsilon(s)}{h} \right\|_{\mathbf{V}^1_{(0)}}^2 \, ds \\ &\leq \frac{\varepsilon}{2} \left\| \frac{\omega_\varepsilon(h) - \omega_0}{h} \right\|_{\mathcal{X}^2}^2 + \frac{1}{2} \left\| \frac{\omega_\varepsilon(h) - \omega_0}{h} \right\|_{0,*}^2 \\ &\quad + \left(K + K_\Gamma + \frac{1}{2} \right) \int_0^t \left\| \frac{\omega_\varepsilon(s + h) - \omega_\varepsilon(s)}{h} \right\|_{\mathcal{X}^2}^2 \, ds \\ &\quad + \frac{1}{2} \int_0^t \left\| \frac{f(s + h) - f(s)}{h} \right\|_{\mathcal{X}^2}^2 \, ds \end{aligned}$$

$$\begin{aligned} &\leq \frac{\varepsilon}{2} \left\| \frac{\omega_\varepsilon(h) - \omega_0}{h} \right\|_{\mathcal{L}^2}^2 + \frac{1}{2} \left\| \frac{\omega_\varepsilon(h) - \omega_0}{h} \right\|_{0,*}^2 + \frac{1}{4} \int_0^t \left\| \frac{\omega_\varepsilon(s+h) - \omega_\varepsilon(s)}{h} \right\|_{\mathcal{V}_{(0)}^1}^2 ds \\ &\quad + C \int_0^t \left\| \frac{\omega_\varepsilon(s+h) - \omega_\varepsilon(s)}{h} \right\|_{0,*}^2 ds + \frac{1}{2} \int_0^t \left\| \frac{f(s+h) - f(s)}{h} \right\|_{\mathcal{L}^2}^2 ds. \end{aligned}$$

The first two terms on the right-hand side can be controlled as in [16, (4.35)]. Integrating (3.8) from 0 to h , taking the inner product with $(\omega_\varepsilon(h) - \omega_0)/h^2$, we obtain

$$\begin{aligned} &\frac{\varepsilon}{2} \left\| \frac{\omega_\varepsilon(h) - \omega_0}{h} \right\|_{\mathcal{L}^2}^2 + \frac{1}{2} \left\| \frac{\omega_\varepsilon(h) - \omega_0}{h} \right\|_{0,*}^2 \\ &\leq - \left(\frac{\omega_\varepsilon(h) - \omega_0}{h}, \frac{1}{h} \int_0^h \mathbf{P}(\partial\Phi_\delta(\varphi_\varepsilon(s)) + \beta_\varepsilon(\varphi_\varepsilon(s)) + \pi(\varphi_\varepsilon(s)) - f(s)) ds \right)_{\mathcal{L}^2} \\ &= -a_L \left(\mathfrak{C}^L \left(\frac{\omega_\varepsilon(h) - \omega_0}{h} \right), \frac{1}{h} \int_0^h \mathbf{P}(\partial\Phi_\delta(\varphi_\varepsilon(s)) + \beta_\varepsilon(\varphi_\varepsilon(s)) + \pi(\varphi_\varepsilon(s)) - f(s)) ds \right) \\ &\leq \frac{1}{4} \left\| \frac{\omega_\varepsilon(h) - \omega_0}{h} \right\|_{0,*}^2 \\ &\quad + C \left\| \frac{1}{h} \int_0^h \mathbf{P}(\partial\Phi_\delta(\varphi_\varepsilon(s)) + \beta_\varepsilon(\varphi_\varepsilon(s)) + \pi(\varphi_\varepsilon(s)) - f(s)) ds \right\|_{\mathcal{H}^1}^2. \end{aligned} \tag{3.65}$$

Thanks to the additional assumptions (A6), (A7), the last term on the right-hand side of (3.65) is bounded for all $\varepsilon \in (0, \varepsilon_0]$. Then, by Grönwall’s lemma, we obtain that

$$t \mapsto \frac{\omega_\varepsilon(t+h) - \omega_\varepsilon(t)}{h} \text{ are bounded in } L^\infty(0, T-h; \mathcal{H}_{(0)}^{-1}) \cap L^2(0, T-h; \mathcal{V}_{(0)}^1),$$

and

$$t \mapsto \varepsilon^{1/2} \frac{\omega_\varepsilon(t+h) - \omega_\varepsilon(t)}{h} \text{ are bounded in } L^\infty(0, T-h; \mathcal{L}^2),$$

uniformly with respect to $\varepsilon \in (0, \varepsilon_0]$. Passing to the limit as $h \rightarrow 0$, we obtain

$$\varepsilon^{1/2} \|\omega'_\varepsilon\|_{L^\infty(0,T;\mathcal{L}^2)} + \|\omega'_\varepsilon\|_{L^\infty(0,T;\mathcal{H}_{(0)}^{-1})} + \|\omega'_\varepsilon\|_{L^2(0,T;\mathcal{V}^1)} \leq M_8, \tag{3.66}$$

where the constant $M_8 > 0$ is independent of $\varepsilon \in (0, \varepsilon_0]$.

Keeping the improved estimate (3.66) and the assumption (A7) in mind, arguing as in Lemmas 3.5–3.6, without integration in time over $(0, T)$ but taking the L^∞ -norm on $(0, T)$, we obtain

$$\begin{aligned} &\|m_\varepsilon\|_{L^\infty(0,T)} + \|\mu_\varepsilon\|_{L^\infty(0,T;\mathcal{H}^1)} \leq M_9, \\ &\|\beta_\varepsilon(\varphi_\varepsilon)\|_{L^\infty(0,T;H)} + \|\beta_{\Gamma,\varepsilon}(\psi_\varepsilon)\|_{L^\infty(0,T;H_\Gamma)} \leq M_9, \\ &\|\omega_\varepsilon\|_{L^\infty(0,T;\mathcal{H}^2)} \leq M_9, \end{aligned}$$

where the constant $M_9 > 0$ is independent of $\varepsilon \in (0, \varepsilon_0]$. Passing to the limit as $\varepsilon \rightarrow 0$, we

obtain the following additional regularities for the solution (ω, μ, ξ) :

$$\begin{aligned} \omega &\in W^{1,\infty}(0, T; \mathcal{H}_{(0)}^{-1}) \cap H^1(0, T; \mathcal{V}_{(0)}^1) \cap L^\infty(0, T; \mathcal{H}^2), \\ \mu &\in L^\infty(0, T; \mathcal{H}^1), \quad \xi \in L^\infty(0, T; \mathcal{L}^2). \end{aligned}$$

Since $\omega' \in L^2(0, T; \mathcal{V}_{(0)}^1)$ and $\mathbf{P}\mu = \mathcal{C}^L(-\omega')$, we can infer from the elliptic estimate (2.10) that $\mu \in L^2(0, T; \mathcal{H}^2)$.

Applying the above regularity properties and integration by parts, the weak formulation (2.26) can be rewritten as

$$\begin{aligned} 0 &= \int_{\Omega} (\partial_t \omega(t) - \Delta \mu(t))y \, dx + \int_{\Gamma} (\partial_t \omega_{\Gamma}(t) - \Delta_{\Gamma} \theta(t) + \partial_{\mathbf{n}} \mu(t))y_{\Gamma} \, dS \\ &\quad + \frac{1}{L} \int_{\Gamma} (\theta(t) - \mu(t) - L \partial_{\mathbf{n}} \mu(t))(y_{\Gamma} - y) \, dS \quad \forall \mathbf{y} \in \mathcal{H}^1, \text{ a.a. } t \in (0, T). \end{aligned} \tag{3.67}$$

Since $\mathbf{y} \in \mathcal{H}^1$ is arbitrary, it easily follows from (3.67) that

$$\begin{aligned} \partial_t \omega &= \Delta \mu, & \text{a.e. in } Q, \\ \partial_t \omega_{\Gamma} &= \Delta_{\Gamma} \theta - \partial_{\mathbf{n}} \mu, & \text{a.e. on } \Sigma, \\ L \partial_{\mathbf{n}} \mu &= \theta - \mu, & \text{a.e. on } \Sigma. \end{aligned}$$

In summary, (φ, μ, ξ) is a strong solution to problem $(S_{L,\delta})$ in the sense of Definition 2.10. The proof is complete. ■

4. Asymptotic limit as $\delta \rightarrow 0$ and well-posedness without surface diffusion

In this section, we investigate the asymptotic behavior of weak solutions to problem $(S_{L,\delta})$ as $\delta \rightarrow 0$, with $L \in (0, +\infty)$ being fixed. Our goal is to show that the limit problem $(S_{L,0})$ without surface diffusion is still well posed.

4.1. Uniform estimates

Let the assumptions of Theorem 2.15 be satisfied. For any $\delta, \varepsilon \in (0, 1)$, we consider the approximating problem (3.8)–(3.9) with data $\{\varphi_0^\delta, \mathbf{f}^\delta\}_{\delta \in (0,1)}$. Let $\omega_\varepsilon^\delta$ be the unique solution given by Proposition 3.1. Besides, we set

$$\varphi_\varepsilon^\delta = \omega_\varepsilon^\delta + \bar{m}_0^\delta \mathbf{1}, \quad \mu_\varepsilon^\delta := \varepsilon \partial_t \omega_\varepsilon^\delta + \partial \Phi_\delta(\omega_\varepsilon^\delta) + \beta_\varepsilon(\varphi_\varepsilon^\delta) + \pi(\varphi_\varepsilon^\delta) - \mathbf{f}^\delta,$$

where $\bar{m}_0^\delta = \bar{m}(\varphi_0^\delta)$. Then, $(\varphi_\varepsilon^\delta, \mu_\varepsilon^\delta)$ is also uniquely determined and satisfies (3.13)–(3.19) with $\{\varphi_0^\delta, \mathbf{f}^\delta\}_{\delta \in (0,1)}$.

As in the proof of Theorem 2.7, when $\varepsilon \rightarrow 0$, $(\varphi_\varepsilon^\delta, \mu_\varepsilon^\delta, \beta_\varepsilon(\varphi_\varepsilon^\delta))$ converges (up to a subsequence) to some limit triplet $(\varphi^\delta, \mu^\delta, \xi^\delta)$, which is a weak solution to problem $(S_{L,\delta})$

with data $\{\varphi_0^\delta, f^\delta\}_{\delta \in (0,1)}$. Since the uniqueness of μ^δ, ξ^δ is not clear (cf., Theorem 2.9), the related convergence should always be understood in the sense of a suitable subsequence. Below we will not relabel the convergent subsequence for the sake of simplicity.

We now derive uniform estimates of the solutions $(\varphi^\delta, \mu^\delta, \xi^\delta)$ with respect to the parameter $\delta \in (0, 1)$.

Lemma 4.1. *There exists $\tilde{\delta} \in (0, 1)$ such that*

$$\sup_{t \in [0, T]} |\bar{m}(\varphi^\delta(t))| \leq M_{10} \quad \forall \delta \in (0, \tilde{\delta}], \quad (4.1)$$

where $M_{10} > 0$ is independent of $\delta \in (0, \tilde{\delta}]$.

Proof. Recalling the mass conservation property (3.20), we have $\bar{m}(\varphi_\varepsilon^\delta(t)) = \bar{m}(\varphi_0^\delta)$ for $t \in [0, T]$. This together with (3.52) implies

$$\bar{m}(\varphi^\delta(t)) = \bar{m}(\varphi_0^\delta) \quad \forall t \in [0, T]. \quad (4.2)$$

On the other hand, it follows from (2.39) that $\lim_{\delta \rightarrow 0} \bar{m}(\varphi_0^\delta) = \bar{m}(\varphi_0) \in \text{Int } D(\beta_\Gamma)$. Then, the conclusion (4.1) easily follows. \blacksquare

Lemma 4.2. *There exists a positive constant M_{11} , independent of $\delta \in (0, \tilde{\delta}]$, such that*

$$\begin{aligned} & \|\varphi^\delta\|_{L^\infty(0, T; \mathcal{H}^1)} + \|\varphi^\delta\|_{L^2(0, T; V)} + \delta^{1/2} \|\nabla_\Gamma \psi^\delta\|_{L^2(0, T; H_\Gamma)} \\ & + \|\xi^\delta\|_{L^1(0, T; L^1(\Omega))} + \|\xi_\Gamma^\delta\|_{L^1(0, T; L^1(\Gamma))} \leq M_{11}. \end{aligned} \quad (4.3)$$

Proof. Recalling (3.21), we have

$$\begin{aligned} & \|\omega_\varepsilon^\delta\|_{L^\infty(0, T; \mathcal{H}^{-1}_0)} + \|\nabla \omega_\varepsilon^\delta\|_{L^2(0, T; H)} + \delta^{1/2} \|\nabla_\Gamma \omega_{\Gamma, \varepsilon}^\delta\|_{L^2(0, T; H_\Gamma)} \\ & + \|\beta_\varepsilon(\varphi_\varepsilon^\delta)\|_{L^1(0, T; L^1(\Omega))} + \|\beta_{\Gamma, \varepsilon}(\psi_\varepsilon^\delta)\|_{L^1(0, T; L^1(\Gamma))} \leq C, \end{aligned}$$

where $C > 0$ is independent of δ, ε , thanks to the assumptions (2.37), (2.38). By the weak lower semicontinuity as $\varepsilon \rightarrow 0$, we find

$$\begin{aligned} & \|\omega^\delta\|_{L^\infty(0, T; \mathcal{H}^{-1}_0)} + \|\nabla \omega^\delta\|_{L^2(0, T; H)} + \delta^{1/2} \|\nabla_\Gamma \omega_\Gamma^\delta\|_{L^2(0, T; H_\Gamma)} \\ & + \|\xi^\delta\|_{L^1(0, T; L^1(\Omega))} + \|\xi_\Gamma^\delta\|_{L^1(0, T; L^1(\Gamma))} \leq C, \end{aligned}$$

where $\omega^\delta = \varphi^\delta - \bar{m}_0^\delta \mathbf{1}$ and $C > 0$ is independent of δ .

The above estimate combined with (4.1) and Poincaré's inequality (2.12) yields the conclusion (4.3). \blacksquare

Lemma 4.3. *There exists a positive constant M_{12} , independent of $\delta \in (0, \tilde{\delta}]$, such that*

$$\begin{aligned} & \|\varphi^\delta\|_{L^\infty(0, T; V)} + \delta^{1/2} \|\nabla_\Gamma \psi^\delta\|_{L^\infty(0, T; H_\Gamma)} + \|\hat{\beta}(\varphi^\delta)\|_{L^\infty(0, T; L^1(\Omega))} \\ & + \|\hat{\beta}_\Gamma(\psi^\delta)\|_{L^\infty(0, T; L^1(\Gamma))} + \|\partial_t \varphi^\delta\|_{L^2(0, T; \mathcal{H}^{-1}_0)} + \|\mathbf{P}\mu^\delta\|_{L^2(0, T; \mathcal{H}^1_{L^0})} \leq M_{12}. \end{aligned} \quad (4.4)$$

Proof. Recalling (3.29) and using the assumptions (2.37), (2.38), we get

$$\begin{aligned} & \frac{1}{4} \|\nabla \omega_\varepsilon^\delta(t)\|_H^2 + \frac{\delta}{2} \|\nabla \omega_{\Gamma,\varepsilon}^\delta(t)\|_{H_\Gamma}^2 + \frac{1}{2} \int_0^t \|\partial_\tau \omega_\varepsilon^\delta(s)\|_{0,*}^2 \, ds \\ & + \int_\Omega \widehat{\beta}_\varepsilon(\varphi_\varepsilon^\delta(t)) \, dx + \int_\Gamma \widehat{\beta}_{\Gamma,\varepsilon}(\psi_\varepsilon^\delta(t)) \, dS \leq C \quad \forall t \in [0, T], \end{aligned}$$

where $C > 0$ is independent of δ, ε . Using (3.33) and passing to the limit as $\varepsilon \rightarrow 0$, we deduce from the weak lower semicontinuity of norms that

$$\begin{aligned} & \|\nabla \varphi^\delta(t)\|_H^2 + \delta \|\nabla \psi_\delta(t)\|_{H_\Gamma}^2 + \int_0^t \|\partial_\tau \varphi^\delta(s)\|_{0,*}^2 \, ds + \int_0^t \|\mathbf{P}\boldsymbol{\mu}^\delta(s)\|_{H_{L,0}^1}^2 \, ds \\ & + \int_\Omega \widehat{\beta}(\varphi^\delta(t)) \, dx + \int_\Gamma \widehat{\beta}_\Gamma(\psi^\delta(t)) \, dS \leq C \quad \text{for a.a. } t \in [0, T]. \end{aligned}$$

The above estimate combined with (4.1) and Poincaré’s inequality (2.12) yields the conclusion (4.4). ■

Remark 4.4. From (4.4), we easily obtain the uniform estimate

$$\|\boldsymbol{\varphi}^\delta\|_{L^\infty(0,T;\mathfrak{V}^1)} + \delta^{1/2} \|\boldsymbol{\varphi}^\delta\|_{L^\infty(0,T;\mathfrak{V}^1)} \leq C.$$

Lemma 4.5. *There exists a positive constant M_{13} , independent of $\delta \in (0, \widetilde{\delta}]$ such that*

$$\|\xi^\delta\|_{L^2(0,T;L^1(\Omega))} + \|\xi_\Gamma^\delta\|_{L^2(0,T;L^1(\Gamma))} + \|\boldsymbol{\mu}^\delta\|_{L^2(0,T;\mathcal{H}^1)} \leq M_{13}. \tag{4.5}$$

Proof. Using the estimates obtained in Lemmas 4.1–4.3, we can get the conclusion (4.5) by the same argument for Lemma 3.5 and passing to the limit as $\varepsilon \rightarrow 0$. ■

Lemma 4.6. *There exists a positive constant M_{14} , independent of $\delta \in (0, \widetilde{\delta}]$, such that*

$$\begin{aligned} & \|\Delta \varphi^\delta\|_{L^2(0,T;H)} + \|\partial_n \varphi^\delta\|_{L^2(0,T;(H^{1/2}(\Gamma)))} + \delta^{1/2} \|\Delta_\Gamma \psi^\delta\|_{L^2(0,T;V_\Gamma')} \\ & + \|\xi^\delta\|_{L^2(0,T;H)} + \|\xi_\Gamma^\delta\|_{L^2(0,T;V_\Gamma')} \leq M_{14}. \end{aligned} \tag{4.6}$$

Proof. By the same argument for (3.38) and using the assumptions (2.37), (2.38), we can deduce that

$$\|\beta_\varepsilon(\varphi_\varepsilon^\delta)\|_{L^2(0,T;H)} + \varrho^{-1/2} \|\beta_\varepsilon(\psi_\varepsilon^\delta)\|_{L^2(0,T;H_\Gamma)} \leq C,$$

where $C > 0$ is independent of δ, ε . Passing to the limit as $\varepsilon \rightarrow 0$, we get

$$\|\xi^\delta\|_{L^2(0,T;H)} + \varrho^{-1/2} \|\xi^\delta\|_{L^2(0,T;H_\Gamma)} \leq C. \tag{4.7}$$

By comparison in (2.27), we infer from the above estimates and the assumption (2.38) that

$$\|\Delta \varphi^\delta\|_{L^2(0,T;H)} \leq C, \tag{4.8}$$

where $C > 0$ is independent of δ . Thanks to the trace theorem [7, Theorem 2.27] and the elliptic regularity theorem [7, Theorem 3.2], we infer from the estimates (4.4), (4.8) that

$$\|\partial_{\mathbf{n}}\varphi^\delta\|_{L^2(0,T;(H^{1/2}(\Gamma))')} \leq C. \tag{4.9}$$

By comparison in (2.28), we infer from the estimates (4.4), (4.5), (4.9) and the assumption (2.38) that

$$\begin{aligned} & \|-\delta\Delta_\Gamma\psi^\delta + \xi_\Gamma^\delta\|_{L^2(0,T;(H^{1/2}(\Gamma))')} \\ & \leq \|\theta^\delta\|_{L^2(0,T;H_\Gamma)} + \|\partial_{\mathbf{n}}\varphi^\delta\|_{L^2(0,T;(H^{1/2}(\Gamma))')} + \|\pi_\Gamma(\psi^\delta)\|_{L^2(0,T;H_\Gamma)} \\ & \quad + \|f_\Gamma^\delta\|_{L^2(0,T;H_\Gamma)} \\ & \leq C. \end{aligned} \tag{4.10}$$

Since $\delta^{1/2}\Delta_\Gamma\psi^\delta$ is uniformly bounded in $L^2(0, T; V'_\Gamma)$ by the estimate (4.3), a direct comparison in (4.10) yields

$$\|\xi_\Gamma^\delta\|_{L^2(0,T;V'_\Gamma)} \leq C. \tag{4.11}$$

Collecting the estimates (4.7), (4.8), (4.9), and (4.11), we arrive at the conclusion (4.6). ■

4.2. Passage to the limit as $\delta \rightarrow 0$

Proof of Theorem 2.15. From the uniform estimates obtained in Lemmas 4.1–4.6, we find that there exists a triplet (φ, μ, ξ) satisfying the regularity properties

$$\begin{aligned} \varphi & \in H^1(0, T; (\mathcal{H}^1)') \cap L^\infty(0, T; \tilde{\mathcal{V}}^1), \quad \Delta\varphi \in L^2(0, T; H), \\ \mu & \in L^2(0, T; \mathcal{H}^1), \\ \xi & \in L^2(0, T; H \times (H^{1/2}(\Gamma))'), \end{aligned}$$

and the following convergence results hold as $\delta \rightarrow 0$ (in the sense of a subsequence):

$$\varphi^\delta \rightharpoonup \varphi \quad \text{weakly star in } L^\infty(0, T; \tilde{\mathcal{V}}^1), \tag{4.12}$$

$$\varphi^\delta \rightarrow \varphi \quad \text{weakly in } H^1(0, T; (\mathcal{H}^1)'), \tag{4.13}$$

$$\Delta\varphi^\delta \rightharpoonup \Delta\varphi \quad \text{weakly in } L^2(0, T; H), \tag{4.14}$$

$$\mu^\delta \rightharpoonup \mu \quad \text{weakly in } L^2(0, T; \mathcal{H}^1), \tag{4.15}$$

$$\xi^\delta \rightharpoonup \xi \quad \text{weakly in } L^2(0, T; H \times V'_\Gamma), \tag{4.16}$$

$$\delta\varphi^\delta \rightarrow \mathbf{0} \quad \text{strongly in } L^\infty(0, T; \mathcal{V}^1), \tag{4.17}$$

$$-\delta\Delta_\Gamma\psi^\delta + \xi_\Gamma^\delta \rightharpoonup \xi_\Gamma \quad \text{weakly in } L^2(0, T; (H^{1/2}(\Gamma))'). \tag{4.18}$$

Here, we note that (4.17) implies $\delta\Delta_\Gamma\psi^\delta \rightarrow 0$ in $L^2(0, T; V'_\Gamma)$; thus, (4.16) and (4.18) are consistent. Next, by Lemma A.1, we can conclude

$$\varphi^\delta \rightarrow \varphi \quad \text{strongly in } C([0, T]; \mathcal{L}^2). \tag{4.19}$$

This implies $\varphi|_{t=0} = \varphi_0$ thanks to the assumption (2.39). Besides, it follows from (4.2) that

$$\bar{m}(\varphi(t)) = \bar{m}(\varphi_0) \quad \forall t \in [0, T].$$

The strong convergence (4.19) combined with the assumption (A3) also gives

$$\begin{aligned} \pi(\varphi^\delta) &\rightarrow \pi(\varphi) \quad \text{strongly in } C([0, T]; H), \\ \pi_\Gamma(\psi^\delta) &\rightarrow \pi_\Gamma(\psi) \quad \text{strongly in } C([0, T]; H_\Gamma). \end{aligned}$$

Passing to the weak limit as $\delta \rightarrow 0$ in (2.26) yields the variational formulation (2.31). Next, we write (3.54) as

$$\begin{aligned} (\boldsymbol{\mu}^\delta(t), \mathbf{z})_{\mathcal{X}^2} &= (\nabla\varphi^\delta(t), \nabla z)_H + (\xi^\delta, z)_H + \langle -\delta\Delta_\Gamma\psi^\delta + \xi_\Gamma^\delta, z_\Gamma \rangle_{(H^{1/2}(\Gamma))', H^{1/2}(\Gamma)} \\ &\quad + (\boldsymbol{\pi}(\varphi^\delta(t)) - \mathbf{f}^\delta(t), \mathbf{z})_{\mathcal{X}^2} \quad \forall \mathbf{z} \in \mathcal{V}^1 \end{aligned}$$

for almost all $t \in (0, T)$. Letting $\delta \rightarrow 0$, we easily recover (2.32). Thanks to the classical results in [4, 6], the weak convergence (4.16) and the strong convergence (4.19) combined with the demi-closedness of the maximal monotone operator β yield (2.33), that is, $\xi \in \beta(\varphi)$ almost everywhere in Q . Moreover, by exactly the same argument as in [18, Section 3.2], we can justify the variational inequality (2.34) for ξ_Γ . According to (4.18) and [18, Remark 2.6], we have $\xi_\Gamma \in \partial J_\Sigma(\psi)$, where

$$J_\Sigma : L^2(0, T; H^{1/2}(\Gamma)) \rightarrow [0, +\infty], \quad J_\Sigma := \begin{cases} \int_\Sigma \widehat{\beta}_\Gamma(\psi) \, dS \, dt & \text{if } \widehat{\beta}_\Gamma(\psi) \in L^1(\Sigma), \\ +\infty & \text{otherwise.} \end{cases}$$

In summary, the limit $(\varphi, \boldsymbol{\mu}, \xi)$ is a weak solution to the system $(S_{L,0})$ in the sense of Definition 2.13. The proof of Theorem 2.15 is complete. ■

4.3. Uniqueness and further properties

Proof of Theorem 2.17. The proof is similar to that for Theorem 2.9. Keeping in mind that $\delta = 0$, in analogy to (3.57), we obtain

$$\begin{aligned} &\frac{1}{2} \frac{d}{ds} \|\boldsymbol{\omega}_1(s) - \boldsymbol{\omega}_2(s)\|_{0,*}^2 + \|\nabla(\omega_1(s) - \omega_2(s))\|_H^2 \\ &\quad + \int_\Omega (\xi_1(s) - \xi_2(s))(\omega_1(s) - \omega_2(s)) \, dx \\ &\quad + \langle \xi_{\Gamma,1}(s) - \xi_{\Gamma,2}(s), \omega_{\Gamma,1}(s) - \omega_{\Gamma,2}(s) \rangle_{(H^{1/2}(\Gamma))', H^{1/2}(\Gamma)} \\ &\leq (K + K_\Gamma) \|\boldsymbol{\omega}_1(s) - \boldsymbol{\omega}_2(s)\|_{\mathcal{X}^2}^2 + \|\mathbf{f}_1(s) - \mathbf{f}_2(s)\|_{(\widetilde{\mathcal{V}}_1)', \|\boldsymbol{\omega}_1(s) - \boldsymbol{\omega}_2(s)\|_{\widetilde{\mathcal{V}}_1}, \end{aligned} \tag{4.20}$$

which holds for almost all $s \in (0, T)$. Since $\boldsymbol{\omega}_1(s) - \boldsymbol{\omega}_2(s) \in \widetilde{\mathcal{V}}_{(0)}^1$ for almost all $s \in (0, T)$, from Poincaré’s inequality (2.12) and the interpolation inequality (2.13), we get

$$\begin{aligned} \|\boldsymbol{\omega}_1(s) - \boldsymbol{\omega}_2(s)\|_{\widetilde{\mathcal{V}}_1} &\leq C \|\nabla(\omega_1(s) - \omega_2(s))\|_H, \\ \|\boldsymbol{\omega}_1(s) - \boldsymbol{\omega}_2(s)\|_{\mathcal{X}^2}^2 &\leq \gamma \|\nabla(\omega_1(s) - \omega_2(s))\|_H^2 + C_\gamma \|\boldsymbol{\omega}_1(s) - \boldsymbol{\omega}_2(s)\|_{0,*}^2 \end{aligned}$$

for any $\gamma > 0$ and almost all $s \in (0, T)$. Besides, it follows from the monotonicity of β , β_Γ that

$$\int_{\Omega} (\xi_1(s) - \xi_2(s))(\omega_1(s) - \omega_2(s)) \, dx + \langle \xi_{\Gamma,1}(s) - \xi_{\Gamma,2}(s), \omega_{\Gamma,1}(s) - \omega_{\Gamma,2}(s) \rangle_{(H^{1/2}(\Gamma))^\gamma, H^{1/2}(\Gamma)} \geq 0 \quad \text{for a.a. } s \in (0, T).$$

Inserting the above inequalities into (4.20), taking γ sufficiently small, then by Grönwall’s lemma, the fact $\bar{m}(\varphi_1) = \bar{m}(\varphi_2)$, Young’s inequality, and Poincaré’s inequality (2.12), we can conclude (2.40). ■

Finally, if the bulk and boundary potentials exhibit the same growth, some further properties of the asymptotic limit as $\delta \rightarrow 0$ can be obtained. More precisely, we have the following corollary.

Corollary 4.7. *Let the assumptions of Theorem 2.15 be satisfied. In addition, we assume that*

(A8) $D(\beta) = D(\beta_\Gamma)$ and there exists a constant $M \geq 1$ such that

$$\frac{1}{M} |\beta_\Gamma^\circ(r)| - M \leq |\beta^\circ(r)| \leq M |\beta_\Gamma^\circ(r)| + M \quad \forall r \in D(\beta).$$

(1) *Improved regularity: the limit triplet (φ, μ, ξ) obtained in Theorem 2.15 satisfies*

$$\varphi \in L^2(0, T; H^{3/2}(\Omega)), \quad \partial_n \varphi \in L^2(0, T; H_\Gamma), \quad \psi \in L^2(0, T; V_\Gamma), \\ \xi_\Gamma \in L^2(0, T; H_\Gamma), \quad \xi_\Gamma \in \beta_\Gamma(\psi) \quad \text{a.e. on } \Sigma.$$

(2) *Refined convergence (in the sense of a subsequence): it holds*

$$\xi_\Gamma^\delta \rightharpoonup \xi \quad \text{weakly in } L^2(0, T; H_\Gamma), \\ \delta \psi^\delta \rightharpoonup 0 \quad \text{weakly in } L^2(0, T; H^{3/2}(\Gamma)), \\ \partial_n \varphi^\delta - \delta \Delta_\Gamma \psi^\delta \rightharpoonup \partial_n \varphi \quad \text{weakly in } L^2(0, T; H_\Gamma)$$

as $\delta \rightarrow 0$ and (2.36) can be replaced by $\theta = \partial_n \varphi + \xi_\Gamma + \pi_\Gamma(\psi) - f_\Gamma$ almost everywhere on Σ .

(3) *Convergence rate: if $\bar{m}(\varphi_0^\delta) \equiv \bar{m}(\varphi_0)$ for $\delta \in (0, 1)$, then: for all $\delta \in (0, 1)$, it holds*

$$\|\varphi^\delta - \varphi\|_{L^\infty(0, T; \mathcal{H}_{(0)}^{-1})} + \|\varphi^\delta - \varphi\|_{L^2(0, T; \tilde{V}^1)} \\ \leq C(\delta^{1/2} + \|\varphi_0^\delta - \varphi_0\|_{0,*} + \|f^\delta - f\|_{L^2(0, T; \mathcal{X}^2)}).$$

Since the proof of Corollary 4.7 follows the same arguments for [18, Theorems 2.10, 2.12], we leave the details to interested readers.

5. Asymptotic limits with respect to kinetic rate: $L \rightarrow 0$ or $L \rightarrow +\infty$

In this section, we study the asymptotic limit of solutions to problem $(S_{L,\delta})$ as $L \rightarrow 0$ or $L \rightarrow +\infty$, with $\delta \in (0, +\infty)$ being fixed.

For any $\varepsilon \in (0, 1)$, we consider the approximating problem (3.8)–(3.9) with data (φ_0, f) . Let ω_ε^L be the unique solution given by Proposition 3.1. We set

$$\varphi_\varepsilon^L = \omega_\varepsilon^L + \bar{m}_0 \mathbf{1}, \quad \mu_\varepsilon^L := \varepsilon \partial_t \omega_\varepsilon^L + \partial \Phi_\delta(\omega_\varepsilon^L) + \beta_\varepsilon(\varphi_\varepsilon^L) + \pi(\varphi_\varepsilon^L) - f,$$

where $\bar{m}_0 = \bar{m}(\varphi_0)$. Then, $(\varphi_\varepsilon^L, \mu_\varepsilon^L)$ is uniquely determined and satisfies (3.13)–(3.19) with data (φ_0, f) .

As in the proof of Theorem 2.7, when $\varepsilon \rightarrow 0$, $(\varphi_\varepsilon^L, \mu_\varepsilon^L, \beta_\varepsilon(\varphi_\varepsilon^L))$ converges to certain limit triplet $(\varphi^L, \mu^L, \xi^L)$, which is a weak solution to problem $(S_{L,\delta})$ with data (φ_0, f) . Since the uniqueness of μ^L, ξ^L is not clear (cf., Theorem 2.9), the related convergence should always be understood in the sense of a suitable subsequence. Below we will not relabel the convergent subsequence for the sake of simplicity.

The proofs of Theorems 2.20 and 2.23 rely on uniform estimates with respect to the kinetic rate L . The first estimate comes from the mass conservation. Recalling (3.20) and passing to the limit $\varepsilon \rightarrow 0$, we get

$$\bar{m}(\varphi^L(t)) = \bar{m}(\varphi_0) = \bar{m}_0 \quad \forall t \in [0, T]. \tag{5.1}$$

Nevertheless, further uniform estimates have to be derived separately for the two different cases $L \rightarrow 0$ and $L \rightarrow +\infty$.

5.1. The case $L \rightarrow 0$

We now proceed to derive uniform estimates for $L \in (0, 1]$.

Lemma 5.1. *There exists a positive constant C_1 , independent of $L \in (0, 1]$, such that*

$$\begin{aligned} & \|\omega^L\|_{L^\infty(0,T;\mathcal{H}_{(0)}^{-1})} + \|\omega^L\|_{L^2(0,T;\mathcal{V}_{(0)}^1)} \\ & + \|\xi^L\|_{L^1(0,T;L^1(\Omega))} + \|\xi_\Gamma^L\|_{L^1(0,T;L^1(\Gamma))} \leq C_1, \end{aligned} \tag{5.2}$$

where $\omega^L = \varphi^L - \bar{m}_0 \mathbf{1}$.

Proof. Recalling the derivation of (3.21), we have

$$\begin{aligned} & \|\omega_\varepsilon^L\|_{L^\infty(0,T;\mathcal{H}_{(0)}^{-1})}^2 + \|\omega_\varepsilon^L\|_{L^2(0,T;\mathcal{V}_{(0)}^1)}^2 + \|\beta_\varepsilon(\varphi_\varepsilon^L)\|_{L^1(0,T;L^1(\Omega))} \\ & + \|\beta_{\Gamma,\varepsilon}(\psi_\varepsilon^L)\|_{L^1(0,T;L^1(\Gamma))} \\ & \leq C(\varepsilon\|\omega_0\|_{\mathcal{H}^2}^2 + \|\omega_0\|_{0,*}^2 + 1), \end{aligned}$$

where $C > 0$ is independent of L, ε . By the weak lower semicontinuity as $\varepsilon \rightarrow 0$, we find

$$\begin{aligned} & \|\omega^L\|_{L^\infty(0,T;\mathcal{H}_{(0)}^{-1})}^2 + \|\omega^L\|_{L^2(0,T;\mathcal{V}_{(0)}^1)}^2 + \|\xi^L\|_{L^1(0,T;L^1(\Omega))} + \|\xi_\Gamma^L\|_{L^1(0,T;L^1(\Gamma))} \\ & \leq C(\|\omega_0\|_{0,*}^2 + 1). \end{aligned} \tag{5.3}$$

Recall that, for any fixed $L \in (0, +\infty)$, the norms $\|\cdot\|_{\mathcal{H}^1}$ and $\|\cdot\|_{\mathcal{H}_{L,0}^1}$ are equivalent on $\mathcal{H}_{L,0}^1$. This fact yields

$$\|\mathbf{y}\|_{\mathcal{X}^2} \leq \|\mathbf{y}\|_{\mathcal{H}^1} \leq \widehat{C} \|\mathbf{y}\|_{\mathcal{H}_{L,0}^1} \leq \widehat{C} \|\mathbf{y}\|_{\mathcal{H}_{L,0}^1} \quad \forall \mathbf{y} \in \mathcal{H}_{L,0}^1 \quad \forall L \in (0, 1], \quad (5.4)$$

where $\widehat{C} > 0$ is independent of $L \in (0, 1]$. As a consequence, it holds

$$\begin{aligned} \|\boldsymbol{\omega}_0\|_{0,*}^2 &= \|\mathcal{S}^L \boldsymbol{\omega}_0\|_{\mathcal{H}_{L,0}^1}^2 = (\boldsymbol{\omega}_0, \mathcal{S}^L \boldsymbol{\omega}_0)_{\mathcal{X}^2} \\ &\leq \|\boldsymbol{\omega}_0\|_{\mathcal{X}^2} \|\mathcal{S}^L \boldsymbol{\omega}_0\|_{\mathcal{X}^2} \leq \widehat{C} \|\boldsymbol{\omega}_0\|_{\mathcal{X}^2} \|\mathcal{S}^L \boldsymbol{\omega}_0\|_{\mathcal{H}_{L,0}^1} = \widehat{C} \|\boldsymbol{\omega}_0\|_{\mathcal{X}^2} \|\boldsymbol{\omega}_0\|_{0,*}. \end{aligned}$$

Hence, $\|\boldsymbol{\omega}_0\|_{0,*} \leq \widehat{C} \|\boldsymbol{\omega}_0\|_{\mathcal{X}^2}$, and thus, it is uniformly bounded for $L \in (0, 1]$. This together with (5.3) yields the conclusion (5.2). ■

Lemma 5.2. *There exists a positive constant C_2 , independent of $L \in (0, 1]$, such that*

$$\begin{aligned} \|\varphi^L\|_{L^\infty(0,T;\mathcal{V}^1)} + \|\widehat{\beta}(\varphi^L)\|_{L^\infty(0,T;L^1(\Omega))} + \|\widehat{\beta}_\Gamma(\psi^L)\|_{L^\infty(0,T;L^1(\Gamma))} \\ + \|\partial_t \varphi^L\|_{L^2(0,T;\mathcal{H}_{(0)}^{-1})} + \|\mathbf{P}\boldsymbol{\mu}^L\|_{L^2(0,T;\mathcal{H}_{L,0}^1)} \leq C_2. \end{aligned} \quad (5.5)$$

Proof. Recalling (3.29) and (3.33), passing to the limit as $\varepsilon \rightarrow 0$, we can deduce the uniform estimate (5.5) from the weak lower semicontinuity, (5.1), (5.2), and Poincaré’s inequality (2.12). ■

Lemma 5.3. *There exists a positive constant C_3 , independent of $L \in (0, 1]$, such that*

$$\|\partial_t \boldsymbol{\varphi}^L\|_{L^2(0,T;(\mathcal{V}^1)')} \leq C_3. \quad (5.6)$$

Proof. From (5.5) and the definition of $\|\cdot\|_{\mathcal{H}_{L,0}^1}$, we find

$$\|\nabla \boldsymbol{\mu}^L\|_{L^2(0,T;H)} + \|\nabla_\Gamma \theta^L\|_{L^2(0,T;H_\Gamma)} + \frac{1}{\sqrt{L}} \|\boldsymbol{\mu}^L - \theta^L\|_{L^2(0,T;H_\Gamma)} \leq C, \quad (5.7)$$

where $C > 0$ is independent of $L \in (0, 1]$. Taking $\mathbf{z} = (z, z_\Gamma) \in \mathcal{V}^1$ in the weak formulation (2.26), we get

$$(\partial_t \boldsymbol{\varphi}^L(t), \mathbf{z})_{(\mathcal{V}^1)', \mathcal{V}^1} + \int_\Omega \nabla \boldsymbol{\mu}^L(t) \cdot \nabla z \, dx + \int_\Gamma \nabla \theta^L(t) \cdot \nabla_\Gamma z_\Gamma \, dS = 0$$

for almost all $t \in (0, T)$. Then, it follows that

$$\begin{aligned} |(\partial_t \boldsymbol{\varphi}^L(t), \mathbf{z})_{(\mathcal{V}^1)', \mathcal{V}^1}| &\leq \|\nabla \boldsymbol{\mu}^L(t)\|_H \|\nabla z\|_H + \|\nabla \theta^L(t)\|_{H_\Gamma} \|\nabla_\Gamma z_\Gamma\|_{H_\Gamma} \\ &\leq (\|\nabla \boldsymbol{\mu}^L(t)\|_H + \|\nabla \theta^L(t)\|_{H_\Gamma}) \|\mathbf{z}\|_{\mathcal{V}^1}, \end{aligned}$$

which together with (5.7) implies the estimate (5.6). ■

Lemma 5.4. *There exists a positive constant C_4 , independent of $L \in (0, 1]$, such that*

$$\|\xi^L\|_{L^2(0,T;L^1(\Omega))} + \|\xi_\Gamma^L\|_{L^2(0,T;L^1(\Gamma))} + \|\mu^L\|_{L^2(0,T;\mathcal{H}^1)} \leq C_4. \tag{5.8}$$

Proof. Keeping the estimates (5.1), (5.2), (5.4), (5.5) in mind, we can obtain the uniform estimate (5.8) with respect to $L \in (0, 1]$ by the same argument for Lemma 3.5 and passing to the limit as $\varepsilon \rightarrow 0$. ■

Lemma 5.5. *There exists a positive constant C_5 , independent of $L \in (0, 1]$, such that*

$$\|\xi^L\|_{L^2(0,T;H)} + \|\xi_\Gamma^L\|_{L^2(0,T;H_\Gamma)} + \|\varphi^L\|_{L^2(0,T;\mathcal{V}^2)} \leq C_5.$$

Proof. To conclude, we apply the same argument for Lemma 3.6 and then pass to the limit as $\varepsilon \rightarrow 0$. The estimate for $\|\varphi^L\|_{L^2(0,T;\mathcal{V}^2)}$ follows from the estimate for $\|\omega^L\|_{L^2(0,T;\mathcal{V}^2)}$, (5.1), and Poincaré’s inequality (2.12). ■

Proof of Theorem 2.20. The existence of a limit triplet $(\varphi^0, \mu^0, \xi^0)$ with expected regularity properties and the (sequential) convergence results (2.44)–(2.48) are guaranteed by (5.1), Lemmas 5.1–5.5, and the same compactness argument as in the proof of Theorem 2.15. By (2.46), (2.48) and the maximal monotonicity of β and β_Γ , we find

$$\xi^0 \in \beta(\varphi^0) \quad \text{a.e. in } Q, \quad \xi_\Gamma^0 \in \beta_\Gamma(\psi^0) \quad \text{a.e. on } \Sigma.$$

Then, taking limit $L \rightarrow 0$ in (2.26), (3.54), we recover (2.41) and

$$(\mu^0(t), z)_{\mathcal{X}^2} = b_\delta(\varphi^0(t), z) + (\xi^0(t) + \pi(\varphi^0(t)) - f(t), z)_{\mathcal{X}^2} \quad \forall z \in \mathcal{V}^1 \tag{5.9}$$

holds for almost all $t \in (0, T)$. Since $z \in \mathcal{V}^1$ is arbitrary, we can conclude (2.42) and (2.43) from (5.9). The inequality (2.49) follows from (5.7). This also implies

$$\mu^0|_\Gamma = \theta^0 \quad \text{a.e. on } \Sigma.$$

Hence, $(\varphi^0, \mu^0, \xi^0)$ is a weak solution of GMS model in the sense of Definition 2.19. The uniqueness of φ^0 has been established in [16, Theorem 2.1]. The proof of Theorem 2.20 is complete. ■

5.2. The case $L \rightarrow +\infty$

We now derive uniform estimates for $L \geq \widehat{L}$, where $\widehat{L} \geq 1$ is a constant to be determined later.

Lemma 5.6. *There exists a positive constant C_6 , independent of $L \geq 1$, such that*

$$\begin{aligned} &\|\partial_t \omega^L\|_{L^2(0,T;\mathcal{H}^{-1}_0)} + \|\varphi^L\|_{L^\infty(0,T;\mathcal{V}^1)} \\ &+ \|\widehat{\beta}(\varphi^L)\|_{L^\infty(0,T;L^1(\Omega))} + \|\widehat{\beta}_\Gamma(\psi^L)\|_{L^\infty(0,T;L^1(\Gamma))} \leq C_6. \end{aligned} \tag{5.10}$$

Proof. Testing (3.8) at time $s \in (0, T)$ by $\partial_t \omega_\varepsilon^L \in \mathcal{L}_{(0)}^2$, with the help of (2.23), we obtain

$$\begin{aligned} & \frac{d}{ds} \left(\Phi_\delta(\omega_\varepsilon^L(s)) + \int_\Omega \widehat{\beta}_\varepsilon(\varphi_\varepsilon^L(s)) \, dx + \int_\Gamma \widehat{\beta}_{\Gamma,\varepsilon}(\psi_\varepsilon^L(s)) \, dS \right) \\ & \quad + \varepsilon \|\partial_t \omega_\varepsilon^L(s)\|_{\mathcal{X}^2}^2 + \|\omega_\varepsilon^L(s)\|_{0,*}^2 \\ & = -(\mathbf{P}\boldsymbol{\pi}(\varphi_\varepsilon^L(s)), \partial_t \omega_\varepsilon^L(s))_{\mathcal{X}^2} + (\mathbf{P}\mathbf{f}(s), \partial_t \omega_\varepsilon^L(s))_{\mathcal{X}^2} \quad \text{for a.a. } s \in (0, T). \end{aligned} \tag{5.11}$$

Similar to (3.27), it holds

$$\begin{aligned} (\mathbf{P}\mathbf{f}(s), \partial_t \omega_\varepsilon^L(s))_{\mathcal{X}^2} &= (\mathbf{P}\mathbf{f}(s), \mathfrak{S}^L(\partial_t \omega_\varepsilon^L(s)))_{\mathcal{H}_{L,0}^1} \\ &\leq \frac{1}{4} \|\partial_t \omega_\varepsilon^L(s)\|_{0,*}^2 + \|\mathbf{f}(s)\|_{\mathcal{V}^1}^2, \end{aligned} \tag{5.12}$$

and

$$\begin{aligned} |(\mathbf{P}\boldsymbol{\pi}(\varphi_\varepsilon^L(s)), \partial_t \omega_\varepsilon^L(s))_{\mathcal{X}^2}| &= |(\mathbf{P}\boldsymbol{\pi}(\varphi_\varepsilon^L(s)), \mathfrak{S}^L(\partial_t \omega_\varepsilon^L(s)))_{\mathcal{H}_{L,0}^1}| \\ &\leq \|\mathbf{P}\boldsymbol{\pi}(\varphi_\varepsilon^L(s))\|_{\mathcal{H}_{L,0}^1} \|\partial_t \omega_\varepsilon^L(s)\|_{0,*} \\ &\leq \frac{1}{4} \|\partial_t \omega_\varepsilon^L(s)\|_{0,*}^2 + \|\mathbf{P}\boldsymbol{\pi}(\varphi_\varepsilon^L(s))\|_{\mathcal{H}_{L,0}^1}^2. \end{aligned} \tag{5.13}$$

Thanks to the additional assumption that π, π_Γ are differentiable on \mathbb{R} , we deduce from (A3), $\varphi_\varepsilon^L = \omega_\varepsilon^L + \bar{m}_0 \mathbf{1}$, and Poincaré’s inequality (2.12) for all $L \geq 1$, it holds

$$\begin{aligned} \|\mathbf{P}\boldsymbol{\pi}(\varphi_\varepsilon^L(s))\|_{\mathcal{H}_{L,0}^1}^2 &\leq \|\nabla \pi(\varphi_\varepsilon^L(s))\|_H^2 + \|\nabla_\Gamma \pi_\Gamma(\psi_\varepsilon^L(s))\|_{H_\Gamma}^2 \\ &\quad + \|\pi(\varphi_\varepsilon^L(s)) - \pi_\Gamma(\psi_\varepsilon^L(s))\|_{H_\Gamma}^2 \\ &\leq K^2 \|\nabla \omega_\varepsilon^L(s)\|_H^2 + K_\Gamma^2 \|\nabla_\Gamma \omega_{\Gamma,\varepsilon}^L(s)\|_{H_\Gamma}^2 + 3K^2 \|\omega_\varepsilon^L(s)\|_H^2 \\ &\quad + 3K_\Gamma^2 \|\omega_{\Gamma,\varepsilon}^L(s)\|_{H_\Gamma}^2 + 6|\Gamma|(|\pi(\bar{m}_0)|^2 + |\pi_\Gamma(\bar{m}_0)|^2) \\ &\leq C \|\omega_\varepsilon^L(s)\|_{\mathcal{V}_{(0)}^1}^2 + C, \end{aligned} \tag{5.14}$$

where $C > 0$ is independent of $L \geq 1$. In view of (5.12), (5.13), (5.14), and (A4), we infer from (5.11) and Grönwall’s lemma that, for all $L \geq 1$, it holds

$$\begin{aligned} & \frac{1}{2} \|\omega_\varepsilon^L(t)\|_{\mathcal{V}_{(0)}^1}^2 + \int_\Omega \widehat{\beta}_\varepsilon(\varphi_\varepsilon^L(t)) \, dx + \int_\Gamma \widehat{\beta}_{\Gamma,\varepsilon}(\psi_\varepsilon^L(t)) \, dS \\ & \quad + \varepsilon \int_0^t \|\partial_t \omega_\varepsilon^L(s)\|_{\mathcal{X}^2}^2 \, ds + \frac{1}{2} \int_0^t \|\partial_t \omega_\varepsilon^L(s)\|_{0,*}^2 \, ds \leq C \quad \forall t \in [0, T], \end{aligned}$$

where $C > 0$ depends on $\|\omega_0\|_{\mathcal{V}_{(0)}^1}^2, \bar{m}_0, \int_\Omega \widehat{\beta}(\varphi_0) \, dx, \int_\Gamma \widehat{\beta}_\Gamma(\psi_0) \, dS, \|\mathbf{f}\|_{L^2(0,T;\mathcal{V}^1)}, \Omega,$ and T . Passing to the limit as $\varepsilon \rightarrow 0$, by the weak lower semicontinuity and $\varphi^L = \omega^L + \bar{m}_0 \mathbf{1}$, we obtain the estimate (5.10). ■

Lemma 5.7. *There exists a positive constant C_7 , independent of $L \geq 1$, such that*

$$\|\partial_t \varphi^L\|_{L^2(0,T;\mathcal{V}')} \leq C_7, \tag{5.15}$$

$$\|\partial_t \psi^L\|_{L^2(0,T;\mathcal{V}'_\Gamma)} \leq C_7. \tag{5.16}$$

Proof. From (5.10) and the definition of $\|\cdot\|_{\mathcal{H}_{L,0}^1}$, we see that, for $L \geq 1$, it holds

$$\|\nabla\mu^L\|_{L^2(0,T;H)} + \|\nabla\Gamma\theta^L\|_{L^2(0,T;H_\Gamma)} + \frac{1}{\sqrt{L}}\|\mu^L - \theta^L\|_{L^2(0,T;H_\Gamma)} \leq C, \quad (5.17)$$

where $C > 0$ is independent of L . For any $z \in V$, taking $\mathbf{z} = (z, 0) \in \mathcal{H}^1$ in (2.26), we obtain

$$\langle \partial_t \varphi^L(t), z \rangle_{V',V} + \int_\Omega \nabla\mu^L(t) \cdot \nabla z \, dx + \frac{1}{L} \int_\Gamma (\mu^L(t) - \theta^L(t))z \, dS = 0 \quad (5.18)$$

for almost all $t \in (0, T)$. It follows that

$$\begin{aligned} |\langle \partial_t \varphi^L(t), z \rangle_{V',V}| &\leq \|\nabla\mu^L(t)\|_H \|\nabla z\|_H + \frac{1}{L} \|\mu^L(t) - \theta^L(t)\|_{H_\Gamma} \|z\|_{H_\Gamma} \\ &\leq C \left(\|\nabla\mu^L(t)\|_H + \frac{1}{\sqrt{L}} \cdot \frac{1}{\sqrt{L}} \|\mu^L(t) - \theta^L(t)\|_{H_\Gamma} \right) \|z\|_V, \end{aligned}$$

which implies

$$\|\partial_t \varphi^L(t)\|_{V'} \leq C \left(\|\nabla\mu^L(t)\|_H + \frac{1}{\sqrt{L}} \cdot \frac{1}{\sqrt{L}} \|\mu^L(t) - \theta^L(t)\|_{H_\Gamma} \right).$$

Then, by (5.17), we get (5.15). Analogously, for any $z_\Gamma \in V_\Gamma$, taking $\mathbf{z} = (0, z_\Gamma) \in \mathcal{H}^1$ in (2.26), we obtain

$$\langle \partial_t \psi^L(t), z_\Gamma \rangle_{V'_\Gamma, V_\Gamma} + \int_\Gamma \nabla_\Gamma \theta^L(t) \cdot \nabla_\Gamma z_\Gamma \, dS - \frac{1}{L} \int_\Gamma (\mu^L(t) - \theta^L(t))z_\Gamma \, dS = 0.$$

By the same reasoning, we arrive at (5.16). ■

Lemma 5.8. *There exists a large constant $\widehat{L} \geq 1$ such that, for all $L \geq \widehat{L}$, it holds*

$$\|\xi^L\|_{L^2(0,T;L^1(\Omega))} + \|\xi_\Gamma^L\|_{L^2(0,T;L^1(\Gamma))} \leq C_8(1 + \|\partial_n \varphi^L\|_{L^2(0,T;H_\Gamma)}), \quad (5.19)$$

where $C_8 > 0$ is independent of L .

Proof. Taking the test function

$$\mathbf{y} = (\mathcal{N}_\Omega(\varphi^L - \langle \varphi^L \rangle_\Omega), \mathcal{N}_\Gamma(\psi^L - \langle \psi^L \rangle_\Gamma)) \in \mathcal{H}^1$$

in (2.26), then we obtain

$$\begin{aligned} 0 &= \langle \partial_t \varphi^L, \mathcal{N}_\Omega(\varphi^L - \langle \varphi^L \rangle_\Omega) \rangle_{V',V} + \langle \partial_t \psi^L, \mathcal{N}_\Gamma(\psi^L - \langle \psi^L \rangle_\Gamma) \rangle_{V'_\Gamma, V_\Gamma} \\ &\quad + \int_\Omega \nabla\mu^L \cdot \nabla \mathcal{N}_\Omega(\varphi^L - \langle \varphi^L \rangle_\Omega) \, dx + \int_\Gamma \nabla_\Gamma \theta^L \cdot \nabla_\Gamma \mathcal{N}_\Gamma(\psi^L - \langle \psi^L \rangle_\Gamma) \, dS \\ &\quad + \frac{1}{L} \int_\Gamma (\mu^L - \theta^L) [\mathcal{N}_\Omega(\varphi^L - \langle \varphi^L \rangle_\Omega) - \mathcal{N}_\Gamma(\psi^L - \langle \psi^L \rangle_\Gamma)] \, dS \end{aligned}$$

$$\begin{aligned}
 &= \langle \partial_t \varphi^L, \mathcal{N}_\Omega(\varphi^L - \langle \varphi^L \rangle_\Omega) \rangle_{V', V} + \langle \partial_t \psi^L, \mathcal{N}_\Gamma(\psi^L - \langle \psi^L \rangle_\Gamma) \rangle_{V'_\Gamma, V_\Gamma} \\
 &\quad + \int_\Omega \mu^L (\varphi^L - \langle \varphi^L \rangle_\Omega) \, dx + \int_\Gamma \theta^L (\psi^L - \langle \psi^L \rangle_\Gamma) \, dS \\
 &\quad + \frac{1}{L} \int_\Gamma (\mu^L - \theta^L) [\mathcal{N}_\Omega(\varphi^L - \langle \varphi^L \rangle_\Omega) - \mathcal{N}_\Gamma(\psi^L - \langle \psi^L \rangle_\Gamma)] \, dS \\
 &=: \sum_{j=1}^5 R_j. \tag{5.20}
 \end{aligned}$$

It follows from the definition of \mathcal{N}_Ω , \mathcal{N}_Γ , Hölder's inequality, and Poincaré's inequalities (2.1), (2.2) that

$$\begin{aligned}
 |R_1| + |R_2| &\leq |\langle \partial_t \varphi^L, \mathcal{N}_\Omega(\varphi^L - \langle \varphi^L \rangle_\Omega) \rangle_{V', V}| + |\langle \partial_t \psi^L, \mathcal{N}_\Gamma(\psi^L - \langle \psi^L \rangle_\Gamma) \rangle_{V'_\Gamma, V_\Gamma}| \\
 &\leq C \|\partial_t \varphi^L\|_{V'} \|\nabla \varphi^L\|_H + C \|\partial_t \psi^L\|_{V'_\Gamma} \|\nabla_\Gamma \psi^L\|_{H_\Gamma}. \tag{5.21}
 \end{aligned}$$

Next, by the trace theorem and the elliptic estimates, we find

$$\begin{aligned}
 |R_5| &= \left| \frac{1}{L} \int_\Gamma (\mu^L - \theta^L) [\mathcal{N}_\Omega(\varphi^L - \langle \varphi^L \rangle_\Omega) - \mathcal{N}_\Gamma(\psi^L - \langle \psi^L \rangle_\Gamma)] \, dS \right| \\
 &\leq \frac{C}{L} \|\mu^L - \theta^L\|_{H_\Gamma} (\|\varphi^L - \langle \varphi^L \rangle_\Omega\|_{V_0^*} + \|\psi^L - \langle \psi^L \rangle_\Gamma\|_{V_{\Gamma_0}^*}) \\
 &\leq \frac{C}{\sqrt{L}} \left(\frac{1}{\sqrt{L}} \|\mu^L - \theta^L\|_{H_\Gamma} \right) (\|\varphi^L\|_H + \|\psi^L\|_{H_\Gamma}). \tag{5.22}
 \end{aligned}$$

The estimates for R_3 , R_4 are more involved. By (2.27) and (2.28), we get

$$\begin{aligned}
 R_3 + R_4 &= \int_\Omega \mu^L (\varphi^L - \langle \varphi^L \rangle_\Omega) \, dx + \int_\Gamma \theta^L (\psi^L - \langle \psi^L \rangle_\Gamma) \, dS \\
 &= \int_\Omega |\nabla \varphi^L|^2 \, dx + \delta \int_\Gamma |\nabla_\Gamma \psi^L|^2 \, dS \\
 &\quad + \int_\Omega \xi^L (\varphi^L - \langle \varphi^L \rangle_\Omega) \, dx + \int_\Gamma \xi_\Gamma^L (\psi^L - \langle \psi^L \rangle_\Gamma) \, dS \\
 &\quad + \int_\Omega (\pi(\varphi^L) - f)(\varphi^L - \langle \varphi^L \rangle_\Omega) \, dx \\
 &\quad + \int_\Gamma (\pi_\Gamma(\psi^L) - f_\Gamma)(\psi^L - \langle \psi^L \rangle_\Gamma) \, dS \\
 &\quad + (\langle \varphi^L \rangle_\Omega - \langle \psi^L \rangle_\Gamma) \int_\Gamma \partial_n \varphi^L \, dS. \tag{5.23}
 \end{aligned}$$

Consider the third and fourth terms on the right-hand side of (5.23):

$$\begin{aligned}
 &\int_\Omega \xi^L (\varphi^L - \langle \varphi^L \rangle_\Omega) \, dx + \int_\Gamma \xi_\Gamma^L (\psi^L - \langle \psi^L \rangle_\Gamma) \, dS \\
 &= \int_\Omega \xi^L (\varphi^L - \langle \varphi_0 \rangle_\Omega) \, dx + \int_\Gamma \xi_\Gamma^L (\psi^L - \langle \psi_0 \rangle_\Gamma) \, dS \\
 &\quad + (\langle \varphi_0 \rangle_\Omega - \langle \varphi^L \rangle_\Omega) \int_\Omega \xi^L \, dx + (\langle \psi_0 \rangle_\Gamma - \langle \psi^L \rangle_\Gamma) \int_\Gamma \xi_\Gamma^L \, dS. \tag{5.24}
 \end{aligned}$$

It follows from (5.17) and (5.18) that

$$\begin{aligned}
 |(\varphi_0)_\Omega - (\varphi^L(t))_\Omega| &= \frac{1}{|\Omega|} \left| \int_0^t \langle \partial_t \varphi^L(s), 1 \rangle_{V',V} ds \right| \\
 &\leq \frac{1}{|\Omega|} \frac{1}{L} \int_0^t \|\mu^L(s) - \theta^L(s)\|_{L^1(\Gamma)} ds \\
 &\leq \frac{|\Gamma|^{1/2} |T|^{1/2}}{|\Omega|} \frac{1}{\sqrt{L}} \left(\frac{1}{\sqrt{L}} \|\mu^L - \theta^L\|_{L^2(0,T;H_\Gamma)} \right) \\
 &\leq \frac{C' |\Gamma|^{1/2} |T|^{1/2}}{|\Omega|} \frac{1}{\sqrt{L}} \quad \forall t \in [0, T],
 \end{aligned} \tag{5.25}$$

and in a similar manner,

$$|(\psi_0)_\Gamma - (\psi^L(t))_\Gamma| \leq \frac{C'' |T|^{1/2}}{|\Gamma|^{1/2}} \frac{1}{\sqrt{L}} \quad \forall t \in [0, T], \tag{5.26}$$

where the constants $C', C'' > 0$ in (5.25), (5.26) are independent of L . Since $(\varphi_0)_\Omega, (\psi_0)_\Gamma \in \text{Int } D(\beta_\Gamma)$, recalling (3.6), we can first work with the approximating solutions φ_ε^L and then pass to the limit as $\varepsilon \rightarrow 0$ to get

$$\begin{aligned}
 \int_\Omega \xi^L (\varphi^L - (\varphi_0)_\Omega) dx &\geq \delta_0^{(1)} \|\xi^L\|_{L^1(\Omega)} - c_1^{(1)} |\Omega|, \\
 \int_\Gamma \xi_\Gamma^L (\psi^L - (\psi_0)_\Gamma) dS &\geq \delta_0^{(2)} \|\xi_\Gamma^L\|_{L^1(\Gamma)} - c_1^{(2)} |\Gamma|.
 \end{aligned}$$

Set $\hat{\delta}_0 = \min\{\delta_0^{(1)}, \delta_0^{(2)}\}, \hat{c}_1 = \max\{c_1^{(1)}, c_1^{(2)}\}$. There exists some $\hat{L} \geq 1$ sufficiently large such that

$$\frac{C' |\Gamma|^{\frac{1}{2}} |T|^{\frac{1}{2}}}{|\Omega|} \frac{1}{\sqrt{L}} \leq \frac{\hat{\delta}_0}{2}, \quad \frac{C'' |T|^{\frac{1}{2}}}{|\Gamma|^{\frac{1}{2}}} \frac{1}{\sqrt{L}} \leq \frac{\hat{\delta}_0}{2} \quad \forall L \geq \hat{L}.$$

Then, it follows from (5.24)–(5.26) that

$$\begin{aligned}
 \int_\Omega \xi^L (\varphi^L - (\varphi^L)_\Omega) dx + \int_\Gamma \xi_\Gamma^L (\psi^L - (\psi^L)_\Gamma) dS \\
 \geq \frac{\hat{\delta}_0}{2} (\|\xi^L\|_{L^1(\Omega)} + \|\xi_\Gamma^L\|_{L^1(\Gamma)}) - \hat{c}_1 (|\Omega| + |\Gamma|).
 \end{aligned} \tag{5.27}$$

Concerning the last three terms on the right-hand side of (5.23), from (A3) and Hölder’s inequality, we get

$$\begin{aligned}
 &\left| \int_\Omega (\pi(\varphi^L) - f)(\varphi^L - (\varphi^L)_\Omega) dx \right| + \left| \int_\Gamma (\pi_\Gamma(\psi^L) - f_\Gamma)(\psi^L - (\psi^L)_\Gamma) dS \right| \\
 &\quad + \left| ((\varphi^L)_\Omega - (\psi^L)_\Gamma) \int_\Gamma \partial_n \varphi^L dS \right| \\
 &\leq C \|\varphi^L\|_{\mathcal{X}^2} (1 + \|\varphi^L\|_{\mathcal{X}^2} + \|f\|_{\mathcal{X}^2} + \|\partial_n \varphi^L\|_{H_\Gamma}).
 \end{aligned} \tag{5.28}$$

Returning to (5.20), owing to the estimates (5.21)–(5.28), we infer from (5.10), (5.15), (5.16), (5.17), and (A4) that

$$\begin{aligned} & \frac{\widehat{\delta}_0}{2} (\|\xi^L\|_{L^2(0,T;L^1(\Omega))} + \|\xi_\Gamma^L\|_{L^2(0,T;L^1(\Gamma))}) \\ & \leq C T^{1/2} (1 + \|\varphi^L\|_{L^\infty(0,T;\mathcal{X}^2)}^2) \\ & \quad + C \|\varphi^L\|_{L^\infty(0,T;\mathcal{X}^2)} (\|f\|_{L^2(0,T;\mathcal{X}^2)} + \|\partial_n \varphi^L\|_{L^2(0,T;H_\Gamma)}) \\ & \quad + \frac{C}{\sqrt{L}} \left(\frac{1}{\sqrt{L}} \|\mu^L - \theta^L\|_{L^2(0,T;H_\Gamma)} \right) \|\varphi^L\|_{L^\infty(0,T;\mathcal{X}^2)} \\ & \quad + C (\|\partial_t \varphi^L\|_{L^2(0,T;V')} + \|\partial_t \psi^L\|_{L^2(0,T;V'_\Gamma)}) \|\varphi^L\|_{L^\infty(0,T;V^1)} \\ & \leq C (1 + \|\partial_n \varphi^L\|_{L^2(0,T;H_\Gamma)}) \quad \forall L \geq \widehat{L}, \end{aligned}$$

where $C > 0$ is independent of $L \in [\widehat{L}, +\infty)$. Thus, the conclusion (5.19) follows. ■

Lemma 5.9. *There exists a positive constant C_9 , independent of $L \geq \widehat{L}$, such that*

$$\|\mu^L\|_{L^2(0,T;\mathcal{H}^1)} \leq C_9 (1 + \|\partial_n \varphi^L\|_{L^2(0,T;H_\Gamma)}). \tag{5.29}$$

Proof. From (2.27) and (2.28), we find

$$\begin{aligned} \langle \mu^L \rangle_\Omega &= \frac{1}{|\Omega|} \left[\int_\Omega (\xi^L + \pi(\varphi^L) - f) \, dx - \int_\Gamma \partial_n \varphi^L \, dS \right], \\ \langle \theta^L \rangle_\Gamma &= \frac{1}{|\Gamma|} \int_\Gamma (\xi_\Gamma^L + \pi(\psi^L) - f_\Gamma + \partial_n \varphi^L) \, dS, \end{aligned}$$

which together with (5.10), (5.19) and (A4) imply

$$\|\langle \mu^L \rangle_\Omega\|_{L^2(0,T)} + \|\langle \theta^L \rangle_\Gamma\|_{L^2(0,T)} \leq C (1 + \|\partial_n \varphi^L\|_{L^2(0,T;H_\Gamma)}). \tag{5.30}$$

By (5.17), (5.30) and Poincaré’s inequalities (2.1), (2.2), we obtain (5.29). ■

Lemma 5.10. *There exists a positive constant C_{10} , independent of $L \geq \widehat{L}$, such that*

$$\begin{aligned} & \|\xi^L\|_{L^2(0,T;H)} + \|\xi_\Gamma^L\|_{L^2(0,T;H_\Gamma)} + \|\varphi^L\|_{L^2(0,T;\mathcal{H}^2)} \\ & \leq C_{10} (1 + \|\partial_n \varphi^L\|_{L^2(0,T;H_\Gamma)}). \end{aligned} \tag{5.31}$$

Proof. To conclude, we apply the same argument for Lemma 3.6 and then pass to the limit as $\varepsilon \rightarrow 0$. The estimate for $\|\varphi^L\|_{L^2(0,T;V^2)}$ follows from the estimate for $\|\omega^L\|_{L^2(0,T;V^2)}$, (5.1) and Poincaré’s inequality (2.12). ■

Proof of Theorem 2.23. To complete the proof, it remains to control $\|\partial_n \varphi^L\|_{L^2(0,T;H_\Gamma)}$. Thanks to the trace theorem and the Ehrling lemma (see Lemma A.2), we find

$$\begin{aligned} \|\partial_n \varphi^L\|_{L^2(0,T;H_\Gamma)} &\leq C \|\varphi^L\|_{L^2(0,T;H^{2-r}(\Omega))} \quad \text{for some } r \in (0, 1/2) \\ &\leq \epsilon \|\varphi^L\|_{L^2(0,T;H^2(\Omega))} + C_\epsilon \|\varphi^L\|_{L^2(0,T;V)}. \end{aligned}$$

Hence, taking $\epsilon > 0$ sufficiently small, we can deduce from (5.10) and (5.31) that

$$\|\varphi^L\|_{L^2(0,T;\mathcal{H}^2)} \leq \frac{1}{2}\|\varphi^L\|_{L^2(0,T;H^2(\Omega))} + C \quad \forall L \geq \widehat{L}.$$

This yields the uniform bound of $\|\varphi^L\|_{L^2(0,T;\mathcal{H}^2)}$ for all $L \geq \widehat{L}$, and thus, $\|\partial_n \varphi^L\|_{L^2(0,T;H_\Gamma)}$ is bounded as well.

Keeping the above uniform estimates in mind, we can take the limit as $L \rightarrow +\infty$ (in the sense of a subsequence) and prove Theorem 2.23 in the same way as for Theorem 2.20. ■

A. Appendix

We report some technical lemmas that have been frequently used in our analysis.

First, we recall the compactness lemma of Aubin–Lions–Simon type (see, for instance, [48] in the case $q > 1$ and [61] when $q = 1$).

Lemma A.1. *Let $X_0 \xrightarrow{c} X_1 \subset X_2$, where X_j are (real) Banach spaces ($j = 0, 1, 2$). Let $1 < p \leq +\infty$, $1 \leq q \leq +\infty$ and I be a bounded subinterval of \mathbb{R} . Then, the sets*

$$\{\varphi \in L^p(I; X_0) : \partial_t \varphi \in L^q(I; X_2)\} \xrightarrow{c} L^p(I; X_1) \quad \text{if } 1 < p < +\infty,$$

and

$$\{\varphi \in L^p(I; X_0) : \partial_t \varphi \in L^q(I; X_2)\} \xrightarrow{c} C(I; X_1) \quad \text{if } p = +\infty, q > 1.$$

The following Ehrling lemma can be found in [48].

Lemma A.2. *Let B_0, B_1, B be three Banach spaces so that B_0 and B_1 are reflexive. Moreover,*

$$B_0 \hookrightarrow B \hookrightarrow B_1.$$

Then, for each $\epsilon > 0$, there exists a positive constant C_ϵ depending on ϵ such that

$$\|z\|_B \leq \epsilon \|z\|_{B_0} + C_\epsilon \|z\|_{B_1} \quad \forall z \in B_0.$$

Acknowledgments. The authors would like to thank the anonymous referees for their careful reading of an initial version of this paper and for many helpful comments that allowed them to improve the presentation.

Funding. The research of the corresponding author H. Wu was partially supported by NNSFC (Grant no. 12071084) and the Shanghai Center for Mathematical Sciences at Fudan University. H. Wu is a member of the Key Laboratory of Mathematics for Nonlinear Sciences (Fudan University), Ministry of Education of China.

References

- [1] D. M. Anderson, G. B. McFadden, and A. A. Wheeler, [Diffuse-interface methods in fluid mechanics](#). In *Annual review of fluid mechanics*, Vol. 30, pp. 139–165, Annu. Rev. Fluid Mech. 30, Annual Reviews, Palo Alto, CA, 1998 Zbl [1398.76051](#) MR [1609626](#)
- [2] E. Bänsch, K. Deckelnick, H. Garcke, and P. Pozzi, [Interfaces: modeling, analysis, numerics](#). Oberwolfach Semin. 51, Birkhäuser/Springer, Cham, 2023 Zbl [07756324](#) MR [4676342](#)
- [3] X. Bao and H. Zhang, [Numerical approximations and error analysis of the Cahn–Hilliard equation with reaction rate dependent dynamic boundary conditions](#). *J. Sci. Comput.* **87** (2021), no. 3, article no. 72 Zbl [1475.65101](#) MR [4248673](#)
- [4] V. Barbu, [Nonlinear differential equations of monotone types in Banach spaces](#). Springer Monogr. Math., Springer, New York, 2010 Zbl [1197.35002](#) MR [2582280](#)
- [5] J. F. Blowey and C. M. Elliott, [The Cahn–Hilliard gradient theory for phase separation with nonsmooth free energy. I. Mathematical analysis](#). *European J. Appl. Math.* **2** (1991), no. 3, 233–280 Zbl [0797.35172](#) MR [1123143](#)
- [6] H. Brézis, [Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert](#). North-Holland Math. Stud. 5, North-Holland, Amsterdam; American Elsevier Publishing, New York, 1973 Zbl [0252.47055](#) MR [0348562](#)
- [7] F. Brezzi and G. Gilardi, [Chapters 1–3 in Finite element handbook](#). McGraw-Hill Book, New York, 1987 Zbl [0638.65076](#) MR [0900813](#)
- [8] J. W. Cahn and J. E. Hilliard, [Free energy of a nonuniform system I. Interfacial free energy](#). *J. Chem. Phys.* **28** (1958), no. 2, 258–267 Zbl [1431.35066](#)
- [9] L. Calatroni and P. Colli, [Global solution to the Allen–Cahn equation with singular potentials and dynamic boundary conditions](#). *Nonlinear Anal.* **79** (2013), 12–27 Zbl [1270.35275](#) MR [3005023](#)
- [10] K. Chelmiński and P. Gwiazda, [Nonhomogeneous initial-boundary value problems for coercive and self-controlling models of monotone type](#). *Contin. Mech. Thermodyn.* **12** (2000), no. 4, 217–234 Zbl [1003.74033](#) MR [1779916](#)
- [11] X. Chen, X. Wang, and X. Xu, [Analysis of the Cahn–Hilliard equation with a relaxation boundary condition modeling the contact angle dynamics](#). *Arch. Ration. Mech. Anal.* **213** (2014), no. 1, 1–24 Zbl [1293.35235](#) MR [3198643](#)
- [12] L. Cherfils, S. Gatti, and A. Miranville, [A variational approach to a Cahn–Hilliard model in a domain with nonpermeable walls](#). *J. Math. Sci. (N.Y.)* **189** (2013), no. 4, 604–636 Zbl [1276.49014](#) MR [3098333](#)
- [13] L. Cherfils, A. Miranville, and S. Zelik, [The Cahn–Hilliard equation with logarithmic potentials](#). *Milan J. Math.* **79** (2011), no. 2, 561–596 Zbl [1250.35129](#) MR [2862028](#)
- [14] L. Cherfils and M. Petcu, [A numerical analysis of the Cahn–Hilliard equation with nonpermeable walls](#). *Numer. Math.* **128** (2014), no. 3, 517–549 Zbl [1325.65133](#) MR [3268846](#)
- [15] P. Colli and T. Fukao, [Cahn–Hilliard equation with dynamic boundary conditions and mass constraint on the boundary](#). *J. Math. Anal. Appl.* **429** (2015), no. 2, 1190–1213 Zbl [1327.35149](#) MR [3342513](#)
- [16] P. Colli and T. Fukao, [Equation and dynamic boundary condition of Cahn–Hilliard type with singular potentials](#). *Nonlinear Anal.* **127** (2015), 413–433 Zbl [1323.35092](#) MR [3392376](#)
- [17] P. Colli and T. Fukao, [Vanishing diffusion in a dynamic boundary condition for the Cahn–Hilliard equation](#). *NoDEA Nonlinear Differential Equations Appl.* **27** (2020), no. 6, article no. 53 Zbl [1452.35080](#) MR [4160936](#)

- [18] P. Colli, T. Fukao, and L. Scarpa, [The Cahn–Hilliard equation with forward-backward dynamic boundary condition via vanishing viscosity](#). *SIAM J. Math. Anal.* **54** (2022), no. 3, 3292–3315 Zbl 1504.35063 MR 4431962
- [19] P. Colli, T. Fukao, and L. Scarpa, [A Cahn–Hilliard system with forward-backward dynamic boundary condition and non-smooth potentials](#). *J. Evol. Equ.* **22** (2022), no. 4, article no. 89 Zbl 1501.35250 MR 4507218
- [20] P. Colli, T. Fukao, and H. Wu, [On a transmission problem for equation and dynamic boundary condition of Cahn–Hilliard type with nonsmooth potentials](#). *Math. Nachr.* **293** (2020), no. 11, 2051–2081 Zbl 1525.35163 MR 4188682
- [21] P. Colli, G. Gilardi, and J. Sprekels, [On the Cahn–Hilliard equation with dynamic boundary conditions and a dominating boundary potential](#). *J. Math. Anal. Appl.* **419** (2014), no. 2, 972–994 Zbl 1295.35248 MR 3225416
- [22] P. Colli, G. Gilardi, and J. Sprekels, [On a Cahn–Hilliard system with convection and dynamic boundary conditions](#). *Ann. Mat. Pura Appl. (4)* **197** (2018), no. 5, 1445–1475 Zbl 1406.35163 MR 3848459
- [23] P. Colli, P. Knopf, G. Schimperna, and A. Signori, [Two-phase flows through porous media described by a Cahn–Hilliard–Brinkman model with dynamic boundary conditions](#). *J. Evol. Equ.* **24** (2024), no. 4, article no. 85 MR 4805011
- [24] P. Colli and A. Visintin, [On a class of doubly nonlinear evolution equations](#). *Comm. Partial Differential Equations* **15** (1990), no. 5, 737–756 Zbl 0707.34053 MR 1070845
- [25] Q. Du and X. Feng, [The phase field method for geometric moving interfaces and their numerical approximations](#). In *Geometric partial differential equations. Part I*, pp. 425–508, Handb. Numer. Anal. 21, Elsevier/North-Holland, Amsterdam, 2020 Zbl 1455.35276 MR 4378430
- [26] G. Dziuk and C. M. Elliott, [Finite element methods for surface PDEs](#). *Acta Numer.* **22** (2013), 289–396 Zbl 1296.65156 MR 3038698
- [27] H. P. Fischer, P. Maass, and W. Dieterich, [Novel surface modes in spinodal decomposition](#). *Phys. Rev. Lett.* **79** (1997), no. 5, 893–896
- [28] T. Fukao and H. Wu, [Separation property and convergence to equilibrium for the equation and dynamic boundary condition of Cahn–Hilliard type with singular potential](#). *Asymptot. Anal.* **124** (2021), no. 3-4, 303–341 Zbl 1502.35110 MR 4303022
- [29] C. G. Gal, [A Cahn–Hilliard model in bounded domains with permeable walls](#). *Math. Methods Appl. Sci.* **29** (2006), no. 17, 2009–2036 Zbl 1109.35057 MR 2268279
- [30] C. G. Gal, M.-Y. Lv, and H. Wu, [On a thermodynamically consistent diffuse interface model for two-phase incompressible flows with non-matched densities: Dynamics of moving contact lines, surface diffusion, and mass transfer](#). 2024, researchGate:RG.2.2.27875.73764
- [31] C. G. Gal and H. Wu, [Asymptotic behavior of a Cahn–Hilliard equation with Wentzell boundary conditions and mass conservation](#). *Discrete Contin. Dyn. Syst.* **22** (2008), no. 4, 1041–1063 Zbl 1158.35052 MR 2434981
- [32] H. Garcke and P. Knopf, [Weak solutions of the Cahn–Hilliard system with dynamic boundary conditions: A gradient flow approach](#). *SIAM J. Math. Anal.* **52** (2020), no. 1, 340–369 Zbl 1429.35114 MR 4054589
- [33] H. Garcke, P. Knopf, and S. Yayla, [Long-time dynamics of the Cahn–Hilliard equation with kinetic rate dependent dynamic boundary conditions](#). *Nonlinear Anal.* **215** (2022), article no. 112619 Zbl 1479.35094 MR 4330180
- [34] G. Gilardi, A. Miranville, and G. Schimperna, [On the Cahn–Hilliard equation with irregular potentials and dynamic boundary conditions](#). *Commun. Pure Appl. Anal.* **8** (2009), no. 3, 881–912 Zbl 1172.35417 MR 2476663

- [35] G. Gilardi, A. Miranville, and G. Schimperna, [Long time behavior of the Cahn–Hilliard equation with irregular potentials and dynamic boundary conditions](#). *Chinese Ann. Math. Ser. B* **31** (2010), no. 5, 679–712 Zbl [1223.35067](#) MR [2726062](#)
- [36] A. Giorgini, M. Grasselli, and A. Miranville, [The Cahn–Hilliard–Oono equation with singular potential](#). *Math. Models Methods Appl. Sci.* **27** (2017), no. 13, 2485–2510 Zbl [1386.35023](#) MR [3714635](#)
- [37] A. Giorgini and P. Knopf, [Two-phase flows with bulk-surface interaction: thermodynamically consistent Navier–Stokes–Cahn–Hilliard models with dynamic boundary conditions](#). *J. Math. Fluid Mech.* **25** (2023), no. 3, article no. 65 Zbl [1529.35383](#) MR [4610947](#)
- [38] G. R. Goldstein, A. Miranville, and G. Schimperna, [A Cahn–Hilliard model in a domain with non-permeable walls](#). *Phys. D* **240** (2011), no. 8, 754–766 Zbl [1228.35051](#) MR [2770968](#)
- [39] P. Gwiazda, J. Skrzeczkowski, and L. Trussardi, [On the rate of convergence of Yosida approximation for the nonlocal Cahn–Hilliard equation](#). *IMA J. Numer. Anal.* (2024), 1–19
- [40] X. Jing and Q. Wang, [Thermodynamically consistent dynamic boundary conditions of phase field models](#). *Commun. Math. Sci.* **21** (2023), no. 3, 859–883 Zbl [1518.35238](#) MR [4562456](#)
- [41] P. Knopf and K. F. Lam, [Convergence of a Robin boundary approximation for a Cahn–Hilliard system with dynamic boundary conditions](#). *Nonlinearity* **33** (2020), no. 8, 4191–4235 Zbl [1461.35099](#) MR [4169658](#)
- [42] P. Knopf, K. F. Lam, C. Liu, and S. Metzger, [Phase-field dynamics with transfer of materials: The Cahn–Hilliard equation with reaction rate dependent dynamic boundary conditions](#). *ESAIM Math. Numer. Anal.* **55** (2021), no. 1, 229–282 Zbl [1470.35173](#) MR [4216838](#)
- [43] P. Knopf and C. Liu, [On second-order and fourth-order elliptic systems consisting of bulk and surface PDEs: Well-posedness, regularity theory and eigenvalue problems](#). *Interfaces Free Bound.* **23** (2021), no. 4, 507–533 Zbl [1485.35162](#) MR [4337445](#)
- [44] P. Knopf and A. Signori, [On the nonlocal Cahn–Hilliard equation with nonlocal dynamic boundary condition and boundary penalization](#). *J. Differential Equations* **280** (2021), 236–291 Zbl [1465.35286](#) MR [4207300](#)
- [45] P. Knopf and J. Stange, [Strong well-posedness and separation properties for a bulk-surface convective Cahn–Hilliard system with singular potentials](#). 2024, arXiv:[2407.14089v1](#)
- [46] P. Knopf and J. Stange, [Well-posedness of a bulk-surface convective Cahn–Hilliard system with dynamic boundary conditions](#). *NoDEA Nonlinear Differential Equations Appl.* **31** (2024), no. 5, article no. 82 MR [4761865](#)
- [47] M. Kubo, [The Cahn–Hilliard equation with time-dependent constraint](#). *Nonlinear Anal.* **75** (2012), no. 14, 5672–5685 Zbl [1250.35130](#) MR [2942945](#)
- [48] J.-L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*. Dunod, Paris; Gauthier-Villars, Paris, 1969 Zbl [0189.40603](#) MR [0259693](#)
- [49] C. Liu and H. Wu, [An energetic variational approach for the Cahn–Hilliard equation with dynamic boundary condition: model derivation and mathematical analysis](#). *Arch. Ration. Mech. Anal.* **233** (2019), no. 1, 167–247 Zbl [1416.35209](#) MR [3974640](#)
- [50] M.-Y. Lv and H. Wu, [On the Cahn–Hilliard equation with kinetic rate dependent dynamic boundary condition and non-smooth potential: Separation property and long-time behavior](#). 2024, arXiv:[2405.20807v1](#)
- [51] M.-Y. Lv and H. Wu, [Separation property and asymptotic behavior for a transmission problem of the bulk-surface coupled Cahn–Hilliard system with singular potentials and its Robin approximation](#). 2024, arXiv:[2407.01986v1](#)

- [52] S. Metzger, [An efficient and convergent finite element scheme for Cahn–Hilliard equations with dynamic boundary conditions](#). *SIAM J. Numer. Anal.* **59** (2021), no. 1, 219–248
Zbl [1458.35345](#) MR [4202510](#)
- [53] A. Miranville, *The Cahn–Hilliard equation. Recent advances and applications*. CBMS-NSF Regional Conf. Ser. in Appl. Math. 95, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2019 Zbl [1446.35001](#) MR [4001523](#)
- [54] A. Miranville and H. Wu, [Long-time behavior of the Cahn–Hilliard equation with dynamic boundary condition](#). *J. Elliptic Parabol. Equ.* **6** (2020), no. 1, 283–309 Zbl [1443.35054](#)
MR [4114353](#)
- [55] A. Miranville and S. Zelik, [Robust exponential attractors for Cahn–Hilliard type equations with singular potentials](#). *Math. Methods Appl. Sci.* **27** (2004), no. 5, 545–582
Zbl [1050.35113](#) MR [2041814](#)
- [56] A. Miranville and S. Zelik, [Exponential attractors for the Cahn–Hilliard equation with dynamic boundary conditions](#). *Math. Methods Appl. Sci.* **28** (2005), no. 6, 709–735 Zbl [1068.35020](#)
MR [2125817](#)
- [57] T. Qian, X.-P. Wang, and P. Sheng, [A variational approach to moving contact line hydrodynamics](#). *J. Fluid Mech.* **564** (2006), 333–360 Zbl [1178.76296](#) MR [2261865](#)
- [58] R. Racke and S. Zheng, [The Cahn–Hilliard equation with dynamic boundary conditions](#). *Adv. Differential Equations* **8** (2003), no. 1, 83–110 Zbl [1035.35050](#) MR [1946559](#)
- [59] L. Scarpa, [Existence and uniqueness of solutions to singular Cahn–Hilliard equations with nonlinear viscosity terms and dynamic boundary conditions](#). *J. Math. Anal. Appl.* **469** (2019), no. 2, 730–764 Zbl [1516.35224](#) MR [3860445](#)
- [60] R. E. Showalter, *Monotone operators in Banach space and nonlinear partial differential equations*. Math. Surveys Monogr. 49, American Mathematical Society, Providence, RI, 1997
Zbl [0870.35004](#) MR [1422252](#)
- [61] J. Simon, [Compact sets in the space \$L^p\(0, T; B\)\$](#) . *Ann. Mat. Pura Appl. (4)* **146** (1987), 65–96
Zbl [0629.46031](#) MR [0916688](#)
- [62] H. Wu, [Convergence to equilibrium for a Cahn–Hilliard model with the Wentzell boundary condition](#). *Asymptot. Anal.* **54** (2007), no. 1-2, 71–92 Zbl [1139.34028](#) MR [2356465](#)
- [63] H. Wu, [A review on the Cahn–Hilliard equation: Classical results and recent advances in dynamic boundary conditions](#). *Electron. Res. Arch.* **30** (2022), no. 8, 2788–2832
Zbl [1517.35002](#) MR [4432003](#)
- [64] H. Wu and S. Zheng, [Convergence to equilibrium for the Cahn–Hilliard equation with dynamic boundary conditions](#). *J. Differential Equations* **204** (2004), no. 2, 511–531 Zbl [1068.35018](#)
MR [2085545](#)

Received 7 February 2024; revised 23 September 2024.

Maoyin Lv

School of Mathematical Sciences, Fudan University, Handan Road 220, 200433 Shanghai,
P. R. China; mylv22@m.fudan.edu.cn

Hao Wu

School of Mathematical Sciences, Fudan University, Handan Road 220, 200433 Shanghai,
P. R. China; haowufd@fudan.edu.cn