The second Picard iteration of NLS on the 2*d* sphere does not regularize Gaussian random initial data

Nicolas Burq, Nicolas Camps, Mickaël Latocca, Chenmin Sun, and Nikolay Tzvetkov

Abstract. We consider the Wick-ordered cubic Schrödinger equation (NLS) posed on the twodimensional sphere, with initial data distributed according to a Gaussian measure. We show that the second Picard iteration does not improve the regularity of the initial data in the scale of the classical Sobolev spaces. This is in sharp contrast with the Wick-ordered NLS on the two-dimensional tori, a model for which we know from the work of Bourgain that the second Picard iteration gains one half-derivative. Our proof relies on identifying a singular part of the nonlinearity. We show that this singular part is responsible for a concentration phenomenon on a large circle (i.e., a stable closed geodesic), which prevents any regularization in the second Picard iteration.

1. Introduction

1.1. Context

The present work is motivated by the study of the influence of the background geometry on the low-regularity well-posedness theory for nonlinear Schrödinger equations (and more generally for dispersive partial differential equations). Due to infinite propagation speed, even the short-time nonlinear evolution is sensitive to the geometry and (at least for certain geometries based on the model case of the 2d sphere) high-frequencies instabilities occur (see, e.g., [13, 16]). We prove in this work that these instabilities persist for randomized initial data, which exposes a fundamental obstruction for extending to the 2d sphere the probabilistic well-posedness theory for NLS that was developed in the case of Euclidean geometries over the past thirty years.

In the case of integrable partial differential equations, the best results concerning lowregularity well-posedness are exploiting fine properties of the Lax pair structures yielding thus results going beyond the scope of applicability of more traditional PDE techniques. Thomas Kappeler was a pioneer in the use of integrability techniques in the context of low-regularity well-posedness of dispersive partial differential equations; see [27, 28, 33]. Integrability methods will not be used in the present paper, but we believe that using

Keywords: Picard iteration.

Mathematics Subject Classification 2020: 35Q41.

random data techniques in the context of integrable partial differential equations is an interesting line of research.

1.1.1. The nonlinear Schrödinger equation on compact surfaces. Given (M, g) a compact Riemannian surface without boundary, the cubic Schrödinger equation (NLS) posed on M reads

$$i\partial_t u + \Delta_g u = \lambda |u|^2 u, \quad (t, x) \in \mathbb{R} \times M,$$
 (NLS)

where $\lambda \in \mathbb{R}$ dictates the attractive ($\lambda < 0$) or repulsive ($\lambda > 0$) nature of the nonlinear interaction. When s > 1, the Sobolev embedding and a fixed-point argument easily solve the Cauchy problem associated to (NLS): for every bounded set $B \subset H^s(M)$, there exist $T_B > 0$ and a unique solution map

$$\Phi: u_0 \in B \mapsto (t \mapsto \Phi^t(u_0)) \in C([-T_B, T_B]; H^s(M)).$$

For $u_0 \in B$, the mapping $t \in [-T_B, T_B] \mapsto \Phi^t(u_0)$ solves (NLS) in the integrated form, through the Duhamel formula, with initial data u_0 :

$$\Phi^{t}(u_{0}) = e^{it\Delta} u_{0} - i\lambda \int_{0}^{t} e^{i(t-t')\Delta} |\Phi^{t'}(u_{0})|^{2} \Phi^{t'}(u_{0}) dt'.$$
(1.1)

Moreover, Φ is uniformly continuous (it is actually analytic). When solving the fixed-point problem (1.1) by a Picard iteration scheme, we write

$$\Phi^{t}(u_{0}) = e^{it\Delta} u_{0} - i\lambda \int_{0}^{t} e^{i(t-t')\Delta} |e^{it'\Delta} u_{0}|^{2} e^{it'\Delta} u_{0} dt' + \text{higher-order expansions.}$$

The first nonlinear term is precisely the second Picard iteration.

Motivated by the conservation laws associated with the equation, the main question is to determine the largest Sobolev space in which the flow map extends (uniformly) continuously. The L^p - properties of the eigenfunctions of $-\Delta$ play a key role in this program.

For general (M, g), the L^p estimates on the eigenfunctions come with some derivative loss. These estimates are well known from the works of Hörmander [32] and Sogge [36] and are recalled in (3.4). They turn out to be optimal for the 2*d* sphere. Then, motivated by the study of nonlinear waves, Burq–Gérard–Tzvetkov [14] obtained semiclassical $L_t^2 L_x^\infty$ Strichartz estimates with $\frac{1}{2}$ -derivative loss and extended the uniform local well-posedness up to $H^s(M)$ with $s > \frac{1}{2}$. So far, it is the lowest common regularity where uniform local well-posedness is known for all (M, g). Besides, in the defocusing case $(\lambda > 0)$, the (coercive) conserved energy combined with the local well-posedness in $H^1(M)$ leads to global well-posedness in $H^s(M)$ for all $s \ge 1$. Hani [30] extended the range of global well-posedness in $H^s(M)$ with $s > \frac{2}{3}$.

In the Euclidean case, when $M = \mathbb{T}^2$ is the flat tori, Bourgain [4,5] proved that the scaling invariance dictates the well-posedness threshold: the flow map extends uniformly continuously in $H^s(\mathbb{T}^2)$ for all s > 0 (see also the very interesting recent work [31] for the global well-posedness in this range). This fails when s < 0 [21,22], and the endpoint

s = 0 is a challenging open problem. The Strichartz estimate due to Bourgain–Demeter [8] extends the local well-posedness result for s > 0 to the case of irrational tori.

Outside the Euclidean case, the only fairly well-understood situation is the case of the 2*d* sphere or more generally of a Zoll surface (a surface on which the geodesic flow is periodic, see [3]). In these geometries, the Cauchy problem is uniformly well posed [15] up to $s > \frac{1}{4}$ and this is optimal [2, 13].

Firstly, the uniform well-posedness result above $H^{\frac{1}{4}}(\mathbb{S}^2)$ follows from a bilinear refinement of the Strichartz estimate due to Burq–Gérard–Tzvetkov [15]. This bilinear estimate results from the localization of the spectrum of the Laplace operator on a Zoll surface combined with a general bilinear estimate on the spectral projectors (which is true on any surface).

Conversely, instabilities in $H^s(\mathbb{S}^2)$ when $s \leq \frac{1}{4}$ arise from eigenfunctions that concentrate on a stable closed geodesic, such as the equator in the case of the sphere (and more generally on a surface with a closed stable geodesic [39]). These eigenfunctions are the highest-weight spherical harmonics and were also used to construct stationary solution in the defocusing case [41]. In the present paper, we use a broader family of spherical harmonics with high weight to evidence instabilities in a probabilistic setting.

1.1.2. Statistical approaches to nonlinear waves. In [17], Burq and Lebeau considered a natural probability measure on the space of spherical harmonics and proved that almost every random orthonormal basis is uniformly bounded in $L^p(\mathbb{S}^2)$, when $p < +\infty$, contrasting with the L^p -bounds discussed above. For instance, the *n*-th highest-weight spherical harmonics have their L^4 -norm growing like $n^{\frac{1}{8}}$ due to concentration around a large circle. Former works [1, 35, 40] also go in the direction of improving the L^p -estimates for generic eigenfunctions.

The natural question is whether this enhanced Sobolev embedding for generic functions on the sphere can lead to probabilistic well-posedness below the deterministic threshold $s = \frac{1}{4}$ or not.

When s is small enough such that instabilities are known to occur in H^s for some initial data, the probabilistic well-posedness theory initiated by Burq–Tzvetkov [18,19] after the pioneering work of Bourgain [7] consists in the construction of full-measure subsets of H^s (for some natural measures) made of initial data that lead to strong solutions. In addition, the obtained probabilistic solution can be seen as the unique limit in H^s of the (deterministic) flow applied to (suitable) smooth approximations of the initial datum.

Additionally, this approach is motivated by the construction of global recurrent solutions using the Gibbs invariant measures (supported in $\bigcap_{\varepsilon>0} H^{-\varepsilon}(M)$) and, more broadly, the study of the transport of Gaussian measures by nonlinear flows. In the Euclidean setting, or more recently when the target space is a Riemannian manifold [11], the probabilistic approach has led to several significant advances. An overview is much beyond the scope of this introduction. Nonetheless, to motivate our main result, we recall the key mechanism driving the probabilistic method.

In the Euclidean case, the probabilistic decoupling between the Fourier modes of the initial datum leads to nonlinear smoothing effects. Namely, the second Picard iteration gains, say, σ -derivatives (for some fixed $\sigma > 0$) for almost every initial data (see [7] and Theorem 1.2). The standard linear-nonlinear decomposition trick consists in solving the fixed-point problem for the re-centered solution

$$v(t) := \Phi^t(u_0) - e^{it\Delta} u_0, \quad v(0) = 0,$$

(formally) solution to (NLS) with a stochastic forcing term (which corresponds to the second Picard iteration in (1.1)) and mixed terms depending both on v and $e^{it\Delta} u_0$. By understanding the mixed terms, one may indeed run a fixed-point argument for v in $H^{s+\sigma}$, provided $s + \sigma$ is greater than the deterministic regularity threshold.

1.2. Set-up and main results of the present work

The current paper is a preliminary step towards a probabilistic well-posedness theory for the cubic nonlinear Schrödinger equation posed on a non-Euclidean compact surface. In this work, we consider randomized initial data distributed according to a Gaussian measure, and we prove that the second Picard iteration does not gain any regularity. This is in sharp contrast with the case of the torus and precludes the construction of strong solutions by using the Bourgain linear-nonlinear decomposition.

In order to make our statement precise, we introduce some notations. We consider the NLS equation on S^2 with a Wick-ordered nonlinearity:

$$i\partial_t u - (-\Delta + 1)u =: |u|^2 u: \quad (t, x) \in \mathbb{R} \times \mathbb{S}^2, \tag{NLS}$$

where Δ is the Laplace–Beltrami operator on \mathbb{S}^2 and the renormalized nonlinearity is¹

$$|u|^2 u: (x) = (|u(x)|^2 - 2||u||^2_{L^2(\mathbb{S}^2)})u(x), \quad x \in \mathbb{S}^2.$$

The norm $L^2(\mathbb{S}^2)$ is associated to the scalar product

$$\langle f \mid g \rangle = \int_{\mathbb{S}^2} f(x) \bar{g}(x) \mathrm{d}\sigma(x),$$

where the Lebesgue measure on \mathbb{S}^2 is normalized such that the sphere has volume 1 (see (2.1)).

On the torus \mathbb{T}^2 , the Wick ordering removes the nonlinear interactions that are not regularizing. Indeed, since the mass is preserved, $||u||_{L^2}^2 u$ is a linear term, which is not regularizing. It is more subtle in the case of the sphere. The reason is that the amplitudes of

¹There is another convention of wick cubic power defined formally as $(|u(x)|^2 - 2\mathbb{E} ||u||^2_{L^2(\mathbb{S}^2)})u(x)$. As $|u(x)|^2 - \mathbb{E} ||u||^2_{L^2(\mathbb{S}^2)}$ is well defined for almost all the initial data we will consider later (distributed according to the Gaussian free field), these two conventions are equivalent.

the eigenfunctions depend in a complicated way on the physical space variable x, whereas, on flat tori, the plane waves have a constant amplitude equal to 1. This point is discussed in Remark 3.4. We stress that if u is a solution to (NLS), then

$$v(t) := e^{it - 2it \|u\|_{L^2}^2} u(t)$$
(1.2)

is a solution to the standard cubic NLS

$$i\partial_t v + \Delta v = |v|^2 v. \tag{1.3}$$

Hence, the regularity properties studied in this article also hold for the solutions of (1.3) thanks to the Gauge transform (1.2). The second Picard iteration of (NLS) reads

$$\mathcal{I}_{\mathbb{S}^2}(t, u_0) = -i \int_0^t e^{i(t-t')(\Delta-1)} \left(: |e^{it'(\Delta-1)} u_0|^2 e^{it'(\Delta-1)} u_0 : \right) dt'$$

Let us turn to the spectral properties of $-\Delta + 1$, as the self-adjoint operator with domain $H^2(\mathbb{S}^2)$. It has a discrete spectrum with eigenvalues

$$\lambda_n^2 = n^2 + n + 1, \quad n \in \mathbb{N}$$

so that $\lambda_n \sim n$ at infinity. The eigenvalues are degenerate in the sense that they have multiplicity 2n + 1, and the eigenspace is spanned by the spherical harmonics of degree n, denoted by $(Y_{n,k})_{|k| \leq n}$. They are the restriction to \mathbb{S}^2 of the harmonic homogeneous polynomials of degree n. We recall some properties in Section 2 and prove a concentration property of many of them on a geodesic circle.

For $n \in \mathbb{N}$, we denote by \mathcal{E}_n the eigenspace

$$\mathscr{E}_n = \ker(-\Delta + 1 - \lambda_n^2 \operatorname{Id}) = \operatorname{span}_{\mathbb{C}} \{\mathbf{b}_{n,k} \mid |k| \le n\},\$$

where $(\mathbf{b}_{n,k})_{|k| \le n}$ is a general orthonormal basis of \mathcal{E}_n . This gives a spectral resolution of $L^2(\mathbb{S}^2)$:

$$L^2(\mathbb{S}^2) = \bigoplus_{n \ge 0} \mathcal{E}_n, \quad \mathcal{E}_n = \pi_n L^2(\mathbb{S}^2),$$

where π_n is the orthogonal projector on \mathcal{E}_n defined by

$$\pi_n = \sum_{|k| \le n} \langle \cdot | \mathbf{b}_{n,k} \rangle \mathbf{b}_{n,k}.$$

The Sobolev norm $H^{s}(\mathbb{S}^{2})$ is equivalent to

$$\left(\sum_{n\geq 0}\lambda_n^{2s}\|\pi_n f\|_{L^2(\mathbb{S}^2)}^2\right)^{\frac{1}{2}}.$$

Given $N \in \mathbb{N}$,

$$P_{\leq N}u = \sum_{n:\,\lambda_n \leq N} \pi_n u$$

is the orthogonal projection on the space spanned by the eigenfunctions with eigenvalues $\leq N$.

We can now define the Gaussian measures. Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, an orthonormal basis $(\mathbf{b}_{n,k})_{n \in \mathbb{N}, |k| \le n}$ made of eigenfunctions of $-\Delta_{\mathbb{S}^2}$, and i.i.d. complex standard Gaussian random variables $(g_{n,k})_{n \in \mathbb{N}, |k| \le n}$, namely,

$$g_{n,k}(\omega) = \frac{1}{\sqrt{2}}(\mathfrak{g}_{n,k}(\omega) + i\mathfrak{h}_{n,k}(\omega)),$$

where $\mathfrak{g}_{n,k}$ and $\mathfrak{h}_{n,k}$ are independent real-valued standard Gaussian random variables on $(\Omega, \mathcal{F}, \mathbb{P})$. Given $\alpha \in \mathbb{R}$, the Gaussian measure μ_{α} is the probability measure induced by the mapping

$$\omega \in \Omega \mapsto \phi_{\alpha}^{\omega} = \sum_{n \ge 0} \lambda_n^{-\alpha} \sum_{|k| \le n} g_{n,k}(\omega) \mathbf{b}_{n,k}.$$
(1.4)

The case $\alpha = 1$ corresponds to the Gaussian free field, which is used to define the formally invariant Gibbs measure. We detail the properties of functions in the support of μ_{α} in Section 3. At this stage, it is important to note that the law of Gaussian measures does not depend on the choice of the orthonormal basis ($\mathbf{b}_{n,k}$). Nevertheless, to evidence some instabilities, we will work with the particular basis made of spherical harmonics ($Y_{n,k}$), presented in Section 2.

Note also that the typical regularity of ϕ^{ω}_{α} is $H^{\alpha-1-0}(\mathbb{S}^2)$ in the sense that, for all $\varepsilon > 0$,

$$\mu_{\alpha}(H^{\alpha-1-\varepsilon}(\mathbb{S}^2)) = 1$$
, but $\mu_{\alpha}(H^{\alpha-1}(\mathbb{S}^2)) = 0$

According to the Weyl's law, this regularity property does not depend on the surface M. In the particular case of the sphere, our main result is that the typical regularity of the second Picard iteration is not better than the regularity of ϕ_{α}^{ω} .

Theorem 1.1. Fix $t \ge 0$ and $\alpha > \frac{1}{2}$. There exist $N_0 \in \mathbb{N}$ and $\eta > 0$ such that, for all $N \ge N_0$,

$$\eta |t| \ln(N)^{\frac{1}{2}} \le \|\mathcal{I}_{\mathbb{S}^2}(t, P_{\le N} \phi^{\omega}_{\alpha})\|_{L^2(\Omega; H^{\alpha - 1}(\mathbb{S}^2))}.$$
(1.5)

We give some comments.

- The logarithmic divergence of the $H^{\alpha-1}(\mathbb{S}^2)$ -norm should be viewed as a lack of regularization for the second Picard iteration, contrasting with the case of the tori, rational or irrational (see Theorem 1.2).
- More precisely, we proved the divergence of quadratic moment of the $H^{\alpha-1}(\mathbb{S}^2)$ norm of the second Picard iteration of the frequency truncated initial data. This is likely to yield the almost sure divergence of this random process, but we do not have a self-contained elementary proof of this result.
- It is not clear whether the threshold $\alpha > \frac{1}{2}$ is technical or not. It appears when we prove that the fully paired interactions are regularizing (so that they do not cancel the divergent term). We refer to [24] for a discussion on the notion of probabilistic criticality, which is heuristically related with this threshold.

Theorem 1.1 indicates that the linear evolution is not a good approximation of the solution, even if the initial data are randomized. In this light, the structure of the probabilistic solution (if it ever exists, in a suitable sense) associated to ϕ_{α}^{ω} cannot be

$$u(t) = e^{it(\Delta-1)}\phi^{\omega}_{\alpha}$$
 + smoother remainder.

In the deterministic setting, the consideration of high-frequency limits in [16] also indicates that on the 2*d* sphere, when $s \leq \frac{1}{4}$, the linear evolution of an initial data in $H^{s}(\mathbb{S}^{2})$ does not approximate well the solution.

While proving Theorem 1.1, we isolate a singular resonant interaction between high and low frequencies. This interaction will play a key role in the probabilistic wellposedness theory we develop in the subsequent work [12]. It is now well established that adapted ansatz in the spirit of paracontrolled calculus allows one to go beyond the linear-nonlinear decomposition of Bourgain [7]. The strategy is to perform an induction on frequency scales and absorb the singular high \times low interactions in a linear operator applied to the high-frequency part of the initial data. We refer, for instance, to the introduction of [10] for discussion on the modern techniques, such as the random averaging operators [24, 38] and the random tensors [23]. The precursors of these methods, for the wave equation, were [9, 29]. Finally, we stress that in both [9, 20] the second Picard iteration is not smoother than the initial data. This was rigorously proved by Oh [34] in the case of the Szegő equation which is covered by the result in [20].

In a second result, proved in Appendix 5, we propose a self-contained proof of the gain of $\frac{1}{2}$ -derivative of the second Picard iteration in the case of irrational tori, for initial data distributed according to the Gaussian free field. We define

$$\mathcal{I}_{\mathbb{T}^2_{\beta}}(t, u_0) = \int_0^t e^{i(t-t')(\Delta_{\beta}-1)} \left(: |e^{it'(\Delta_{\beta}-1)} u_0|^2 e^{it'(\Delta_{\beta}-1)} u_0 : \right) dt',$$

where, for $\beta > 0$, the rescaled Laplacian $-\Delta_{\beta}$ is the rescaled Laplacian:

$$-\Delta_{\beta} := \partial_{x_1}^2 + \beta^2 \partial_{x_2}^2.$$

On the torus, the Gaussian free field is induced by the random variable

$$\phi_1^{\omega}(x) = \sum_{n \in \mathbb{Z}^2} \frac{1}{\langle n \rangle} g_n(\omega) e^{i n \cdot x},$$

where $\langle n \rangle = (1 + Q(n))^{1/2}$, with for $n = (k, m) \in \mathbb{Z}^2$, $Q(n) = k^2 + \beta^2 m^2$. We have that $\phi_1^{\omega} \in H^{0-}(\mathbb{T}^2_{\beta})$ for almost every ω .

Theorem 1.2 (Regularity of the first iteration on tori). For $\alpha = 1$, and for all $\varepsilon > 0$, there exists C > 0 such that, for all $N \in \mathbb{N}$ and $t \in \mathbb{R}$,

$$\left\|\mathcal{I}_{\mathbb{T}_{\beta}^{2}}(t, P_{\leq N}\phi_{1}^{\omega})\right\|_{L^{2}_{\omega}(\Omega; H^{\frac{1}{2}-\varepsilon}(\mathbb{T}^{2}))} \leq C.$$
(1.6)

This result is implicitly contained in [7] (see also [25]). For completeness and to put Theorem 1.1 into context, we give a short proof.

Organization of the article

The proof of Theorem 1.1 in the case of the sphere is given in Section 4 and mostly relies on a quantitative concentration property, which is proven in Section 2.2. We also recall or prove preliminary properties on the spherical harmonics in Section 2, and on Gaussian measures on S^2 in Section 3 together with some preparations on the nonlinearity. Finally, we prove Theorem 1.2 in Section 5.

2. Spherical harmonics

2.1. Spherical harmonics

The polar (colatitudinal) coordinate is $\theta \in (0, \pi)$, the azimuthal (longitudinal) coordinate is $\varphi \in (0, 2\pi)$, and $\rho \in \mathbb{R}^+$ is the radial distance so that $\mathbb{S}^2 = \{x \in \mathbb{R}^3 \mid |\rho| = 1\}$. We have

$$(x_1, x_2, x_3) = (\rho \sin(\theta) \cos(\varphi), \rho \sin(\theta) \sin(\varphi), \rho \cos(\theta)).$$

With this coordinates system, the Lebesgue measure is

$$d\sigma = \frac{1}{4\pi} \sin(\theta) d\theta d\varphi, \qquad (2.1)$$

and the Hermitian scalar product is

$$\langle f \mid g \rangle_{L^2(\mathbb{S}^2)} = \frac{1}{4\pi} \int_0^{2\pi} \int_0^{\pi} f(\theta, \varphi) \overline{g(\theta, \varphi)} \sin(\theta) d\theta d\varphi.$$

With this normalization, S^2 has volume 1. Note that this coordinate system is singular on the axis (Ox_3). The (shifted) Laplace–Beltrami operator on the sphere reads

$$-\Delta_{\mathbb{S}^2} + 1 = -\frac{1}{\sin^2\theta} \frac{\partial^2}{\partial\varphi^2} - \frac{1}{\sin\theta} \frac{\partial}{\partial\theta} \left(\sin\theta \frac{\partial}{\partial\theta}\right) + 1.$$
(2.2)

It is a self-adjoint operator with domain $H^2(\mathbb{S}^2)$, and it has discrete spectrum. The eigenvalues are

$$\lambda_n^2 = n^2 + n + 1, \quad n \in \mathbb{N},$$

with multiplicity 2n + 1. The normalized spherical harmonics, defined as the restrictions to \mathbb{S}^2 of the harmonic homogeneous polynomials, form a particular orthonormal basis of eigenfunction. For $n \in \mathbb{N}$ and $k \in \{-n, \ldots, n\}$, we denote by $Y_{n,k}$ the spherical harmonics of degree *n* and order *k*. In spherical coordinates, we have

$$Y_{n,k}(\theta,\varphi) = e^{ik\varphi} v_{n,k}(\theta), \quad v_{n,k}(\theta) = c_{n,k} L_{n,k}(\cos(\theta)), \tag{2.3}$$

where $L_{n,k}(\cos(\theta))$ is the associated Legendre function of degree *n* and order *k*, and $c_{n,k}$ is a normalization constant

$$c_{n,k} = \sqrt{(2n+1)\frac{(n-k)!}{(n+k)!}}$$

Proposition 2.1. Spherical harmonics $(Y_{n,k})_{1 \le n, |k| \le n}$ form an orthonormal basis of $L^2(\mathbb{S}^2)$ made of eigenfunctions of the Laplace operator on \mathbb{S}^2 . They satisfy

$$(-\Delta_{\mathbb{S}^2}+1)Y_{n,k}=\lambda_n^2Y_{n,k}, \quad n\in\mathbb{N}^*,\ k\in\{-n,\ldots,n\}$$

with $\lambda_n^2 = n^2 + n + 1$.

We refer to [36] and [37, Chapter IV] for a detailed analysis of the spherical harmonics.

2.2. Concentration of spherical harmonics with high order

It follows from the expression of the Laplace operator on the sphere (2.2) that $v_{n,k}$ is solution on $(0, \pi)$ to

$$-\left(\sin(\theta)\frac{\mathrm{d}}{\mathrm{d}\theta}\right)^2 v_{n,k}(\theta) + (k^2 - n(n+1)\sin^2(\theta))v_{n,k}(\theta) = 0.$$
(2.4)

The next proposition claims that in the high-frequency regime $n \to \infty$ a large family of spherical harmonics with high weights (when |k| is close to *n*) concentrate their mass near the equator (corresponding to the region $\theta = \frac{\pi}{2}$). Given $\delta \in (0, 1)$, we denote

$$\mathcal{C}_{\delta} := \left\{ x = (\theta, \varphi) \in \mathbb{S}^2 \mid \delta < |\cos(\theta)| \right\}.$$
(2.5)

Proposition 2.2. For all $\delta > 0$, $n \ge 1$, and $k \in \mathbb{Z}$ such that

$$n(n+1)(1-\delta^2) \le k^2 \le n^2, \tag{2.6}$$

we have

$$\|\mathbf{1}_{\mathcal{C}_{2\delta}}(x)Y_{n,k}(x)\|_{L^2_x(\mathbb{S}^2)} \lesssim_{\delta} \frac{1}{n}$$

The proof relies on utilizing semiclassical functional calculus in the high-frequency regime to quantitatively exploit the ellipticity of equation (2.4) away from the equator. A different approach, based on ordinary differential equations, can be found in [26, Section 3], where families of spherical harmonics with high weight $|k| \sim n$ are used to saturate certain L^p spectral cluster bounds.

Proof. In order to use semiclassical functional calculus and prove this proposition, we will extend $v_{n,k}$ to a function defined on the whole real line \mathbb{R} . For this purpose, we make the change of variable

$$f: \quad (0,\pi) \to \mathbb{R}$$
$$\theta \mapsto \tanh^{-1}(\cos(\theta)).$$

Note that f is a C^{∞} -diffeomorphism and that we have the identity: for all $\theta \in (0, \pi)$,

$$\sin^2(\theta) = 1 - \cos^2(\theta) = 1 - \tanh^2(f(\theta)) = \frac{1}{(\cosh \circ f(\theta))^2}$$

This yields

$$f'(\theta) = -\frac{1}{\sin(\theta)}, \quad (f^{-1})'(y) = -\frac{1}{\cosh(y)},$$
 (2.7)

which in turn implies that

$$\int_{-\infty}^{\infty} \frac{1}{\cosh(y)^2} |v_{n,k} \circ f^{-1}(y)|^2 dy = \int_0^{\pi} |v_{n,k}(\theta)|^2 \sin(\theta) d\theta = \frac{1}{2\pi}.$$
 (2.8)

We set

$$\tilde{v}_{n,k}(y) = \frac{(2\pi)^{\frac{1}{2}}}{\cosh(y)} v_{n,k} \circ f^{-1}(y), \quad y \in \mathbb{R},$$

so that

$$\int_{-\infty}^{\infty} |\tilde{v}_{n,k}(y)|^2 dy = 1.$$

In what follows, we may abuse notations and write $\theta = f^{-1}(y)$ for $y \in \mathbb{R}$. We have from the chain rule and from (2.7) that

$$\frac{\mathrm{d}}{\mathrm{d}y}\tilde{v}_{n,k}(y) = -\tanh(y)\tilde{v}_{n,k}(y) - \frac{(2\pi)^{\frac{1}{2}}}{\cosh(y)}\sin(\theta)\frac{\mathrm{d}}{\mathrm{d}\theta}v_{n,k}(\theta).$$
(2.9)

We deduce from this that

$$\left\|\frac{\mathrm{d}}{\mathrm{d}y}\tilde{v}_{n,k}\right\|_{L^2(\mathbb{R})} \lesssim 1.$$
(2.10)

Indeed,

$$\|\tanh \tilde{v}_{n,k}\|_{L^2(\mathbb{R})} \le \|\tilde{v}_{n,k}\|_{L^2(\mathbb{R})} \le 1,$$

and

$$\left\|\frac{1}{\cosh(y)}\sin(\theta)\frac{\mathrm{d}}{\mathrm{d}\theta}v_{n,k}(\theta)\right\|_{L^{2}(\mathbb{R})}^{2} = \int_{\mathbb{R}}\sin^{2}(\theta)\frac{1}{\cosh(y)}|v_{n,k}'\circ f^{-1}(y)|^{2}|(f^{-1})'(y)|\mathrm{d}y$$
$$= \int_{0}^{\pi}\sin^{3}(\theta)|v_{n,k}'(\theta)|^{2}\mathrm{d}\theta \lesssim 1.$$

Moreover, we deduce from (2.9) that

$$\frac{\mathrm{d}^2}{\mathrm{d}y^2}\tilde{v}_{n,k}(y) = (\tanh^2(y) - 1)\tilde{v}_{n,k}(y) - \tanh(y)\frac{\mathrm{d}}{\mathrm{d}y}\tilde{v}_{n,k}(y) + \frac{(2\pi)^{\frac{1}{2}}}{\cosh(y)} \bigg(2\cos(\theta)\sin(\theta)\frac{\mathrm{d}}{\mathrm{d}\theta}v_{n,k}(\theta) + \sin^2(\theta)\frac{\mathrm{d}^2}{\mathrm{d}\theta^2}v_{n,k}(\theta)\bigg).$$

Plugging in expression (2.9), we deduce that

$$\frac{\mathrm{d}^2}{\mathrm{d}y^2}\tilde{v}_{n,k}(y) = -\left(1 + 2\cos(\theta)\frac{\mathrm{d}}{\mathrm{d}y}\right)\tilde{v}_{n,k}(y) + \frac{(2\pi)^{\frac{1}{2}}}{\cosh(y)}\left(\sin(\theta)\frac{\mathrm{d}}{\mathrm{d}\theta}\right)^2 v_{n,k}(\theta).$$

Therefore, we conclude form (2.4) that $\tilde{v}_{n,k} \in C^{\infty}(\mathbb{R})$ solves

$$\frac{d^2}{dy^2}\tilde{v}_{n,k} + \left(1 + 2\cos(\theta)\frac{d}{dy}\right)\tilde{v}_{n,k} - (k^2 - n(n+1)\sin^2(\theta))\tilde{v}_{n,k} = 0.$$
(2.11)

We can now introduce the semiclassical parameter in the high-frequency regime $n \rightarrow \infty$:

$$h := (n(n+1))^{-\frac{1}{2}}, \quad h \sim_{n \to \infty} n^{-1}.$$

For the rest of the proof, we denote

 $\alpha := |k|h.$

This parameter should not be confused with the parameter α in definition (1.4) of the initial data, which is not involved in this proof. Multiplying equation (2.11) by h^2 gives

$$-h^{2}\frac{\mathrm{d}^{2}}{\mathrm{d}y^{2}}\tilde{v}_{n,k}(y) + (\alpha^{2} - \sin^{2}(\theta(y)))\tilde{v}_{n,k}(y) - 2h^{2}\cos(\theta(y))\frac{\mathrm{d}}{\mathrm{d}y}\tilde{v}_{n,k}(y) - h^{2} = 0.$$
(2.12)

We reformulate (2.12) as follows:

(2.12)
$$\iff$$
 $P_{\alpha}(y,hD_y)(\tilde{v}_{n,k}) = 2h^2\cos(\theta(y))\frac{d}{dy}\tilde{v}_{n,k}(y) + h^2,$

where P_{α} is a differential operator of order 2 with symbol

$$p_{\alpha}(y,\xi) = \xi^2 + \alpha^2 - \sin^2(\theta(y)).$$

We deduce from (2.10) that

$$\|\mathbf{P}_{\alpha}(y,hD_{y})\tilde{v}_{n,k}\|_{L^{2}(\mathbb{R})} \lesssim h^{2}.$$
(2.13)

Consider

Char(P_{α}) = { $(y, \xi) \in \mathbb{R}^2 | p_{\alpha}(y, \xi) = 0$ } = { $(y, \xi) \in \mathbb{R}^2 | \xi^2 + \alpha^2 - \sin^2(\theta(y)) = 0$ }, and set, for $\delta \in (0, 1)$,

$$\overline{\mathcal{C}_{\delta}} := \left\{ y \in \mathbb{R} \mid \delta < |\cos(\theta(y))| \right\}.$$
(2.14)

Observe that, for all $\delta \in (0, 1)$,

$$\sqrt{1-\delta^2} \leq \alpha \implies \operatorname{Char}(\mathbf{P}_{\alpha}) \subset \mathbb{R} \setminus \widetilde{\mathbb{C}_{\delta}}.$$

Indeed,

$$\begin{cases} \sqrt{1-\delta^2} \le \alpha, \\ \xi^2 + \alpha^2 - \sin^2(\theta) = 0 \end{cases} \implies 1-\delta^2 \le \sin^2(\theta(y)) \implies |\cos(\theta(y))| \le \delta. \end{cases}$$

Hence, in the regime where $\sqrt{1-\delta^2} \le \alpha$, Char(P_{α}) concentrates near the equator as δ goes to 0. By the use of semiclassical functional calculus, we will deduce from this that $\tilde{v}_{n,k}$ also concentrates its mass near the equator in this regime.

Lemma 2.3. Let $\delta \in (0, 1)$, and suppose that $\sqrt{1 - \delta^2} \leq \alpha$. Then, for all $\chi \in C_c^{\infty}(\mathbb{R})$ such that

$$\chi \equiv 1 \text{ on } \widetilde{\mathbb{C}}_{2\delta}, \quad \text{supp } \chi \subset \widetilde{\mathbb{C}}_{\frac{3}{2}\delta},$$

where $\widetilde{\mathbb{C}}_{\delta}$ is defined in (2.14), there exists a bounded operator $Q_{\alpha,h}$ such that, for all $h \in (0, 1)$,

$$\|\mathbf{Q}_{\alpha,h}\|_{L^2(\mathbb{R})\to L^2(\mathbb{R})}\lesssim_{\delta} 1,$$

and

$$\|\mathbf{Q}_{\alpha,h}\circ\mathbf{P}_{\alpha}(y,hD_{y})-\chi(y)\|_{L^{2}(\mathbb{R})\to L^{2}(\mathbb{R})}\lesssim_{\delta}h$$

Proof. Fix $\delta \in (0, 1)$ and suppose that $\sqrt{1 - \delta^2} \le \alpha$. The key point is that $P_{\alpha}(y, hD_y)$ is elliptic in the region $\widetilde{C}_{2\delta}$, and the proof follows from the standard parametrix construction of elliptic operators. Since we only need to invert $P_{\alpha}(y, hD_y)$ in the elliptic region up to order 1 in *h*, we propose a self-contained proof.

We introduce the symbol

$$q(y,\xi;\alpha) := \frac{\chi(y)}{p_{\alpha}(y,\xi)}.$$

Recall that for all $y \in \text{supp}(\chi), \xi \in \mathbb{R}$, and $\sqrt{1-\delta^2} \le \alpha$ we have

$$p_{\alpha}(y,\xi) = \xi^{2} + \alpha^{2} - \sin^{2}(\theta(y))$$

$$\geq 1 - \delta^{2} - \sin^{2}(\theta(y))$$

$$= \cos^{2}(\theta(y)) - \delta^{2}$$

$$\geq \delta^{2}.$$
(2.15)

The semiclassical pseudodifferential operator $Q_{\alpha}(y, hD_y)$ associated with *q* is defined via its Schwartz kernel as follows:

$$Q_{\alpha}(y,hD_{y})[f](y) = \int_{\mathbb{R}} K(y,z)f(z)dz, \quad K(y,z) = \frac{1}{2\pi h} \int_{\mathbb{R}} e^{\frac{i(y-z)\xi}{h}} q(y,\xi;\alpha)d\xi.$$

Observe that

$$(y-z)e^{\frac{i(y-z)\xi}{\hbar}} = -i\hbar\partial_{\xi}\left(e^{\frac{i(y-z)\xi}{\hbar}}\right).$$
(2.16)

Hence, integrating by parts twice yields that

$$(y-z)^2 K(y,z) = -\frac{h}{2\pi} \int_{\mathbb{R}} e^{\frac{i(y-z)\xi}{h}} \,\partial_{\xi}^2 q(y,\xi;\alpha) \mathrm{d}\xi,$$

It follows from (2.15) that, for all $k \in \mathbb{N}$, α , δ such that $\sqrt{1 - \delta^2} < \alpha \le 1$ and for all $\xi \in \mathbb{R}$,

$$\sup_{y\in\mathbb{R}}|\partial_{\xi}^{k}q(y,\xi,\alpha)|\lesssim_{k}\delta^{-2k}\langle\xi\rangle^{-2k}.$$

Hence,

$$\sup_{(y,z)\in\mathbb{R}^2} (y-z)^2 |K(y,z)| \lesssim h\delta^{-4} \bigg(\int_{\mathbb{R}} \langle \xi \rangle^{-4} \mathrm{d}\xi \bigg) \lesssim h\delta^{-4}.$$

Moreover, we have the trivial bound

$$\sup_{(y,z)\in\mathbb{R}^2} |K(y,z)| \le h^{-1}\delta^{-2} \bigg(\int_{\mathbb{R}} \langle \xi \rangle^{-2} \mathrm{d}\xi \bigg) \lesssim h^{-1}\delta^{-4}.$$

Therefore,

$$\begin{split} \sup_{z \in \mathbb{R}} \int_{\mathbb{R}} |K(y, z)| \mathrm{d}y &\lesssim \delta^{-4} \sup_{z \in \mathbb{R}} \int_{\mathbb{R}} \min\left(\frac{h}{(y - z)^2}, \frac{1}{h}\right) \mathrm{d}y \\ &\lesssim \delta^{-4} \left(\int_{|s| \le h} \frac{1}{h} \mathrm{d}s + \int_{|s| > h} \frac{h}{s^2} \mathrm{d}s \right) \\ &\lesssim \delta^{-4}. \end{split}$$

Since the computations are the same when changing the role of y and z, we prove that when $\sqrt{1-\delta^2} \le \alpha$, then

$$\sup_{y\in\mathbb{R}}\int_{\mathbb{R}}|K(y,z)|\mathrm{d} z+\sup_{z\in\mathbb{R}}\int_{\mathbb{R}}|K(y,z)|\mathrm{d} y\lesssim\delta^{-4},$$

uniformly in $h \in (0, 1)$. It follows from Schur's test that $Q_{\alpha}(y, hD_y)$ is bounded on $L^2(\mathbb{R})$, uniformly in $h \in (0, 1)$, with

$$\|\mathbf{Q}_{\alpha}(y,hD_{y})\|_{L^{2}(\mathbb{R})\to L^{2}(\mathbb{R})} \lesssim \delta^{-4}.$$
(2.17)

The Schwartz kernel of the operator

$$\mathbf{R}_{\alpha}(y,hD_{y}) := \mathbf{Q}_{\alpha}(y,hD_{y}) \circ \mathbf{P}_{\alpha}(y,hD_{y}) - \chi(y)$$

is given by

$$\mathcal{K}(y,z) = (\phi(z) - \phi(y))\chi(y) \int_{\mathbb{R}} e^{\frac{i(y-z)\xi}{h}} \frac{1}{p_{\alpha}(y,\xi)} \frac{d\xi}{2\pi h},$$

where

$$\phi(y) := \sin^2(\theta(y)).$$

We write

$$\varphi(y,z) := \int_0^1 \phi'(tz + (1-t)y) \mathrm{d}t$$

so that

$$\phi(y) - \phi(z) = (y - z)\varphi(y, z).$$

It follows from (2.16) and integration by parts that

$$\mathcal{K}(y,z) = -i\varphi(y,z)\chi(y)\int_{\mathbb{R}}\partial_{\xi}\left(\frac{1}{p_{\alpha}(y,\xi)}\right)e^{\frac{i(y-z)\xi}{\hbar}}\frac{d\xi}{2\pi}.$$

We proceed with the estimate on the kernel of Q_{α} , and we integrate by parts twice (using (2.16)) to get

$$|\mathcal{K}(y,z)| \lesssim \delta^{-6} \min\left(1, \frac{h^2}{|y-z|^2}\right).$$

We deduce

$$\sup_{z \in \mathbb{R}} \int_{\mathbb{R}} |\mathcal{K}(y, z)| \mathrm{d}y + \sup_{y \in \mathbb{R}} \int_{\mathbb{R}} |\mathcal{K}(y, z)| \mathrm{d}z \lesssim \delta^{-6}h,$$

and we conclude from Schur's test that

$$\|\mathbf{R}_{\alpha}(y, hD_{y})\|_{L^{2} \to L^{2}} \lesssim \delta^{-6}h.$$

This concludes the proof of Lemma 2.3 with an implicit constant $\lesssim \delta^{-6}$.

We now complete the proof of Proposition 2.2 as follows. We have from Lemma 2.3, from (2.17), and from (2.13) that

$$\|\chi \tilde{v}_{n,k}\|_{L^2(\mathbb{R})} = \|\mathbf{Q}_{\alpha}(y,hD_y)\mathbf{P}_{\alpha}(y,hD_y)\tilde{v}_{n,k} - \mathbf{R}_{\alpha}(y,hD_y)\tilde{v}_{n,k}\|_{L^2(\mathbb{R})} \lesssim_{\delta} h.$$

Hence, according to (2.8),

$$\|Y_{n,k}\|_{L^2(\mathcal{C}_{2\delta})} = \|\tilde{v}_{n,k}\|_{L^2(\widetilde{\mathcal{C}_{2\delta}})} \le \|\chi\tilde{v}_{n,k}\|_{L^2(\mathbb{R})} \le C_{\delta}h,$$

which completes the proof of Proposition 2.2.

3. Preparations

3.1. Gaussian measures

Recall that the Gaussian measure μ_{α} of parameter $\alpha \in \mathbb{R}$ is defined in (1.4). To present some properties of the random functions $\phi_{\alpha}(\omega)$ in the support of μ_{α} , we need to first recall the local Weyl law for $-\Delta_{\mathbb{S}^2}$.

Lemma 3.1 (Local Weyl's law). For every $n \in \mathbb{N}$, every orthonormal basis $(\mathbf{b}_{n,k})$ of \mathcal{E}_n , and every $x \in \mathbb{S}^2$, we have

$$\frac{1}{2n+1} \sum_{|k| \le n} |\mathbf{b}_{n,k}(x)|^2 = 1.$$
(3.1)

We detail the very concise proof of this lemma.

Proof. Given $n \in \mathbb{N}$ and $(\mathbf{b}_{n,k})_{|k| \le n}$ an orthonormal basis of \mathcal{E}_n , the integral kernel of the orthogonal projector π_n onto \mathcal{E}_n is

$$K_n(x, y) = \sum_{|k| \le n} \mathbf{b}_{n,k}(x) \overline{\mathbf{b}_{n,k}(y)}, \quad (x, y) \in (\mathbb{S}^2)^2.$$

It follows from the rotational invariance of the operator $-\Delta_{\mathbb{S}^2}$ that, for all $R \in SO_3$,

$$\pi_n \circ R = R \circ \pi_n$$

which in turns implies that, for every $(x, y) \in (\mathbb{S}^2)^2$ and $R \in SO_3$,

$$K_n(Rx, y) = K_n(x, R^{-1}y)$$

In particular, for all $x \in \mathbb{S}^2$,

$$K_n(Rx, Rx) = K_n(x, x).$$

Since SO_3 acts transitively on \mathbb{S}^2 , we deduce that

$$x \in \mathbb{S}^2 \mapsto K_n(x, x) = \sum_{|k| \le n} |\mathbf{b}_{n,k}(x)|^2$$

is a constant function. Integrating over $x \in S^2$ and using the assumption that the spherical harmonics are normalized, we conclude the proof of the identity (3.1).

Let us now reorganize the terms in series (1.4), which defines the Gaussian measure μ_{α} . Given an orthonormal basis (**b**_{*n*,*k*}), we group the eigenfunctions in clusters of same eigenvalue, with multiplicity 2n + 1, and write

$$\phi_{\alpha}^{\omega} = \sum_{n \ge 0} \lambda_n^{-\alpha} \sum_{|k| \le n} g_{n,k}(\omega) \mathbf{b}_{n,k} = \sum_{n \ge 0} \widetilde{\lambda}_n^{-(\alpha - \frac{1}{2})} e_n^{\omega},$$

where, for $n \in \mathbb{N}$,

$$e_n^{\omega} = \frac{1}{\sqrt{2n+1}} \sum_{|k| \le n} g_{n,k}(\omega) \mathbf{b}_{n,k}, \qquad (3.2)$$

and $\tilde{\lambda}_n = \lambda_n (\frac{2n+1}{\lambda_n})^{\frac{1}{2}}$. Since λ_n and $\tilde{\lambda}_n$ have the same asymptotic up to a factor 2, we abuse notations and we keep writing

$$\lambda_n \approx \widetilde{\lambda}_n$$

For every $n \ge 0$, e_n^{ω} can be seen as a Gaussian vector on \mathcal{E}_n , with

$$\mathbb{E}(e_n^{\omega}) = 0, \quad \operatorname{Cov}(e_n^{\omega}) = \frac{1}{2n+1} \operatorname{Id}_{\mathcal{E}_n},$$

where $\text{Cov}(e_n^{\omega})$ is the covariance matrix of the Gaussian vector e_n^{ω} . We recall in the next lemma that the law of a complex standard Gaussian vector is invariant under the action of the unitary group.

Lemma 3.2. For all $X \sim \mathcal{N}_{\mathbb{C}}(0, \frac{1}{2n+1} \mathrm{Id}_{\mathcal{E}_n})$ and $A \in U(\mathcal{E}_n)$, the unitary group of \mathcal{E}_n , we have

$$\mathcal{L}(AX) = \mathcal{L}(X).$$

Proof. For *Y* a random variable, we denote by φ_Y its characteristic function. For all $\xi \in \mathcal{E}_n$,

$$\varphi_{AX}(\xi) = \mathbb{E}[e^{i\langle\xi|AX\rangle}] = \mathbb{E}[e^{i\langle A^*\xi|X\rangle}] = \varphi_X(A^*\xi)$$
$$= e^{-\frac{1}{2n+1}\|A^*\xi\|_{L^2}^2} = e^{-\frac{1}{2n+1}\|\xi\|_{L^2}^2} = \varphi_X(\xi).$$

This proves the lemma.

A consequence of the above lemma is that the law of e_n^{ω} —and therefore the law μ_{α} —does not depend on the choice of the orthonormal eigenbasis of \mathcal{E}_n . Alternatively, we could fix x and view $e_n^{\omega}(x)$ as a complex standard Gaussian random variable. Indeed, $\mathbb{E}[e_n^{\omega}(x)] = 0$ and, according to (3.1),

$$\operatorname{Var}(e_n^{\omega}(x)) = \frac{1}{2n+1} \sum_{|k| \le n} |\mathbf{b}_{n,k}(x)|^2 = 1.$$

In particular, the law of $e_n^{\omega}(x)$ does not depend on the point x on the sphere.

Let us now recall the elementary property of complex standard Gaussian variables: for all n, k, we have

$$\mathbb{E}[g_{n,k}] = \mathbb{E}[g_{n,k}^2] = \mathbb{E}[g_{n,k}|g_{n,k}|^2] = 0$$

In particular, we deduce from the above identities and from the definition

$$e_n^{\omega}(x) = (2n+1)^{-\frac{1}{2}} \sum_{|k| \le n} g_{n,k}(\omega) \mathbf{b}_{n,k}(x)$$

that, for fixed $n \ge 0$ and $(x, y) \in (\mathbb{S}^2)^2$,

$$\mathbb{E}[e_n^{\omega}(x)|e_n^{\omega}(y)|^2] = \mathbb{E}[e_n^{\omega}(x)||e_n^{\omega}||_{L^2(\mathbb{S}^2)}^2] = 0.$$
(3.3)

For fixed $x \in \mathbb{S}^2$, we will use the fact that $e_n^{\omega}(x)$ is \mathcal{B}_n -measurable, where \mathcal{B}_n is the σ algebra generated by $(g_{n,k})_{|k| \leq n}$. The independence of \mathcal{B}_n from $\mathcal{B}_{n'}$ whenever $n \neq n'$ plays an important role in the multilinear estimates involving the random functions e_n^{ω} .

3.2. L^p -bounds

We recall the eigenfunctions estimates due to Sogge [36]. There exists C > 0 such that, for all $n \ge 0$ and $f \in L^2(\mathbb{S}^2)$,

$$\|\pi_n f\|_{L^p(\mathbb{S}^2)} \le C \|\pi_n f\|_{L^2(\mathbb{S}^2)} \begin{cases} \lambda_n^{\frac{1}{2}(\frac{1}{2} - \frac{1}{p})}, & 2 \le p \le 6, \\ \lambda_n^{\frac{1}{2} - \frac{2}{p}}, & 6 \le p \le \infty. \end{cases}$$
(3.4)

In contrast, we prove that the $L^p(\mathbb{S}^2)$ -norm of a $L^2(\mathbb{S}^2)$ -normalized Gaussian spherical harmonic e_n^{ω} has its moment of order p uniformly bounded in n.

Lemma 3.3. There exists C > 0 such that, for all $p \ge 2$ and $n \ge 0$,

$$\|e_n^{\omega}\|_{L^p_{\omega}(\Omega; L^p_x(\mathbb{S}^2))} \le C\sqrt{p}.$$
(3.5)

Proof. We deduce from Lemma 3.1 and from Khintchine's inequality that there exists C > 0 such that, for all $n \ge 0$, $x \in \mathbb{S}^2$, $e_n^{\omega}(x)$, and $p \ge 2$,

$$\|e_n^{\omega}(x)\|_{L^p_{\omega}(\Omega)} \le C\sqrt{p}.$$

Subsequently, we deduce from Fubini's theorem that

$$\mathbb{E}\left[\left\|e_{n}^{\omega}\right\|_{L_{x}^{p}(\mathbb{S}^{2})}^{p}\right]^{\frac{1}{p}} = \left(\int_{\mathbb{S}^{2}}\left\|e_{n}^{\omega}(x)\right\|_{L_{\omega}^{p}}^{p}\mathrm{d}\sigma(x)\right)^{\frac{1}{p}} \leq C\operatorname{vol}(\mathbb{S}^{2})^{\frac{1}{p}}\sqrt{p} = C\sqrt{p},$$

since we normalized the Lebesgue measure such that $vol(S^2) = 1$. This concludes the proof of Lemma 3.3.

In [17], the authors prove some more precise large deviation estimates, using the concentration of the measure.

3.3. Decomposition of the nonlinearity

We decompose the Wick cubic power into three parts:

$$|u|^{2}u := |u|^{2}u - 2||u||_{L^{2}(\mathbb{S}^{2})}^{2}u$$
$$= \mathcal{N}_{1}(u) + \mathcal{N}_{2}(u) + \mathcal{N}_{3}(u),$$

where

$$\mathcal{N}_{1}(u) = \sum_{n_{1},n_{2},n_{3}} \mathbf{1}_{n_{2} \neq n_{1},n_{2} \neq n_{3}} \pi_{n_{1}} u \pi_{n_{2}} \bar{u} \pi_{n_{3}} u,$$

$$\mathcal{N}_{2}(u) = 2 \sum_{n_{1},n_{2}} \mathbf{1}_{n_{1} \neq n_{2}} \left(|\pi_{n_{2}}u|^{2} - ||\pi_{n_{2}}u||^{2}_{L^{2}(\mathbb{S}^{2})} \right) \pi_{n_{1}} u,$$

$$\mathcal{N}_{3}(u) = \sum_{n} |\pi_{n}u|^{2} \pi_{n}u - 2 ||\pi_{n}u||^{2}_{L^{2}(\mathbb{S}^{2})} \pi_{n}u.$$

We isolate the resonant interaction from \mathcal{N}_2 :

$$\mathcal{N}_{2,\mathrm{res}}(u) := 2 \sum_{n_1,n_2} \mathbf{1}_{n_1 \neq n_2} \pi_{n_1} \big(\pi_{n_1} u (|\pi_{n_2} u|^2 - ||\pi_{n_2} u||^2_{L^2(\mathbb{S}^2)}) \big).$$

As we will see in the next section, the term $\mathcal{N}_{2,res}$ is responsible for the divergence of the second Picard iteration (1.5) claimed in our main theorem.

Remark 3.4. In the case of the torus \mathbb{T}^2 , where the plane waves $e^{in \cdot x}$ have constant amplitude, we have that, for all $n \in \mathbb{Z}^2$ and $x \in \mathbb{T}^2$,

$$|e^{in \cdot x}|^2 - ||e^{in \cdot x}||^2_{L^2(\mathbb{T}^d)} = 0$$

so that the term N_2 does not exist after the Wick ordering. On the sphere, however, the variable x in the physical space has a role to play and pointwise in x, the *Wick square*

$$|e_n^{\omega}(x)|^2 - ||e_n^{\omega}||^2_{L^2(\mathbb{S})}$$

has no reason to vanish. Still, it does vanish on average (both in space and in probability measure):

$$\int_{\mathbb{S}^2} |e_n^{\omega}(x)|^2 - \|e_n^{\omega}\|_{L^2(\mathbb{S})}^2 \mathrm{d}\sigma(x) = \int_{\Omega} |e_n^{\omega}(x)|^2 - \|e_n^{\omega}\|_{L^2(\mathbb{S})}^2 \mathrm{d}\mathbb{P}(\omega) = 0.$$

3.4. The second Picard iteration

Fix $\alpha \in \mathbb{R}$, and recall that the Gaussian measure μ_{α} is induced by the random variable

$$\phi_{\alpha}^{\omega} = \sum_{n \ge 0} \lambda_n^{-(\alpha - \frac{1}{2})} e_n^{\omega},$$

where the Gaussian spherical harmonics e_n^{ω} are defined in (3.2). We denote by $u^{\omega}(t)$ the linear evolution

$$u^{\omega}(t,x) = e^{it(\Delta-1)} \phi^{\omega}_{\alpha} = \sum_{n \ge 0} e^{-it\lambda_n^2} \lambda_n^{-(\alpha-\frac{1}{2})} e^{\omega}_n(x).$$

In order to make sense to the series in $L^{\infty}(\mathbb{R}; H^{s}(\mathbb{S}^{2}))$ for any $s \in \mathbb{R}$, we also consider the finite-dimensional approximation

$$P_{\leq N}u^{\omega}(t,x) = \sum_{n : \lambda_n \leq N} e^{-it\lambda_n^2} \lambda_n^{-(\alpha - \frac{1}{2})} e_n^{\omega}(x).$$

For $N \in \mathbb{N}$, dyadic, set

$$\Gamma_N = \left\{ \vec{n} = (n_0, n_1, n_2, n_3) \in \mathbb{N}^4 \mid n_i \le N \text{ for } i \in \{1, 2, 3\} \right\}$$

and

$$\Gamma_N^{(1)} := \{ \vec{n} \in \Gamma_N \mid n_2 \neq n_1, n_2 \neq n_3 \},\$$

$$\Gamma_N^{(2)} := \{ \vec{n} \in \Gamma_N \mid n_2 = n_3, n_2 \neq n_1 \},\$$

$$\Gamma_N^{(3)} := \{ \vec{n} \in \Gamma_N \mid n_1 = n_2 = n_3 \}.$$

For $\vec{n} \in \Gamma_N$, we denote the resonance function

$$\Omega(\vec{n}) = \lambda_{n_0}^2 - \lambda_{n_1}^2 + \lambda_{n_2}^2 - \lambda_{n_3}^2.$$

Then, we have

$$\begin{split} \mathcal{I}_{\mathbb{S}^{2}}(t, P_{\leq N}u^{\omega}) &= -i \int_{0}^{t} e^{i(t-t')(\Delta-1)} \mathcal{N}(P_{\leq N}u^{\omega}(t')) dt' \\ &= -i \int_{0}^{t} e^{i(t-t')(\Delta-1)} \big(\mathcal{N}_{1}(P_{\leq N}u^{\omega}(t')) \\ &+ \mathcal{N}_{2}(P_{\leq N}u^{\omega}(t')) + \mathcal{N}_{3}(P_{\leq N}u^{\omega}(t')) \big) dt' \\ &= \mathrm{I} + \mathrm{II} + \mathrm{III}, \end{split}$$
(3.6)

where

$$I_{N}(t,x) = \sum_{\vec{n}\in\Gamma_{N}^{(1)}} e^{-it\lambda_{n_{0}}^{2}} \left(\frac{e^{-it\Omega(\vec{n})}-1}{\Omega(\vec{n})}\right) (\lambda_{n_{1}}\lambda_{n_{2}}\lambda_{n_{3}})^{-(\alpha-\frac{1}{2})} \pi_{n_{0}} \left(e_{n_{1}}^{\omega} \bar{e}_{n_{2}}^{\omega} e_{n_{3}}^{\omega}\right), (3.7)$$

$$\Pi_{N}(t,x) = \sum_{\vec{n}\in\Gamma_{N}^{(2)}} e^{-it\lambda_{n_{0}}^{2}} \left(\frac{e^{-it\Omega(\vec{n})}-1}{\Omega(\vec{n})}\right) (\lambda_{n_{1}}\lambda_{n_{2}}^{2})^{-(\alpha-\frac{1}{2})} \pi_{n_{0}} \times \left(e_{n_{1}}^{\omega} (|e_{n_{2}}^{\omega}|^{2} - ||e_{n_{2}}^{\omega}||_{L^{2}\mathbb{S}^{2}}^{2})\right), \qquad (3.8)$$

$$\Pi_{N}(t,x) = \sum_{\vec{n}\in\Gamma^{(3)}} e^{-it\lambda_{n_{0}}^{2}} \left(\frac{e^{-it\Omega(\vec{n})}-1}{\Omega(\vec{n})}\right) \lambda_{n_{1}}^{-3(\alpha-\frac{1}{2})} \pi_{n_{0}} \left(|e_{n_{1}}^{\omega}|^{2} e_{n_{1}}^{\omega}\right)$$

$$-2\sum_{n\geq 0} e^{-it\lambda_{n_0}^2} \lambda_n^{-3(\alpha-\frac{1}{2})} \|e_n^{\omega}\|_{L^2(\mathbb{S}^2)}^2 e_n^{\omega}.$$
(3.9)

4. Proof of the main theorem

Given $\alpha \in \mathbb{R}$, our goal is to show a logarithmic divergence in N of the $H^{\alpha-1}(\mathbb{S}^2)$ -norm of (3.6).

4.1. Isolating the divergent term

We first isolate the singular contribution (3.8) from (3.7) by using probabilistic independence, and the ingredients such as (3.3) discussed at the end of Section 3.1.

Lemma 4.1. For every α , N, and t, we have

$$\begin{aligned} \left\| \| \mathrm{II}_{N}(t) \|_{L^{2}(\Omega; H^{\alpha-1}(\mathbb{S}^{2}))} - \| \mathrm{III}_{N}(t) \|_{L^{2}(\Omega; H^{\alpha-1}(\mathbb{S}^{2}))} \right\| \\ & \leq \| \mathcal{I}_{\mathbb{S}^{2}}(t, P_{\leq N} u^{\omega}) \|_{L^{2}(\Omega; H^{\alpha-1}(\mathbb{S}^{2}))}, \end{aligned}$$

where the notations are introduced in Section 3.4

Proof. For simplicity, we consider the case $\alpha = 1$. The other cases are the same up to applying $\langle \Delta \rangle^{\frac{\alpha-1}{2}}$ to every term. We have

$$\begin{aligned} \|\mathcal{I}_{\mathbb{S}^{2}}(t, P_{\leq N}u^{\omega})\|_{L^{2}(\Omega; L^{2}(\mathbb{S}^{2}))}^{2} &= \|\mathrm{I}_{N}(t)\|_{L^{2}(\Omega; L^{2}(\mathbb{S}^{2}))}^{2} + \|\mathrm{II}_{N}(t) + \mathrm{III}_{N}(t)\|_{L^{2}(\Omega; L^{2}(\mathbb{S}^{2}))}^{2} \\ &+ 2\mathbb{E}[\langle\mathrm{I}_{N}(t)|\mathrm{II}_{N}(t) + \mathrm{III}_{N}(t)\rangle_{L^{2}(\mathbb{S}^{2})}]. \end{aligned}$$

We show that, for every t and N,

$$\mathbb{E}[\langle \mathbf{I}_N(t) | \mathbf{II}_N(t) \rangle_{L^2(\mathbb{S}^2)}] = \mathbb{E}[\langle \mathbf{I}_N(t) | \mathbf{III}_N(t) \rangle_{L^2(\mathbb{S}^2)}] = 0.$$
(4.1)

Expanding the scalar product, we have

$$\begin{split} &\mathbb{E}[\langle \mathbf{I}_{N}(t) | \mathbf{II}_{N}(t) \rangle_{L^{2}(\mathbb{S}^{2})}] \\ &= \sum_{\vec{n} \in \Gamma_{N}^{(1)}} \sum_{\vec{n}' \in \Gamma_{N}^{(2)}} \mathbf{1}_{n_{0} = n'_{0}} \mathbb{E}\Big[\langle \pi_{n_{0}}(e_{n_{1}}^{\omega} \bar{e}_{n_{2}}^{\omega} e_{n_{3}}^{\omega}) \mid \bar{e}_{n'_{1}}^{\omega} (|e_{n'_{2}}^{\omega}|^{2} - ||e_{n'_{2}}^{\omega}||_{L^{2}(\mathbb{S}^{2})}^{2}) \rangle_{L^{2}(\mathbb{S}^{2})} \Big] \\ &\times \Big(\frac{e^{-it\Omega(\vec{n})} - 1}{\Omega(\vec{n})} \Big) (\lambda_{n_{1}}\lambda_{n_{2}}\lambda_{n_{3}})^{-(\alpha - \frac{1}{2})} \Big(\frac{e^{it\Omega(\vec{n}')} - 1}{\Omega(\vec{n}')} \Big) (\lambda_{n'_{1}}\lambda_{n'_{2}}^{2})^{-(\alpha - \frac{1}{2})}. \end{split}$$

Using the integral kernel K_{n_0} of π_{n_0} , we observe that, for every $(\vec{n}, \vec{n}') \in (\Gamma_N^{(1)} \times \Gamma_N^{(2)})$,

$$\mathbb{E}\left[\left\langle \pi_{n_{0}}(e_{n_{1}}^{\omega}\bar{e}_{n_{2}}^{\omega}e_{n_{3}}^{\omega}) \left| \bar{e}_{n_{1}'}^{\omega}(|e_{n_{2}}^{\omega}|^{2} - \|e_{n_{2}}^{\omega}\|_{L^{2}(\mathbb{S}^{2})}^{2}) \right\rangle_{L^{2}(\mathbb{S}^{2})} \right]$$

$$= \int_{(\mathbb{S}^{2})^{2}} K_{n_{0}}(x, y) \mathbb{E}\left[e_{n_{1}}^{\omega}(x)\bar{e}_{n_{2}}^{\omega}(x)e_{n_{3}}^{\omega}(x)\bar{e}_{n_{1}'}^{\omega}(y)(|e_{n_{2}'}^{\omega}(y)|^{2} - \|e_{n_{2}'}^{\omega}\|_{L^{2}(\mathbb{S}^{2})}^{2})\right] \mathrm{d}\sigma(x)\mathrm{d}\sigma(y).$$

We see from the non-pairing condition $n_2 \neq n_1, n_3$ of $\vec{n} \in \Gamma_N^{(1)}$ and from (3.3) that, for fixed x and y,

$$\mathbb{E}\Big[e_{n_1}^{\omega}(x)\bar{e}_{n_2}^{\omega}(x)e_{n_3}^{\omega}(x)\bar{e}_{n_1'}^{\omega}(y)(|e_{n_2'}^{\omega}(y)|^2 - \|e_{n_2'}^{\omega}\|_{L^2(\mathbb{S}^2)}^2)\Big] = 0.$$

This gives the first equality in (4.1). The second follows analogously, using that, for all $\vec{n} \in \Gamma_N^{(1)}$ and $\vec{n}' \in \Gamma_N^{(3)}$, and for all x, y in \mathbb{S}^2 ,

$$\begin{split} & \mathbb{E}\Big[e_{n_1}^{\omega}(x)\bar{e}_{n_2}^{\omega}(x)e_{n_3}^{\omega}(x)\bar{e}_{n_1'}^{\omega}(y)|e_{n_1'}^{\omega}(y)|^2\Big] \\ & = \mathbb{E}\Big[e_{n_1}^{\omega}(x)\bar{e}_{n_2}^{\omega}(x)e_{n_3}^{\omega}(x)\bar{e}_{n_1'}^{\omega}(y)\|e_{n_1'}^{\omega}(y)\|_{L^2(\mathbb{S}^2)}^2\Big] = 0. \end{split}$$

This proves (4.1). We deduce that

$$\|\mathcal{I}_{\mathbb{S}^{2}}(t, P_{\leq N}u^{\omega})\|_{L^{2}(\Omega; L^{2}(\mathbb{S}^{2}))}^{2} = \|\mathbf{I}_{N}(t)\|_{L^{2}(\Omega; L^{2}(\mathbb{S}^{2}))}^{2} + \|\mathbf{II}_{N}(t) + \mathbf{III}_{N}(t)\|_{L^{2}(\Omega; L^{2}(\mathbb{S}^{2}))}^{2},$$

and we conclude from the triangle inequality that

$$\begin{split} \|\mathcal{I}_{\mathbb{S}^{2}}(t,P_{\leq N}u^{\omega})\|_{L^{2}(\Omega;L^{2}(\mathbb{S}^{2}))} &\geq \|\Pi_{N}(t)+\Pi_{N}(t)\|_{L^{2}(\Omega;L^{2}(\mathbb{S}^{2}))} \\ &\geq \left|\|\Pi_{N}(t)\|_{L^{2}(\Omega;H^{\alpha-1}(\mathbb{S}^{2}))}-\|\Pi\Pi_{N}(t)\|_{L^{2}(\Omega;H^{\alpha-1}(\mathbb{S}^{2}))}\right|. \end{split}$$

This concludes the proof of Lemma 4.1.

Let us now show that the term III_N , written in (3.9), gains one derivative with respect to the initial data ϕ_{α} . This implies in particular that it is bounded in $L^2(\Omega; H^{\alpha-1}(\mathbb{S}^2))$ uniformly in N.

Lemma 4.2. For every α and s such that $s < 3\alpha - 2$,

$$\sup_{N} \mathbb{E} \left[\| \mathrm{III}_{N}(t) \|_{H^{s}(\mathbb{S}^{2})}^{2} \right] < +\infty.$$

In particular, when $\alpha > \frac{1}{2}$, then

$$\sup_{N} \mathbb{E} \Big[\| \mathrm{III}_{N}(t) \|_{H^{\alpha-1}(\mathbb{S}^{2})}^{2} \Big] < +\infty.$$

Proof. Fix $N \in 2^{\mathbb{N}}$. By Minkowski and the structure of $III_N(t)$, we have

$$\left(\mathbb{E}\left[\left\|\mathrm{III}_{N}(t)\right\|_{H^{s}(\mathbb{S}^{2})}^{2}\right]\right)^{\frac{1}{2}} \leq \sum_{\substack{K \leq N \\ K \text{ dyadic}}} \left(\mathbb{E}\left[\left\|\mathrm{III}(\mathrm{P}_{K} u^{\omega})(t)\right\|_{H^{s}(\mathbb{S}^{2})}^{2}\right]\right)^{\frac{1}{2}}.$$

It suffices to show that there exists $\delta > 0$, depending on *s* and α , such that, for all *K*,

$$\mathbb{E}\left[\|\mathrm{III}_N(\mathsf{P}_K \, u^{\omega})(t)\|_{H^s(\mathbb{S}^2)}^2\right] \lesssim K^{-\delta}.$$

We have

$$\mathbb{E}\left[\left\|\mathrm{III}_{N}(\mathsf{P}_{K}\,u^{\omega})(t)\right\|_{H^{s}(\mathbb{S}^{2})}^{2}\right] = \sum_{n_{0}}\lambda_{n_{0}}^{2s}\mathbb{E}\left[\left\|\pi_{n_{0}}\sum_{n_{1}\sim K}\left(\frac{\mathrm{e}^{-it(\lambda_{n_{0}}^{2}-\lambda_{n_{1}}^{2})}-1}{\lambda_{n_{0}}^{2}-\lambda_{n_{1}}^{2}}\right)\lambda_{n_{1}}^{-3(\alpha-\frac{1}{2})}|e_{n_{1}}^{\omega}|^{2}e_{n_{1}}^{\omega}\right\|_{L^{2}(\mathbb{S}^{2})}^{2}\right].$$
 (4.2)

By some degree considerations, we see that the terms contribute only when $n_0 \le 3n_1$. Moreover, expanding the square, we see that

$$\begin{aligned} |(4.2)| &\lesssim \sum_{\lambda_{n_0} \lesssim K} \sum_{n_1, n_1' \sim K} \lambda_{n_0}^{2s} (\lambda_{n_1} \lambda_{n_1'})^{-3(\alpha - \frac{1}{2})} \\ &\times \left| \int_{(\mathbb{S}^2)^2} K_{n_0}(x, y) \mathbb{E}[e_{n_1}^{\omega}(x)|e_{n_1}^{\omega}(x)|^2 \bar{e}_{n_1'}^{\omega}(y)|e_{n_1'}^{\omega}(y)|^2] \mathrm{d}\sigma(x) \mathrm{d}\sigma(y) \right|. \end{aligned}$$

Using the independence of $e_{n_1}(x)$ and $e_{n'_1}(y)$ when $n_1 \neq n'_1$, we see that only the terms with $n_1 = n'_1$ contribute. Hence,

$$\begin{aligned} |(4.2)| &\lesssim K^{2s} \sum_{n_0 \lesssim K} \sum_{n_1 \sim K} \lambda_{n_1}^{-6(\alpha - \frac{1}{2})} \\ &\times \left| \mathbb{E} \left[\int_{(\mathbb{S}^2)^2} K_{n_0}(x, y) |e_{n_1}^{\omega}(x)|^2 |e_{n_1}^{\omega}(y)|^2 e_{n_1}^{\omega}(x) \bar{e}_{n_1}^{\omega}(y) \mathrm{d}\sigma(x) \mathrm{d}\sigma(y) \right] \right| \\ &\lesssim K^{2s} \sum_{n_0 \lesssim K} \sum_{n_1 \sim K} \lambda_{n_1}^{-6(\alpha - \frac{1}{2})} \mathbb{E} \|\pi_{n_0}(|e_{n_1}^{\omega}|^2 e_{n_1}^{\omega})\|_{L^2(\mathbb{S}^2)}^2 \\ &\lesssim K^{2s} \sum_{n_1 \sim K} \lambda_{n_1}^{-6(\alpha - \frac{1}{2})} \mathbb{E} \|e_{n_1}^{\omega}\|_{L^6(\mathbb{S}^2)}^6. \end{aligned}$$

According to (3.5), we conclude that

$$|(4.2)| \lesssim K^{2s} \sum_{n_1 \sim K} \lambda_{n_1}^{-6(\alpha - \frac{1}{2})} \lesssim K^{2s + 1 - 6(\alpha - \frac{1}{2})},$$

which is conclusive when

$$s < 3\alpha - 2 = \alpha - 1 + 2\left(\alpha - \frac{1}{2}\right).$$

This concludes the proof of Lemma 4.2.

At this stage, we proved in Lemma 4.1 and in Lemma 4.2 that there exists C > 0 such that, for all $N \ge 1$, $\alpha > \frac{1}{2}$, $t \in \mathbb{R}$, and $s \le 3\alpha - 2$,

$$\| \Pi_N(t) \|_{L^2(\Omega; H^{\alpha-1}(\mathbb{S}^2))} - C \le \| \mathcal{I}_{\mathbb{S}^2}(t, P_{\le N} u_\alpha^{\omega}) \|_{L^2(\Omega; H^{\alpha-1}(\mathbb{S}^2))}.$$
(4.3)

It remains to show the divergence of the term on the left-hand side.

4.2. Proof of the divergence claimed in Theorem 1.1

In order to demonstrate the divergence asserted in Theorem 1.1, we exploit the concentration property associated with high-order spherical harmonics, as written in Proposition 2.2.

Proposition 4.3. There exist $\eta > 0$ and $N_0 \ge 0$ such that for all $N \ge N_0$ and $t \in \mathbb{R}$ we have

$$|t|\eta \log(N)^{\frac{1}{2}} \le \|\mathrm{II}_{N}(t)\|_{L^{2}(\Omega; H^{\alpha-1}(\mathbb{S}^{2}))}.$$
(4.4)

Proof. Recall that we defined in (3.8)

$$\Pi_N(t,x) = \sum_{\vec{n}\in\Gamma_N^{(2)}} \left(\frac{e^{-it\Omega(\vec{n})}-1}{\Omega(\vec{n})}\right) (\lambda_{n_1}\lambda_{n_2}^2)^{-(\alpha-\frac{1}{2})} \pi_{n_0} \left(e_{n_1}^{\omega}(|e_{n_2}^{\omega}|^2 - \|e_{n_2}^{\omega}\|_{L^2\mathbb{S}^2}^2)\right).$$

Note that when $\vec{n} = (n, n_1, n_2, n_3) \in \Gamma_N^{(2)}$, then the resonant function is $\Omega(\vec{n}) = \lambda_n^2 - \lambda_{n_1}^2$. We expand $\Pi_N(t)$ on the orthonormal basis of $L^2(\mathbb{S}^2)$ made of spherical harmonics:

$$\begin{aligned} \Pi_N(t) &= \sum_{n \leq N} \sum_{|k| \leq n} Y_{n,k} \sum_{\substack{n_1, n_2 \leq N \\ n_1, n_2 \leq N}} \mathbf{1}_{n_1 \neq n_2} (\lambda_{n_1} \lambda_{n_2}^2)^{-(\alpha - \frac{1}{2})} \frac{e^{-it(\lambda_n^2 - \lambda_{n_1}^2)} - 1}{\lambda_n^2 - \lambda_{n_1}^2} \\ &\times \left(\int_{\mathbb{S}^2} e_{n_1}^{\omega}(x) (|e_{n_2}^{\omega}(x)|^2 - \|e_{n_2}^{\omega}\|_{L^2(\mathbb{S}^2)}^2) \overline{Y_{n,k}(x)} \mathrm{d}\sigma(x) \right). \end{aligned}$$

Remark 4.4. Note that since we truncated the initial data at frequency N, we can deduce from degree considerations on the spherical harmonics that only the modes with $n \leq N$ contribute to the above sum.

By applying Plancherel's formula and using the fact that $e^{it(\Delta_{\mathbb{S}^2}-1)}$ is a unitary operator on $L^2(\mathbb{S}^2)$, we obtain

$$\|\mathrm{II}_{N}(t)\|_{H^{\alpha-1}(\mathbb{S}^{2})}^{2} = \sum_{n \leq N} \lambda_{n}^{2(\alpha-1)} \sum_{|k| \leq n} \left| \sum_{n_{1}, n_{2} \leq N} \mathbf{1}_{n_{1} \neq n_{2}} (\lambda_{n_{1}} \lambda_{n_{2}}^{2})^{-(\alpha-\frac{1}{2})} \frac{e^{-it(\lambda_{n}^{2}-\lambda_{n_{1}}^{2})} - 1}{\lambda_{n}^{2} - \lambda_{n_{1}}^{2}} \right| \\ \times \left(\int_{\mathbb{S}^{2}} e_{n_{1}}^{\omega}(x) (|e_{n_{2}}^{\omega}(x)|^{2} - ||e_{n_{2}}^{\omega}||_{L^{2}(\mathbb{S}^{2})}^{2}) \overline{Y_{n,k}(x)} \mathrm{d}\sigma(x) \right) \Big|^{2}.$$

By expressing $e_{n_1}^{\omega}$ in the basis made of spherical harmonics (Y_{n_1,k_1}) (recall that the law of $e_{n_1}^{\omega}$ does not depend on the choice of the orthonormal basis), one deduces from the independence and from (3.3) that

$$\mathbb{E} \left| \sum_{n_{1},n_{2} \leq N} \mathbf{1}_{n_{1} \neq n_{2}} (\lambda_{n_{1}} \lambda_{n_{2}}^{2})^{-(\alpha - \frac{1}{2})} \frac{e^{-it(\lambda_{n}^{2} - \lambda_{n_{1}}^{2})} - 1}{\lambda_{n}^{2} - \lambda_{n_{1}}^{2}} \\
\times \left(\int_{\mathbb{S}^{2}} e_{n_{1}}^{\omega}(x) (|e_{n_{2}}^{\omega}(x)|^{2} - ||e_{n_{2}}^{\omega}||_{L^{2}(\mathbb{S}^{2})}^{2}) \overline{Y_{n,k}(x)} d\sigma(x) \right) \right|^{2} \\
= \sum_{n_{1},n_{2} \leq N} \mathbf{1}_{n_{1} \neq n_{2}} (\lambda_{n_{1}} \lambda_{n_{2}}^{2})^{-2(\alpha - \frac{1}{2})} (2n_{1} + 1)^{-1} \left| \frac{e^{-it(\lambda_{n}^{2} - \lambda_{n_{1}}^{2})} - 1}{\lambda_{n}^{2} - \lambda_{n_{1}}^{2}} \right|^{2} \\
\times \sum_{|k_{1}| \leq n_{1}} \mathbb{E} \left| \int_{\mathbb{S}^{2}} (|e_{n_{2}}^{\omega}(x)|^{2} - ||e_{n_{2}}^{\omega}||_{L^{2}(\mathbb{S}^{2})}^{2}) g_{n_{1},k_{1}}(\omega) Y_{n_{1},k_{1}}(x) \overline{Y_{n,k}(x)} d\sigma(x) \right|^{2}. \quad (4.5)$$

Let us justify why only the terms with $n_1 = n'_1$ and $n_2 = n'_2$ contributed to the double sum over (n_1, n'_1, n_2, n'_2) . Fix $(x, y) \in (\mathbb{S}^2)^2$. We distinguish three cases.

Case 1: $n_1 \neq n'_1$ and $n_2 \neq n'_2$. Since, by assumption, $n'_1 \neq n'_2$ and $n_1 \neq n_2$, we have from the independence of \mathcal{B}_{n_1} and $\mathcal{B}_{n'_2}$ from $\mathcal{B}_{n'_1}$ and \mathcal{B}_{n_2} that

$$\mathbb{E}\Big[e_{n_1}^{\omega}(x)\Big(|e_{n_2}^{\omega}(x)|^2 - \|e_{n_2}^{\omega}\|_{L^2(\mathbb{S}^2)}^2\Big)\overline{e_{n_1'}^{\omega}(y)}\Big(|e_{n_2'}^{\omega}(y)|^2 - \|e_{n_2'}^{\omega}\|_{L^2(\mathbb{S}^2)}^2\Big)\Big] \\ = \mathbb{E}\Big[e_{n_1}^{\omega}(x)\Big(|e_{n_2'}^{\omega}(y)|^2 - \|e_{n_2'}^{\omega}\|_{L^2(\mathbb{S}^2)}^2\Big)\Big]\mathbb{E}\Big[\overline{e_{n_1'}^{\omega}(y)}\Big(|e_{n_2}^{\omega}(x)|^2 - \|e_{n_2}^{\omega}\|_{L^2(\mathbb{S}^2)}^2\Big)\Big] = 0,$$

where we used the identity (3.3).

Case 2: $n_1 \neq n'_1$ and $n_2 = n'_2$. In this case, we have from the independence of \mathcal{B}_{n_1} from $\mathcal{B}_{n'_1}$ and \mathcal{B}_{n_2} that

$$\mathbb{E}\Big[e_{n_1}^{\omega}(x)\big(|e_{n_2}^{\omega}(y)|^2 - ||e_{n_2}^{\omega}||_{L^2(\mathbb{S}^2)}^2\big)\overline{e_{n_1}^{\omega}(y)}\big(|e_{n_2}^{\omega}(y)|^2 - ||e_{n_2}^{\omega}||_{L^2(\mathbb{S}^2)}^2\big)\Big] \\ = \mathbb{E}[e_{n_1}^{\omega}(x)]\mathbb{E}\Big[\big(|e_{n_2}^{\omega}(y)|^2 - ||e_{n_2}^{\omega}||_{L^2(\mathbb{S}^2)}^2\big)\overline{e_{n_1}^{\omega}(y)}\big(|e_{n_2}^{\omega}(y)|^2 - ||e_{n_2}^{\omega}||_{L^2(\mathbb{S}^2)}^2\big)\Big] = 0,$$

where we used that $e_{n_1}^{\omega}(x)$ has zero average.

Case 3: $n_1 = n'_1$ and $n_2 \neq n'_2$. By assumption, $n_1 \neq n_2$ and we deduce from the independence of \mathcal{B}_{n_2} from \mathcal{B}_{n_1} and $\mathcal{B}_{n'_2}$ that

$$\begin{split} & \mathbb{E}\Big[e_{n_1}^{\omega}(x)\big(|e_{n_2}^{\omega}(x)|^2 - \|e_{n_2}^{\omega}\|_{L^2(\mathbb{S}^2)}^2\big)\overline{e_{n_1}^{\omega}(y)}\big(|e_{n'_2}^{\omega}(y)|^2 - \|e_{n'_2}^{\omega}\|_{L^2(\mathbb{S}^2)}^2\big)\Big] \\ & = \mathbb{E}\Big[|e_{n_2}^{\omega}(x)|^2 - \|e_{n_2}^{\omega}\|_{L^2(\mathbb{S}^2)}^2\Big]\mathbb{E}\Big[e_{n_1}^{\omega}(x)\overline{e_{n_1}^{\omega}(y)}\big(|e_{n'_2}^{\omega}(y)|^2 - \|e_{n'_2}^{\omega}\|_{L^2(\mathbb{S}^2)}^2\big)\Big] = 0, \end{split}$$

where we used that the first term in the second line vanishes according to (3.3).

This proves that only $n_1 = n'_1$ and $n_2 = n'_2$ contributed to the double sum.

We now estimate from below (4.5), keeping only the $high \times low$ frequency interactions, namely, when $n_2 = 1$, which are resonant at $n = n_1$, and with $k = k_1$. Uniformly in n, k with $n \ge 2$, we have

$$(4.5) \gtrsim t^2 n^{-1} \lambda_n^{-2(\alpha-\frac{1}{2})} \mathbb{E} \left| \int_{\mathbb{S}^2} \left(|e_1^{\omega}(x)|^2 - ||e_1^{\omega}||_{L^2(\mathbb{S}^2)}^2 \right) |Y_{n,k}(x)|^2 \mathrm{d}\sigma(x) \right|^2 \\ \sim t^2 n^{-2} \lambda_n^{-2(\alpha-1)} \mathbb{E} \left| \int_{\mathbb{S}^2} \left(|e_1^{\omega}(x)|^2 - ||e_1^{\omega}||_{L^2(\mathbb{S}^2)}^2 \right) |Y_{n,k}(x)|^2 \mathrm{d}\sigma(x) \right|^2,$$

where we used that $\lambda_n \sim cn$. We deduce from (3.8) and (4.5) that

$$\mathbb{E} \|\Pi_{N}(t)\|_{H^{\alpha-1}}^{2} \gtrsim t^{2} \sum_{2 \le n \lesssim N} n^{-2} \sum_{|k| \le n} \mathbb{E} \left| \int_{\mathbb{S}^{2}} \left(|e_{1}^{\omega}(x)|^{2} - \|e_{1}^{\omega}\|_{L^{2}(\mathbb{S}^{2})}^{2} \right) |Y_{n,k}(x)|^{2} \mathrm{d}\sigma(x) \right|^{2}.$$
(4.6)

Then, we use the explicit expression of e_1^{ω} , and we remove a well-chosen subset from Ω (such that the remainder still has positive measure) in order to exploit the concentration of $Y_{1,1}$ near the equator. We have, for $\omega \in \Omega$ and $x \in \mathbb{S}^2$,

$$e_1^{\omega}(x) = \frac{1}{\sqrt{3}} \Big(g_{1,1}(\omega) Y_{1,1}(x) + g_{1,-1}(\omega) Y_{1,-1}(x) + g_{1,0}(\omega) Y_{1,0}(x) \Big).$$

According to (2.3), we can express in spherical coordinates

$$Y_{1,1}(\theta,\varphi) = -\sqrt{\frac{3}{2}}\sin(\theta) e^{i\varphi}.$$

Then, given $0 < \varepsilon \ll 1$, we define

$$S_{\varepsilon} = \{ \omega \in \Omega \mid 1 \le |g_{1,1}(\omega)|^2 \le 100, \ |g_{1,-1}(\omega)|^2 + |g_{1,0}(\omega)|^2 \le \varepsilon^2 \}.$$

Note that $\mathbb{P}(S_{\varepsilon}) > 0$. We write

$$|e_1^{\omega}(x)|^2 - ||e_1^{\omega}||_{L^2(\mathbb{S}^2)}^2 = \frac{1}{3}|g_{1,1}(\omega)|^2 \left(\frac{3}{2}\sin^2(\theta) - 1\right) + r(x,\omega),$$

where there exists C > 0 such that function r satisfies, for all $x \in S^2$ and $\omega \in S_{\varepsilon}$,

$$|r(x,\omega)| \le C\varepsilon. \tag{4.7}$$

We observe that, for all $\delta > 0$,

$$x = (\theta, \varphi) \in \mathbb{S}^2 \setminus \mathbb{C}_{2\delta} \implies \frac{3}{2}\sin^2(\theta) - 1 \ge \frac{1}{2}(1 - 12\delta^2), \tag{4.8}$$

where \mathcal{C}_{δ} is defined in (2.5). Fix $\delta > 0$, $\omega \in S_{\varepsilon}$ and set

$$\begin{aligned} a_{n,k}(\omega,\delta) &= \frac{1}{3} |g_{1,1}(\omega)|^2 \int_{\mathbb{S}^2} (1 - \mathbf{1}_{\mathcal{C}_{2\delta}}(x)) \left(\frac{3}{2} \sin^2(\theta) - 1\right) |Y_{n,k}(x)|^2 d\sigma(x) \\ b_{n,k}(\omega,\delta) &= \frac{1}{3} |g_{1,1}(\omega)|^2 \int_{\mathbb{S}^2} \mathbf{1}_{\mathcal{C}_{2\delta}}(x) \left(\frac{3}{2} \sin^2(\theta) - 1\right) |Y_{n,k}(x)|^2 d\sigma(x) \\ &+ \int_{\mathbb{S}^2} r(x,\omega) |Y_{n,k}(x)|^2 d\sigma(x) \end{aligned}$$

so that

$$\int_{\mathbb{S}^2} \left(|e_1^{\omega}(x)|^2 - \|e_1^{\omega}\|_{L^2(\mathbb{S}^2)}^2 \right) |Y_{n,k}(x)|^2 \mathrm{d}\sigma(x) = a_{n,k}(\omega,\delta) + b_{n,k}(\omega,\delta)$$

Then, we deduce from the inequality $(a + b)^2 \ge \frac{a^2}{2} - b^2$ (for $a, b \in \mathbb{R}$) that, for all ω , n, k,

$$\mathbb{E}\left[\mathbf{1}_{S_{\varepsilon}}(\omega)\left|\int_{\mathbb{S}^{2}}\left(|e_{1}^{\omega}(x)|^{2}-\|e_{1}^{\omega}\|_{L^{2}(\mathbb{S}^{2})}\right)|Y_{n,k}(x)|^{2}\mathrm{d}\sigma(x)\right|^{2}\right]$$

$$\geq\frac{1}{2}\mathbb{E}[\mathbf{1}_{S_{\varepsilon}}(\omega)a_{n,k}(\omega)^{2}]-\mathbb{E}[\mathbf{1}_{S_{\varepsilon}}(\omega)b_{n,k}(\omega)^{2}].$$
(4.9)

Moreover, for all $\varepsilon, \delta > 0, n, k$ and $\omega \in S_{\varepsilon}$, we have form (4.8) that

$$a_{n,k}(\omega,\delta) \ge \frac{1}{6} |g_{1,1}(\omega)|^2 (1-12\delta^2) \int_{\mathbb{S}^2} (1-\mathbf{1}_{\mathcal{C}_{2\delta}}(x)) |Y_{n,k}(x)|^2 \mathrm{d}\sigma(x).$$

On the other hand, it follows from (4.7) and from the fact that $\frac{3}{2}\sin(\theta)^2 - 1 \le \frac{1}{2}$ when $x = (\theta, \varphi) \in \mathbb{C}_{\varepsilon}$ that

$$|b_{n,k}(\omega,\delta)| \leq \frac{1}{12} |g_{1,1}(\omega)|^2 \int_{\mathbb{S}^2} \mathbf{1}_{\mathcal{C}_{2\delta}}(x) |Y_{n,k}(x)|^2 \mathrm{d}\sigma(x) + C\varepsilon.$$

In Section 2.2, we prove that in the regime where *n* is large and $n(n + 1)(1 - \delta^2) \le k^2 \le n^2$ (see (2.6)) the spherical harmonics $\{Y_{n,k}\}$ concentrate their mass near the equator, precisely in the region outside $C_{2\delta}$. Applying Proposition 2.2, we obtain that there exists C_{δ} such that, for all $\delta > 0$, $\omega \in S_{\varepsilon}$, $n \ge 2$ and *k* as in (2.6),

$$a_{n,k}(\omega) \ge \frac{1}{6} |g_{1,1}(\omega)|^2 (1 - 12\delta^2) - \frac{C_{\delta}}{n} |g_{1,1}(\omega)|^2,$$

and

$$|b_{n,k}(\omega)| \leq \frac{C_{\delta}}{n} |g_{1,1}(\omega)|^2 + C\varepsilon.$$

Let $\delta > 0$ and $N \in \mathbb{N}$. We set

$$\Lambda_{N,\delta} := \{ (n,k) \in \mathbb{N}^2, \ N \le n, \ n(n+1)(1-\delta^2) \le k^2 \le n^2 \}.$$

We conclude from (4.9) that, when $\delta > 0$ is sufficiently small, there exists $N_{\delta} \ge 2$ such that, for all $(n, k) \in \Lambda_{N_{\delta}, \delta}$ and $\varepsilon > 0$ sufficiently small,

$$\mathbb{E} \left| \int_{\mathbb{S}^2} \left(|e_1^{\omega}(x)|^2 - \|e_1^{\omega}\|_{L^2(\mathbb{S}^2)}^2 \right) |Y_{n,k}(x)|^2 \mathrm{d}\sigma(x) \right|^2 \\ \ge \mathbb{E} \left[\mathbf{1}_{S_{\varepsilon}}(\omega) \left| \int_{\mathbb{S}^2} \left(|e_1^{\omega}(x)|^2 - \|e_1^{\omega}\|_{L^2(\mathbb{S}^2)}^2 \right) |Y_{n,k}(x)|^2 \mathrm{d}\sigma(x) \right|^2 \right] \ge \frac{1}{100} \mathbb{P}(S_{\varepsilon}).$$

Plugging the above estimate in (4.6), we conclude that there exist $\delta > 0$ and N_{δ} such that, for all $N \ge N_{\delta}$ and $\varepsilon > 0$ small enough,

$$\mathbb{E}\|\mathrm{II}_{N}(t)\|_{H^{\alpha-1}(\mathbb{S}^{2})}^{2} \gtrsim t^{2} \sum_{N_{\delta} \leq n \leq N} n^{-2} \#\{k : (n,k) \in \Lambda_{N_{\delta},\delta}\} \frac{1}{100} \mathbb{P}(S_{\varepsilon}) \gtrsim_{\delta,\varepsilon} t^{2} \log(N).$$

At this stage, δ and ε are small but fixed positive constants. This completes the proof of Proposition 4.3.

4.3. Conclusion

In order to complete the proof of Theorem 1.1, it suffices to inject (4.4) into (4.3). This yields the existence of $C, \eta > 0$ and a sufficiently large integer N_0 such that, for all t and for all $N \ge N_0$,

$$\|\mathcal{I}_{\mathbb{S}^{2}}(t, P_{\leq N}u_{\alpha}^{\omega})\|_{L^{2}(\Omega; H^{\alpha-1}(\mathbb{S}^{2}))} \geq |t|\eta \log(N)^{\frac{1}{2}} - C.$$

Taking N_0 large enough compared to C gives

$$\|\mathcal{I}_{\mathbb{S}^{2}}(t, P_{\leq N}u_{\alpha}^{\omega})\|_{L^{2}(\Omega; H^{\alpha-1}(\mathbb{S}^{2}))} \geq |t|\frac{\eta}{2}\log(N)^{\frac{1}{2}}.$$

This completes the proof of Theorem 1.1.

5. Regularity of the first iteration on tori

In this appendix, we consider the cubic NLS on the general torus

$$\mathbb{T}_{\beta}^2 = \mathbb{R}^2 / (2\pi\mathbb{Z})^2,$$

endowed with the metric $g = dx_1^2 + \beta^{-2}dx_2^2$ for some $\beta > 0$. The cubic NLS on \mathbb{T}_{β}^2 is written as

$$i\partial_t v + \Delta_\beta v = |v|^2 v, \tag{5.1}$$

where $\Delta_{\beta} = \partial_{x_1}^2 + \beta^2 \partial_{x_2}^2$. Hence, from now on, we will only concentrate on (5.1). For $n = (k, m), n' = (k', m') \in \mathbb{Z}^2$, we denote by $Q(n, n') = kk' + \beta^2 mm'$ the associate quadratic form and Q(n) := Q(n, n). Then,

$$e^{it\Delta_{\beta}} = \mathcal{F}_{x}^{-1} e^{-itQ(\cdot)} \mathcal{F}_{x}$$

In light of the Gaussian measure (1.4) defined in the case of \mathbb{S}^2 , we consider initial data of the form

$$\phi_1^{\omega}(x) = \sum_{n \in \mathbb{Z}^2} \frac{1}{\langle n \rangle} g_n^{\omega} e^{in \cdot x},$$

where $\langle n \rangle = (1 + Q(n))^{\frac{1}{2}}$, and

$$u^{\omega}(t,x) = \sum_{n \in \mathbb{Z}^2} \frac{e^{itQ(n)}}{\langle n \rangle} g_n^{\omega} e^{in \cdot x} \,.$$

We write the Wick-ordered nonlinearity

$$\mathcal{N}(u^{\omega})(t) = \sum_{\substack{(n_1, n_2, n_3) \in \mathbb{Z}^2 \\ n_2 \neq n_1, n_3}} \frac{e^{it(Q(n_1) - Q(n_2) + Q(n_3))}}{\langle n_1 \rangle \langle n_2 \rangle \langle n_3 \rangle} g_{n_1}(\omega) \overline{g_{n_2}(\omega)} g_{n_3}(\omega) e^{i(n_1 - n_2 + n_3) \cdot x}$$
$$= \sum_{\substack{(n_1, n_2, n_3, n) \in \Gamma}} \frac{e^{it\Phi(n)}}{\langle n_1 \rangle \langle n_2 \rangle \langle n_3 \rangle} g_{n_1}(\omega) \overline{g_{n_2}(\omega)} g_{n_3}(\omega) e^{in \cdot x},$$

where the constraint set is

$$\Gamma = \{ (n_1, n_2, n_3, n) \in (\mathbb{Z}^2)^4 : n = n_1 - n_2 + n_3, n_2 \neq n_1, n_2 \neq n_3 \}.$$

And the resonant function Φ defined on Γ is

$$\Phi(n) := Q(n_1) - Q(n_2) + Q(n_3) - Q(n_1 - n_2 + n_3) = 2Q(n_1 - n_2, n_2 - n_3).$$

The second Picard iteration is written as

$$\mathcal{I}_{\mathbb{T}^2_{\beta}}(t,\phi^{\omega}) = \int_0^t e^{i(t-t')\Delta_{\beta}} \mathcal{N}(u^{\omega}(t')) dt'.$$

We will now prove the gain of regularity (1.6) claimed in Theorem 1.1.

Proof. For fixed t, direct computation, independence assumption as well as the observation that $\mathbb{E}[g^2] = 0$ for standard complex Guassian functions yield that

$$\mathbb{E}\left[\left\|\mathcal{I}_{\mathbb{T}^{2}_{\beta}}(t,\phi^{\omega})\right\|_{H^{s}_{x}}^{2}\right] \sim \sum_{n=(n_{1},n_{2},n_{3})\in\Gamma} \frac{\langle n_{1}-n_{2}+n_{3}\rangle^{2s}}{\langle n_{1}\rangle^{2}\langle n_{2}\rangle^{2}\langle n_{3}\rangle^{3}\langle\Phi(n)\rangle^{2}}.$$
(5.2)

To compute the right-hand side of (5.2), we decompose dyadically $|n_j| \in N_j$, and assume that $N^{(1)} \ge N^{(2)} \ge N^{(3)}$ is the non-increasing order of N_1, N_2, N_3 . It would be sufficient to obtain an inequality of the form

$$(N_1 N_2 N_3)^{-2} \sum_{\substack{|n_j| \sim N_j, \, j = 1, 2, 3 \\ n_2 \neq n_1, n_3}} \frac{\langle n_1 - n_2 + n_3 \rangle^{2s}}{\langle Q(n_1 - n_2, n_2 - n_3) \rangle^2} \lesssim (N^{(1)})^{-\delta}.$$
(5.3)

The left-hand side of (5.3) can be bounded by

$$(N_1 N_2 N_3)^{-2} \sum_{l \in \mathbb{Z}} \frac{(N^{(1)})^{2s}}{\langle l \rangle^2} \sum_{\substack{|n_j| \in N_j \\ n_2 \neq n_1, n_3}} \mathbf{1}_{|\mathcal{Q}(n_1 - n_2, n_2 - n_3) - l| \le \delta}$$
(5.4)

$$\leq (N_1 N_2 N_3)^{-2} (N^{(1)})^{2s} \sup_{\substack{l \in \mathbb{Z} \\ n_2 \neq n_1, n_3}} \sum_{\substack{|n_j| \in N_j \\ n_2 \neq n_1, n_3}} \mathbf{1}_{|\mathcal{Q}(n_1 - n_2, n_2 - n_3) - l| \leq \delta}$$
(5.5)

for some number $0 < \delta < 1$ to be fixed later.

Let us now estimate the quantity (for fixed $l \in \mathbb{Z}$ and $\delta < 1/4$)

$$M_{N_1,N_2,N_3} := \sum_{\substack{|n_j| \sim N_j \\ n_2 \neq n_1, n_3}} \mathbf{1}_{|\mathcal{Q}(n_1 - n_2, n_2 - n_3) - l| \le \delta}.$$
(5.6)

Case 1: $N_1 \gg N_2$, N_3 . In this case, (5.6) can be majorized by

$$(N_2N_3)^2 \sup_{\substack{n_2,n_3:n_2 \neq n_3}} \sum_{\substack{n_1:|n_1| \sim N_1, n_1 \neq n_2}} \mathbf{1}_{|\mathcal{Q}(n_1 - n_2, n_2 - n_3) - l| \leq \delta}.$$

For fixed n_2 , n_3 , we denote by $m_1 = n_1 - n_2$, $m_0 = n_2 - n_3$, $|m_1| \sim N_1 \gg |m_0|$. It is sufficient to estimate the number of $m_1 \in \mathbb{Z}^2$, $|m_1| \sim N_1$ such that $|Q(m_1, m_0) - l| \leq \delta$. Denote by $m_0 = (\xi_0, \eta_0)$, if $\eta_0 = 0$ or $\xi_0 = 0$, $Q(m_1, m_0) \in \beta^2 \mathbb{Z}$ or $Q(m_1, m_0) \in \mathbb{Z}$, respectively. Then, $Q(m_1, m_0)$ can only take a discrete number of values. In these situations, we have (the same dimension treatment as in [6])

$$#\{m_1 \in \mathbb{Z}^2 : |m_1| \sim N_1, |Q(m_1, m_0) - l| \le \delta\} \lesssim N_1.$$

Therefore, without loss of generality, we may assume that both ξ_0 and η_0 are non-zero. Denote by

$$m_{\beta} = (\xi_0, \beta^2 \eta_0) = \text{diag}(1, \beta^2) m_0.$$

Then, it is reduced to estimate the cardinality of the set

$$S_{\delta} := \left\{ z \in \mathbb{Z}^2 : \left| z \cdot \frac{m_{\beta}}{|m_{\beta}|} - \frac{l}{|m_{\beta}|} \right| \le \frac{\delta}{|m_{\beta}|} \right\}$$

We observe that S_{δ} is a rectangle with side length $\sim N_1$ and width $\leq 2\delta$. Since $2\delta < 1/2$, we see that

$$\#S_{\delta} \lesssim N_1$$

Thus, the contribution of (5.4) in the sum is less than N_1^{2s-1} , and the associated dyadic summation over $N_1 \gg N_2$, N_3 converges provided that $s < \frac{1}{2}$.

Case 2: $N_2 \gg N_1$, N_3 . In this case, $|Q(n_1 - n_2, n_2 - n_3)| \sim N_2^2$. Coming back to (5.4), the range of the sum of l is $|l| \ge N_2^2$. Hence, by the crude estimate

$$M_{N_1,N_2,N_3} \lesssim (N_1 N_2 N_3)^3$$

the contribution of (5.4) is bounded by

$$\sum_{l|\geq N_2^2} \frac{N_2^{2s}}{l^2} \lesssim N_2^{2s-2}.$$

Then, associated dyadic summation over $N_2 \gg N_1$, N_3 converges, provided that s < 1.

For the remaining situations, the argument is the same as for Case 1. For example, if $N_1 \sim N_2 \gg N_3$ ($N_2 \sim N_3 \gg N_1$), we can fix n_2 , n_3 (n_1 , n_2), and do the same manipulation as in Case 1.

This completes the proof of (1.6) in Theorem 1.1.

Acknowledgments. The authors would like to thank the anonymous referee for the careful reading of the manuscript and for the valuable comments and suggestions.

Funding. This research was supported by the European research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant agreement 101097172 - GEOEDP). C.S and N.T. were partially supported by the ANR project Smooth ANR-22-CE40-0017. N.C. benefited from the support of the Center Henri Lebesgue ANR-11-LABX-0020-0, the region Pays de la Loire through the project MasCan, and the ANR project KEN ANR-22-CE40-0016.

References

- A. Ayache and N. Tzvetkov, L^p properties for Gaussian random series. Trans. Amer. Math. Soc. 360 (2008), no. 8, 4425–4439 Zbl 1145.60019 MR 2395179
- [2] V. Banica, On the nonlinear Schrödinger dynamics on S². J. Math. Pures Appl. (9) 83 (2004), no. 1, 77–98 Zbl 1084.35082 MR 2032582
- [3] A. L. Besse, Manifolds all of whose geodesics are closed. Ergeb. Math. Grenzgeb. 93, Springer, Berlin, 1978 Zbl 0387.53010 MR 0496885
- [4] J. Bourgain, Exponential sums and nonlinear Schrödinger equations. Geom. Funct. Anal. 3 (1993), no. 2, 157–178 Zbl 0787.35096 MR 1209300
- [5] J. Bourgain, Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. I. Schrödinger equations. *Geom. Funct. Anal.* 3 (1993), no. 2, 107–156 Zbl 0787.35097 MR 1209299
- [6] J. Bourgain, Construction of approximative and almost periodic solutions of perturbed linear Schrödinger and wave equations. *Geom. Funct. Anal.* 6 (1996), no. 2, 201–230 Zbl 0872.35007 MR 1384610
- J. Bourgain, Invariant measures for the 2D-defocusing nonlinear Schrödinger equation. Comm. Math. Phys. 176 (1996), no. 2, 421–445 Zbl 0852.35131 MR 1374420
- [8] J. Bourgain and C. Demeter, The proof of the l² decoupling conjecture. Ann. of Math. (2) 182 (2015), no. 1, 351–389 Zbl 1322.42014 MR 3374964
- [9] B. Bringmann, Almost sure local well-posedness for a derivative nonlinear wave equation. Int. Math. Res. Not. IMRN 2021 (2021), no. 11, 8657–8697 Zbl 1473.35361 MR 4266148
- [10] B. Bringmann, Y. Deng, A. R. Nahmod, and H. Yue, Invariant Gibbs measures for the three dimensional cubic nonlinear wave equation. *Invent. Math.* 236 (2024), no. 3, 1133–1411 Zbl 1541.35502 MR 4743517
- [11] B. Bringmann, J. Lührmann, and G. Staffilani, The wave maps equation and Brownian paths. *Comm. Math. Phys.* 405 (2024), no. 3, article no. 60 Zbl 1534.35261 MR 4709099
- [12] N. Burq, N. Camps, C. Sun, and N. Tzvetkov, Probabilistic well-posedeness for the nonlinear Schrödinger equation on the 2*d* sphere I: Positive regularities. 2024, arXiv:2404.18229v1

- [13] N. Burq, P. Gérard, and N. Tzvetkov, An instability property of the nonlinear Schrödinger equation on S^d. Math. Res. Lett. 9 (2002), no. 2-3, 323–335 Zbl 1003.35113 MR 1909648
- [14] N. Burq, P. Gérard, and N. Tzvetkov, Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds. *Amer. J. Math.* 126 (2004), no. 3, 569–605 Zbl 1067.58027 MR 2058384
- [15] N. Burq, P. Gérard, and N. Tzvetkov, Bilinear eigenfunction estimates and the nonlinear Schrödinger equation on surfaces. *Invent. Math.* 159 (2005), no. 1, 187–223 Zbl 1092.35099 MR 2142336
- [16] N. Burq, P. Gérard, and N. Tzvetkov, High frequency solutions of the nonlinear Schrödinger equation on surfaces. *Quart. Appl. Math.* 68 (2010), no. 1, 61–71 Zbl 1187.35232 MR 2598880
- [17] N. Burq and G. Lebeau, Injections de Sobolev probabilistes et applications. Ann. Sci. Éc. Norm. Supér. (4) 46 (2013), no. 6, 917–962 Zbl 1296.46031 MR 3134684
- [18] N. Burq and N. Tzvetkov, Random data Cauchy theory for supercritical wave equations. I. Local theory. *Invent. Math.* **173** (2008), no. 3, 449–475 Zbl 1156.35062 MR 2425133
- [19] N. Burq and N. Tzvetkov, Random data Cauchy theory for supercritical wave equations. II. A global existence result. *Invent. Math.* 173 (2008), no. 3, 477–496 Zbl 1187.35233 MR 2425134
- [20] N. Camps, L. Gassot, and S. Ibrahim, Refined probabilistic local well-posedness for a cubic Schrödinger half-wave equation. J. Differential Equations 380 (2024), 443–490 Zbl 1532.35416 MR 4666216
- [21] M. Christ, J. Colliander, and T. Tao, Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations. *Amer. J. Math.* **125** (2003), no. 6, 1235–1293 Zbl 1048.35101 MR 2018661
- [22] M. Christ, J. Colliander, and T. Tao, Ill-posedness for nonlinear Schrodinger and wave equations. 2003, arXiv:math/0311048
- [23] Y. Deng, A. R. Nahmod, and H. Yue, Random tensors, propagation of randomness, and nonlinear dispersive equations. *Invent. Math.* 228 (2022), no. 2, 539–686 Zbl 1506.35208 MR 4411729
- [24] Y. Deng, A. R. Nahmod, and H. Yue, Invariant Gibbs measures and global strong solutions for nonlinear Schrödinger equations in dimension two. Ann. of Math. (2) 200 (2024), no. 2, 399–486 MR 4792067
- [25] C. Fan, Y. Ou, G. Staffilani, and H. Wang, 2D-defocusing nonlinear Schrödinger equation with random data on irrational tori. *Stoch. Partial Differ. Equ. Anal. Comput.* 9 (2021), no. 1, 142–206 Zbl 1479.35269 MR 4218790
- [26] R. L. Frank and J. Sabin, Spectral cluster bounds for orthonormal systems and oscillatory integral operators in Schatten spaces. Adv. Math. 317 (2017), 157–192 Zbl 1373.58015 MR 3682666
- [27] P. Gérard and T. Kappeler, On the integrability of the Benjamin–Ono equation on the torus. *Comm. Pure Appl. Math.* 74 (2021), no. 8, 1685–1747 Zbl 1471.35354 MR 4275336
- [28] P. Gérard, T. Kappeler, and P. Topalov, Sharp well-posedness results of the Benjamin–Ono equation in H^s(T, ℝ) and qualitative properties of its solutions. Acta Math. 231 (2023), no. 1, 31–88 Zbl 1533.35061 MR 4652410
- [29] M. Gubinelli, H. Koch, and T. Oh, Paracontrolled approach to the three-dimensional stochastic nonlinear wave equation with quadratic nonlinearity. J. Eur. Math. Soc. (JEMS) 26 (2024), no. 3, 817–874 Zbl 1537.35434 MR 4721025

- [30] Z. Hani, Global well-posedness of the cubic nonlinear Schrödinger equation on closed manifolds. *Comm. Partial Differential Equations* 37 (2012), no. 7, 1186–1236 Zbl 1256.35136 MR 2942981
- [31] S. Herr and B. Kwak, Strichartz estimates and global well-posedness of the cubic NLS on T². Forum Math. Pi 12 (2024), article no. e14 Zbl 07919006 MR 4794808
- [32] L. Hörmander, The spectral function of an elliptic operator. Acta Math. 121 (1968), 193–218
 Zbl 0164.13201 MR 0609014
- [33] T. Kappeler and P. Topalov, Global wellposedness of KdV in H⁻¹(T, ℝ). Duke Math. J. 135 (2006), no. 2, 327–360 Zbl 1106.35081 MR 2267286
- [34] T. Oh, Remarks on nonlinear smoothing under randomization for the periodic KdV and the cubic Szegő equation. *Funkcial. Ekvac.* 54 (2011), no. 3, 335–365 Zbl 1387.35637 MR 2918143
- [35] B. Shiffman and S. Zelditch, Random polynomials of high degree and Levy concentration of measure. Asian J. Math. 7 (2003), no. 4, 627–646 Zbl 1060.32012 MR 2074895
- [36] C. D. Sogge, Oscillatory integrals and spherical harmonics. *Duke Math. J.* 53 (1986), no. 1, 43–65 Zbl 0636.42018 MR 0835795
- [37] E. M. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*. Princeton Math. Ser. 32, Princeton University Press, Princeton, NJ, 1971 Zbl 0232.42007 MR 0304972
- [38] C. Sun and N. Tzvetkov, Refined probabilistic global well-posedness for the weakly dispersive NLS. Nonlinear Anal. 213 (2021), article no. 112530 Zbl 1479.35824 MR 4299837
- [39] L. Thomann, The WKB method and geometric instability for nonlinear Schrödinger equations on surfaces. *Bull. Soc. Math. France* 136 (2008), no. 2, 167–193 Zbl 1161.35050 MR 2415340
- [40] J. M. VanderKam, L^{∞} norms and quantum ergodicity on the sphere. *Internat. Math. Res. Notices* (1997), no. 7, 329–347 Zbl 0877.58056 MR 1440572
- [41] A. Weinstein, Nonlinear stabilization of quasimodes. In Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), pp. 301–318, Proc. Sympos. Pure Math. XXXVI, American Mathematical Society, Providence, RI, 1980 MR 0573443

Received 31 January 2024; revised 2 September 2024.

Nicolas Burq

Laboratoire de Mathématique d'Orsay, UMR CNRS 8628 and Institut Universitaire de France, Université Paris-Saclay, Bâtiment 307, 307 Rue Michel Magat, 91400 Orsay, France; nicolas.burq@universite-paris-saclay.fr

Nicolas Camps

Institut de recherche mathématique de Rennes, UMR CNRS 6625, Université de Rennes, Campus de Beaulieu, bâtiments 22 et 23, 263 avenue du Général Leclerc, CS 74205, 35042 Rennes, France; nicolas.camps@univ-rennes.fr

Mickaël Latocca

Laboratoire de Mathématiques et de Modélisation d'Évry, UMR CNRS 8071, Université d'Evry Val d'Essonne, 23 Boulevard de France, 91037 Évry, France; mickael.latocca@univ-evry.fr

Chenmin Sun

Laboratoire d'Analyse et de Mathématiques Appliquées, UMR CNRS 8050, Université Paris-Est Créteil, Bâtiment P3 - 4ème étage 61, avenue du Général de Gaulle, 94010 Créteil, France; chenmin.sun@cnrs.fr

Nikolay Tzvetkov

Unité de Mathématiques Pures et Appliquées, UMR CNRS 5669, École Normale Supérieure de Lyon, 46 Allée d'Italie, 69007 Lyon, France; nikolay.tzvetkov@ens-lyon.fr