

Finite simple characteristic quotients of the free group of rank 2

William Y. Chen, Alexander Lubotzky, and Pham Huu Tiep

Abstract. In this paper, we describe how to explicitly construct infinitely many finite simple groups as characteristic quotients of the rank 2 free group F_2 . This shows that a “baby” version of the Wiegold conjecture [in: *Geometry, Rigidity, and Group Actions* (2011), 609–643] fails for F_2 and provides counterexamples to two conjectures in the theory of noncongruence subgroups of $\mathrm{SL}_2(\mathbb{Z})$ by Chen [Math. Ann. 371 (2018), 41–126]. Our main result explicitly produces, for every prime power $q \geq 7$, the groups $\mathrm{SL}_3(\mathbb{F}_q)$ and $\mathrm{SU}_3(\mathbb{F}_q)$ as characteristic quotients of F_2 . Our strategy is to study specializations of the Burau representation for the braid group B_4 , exploiting an exceptional relationship between F_2 and B_4 first observed by Dyer, Formanek, and Grossman [Arch. Math. (Basel) 38 (1982), 404–409]. Weisfeiler’s strong approximation theorem guarantees that our specializations are surjective for infinitely many primes, but they are not effective. To make our result effective, we give another proof of surjectivity via a careful analysis of the maximal subgroup structures of $\mathrm{SL}_3(\mathbb{F}_q)$ and $\mathrm{SU}_3(\mathbb{F}_q)$. These examples are minimal in the sense that no finite simple group of the form $\mathrm{PSL}_2(\mathbb{F}_q)$ appears as a characteristic quotient of F_2 .

1. Introduction

Let F_n be the free group on n generators, G a finite group, and $\mathrm{Epi}(F_n, G)$ the set of epimorphisms from F_n onto G . The group $\mathrm{Aut}(G) \times \mathrm{Aut}(F_n)$ acts on $\mathrm{Epi}(F_n, G)$ as follows: for $\alpha \in \mathrm{Aut}(F_n)$, $\beta \in \mathrm{Aut}(G)$, and $\varphi \in \mathrm{Epi}(F_n, G)$,

$$(\beta, \alpha) \cdot \varphi := \beta \circ \varphi \circ \alpha^{-1}.$$

A long-standing conjecture attributed to Wiegold [40, Conjecture 2.5.4] asserts that if $n \geq 3$, G is a finite simple group, then this action is transitive. As $\mathrm{Epi}(F_n, G)/\mathrm{Aut}(G)$ can be identified with the set

$$\mathcal{N}_n(G) := \{N \triangleleft G \mid F_n/N \cong G\}$$

Mathematics Subject Classification 2020: 20E05 (primary); 20G40, 20F36, 05E16, 14H10 (secondary).

Keywords: braid group, elliptic curves, Burau representation, characteristic quotient, strong approximation, noncongruence subgroup.

of normal subgroups of F_n with quotient isomorphic to G , this conjecture can be reformulated as follows.

Conjecture 1.1 (Wiegold). If $n \geq 3$ and G is a finite simple group, then $\text{Aut}(F_n)$ acts transitively on $\mathcal{N}_n(G)$.

In the language of group presentations, the conjecture roughly says that, for any $n \geq 3$, any finite simple group G has a unique presentation as an n -generated group. It has been known for a long time that the statement of the conjecture is false for $n = 2$. In fact, in 1951, B. H. Neumann and H. Neumann noticed that already for $G = A_5$, $\text{Aut}(F_2)$ acts with two orbits on $\mathcal{N}_2(A_5)$ [39]. A more quantitative statement was proved by Garion and Shalev, who showed that $|\mathcal{N}_2(G)/\text{Aut}(F_2)| \rightarrow \infty$ for any infinite family of nonabelian finite simple groups G ; i.e., the number of orbits goes to infinity as the order of the group goes to infinity.¹

In [30], the second author suggested a “baby” version of the Wiegold conjecture, which seems to be of independent interest.

Conjecture 1.2 (“Baby Wiegold”, Lubotzky [30]). For $n \geq 3$ and G a finite simple group, $\text{Aut}(F_n)$ has no fixed points on $\mathcal{N}_n(G)$.

It is easy to see that the “Baby Wiegold conjecture” is equivalent to the following conjecture.

Conjecture 1.3. For $n \geq 3$, F_n does not have a characteristic finite index subgroup R for which F_n/R is a finite simple group.

The goal of this note is to show that even the baby Wiegold conjecture fails for $n = 2$.

Theorem 1.4 (See Section 3.2). *The free group on two generators has infinitely many characteristic subgroups N with F_2/N being a finite simple group. In particular, the Baby Wiegold conjecture fails for $n = 2$.*

The strong failure of Wiegold’s conjecture in the case $n = 2$ as shown in [15, 19] can be explained as follows: it is a classical result of Nielsen that if a, b and a', b' are two bases for F_2 , then the commutator $[a, b]$ is conjugate to $[a', b']^{\pm 1}$. It follows that for $\varphi \in \text{Epi}(F_2, G)$, the union of the conjugacy classes of $[\varphi(a), \varphi(b)]^{\pm 1}$ is an invariant of the $\text{Aut}(F_2)$ -action. Since almost every pair of elements of a finite simple group G generates, it is natural to expect that the number of conjugacy classes of commutators of generating pairs goes to ∞ as $|G| \rightarrow \infty$; a main achievement of [19] is making

¹The case where G ranges over groups of type $\text{PSL}_2(\mathbb{F}_q)$ was already known to Evans [15].

this idea precise. However, their method of proof it does not give any information on fixed points, and so, does not say anything about baby Wiegold.

In this paper, we take a very different route. In [30], a connection was established between Wiegold's conjecture and the following well-known conjecture.

Conjecture 1.5. For every finite dimensional linear representation ρ of $\text{Aut}(F_n)$, $n \geq 3$, $\rho(\text{Inn}(F_n))$ is virtually solvable.

In fact, it was shown in [30] that Wiegold implies Conjecture 1.5 when the Zariski closure of $\rho(\text{Aut}(F_n))$ is connected (see the more recent manuscript of Gelander [20]). Conjecture 1.5 fails for $\text{Aut}(F_2)$; in fact, $\text{Aut}(F_2)$ is known to have a faithful linear representation. Indeed, in [14], Dyer, Formanek, and Grossman showed that $\text{Aut}(F_2)$ has a faithful representation if and only if the braid group B_4 does. At the time, the problem for B_4 was an open problem, but later it was shown in [4, 26, 27] that all braid groups are linear, via the so-called ‘‘Lawrence–Krammer representation’’. Consequently, $\text{Aut}(F_2)$ is also linear.

In this paper, we will use the related *Burau representation* $\rho : B_4 \rightarrow \text{GL}_3(\mathbb{Z}[q, q^{-1}])$ to produce infinitely many finite simple groups as characteristic quotients of F_2 , thereby proving Theorem 1.4 and showing that baby Wiegold fails for F_2 . A key tool is the strong approximation theorem of Weisfeiler and Pink, which allows us to deduce that the Burau representation is surjective onto the k_0 -valued points of a certain algebraic group, where k_0 is an appropriate finite field quotient of the ring $\mathbb{Z}[q + q^{-1}]$. A similar strategy was employed by Funar and Lochak [17] to show that for every $g \geq 2$, the fundamental group Π_g of a closed surface of genus g has infinitely many characteristic finite simple quotients. For them, the Burau representation is replaced by quantum representations of mapping class groups.

The strategy using strong approximation is quite general, and tells us which types of finite simple groups we can expect to produce as characteristic quotients of F_2 . However, it requires us to discard an indeterminate finite set of primes, and hence, the method does not specify even a single group as a characteristic quotient of F_2 . Because of this, we were motivated to prove the following more precise result.

Theorem 1.6 (See Section 4). *Let q be any prime power ≥ 7 . Then, the finite simple groups $\text{PSL}_3(\mathbb{F}_q)$ and $\text{PSU}_3(\mathbb{F}_q)$ are both characteristic quotients of F_2 . The finite simple groups $\text{PSU}_3(\mathbb{F}_4)$ and $\text{PSU}_3(\mathbb{F}_5)$ are also characteristic quotients of F_2 . The cases $\text{PSL}_3(\mathbb{F}_q)$ ($q = 2, 3, 4, 5$) and $\text{PSU}_3(\mathbb{F}_q)$ ($q = 2, 3$) are not characteristic quotients of F_2 .*

Here, $\text{PSU}_3(\mathbb{F}_q)$ denotes the projective special unitary group of a 3-dimensional Hermitian space over \mathbb{F}_{q^2} relative to the \mathbb{F}_q -linear involution $a \mapsto a^q$. In the proof of Theorem 1.6, the surjectivity provided by strong approximation is replaced by a

careful analysis of the maximal subgroup structure² of $\mathrm{SL}_3(\mathbb{F}_q)$ and $\mathrm{SU}_3(\mathbb{F}_q)$; moreover, these characteristic quotients are constructed using *explicit* specializations of the Burau representation. We have checked using GAP [18] that $\mathrm{PSU}_3(\mathbb{F}_4)$ is the smallest finite simple characteristic quotient of F_2 .

An overview of the strong approximation strategy is given in Proposition 3.4, where we give general conditions under which a representation of $\mathrm{Aut}(F_2)$ would give infinitely many concrete counterexamples to baby Wiegold for $n = 2$. We expect that there are many representations satisfying such conditions, though in this paper, we only consider those coming from various specializations of the Burau representation of B_4 . Other candidates include the Jones representation [25], and the Lawrence–Krammer representation (see [3] and [47, Section 3]).³ In particular, the counterexamples G that we get are all of bounded Lie rank. It would be interesting to find examples of unbounded rank. Another question of interest is whether there exist an alternating group A_n , which appears as a characteristic quotient of F_2 . Our method cannot answer this last problem.

We will also show that $\mathrm{PSL}_2(\mathbb{F}_q)$ is never a characteristic quotient of F_2 (Corollary 5.4), so in some sense the quotients $\mathrm{PSL}_3(\mathbb{F}_q)$, $\mathrm{PSU}_3(\mathbb{F}_q)$ that we find are the simplest finite simple groups of Lie type which appear as characteristic quotients of F_2 . Our strategy for this is to show that the fixed points of the $\mathrm{Out}(F_2)$ -action on the \mathbb{F}_q -points of the character variety for SL_2 -representations of F_2 correspond to nonsurjective representations. This approach is made possible by the known integral structure of this character variety. A similar understanding for the SL_3 - and SU_3 -character varieties would in theory allow for a similar approach to Theorem 1.6, though the equations would be notably more complicated [29, 37, 45], and it would not shed any light on where these fixed points come from.

Finally, our main theorem provides counterexamples to two conjectures in the theory of noncongruence modular curves [8, Conjectures 4.4.1, 4.4.15]. Let $\mathrm{Aut}^+(F_2)$ be the subgroup inducing determinant 1 automorphisms of the abelianization \mathbb{Z}^2 . The conjectures would assert that for nonsolvable groups G , the $\mathrm{Aut}^+(F_2)$ -stabilizers of any surjection $\varphi : F_2 \rightarrow G$ should map to a *noncongruence* subgroup of $\mathrm{SL}_2(\mathbb{Z})$. In Section 6, we explain why this is a reasonable conjecture, and why our Theorem 1.6 provides counterexamples.

Organization of the paper. In Section 2, we introduce the Burau representation of the braid group B_4 and the associated Hermitian form and unitary group, and show how it can be used to produce characteristic quotients of F_2 .

²This was worked out by Mitchell and Hartley in the early 1900s and does not use the classification of finite simple groups.

³In [3], it is noted that the description of the Lawrence–Krammer representation in the earlier paper [4] has a sign error.

In Section 3, we describe Weisfeiler and Pink's strong approximation theorem, and show, following [30], how it can be used to produce infinitely many finite linear groups as invariant quotients of a group F from a single characteristic 0 representation of $\text{Aut}(F)$. We apply this to the Burau representation to deduce Theorem 1.4.

In Section 4, we prove the stronger Theorem 1.6 by carefully choosing specializations of the Burau representation, exploiting the known structure of the maximal subgroups of $\text{SL}_3(\mathbb{F}_q)$ and $\text{SU}_3(\mathbb{F}_q)$.

In Section 5, we will show that $\text{PSL}_2(\mathbb{F}_q)$ is never a characteristic quotient of F_2 (for any prime power q).

In Section 6, we describe how our main theorem provides counterexamples to two conjectures of [8] in the theory of noncongruence modular curves.

2. Invariant quotients and the Burau representation

2.1. Generalities on invariant quotients

Let F be a normal subgroup of a group A . The conjugation action of A on F yields a homomorphism $h : A \rightarrow \text{Aut}(F)$. We say that a homomorphism of abstract groups $\varphi : F \rightarrow G$ is $h(A)$ -invariant (or A -invariant) if for every $\alpha \in h(A) \subset \text{Aut}(F)$,

$$\varphi \circ \alpha = \sigma \circ \varphi$$

for some $\sigma \in \text{Aut}(G)$, and *strictly A -invariant* if we can, moreover, take $\sigma \in \text{Inn}(G)$. If φ is surjective, then whenever σ exists, it is uniquely determined by α , in which case, we will write $\sigma = \varphi_*(\alpha)$. For A -invariant quotients φ , the map

$$\begin{aligned} \varphi_* : A &\rightarrow \text{Aut}(G), \\ \alpha &\mapsto \sigma \end{aligned}$$

is a homomorphism.

If $h(A) = \text{Aut}(F)$, then a homomorphism $\varphi : F \rightarrow G$ is A -invariant if and only if $\ker(\varphi)$ is a characteristic subgroup of F ; in this case, we say that φ is a characteristic homomorphism, or a characteristic quotient if φ is moreover surjective. We will make use of the following criterion for producing A -invariant homomorphisms.

Proposition 2.1. *With the above notation, if G sits as a subgroup inside a group \tilde{G} such that $\varphi : F \rightarrow G$ extends to a map $\tilde{\varphi} : A \rightarrow \tilde{G}$, then φ is A -invariant. If φ is surjective and $Z(G) = 1$, then the converse is also true.*

Proof. The forward direction is clear. For the converse, take $G \hookrightarrow \tilde{G}$ to be $G \cong \text{Inn}(G) \hookrightarrow \text{Aut}(G)$. ■

2.2. The braid group B_4 and the free subgroup F

Recall that the braid group B_n on n strands is the group generated by $\sigma_1, \dots, \sigma_{n-1}$ subject to the following relations:

- $[\sigma_i, \sigma_j] = 1$ if $|i - j| \geq 2$,
- $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$ if $1 \leq i \leq n - 2$.

The center of B_n is cyclic of infinite order generated by $(\sigma_1 \sigma_2 \cdots \sigma_{n-1})^n$, and has abelianization \mathbb{Z} , given by the word length map $\ell : B_n \rightarrow \mathbb{Z}$ defined by $\ell(\sigma_i) = 1$ for each i .

Let F_n denote the free group on the generators x_1, \dots, x_n . There is a well-known action of B_n on F_n given by $\sigma_i(x_1, \dots, x_n) = (x_1, \dots, x_{i-1}, x_{i+1}, x_{i+1}^{-1} x_i x_{i+1}, x_{i+2}, \dots, x_n)$. In this paper, we make use of a different representation, which seems to be exceptional for the case $n = 4$. From now on, we let x, y, F denote the following objects inside B_4 :

$$x := \sigma_1 \sigma_3^{-1}, \quad y := \sigma_2 \sigma_1 \sigma_3^{-1} \sigma_2^{-1}, \quad F := \langle x, y \rangle.$$

Then, x, y are free generators of F , and F is a *normal* subgroup of B_4 [14, p. 406]. The action of B_4 on F by conjugation defines a homomorphism

$$\begin{aligned} \xi : B_4 &\rightarrow \text{Aut}(F), \\ \sigma_1 &\mapsto (x, y) \mapsto (x, yx^{-1}), \\ \sigma_2 &\mapsto (x, y) \mapsto (y, yx^{-1}y), \\ \sigma_3 &\mapsto (x, y) \mapsto (x, x^{-1}y). \end{aligned}$$

The image of ξ is the group of automorphisms of F which act with determinant 1 on the abelianization $F^{\text{ab}} \cong \mathbb{Z}^2$, denoted by $\text{Aut}^+(F)$, and the kernel is exactly the center $Z(B_4)$ [14, p. 406]. Thus, we have an exact sequence

$$1 \rightarrow Z(B_4) \hookrightarrow B_4 \xrightarrow{\xi} \text{Aut}^+(F) \rightarrow 1, \quad (1)$$

where we recall that $Z(B_4) = \langle (\sigma_1 \sigma_2 \sigma_3)^4 \rangle \cong \mathbb{Z}$. In terms of invariant quotients, we have the following proposition.

Proposition 2.2. *Let $\varphi : B_4 \rightarrow G$ be a homomorphism of groups. Then, the restriction $\varphi|_F : F \rightarrow \varphi(F)$ is $\text{Aut}^+(F)$ -invariant.*

2.2.1. Remarks on canonical locally constant sheaves on surfaces. Geometrically, Proposition 2.2 implies that representations of B_4 correspond to *canonical* geometric objects on punctured elliptic curves. We first explain this for covering spaces. Let S_g be a closed surface of genus g , and let $S_{g,n}$ be an n -punctured S_g . For $g = 1$, the fundamental group $\pi_1(S_{1,1})$ is free of rank 2, and we will identify $\pi_1(S_{1,1})$ with F . By

covering space theory, the map $\varphi|_F : F \rightarrow \varphi(F)$ determines a $\varphi(F)$ -Galois covering space $C \rightarrow S_{1,1}$; here, $\varphi|_F$ is the *monodromy representation* attached to $C \rightarrow S_{1,1}$. If $S'_{1,1}$ is another punctured torus, then by choosing an orientation-preserving homeomorphism $f : S'_{1,1} \rightarrow S_{1,1}$, we can pull back C to obtain a $\varphi(F)$ -Galois covering f^*C of $S'_{1,1}$, and the isomorphism class of the pullback f^*C depends only on the homotopy class of f . The group of homotopy classes of orientation-preserving self-homeomorphisms of $S_{1,1}$ is the *mapping class group* $\Gamma(S_{1,1})$. If the isomorphism class of C is invariant under pulling back by elements of $\Gamma(S_{1,1})$, then the isomorphism class of f^*C is *independent* of the choice of homeomorphism f . At the level of the homomorphism $\varphi|_F$, this is equivalent to the condition that for any $\gamma \in \Gamma(S_{1,1})$, there exists a $\sigma \in \text{Aut}(\varphi(F))$ such that

$$\psi \circ \gamma_* \sim \sigma \circ \psi,$$

where γ_* denotes the outer automorphism of $F \cong \pi_1(S_{1,1})$ induced by $\Gamma(S_{1,1})$, and \sim denotes “is conjugate by some element of G ”. Since the outer action of $\Gamma(S_{1,1})$ on F factors through $\text{Out}^+(F)$, Proposition 2.2 implies that this condition is satisfied.⁴ Thus, the existence of the homomorphism $\varphi : B_4 \rightarrow G$ implies that every punctured torus (in particular, every punctured elliptic curve) has a *canonical* $\varphi(F)$ -Galois cover.

If $G \leq \text{GL}_n(k)$ is a linear group over some ring k , then $\varphi|_F$ can also be viewed as a k -local system of rank n , and the same argument implies that every elliptic curve has a *canonical* k -local system of rank n . The relation between the two perspectives is that covering spaces and local systems are both examples of *locally constant sheaves*.⁵

In genus $g \geq 2$, the mapping class group $\Gamma(S_{g,1})$ plays the role for S_g that B_4 did for $S_{1,1}$. In particular, there is an isomorphism $\Gamma(S_{g,1}) \cong \text{Aut}^+(\pi_1(S_g))$ [16, p. 235], and hence, representations of $\Gamma(S_{g,1})$ correspond to canonical locally constant sheaves on S_g . This gives a geometric interpretation to a result of Funar and Lochak [17, Theorem 1.4].

2.3. The unitary group

Let R denote the ring $\mathbb{Z}[q, q^{-1}]$, and let

$$R_0 := \mathbb{Z}[q + q^{-1}] \subset R.$$

⁴In fact, the outer representation $\Gamma(E^\circ) \rightarrow \text{Out}^+(F)$ is an *isomorphism* [16, Section 8.2.7].

⁵In general, for a connected and locally simply connected topological space X , locally constant sheaves on X are determined by representations of the fundamental group $\rho : \pi_1(E^\circ) \rightarrow \text{Aut}(S)$ for some object S . If S is a set, $\text{Aut}(S)$ is a permutation group, and ρ corresponds to a covering space (a locally constant sheaf of sets). If S is a group, ρ corresponds to an S -local system (a locally constant sheaf of groups). See [48, Section 2.5–2.6].

Definition 2.3 (The Hermitian form h [49, Section 3]). The ring $R = \mathbb{Z}[q, q^{-1}]$ admits an involution given by the unique \mathbb{Z} -algebra automorphism sending $q \mapsto q^{-1}$. We will often denote this involution by $r \mapsto \bar{r}$ for $r \in R$; we will also sometimes call it *conjugation*. The subring of fixed points under the involution is $R_0 = \mathbb{Z}[q + q^{-1}]$. Since $R \cong R_0[t]/(t^2 - (q + q^{-1})t + 1)$, R is a free R_0 -module of rank 2. Let h denote the sesquilinear form antilinear in the second coordinate (relative to the involution) on the free R -module R^3 given by the matrix

$$H := \begin{bmatrix} \frac{(q+1)^2}{q} & -q-1 & 0 \\ -q^{-1}-1 & \frac{(q+1)^2}{q} & -q-1 \\ 0 & -q^{-1}-1 & \frac{(q+1)^2}{q} \end{bmatrix}.$$

Thus, we have $h(v, w) = v^t H \bar{w}$ for all $v, w \in R^3$. Since $H^t = \bar{H}$, we see that h is a Hermitian form.

The matrix H has determinant [49, Lemma 13]

$$\det(H) = \left(\frac{q+1}{q}\right)^3 \frac{q^4 - 1}{q - 1}. \quad (2)$$

The unitary group for h is by definition the group of isometries of the Hermitian space (R^3, h) . As a group scheme, this group should be understood as a subgroup of the restriction of scalars $\text{Res}_{R/R_0} \text{GL}_{3,R}$.

Definition 2.4. The unitary group U_h is the group scheme over R_0 whose B -points (for any R_0 -algebra B) are

$$U_h(B) = \{g \in \text{GL}_3(R \otimes_{R_0} B) \mid h(gv, gw) = h(v, w) \text{ for all } v, w \in (R \otimes_{R_0} B)^3\}.$$

Here, the involution on $R \otimes_{R_0} B$ is induced from R/R_0 . Let SU_h be the subgroup scheme consisting of matrices with determinant 1. If B is any R_0 algebra, let $U_{h,B} := U_h \times_{R_0} B$ denote the base change of U_h to B . For more details on unitary groups, see [49, Section 3], [42, Section 2.3.3], and [11, Exercise 4.4.5].

Recall that the standard unitary group $U(n)$ can be viewed as a real form of GL_n . Here is the analogous result in our case.

Proposition 2.5. *Let k be an R -algebra which is a field such that the images of q, q^{-1} are distinct in k . The base change of U_h to the R_0 -algebra k is isomorphic to $\text{GL}_{3,k}$, and the base change of SU_h to k is isomorphic to $\text{SL}_{3,k}$. In particular, this applies to $k = \mathbb{Q}(q)$.*

Remark 2.6. The assumption that q, q^{-1} are distinct in k is equivalent to assuming that $q \neq \pm 1 \in k$. In particular, the result still holds if q maps to a primitive 4th root of unity, even though h is degenerate in this case.

Proof. For any R_0 -algebra B , $U_{h,R_0}(B)$ is the subgroup of matrices $g \in \mathrm{GL}_3(B \otimes_{R_0} R)$ satisfying $g^t H \bar{g} = H$, where the conjugation on g comes from the conjugation on $B \otimes_{R_0} R$ which in turn is induced from that of R . Similarly, the elements of $\mathrm{SU}_{h,R_0}(B)$ are the matrices in $\mathrm{SL}_3(B \otimes_{R_0} R)$ satisfying the same relation. If B is any k -algebra and q_B denotes the image of q in B , then by our assumption on k , we have $q_B \neq q_B^{-1}$. It follows that $B \otimes_{R_0} R \cong B \times B$, with conjugation given by $(x, y) \mapsto (y, x)$. Writing $g \in \mathrm{GL}_3(B \times B) = \mathrm{GL}_3(B) \times \mathrm{GL}_3(B)$ as (g_1, g_2) , the relation $g^t H \bar{g} = H$ becomes $(g_1^t, g_2^t)(H, \bar{H})(g_2, g_1) = (H, \bar{H})$. This implies $g_1^t H g_2 = H$, and $g_2^t \bar{H} g_1 = \bar{H}$, but since $\bar{H} = H^t$, these two relations are the same, so we are left with the single relation $g_1^t H g_2 = H$. Thus, we can take g_1 to be anything in $\mathrm{GL}_3(B)$, and then g_2 is forced to be $H^{-1} g_1^{-t} H$. This shows that $U_{h,k}(B) = \mathrm{GL}_3(B)$ and $\mathrm{SU}_{h,k}(B) = \mathrm{SL}_3(B)$ as desired. ■

2.4. The representation

The Burau representation arises from the monodromy action on the middle homology of a certain manifold, which is isomorphic to R^3 as abelian groups. The Hermitian form h is constructed using the intersection form on this homology group, relative to which the Burau representation acts via isometries. An explicit description is given below. See [49], [47, Section 2.2] for more details.

Definition 2.7. The Burau representation for B_4 is the representation

$$\rho := \rho_{B,4} : B_4 \rightarrow U_h(R_0) \subset \mathrm{GL}_3(R)$$

given by

$$(\sigma_1, \sigma_2, \sigma_3) \mapsto \left(\left[\begin{array}{ccc} -q & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right], \left[\begin{array}{ccc} 1 & 0 & 0 \\ q & -q & 1 \\ 0 & 0 & 1 \end{array} \right], \left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & q & -q \end{array} \right] \right).$$

One can easily calculate the following:

$$\rho(x) = \begin{bmatrix} -q & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & -q^{-1} \end{bmatrix}, \quad \rho(y) = \begin{bmatrix} 1-q & -q^{-1} & q^{-1} \\ 1-q^2 & -q^{-1} & 0 \\ 1 & -q^{-1} & 0 \end{bmatrix}.$$

We note that $\sigma_1, \sigma_2, \sigma_3$ are all conjugate in B_4 , and hence, $x = \sigma_1 \sigma_3^{-1}$ and its conjugate $y = \sigma_2 \sigma_1 \sigma_3^{-1} \sigma_2^{-1}$ must have determinant 1 under any representation. In particular, the Burau representation restricts to a representation

$$\rho|_F : F \rightarrow \mathrm{SU}_h(R_0).$$

If k is any $\mathbb{Z}[q, q^{-1}]$ -algebra where $q + 1$ is invertible, the vectors

$$v_1 := \begin{bmatrix} 1 \\ q+1 \\ q \end{bmatrix}, \quad v_2 := e_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad v_3 := e_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

give an eigenbasis of k^3 for $\rho(x)$ with corresponding eigenvalues $1, -q, -q^{-1}$, respectively. It follows that $\rho(y)$ has the same eigenvalues, with eigenbasis $\sigma_2 v_1, \sigma_2 v_2, \sigma_2 v_3$. Since $\rho(x)$ has distinct eigenvalues over $\mathbb{Q}(q)$, $\{v_1, v_2, v_3\}$ is orthogonal for h in the sense that $h(v_i, v_j) = 0$ if $i \neq j$. Let $O := [v_1 \ v_2 \ v_3]$, and let k be any $\mathbb{Z}[q, q^{-1}]$ -algebra where $q + 1$ is invertible. Then, $\{v_1, v_2, v_3\}$ is a basis of k^3 , and relative to this basis, the Hermitian form h on k^3 is given by the matrix

$$D := O^t H \bar{O} = \begin{bmatrix} (q + q^{-1})^2 + 2(q + q^{-1}) & 0 & 0 \\ 0 & \frac{(q+1)^2}{q} & 0 \\ 0 & 0 & \frac{(q+1)^2}{q} \end{bmatrix}.$$

2.5. Bootstrapping $\text{Aut}^+(F)$ -invariance to $\text{Aut}(F)$ -invariance

For an R_0 -algebra k_0 , let ρ_{k_0} denote the composition

$$\rho_{k_0} : B_4 \xrightarrow{\rho} U_h(R_0) \rightarrow U_h(k_0),$$

where ρ is the Burau representation. Via Proposition 2.2, for every R_0 -algebra k_0 , ρ_{k_0} gives rise to the $\text{Aut}^+(F)$ -invariant map

$$\rho_{k_0}|_F : F \rightarrow \text{SU}_h(k_0).$$

We will show that if $q + 1$ is invertible in $k_0 \otimes_{R_0} R$, then $\rho_{k_0}|_F$ is in fact characteristic, see Proposition 2.8. Since the unique nontrivial coset of $\text{Aut}(F)/\text{Aut}^+(F)$ consists of automorphisms of determinant -1 , it suffices to show that for the automorphism $\alpha \in \text{Aut}(F)$ given by

$$\alpha(x) = x^{-1}, \quad \alpha(y) = y,$$

there exists an automorphism $\bar{\alpha} \in \text{Aut}(\text{SU}(k_0))$ such that

$$\rho_{k_0}|_F \circ \alpha = \bar{\alpha} \circ \rho_{k_0}|_F.$$

If $X := \rho_{k_0}(x)$ and $Y := \rho_{k_0}(y)$, then we wish to find an automorphism $\bar{\alpha}$ of $\text{SU}_h(k_0)$ sending $(X, Y) \mapsto (X^{-1}, Y)$.

The easiest situation is if one can take $\bar{\alpha}$ to be inner, but it turns out this is impossible because α fails to preserve traces;⁶ thus, we are led to consider non-inner $\bar{\alpha}$. Over \mathbb{C} , $\text{Out}(\text{SU}_h(\mathbb{C})) \cong \mathbb{Z}/2$, the nontrivial class being represented by “complex conjugation with respect to a real structure”.⁷ To describe the desired automorphism $\bar{\alpha} \in \text{SU}_h(k_0)$, we begin by describing the arithmetic version of this conjugation.

Let $k := k_0 \otimes_{R_0} R$, so k is a free k_0 -module of rank 2, and $k_0 \subset k$ is the subalgebra fixed by the involution. We make the assumption that $q + 1$ is invertible in k . Then, the vectors $\{v_1, v_2, v_3\}$ described in Section 2.4 give an eigenbasis in k^3 for X . This is an orthogonal basis of k^3 relative to the Hermitian form h . Let $I := q - q^{-1} \in k$, and let V_0 be the k_0 -span of $\{v_1, v_2, v_3\}$; this is a real structure in the sense that h takes values in k_0 when restricted to V_0 . Thus, $k^3 = V_0 \oplus IV_0$. Define J to be the k_0 -linear automorphism of k^3 given by

$$J := \text{id}_{V_0} \oplus (-\text{id}_{IV_0}). \quad (3)$$

In particular, $Jv_i = v_i$ and $JIV_i = -IV_i$ for all $1 \leq i \leq 3$. Thus, J is the analog of “complex conjugation” relative to the real structure V_0 . Define $\sigma_J \in \text{Aut}(U_h(k_0))$ by

$$\sigma_J(B) = JBJ^{-1} = JBJ.$$

Then, σ_J represents a nontrivial class of $\text{Out}(U_h(k_0))$.⁸ Since v_1, v_2, v_3 is an eigenbasis of X with eigenvalues $1, -q, -q^{-1}$, respectively, we find that v_1, v_2, v_3 are eigenvectors of JXJ with eigenvalues $1, -q^{-1}, -q$, so $\sigma_J(X) = JXJ = X^{-1}$. It is not quite true that $\sigma_J(Y) = Y$, but it turns out this can be fixed by composing by the inner automorphism given by a suitable matrix centralizing X .

Proposition 2.8. *Let k_0 be any R_0 -algebra such that $q + 1$ is invertible in $k := k_0 \otimes_{R_0} R$. Let J be the k_0 -linear endomorphism of k^3 described in (3). Let δ be the matrix*

$$\delta = \begin{bmatrix} q^{-1} & -q^{-2} & 0 \\ 0 & -q^{-2} & 0 \\ 0 & -q^{-2} & q^{-3} \end{bmatrix}.$$

Let $\bar{\alpha}$ be the automorphism of $U_h(k_0) \subset \text{GL}_3(k)$ given by

$$\bar{\alpha}(B) = \delta^{-1}JBJ\delta \quad \text{for } B \in U_h(k_0).$$

⁶The shortest word $w \in F$ we could find for which $\text{tr } \rho(w) \neq \text{tr } \rho(\alpha(w))$ is $w = xyx^{-2}y^2$. In this case, $\text{tr}(\rho(xy x^{-2} y^2)) = q^5 - 2q^4 + q^3 + q^2 - 4q + 4 - 2q^{-1} - q^{-2} + 2q^{-3} - q^{-4}$ but $\text{tr}(\rho(\alpha(xy x^{-2} y^2))) = -q^4 + 2q^3 - q^2 - 2q + 4 - 4q^{-1} + q^{-2} + q^{-3} - 2q^{-4} + q^{-5}$.

⁷Since SU_h is simple and simply connected, its outer automorphisms come from automorphisms of Dynkin diagram. Since its Dynkin diagram is a line, we have $\text{Out}(\text{SU}_h) \cong \mathbb{Z}/2$.

⁸More generally, every real structure V_0 on the Hermitian space (k^3, h) determines, via this construction, an automorphism σ_J of $\text{SU}_h(k_0)$.

Then, $\bar{\alpha}(X) = X^{-1}$ and $\bar{\alpha}(Y) = Y$. In particular, $\rho_{k_0}|_F \circ \alpha = \bar{\alpha} \circ \rho_{k_0}|_F$, and $\rho_{k_0}|_F : F \rightarrow \mathrm{SU}(k_0)$ is characteristic.

Proof. The key point is to find the appropriate matrix δ . One can check that our δ centralizes X , and satisfies the advertised properties. We also note that

$$\delta v_1 = -q^{-2}v_1, \quad \delta v_2 = q^{-1}v_2, \quad \delta v_3 = q^{-3}v_3,$$

so δ is invertible for any R_0 -algebra k_0 . The process of finding δ is a somewhat laborious calculation, which we omit. ■

Remark 2.9. The existence of the matrix δ can also be checked by showing that the traces of $\rho_{k_0}|_F(w)$ and $(\rho_{k_0}|_F \circ \alpha)(w)$ agree for sufficiently long words $w \in F$ [44, Theorems 3.3, 3.4], though this does not shed any light on *why* it should exist. A better understanding of this would give guidance on when one can expect a general representation of B_4 to lead to characteristic quotients of F .

3. Characteristic quotients of F via strong approximation

The Burau representation gives an $\mathrm{Aut}^+(F)$ -invariant map $\rho_{k_0}|_F : F \rightarrow \mathrm{SU}_h(k_0)$ for any R_0 -algebra k_0 . In Proposition 2.8, we showed that this map is moreover characteristic. To obtain finite simple characteristic quotients of F , it remains to understand the image of $\rho_{k_0}|_F$. We will show that for suitable choices of finite fields k_0 , the map is surjective onto $\mathrm{SU}_h(k_0)$. Taking the quotient by the center, we obtain simple characteristic quotients of F .

In this section, we present a rather general method using the strong approximation theorems of Weisfeiler and Pink. For most number fields K , this establishes surjectivity for all but finitely many residue fields k_0 of K . This gives us infinitely many simple characteristic quotients, thus proving Theorem 1.4, but at the cost of a lack of specificity due to the mysterious finite collection of primes that must be excluded at every application of the strong approximation theorem.

In Section 4, we describe a more direct method: by cleverly choosing specializations of q to generators of various finite fields and exploiting the known structure of the maximal subgroups of $\mathrm{SL}_3(\mathbb{F}_q)$ and $\mathrm{SU}_3(\mathbb{F}_q)$, we are able to prove the more general Theorem 1.6.

3.1. Algebraic groups and strong approximation

We recall some basic definitions of reductive group schemes. An algebraic group G over a field k is a smooth affine group scheme over k . It is reductive if $G_{\bar{k}}$ has trivial

unipotent radical, semisimple if $G_{\bar{k}}$ has trivial radical, simple if $G_{\bar{k}}$ has no proper nontrivial connected normal subgroups, and simply connected if any central isogeny from a connected algebraic group to G is trivial. When discussing reductive group schemes, we follow the conventions of Conrad [11, Definition 3.1.1].

Definition 3.1 (Algebraic group schemes). Let S be a scheme. An algebraic group scheme over S is a smooth group scheme affine over S (sometimes, we will say a “smooth affine S -group scheme”). It is reductive if its geometric fibers are connected reductive algebraic groups in the usual sense. It is simple (resp., semisimple, simply connected) if its geometric fibers are simple (resp., semisimple, simply connected).

If v is a place of a number field K , let K_v denote the completion of K at v , and if v is a finite place, let \mathcal{O}_v (resp., k_v) be the ring of integers (resp., residue field) of K_v . Recall that for a set of places S , \mathbb{A}_K^S denotes the ring of adèles away from S : the restricted product $\prod'_{v \notin S} K_v$. Given Proposition 2.1, our general method for producing finite simple characteristic quotients is based on the following strong approximation theorem due to Weisfeiler and Pink.

Theorem 3.2 (Strong approximation for linear groups). *Let K be a number field, let G be a simple, simply connected linear algebraic group over K , and let $\Lambda \subset G(K)$ be a finitely generated Zariski dense subgroup. Let S denote a set of places v of K for which the image of Λ in $G(\mathbb{A}_K^S)$ is not discrete. Suppose K is generated over \mathbb{Q} by the traces $\text{tr Ad } \gamma$. Then, the closure of Λ in $G(\mathbb{A}_K^S)$ is open in $G(\mathbb{A}_K^S)$.*

This theorem is a special case of the results of Weisfeiler [50] and Pink [41, Theorem 0.3].⁹ The condition that K is generated by the traces $\text{tr Ad } \gamma$ exists essentially to guarantee that Λ is not a subgroup of $G(L)$ for some subfield $L \subset K$.¹⁰

Keeping the notation and assumptions as Theorem 3.2, let us further choose an embedding of G into $\text{GL}_{n,K}$. Relative to this embedding, define

$$G(\mathcal{O}_v) := G(K) \cap \text{GL}_n(\mathcal{O}_v),$$

and define $G(k_v)$ to be its image in $\text{GL}_n(k_v)$. For varying v , these groups are related as follows: if \mathcal{G} denotes the schematic closure of G inside $\text{GL}_{n,\mathcal{O}_K}$, then \mathcal{G} admits a unique structure of an affine flat group scheme over \mathcal{O}_K with generic fiber G and $\mathcal{G}(\mathcal{O}_K) = G(K) \cap \text{GL}_n(\mathcal{O}_K)$ (see Proposition A.1 in the appendix). In particular, for varying places v , the groups $G(\mathcal{O}_v)$ (resp., $G(k_v)$) are all obtained as the \mathcal{O}_v (resp., k_v) valued points of the single group scheme \mathcal{G} over \mathcal{O}_K . Moreover, there exists $b \in \mathcal{O}_K$ such that the geometric fibers of $\mathcal{G}_{\mathcal{O}_K[1/b]}$ over $\mathcal{O}_K[1/b]$ are connected reductive

⁹Also, see [31, Window 9, Theorem 2] for a discussion, and [34] for a somewhat weaker result.

¹⁰This ensures that [41, Assumption 0.1] is satisfied.

with identical root data.¹¹ In particular, since G is simple and simply connected, so is $\mathcal{G}_{\mathcal{O}_K[1/b]}$.

Corollary 3.3. *For all but finitely many places v of K , Λ is contained in $G(\mathcal{O}_v)$, and the reduction map $G(\mathcal{O}_v) \rightarrow G(k_v)$ maps Λ surjectively onto $G(k_v)$.*

Proof. Choose generators $\lambda_1, \dots, \lambda_r$ for Λ . There is a finite set of places S_1 , containing all infinite places, such that the images of λ_i have v -integral coordinates in GL_n for every $v \notin S_1$. In particular, Λ is contained in the compact group

$$G\left(\prod_{v \notin S_1} \mathcal{O}_v\right) \subset G(\mathbb{A}_K^{S_1}).$$

Here, this containment should be viewed as occurring inside $\mathrm{GL}_n(\mathbb{A}_K^{S_1})$. Thus, the infinitude of Λ implies that its image in $G(\mathbb{A}_K^{S_1})$ is not discrete. Applying the strong approximation theorem with $S := S_1$, we deduce that Λ is dense in an open subgroup of $G(\mathbb{A}_K^{S_1})$, and hence, there is a finite set of places S_2 such that Λ is dense in $G(\mathcal{O}_v)$ for all $v \notin S_1 \cup S_2$. In particular, possibly excluding the places above b (see preceding paragraph), Hensel's lemma implies that Λ surjects onto the finite group $G(k_v)$ [43, Theorem 3.5.62]. ■

Combining Theorem 3.2 and Proposition 2.1, we arrive at a rather general method for producing invariant quotients.

Proposition 3.4. *Let A be a finitely generated group, and $F \trianglelefteq A$ a finitely generated normal subgroup. Let K be a number field, and let G be a connected, simply connected, simple algebraic group over K embedded inside an algebraic group H . Let*

$$\rho : A \rightarrow H(\bar{K})$$

be a homomorphism. Assume that

- (a) K is generated over \mathbb{Q} by the traces $\mathrm{tr} \mathrm{Ad} \gamma$ for $\gamma \in F$,
- (b) $G \subset \overline{\rho(A)}^{\mathrm{zar}}$,
- (c) $\rho(F) \subset G(K)$,
- (d) $\rho(F) \not\subset Z(G)(K)$.

¹¹More precisely, since the smooth locus is open and G is smooth, \mathcal{G} is smooth over a dense open of $\mathrm{Spec} \mathcal{O}_K$. Next, since the generic fiber is reductive, passing to a possibly smaller dense open, we may assume that the identity components of the fibers are all reductive [11, Proposition 3.1.9], which further implies that all fibers are connected [11, Proposition 3.1.12]. Since \mathcal{G} étale locally admits a split maximal torus, we see that the root datum of geometric fibers of \mathcal{G} are locally constant.

Then, for all but finitely many places v of K , $G(k_v)$, and hence, the simple group $G(k_v)/Z(G(k_v))$ is an A -invariant quotient of F .

Proof. Since G is simple, finite normal subgroups of G are central, so the noncentrality hypothesis on $\rho(F)$ implies that it is infinite, and (b) implies that $\overline{\rho(F)}^{\text{zar}}$ is normal inside G . In particular, $\overline{\rho(F)}^{\text{zar}}$ is a positive-dimensional normal subgroup of G , and hence, it is the entirety of G , so $\rho(F)$ is a finitely generated Zariski dense subgroup of $G(K)$. Fixing an embedding $G \hookrightarrow \text{GL}_{n,K}$, strong approximation implies that for all but finitely many places v of K , $\rho(F) \subset G(\mathcal{O}_v)$ and surjects onto $G(k_v)$, as desired. Possibly excluding finitely many additional primes, $\rho|_F : F \rightarrow G(k_v)$ extends to a map $A \rightarrow \overline{\rho(A)}^{\text{zar}}(k_v)$. It follows from Proposition 2.1 that $\rho|_F : F \rightarrow G(k_v)$ is A -invariant, and so, is its composition with the characteristic quotient $G(k_v) \rightarrow G(k_v)/Z(G(k_v))$. Finally, we note for all but finitely many places v , $G(k_v)/Z(G(k_v))$ is simple [33, Theorem 24.17]. ■

Remark 3.5. Let S_g be a closed surface of genus $g \geq 2$, and let $S_{g,1}$ be a once-punctured S_g . Funar and Lochak's [17, Theorem 1.4] can be understood as applying the above proposition to certain quantum representations of the mapping class group $\Gamma_{g,1}$ of $S_{g,1}$. Using the isomorphism

$$\Gamma_{g,1} \cong \text{Aut}^+(\pi_1(S_g))$$

(see [16, p. 235]), they obtain infinitely many finite simple characteristic quotients of $\pi_1(S_g)$ for every $g \geq 2$.

3.2. Specializations of the Burau representation at roots of unity

For a nonzero complex number $t \in \mathbb{C}^\times$, let ρ_t , h_t , H_t , and D_t denote the specializations of ρ , h , H , D by simply setting $q = t$. If $t \in \mathbb{C}^\times - \{\pm 1\}$, we will write $U_{h,t}$ for the base change

$$U_{h,\mathbb{Z}[t+t^{-1}]} = U_h \times_{R_0} \mathbb{Z}[t+t^{-1}],$$

where $\mathbb{Z}[t+t^{-1}] \subset \mathbb{C}$ is given the R_0 -algebra structure sending $q + q^{-1} \mapsto t + t^{-1}$. Similarly, $\text{SU}_{h,t}$ is the subgroup of matrices of determinant 1. When t lies on the complex unit circle, the behavior of h_t can be summarized as follows.

Proposition 3.6. *If t is a complex number with absolute value $|t| = 1$. Then, h_t is a Hermitian form on \mathbb{C}^3 relative to complex conjugation. We view \mathbb{R} as a $\mathbb{Z}[q + q^{-1}]$ -algebra via the map $q + q^{-1} \mapsto t + t^{-1}$. Then, the following statements hold.*

- (a) h_t is nondegenerate if and only if $t \notin \{-1, i, -i\}$.
- (b) If $0 < \text{Re } t < 1$, then h_t is positive definite, and $U_{h,t}(\mathbb{R}) \cong \text{U}(3)$ is compact.
- (c) If $-1 < \text{Re } t < 0$, then h_t is of indefinite type $(2, 1)$, and $\text{SU}_{h,t}(\mathbb{R}) \cong \text{SU}(2, 1)$ is noncompact.

Proof. The noncompactness of $SU(2, 1)$ can be seen directly from the observation that it contains every matrix of the form

$$\begin{bmatrix} \sqrt{x^2 + 1} & 0 & x \\ 0 & 1 & 0 \\ x & 0 & \sqrt{x^2 + 1} \end{bmatrix}, \quad x \in \mathbb{R},$$

and hence, it is unbounded, hence noncompact. Everything else follows from the formulas for $\det(H)$ and D given in Section 2.4. \blacksquare

Remark 3.7. The notation $U_{h,t}$ is intuitive, but can be misleading. For example, if $t = 1$, $h_t = h_1$ is a nondegenerate Hermitian form on \mathbb{C}^3 , but $U_{h,t}(\mathbb{R})$ is not the unitary group of h_1 . The issue is that in this case $U_{h,1}(\mathbb{R})$ is defined to be a subgroup of 3×3 matrices with coefficients in the non-étale algebra $\mathbb{R} \otimes_{R_0} R \cong \mathbb{R}[T]/(T - 1)^2$. One can show that this is not even a reductive group (see Proposition 3.12 (c)).

Remark 3.8. The element $t + t^{-1}$ may not be integral. In particular, after specialization, the group scheme $U_{h,t}$ may not surject onto $\text{Spec } \mathbb{Z}$.

In [49], Venkataramana studied the images of the specializations of the Burau representation (for various braid groups) when q is a root of unity. Let $\zeta_d := e^{2\pi i/d}$. Then, the groups U_{h,ζ_d} , SU_{h,ζ_d} , PU_{h,ζ_d} are defined over $\mathbb{Z}[\zeta_d + \zeta_d^{-1}]$. In particular, for $d = 3, 4, 6$, these groups are defined over \mathbb{Z} .

Theorem 3.9 (Venkataramana, McMullen, Deligne–Mostow). *For $d = 3, 4, 6$, the image of ρ_{ζ_d} has finite index in $U_{h,\zeta_d}(\mathbb{Z})$.*

Proof. This is [49, Theorem 3], though in the case $d = 3$, this is due to McMullen [35, Theorem 10.3], and Deligne–Mostow [12]. \blacksquare

For other values of d , the image of ρ_{ζ_d} can fail to be arithmetic see [35, Corollary 11.8] for the case $d = 18$. Note that h_{ζ_4} is degenerate and $SU_{h,\zeta_6}(\mathbb{Z})$ is finite, since it is a lattice inside the compact group $SU_{h,\zeta_6}(\mathbb{R})$. Thus, for the purpose of constructing finite simple characteristic quotients, we are led to consider the specialization at ζ_3 .

Proposition 3.10. *The Zariski closure of $\rho_{\zeta_3}(B_4)$ contains SU_{h,ζ_3} .*

Proof. Since $U_{h,\zeta_3}(\mathbb{Z}) \subset GL_3(\mathbb{Z}[\zeta_3])$, the image of the determinant is contained in the finite group $\mathbb{Z}[\zeta_3]^\times$. It follows that $H := \rho_{\zeta_3}^{-1}(SU_{h,\zeta_3}(\mathbb{Z}))$ has finite index in B_4 , and hence, the image of H is a lattice in $SU_{h,\zeta_3}(\mathbb{Z})$. Since $SU_{h,\zeta_3}(\mathbb{R}) \cong SU(2, 1)$ is noncompact and simple, the Borel density theorem implies that $\rho_{\zeta_3}(H)$ is Zariski dense in SU_{h,ζ_3} [36, Corollary 4.5.6]. \blacksquare

Since $\rho(F)$ is not central in SU_{h,ξ_3} , applying strong approximation, we find the following theorem.

Theorem 3.11 (characteristic quotients of degree 1). *For all but finitely many primes p , $SU_{h,\xi_3}(\mathbb{F}_p)$ is a characteristic quotient of F_2 .*

Proof. The groups U_{h,ξ_3} , SU_{h,ξ_3} are group schemes over $\mathbb{Z}[\zeta_3 + \zeta_3^{-1}] = \mathbb{Z}$. Applying Proposition 3.4, to the Burau representation, taking $H = U_{h,\xi_3,\mathbb{Q}}$, $G = SU_{h,\xi_3,\mathbb{Q}}$, shows that $SU_{h,\xi_3}(\mathbb{F}_p)$ is an $\text{Aut}^+(F)$ -invariant quotient. Proposition 2.8 shows that this is in fact a characteristic quotient. ■

The finite group $SU_{h,\xi_3}(\mathbb{F}_p)$ can be described in more familiar terms as follows. Recall that over a finite field \mathbb{F}_q , up to isometry there is a unique Hermitian form on $(\mathbb{F}_{q^2})^3$ relative to the extension $\mathbb{F}_{q^2}/\mathbb{F}_q$ [24, Corollary 10.4]. Let $U_3(\mathbb{F}_q) \subset GL_3(\mathbb{F}_{q^2})$ denote the corresponding group of isometries, and let $SU_3(\mathbb{F}_q)$ be the subgroup of determinant 1 matrices. The following proposition completes the proof of Theorem 1.4.

Proposition 3.12. *For a prime p , the finite group $SU_{h,\xi_3}(\mathbb{F}_p)$ is described as follows.*

- (a) *If $p \equiv 1 \pmod{3}$, then $U_{h,\xi_3}(\mathbb{F}_p) \cong GL_3(\mathbb{F}_p)$, and $SU_{h,\xi_3}(\mathbb{F}_p) \cong SL_3(\mathbb{F}_p)$.*
- (b) *If $p \equiv 2 \pmod{3}$, then $U_{h,\xi_3}(\mathbb{F}_p) \cong U_3(\mathbb{F}_p)$, and $SU_{h,\xi_3}(\mathbb{F}_p) \cong SU_3(\mathbb{F}_p) \subset SL_3(\mathbb{F}_{p^2})$.*
- (c) *If $p = 3$, then $U_{h,\xi_3}(\mathbb{F}_3)$ is an extension of the orthogonal group $O_3(\mathbb{F}_3)$ by the additive group \mathbb{F}_3^6 .*

Proof. At primes $p \equiv 1 \pmod{3}$, the statement follows from Proposition 2.5 where $k = \mathbb{F}_p$, with R -algebra structure given by sending q to an element of order 3.

At primes $p \equiv 2 \pmod{3}$, $\mathbb{Z}[\zeta_3] \times_{\mathbb{Z}} \mathbb{F}_p \cong \mathbb{F}_{p^2}$, so U_{h,ξ_3,\mathbb{F}_p} is the usual unitary group relative to the extension $\mathbb{F}_{p^2}/\mathbb{F}_p$.

At $p = 3$, $\mathbb{Z}[\zeta_3] \otimes_{\mathbb{Z}} \mathbb{F}_3 = \mathbb{F}_3[x]/(x^2 + x + 1)$, where x is the image of $\zeta_3 \otimes 1$. Let $\varepsilon := x - 1$, then this becomes the ring of dual numbers $\mathbb{F}_3[\varepsilon]$ (where $\varepsilon^2 = 0$), with conjugation given by $\varepsilon \mapsto -\varepsilon$. To simplify calculations we may use the orthogonal basis $\{v_1, v_2, v_3\}$ given in Section 2.4 to replace H with the diagonal matrix $D = D_{\zeta_3} = \text{diag}(-1, 1, 1)$. For any field k , write $g \in GL_3(k[\varepsilon])$ as $g_0 + \varepsilon g_1$, where $g_0 \in GL_3(k)$ and $g_1 \in M_3(k)$. The relation $g^t D \bar{g} = D$ becomes

$$g_0^t D g_0 + \varepsilon(g_1^t D g_0 - g_0^t D g_1) = D.$$

Thus, we have $g_0^t D g_0 = D$, and $g_1^t D g_0 = g_0^t D g_1$. The first relation asserts that g_0 lies in the orthogonal group for D , and the second asserts that $g_0^t D g_1$ is symmetric. For fixed g_0 , this is a linear equation in the coordinates of g_1 , hence defines an affine space. For $g_0 = I_3$, this defines an additive group of dimension 6, so the map

$U_{h,\zeta_3,\mathbb{F}_3} \rightarrow O_{3,\mathbb{F}_3}$ sending $g \mapsto g_0$ realizes $U_{h,\zeta_3}(\mathbb{F}_3)$ as an extension of $O_3(\mathbb{F}_3)$ by \mathbb{F}_3^6 . ■

Remark 3.13. By the discussion in Section 2.2.1, the Burau representation ρ corresponds to a *canonical* rank 3 R -local system on punctured elliptic curves with infinite monodromy (i.e., $\rho|_F$ has infinite image). This infinitude was critical in the proof of Theorem 3.11. We note that in general, low rank canonical local systems with infinite monodromy seem to be, in some sense, quite rare: in a recent preprint of Landesman and Litt [28], they show that if $r < \sqrt{g+1}$, then a canonical \mathbb{C} -local system of rank r on $S_{g,n}$ has *finite monodromy* (i.e., the associated representation of $\pi_1(S_{g,n})$ has *finite image* in $GL_r(\mathbb{C})$).

3.3. Specialization at points on the unit circle which are not roots of unity

In the previous section, we showed that infinitely many groups of the form $SU_3(\mathbb{F}_p)$ and $SL_3(\mathbb{F}_p)$ are characteristic quotients of F_2 . In Section 4, we will show the much stronger fact (Theorem 4.3) that implies that for every $q \geq 7$, both $SU_3(\mathbb{F}_q)$ and $SL_3(\mathbb{F}_q)$ are finite simple characteristic quotients of F_2 . However, because strong approximation forces us to discard an indeterminate finite set of primes, we cannot hope to prove such a result using strong approximation. Nonetheless, the method of strong approximation provides a general framework for understanding why something like Theorem 4.3 could be true. In this section, we illustrate this with an intermediate result (Theorem 3.15), obtained by considering more general specializations of the Burau representation.

Theorem 3.14 ([47, Theorem 1]). *For any complex number t which lies on the unit circle, is not a root of unity, and is sufficiently close to 1, $\rho_t(B_4)$ is dense in the Euclidean topology on $U_{h,t}(\mathbb{R}) \subset GL_3(\mathbb{C})$.*

Since $\rho_t(B_4)$ is dense in the Euclidean topology on $U_{h,t}(\mathbb{R})$, it is also Zariski dense. This leads us to the following generalization of Theorem 3.11.

Theorem 3.15 (Aut⁺(F)-Invariant quotients of arbitrary degree). *Let $d \geq 1$ be an integer. Then, the following statements hold.*

- (a) *For infinitely many primes p , $SL_3(\mathbb{F}_{p^d})$ is a characteristic quotient of F .*
- (b) *For infinitely many primes p , $SU_3(\mathbb{F}_{p^d})$ is a characteristic quotient of F .*

Unlike the specialization at ζ_3 , where we work over the explicit ring $\mathbb{Z}[\zeta_3 + \zeta_3^{-1}] = \mathbb{Z}$, here, for every p^d , we must effectively describe an infinite sequence of algebraic numbers t on the unit circle converging to 1 for which the associated number fields have a prime of inertia degree d over p . Moreover, there is also the added complexity of dealing with the trace field $\mathbb{Q}(\text{tr Ad } \rho_t(F))$, which in the $t = \zeta_3$ case

was just \mathbb{Q} ; ideally, this trace field is just $\mathbb{Q}(t + t^{-1})$, but we do not know if this is true. However, it is not difficult to show that it contains $\mathbb{Q}(t^3 + t^{-3})$, and that for appropriately chosen t , we have $\mathbb{Q}(t + t^{-1}) = \mathbb{Q}(t^3 + t^{-3})$. We begin with a few lemmas.

Lemma 3.16. *The trace field $\mathbb{Q}(\text{tr Ad } \rho(\gamma) \mid \gamma \in F)$ contains $\mathbb{Q}(q^3 + q^{-3})$.*

Proof. The trace of the adjoint action of $\gamma \in \text{SU}_h(\mathbb{Q}(q + q^{-1}))$ on the Lie algebra of $\text{SU}_h, \mathbb{Q}(q + q^{-1})$ can be calculated after base change to $\mathbb{Q}(q)$, in which case it can be identified with the trace of the image of γ in $\text{SU}_h(\mathbb{Q}(q)) \cong \text{SL}_3(\mathbb{Q}(q))$ acting on the Lie algebra of $\text{SL}_3, \mathbb{Q}(q)$. This latter Lie algebra consists of the trace 0 matrices, and the adjoint action of γ on a trace zero matrix X is simply given by conjugation by γ . An explicit calculation shows that

$$\begin{aligned} \text{tr Ad } \rho(x) &= q^2 - 2q + 2 - 2q^{-1} + q^{-2}, \\ \text{tr Ad } \rho(x^2) &= q^4 + 2q^2 + 2 + 2q^{-2} + q^{-4}. \end{aligned}$$

From this, one calculates that

$$(\text{tr Ad } \rho(x))^2 - \text{tr Ad } \rho(x^2) - 6 \text{tr Ad } \rho(x) = -4q^3 - 4q^{-3}.$$

It follows that the trace field contains $\mathbb{Q}(q^3 + q^{-3})$. ■

Lemma 3.17. *Let $t \in \bar{\mathbb{Q}}$ satisfy $|t| = 1$. Suppose $\mathbb{Q}(t + t^{-1})$ is Galois over \mathbb{Q} and that $t + t^{-1}$ has nonzero trace over \mathbb{Q} . Then, $\mathbb{Q}(t + t^{-1}) = \mathbb{Q}(t^3 + t^{-3})$.*

Proof. The result is clear if $t = \pm 1$, so we may assume $t \neq \pm 1$. Let $s_i := t^i + t^{-i}$. Let $d := [\mathbb{Q}(s_1) : \mathbb{Q}]$. We have the relation $s_3 = s_1^3 - 3s_1$, or equivalently, s_1 is a root of the polynomial $f(X) := X^3 - 3X - s_3 \in \mathbb{Q}(s_3)[X]$. Since $\mathbb{Q}(s_1)/\mathbb{Q}(s_3)$ is Galois, either $f(X)$ is irreducible over $\mathbb{Q}(s_3)$, or $f(X)$ is totally split. Moreover, it is irreducible if and only if $\deg_{\mathbb{Q}}(s_3) = d/3$. We will show that the trace condition on s_1 implies that this is impossible, and hence, $f(X)$ must be totally split, which is to say that $\mathbb{Q}(s_1) = \mathbb{Q}(s_3)$.

If d is not divisible by 3, then $f(X)$ cannot be irreducible, so we may assume $d = 3m$ for some integer m . In this case, $f(X)$ is irreducible if and only if $\deg_{\mathbb{Q}}(s_3) = m$, in which case the minimal polynomial of s_3 gives a \mathbb{Q} -linear dependence relation between $1, s_3, s_3^2, \dots, s_3^m$. Using the relation $s_3 = s_1^3 - 3s_1$, we find that for any $k \geq 1$,

$$s_3^k = (s_1^3 - 3s_1)^k = s_1^{3k} + \binom{k}{1}(-3)s_1^{3k-2} + \binom{k}{2}(-3)^2 s_1^{3k-4} + \dots + \binom{k}{k}(-3)^k s_1^k. \quad (4)$$

We will show that $1, s_3, \dots, s_3^m$ are \mathbb{Q} -linearly independent. Since $1, s_1, \dots, s_1^{d-1}$ is a \mathbb{Q} -basis of $\mathbb{Q}(s_1)$, clearly $1, s_3, \dots, s_3^{m-1}$ are \mathbb{Q} -linearly independent. Using the

minimal polynomial of s_1 to write $s_1^{3m} = s_1^d$ as a linear combination of $1, s_1, \dots, s_1^{d-1}$, the coefficient of s_1^{d-1} is equal to $\text{tr}_{\mathbb{Q}(s_1)/\mathbb{Q}}(s_1)$, and hence nonzero. Using (4), this implies that the coefficient of s_1^{d-1} in s_3^m is nonzero, and hence, $1, s_3, s_3^2, \dots, s_3^m$ is \mathbb{Q} -linearly independent. Thus, $f(X)$ is totally split, and hence, $\mathbb{Q}(s_1) = \mathbb{Q}(s_3)$. ■

Proof of Theorem 3.15. For integers $n, k \geq 1$, let ζ be a primitive n th root of unity, and let

$$t_{\zeta,k} := \frac{\zeta + k}{\zeta^{-1} + k}.$$

Clearly, $t_{\zeta,k} \in \mathbb{Q}(\zeta_n)$, $|t_{\zeta,k}| = 1$, $\arg(t_{\zeta,k}) = 2 \arg(\zeta + k)$, and $t_{\zeta,k} \rightarrow 1$ as $k \rightarrow \infty$. Moreover, for all k sufficiently large, we have the following.

- (a) $t_{\zeta,k}$ generates the cyclotomic field $\mathbb{Q}(\zeta_n)$.
Indeed, the map $\zeta \mapsto t_{\zeta,k}$ is $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -equivariant, and for large k the map $\zeta \mapsto t_{\zeta,k}$ is injective on primitive n th roots of unity. This shows that the Galois orbit of $t_{\zeta,k}$ has size $\phi(n) = [\mathbb{Q}(\zeta_n) : \mathbb{Q}]$.
- (b) $s_{\zeta,k} := t_{\zeta,k} + t_{\zeta,k}^{-1}$ generates the maximal totally real subfield of $\mathbb{Q}(\zeta_n)$, and has nonzero trace over \mathbb{Q} .
Indeed, $\text{tr}_{\mathbb{Q}}(s_{\zeta,k}) = 2 \text{tr}_{\mathbb{Q}} t_{\zeta,k}$, which is nonzero since for $k > 2$, each Galois conjugate of $t_{\zeta,k}$ has positive real part. If $n \leq 2$, the claim is trivial, so assume $n \geq 3$. Then, as ζ runs over all $\frac{\phi(n)}{2}$ primitive n th roots of unity with positive imaginary part, for sufficiently¹² large k , the real parts of $t_{\zeta,k}$ are all distinct, and hence, the elements $s_{\zeta,k}$ are all distinct. Thus, $s_{\zeta,k}$ is a real number of degree $\geq \frac{\phi(n)}{2}$, and hence, it must generate the maximal totally real subfield of $\mathbb{Q}(\zeta_n)$.

Thus, for fixed n , let $\zeta_n := e^{2\pi i/n}$. Then, choosing k large enough, we may apply Stoimenow's Theorem 3.14 to deduce that the representation

$$\rho_{t_{\zeta_n,k}} : B_4 \rightarrow U_{h,t_{\zeta_n,k}}(\mathbb{Q}(s_{\zeta_n,k}))$$

is Zariski dense. Since $\mathbb{Q}(s_{\zeta_n,k})$ is Galois over \mathbb{Q} with nonzero trace, Lemmas 3.16 and 3.17 imply that the trace field of $\rho_{t_{\zeta_n,k}}$ is equal to $\mathbb{Q}(s_{\zeta_n,k})$. Propositions 3.4 and 2.8 then imply that for all but finitely many primes \mathfrak{p} of $\mathbb{Q}(s_{\zeta_n,k})$ with residue field κ , the finite group $\text{SU}_{h,t_{\zeta_n,k}}(\kappa)$ is a characteristic quotient of F .

To complete the proof, we want to show that we can choose n (and $k \gg 0$) such that amongst the primes \mathfrak{p} of $\mathbb{Q}(s_{\zeta_n,k})$ with residue field of degree d over the prime

¹²Let S denote the set of primitive n th roots of 1 with positive imaginary part. Here, we should choose k so that every line passing through $\zeta + k, \zeta' + k$ for any pair $\zeta, \zeta' \in S$ does not pass through $0 \in \mathbb{C}$. This can always be done for k large enough relative to n .

field, infinitely many split (resp., remain inert). To do this, let $p > 3$ be a prime satisfying $p \equiv 1 \pmod{2d}$, let $n := 3p$, and k sufficiently large as above. The extension $\mathbb{Q}(t_{\xi_n, k})/\mathbb{Q}(s_{\xi_n, k})$ induces the following surjection of Galois groups:

$$\mathrm{Gal}(\mathbb{Q}(t_{\xi_n, k})/\mathbb{Q}) \cong (\mathbb{Z}/n)^\times \rightarrow (\mathbb{Z}/n)^\times / \{\pm 1\} \cong \mathrm{Gal}(\mathbb{Q}(s_{\xi_n, k})/\mathbb{Q}).$$

Note $(\mathbb{Z}/n)^\times \cong (\mathbb{Z}/p)^\times \times (\mathbb{Z}/3)^\times$. Let $x \in (\mathbb{Z}/p)^\times$ be an element of order $2d$, and let $\alpha \in \mathrm{Gal}(\mathbb{Q}(t_{\xi_n, k})/\mathbb{Q}) \cong (\mathbb{Z}/n)^\times$ be the element corresponding to $(x, -1) \in (\mathbb{Z}/p)^\times \times (\mathbb{Z}/3)^\times$. Then, α has order $2d$ but maps to an element of order d in $\mathrm{Gal}(\mathbb{Q}(s_{\xi_n, k})/\mathbb{Q}) \cong (\mathbb{Z}/n)^\times / \{\pm 1\}$. By Chebotarev density, infinitely many primes \mathfrak{p} of $\mathbb{Q}(s_{\xi_n, k})$ have Frobenius element α ; such primes have residue degree d and remain inert in $\mathbb{Q}(t_{\xi_n, k})$. Using similar arguments as in Proposition 3.12, for all but finitely many such primes, we have $\mathrm{SU}_{h, t_{\xi_n, k}}(\kappa) \cong \mathrm{SU}_3(\mathbb{F}_{p^d})$, where $\kappa \cong \mathbb{F}_{p^d}$ denotes the residue field of \mathfrak{p} .

To obtain quotients of type SL_3 , the element α^2 of $\mathrm{Gal}(\mathbb{Q}(t_{\xi_n, k})/\mathbb{Q}) \cong (\mathbb{Z}/n)^\times$ (corresponding to $(x^2, 1)$) has order d in $\mathrm{Gal}(\mathbb{Q}(t_{\xi_n, k}))$ and its image in $\mathrm{Gal}(\mathbb{Q}(s_{\xi_n, k}))$ also has order d . Again, by Chebotarev density, we obtain infinitely many primes with Frobenius element α^2 . For such primes, we have $\mathrm{SU}_{h, t_{\xi_n, k}}(\kappa) \cong \mathrm{SL}_3(\mathbb{F}_{p^d})$. ■

4. Proof of Theorem 1.6

Recall that $R_0 := \mathbb{Z}[q + q^{-1}]$ is the subring of $R := \mathbb{Z}[q^{\pm 1}]$ fixed by the involution sending $q \mapsto q^{-1}$. The Burau representation

$$\rho : B_4 \rightarrow U_h(R_0) \subset \mathrm{GL}_3(R)$$

sends $x, y \in F := \langle x, y \rangle \subset B_4$ to the matrices

$$X := \rho(x) = \begin{bmatrix} -q & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & -q^{-1} \end{bmatrix}, \quad Y := \rho(y) = \begin{bmatrix} 1 - q & -q^{-1} & q^{-1} \\ 1 - q^2 & -q^{-1} & 0 \\ 1 & -q^{-1} & 0 \end{bmatrix}.$$

These matrices have eigenvalues $1, -q, -q^{-1}$, and $Y = S_2 X S_2^{-1}$, where

$$S_2 := \rho(\sigma_2) = \begin{bmatrix} 1 & 0 & 0 \\ q & -q & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

To prove Theorem 1.6, we will show how to obtain, for all sufficiently large finite fields k_0 , suitable specializations of the Burau representation which give $\mathrm{SL}_3(k_0)$ and $\mathrm{SU}_3(k_0)$ as characteristic quotients of F . In our formalism, such specializations should be understood as a base change. We recall the setup.

Let $r := q + q^{-1} \in R_0$. Thus, R is free of rank 2 over $R_0 = \mathbb{Z}[r]$, generated by a root of the polynomial $f(T) := T^2 - rT + 1 \in R_0[T]$. Let k_0 be a finite field of characteristic p . For $s \in k_0$, let

$$\mathrm{SU}_{h_s} := \mathrm{SU}_h \times_{R_0} k_0$$

be the base change of SU_h to the R_0 -algebra k_0 with R_0 -algebra structure given by the map sending $r \mapsto s$. Let k be the quadratic k_0 -algebra $k := R \otimes_{R_0} k_0 = k_0[T]/f(T)_{r:=s}$, with involution inherited from R/R_0 . Let $t \in k$ be a fixed root of $f(T)_{r:=s}$; this is guaranteed to be a unit, and the other root is t^{-1} . Thus, t, s satisfy the relation

$$s = t + t^{-1}.$$

To prove Theorem 1.6, we first need to understand for which values of s do the finite groups $\mathrm{SU}_{h_s}(k_0)$ give groups of type $\mathrm{SL}_3(\mathbb{F}_q)$ or $\mathrm{SU}_3(\mathbb{F}_q)$.

Lemma 4.1. *Let k_0 be a finite field. Then,*

$$\mathrm{SU}_{h_s}(k_0) \cong \begin{cases} \mathrm{SL}_3(k_0) & \text{if } T^2 - sT + 1 \text{ splits into coprime factors over } k_0, \\ \mathrm{SU}_3(k_0) & \text{if } T^2 - sT + 1 \text{ is irreducible over } k_0 \text{ and } t^2 + 1 \neq 0. \end{cases}$$

If k_0 has odd characteristic, $T^2 - sT + 1$ splits into coprime factors if and only if $s^2 - 4$ is a square in k_0^\times , and is irreducible if $s^2 - 4$ is a nonsquare. If k_0 has characteristic 2, $T^2 - sT + 1$ splits into coprime factors over k_0 if and only if $s \neq 0$ and $\mathrm{Tr}_{k_0/\mathbb{F}_2}(s^{-1}) = 0$, and is irreducible if and only if $s \neq 0$ and $\mathrm{Tr}_{k_0/\mathbb{F}_2}(s^{-1}) = 1$.

Remark 4.2. If $T^2 - sT + 1$ is a square, $k \cong k_0[\varepsilon]/\varepsilon^2$, in which case by arguments similar to that used in Proposition 3.12, we find that SU_{h_s} is nonreductive.

Proof. If $f(T)_{r:=s} = T^2 - sT + 1$ splits into distinct factors, then $k \cong k_0 \times k_0$ and by Proposition 2.5, we have $\mathrm{SU}_{h_s} \cong \mathrm{SL}_{3,k_0}$. If $T^2 - sT + 1$ is irreducible, then $k = k_0(t)$ is a field, and SU_{h_s} is the special unitary group of the Hermitian space (k^3, h_s) . From (2), we find that h_s is nondegenerate if and only if $(t+1)(t^2+1) \neq 0$. Since t generates the extension k/k_0 , $t \neq -1$, so as long as $t^2+1 \neq 0$, h_s is nondegenerate, in which case $\mathrm{SU}_{h_s}(k_0) \cong \mathrm{SU}_3(k_0)$.

In odd characteristic, the discriminant of $T^2 - sT + 1$ is $s^2 - 4$, and hence, it splits into coprime factors if and only if $s^2 - 4$ is a square in k_0^\times , and is irreducible if $s^2 - 4$ is a nonsquare. In characteristic 2, $T^2 - sT + 1$ is a square if and only if $s = 0$. If $s \neq 0$, then by the change of variable $X = T/s$, $T^2 - sT + 1$ is split over k_0 if and only if $g(X) := X^2 + X + s^{-2}$ is split, which happens if and only if s^{-2} can be written as $a^2 + a$ for some $a \in k_0$. The map $k_0 \rightarrow k_0$ sending $x \mapsto x^2 + x$ is \mathbb{F}_2 -linear, and its image is precisely the kernel of $\mathrm{Tr}_{k_0/\mathbb{F}_2} : k_0 \rightarrow \mathbb{F}_2$. Thus, we

find that $T^2 - sT + 1$ splits into coprime factors over k_0 if and only if $s \neq 0$ and $\text{Tr}_{k_0/\mathbb{F}_2}(s^{-1}) = 0$, and is irreducible if $s \neq 0$ and $\text{Tr}_{k_0/\mathbb{F}_2}(s^{-1}) = 1$. ■

Our main Theorem 1.6 is implied by the following two results.

Theorem 4.3. *Let $k_0 := \mathbb{F}_{p^d}$ be a finite field such that $p^d \geq 9$. Then, there exist elements $s, s' \in k_0$ satisfying $t^4 \neq 1$, where $s = t + t^{-1}$, and $(t')^4 \neq 1$, where $s' = t' + t'^{-1}$ such that*

- (a) $T^2 - sT + 1$ splits into coprime factors over k_0 and $\text{SU}_{h_s}(k_0) \cong \text{SL}_3(k_0)$ is generated by the images of X, Y ,
- (b) $T^2 - s'T + 1$ is irreducible over k_0 and $\text{SU}_{h_{s'}}(k_0) \cong \text{SU}_3(k_0)$ is generated by the images of X, Y .

Moreover, (a) is also attainable for $k_0 = \mathbb{F}_8$, and (b) is attainable for $k_0 = \mathbb{F}_4$.

The remaining cases of Theorem 1.6 were checked using GAP [18].

Proposition 4.4. *For a prime power q , let Z_q denote a generator of the multiplicative group \mathbb{F}_q^\times .*

- (a) *Let $t := Z_{64}^7$, and $s := t + t^{-1} \in \mathbb{F}_8$. Then, $\text{SU}_{h_s}(\mathbb{F}_8) \cong \text{SU}_3(\mathbb{F}_8)$ is generated by the images of X, Y .*
- (b) *Let $t := Z_{49}^6$, and $s := t + t^{-1} \in \mathbb{F}_7$. Then, $\text{SU}_{h_s}(\mathbb{F}_7) \cong \text{SU}_3(\mathbb{F}_7)$ is generated by the images of X, Y .*
- (c) *Let $t := Z_7^2$, and $s := t + t^{-1} \in \mathbb{F}_7$. Then, $\text{SU}_{h_s}(\mathbb{F}_7) \cong \text{SL}_3(\mathbb{F}_7)$ is generated by the images of X, Y .*
- (d) *Let $t := Z_{25}^8$, and $s := t + t^{-1} \in \mathbb{F}_5$. Then, $\text{SU}_{h_s}(\mathbb{F}_5) \cong \text{SU}_3(\mathbb{F}_5)$ is generated by the images of X, Y .*
- (e) *The groups $\text{PSL}_3(\mathbb{F}_2), \text{PSL}_3(\mathbb{F}_3), \text{PSL}_3(\mathbb{F}_4), \text{PSL}_3(\mathbb{F}_5), \text{PSU}_3(\mathbb{F}_2), \text{PSU}_3(\mathbb{F}_3)$ are not characteristic quotients of F_2 .*

Proof. The specializations (a)–(d) were checked explicitly in GAP [18]. For each group G in (e), we checked that for every surjection $\varphi : F_2 \rightarrow G$, the class of φ mod $\text{Aut}(G)$ has a proper stabilizer in $\text{Aut}(F_2)$. ■

Proof of Theorem 4.3. For $a, d \in \mathbb{Z}_{\geq 2}$, we say that a prime divisor $\ell \mid a^d - 1$ is *primitive* if ℓ does not divide $\prod_{i=1}^{d-1} (a^i - 1)$. By a result of Zsigmondy [51], for any $a, d \in \mathbb{Z}_{\geq 2}$, $a^d - 1$ always admits a primitive prime divisor unless $(a, d) = (2, 6)$ or $d = 2$ and $a + 1$ is a power of 2. Any such prime divisor ℓ satisfies $\ell \geq d + 1$. We will use this result repeatedly without mention.

We will also often use X, Y to denote their images in $\text{SU}_{h_s}(k_0)$ via the relevant specialization, and similarly, we will use $v_1 = (1, q + 1, q), v_2 = e_1, v_3 = e_3$ to denote the eigenbasis for X defined in Section 2.4, with respective eigenvalues $1, -q, -q^{-1}$.

We begin with (a). Here, we assume that

$$Q := p^d \neq 2, 3, 5, 8, \quad (5)$$

and set

$$\kappa := \gcd(2, p - 1).$$

(a1) We can find $t' \in \mathbb{F}_{p^{2d}}^\times$ such that $-q = -t'$ has order $Q + 1$. With this choice,

$$|X| = |Y| = Q + 1 \geq 5.$$

The spectrum of X in fact implies that

$$\text{the smallest } j \geq 1 \text{ such that } X^j \text{ acts as a scalar is } Q + 1. \quad (6)$$

If $p = 2$, then $t' = q = -q$ has order $Q + 1 \geq 5$ by (5). If $p > 2$ and $Q \equiv 1 \pmod{4}$, then $t' = q$ has order $(Q + 1)/2 \geq 5$ by (5). If $p > 2$ but $Q \equiv 3 \pmod{4}$, then $t = q'$ has order $Q + 1 \geq 8$ by (5). Thus, in all cases, we have

$$(t')^4 \neq 1, \quad (t')^{Q-1} \neq 1. \quad (7)$$

We also have $q^{-1} = q^Q$, and so,

$$s' = t' + (t')^{-1} = q + q^{-1} = q + q^{p^d} \in \mathbb{F}_{p^d} = k_0.$$

Furthermore, if $p > 2$ then $(s')^2 - 4 = (t' - (t')^{-1})^2$ is a non-square in k_0 . Otherwise, we would have $t' - (t')^{-1} \in k_0$, $s' = t' + (t')^{-1} \in k_0$, and so, $q = t' \in k_0$. This would imply $(t')^{Q-1} = 1$, contradicting (7). If $p = 2$, then the polynomial $T^2 - s'T + 1$ has roots t, t^{-1} which are not in k_0 , and hence, it does not split over k_0 .

Note that, under the extra assumption

$$d \geq 2 \quad \text{and} \quad (p, d) \neq (2, 3), \quad (8)$$

$p^{2d} - 1$ admits a primitive prime divisor

$$\ell \geq 2d + 1 \geq 5,$$

which certainly divides $Q + 1$.

(a2) The preceding analysis in our paper shows that, with this choice $q = t'$,

$$X, Y \in G := \text{SU}_{h_{s'}}(k_0) \cong \text{SU}_3(k_0).$$

We can also verify this independently. Consider the basis (v_1, e_1, e_3) of $k^3 = \mathbb{F}_{p^{2d}}^3$ in which X and Y act via the matrices

$$X' := \begin{pmatrix} 1 & 0 & 0 \\ 0 & -q & 0 \\ 0 & 0 & -q^{-1} \end{pmatrix} \quad \text{and} \quad Y' := \begin{pmatrix} 1 - q - q^{-1} & 1 - q & 0 \\ 0 & 0 & q^{-1} \\ q^2 - q + 1 - q^{-1} & q^2 - q + 1 & 0 \end{pmatrix},$$

respectively. Then, X' and Y' preserve the Hermitian form $u \circ w$ on k^3 (semi-linear on w , i.e., $u \circ (\lambda w) = \lambda^{p^d} (u \circ w)$), with Gram matrix

$$\begin{pmatrix} q + q^{-1} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

in this basis, which is $1/(q + 2 + q^{-1})$ times the form h (represented by the matrix D in Section 2.4).

(a3) We will now show that

$$\langle X, Y \rangle = G \cong \mathrm{SU}_3(k_0).$$

Assume the contrary:

$$\langle X, Y \rangle \leq M$$

for some maximal subgroup M of G . We now apply the classification of maximal subgroups of $G \cong \mathrm{SU}_3(Q)$, as listed in [5, Tables 8.5, 8.6], to M .

Note that, when $Q = p^d = p$, we have $p \geq 7$ by (5), and so, the order of X in $G/Z(G)$ is $Q + 1 \geq 8$ by (6). This implies that M does not belong to class \mathcal{S} in [5, Table 8.6].

(a4) By our choice, $q^4 \neq 0, 1$. In particular, the eigenvalues $1, -q, -q^{-1}$ are pairwise distinct, and thus any line in k^3 , that is, X -invariant must be spanned by e_1, v_1 , or e_3 . (Here, we have specialized $k \mapsto \mathbb{F}_{p^{2d}}$.) Note that $Y(e_1) \notin ke_1, Y(e_3) \notin ke_3$. Next,

$$Y(v) = (1 - q - q^{-1})e_1 + (-q^2 - q^{-1})e_2 + (-q^{-1})e_3.$$

So, if $Y(v) \in kv$, we must have that $Y(v) = -q^{-2}v$, and hence, $-q^2 - q^{-1} = -q^{-1} - q^{-2}$, i.e., $q^4 = 1$, a contradiction. We have shown that there is no line in k^3 , that is, invariant under both X and Y . In particular, M fixes no line in k^3 , and hence, cannot belong to any member of class \mathcal{C}_1 in [5, Table 8.5].

(a5) Suppose M is a member of class \mathcal{C}_2 in [5, Table 8.5], then $M \cong C_{Q+1}^2 \rtimes \mathfrak{S}_3$, the stabilizer of a decomposition of k^3 into an orthogonal sum of three anisotropic lines. Any element of M that permutes these three lines cyclically have central order 3 in G . By (6), this is not the case for X . So, X , being an element in M , must fix at least one of these three lines, again a contradiction.

By assumption, $Q \neq 2, 8$, which implies that $Q + 1$ does not divide $3(Q^2 - Q + 1)$. Since $|X| = Q + 1$, this shows that M cannot belong to class \mathcal{C}_3 in [5, Table 8.5].

Suppose M belongs to class \mathcal{C}_6 in [5, Table 8.5]. Then, $M \leq 3_+^{1+2} \rtimes 2A_4 < \mathrm{SU}_4(2)$. In particular, X , as an element of M , must have order dividing 9 or 12, and hence, $Q \in \{2, 3, 5, 8, 11\}$. Using (5), we are left with $Q = 11$, in which case X has order 12. However, any element of order 12 of $3_+^{1+2} \rtimes 2A_4$ does not possess an

eigenvalue 1 in any 3-dimensional irreducible representation over \mathbb{C} or $\overline{\mathbb{F}}_{11}$ (as one can check using GAP [18]).

Suppose M is a member of type SU_3 in class \mathcal{C}_5 in [5, Table 8.5]. In this case, $Q = Q_0^m$ for some prime $m \geq 3$, and so, (8) holds. In this case, $p^{2d} - 1$ has a primitive prime divisor $\ell \geq 5$ which divides $|X| = Q + 1$. Now, the element $X' := X^{(Q+1)/\ell}$ of order ℓ must belong to $SU_3(Q_0)$. The primitivity of ℓ implies that $|X'| = \ell$ does not divide $Q_0^2 - 1 = p^{2d/m} - 1$. Hence, ℓ must divide $Q_0^3 + 1 = p^{3d/m} + 1$ (and so, $m = 3$). In such a case, any element of order ℓ in $SU_3(Q_0)$ has spectrum of the form $\{\lambda, \lambda^{-Q_0}, \lambda^{Q_0^2}\}$ for some $1 \neq \lambda \in \overline{\mathbb{F}}_p^\times$. But this is a contradiction, since 1 is still an eigenvalue of X' .

(a6) Thus, $p > 2$ and M is a member of type SO_3 in class \mathcal{C}_5 in [5, Table 8.5]:

$$M \cong C_a \times SO_3(Q),$$

where $C_a = \langle z \rangle$ has order $a \mid \gcd(3, Q^2 - 1)$, and z acts as a scalar α on k^3 . In this case, observe that

$$X, Y \in SO_3(Q). \quad (9)$$

Indeed, suppose, for instance, that $X = zX_1$ with $X_1 \in SO_3(Q)$ but $\alpha \neq 1$. As z is central of order coprime to p and $|X| = Q + 1$, we see that X_1 is a semisimple element in $SO_3(Q)$. As such, X_1 has spectrum $\{1, \beta, \beta^{-1}\}$ on k^3 for some $\beta \in \overline{\mathbb{F}}_p^\times$. Thus,

$$\{1, -q, -q^{-1}\} = \{\alpha, \alpha\beta, \alpha\beta^{-1}\}.$$

But $\alpha \neq 1$, so α must be either $-q$ or $-q^{-1}$, and so, $3 = |\alpha| = Q + 1$ by the choice of $-q = -t'$, a contradiction.

The inclusion (9) implies that k^3 supports a non-degenerate symmetric bilinear form (\cdot, \cdot) which is invariant under both X and Y . Recall that e_1, e_3 , and v_1 are eigenvectors of X with eigenvalues $-q, -q^{-1}$, and 1, respectively. As $q^2 \neq 1$, we must have

$$(e_1, e_1) = (e_3, e_3) = 0.$$

Similarly, the eigenvectors $S(e_1) = e_1 + qe_2$ and $S(e_3) = e_2 + e_3$ of Y are isotropic, whence

$$(e_1, e_2) = -(q/2)(e_2, e_2), \quad (e_2, e_3) = -(1/2)(e_2, e_2).$$

We also have $v_1 \perp e_3$, which implies

$$(e_1, e_3) = -(q + 1)(e_2, e_3) = ((q + 1)/2)(e_2, e_2).$$

Thus, denoting $(e_2, e_2) = 2\varepsilon$ we see that the Gram matrix of (\cdot, \cdot) in the basis (e_1, e_2, e_3) of k^3 is

$$\Gamma := \varepsilon \begin{pmatrix} 0 & -q & q + 1 \\ -q & 2 & -1 \\ q + 1 & -1 & 0 \end{pmatrix};$$

in particular, $\varepsilon \neq 0$. Comparing the $(2, 3)$ -entries of Γ and ${}^t Y \Gamma Y$, we see that $-q^{-2} = -1$, i.e., $q^2 = 1$, again a contradiction.

We have therefore shown that $\langle X, Y \rangle = \text{SU}_{h_s}(k_0)$.

We now consider part (b). Here, we assume that

$$Q := p^d \neq 2, 3, 4, 5, 7. \quad (10)$$

(b1) We can find $t \in \mathbb{F}_{p^{2d}}^\times$ such that $-q = -t$ has order $Q - 1$. With this choice,

$$|X| = |Y| = Q - 1 \geq 7.$$

The spectrum of X in fact implies that

$$\text{the smallest } j \geq 1 \text{ such that } X^j \text{ acts as a scalar is } Q - 1. \quad (11)$$

If $Q \equiv 1 \pmod{4}$ or $2|Q$, then $t = q$ has order $Q - 1 \geq 7$ by (10). If $Q \equiv 3 \pmod{4}$, then $t = q$ has order $(Q - 1)/2 \geq 5$ by (10). Thus, in all cases, we have

$$t^4 \neq 1.$$

Furthermore,

$$s = t + t^{-1} = q + q^{-1} \in \mathbb{F}_{p^d} = k_0,$$

and $s^2 - 4 = (t - t^{-1})^2$ is a square in k_0 . If $p = 2$, then the polynomial $T^2 - sT + 1$ has roots t, t^{-1} which are in k_0 , and hence, it splits over k_0 .

(b2) With the choice $q = t$ made above, clearly, we have

$$X, Y \in H := \text{SL}_3(k_0).$$

We will now show that

$$\langle X, Y \rangle = H.$$

Assume the contrary:

$$\langle X, Y \rangle \leq M$$

for some maximal subgroup M of H . We now apply the classification of maximal subgroups of $H \cong \text{SL}_3(Q)$, as listed in [5, Tables 8.3, 8.4], to M .

(b3) Note that the order of X in $G/Z(G)$ is $Q - 1 \geq 7$ by (11). This implies that M does not belong to class \mathcal{S} in [5, Table 8.4]. (Indeed, the possibility $M = C_{\text{gcd}(3, Q-1)} \times \text{PSL}_2(7)$ can occur only when $Q = p \equiv 1, 2, 4 \pmod{7}$, which, together with (10), implies that $Q - 1 \geq 10$.)

Next, by our choice, $q^4 \neq 0, 1$. Hence, the arguments in part (a4) show that there is no line in k^3 , that is, invariant under both X and Y . Here, we specialize $k \mapsto \mathbb{F}_Q = \mathbb{F}_{p^d}$. Suppose that $\langle X, Y \rangle$ fixes a plane $U \subset k^3$. Since X acts on U and its eigenvalues

all belong to k_0 , two of these eigenvalues occur in U . If one of these eigenvalues is $-q^{-1}$, then $e_3 \in U$. In this case, $q^{-1}e_1 = Y(e_3) \in U$, so $e_1 \in U$. In turn, this implies that

$$(1 - q^2)e_2 = Y(e_1) - (1 - q)e_1 - e_3 \in U,$$

so $U = k^3$, a contradiction. Thus, $X|_U$ affords the eigenvalues $-q$ and 1 , and hence, e_1, v_1 span U . But

$$Y(e_1) = (1 - q)e_1 + (1 - q^2)e_2 + e_3$$

is not a linear combination of e_1 and v_1 , and so, $Y(e_1) \notin U$, again a contradiction. We have shown that M cannot fix any line or plane in k^3 , and hence, cannot belong to members of class \mathcal{C}_1 in [5, Table 8.3].

Suppose M is a member of class \mathcal{C}_2 in [5, Table 8.3], then $M \cong C_{Q-1}^2 \rtimes \mathfrak{S}_3$, the stabilizer of a decomposition of k^3 into a direct sum of three lines. Any element of M that permutes these three lines cyclically have central order 3 in G . By (11), this is not the case for X . So, X , being an element in M , must fix at least one of these three lines, again a contradiction.

By assumption, $(Q - 1) \nmid 9$, which implies that $Q - 1$ does not divide $3(Q^2 + Q + 1)$. Since $|X| = Q - 1$, this shows that M cannot belong to class \mathcal{C}_3 in [5, Table 8.3].

Suppose M belongs to class \mathcal{C}_6 in [5, Table 8.3]. Then, $M \leq 3_+^{1+2} \rtimes 2A_4 < \text{SU}_4(2)$. In particular, X , as an element of M , must have order dividing 9 or 12, and hence, $Q \in \{2, 3, 4, 5, 7, 13\}$. Using (10), we are left with $Q = 13$, in which case X has order 12. However, any element of order 12 of $3_+^{1+2} \rtimes 2A_4$ does not possess an eigenvalue 1 in any 3-dimensional irreducible representation over \mathbb{C} or $\overline{\mathbb{F}}_{13}$.

(b4) Suppose M is a member of type SL_3 in class \mathcal{C}_5 in [5, Table 8.3]. Then, $p^d = Q = Q_0^m$ for some prime m ; in particular, $d \geq 2$. Also, set

$$e := \gcd(3, Q - 1).$$

Then, the element X^e belongs to $\text{SL}_3(Q_0)$, and admits eigenvalues 1 , and $(-q)^e$, $(-q^{-1})^e$ of order $(Q - 1)/e$. Consider any maximal torus T of $\text{SL}_3(Q_0)$ that contains the semisimple element X^e . If $T \cong C_{Q_0-1}^2$, then $(Q - 1)/e$ divides

$$Q_0 - 1 = p^{d/m} - 1 \leq p^{d/2} - 1 < \frac{p^d - 1}{3} \leq \frac{Q - 1}{e},$$

a contradiction. Suppose $T \cong C_{Q_0^2+Q_0+1}$. Then, any element in T has spectrum of the form $\{\lambda, \lambda^{Q_0}, \lambda^{Q_0^2}\}$ for some $1 \neq \lambda \in \overline{\mathbb{F}}_p^\times$. As 1 is an eigenvalue of $X^e \in T$, we see that the corresponding λ for X^e must be 1, and so, $X^e = \text{Id}$, again a contradiction. In the remaining cases, $T \cong C_{Q_0-1}$, which consists of elements with spectrum of the

form $\{\mu, \mu^{Q_0}, \mu^{-Q_0-1}\}$ for some $\mu \in \overline{\mathbb{F}_p}^\times$. As 1 is an eigenvalue of multiplicity one for X^e , we get $\mu^{Q_0+1} = 1$ for the corresponding μ for X^e . If $Q \neq 16$, then all eigenvalues of X must be of order dividing

$$e(Q_0 + 1) \leq e(p^{d/2} + 1) < p^d - 1,$$

which is a contradiction since the eigenvalue $-q$ of X has order $Q - 1$. If $Q = 16$, then $M = \mathrm{SL}_3(4)$, and one can check that no element of order $Q - 1 = 15$ can have the same spectrum as that of X , again a contradiction.

(b5) Suppose M is a member of type SU_3 in class \mathcal{C}_8 in [5, Table 8.3]. Then, $p^d = Q = Q_0^2$ and $M \cong \mathrm{SU}_3(Q_0) \times C_f$, where

$$f := \gcd(3, Q_0 - 1).$$

By (10), we have $Q_0 \neq 2$. Assume in addition that $Q_0 \neq 4$. Now, the element X^f belongs to $\mathrm{SU}_3(Q_0)$, and admits eigenvalues 1, and $(-q)^f, (-q^{-1})^f$ of order $(Q - 1)/f$. Consider any maximal torus T of $\mathrm{SU}_3(Q_0)$ that contains the semisimple element X^f . If $T \cong C_{Q_0+1}^2$, then $(Q - 1)/f = (Q_0^2 - 1)/f$ divides $Q_0 + 1$, and so, $Q_0 = 2$ or 4 , a contradiction. Suppose $T \cong C_{Q_0^2-Q_0+1}$. Then, any element in T has spectrum of the form $\{\lambda, \lambda^{-Q_0}, \lambda^{Q_0^2}\}$ for some $1 \neq \lambda \in \overline{\mathbb{F}_p}^\times$. As 1 is an eigenvalue of $X^f \in T$, we see that the corresponding λ for X^f must be 1, and so, $X^f = \mathrm{Id}$, again a contradiction. In the remaining cases, $T \cong C_{Q_0^2-1}$, which consists of elements with spectrum of the form $\{\mu, \mu^{-Q_0}, \mu^{Q_0-1}\}$ for some $\mu \in \overline{\mathbb{F}_p}^\times$. As 1 is an eigenvalue of multiplicity one for X^f , we get $\mu^{Q_0-1} = 1$ for the corresponding μ for X^f . But then all eigenvalues of X must be of order dividing

$$f(Q_0 - 1) \leq 3(Q_0 - 1) < Q_0^2 - 1,$$

a contradiction since the eigenvalue $-q$ of X has order $Q - 1$.

Now, we consider the case $Q_0 = 4$, so that $Q = 16$ and

$$M = \langle t \rangle \times \mathrm{SU}_3(4),$$

where t acts as a scalar β of order 3 on k^3 . Then, we can write $X = t^i X_1$, where X_1 is a semisimple element in $\mathrm{SU}_3(4)$ and $i \in \mathbb{Z}$. Consider any maximal torus T of $\mathrm{SU}_3(4)$ that contains X_1 . If $T \cong C_5^2$, then X_1 is conjugate to $\mathrm{diag}(\gamma_1, \gamma_2, \gamma_3)$, where $\gamma_j^5 = 1$. Since 1 is an eigenvalue of X , we may assume that $\beta^i \gamma_1 = 1$, which implies that $\beta^i = \gamma_1 = 1$ and $X = X_1$. But in this case, no eigenvalue of X can have order 15, a contradiction. Suppose $T \cong C_{13}$. Then, any element in T has spectrum of the form $\{\lambda, \lambda^{-Q_0}, \lambda^{Q_0^2}\}$ for some $1 \neq \lambda \in \overline{\mathbb{F}_2}^\times$ and $\lambda^{13} = 1$. As 1 is an eigenvalue of $X = t^i X_1$, we see that the corresponding λ for X_1 must be 1 and $\beta^i = 1$, and so, $X = \mathrm{Id}$, again a contradiction. In the remaining cases, $T \cong C_{15}$, which consists of

elements with spectrum of the form $\{\mu, \mu^{-4}, \mu^3\}$ for some $\mu \in \overline{\mathbb{F}_2}^\times$ with $\mu^{15} = 1$. As 1 is an eigenvalue of multiplicity one for $X = t^i X_1$, for the corresponding μ for X_1 , we have

$$\beta^i \mu = 1 \quad \text{or} \quad \beta^i \mu^4 = 1 \quad \text{or} \quad \beta^i \mu^3 = 1.$$

Since $|\beta| = 3$, we get $\mu^9 = 1$, and hence, $\mu^3 = 1$. But then all eigenvalues of X must be of order dividing 3, a contradiction since the eigenvalue $-q$ of X has order $Q - 1 = 15$.

(b6) The only remaining possibility is that $p > 2$ and M is a member of type SO_3 in class \mathcal{C}_8 in [5, Table 8.3]:

$$M \cong \text{SO}_3(Q) \times C_{\gcd(3, Q-1)}.$$

This is, however, ruled out by the same arguments as in part (a6). Hence, we have shown that $\langle X, Y \rangle = \text{SL}_3(k_0)$. ■

5. $\text{PSL}_2(\mathbb{F}_q)$ is not a characteristic quotient of F_2

In this section, we show that for any prime power q , $\text{PSL}_2(\mathbb{F}_q)$ is not a characteristic quotient of F_2 . In particular, this shows that the groups $\text{PSL}_3(\mathbb{F}_q)$, $\text{PSU}_3(\mathbb{F}_q)$ that we find are the lowest (untwisted Lie) rank finite simple groups of Lie type that are characteristic quotients of F_2 . The meat of the proof is a short explicit calculation, though first we will recall the relevant formalism. For more details, see [9, Section 5.2].

Let SL_2 denote the affine group scheme over \mathbb{Z} with ring $A = \mathbb{Z}[a, b, c, d]/(ad - bc - 1)$. Representations of F_2 in SL_2 form a scheme $\text{Hom}(F_2, \text{SL}_2) \cong \text{SL}_2 \times \text{SL}_2 \cong \text{Spec } A \otimes_{\mathbb{Z}} A$ whose k -valued points, for any ring k , are exactly the representations of F_2 in $\text{SL}_2(k)$. Letting superscripts denote fixed points, the quotient

$$X_{\text{SL}_2} := \text{Hom}(F_2, \text{SL}_2) // \text{GL}_2 = \text{Spec}(A \otimes_{\mathbb{Z}} A)^{\text{GL}_2}$$

by the conjugation action of GL_2 (equivalently, the diagonal action by conjugation on $\text{SL}_2 \times \text{SL}_2$) is the (integral) *character variety* for SL_2 -representations of F_2 . The natural action of $\text{Aut}(F_2)$ on $\text{Hom}(F_2, \text{SL}_2)$ induces an action of $\text{Out}(F_2)$ on the character variety X_{SL_2} . Much of the theory of SL_2 -representations of F_2 follows from the following fundamental result.

Theorem 5.1 (Brumfiel–Hilden). *The map $\text{Tr} : X_{\text{SL}_2} \rightarrow \mathbb{A}_{\mathbb{Z}}^3$ sending the class of a representation $\varphi : F_2 \rightarrow \text{SL}_2(k)$ (for any ring k) to the point $(\text{tr } \varphi(a), \text{tr } \varphi(b), \text{tr } \varphi(ab)) \in k^3$ is an isomorphism.*

Proof. The statement over \mathbb{C} was known to Fricke and Vogt, see [21, 22]. The statement over \mathbb{Z} is due to Brumfiel–Hilden [6, Propositions 3.5 and 9.1 (ii)], also see [9, Theorem 5.2.1]. ■

Beware that while GL_2 -equivalent representations give rise to the same point in the character variety, in general, the preimage of a point in X_{SL_2} is the union of multiple GL_2 -orbits in $\mathrm{Hom}(F_2, \mathrm{SL}_2)$. These difficulties disappear if we consider points valued over finite fields corresponding to absolutely irreducible representations.

Theorem 5.2. *Let $Z \subset \mathbb{A}_{\mathbb{Z}}^3$ denote the hypersurface given by $x^2 + y^2 + z^2 - xyz = 4$. For any finite field \mathbb{F}_q , the map $\mathrm{Tr} : X_{\mathrm{SL}_2} \xrightarrow{\sim} \mathbb{A}_{\mathbb{Z}}^3$ induces a bijection*

$$\{\text{Absolutely irreducible representations } \varphi : F_2 \rightarrow \mathrm{SL}_2(\mathbb{F}_q)\} / \mathrm{GL}_2(\mathbb{F}_q) \xrightarrow{\sim} \mathbb{F}_q^3 \setminus Z(\mathbb{F}_q).$$

Proof. See [9, Theorem 5.2.10]. ■

Since the conjugation action of $\mathrm{GL}_2(\mathbb{F}_q)$ on $\mathrm{SL}_2(\mathbb{F}_q)$ induces the full action of $\mathrm{Aut}(\mathrm{SL}_2(\mathbb{F}_q))$ and the action of $\mathrm{Out}(F_2)$ preserves (absolute) irreducibility, we find that characteristic quotients $F_2 \twoheadrightarrow \mathrm{SL}_2(\mathbb{F}_q)$ correspond to $\mathrm{Out}(F_2)$ -fixed points on $\mathbb{F}_q^3 \setminus Z(\mathbb{F}_q)$.

Since two maps to $\mathrm{SL}_2(\mathbb{F}_q)$ give rise to the same map to $\mathrm{PSL}_2(\mathbb{F}_q)$ if and only if the images of a, b under the two maps differ by $\pm I$, it follows that $\mathrm{Epi}(F_2, \mathrm{PSL}_2(\mathbb{F}_q)) / \mathrm{Aut}(\mathrm{PSL}_2(\mathbb{F}_q))$ can be identified with a subset of the orbit space $(\mathbb{F}_q^3 \setminus Z(\mathbb{F}_q)) / V$, where $V \cong \mathbb{Z}/2 \times \mathbb{Z}/2$ acts by negating two of the coordinates (note that V preserves Z). Accordingly, characteristic quotients $F_2 \twoheadrightarrow \mathrm{PSL}_2(\mathbb{F}_q)$ correspond to $\mathrm{Out}(F_2)$ -fixed points on $(\mathbb{F}_q^3 \setminus Z(\mathbb{F}_q)) / V$.

Theorem 5.3. *Let k be a ring. The action of $\mathrm{Out}(F_2)$ on $k^3 \cong X_{\mathrm{SL}_2}(k)$ descends to an action on the orbit space k^3 / V , where $V \cong \mathbb{Z}/2 \times \mathbb{Z}/2$ acts by negating two coordinates. The only points in k^3 / V fixed by $\mathrm{Out}(F_2)$ are the V -orbits of $(0, 0, 0)$ and $(2, 2, 2)$.*

Proof. The group $\mathrm{Aut}(F_2)$ is generated by three elements r, s, t . The definitions of r, s, t and the corresponding automorphisms of $\mathbb{A}_{\mathbb{Z}}^3 \cong X_{\mathrm{SL}_2}$ are given as follows (see [9, Lemma 5.2.4]):

$$\begin{array}{lll} r : (a, b) \mapsto (a^{-1}, b) & & \mathrm{Tr}_*(r) : (x, y, z) \mapsto (x, y, xy - z) \\ s : (a, b) \mapsto (b, a) & \xrightarrow{\mathrm{Tr}_*} & \mathrm{Tr}_*(s) : (x, y, z) \mapsto (y, x, z) \\ t : (a, b) \mapsto (a^{-1}, ab) & & \mathrm{Tr}_*(t) : (x, y, z) \mapsto (x, z, y). \end{array}$$

From this, we find that s, t -invariance implies that $\pm X = \pm Y = \pm Z$. Assuming this, r -invariance additionally implies that $X \in \{0, -2, 2\}$. One checks that the V -orbit of $(-2, 2, 2)$ is not r -invariant, and hence, the only fixed points are the orbits $V \cdot (0, 0, 0)$ and $V \cdot (2, 2, 2)$. ■

In the proof above, beware that while $\mathrm{Aut}(F_2)$ acts on the *right* on X_{SL_2} , the equations for $\mathrm{Tr}_*(r), \mathrm{Tr}_*(s), \mathrm{Tr}_*(t)$ describe a *left* action on \mathbb{A}^3 . Thus, Tr_* is an *anti*-homomorphism $\mathrm{Aut}(F_2) \rightarrow \mathrm{Aut}(\mathbb{A}^3)$.

Corollary 5.4. *For any prime power q , neither $\mathrm{SL}_2(\mathbb{F}_q)$ nor $\mathrm{PSL}_2(\mathbb{F}_q)$ is a characteristic quotient of F_2 .*

Proof. It suffices to show that $\mathrm{PSL}_2(\mathbb{F}_q)$ is not a characteristic quotient. For this, by Theorem 5.3, it suffices to note that the V -orbits of $(0, 0, 0)$ and $(2, 2, 2)$ cannot correspond to *surjective* representations $\varphi : F_2 \rightarrow \mathrm{PSL}_2(\mathbb{F}_q)$. Since $(2, 2, 2)$ satisfies $x^2 + y^2 + z^2 - xyz = 4$, the corresponding SL_2 -representation is not absolutely irreducible, and hence, it cannot be surjective. This also handles $(0, 0, 0)$ for even q , so it remains to consider $(0, 0, 0)$ for q odd.

Note that a trace 0 matrix $A \in \mathrm{SL}_2(\mathbb{F}_q)$ has characteristic polynomial $T^2 + 1$. Thus, A has order 4 and $A^2 = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$, so that A has order 2 in $\mathrm{PSL}_2(\mathbb{F}_q)$. It follows that for q odd, the point $(0, 0, 0)$ corresponds to a representation whose image in $\mathrm{PSL}_2(\mathbb{F}_q)$ is isomorphic to $\mathbb{Z}/2 \times \mathbb{Z}/2$, and hence, it is not surjective. ■

6. Relation with noncongruence modular curves

In this section, we explain how our main theorem disproves two conjectures in the theory of noncongruence subgroups of $\mathrm{SL}_2(\mathbb{Z})$. Recall that a subgroup $\Gamma \leq \mathrm{SL}_2(\mathbb{Z})$ is *congruence* if it contains $\Gamma(n) := \ker(\mathrm{SL}_2(\mathbb{Z}) \rightarrow \mathrm{SL}_2(\mathbb{Z}/n))$ for some integer $n \geq 1$. Otherwise, it is *noncongruence*. Let us define the *congruence degree* of a subgroup $\Gamma \leq \mathrm{SL}_2(\mathbb{Z})$ to be the index inside $\mathrm{SL}_2(\mathbb{Z})$ of the *congruence closure* Γ^c of Γ ; this Γ^c is the intersection of all congruence subgroups containing Γ . Noncongruence subgroups of congruence degree 1 are *totally noncongruent* [8, Section 4.4]. Since $\mathrm{SL}_2(\mathbb{Z})$ admits the free product $\mathrm{PSL}_2(\mathbb{Z}) \cong \mathbb{Z}/2 * \mathbb{Z}/3$ as a quotient, it follows that $\mathrm{SL}_2(\mathbb{Z})$ contains many finite-index noncongruence subgroups.¹³

In the theory of modular curves, many congruence subgroups can be understood as the $\mathrm{SL}_2(\mathbb{Z})$ -stabilizer of elements of $\mathrm{Epi}(\mathbb{Z}^2, G)$, where G is a finite quotient of \mathbb{Z}^2 . Identifying \mathbb{Z}^2 as the fundamental group of an elliptic curve, this leads to a moduli interpretation of the modular curve \mathcal{H}/Γ , where \mathcal{H} denotes the complex upper half-plane, and $\Gamma \leq \mathrm{SL}_2(\mathbb{Z})$ is a congruence subgroup acting on \mathcal{H} by mobius transformations. For example, if $\varphi : \mathbb{Z}^2 \rightarrow \mathbb{Z}/n$ is a surjection with stabilizer $\Gamma_\varphi := \mathrm{Stab}_{\mathrm{SL}_2(\mathbb{Z})}(\varphi)$, then \mathcal{H}/Γ is the moduli space of elliptic curves E equipped with a point of order n (equivalently, a \mathbb{Z}/n -cover $E' \rightarrow E$), see [8, Proposition 2.2.12] or [13, Section 1.5]. More generally, congruence modular curves are moduli spaces of elliptic curves with *abelian covers*. Such abelian covers can be equivalently described by torsion data.

¹³If G is a finite group which has a nonabelian simple composition factor not of type $\mathrm{PSL}_2(\mathbb{F}_p)$, then the kernel of any surjection $\mathrm{SL}_2(\mathbb{Z}) \rightarrow G$ is noncongruence.

By [32, Corollary N4], the abelianization map $F_2 \rightarrow \mathbb{Z}^2$ induces a rather exceptional isomorphism $\text{Out}(F_2) \cong \text{GL}_2(\mathbb{Z})$, restricting to an isomorphism $\text{Out}^+(F_2) \cong \text{SL}_2(\mathbb{Z})$. It follows that the action of $\text{SL}_2(\mathbb{Z})$ on $\text{Epi}(\mathbb{Z}^2, G)$ is a special case of the more general action

$$\text{SL}_2(\mathbb{Z}) \cong \text{Out}^+(F_2) \curvearrowright \text{Epi}(F_2, G) / \text{Inn}(G).$$

If $\varphi : F_2 \rightarrow G$ is a surjection, let Γ_φ denote the $\text{SL}_2(\mathbb{Z}) \cong \text{Out}^+(F_2)$ -stabilizer of its conjugacy class $\varphi \pmod{\text{Inn}(G)}$. Viewing F_2 as the fundamental group of an elliptic curve with the origin removed, it was shown in [8] that $\mathcal{H} / \Gamma_\varphi$ is a moduli space of elliptic curves with branched G -Galois covers, only ramified above the origin. By a result of Asada, every finite index subgroup of $\text{SL}_2(\mathbb{Z})$ contains Γ_φ for some appropriate finite group G and surjection $\varphi : F_2 \rightarrow G$. This endows every noncongruence modular curve with a moduli interpretation. In some sense, this puts noncongruence modular curves on a similar footing to congruence modular curves, and opens the door to a deeper investigation of the noncongruence side, guided by the wildly successful congruence theory.

A fundamental question that arises in this theory is: when does a surjection $\varphi : F_2 \rightarrow G$ have a congruence or noncongruence stabilizer? In the former case, we say that φ is congruence, and in the latter that φ is noncongruence. It is easy to see that if G is abelian, then every surjection is congruence. In [10], it is also shown that the same is true if G is metabelian. In [8, Theorem 4.4.10], it was shown that if x, y are generators of F_2 , then if G is not the trivial group and the orders of $\varphi(x), \varphi(y), \varphi(xy)$ are pairwise coprime, then Γ_φ is noncongruence, and its congruence closure¹⁴ is $\text{SL}_2(\mathbb{Z})$. These results support the following philosophy.

Philosophy 1. *Surjections onto highly nonabelian groups G should tend to have highly noncongruence stabilizers.*

In [8], the author attempted to make this philosophy more precise via a pair of conjectures. First, [8, Conjecture 4.4.10] asserts that for a fixed finite group G , either every surjection onto G is congruence, or they are all noncongruence. Second, [8, Conjecture 4.4.1] asserts that for nonsolvable G , all surjections onto G should be noncongruence. While supported by computational data at the time,¹⁵ both conjectures are false.

At the time of writing of this paper, all stabilizers of surjections onto nonabelian finite simple groups of order $\leq |J_1| = 175560$ (J_1 is the Janko group) have been

¹⁴The congruence closure of finite index a subgroup Γ of $\text{SL}_2(\mathbb{Z})$ is the intersection of all congruence subgroups containing Γ .

¹⁵At the time, all stabilizers of surjections onto groups of order ≤ 255 and simple groups of order $\leq |\text{Sz}(8)| = 29120$ had been computed. See [8, Appendix B].

computed. Amongst these, there are only two $\text{Aut}^+(F_2) \times \text{Aut}(G)$ -orbits of congruence surjections, both coming from characteristic quotients obtained via the Burau representation. These give rise to four $\text{Aut}^+(F_2)$ -orbits of congruence surjections onto $\text{PSU}_3(\mathbb{F}_4)$ (the remaining 29 being noncongruence), and two $\text{Aut}^+(F_2)$ -orbits of congruence surjections onto $\text{PSU}_3(\mathbb{F}_5)$ (the remaining 67 being noncongruence). Amongst simple groups, these are smallest counterexamples to [8, Conjectures 4.4.1, 4.4.10]. Theorem 1.6 in fact produces infinitely many counterexamples to [8, Conjecture 4.4.1]: for primes p , groups G of type $\text{PSL}_3(\mathbb{F}_p)$ or $\text{PSU}_3(\mathbb{F}_p)$ have $\text{Out}(G) \leq 6$, and hence, characteristic surjections onto such groups have $\text{SL}_2(\mathbb{Z})$ -stabilizers of index ≤ 6 . Since all subgroups of index ≤ 6 are congruence, the characteristic quotients onto such groups produced by Theorem 1.6 are all congruence.

Other than the 6 conjugacy classes of characteristic quotients onto $\text{PSU}_3(\mathbb{F}_4)$, $\text{PSU}_3(\mathbb{F}_5)$, the remaining 1328 $\text{Aut}^+(F_2)$ -orbits of surjections onto simple groups of order $\leq |J_1| = 175560$ are all noncongruence with congruence degree ≤ 3 . This leads to the natural question.

Question 6.1. Is there a *global* upper bound for the congruence degrees of stabilizers of surjections onto nonabelian finite simple groups?

We have also checked that amongst the 60 groups G of order ≤ 255 with solvable length ≥ 4 , all surjections onto G are noncongruence [8, Appendix B.1].

Question 6.2. Is there a lower bound on solvable length that can guarantee either the existence of noncongruence surjections or the nonexistence of congruence surjections?

In light of our results, we propose the weakening of [8, Conjecture 4.4.1].

Conjecture 6.3. If G is a nonabelian finite simple group, then there exists a surjection $\varphi : F_2 \rightarrow G$ such that Γ_φ is noncongruence. In other words, the kernel of the $\text{SL}_2(\mathbb{Z})$ action on $\text{Epi}(F_2, G)/\text{Inn}(G)$ is noncongruence.

This conjecture, which we will address in a sequel paper, still does not capture the empirical rareness of congruence surjections onto simple groups G . Indeed, it is unclear how to formulate a statement to this end that might be true. The fact that the line between congruence and noncongruence is so difficult to pin down in this setting contrasts with the clean characterizations of noncongruence subgroups given by the Hecke theory [2] and the unbounded denominators property for noncongruence modular forms [7]. It would be very interesting to see if a moduli interpretation for Hecke operators on noncongruence modular curves could shed light on this problem.

Geometrically, our main Theorem 1.6 implies a strange correspondence between abelian covers (or torsion points) on an elliptic curve (E, O) and $\text{PSL}_3(\mathbb{F}_p)$ -covers (or $\text{PSU}_3(\mathbb{F}_p)$ -covers of E ramified above the origin O . Our methods imply a relation

between such examples and Zariski dense representations of B_4 whose restrictions to F have large image. We end by noting that an elliptic curve (E, O) simultaneously determines an abelian variety, as well as a hyperbolic curve $E^\circ := E \setminus O$. Hyperbolic curves are also examples of *anabelian varieties*, in the sense of Grothendieck's anabelian geometry [23, 38], which are distinguished by the property that their structure as curves over a number field K is completely determined by the action of $\text{Gal}(\bar{\mathbb{Q}}/K)$ on their profinite fundamental group. The results of [1, 8] suggest that while congruence modular curves capture the essence of elliptic curves as abelian varieties, noncongruence modular curves capture their essence as anabelian curves. The problem of making precise the Philosophy 1 amounts to mapping out the boundary where the congruence theory stops and the noncongruence theory begins.

A. Appendix

A.1. Algebraic groups embedded in GL_n

Let R be a Dedekind domain with fraction field K . If G is an algebraic group over K equipped with a (homomorphic) embedding $G \hookrightarrow \text{GL}_{n,K}$ then we may define $G(R) := G(K) \cap \text{GL}_n(R)$ as the set of R -points of $\text{GL}_{n,R}$ whose intersection (as a closed subscheme) with the generic fiber $\text{GL}_{n,K}$ lies in G . Sometimes, it is nice to be able to consider $G(R)$ as the actual R -points of an algebraic group scheme over R . Indeed, this can be done by taking schematic closures.

Proposition A.1. *Let R be a Dedekind domain with fraction field K . Let H be a reduced flat affine group scheme over R , and let G be a reduced affine K -group scheme equipped with a homomorphic closed immersion $G \hookrightarrow H_K$ (for example, one might want to take $H = \text{GL}_{n,R}$). Let \bar{G} denote the schematic closure of G inside H , defined as the scheme theoretic image [46, 01R5] of the composition $G \hookrightarrow H_K \hookrightarrow H$. Then, \bar{G} is an R -flat closed subscheme of H with a unique structure of an R -group scheme making the morphisms $G \rightarrow \bar{G} \rightarrow H$ homomorphisms of R -group schemes. Moreover,*

$$\bar{G}_K := \bar{G} \times_{\text{Spec } R} \text{Spec } K$$

is canonically isomorphic to G , and we have

$$G(K) \cap H(R) = \bar{G}(R).$$

Note that since G, H are affine, the map $G \hookrightarrow H$ is affine, and hence, quasi-compact [46, 01S8]. Since G is reduced, the scheme theoretic image is the closure

of the image with the reduced induced structure [46, 056B].¹⁶ Thus, the schematic closure is just the topological closure with the reduced induced structure.

Proof. The statement (in a somewhat weaker form) is [11, Section 2], and the proof is due to Damian Rössler. Let $H = \text{Spec } A$, and $G = \text{Spec } B$, so we have by assumption a surjection $f : A_K := A \otimes_R K \rightarrow B$ coming from $G \hookrightarrow H_K$ and an injection $g : A \hookrightarrow A_K$ coming from $H_K \hookrightarrow H$. Let $\mu := f \circ g : A \rightarrow B$, then by definition we have $\bar{G} = \text{Spec } \mu(A)$. Since $\mu(A)$ is a subring of the R -flat module B , it is torsion-free, hence flat since R is Dedekind. It remains to show that $\mu(A)$ admits the structure of a Hopf algebra compatible with those of A, B . Let m_A, m_B be the comultiplications of A, B . Then, viewing B as a group scheme over R , we are looking to fill in the dotted arrow in the diagram

$$\begin{array}{ccccc}
 A & \xrightarrow{\mu} & \mu(A) & \hookrightarrow & B \\
 \downarrow m_A & & \downarrow & & \downarrow m_B \\
 A \otimes_R A & \xrightarrow{\mu \otimes \mu} & \mu(A) \otimes_R \mu(A) & \xrightarrow{h} & B \otimes_R B.
 \end{array}$$

Indeed, any such arrow would result in a Hopf algebra structure making the maps $G = \text{Spec } B \rightarrow \text{Spec } \mu(A) \rightarrow \text{Spec } A = H$ homomorphisms of R -group schemes (use the left half of the diagram, noting that $A \rightarrow \mu(A)$ is surjective). For such a dashed arrow to exist, it clearly suffices to check that the map h is injective. Set $M := \mu(A) \otimes_R \mu(A)$. Note that $\mu(A) \otimes_R K = B$, and since $K \otimes_R K = K$, $M \otimes_R K = B \otimes_R B$, so the map h can be identified with the canonical map $M \rightarrow M \otimes_R K$. Identifying $M \otimes_R K$ with the localization $(R - \{0\})^{-1}M$, we find that $\ker(M \rightarrow M \otimes_R K)$ is the R -torsion submodule of M , and so, the map h is injective if and only if $\mu(A) \otimes_R \mu(A)$ is torsion-free, which holds for us since $\mu(A)$ is R -flat. In this case, it is clear that the dashed arrow is even unique, so the group structure on \bar{G} is uniquely determined by $G \hookrightarrow H$.

Finally, let $\eta : \text{Spec } K \rightarrow \text{Spec } R$ be induced by the inclusion $R \subset K$. Given $g \in \bar{G}(R)$ with $g : \text{Spec } R \rightarrow \bar{G}$, the composition $\text{Spec } R \xrightarrow{g} \bar{G} \rightarrow H$ gives an R -point of H , and the composition $\text{Spec } K \xrightarrow{\eta} \text{Spec } R \xrightarrow{g} \bar{G}$ gives a K -point of \bar{G} , and hence, a K -point of G . The latter construction defines a map $\bar{G}(R) \rightarrow G(K)$ which is injective because \bar{G} is separated (since it is affine). Thus, we may view $\bar{G}(R)$ and $G(K) \cap H(R)$ as subgroups of $G(K)$, with $\bar{G}(R) \subset G(K) \cap H(R)$. To show they are equal, suppose we have an R -point $x : \text{Spec } R \rightarrow H$ such that $x \circ \eta \in G(K)$. Then, x defines an irreducible closed subscheme of H inside which the image of $x \circ \eta$ (a point of G)

¹⁶We think the referenced lemma also requires the quasi-compactness of “ f ”, even if it is not stated.

is dense. Since \bar{G} is the closure of G , \bar{G} contains the image of x . Since everything in sight is reduced, we conclude that x factors through \bar{G} [46, 0356]; thus $\bar{G}(R) = G(K) \cap H(R)$ as desired. ■

Remark A.2. If $G \hookrightarrow H_K \hookrightarrow H$ is given on rings by $\mu : A \rightarrow A_K \rightarrow B$, then the proof of the proposition shows that the schematic closure $\bar{G} = \text{Spec } \mu(A)$. If there exists an ideal $I \subset A$ such that $A_K \rightarrow B$ is given by $A_K \rightarrow A_K/IA_K$, then we would have $\mu(A) = A/I$ (note that $IA_K = I \otimes_R K$, and use exactness of $\otimes_R K$). In practice, this means that if G is cut out inside H_K by equations “with coefficients in R ”, then \bar{G} is cut out inside H by the same equations. In particular, the scheme cut out by those equations is flat and finite type over R .

Acknowledgments. This paper originated from discussions held at the Institute for Advanced Study in Princeton during the Fall of 2021. We thank the IAS for its hospitality. We also thank Gabriel Navarro for some helpful discussions, and Danny Neftin for his contributions during the early stages of the paper. We would also like to thank the referee for valuable remarks, which in particular inspired the results in Section 5.

Funding. A.L. was supported by the European Research Council (ERC) under the European Union’s Horizon 2020 (Grant agreement no. 882751). P.H.T. gratefully acknowledges the support of the NSF (Grants nos. DMS-1840702, DMS-2200850), the Simons Foundation, and the Joshua Barlaz Chair in Mathematics.

References

- [1] M. Asada, [The faithfulness of the monodromy representations associated with certain families of algebraic curves](#). *J. Pure Appl. Algebra* **159** (2001), no. 2-3, 123–147
Zbl [1045.14013](#) MR [1828935](#)
- [2] G. Berger, Hecke operators on noncongruence subgroups. *C. R. Acad. Sci. Paris Sér. I Math.* **319** (1994), no. 9, 915–919 Zbl [0815.11024](#) MR [1302789](#)
- [3] S. Bigelow, [The Lawrence–Krammer representation](#). In *Topology and geometry of manifolds (Athens, GA, 2001)*, pp. 51–68, Proc. Sympos. Pure Math. 71, American Mathematical Society, Providence, RI, 2003 Zbl [1061.20034](#) MR [2024629](#)
- [4] S. J. Bigelow, [Braid groups are linear](#). *J. Amer. Math. Soc.* **14** (2001), no. 2, 471–486
Zbl [0988.20021](#) MR [1815219](#)
- [5] J. N. Bray, D. F. Holt, and C. M. Roney-Dougal, [The maximal subgroups of the low-dimensional finite classical groups](#). London Math. Soc. Lecture Note Ser. 407, Cambridge University Press, Cambridge, 2013 Zbl [1303.20053](#) MR [3098485](#)
- [6] G. W. Brumfiel and H. M. Hilden, [SL\(2\) representations of finitely presented groups](#). Contemp. Math. 187, American Mathematical Society, Providence, RI, 1995
Zbl [0838.20006](#) MR [1339764](#)

- [7] F. Calegari, V. Dimitrov, and Y. Tang, [The unbounded denominators conjecture](#). *J. Amer. Math. Soc.* **38** (2025), no. 3, 627–702 Zbl [08026280](#) MR [4890656](#)
- [8] W. Y. Chen, [Moduli interpretations for noncongruence modular curves](#). *Math. Ann.* **371** (2018), no. 1-2, 41–126 Zbl [1454.11110](#) MR [3788845](#)
- [9] W. Y. Chen, [Nonabelian level structures, Nielsen equivalence, and Markoff triples](#). *Ann. of Math. (2)* **199** (2024), no. 1, 301–443 Zbl [7782633](#) MR [4681147](#)
- [10] W. Y. Chen and P. Deligne, [Arithmetic monodromy actions on pro-metabelian fundamental groups of once-punctured elliptic curves](#). 2017, arXiv:[1710.05532v1](#)
- [11] B. Conrad, [Reductive group schemes](#). In *Autour des schémas en groupes. Vol. I*, pp. 93–444, Panor. Synthèses 42/43, Société mathématique de France, Paris, 2014 Zbl [1349.14151](#) MR [3362641](#)
- [12] P. Deligne and G. D. Mostow, [Monodromy of hypergeometric functions and nonlattice integral monodromy](#). *Inst. Hautes Études Sci. Publ. Math.* (1986), no. 63, 5–89 Zbl [0615.22008](#) MR [0849651](#)
- [13] F. Diamond and J. Shurman, [A first course in modular forms](#). Grad. Texts in Math. 228, Springer, New York, 2005 Zbl [1062.11022](#) MR [2112196](#)
- [14] J. L. Dyer, E. Formanek, and E. K. Grossman, [On the linearity of automorphism groups of free groups](#). *Arch. Math. (Basel)* **38** (1982), no. 5, 404–409 Zbl [0476.20025](#) MR [0666911](#)
- [15] M. J. Evans, [Problems concerning generating sets for groups](#). Ph.D. thesis, Cardiff university, 1985
- [16] B. Farb and D. Margalit, [A primer on mapping class groups](#). Princeton Math. Ser. 49, Princeton University Press, Princeton, NJ, 2012 Zbl [1245.57002](#) MR [2850125](#)
- [17] L. Funar and P. Lochak, [Profinite completions of Burnside-type quotients of surface groups](#). *Comm. Math. Phys.* **360** (2018), no. 3, 1061–1082 Zbl [1404.20021](#) MR [3803818](#)
- [18] The GAP Group, [GAP—groups, algorithms, and programming. Version 4.12.2](#). 2022
- [19] S. Garion and A. Shalev, [Commutator maps, measure preservation, and \$T\$ -systems](#). *Trans. Amer. Math. Soc.* **361** (2009), no. 9, 4631–4651 Zbl [1182.20015](#) MR [2506422](#)
- [20] T. Gelander, [Aut\(\$F_n\$ \) actions on representation spaces](#). *J. Algebra* **656** (2024), 206–225 Zbl [1552.22021](#) MR [4759502](#)
- [21] W. M. Goldman, [Trace coordinates on Fricke spaces of some simple hyperbolic surfaces](#). In *Handbook of Teichmüller theory. Vol. II*, pp. 611–684, IRMA Lect. Math. Theor. Phys. 13, European Mathematical Society, Zürich, 2009 Zbl [1175.30043](#) MR [2497777](#)
- [22] W. M. Goldman, [An exposition of results of Fricke](#). [v1] 2004, [v2] 2005, arXiv:[math/0402103v2](#)
- [23] A. Grothendieck, [Esquisse d’un programme](#). In *Geometric Galois actions, I*, pp. 5–48, London Math. Soc. Lecture Note Ser. 242, Cambridge University Press, Cambridge, 1997 Zbl [0901.14001](#) MR [1483107](#)
- [24] L. C. Grove, [Classical groups and geometric algebra](#). Grad. Stud. Math. 39, American Mathematical Society, Providence, RI, 2002 Zbl [0990.20001](#) MR [1859189](#)
- [25] V. F. R. Jones, [Hecke algebra representations of braid groups and link polynomials](#). *Ann. of Math. (2)* **126** (1987), no. 2, 335–388 Zbl [0631.57005](#) MR [0908150](#)

- [26] D. Krammer, [The braid group \$B_4\$ is linear](#). *Invent. Math.* **142** (2000), no. 3, 451–486
Zbl [0988.20023](#) MR [1804157](#)
- [27] D. Krammer, [Braid groups are linear](#). *Ann. of Math. (2)* **155** (2002), no. 1, 131–156
Zbl [1020.20025](#) MR [1888796](#)
- [28] A. Landesman and D. Litt, [Canonical representations of surface groups](#). *Ann. of Math. (2)* **199** (2024), no. 2, 823–897 Zbl [07836052](#) MR [4717077](#)
- [29] S. Lawton, [Generators, relations and symmetries in pairs of \$3 \times 3\$ unimodular matrices](#). *J. Algebra* **313** (2007), no. 2, 782–801 Zbl [1119.13004](#) MR [2329569](#)
- [30] A. Lubotzky, Dynamics of $\text{Aut}(F_N)$ actions on group presentations and representations. In *Geometry, rigidity, and group actions*, pp. 609–643, Chicago Lectures in Math., The University of Chicago Press, Chicago, IL, 2011 Zbl [1266.20045](#) MR [2807845](#)
- [31] A. Lubotzky and D. Segal, [Subgroup growth](#). Progr. Math. 212, Birkhäuser, Basel, 2003
Zbl [1071.20033](#) MR [1978431](#)
- [32] W. Magnus, A. Karrass, and D. Solitar, *Combinatorial group theory*. 2nd edn., Dover, Mineola, NY, 2004 Zbl [1130.20307](#) MR [2109550](#)
- [33] G. Malle and D. Testerman, [Linear algebraic groups and finite groups of Lie type](#). Cambridge Stud. Adv. Math. 133, Cambridge University Press, Cambridge, 2011
Zbl [1256.20045](#) MR [2850737](#)
- [34] C. R. Matthews, L. N. Vaserstein, and B. Weisfeiler, [Congruence properties of Zariski-dense subgroups. I](#). *Proc. London Math. Soc. (3)* **48** (1984), no. 3, 514–532
Zbl [0551.20029](#) MR [0735226](#)
- [35] C. T. McMullen, [Braid groups and Hodge theory](#). *Math. Ann.* **355** (2013), no. 3, 893–946
Zbl [1290.30050](#) MR [3020148](#)
- [36] D. W. Morris, *Introduction to arithmetic groups*. Deductive Press, 2015 Zbl [1319.22007](#)
MR [3307755](#)
- [37] K. Nakamoto, [The structure of the invariant ring of two matrices of degree 3](#). *J. Pure Appl. Algebra* **166** (2002), no. 1-2, 125–148 Zbl [1001.15022](#) MR [1868542](#)
- [38] H. Nakamura, A. Tamagawa, and S. Mochizuki, The Grothendieck conjecture on the fundamental groups of algebraic curves (in Japanese). *Sūgaku* **50** (1998), no. 2, 113–129. English translation. *Sugaku Expo.* **14** (2001), no. 1, 31–53 Zbl [0943.14014](#)
MR [1648427](#)
- [39] B. H. Neumann and H. Neumann, [Zwei Klassen charakteristischer Untergruppen und ihre Faktorgruppen](#). *Math. Nachr.* **4** (1951), 106–125 Zbl [0042.02102](#) MR [0040297](#)
- [40] I. Pak, [What do we know about the product replacement algorithm?](#) In *Groups and computation, III (Columbus, OH, 1999)*, pp. 301–347, Ohio State Univ. Math. Res. Inst. Publ. 8, de Gruyter, Berlin, 2001 Zbl [0986.68172](#) MR [1829489](#)
- [41] R. Pink, [Strong approximation for Zariski dense subgroups over arbitrary global fields](#). *Comment. Math. Helv.* **75** (2000), no. 4, 608–643 Zbl [0981.20036](#) MR [1789179](#)
- [42] V. Platonov and A. Rapinchuk, *Algebraic groups and number theory*. Pure Appl. Math. 139, Academic Press, Boston, MA, 1994 Zbl [0841.20046](#) MR [1278263](#)
- [43] B. Poonen, [Rational points on varieties](#). Grad. Stud. Math. 186, American Mathematical Society, Providence, RI, 2017 Zbl [1387.14004](#) MR [3729254](#)

- [44] C. Procesi, [The invariant theory of \$n \times n\$ matrices](#). *Advances in Math.* **19** (1976), no. 3, 306–381 Zbl [0331.15021](#) MR [0419491](#)
- [45] A. S. Sikora, [\$SL_n\$ -character varieties as spaces of graphs](#). *Trans. Amer. Math. Soc.* **353** (2001), no. 7, 2773–2804 Zbl [1046.57014](#) MR [1828473](#)
- [46] Stacks Project Authors, *The Stacks project*. 2022, available at stacks.math.columbia.edu, visited on 3 May 2025
- [47] A. Stoimenow, [Non-conjugate braids with the same closure link from density of representations](#). *J. Math. Pures Appl. (9)* **94** (2010), no. 5, 470–496 Zbl [1216.57002](#) MR [2732925](#)
- [48] T. Szamuely, *Galois groups and fundamental groups*. Cambridge Stud. Adv. Math. 117, Cambridge University Press, Cambridge, 2009 Zbl [1189.14002](#) MR [2548205](#)
- [49] T. N. Venkataramana, [Image of the Burau representation at \$d\$ -th roots of unity](#). *Ann. of Math. (2)* **179** (2014), no. 3, 1041–1083 Zbl [1345.20051](#) MR [3171758](#)
- [50] B. Weisfeiler, [Strong approximation for Zariski-dense subgroups of semisimple algebraic groups](#). *Ann. of Math. (2)* **120** (1984), no. 2, 271–315 Zbl [0568.14025](#) MR [0763908](#)
- [51] K. Zsigmondy, [Zur Theorie der Potenzreste](#). *Monatsh. Math. Phys.* **3** (1892), no. 1, 265–284 Zbl [24.0176.02](#) MR [1546236](#)

Received 28 August 2023.

William Y. Chen (corresponding author)

Department of Mathematics, University of Illinois Urbana-Champaign, 273 Altgeld Hall, 1409 W. Green Street, Urbana, IL 61801, USA; oxeimon@gmail.com

Alexander Lubotzky

Faculty of Mathematics and Computer Science, The Weizmann Institute of Science, 234 Herzl Street, 7610001 Rehovot, Israel; alex.lubotzky@mail.huji.ac.il

Pham Huu Tiep

Department of Mathematics, Hill Center for the Mathematical Sciences, Rutgers University, 110 Frelinghuysen Rd, Piscataway, NJ 08854-8019, USA; pht19@math.rutgers.edu