



Oscillatory integrals and weighted gradient flows

Michael Greenblatt

Abstract. We investigate estimating scalar oscillatory integrals by integrating by parts in directions based on $(x_1 \partial_{x_1} f(x), \dots, x_n \partial_{x_n} f(x))$, where $f(x)$ is the phase function. We prove a theorem which provides estimates that are uniform with respect to linear perturbations of the phase and investigate some consequences. When the phase function is quasi-homogeneous the theorem gives estimates for the associated surface measure Fourier transforms that are generally not too far off from being sharp. In addition, the theorem provides a new proof, up to endpoints, that the well-known oscillatory integral estimates of Varchenko when the Newton polyhedron of the phase function is nondegenerate extend to corresponding bounds for surface measure Fourier transforms when the index is less than $1/2$. A sharp version of this was originally proven by the author [J. Funct. Anal. 262 (2012), no. 5, 2314–2348].

1. Background and theorem statements

We consider oscillatory integrals of the form

$$(1.1) \quad I(\lambda) = \int e^{i\lambda f(x_1, \dots, x_n)} \phi(x_1, \dots, x_n) dx_1 \cdots dx_n.$$

Here $f(x)$ is a real analytic function defined on a bounded neighborhood U of the origin, $\phi(x)$ is a C^1 real-valued function supported in U , and λ is a real parameter. Often one seeks estimates of the form $|I(\lambda)| \leq h(|\lambda|)$, where h is an appropriately decreasing function, such as a function of the form $C(1 + |\lambda|)^{-s}$, for $C, s > 0$. We will always assume that $\nabla f(0) = 0$ to ensure we are in a nontrivial situation. By subtracting a constant from f , without loss of generality we may also assume that $f(0) = 0$.

A canonical example of where oscillatory integrals (1.1) show up is in the analysis of Fourier transforms of surface measures. If S is a surface in \mathbb{R}^{n+1} that is given by the graph of a real analytic $f(x_1, \dots, x_n)$ on a bounded neighborhood U of the origin and $\phi(x)$ is a real-valued C^1 function supported in U , then the Fourier transform of the Euclidean surface measure on S localized through $\phi(x)$, which we denote by μ , is given by

$$(1.2) \quad \hat{\mu}(\lambda_1, \dots, \lambda_{n+1}) = \int e^{-i\lambda_{n+1} f(x_1, \dots, x_n) - i\lambda_1 x_1 - \cdots - i\lambda_n x_n} \phi(x_1, \dots, x_n) dx_1 \cdots dx_n.$$

Mathematics Subject Classification 2020: 42A38 (primary); 41A60, 42B99 (secondary).

Keywords: oscillatory integral, weighted gradient flow.

We always assume the surface has been translated and rotated so that like before, $f(0) = 0$ and $\nabla f(0) = 0$. This time the goal is to find estimates of the form $|\hat{\mu}(\lambda)| \leq h(|\lambda|)$ for appropriate decreasing h . Since Fourier transforms of surface measures appear in a range of subjects including restriction problems, maximal averages, lattice point discrepancy, and more, estimates of this form can help lead to developments in those subjects. We refer to Chapter 8 of [20] for an introduction to the connections between surface measure Fourier transforms and restriction problems, and we refer to [11] for background material on the connections between maximal averages, lattice point discrepancy, and surface measure Fourier transforms. The paper [11] also describes various results on surface measure Fourier transforms.

Since the phase function in (1.2) is a linear perturbation of that of (1.1), one approach to proving such bounds on $|\hat{\mu}(\lambda)|$ is to provide bounds on $|I(\lambda)|$ that are uniform under linear perturbations of the phase. In this paper, we will describe one method of doing that.

To motivate what we will be doing, observe that one approach to the analysis of oscillatory integrals (1.1)–(1.2) is to appropriately divide the domain of integration into curves, do an appropriate integration by parts on each curve, and then integrate the result in the remaining $n - 1$ dimensions. The hope would be that if the curves are properly chosen then one could obtain desirable functions $h(|\lambda|)$ bounding the overall oscillatory integral. A clue on how to select these curves is given by how oscillatory integral decay is often connected to bounds on sublevel set measures. Namely, in many situations, the supremum of the ε for which (1.1) satisfies a bound $|I(\lambda)| \leq C(1 + |\lambda|)^{-\varepsilon}$ for some $C > 0$ is the same as the supremum of the ε for which there is a constant $C > 0$ such that the following holds for all $s > 0$:

$$(1.3) \quad m(\{x \in U : |f(x)| < s\}) \leq Cs^\varepsilon.$$

Here m is Lebesgue measure. This suggests that one might get good results if the curves on which one performs the integrations by parts are perpendicular to the boundaries of these sublevel sets. In other words, one might choose these curves to be tangent to ∇f . This idea is further bolstered by the fact that $f(x)$ increases or decreases fastest in the direction of the gradient, so that the phase oscillates fastest in the directions of such curves.

However, the above is not the whole story, since even if the phase is oscillating quickly in directions tangent to a curve, if the second derivative of the phase is also large, these quick oscillations might not have the desired effect; the curve might head rapidly into region where the phase has a stationary point. Thus to add some flexibility to our activities, we consider not just curves whose tangents are in the direction of ∇f , but also in directions $(a_1(x)\partial_{x_1}f(x), \dots, a_n(x)\partial_{x_n}f(x))$, where the weights $a_i(x)$ are real analytic or even quotients of real analytic functions.

Using directions of this form have an additional advantage. If one replaces $f(x)$ by $f(x) + b \cdot x$ for some $b \in \mathbb{R}^n$, then $a_i(x)\partial f/\partial x_i$ becomes $a_i(x)(\partial f/\partial x_i + b_i)$. Using resolution of singularities one can show that, generally speaking, for the type of $a_i(x)$ under discussion, if one has sublevel set measure estimates of the form

$$m\left(\left\{x \in U : \sum_{i=1}^n \left|a_i(x) \frac{\partial f}{\partial x_i}\right| < s\right\}\right) \leq Cs^\varepsilon$$

for some $C > 0$ and $0 < \varepsilon < 1$, then if U is sufficiently small, the same will hold when each $a_i(x)\partial f/\partial x_i$ is replaced by $a_i(x)(\partial f/\partial x_i + b_i)$. This will allow us to state our theorems in terms of the optimal ε for which sublevel set bounds of the form (1.3) hold for certain functions of the form $\sum_{i=1}^n |\tilde{a}_i(x)\partial f/\partial x_i|$ (where the $\tilde{a}_i(x)$ are slightly different from the $a_i(x)$.) Hence we will have a way of estimating $|I(\lambda)|$ that is uniform under linear perturbations, providing a way of bounding $|\hat{\mu}(\lambda)|$.

To motivate possible choices of the weight functions $a_i(x)$, we consider the case where $f(x)$ is a monomial $ax_1^{\alpha_1} \cdots x_n^{\alpha_n}$. When $\alpha_i > 0$, the effect of taking an x_i derivative on $f(x)$ is to multiply it by a constant times x_i^{-1} . This suggests that for “balance”, one might choose $a_i(x) = x_i$, so that the directions in which one integrates by parts are of the form $(x_1 \partial_{x_1} f(x), \dots, x_n \partial_{x_n} f(x))$. Our main theorem, Theorem 1.1, will be based on using such weights. We will then see in Section 2.1 that this theorem provides good bounds for $|\hat{\mu}(\lambda)|$ when $f(x)$ is a quasi-homogeneous polynomial. Then, in Section 2.2, we will that Theorem 1.1 can be used to show, up to endpoints, that when $f(x)$ has nondegenerate Newton polyhedron in the sense of [21], the optimal estimates for $|I(\lambda)|$ in [21] extend to analogous bounds for surface measure Fourier transforms. This was earlier shown (including endpoints) in [9].

Other choices of $a_i(x)$ also lead to results that might be of some interest, but for simplicity of exposition, we are only focusing on the weights $a_i(x) = x_i$ in this paper.

We now come to our theorem, whose proof is based on using the above weighted gradient flow. While it gives especially desirable results in the above situations, the theorem holds generally.

Theorem 1.1. *Suppose $\phi(x)$ is C^1 and $f(x)$ is real analytic with $f(0)=0$ and $\nabla f(0)=0$. Suppose W is a bounded neighborhood of the origin such that if $\varepsilon > 0$ is such that for some $C > 0$ and all $s > 0$ one has the sublevel set measure estimate*

$$(1.4) \quad m\left(\left\{x \in W : \frac{\sum_{i=1}^n |x_i \partial f/\partial x_i|}{\prod_{i=1}^n |x_i|} < s\right\}\right) \leq Cs^\varepsilon.$$

Then if $U \subset W$ is a sufficiently small ball centered at the origin, the following hold.

- (1) *For all $\delta < \varepsilon/(\varepsilon + 1)$, there is a constant A such that $|I(\lambda)| \leq A(1 + |\lambda|)^{-\delta}$ whenever ϕ is supported in U . Here A depends on f , ϕ , and δ .*
- (2) *For all $\delta < \min(\varepsilon/(\varepsilon + 1), 1/2)$, there is a constant B such that one has $|\hat{\mu}(\lambda)| \leq B(1 + |\lambda|)^{-\delta}$ whenever ϕ is supported in U . Here B depends on f , ϕ , and δ .*

An examination of the proof of Theorem 1.1 reveals that the condition that ϕ is C^1 can actually be weakened to requiring that there is a constant M for which $|\phi(x)| \leq M$ and $|\partial_{x_i} \phi(x)| \leq M/|x_i|$ for each i . On the other hand, if ϕ is smooth, it turns out that if U is a small enough neighborhood of the origin, if ε is such that the sublevel set bounds (1.3) hold for some C , then one automatically has $|I(\lambda)| \leq C'(1 + |\lambda|)^{-\varepsilon}$ for some C' depending on f , ϕ , and U . If in addition $\phi(x)$ is nonnegative with $\phi(0) > 0$, then the supremum of the ε for which such an estimate $|I(\lambda)| \leq C'(1 + |\lambda|)^{-\varepsilon}$ holds is actually equal to the supremum of the ε for which (1.3) holds, unless the former supremum is a negative integer. We refer to Chapter 6 of [1] for more information about these matters. This leads to the following corollary to Theorem 1.1, which may be of interest in its own right.

Corollary 1.2. *If W is a bounded neighborhood of the origin and ε_1 denotes the supremum of the ε for which (1.4) holds for some C , then if $U \subset W$ is a sufficiently small neighborhood of the origin and ε_2 denotes the supremum of the ε for which (1.3) holds for some C , then $\varepsilon_1/(\varepsilon_1 + 1) \leq \varepsilon_2$.*

Although we will not prove it here, using resolution of singularities one can show that ε_1 is independent of W and ε_2 is independent of U if W and U are sufficiently small neighborhoods of the origin, so that one can take $U = W$ in Corollary 1.2.

There has been quite a bit of work done on scalar oscillatory integrals of the form (1.1). In addition to [21], some notable examples include the papers [3, 6, 7, 13, 19]. For the surface measure Fourier transforms (1.2), there has also been a lot of work done, in part due to their connections with maximal averages, lattice point discrepancy, and other areas. Much of the effort in this area has focused on either convex surfaces or the two-dimensional case. We mention the references [2, 4, 5, 18] for the convex situation, and [16, 17] for the two-dimensional situation.

Gradient flows for scalar oscillatory integrals are often used in fields such as physics when putting the method of steepest descent into effect. In addition, gradient flows appear in various areas of mathematics, including partial differential equations, optimization, and more applied fields like computer vision and machine learning. The author does not know of any specific connection between this paper and the work in the above subjects, but it might be a direction worth exploring.

2. Consequences of Theorem 1.1

2.1. Quasi-homogeneous functions

A polynomial $p(x_1, \dots, x_n) = \sum_{\alpha} c_{\alpha} x^{\alpha}$ is said to be quasi-homogeneous if there are positive rational numbers k_1, \dots, k_n such for any $\alpha = (\alpha_1, \dots, \alpha_n)$ for which $c_{\alpha} \neq 0$, one has $\sum_{i=1}^n k_i \alpha_i = 1$. An equivalent statement is that $p(t^{k_1} x_1, \dots, t^{k_n} x_n) = t p(x_1, \dots, x_n)$ for all $(x_1, \dots, x_n) \in \mathbb{R}^n$ and all $t > 0$, and this definition extends the former definition to non-polynomials. A canonical example of a quasi-homogeneous polynomial is $x_1^{l_1} + \dots + x_n^{l_n}$, where one has $k_i = 1/l_i$ for each i .

Note that if $f(x)$ is a quasi-homogeneous polynomial, the function $\sum_{i=1}^n |x_i \partial f / \partial x_i|$ appearing in Theorem 1.1 is also quasi-homogeneous, with the same (k_1, \dots, k_n) . Given the nature of the statement of Theorem 1.1, it makes sense that we would want to understand the growth rate of the measure of the sublevel sets of quasi-homogeneous functions. Suppose $f(x)$ is a quasi-homogeneous function and k_1, \dots, k_n are as above. Let V be a bounded neighborhood of the origin, and let ε_0 be defined by the supremum of the ε such that there is a constant $C > 0$ such that

$$(2.1) \quad m(\{x \in V : |f(x)| < s\}) \leq C s^{\varepsilon}.$$

Here, as before, m denotes Lebesgue measure, and we will always work with functions for which $\varepsilon_0 > 0$. Note that an equivalent definition of ε_0 is the supremum of the ε such that

$$(2.2) \quad \int_V |f(x)|^{-\varepsilon} dx < \infty.$$

By the quasihomogeneity of f , the number ε_0 is independent of V . In particular, one may take V to be the box $\{x : -1 < x_i < 1 \text{ for all } i\}$, which we do henceforth. If we change variables in (2.2) from x to y , where $x_1 = y_1$ and $x_i = (\operatorname{sgn} x_i)|y_i|^{k_i/k_1}$ for $i > 1$, the integral in (2.2) becomes a constant times

$$(2.3) \quad \int_V |\tilde{f}(y)|^{-\varepsilon} \prod_{i=1}^n |y_i|^{k_i/k_1-1} dy.$$

Here $\tilde{f}(y)$ is the function $f(x)$ in the y coordinates, which has the key property that it is homogeneous of degree $1/k_1$. By conversion to polar coordinates, (2.3) is finite when two requirements are met. First, we need that $-\varepsilon/k_1 + \sum_{i=1}^n (k_i/k_1 - 1) > -n$. Secondly, we need that $|\tilde{f}(y)|^{-\varepsilon}$ integrates to a finite value over the boundary sides of V , which is equivalent to $|f(x)|^{-\varepsilon}$ integrating to a finite value over the boundary sides of V . The first condition translates into $\varepsilon < \sum_{i=1}^n k_i$. Thus ε_0 is the supremum of all ε such that $\varepsilon < \sum_{i=1}^n k_i$ and such that for each i and $a = 1, -1$, we have

$$(2.4) \quad \int_{\{x: x_i=a, -1 < x_j < 1 \text{ for all } j \neq i\}} |f(x)|^{-\varepsilon} dx_1 \cdots dx_{i-1} dx_{i+1} \cdots dx_n < \infty.$$

We now investigate what Theorem 1.1 says in the quasi-homogeneous case. Assume that $f(x)$ is a quasi-homogeneous polynomial. Notice that the function $\frac{\sum_{i=1}^n |x_i \partial f / \partial x_i|}{\prod_{i=1}^n |x_i|}$ appearing in Theorem 1.1 satisfies

$$(2.5) \quad \frac{\sum_{i=1}^n |x_i \partial f / \partial x_i|}{\prod_{i=1}^n |x_i|} \geq \sum_{i=1}^n \left| x_i \frac{\partial f}{\partial x_i} \right| \geq C \left| \sum_{i=1}^n k_i x_i \frac{\partial f}{\partial x_i} \right| = C |f(x)|.$$

Thus whenever we are in a situation where (2.1) holds on some bounded neighborhood V of the origin, the same will be true if we replace $|f(x)|$ by $\frac{\sum_{i=1}^n |x_i \partial f / \partial x_i|}{\prod_{i=1}^n |x_i|}$. In particular, this holds for every $\varepsilon < \varepsilon_0$. Thus part (2) of Theorem 1.1 says that in the quasi-homogeneous case, one has that $|\hat{\mu}(\lambda)| \leq B(1 + |\lambda|)^{-\delta}$ holds for all $\delta < \min(\varepsilon_0/(\varepsilon_0 + 1), 1/2)$. So if $\varepsilon_0 < 1$, whereas ε_0 gives the supremal exponent for the scalar oscillatory integral $I(\lambda)$ due to the connection between oscillatory integrals and sublevel set measure growth, the supremal exponent for the surface measure Fourier transform is at least $\varepsilon_0/(\varepsilon_0 + 1)$. Hence the true exponent lies somewhere in the interval $[\varepsilon_0/(\varepsilon_0 + 1), \varepsilon_0]$. This can be a substantial improvement over simply using stationary phase or the Van der Corput lemma along curves $(c_1 t^{k_1}, \dots, c_n t^{k_n})$ as in [10], where one can typically get an exponent no better than $1/(n + 1)$.

It might occur to one that since the estimates in (2.5) are not that refined, we might be able to get better results by using more careful estimates than those of (2.5). It turns out that this often is the case if the condition $\varepsilon < \sum_{i=1}^n k_i$ is more stringent than the ones in (2.4), so that $\varepsilon_0 = \sum_{i=1}^n k_i$. One can show that in many such situations, one will have that the exponent given by part (2) of Theorem 1.1 is $\min(\sum_{i=1}^n k_i, 1/2)$, so that there is no reduction in the exponent when it is less than $1/2$. On the other hand, if the conditions of (2.4) are more stringent than the condition $\varepsilon < \sum_{i=1}^n k_i$, then often one can show that Theorem 1.1 gives no better exponent than the $\min(\varepsilon_0/(\varepsilon_0 + 1), 1/2)$ given above, while the true exponent can be as large as ε_0 .

2.2. Estimates in terms of the Newton polyhedron

We first provide some relevant terminology.

Definition 2.1. Let $f(x)$ be a smooth function defined on a neighborhood of the origin in \mathbb{R}^n , and let $f(x) = \sum_{\alpha} f_{\alpha} x^{\alpha}$ denote the Taylor expansion of $f(x)$ at the origin. For any α for which $f_{\alpha} \neq 0$, let Q_{α} be the octant $\{x \in \mathbb{R}^n : x_i \geq \alpha_i \text{ for all } i\}$. Then the *Newton polyhedron* $N(f)$ of $f(x)$ is defined to be the convex hull of all Q_{α} .

A Newton polyhedron may contain faces of dimensions zero through $n - 1$ (a vertex is considered to be a compact face of dimension zero.) These faces can be either compact or unbounded. In this paper, as in earlier work like [8, 9, 21], an important role is played by the following functions, defined for compact faces of the Newton polyhedron.

Definition 2.2. Suppose F is a compact face of $N(f)$. Then if $f(x) = \sum_{\alpha} f_{\alpha} x^{\alpha}$ denotes the Taylor expansion of f like above, we define $f_F(x) = \sum_{\alpha \in F} f_{\alpha} x^{\alpha}$.

Definition 2.3. The Newton polyhedron of $f(x)$ is said to be nondegenerate if for each compact face F of $N(f)$, the function $\nabla f_F(x)$ is nonvanishing on $(\mathbb{R} - \{0\})^n$.

Definition 2.4. Assume $N(f)$ is nonempty. Then the *Newton distance* $d(f)$ of $f(x)$ is defined to be $\inf\{t : (t, t, \dots, t, t) \in N(f)\}$.

The following is a well-known theorem of Varchenko [21]. We include the condition $|\lambda| \geq 2$ in the statement since the result is immediate for $|\lambda| < 2$, and we want to avoid situations where $\ln |\lambda|$ is near zero.

Theorem 2.5 (Varchenko). *Suppose $f(x)$ is real analytic on a neighborhood of the origin with $f(0) = 0$ such that $N(f)$ is nondegenerate. Let k denote the dimension of the face of $N(f)$ intersecting the line $x_1 = \dots = x_n$ in its interior. There is a neighborhood U of the origin such that if ϕ is supported in U , then the following hold.*

- (1) *There is a constant $C > 0$, depending on f and ϕ , such that for all $|\lambda| \geq 2$, one has*

$$|I(\lambda)| \leq C |\lambda|^{-1/d(f)} (\ln |\lambda|)^{n-1-k}.$$

- (2) *If $\phi(x)$ is nonnegative, $\phi(0) > 0$, and $d(f) > 1$, then there is a $C' > 0$ depending on f and ϕ such that if $|\lambda|$ is sufficiently large, one has*

$$|I(\lambda)| \geq C' |\lambda|^{-1/d(f)} (\ln |\lambda|)^{n-1-k}$$

In [9], among other things, Theorem 2.5 was extended to surface measure Fourier transforms when $d(f) > 2$. This follows from parts (a) and (b) of the following consequence of Theorem 1.5 of [9].

Theorem 2.6 (Theorem 1.5 of [9]). *Suppose $f(x)$ is real analytic on a neighborhood of the origin with $f(0) = 0$ and $\nabla f(0) = 0$. Let k denote the dimension of the face of $N(f)$ intersecting the line $x_1 = \dots = x_n$ in its interior. There is a neighborhood U of the origin such that if ϕ is supported in U , the following hold for $|\lambda| \geq 2$, where C denotes a constant depending on f and ϕ .*

- (a) *If $d(f) < 2$, and each zero of each $f_F(x)$ on $(\mathbb{R} - \{0\})^n$ has order at most 2, then there is a constant C such that $|\hat{\mu}(\lambda)| \leq C |\lambda|^{-1/2}$.*

- (b) If $d(f) \geq 2$ and each zero of each $f_F(x)$ on $(\mathbb{R} - \{0\})^n$ has order at most $d(f)$, then there is a constant C such that $|\hat{\mu}(\lambda)| \leq C|\lambda|^{-1/d(f)} (\ln |\lambda|)^{n-k}$. If $d(f)$ is not an integer, the exponent $n - k$ can be improved to $n - 1 - k$.
- (c) If the maximum order m of any zero of any $f_F(x)$ on $(\mathbb{R} - \{0\})^n$ satisfies $m > \max(d(f), 2)$, then there is a constant C such that $|\hat{\mu}(\lambda)| \leq C|\lambda|^{-1/m}$.

To be clear, the order of a zero of $f_F(x)$ at some x_0 here means the order of vanishing of the Taylor series of $f_F(x)$ at x_0 . Also, the nondegeneracy condition in Theorem 2.5 is equivalent to the statement that the zeroes of each $f_F(x)$ on $(\mathbb{R} - \{0\})^n$ are of order at most one. It may be worth pointing out that since the $f_F(x)$ here are quasi-homogeneous polynomials, their zeroes in $(\mathbb{R} - \{0\})^n$ are never isolated.

We now investigate what Theorem 1.1 says in the case where $f(x)$ has nondegenerate Newton polyhedron. The statement (1.4) is equivalent to the statement that

$$(2.6) \quad m\left(\left\{x \in W : \frac{\sum_{i=1}^n (x_i \partial f / \partial x_i)^2}{\prod_{i=1}^n x_i^2} < s\right\}\right) \leq C s^{\varepsilon/2}.$$

Let

$$g(x) = \sum_{i=1}^n \left(x_i \frac{\partial f}{\partial x_i}\right)^2.$$

Then the Newton polyhedron $N(g)$ is the double $2N(f) = \{2x : x \in N(f)\}$, the faces F' of $N(g)$ are the sets $\{2x : x \in F\}$ for faces F of $N(f)$. The statement that $N(f)$ is nondegenerate, namely that for each compact face F of $N(f)$ the function $\nabla f_F(x)$ is non-vanishing on $(\mathbb{R} - \{0\})^n$, translates into the statement that each compact face F' of $N(g)$, $g_{F'}(x)$ has no zeroes at all in $(\mathbb{R} - \{0\})^n$.

Suppose $d(f) > 1$, so that $d(g) > 2$. Let

$$h(x) = \frac{\sum_{i=1}^n (x_i \partial f / \partial x_i)^2}{\prod_{i=1}^n x_i^2} = \frac{g(x)}{\prod_{i=1}^n x_i^2}.$$

Then one can define $N(h)$, $d(h)$, and $h_F(x)$ for faces F of $N(h)$ analogously to Definitions 2.2–2.4. So we have $d(h) = d(g) - 2 = 2d(f) - 2 > 0$. Due to the analogous statement holding for $g(x)$, if F is a compact face of $N(h)$, then the function $h_F(x)$ has no zeroes in $(\mathbb{R} - \{0\})^n$.

Note that the vertices of $N(h)$ may now have components as low as -2 . Nonetheless, many of the arguments of [8] extend to $h(x)$, in particular the proof of Theorem 1.2 of [8], which implies that since each for each compact face F of $N(h)$ the function $h_F(x)$ has no zeroes in $(\mathbb{R} - \{0\})^n$, equation (2.6) holds for all $\varepsilon/2 < 1/d(h) = 1/(2d(f) - 2)$. This can also be shown using toric resolution of singularities, similarly to the arguments in [1]. Consequently, (1.4) holds for any $\varepsilon < 1/(d(f) - 1)$, or equivalently, when

$$\frac{\varepsilon}{\varepsilon + 1} < \frac{\frac{1}{d(f)-1}}{\frac{1}{d(f)-1} + 1} = \frac{1}{d(f)}.$$

As a result, the second part of Theorem 1.1 says that $|\hat{\mu}(\lambda)| \leq B(1 + |\lambda|)^{-\delta}$ for each $\delta < \min(1/2, 1/d(f))$. Up to endpoints, these are the estimates provided by Theorem 2.6.

Such estimates are best possible since by taking $\lambda_k = 0$ for $k < n + 1$, one reduces to the oscillatory integral $I(\lambda)$ for which one has sharpness by [21] (part (2) of Theorem 2.5 here). Thus we see that Theorem 1.1 provides another approach to proving such estimates, modulo endpoints.

3. The proof of Theorem 1.1

3.1. Some preliminary lemmas

We will make use of the following relatively easy lemma, which follows from Lemma 3.2 of [12].

Lemma 3.1 (Lemma 3.2 of [12]). *Let (E, μ) be a finite measure space and suppose $g(x)$ is a measurable function on E such that for some positive constants C and δ , for all $t > 0$ one has $\mu(\{x \in E : |g(x)| < t\}) \leq Ct^\delta$. There is a constant $D_\delta > 0$ such that the following holds for all $M \neq 0$.*

- If $\delta < 1$, then $\int_E \min(1, |Mg|^{-1}) d\mu < CD_\delta |M|^{-\delta}$.
- If $\delta = 1$, then $\int_E \min(1, |Mg|^{-1}) d\mu < CD_1(1 + \log_+ |M|)|M|^{-1} + \mu(E)|M|^{-1}$.
- If $\delta > 1$, then $\int_E \min(1, |Mg|^{-1}) d\mu < C(|M|^{-\delta} + D_1|M|^{-1}) + \mu(E)|M|^{-1}$.

For the non-polynomial case, we will also need the following lemma from [12].

Lemma 3.2 (Corollary 2.1.2 of [12]). *Suppose $f_1(y_1, \dots, y_m), \dots, f_l(y_1, \dots, y_m)$ are real analytic functions on a neighborhood of the origin, none identically zero. Then there are an $m - 1$ dimensional ball $B_{m-1}(0, \eta)$ and a positive integer p such that for each s_1, \dots, s_l and each $(y_1, \dots, y_{m-1}) \in B_{m-1}(0, \eta)$, the set*

$$\{y_m : |y_m| < \eta \text{ and } f_i(y_1, \dots, y_m) < s_i \text{ for each } i\}$$

consists of at most p intervals.

3.2. The beginning of the proof of Theorem 1.1

The size of the domain U will be determined by our arguments; at certain junctures, U will have to be sufficiently small for the arguments to be valid. Also, we will always prove bounds of the form $C|\lambda|^{-\delta}$ for $|\lambda| \geq 2$ rather than $C(1 + |\lambda|)^{-\delta}$ for all λ , since the latter will always hold for $|\lambda| < 2$ simply by taking absolute values inside the integral and integrating. The exposition is somewhat easier if we prove estimates in the former form. In addition, we will always be bounding $|\hat{\mu}(\lambda)|$; bounds for $|I(\lambda)|$ will follow by setting $\lambda_k = 0$ for $k < n + 1$.

To start the proof, we observe that we may assume that $|(\lambda_1, \dots, \lambda_n)| < |\lambda_{n+1}|$, for if U is sufficiently small, if $|(\lambda_1, \dots, \lambda_n)| \geq |\lambda_{n+1}|$, then the gradient of the phase function is of magnitude at least $C|\lambda|$, and one may obtain a bound of $C'|\lambda|^{-1}$ simply by integrating by parts, better than the estimates we need. Hence in our arguments we will always assume that $|(\lambda_1, \dots, \lambda_n)| < |\lambda_{n+1}|$. This in particular implies that $|\lambda_{n+1}| \geq \frac{1}{\sqrt{2}}|\lambda|$.

We next define the sets U_i by

$$(3.1) \quad U_i = \{x \in U : |x_i \partial_{x_i} f(x)| > |x_j \partial_{x_j} f(x)| \text{ for } j \neq i\}.$$

As long as no two functions $x_j \partial_{x_j} f(x)$ are the same, up to a set of measure zero we will have $\bigcup_{i=1}^n U_i = U$. In the rare event that two functions $x_j \partial_{x_j} f(x)$ are in fact the same, we simply remove redundant $x_i \partial_{x_i} f(x)$ from the list when defining the U_i in (3.1), so that we always have $\bigcup_i U_i = U$ up to a set of measure zero. The idea now is that since we are trying to integrate in directions based on the weighted gradient flow along $(x_1 \partial_{x_1} f(x), \dots, x_n \partial_{x_n} f(x))$, on each U_i we will integrate by parts in the x_i direction, since in this direction $|x_i \partial_{x_i} f(x)|$ is at least $c|(x_1 \partial_{x_1} f(x), \dots, x_n \partial_{x_n} f(x))|$ for $c = n^{-1/2}$. To this end, we correspondingly define the integrals $I_i(\lambda)$ by

$$(3.2) \quad I_i(\lambda) = \int_{U_i} e^{-i\lambda_{n+1}f(x_1, \dots, x_n) - i\lambda_1 x_1 - \dots - i\lambda_n x_n} \phi(x_1, \dots, x_n) dx_1 \cdots dx_n.$$

Since $\bigcup_i U_i = U$ up to a set of measure zero, we have $\hat{\mu}(\lambda) = \sum_i I_i(\lambda)$, and in order to prove Theorem 1.1, it suffices to show that each $|I_i(\lambda)|$ is bounded by the appropriate of $A|\lambda|^{-\delta}$ or $B|\lambda|^{-\delta}$ as in the statement of the theorem.

Let

$$P(x) = f(x_1, \dots, x_n) + \frac{\lambda_1}{\lambda_{n+1}} x_1 + \dots + \frac{\lambda_n}{\lambda_{n+1}} x_n,$$

so that the phase function in (3.2) is given by $\lambda_{n+1} P(x)$. The function $P(x)$ is sensible to focus on here since $|\lambda_{n+1}| \sim |\lambda|$, and for our arguments it is helpful to view the phase function as a perturbation of the phase function when $\lambda_k = 0$ for all $k < n + 1$.

Where ε is such that (1.4) holds, we write $U_i = D_1 \cup D_2$, where

$$(3.3) \quad \begin{aligned} D_1 &= \left\{ x \in U_i : |\partial_{x_i} P(x)| \leq |\lambda|^{-1/(\varepsilon+1)} \prod_{j \neq i} |x_j| \right\}, \\ D_2 &= \left\{ x \in U_i : |\partial_{x_i} P(x)| > |\lambda|^{-1/(\varepsilon+1)} \prod_{j \neq i} |x_j| \right\}. \end{aligned}$$

We correspondingly write $I_i(\lambda) = J_1(\lambda) + J_2(\lambda)$, where

$$(3.4) \quad J_1(\lambda) = \int_{D_1} e^{-i\lambda_{n+1}f(x_1, \dots, x_n) - i\lambda_1 x_1 - \dots - i\lambda_n x_n} \phi(x_1, \dots, x_n) dx_1 \cdots dx_n,$$

$$(3.5) \quad J_2(\lambda) = \int_{D_2} e^{-i\lambda_{n+1}f(x_1, \dots, x_n) - i\lambda_1 x_1 - \dots - i\lambda_n x_n} \phi(x_1, \dots, x_n) dx_1 \cdots dx_n.$$

To bound $|J_1(\lambda)|$, we will simply take absolute values of the integrand and integrate in all variables, but it will take some effort to properly analyze the result. To bound $|J_2(\lambda)|$, we will perform integrations by parts in the x_i variable. For this, we will need that each domain of integration in the x_i variable consists of a number of intervals that is uniformly bounded. When $f(x)$ is a polynomial, this is immediate, and for general real analytic $f(x)$, this follows from applying Lemma 3.2 as follows.

We take the y_m variable in that lemma to be the x_i variable here, and the remaining y_k variables to be the x_j for $j \neq i$ along with two additional variables, which we call z_1 and z_2 . We take the s_i of the lemma to be zero in all cases, and we let the $f_i(y)$ of the lemma to be the functions $(x_j \partial_{x_j} f(x))^2 - (x_i \partial_{x_i} f(x))^2$ for $j \neq i$, along with the functions $z_1 \prod_{j \neq i} x_j^2 - (\partial_{x_i} f(x) + z_2)^2$, which we need for $z_1 = |\lambda|^{-2/(\varepsilon+1)}$ and $z_2 = \lambda_i / \lambda_{n+1}$.

Although Lemma 3.2 is a local statement, by compactness one may extend the above application of Lemma 3.2 to the whole domain of integration here, so that there exists an N such that for all $|\lambda| \geq 2$ and all x_j for $j \neq i$, the x_i domain of integration of $J_2(\lambda)$ consists of at most N intervals.

We will bound $|J_1(\lambda)|$ and $|J_2(\lambda)|$ separately, starting with $|J_2(\lambda)|$.

3.3. The analysis of $J_2(\lambda)$

Since the exponents δ appearing in Theorem 1.1 satisfy $\delta < 1$, we can remove the regions where $|x_j| \leq |\lambda|^{-1}$ for some j from the domains of integration when bounding $|J_2(\lambda)|$. In other words, it suffices to bound $|J'_2(\lambda)|$, where $J'_2(\lambda)$ is given by

$$(3.6) \quad J'_2(\lambda) = \int_{\{x \in D_2 : |x_j| > |\lambda|^{-1} \text{ for all } j\}} e^{-i\lambda_{n+1}f(x) - i\lambda_1x_1 - \dots - i\lambda_nx_n} \phi(x) dx$$

For each $k = (k_1, \dots, k_n)$ such that the set $\{x : 2^{-k_j-1} < |x_j| \leq 2^{-k_j} \text{ for all } j\}$ intersects the domain of integration of (3.6), we define $J_{k,2}(\lambda)$ by

$$(3.7) \quad J_{k,2}(\lambda) = \int_{\{x \in D_2 : |x_j| > |\lambda|^{-1}, 2^{-k_j-1} < |x_j| \leq 2^{-k_j} \text{ for all } j\}} e^{-i\lambda_{n+1}f(x) - i\lambda_1x_1 - \dots - i\lambda_nx_n} \phi(x) dx.$$

Hence $J'_2(\lambda) = \sum_k J_{k,2}(\lambda)$. Without loss of generality, we may assume U is a subset of the unit ball, so that each $k_j \geq 0$ and there at most $C(\ln |\lambda|)^n$ terms $J_{k,2}(\lambda)$ to consider.

In accordance with the weighted gradient flow concept, since we are on U_i , where $|x_i \partial_{x_i} f(x)| \geq n^{-1/2} |(x_1 \partial_{x_1} f(x), \dots, x_n \partial_{x_n} f(x))|$, a given term $J_{k,2}(\lambda)$ will be analyzed using an integration by parts in the x_i direction.

In (3.7), we now integrate by parts in the x_i variable over in any of the at most N intervals of integration for fixed other variables. Suppose $L = [l_1, l_2]$ is one such interval. Then the exponential appearing in (3.7) can be written as $e^{-i\lambda_{n+1}P(x)}$, which we may write as $-i\lambda_{n+1} \partial_{x_i} P(x) (e^{-i\lambda_{n+1}P(x)} / (-i\lambda_{n+1} \partial_{x_i} P(x)))$. Note that the denominator $-i\lambda_{n+1} \partial_{x_i} P(x)$ is never zero by the definition of D_2 . We integrate by parts in x_i over L , integrating $-i\lambda_{n+1} \partial_{x_i} P(x) e^{-i\lambda_{n+1}P(x)}$ to $e^{-i\lambda_{n+1}P(x)}$ and differentiating the rest. Taking absolute values in the result, we get the following, where $P_0(x_i)$ denotes $P(x)$ as a function of x_i with all other variables fixed:

$$(3.8) \quad \left| \int_L e^{-i\lambda_{n+1}f(x_1, \dots, x_n) - i\lambda_1x_1 - \dots - i\lambda_nx_n} \phi(x_1, \dots, x_n) dx_i \right| \\ \leq C \frac{1}{|\lambda_{n+1}|} \left(\frac{1}{|\partial_{x_i} P_0(l_1)|} + \frac{1}{|\partial_{x_i} P_0(l_2)|} + \int_L \frac{|\partial_{x_i x_i}^2 P_0(x_i)|}{(\partial_{x_i} P_0(x_i))^2} dx_i + \int_L \frac{1}{|\partial_{x_i} P_0(x_i)|} dx_i \right).$$

We would like to integrate the first integral in (3.8) back to get terms similar to the endpoint terms of (3.8). For this to work, we need that $\partial_{x_i x_i}^2 P_0(x_i)$ changes sign at boundedly many points in a given interval L . Since $\partial_{x_i x_i}^2 P_0(x_i) = \partial_{x_i x_i}^2 f(x_1, \dots, x_n)$, we may show this by once again invoking Lemma 3.2, this time for one function $-(\partial_{x_i x_i}^2 f(x_1, \dots, x_n))^2$, letting the y_m variable be x_i and y_1, \dots, y_{m-1} the remaining x_j variables, and $s_1 = 0$. Again, due to compactness, it suffices to have the local result of Lemma 3.2 (or alternatively,

one could just assume U is a sufficiently small neighborhood of the origin.) In any event, there is an N_0 such that $\partial_{x_i x_i}^2 P_0(x_i)$ changes sign at most N_0 times on any interval L . Consequently, one may write the integral in (3.8) as the union of at most N_0 intervals on which $\partial_{x_i x_i}^2 P_0(x_i)$ is either nonnegative or nonpositive, and integrate back to terms similar to the endpoint terms. Specifically, one may bound (3.8) by

$$(3.9) \quad C \frac{1}{|\lambda_{n+1}|} \left(\frac{1}{|\partial_{x_i} P_0(l_1)|} + \frac{1}{|\partial_{x_i} P_0(l_2)|} + \sum_l \frac{1}{|\partial_{x_i} P_0(j_l)|} + \int_L \frac{1}{|\partial_{x_i} P_0(x_i)|} dx_i \right).$$

Here there are at most N_0 terms in the sum of (3.9). Since the domain of integration of $J_{k,2}$ consists of points where $|\partial_{x_i} P(x)| > |\lambda|^{-1/(\varepsilon+1)} \prod_{j \neq i} |x_j|$, and since $|\lambda_{n+1}| \geq \frac{1}{\sqrt{2}} |\lambda|$ in the situation at hand, we have that (3.9) is bounded by

$$(3.10) \quad C' |\lambda|^{-\varepsilon/(\varepsilon+1)} \left(\prod_{j \neq i} |x_j| \right)^{-1}.$$

Equation (3.10) bounds the x_i integral over a single interval L . There are at most N such intervals, so the overall integral of (3.7) in the x_i direction is also bounded by a constant times $|\lambda|^{-\varepsilon/(\varepsilon+1)} (\prod_{j \neq i} |x_j|)^{-1}$. If one integrates in the remaining $n-1$ variables, keeping in mind that the size of the domain of integration in each x_j direction for $j \neq i$ is bounded by a constant times $|x_j|$, we obtain that

$$|J_{k,2}(\lambda)| \leq C'' |\lambda|^{-\varepsilon/(\varepsilon+1)}.$$

Because there are at most a constant times $(\ln |\lambda|)^n$ possible values of k , the sum of all $|J_{k,2}(\lambda)|$ is at most $C_3 (\ln |\lambda|)^n |\lambda|^{-\varepsilon/(\varepsilon+1)}$. This is less than $C_4 |\lambda|^{-\delta}$ for any $\delta < \varepsilon/(\varepsilon+1)$, so $|J_2(\lambda)| \leq \sum_k |J_{k,2}(\lambda)|$ will always satisfy the bounds needed for Theorem 1.1. Thus we may focus our attention henceforth on bounding $|J_1(\lambda)|$, which we must show is bounded by the appropriate quantity in Theorem 1.1.

3.4. The analysis of $J_1(\lambda)$

Taking absolute values of the integrand in (3.4) and integrating (recalling the definition (3.3) of D_1) leads to

$$(3.11) \quad |J_1(\lambda)| \leq C m \left(\left\{ x \in U_i : |x_i \partial_{x_i} P(x)| \leq |\lambda|^{-1/(\varepsilon+1)} \prod_{j=1}^n |x_j| \right\} \right).$$

Here, as before, m denotes Lebesgue measure. Suppose we are in the special case where $\lambda_k = 0$ for $k < n+1$, the setting of part (1) of Theorem 1.1. Here $P(x) = f(x)$. Since U_i is the set of points where $|x_i \partial_{x_i} f(x)| > |x_j \partial_{x_j} f(x)|$ for all $j \neq i$, (3.11) leads to

$$(3.12) \quad |J_1(\lambda)| \leq C m \left(\left\{ x \in U : \sum_{j=1}^n |x_j \partial_{x_j} f(x)| \leq n |\lambda|^{-1/(\varepsilon+1)} \prod_{j=1}^n |x_j| \right\} \right).$$

By the assumption (1.4), we have that (3.12) is bounded by a constant times $|\lambda|^{-\varepsilon/(\varepsilon+1)}$. This is better than what is needed for part (1) of Theorem 1.1, so we have now shown part (1) of Theorem 1.1.

We focus our attention on proving part (2) of the theorem. We will show that if the neighborhood U of the origin is sufficiently small, the right-hand side of (3.11) is bounded by $C|\lambda|^{-\delta}$ whenever $\delta < \min(\varepsilon/(\varepsilon + 1), 1/2)$.

Since $\partial_{x_i} P(x) = \partial_{x_i} f(x) - c$, where $c = \lambda_i/\lambda_{n+1}$ satisfies $|c| \leq 1$, we can rewrite the measure on the right-hand side of (3.11) as

$$(3.13) \quad m\left(\left\{x \in U_i : \left|\frac{\partial_{x_i} f(x) - c}{\prod_{j \neq i} x_j}\right| \leq |\lambda|^{-1/(\varepsilon+1)}\right\}\right).$$

The idea behind bounding (3.13) is that using resolution of singularities, one can show that if $\varepsilon < 1$, then the measure in (3.13) is bounded by the measure when $c = 0$, while if $\varepsilon \geq 1$, one at least has a uniform bound of $C|\lambda|^{-1/(\varepsilon+1)}$. In the former case, one uses the $c = 0$ case exactly above to get an overall bound of $C|\lambda|^{-\varepsilon/(\varepsilon+1)}$, while in the latter case, since $\varepsilon \geq 1$, (1.4) holds if one replaces $\varepsilon = 1$, and thus one can use the $\varepsilon = 1$ case to say that we have an overall bound of $C|\lambda|^{-1/2}$.

We use resolution of singularities, first proved by Hironaka [14, 15], as follows. Let W be as in Theorem 1.1. Assuming U is a sufficiently small neighborhood of the origin, by resolution of singularities there exists an open $U' \subset W$ containing the closure \bar{U} and a finite collection $\{g_l\}_{l=1}^L$ of proper real analytic mappings $g_l: V_l \rightarrow U'$, where each V_l is open and contains $\bar{g}_l^{-1}(U)$, with the following properties.

Each g_l is one to one outside a set of measure zero. Let $\{h_m(x)\}_{m=1}^M$ denote the list of functions consisting of each $\partial_{x_j} f(x)$, each nonzero difference $(x_j \partial_{x_j} f(x))^2 - (x_k \partial_{x_k} f(x))^2$, and each coordinate function x_j . Then on V_l each function $h_m \circ g_l(x)$ and the Jacobian determinant $\text{Jac}_l(x)$ of each $g_l(x)$ is of the form $a_{lm}(x)p_{lm}(x)$, where $p_{lm}(x)$ is a monomial and $a_{lm}(x)$ never vanishes; in fact, one will have $|a_{lm}(x)| > \delta$ for some $\delta > 0$. For any bounded measurable function $F(x)$ on U , one has

$$(3.14) \quad \left| \int_U F(x) dx \right| \leq \sum_{l=1}^L \int_{g_l^{-1}(U)} |F \circ g_l(x) \text{Jac}_l(x)| dx.$$

Because of our assumption that $\nabla f(0) = 0$, the monomials $p_{lm}(x)$ are never just the constant monomial 1.

Let m_0 be such that $h_{m_0}(x) = \partial_{x_i} f(x)$ for the i being used above, and let j_0 be any index such that $p_{lm_0}(x)$ contains x_{j_0} to at least the first power. Then for some $d \neq 0$, one has

$$\partial_{x_{j_0}}(a_{lm_0}(x)p_{lm_0}(x)) = \left(\partial_{x_{j_0}} a_{lm_0}(x) + \frac{d}{x_{j_0}} a_{lm_0}(x) \right) p_{lm_0}(x).$$

As long as U is a sufficiently small neighborhood of the origin, the term $\frac{d}{x_{j_0}} a_{lm_0}(x)$ will dominate $\partial_{x_{j_0}} a_{lm_0}(x)$ in absolute value for at least one choice of j_0 , since we will be near enough to $g_l^{-1}(0)$. So we may work under the assumption that at each x , there is always some j_0 and some constant $d' > 0$ so that we have

$$(3.15) \quad |\partial_{x_{j_0}}(a_{lm_0}(x)p_{lm_0}(x))| \geq \frac{d'}{|x_{j_0}|} |a_{lm_0}(x)p_{lm_0}(x)|.$$

In order to bound (3.13), we will apply (3.14), letting $F(x)$ be the characteristic function of

$$\left\{ x \in U_i : \left| \frac{(\partial_{x_i} f(x) - c)}{\prod_{j \neq i} x_j} \right| \leq |\lambda|^{-1/(\varepsilon+1)} \right\},$$

or equivalently, the characteristic function of

$$\left\{ x \in U_i : c - |\lambda|^{-1/(\varepsilon+1)} \prod_{j \neq i} |x_j| < \partial_{x_i} f(x) < c + |\lambda|^{-1/(\varepsilon+1)} \prod_{j \neq i} |x_j| \right\}.$$

Let $q_i(x) = \prod_{j \neq i} x_j$. Then $q_i \circ g_l(x)$ is of the form $b_{il}(x)r_{il}(x)$, where $r_{il}(x)$ is a monomial and $|b_{il}(x)| > \delta$ for some positive δ . Then $F \circ g_l(x)$ the characteristic function of

$$(3.16) \quad \left\{ x \in g_l^{-1}(U_i) : c - |\lambda|^{-1/(\varepsilon+1)} |b_{il}(x)r_{il}(x)| < a_{lm_0}(x)p_{lm_0}(x) < c + |\lambda|^{-1/(\varepsilon+1)} |b_{il}(x)r_{il}(x)| \right\}.$$

Denote the set in (3.16) by $E_{il\lambda c}$. Then for the above $F(x)$, equation (3.14) leads to

$$(3.17) \quad m\left(\left\{ x \in U_i : \left| \frac{\partial_{x_i} f(x) - c}{\prod_{j \neq i} x_j} \right| \leq |\lambda|^{-1/(\varepsilon+1)} \right\}\right) \leq \sum_{l=1}^L \int_{E_{il\lambda c}} |\text{Jac}_l(x)| dx.$$

Thus we turn our attention to bounding a given term $\int_{E_{il\lambda c}} |\text{Jac}_l(x)| dx$. We examine the intersection of the domain of integration $E_{il\lambda c}$ with a given dyadic rectangle which we denote by R . Because $\text{Jac}_l(x)$, $b_{il}(x)r_{il}(x)$, and $a_{lm_0}(x)p_{lm_0}(x)$ are comparable to monomials, on $E_{il\lambda c} \cap R$, the functions $|\text{Jac}_l(x)|$, $|b_{il}(x)r_{il}(x)|$, and $|a_{lm_0}(x)p_{lm_0}(x)|$ are within a constant factor of $|\text{Jac}_l(x^*)|$, $|b_{il}(x^*)r_{il}(x^*)|$, and $|a_{lm_0}(x^*)p_{lm_0}(x^*)|$, respectively, where $x^* \in E_{il\lambda c} \cap R$ is fixed.

As long as the neighborhood of U is sufficiently small, not only does (3.15) hold, but also for each R there will necessarily be a single j_0 for which (3.15) holds throughout R , since we will be close enough to $g_l^{-1}(0)$ for this to be true. By (3.15) and (3.16), the width of the x_{j_0} cross section of $E_{il\lambda c} \cap R$ is bounded by

$$C |\lambda|^{-1/(\varepsilon+1)} |b_{il}(x^*)r_{il}(x^*)| \left| \frac{a_{lm_0}(x^*)p_{lm_0}(x^*)}{x_{j_0}^*} \right|^{-1}.$$

The width of this cross section is also trivially bounded by $x_{j_0}^*$. These can be combined by saying the width of this cross section is bounded by the following, where R_0 denotes the cross section of R in the x_{j_0} direction:

$$C \int_{R_0} \min(1, |\lambda|^{-1/(\varepsilon+1)} |b_{il}(x^*)r_{il}(x^*)| |a_{lm_0}(x^*)p_{lm_0}(x^*)|^{-1}) dx_{j_0}.$$

Integrating this in the remaining $n - 1$ variables and inserting the Jacobian factor, we see that

$$(3.18) \quad \int_{E_{il\lambda c} \cap R} |\text{Jac}_l(x)| dx \leq C \int_R |\text{Jac}_l(x^*)| \min(1, |\lambda|^{-1/(\varepsilon+1)} |b_{il}(x^*)r_{il}(x^*)| |a_{lm_0}(x^*)p_{lm_0}(x^*)|^{-1}) dx.$$

Since the various factors in (3.18) stay within a bounded factor on R , we may replace the bound (3.18) by

$$(3.19) \quad \int_{E_{il\lambda c} \cap R} |\text{Jac}_l(x)| dx \leq C \int_R |\text{Jac}_l(x)| \min(1, |\lambda|^{-1/(\varepsilon+1)} |b_{il}(x) r_{il}(x)| |a_{lm_0}(x) p_{lm_0}(x)|^{-1}) dx.$$

We next add (3.19) over all R intersecting $g_l^{-1}(U_i)$. Letting W_{il} denote the union of all R intersecting $g_l^{-1}(U_i)$, we get

$$(3.20) \quad \int_{E_{il\lambda c}} |\text{Jac}_l(x)| dx \leq C \int_{W_{il}} |\text{Jac}_l(x)| \min(1, |\lambda|^{-1/(\varepsilon+1)} |b_{il}(x) r_{il}(x)| |a_{lm_0}(x) p_{lm_0}(x)|^{-1}) dx.$$

By shrinking U if necessary, we can always assume that $W_{il} \subset g_l^{-1}(U')$, where U' is the open set containing \bar{U} of the resolution of singularities procedure. Thus (3.20) becomes

$$(3.21) \quad \int_{E_{il\lambda c}} |\text{Jac}_l(x)| dx \leq C \int_{g_l^{-1}(U')} |\text{Jac}_l(x)| \min(1, |\lambda|^{-1/(\varepsilon+1)} |b_{il}(x) r_{il}(x)| |a_{lm_0}(x) p_{lm_0}(x)|^{-1}) dx.$$

Now going back into the original coordinates using the coordinate change map g_l , (3.21) leads to

$$(3.22) \quad \int_{E_{il\lambda c}} |\text{Jac}_l(x)| dx \leq C \int_{U'} \min(1, |\lambda|^{-1/(\varepsilon+1)} \left(\prod_{j \neq i} |x_j| \right) |\partial_{x_i} f(x)|^{-1}) dx.$$

We can refine (3.22) in the following way. Since each nonzero difference of the form $(x_j \partial_{x_j} f(x))^2 - (x_k \partial_{x_k} f(x))^2$ was monomialized, in particular, each difference of the form $(x_i \partial_{x_i} f(x))^2 - (x_j \partial_{x_j} f(x))^2$ was monomialized for each j . Thus in the blown up coordinates of $g_l^{-1}(U')$, the set of points where $|x_i \partial_{x_i} f(x)| > |x_j \partial_{x_j} f(x)|$ for a given j , which is the same as the set where $(x_i \partial_{x_i} f(x))^2 - (x_j \partial_{x_j} f(x))^2 > 0$, consists of the points in $g_l^{-1}(U')$ where a certain monomial is positive. This consists of the set of points in $g_l^{-1}(U')$ that are in a certain collection of octants. Thus in the blown up coordinates, the set of points in $g_l^{-1}(U')$ where $|x_i \partial_{x_i} f(x)| > |x_j \partial_{x_j} f(x)|$ for all j are also the points in $g_l^{-1}(U')$ that are in certain octants. Furthermore, the above dyadic rectangles R will be contained in these octants. Hence the relation $|x_i \partial_{x_i} f(x)| > |x_j \partial_{x_j} f(x)|$ for $j \neq i$ will still hold when we return the union of all such R to the original coordinates as in (3.22). Thus we may amend (3.22) to

$$(3.23) \quad \int_{E_{il\lambda c}} |\text{Jac}_l(x)| dx \leq C \int_{\{x \in U' : |x_i \partial_{x_i} f(x)| > |x_j \partial_{x_j} f(x)| \text{ for } j \neq i\}} \min \left(1, |\lambda|^{-1/(\varepsilon+1)} \left(\prod_{j \neq i} |x_j| \right) |\partial_{x_i} f(x)|^{-1} \right) dx.$$

Since

$$\left(\prod_{j \neq i} |x_j|\right) |\partial_{x_i} f(x)|^{-1} = \left(\frac{|x_i \partial_{x_i} f(x)|}{\prod_{j=1}^n |x_j|}\right)^{-1} \quad \text{and} \quad |x_i \partial_{x_i} f(x)| > |x_j \partial_{x_j} f(x)|$$

for all $j \neq i$ in the domain of integration of (3.23), we may let

$$r(x) = \frac{\sum_{i=1}^n |x_i \partial f / \partial x_i|}{\prod_{i=1}^n |x_i|}$$

(the function in Theorem 1.1), and then (3.23) implies

$$\int_{E_{il\lambda c}} |\text{Jac}_l(x)| dx \leq C \int_{U'} \min(1, |\lambda|^{-1/(\varepsilon+1)} |r(x)|^{-1}) dx.$$

Combining with (3.17) leads to

(3.24)

$$m\left(\left\{x \in U_i : \left|\frac{\partial_{x_i} f(x) - c}{\prod_{j \neq i} x_j}\right| \leq |\lambda|^{-1/(\varepsilon+1)}\right\}\right) \leq C' \int_{U'} \min(1, |\lambda|^{-1/(\varepsilon+1)} |r(x)|^{-1}) dx.$$

Since the left-hand side of (3.24) is the right-hand side of (3.13), which by (3.11) is an upper bound for $|J_1(\lambda)|$, we conclude that

$$(3.25) \quad |J_1(\lambda)| \leq C' \int_{U'} \min(1, |\lambda|^{-1/(\varepsilon+1)} |r(x)|^{-1}) dx.$$

We now apply Lemma 3.1, using the definition (1.4) of ε and the fact that $U' \subset W$. If $\varepsilon < 1$, the first part of Lemma 3.1 gives that the right-hand side of (3.25) is bounded by a constant times $|\lambda|^{-\varepsilon/(\varepsilon+1)}$, which provides the bound we need for $|J_1(\lambda)|$ to give the correct exponent in part (2) of Theorem 1.1. On the other hand, if $\varepsilon \geq 1$, then (1.4) still holds for $\varepsilon = 1$, in which case the second part of Lemma 3.1 gives a bound of $C' |\lambda|^{-1/2} \ln |\lambda|$, which is better than $C'' |\lambda|^{-\delta}$ for all $\delta < 1/2$, which is what we need for part (2) of Theorem 1.1 when $\varepsilon \geq 1$. Thus we see that in all cases that $|J_1(\lambda)|$ is bounded by the appropriate quantity in part (2) of Theorem 1.1. This completes the proof of Theorem 1.1.

Funding. This work was supported by a grant from the Simons Foundation.

References

- [1] Arnol'd, V. I., Guseĭn-Zade, S. M. and Varchenko, A. N.: *Singularities of differentiable maps. Vol. II*. Monogr. Math. 83, Birkhäuser, Boston, MA, 1988. Zbl 0659.58002 MR 0966191
- [2] Bak, J.-G., McMichael, D., Vance, J. and Wainger, S.: *Fourier transforms of surface area measure on convex surfaces in \mathbb{R}^3* . *Amer. J. Math.* **111** (1989), no. 4, 633–668. Zbl 0687.49028 MR 1011552
- [3] Basu, S., Guo, S., Zhang, R. and Zorin-Kranich, P.: *A stationary set method for estimating oscillatory integrals*. To appear in *J. Eur. Math. Soc.*, published online first 2024, DOI 10.4171/jems/1506.
- [4] Brandolini, L., Hofmann, S. and Iosevich, A.: *Sharp rate of average decay of the Fourier transform of a bounded set*. *Geom. Funct. Anal.* **13** (2003), no. 4, 671–680. Zbl 1032.42017 MR 2006553

- [5] Bruna, J., Nagel, A. and Wainger, S.: [Convex hypersurfaces and Fourier transforms](#). *Ann. of Math. (2)* **127** (1988), no. 2, 333–365. Zbl [0666.42010](#) MR [0932301](#)
- [6] Carbery, A., Christ, M. and Wright, J.: [Multidimensional van der Corput and sublevel set estimates](#). *J. Amer. Math. Soc.* **12** (1999), no. 4, 981–1015. Zbl [0938.42008](#) MR [1683156](#)
- [7] Green, J.: [Lower bounds on \$L^p\$ quasi-norms and the uniform sublevel set problem](#). *Mathematika* **67** (2021), no. 2, 296–323. Zbl [07738343](#) MR [4220992](#)
- [8] Greenblatt, M.: [Oscillatory integral decay, sublevel set growth, and the Newton polyhedron](#). *Math. Ann.* **346** (2010), no. 4, 857–895. Zbl [1192.14005](#) MR [2587095](#)
- [9] Greenblatt, M.: [Maximal averages over hypersurfaces and the Newton polyhedron](#). *J. Funct. Anal.* **262** (2012), no. 5, 2314–2348. Zbl [1243.42026](#) MR [2876407](#)
- [10] Greenblatt, M.: [Fourier transforms of irregular mixed homogeneous hypersurface measures](#). *Math. Nachr.* **291** (2018), no. 7, 1075–1087. Zbl [1394.42005](#) MR [3807636](#)
- [11] Greenblatt, M.: [Fourier transforms of indicator functions, lattice point discrepancy, and the stability of integrals](#). *Math. Ann.* **380** (2021), no. 3-4, 1959–1990. Zbl [1471.42021](#) MR [4297203](#)
- [12] Greenblatt, M.: [A method for bounding oscillatory integrals in terms of non-oscillatory integrals](#). Preprint, arXiv: [2210.09093v3](#), 2024.
- [13] Gressman, P. T.: [Scalar oscillatory integrals in smooth spaces of homogeneous type](#). *Rev. Mat. Iberoam.* **31** (2015), no. 1, 215–244. Zbl [1321.42030](#) MR [3320838](#)
- [14] Hironaka, H.: [Resolution of singularities of an algebraic variety over a field of characteristic zero. I](#). *Ann. of Math. (2)* **79** (1964), 109–203. Zbl [0122.38603](#) MR [0199184](#)
- [15] Hironaka, H.: [Resolution of singularities of an algebraic variety over a field of characteristic zero. II](#). *Ann. of Math. (2)* **79** (1964), 205–326. Zbl [1420.14031](#) MR [0199184](#)
- [16] Ikromov, I. A., Kempe, M. and Müller, D.: [Estimates for maximal functions associated with hypersurfaces in \$\mathbb{R}^3\$ and related problems of harmonic analysis](#). *Acta Math.* **204** (2010), no. 2, 151–271. Zbl [1223.42011](#) MR [2653054](#)
- [17] Ikromov, I. A. and Müller, D.: [Uniform estimates for the Fourier transform of surface carried measures in \$\mathbb{R}^3\$ and an application to Fourier restriction](#). *J. Fourier Anal. Appl.* **17** (2011), no. 6, 1292–1332. Zbl [1241.42009](#) MR [2854839](#)
- [18] Nagel, A., Seeger, A. and Wainger, S.: [Averages over convex hypersurfaces](#). *Amer. J. Math.* **115** (1993), no. 4, 903–927. Zbl [0832.42010](#) MR [1231151](#)
- [19] Phong, D. H., Stein, E. M. and Sturm, J. A.: [On the growth and stability of real-analytic functions](#). *Amer. J. Math.* **121** (1999), no. 3, 519–554. Zbl [1015.26031](#) MR [1738409](#)
- [20] Stein, E. M.: [Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals](#). Princeton Math. Ser. 43, Princeton University Press, Princeton, NJ, 1993. Zbl [0821.42001](#) MR [1232192](#)
- [21] Varčenko, A. N.: [Newton polyhedra and estimates of oscillatory integrals](#). *Funct. Anal. Appl.* **10** (1977), 175–196. Zbl [0351.32011](#) MR [0422257](#)

Received April 5, 2024; revised September 25, 2024.

Michael Greenblatt

Department of Math, Statistics, and Computer Science, University of Illinois at Chicago
 851 S. Morgan St, Chicago, IL 60607-7045, USA;
greenbla@uic.edu