



Christopher Lutsko · Athanasios Sourmelidis · Niclas Technau

Pair correlation of the fractional parts of αn^θ

Received 8 January 2022; revised 15 February 2023

Abstract. Fix $\alpha, \theta > 0$ and consider the sequence $(\alpha n^\theta \bmod 1)_{n \geq 1}$. Since the seminal work of Rudnick–Sarnak (1998) and due to the Berry–Tabor conjecture in quantum chaos, the fine-scale properties of these dilated monomial sequences have been intensively studied. In this paper, we show that for $\theta < 14/41$ and $\alpha > 0$, the pair correlation function is Poissonian. While (for a given $\theta \neq 1$) this strong pseudo-randomness property has been proven for almost all values of α , there are next-to-no instances where this has been proven for explicit α . Our result holds for all $\alpha > 0$ and relies solely on classical Fourier analytic techniques. This addresses (in the sharpest possible way) a problem posed by Aistleitner–El-Baz–Munsch (2021).

Keywords: local statistics, sequences modulo 1, exponential sums, van der Corput’s method, Weyl’s differencing method.

1. Introduction

Let $x = (x_n)_{n \geq 1}$ be a sequence on the unit interval $[0, 1)$. The *pair correlation function* of x measures the correlation between points in the initial segment $\{x_n : n \leq N\}$ on the scale of the mean spacing, $1/N$, and is defined by

$$R(x, N, f) := \frac{1}{N} \sum_{i \neq j \leq N} \sum_{k \in \mathbb{Z}} f(N(x_i - x_j + k)),$$

where $f \in \mathcal{C}_c^\infty(\mathbb{R})$ is a compactly supported, C^∞ -function. The sequence x is said to have *Poissonian pair correlation* if the pair correlation function converges to the integral of f (over \mathbb{R}) as $N \rightarrow \infty$, just as one would expect for uniformly distributed and

Christopher Lutsko: Department of Mathematics, Rutgers University, Hill Center - Busch Campus, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA; Institut für Mathematik, Universität Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland; christopher.lutsko@math.uzh.ch

Athanasios Sourmelidis: Institute of Analysis and Number Theory, Graz University of Technology, Steyrergasse 30, 8010 Graz, Austria; sourmelidis@math.tugraz.at

Niclas Technau: Department of Mathematics, University of Wisconsin–Madison, 480 Lincoln Drive, Madison, WI 53706, USA; niclastechnau@gmail.com

Mathematics Subject Classification 2020: 11K06 (primary); 11L07 (secondary).

independent random variables. That is, the sequence x has Poissonian pair correlation if for all $f \in \mathcal{C}_c^\infty(\mathbb{R})$,

$$\lim_{N \rightarrow \infty} R(x, N, f) = \int_{\mathbb{R}} f(t) dt.$$

The notion of Poissonian pair correlation defines a strong measure of pseudo-randomness and is a basic concept in quantum chaos. Unsurprisingly, various efforts have been made [2, 6, 12, 21, 27, 31] to study the pair correlation function of monomial sequences

$$(\alpha n^\theta \bmod 1)_{n \geq 1}, \quad (1.1)$$

where $\theta > 0$ and $\alpha > 0$. However, little progress has been made to verify that the pair correlation of such monomial sequences is Poissonian (under explicit conditions on α, θ). We present the state of the art for (1.1) in Section 1.1. In this paper, we prove the first general and explicit result showing that such monomial sequences exhibit Poissonian pair correlation.

Theorem 1.1. *If $\theta \in (0, 14/41)$ and $\alpha > 0$, then (1.1) has Poissonian pair correlation.*

Remark. (1) Our method applies to higher level correlations, although this generalization is not straightforward as it requires a genuinely multidimensional approach (see [17, 18]). Moreover, the only arithmetic input of our method are exponential sum bounds. Thus, with some modification, the method can be extended to more general sequences satisfying certain growth conditions.

(2) The method of proof allows one to show that the pair correlation function converges to $\int_{\mathbb{R}} f(x) dx$ with a polynomially decaying error in N which is uniform for all α in a fixed compact interval.

(3) When $\theta = 1/3$ and $\alpha^3 \in \mathbb{Q}$, then the triple correlation is not Poissonian (because of the cubes n^3). Thus, Theorem 1.1 gives an example of a sequence whose pair correlation is Poissonian, but whose triple correlation is not.

Organization of the paper. Section 1.1 presents a brief history of these monomial sequences. Section 1.2 sketches the proof of Theorem 1.1, and Section 1.3 provides a heuristic argument which indicates the limitations of our method. In Section 2, we collect lemmas, reducing matters to bounding certain exponential sums. Finally, in Section 3 we prove Theorem 1.1.

1.1. Background

The study of monomial sequences dates back to Weyl [33], who used them in his study of the uniform distribution (see [14] or [7]). More recently, there has been renewed interest in these sequences. In part, this is due to the well-known Berry–Tabor conjecture [4], which hypothesizes a link between the pseudo-randomness properties of energy levels and dynamics of quantum systems. For more details, see either of the following review papers [19, 25].

The holy grail of this field is to find circumstances for which a sequence has Poissonian gap statistics. That is, consider the distribution of gaps between neighboring first N elements of the sequence – scaled to have average 1 – then we say the sequence exhibits Poissonian gap statistics if this (finite) distribution converges to the exponential distribution, as one would expect for independent random variables. While the aforementioned behavior is conjectured in many instances, it is truly challenging to prove. Thus, mathematicians have turned to weaker measures of pseudo-randomness. In particular, there has been a lot of recent work on the pair correlation. Indeed, if one could show that the m -level correlation converges to the expected value for independent random variables (for every $m \geq 2$) then, by the method of moments, one can infer that the sequence has Poissonian gap statistics.

If we consider the random variable counting the number of sequence elements in a randomly shifted set of size comparable to $1/N$, then the m -level correlations arise from the moments of this variable. Thus, the m -level correlations are natural measures of pseudo-randomness in their own right. We refer to [20] for further discussion.

1.1.1. Pair correlation of deterministic sequences. Those few deterministic sequences, whose pair correlation functions are known to be Poissonian, either require the presence of particularly strong arithmetic structure, or tools from homogeneous dynamics to apply. An example of the former is the work of Kurlberg and Rudnick [15] on the (appropriately normalized) spacing of the quadratic residues of a highly composite modulus. In fact, they show that the gap statistics are Poissonian. However, this setting requires the use of arithmetic tools which cannot be relied on in our situation.

On the homogeneous dynamics side, Elkies and McMullen [9] established a remarkable link between (1.1), for $(\theta, \alpha) = (1/2, 1)$, and flows on the modular surface $\mathrm{SL}_2(\mathbb{R})/\mathrm{SL}_2(\mathbb{Z})$. They used this connection, and tools from homogeneous dynamics, to establish that the corresponding gap distribution is *not* Poissonian. Surprisingly, El-Baz, Marklof, and Vinogradov [8] then exploited the said relationship further to show that, if one removed the squares, the pair correlation is *Poissonian*. However, the connection to homogeneous dynamics requires a particular scaling property which only holds when $\theta = 1/2$ and $\alpha^2 \in \mathbb{Q}$. Indeed, for $\alpha^2 \notin \mathbb{Q}$ it is conjectured [9] the gap statistics are Poissonian.

For the sequence $(\alpha n \bmod 1)_{n \geq 1}$, the three gap theorem (also known as the Steinhaus conjecture) states that the size of the gaps between neighboring points, at any time N , form a set of cardinality at most 3. Hence, the local statistics are certainly not Poissonian. For background see [22, 23].

1.1.2. Metric Poissonian pair correlation. Generally speaking, it is believed that, given a $\theta > 0$, the pseudo-random properties of (1.1) are determined by the Diophantine properties of α (e.g., see [27, Remark 1.2]). However, in the absence of methods to prove Poissonian pair correlation for explicit values of α , Rudnick and Sarnak [27] introduced the concept of *metric Poissonian pair correlation*. Namely, a general sequence $(x_n)_{n \geq 1}$ has metric Poissonian pair correlation, if the dilated sequence $(\alpha x_n \bmod 1)_{n \geq 1}$ has Poissonian pair correlation for all $\alpha > 0$ outside of a (Lebesgue) null set.

For $\theta \in \mathbb{N}_{>1}$, Rudnick and Sarnak [27] proved metric Poissonian pair correlation of (1.1) in the late 90s. The case of a non-integer $\theta > 1$ was only recently settled by Aistleitner, El-Baz, and Munsch [1]. The regime $0 < \theta < 1$ was addressed by Rudnick and the third named author [30].

Special attention has been given to the quadratic case, $\theta = 2$, due to its connection with quantum chaos and the boxed harmonic oscillator. Here, Heath-Brown [12] gave an algorithmic construction of a dense set of α for which the pair correlation is Poissonian. Moreover, there have been some results for longer-range correlations [16, 32], convergence along sparse subsequences [10, 28], and minimal gaps [24, 26, 34]. However, finding explicit α for which the pair correlation is Poissonian remains out of reach.

Finally, it is worth noting that the metric Poisson pair correlation theory has been generalized beyond monomial sequences and exploits some deep connections to additive combinatorics [2, 5]. However, this connection is beyond the scope of this paper.

Notation. Throughout, we use the usual Bachmann–Landau notation: for functions $f, g: X \rightarrow \mathbb{R}$, defined on some set X , we write $f \ll g$ (or $f = O(g)$) to denote that there exists a constant $C > 0$ such that $|f(x)| \leq C|g(x)|$ for all $x \in X$. Moreover, let $f \asymp g$ denote $f \ll g$ and $g \ll f$. Furthermore, let $f = o(g)$ denote that $\frac{f(x)}{g(x)} \rightarrow 0$.

Throughout, we denote $e(x) = e^{2\pi i x}$, and \hat{f} is the Fourier transform (on \mathbb{R}) of f . All of the sums which appear range over integers, in the indicated interval. As $\alpha, \varepsilon, \theta$, and f are considered fixed, we suppress any dependence in the implied constants. Moreover, for ease of notation, $\varepsilon > 0$ may vary from line to line by a bounded constant. Further, we will frequently encounter the exponent

$$\Theta := \frac{1}{1 - \theta}.$$

1.2. Idea of the proof

The proof relies on a well-known Fourier decomposition. First, we include the diagonal term in the pair correlation function, to simplify technicalities. Thus, define

$$\tilde{R}(N, f) := \frac{1}{N} \sum_{y \in [1, N]^2} \sum_{k \in \mathbb{Z}} f(N(\alpha y_1^\theta - \alpha y_2^\theta + k)).$$

Note that Theorem 1.1 is equivalent to showing that (as $N \rightarrow \infty$)

$$\tilde{R}(N, f) = \int_{\mathbb{R}} f(x) dx + f(0) + o(1).$$

By the Poisson summation formula,

$$\tilde{R}(N, f) = \hat{f}(0) + \frac{1}{N^2} \sum_{|k| \in [1, N^{1+\varepsilon}]} \left| \hat{f}\left(\frac{k}{N}\right) \right| \sum_{y \in [1, N]} e(\alpha k y^\theta) + o(1)$$

for $\varepsilon > 0$, where the $o(1)$ -error comes from the fast decay of \hat{f} . Note that f can be decomposed into a sum of an even and an odd function. Further, the Fourier coefficients

of the odd part cancel out, and the Fourier coefficients of the even part are even functions themselves. Thus, without loss of generality we may assume f is even. Hence, it suffices to show that

$$\mathcal{E}(N) := \frac{2}{N^2} \sum_{k \in [1, N^{1+\varepsilon}]} \left| \hat{f}\left(\frac{k}{N}\right) \right| \left| \sum_{y \in [1, N]} e(\alpha k y^\theta) \right|^2 = f(0) + o(1). \quad (1.2)$$

To achieve the desired bound requires a detailed analysis of the exponential sums in (1.2). We argue in, roughly, two steps: first, we decompose the innermost summation, and apply van der Corput's B-process to obtain a saving in the y -summation. Second, we expand the square and use some analytic tricks to reduce the estimates to exponential sums over k . Now to obtain a saving in the k -summation, we again use the B-process coupled with other estimates (such as Weyl differencing).

1.3. Heuristic

After applying the B-process, interchanging the order of summation, extracting the main terms and dealing with the error terms, our task is the following. We need to show that

$$\text{Err} := \frac{2}{N^2} \sum_{\substack{r_1, r_2 > cN^\theta \\ r_1 \neq r_2}}^N \frac{1}{(r_1 r_2)^{(\Theta+1)/2}} \sum_{k \in [1, N^{1+\varepsilon}]} \hat{f}\left(\frac{k}{N}\right) k^\Theta e(\gamma(\mathbf{r})k^\Theta)$$

is $o(1)$ as $N \rightarrow \infty$, where $\gamma(\mathbf{r}) = \beta(r_2^{1-\Theta} - r_1^{1-\Theta})$, and β and c depend only on θ and α . Now we apply partial summation to reduce matters to estimating

$$\left| \sum_{k \in [1, N^{1+\varepsilon}]} e(\gamma(\mathbf{r})k^\Theta) \right|.$$

If we had square root cancellation for this sum – uniformly in $\gamma(\mathbf{r})$ – then our method yields

$$\text{Err} \approx N^{\theta-1/2+\varepsilon}.$$

In other words, even with optimal bounds, we cannot hope to go past the barrier $\theta = 1/2 - \varepsilon$. To move past this barrier, our analytic method would require taking advantage of the cancellation between exponential sums for different values of \mathbf{r} . This seems to be well beyond current technology.

Interestingly, if we consider instead the triple correlation function, the natural barrier to our methods turns out to be $\theta < 1/3$. In fact, as we consider higher and higher correlations, that barrier goes to 0.

1.4. Preliminaries

The following two results are fundamental in the modern study of exponential sums. First, we recall an application of Weyl's differencing method (called the A-process, see [11, Theorem 2.9]).

Theorem 1.2 (A-process). *Let $l \geq 0$ be an integer and let $M > 2$. Suppose $\phi: [a, b] \rightarrow \mathbb{R}$ has $l + 2$ continuous derivatives on $[a, b] \subseteq [M, CM)$, where $C > 1$ is some fixed constant, and assume there exists a constant $F > 0$ such that*

$$\phi^{(r)}(x) \asymp FM^{-r} \quad (1.3)$$

for $r = 1, \dots, l + 2$. Then

$$\sum_{x \in [a, b]} e(\phi(x)) \ll F^{1/(4L-2)} M^{1-(l+2)/(4L-2)} + F^{-1} M, \quad (1.4)$$

where $L := 2^l$. The implicit constant in (1.4) depends on the choice of l and the implicit constant(s) in (1.3).

Further, we will use van der Corput's B-process, which follows from the Poisson summation and a stationary phase argument (see [13, Theorem 8.16]).

Theorem 1.3 (B-process). *Let $\phi: [A, B] \rightarrow \mathbb{R}$ be a C^4 -function so that there are $\Lambda > 0$ and $\eta \geq 1$ with*

$$\Lambda \leq \phi^{(2)}(x) < \eta\Lambda, \quad |\phi^{(3)}(x)| < \frac{\eta\Lambda}{B-A}, \quad |\phi^{(4)}(x)| < \frac{\eta\Lambda}{(B-A)^2}$$

for all $x \in [A, B]$. Let also

$$a = \phi'(A) \quad \text{and} \quad b = \phi'(B).$$

Then

$$\sum_{n \in [A, B]} e(\phi(n)) = e\left(\frac{1}{8}\right) \sum_{m \in [a, b]} \frac{e(\phi(x_m) - mx_m)}{\sqrt{\phi^{(2)}(x_m)}} + \omega_\phi(A, B),$$

where x_m denotes the unique solution to $\phi'(x) = m$. Furthermore,

$$\omega_\phi(A, B) \ll \Lambda^{-1/2} + \eta^2 \log(b - a + 1),$$

where the implied constant is absolute.

We will often need to bound weighted exponential sums. To reduce these estimates to bounding unweighted sums, we use partial summation.

Lemma 1.4. *Let $(a_s)_s$ and $(b_s)_s$ be sequences of complex numbers. Fix a constant $c > 1$. If $T > 0$ is such that*

$$|b_s - b_{s+1}| \leq \frac{T}{s},$$

then

$$\left| \sum_{S \leq s < \tilde{S}} a_s b_s \right| \leq \left(\max_{S \leq s \leq cS} |b_s| + O(T) \right) \max_{S \leq \tilde{S} \leq cS} \left| \sum_{S \leq s < \tilde{S}} a_s \right|$$

for any positive integers S and \tilde{S} satisfying $S \leq \tilde{S} \leq cS$.

2. Reducing to exponential sum bounds

2.1. Decomposing the sums and applying the B-process

Now consider the term $\mathcal{E}(N)$ defined in (1.2). We shall apply the B-process (Theorem 1.3) to the exponential sum in $\mathcal{E}(N)$, but presently we do not have sufficient control on the derivative of $y \mapsto \alpha k y^\theta$. To gain control, we use several decompositions. First, we assume (without loss of generality) that

$$N = N_Q := Q^\Gamma,$$

for some fixed $\Gamma > 0$, which will be chosen to be sufficiently large in our proof (in a way depending on θ), as it is enough to prove that the correlations converge along such a subsequence, see [29, Lemma 3.1]. Thus, we decompose the inner summation into the pieces

$$E_q(k) := \sum_{y \in [N_q, N_{q+1})} e(\alpha k y^\theta),$$

where $N_q := q^\Gamma$. To catch the largest term, we set $N_{Q+1} = N_Q + 1$. Thus,

$$\mathcal{E}(N) = \frac{2}{N^2} \sum_{k \in [1, N^{1+\varepsilon})} \hat{f}\left(\frac{k}{N}\right) \left| \sum_{q \leq Q} E_q(k) \right|^2. \quad (2.1)$$

Now, the next lemma shows that we can replace each $E_q(k)$ by

$$E_q^{(B)}(k) := c_1 \sum_{r \in \mathcal{R}_q(k)} \frac{k^{\Theta/2}}{r^{(\Theta+1)/2}} e(\beta k^\Theta r^{1-\Theta}),$$

where

$$\mathcal{R}_q(k) := \alpha \theta k (N_{q+1}^{\theta-1}, N_q^{\theta-1}],$$

which is the main term after applying the B-process to $E_q(k)$; the constants c_1 and β are defined by

$$c_1 := e\left(-\frac{1}{8}\right) \sqrt{\Theta(\alpha\theta)^\Theta} \quad \text{and} \quad \beta := \alpha^\Theta (\theta^{1-\Theta} - \theta^\Theta).$$

For later reference, let

$$\mathcal{R}(k) := \bigcup_{q \in [1, Q]} \mathcal{R}_q(k).$$

Let

$$\mathcal{E}^{(B)}(N, \hat{f}) := \frac{2}{N^2} \sum_{q \in [1, Q]^2} \sum_{k \in [1, N^{1+\varepsilon})} \hat{f}\left(\frac{k}{N}\right) E_{q_1}^{(B)}(k) \overline{E_{q_2}^{(B)}(k)}. \quad (2.2)$$

Lemma 2.1. *Let $\mathcal{E}(N)$ and $\mathcal{E}^{(B)}(N, \hat{f})$ be defined as in (2.1) and (2.2), respectively. Then*

$$\mathcal{E}(N) = \mathcal{E}^{(B)}(N, \hat{f}) + O(N^{-\theta+\varepsilon} + |\mathcal{E}^{(B)}(N, |\hat{f}|)|^{1/2} N^{-\theta/2+\varepsilon}).$$

Proof. First, we apply Theorem 1.3 to each $E_q(k)$, $q \leq Q$, with $\Lambda = k\alpha\theta(1-\theta)N_{q+1}^{\theta-2}$ and $\eta = 2^{5\Gamma}$. Hence,

$$|E_q(k) - E_q^{(B)}(k)| \ll \frac{N_{q+1}}{\sqrt{N_{q+1}^\theta k}} + \log N \ll N^{1-\theta/2} k^{-1/2},$$

with the implied constant being uniform in q and k . Thus,

$$\mathcal{E}(N) = \frac{2}{N^2} \sum_{k \in [1, N^{1+\varepsilon})} \hat{f}\left(\frac{k}{N}\right) \left| \sum_{q \leq Q} E_q^{(B)}(k) + O(N^{1-\theta/2+1/\Gamma} k^{-1/2}) \right|^2.$$

Taking $\Gamma > 1/\varepsilon$ ensures that in the above error term $N^{1/\Gamma}$ can be replaced by N^ε . Squaring out and applying the Cauchy–Schwarz inequality yields now the lemma. ■

2.2. The diagonal

Presently, our goal is to establish (1.2). In view of Lemma 2.1, it suffices to prove that

$$\mathcal{E}^{(B)}(N, \hat{f}) = f(0) + o(1) \quad \text{and} \quad \mathcal{E}^{(B)}(N, |\hat{f}|) = O(1), \quad N \rightarrow \infty,$$

for in that case (1.2) is true and Theorem 1.1 will follow. Computing the estimates of $\mathcal{E}^{(B)}(N, \hat{f})$ and $\mathcal{E}^{(B)}(N, |\hat{f}|)$ is done by completely analogous way and, therefore, we give detailed proofs only for $\mathcal{E}^{(B)}(N, \hat{f})$.

The main term, $f(0)$, will come from the diagonal term when expanding the square in $\mathcal{E}^{(B)}(N, \hat{f})$. That is, in (2.2) we square out and consider the term

$$\mathcal{D}(N, \hat{f}) := \frac{2|c_1|^2}{N^2} \sum_{k \in [1, N^{1+\varepsilon})} \hat{f}\left(\frac{k}{N}\right) \sum_{r \in \mathcal{R}(k)} \frac{k^\Theta}{r^{\Theta+1}}.$$

Lemma 2.2. *If $\theta \in (0, 1)$, then*

$$\mathcal{D}(N, \hat{f}) = f(0) + o(1) \quad \text{and} \quad \mathcal{D}(N, |\hat{f}|) = O(1), \quad N \rightarrow \infty. \quad (2.3)$$

Proof. By a Riemann integral argument (see, for example, [3, Theorem 3.2]), the sum

$$\sum_{r \in \mathcal{R}(k)} \frac{1}{r^{\Theta+1}} = \frac{(\alpha\theta k)^{-\Theta} - (\alpha\theta k(N+1)^{\theta-1})^{-\Theta}}{-\Theta} + O((\alpha\theta k N^{\theta-1})^{-\Theta-1}).$$

Recall that $|c_1|^2 = \Theta(\alpha\theta)^\Theta$. Thus,

$$\begin{aligned} \mathcal{D}(N, \hat{f}) &= \frac{2}{N^2} \sum_{k \in [1, N^{1+\varepsilon})} \hat{f}\left(\frac{k}{N}\right) \left(N + O\left(\frac{N^{2-\theta}}{k}\right) \right) \\ &= \frac{2}{N} \sum_{k \in [1, N^{1+\varepsilon})} \hat{f}\left(\frac{k}{N}\right) + O(N^{-\theta+\varepsilon}). \end{aligned}$$

Now (2.3) follows by the Poisson summation formula and the fact that f is an even function. The second statement of the lemma follows as above, only this time we employ the rapid decay of \hat{f} to give an upper bound for $\mathcal{D}(N, |\hat{f}|)$. ■

2.3. Partial summation

Lemmas 2.1 and 2.2 reduce the problem to estimating $\mathcal{E}^{(B)}(N, g) - \mathcal{D}(N, g)$ for $g = \widehat{f}$ and $|\widehat{f}|$. Estimating these errors requires a second application of the B -process, this time to the k -variable. In order to have adequate control on the derivative of $y \mapsto \gamma(\mathbf{r})y^\Theta$ in the next section, we introduce a second decomposition. In particular, let $U \in \mathbb{N}$ be such that $e^U \leq N^{1+\varepsilon} < e^{U+1}$. Then we may decompose the sum in k into a sum over intervals $[e^u, e^{u+1})$, the last one being $[e^U, N^{1+\varepsilon})$. Let

$$\mathcal{E}_{\mathbf{q},u}^{(B)}(N, g) := \frac{2}{N^2} \sum_{k \in [e^u, e^{u+1})} g\left(\frac{k}{N}\right) E_{q_1}^{(B)}(k) \overline{E_{q_2}^{(B)}(k)}.$$

The next lemma reduces matters further to bounding the unweighted exponential sums

$$S(\gamma, \mathcal{I}) := \sum_{k \in \mathcal{I}} e(\gamma k^\Theta)$$

requiring the bound to be uniform in the size of γ , and the interval \mathcal{I} . Therefore, we introduce

$$\widetilde{S}_{\Lambda_1, \Lambda_2}(u) := \sup_{\gamma \in [\Lambda_1, \Lambda_2]} \sup_{\mathcal{I} \subseteq [e^u, e^{u+1})} |S(\gamma, \mathcal{I})|.$$

With this maximal operator at hand, we have the following.

Lemma 2.3. *Let $q_1 \leq q_2$ and $g = \widehat{f}$ or $|\widehat{f}|$. Then*

$$\begin{aligned} \mathcal{E}_{\mathbf{q},u}^{(B)}(N, g) &= 1_{\{q_1=q_2\}} D_{u,q_1}(N, g) \\ &\quad + O\left(N^{-2+\varepsilon} \sum_{j \in \mathcal{J}_{u,q_1}} e^j \sqrt{N_{q_2}^\theta N_{q_1}^{2-\theta}} \widetilde{S}_{\Lambda_1(j), \Lambda_2(u)}(u)\right), \end{aligned}$$

where $1_{\{P\}}$ is 1 if the property P is satisfied and 0 otherwise,

$$\begin{aligned} D_{u,q}(N, g) &:= \frac{2|c_1|^2}{N^2} \sum_{k \in [e^u, e^{u+1})} g\left(\frac{k}{N}\right) \sum_{r \in \mathcal{R}_q(k)} \frac{k^\Theta}{r^{\Theta+1}}, \\ \Lambda_1(j) &:= C_1 e^{j-u\Theta} N_{q_1}, \quad \Lambda_2(u) := C_2 e^{u(1-\Theta)} N_{q_2}^\theta, \\ \mathcal{J}_{u,q_1} &:= [0, C_3 + u - (1-\theta)\Gamma \log q_1) \end{aligned}$$

for some constants $C_1, C_2, C_3 > 0$.

Proof. For brevity, in this proof, let

$$\gamma(\mathbf{r}) := \beta(r_2^{1-\Theta} - r_1^{1-\Theta}).$$

We also consider only the case $g = \widehat{f}$ since the second case follows by repeating the same arguments. With this notation, we have

$$\mathcal{E}_{\mathbf{q},u}^{(B)}(N, \widehat{f}) = \frac{2|c_1|^2}{N^2} \sum_{k \in [e^u, e^{u+1})} \widehat{f}\left(\frac{k}{N}\right) \sum_{\substack{r_i \in \mathcal{R}_{q_i}(k) \\ i=1,2}} \frac{k^\Theta}{(r_1 r_2)^{(\Theta+1)/2}} e(-\gamma(\mathbf{r})k^\Theta).$$

Thus, the r_i which appear in the overall sum all fall within the ranges

$$\mathcal{R}_{q_i,u} := \left(\frac{\alpha\theta e^u}{N_{q_i+1}^{1-\theta}}, \frac{\alpha\theta e^{u+1}}{N_{q_i}^{1-\theta}} \right), \quad i = 1, 2.$$

Now we interchange the r and k summations. For each choice of r_1 and r_2 , we have that

$$k \in \mathcal{K}_{\mathbf{q}}(\mathbf{r}) := \frac{1}{\alpha\theta} [\max(r_1 N_{q_1}^{1-\theta}, r_2 N_{q_2}^{1-\theta}, \alpha\theta e^u), \min(r_1 N_{q_1+1}^{1-\theta}, r_2 N_{q_2+1}^{1-\theta}, \alpha\theta e^{u+1})].$$

Note that this interval may sometimes be empty. With that,

$$\mathcal{E}_{\mathbf{q},u}^{(B)}(N, \hat{f}) = \frac{2|c_1|^2}{N^2} \sum_{\substack{r_i \in \mathcal{R}_{q_i,u} \\ i=1,2}} (r_1 r_2)^{-(\Theta+1)/2} \sum_{k \in \mathcal{K}_{\mathbf{q}}(\mathbf{r})} \hat{f}\left(\frac{k}{N}\right) k^{\Theta} e^{(-\gamma(\mathbf{r})k^{\Theta})}.$$

Next, we remove the weights via partial summation, see Lemma 1.4. Let $w_k = \hat{f}\left(\frac{k}{N}\right) k^{\Theta}$, we first show

$$|w_k - w_{k+1}| \ll \frac{N^{\varepsilon} e^{\Theta u}}{k} \quad (2.4)$$

for any $k \in \mathcal{K}_{\mathbf{q}}(\mathbf{r})$ with the implied constant being absolute. The mean value theorem implies

$$\begin{aligned} |w_k - w_{k+1}| &\leq \left| \hat{f}\left(\frac{k}{N}\right) \right| |k^{\Theta} - (k+1)^{\Theta}| + \left| \hat{f}\left(\frac{k}{N}\right) - \hat{f}\left(\frac{k+1}{N}\right) \right| (k+1)^{\Theta} \\ &\ll \frac{|w_k|}{k} + \frac{e^{\Theta u}}{N}. \end{aligned}$$

Using that $k \ll N^{1+\varepsilon}$ yields (2.4).

Note that $\mathcal{K}_{\mathbf{q}}(\mathbf{r}) \subset [e^u, e^{u+1})$. Thus, Lemma 1.4 is applicable and, in combination with (2.4), yields

$$\sum_{k \in \mathcal{K}_{\mathbf{q}}(\mathbf{r})} w_k e^{(-\gamma(\mathbf{r})k^{\Theta})} \ll N^{\varepsilon} e^{\Theta u} \max_{K \in \mathcal{K}_{\mathbf{q}}(\mathbf{r})} \left| \sum_{k \in \mathcal{K}_{\mathbf{q}}(\mathbf{r}) \cap [1, K]} e^{(\gamma(\mathbf{r})k^{\Theta})} \right|.$$

Note that each $r_i \in \mathcal{R}_{q_i,u}$ satisfies $r_i \asymp e^u N_{q_i}^{-(1-\theta)}$ for $i = 1, 2$.

Reducing matters to exponential sums requires control of the difference $\gamma(\mathbf{r})$; thus, let $\mathcal{T}(j)$ denote the set of pairs $(r_1, r_2) \in \mathcal{R}_{q_1,u} \times \mathcal{R}_{q_2,u}$ satisfying $e^j \leq |r_2 - r_1| < e^{j+1}$. Recall $q_1 \leq q_2$. From now on, we assume $r_2 < r_1$, since the case $r_2 > r_1$ can be done in exactly the same way and therefore shall not be discussed in detail. Observe also that $q_1 < q_2$ and $r_1 = r_2$ implies that $\mathcal{K}_{\mathbf{q}}(\mathbf{r}) = \emptyset$, i.e., this case gives zero contribution. Assume now $q_1 < q_2$. Then

$$\begin{aligned} \gamma(\mathbf{r}) &= \beta(\Theta - 1) \int_{r_2}^{r_1} \tau^{-\Theta} d\tau \gg (r_1 - r_2) r_1^{-\Theta} \gg (r_1 - r_2) \left(\frac{N_{q_1}^{1-\theta}}{\alpha\theta e^{u+1}} \right)^{\Theta} \\ &\gg (r_1 - r_2) e^{-u\Theta} N_{q_1}. \end{aligned}$$

On the other hand,

$$\begin{aligned}\gamma(\mathbf{r}) &= \beta(r_2^{1-\Theta} - r_1^{1-\Theta}) \ll \max(r_1^{1-\Theta}, r_2^{1-\Theta}) = \min(r_1, r_2)^{1-\Theta} \\ &\ll \min(e^u N_{q_1}^{-(1-\theta)}, e^u N_{q_2}^{-(1-\theta)})^{1-\Theta} = e^{u(1-\Theta)} N_{q_2}^\theta.\end{aligned}$$

Thus, provided $\mathbf{r} \in \mathcal{T}(j)$ we deduce that $\gamma(\mathbf{r}) \in [\Lambda_1(j), \Lambda_2(u)]$.

Moreover, the range of j can be constrained by

$$e^j \asymp r_1 - r_2 \ll r_1 \ll e^u N_{q_1}^{-(1-\theta)}$$

which implies that $j \in \mathcal{J}_{u, q_1}$. Thus, we have the following bound:

$$\begin{aligned}\mathcal{E}_{\mathbf{q}, u}^{(B)}(N, \hat{f}) &\ll N^{-2+\varepsilon} \sum_{j \in \mathcal{J}_{u, q_1}} \left(\frac{e^u}{N_{q_1}^{1-\theta}} \frac{e^u}{N_{q_2}^{1-\theta}} \right)^{-(\Theta+1)/2} e^{\Theta u} \#\mathcal{T}(j) \tilde{S}_{\Lambda_1(j), \Lambda_2(u)}(u) \\ &\ll N^{-2+\varepsilon} \sum_{j \in \mathcal{J}_{u, q_1}} \left(\frac{e^u}{N_{q_1}^{1-\theta}} \frac{e^u}{N_{q_2}^{1-\theta}} \right)^{-(\Theta+1)/2} e^{\Theta u} \frac{e^u}{N_{q_2}^{1-\theta}} e^j \tilde{S}_{\Lambda_1(j), \Lambda_2(u)}(u).\end{aligned}$$

We observe that

$$\left(\frac{e^u}{N_{q_1}^{1-\theta}} \frac{e^u}{N_{q_2}^{1-\theta}} \right)^{-(\Theta+1)/2} e^{\Theta u} \frac{e^u}{N_{q_2}^{1-\theta}} e^j = e^j \sqrt{N_{q_2}^\theta N_{q_1}^{2-\theta}}.$$

Overall, we infer, for $q_1 < q_2$, that

$$\mathcal{E}_{\mathbf{q}, u}^{(B)}(N, \hat{f}) \ll N^{-2+\varepsilon} \sum_{j \in \mathcal{J}_{u, q_1}} e^j \sqrt{N_{q_2}^\theta N_{q_1}^{2-\theta}} \tilde{S}_{\Lambda_1(j), \Lambda_2(u)}(u).$$

If $q_1 = q_2$, we remove the diagonal $r_1 = r_2$, which corresponds to the term $D_{u, q_1}(N, \hat{f})$, and apply exactly the same bound to the off-diagonal. ■

3. Proof of the main theorem

3.1. Exponential sum bounds

Thanks to Lemmas 2.1, 2.2, and 2.3, we will show that Theorem 1.1 can be deduced from the next lemma.

Lemma 3.1. *Assume $q_1 \leq q_2$. Then we have the following bound:*

$$\tilde{S}_{\Lambda_1(j), \Lambda_2(u)}(u) \ll e^{8u/15} N_{q_2}^{11\theta/30} + e^{u-j/2} N_{q_1}^{-1/2}. \quad (3.1)$$

Proof. First, we apply the B-process (Theorem 1.3) in the k variable. Here k is in an interval $[A, B]$ of size e^u , and the phase function $\phi(k) = \gamma k^\Theta$ satisfies $\phi(k) \asymp \gamma e^{u\Theta}$, as well as $\phi'(k) \asymp \gamma e^{u(\Theta-1)}$ and $\phi''(k) \asymp \gamma e^{u(\Theta-2)}$, where $\gamma \in [\Lambda_1(j), \Lambda_2(u)]$. Applying Theorem 1.3 and a trivial estimate gives one bound (which will suffice for $\theta < 1/3$).

In fact, we require slightly more than the B-process to move past $\theta = 1/3$. Thus, we will apply the B-process precisely, and then use partial summation and the A-process to bound the resulting sum. First, set $\vartheta = 1/(1 - \Theta)$ and apply Theorem 1.3 to conclude

$$\sum_{k \in [e^u, e^{u+1})} e(\gamma k^\Theta) = c_3 \sum_{h \in [a, b)} \sqrt{\frac{\gamma^\vartheta}{h^{\vartheta+1}}} e(c_4 \gamma^\vartheta h^{1-\vartheta}) + O(e^{u-j/2} N_{q_1}^{-1/2} + \log(N_{q_2})), \quad (3.2)$$

where $a < b$ are positive integers of size $\gamma e^{u(\Theta-1)}$, c_3 and c_4 are complex and real non-zero constants, respectively. A trivial estimate implies that we can assume $b - a \geq 10$.

By exploiting partial summation, we may apply Lemma 1.4 to the main term in (3.2). Thus, to prove (3.1), it suffices to bound

$$\frac{e^u}{\gamma^{1/2} e^{\Theta u/2}} \sum_{h \in [a, b)} e(c_4 \gamma^\vartheta h^{1-\vartheta}) \quad (3.3)$$

for $a < b$ being of size $\gamma e^{u(\Theta-1)}$ and such that $b - a \geq 10$. To that end, we use Theorem 1.2 for an arbitrary integer l . In that notation, let $F \asymp \gamma e^{u\Theta}$, and $M \asymp \gamma e^{u(\Theta-1)}$. Thus, we conclude that (recall $L = 2^l$)

$$\sum_{h \in [a, b)} e(c_4 \gamma^\vartheta h^{1-\vartheta}) \ll (\gamma e^{u\Theta})^{1/(4L-2)} (\gamma e^{u(\Theta-1)})^{1-(l+2)/(4L-2)} + e^{-u}.$$

Inserting this into (3.3) and using that $e^{j-u\Theta} N_{q_1} \ll \gamma \ll e^{u(1-\Theta)} N_{q_2}^\theta$, we obtain

$$\frac{e^u}{\gamma^{1/2} e^{\Theta u/2}} \sum_{h \in [a, b)} e(c_4 \gamma^\vartheta h^{1-\vartheta}) \ll e^u (e^u N_{q_2}^\theta)^{1/(4L-2)-1/2} (N_{q_2}^\theta)^{1-(l+2)/(4L-2)} + e^{-j/2} N_{q_1}^{-1/2},$$

which, on rearranging and choosing $l = 3$, gives (3.1). ■

3.2. Proof of Theorem 1.1

Recall that, after applying Lemma 2.1, our goal is to show

$$\mathcal{E}^{(B)}(N, \hat{f}) = \sum_{\mathbf{q} \in [1, Q]^2} \sum_{u \leq U} \mathcal{E}_{\mathbf{q}, u}^{(B)}(N, \hat{f}) = f(0) + o(1) \quad \text{and} \quad \mathcal{E}^{(B)}(N, |\hat{f}|) = O(1)$$

when $N \rightarrow \infty$.

In view of (3.1) (a similar calculation shows that the second error term in (3.1) gives a negligible contribution), we have for $q_1 \leq q_2$ that

$$\begin{aligned} N^{\varepsilon-2} \sqrt{N_{q_2}^\theta N_{q_1}^{2-\theta}} \sum_{j \in \mathcal{J}_{u, q_1}} e^j \tilde{S}_{\Lambda_1(j), \Lambda_2(u)}(u) \\ \ll N^{\varepsilon-2} \sqrt{N_{q_2}^\theta N_{q_1}^{2-\theta}} \sum_{j \in \mathcal{J}_{u, q_1}} e^j e^{8u/15} N_{q_2}^{11\theta/30} \ll N^{\varepsilon-2} \sqrt{N_{q_2}^\theta N_{q_1}^\theta} e^u e^{8u/15} N_{q_2}^{11\theta/30} \\ \ll N^{(41\theta-14)/30+\varepsilon}. \end{aligned}$$

The case $q_2 < q_1$ can be treated in the same way by interchanging the roles of q_1 and q_2 . Therefore, taking into account Lemmas 2.2 and 2.3, we obtain that

$$\begin{aligned} \sum_{q \in [1, Q]^2} \sum_{u \leq U} \mathcal{E}_{q, u}^{(B)}(N, \hat{f}) &= \mathcal{D}(N, \hat{f}) + O\left(N^{(41\theta-14)/30+\varepsilon} \sum_{q \in [1, Q]^2} \sum_{u \leq U} 1\right) \\ &= f(0) + o(1) + O(N^{(41\theta-14)/30+2/\Gamma+\varepsilon}), \quad N \rightarrow \infty. \end{aligned}$$

In a similar fashion,

$$\mathcal{E}^{(B)}(N, |\hat{f}|) \ll 1 + N^{(41\theta-14)/30+2/\Gamma+\varepsilon}.$$

Thus, for any $\theta < 14/41$ the theorem follows by choosing Γ large enough and ε small enough.

Acknowledgments. The authors would like to thank Jens Marklof and Zeev Rudnick for comments on a earlier draft and Christoph Aistleitner, Daniel El-Baz and Marc Munsch for discussions and comments. We are especially grateful to the anonymous referees for their thorough reading and many insightful comments.

Funding. AS was supported by the Austrian Science Fund (FWF): project M 3246-N. NT was supported by a Schrödinger Fellowship of the Austrian Science Fund (FWF): project J 4464-N.

References

- [1] Aistleitner, C., El-Baz, D., Munsch, M.: [A pair correlation problem, and counting lattice points with the zeta function](#). *Geom. Funct. Anal.* **31**, 483–512 (2021) Zbl 07405882 MR 4311578
- [2] Aistleitner, C., Larcher, G., Lewko, M.: [Additive energy and the Hausdorff dimension of the exceptional set in metric pair correlation problems \(with an appendix by Jean Bourgain\)](#). *Israel J. Math.* **222**, 463–485 (2017) Zbl 1388.11043 MR 3736514
- [3] Apostol, T. M.: [Introduction to analytic number theory](#). Undergrad. Texts Math., Springer, New York (1976) Zbl 0335.10001 MR 434929
- [4] Berry, M. V., Tabor, M.: [Level clustering in the regular spectrum](#). *Proc. Roy. Soc. Lond. Ser. A* **356**, 375–394 (1977) Zbl 1119.81395
- [5] Bloom, T. F., Walker, A.: [GCD sums and sum-product estimates](#). *Israel J. Math.* **235**, 1–11 (2020) Zbl 1450.11094 MR 4068774
- [6] Boca, F. P., Zaharescu, A.: [Pair correlation of values of rational functions \(mod \$p\$ \)](#). *Duke Math. J.* **105**, 267–307 (2000) Zbl 1017.11037 MR 1793613
- [7] Drmota, M., Tichy, R. F.: [Sequences, discrepancies and applications](#). Lecture Notes in Math. 1651, Springer, Berlin (1997) Zbl 0877.11043 MR 1470456
- [8] El-Baz, D., Marklof, J., Vinogradov, I.: [The two-point correlation function of the fractional parts of \$\sqrt{n}\$ is Poisson](#). *Proc. Amer. Math. Soc.* **143**, 2815–2828 (2015) Zbl 1318.11091 MR 3336607
- [9] Elkies, N. D., McMullen, C. T.: [Gaps in \$\sqrt{n} \bmod 1\$ and ergodic theory](#). *Duke Math. J.* **123**, 95–139 (2004) Zbl 1063.11020 MR 2060024
- [10] Fassina, M., Kim, S., Zaharescu, A.: [The distribution of spacings between the fractional parts of \$n^d \alpha\$](#) . *Int. Math. Res. Not. IMRN* **2022**, 10555–10585 (2022) Zbl 1512.11059 MR 4452436
- [11] Graham, S. W., Kolesnik, G.: [Van der Corput’s method of exponential sums](#). London Math. Soc. Lecture Note Ser. 126, Cambridge University Press, Cambridge (1991) Zbl 0713.11001 MR 1145488

- [12] Heath-Brown, D. R.: [Pair correlation for fractional parts of \$\alpha n^2\$](#) . Math. Proc. Cambridge Philos. Soc. **148**, 385–407 (2010) Zbl [1239.11081](#) MR [2609298](#)
- [13] Iwaniec, H., Kowalski, E.: [Analytic number theory](#). Amer. Math. Soc. Colloq. Publ. 53, American Mathematical Society, Providence, RI (2004) Zbl [1059.11001](#) MR [2061214](#)
- [14] Kuipers, L., Niederreiter, H.: Uniform distribution of sequences. Pure Appl. Math., Wiley-Interscience, New York (1974) Zbl [0281.10001](#) MR [419394](#)
- [15] Kurlberg, P., Rudnick, Z.: [The distribution of spacings between quadratic residues](#). Duke Math. J. **100**, 211–242 (1999) Zbl [0985.11038](#) MR [1722952](#)
- [16] Lutsko, C.: [Long-range correlations of sequences modulo 1](#). J. Number Theory **234**, 333–348 (2022) Zbl [1487.11074](#) MR [4370539](#)
- [17] Lutsko, C., Technau, N.: [Correlations of the fractional parts of \$\alpha n^\theta\$](#) . arXiv:[2112.11524](#) (2021)
- [18] Lutsko, C., Technau, N.: [Full Poissonian local statistics of slowly growing sequences](#). arXiv:[2206.07809](#) (2024)
- [19] Marklof, J.: The Berry–Tabor conjecture. In: European Congress of Mathematics, Vol. II (Barcelona, 2000), Progr. Math. 202, Birkhäuser, Basel, 421–427 (2001) Zbl [1023.81011](#) MR [1905381](#)
- [20] Marklof, J.: [Distribution modulo one and Ratner’s theorem](#). In: Equidistribution in number theory, an introduction, NATO Sci. Ser. II Math. Phys. Chem. 237, Springer, Dordrecht, 217–244 (2007) Zbl [1181.11048](#) MR [2290501](#)
- [21] Marklof, J., Strömbergsson, A.: [Equidistribution of Kronecker sequences along closed horocycles](#). Geom. Funct. Anal. **13**, 1239–1280 (2003) Zbl [1048.37009](#) MR [2033838](#)
- [22] Marklof, J., Strömbergsson, A.: [The three gap theorem and the space of lattices](#). Amer. Math. Monthly **124**, 741–745 (2017) Zbl [1391.11092](#) MR [3706822](#)
- [23] Mukherjee, M., Karner, G.: Irrational numbers of constant type — a new characterization. New York J. Math. **4**, 31–34 (1998) Zbl [0893.11004](#) MR [1609027](#)
- [24] Regavim, S.: [Minimal gaps and additive energy in real-valued sequences](#). Q. J. Math. **74**, 825–866 (2023) Zbl [07795668](#) MR [4642240](#)
- [25] Rudnick, Z.: What is . . . quantum chaos? Notices Amer. Math. Soc. **55**, 32–34 (2008) Zbl [1153.81502](#) MR [2373795](#)
- [26] Rudnick, Z.: [A metric theory of minimal gaps](#). Mathematika **64**, 628–636 (2018) Zbl [1451.11085](#) MR [3810030](#)
- [27] Rudnick, Z., Sarnak, P.: [The pair correlation function of fractional parts of polynomials](#). Comm. Math. Phys. **194**, 61–70 (1998) Zbl [0919.11052](#) MR [1628282](#)
- [28] Rudnick, Z., Sarnak, P., Zaharescu, A.: [The distribution of spacings between the fractional parts of \$n^2\alpha\$](#) . Invent. Math. **145**, 37–57 (2001) Zbl [1006.11041](#) MR [1839285](#)
- [29] Rudnick, Z., Technau, N.: [The metric theory of the pair correlation function of real-valued lacunary sequences](#). Illinois J. Math. **64**, 583–594 (2020) Zbl [1464.11069](#) MR [4164447](#)
- [30] Rudnick, Z., Technau, N.: [The metric theory of the pair correlation function for small non-integer powers](#). J. Lond. Math. Soc. (2) **106**, 2752–2772 (2022) Zbl [07730876](#) MR [4498566](#)
- [31] Rudnick, Z., Zaharescu, A.: [The distribution of spacings between fractional parts of lacunary sequences](#). Forum Math. **14**, 691–712 (2002) Zbl [1168.11317](#) MR [1924774](#)
- [32] Technau, N., Walker, A.: [On the triple correlations of fractional parts of \$n^2\alpha\$](#) . Canad. J. Math. **74**, 1347–1384 (2022) Zbl [1506.11102](#) MR [4504666](#)
- [33] Weyl, H.: [Über die Gleichverteilung von Zahlen mod. Eins](#). Math. Ann. **77**, 313–352 (1916) Zbl [46.0278.06](#) MR [1511862](#)
- [34] Zaharescu, A.: [Small values of \$n^2\alpha \pmod{1}\$](#) . Invent. Math. **121**, 379–388 (1995) Zbl [0827.11040](#) MR [1346212](#)