

Divergence-free framings of three-manifolds via eigenspinors

Francesco Lin

Abstract. Gromov used convex integration to prove that any closed orientable three-manifold equipped with a volume form admits three divergence-free vector fields which are linearly independent at every point. We provide an alternative proof of this using geometric properties of eigenspinors in three dimensions. In fact, our proof shows that for any Riemannian metric, one can find three divergence-free vector fields such that at every point they are orthogonal and have the same non-zero length.

Dedicated to Paolo Lisca in the occasion of his 60th birthday.

The following classical result of Stiefel is fundamental in three-manifold topology.

Theorem 1 ([16]). *Every closed orientable three-manifold Y admits a framing, i.e., three vector fields X_1, X_2 and X_3 which are linearly independent at every point.*

The hardest part of the standard proofs of such result is to establish that the second Stiefel–Whitney class $w_2(TM)$ vanishes; after this, it follows from obstruction theory because $\pi_2(\mathrm{SO}(3)) = 0$ (see [13, Chapter 12]). For alternative ‘bare hands’ proofs, see [1].

It is natural to ask whether, in the presence of an additional geometric structure on Y , the framing can be chosen to be compatible with it. In this direction, we have the following result of Gromov.

Theorem 2 ([6, p. 182]). *Every closed orientable three-manifold Y equipped with a volume form Ω admits a framing X_1, X_2 and X_3 consisting of divergence-free vector fields.*

Recall that the divergence $\mathrm{div}(X)$ of a vector field X (with respect to the volume form Ω) is defined in terms of the Lie derivative by

$$\mathcal{L}_X \Omega = \mathrm{div}(X) \Omega;$$

a vector field is *divergence-free* if its divergence vanishes, or equivalently if its associated flow is volume-preserving.

If one fixes a Riemannian metric g on Y (and considers the volume form $\Omega = d\text{vol}_g$), the following is a very natural question with implications in hyperkähler geometry due to Bryant (see also [5]).

Question 1 ([2, Remark 3]). *Which closed orientable Riemannian three manifolds (Y, g) admit a divergence-free framing X_1, X_2 and X_3 which is orthonormal at every point?*

Our main goal is to show that if one relaxes the condition of orthonormality to orthonormality up to scaling then such a framing can always be found.

Theorem 3. *Every closed orientable three-manifold Y equipped with a Riemannian metric g admits a framing X_1, X_2 and X_3 consisting of divergence-free vector fields so that at every point p in Y , $X_1(p), X_2(p)$ and $X_3(p)$ are orthogonal and have the same length.*

This recovers Gromov's result because any volume form Ω is the volume form of some Riemannian metric. While Gromov's proof is based on h-principles and in particular convex integration techniques (see also [3, Chapter 20] for an exposition), our approach is based on elliptic PDEs (and inspired by Seiberg–Witten theory), in the sense that it uses geometric properties of eigenspinors in dimension three. It is not clear whether the convex integration approach can be adapted to prove Theorem 3; notice that the geometric setup of our result is much more rigid because it involves three differential equations in four (rather than nine) variables.

Preliminaries on spin Dirac operators

We begin by recalling some basic facts in spin geometry; we refer the reader to [14] for a general discussion and [10] for a treatment specific for our three-dimensional needs. We will begin by choosing a spin structure on Y , which exists because TY is trivial (Theorem 1 above). Now $\text{Spin}(3) = \text{SU}(2)$, and the spinor representation is given by the natural vector representation on \mathbb{C}^2 . We denote the associated (rank 2 hermitian) spinor bundle by $S \rightarrow Y$; this is equipped with the spin connection ∇ . The associated Clifford multiplication provides an identification

$$\rho: TY \rightarrow \mathfrak{su}(S)$$

such that for each oriented orthonormal frame e_1, e_2, e_3 at a point p , we can find a basis of S_p such that $\rho(e_i) = \sigma_i$ where

$$\sigma_1 = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}, \quad \sigma_2 = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}, \quad (1)$$

are the Pauli matrices. The spin Dirac operator

$$D: \Gamma(S) \rightarrow \Gamma(S)$$

is given by the composition

$$\Gamma(S) \xrightarrow{\nabla} \Gamma(T^*Y \otimes S) \xrightarrow{\rho} \Gamma(S),$$

where we extended ρ to 1-forms via the musical isomorphism. The spin Dirac operator is a first-order elliptic formally self-adjoint operator, and therefore (given that Y is closed) diagonalizable in L^2 with real discrete spectrum infinite in both directions. We will be particularly interested in its eigenspinors, i.e., non-zero solutions to the eigenvalue equation

$$D\Psi = \lambda\Psi, \quad (2)$$

especially in the situation of $\lambda \neq 0$.

The quadratic map

Given any section $\Psi \in \Gamma(S)$ we can consider the traceless hermitian endomorphism $(\Psi\Psi^*)_0 \in \Gamma(i\mathfrak{su}(S))$. Choosing coordinates so that Clifford multiplication is given by (1), if $\Psi = (\alpha, \beta)$, then

$$(\Psi\Psi^*)_0 = \begin{bmatrix} \frac{1}{2}(|\alpha|^2 - |\beta|^2) & \alpha\bar{\beta} \\ \bar{\alpha}\beta & \frac{1}{2}(|\beta|^2 - |\alpha|^2) \end{bmatrix}.$$

The key computation for our purposes is the following.

Lemma 4. *If Ψ is an eigenspinor, then the vector field $X := \rho^{-1}(i(\Psi\Psi^*)_0)$ is divergence-free.*

Remark 1. While our inspiration for considering the quantity X and the Lemma comes from Seiberg–Witten theory, both appear very classically in physics when considering the conserved current for the Dirac eigenvalue equation in three-dimensions, cf. [15, Chapter 8]. Furthermore, the result also readily follows from the (more general) computations in the proof that the Dirac operator is formally-self-adjoint, see [14, Proposition 3.11].

Proof. We will check that the statement holds at any fixed p in Y . Fix a local orthonormal frame e_i , which we assume to be synchronous at p , i.e., $\nabla_{e_i}e_j(p) = 0$. Using (1) we see that

$$X = \rho^{-1}(i(\Psi\Psi^*)_0) = \frac{1}{2}(|\alpha|^2 - |\beta|^2)e_1 + \text{Im}(\alpha\bar{\beta})e_2 + \text{Re}(\alpha\bar{\beta})e_3. \quad (3)$$

Because the framing is synchronous at p , in these coordinates the Dirac operator at p is

$$D\Psi = \sum_{i=1}^3 \sigma_i \cdot \nabla_i \Psi,$$

where ∇_i is the standard derivative in the direction e_i , so that the eigenvalue equation (2) is written explicitly in terms of $\Psi = (\alpha, \beta)$ at the point p as

$$i\nabla_1\alpha - \nabla_2\beta + i\nabla_3\beta = \lambda\alpha, \quad \nabla_2\alpha + i\nabla_3\alpha - i\nabla_1\beta = \lambda\beta.$$

Because the frame is synchronous at p , we also have that for a vector field $Y = \sum_{i=1}^3 Y_i e_i$ the divergence at p is given by

$$\text{div}(Y) = \sum_{i=1}^3 \nabla_i Y_i.$$

Applying this to the vector field (3), we compute directly that the divergence of X at p is

$$\begin{aligned}
& \operatorname{Re}(\bar{\alpha}(\nabla_1 \alpha)) - \operatorname{Re}(\bar{\beta}(\nabla_1 \beta)) + \operatorname{Im}((\nabla_2 \alpha)\bar{\beta}) + \operatorname{Im}(\alpha(\nabla_2 \bar{\beta})) \\
& \quad + \operatorname{Re}((\nabla_3 \alpha)\bar{\beta}) + \operatorname{Re}(\alpha(\nabla_3 \bar{\beta})) \\
& = \operatorname{Re}((\nabla_1 \alpha)\bar{\alpha}) - \operatorname{Re}((\nabla_1 \beta)\bar{\beta}) - \operatorname{Re}((i \nabla_2 \alpha)\bar{\beta}) + \operatorname{Re}((i \nabla_2 \beta)\bar{\alpha}) \\
& \quad + \operatorname{Re}((\nabla_3 \alpha)\bar{\beta}) + \operatorname{Re}((\nabla_3 \beta)\bar{\alpha}) \\
& = \operatorname{Re}((-\nabla_1 \beta - i \nabla_2 \alpha + \nabla_3 \alpha)\bar{\beta}) + \operatorname{Re}((\nabla_1 \alpha + i \nabla_2 \beta + \nabla_3 \beta)\bar{\alpha}) \\
& = \operatorname{Re}((-i \lambda \beta)\bar{\beta}) + \operatorname{Re}((-i \lambda \alpha)\bar{\alpha}) = 0,
\end{aligned}$$

where in the first equality we used $\operatorname{Im}(z) = -\operatorname{Re}(iz)$, in the third equality we used the eigenvalue equations above, and in the last equality we used that λ is real. ■

Remark 2. The result is still true if we consider more generally spin^c Dirac operators D_B (as it is customary in Seiberg–Witten theory). Indeed, we performed the computation pointwise, and any spin^c connection B can be made into the spin connection at a point via a gauge transformation. Furthermore, we can also allow λ to be any real valued function on Y .

The quaternionic structure

A fundamental feature of the spin Dirac operator in three dimensions is its additional quaternionic structure (see for example [11, Chapter I.4] for more details). Namely, we can identify the spinor representation as

$$\begin{aligned}
\mathbb{C}^2 &\equiv \mathbb{H} \\
(v, w) &\mapsto v + jw
\end{aligned}$$

and consider the right action of \mathbb{H} by multiplication; in particular, complex scalars act as usual while the action of j under identification is given by

$$(v, w) \cdot j = (-\bar{w}, \bar{v}).$$

This induces a complex antilinear map squaring to -1 on the spinor bundle $S \rightarrow Y$ (i.e., a quaternionic structure) which we still denote by j . The spin Dirac operator D is compatible with this action in the sense that

$$D(\Psi \cdot j) = (D\Psi) \cdot j.$$

In particular, its eigenspaces are naturally equipped with a quaternionic structure (hence are even dimensional as complex vector spaces). In what follows, we will say that an eigenvalue D is *simple* if the corresponding eigenspace is one dimensional over \mathbb{H} .

Geometry of eigenspinors

With this in mind, we will now state the two main results [4, 7] about the geometry of eigenspinors on three-manifolds that will be fundamental for our purposes: informally

speaking, for a generic metric the spin Dirac operator has no kernel and only simple eigenvalues; furthermore all eigenspinors are nowhere vanishing. Intuitively speaking, the latter should be expected as the spinor bundle $S \rightarrow Y$ has real rank 4. Of course, the proof of such results is quite technical in nature as the Dirac operator depends on the metric in a complicated way. Furthermore, we will need the following more refined version for our purposes.

Theorem 5 ([4, 7]). *Consider a closed three-manifold Y equipped with a Riemannian metric g and a spin structure. Then for a generic metric g' conformal to g , all non-zero eigenvalues of the spin Dirac operator are simple, and all eigenspinors corresponding to non-zero eigenvalues are nowhere vanishing.*

It is important in the statement to focus on *non-zero* eigenvalues, because the kernel of D (i.e., the space of harmonic spinors) is conformally invariant [8]. On the other hand, for a generic metric (not necessarily conformal to a given one) the kernel is trivial [12]. Notice that while the main statements of [4, 7] concern the space of all metrics, the proof is based on a careful analysis of a given conformal class; in particular the result we stated consists of [4, Remark 1.3] and [7, Theorem 4.3]. Finally, for our purposes we will only need the statement that non-harmonic eigenspinors have no zeroes, but we emphasized the role of simple eigenvalues as it is an assumption in its proof.

Proof of the main result

Fix a spin structure and choose a metric g' conformal to g such that the conclusion of Theorem 5 holds, and consider an eigenspinor Ψ' corresponding to an eigenvalue $\lambda' \neq 0$. Using the quaternionic structure, we consider the three λ' -eigenspinors

$$\Psi'_1 = \Psi', \quad \Psi'_2 = \Psi' \cdot \frac{1+k}{\sqrt{2}}, \quad \Psi'_3 = \Psi' \cdot \frac{1+j}{\sqrt{2}}$$

all of which are nowhere vanishing (here $k = ij \in \mathbb{H}$). By Lemma 4 the quadratic map associates to them nowhere-vanishing vector fields X'_1, X'_2, X'_3 which are divergence-free (with respect to $d\text{vol}_{g'}$). Furthermore, they are readily checked to be orthogonal and to have the same length with respect to g' at every point. Indeed, we can identify $S'_p \equiv \mathbb{C}^2 \equiv \mathbb{H}$ by setting

$$\Psi' \equiv (a, 0) \quad \text{and} \quad \Psi' \cdot j \equiv (0, a) \quad \text{where} \quad a = |\Psi'(p)| \in \mathbb{R}^{>0}.$$

This determines a g' -orthonormal basis of $T_p Y$ (denoted by $\{e'_i\}$) via the identification (1). Then we have that at the point we can identify the three spinors as

$$\Psi'_1 = (a, 0), \quad \Psi'_2 = \left(\frac{a}{\sqrt{2}}, \frac{-ia}{\sqrt{2}} \right), \quad \Psi'_3 = \left(\frac{a}{\sqrt{2}}, \frac{a}{\sqrt{2}} \right),$$

which correspond via the quadratic map to the vectors

$$X'_1 = \frac{a^2}{2} e'_1, \quad X'_2 = \frac{a^2}{2} e'_2, \quad X'_3 = \frac{a^2}{2} e'_3$$

respectively. Finally, we can write $g = f^2 g'$ for some positive function f , and the vector fields

$$X_i := \frac{1}{f^3} X'_i$$

are divergence-free with respect to $d \operatorname{vol}_g = f^3 d \operatorname{vol}_{g'}$ because by Cartan's formula

$$\mathcal{L}_X \Omega = d(\iota_X \Omega) + \iota_X d\Omega = d(\iota_X \Omega)$$

the vector field X is divergence-free with respect to Ω if and only if the 2-form $\iota_X \Omega$ is closed. ■

Remark 3. Notice that the proofs of Theorems 2 and 3 both take as input Theorem 1. Indeed, a key ingredient in our proof is the existence of a spin structure, which is equivalent to $w_2(TY) = 0$. On the other hand, Gromov's approach shows that any framing of Y is homotopic (through framings) to a framing by divergence-free vector fields. It is an interesting question to understand which homotopy classes of framings admit representatives as in Theorem 3. Referring to [9] for details, given a framing all other ones are classified up to homotopy by the set of homotopy classes $[Y, \operatorname{SO}(3)]$. To a homotopy class one can associate an element

$$\operatorname{Hom}(\pi_1(Y), \pi_1(\operatorname{SO}(3))) = H^1(Y; \mathbb{Z}/2)$$

which corresponds to the underlying spin structure. Our proof shows that any spin structure admits a framing as in Theorem 3. On the other hand, the homotopy classes inducing the same spin structure form an affine space over

$$H^3(Y; \pi_3(\operatorname{SO}(3))) = \mathbb{Z},$$

and it is not clear from our approach whether all of them can be realized. More in general, it is an interesting question to understand the topological features of eigenspinors on three-manifolds for generic metrics.

Acknowledgments. The author thanks Robert Bryant and Yakov Eliashberg for some useful comments.

Funding. This work was partially supported by NSF grant DMS-2203498.

References

- [1] R. Benedetti and P. Lisca, [Framing 3-manifolds with bare hands](#). *Enseign. Math.* **64** (2018), no. 3-4, 395–413 Zbl [1426.57044](#) MR [3987151](#)
- [2] R. L. Bryant, [Non-embedding and non-extension results in special holonomy](#). In *The many facets of geometry*, pp. 346–367, Oxford University Press, Oxford, 2010 Zbl [1221.53084](#) MR [2681703](#)

- [3] K. Cieliebak, Y. Eliashberg, and N. Mishachev, *Introduction to the h-principle*. 2nd edn., Grad. Stud. Math. 239, American Mathematical Society, Providence, RI, 2024 Zbl [1531.58008](#) MR [4677522](#)
- [4] M. Dahl, *Dirac eigenvalues for generic metrics on three-manifolds*. *Ann. Global Anal. Geom.* **24** (2003), no. 1, 95–100 Zbl [1035.53065](#) MR [1990087](#)
- [5] J. Fine, J. D. Lotay, and M. Singer, *The space of hyperkähler metrics on a 4-manifold with boundary*. *Forum Math. Sigma* **5** (2017), article no. e6 Zbl [1375.53061](#) MR [3631268](#)
- [6] M. Gromov, *Partial differential relations*. *Ergeb. Math. Grenzgeb.* (3) 9, Springer, Berlin, 1986 Zbl [0651.53001](#) MR [0864505](#)
- [7] A. Hermann, *Zero sets of eigenspinors for generic metrics*. *Comm. Anal. Geom.* **22** (2014), no. 2, 177–218 Zbl [1300.53051](#) MR [3210753](#)
- [8] N. Hitchin, *Harmonic spinors*. *Advances in Math.* **14** (1974), 1–55 Zbl [0284.58016](#) MR [0358873](#)
- [9] R. Kirby and P. Melvin, Canonical framings for 3-manifolds. In *Proceedings of 6th Gökova Geometry-Topology Conference*. *Turkish J. Math.* **23** (1999), no. 1, 89–115 Zbl [0947.57020](#) MR [1701641](#)
- [10] P. Kronheimer and T. Mrowka, *Monopoles and three-manifolds*. New Math. Monogr. 10, Cambridge University Press, Cambridge, 2007 Zbl [1158.57002](#) MR [2388043](#)
- [11] H. B. Lawson, Jr. and M.-L. Michelsohn, *Spin geometry*. Princeton Math. Ser. 38, Princeton University Press, Princeton, NJ, 1989 Zbl [0688.57001](#) MR [1031992](#)
- [12] S. Maier, *Generic metrics and connections on Spin- and Spin^c-manifolds*. *Comm. Math. Phys.* **188** (1997), no. 2, 407–437 Zbl [0899.53036](#) MR [1471821](#)
- [13] J. W. Milnor and J. D. Stasheff, *Characteristic classes*. Ann. of Math. Stud. 76, Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1974 Zbl [0298.57008](#) MR [0440554](#)
- [14] J. Roe, *Elliptic operators, topology and asymptotic methods*. 2nd edn., Pitman Res. Notes Math. Ser. 395, Longman, Harlow, 1998 Zbl [0919.58060](#) MR [1670907](#)
- [15] J. J. Sakurai and J. Napolitano, *Modern quantum mechanics*. 3rd edn., Cambridge University Press, Cambridge, 2020 Zbl [1444.81001](#)
- [16] E. Stiefel, *Richtungsfelder und Fernparallelismus in n -dimensionalen Mannigfaltigkeiten*. *Comment. Math. Helv.* **8** (1935), no. 1, 305–353 Zbl [62.0662.02](#) MR [1509530](#)

Communicated by Christian Bär

Received 16 July 2024; revised 4 October 2024.

Francesco Lin

Department of Mathematics, Columbia University, 2990 Broadway, New York, NY 10027, USA;
flin@math.columbia.edu