

On the partial regularity of weak solutions for the magnetohydrodynamics system in \mathbb{R}^4

Zhongbao Zuo

Abstract. In this paper, the partial regularity of the weak solutions to the magnetohydrodynamics (MHD) system in \mathbb{R}^4 is studied. In order to tackle the lack of compactness arising in the spatially high-dimensional setting, inspired by Wu [Arch. Rational Mech. Anal. 239 (2021), 1771–1808], we use the defect measures and prove the existence of partially regular weak solutions (satisfying certain local energy inequality) to the 4-dimensional MHD system. As an application, we obtain that the 2-dimensional Hausdorff dimension of singular sets of these weak solutions is finite.

1. Introduction

In this paper, we study partial regularity results for the 4D incompressible MHD equations, which are given by the following system:

$$\begin{cases} \partial_t u - \Delta u + u \cdot \nabla u - B \cdot \nabla B + \nabla \Pi = 0, \\ \partial_t B - \Delta B + u \cdot \nabla B - B \cdot \nabla u = 0, \\ \operatorname{div} u = \operatorname{div} B = 0. \end{cases} \quad \text{in } \mathbb{R}^4 \times [0, \infty), \quad (1.1)$$

Here, the unknown variable u denotes fluid velocity field, B represents the magnetic field, and $\Pi = \pi + \frac{1}{2}B^2$ signifies the total pressure, respectively. The MHD system (1.1) plays a crucial role in the dynamics of electrically conducting fluids, such as plasmas (see [2], for example). Because of the nonlinear interaction between the fluid velocity field and the magnetic field, the MHD system can accommodate much richer phenomena than the Navier–Stokes system. One important example is that the magnetic field can actually stabilize the fluid motion [1]. There have been extensive studies on various topics concerning the MHD system (see, e.g., [4, 6, 9, 12–14, 19, 28] and references therein). The global weak solutions and the local strong solutions to the 3D MHD system were constructed by Duvaut and Lions [9] and Sermange and Teman [24]. Meanwhile, as in the classical incompressible Navier–Stokes equations (see [3, 21, 22, 26, 27]), partial regularity of suitable weak solutions to the 3D MHD system was investigated by He and Xin in [13]. Wang and Zhang [28] removed the magnetic field hypothesis of the regularity criteria

for the suitable weak solutions to the 3D MHD system. The results obtained in [13, 28] indicate that the velocity field plays a more dominant role than the magnetic field does on the regularity of solutions to the MHD system. Recently, by introducing the notion of dissipative solutions, Chamorro and He [5] weakened the hypothesis on the pressure and obtained a generalization of C-K-N type theorem for weak solutions of MHD system (1.1).

The partial regularity of weak solutions satisfying the local energy inequality was originated from Scheffer [21–23] for the Navier–Stokes equations. And the author studied the Hausdorff measure of potential singular points set of the solutions to the 3D and 4D Navier–Stokes equations. Later, Caffarelli, Kohn, and Nirenberg [3] proved that the one-dimensional Hausdorff measure of potential singular set of suitable weak solutions to the Navier–Stokes equations is zero. Since then, there have been extensive studies on the partial regularity of solutions to the Navier–Stokes equations, MHD equations, and other hydrodynamic models; see [15–17, 27], for example. For the high-dimensional Navier–Stokes equations, Dong and Du [7] proved that the 2-dimensional Hausdorff measure of the set of singular points at the first blow-up time is zero. As in [3] for the dimensional analysis of the Navier–Stokes equations, time is equivalent to two space dimensions. Later, Dong and Strain [8] proved the partial regularity theory of suitable weak solutions of the 6D stationary Navier–Stokes equations. Wang and Wu [29] gave a unified proof on the results of [3, 7, 8]. Recently, by introducing the defect measure, Wu [30, 31] constructed partially regular weak solutions which satisfy local energy inequalities to the non-stationary incompressible Navier–Stokes equations in \mathbb{R}^4 and stationary Navier–Stokes equations in \mathbb{R}^6 , whose singular sets have a local finite 2-dimensional parabolic Hausdorff measure. In context of MHD equations in high-dimensional space, Han and He [11] studied the 4-dimensional non-stationary MHD equations (1.1) interior partial regularity. Recently, Gu [10] and Liu–Wang [18] also obtained some boundary regularity criteria for the 4D non-stationary MHD equations and 6D stationary MHD equations. Motivated by [3, 30] for Navier–Stokes equations, we obtained the existence of partially regular weak solutions for the 4D MHD equations and proved that the 2-dimensional parabolic Hausdorff measure of singular set is finite. Before presenting our main result, we first give the definition of parabolic Hausdorff dimension.

Definition 1.1. Given a set $D \subset \mathbb{R}^4 \times \mathbb{R}$, for a fixed positive real number s , s -dimensional parabolic Hausdorff measure is defined as

$$\mathcal{H}^s(D) = \lim_{\delta \rightarrow 0^+} \mathcal{H}_\delta^s(D),$$

where

$$\mathcal{H}_\delta^s(D) = \inf \left\{ \sum_{i=1}^{\infty} r_i^s \mid D \subset \bigcup_{i \in \mathbb{N}^+} Q_{r_i}(z_0), 0 < r_i < \delta, z_0 = (x_0, t_0) \in \mathbb{R}^4 \times \mathbb{R} \right\}.$$

Here, $Q_r(z_0)$ is centered parabolic cylinder defined by

$$Q_r(z_0) := B_r(x_0) \times \left(t_0 - \frac{r^2}{2}, t_0 + \frac{r^2}{2} \right).$$

The main result of the present paper is the following statement.

Theorem 1.1. *For a fixed time $T > 0$, there exists a weak solution set $(u, B, \Pi, \lambda, \omega)$ for the non-stationary MHD system (1.1) in $\mathbb{R}^4 \times [0, T]$, which satisfies the local energy inequalities (2.11) and (2.12). Moreover, the 2-dimensional Hausdorff dimension of singular set S of weak solutions to (1.1) is finite.*

2. Existence of weak solutions set and local energy inequalities

In order to construct generalized weak solutions to the non-stationary MHD system (1.1), we consider the following regularized MHD system:

$$\begin{cases} \partial_t u_k - \Delta u_k + [(\chi_k * u_k) \cdot \nabla] u_k - [(\chi_k * B_k) \cdot \nabla] B_k + \nabla \Pi = 0, \\ \operatorname{div} u_k = \operatorname{div} B_k = 0, \\ \partial_t B_k - \Delta B_k + [(\chi_k * u_k) \cdot \nabla] B_k - [(\chi_k * B_k) \cdot \nabla] u_k = 0, \\ u_k(\cdot, 0) = u_0, \quad B_k(\cdot, 0) = B_0. \end{cases} \quad (2.1)$$

Here, $\{\chi_k\}_{k \in \mathbb{N}} \subset C_c^\infty(\mathbb{R}^4)$ denotes the standard mollifiers.

Lemma 2.1. *For the regularized non-stationary MHD system (2.1), we have a sequence $\{(u_k, B_k, \Pi_k)\}_{k \in \mathbb{N}} \subset L_t^\infty L_x^2 \cap L_t^2 H_x^1(\mathbb{R}^4 \times [0, T]) \times L_t^\infty L_x^2 \cap L_t^2 H_x^1(\mathbb{R}^4 \times [0, T]) \times L^{\frac{3}{2}}(\mathbb{R}^4 \times [0, T])$ such that (u_k, B_k, Π_k) is a weak solution to the system (2.1). Moreover,*

- (1) $\{u_k\}_{k \in \mathbb{N}}$ and $\{B_k\}_{k \in \mathbb{N}}$ are uniformly bounded in $L_t^\infty L_x^2 \cap L_t^2 H_x^1(\mathbb{R}^4 \times [0, T])$;
- (2) $\{\Pi_k\}_{k \in \mathbb{N}}$ is uniformly bounded in $L^{\frac{3}{2}}(\mathbb{R}^4 \times [0, T])$;
- (3) $\{\partial_t u_k\}_{k \in \mathbb{N}}$ and $\{\partial_t B_k\}_{k \in \mathbb{N}}$ are uniformly bounded in $L_t^1 H_{x, \text{loc}}^{-1}(\mathbb{R}^4 \times [0, T])$.

Therefore, there exists a triple (u, B, Π) such that

$$\begin{aligned} u_k &\rightharpoonup u, \quad B_k \rightharpoonup B && \text{weakly in } L_t^2 H_x^1(\mathbb{R}^4 \times [0, T]), \\ u_k &\rightharpoonup u, \quad B_k \rightharpoonup B && \text{weakly-}^* \text{ in } L_t^\infty L_x^2(\mathbb{R}^4 \times [0, T]), \\ \Pi_k &\rightharpoonup \Pi && \text{weakly in } L^{\frac{3}{2}}(\mathbb{R}^4 \times [0, T]). \end{aligned} \quad (2.2)$$

Additionally, for any bounded smooth function ϕ with bounded derivatives, this sequence satisfies the following local energy inequality:

$$\begin{aligned} &\int_{\mathbb{R}^4} (|u_k(t)|^2 + |B_k(t)|^2) \phi(t) dx + \int_0^t \int_{\mathbb{R}^4} (|\nabla u_k|^2 + |\nabla B_k|^2) \phi dx ds \\ &\leq \int_0^t \int_{\mathbb{R}^4} (|u_k|^2 + |B_k|^2) |\partial_t \phi + \Delta \phi| dx ds + \int_0^t \int_{\mathbb{R}^4} (|u_k|^2 + |B_k|^2) (\tilde{u}_k \cdot \nabla) \phi dx ds \\ &\quad + 2 \int_0^t \int_{\mathbb{R}^4} \Pi_k (u_k \cdot \nabla) \phi dx ds - \int_0^t \int_{\mathbb{R}^4} (\tilde{B}_k \cdot \nabla \phi) (u_k \cdot B_k) dx ds \\ &\quad + \int_{\mathbb{R}^4} (|u_0|^2 + |B_0|^2) \phi(0) dx, \end{aligned} \quad (2.3)$$

where $\tilde{u}_k := \chi_k * u_k$, $\tilde{B}_k := \chi_k * B_k$.

Proof. The existence of weak solutions u and B can be obtained by a standard Galerkin method (see [20, Theorems 4.4 and 14.1], for example). The existence of Π can be proved by Calderon–Zygmund theory. It is noteworthy that $u_k\phi$ and $B_k\phi$ are admissible test functions in the regularized system (2.1). Multiplying equation (2.1)₁ by $u_k\phi$, integrating the resulting equation over \mathbb{R}^4 , and using integration by parts and incompressible conditions, we get that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^4} |u_k|^2 \phi dx + \int_{\mathbb{R}^4} |\nabla u_k|^2 \phi dx \\ &= \frac{1}{2} \int_{\mathbb{R}^4} |u_k|^2 (\partial_t + \Delta \phi) dx + \frac{1}{2} \int_{\mathbb{R}^4} |u_k|^2 \tilde{u}_k \cdot \nabla \phi dx + \int_{\mathbb{R}^4} \Pi u_k \cdot \nabla \phi dx \\ & \quad - \int_{\mathbb{R}^4} (\tilde{B}_k \cdot \nabla) u_k \cdot B_k \phi + (\tilde{B}_k \cdot \nabla \phi) (u_k \cdot B_k) dx. \end{aligned} \quad (2.4)$$

Next, multiplying equation (2.1)₃ by $B_k\phi$, integrating the resulting equation over \mathbb{R}^4 , and applying integration by parts and incompressible conditions again, we obtain that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^4} |B_k|^2 \phi dx + \int_{\mathbb{R}^4} |\nabla B_k|^2 \phi dx \\ &= \frac{1}{2} \int_{\mathbb{R}^4} |B_k|^2 (\partial_t + \Delta \phi) dx + \frac{1}{2} \int_{\mathbb{R}^4} |B_k|^2 \tilde{u}_k \cdot \nabla \phi dx + \int_{\mathbb{R}^4} (\tilde{B}_k \cdot \nabla) u_k \cdot B_k \phi dx. \end{aligned} \quad (2.5)$$

Hence, by adding (2.4) and (2.5) and after integrating over $[0, t]$, we yield the local energy inequality (2.3). For the uniform boundedness of $\{\partial_t u_k\}_{k \in \mathbb{N}}$ and $\{\partial_t B_k\}_{k \in \mathbb{N}}$, applying duality theory, we know that, for almost every $t \in [0, T]$, the weak formulations (2.1) for u_k and B_k are equivalent to

$$\begin{aligned} & \langle \partial_t u_k, \xi \rangle_{H_x^{-1} \times H_x^1} \\ &= - \int_{\mathbb{R}^4} \nabla u_k \nabla \xi + (\tilde{u}_k \cdot \nabla) u_k \xi - (\tilde{B}_k \cdot \nabla) B_k \xi dx, \quad \forall \xi \in H_x^1 \text{ with } \operatorname{div} \xi = 0 \end{aligned}$$

and

$$\langle \partial_t B_k, \eta \rangle_{H_x^{-1} \times H_x^1} = - \int_{\mathbb{R}^4} \nabla B_k \nabla \eta + (\tilde{u}_k \cdot \nabla) B_k \eta - (\tilde{B}_k \cdot \nabla) u_k \eta dx, \quad \forall \eta \in H_x^1.$$

For every $\xi \in C_c^\infty(\Omega)$ with $\Omega \subset \subset \mathbb{R}^4$, the Hölder inequality and the Sobolev embedding inequality imply that

$$\begin{aligned} & \left| \int_{\mathbb{R}^4} \nabla u_k \nabla \xi + (\tilde{u}_k \cdot \nabla) u_k \xi - (\tilde{B}_k \cdot \nabla) B_k \xi dx \right| \\ & \leq \|\nabla u_k\|_{L_x^2} \|\nabla \xi\|_{L_x^2} + (\|\nabla u_k\|_{L_x^2} \|u_k\|_{L_x^4} + \|\nabla B_k\|_{L_x^2} \|B_k\|_{L_x^4}) \|\xi\|_{L_x^4} \\ & \leq C(\|u_k\|_{H_x^1}^2 + \|u_k\|_{H_x^1} + \|B_k\|_{H_x^1}^2 + \|B_k\|_{H_x^1}) \|\xi\|_{H_x^1}. \end{aligned}$$

Integrating in time yields that $\{\partial_k u_k\}_{k \in \mathbb{N}}$ are uniformly bounded in $L_t^1 H_{x, \text{loc}}^{-1}(\mathbb{R}^4 \times [0, T])$. Applying the same procedure, we can obtain the uniform boundedness of $\{\partial_t B_k\}_{k \in \mathbb{N}}$ in $L_t^1 H_{x, \text{loc}}^{-1}(\mathbb{R}^4 \times [0, T])$. ■

Next, we prove that certain measures in the limit are relatively compact. We recall that a collection of measures $\{\mu_i\}_{i \in \Lambda}$ on \mathbb{R}^N is called tight if, for any $\varepsilon > 0$, there exists a compact set $\Omega_\varepsilon \subset \mathbb{R}^N$ such that $\mu_i(\mathbb{R}^N \setminus \Omega_\varepsilon) < \varepsilon$ for any $i \in \Lambda$.

Lemma 2.2. *Let the assumptions be as in Lemma 2.1; then, $\{(|\nabla u_k|^2 + |\nabla B_k|^2) dx dt\}_{k \in \mathbb{N}}$, $\{(|u_k|^2 + |B_k|^2) dx dt\}_{k \in \mathbb{N}}$, and $\{(|u_k|^3 + |B_k|^3) dx dt\}_{k \in \mathbb{N}}$ are tight in the sense of measures.*

Proof. For fixed $r > 0$, we consider a smooth cutoff function $\xi \in C_c^\infty(\mathbb{R}^4)$ such that $0 \leq \xi \leq 1$, $\xi|_{B_r} = 0$, $\xi|_{\mathbb{R}^4 \setminus B_{2r}} = 1$, satisfying property $|\nabla \xi|^2 + |\nabla^2 \xi| \leq C r^{-2}$. Testing the regularity non-stationary MHD system (2.1)₁ with $u_k \xi$ and (2.1)₃ with $B_k \xi$, respectively, we get

$$\begin{aligned} & \sup_t \int_{\mathbb{R}^4} (|u_k(t)|^2 + |B_k(t)|^2) \xi dx - \int_{\mathbb{R}^4} (|u_0|^2 + |B_0|^2) \xi dx \\ & \quad + \int_0^T \int_{\mathbb{R}^4} (|\nabla u_k|^2 + |\nabla B_k|^2) \xi dx dt \\ & \leq \int_0^T \int_{\mathbb{R}^4} (|u_k|^2 + |B_k|^2) |\Delta \xi| dx dt + \int_0^T \int_{\mathbb{R}^4} (|u_k|^2 + |B_k|^2) |\tilde{u}_k| |\nabla \xi| dx dt \\ & \quad + 2 \int_0^T \int_{\mathbb{R}^4} |\Pi_k u_k| |\nabla \xi| dx dt + \int_0^T \int_{\mathbb{R}^4} |\tilde{B}_k| |\nabla \xi| |u_k| |B_k| dx dt. \end{aligned}$$

According to the properties of test function and the Hölder inequality, we have

$$\begin{aligned} & \sup_t \int_{\mathbb{R}^4 \setminus B_{2r}} (|u_k(t)|^2 + |B_k(t)|^2) dx - \int_{\mathbb{R}^4 \setminus B_r} (|u_0|^2 + |B_0|^2) dx \\ & \quad + \int_0^T \int_{\mathbb{R}^4 \setminus B_{2r}} (|\nabla u_k|^2 + |\nabla B_k|^2) dx dt \\ & \leq C r^{-2} \int_0^T \int_{B_{2r} \setminus B_r} (|u_k|^2 + |B_k|^2) dx dt \\ & \quad + C r^{-1} \int_0^T \int_{B_{2r} \setminus B_r} (|u_k|^2 + |B_k|^2) |\tilde{u}_k| dx dt \\ & \quad + C r^{-1} \int_0^T \int_{B_{2r} \setminus B_r} |\Pi_k u_k| dx dt + C r^{-1} \int_0^T \int_{B_{2r} \setminus B_r} |\tilde{B}_k| |u_k| |B_k| dx dt \\ & \leq C r^{-\frac{2}{3}} T^{\frac{1}{3}} + C r^{-1}, \end{aligned}$$

where we have used the fact that $\{u_k\}_{k \in \mathbb{N}}$ and $\{B_k\}_{k \in \mathbb{N}}$ are uniformly bounded in $L^3(\mathbb{R}^3 \times [0, T])$ and $\{\Pi\}_{k \in \mathbb{N}}$ is uniformly bounded in $L^{\frac{3}{2}}(\mathbb{R}^3 \times [0, T])$. Letting r be arbitrarily large

implies the tightness of $\{(|\nabla u_k|^2 + |\nabla B_k|^2)dxdt\}_{k \in \mathbb{N}}$ and $\{(|u_k(t)|^2 + |B_k(t)|^2)dx\}_{k \in \mathbb{N}}$ uniformly in t , which implies the tightness of $\{(|u_k|^2 + |B_k|^2)dxdt\}_{k \in \mathbb{N}}$. On the other hand, applying Sobolev inequality, we have

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^4} (|u_k \xi|^3 + |B_k \xi|^3) dx dt \\
& \leq \|u_k \xi\|_{L_t^\infty L_x^2} \int_0^T \int_{\mathbb{R}^4} |\nabla(u_k \xi)|^2 dx dt + \|B_k \xi\|_{L_t^\infty L_x^2} \int_0^T \int_{\mathbb{R}^4} |\nabla(B_k \xi)|^2 dx dt \\
& \leq C \|u_k \xi\|_{L_t^\infty L_x^2} \left(\int_0^T \int_{\mathbb{R}^4} |\nabla u_k|^2 |\xi|^2 dx dt + \int_0^T \int_{\mathbb{R}^4} |u_k|^2 |\nabla \xi|^2 dx dt \right) \\
& \quad + C \|B_k \xi\|_{L_t^\infty L_x^2} \left(\int_0^T \int_{\mathbb{R}^4} |\nabla B_k|^2 |\xi|^2 dx dt + \int_0^T \int_{\mathbb{R}^4} |B_k|^2 |\nabla \xi|^2 dx dt \right) \\
& \leq C (\|u_k \xi\|_{L_t^\infty L_x^2} + \|B_k \xi\|_{L_t^\infty L_x^2}) \int_0^T \int_{\mathbb{R}^4} (|\nabla u_k|^2 + |\nabla B_k|^2) |\xi|^2 dx dt \\
& \quad + C (\|u_k \xi\|_{L_t^\infty L_x^2} + \|B_k \xi\|_{L_t^\infty L_x^2}) r^{-\frac{2}{3}} T^{\frac{1}{3}} \left(\int_0^T \int_{\mathbb{R}^4} (|u_k|^3 + |B_k|^3) dx dt \right)^{\frac{2}{3}}.
\end{aligned}$$

Combining the tightness of $\{(|\nabla u_k|^2 + |\nabla B_k|^2)dxdt\}_{k \in \mathbb{N}}$ with the uniform boundedness of u_k and B_k in the natural energy space implies the tightness of $\{(|u_k|^3 + |B_k|^3)dxdt\}_{k \in \mathbb{N}}$. \blacksquare

In order to obtain local energy inequalities for the weak limit (u, B, Π) , we will pass to the limit $k \rightarrow \infty$ in the local energy inequalities (2.3).

Lemma 2.3. *Given a bounded sequence $\{u_k\}_{k \in \mathbb{N}}$,*

$$\{B_k\}_{k \in \mathbb{N}} \subset L_t^\infty L_x^2 \cap L_t^2 H_x^1(\mathbb{R}^4 \times [0, T]),$$

let (u, B) be given by the limit in (2.2). Assume that u_k converges to u and B_k converges to B , respectively, in $L_{\text{loc}}^1(\mathbb{R}^4 \times [0, T])$. Suppose that $\mu_k = (|\nabla u_k|^2 + |\nabla B_k|^2)dxdt \rightarrow \mu$, $\nu_k = (|u_k|^3 + |B_k|^3)dxdt \rightarrow \nu$ weakly in the sense of measures, where μ and ν are bounded nonnegative measures on the $\mathbb{R}^4 \times [0, T]$. Then, there exist nonnegative finite measures λ and ω such that, for any $\eta \in C_c^\infty(\mathbb{R}^4 \times [0, T])$,

$$\int_0^T \int_{\mathbb{R}^4} \eta d\mu = \int_0^T \int_{\mathbb{R}^4} \eta (|\nabla u|^2 + |\nabla B|^2) dx dt + \int_0^T \int_{\mathbb{R}^4} \eta d\lambda, \quad (2.6)$$

$$\int_0^T \int_{\mathbb{R}^4} \eta d\nu = \int_0^T \int_{\mathbb{R}^4} \eta (|u|^3 + |B|^3) dx dt + \int_0^T \int_{\mathbb{R}^4} \eta d\omega. \quad (2.7)$$

Moreover, $\omega \ll \lambda$, and for any open subdomain Ω of $\mathbb{R}^4 \times [0, T]$,

$$\int_\Omega d\omega \leq C \liminf_{k \rightarrow \infty} (\|u_k - u\|_{L_t^\infty L_x^2(\Omega)} + \|B_k - B\|_{L_t^\infty L_x^2(\Omega)}) \int_\Omega d\lambda. \quad (2.8)$$

In particular, the Radon–Nikodym derivative satisfies

$$\frac{d\omega}{d\lambda} \leq C \lim_{r \rightarrow 0} \liminf_{k \rightarrow \infty} (\|u_k - u\|_{L_t^\infty L_x^2(Q_r(x_0, t_0))} + \|B_k - B\|_{L_t^\infty L_x^2(Q_r(x_0, t_0))}),$$

where $Q_r(x_0, t_0) := B_r(x_0) \times (t_0 - \frac{r^2}{2}, t_0 + \frac{r^2}{2})$.

Proof. We let $v_k = u_k - u \in L_t^\infty L_x^2 \cap L_t^2 H_x^1$, $A_k = B_k - B \in L_t^\infty L_x^2 \cap L_t^2 H_x^1$. Combining Simon compactness theory (see [25]) with Lemma 2.1 implies that

$$\begin{aligned} v_k &\rightarrow 0, A_k \rightarrow 0 \quad \text{strongly in } L_{t,x}^2, \text{ locally in space,} \\ v_k &\rightarrow 0, A_k \rightarrow 0 \quad \text{weakly in } L_t^2 H_x^1, \\ v_k &\rightarrow 0, A_k \rightarrow 0 \quad \text{weakly-* in } L_t^\infty L_x^2. \end{aligned}$$

Defining $\lambda_k := (|\nabla v_k|^2 + |\nabla A_k|^2)dxdt$, we claim that $\{\lambda_k\}_{k \in \mathbb{N}}$ is tight. Indeed, for any compactness subset $\Omega \subset \mathbb{R}^4$, let $D^c := (\mathbb{R}^4 \setminus \Omega) \times [0, T]$. Then,

$$\begin{aligned} \|\nabla v_k\|_{L^2(D^c)} &\leq \|\nabla u_k\|_{L^2(D^c)} + \|\nabla u\|_{L^2(D^c)}, \\ \|\nabla A_k\|_{L^2(D^c)} &\leq \|\nabla B_k\|_{L^2(D^c)} + \|\nabla B\|_{L^2(D^c)}. \end{aligned}$$

Due to the weak convergence of $\{\mu_k\}_{k \in \mathbb{N}}$, we know that $\{\mu_k\}_{k \in \mathbb{N}}$ is tight; hence, $\|\nabla v_k\|_{L^2(D^c)}$ and $\|\nabla A_k\|_{L^2(D^c)}$ are arbitrarily small given Ω large enough. Thus, we can extract a weakly convergent subsequence with a limit denoted by λ . For any $\eta \in C_c^\infty(\mathbb{R}^4 \times [0, T])$, we have

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^4} \eta d\mu &= \lim_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \eta d\mu_k \\ &= \lim_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \eta (|\nabla u_k|^2 + |\nabla B_k|^2) dxdt \\ &= \int_0^T \int_{\mathbb{R}^4} \eta (|\nabla u|^2 + |\nabla B|^2) dxdt \\ &\quad + \lim_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \eta (|\nabla v_k|^2 + |\nabla A_k|^2) dxdt \\ &= \int_0^T \int_{\mathbb{R}^4} \eta (|\nabla u|^2 + |\nabla B|^2) dxdt + \int_0^T \int_{\mathbb{R}^4} \eta d\lambda, \end{aligned}$$

where we have used the fact that interaction term vanishes since $u_k \rightarrow u$ and $B_k \rightarrow B$ weakly in $L_t^2 H_x^1(\mathbb{R}^4 \times [0, T])$. Let

$$\omega_k := (|v_k|^3 + |A_k|^3)dxdt \rightarrow \omega$$

weakly in the sense of measures. A similar process verifies (2.7). On the other hand, for any $\eta \in C_c^\infty(\mathbb{R}^4 \times [0, T])$, using interpolation inequality and Sobolev embedding inequality,

we have

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^4} |\eta|^3 d\omega \\
&= \lim_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} |\eta|^3 d\omega_k \\
&= \lim_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} |v_k \eta|^3 + |A_k \eta|^3 dx dt \\
&\leq \liminf_{k \rightarrow \infty} \sup_{0 < t < T} (\|v_k \eta\|_{L_x^2} + \|A_k \eta\|_{L_x^2}) \int_0^T \|v_k \eta\|_{L_x^4}^2 + \|A_k \eta\|_{L_x^4}^2 dt \\
&\leq C \liminf_{k \rightarrow \infty} \sup_{0 < t < T} (\|v_k \eta\|_{L_x^2} + \|A_k \eta\|_{L_x^2}) \int_0^T \int_{\mathbb{R}^4} |\nabla(v_k \eta)|^2 + |\nabla(A_k \eta)|^2 dx dt \\
&\leq C \liminf_{k \rightarrow \infty} \sup_{0 < t < T} (\|v_k \eta\|_{L_x^2} + \|A_k \eta\|_{L_x^2}) \int_0^T \int_{\mathbb{R}^4} |\eta|^2 |\nabla v_k|^2 + |\eta|^2 |\nabla A_k|^2 dx dt \\
&\leq C \liminf_{k \rightarrow \infty} \sup_{0 < t < T} (\|v_k \eta\|_{L_x^2} + \|A_k \eta\|_{L_x^2}) \int_0^T \int_{\mathbb{R}^4} |\eta|^2 d\lambda.
\end{aligned}$$

Notice that, in the third inequality, the terms converge to zero when at least one derivative hits η . Using smooth functions to approximate the indicator function Ω yields the inequality (2.8). Hence, ω is absolutely continuous with respect to λ , and by Radon–Nikodym theorem, we have

$$\frac{d\omega}{d\lambda} \in L^1(\mathbb{R}^4 \times [0, T]; \lambda)$$

with, for any $(x_0, t_0) \in \mathbb{R}^4 \times (0, T)$,

$$\frac{d\omega}{d\lambda}(x_0, t_0) \leq C \lim_{r \rightarrow 0} \liminf_{k \rightarrow \infty} (\|v_k\|_{L_t^\infty L_x^2(Q_r(x_0, t_0))} + \|B_k\|_{L_t^\infty L_x^2(Q_r(x_0, t_0))}). \quad \blacksquare$$

Using the parabolic concentration-compactness framework in Lemma 2.3, we can define the weak solution sets involving concentration measures.

Definition 2.1. $(u, B, \Pi, \lambda, \omega)$ is a weak solution set of the MHD system (1.1) if

- (1) u , B , and Π are obtained as weak limits of the weak solutions $\{(u_k, B_k, \Pi_k)\}_{k \in \mathbb{N}}$ of the regularized MHD system (2.1);
- (2) λ and ω are obtained as weak limits of the measures in Lemma 2.3.

In order to obtain local energy inequalities for weak solution sets, we need the following lemma, which shows that the concentration in $|u\Pi|dxdt$ is localizable and comparable to the concentration in $|u|^3dxdt$.

Lemma 2.4. Assume that $\{(u_k, B_k, \Pi_k)\}_{k \in \mathbb{N}}$ are the solutions of the regularized system (2.1) and $(u, B, \Pi, \lambda, \omega)$ is the corresponding weak solution set; then,

$$\limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \xi |u_k(\Pi_k - \gamma) - u(\Pi - \gamma)| dx dt \leq C \int_0^T \int_{\mathbb{R}^4} \xi d\omega$$

for any $\psi \in C_c^\infty(\mathbb{R}^4 \times [0, T])$ and $\gamma \in \mathbb{R}$ with $\psi \geq 0$.

Proof. According to the regularized system (2.1)₁, we see that Π satisfies Poisson equation formally:

$$-\Delta \pi_k = \partial_i \partial_j (\tilde{u}_k^i u_k^j - \tilde{B}_k^i B_k^j).$$

Then, we localize the above equation with an arbitrary Lipschitz function $\psi \in C^{0,1}(\mathbb{R}^4)$, which yields

$$\begin{aligned} -\Delta(\Pi_k \psi) &= \psi \partial_i \partial_j (\tilde{u}_k^i u_k^j - \tilde{B}_k^i B_k^j) - \operatorname{div}(\Pi_k \nabla \psi) - \nabla \Pi_k \cdot \nabla \psi \\ &= \partial_i \partial_j [\psi (\tilde{u}_k^i u_k^j - \tilde{B}_k^i B_k^j)] - \partial_i [(\tilde{u}_k^i u_k^j - \tilde{B}_k^i B_k^j) \partial_j \psi] \\ &\quad - \partial_j (\tilde{u}_k^i u_k^j - \tilde{B}_k^i B_k^j) \partial_i \psi - \operatorname{div}(\Pi_k \nabla \psi) - \nabla \Pi_k \cdot \nabla \psi \\ &= \partial_i \partial_j [\psi (\tilde{u}_k^i u_k^j - \tilde{B}_k^i B_k^j)] \\ &\quad - \partial_i [(\tilde{u}_k^i u_k^j - \tilde{B}_k^i B_k^j) \partial_j \psi + \Pi_k \nabla \psi] \\ &\quad - \partial_j (\tilde{u}_k^i u_k^j - \tilde{B}_k^i B_k^j) \partial_i \psi - \nabla \Pi_k \cdot \nabla \psi \\ &:= -\Delta \Pi_k^1 - \Delta \Pi_k^2 - \Delta \Pi_k^3. \end{aligned} \quad (2.9)$$

Similarly, we decompose $\Pi \psi$ in a similar way. From the Calderon–Zygmund theory, we get

$$\begin{aligned} &\|\Pi_k^1(t) - \Pi^1(t)\|_{L_x^{\frac{3}{2}}} \\ &\leq C \|\psi \tilde{u}_k^i(t) u_k^j(t) - \psi \tilde{u}^i(t) u^j(t)\|_{L_x^{\frac{3}{2}}} + C \|\psi \tilde{B}_k^i(t) B_k^j(t) - \psi \tilde{B}^i(t) B^j(t)\|_{L_x^{\frac{3}{2}}} \\ &\leq C \|\psi (\tilde{u}_k^i(t) - \tilde{u}^i(t)) (u_k^j(t) - u^j(t))\|_{L_x^{\frac{3}{2}}} + C \|\psi \tilde{u}^i(t) (u_k^j(t) - u^j(t))\|_{L_x^{\frac{3}{2}}} \\ &\quad + C \|\psi u^j(t) (\tilde{u}_k^i(t) - \tilde{u}^i(t))\|_{L_x^{\frac{3}{2}}} + \|\psi (\tilde{B}_k^i(t) - \tilde{B}^i(t)) (B_k^j(t) - B^j(t))\|_{L_x^{\frac{3}{2}}} \\ &\quad + C \|\psi \tilde{B}^i(t) (B_k^j(t) - B^j(t))\|_{L_x^{\frac{3}{2}}} + C \|\psi B^j(t) (\tilde{B}_k^i(t) - \tilde{B}^i(t))\|_{L_x^{\frac{3}{2}}} \\ &\leq C \|\psi^{\frac{1}{2}} (u_k^j(t) - u^j(t))\|_{L_x^3}^2 + C \|\psi \tilde{u}^i(t) (u_k^j(t) - u^j(t))\|_{L_x^{\frac{3}{2}}} \\ &\quad + C \|\psi u^j(t) (\tilde{u}_k^i(t) - \tilde{u}^i(t))\|_{L_x^{\frac{3}{2}}} + \|\psi^{\frac{1}{2}} (B_k^j(t) - B^j(t))\|_{L_x^3}^2 \\ &\quad + C \|\psi \tilde{B}^i(t) (B_k^j(t) - B^j(t))\|_{L_x^{\frac{3}{2}}} + C \|\psi B^j(t) (\tilde{B}_k^i(t) - \tilde{B}^i(t))\|_{L_x^{\frac{3}{2}}}. \end{aligned}$$

Since $\omega_k \rightarrow \omega$ weakly, $|u_k^j - u^j|^{\frac{3}{2}}$ and $|B_k^j - B^j|^{\frac{3}{2}}$ are uniformly integrable with respect to $|\psi|^{\frac{3}{2}} |\tilde{u}^i|^{\frac{3}{2}} dx dt$ and $|\psi|^{\frac{3}{2}} |\tilde{B}^i|^{\frac{3}{2}} dx dt$, respectively, we have

$$\begin{aligned} &\limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} |\Pi_k^1 - \Pi^1|^{\frac{3}{2}} dx dt \\ &\leq C \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} |\psi|^{\frac{3}{2}} (|u_k - u|^3 + |B_k - B|^3) dx dt \\ &\quad + C \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} |\psi|^{\frac{3}{2}} (|\tilde{u}^i (u_k^j - u^j)|^{\frac{3}{2}} + |\tilde{B}^i (B_k^j - B^j)|^{\frac{3}{2}}) dx dt \end{aligned}$$

$$\begin{aligned}
& + C \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} |\psi|^{\frac{3}{2}} (|u^j (\tilde{u}_k^i - \tilde{u}^i)|^{\frac{3}{2}} + |B^j (\tilde{B}_k^i - \tilde{B}^i)|^{\frac{3}{2}}) dx dt \\
& \leq C \int_0^T \int_{\mathbb{R}^4} |\psi|^{\frac{3}{2}} d\omega,
\end{aligned}$$

where we have used the Vitali's convergence theorem for the second and third lines:

$$\begin{aligned}
& \|\Pi_k^2 - \Pi^2\|_{L_{t,x}^{\frac{3}{2}}} \\
& = \|(-\Delta)^{-1}[-\operatorname{div}((u_k u_k^j - u u^j) \partial_j \psi - (B_k B_k^j - B B^j) \partial_j \psi + (\Pi_k - \Pi) \nabla \psi)]\|_{L_{t,x}^{\frac{3}{2}}} \\
& \leq C \|(u_k u_k^j - u u^j) \partial_j \psi - (B_k B_k^j - B B^j) \partial_j \psi + (\Pi_k - \Pi) \nabla \psi\|_{L_t^{\frac{3}{2}} L_x^{\frac{12}{11}}} \\
& \leq C \|\nabla \psi\|_{L^\infty} (\|u_k u_k^j - u u^j\|_{L_t^{\frac{3}{2}} L_x^{\frac{12}{11}}} + \|B_k B_k^j - B B^j\|_{L_t^{\frac{3}{2}} L_x^{\frac{12}{11}}} + \|\Pi_k - \Pi\|_{L_t^{\frac{3}{2}} L_x^{\frac{12}{11}}}) \\
& \leq C \|\nabla \psi\|_{L^\infty} (\|u_k - u\|_{L_t^3 L_x^{\frac{24}{11}}} + \|B_k - B\|_{L_t^3 L_x^{\frac{24}{11}}} + \|\Pi_k - \Pi\|_{L_t^{\frac{3}{2}} L_x^{\frac{12}{11}}}).
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \|\Pi_k^3 - \Pi^3\|_{L_{t,x}^{\frac{3}{2}}} \\
& \leq C \|\nabla \psi\|_{L^\infty} (\|u_k - u\|_{L_t^3 L_x^{\frac{24}{11}}} + \|B_k - B\|_{L_t^3 L_x^{\frac{24}{11}}} + \|\Pi_k - \Pi\|_{L_t^{\frac{3}{2}} L_x^{\frac{12}{11}}}).
\end{aligned}$$

Since for any $f \in L_t^\infty L_x^2 \cap L_t^2 H_x^1$, with the help of interpolation inequality and Sobolev inequality, we have

$$\begin{aligned}
\int_0^T \|f(t)\|_{L_x^{\frac{24}{11}}}^3 dt & = \int_0^T \|f(t)\|_{L_x^{\frac{24}{11}}} \cdot \|f(t)\|_{L_x^{\frac{24}{11}}}^2 dt \\
& \leq \|f\|_{L_t^{12} L_x^{\frac{24}{11}}} \|f\|_{L_{t,x}^{\frac{24}{11}}}^2 \leq \|f\|_{L_t^{12} L_x^{\frac{24}{11}}} \|f\|_{L_{t,x}^2}^{\frac{3}{2}} \|f\|_{L_{t,x}^{\frac{1}{2}}}^{\frac{1}{2}} \\
& \leq \|f\|_{L_{t,x}^2}^{\frac{3}{2}} \|f\|_{L_{t,x}^3}^{\frac{1}{2}} \left(\int_0^T \|f\|_{L_x^{10}}^{10} \|f\|_{L_x^4}^2 dt \right)^{\frac{1}{12}} \\
& \leq \|f\|_{L_{t,x}^2}^{\frac{3}{2}} \|f\|_{L_{t,x}^3}^{\frac{1}{2}} \|f\|_{L_t^\infty L_x^2}^{\frac{5}{6}} \|\nabla f\|_{L_{t,x}^2}^{\frac{1}{6}}.
\end{aligned}$$

Hence, taking $f = u_k - u$ or $f = B_k - B$, we get that

$$\limsup_{k \rightarrow \infty} \|u_k - u\|_{L_t^3 L_x^{\frac{24}{11}}} = 0, \quad \limsup_{k \rightarrow \infty} \|B_k - B\|_{L_t^3 L_x^{\frac{24}{11}}} = 0.$$

Combining Calderon–Zygmund theory, this yields

$$\limsup_{k \rightarrow \infty} \|\Pi_k - \Pi\|_{L_t^3 L_x^{\frac{12}{11}}} = 0. \quad (2.10)$$

Collecting (2.9)–(2.10), we know that

$$\limsup_{k \rightarrow \infty} \|(\Pi_k - \Pi) \psi\|_{L_{t,x}^{\frac{3}{2}}}^{\frac{3}{2}} \leq \limsup_{k \rightarrow \infty} \sum_{l=1}^3 \|\Pi_k^l - \Pi^l\|_{L_{t,x}^{\frac{3}{2}}}^{\frac{3}{2}} \leq \int_0^T \int_{\mathbb{R}^4} |\psi|^{\frac{3}{2}} d\omega.$$

Thus, applying Vitali's convergence theorem and taking $\xi = \psi^{\frac{2}{3}}$, we get

$$\begin{aligned}
& \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \xi |u_k(\Pi_k - \gamma) - u(\Pi - \gamma)| dx dt \\
& \leq \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \xi |u_k| |\Pi_k - p| dx dt + \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \xi |u_k - u| |\Pi - \gamma| dx dt \\
& \leq \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \xi |u_k - u| |\Pi_k - p| dx dt + \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \xi |u| |\Pi_k - p| dx dt \\
& \leq C \int_0^T \int_{\mathbb{R}^4} \xi d\omega.
\end{aligned}$$

Hence, we complete the proof of Lemma 2.4. ■

Now, we can prove the following local energy inequalities.

Proposition 2.1. *Let the assumptions be as in Lemma 2.1; then, for any nonnegative cut-off functions $\eta \in C_c^\infty(\mathbb{R}^4 \times [0, T])$ with $\eta(\cdot, 0) = 0$ and $\{\eta_i\}_{1 \leq i \leq n} \subset C_c^\infty(\mathbb{R}^4 \times [0, T])$ with $\eta = \sum_{i=1}^n \eta_i$, any functions $\{\gamma_i\}_{1 \leq i \leq n} \subset L^{\frac{3}{2}}([0, T], \mathbb{R})$ and $\beta \in L^3([0, T]; \mathbb{R}^4)$, the following local energy inequalities hold:*

$$\begin{aligned}
& \limsup_{k \rightarrow \infty} \sup_t \int_{\mathbb{R}^4} (|u_k(t)|^2 + |B_k(t)|^2) \eta(t) dx \\
& + \int_0^T \int_{\mathbb{R}^4} (|\nabla u|^2 + |\nabla B|^2) \eta dx dt + \int_0^T \int_{\mathbb{R}^4} \eta d\lambda \\
& \leq \int_0^T \int_{\mathbb{R}^4} (|u|^2 + |B|^2) |\partial_t \eta + \Delta \eta| dx dt + 2 \sum_{i=1}^n \int_0^T \int_{\mathbb{R}^4} (|u|^3 + |B|^3) |\nabla \eta_i| dx dt \\
& + 3 \sum_{i=1}^n \int_0^T \int_{\mathbb{R}^4} |\nabla \eta_i| d\omega + 2 \sum_{i=1}^n \int_0^T \int_{\mathbb{R}^4} |\nabla \eta_i| |\Pi - \gamma_i|^{\frac{3}{2}} dx dt; \tag{2.11}
\end{aligned}$$

$$\begin{aligned}
& \limsup_{k \rightarrow \infty} \sup_t \int_{\mathbb{R}^4} (|u_k(t)|^2 + |B_k(t)|^2) \eta(t) dx \\
& + \int_0^T \int_{\mathbb{R}^4} (|\nabla u|^2 + |\nabla B|^2) \eta dx dt + \int_0^T \int_{\mathbb{R}^4} \eta d\lambda \\
& \leq \int_0^T \int_{\mathbb{R}^4} (|u|^2 + |B|^2) |\partial_t \eta + \Delta \eta| dx dt + \int_0^T \int_{\mathbb{R}^4} (|u|^2 + |B|^2) (u \cdot \nabla) \eta dx dt \\
& + 2 \int_0^T \int_{\mathbb{R}^4} (|u - \beta|^3 + |B - \beta|^3) |\nabla \eta| dx dt + 3 \int_0^T \int_{\mathbb{R}^4} |\nabla \eta| d\omega \\
& + 2 \int_0^T \int_{\mathbb{R}^4} \Pi (u \cdot \nabla) \eta dx dt. \tag{2.12}
\end{aligned}$$

Proof. Substituting the test function η defined above into the local energy inequality (2.3) yields

$$\begin{aligned} & \int_{\mathbb{R}^4} (|u_k(t)|^2 + |B_k(t)|^2) \eta(t) dx + \int_0^t \int_{\mathbb{R}^4} (|\nabla u_k|^2 + |\nabla B_k|^2) \eta dx dt \\ & \leq \int_0^t \int_{\mathbb{R}^4} (|u_k|^2 + |B_k|^2) |\partial_t \eta + \Delta \eta| dx dt + \int_0^t \int_{\mathbb{R}^4} (|u_k|^2 + |B_k|^2) (\tilde{u}_k \cdot \nabla) \eta dx dt \\ & \quad + 2 \int_0^t \int_{\mathbb{R}^4} \Pi_k(u_k \cdot \nabla) \eta dx dt - \int_0^t \int_{\mathbb{R}^4} (\tilde{B}_k \cdot \nabla \eta) (u_k \cdot B_k) dx dt. \end{aligned} \quad (2.13)$$

The convergence of the second term can be obtained by Lemma 2.2 and (2.6). Since $u_k \rightarrow u$ and $B_k \rightarrow B$ in $L^2(\mathbb{R}^4 \times [0, T])$, then the convergence of the third term is straightforward. On the other hand, we note that

$$\int_0^T \int_{\mathbb{R}^4} |\tilde{u}_k|^3 |\nabla \eta| dx dt = \|g_1 + g_2\|_{L^3}^3,$$

where

$$\begin{aligned} g_1(x, t) &= \int_{\mathbb{R}^4} u_k(x - y, t) \chi_k(y) (|\nabla \eta(x, t)|^{\frac{1}{3}} - |\nabla \eta(x - y, t)|^{\frac{1}{3}}) dy, \\ g_2(x, t) &= \int_{\mathbb{R}^4} u_k(x - y, t) \chi_k(y) |\nabla \eta(x - y, t)|^{\frac{1}{3}} dy. \end{aligned}$$

For g_1 , using Young's inequality for convolution, we have

$$\begin{aligned} \|g_1\|_{L^3} &\leq \left\| \int_{\mathbb{R}^4} u_k(x - y, t) \chi_k(y) \frac{|\nabla \eta(x, t)|^{\frac{1}{3}} - |\nabla \eta(x - y, t)|^{\frac{1}{3}}}{|y|^{\frac{1}{3}}} d_k^{\frac{1}{3}} dy \right\|_{L^3} \\ &\leq C d_k^{\frac{1}{3}} \|\eta\|_{C^2} \|\tilde{u}\|_{L^3} \\ &\leq C d_k^{\frac{1}{3}} \|\eta\|_{C^2} \|u_k\|_{L^3}, \end{aligned}$$

where $d_k := \text{diam}(\text{supp } \chi_k) \rightarrow 0$ when $k \rightarrow \infty$. For g_2 , using Young's inequality for convolution again, we have

$$\|g_2\|_{L^3} = \|(u_k |\nabla \eta|^{\frac{1}{3}}) * \chi_k\|_{L^3} \leq \|u_k |\nabla \eta|^{\frac{1}{3}}\|_{L^3}.$$

Hence, combining the estimates of g_1 and g_2 , we have

$$\begin{aligned} & \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} |u_k|^2 (\tilde{u}_k \cdot \nabla) \eta dx dt \\ & \leq C \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} |u_k|^3 |\nabla \eta| dx dt + C \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} |\tilde{u}_k|^3 |\nabla \eta| dx dt \\ & \leq \int_0^T \int_{\mathbb{R}^4} |u|^3 |\nabla \eta| dx dt + \int_0^T \int_{\mathbb{R}^4} |\nabla \eta| d\omega. \end{aligned}$$

Going through a similar process, we have

$$\begin{aligned} & \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} (\tilde{B}_k \cdot \nabla \eta)(u_k \cdot B_k) dx dt \\ & \leq C \int_0^T \int_{\mathbb{R}^4} |B|^3 |\nabla \eta| dx dt + C \int_0^T \int_{\mathbb{R}^4} |\nabla \eta| d\omega. \end{aligned}$$

For the term involving pressure, with the help of incompressible condition and Lemma 2.4, we have

$$\begin{aligned} \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \Pi_k(u_k \cdot \nabla) \eta dx dt &= \sum_{i=1}^n \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \Pi_k(u_k \cdot \nabla) \eta_i dx dt \\ &= \sum_{i=1}^n \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \Pi_k(u_k \cdot \nabla) \eta_i dx dt \\ &\leq C \sum_{i=1}^n \int_0^T \int_{\mathbb{R}^4} |u|^3 |\nabla \eta_i| dx dt \\ &\quad + \sum_{i=1}^n \int_0^T \int_{\mathbb{R}^4} |\nabla \eta_i| d\omega \\ &\quad + \sum_{i=1}^n C \int_0^T \int_{\mathbb{R}^4} |\nabla \eta_i| |\Pi - \gamma_i|^{\frac{3}{2}} dx dt. \quad (2.14) \end{aligned}$$

Combining (2.13)–(2.14), we complete the proof of (2.11). For the local energy inequality (2.12), from incompressible conditions, we have

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^4} (|u_k|^2 + |B_k|^2)(\tilde{u}_k \cdot \nabla) \eta dx dt + \int_0^T \int_{\mathbb{R}^4} (\tilde{B}_k \cdot \nabla \eta)(u_k \cdot B_k) dx dt \\ & \leq C \int_0^T \int_{\mathbb{R}^4} |u_k - \beta + \beta|^2 (\tilde{u}_k \cdot \nabla) \eta dx dt + C \int_0^T \int_{\mathbb{R}^4} |B_k - \beta + \beta|^2 (\tilde{u}_k \cdot \nabla) \eta dx dt \\ & \leq C \int_0^T \int_{\mathbb{R}^4} |u_k - \beta|^2 [(\tilde{u}_k - \beta) \cdot \nabla] \eta dx dt + C \int_0^T \int_{\mathbb{R}^4} |u_k - \beta|^2 (\beta \cdot \nabla) \eta dx dt \\ & \quad + C \int_0^T \int_{\mathbb{R}^4} [(u_k - \beta) \cdot \beta] (\tilde{u}_k \cdot \nabla) \eta dx dt + C \int_0^T \int_{\mathbb{R}^4} |B_k - \beta|^2 [(\tilde{u}_k - \beta) \cdot \nabla] \eta dx dt \\ & \quad + C \int_0^T \int_{\mathbb{R}^4} |B_k - \beta|^2 (\beta \cdot \nabla) \eta dx dt + C \int_0^T \int_{\mathbb{R}^4} [(B_k - \beta) \cdot \beta] (\tilde{u}_k \cdot \nabla) \eta dx dt. \end{aligned} \quad (2.15)$$

Since

$$\tilde{u}_k - \beta = (u_k - \beta) * \chi_k \quad \text{and} \quad \tilde{B}_k - \beta = (B_k - \beta) * \chi_k,$$

applying Young's inequality for convolution yields

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^4} |u_k - \beta|^2 [(\tilde{u}_k - \beta) \cdot \nabla] \eta dx dt + \int_0^T \int_{\mathbb{R}^4} |B_k - \beta|^2 [(\tilde{u}_k - \beta) \cdot \nabla] \eta dx dt \\
& \leq \int_0^T \int_{\mathbb{R}^4} (|u_k - u|^3 + |B_k - B|^3) |\nabla \eta| dx dt \\
& \quad + \int_0^T \int_{\mathbb{R}^4} (|u - \beta|^3 + |B - \beta|^3) |\nabla \eta| dx dt \\
& \quad + C \int_0^T \int_{\mathbb{R}^4} |u_k - u|^2 |u - \beta| |\nabla \eta| dx dt \\
& \quad + C \int_0^T \int_{\mathbb{R}^4} |u_k - u| |u - \beta|^2 |\nabla \eta| dx dt \\
& \quad + C \int_0^T \int_{\mathbb{R}^4} |B_k - B|^2 |B - \beta| |\nabla \eta| dx dt \\
& \quad + C \int_0^T \int_{\mathbb{R}^4} |B_k - B| |B - \beta|^2 |\nabla \eta| dx dt \\
& \leq C \int_0^T \int_{\mathbb{R}^4} (|u_k - u|^3 + |B_k - B|^3) |\nabla \eta| dx dt \\
& \quad + C \int_0^T \int_{\mathbb{R}^4} (|u - \beta|^3 + |B - \beta|^3) |\nabla \eta| dx dt \\
& \rightarrow C \int_0^T \int_{\mathbb{R}^4} |\nabla \eta| d\omega + C \int_0^T \int_{\mathbb{R}^4} (|u - \beta|^3 + |B - \beta|^3) |\nabla \eta| dx dt \text{ as } (k \rightarrow \infty).
\end{aligned}$$

Now, we can pass $k \rightarrow \infty$ in (2.15), which yields

$$\begin{aligned}
& \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} (|u_k|^2 + |B_k|^2) (\tilde{u}_k \cdot \nabla) \eta dx dt \\
& \quad + \limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} (\tilde{B}_k \cdot \nabla \eta) (u_k \cdot B_k) dx dt \\
& \leq C \int_0^T \int_{\mathbb{R}^4} |\nabla \eta| d\omega + C \int_0^T \int_{\mathbb{R}^4} (|u - \beta|^3 + |B - \beta|^3) |\nabla \eta| dx dt \\
& \quad + C \int_0^T \int_{\mathbb{R}^4} (|u - \beta|^2 + |B - \beta|^2) (\beta \cdot \nabla) \eta dx dt \\
& \quad + C \int_0^T \int_{\mathbb{R}^4} [(u - \beta) \cdot \beta + (B - \beta) \cdot \beta] (u \cdot \nabla) \eta dx dt \\
& \leq C \int_0^T \int_{\mathbb{R}^4} |\nabla \eta| d\omega + C \int_0^T \int_{\mathbb{R}^4} (|u - \beta|^3 + |B - \beta|^3) |\nabla \eta| dx dt \\
& \quad + C \int_0^T \int_{\mathbb{R}^4} (|u|^2 + |B|^2) (u \cdot \nabla) \eta dx dt
\end{aligned}$$

$$\begin{aligned}
 & + C \int_0^T \int_{\mathbb{R}^4} (|u - \beta|^2 + |B - \beta|^2) [(u - \beta) \cdot \eta] \eta dx dt \\
 & \leq C \int_0^T \int_{\mathbb{R}^4} |\nabla \eta| d\omega + C \int_0^T \int_{\mathbb{R}^4} (|u - \beta|^3 + |B - \beta|^3) |\nabla \eta| dx dt \\
 & \quad + C \int_0^T \int_{\mathbb{R}^4} (|u|^2 + |B|^2) (u \cdot \nabla) \eta dx dt.
 \end{aligned}$$

Finally, using Lemma 2.4 yields

$$\limsup_{k \rightarrow \infty} \int_0^T \int_{\mathbb{R}^4} \Pi_k (u_k \cdot \nabla) \eta dx dt \leq \int_0^T \int_{\mathbb{R}^4} \Pi (u \cdot \nabla) \eta dx dt + \int_0^T \int_{\mathbb{R}^4} |\nabla \eta| d\omega.$$

Hence, we complete the proof of Proposition 2.1. ■

3. Partial regularity theory: Part I

This section is devoted to establishing an ε -regularity criterion for weak solution sets $(u, B, \Pi, \lambda, \omega)$ of (1.1) in terms of L^3 -norm of (u, B) . The argument we will present is based on an iterative method. Firstly, as mentioned in [30], the weak solution sets have the following scaling property.

Lemma 3.1. *If $(u, B, \Pi, \lambda, \omega)$ is a weak solution set of the MHD system (1.1), then, for any $r > 0$, the scaled set $(u_r, B_r, \Pi_r, \lambda_r, \omega_r)$ is also a weak solution set of (1.1), where $u_r, B_r, \Pi_r, \lambda_r$, and ω_r are defined as*

$$\begin{aligned}
 u_r(x, t) &= ru(rx, r^2t), \quad B_r(x, t) = rB(rx, r^2t), \quad \Pi_r(x, t) = r^2\Pi(rx, r^2t), \\
 \int_0^T \int_{\mathbb{R}^4} d\lambda_r &:= \frac{1}{r^2} \iint_{\{(rx, r^2t) | (x, t) \in \Omega\}} d\lambda, \quad \int_0^T \int_{\mathbb{R}^4} d\omega_r := \frac{1}{r^3} \iint_{\{(rx, r^2t) | (x, t) \in \Omega\}} d\omega
 \end{aligned}$$

for any $\omega \subset \mathbb{R}^4 \times \mathbb{R}$.

Proposition 3.1. *There exist an $r > 0$ and an absolute positive constant $\varepsilon > 0$ such that if a weak solution set $(u, B, \Pi, \lambda, \omega)$ of the MHD system (1.1) satisfies*

$$\frac{1}{r^3} \int_{Q_r(z_0)} |u|^3 + |B|^3 + |\Pi|^{\frac{3}{2}} dx dt + \frac{1}{r^3} \int_{Q_r(z_0)} d\omega \leq \varepsilon,$$

then $z_0 = (x_0, t_0)$ is a regular point and $Q_r(z_0) := B_r(x_0) \times (t_0 - r^2, t_0)$ denote the parabolic cylinder centered at $z_0 = (x_0, t_0)$.

For the proof of Proposition 3.1, we need two auxiliary lemmas, which play an important role in the proof of Proposition 3.1. Before stating lemmas, we introduce the following

dimensionless quantities due to the scaling property of system (1.1):

$$\begin{aligned}
 A(r) &= \limsup_{k \rightarrow \infty} \sup_{t_0 - r^2 < t < t_0} \frac{1}{r^2} \int_{B_r(x_0)} |u_k(t)|^2 + |B_k(t)|^2 dx, \\
 B(r) &= \frac{1}{r^2} \int_{Q_r(z_0)} |\nabla u|^2 + |\nabla B|^2 dx dt, \quad \tilde{B}(r) = \frac{1}{r^2} \int_{Q_r(z_0)} d\lambda, \\
 C(r) &= \frac{1}{r^3} \int_{Q_r(z_0)} |u|^3 + |B|^3 dx dt, \quad \tilde{C}(r) = \frac{1}{r^3} \int_{Q_r(z_0)} d\omega, \\
 \bar{C}(r) &= \frac{1}{r^3} \int_{Q_r(z_0)} |u - u_r|^3 + |B - B_r|^3 dx dt, \\
 D(r) &= \frac{1}{r^3} \int_{Q_r(z_0)} |\Pi|^{\frac{3}{2}} dx dt, \quad \bar{D}(r) = \frac{1}{r^3} \int_{Q_{-r}(z_0)} |\Pi - \Pi_r|^{\frac{3}{2}} dx dt,
 \end{aligned}$$

where $f_r(t) := \frac{1}{|B_r(x_0)|} \int_{B_r(x_0)} f(x, t) dx$ is the integral average of $f(x, t)$ on $B_r(x_0)$. By using iteration method, we can state that u and B are locally bounded if u , B , Π , and concentration measure ω satisfy a local smallness condition.

Lemma 3.2. *Assume that $(u, B, \Pi, \lambda, \omega)$ is a weak solution set of the MHD system (1.1) in $Q_r(z_0)$; then, there exists a universal positive constant $C > 0$ such that, for any $r > 0$,*

$$C(r) \leq CA^{\frac{3}{2}}(r) + CB^{\frac{3}{2}}(r), \quad (3.1)$$

$$\tilde{C}(r) \leq CA^{\frac{1}{2}}(r) \tilde{B}(r), \quad (3.2)$$

$$\bar{C}(r) \leq CA^{\frac{1}{2}}(r) B(r). \quad (3.3)$$

Proof. By using interpolation inequality and Sobolev inequality, we have

$$\begin{aligned}
 &\|u - u_r\|_{L^3(B_r(x_0))}^3 + \|B - B_r\|_{L^3(B_r(x_0))}^3 \\
 &\leq C \|u - u_r\|_{L^4(B_r(x_0))}^2 \|u - u_r\|_{L^2(B_r(x_0))} \\
 &\quad + C \|B - B_r\|_{L^4(B_r(x_0))}^2 \|B - B_r\|_{L^2(B_r(x_0))} \\
 &\leq \|\nabla u\|_{L^2(B_r(x_0))}^2 \|u\|_{L^2(B_r(x_0))} + \|\nabla B\|_{L^2(B_r(x_0))}^2 \|B\|_{L^2(B_r(x_0))}.
 \end{aligned}$$

Then, integrating in time, we get

$$\int_{Q_r(z_0)} |u - u_r|^3 + |B - B_r|^3 dx dt \leq Cr^3 A^{\frac{1}{2}}(r) B(r),$$

where we have used the lower semi-continuity of weak-* convergence, this is, (3.3). Noting that $\|f\|_{L^4(B_r(x_0))} \leq C \|f\|_{L^2(B_r(x_0))} + \|\nabla f\|_{L^2(B_r(x_0))}$, by a direct calculation, we get (3.1). The inequality (3.2) follows directly from Lemma 2.3. ■

In order to prove Proposition 3.1, we will define a quantity $E(r)$, which is not scale-invariant: $E(r) = \frac{1}{r^{\frac{5}{2}}} \int_{Q_r(z_0)} |\Pi - \Pi_r|^{\frac{3}{2}} dx dt = r^{\frac{1}{2}} \bar{D}(r)$.

Lemma 3.3. *Suppose that $(u, B, \Pi, \lambda, \omega)$ is a weak solution set of the MHD system (1.1) in $Q_\rho(z_0)$. Then, there exists a universal positive constant $C > 0$ such that, for any $0 < r \leq \frac{\rho}{2}$,*

$$\begin{aligned} E(r) &\leq C \frac{1}{r^{\frac{5}{2}}} \int_{Q_{2r}(z_0)} |u|^3 + |B|^3 dx dt \\ &\quad + C r^5 \left(\sup_{t_0 - r^2 < t < t_0} \int_{2r < |y - x_0| < \rho} \frac{|u|^2 + |B|^2}{|y - x_0|^5} dy \right)^{\frac{3}{2}} \\ &\quad + C \frac{r^3}{\rho^{\frac{11}{2}}} \int_{Q_\rho(z_0)} |u|^3 + |B|^3 + |\Pi|^{\frac{3}{2}} dx dt. \end{aligned} \quad (3.4)$$

Proof. We consider a smooth cutoff function $\phi \in C_c^\infty(\mathbb{R}^4)$ such that $0 \leq \phi \leq 1$, $\phi \equiv 1$ in $B_{\frac{3}{4}\rho}(x_0)$, $\phi \equiv 0$ in $\mathbb{R}^4 \setminus B_\rho(x_0)$, satisfying property $|\nabla \phi|^2 + |\nabla^2 \phi| \leq C \frac{1}{\rho^2}$. Then, similar to (2.9) for the local decomposition for pressure, we have

$$\begin{aligned} \Pi(x, t)\phi(x) &= \frac{1}{4\pi^2} \int_{\mathbb{R}^4} \frac{1}{|x - y|^2} (\eta \partial_i \partial_j (u_i u_j - B_i B_j) - 2\nabla \phi \cdot \nabla \Pi - \Pi \Delta \phi) dy \\ &= \frac{1}{4\pi^2} \int_{B_{2r}(x_0)} (u_i u_j - B_i B_j) \phi \partial_i \partial_j \left(\frac{1}{|x - y|^2} \right) dy \\ &\quad + \frac{1}{4\pi^2} \int_{B_\rho(x_0) \setminus B_{2r}(x_0)} (u_i u_j - B_i B_j) \phi \partial_i \partial_j \left(\frac{1}{|x - y|^2} \right) dy \\ &\quad + \frac{1}{4\pi^2} \int_{B_\rho(x_0)} (u_i u_j - B_i B_j) \left(\frac{\partial_i \partial_j \phi}{|x - y|^2} + \partial_j \phi \frac{4(x_i - y_i)}{|x - y|^4} \right) dy \\ &\quad + \frac{1}{4\pi^2} \int_{\mathbb{R}^4} \Pi \left(\frac{\Delta \phi}{|x - y|^2} + \frac{4(x - y) \cdot \nabla \phi}{|x - y|^4} \right) dy \\ &:= \Pi_1(x, t) + \Pi_2(x, t) + \Pi_3(x, t) + \Pi_4(x, t). \end{aligned} \quad (3.5)$$

Hence, the term $E(r)$ can be decomposed into four terms involving $\Pi_1, \Pi_2, \Pi_3, \Pi_4$, respectively:

$$E(r) \leq \sum_{l=1}^4 \frac{1}{r^{\frac{5}{2}}} \int_{Q_r(z_0)} |\Pi_l - \Pi_{l,r}|^{\frac{3}{2}} dx dt. \quad (3.6)$$

Using the Calderon–Zygmund theory for Π_1 , we get

$$\int_{B_r(x_0)} |\Pi_1|^{\frac{3}{2}} dx \leq C \int_{B_{2r}(x_0)} |u|^3 + |B|^3 dx,$$

which implies

$$\int_{Q_r(z_0)} |\Pi_1 - \Pi_{1,r}|^{\frac{3}{2}} dx dt \leq C \int_{Q_{2r}(z_0)} |u|^3 + |B|^3 dx dt.$$

For Π_2 , it is noteworthy that $2|x - y| > |y|$ when $x \in B_r(x_0)$ and $y \in B_{2r}^c(x_0)$; thus, for any $(x, t) \in Q_r(z_0)$,

$$|\nabla \Pi_2(x, t)| \leq C \int_{2r < |y| < \rho} \frac{\phi(|u|^2 + |B|^2)}{|x - y|^5} dx \leq C \int_{2r < |y| < \rho} \frac{|u|^2 + |B|^2}{|y|^5} dx,$$

by which one has

$$\begin{aligned} \int_{Q_r(z_0)} |\Pi_2 - \Pi_{2,r}|^{\frac{3}{2}} dx dt &\leq C r^4 \int_{t_0-r^2}^{t_0} \|\Pi_2 - \Pi_{2,r}\|_{L^\infty(B_r(x_0))}^{\frac{3}{2}} dt \\ &\leq C r^{\frac{11}{2}} \int_{t_0-r^2}^{t_0} \|\nabla \Pi_2\|_{L^\infty(B_r(x_0))}^{\frac{3}{2}} dt \\ &\leq C r^{\frac{15}{2}} \left(\sup_{t_0-r^2 < t < t_0} \int_{2r < |y| < \rho} \frac{|u|^2 + |B|^2}{|y|^5} dx \right)^{\frac{3}{2}}. \end{aligned}$$

Similarly, we note that $\nabla \phi = \nabla^2 \phi = 0$ in $B_{\frac{3}{4}}(x_0)$, and $|x - y| > \frac{\rho}{4}$ when $x \in B_r(x_0)$, $y \in B_\rho(x_0) \setminus B_{\frac{3\rho}{4}}(x_0)$. Thus, for any $(x, t) \in Q_r(z_0)$,

$$\begin{aligned} |\nabla \Pi_3| &\leq C \int_{B_\rho(x_0) \setminus B_{\frac{3}{4}}\rho(x_0)} (|u|^2 + |B|^2) \left(\frac{|\nabla \phi|}{|x - y|^3} + \frac{|\nabla \phi|}{|x - y|^4} \right) dy \\ &\leq C \frac{1}{\rho^5} \int_{B_\rho(x_0) \setminus B_{\frac{3}{4}}\rho(x_0)} (|u|^2 + |B|^2) dy. \\ |\nabla \Pi_4| &\leq C \int_{B_\rho(x_0) \setminus B_{\frac{3}{4}}\rho(x_0)} |\Pi| \left(\frac{|\nabla \phi|}{|x - y|^3} + \frac{|\nabla \phi|}{|x - y|^4} \right) dy \\ &\leq C \frac{1}{\rho^5} \int_{B_\rho(x_0) \setminus B_{\frac{3}{4}}\rho(x_0)} |\Pi| dy. \end{aligned}$$

Hence,

$$\begin{aligned} \sum_{l=3}^4 \int_{Q_r(z_0)} |\Pi_l - \Pi_{l,r}|^{\frac{3}{2}} dx dt &\leq C \sum_{l=3}^4 r^4 \int_{t_0-r^2}^{t_0} \|\Pi_l - \Pi_{l,r}\|_{L^\infty(B_r(x_0))}^{\frac{3}{2}} dt \\ &\leq C \sum_{l=3}^4 r^{\frac{11}{2}} \int_{Q_\rho(z_0)} \|\nabla \Pi_l\|_{L^\infty(B_r(x_0))}^{\frac{3}{2}} dt \\ &\leq C \left(\frac{r}{\rho}\right)^{\frac{11}{2}} \int_{Q_\rho(z_0)} |u|^3 + |B|^3 + |\Pi|^{\frac{3}{2}} dx dt. \quad (3.7) \end{aligned}$$

Combining (3.6) with (3.7) yields the inequality (3.4). \blacksquare

Proof of Proposition 3.1. We let $r_n = 2^{-n}$ with $n \geq 2$. The method is to iteratively prove the following estimates:

$$C(r_n) + \tilde{C}(r_n) + E(r_n) \leq \varepsilon^{\frac{2}{3}} r_n^3, \quad (3.8)$$

$$A(r_n) + B(r_n) + \tilde{B}(r_n) \leq C r_n^2 \varepsilon^{\frac{2}{3}}. \quad (3.9)$$

First, we claim that the inequality (3.8)₁ holds. Indeed, the Hölder inequality yields

$$\begin{aligned} & C(r_1) + \tilde{C}(r_1) + E(r_1) \\ & \leq 8 \int_{Q_{\frac{1}{2}}(z_0)} |u|^3 + |B|^3 dx dt + 8 \int_{Q_{\frac{1}{2}}(z_0)} d\omega + 16 \int_{Q_{\frac{1}{2}}(z_0)} |\Pi|^{\frac{3}{2}} dx dt. \end{aligned}$$

Then, we take $\varepsilon \leq 2^{-21}$ and apply the smallness assumption, which implies that

$$C(r_1) + \tilde{C}(r_1) + E(r_1) \leq 16\varepsilon \leq \varepsilon^{\frac{2}{3}} r_1^3.$$

Claim 1: $\{(3.8)_k\}_{2 \leq k \leq n}$ implies (3.9)_{n+1}. Let ϕ_n be the localized solution of the backward heat equation

$$\phi_n(x, t) = \frac{\chi(x, t)}{(r_n^2 - t)^2} \exp\left(-\frac{|x|^2}{4(r_n^2 - t)}\right),$$

where χ is a cutoff function such that $\chi \equiv 1$ in $Q_{\frac{1}{4}}(z_0)$ and $\chi \equiv 0$ in $\mathbb{R}^4 \times (-\infty, 0) \setminus Q_{\frac{1}{3}}(z_0)$.

A direct computation yields that ϕ_n satisfies the following properties:

$$\begin{aligned} & \partial_t \phi_n + \Delta \phi_n = 0 \text{ in } Q_{\frac{1}{4}}(z_0), \quad |\partial_t \phi_n + \Delta \phi_n| \leq C \text{ in } \mathbb{R}^4 \times (-\infty, 0), \\ & C^{-1} r_n^{-4} \leq \phi_n \leq C r_n^{-4} \quad \text{and} \quad |\nabla \phi_n| \leq C r_n^{-5} \text{ in } Q_{r_n}(z_0), \\ & \phi_n \leq C r_k^{-4} \quad \text{and} \quad |\nabla \phi_n| \leq C r_k^{-5} \text{ in } Q_{r_{k-1}}(z_0) \setminus Q_{r_k}(z_0) \text{ for any } 2 \leq k \leq n. \end{aligned}$$

Next, we define the smooth cutoff function $\{\eta_k\}_{k \in \mathbb{N}}$ such that

$$\eta_k \equiv 1 \text{ in } Q_{r_{\frac{7}{8}k}}(z_0), \quad \eta_k \equiv 0 \text{ in } \mathbb{R}^4 \times (-\infty, 0) \setminus Q_{r_k}(z_0), \quad |\nabla \eta_k| \leq C r_k^{-1}.$$

Then, we define $\varphi_k := \phi_n(\eta_k - \eta_{k+1})$ for $1 \leq k \leq n-1$ and $\varphi_n := \phi_n \eta_n$. It is easy to see that $\phi_n = \sum_{k=1}^n \varphi_k$ and

$$|\nabla \varphi_k| = |\phi_n \nabla \eta_k + \eta_k \nabla \phi_n| \leq C r_k^{-5} \quad \text{for any } k \leq n.$$

Substituting ϕ_n into the local energy inequality (2.11) yields

$$\begin{aligned} & \limsup_{k \rightarrow \infty} \sup_t \int_{B_{\frac{1}{2}}(x_0)} (|u_k(t)|^2 + |B_k(t)|^2) \phi_n dx \\ & + \int_{Q_{\frac{1}{2}}(z_0)} (|\nabla u|^2 + |\nabla B|^2) \phi_n dx dt + \int_{Q_{\frac{1}{2}}(z_0)} \phi_n d\lambda \\ & \leq \int_{Q_{\frac{1}{2}}(z_0)} (|u|^2 + |B|^2) |\partial_t \phi_n + \Delta \phi_n| dx dt \\ & + 2 \sum_{k=1}^n \int_{Q_{\frac{1}{2}}(z_0)} |\nabla \varphi_k| [(|u|^3 + |B|^3) dx dt + d\omega] \\ & + 2 \sum_{k=1}^n \int_{Q_{\frac{1}{2}}(z_0)} |\nabla \varphi_k| |\Pi - \Pi_{r_k}|^{\frac{3}{2}} dx dt \\ & := I_1 + I_2 + I_3. \end{aligned} \tag{3.10}$$

With the properties of ϕ_n , we deduce that

$$C^{-1}r_{n+1}^{-2}(A(r_{n+1}) + B(r_{n+1}) + \tilde{B}(r_{n+1})) \leq I_1 + I_2 + I_3.$$

For I_1 , applying the Hölder inequality and the smallness assumption yields

$$\begin{aligned} I_1 &\leq C \int_{Q_{\frac{1}{2}}(z_0)} |u|^2 + |B|^2 dx dt \\ &\leq C \left(\int_{Q_{\frac{1}{2}}(z_0)} 1 dx dt \right)^{\frac{1}{3}} \left(\int_{Q_{\frac{1}{2}}(z_0)} |u|^3 + |B|^3 dx dt \right)^{\frac{2}{3}} \\ &\leq C \left(\int_{Q_1(z_0)} |u|^3 + |B|^3 dx dt \right)^{\frac{2}{3}} \\ &\leq C \varepsilon^{\frac{2}{3}}, \\ I_2 &= \sum_{k=2}^n \int_{Q_{r_{k-1}}(z_0) \setminus Q_{r_k}(z_0)} |\nabla \varphi_k| (|u|^3 + |B|^3) dx dt + d\omega \\ &\quad + \int_{Q_{r_n}(z_0)} |\nabla \varphi_k| (|u|^3 + |B|^3) dx dt + d\omega \\ &\leq C \sum_{k=2}^n \frac{1}{r_k^5} \int_{Q_{r_{k-1}}(z_0) \setminus Q_{r_k}(z_0)} (|u|^3 + |B|^3) dx dt + d\omega \\ &\quad + C \frac{1}{r_n^5} \int_{Q_{r_n}(z_0)} (|u|^3 + |B|^3) dx dt + d\omega \\ &\leq C \sum_{k=2}^n \frac{1}{r_{k-1}^5} \int_{Q_{r_{k-1}}(z_0)} (|u|^3 + |B|^3) dx dt + d\omega \\ &\quad + C \frac{1}{r_n^5} \int_{Q_{r_n}(z_0)} (|u|^3 + |B|^3) dx dt + d\omega \\ &\leq C \sum_{k=1}^n r_k \varepsilon^{\frac{2}{3}}, \\ I_3 &\leq C \sum_{k=2}^n \frac{1}{r_k^5} \int_{Q_{r_{k-1}}(z_0) \setminus Q_{r_k}(z_0)} |\Pi - \Pi_{r_k}|^{\frac{3}{2}} dx dt \\ &\quad + C \frac{1}{r_n^5} \int_{Q_{r_n}(z_0)} |\Pi - \Pi_{r_n}|^{\frac{3}{2}} dx dt \\ &\leq C \sum_{k=1}^n r_k^{\frac{1}{2}} \varepsilon^{\frac{2}{3}}. \end{aligned}$$

Combining the estimates for I_1, I_2, I_3 and the local energy inequality (3.10), we deduce that (3.9)_{n+1} holds.

Claim 2: $\{(3.9)_k\}_{2 \leq k \leq n}$ implies $(3.8)_n$. Lemma 3.2 yields that, for any $2 \leq k \leq n$,

$$C(r_k) + \tilde{C}(r_k) \leq CA^{\frac{3}{2}}(r_k) + CB^{\frac{3}{2}}(r_k) + C\tilde{B}(r_k) \leq C\epsilon r_k^3. \quad (3.11)$$

Taking $\rho = r_1$, $r = r_n$ and using Lemmas 3.2 and 3.3, we have

$$\begin{aligned} E(r) &\leq C \frac{1}{r_n^{\frac{5}{2}}} \int_{Q_{r_{n-1}}(z_0)} |u|^3 + |B|^3 dx dt \\ &\quad + C r_n^5 \left(\sup_{t_0 - r_n^2 < t < t_0} \int_{2r_n < |y| < r_1} \frac{|u|^2 + |B|^2}{|y|^5} dy \right)^{\frac{3}{2}} \\ &\quad + C \frac{r_n^3}{r_1^{\frac{11}{2}}} \int_{Q_{r_1}(z_0)} |u|^3 + |B|^3 + |\Pi|^{\frac{3}{2}} dx dt \\ &\leq C r_n^{\frac{1}{2}} C(r_{n-1}) + C r_n^5 \left(\sup_{t_0 - r_n^2 < t < t_0} \int_{2r_n < |y| < r_1} \frac{|u|^2 + |B|^2}{|y|^5} dy \right)^{\frac{3}{2}} + C \epsilon r_n^3 \\ &\leq C \epsilon r_n^3 + C r_n^5 \left(\sup_{t_0 - r_n^2 < t < t_0} \int_{2r_n < |y| < r_1} \frac{|u|^2 + |B|^2}{|y|^5} dy \right)^{\frac{3}{2}}. \end{aligned}$$

It is noteworthy that

$$\begin{aligned} \sup_{t_0 - r_n^2 < t < t_0} \int_{2r_n < |y| < r_1} \frac{|u|^2 + |B|^2}{|y|^5} dy &\leq \sum_{k=2}^{n-1} \sup_{t_0 - r_{k-1}^2 < t < t_0} \int_{r_k < |y| < r_{k-1}} \frac{|u|^2 + |B|^2}{|y|^5} dy \\ &\leq C \sum_{k=2}^{n-1} \frac{1}{r_k^3} A(r_{k-1}) \leq C \epsilon^{\frac{2}{3}} \sum_{k=2}^{n-1} \frac{1}{r_k} \leq C \epsilon^{\frac{2}{3}} r_n^{-1}. \end{aligned} \quad (3.12)$$

Hence, from (3.11)–(3.12), we can deduce that

$$C(r_n) + \tilde{C}(r_n) + E(r_n) \leq C \epsilon r_n^3 \leq \epsilon^{\frac{2}{3}} r_n^3.$$

Now, we can see that $(3.9)_k$ holds for any $k \geq 2$. Hence, using Lebesgue convergence theorem, we get that $z_0 = (x_0, t_0)$ is a regular point. ■

4. Partial regularity theory: Part II

With the help of Proposition 3.1, we can prove the following partial regularity result, which plays a vital role in the proof of Theorem 1.1.

Proposition 4.1. *Assume that $(u, B, \Pi, \lambda, \omega)$ is a weak solution set of the MHD system (1.1) in some cylinder $Q_\rho(z_0)$. Then, there exists a universal positive constant $\epsilon_0 > 0$*

such that if

$$\limsup_{r \rightarrow 0} \frac{1}{r^2} \int_{Q_r(z_0)} (|\nabla u|^2 + |\nabla B|^2) dx dt + d\lambda \leq \varepsilon_0,$$

then $\|u\|_{L^\infty(Q_r(z_0))} < C r_0^{-1}$ and $\|B\|_{L^\infty(Q_r(z_0))} < C r_0^{-1}$ for some $0 < r_0 < \rho$.

For the proof of Proposition 4.1, we also need another decay estimate for the pressure Π .

Lemma 4.1. Assume that $(u, B, \Pi, \lambda, \omega)$ is a weak solution set of the MHD system (1.1) in $Q_r(z_0)$. Then, there exists a universal positive constant $C > 0$ such that, for $r > 0$ and $\theta \in (0, \frac{1}{2}]$,

$$D(\theta r) \leq C \theta^{-3} A^{\frac{1}{2}}(r) B(r) + C \theta D(r).$$

Proof. We consider a smooth cutoff function $\psi \in C_c^\infty(\mathbb{R}^4)$ such that $0 \leq \psi \leq 1$ and $\psi \equiv 1$ in $B_{\frac{3}{4}}(x_0)$, $\psi \equiv 0$ in $\mathbb{R}^4 \setminus B_r(x_0)$, $|\nabla \psi|^2 + |\nabla^2 \psi| \leq C \frac{1}{r^2}$. The pressure equation can be written as

$$-\Delta \Pi = \partial_i \partial_j [(u_i - u_{i,r})(u_j - u_{j,r}) - (B_i - B_{i,r})(B_j - B_{j,r})].$$

We can localize this equation like (3.5) to obtain

$$\Pi(x, t) \psi(x) = \widetilde{\Pi}_1(x, t) + \widetilde{\Pi}_2(x, t) + \widetilde{\Pi}_3(x, t),$$

where

$$\begin{aligned} \widetilde{\Pi}_1(x, t) &= \frac{1}{4\pi^2} \int_{B_r(x_0)} [(u_i - u_{i,r})(u_j - u_{j,r}) - (B_i - B_{i,r})(B_j - B_{j,r})] \\ &\quad \times \psi \partial_i \partial_j \left(\frac{1}{|x - y|^2} \right) dy, \\ \widetilde{\Pi}_2(x, t) &= \frac{1}{4\pi^2} \int_{B_r(x_0)} [(u_i - u_{i,r})(u_j - u_{j,r}) - (B_i - B_{i,r})(B_j - B_{j,r})] \\ &\quad \times \left(\frac{\partial_i \partial_j \psi}{|x - y|^2} + \partial_j \psi \frac{4(x_i - y_i)}{|x - y|^4} \right) dy, \\ \widetilde{\Pi}_3(x, t) &= \frac{1}{4\pi^2} \int_{B_r(x_0)} \Pi \left(\frac{\Delta \psi}{|x - y|^2} + \frac{4(x - y) \cdot \nabla \psi}{|x - y|^4} \right) dy. \end{aligned}$$

For $\widetilde{\Pi}_1$, the Calderon–Zygmund theory yields

$$\begin{aligned} \int_{B_{\theta r}(x_0)} |\widetilde{\Pi}_1|^{\frac{3}{2}} dx dt &\leq C \int_{B_r(x_0)} |(u_i - u_{i,r})(u_j - u_{j,r})|^{\frac{3}{2}} \\ &\quad + |(B_i - B_{i,r})(B_j - B_{j,r})|^{\frac{3}{2}} dx \\ &\leq C \int_{B_r(x_0)} |u - u_r|^3 + |B - B_r|^3 dx. \end{aligned}$$

Integrating in time gives

$$\int_{Q_{\theta r}(z_0)} |\widetilde{\Pi}_1|^{\frac{3}{2}} dx dt \leq C \int_{Q_r(z_0)} |u - u_r|^3 + |B - B_r|^3 dx dt.$$

For $\widetilde{\Pi}_2$, we note that $\nabla \psi$ is supported in $B_r(x_0) \setminus B_{\frac{3}{4}r}(x_0)$. Then, for $x \in B_{\theta r}(x_0)$ and $|x - y| > \frac{r}{4}$, combining the properties of ψ , we know that

$$|\widetilde{\Pi}_2| \leq C \frac{1}{r^4} \int_{B_r(x_0)} |u - u_r|^2 + |B - B_r|^2 dx,$$

which yields that

$$\begin{aligned} \int_{Q_{\theta r}(z_0)} |\widetilde{\Pi}_2|^{\frac{3}{2}} dx dt &\leq C(\theta r)^4 \int_{t_0 - (\theta r)^2}^{t_0} \|\widetilde{\Pi}_2\|_{L^\infty} dt \\ &\leq C\theta^4 \int_{Q_r(z_0)} |u - u_r|^3 + |B - B_r|^3 dx dt. \end{aligned}$$

Similarly, for $\widetilde{\Pi}_3$, we have

$$\int_{Q_{\theta r}(z_0)} |\widetilde{\Pi}_3|^{\frac{3}{2}} dx dt \leq C\theta^4 \int_{Q_r(z_0)} |\Pi|^{\frac{3}{2}} dx dt.$$

Collecting all estimates for $\widetilde{\Pi}_1$, $\widetilde{\Pi}_2$, $\widetilde{\Pi}_3$ and applying Lemma 3.2, we complete the proof of Lemma 4.1. \blacksquare

Proof of Proposition 4.1. According to Proposition 3.1, we need only to prove that, for $r_1 > 0$,

$$\frac{1}{r_1^3} \int_{Q_{r_1}(z_0)} |u|^3 + |B|^3 + |\Pi|^{\frac{3}{2}} dx dt + \frac{1}{r_1^3} \int_{Q_{r_1}(z_0)} d\omega \leq \varepsilon. \quad (4.1)$$

For fixed $r > 0$ and any $0 < \theta \leq \frac{1}{2}$, we consider the cutoff function

$$\phi_\theta(x, t) = \frac{1}{[(\theta r)^2 - t]} \exp\left(-\frac{|x|^2}{4[(\theta r)^2 - t]}\right) \chi\left(\frac{x}{r}, \frac{t}{r^2}\right)(x, t) \in \mathbb{R}^4 \times (-\infty, 0),$$

where $\chi \in C_c^\infty(B_1(x_0) \times (-1, 1))$ is a cutoff function such that

$$\chi \equiv 1 \quad \text{in } B_{\frac{1}{2}}(x_0) \times \left(-\frac{1}{4}, \frac{1}{4}\right),$$

satisfying the following properties:

$$\begin{aligned} C^{-1}(\theta r)^{-4} &\leq \phi_\theta \leq C(\theta r)^{-4} \quad \text{in } Q_{\theta r}(z_0), \\ \phi_\theta &\leq C(\theta r)^{-4}, \quad |\nabla \phi_\theta| \leq C(\theta r)^{-5}, \quad |\partial_t \phi_\theta + \Delta \phi_\theta| \leq C r^{-6} \quad \text{in } Q_r(z_0). \end{aligned}$$

Substituting ϕ_θ into the local energy inequality (2.12) yields that

$$\begin{aligned}
& \frac{1}{C(\theta r)^2} \limsup_{k \rightarrow \infty} \sup_{t_0 - (\theta r)^2 < t < t_0} \int_{B_{\theta r}(x_0)} |u_k|^2 + |B_k|^2 dx \\
& + \frac{1}{C(\theta r)^2} \int_{Q_{\theta r}(z_0)} (|\nabla u|^2 + |\nabla B|^2) dx dt + d\lambda \\
& \leq (\theta r)^2 \int_{Q_r(z_0)} (|u|^2 + |B|^2) (\partial_t \phi_\theta + \Delta \phi_\theta) dx dt \\
& + (\theta r)^2 \int_{Q_r(z_0)} (|u|^2 + |B|^2) u \cdot \nabla \phi_\theta dx dt \\
& + (\theta r)^2 \int_{Q_r(z_0)} |\nabla \phi_\theta| [(|u - u_r|^3 + |B - B_r|^3) dx dt + d\omega] \\
& + (\theta r)^2 \int_{Q_r(z_0)} \Pi u \cdot \nabla \phi_\theta dx dt \\
& := K_1 + K_2 + K_3 + K_4.
\end{aligned} \tag{4.2}$$

Applying the Hölder inequality and the interpolation inequality in Lemma 3.2, we have

$$K_1 \leq C\theta^2 C^{\frac{2}{3}}(r), \quad K_3 \leq C\theta^{-3} [\tilde{C}(r) + \bar{C}(r)], \quad K_4 \leq C\theta^{-3} D^{\frac{2}{3}}(r) C^{\frac{1}{3}}. \tag{4.3}$$

For K_2 , using the property of ϕ_θ and integration by parts, we know that

$$\begin{aligned}
K_2 &= (\theta r)^2 \int_{Q_r(z_0)} (|u|^2 + |B|^2) (u - u_r) \cdot \nabla \phi_\theta dx dt \\
&+ (\theta r)^2 \int_{Q_r(z_0)} (|u|^2 + |B|^2) u_r \cdot \nabla \phi_\theta dx dt \\
&= (\theta r)^2 \int_{Q_r(z_0)} (|u - u_r|^2 + |B - B_r|^2) (u - u_r) \cdot \nabla \phi_\theta dx dt \\
&+ 2(\theta r)^2 \int_{Q_r(z_0)} [u \cdot u_r (u - u_r) + B \cdot B_r (u - u_r)] \cdot \nabla \phi_\theta dx dt \\
&- (\theta r)^2 \int_{Q_r(z_0)} (|u_r|^2 + |B_r|^2) (u - u_r) \cdot \nabla \phi_\theta dx dt \\
&+ (\theta r)^2 \int_{Q_r(z_0)} (|u|^2 + |B|^2) u_r \cdot \nabla \phi_\theta dx dt \\
&\leq C(\theta r)^{-3} \int_{Q_r(z_0)} |u - u_r|^3 + |B - B_r|^3 dx dt \\
&- 2(\theta r)^2 \int_{Q_r(z_0)} \phi_\theta [(u - u_r) \cdot \nabla u \cdot u_r + (u - u_r) \cdot \nabla B \cdot B_r] dx dt \\
&- 2(\theta r)^2 \int_{Q_r(z_0)} \phi_\theta [u \cdot (u_r \cdot \nabla) u + B \cdot (u_r \cdot \nabla) B] dx dt,
\end{aligned} \tag{4.4}$$

where we have used the fact that u and $u - u_r$ are divergence-free. Furthermore,

$$\begin{aligned}
& \int_{Q_r(z_0)} \phi_\theta [(u - u_r) \cdot \nabla u \cdot u_r + (u - u_r) \cdot \nabla B \cdot B_r] dx dt \\
& \leq \frac{C}{(\theta r)^4} \int_{t_0-r^2}^{t_0} \frac{1}{r^4} \int_{B_r(x_0)} |u| + |B| dx \int_{B_r(x_0)} |u - u_r| (|\nabla u| + |\nabla B|) dx dt \\
& \leq \frac{C}{r^{\theta^4 \frac{16}{3}}} \int_{t_0-r^2}^{t_0} \left(\int_{B_r(x_0)} |u|^3 + |B|^3 dx \right)^{\frac{1}{3}} \left(\int_{B_r(x_0)} |u - u_r|^2 dx \right)^{\frac{1}{2}} \\
& \quad \times \left(\int_{B_r(x_0)} |\nabla u|^2 + |\nabla B|^2 dx \right)^{\frac{1}{2}} dt \\
& \leq \frac{C}{\theta^4 r^5} \left(\int_{Q_r(z_0)} |u|^3 + |B|^3 dx dt \right)^{\frac{1}{3}} \left(\int_{Q_r(z_0)} |\nabla u|^2 + |\nabla B|^2 dx dt \right)^{\frac{1}{2}} \\
& \quad \times \left(\sup_{t_0-r^2 < t < t_0} \int_{B_r(x_0)} |u|^2 dx \right)^{\frac{1}{2}}.
\end{aligned}$$

Similarly, using the Hölder equality, we have

$$\begin{aligned}
& \int_{Q_r(z_0)} \phi_\theta [u \cdot (u_r \cdot \nabla) u + B \cdot (u_r \cdot \nabla) B] dx dt \\
& \leq \frac{C}{\theta^4 r^5} \left(\int_{Q_r(z_0)} |u|^3 + |B|^3 dx dt \right)^{\frac{1}{3}} \left(\int_{Q_r(z_0)} |\nabla u|^2 + |\nabla B|^2 dx dt \right)^{\frac{1}{2}} \\
& \quad \times \left(\sup_{t_0-r^2 < t < t_0} \int_{B_r(x_0)} |u|^2 dx \right)^{\frac{1}{2}}. \tag{4.5}
\end{aligned}$$

Combining (4.4)–(4.5), we get

$$K_2 \leq C \theta^{-2} C^{\frac{1}{3}}(r) A^{\frac{1}{2}}(r) B^{\frac{1}{2}}(r) + C \theta^{-3} \bar{C}(r). \tag{4.6}$$

Substituting (4.3) and (4.6) into (4.2) and using Lemma 3.2, we get

$$\begin{aligned}
& A(\theta r) + B(\theta r) + \tilde{B}(\theta r) \\
& \leq C(\theta^2 C^{\frac{2}{3}}(r) + \theta^{-3} D^{\frac{2}{3}}(r) C^{\frac{1}{3}}(r) + \theta^{-3} [\bar{C}(r) + \tilde{C}(r)] + \theta^{-2} C^{\frac{1}{3}} A^{\frac{1}{2}}(r) B^{\frac{1}{2}}(r)) \\
& \leq C[\theta^2 C^{\frac{2}{3}}(r) + \theta^{-8} D^{\frac{4}{3}}(r) + \theta^{-3} A^{\frac{1}{2}}(r)(B(r) + \tilde{B}(r)) + \theta^{-6} A(r)B(r)] \\
& \leq C[\theta^2 A(r) + \theta^2 [B(r) + \tilde{B}(r)] + \theta^{-8} A(r)[B(r) + \tilde{B}(r)] \\
& \quad + \theta^{-6} A(r)B(r) + \theta^{-8} D^{\frac{4}{3}}(r)]. \tag{4.7}
\end{aligned}$$

On the other hand, from Lemma 4.1, we get

$$\begin{aligned}
D^{\frac{4}{3}}(\theta r) & \leq C \theta^{-4} A^{\frac{2}{3}}(r) B^{\frac{4}{3}}(r) + \theta^{\frac{4}{3}} D^{\frac{4}{3}}(r) \\
& \leq C \theta^{-12} A(r)B(r) + \theta^{12} B^2(r) + C \theta^{\frac{4}{3}} D^{\frac{4}{3}}(r). \tag{4.8}
\end{aligned}$$

Combining (4.7) and (4.8) yields

$$\begin{aligned} & A^{\theta r} + \theta^{-9} D^{\frac{4}{3}}(\theta r) + B(\theta r) + \tilde{B}(\theta r) \\ & \leq C[\theta^2 A(r) + \theta^2[B(r) + \tilde{B}(r)] + \theta^3 B^2(r) \\ & \quad + (1 + \theta^{-15} + \theta^{-2})\theta^{-6} A(r)[B(r) + \tilde{B}(r)] + (\theta + \theta^{\frac{4}{3}})\theta^{-9} D^{\frac{4}{3}}(r)]. \end{aligned} \quad (4.9)$$

We can fix $\theta \in (0, \frac{1}{2}]$ such that

$$C(\theta + \theta^{\frac{4}{3}} + \theta^2) \leq \frac{1}{4}.$$

According to the smallness assumption, we deduce that there exists an $r' > 0$ such that, for any $0 < r \leq r'$, we have

$$B(r) + \tilde{B}(r) \leq 2\varepsilon_0.$$

Then, we take ε_0 sufficiently small such that

$$C(1 + \theta^{-10} + \theta^{-2})\theta^{-6}\varepsilon_0 \leq \frac{1}{4}.$$

Letting $G(r) = A(r) + \theta^{-9} D^{\frac{4}{3}}(r) + B(r) + \tilde{B}(r)$, then, for any $0 < r \leq r'$, from (4.9) we know that

$$G(\theta r) \leq \frac{1}{2}G(r) + \frac{\varepsilon_0}{2}.$$

Iterating this inequality yields

$$A(\theta^k r') + \theta^{-9} D^{\frac{4}{3}}(\theta^k r') + B(\theta^k r') + \tilde{B}(\theta^k r') = G(\theta^k r') \leq \frac{1}{2}G(r') + \varepsilon_0.$$

Then, there exists some $r_1 > 0$ such that

$$A(r_1) + B(r_1) + \tilde{B}(r_1) + \theta^{-9} D^{\frac{4}{3}}(r_1) \leq 4\varepsilon_0.$$

Using Lemma 3.2 again, we can bound $C(r_1) + \tilde{C}(r_1)$ with $A(r_1) + B(r_1) + \tilde{B}(r_1)$. Then, we can impose additional condition on ε_0 to ensure (4.1). Hence, we complete the proof of Proposition 4.1. \blacksquare

Proof of Theorem 1.1. Without loss of generality, we may suppose the S is bounded. Let V be a parabolic neighborhood of S and fix $\delta > 0$. According to Proposition 4.1, for each $(x, t) \in S$, we can choose $Q_r(x, t) \subset V$ with $r < \delta$ such that

$$\frac{1}{r^2} \int_{Q_r(z_0)} (|\nabla u|^2 + |\nabla B|^2) dx dt + d\lambda > \varepsilon_0.$$

Since S is bounded, we can use Vitali covering lemma to obtain a family of finite disjoint parabolic cylinders $\{Q_{r_i}(x_i, t_i)\}_{i \in \Lambda}$ such that

$$S \subset \bigcup_{i \in \Lambda} Q_{5r_i}(x_i, t_i).$$

Then,

$$\begin{aligned}\sum_{i \in \Lambda} r_i^2 &\leq \frac{1}{\varepsilon_0} \sum_{i \in \Lambda} \int_{Q_{r_i}(x_i, t_i)} (|\nabla u|^2 + |\nabla B|^2) dx dt + d\lambda \\ &\leq \frac{1}{\varepsilon_0} \int_V (|\nabla u|^2 + |\nabla B|^2) dx dt + d\lambda.\end{aligned}$$

Since δ is arbitrary, we conclude that S has Lebesgue measure zero and

$$\mathcal{H}^2(S) \leq \frac{5}{\varepsilon_0} \int_V (|\nabla u|^2 + |\nabla B|^2) dx dt + d\lambda$$

for every neighborhood V of S . From the fact that $|\nabla u|^2$ and $|\nabla B|^2$ are integrable functions, hence, we finish the proof of Theorem 1.1. \blacksquare

Conflicts of interest. The author declares that there are no conflicts of interest related to this work.

References

- [1] R. Beekie, S. Friedlander, and V. Vicol, [On Moffatt's magnetic relaxation equations](#). *Comm. Math. Phys.* **390** (2022), no. 3, 1311–1339 Zbl 1508.35058 MR 4389083
- [2] D. Biskamp, [Nonlinear magnetohydrodynamics](#). Cambridge Monogr. Plasma Phys. 1, Cambridge University Press, Cambridge, 1993 MR 1250152
- [3] L. Caffarelli, R. Kohn, and L. Nirenberg, [Partial regularity of suitable weak solutions of the Navier–Stokes equations](#). *Comm. Pure Appl. Math.* **35** (1982), no. 6, 771–831 Zbl 0509.35067 MR 0673830
- [4] C. Cao, J. Wu, and B. Yuan, [The 2D incompressible magnetohydrodynamics equations with only magnetic diffusion](#). *SIAM J. Math. Anal.* **46** (2014), no. 1, 588–602 Zbl 1293.35233 MR 3163239
- [5] D. Chamorro and J. He, [Regularity theory for the dissipative solutions of the magnetohydrodynamic equations](#). *SIAM J. Math. Anal.* **53** (2021), no. 5, 5288–5321 Zbl 1473.76074 MR 4315481
- [6] Q. Chen, C. Miao, and Z. Zhang, [The Beale–Kato–Majda criterion for the 3D magnetohydrodynamics equations](#). *Comm. Math. Phys.* **275** (2007), no. 3, 861–872 Zbl 1138.76066 MR 2336368
- [7] H. Dong and D. Du, [Partial regularity of solutions to the four-dimensional Navier–Stokes equations at the first blow-up time](#). *Comm. Math. Phys.* **273** (2007), no. 3, 785–801 Zbl 1156.35442 MR 2318865
- [8] H. Dong and R. M. Strain, [On partial regularity of steady-state solutions to the 6D Navier–Stokes equations](#). *Indiana Univ. Math. J.* **61** (2012), no. 6, 2211–2229 Zbl 1286.35193 MR 3129108
- [9] G. Duvaut and J.-L. Lions, [Inéquations en thermoélasticité et magnétohydrodynamique](#). *Arch. Rational Mech. Anal.* **46** (1972), 241–279 Zbl 0264.73027 MR 0346289
- [10] X. Gu, [Regularity criteria for suitable weak solutions to the four dimensional incompressible magneto-hydrodynamic equations near boundary](#). *J. Differential Equations* **259** (2015), no. 4, 1354–1378 Zbl 1317.35198 MR 3345854

- [11] P. Han and C. He, [Partial regularity of suitable weak solutions to the four-dimensional incompressible magneto-hydrodynamic equations](#). *Math. Methods Appl. Sci.* **35** (2012), no. 11, 1335–1355 Zbl 1256.35081 MR 2945856
- [12] C. He and Z. Xin, [On the regularity of weak solutions to the magnetohydrodynamic equations](#). *J. Differential Equations* **213** (2005), no. 2, 235–254 Zbl 1072.35154 MR 2142366
- [13] C. He and Z. Xin, [Partial regularity of suitable weak solutions to the incompressible magneto-hydrodynamic equations](#). *J. Funct. Anal.* **227** (2005), no. 1, 113–152 Zbl 1083.35110 MR 2165089
- [14] Q. Jiu and Y. Wang, [Remarks on partial regularity for suitable weak solutions of the incompressible magnetohydrodynamic equations](#). *J. Math. Anal. Appl.* **409** (2014), no. 2, 1052–1065 Zbl 1306.35096 MR 3103218
- [15] O. A. Ladyzhenskaya and G. A. Seregin, [On partial regularity of suitable weak solutions to the three-dimensional Navier–Stokes equations](#). *J. Math. Fluid Mech.* **1** (1999), no. 4, 356–387 Zbl 0954.35129 MR 1738171
- [16] F. Lin, [A new proof of the Caffarelli–Kohn–Nirenberg theorem](#). *Comm. Pure Appl. Math.* **51** (1998), no. 3, 241–257 Zbl 0958.35102 MR 1488514
- [17] F.-H. Lin and C. Liu, [Partial regularity of the dynamic system modeling the flow of liquid crystals](#). *Discrete Contin. Dynam. Systems* **2** (1996), no. 1, 1–22 Zbl 0948.35098 MR 1367385
- [18] J. Liu and W. Wang, [Boundary regularity criteria for the 6D steady Navier–Stokes and MHD equations](#). *J. Differential Equations* **264** (2018), no. 3, 2351–2376 Zbl 1378.35221 MR 3721431
- [19] W. Ren and G. Wu, [Partial regularity for the 3D magneto-hydrodynamics system with hyperdissipation](#). *Acta Math. Sin. (Engl. Ser.)* **31** (2015), no. 7, 1097–1112 Zbl 1320.35129 MR 3360775
- [20] J. C. Robinson, J. L. Rodrigo, and W. Sadowski, [The three-dimensional Navier–Stokes equations. Classical theory](#). Cambridge Stud. Adv. Math. 157, Cambridge University Press, Cambridge, 2016 Zbl 1358.35002 MR 3616490
- [21] V. Scheffer, [Partial regularity of solutions to the Navier–Stokes equations](#). *Pacific J. Math.* **66** (1976), no. 2, 535–552 Zbl 0325.35064 MR 0454426
- [22] V. Scheffer, [Hausdorff measure and the Navier–Stokes equations](#). *Comm. Math. Phys.* **55** (1977), no. 2, 97–112 Zbl 0357.35071 MR 0510154
- [23] V. Scheffer, [The Navier–Stokes equations in space dimension four](#). *Comm. Math. Phys.* **61** (1978), no. 1, 41–68 Zbl 0403.35088 MR 0501249
- [24] M. Sermange and R. Temam, [Some mathematical questions related to the MHD equations](#). *Comm. Pure Appl. Math.* **36** (1983), no. 5, 635–664 Zbl 0524.76099 MR 0716200
- [25] J. Simon, [Compact sets in the space \$L^p\(0, T; B\)\$](#) . *Ann. Mat. Pura Appl. (4)* **146** (1987), 65–96 Zbl 0629.46031 MR 0916688
- [26] M. Struwe, [On partial regularity results for the Navier–Stokes equations](#). *Comm. Pure Appl. Math.* **41** (1988), no. 4, 437–458 Zbl 0632.76034 MR 0933230
- [27] A. F. Vasseur, [A new proof of partial regularity of solutions to Navier–Stokes equations](#). *NoDEA Nonlinear Differential Equations Appl.* **14** (2007), no. 5-6, 753–785 Zbl 1142.35066 MR 2374209
- [28] W. Wang and Z. Zhang, [On the interior regularity criteria for suitable weak solutions of the magnetohydrodynamics equations](#). *SIAM J. Math. Anal.* **45** (2013), no. 5, 2666–2677 Zbl 1284.35104 MR 3097270

- [29] Y. Wang and G. Wu, [A unified proof on the partial regularity for suitable weak solutions of non-stationary and stationary Navier–Stokes equations](#). *J. Differential Equations* **256** (2014), no. 3, 1224–1249 Zbl [1283.35069](#) MR [3128938](#)
- [30] B. Wu, [Partially regular weak solutions of the Navier–Stokes equations in \$\mathbb{R}^4 \times \[0, \infty\[\$](#) . *Arch. Ration. Mech. Anal.* **239** (2021), no. 3, 1771–1808 Zbl [1513.35440](#) MR [4215201](#)
- [31] B. Wu, [Partially regular weak solutions of the stationary Navier–Stokes equations in dimension 6](#). *Calc. Var. Partial Differential Equations* **61** (2022), no. 4, article no. 152 Zbl [1491.35324](#) MR [4438343](#)

Received 14 October 2023; revised 13 March 2025.

Zhongbao Zuo

School of Mathematics and Statistics, Central South University, 410083 Changsha, P. R. China;
zuozhongbao0927@163.com