



Jean-François Bony · Dorian Le Peutrec · Laurent Michel

Eyring–Kramers law for Fokker–Planck type differential operators

Received January 18, 2022; revised October 28, 2023

Abstract. We consider Fokker–Planck type differential operators associated with general Langevin processes admitting a Gibbs stationary distribution. Under assumptions ensuring suitable resolvent estimates, we prove Eyring–Kramers formulas for the bottom of the spectrum of these operators in the low temperature regime. Our approach is based on the construction of sharp Gaussian quasimodes and avoids supersymmetry or PT-symmetry assumptions.

Keywords: metastability, spectral asymptotics, infinitesimal generator, tunneling effect.

1. Introduction

1.1. Motivations

Let P be the real semiclassical second order differential operator

$$P = -h \operatorname{div} \circ A \circ h \nabla + \frac{1}{2}(b \cdot h \nabla + h \operatorname{div} \circ b) + c, \quad (1.1)$$

where the matrix field A , the vector field b and the function c depend smoothly on $x \in \mathbb{R}^d$, and $h > 0$ is a small parameter. We assume that the matrix field A is pointwise symmetric and positive semidefinite and the function c is nonnegative. In stochastic analysis, such operators arise naturally as the generators of time homogeneous Langevin processes

$$dX_t = \xi(X_t) + \sqrt{2h\sigma(X_t)} dB_t, \quad (1.2)$$

where (B_t) denotes the Brownian motion on \mathbb{R}^d , the vector field ξ is the drift coefficient, the matrix field σ is the diffusion coefficient and the parameter h is proportional

Jean-François Bony: Institut Mathématique de Bordeaux, Université de Bordeaux, 33405 Talence, France; jean-francois.bony@math.u-bordeaux.fr

Dorian Le Peutrec: Laboratoire de Mathématiques Jean Leray, Université de Nantes, 92208 Nantes, France; dorian.lepeutrec@univ-nantes.fr

Laurent Michel: Institut Mathématique de Bordeaux, Université de Bordeaux, 33405 Talence, France; laurent.michel@math.u-bordeaux.fr

Mathematics Subject Classification 2020: 35P05 (primary); 35P20, 60J60 (secondary).

to the temperature of the system. Given any test function φ , the expectation $u(t, x) = \mathbb{E}(\varphi(X_t) \mid X_0 = x)$ solves the Fokker–Planck equation

$$\begin{cases} \partial_t u + \mathcal{L}u = 0, \\ u|_{t=0} = \varphi, \end{cases} \quad (1.3)$$

where

$$\mathcal{L} = -h \sum_{i,j=1}^d a_{i,j} \partial_{x_i} \partial_{x_j} - \sum_{k=1}^d \xi_k \partial_{x_k},$$

with $A = (a_{i,j}) = \sigma \sigma^t$. Up to the multiplicative factor h , this operator has the form (1.1) for some suitable b and c . Denoting by \mathcal{L}^\dagger the formal $L^2(dx)$ adjoint of \mathcal{L} , (1.3) is equivalent to saying that the probability density $\varrho(t, \cdot)$ of the process (X_t) is a solution of the adjoint equation

$$\partial_t \varrho = \mathcal{L}^\dagger \varrho.$$

Among many examples of such operators, let us mention two cases of particular interest.

Taking $\xi = -\nabla f$ for some smooth function f on \mathbb{R}^d and $\sigma = \text{Id}_{\mathbb{R}^d}$, the generator \mathcal{L} of the overdamped Langevin process

$$dX_t = -\nabla f(X_t) + \sqrt{2h} dB_t \quad (1.4)$$

can be written as

$$\mathcal{L} = \mathcal{L}_{\text{KS}} = -h\Delta + \nabla f \cdot \nabla, \quad (1.5)$$

which is sometimes called the Kramers–Smoluchowski operator. Depending on the field of research, this operator is also known as the weighted Laplacian or Bakry–Émery Laplacian and is unitarily equivalent to the Witten Laplacian.

Another famous example is given by the case where $\sigma : \mathbb{R}^{2d} \rightarrow \mathbb{R}^{2d}$ is the projection onto the subspace $0 \oplus \mathbb{R}^d$, $\sigma(x, v) = (0, v)$, and $\xi : \mathbb{R}^{2d} \rightarrow \mathbb{R}^{2d}$, defined by $\xi(x, v) = (v, -\nabla_x V - v)$, is related to the energy function $f(x, v) = \frac{1}{2}|v|^2 + V(x)$ depending on a smooth potential V on \mathbb{R}^d . The associated Langevin equation reads

$$\begin{cases} dx_t = v_t dt, \\ dv_t = -\nabla_x V(x_t) dt - v_t dt + \sqrt{2h} dB_t, \end{cases} \quad (1.6)$$

where (B_t) is the Brownian motion on \mathbb{R}^d . The associated generator is the Kramers–Fokker–Planck operator

$$\mathcal{L} = \mathcal{L}_{\text{KFP}} = -v \cdot \nabla_x + \nabla_x V \cdot \nabla_v + v \cdot \nabla_v - h\Delta_v, \quad (1.7)$$

where Δ_v is the Laplace operator in the v variable only.

The study of the operators \mathcal{L}_{KS} and \mathcal{L}_{KFP} has been the subject of many works in the last decades. It is particularly motivated by its applications to computational physics. The above processes are indeed ergodic with respect to their Gibbs measure and can thus be used to sample this distribution. We refer to [22] for details on these topics.

From a theoretical point of view, the study of the qualitative properties (well-posedness, asymptotic behavior) as well as of the quantitative properties (precise spectral asymptotics) of the Fokker–Planck equation (1.3) has recently seen a major progress on the impulse of microlocal techniques. When the matrix field A is positive definite, the operator P is elliptic and standard tools apply to prove general properties on the operator P (maximal accretivity, compactness of the resolvent). When A is not invertible, the operator P is not elliptic anymore (it is sometimes called a degenerate diffusion) and advances have recently been made using hypoelliptic methods in the spirit of Hörmander. For the Kramers–Fokker–Planck operator \mathcal{L}_{KFP} , exponential convergence to equilibrium was proved in [26] and explicit rate of decay in the nonsemiclassical setting was given in [16]. More generally, hypocoercive methods developed for various kinetic models provide now robust tools to prove return to equilibrium and spectral gap estimates (see [27] for an overview).

In the semiclassical setting $h \rightarrow 0$, computing sharp spectral asymptotics for the low-lying spectrum of P is a classical problem having a long history. In the elliptic self-adjoint case, i.e. when A is uniformly positive definite and $b = 0$, the low-lying eigenvalues of P are localized near the absolute minimum value of the zeroth order part of P , that is, the minimum value of the function c . Moreover, the harmonic and WKB approximations of P near the absolute minima of c yield spectral expansions in powers of h of the low-lying eigenvalues of P (see [7, Chapters 3 and 4] for a detailed study in the case of Schrödinger operators).

However, in certain situations, these expansions are identical, and thus do not permit one to discriminate between these low-lying eigenvalues. This is for instance the case for Witten Laplacians associated with a confining Morse function f (in this case, the corresponding function c also depends on h), for which we know from the early works of Witten [28] and Helffer–Sjöstrand [10] that P admits exponentially small eigenvalues (that is, of order $\mathcal{O}(e^{-C/h})$ for some $C > 0$, and hence indistinguishable on the basis of their expansions in powers of h), in one-to-one correspondence with the minima of f , and that the rest of the spectrum is above $\varepsilon_* h$ for some $\varepsilon_* > 0$.

Up to a unitary conjugation, the Witten Laplacian is nothing other than the Kramers–Smoluchowski operator (1.5) and its small eigenvalues govern the time of return to equilibrium for the overdamped Langevin process (1.4). The computation of these eigenvalues is a historical problem which goes back at least to Eyring [8] and Kramers [18]. In the early 2000s, sharp asymptotics of these small eigenvalues were obtained by Bovier–Eckhoff–Gayraud–Klein [4] and Helffer–Klein–Nier [9]. Known as Eyring–Kramers formulas, such spectral asymptotics were also obtained recently in [20] in elliptic non-self-adjoint settings, associated with nonreversible overdamped Langevin processes generalizing (1.4). Concerning the transition times of these processes, Eyring–Kramers formulas have been established in both reversible and nonreversible settings in [3] and [2, 19, 21] respectively under similar assumptions. We also refer to [1] for a nice introduction to these topics.

In the nonelliptic case, major progress in the analysis of the operator P was made by Hérau, Hitrik and Sjöstrand in a series of works. In [13], the authors proved resolvent

estimates and established harmonic approximation of the spectrum under dynamical assumptions on the symbol of the operator P . In [14], they applied these results to operators satisfying additional symmetries (supersymmetry and PT-symmetry). Under these assumptions, the operator P admits a natural Gibbs stationary distribution $e^{-f/h}$ and the authors proved spectral Eyring–Kramers formulas in this setting, where the small eigenvalues govern the time of return to equilibrium for the Langevin process (1.6).

Though they are satisfied by many interesting examples (like Kramers–Fokker–Planck operators), the assumptions of supersymmetry and PT-symmetry do not look necessary to prove sharp spectral asymptotics, as shown in [20]. The aim of the present paper is to prove spectral Eyring–Kramers formulas for general operators P satisfying the assumptions of [13] and admitting an explicit Gibbs stationary distribution, but without the additional symmetry assumptions of [14].

1.2. Framework and results

Let $P = P(x, h\partial_x, h)$ be a semiclassical second order differential operator on \mathbb{R}^d , $d \geq 1$, with smooth real coefficients,

$$P = - \sum_{i,j=1}^d h\partial_{x_i} \circ a_{i,j}(x, h) \circ h\partial_{x_j} + \frac{1}{2} \sum_{j=1}^d (b_j(x, h) \circ h\partial_{x_j} + h\partial_{x_j} \circ b_j(x, h)) + c(x, h), \quad (1.8)$$

where the real functions $a_{i,j}$, b_j , and c depend smoothly on $x \in \mathbb{R}^d$, $a_{i,j} = a_{j,i}$ for all $i, j = 1, \dots, d$, and $h \in]0, 1]$ denotes the semiclassical parameter. We suppose that the coefficients of P satisfy the following growth condition at infinity:

$$\begin{aligned} \forall |\alpha| \geq 0, \quad \partial_x^\alpha a_{i,j}(x, h) &= \mathcal{O}(1), \\ \forall |\alpha| \geq 1, \quad \partial_x^\alpha b_j(x, h) &= \mathcal{O}(1), \\ \forall |\alpha| \geq 2, \quad \partial_x^\alpha c(x, h) &= \mathcal{O}(1), \end{aligned} \quad (1.9)$$

uniformly with respect to h , where the $\mathcal{O}(1)$'s may depend on α . We assume moreover that the above coefficients admit classical expansions $a_{i,j}(x, h) \sim \sum_{k \in \mathbb{N}} h^k a_{i,j}^k(x)$, $b_j(x, h) \sim \sum_{k \in \mathbb{N}} h^k b_j^k(x)$ and $c(x, h) \sim \sum_{k \in \mathbb{N}} h^k c^k(x)$ in the sense that

$$\partial_x^\alpha \left(e(x, h) - \sum_{0 \leq k \leq K} e^k(x) h^k \right) = \mathcal{O}(h^{K+1}) \quad (1.10)$$

for all $\alpha \in \mathbb{N}^d$, $K \in \mathbb{N}$ and $e \in \{a_{i,j}, b_j, c\}$. This yields classical expansions for the matrix field $A(x, h) = (a_{i,j}(x, h))_{i,j} \sim \sum_k h^k A^k(x)$ and the vector field $b(x, h) \sim \sum_k h^k b^k(x)$. Considering symbols which have a classical expansion allows us to deal with, e.g., Witten Laplacians and Kramers–Fokker–Planck operators, which naturally have subprincipal

terms. Finally, we also assume partial positivity of the symbols of the operator P : for all $x \in \mathbb{R}^d$ and $h \in]0, 1]$,

$$c^0(x) \geq 0 \quad \text{and} \quad A(x, h) = (a_{i,j}(x, h))_{i,j} \text{ is positive semidefinite.} \quad (1.11)$$

Such operators were studied in detail in [13], where the authors establish resolvent estimates together with spectral asymptotics. In particular, the graph closure of the operator P initially defined on the Schwartz space $\mathcal{S}(\mathbb{R}^d)$, still denoted by P , is maximal quasi-accretive (more precisely, $P + Ch$ is maximal accretive for some $C > 0$) and has domain

$$\mathcal{D}(P) = \{u \in L^2(\mathbb{R}^d); Pu \in L^2(\mathbb{R}^d)\}, \quad (1.12)$$

by [13, Proposition 3.1 and Corollary 3.2]. The same properties also hold for the adjoint P^* mutatis mutandis. Throughout the paper, we assume (1.8)–(1.11) without explicit mention.

Let us introduce the symbol $p(x, \xi, h)$ of P in the semiclassical Weyl quantization. It satisfies

$$p(x, \xi, h) = \xi \cdot A(x, h)\xi + i b(x, h) \cdot \xi + c(x, h) + \frac{h^2}{4} \sum_{i,j=1}^d \partial_{x_i} \partial_{x_j} a_{i,j}(x, h), \quad (1.13)$$

where, throughout the paper, $x \cdot y$ denotes the usual scalar product of x and y in \mathbb{R}^d (in order to facilitate the reading, we will also sometimes use the notation $\langle x, y \rangle = x \cdot y$). The symbol admits a classical expansion $p(x, \xi, h) \sim \sum_k h^k p^k(x, \xi)$ and the principal symbol p^0 is given by $p^0 = p_2^0 + i p_1^0 + p_0^0$ with $p_2^0(x, \xi) = \xi \cdot A^0(x)\xi$, $p_1^0(x, \xi) = b^0(x) \cdot \xi$ and $p_0^0(x) = c^0(x)$. In order to lighten the notation, we will drop the superscript 0 when it is unambiguous. Consider the symbol

$$\tilde{p}(x, \xi) = p_0^0(x) + \frac{p_2^0(x, \xi)}{\langle \xi \rangle^2}, \quad (1.14)$$

where $\langle \xi \rangle^2 = 1 + |\xi|^2$. Thanks to (1.11), one has $p_0^0, p_2^0 \geq 0$ and hence $\tilde{p} \geq 0$. Given $T > 0$, let us define the symbol $\langle \tilde{p} \rangle_T$ by

$$\langle \tilde{p} \rangle_T = \frac{1}{2T} \int_{-T}^T \tilde{p} \circ e^{tH_{p_1^0}} dt, \quad (1.15)$$

where $H_{p_1^0} = \partial_\xi p_1^0 \cdot \partial_x - \partial_x p_1^0 \cdot \partial_\xi$ denotes the Hamilton vector field associated with the symbol p_1^0 . The critical set associated with p is defined by

$$\mathcal{C} = \{(x, 0) \in T^*\mathbb{R}^d; b^0(x) = 0 \text{ and } c^0(x) = 0\}. \quad (1.16)$$

As in [13], we suppose that the set \mathcal{C} is finite, $\mathcal{C} = \{\rho_1, \dots, \rho_N\}$, and for some fixed $T > 0$ (see [13, (4.21) and (4.23)]) there exists a constant $C > 0$ such that

$$\langle \tilde{p} \rangle_T(\rho) \geq \frac{1}{C} |\rho - \rho_j|^2 \quad \text{for all } \rho \text{ near any } \rho_j, \quad (\text{Harmo})$$

and, for any neighborhood U of $\pi_x \mathcal{C}$,

$$\exists C > 0, \forall x \in \mathbb{R}^d \setminus U, \quad \text{meas} \left\{ t \in [-T, T]; c^0(\exp(tb^0 \cdot \nabla)(x)) \geq \frac{1}{C} \right\} \geq \frac{1}{C}. \quad (\text{Hypo})$$

Assumption **(Harmo)** may look difficult to check in applications, but Corollary 2.4 and Remark 2.5 provide a concrete criterion to verify it. Observe also that **(Hypo)** holds true for instance when c^0 is uniformly positive outside each neighborhood of $\pi_x \mathcal{C}$. Note that it is not necessary to assume (4.22) of [13] here since this is a consequence of **(Harmo)** and **(Hypo)**, as explained in [13, p. 223].

Under these assumptions, Hérau, Hitrik and Sjöstrand obtained spectral information that we sum up below. For this purpose, we introduce the fundamental matrix F_{p^0} of the symbol p^0 at a critical point $\rho \in \mathcal{C}$ (see (1.16)) as the linearization of the Hamilton field H_{p^0} at ρ . Its eigenvalues are of the form $\pm \lambda_{\rho, \ell}$, $1 \leq \ell \leq d$, with $\text{Im } \lambda_{\rho, \ell} > 0$. We use the notation

$$\widetilde{\text{tr}}(p, \rho) = -i \sum_{\ell=1}^d \lambda_{\rho, \ell} + 2c^1(\pi_x \rho).$$

The following results from [13, Proposition 7.2, Theorems 8.3, 8.4].

Theorem 1 (Hérau–Hitrik–Sjöstrand). *Assume that **(Harmo)** and **(Hypo)** hold true. For any $B > 0$, there exists $C > 0$ such that for h small enough, the operator P has no spectrum in*

$$\{z \in \mathbb{C}; \text{Re } z < Bh \text{ and } |\text{Im } z| > Ch\}.$$

Moreover, for any $B > 0$, there exists $\alpha > 0$ such that, for h small enough, the spectrum of P in $D(0, Bh)$ is discrete and made up of eigenvalues (with multiplicity) of the form

$$\mu_{\rho, k}(h) = h(\mu_{\rho, k}^0 + \mathcal{O}(h^\alpha)),$$

where the $(\mu_{\rho, k}^0)_{\rho \in \mathcal{C}, k \in \mathbb{N}}$ are all the possible numbers of the form

$$\mu_{\rho, k}^0 = \frac{1}{i} \sum_{\ell=1}^d \nu_{\rho, k, \ell} \lambda_{\rho, \ell} + \frac{1}{2} \widetilde{\text{tr}}(p, \rho) \quad \text{with } \nu_{\rho, k, \ell} \in \mathbb{N}.$$

Finally, for every $B > 0$, there exists $C > 0$ such that

$$\|(P - z)^{-1}\| \leq C/h \quad (1.17)$$

for all h small enough and $z \in \mathbb{C}$ such that $\text{Re } z < Bh$ and $\text{dist}(z, \sigma(P)) \geq h/B$.

In addition, they showed that the remainder terms $\mathcal{O}(h^\alpha)$ have a classical expansion in fractional powers of h . It is assumed in [13] that the coefficients $a_{i,j}, b_j, c$ of the operator P (see (1.8)) do not depend on h , but a direct adaptation to our setting (using in particular (1.10)) gives Theorem 1. It turns out that in many situations, some leading coefficients $\mu_{\rho, k}^0$ vanish and one aims at having a more accurate description of the spectrum. This is the case for instance for Witten Laplacians or Kramers–Fokker–Planck operators,

which both admit an invariant distribution. In the present paper, we consider the situation where the operator P satisfies the assumptions of Theorem 1 and there exists a smooth function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ such that

$$e^{-f/h} \in L^2(\mathbb{R}^d), \quad \lim_{|x| \rightarrow +\infty} f(x) = +\infty, \quad (\text{Confin})$$

with

$$P(e^{-f/h}) = 0 \quad \text{and} \quad P^*(e^{-f/h}) = 0, \quad (\text{Gibbs})$$

in the sense of distributions. In particular, (1.12) shows that $e^{-f/h} \in \mathcal{D}(P) \cap \mathcal{D}(P^*)$. Moreover, we will assume that

$$f \text{ is a Morse function with a finite number of critical points.} \quad (\text{Morse})$$

We denote by \mathcal{U} the set of critical points of the Morse function f and by $\mathcal{U}^{(j)}$ the set of its critical points of index $j = 0, \dots, d$ (that is, the set of those critical points \mathbf{u} such that $\text{Hess } f(\mathbf{u})$ has signature $(d - j, j)$). Moreover, we denote by $n_0 := \#\mathcal{U}^{(0)}$ the number of local minimum points of f , by $H(x) := \text{Hess } f(x)$ the Hessian matrix of f at $x \in \mathbb{R}^d$, and we call the elements of $\mathcal{U}^{(1)}$ saddle points.

For $j \in \{0, 1, 2\}$, let us denote by P_j the j th order part of the operator $P = P_2 + P_1 + P_0$ with $P_2 = -h \operatorname{div} \circ A \circ h \nabla$ formally self-adjoint, $P_1 = \frac{1}{2}(b \cdot h \nabla + h \operatorname{div} \circ b)$ formally anti-adjoint, and $P_0 = c$ formally self-adjoint. It then follows from (Gibbs) that $P_1(e^{-f/h}) = 0$ and $(P_2 + P_0)(e^{-f/h}) = 0$. Using the classical expansions of the coefficients and looking at the terms of order 0 in the expansion, we obtain the following eikonal equations: for all $x \in \mathbb{R}^d$,

$$\langle A^0(x) \nabla f(x), \nabla f(x) \rangle_{\mathbb{R}^d} = c^0(x), \quad (1.18)$$

$$\langle b^0(x), \nabla f(x) \rangle_{\mathbb{R}^d} = 0. \quad (1.19)$$

The first consequence of these identities is the following lemma whose proof is postponed to the next section.

Lemma 1.1. *If (Gibbs) and (Morse) hold true, then $\mathcal{U} \times \{0\} \subset \mathcal{C}$. If in addition (Harmo) is satisfied, then $\mathcal{C} = \mathcal{U} \times \{0\}$.*

Using this lemma, we obtain our first localization result for the spectrum of P . Its proof will also be given in the next section. Note that the eigenvalues are counted according to their algebraic multiplicities throughout the paper.

Proposition 1.2. *Assume the hypotheses of Theorem 1, (Gibbs) and (Morse). There exist $h_0, \varepsilon_* > 0$ such that, for every $h \in]0, h_0]$, P has exactly n_0 eigenvalues in $\{z \in \mathbb{C}; \operatorname{Re} z < \varepsilon_* h\}$. Moreover, these eigenvalues are of order $\mathcal{O}(h^{1+\alpha})$, where $\alpha > 0$ is given by Theorem 1.*

The aim of this paper is to give sharp asymptotics of these n_0 small eigenvalues of P . For this purpose, we recall the general labeling of minimum points introduced in [9] and generalized in [14]. The presentation below originates from [20, 24], where extra material can be found. The main ingredient is the notion of separating saddle point which is defined

as follows. Note that, for a saddle point $\mathbf{s} \in \mathcal{U}^{(1)}$ of f and $r > 0$ small enough, the set

$$\{x \in D(\mathbf{s}, r); f(x) < f(\mathbf{s})\}$$

has exactly two connected components $C_j(\mathbf{s}, r)$, $j = 1, 2$. Observe that this set is empty when $\mathbf{s} \in \mathcal{U}^{(0)}$ and connected when $\mathbf{s} \in \mathbb{R}^d \setminus (\mathcal{U}^{(0)} \cup \mathcal{U}^{(1)})$. The following definition comes from [14, Definition 4.1].

Definition 1.3. We say that $\mathbf{s} \in \mathcal{U}^{(1)}$ is a separating saddle point of f if, for every $r > 0$ small enough, $C_1(\mathbf{s}, r)$ and $C_2(\mathbf{s}, r)$ are contained in two different connected components of $\{x \in \mathbb{R}^d; f(x) < f(\mathbf{s})\}$. We will denote by $\mathcal{V}^{(1)}$ the set of these points.

We say that $\sigma \in \mathbb{R}$ is a separating saddle value of f if $\sigma = f(\mathbf{s})$ for some $\mathbf{s} \in \mathcal{V}^{(1)}$.

We say that $E \subset \mathbb{R}^d$ is a critical component of f if there exists $\sigma \in f(\mathcal{V}^{(1)})$ such that E is a connected component of $\{f < \sigma\}$ and $\partial E \cap \mathcal{V}^{(1)} \neq \emptyset$.

Let us now describe the labeling procedure of [14]. Assume that $f(x) \rightarrow +\infty$ when $|x| \rightarrow +\infty$ and that f satisfies (Morse). The set $f(\mathcal{V}^{(1)})$ is then finite. We denote by $\sigma_2 > \sigma_3 > \dots > \sigma_N$ its elements and, for convenience, we also introduce a fictive infinite saddle value $\sigma_1 = +\infty$. Starting from σ_1 , we will recursively associate to each σ_i a finite family $(\mathbf{m}_{i,j})_j$ of local minimum points and a finite family $(E_{i,j})_j$ of critical components:

- Let $X_{\sigma_1} = \{x \in \mathbb{R}^d; f(x) < \sigma_1 = +\infty\} = \mathbb{R}^d$. We let $\mathbf{m}_{1,1}$ be any global minimum point of f (not necessarily unique) and $E_{1,1} = \mathbb{R}^d$. In the following, we will write $\underline{\mathbf{m}} = \mathbf{m}_{1,1}$.
- Next, we consider $X_{\sigma_2} = \{x \in \mathbb{R}^d; f(x) < \sigma_2\}$. It is the union of finitely many connected components. Exactly one of these components contains $\mathbf{m}_{1,1}$ and the other components are denoted by $E_{2,1}, \dots, E_{2,N_2}$. They are all critical and, in each component $E_{2,j}$, we pick up a point $\mathbf{m}_{2,j}$ which is a global minimum point of $f|_{E_{2,j}}$.
- Suppose now that the families $(\mathbf{m}_{k,j})_j$ and $(E_{k,j})_j$ have been constructed until rank $k = i - 1$. The set $X_{\sigma_i} = \{x \in \mathbb{R}^d; f(x) < \sigma_i\}$ has again finitely many connected components and we label $E_{i,j}$, $j = 1, \dots, N_i$, those of them that do not contain any $\mathbf{m}_{k,\ell}$ with $k < i$. They are all critical and, in each $E_{i,j}$, we pick up a point $\mathbf{m}_{i,j}$ which is a global minimum of point $f|_{E_{i,j}}$.

At the end of this procedure, all the minimum points have been labeled (see Fig. 1.1).

We now recall some useful constructions of [20, 24]. Throughout, we denote by \mathbf{s}_1 a fictive saddle point such that $f(\mathbf{s}_1) = \sigma_1 = +\infty$ and, for any set A , $\mathcal{P}(A)$ denotes the power set of A . From the above labeling, we define two mappings

$$E : \mathcal{U}^{(0)} \rightarrow \mathcal{P}(\mathbb{R}^d) \quad \text{and} \quad \mathbf{j} : \mathcal{U}^{(0)} \rightarrow \mathcal{P}(\mathcal{V}^{(1)} \cup \{\mathbf{s}_1\})$$

as follows: for every $i \in \{1, \dots, N\}$ and $j \in \{1, \dots, N_i\}$,

$$E(\mathbf{m}_{i,j}) := E_{i,j}, \tag{1.20}$$

and

$$\mathbf{j}(\underline{\mathbf{m}}) := \{\mathbf{s}_1\} \quad \text{and} \quad \mathbf{j}(\mathbf{m}_{i,j}) := \partial E_{i,j} \cap \mathcal{V}^{(1)} \text{ for } i \geq 2. \tag{1.21}$$

In particular, $E(\underline{\mathbf{m}}) = \mathbb{R}^d$ and, for all $i \in \{1, \dots, N\}$ and $j \in \{1, \dots, N_i\}$, one has $\emptyset \neq \mathbf{j}(\mathbf{m}_{i,j}) \subset \{f = \sigma_i\}$. We then define the mappings

$$\sigma : \mathcal{U}^{(0)} \rightarrow f(\mathcal{V}^{(1)}) \cup \{\sigma_1\} \quad \text{and} \quad S : \mathcal{U}^{(0)} \rightarrow]0, +\infty],$$

by

$$\forall \mathbf{m} \in \mathcal{U}^{(0)}, \quad \sigma(\mathbf{m}) := f(\mathbf{j}(\mathbf{m})) \quad \text{and} \quad S(\mathbf{m}) := \sigma(\mathbf{m}) - f(\mathbf{m}), \quad (1.22)$$

where, with a slight abuse of notation, we have identified the set $f(\mathbf{j}(\mathbf{m}))$ with its unique element. Note that $S(\mathbf{m}) = +\infty$ if and only if $\mathbf{m} = \underline{\mathbf{m}}$.




Fig. 1.1. An example of the labeling procedure.

We are now in a position to introduce our last assumption. In addition to (Gibbs), (Confin), and (Morse), we assume the following:

- ★ for any $\mathbf{m} \in \mathcal{U}^{(0)}$, \mathbf{m} is the unique global minimum point of $f|_{E(\mathbf{m})}$,
 - ★ for all $\mathbf{m} \neq \mathbf{m}'$ in $\mathcal{U}^{(0)}$, $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset$.
- (Gener)

In particular, (Gener) implies that f attains its global minimum only at $\underline{\mathbf{m}} \in \mathcal{U}^{(0)}$. This assumption is a generalization of [14, Assumption 5.1], which was already used in [20]. In Section 6, we discuss how to remove this hypothesis and deal with the general case, in the spirit of [24].

In order to state our main result, we need the following lemma which is proved in Section 2. Throughout the paper, we denote $\mathbb{C}_{\pm} = \{z \in \mathbb{C}; \pm \operatorname{Re} z > 0\}$, and M^t is the transpose of any matrix M .

Lemma 1.4. Assume (Harmo), (Gibbs), and (Morse), and let $k \in \{0, \dots, d\}$. Let $\mathbf{u} \in \mathcal{U}^{(k)}$ be a critical point of f of index k . Denote $B(\mathbf{u}) = db^0(\mathbf{u})$ and recall that $H(\mathbf{u})$ is the Hessian matrix of f at \mathbf{u} . Then

- (i) the matrix $\Lambda(\mathbf{u}) := 2H(\mathbf{u})A^0(\mathbf{u}) + B^t(\mathbf{u})$ admits exactly k eigenvalues in \mathbb{C}_{-} and $d - k$ eigenvalues in \mathbb{C}_{+} ,
- (ii) if $k = 1$, its unique eigenvalue $\mu(\mathbf{u})$ in \mathbb{C}_{-} is real (and thus $\mu(\mathbf{u}) < 0$).

We can now state our main result.

Theorem 2. *Suppose that the assumptions of Theorem 1 are satisfied. Assume also that (Gibbs), (Confin), (Morse) and (Gener) hold true. Let ε_* be given by Proposition 1.2. There exists $h_0 > 0$ such that, for all $h \in]0, h_0]$, counting the eigenvalues with multiplicity, one has*

$$\sigma(P) \cap \{\operatorname{Re} z < \varepsilon_* h\} = \{\lambda(\mathbf{m}, h); \mathbf{m} \in \mathcal{U}^{(0)}\},$$

where $\lambda(\underline{\mathbf{m}}, h) = 0$ and, for all $\mathbf{m} \neq \underline{\mathbf{m}}$, $\lambda(\mathbf{m}, h)$ satisfies the Eyring–Kramers type formula

$$\lambda(\mathbf{m}, h) = z(\mathbf{m}) h e^{-2S(\mathbf{m})/h} a(h), \quad (1.23)$$

where $a(h) \in \mathbb{C}$ admits a classical expansion $a(h) \sim 1 + \sum_{j \geq 1} a_j h^j$ with every a_j real. Here, $S : \mathcal{U}^{(0)} \rightarrow]0, +\infty]$ is defined in (1.22) and, for every $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$,

$$z(\mathbf{m}) = \frac{(\det \operatorname{Hess} f(\mathbf{m}))^{1/2}}{2\pi} \left(\sum_{\mathbf{s} \in \mathbf{j}(\mathbf{m})} \frac{|\mu(\mathbf{s})|}{|\det \operatorname{Hess} f(\mathbf{s})|^{1/2}} \right), \quad (1.24)$$

where $\mathbf{j} : \mathcal{U}^{(0)} \rightarrow \mathcal{P}(\mathcal{V}^{(1)} \cup \{\mathbf{s}_1\})$ is defined in (1.21) and $\mu(\mathbf{s})$ is given by Lemma 1.4.

Let us make some comments on the above result. In [14], the authors studied the case where the operator satisfies some supersymmetry property. More precisely, they assumed the existence of a smooth matrix-valued function $G(x)$ such that $P = \Delta_{f,G}$, where

$$\Delta_{f,G} = d_f^* \circ G(x) \circ d_f, \quad (1.25)$$

and d_f denotes the semiclassical Hodge derivative twisted “à la Witten”: $d_f = e^{-f/h} \circ h d \circ e^{f/h}$ for some smooth function f . Under suitable assumptions on f and G , $\Delta_{f,G}$ satisfies the general hypotheses of Theorem 1. Moreover, one has obviously $\Delta_{f,G}(e^{-f/h}) = \Delta_{f,G}^*(e^{-f/h}) = 0$. Assuming additionally that f is a Morse function, the authors proved that the smallest eigenvalues of $\Delta_{f,G}$ are exponentially small with respect to h , and established Eyring–Kramers type formulas under suitable topological assumptions (see [14, Theorem 5.10, Proposition 6.7 and (6.71)]). In their paper, the supersymmetry assumption is fundamental since, combined with the PT-symmetry property, it permits following the strategy used by Helffer–Klein–Nier [9] in the supersymmetric framework of Witten Laplacians. More recently, the last two authors [20] studied the case of nonreversible diffusions

$$P = \Delta_f + b \cdot d_f, \quad (1.26)$$

where $\Delta_f = -h^2 \Delta + |\nabla f|^2 - h \Delta f$ denotes the Witten Laplacian and b is a vector field satisfying $\operatorname{div} b = 0$ and $b \cdot \nabla f = 0$. In this setting, which is not supersymmetric in general, the authors built accurate quasimodes and used them to prove Eyring–Kramers asymptotics.

The interest of the approach developed in the present paper is in dealing simultaneously with all these models without assuming additional symmetry. In particular, Theorem 2 applies to both (1.25) and (1.26). Moreover, in Appendix B we give examples of operators P satisfying our assumption but which do not enjoy a nice supersymmetric structure (1.25). Compared to the results of [14], our approach also has the advantage of

giving formulas which are completely explicit in terms of the coefficients of the operator and of the function f . Moreover, compared to the result of [20], we obtain a full asymptotic expansion of the prefactor $z(\mathbf{m})a(h)$. The proof relies on the resolvent estimates of [13] and on the construction of accurate quasimodes near the saddle points of f . These constructions, inspired by [3, 6, 20], are the main novelty of this paper. To be more precise, we generalize the use of Gaussian cut-off functions introduced in [20] by using geometric constructions, which lead to complete asymptotic expansions.

Theorem 2 permits us to determine the long time behavior of the solutions of the evolution equation associated with P ,

$$\begin{cases} h\partial_t u + Pu = 0, \\ u|_{t=0} = u_0. \end{cases} \quad (1.27)$$

Under our assumptions, P is better than maximal quasi-accretive. More precisely, we have the following.

Remark 1.5. If (Gibbs) holds true, then P is maximal accretive. Indeed, the same computation as the one leading to (5.2) below shows that

$$\forall u \in C_0^\infty(\mathbb{R}^d), \quad \operatorname{Re} \langle Pu, u \rangle = h^2 \langle e^{-f/h} A \nabla(e^{f/h} u), e^{-f/h} \nabla(e^{f/h} u) \rangle \geq 0.$$

Therefore the Cauchy problem (1.27) has, for every $u_0 \in L^2(\mathbb{R}^d)$, a unique weak solution in $C^0([0, +\infty[; L^2(\mathbb{R}^d))$, denoted by $u(t) = e^{-tP/h} u_0$. Modulo conjugation by $e^{-f/h}$, (1.27) is the Fokker–Planck equation (1.3) in the case of the general Langevin process (1.2).

We first state the spectral expansion of the propagator.

Corollary 1.6. *In the setting of Theorem 2, there exist $C, h_0, \varepsilon > 0$ such that, for all $h \in]0, h_0]$ and $u_0 \in L^2(\mathbb{R}^d)$, there exists $(u_{\lambda,n})_{\lambda,n} \subset L^2(\mathbb{R}^d)$ such that the solution $u(t)$ of (1.27) satisfies*

$$\forall t \geq 0, \quad \left\| u(t) - \sum_{\substack{\lambda \in \{\lambda(\mathbf{m}, h); \mathbf{m} \in \mathcal{U}^{(0)}\} \\ 0 \leq n \leq \operatorname{mult}(\lambda) - 1}} u_{\lambda,n} t^n e^{-\lambda t/h} \right\| \leq C e^{-\varepsilon t} \|u_0\|. \quad (1.28)$$

Moreover, for all $N \in \mathbb{N}$, there exist $C_N, h_0 > 0$ such that, for all $h \in]0, h_0]$ and $u_0 \in L^2(\mathbb{R}^d)$, the solution $u(t)$ of (1.27) satisfies

$$\forall t \geq 0, \quad \left\| u(t) - \frac{\langle e^{-f/h}, u_0 \rangle}{\|e^{-f/h}\|^2} e^{-f/h} \right\| \leq C_N e^{-t \min_{\mathbf{m} \neq \mathbf{m}} \operatorname{Re}(\lambda(\mathbf{m}, h))(1-h^N)/h} \|u_0\|. \quad (1.29)$$

The double sum appearing in (1.28) is nothing but $e^{-tP/h} \Pi_h$, where Π_h denotes the spectral projector of P associated with its n_0 exponentially small eigenvalues. In particular, when the $\lambda(\mathbf{m}, h)$ are pairwise distinct, (1.28) reads

$$u(t) = \sum_{\mathbf{m} \in \mathcal{U}^{(0)}} \Pi_{\mathbf{m}}(u_0) e^{-\lambda(\mathbf{m}, h)t/h} + \mathcal{O}(e^{-\varepsilon t}) \|u_0\|, \quad (1.30)$$

where $\Pi_{\mathbf{m}}$ is the rank-one spectral projector of P associated with the eigenvalue $\lambda(\mathbf{m}, h)$.

Estimate (1.29) implies that $u_{0,0} = \Pi_{\underline{\mathbf{m}}} u_0 = \|e^{-f/h}\|^{-2} \langle e^{-f/h}, u_0 \rangle e^{-f/h}$ and $\text{mult}(0) = 1$, where $\Pi_{\underline{\mathbf{m}}}$ is the (orthogonal) spectral projector of P associated with the eigenvalue 0 with the corresponding eigenspace $e^{-f/h} \mathbb{C}$. Equation (1.29) is a return-to-equilibrium formula generalizing [20, Theorem 1.11]. We see that the spectral gap (that is, $\min_{\mathbf{m} \neq \underline{\mathbf{m}}} \text{Re}(\lambda(\mathbf{m}, h))$) indeed gives the rate of convergence to the equilibrium state modulo $\mathcal{O}(h^\infty)$. In addition, when there exists precisely one $\mathbf{m}^* \in \mathcal{U}^{(0)} \setminus \{\mathbf{m}\}$ such that

$$\lambda(\mathbf{m}^*, h) = \min_{\mathbf{m} \neq \underline{\mathbf{m}}} \text{Re}(\lambda(\mathbf{m}, h))(1 + \mathcal{O}(h^\infty)),$$

the eigenvalue $\lambda(\mathbf{m}^*, h)$ is simple and real and $\min_{\mathbf{m} \neq \underline{\mathbf{m}}} \text{Re}(\lambda(\mathbf{m}, h))(1 - h^N)$ can be replaced by $\lambda(\mathbf{m}^*, h)$ in (1.29).

One can also show the metastable behavior of the solutions of (1.27).

Corollary 1.7 (Metastability). *In the setting of Theorem 2, let $S_1 < \dots < S_K = +\infty$ denote the increasing sequence of the $S(\mathbf{m})$'s defined in (1.22) and let Π_k^\leq be the spectral projector of P associated with its eigenvalues of modulus less than $e^{-2S_k/h}$. For two positive functions $g_\pm(h)$ such that $g_-(h) = \mathcal{O}(h^\infty)$ and $g_+^{-1}(h) = o(|\ln h|^{-1})$, we define*

$$t_0^+ = g_+(h) \quad \text{and} \quad \forall 1 \leq k \leq K, \quad t_k^\pm = g_\pm(h) e^{2S_k/h}$$

(in particular, $t_K^- = +\infty$). Then, for every h small enough, the solution $u(t)$ of (1.27) satisfies

$$\forall t_{k-1}^+ \leq t \leq t_k^-, \quad u(t) = \Pi_k^\leq u_0 + \mathcal{O}(h^\infty) \|u_0\|, \quad (1.31)$$

uniformly with respect to t , $1 \leq k \leq K$ and $u_0 \in L^2(\mathbb{R}^d)$.

In other words, $e^{-tP/h} \approx \Pi_k^\leq$ in the time interval $[t_{k-1}^+, t_k^-]$, whereas transitions occur around the times $t_k = e^{2S_k/h}$. In this result, one can take for instance $g_-(h) = e^{-\delta/h}$ with $\delta > 0$, and $g_+(h) = |\ln h|^2$. Note that $\Pi_j^\leq \Pi_k^\leq = \Pi_{\max(j,k)}^\leq$, $\Pi_1^\leq = \Pi_h$ is the spectral projector of P associated with its n_0 exponentially small eigenvalues, and $\Pi_K^\leq = \Pi_{\underline{\mathbf{m}}}$ is the orthogonal projector on $e^{-f/h} \mathbb{C}$. The proofs of Corollaries 1.6 and 1.7 are given at the end of Section 5.

We conclude this introduction by applying Theorem 2 to a generalization of the Kramers–Fokker–Planck operator defined in (1.7). For two smooth functions $V(x)$ and $W(v)$ and a friction coefficient $\gamma > 0$, the generator associated with the SDE

$$\begin{cases} dx_t = \partial_v W(v_t) dt, \\ dv_t = -\partial_x V(x_t) dt - \gamma \partial_v W(v_t) dt + \sqrt{2\gamma h} dB_t, \end{cases} \quad (1.32)$$

is given by

$$\mathcal{L} = -\partial_v W \cdot \partial_x + \partial_x V \cdot \partial_v + \gamma \partial_v W \cdot \partial_v - \gamma h \Delta_v. \quad (1.33)$$

This is an example of a Hamiltonian SDE discussed in [22, Section 2.2.3]. Defining $M_h(x, v) := e^{-(V(x) + W(v))/h}$, we have

$$\mathcal{L}(1) = 0 \quad \text{and} \quad \mathcal{L}^\dagger(M_h) = 0.$$

Hence, $P := M_h^{-1/2} h \mathcal{L}^\dagger M_h^{1/2}$ satisfies

$$P(e^{-f/h}) = 0 \quad \text{and} \quad P^*(e^{-f/h}) = 0,$$

where $f(x, v) := (V(x) + W(v))/2$. Moreover, an immediate computation shows that

$$P = \partial_v W \cdot h \partial_x - \partial_x V \cdot h \partial_v + \gamma \Delta_{W/2}, \quad (1.34)$$

where $\Delta_{W/2} = (-h \partial_v + \partial_v W/2) \circ (h \partial_v + \partial_v W/2)$ is the Witten Laplacian in the variable v associated with the function $W(v)/2$. Thus, P has the form (1.8) with

$$A(x, v) = \begin{pmatrix} 0 & 0 \\ 0 & \gamma \text{Id} \end{pmatrix}, \quad b(x, v) = \begin{pmatrix} \partial_v W(v) \\ -\partial_x V(x) \end{pmatrix}, \quad c(x, v) = \frac{\gamma}{4} |\nabla_v W|^2 - h \frac{\gamma}{2} \Delta W.$$

Of course, f satisfies (Morse) if and only if V and W do. In that case, Corollary 2.4, Remark 2.5 and

$$B := db^0 = \begin{pmatrix} 0 & \text{Hess}_v W \\ -\text{Hess}_x V & 0 \end{pmatrix}$$

show that (Harmo) is satisfied. Under some additional growth assumptions on V and W at infinity, one can verify that (Confin) and (Hypo) hold true. At a critical point \mathbf{u} , the matrix $\Lambda(\mathbf{u})$ of Lemma 1.4 is given by

$$\Lambda(\mathbf{u}) = \begin{pmatrix} 0 & -\text{Hess}_x V \\ \text{Hess}_v W & \gamma \text{Hess}_v W \end{pmatrix}.$$

The setting of (1.7) corresponds to $W(v) = v^2/2$. In that case, the saddle points of f are of the form $\mathbf{s} = (\mathbf{x}, 0)$, where \mathbf{x} is a saddle point of V , and the unique eigenvalue of $\Lambda(\mathbf{s})$ with negative real part is

$$\mu(\mathbf{s}) = \frac{1}{2}(\gamma - \sqrt{\gamma^2 - 4\lambda_1}), \quad (1.35)$$

where λ_1 is the unique negative eigenvalue of $\text{Hess}_x V(\mathbf{x})$. This yields an explicit expression of the prefactor $z(\mathbf{m})$ in (1.24) and we observe that it has the same form as the one obtained in [14, (6.67)]. Observe also that $|\mu(\mathbf{s})| < |\lambda_1|$ if and only if $\gamma > 1 + \lambda_1$. Thus, the rate of convergence to the equilibrium for the Langevin process (1.6) (whose generator is the Kramers–Fokker–Planck operator, see (1.7)) is smaller than for the overdamped Langevin process (1.4) (generated by the Witten Laplacian, see the lines above (1.5)) if and only if this condition holds. Note that this discussion is more generally valid if W admits a unique critical point at $v = \mathbf{v}$ with $\text{Hess}_v W(\mathbf{v}) = \text{Id}$. In particular, it is easy to choose W such that P is not PT-symmetric, which provides a setting covered by Theorem 2 but which cannot be treated using [14].

More sophisticated Langevin equations have been considered in the literature, as the generalized Langevin equation in [25] (see also references therein). Our results enable one to compute in the low temperature regime the low-lying eigenvalues of the different generators treated in [25].

The rest of this paper is organized as follows. In Section 2, we derive algebraic and geometric results from our assumptions. This leads to the proofs of Lemmas 1.1 and 1.4,

and of the rough localization of the spectrum of P stated in Proposition 1.2. Section 3 is devoted to the construction of new Ansätze of the eigenmodes of P near the saddle points of f . These Ansätze are then used in Section 4 to construct global quasimodes. Afterwards, in Section 5, we prove our main results, Theorem 2 and Corollaries 1.6 and 1.7. The aim of Section 6 is then to show that Theorem 2 can be generalized to an arbitrary Morse function f , that is, without assuming (Gener). A vague statement is given in Theorem 3, while precise ones are Theorems 5 and 6. Let us also recall here that we are not working with symmetry assumptions such as supersymmetry or PT-symmetry. This prevents us from concluding as in [9] (and in many other later works such as, e.g., [14, 24]) after reduction of the problem to the computation of the eigenvalues of a square matrix of size n_0 . To handle this computation, we then make crucial use of the Schur complement method, as in [20], and refine [20, Theorem A.4] in Theorem 4. We believe that Theorem 4 is of independent interest and may be used in other contexts.

2. Preliminary results

In this section, we prove some preparatory geometric results and use them to show the rough localization of the spectrum of P stated in Proposition 1.2. For a critical point \mathbf{u} of f , we use the abbreviations

$$A^0 = A^0(\mathbf{u}), \quad B = db^0(\mathbf{u}), \quad H = \text{Hess } f(\mathbf{u}).$$

2.1. Analysis of the critical set

The aim of this section is to prove Lemma 1.1 and to discuss the assumptions of Section 1.2. We start with a microlocal observation.

Lemma 2.1. (i) Assume (Harmo) and let $(\mathbf{u}, 0) \in \mathcal{C}$. Then

$$\text{the real symmetric matrix } \int_{-T}^T e^{-tB} A^0 e^{-tB^t} dt \text{ is positive definite.} \quad (2.1)$$

(ii) If (2.1) holds true, then $\ker(A^0) \cap \ker(B^t - z) = \{0\}$ for every $z \in \mathbb{C}$.

Proof. Since the Hamilton vector field $H_{p_1^0}$ at (\mathbf{u}, ξ) equals $(0, -B^t \xi)$ when $\xi \in \mathbb{R}^d$, it follows that, for every $\xi \in \mathbb{R}^d$ (see (1.14) and (1.15)),

$$\begin{aligned} \langle \tilde{p} \rangle_T(\mathbf{u}, \xi) &= \frac{1}{2T} \int_{-T}^T \tilde{p}(\mathbf{u}, e^{-tB^t} \xi) dt = \frac{1}{2T} \int_{-T}^T \frac{p_2^0(\mathbf{u}, e^{-tB^t} \xi)}{\langle e^{-tB^t} \xi \rangle^2} dt \\ &\leq \frac{1}{2T} \int_{-T}^T \langle A^0 e^{-tB^t} \xi, e^{-tB^t} \xi \rangle dt, \end{aligned} \quad (2.2)$$

where we have used $(\mathbf{u}, 0) \in \mathcal{C}$ and thus $c(\mathbf{u}) = 0$ to obtain the second equality. Since (Harmo) implies $\langle \tilde{p} \rangle_T(\mathbf{u}, \xi) \geq \frac{1}{C} |\xi|^2$ for some constant $C > 0$ and every ξ small enough, part (i) of Lemma 2.1 is then an immediate consequence of (2.2).

To prove the second part of Lemma 2.1, let $\eta \in \mathbb{C}^d$ belong to $\ker(A^0) \cap \ker(B^t - z)$ for some $z \in \mathbb{C}$. Then $e^{-tB^t} \eta = e^{-tz} \eta$, and thus $\eta \in \ker(\int_{-T}^T e^{-tB} A^0 e^{-tB^t} dt) = \{0\}$. ■

For a not necessarily Morse function f (that is, without assuming (Morse)), \mathcal{U} (resp. $\tilde{\mathcal{U}}$) denotes the set of critical (resp. nondegenerate critical) points of f .

Lemma 2.2. (i) If (Gibbs) holds true, then

$$\tilde{\mathcal{U}} \times \{0\} \subset \mathcal{C}. \quad (2.3)$$

(ii) If (Gibbs) and (2.1) hold true, then

$$\mathcal{C} \subset \mathcal{U} \times \{0\}. \quad (2.4)$$

Proof. Suppose that $\mathbf{u} \in \tilde{\mathcal{U}}$ and assume without loss of generality that $\mathbf{u} = 0$. Thanks to (1.18), we have $c^0(0) = 0$. Moreover, near the origin,

$$\nabla f(x) = Hx + \mathcal{O}(x^2) \quad \text{and} \quad b^0(x) = b^0(0) + \mathcal{O}(x),$$

which, combined with (1.19), yields

$$\forall x \in \mathbb{R}^d, \quad \langle b^0(0), Hx \rangle = 0,$$

and thus $Hb^0(0) = 0$. Since $\mathbf{u} = 0$ is assumed to be nondegenerate, H is invertible and hence $b^0(0) = 0$. This proves (2.3).

We now show (2.4). Let $(\mathbf{u}, 0) \in \mathcal{C}$ and assume without loss of generality that $\mathbf{u} = 0$. Then $b^0(0) = 0$ and $c^0(0) = 0$. Denoting $\eta := \nabla f(0)$, it follows from (1.18) that $\langle A^0 \eta, \eta \rangle = 0$. Since A^0 is positive semidefinite, this implies $A^0 \eta = 0$. On the other hand, (1.19) yields

$$\forall x \in \mathbb{R}^d, \quad 0 = \langle b^0(0) + Bx + \mathcal{O}(x^2), \eta + \mathcal{O}(x) \rangle = \langle Bx + \mathcal{O}(x^2), \eta + \mathcal{O}(x) \rangle,$$

and hence $B^t \eta = 0$. It follows that $\eta \in \ker(A^0) \cap \ker(B^t)$ and so $\eta = 0$ thanks to Lemma 2.1 (ii). This completes the proof of (2.4). ■

Proof of Lemma 1.1. We can deduce Lemma 1.1 from Lemma 2.2. Indeed, the sets $\tilde{\mathcal{U}}$ and \mathcal{U} coincide when f is a Morse function. Then Lemma 2.2 (i) provides the first statement of Lemma 1.1. On the other hand, (Harmo) and (Gibbs) imply (2.1) from Lemma 2.1 (i) and hence (2.4). Thus, Lemma 2.2 gives $\mathcal{C} = \mathcal{U} \times \{0\}$ in that case, finishing the proof of Lemma 1.1. ■

Lemma 2.3. Assume (Gibbs) and let $\mathbf{u} \in \mathcal{U}$ with $b(\mathbf{u}) = 0$. Then the matrix $B^t H$ is antisymmetric. If moreover the critical point \mathbf{u} is nondegenerate (i.e. $\mathbf{u} \in \tilde{\mathcal{U}}$), the matrix $J := H^{-1} B^t$ is also antisymmetric.

Proof. As before, we assume for simplicity that $\mathbf{u} = 0$. Performing a Taylor expansion of the identity $b(x) \cdot \nabla f(x) = 0$ (see (1.19)), we deduce $Bx \cdot Hx = 0$ for every $x \in \mathbb{R}^d$.

Therefore, $x \cdot B^t H x = 0$ for every $x \in \mathbb{R}^d$, which implies that the matrix $\tilde{J} = B^t H$ is antisymmetric.

Moreover, if in addition H is invertible, then $J = H^{-1} B^t = H^{-1} \tilde{J} H^{-1}$ is antisymmetric since H is symmetric. ■

Corollary 2.4. Assume (Gibbs). Then condition (Harmo) is satisfied if and only if, for every $(\mathbf{u}, 0) \in \mathcal{C}$,

H is invertible and the symmetric matrix $\int_{-T}^T e^{-tB} A^0 e^{-tB^t} dt$ is positive definite.

Remark 2.5. Mimicking the Kalman criterion for controllability of finite-dimensional systems (see [5, Theorem 1.16]), one can show that $\int_{-T}^T e^{-tB} A^0 e^{-tB^t} dt$ is positive definite if and only if

$$\bigoplus_{n=0}^{d-1} \text{Im}(B^n A^0) = \mathbb{R}^d \quad \text{or} \quad \bigcap_{n=0}^{d-1} \ker(A^0 (B^t)^n) = \{0\}.$$

Proof of Corollary 2.4. We first prove a formula for $\langle \tilde{p} \rangle_T$ near $(\mathbf{u}, 0) \in \mathcal{C}$ with $\mathbf{u} \in \mathcal{U}$. Without loss of generality, we can assume that $\mathbf{u} = 0$. Denoting (x, ξ) by ρ , (1.14) and (1.18) yield, near $(0, 0)$,

$$\begin{aligned} \tilde{p}(\rho) &= p_2^0(x, \xi) + c^0(x) + \mathcal{O}(\rho^3) \\ &= A^0 \xi \cdot \xi + A^0 H x \cdot H x + \mathcal{O}(\rho^3). \end{aligned}$$

On the other hand, since $0 = (0, 0)$ is a critical point of $H_{p_1^0}$, we have $d \exp(t H_{p_1^0})(0) = \exp(t F_{p_1^0})$, where $F_{p_1^0}$ is the linearization of $H_{p_1^0}$ at 0. Using $p_1^0(x, \xi) = B x \cdot \xi + \mathcal{O}(\rho^3)$, we obtain $F_{p_1^0}(x, \xi) = (B x, -B^t \xi)$. Consequently, near $(0, 0)$,

$$\begin{aligned} \langle \tilde{p} \rangle_T(\rho) &= \frac{1}{2T} \int_{-T}^T (\langle A^0 H e^{tB} x, H e^{tB} x \rangle + \langle A^0 e^{-tB^t} \xi, e^{-tB^t} \xi \rangle + \mathcal{O}(\rho^3)) dt \\ &= \frac{1}{2T} \int_{-T}^T (\langle A^0 H e^{tB} x, H e^{tB} x \rangle + \langle A^0 e^{-tB^t} \xi, e^{-tB^t} \xi \rangle) dt + \mathcal{O}(\rho^3), \end{aligned}$$

from (1.15). Since moreover $B^t H$ is antisymmetric according to Lemma 2.3, we have $HB = -B^t H$ and hence $H e^{tB} = e^{-tB^t} H$. This leads to

$$\langle \tilde{p} \rangle_T(\rho) = \frac{1}{2T} \int_{-T}^T (\langle A^0 e^{-tB^t} H x, e^{-tB^t} H x \rangle + \langle A^0 e^{-tB^t} \xi, e^{-tB^t} \xi \rangle) dt + \mathcal{O}(\rho^3). \quad (2.5)$$

Let us now prove the corollary and consider $(\mathbf{u}, 0) \in \mathcal{C}$. If (Harmo) holds, Lemmas 2.1 (i) and 2.2 (ii) imply that $\mathbf{u} \in \mathcal{U}$. Then, we can apply (2.5), which, combined with (Harmo), shows that the matrices

$$H \left(\int_{-T}^T e^{-tB} A^0 e^{-tB^t} dt \right) H \quad \text{and} \quad \int_{-T}^T e^{-tB} A^0 e^{-tB^t} dt$$
 are positive definite.

This is equivalent to

$$H \text{ is invertible and the symmetric matrix } \int_{-T}^T e^{-tB} A^0 e^{-tB^t} dt \text{ is positive definite,} \quad (2.6)$$

proving the direct implication of Corollary 2.4. On the other hand, if (2.6) holds true, then (2.1) holds true and Lemma 2.2 (ii) gives $\mathbf{u} \in \mathcal{U}$. Then (2.5) and (2.6) imply (Harmo), proving the converse implication. ■

2.2. The spectrum of the matrix Λ

The aim of this section is to provide information on the spectrum of the matrix $\Lambda = 2HA^0 + B^t$ and to prove Lemma 1.4. We start with the following result, where $J = H^{-1}B^t$ is antisymmetric according to Lemma 2.3.

Lemma 2.6. Assume (Harmo), (Gibbs) and (Morse). For any $\mathbf{u} \in \mathcal{U}$ and $r \in [0, 1]$, the matrix

$$\Lambda_r := H(r(2A^0 + J) + (1-r)\text{Id})$$

has no eigenvalue on the imaginary axis $\{\text{Re } z = 0\}$.

Proof. Suppose that there exists $v \in \mathbb{C}^d$ such that $\Lambda_r v = zv$ with $\text{Re } z = 0$. Then, using $\langle v, H^{-1}v \rangle_{\mathbb{C}^d} \in \mathbb{R}$, we get

$$\text{Re } \langle \Lambda_r v, H^{-1}v \rangle_{\mathbb{C}^d} = \text{Re}(z \langle v, H^{-1}v \rangle_{\mathbb{C}^d}) = 0. \quad (2.7)$$

On the other hand,

$$\begin{aligned} \text{Re } \langle \Lambda_r v, H^{-1}v \rangle_{\mathbb{C}^d} &= 2r \text{Re } \langle A^0 v, v \rangle_{\mathbb{C}^d} + r \text{Re } \langle Jv, v \rangle_{\mathbb{C}^d} + (1-r) \text{Re } \langle v, v \rangle_{\mathbb{C}^d} \\ &= 2r \langle A^0 v, v \rangle_{\mathbb{C}^d} + (1-r) \|v\|^2. \end{aligned} \quad (2.8)$$

For $r \in [0, 1[$, (2.7) and (2.8) imply $v = 0$ and the lemma follows in that case. Assume now that $r = 1$. Relations (2.7) and (2.8) yield $\langle A^0 v, v \rangle = 0$ and so $A^0 v = 0$ since A^0 is positive semidefinite. Thus, the eigenvalue equation $\Lambda v = zv$ reads $B^t v = zv$. Hence, $v \in \ker(A^0) \cap \ker(B^t - z)$ and thus $v = 0$ according to Lemma 2.1, which completes the proof in the case $r = 1$. ■

We now give the proof of Lemma 1.4. Let $k \in \{0, \dots, d\}$ and let $\mathbf{u} \in \mathcal{U}^{(k)}$. For $r = 0$, $\Lambda_0 = H$ has exactly k eigenvalues in $\{\text{Re } z < 0\}$ since \mathbf{u} is a critical point of index k . As the eigenvalues of Λ_r are continuous with respect to r and cannot cross the imaginary axis from Lemma 2.6, $\Lambda_1 = \Lambda$ has exactly k eigenvalues in $\{\text{Re } z < 0\}$ and no eigenvalue on the imaginary axis. This proves (i) of the lemma.

Suppose now that $k = 1$ and let $\mu(\mathbf{u})$ denote the unique eigenvalue of Λ in $\{\text{Re } z < 0\}$. Since Λ is a real matrix, its set of eigenvalues is stable by complex conjugation and hence $\mu(\mathbf{u}) \in \mathbb{R}$. This proves (ii).

2.3. Rough localization of the spectrum

In this section, we prove Proposition 1.2. The arguments are close to those of [14, Section 2.2.2]. Thanks to Lemma 1.1, one has $\mathcal{C} = \mathcal{U} \times \{0\}$ and hence it suffices to show that, for any $\rho = (\mathbf{u}, 0) \in \mathcal{U} \times \{0\}$, the numbers $\mu_{\rho,0}^0$ of Theorem 1 have the following property:

$$\begin{cases} \mu_{\rho,0}^0 = 0 & \text{for all } \mathbf{u} \in \mathcal{U}^{(0)}, \\ \operatorname{Re} \mu_{\rho,0}^0 > 0 & \text{for all } \mathbf{u} \in \mathcal{U}^{(k)}, k \geq 1, \end{cases} \quad (2.9)$$

where we recall that $\mu_{\rho,0}^0 = \frac{1}{2} \widetilde{\operatorname{tr}}(p, \rho)$ with $\widetilde{\operatorname{tr}}(p, \rho) = -i \sum_{\ell=1}^d \lambda_{\rho,\ell} + 2c^1(\mathbf{u})$, and $\pm \lambda_{\rho,\ell}$, $\ell \in \{1, \dots, d\}$, denote the eigenvalues of the fundamental matrix F_{p^0} of p^0 at $\rho = (\mathbf{u}, 0)$ with the convention $\operatorname{Im} \lambda_{\rho,\ell} > 0$ (see the paragraph above Theorem 1). From now on, we take $\mathbf{u} \in \mathcal{U}^{(k)}$, $k \geq 0$, and we suppose again that $\mathbf{u} = 0$ in order to lighten the notation. Thanks to (1.18), near $(0, 0)$ one has

$$\begin{aligned} p^0(x, \xi) &= \langle A^0(x) \xi, \xi \rangle + i b^0(x) \cdot \xi + c^0(x) \\ &= \langle A^0(x) \xi, \xi \rangle + i b^0(x) \cdot \xi + \langle A^0(x) \nabla f(x), \nabla f(x) \rangle \\ &= \langle A^0 \xi, \xi \rangle + i B x \cdot \xi + \langle A^0 H x, H x \rangle + \mathcal{O}((x, \xi)^3). \end{aligned}$$

Then

$$F_{p^0} = \begin{pmatrix} iB & 2A^0 \\ -2HA^0H & -iB^t \end{pmatrix}. \quad (2.10)$$

Using the identity $B^t H = -HB$ which follows from Lemma 2.3, a direct computation shows that $L_{\pm} = \{(X, \pm i H X); X \in \mathbb{C}^d\}$ are vector spaces stable under F_{p^0} , that the map F_{p^0} restricted to L_{\pm} acts like

$$F_{\pm} = i(\pm 2A^0 H + B),$$

and that $\mathbb{C}^{2d} = L_+ \oplus L_-$ since H is invertible. In particular, $\sigma(F_{p^0}) = \sigma(F_+) \cup \sigma(F_-)$. It follows moreover from Lemma 1.4 that $F_+ = i(2HA^0 + B^t)^t$ has k eigenvalues $\lambda_1^+, \dots, \lambda_k^+$ in $\{\operatorname{Im} z < 0\}$ and $d - k$ eigenvalues $\lambda_{k+1}^+, \dots, \lambda_d^+$ in $\{\operatorname{Im} z > 0\}$. In addition, using again $B^t H = -HB$, we get

$$\begin{aligned} F_- &= i(-2A^0 H + B) = iH^{-1}(-2HA^0 + HBH^{-1})H \\ &= -H^{-1}i(2HA^0 + B^t)H = -H^{-1}(F_+)^t H. \end{aligned}$$

Hence, $\sigma(F_-) = -\sigma(F_+)$, which proves that

$$\sigma(F_{p^0}) \cap \{\operatorname{Im} z > 0\} = \{-\lambda_1^+, \dots, -\lambda_k^+\} \cup \{\lambda_{k+1}^+, \dots, \lambda_d^+\} = \{\lambda_{\rho,1}, \dots, \lambda_{\rho,d}\}. \quad (2.11)$$

On the other hand, let us recall that (Gibbs) implies $(-h \operatorname{div} \circ A \circ h \nabla + c)(e^{-f/h}) = 0$ (see the lines above (1.18)). Using the classical expansions of the coefficients and looking at the terms of order 1 then shows that

$$2c^1(\mathbf{u}) = -2 \sum_{j,\ell=1}^d A_{j,\ell}^0(\mathbf{u}) \partial_{x_j} \partial_{x_\ell} f(\mathbf{u}) = -\operatorname{tr}(2A^0 H).$$

Since $\operatorname{tr}(B) = \operatorname{tr}(-H^{-1}B^tH) = -\operatorname{tr}(B^t) = -\operatorname{tr}(B) = 0$ thanks to $B^tH = -HB$, we get

$$2c^1(\mathbf{u}) = -\operatorname{tr}(2A^0H + B) = i \operatorname{tr}(F_+). \quad (2.12)$$

Combining (2.11) and (2.12) yields

$$\widetilde{\operatorname{tr}}(p, \rho) = -i \sum_{\ell=1}^d \lambda_{\rho, \ell} + 2c^1(\mathbf{u}) = i \sum_{j=1}^k \lambda_j^+ - i \sum_{j=k+1}^d \lambda_j^+ + i \sum_{j=1}^d \lambda_j^+ = 2i \sum_{j=1}^k \lambda_j^+.$$

By definition, this last quantity vanishes when $k = 0$ and has a positive real part when $k \geq 1$. This proves (2.9).

3. Quasimodal constructions near the saddle points

In this part, we assume the hypotheses of Theorem 1, (Gibbs) and (Morse).

3.1. A first step in the construction of quasimodes

Given $\mathbf{s} \in \mathcal{U}^{(1)}$, we look for an approximate solution to the equation $Pu = 0$ in a neighborhood V of \mathbf{s} of the form

$$u(x) = v(x, h)e^{-f(x)/h}$$

with a function v of the form

$$v(x, h) = \int_0^{\ell(x, h)} \zeta(s/\tau) e^{-s^2/(2h)} ds, \quad (3.1)$$

where the function $\ell(x, h) \in C^\infty(V)$ has a classical expansion $\ell(x, h) \sim \sum_{j \geq 0} h^j \ell_j(x)$ in $C^\infty(V)$ with $\ell_0 \not\equiv 0$. Here, ζ denotes a fixed smooth even function equal to 1 on $[-1, 1]$ and supported in $[-2, 2]$, and $\tau > 0$ is a small parameter which will be fixed later. The object of this section is to construct the function ℓ .

Lemma 3.1. *One has*

$$P(v e^{-f/h}) = (w + r) e^{-(f + \ell^2/2)/h},$$

where

$$w = h(2A\nabla\ell \cdot \nabla f + (A\nabla\ell \cdot \nabla\ell)\ell + b \cdot \nabla\ell) - h^2 \operatorname{div}(A\nabla\ell),$$

the function r and all its derivatives are (locally) bounded, uniformly with respect to h , and $\operatorname{supp}(r) \subset \{|\ell| \geq \tau\}$. Here, we recall that all the functions above depend on x and h . Moreover, w admits a classical expansion $w \sim \sum_{j \geq 1} h^j w_j$ with

$$w_1 = 2A^0 \nabla f \cdot \nabla \ell_0 + (A^0 \nabla \ell_0 \cdot \nabla \ell_0) \ell_0 + b^0 \cdot \nabla \ell_0,$$

and, for all $j \geq 1$,

$$w_{j+1} = 2A^0 \nabla f \cdot \nabla \ell_j + (A^0 \nabla \ell_0 \cdot \nabla \ell_0) \ell_j + 2(A^0 \nabla \ell_0 \cdot \nabla \ell_j) \ell_0 + b^0 \cdot \nabla \ell_j + R_j(x, \partial^\alpha \ell_k),$$

where R_j is a polynomial in $\partial^\alpha \ell_k$, $|\alpha| \leq 2$, and $k \in \{0, \dots, j-1\}$, with smooth coefficients.

Proof. Throughout the proof, we write $\zeta(\cdot)$ instead of $\zeta(\cdot/\tau)$. Recall that $P = P_2 + P_1 + P_0$ with $P_0 = c$, $P_1 = \frac{1}{2}(b \cdot h \nabla + h \operatorname{div} \circ b)$ and $P_2 = -h \operatorname{div} \circ A \circ h \nabla$. Of course,

$$[P_0, v] = 0. \quad (3.2)$$

Using (3.1), one gets

$$[P_1, v] = h(b \cdot \nabla \ell) \zeta(\ell) e^{-\ell^2/(2h)} = h((b \cdot \nabla \ell) + r_1) e^{-\ell^2/(2h)}, \quad (3.3)$$

with $r_1 = (b \cdot \nabla \ell)(\zeta(\ell) - 1)$. In particular, r_1 and all its derivatives are (locally) bounded, uniformly with respect to h , and $\operatorname{supp}(r_1) \subset \{|\ell| \geq \tau\}$. On the other hand, since the matrix A is symmetric, for any smooth function ψ one has

$$P_2(v\psi) = -h \operatorname{div}(Ah \nabla(v\psi)) = \psi P_2 v + v P_2 \psi - 2Ah \nabla v \cdot h \nabla \psi.$$

Using again (3.1), this yields

$$[P_2, v] = e^{-\ell^2/(2h)} (-2h \zeta(\ell) A \nabla \ell \cdot h \nabla - h^2 \operatorname{div}(\zeta(\ell) A \nabla \ell) + h \ell \zeta(\ell) A \nabla \ell \cdot \nabla \ell),$$

and hence

$$\begin{aligned} [P_2, v](e^{-f/h}) &= (h \zeta(\ell) (2A \nabla \ell \cdot \nabla f + \ell A \nabla \ell \cdot \nabla \ell) - h^2 \operatorname{div}(\zeta(\ell) A \nabla \ell)) e^{-(f+\ell^2/2)/h} \\ &= (h(2A \nabla \ell \cdot \nabla f + \ell A \nabla \ell \cdot \nabla \ell) - h^2 \operatorname{div}(A \nabla \ell) + r_2) e^{-(f+\ell^2/2)/h} \end{aligned}$$

with $r_2 = h(\zeta(\ell) - 1)(2A \nabla \ell \cdot \nabla f + \ell A \nabla \ell \cdot \nabla \ell) - h^2 \operatorname{div}((\zeta(\ell) - 1)A \nabla \ell)$. In particular, r_2 and all its derivatives are (locally) bounded, uniformly with respect to h , and $\operatorname{supp}(r_2) \subset \{|\ell| \geq \tau\}$. Combining this identity with (3.2) and (3.3), and using the relation $P(v e^{-f/h}) = [P, v](e^{-f/h})$ implied by (Gibbs), we obtain the first part of the statement. Moreover, since the coefficients of P and the function ℓ have classical expansions, so does w . Plugging the expansions of A , b , c , and ℓ into the expression of w and identifying the powers of h , we obtain the formulas for the w_j . ■

In order to construct accurate quasimodes, we have to find smooth functions ℓ_j , $j \geq 0$, with $\ell_0 \not\equiv 0$ and such that the above w_{j+1} vanish. The equation on ℓ_0 is

$$2A^0 \nabla f \cdot \nabla \ell_0 + (A^0 \nabla \ell_0 \cdot \nabla \ell_0) \ell_0 + b^0 \cdot \nabla \ell_0 = 0, \quad (3.4)$$

and the equations on the ℓ_j , $j \geq 1$, are

$$2A^0 \nabla f \cdot \nabla \ell_j + (A^0 \nabla \ell_0 \cdot \nabla \ell_0) \ell_j + 2(A^0 \nabla \ell_0 \cdot \nabla \ell_j) \ell_0 + b^0 \cdot \nabla \ell_j = -R_j. \quad (3.5)$$

By analogy with the usual WKB method, we call (3.4) the eikonal equation and (3.5) the transport equations.

3.2. Resolution of the eikonal equation (3.4)

Consider the complexified symbol

$$q(x, \xi) = -p^0(x, i\xi) = \xi \cdot A^0(x)\xi + b^0(x) \cdot \xi - c^0(x) \in \mathbb{R}, \quad (3.6)$$

and let, for some saddle point $\mathbf{s} \in \mathcal{U}^{(1)}$, Λ_+ (resp. Λ_-) be the stable outgoing (resp. incoming) manifold of the H_q flow passing through $\rho_s = (\mathbf{s}, 0) \in \mathcal{C}$. It is proved in [13, Lemma 8.1] that Λ_{\pm} are Lagrangian manifolds which project nicely on the x -space. Hence, there exist smooth functions ϕ_{\pm} defined in a neighborhood of ρ_s such that $\phi_{\pm}(\mathbf{s}) = 0$ and, near ρ_s ,

$$\Lambda_{\pm} = \{(x, \xi) \in T^*\mathbb{R}^d; \xi = \nabla\phi_{\pm}(x)\}.$$

Moreover, $\Lambda_{\pm} \subset \{(x, \xi) \in T^*\mathbb{R}^d; q(x, \xi) = q(\rho_s) = 0\}$ and, according to [13, Proposition 8.2], $\pm \text{Hess } \phi_{\pm}(\mathbf{s}) > 0$. Summing up, the following properties of the functions ϕ_{\pm} will be used:

$$\phi_{\pm}(\mathbf{s}) = 0, \quad \nabla\phi_{\pm}(\mathbf{s}) = 0, \quad \pm \text{Hess } \phi_{\pm}(\mathbf{s}) > 0. \quad (3.7)$$

Lemma 3.2. *There exists a neighborhood V of \mathbf{s} and a smooth function $\ell_{s,0} \in C^{\infty}(V)$ such that*

$$\forall x \in V, \quad \phi_+(x) = f(x) - f(\mathbf{s}) + \frac{\ell_{s,0}^2(x)}{2}.$$

Moreover, $\nabla\ell_{s,0}(\mathbf{s}) \neq 0$.

Proof. This lemma comes from an observation of [13, (11.20)] and we follow this approach. Let $\Lambda_f = \{(x, \nabla f(x))\} \subset T^*\mathbb{R}^d$ denote the Lagrangian manifold associated with the phase function f . The eikonal equations (1.18) and (1.19) imply $q(x, \nabla f(x)) = 0$ and so Λ_f is stable by the H_q flow. In particular, the tangent space $T_{\rho_s}\Lambda_f$ at ρ_s is stable by F_q , the linearization of H_q at ρ_s . Moreover, a direct computation and (2.10) show that

$$F_q = \begin{pmatrix} B & 2A^0 \\ 2HA^0H & -B^t \end{pmatrix} = -i \begin{pmatrix} 1 & 0 \\ 0 & -i \end{pmatrix} F_{p^0} \begin{pmatrix} 1 & 0 \\ 0 & -i \end{pmatrix}^*.$$

In particular, the discussion below (2.10) implies that F_q has no eigenvalue on the imaginary axis. Let k_{\pm} be the number of eigenvalues of F_q restricted to $T_{\rho_s}\Lambda_f$ with positive/negative real part. Then $k_+ + k_- = d$.

Let K_{\pm} be the stable outgoing/incoming submanifold of Λ_f given by the Hamiltonian vector field H_q restricted to Λ_f . Then K_{\pm} has dimension k_{\pm} and K_{\pm} projects nicely on the x -space

$$K_{\pm} = \Lambda_{\pm} \cap \Lambda_f \quad \text{and} \quad T_{\rho_s}K_{\pm} = T_{\rho_s}\Lambda_{\pm} \cap T_{\rho_s}\Lambda_f. \quad (3.8)$$

On account of $\nabla\phi_{\pm} = \nabla f$ on $\pi_x(K_{\pm})$, we get

$$\forall x \in \pi_x(K_{\pm}), \quad \phi_{\pm}(x) = f(x) - f(\mathbf{s}). \quad (3.9)$$

Since \mathbf{s} is a saddle point of f , its Hessian has signature $(d-1, 1)$. Thus, (3.7) and (3.9) imply that $k_+ = d-1$ and $k_- = 1$. Eventually, using (3.9) again, we have on $T_s\pi_x(K_-)$

$$\text{Hess}(\phi_+ - f) = \text{Hess } \phi_+ - \text{Hess } \phi_- > 0.$$

Thus, in a neighborhood of \mathbf{s} , $g := \phi_+ - f + f(\mathbf{s})$ is a nonnegative function which vanishes to order 2 on $\pi_x(K_+)$.

Let us now construct a square root of g . After a local change of coordinates $x \mapsto (y, z) \in \mathbb{R}^{d-1} \times \mathbb{R}$ mapping \mathbf{s} to 0, we can assume that $\pi_x(K_+) = \{(y, z); z = 0\}$. Near 0, $g(y, 0) = 0$ from (3.9), $\partial_z g(y, 0) = 0$ from (3.8), and $\partial_{z,z}^2 g(y, 0) > 0$ from the last sentence of the previous paragraph. Thus, the Taylor formula gives

$$g(y, z) = z^2 \int_0^1 (1-t) \partial_{z,z}^2 g(y, tz) dt,$$

which leads to

$$\phi_+ = f - f(\mathbf{s}) + \frac{\ell_{\mathbf{s},0}^2}{2} \quad \text{with} \quad \ell_{\mathbf{s},0}(y, z) = z \left(2 \int_0^1 (1-t) \partial_{z,z}^2 g(y, tz) dt \right)^{1/2}.$$

Since the quantity under the square root is positive when evaluated at $z = 0$, $\ell_{\mathbf{s},0}$ is a smooth function in the vicinity of \mathbf{s} and $\nabla \ell_{\mathbf{s},0}(\mathbf{s}) \neq 0$. ■

Lemma 3.3. *Let $\mathbf{s} \in \mathcal{U}^{(1)}$. The function $\ell_{\mathbf{s},0}$ defined by Lemma 3.2 solves (3.4) in a neighborhood of \mathbf{s} . Moreover, $\eta(\mathbf{s}) := \nabla \ell_{\mathbf{s},0}(\mathbf{s})$ is an eigenvector of the matrix $\Lambda(\mathbf{s}) = 2H(\mathbf{s})A^0(\mathbf{s}) + B^t(\mathbf{s})$ associated with the negative eigenvalue $\mu(\mathbf{s})$. Finally,*

$$\mu(\mathbf{s}) = -A^0(\mathbf{s})\eta(\mathbf{s}) \cdot \eta(\mathbf{s}), \quad (3.10)$$

and

$$\det \text{Hess}\left(f + \frac{1}{2}\ell_{\mathbf{s},0}^2\right)(\mathbf{s}) = -\det H(\mathbf{s}). \quad (3.11)$$

In particular, $A^0(\mathbf{s})\eta(\mathbf{s}) \cdot \eta(\mathbf{s}) > 0$.

Proof. We drop the index \mathbf{s} and write ℓ_0 instead of $\ell_{\mathbf{s},0}$. By definition, ϕ_+ is a solution of the eikonal equation

$$A^0 \nabla \phi_+ \cdot \nabla \phi_+ + b^0 \cdot \nabla \phi_+ - c^0 = 0. \quad (3.12)$$

Since $\phi_+ = f - f(\mathbf{s}) + \frac{1}{2}\ell_0^2$, this implies

$$A^0 \nabla f \cdot \nabla f + 2\ell_0 A^0 \nabla f \cdot \nabla \ell_0 + \ell_0^2 A^0 \nabla \ell_0 \cdot \nabla \ell_0 + b^0 \cdot \nabla f + \ell_0 b^0 \cdot \nabla \ell_0 - c^0 = 0.$$

Since f is also a solution of (3.12) by (1.18) and (1.19), it follows that

$$2\ell_0 A^0 \nabla f \cdot \nabla \ell_0 + \ell_0^2 A^0 \nabla \ell_0 \cdot \nabla \ell_0 + \ell_0 b^0 \cdot \nabla \ell_0 = 0,$$

which gives (3.4) by dividing by ℓ_0 (this is allowed since $\nabla \ell_0(\mathbf{s}) \neq 0$ implies $\ell_0 \neq 0$ a.e. around \mathbf{s}). Moreover, the Taylor expansion of (3.4) at \mathbf{s} gives, for every x around \mathbf{s} ,

$$2A^0(\mathbf{s})H(\mathbf{s})(x - \mathbf{s}) \cdot \eta(\mathbf{s}) + (A^0(\mathbf{s})\eta(\mathbf{s}) \cdot \eta(\mathbf{s}))(\eta(\mathbf{s}) \cdot (x - \mathbf{s})) + db^0(\mathbf{s})(x - \mathbf{s}) \cdot \eta(\mathbf{s}) = 0,$$

or equivalently

$$(x - \mathbf{s}) \cdot \left((2H(\mathbf{s})A^0(\mathbf{s}) + B^t(\mathbf{s}) + A^0(\mathbf{s})\eta(\mathbf{s}) \cdot \eta(\mathbf{s}))\eta(\mathbf{s}) \right) = 0.$$

It follows that

$$(2H(\mathbf{s})A^0(\mathbf{s}) + B^t(\mathbf{s}) + A^0(\mathbf{s})\eta(\mathbf{s}) \cdot \eta(\mathbf{s}))\eta(\mathbf{s}) = 0, \quad (3.13)$$

which shows that $\eta(\mathbf{s})$ is an eigenvector of $\Lambda(\mathbf{s}) = 2H(\mathbf{s})A^0(\mathbf{s}) + B^t(\mathbf{s})$ associated with the eigenvalue $-A^0(\mathbf{s})\eta(\mathbf{s}) \cdot \eta(\mathbf{s})$. Since $A^0(\mathbf{s})$ is positive semidefinite, this eigenvalue is nonpositive. Then Lemma 1.4 implies that $-A^0(\mathbf{s})\eta(\mathbf{s}) \cdot \eta(\mathbf{s}) = \mu(\mathbf{s}) < 0$.

It remains to prove (3.11). We drop the dependence on \mathbf{s} in the following when it is unambiguous. By definition of ℓ_0 , one has

$$\text{Hess}\left(f + \frac{1}{2}\ell_0^2\right)(\mathbf{s}) = H + \Pi_\eta,$$

where $\Pi_\eta x := \langle x, \eta \rangle \eta$. Hence, (3.11) is equivalent to

$$\det(\text{Id} + H^{-1}\Pi_\eta) = -1. \quad (3.14)$$

We first observe that η^\perp is stable by $E := \text{Id} + H^{-1}\Pi_\eta$, and $E|_{\eta^\perp} = \text{Id}$. On the other hand,

$$\langle E\eta, \eta \rangle = \|\eta\|^2(1 + \langle H^{-1}\eta, \eta \rangle). \quad (3.15)$$

But $H(2A^0 + J)\eta = \Lambda\eta = \mu\eta$ with $J = H^{-1}B^t$ gives $\langle (2A^0 + J)\eta, \eta \rangle = \mu\langle H^{-1}\eta, \eta \rangle$. Since J is antisymmetric by Lemma 2.3 and thanks to (3.10), this implies

$$\langle H^{-1}\eta, \eta \rangle = \frac{2}{\mu}\langle A^0\eta, \eta \rangle = -2.$$

Plugging this identity into (3.15), we get $\langle E\eta, \eta \rangle = -\|\eta\|^2$. Choosing a basis (e_2, \dots, e_d) of η^\perp and computing the matrix of E in the basis (η, e_2, \dots, e_d) , the above discussion implies (3.14). \blacksquare

3.3. Resolution of the transport equations (3.5)

Lemma 3.4. *There exists an open neighborhood V of \mathbf{s} and some smooth functions $\ell_{\mathbf{s},j} \in C^\infty(V)$ such that $\ell_{\mathbf{s},j}$ solves (3.5) for all $j \geq 1$.*

Proof. Since, for all $j \geq 1$, R_j only depends on the $\ell_{\mathbf{s},k}$ with $0 \leq k \leq j-1$, we can solve the equations (3.5) by induction. It thus suffices to show that there exists an open neighborhood V of \mathbf{s} such that, for any smooth function f , there exists $u \in C^\infty(V)$ satisfying

$$\mathcal{L}u = f, \quad (3.16)$$

where \mathcal{L} is the transport operator defined by

$$\mathcal{L}u = 2A^0\nabla f \cdot \nabla u + (A^0\nabla\ell_{\mathbf{s},0} \cdot \nabla\ell_{\mathbf{s},0})u + 2\ell_{\mathbf{s},0}(A^0\nabla\ell_{\mathbf{s},0} \cdot \nabla u) + b^0 \cdot \nabla u. \quad (3.17)$$

Assume for simplicity that $\mathbf{s} = 0$. We first look for a formal solution in powers of x . Given $m \in \mathbb{N}$, we denote by $\mathcal{P}_{\text{hom}}^m$ the set of homogeneous polynomials of degree m and

we write $f(x) \simeq \sum_{m \in \mathbb{N}} f_m$ with $f_m \in \mathcal{P}_{\text{hom}}^m$. We recall that $\nabla \ell_{s,0}(s) = \eta(s)$ is an eigenvector of $\Lambda(s) = 2H(s)A^0(s) + B^t(s)$ associated with its sole negative eigenvalue $\mu(s)$ (see Lemma 1.4). Then \mathcal{L} decomposes into

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{>},$$

where $\mathcal{L}_{>}(p) = \mathcal{O}(x^{m+1})$ for all $p \in \mathcal{P}_{\text{hom}}^m$ and $\mathcal{L}_0 : \mathcal{P}_{\text{hom}}^m \rightarrow \mathcal{P}_{\text{hom}}^m$ is given by

$$\mathcal{L}_0 = 2A^0(s)H(s)x \cdot \nabla + A^0(s)\eta(s) \cdot \eta(s) + 2\langle \eta(s), x \rangle A^0(s)\eta(s) \cdot \nabla + B(s)x \cdot \nabla,$$

which we can rewrite, since $\mu(s) = -A^0(s)\eta(s) \cdot \eta(s)$ by Lemma 3.3, as

$$\mathcal{L}_0 = (2A^0(s)H(s) + B(s) + 2A^0(s)\Pi_\eta)x \cdot \nabla - \mu(s), \quad (3.18)$$

where $\Pi_\eta y := \langle y, \eta(s) \rangle \eta(s)$. We shall prove that \mathcal{L}_0 is invertible on $\mathcal{P}_{\text{hom}}^m$ for any $m \geq 0$. Let us denote $\Upsilon = 2A^0(s)H(s) + B(s) + 2A^0(s)\Pi_\eta = \Lambda^t(s) + 2A^0(s)\Pi_\eta$. One has

$$\begin{aligned} \Upsilon^t \eta(s) &= \Lambda(s)\eta(s) + 2\Pi_\eta A^0(s)\eta(s) = \mu(s)\eta(s) + 2\langle A^0(s)\eta(s), \eta(s) \rangle \eta(s) \\ &= -\mu(s)\eta(s). \end{aligned}$$

Choosing some vectors e_2, \dots, e_d such that $\mathcal{B} = (\eta(s), e_2, \dots, e_d)$ is a basis of \mathbb{C}^d and the matrix M of $\Lambda(s)$ in the basis \mathcal{B} is upper triangular, it follows that the matrix M' of Υ^t in \mathcal{B} is also upper triangular, with the same diagonal entries as M , except that the first diagonal entry $\mu(s)$ is replaced by $-\mu(s)$ (actually, only the first lines of M and M' differ). Since $\mu(s)$ is the only eigenvalue of $\Lambda(s)$ with nonpositive real part, the spectrum of Υ^t is contained in $\{\text{Re } z > 0\}$. Thanks to Lemma A.1 in the appendix, this implies that the spectrum of $\Upsilon x \cdot \nabla : \mathcal{P}_{\text{hom}}^m \rightarrow \mathcal{P}_{\text{hom}}^m$ is contained in $\{\text{Re } z > 0\}$ for every $m > 0$ and hence $\mathcal{L}_0 = \Upsilon x \cdot \nabla - \mu(s)$ is invertible on $\mathcal{P}_{\text{hom}}^m$ for every $m \geq 0$ (note that $\mathcal{L}_0 = -\mu(s)$ on $\mathcal{P}_{\text{hom}}^0$). Using this property, we can solve the transport equation $\mathcal{L}u = f$ by the method of [7, Chapter 3]. We recall briefly the successive steps. We first find a formal solution \tilde{u} to the equation

$$\mathcal{L}\tilde{u} = \tilde{f}, \quad (3.19)$$

where \tilde{f} denotes the formal power expansion of f : $\tilde{f} \simeq \sum_k \tilde{f}_k$ with $\tilde{f}_k \in \mathcal{P}_{\text{hom}}^k$. We look for \tilde{u} of the form $\tilde{u} \simeq \sum_k \tilde{u}_k$ with $\tilde{u}_k \in \mathcal{P}_{\text{hom}}^k$. Since \mathcal{L}_0 is invertible, there exists $\tilde{u}_0 \in \mathcal{P}_{\text{hom}}^0$ solving $\mathcal{L}_0\tilde{u}_0 = \tilde{f}_0$. Then we choose $\tilde{u}_1 \in \mathcal{P}_{\text{hom}}^1$ solving

$$\mathcal{L}_0\tilde{u}_1 = \tilde{f}_1 + r_1,$$

where r_1 denotes the homogeneous part of degree 1 of $-\mathcal{L}_{>}(\tilde{u}_0)$. Iterating this procedure, we obtain a formal solution to (3.19). Using this formal solution and a Borel procedure, we construct a smooth function \bar{u} such that \bar{u} and \tilde{u} have the same Taylor expansion at the origin. As a consequence, $\mathcal{L}\bar{u} = f + \mathcal{O}(x^\infty)$. The last step consists in showing that for every $g = \mathcal{O}(x^\infty)$, there exists a solution $v = \mathcal{O}(x^\infty)$ to $\mathcal{L}v = g$. This can be done by using the method of characteristics and the fact that the spectrum of Υ is contained in $\{\text{Re } z > 0\}$; we refer to [7, proof of Proposition 3.5] for details. Then, taking \bar{u}_∞ satisfying $\mathcal{L}\bar{u}_\infty = f - \mathcal{L}\bar{u} = \mathcal{O}(x^\infty)$, we find that $u := \bar{u} + \bar{u}_\infty$ is a true solution of (3.16) and

[7, Proposition 3.5] shows that the neighborhood V of \mathbf{s} where u is defined can be chosen independently of f . ■

Using the preceding result and a Borel procedure, we get the following.

Proposition 3.5. *For any $\mathbf{s} \in \mathcal{U}^{(1)}$, there exists a smooth function $x \mapsto \ell_{\mathbf{s}}(x, h)$ defined in a neighborhood $V_{\mathbf{s}}$ of \mathbf{s} such that*

- (i) $\ell_{\mathbf{s}}$ admits a classical expansion $\ell_{\mathbf{s}} \sim \sum_k h^k \ell_{\mathbf{s},k}$,
- (ii) $(2A\nabla \ell_{\mathbf{s}} \cdot \nabla f + (A\nabla \ell_{\mathbf{s}} \cdot \nabla \ell_{\mathbf{s}})\ell_{\mathbf{s}} + b \cdot \nabla \ell_{\mathbf{s}}) - h \operatorname{div}(A\nabla \ell_{\mathbf{s}}) = \mathcal{O}(h^\infty)$, uniformly with respect to x in $V_{\mathbf{s}}$,
- (iii) $\ell_{\mathbf{s},0}(x) = (x - \mathbf{s}) \cdot \eta(\mathbf{s}) + \mathcal{O}(|x - \mathbf{s}|^2)$.

Note that the function $x \mapsto -\ell_{\mathbf{s}}(x, h)$ also satisfies Proposition 3.5. More precisely, (i) and (ii) hold without modification, while $\eta(\mathbf{s})$ has to be replaced by $-\eta(\mathbf{s}) = -\nabla \ell_{\mathbf{s},0}(\mathbf{s})$ in (iii). At this point, we do not specify which function ($\ell_{\mathbf{s}}$ or $-\ell_{\mathbf{s}}$) will be used later.

4. Global construction of quasimodes

In this section, which is an adaptation of [20, Section 4A], we assume the hypotheses of Theorem 1, (Gibbs), (Confin), and (Morse). We refer the reader to the notation following Definition 1.3 and introduce some additional topological objects. Given $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$, one has $\sigma(\mathbf{m}) = \sigma_i$ for some $i \geq 2$. Moreover, since $\sigma_{i-1} > \sigma_i$, there exists a unique connected component of $\{f < \sigma_{i-1}\}$ that contains \mathbf{m} (observe that this component is not necessarily critical). We denote that component by $E_-(\mathbf{m})$, and by

$$E_- : \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\} \rightarrow \mathcal{P}(\mathbb{R}^d) \quad (4.1)$$

the corresponding map. It follows from [24, Remark 2.2] that, for any $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$, there exists a unique $\widehat{\mathbf{m}} \in E_-(\mathbf{m}) \cap \mathcal{U}^{(0)}$ such that $\sigma(\widehat{\mathbf{m}}) > \sigma(\mathbf{m})$. In particular, $\mathbf{m} \in E_-(\mathbf{m}) \subset E(\widehat{\mathbf{m}})$, and thus

$$\forall \mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}, \quad f(\widehat{\mathbf{m}}) \leq f(\mathbf{m}). \quad (4.2)$$




Fig. 4.1. The geometric setting associated to $\mathbf{m}_{4,1}$.

We denote by $\widehat{E}(\mathbf{m})$ the connected component of $\{f < \sigma(\mathbf{m})\}$ containing $\widehat{\mathbf{m}}$. Then $\widehat{E}(\mathbf{m}) \subset E_-(\mathbf{m})$ and $\widehat{E}(\mathbf{m})$ is a critical component (see Definition 1.3). We denote by $\widehat{E} : \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\} \rightarrow \mathbb{C}$ and $\widehat{\mathbf{m}} : \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\} \rightarrow \mathcal{U}^{(0)}$ the corresponding maps (see Fig. 4.1).

Fix $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$. For every $\mathbf{s} \in \mathbf{j}(\mathbf{m})$, one has $f(\mathbf{s}) = \sigma(\mathbf{m})$. For any $\tau, \delta > 0$, we define

$$\begin{aligned} \mathcal{B}_{\mathbf{s}, \tau, \delta} &:= \{f \leq \sigma(\mathbf{m}) + \delta\} \cap \{x \in \mathbb{R}^d; |\eta(\mathbf{s}) \cdot (x - \mathbf{s})| \leq \tau\}, \\ \mathcal{C}_{\mathbf{s}, \tau, \delta} &:= \text{the connected component of } \mathcal{B}_{\mathbf{s}, \tau, \delta} \text{ containing } \mathbf{s}, \end{aligned} \quad (4.3)$$

where $\eta(\mathbf{s})$ has been defined in Lemma 3.3. We recall that $\eta(\mathbf{s})$ is an eigenvector of the matrix $\Lambda(\mathbf{s}) = 2H(\mathbf{s})A^0(\mathbf{s}) + B^t(\mathbf{s})$ associated with its only negative eigenvalue $\mu(\mathbf{s})$, which has multiplicity 1 (see Lemma 1.4). Moreover, from (3.10) we get the normalization condition $A^0(\mathbf{s})\eta(\mathbf{s}) \cdot \eta(\mathbf{s}) = -\mu(\mathbf{s})$. Observe that this condition is not the same as in [20], where $\|\eta(\mathbf{s})\| = 1$ is imposed. Let us also define

$$E_{\mathbf{m}, \tau, \delta} := (E_-(\mathbf{m}) \cap \{f < \sigma(\mathbf{m}) + \delta\}) \setminus \bigcup_{\mathbf{s} \in \mathbf{j}(\mathbf{m})} \mathcal{C}_{\mathbf{s}, \tau, \delta}, \quad (4.4)$$

where $E_-(\mathbf{m})$ is defined by (4.1).

According to the geometry of the Morse function f around $\partial E(\mathbf{m})$ and to the lemmas of Section 3.2, we have the following result.

Lemma 4.1. *For any $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$ and $\mathbf{s} \in \mathbf{j}(\mathbf{m})$, there exists a neighborhood V of \mathbf{s} such that*

$$\forall x \in V \setminus \{\mathbf{s}\}, \quad x - \mathbf{s} \in \eta(\mathbf{s})^\perp \implies f(x) > f(\mathbf{s}).$$

It follows that, for $\tau_0, \delta_0 > 0$ sufficiently small and every $\tau \in]0, \tau_0]$ and $\delta \in]0, \delta_0]$, there exists a connected component of $E_{\mathbf{m}, 3\tau, 3\delta}$ containing $\mathcal{U}^{(0)} \cap E(\mathbf{m})$ and disjoint from $\mathcal{U}^{(0)} \cap (E_-(\mathbf{m}) \setminus E(\mathbf{m}))$. We will denote by $E_{\mathbf{m}, 3\tau, 3\delta}^+$ this component and by $E_{\mathbf{m}, 3\tau, 3\delta}^-$ its complement in $E_{\mathbf{m}, 3\tau, 3\delta}$.

Remark 4.2. The above set $E_{\mathbf{m}, 3\tau, 3\delta}^-$ contains $\widehat{\mathbf{m}}$ but is not connected in general. However, when \mathbf{m} satisfies $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset$ for every $\mathbf{m}' \in \mathcal{U}^{(0)} \setminus \{\mathbf{m}\}$, one has, owing to [20, Remark 1.7 and Section 4A], $\mathbf{j}(\mathbf{m}) = \partial \widehat{E}(\mathbf{m}) \cap \partial E(\mathbf{m})$. In that case, the set $E_{\mathbf{m}, 3\tau, 3\delta}^-$ is connected.

Proof of Lemma 4.1. Without loss of generality, we can assume that $\mathbf{s} = 0$ and $f(\mathbf{s}) = 0$. Thanks to Lemma 3.2 and Proposition 3.5, one has

$$\phi_+(x) = f(x) + \frac{\ell_{\mathbf{s}, 0}^2(x)}{2} = f(x) + \frac{\langle x, \eta(\mathbf{s}) \rangle^2}{2} + \mathcal{O}(x^3),$$

near $\mathbf{s} = 0$. This implies that, for all $x \in \eta(\mathbf{s})^\perp$, we have

$$f(x) = \phi_+(x) + \mathcal{O}(x^3).$$

Since Hess $\phi_+(\mathbf{s})$ is positive definite by (3.7), the conclusion follows. ■




Fig. 4.2. Representation of the Morse function f near a point $\mathbf{s} \in \mathbf{j}(\mathbf{m}) \cap \partial \widehat{E}(\mathbf{m})$ when the latter set is nonempty. Here, $\eta_1(\mathbf{s})$ denotes an eigenvector of $\text{Hess } f(\mathbf{s})$ associated with its negative eigenvalue.

Let us now define, for $h > 0$ and $\tau_0, \delta_0 > 0$ small enough, the function $v_{\mathbf{m},h}$ on the sublevel set $E_-(\mathbf{m}) \cap \{f < \sigma(\mathbf{m}) + 3\delta_0\}$ (see (4.1)) as follows. On the disjoint open sets $E_{\mathbf{m},3\tau_0,3\delta_0}^+$ and $E_{\mathbf{m},3\tau_0,3\delta_0}^-$ introduced in Lemma 4.1, we set

$$v_{\mathbf{m},h}(x) := \begin{cases} +1 & \text{for } x \in E_{\mathbf{m},3\tau_0,3\delta_0}^+, \\ -1 & \text{for } x \in E_{\mathbf{m},3\tau_0,3\delta_0}^-. \end{cases} \quad (4.5)$$

In addition, for every $\mathbf{s} \in \mathbf{j}(\mathbf{m})$ and $x \in \mathcal{C}_{\mathbf{s},3\tau_0,3\delta_0}$ (see (4.3)), we set

$$v_{\mathbf{m},h}(x) := C_{\mathbf{s},h}^{-1} \int_0^{\ell_{\mathbf{s}}(x,h)} \zeta(r/\tau_0) e^{-r^2/(2h)} dr, \quad (4.6)$$

where the function $\ell_{\mathbf{s}}$ is given by Proposition 3.5, and its sign (see the discussion below Proposition 3.5) is chosen so that there exists a neighborhood V of \mathbf{s} such that $E(\mathbf{m}) \cap V$ is included in the half-plane $\{\eta(\mathbf{s}) \cdot (x - \mathbf{s}) > 0\}$ (see Lemma 4.1, Figs. 4.2 and 4.3), $\zeta \in C^\infty(\mathbb{R}; [0, 1])$ is even and satisfies $\zeta = 1$ on $[-1, 1]$, $\zeta(r) = 0$ for $|r| \geq 2$, and

$$C_{\mathbf{s},h} := \frac{1}{2} \int_{-\infty}^{+\infty} \zeta(r/\tau_0) e^{-r^2/(2h)} dr.$$

In particular,

$$\exists \gamma > 0, \quad C_{\mathbf{s},h}^{-1} = \sqrt{\frac{2}{\pi h}} (1 + \mathcal{O}(e^{-\gamma/h})). \quad (4.7)$$

Note also that, for every $\tau_0 > 0$ and then for $\delta_0 > 0$ small enough, thanks to Proposition 3.5 and to the definitions (4.5) and (4.6), and since the sets $E_{\mathbf{m},3\tau_0,3\delta_0}^+$, $E_{\mathbf{m},3\tau_0,3\delta_0}^-$ and $\mathcal{C}_{\mathbf{s},3\tau_0,3\delta_0}$, $\mathbf{s} \in \mathbf{j}(\mathbf{m})$, are mutually disjoint (see Lemma 4.1), $v_{\mathbf{m},h}$ is well defined and is C^∞ on $E_-(\mathbf{m}) \cap \{f < \sigma(\mathbf{m}) + 3\delta_0\}$ for $h > 0$ small enough.




Fig. 4.3. The support of the function $v_{\mathbf{m},h}$

Consider now a smooth function $\theta_{\mathbf{m}}$ such that

$$\theta_{\mathbf{m}}(x) := \begin{cases} 1 & \text{for } x \in \{f \leq \sigma(\mathbf{m}) + \frac{3}{2}\delta_0\} \cap E_-(\mathbf{m}), \\ 0 & \text{for } x \in \mathbb{R}^d \setminus (\{f < \sigma(\mathbf{m}) + \frac{7}{4}\delta_0\} \cap E_-(\mathbf{m})). \end{cases} \quad (4.8)$$

The function $\theta_{\mathbf{m}}v_{\mathbf{m},h}$ belongs to $C_c^\infty(\mathbb{R}^d; [-1, 1])$ and

$$\text{supp}(\theta_{\mathbf{m}}v_{\mathbf{m},h}) \subset E_-(\mathbf{m}) \cap \{f < \sigma(\mathbf{m}) + 2\delta_0\}.$$

Definition 4.3. For $\tau_0 > 0$ and then for $\delta_0, h > 0$ small enough, define

$$\begin{cases} \psi_{\mathbf{m},h}(x) := e^{-\frac{f(x)-f(\mathbf{m})}{h}}, \\ \psi_{\mathbf{m},h}(x) := \theta_{\mathbf{m}}(x)(v_{\mathbf{m},h}(x) + 1)e^{-\frac{f(x)-f(\mathbf{m})}{h}} \quad \text{for } \mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}. \end{cases} \quad (4.9)$$

We then define, for any $\mathbf{m} \in \mathcal{U}^{(0)}$, the quasimode $\varphi_{\mathbf{m},h}$ by

$$\varphi_{\mathbf{m},h} := \frac{\psi_{\mathbf{m},h}}{\|\psi_{\mathbf{m},h}\|_{L^2}}.$$

Note that, for every τ_0, δ_0 , we have $h > 0$ so that the above definition makes sense, $P\varphi_{\mathbf{m},h} = 0$ and, for every $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$, the quasimodes $\psi_{\mathbf{m},h}$ and $\varphi_{\mathbf{m},h}$ belong to $C_c^\infty(\mathbb{R}^d; \mathbb{R}^+)$ with supports included in $E_-(\mathbf{m}) \cap \{f < \sigma(\mathbf{m}) + 2\delta_0\}$. We have more precisely the following lemma resulting from the previous construction.

Lemma 4.4. *For every $\mathbf{m} \in \mathcal{U}^{(0)}$ and $\varepsilon > 0$, there exists $\tau_0 > 0$ and then $\delta_0 > 0$ small enough such that, for every $h > 0$ small enough,*

- (i) $\text{supp}(\psi_{\mathbf{m},h}) \subset \overline{E(\mathbf{m})} + D(0, \varepsilon),$

(ii) when $\mathbf{m} \neq \underline{\mathbf{m}}$, there exists a neighborhood V_{τ_0, δ_0} of $\overline{E(\mathbf{m})}$ such that

$$V_{\tau_0, \delta_0} \setminus \bigcup_{s \in \mathbf{j}(\mathbf{m})} \mathcal{C}_{s, 3\tau_0, 3\delta_0} \subset \{\theta_{\mathbf{m}} v_{\mathbf{m}, h} = 1\},$$

(iii) when $\mathbf{m} \neq \underline{\mathbf{m}}$,

$$\forall x \in \text{supp}(\nabla(\theta_{\mathbf{m}}(v_{\mathbf{m}} + 1))), \quad \left(f(x) < \sigma(\mathbf{m}) + \frac{3}{2}\delta_0 \Rightarrow x \in \bigcup_{s \in \mathbf{j}(\mathbf{m})} \mathcal{C}_{s, 3\tau_0, 3\delta_0} \right).$$

Let moreover $\mathbf{m}' \subset \mathcal{U}^{(0)}$ with $\mathbf{m} \neq \mathbf{m}'$. For every $\tau_0 > 0$ and then $\delta_0 > 0$, one has, for $h > 0$ small enough,

(iv) if $\sigma(\mathbf{m}) = \sigma(\mathbf{m}')$ and $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset$, then $\text{supp}(\psi_{\mathbf{m}, h}) \cap \text{supp}(\psi_{\mathbf{m}', h}) = \emptyset$,

(v) if $\sigma(\mathbf{m}) > \sigma(\mathbf{m}')$, then either

- ★ $\text{supp}(\psi_{\mathbf{m}, h}) \cap \text{supp}(\psi_{\mathbf{m}', h}) = \emptyset$, or
- ★ $\psi_{\mathbf{m}, h} = 2e^{-(f-f(\mathbf{m}))/h}$ on $\text{supp}(\psi_{\mathbf{m}', h})$.

Proof. Points (i)–(iii) follow from the construction of the quasimodes $\varphi_{\mathbf{m}, h}$ in Definition 4.3 for $\mathbf{m} \in \mathcal{U}^{(0)}$: see indeed (4.5), (4.6) and (4.8). Let us now prove the last two points.

If $\sigma(\mathbf{m}) = \sigma(\mathbf{m}')$ and $\mathbf{m} \neq \mathbf{m}'$, note first that \mathbf{m} and \mathbf{m}' differ from $\underline{\mathbf{m}}$ since $\sigma(\mathbf{m}) = +\infty$ if and only if $\mathbf{m} = \underline{\mathbf{m}}$. Moreover, $\sigma(\mathbf{m}) = \sigma(\mathbf{m}')$ and $\mathbf{m} \neq \mathbf{m}'$ imply $E(\mathbf{m}) \cap E(\mathbf{m}') = \emptyset$. Indeed, $E(\mathbf{m}) \cap E(\mathbf{m}') \neq \emptyset$ would imply that $E(\mathbf{m})$ and $E(\mathbf{m}')$ are the same connected component of $\{f < \sigma(\mathbf{m}) = \sigma(\mathbf{m}')\}$, in contradiction with the construction of E . When in addition $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset$, we have

$$\overline{E(\mathbf{m})} \cap \overline{E(\mathbf{m}')} = \partial E(\mathbf{m}) \cap \partial E(\mathbf{m}') = \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset.$$

Combined with Lemma 4.4 (i) with $\varepsilon > 0$ sufficiently small, this implies $\text{supp}(\psi_{\mathbf{m}, h}) \cap \text{supp}(\psi_{\mathbf{m}', h}) = \emptyset$.

If $\sigma(\mathbf{m}) > \sigma(\mathbf{m}')$ and $\mathbf{m}' \notin E(\mathbf{m})$, we have $\overline{E(\mathbf{m})} \cap \overline{E(\mathbf{m}')} = \emptyset$, and again, according to (i), $\text{supp}(\psi_{\mathbf{m}, h}) \cap \text{supp}(\psi_{\mathbf{m}', h}) = \emptyset$ for every $\tau_0 > 0$ and then $\delta_0 > 0$ small enough. Lastly, if $\sigma(\mathbf{m}) > \sigma(\mathbf{m}')$ and $\mathbf{m}' \in E(\mathbf{m})$, then $\overline{E(\mathbf{m}')} \subset E_-(\mathbf{m}') \subset E(\mathbf{m})$ and so, according to (ii), $\psi_{\mathbf{m}, h} = 2e^{-(f-f(\mathbf{m}))/h}$ on $\text{supp}(\psi_{\mathbf{m}', h})$ for every $\tau_0 > 0$ and then $\delta_0 > 0$ small enough. ■

5. Proof of the main results

We will use the following notation. For two families $a = (a_h)_{h \in [0, h_0]}$ and $b = (b_h)_{h \in [0, h_0]}$ of numbers, we write $a \in \mathcal{E}_{\text{cl}}(b)$ if there exists a family $(c_h)_{h \in [0, h_0]}$ such that, for every $h \in [0, h_0]$,

$$a_h = b_h c_h \text{ and } c_h \text{ admits a classical expansion } c_h \sim \sum_{j \geq 0} c_j h^j \text{ with } c_0 = 1.$$

We also write $D_{\mathbf{u}} = |\det \text{Hess}(f)(\mathbf{u})|^{1/2}$ for any $\mathbf{u} \in \mathcal{U}$.

5.1. Computation of interaction coefficients

Proposition 5.1. *Under the assumptions of Theorem 2 but with only the first part of (Gener), we have the following estimates for every $\tau_0 > 0$ and then $\delta_0 > 0$ small enough: there exists $C > 0$ such that, for all $\mathbf{m}, \mathbf{m}' \in \mathcal{U}^{(0)}$ and for $h > 0$ small enough,*

- (i) $\langle \varphi_{\mathbf{m},h}, \varphi_{\mathbf{m}',h} \rangle = \delta_{\mathbf{m},\mathbf{m}'} + \mathcal{O}(e^{-C/h}),$
- (ii) $\langle P\varphi_{\mathbf{m},h}, \varphi_{\mathbf{m},h} \rangle \in \mathcal{E}_{\text{cl}} \left(h e^{-2S(\mathbf{m})/h} \sum_{\mathbf{s} \in \mathbf{j}(\mathbf{m})} \frac{|\mu(\mathbf{s})|}{2\pi} \frac{D_{\mathbf{m}}}{D_{\mathbf{s}}} \right),$
- (iii) $\|P\varphi_{\mathbf{m},h}\|^2 = \mathcal{O}(h^\infty) \langle P\varphi_{\mathbf{m},h}, \varphi_{\mathbf{m},h} \rangle,$
- (iv) $\|P^*\varphi_{\mathbf{m},h}\|^2 = \mathcal{O}(h) \langle P\varphi_{\mathbf{m},h}, \varphi_{\mathbf{m},h} \rangle.$

The statements of this proposition are very close to those of [20, Propositions 4.4 to 4.6]. The difference is that we have a classical expansion in (ii) and the multiplicative error in (iii) is of order $\mathcal{O}(h^\infty)$ instead of $\mathcal{O}(h^2)$. This is due to the fact that our quasimodal constructions are sharper.

Remark 5.2. When, in comparison with the assumptions of Proposition 5.1, the first part of (Gener) is not satisfied, items (iii) and (iv) of Proposition 5.1 still hold, while item (ii) becomes

$$\langle P\varphi_{\mathbf{m},h}, \varphi_{\mathbf{m},h} \rangle \in \mathcal{E}_{\text{cl}} \left(h e^{-2S(\mathbf{m})/h} \sum_{\mathbf{s} \in \mathbf{j}(\mathbf{m})} \frac{|\mu(\mathbf{s})|}{2\pi} \frac{(\sum_{\mathbf{m}' \in \arg\min_{E(\mathbf{m})} f} D_{\mathbf{m}'}^{-1})^{-1}}{D_{\mathbf{s}}} \right).$$

However, the quasi-orthonormality of the family $(\varphi_{\mathbf{m},h})_{\mathbf{m} \in \mathcal{U}^{(0)}}$ stated in (i) is not satisfied anymore when the first part of (Gener) does not hold. Indeed, if $\mathbf{m} \neq \mathbf{m}' \in \mathcal{U}^{(0)}$ satisfy $f(\mathbf{m}) = f(\mathbf{m}')$ and $\mathbf{m}' \in E(\mathbf{m})$, then $\sigma(\mathbf{m}) > \sigma(\mathbf{m}')$ and it follows from Lemma 4.4 (v) that $\psi_{\mathbf{m},h} = 2e^{-(f-f(\mathbf{m}'))/h}$ on $\text{supp}(\psi_{\mathbf{m}',h})$. Arguments similar to those in the proof below show that there exists $c \in]0, 1[$ such that $\langle \varphi_{\mathbf{m},h}, \varphi_{\mathbf{m}',h} \rangle \sim c$ when $h \rightarrow 0^+$.

Proof of Proposition 5.1. The proof follows the proofs of [20, Propositions 4.4 to 4.6] very closely. We just sketch it briefly and drop the index h in order to lighten the notation.

First, we recall that, for any $\mathbf{m} \in \mathcal{U}^{(0)}$, $\varphi_{\mathbf{m}} = \psi_{\mathbf{m}} / \|\psi_{\mathbf{m}}\|$ with $\psi_{\mathbf{m}}$ given by (4.9). Moreover, according to the first part of (Gener), f uniquely attains its absolute minimum on $\overline{E(\mathbf{m})}$ at \mathbf{m} , and hence on $\text{supp}(\psi_{\mathbf{m}})$ for every $\tau_0, \delta_0 > 0$ small enough (see Lemma 4.4 (i)). By the standard Laplace method, we then easily get, for any $\mathbf{m} \in \mathcal{U}^{(0)}$,

$$\|\psi_{\mathbf{m}}\| \in \mathcal{E}_{\text{cl}}(2(\pi h)^{d/4} D_{\mathbf{m}}^{-1/2}). \quad (5.1)$$

Let us now prove (i). First, by definition, we have $\langle \varphi_{\mathbf{m}}, \varphi_{\mathbf{m}} \rangle = 1$ for every $\mathbf{m} \in \mathcal{U}^{(0)}$. Moreover, for every $\mathbf{m} \neq \mathbf{m}' \in \mathcal{U}^{(0)}$, we are in one of the following three cases.

- (a) *The case where $\sigma(\mathbf{m}) = \sigma(\mathbf{m}')$ and $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset$.* We then deduce from Lemma 4.4 (iv) that $\text{supp}(\psi_{\mathbf{m}}) \cap \text{supp}(\psi_{\mathbf{m}'}) = \emptyset$ and so $\langle \varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle = 0$.

- (b) *The case where $\sigma(\mathbf{m}) = \sigma(\mathbf{m}')$ and $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') \neq \emptyset$* (note that this does not occur when the second part of (Gener) is satisfied). Then $\overline{E(\mathbf{m})} \cap \overline{E(\mathbf{m}')} = \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') \neq \emptyset$ and we deduce from the construction of the $\varphi_{\mathbf{m}}$, $\mathbf{m} \in \mathcal{U}^{(0)}$, that $\varphi_{\mathbf{m}\varphi_{\mathbf{m}'}}$ is supported in $\bigcup_{s \in \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}')} \mathcal{C}_{s, 3\tau_0, 3\delta_0}$ (see indeed (4.5), (4.6), (4.8) and Definition 4.3). Since $|\psi_{\mathbf{m}}| \leq 2e^{-(f-f(\mathbf{m}))/h}$, it follows that

$$|\langle \varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle| \leq \sum_{s \in \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}')} \frac{4}{\|\psi_{\mathbf{m}}\| \|\psi_{\mathbf{m}'}\|} \langle e^{-(f-f(\mathbf{m}))/h}, e^{-(f-f(\mathbf{m}'))/h} \rangle_{L^2(\mathcal{C}_{s, 3\tau_0, 3\delta_0})}.$$

Using (5.1) and $f(\mathbf{j}(\mathbf{m})) = f(\mathbf{j}(\mathbf{m}')) > \max(f(\mathbf{m}), f(\mathbf{m}'))$, the last relation implies (i) for $\tau_0 > 0$ and then $\delta_0 > 0$ small enough.

- (c) *The case where up to switching \mathbf{m} and \mathbf{m}' , $\sigma(\mathbf{m}) > \sigma(\mathbf{m}')$* . Here, according to Lemma 4.4 (v), either $\text{supp}(\psi_{\mathbf{m}}) \cap \text{supp}(\psi_{\mathbf{m}'}') = \emptyset$, in which case $\langle \varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'}' \rangle = 0$, or $\psi_{\mathbf{m}} = 2e^{-(f-f(\mathbf{m}))/h}$ on $\text{supp}(\psi_{\mathbf{m}'}')$, in which case, since $f \geq f(\mathbf{m}')$ on $\text{supp}(\psi_{\mathbf{m}'}')$, the Cauchy–Schwarz inequality gives

$$\langle \varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'}' \rangle = \frac{2}{\|\psi_{\mathbf{m}}\|} \langle e^{-(f-f(\mathbf{m}))/h}, \varphi_{\mathbf{m}'}' \rangle_{L^2(\text{supp}(\psi_{\mathbf{m}'}'))} = \frac{1}{\|\psi_{\mathbf{m}}\|} \mathcal{O}(e^{-(f(\mathbf{m}')-f(\mathbf{m}))/h}).$$

Relation (i) follows easily, using (5.1) and the relation $f(\mathbf{m}') > f(\mathbf{m})$ implied by the first part of (Gener).

In order to prove the remaining points (ii) to (iv) of Proposition 5.1, let us write $\psi_{\mathbf{m}} = \tilde{v}_{\mathbf{m}} \tilde{\psi}_{\mathbf{m}}$ with $\tilde{v}_{\mathbf{m}} = \theta_{\mathbf{m}}(1 + v_{\mathbf{m}})$ and $\tilde{\psi}_{\mathbf{m}} = e^{-(f-f(\mathbf{m}))/h}$. Using this notation and the decomposition $P = P_2 + P_1 + P_0$ with $P_0 = c$, $P_1 = \frac{1}{2}(b \cdot h \nabla + h \text{div} \circ b)$, and $P_2 = -h \text{div} \circ A \circ h \nabla$, we get

$$\begin{aligned} \langle P \psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle &= \langle (P_2 + P_0) \psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle = \langle Ah \nabla(\tilde{v}_{\mathbf{m}} \tilde{\psi}_{\mathbf{m}}), h \nabla(\tilde{v}_{\mathbf{m}} \tilde{\psi}_{\mathbf{m}}) \rangle + \langle P_0 \psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle \\ &= h^2 \langle \tilde{v}_{\mathbf{m}} A \nabla \tilde{\psi}_{\mathbf{m}}, \tilde{v}_{\mathbf{m}} \nabla \tilde{\psi}_{\mathbf{m}} \rangle + h^2 \langle \tilde{\psi}_{\mathbf{m}} A \nabla \tilde{v}_{\mathbf{m}}, \tilde{\psi}_{\mathbf{m}} \nabla \tilde{v}_{\mathbf{m}} \rangle \\ &\quad + 2h^2 \langle \tilde{\psi}_{\mathbf{m}} A \nabla \tilde{v}_{\mathbf{m}}, \tilde{v}_{\mathbf{m}} \nabla \tilde{\psi}_{\mathbf{m}} \rangle + \langle P_0 \psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle \\ &= -h^2 \langle \text{div}(\tilde{v}_{\mathbf{m}}^2 A \nabla \tilde{\psi}_{\mathbf{m}}), \tilde{\psi}_{\mathbf{m}} \rangle + h^2 \langle \tilde{\psi}_{\mathbf{m}} A \nabla \tilde{v}_{\mathbf{m}}, \tilde{\psi}_{\mathbf{m}} \nabla \tilde{v}_{\mathbf{m}} \rangle \\ &\quad + 2h^2 \langle \tilde{\psi}_{\mathbf{m}} A \nabla \tilde{v}_{\mathbf{m}}, \tilde{v}_{\mathbf{m}} \nabla \tilde{\psi}_{\mathbf{m}} \rangle + \langle P_0 \psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle \\ &= \langle \tilde{v}_{\mathbf{m}}^2 P_2 \tilde{\psi}_{\mathbf{m}}, \tilde{\psi}_{\mathbf{m}} \rangle + h^2 \langle \tilde{\psi}_{\mathbf{m}} A \nabla \tilde{v}_{\mathbf{m}}, \tilde{\psi}_{\mathbf{m}} \nabla \tilde{v}_{\mathbf{m}} \rangle + \langle \tilde{v}_{\mathbf{m}}^2 P_0 \tilde{\psi}_{\mathbf{m}}, \tilde{\psi}_{\mathbf{m}} \rangle, \end{aligned}$$

and since $(P_2 + P_0)(\tilde{\psi}_{\mathbf{m}}) = 0$, this implies

$$\langle P \psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle = h^2 \langle \tilde{\psi}_{\mathbf{m}} A \nabla \tilde{v}_{\mathbf{m}}, \tilde{\psi}_{\mathbf{m}} \nabla \tilde{v}_{\mathbf{m}} \rangle. \quad (5.2)$$

Since $f - f(\mathbf{m}) > S(\mathbf{m}) + \delta_0$ on $\text{supp}(\nabla \theta_{\mathbf{m}})$ (see (4.8)), it follows that

$$\langle P \psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle = h^2 \int \theta_{\mathbf{m}}^2 A \nabla v_{\mathbf{m}} \cdot \nabla v_{\mathbf{m}} e^{-2(f-f(\mathbf{m}))/h} dx + \mathcal{O}(e^{-2(S(\mathbf{m})+\delta_0)/h}).$$

On the other hand, thanks to (4.6), on $\bigcup_{s \in \mathbf{j}(\mathbf{m})} \mathcal{C}_{s, 3\tau_0, 3\delta_0}$ we have

$$\nabla v_{\mathbf{m}} = \sum_{s \in \mathbf{j}(\mathbf{m})} \frac{1}{C_{s,h}} \zeta(\ell_s/\tau_0) e^{-\ell_s^2/(2h)} \nabla \ell_s, \quad (5.3)$$

which yields, since $\nabla v_{\mathbf{m}} = 0$ on $E_{\mathbf{m}, 3\tau_0, 3\delta_0}^+ \cup E_{\mathbf{m}, 3\tau_0, 3\delta_0}^-$ by (4.5),

$$\langle P\psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle = h^2 \sum_{\mathbf{s} \in \mathbf{j}(\mathbf{m})} \frac{1}{C_{\mathbf{s}, h}^2} \int_{\mathcal{C}_{\mathbf{s}, 3\tau_0, 3\delta_0}} \theta_{\mathbf{m}}^2 \zeta(\ell_{\mathbf{s}}/\tau_0)^2 A \nabla \ell_{\mathbf{s}} \cdot \nabla \ell_{\mathbf{s}} e^{-2(f + \ell_{\mathbf{s}}^2/2 - f(\mathbf{m}))/h} dx + \mathcal{O}(e^{-2(S(\mathbf{m}) + \delta_0)/h}).$$

The first term of the right hand side can now be computed as in [20, proof of Proposition 4.5], the only difference being that here A and $\ell_{\mathbf{s}}$ depend on h and admit a classical expansion with respect to h . More precisely, since

$$f + \frac{\ell_{\mathbf{s}}^2}{2} - f(\mathbf{m}) = f + \frac{\ell_{\mathbf{s}, 0}^2}{2} - f(\mathbf{m}) + \ell_{\mathbf{s}, 0} \ell_{\mathbf{s}, 1} h + \mathcal{O}(h^2),$$

where, according to Lemma 3.2 and to (3.7), the function $f + \frac{\ell_{\mathbf{s}, 0}^2}{2}$ satisfies

$$\left(f + \frac{\ell_{\mathbf{s}, 0}^2}{2}\right)(\mathbf{s}) = f(\mathbf{s}) = \sigma(\mathbf{m}), \quad \nabla \left(f + \frac{\ell_{\mathbf{s}, 0}^2}{2}\right)(\mathbf{s}) = 0, \quad \text{Hess}\left(f + \frac{1}{2}\ell_{\mathbf{s}, 0}^2\right)(\mathbf{s}) > 0, \quad (5.4)$$

we can apply the Laplace method with the phase function $-2(f + \ell_{\mathbf{s}, 0}^2/2 - f(\mathbf{m}))$. Using $\ell_{\mathbf{s}, 0}(\mathbf{s}) = 0$ and (4.7), this yields

$$\langle P\psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle \in \mathcal{E}_{\text{cl}} \left(2h/\pi(\pi h)^{d/2} \sum_{\mathbf{s} \in \mathbf{j}(\mathbf{m})} (A^0 \nabla \ell_{\mathbf{s}, 0} \cdot \nabla \ell_{\mathbf{s}, 0})(\mathbf{s}) \times (\det \text{Hess}(f + \frac{1}{2}\ell_{\mathbf{s}, 0}^2)(\mathbf{s}))^{-1/2} e^{-2S(\mathbf{m})/h} \right).$$

Moreover, thanks to Lemma 3.3,

$$(A^0 \nabla \ell_{\mathbf{s}, 0} \cdot \nabla \ell_{\mathbf{s}, 0})(\mathbf{s}) = |\mu(\mathbf{s})| \quad \text{and} \quad \det \text{Hess}(f + \frac{1}{2}\ell_{\mathbf{s}, 0}^2)(\mathbf{s}) = |\det \text{Hess}(f)(\mathbf{s})|,$$

and hence

$$\langle P\psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle \in \mathcal{E}_{\text{cl}} \left(\frac{2h}{\pi} (\pi h)^{d/2} e^{-2S(\mathbf{m})/h} \sum_{\mathbf{s} \in \mathbf{j}(\mathbf{m})} |\mu(\mathbf{s})| D_{\mathbf{s}}^{-1} \right). \quad (5.5)$$

Combining (5.5) with (5.1) proves (ii).

Let us now prove (iii). Since $P(e^{-f/h}) = 0$, one has

$$\begin{aligned} P\psi_{\mathbf{m}} &= P(\theta_{\mathbf{m}}(v_{\mathbf{m}} + 1)e^{-(f-f(\mathbf{m}))/h}) \\ &= [P, \theta_{\mathbf{m}}](v_{\mathbf{m}} + 1)e^{-(f-f(\mathbf{m}))/h} + \theta_{\mathbf{m}}P(v_{\mathbf{m}}e^{-(f-f(\mathbf{m}))/h}), \end{aligned}$$

and since $f - f(\mathbf{m}) > S(\mathbf{m}) + \delta_0$ on $\text{supp}(\nabla \theta_{\mathbf{m}})$ and on $\text{supp}(1 - \theta_{\mathbf{m}}) \cap \text{supp}(\theta_{\mathbf{m}})$, this implies

$$\begin{aligned} \|P\psi_{\mathbf{m}}\|^2 &= \|\theta_{\mathbf{m}}P(v_{\mathbf{m}}e^{-(f-f(\mathbf{m}))/h})\|^2 + \mathcal{O}(e^{-2(S(\mathbf{m}) + \delta_0)/h}) \\ &= \|P(v_{\mathbf{m}}e^{-(f-f(\mathbf{m}))/h})\|_{L^2(\text{supp}(\theta_{\mathbf{m}}))}^2 + \mathcal{O}(e^{-2(S(\mathbf{m}) + \delta_0)/h}). \end{aligned} \quad (5.6)$$

On the other hand, on $\text{supp}(\theta_{\mathbf{m}})$, $P(v_{\mathbf{m}}e^{-(f-f(\mathbf{m}))/h})$ is supported in $\bigcup_{\mathbf{s} \in \mathbf{j}(\mathbf{m})} \mathcal{C}_{\mathbf{s}, 3\tau_0, 3\delta_0}$ by (4.5) and, on any $\mathcal{C}_{\mathbf{s}, 3\tau_0, 3\delta_0}$, one has (see (4.6))

$$P(v_{\mathbf{m}}e^{-(f-f(\mathbf{m}))/h}) = C_{\mathbf{s}, h}^{-1}(w+r)e^{-(f-f(\mathbf{m})+\ell_{\mathbf{s}}^2/2)/h},$$

where w and r are given by Lemma 3.1. Since $f + \ell_{\mathbf{s}}^2/2 = f + \ell_{\mathbf{s},0}^2/2 + \mathcal{O}(h)$ and the Hessian of $f + \ell_{\mathbf{s},0}^2/2$ at \mathbf{s} is positive definite (see (5.4)), and r is supported away from \mathbf{s} , one has, for some $\delta > 0$,

$$\|re^{-(f-f(\mathbf{m})+\ell_{\mathbf{s}}^2/2)/h}\|_{L^2(\mathcal{C}_{\mathbf{s}, 3\tau_0, 3\delta_0})}^2 = \mathcal{O}(e^{-2(S(\mathbf{m})+\delta)/h}) = \mathcal{O}(h^\infty)\langle P\psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle,$$

where we have used (5.5) to obtain the last equality. Moreover, thanks to Proposition 3.5 (and to Lemma 3.1), one also has

$$\|we^{-(f-f(\mathbf{m})+\ell_{\mathbf{s}}^2/2)/h}\|_{L^2(\mathcal{C}_{\mathbf{s}, 3\tau_0, 3\delta_0})}^2 = \mathcal{O}(h^\infty)e^{-2S(\mathbf{m})/h} = \mathcal{O}(h^\infty)\langle P\psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle.$$

These two estimates and (5.6) show that $\|P\psi_{\mathbf{m}}\|^2 = \mathcal{O}(h^\infty)\langle P\psi_{\mathbf{m}}, \psi_{\mathbf{m}} \rangle$, which proves (iii).

The proof of (iv) is similar and left to the reader. ■

5.2. Proof of Theorem 2

Until the end of this section, the local minimum points $\mathbf{m}_1, \dots, \mathbf{m}_{n_0}$ of f are labeled so that $(S(\mathbf{m}_j))_{j=1}^{n_0}$ is nondecreasing (see (1.22)). That is,

$$\text{for all } j \in \{1, \dots, n_0 - 1\}, \quad S(\mathbf{m}_{j+1}) \geq S(\mathbf{m}_j) \text{ and } S_{\mathbf{m}_{n_0}} = +\infty. \quad (5.7)$$

For $j \in \{1, \dots, n_0\}$, we will also denote for brevity

$$S_j := S(\mathbf{m}_j), \quad \varphi_j := \varphi_{\mathbf{m}_j, h}, \quad \tilde{\lambda}_j := \langle P\varphi_j, \varphi_j \rangle. \quad (5.8)$$

From Proposition 5.1, one knows that, for all $j \in \{1, \dots, n_0\}$,

$$\tilde{\lambda}_j \in \mathcal{E}_{\text{cl}}\left(he^{-2S_j/h} \sum_{\mathbf{s} \in \mathbf{j}(\mathbf{m}_j)} \frac{|\mu(\mathbf{s})|}{2\pi} \frac{D_{\mathbf{m}_j}}{D_{\mathbf{s}}} \right) \quad (5.9)$$

and, for $j, k \in \{1, \dots, n_0\}$,

$$\langle \varphi_j, \varphi_k \rangle = \delta_{j,k} + \mathcal{O}(e^{-C/h}), \quad \|P\varphi_j\| = \mathcal{O}(h^\infty \sqrt{\tilde{\lambda}_j}), \quad \|P^*\varphi_j\| = \mathcal{O}(h^{1/2} \sqrt{\tilde{\lambda}_j}). \quad (5.10)$$

Using in addition Lemma 4.4 (iv, v) together with the second part of (Gener) (see [20, proof of Lemma 4.7] for details), we also have

$$\langle P\varphi_j, \varphi_k \rangle = \delta_{j,k} \tilde{\lambda}_j. \quad (5.11)$$

We then introduce the spectral projector

$$\Pi_h = \frac{1}{2i\pi} \int_{\partial D(0, \varepsilon_* h/2)} (z - P)^{-1} dz, \quad (5.12)$$

with ε_* given by Proposition 1.2. Working as in [20, proof of Lemma 4.9], one deduces from the last two estimates of (5.10) and from the resolvent estimate (1.17) of Theorem 1 that

$$(1 - \Pi_h)\varphi_j = \mathcal{O}(h^\infty \sqrt{\tilde{\lambda}_j}) \quad \text{and} \quad (1 - \Pi_h^*)\varphi_j = \mathcal{O}(h^{-1/2} \sqrt{\tilde{\lambda}_j}). \quad (5.13)$$

The estimates (5.10), (5.11), and (5.13) easily imply the following proposition, whose proof is the same as the proof of [20, Proposition 4.10].

Proposition 5.3. *For every $j \in \{1, \dots, n_0\}$ and $h > 0$ small enough, define $v_j := \Pi_h \varphi_j$. Then there exists $c > 0$ such that, for all $j, k \in \{1, \dots, n_0\}$,*

$$\langle v_j, v_k \rangle = \delta_{j,k} + \mathcal{O}(e^{-c/h}), \quad (5.14)$$

$$\langle P v_j, v_k \rangle = \delta_{j,k} \tilde{\lambda}_j + \mathcal{O}(h^\infty \sqrt{\tilde{\lambda}_j \tilde{\lambda}_k}). \quad (5.15)$$

In particular, it follows from (5.14) that for every $h > 0$ small enough, (v_1, \dots, v_{n_0}) is a basis of $\text{Ran } \Pi_h$.

The end of the proof of Theorem 2 follows line for line [20, proof of Theorem 1.9, pp. 39–42]. First, we orthonormalize the basis $(v_{n_0-j+1})_{j=1}^{n_0}$ to a basis $(e_{n_0-j+1})_{j=1}^{n_0}$ by the Gram–Schmidt process. Thanks to (5.14), we obtain an orthonormal basis of $\text{Ran } \Pi_h$ such that, for every $j \in \{1, \dots, n_0\}$,

$$e_{n_0-j+1} = v_{n_0-j+1} + \mathcal{O}(e^{-C/h})$$

(see [20, Lemma 4.11]). We then show, using (5.15) and the above labeling of the basis $(e_{n_0-j+1})_{j=1}^{n_0}$ starting from e_{n_0} , that, for all $j, k \in \{1, \dots, n_0\}$,

$$\langle P e_{n_0-j+1}, e_{n_0-k+1} \rangle = \delta_{j,k} \tilde{\lambda}_{n_0-j+1} + \mathcal{O}(h^\infty \sqrt{\tilde{\lambda}_{n_0-j+1} \tilde{\lambda}_{n_0-k+1}}) \quad (5.16)$$

(see [20, Proposition 4.12]). We can then compute the eigenvalues of the matrix M_h of $P|_{\text{Ran } \Pi_h}$ in this basis, using the so-called graded structure of M_h (see [20, pp. 41–42 and Theorem A.4] or Theorem 4 of the next section). Since $\tilde{\lambda}_j$ (or more precisely $e^{2S_j/h} \tilde{\lambda}_j$, see (5.9)) admits a classical expansion, and thanks to the h^∞ multiplicative errors in (5.16), we obtain the announced result.

5.3. Proof of Corollaries 1.6 and 1.7

Recall that Π_h , the spectral projector of P associated with the n_0 exponentially small eigenvalues of P , has been defined in (5.12). Since $\partial D(0, \varepsilon_* h/2)$ is at a distance of order h from the spectrum of P (see Proposition 1.2), (1.17) implies that there exists $C > 0$ such that, for every h small enough,

$$\|\Pi_h\| \leq C. \quad (5.17)$$

Then the estimates (1.17) and (5.17) imply that $(P - z)^{-1}(1 - \Pi_h)$ is of order h^{-1} in $\{\operatorname{Re} z < 2\varepsilon_* h/3\} \setminus D(0, \varepsilon_* h/2)$. Since this operator-valued function is holomorphic in $\{\operatorname{Re} z < 2\varepsilon_* h/3\}$, the maximum principle yields

$$\|(P - z)^{-1}(1 - \Pi_h)\| \leq Ch^{-1} \quad (5.18)$$

for $\operatorname{Re} z < 2\varepsilon_* h/3$ and h small enough.

The solution of (1.27) can be written as

$$u(t) = e^{-tP/h}u_0 = e^{-tP/h}\Pi_h u_0 + e^{-tP/h}(1 - \Pi_h)u_0. \quad (5.19)$$

Let $Q : \operatorname{Im}(1 - \Pi_h) \rightarrow \operatorname{Im}(1 - \Pi_h)$ be the operator P restricted to the Hilbert space $\operatorname{Im}(1 - \Pi_h)$. Since P is maximal accretive, so is Q and thus $\|e^{-tQ/h}\| \leq 1$. Moreover, (5.18) shows that $\|(Q - z)^{-1}\| \leq Ch^{-1}$ for $\operatorname{Re} z < 2\varepsilon_* h/3$. To estimate the last term in (5.19), we use a Gearhart–Prüss type inequality with an explicit bound. More precisely, [12, Theorem 1.4] (see [11, Proposition 2.1] for more details) gives, for some $C > 0$ and all $t \geq 0$,

$$\|e^{-tQ/h}\| \leq \left(1 + 2\frac{\varepsilon_* h}{2} \sup_{\operatorname{Re} z = \varepsilon_* h/2} \|(Q - z)^{-1}\|\right) e^{-t\frac{\varepsilon_* h}{2}/h} \leq Ce^{-t\varepsilon_*/2}.$$

Combined with (5.17), this implies the existence of $C > 0$ such that, for all $t \geq 0$,

$$\|e^{-tP/h}(1 - \Pi_h)u_0\| \leq Ce^{-\varepsilon t}\|u_0\|, \quad (5.20)$$

where $\varepsilon = \varepsilon_*/2$. On the other hand, P restricted to $\operatorname{Im} \Pi_h$ is a matrix of size n_0 whose eigenvalues are the $\lambda(\mathbf{m}, h)$'s. Then (5.19), (5.20), and the usual formula for the exponential of a matrix applied to $e^{-tP/h}\Pi_h u_0$ provide (1.28). Moreover, using (5.20) instead of the argument of [20, p. 43], the proof of (1.29) is similar to the one of [20, Theorem 1.11], except we have here to apply Theorem 4 instead of [20, Theorem A.4], and we omit the details. Summing up, we have just shown Corollary 1.6.

Let us now prove Corollary 1.7. For $R > 1$, we define the balls

$$\forall k \in \{1, \dots, K-1\}, \quad D_k = D((R + R^{-1})he^{-2S_k/h}, Rhe^{-2S_k/h}),$$

and $D_K = D(0, R^{-1}he^{-2S_{K-1}/h})$. For R fixed large enough and every h small enough, each exponentially small eigenvalue of P belongs to exactly one of the disjoint sets D_k from Theorem 2. Moreover, ∂D_k is at distance of order $he^{-2S_k/h}$ (resp. $he^{-2S_{K-1}/h}$) from the spectrum of P for $k \in \{1, \dots, K-1\}$ (resp. $k = K$). Using (6.18) to estimate the resolvent of P on the image of Π_h and (5.18) to control the contribution on the image of $1 - \Pi_h$, we get

$$\forall z \in \partial D_k, \quad \|(P - z)^{-1}\| \leq \begin{cases} Ch^{-1}e^{2S_k/h} & \text{for } k \in \{1, \dots, K-1\}, \\ Ch^{-1}e^{2S_{K-1}/h} & \text{for } k = K. \end{cases} \quad (5.21)$$

In particular, the spectral projector associated with the eigenvalues of order $he^{-2S_k/h}$,

$$\tilde{\Pi}_k = \frac{1}{2i\pi} \int_{\partial D_k} (z - P)^{-1} dz, \quad (5.22)$$

is well defined and satisfies $\|\tilde{\Pi}_k\| \leq C$.

We can now decompose

$$e^{-tP/h}\Pi_h = \sum_{1 \leq k \leq K} e^{-tP/h}\tilde{\Pi}_k. \quad (5.23)$$

For $k \in \{1, \dots, K-1\}$ and $0 \leq t \leq t_k^-$, (5.21) and $t_k^- e^{-2S_k/h} = \mathcal{O}(h^\infty)$ imply

$$\begin{aligned} e^{-tP/h}\tilde{\Pi}_k &= \frac{1}{2i\pi} \int_{\partial D_k} e^{-tz/h} (z-P)^{-1} dz \\ &= \frac{1}{2i\pi} \int_{\partial D_k} (z-P)^{-1} dz + \frac{1}{2i\pi} \int_{\partial D_k} (e^{-tz/h} - 1)(z-P)^{-1} dz \\ &= \tilde{\Pi}_k + \int_{\partial D_k} \mathcal{O}(t|z|/h) \|(P-z)^{-1}\| dz \\ &= \tilde{\Pi}_k + \mathcal{O}(t h e^{-2S_k/h}/h) \\ &= \tilde{\Pi}_k + \mathcal{O}(h^\infty). \end{aligned} \quad (5.24)$$

On the contrary, for $t_k^+ \leq t$, (5.21) and $e^{-t_k^+} e^{-2S_k/h/R} = \mathcal{O}(h^\infty)$ give

$$\begin{aligned} e^{-tP/h}\tilde{\Pi}_k &= \frac{1}{2i\pi} \int_{\partial D_k} e^{-tz/h} (z-P)^{-1} dz \\ &= \mathcal{O}\left(\int_{\partial D_k} e^{-t \operatorname{Re} z/h} \|(P-z)^{-1}\| dz\right) \\ &= \mathcal{O}(e^{-tR^{-1} h e^{-2S_k/h}/h}) \int_{\partial D_k} \|(P-z)^{-1}\| dz \\ &= \mathcal{O}(h^\infty). \end{aligned} \quad (5.25)$$

Lastly, $e^{-tP/h}\tilde{\Pi}_K = \tilde{\Pi}_K$ since $\tilde{\Pi}_K$ is the rank 1 spectral projector associated with the eigenvalue 0. On the other hand, since $e^{-\varepsilon t_0^+} = \mathcal{O}(h^\infty)$, (5.20) becomes

$$\|e^{-tP/h}(1 - \Pi_h)\| = \mathcal{O}(h^\infty) \quad (5.26)$$

for $t \geq t_0^+$. Summing up, Corollary 1.7 is a direct consequence of formulas (5.19) and (5.23) together with the relation $\Pi_k^{\leq} = \sum_{k \leq j \leq K} \tilde{\Pi}_j$ and the estimates (5.24)–(5.26).

6. Generalization

In this part, we briefly explain how one can drop the assumption (Gener) and treat the general case in the spirit of [24]. This requires introducing some additional material of [24].

Definition 6.1. Let $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$. We say that \mathbf{m} is of type I when $f(\widehat{\mathbf{m}}) < f(\mathbf{m})$, and of type II when $f(\widehat{\mathbf{m}}) = f(\mathbf{m})$. We will also denote

$$\begin{aligned} \mathcal{U}^{(0),I} &:= \{\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}; \mathbf{m} \text{ is of type I}\}, \\ \mathcal{U}^{(0),II} &:= \{\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}; \mathbf{m} \text{ is of type II}\}. \end{aligned}$$

We have clearly the following disjoint union $\mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\} = \mathcal{U}^{(0),I} \sqcup \mathcal{U}^{(0),II}$.

Given $\sigma \in \Sigma$, let $\Omega_\sigma := \{E(\mathbf{m}); \mathbf{m} \in \sigma^{-1}(\sigma)\} \cup \{\widehat{E}(\mathbf{m}); \mathbf{m} \in \sigma^{-1}(\sigma) \cap \mathcal{U}^{(0),II}\}$.

Definition 6.2. We define an equivalence relation \mathcal{R} on $\mathcal{U}^{(0)}$ by $\mathbf{m}\mathcal{R}\mathbf{m}'$ if and only if

$$\begin{cases} \sigma(\mathbf{m}) = \sigma(\mathbf{m}') = \sigma, \\ \exists \omega_1, \dots, \omega_K \in \Omega_\sigma, \mathbf{m} \in \omega_1, \mathbf{m}' \in \omega_K, \text{ and } \forall k = 1, \dots, K-1, \overline{\omega_k} \cap \overline{\omega_{k+1}} \neq \emptyset. \end{cases}$$

We denote by $\text{Cl}(\mathbf{m})$ the equivalence class of \mathbf{m} for the relation \mathcal{R} . Observe that $\text{Cl}(\underline{\mathbf{m}}) = \{\underline{\mathbf{m}}\}$ since $\underline{\mathbf{m}}$ is the only $\mathbf{m} \in \mathcal{U}^{(0)}$ such that $\sigma(\mathbf{m}) = +\infty$. Let \mathcal{A} denote the (finite) set of equivalence classes for \mathcal{R} and, for $\alpha \in \mathcal{A}$, let $\mathcal{U}_\alpha^{(0)}$ be the set of the elements of the class α . Then evidently

$$\mathcal{U}^{(0)} = \bigsqcup_{\alpha \in \mathcal{A}} \mathcal{U}_\alpha^{(0)} = \{\underline{\mathbf{m}}\} \sqcup \bigsqcup_{\alpha \in \mathcal{A} \setminus \{\text{Cl}(\underline{\mathbf{m}})\}} \mathcal{U}_\alpha^{(0)}.$$

In the theorem below, we sum up in a rather vague way the description of the small eigenvalues of P . Precise statements are given in Theorems 5 and 6.

Theorem 3. Suppose that the assumptions of Theorem 1 are satisfied and that (Gibbs), (Confin), and (Morse) hold true. Let ε_* be given by Proposition 1.2. Then, for $h > 0$ small enough, there exists a bijection, taking into account multiplicities,

$$\beta : \{0\} \cup \bigcup_{\alpha \in \mathcal{A} \setminus \{\text{Cl}(\underline{\mathbf{m}})\}} \bigcup_{j=1}^p h e^{-2S_j/h} \sigma(M_{\alpha,j}) \rightarrow \sigma(P) \cap \{\text{Re } z < \varepsilon_* h\},$$

with $\beta(z) = z + \mathcal{O}(h^\infty |z|)$, for some symmetric positive definite matrices $M_{\alpha,j}$ having a classical expansion in powers of h with explicit invertible leading term and for some labeling $(S_j)_{j=1}^p$ of $S(\mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\})$.

Here, the set $h e^{-2S_j/h} \sigma(M_{\alpha,j})$ is empty whenever $S_j \notin S(\mathcal{U}_\alpha^{(0)})$. The general strategy to prove Theorem 3 (leading to an explicit expression for the matrices $M_{\alpha,j}$) is to combine the quasimodal constructions near the saddle points developed in the preceding section together with the topological classification of the saddle points introduced in [24]. In that work, the construction of the quasimode $\varphi_{\mathbf{m}}$ depends on the fact that \mathbf{m} is of type I or II. In order to lighten the presentation, we assume from now on that every point \mathbf{m} is of type I and we will prove Theorem 3 under this assumption, i.e. when $\mathcal{U}^{(0),II} = \emptyset$. In that case, the leading term of $M_{\alpha,j}$ is given in Theorem 6 below, which makes explicit the statement of Theorem 3 when $\mathcal{U}^{(0),II} = \emptyset$. The reader may check that the construction below can be adapted to the case of type II points as in [24].

Remark 6.3. Note that $\mathcal{U}^{(0),II} = \emptyset$ if and only if $f(\widehat{\mathbf{m}}) < f(\mathbf{m})$ for every $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$ (see Definition 6.1). It follows that $\mathcal{U}^{(0),II} = \emptyset$ if and only if the first part of (Gener) is satisfied. Indeed, if \mathbf{m} is the unique global minimum of $f|_{E(\mathbf{m})}$ for every $\mathbf{m} \in \mathcal{U}^{(0)}$, then, for every $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$, the relation $\mathbf{m} \in E(\widehat{\mathbf{m}})$ implies $f(\widehat{\mathbf{m}}) < f(\mathbf{m})$. Conversely, by contraposition, assume the existence of $\widehat{\mathbf{m}}' \neq \mathbf{m}' \in \mathcal{U}^{(0)}$ such that $\mathbf{m}' \in E(\mathbf{m})$ and $f(\mathbf{m}') = f(\mathbf{m})$. Then $\sigma(\mathbf{m}) > \sigma(\mathbf{m}')$ and thus $\widehat{\mathbf{m}}' \in E(\mathbf{m})$, which implies $f(\mathbf{m}) \leq f(\widehat{\mathbf{m}}') \leq f(\mathbf{m}')$ (see (4.2)), and then $f(\mathbf{m}') = f(\widehat{\mathbf{m}}')$, i.e. $\mathbf{m}' \in \mathcal{U}^{(0),II} \neq \emptyset$.

In particular, the statement of Proposition 5.1 is valid when $\mathcal{U}^{(0),II} = \emptyset$. This will be used later.

6.1. Quasimodal constructions for type I minima

Let $(\psi_{\mathbf{m}})_{\mathbf{m} \in \mathcal{U}^{(0)}}$ denote the family of quasimodes of Definition 4.3. We recall that when $\mathbf{m} \neq \bar{\mathbf{m}}$,

$$\psi_{\mathbf{m}}(x) = \theta_{\mathbf{m}}(x)(v_{\mathbf{m}}(x) + 1)e^{-(f(x)-f(\mathbf{m}))/h},$$

with $\theta_{\mathbf{m}}$ and $v_{\mathbf{m}}$ defined by (4.5), (4.6), and (4.8) (here and throughout we drop the subscript h to lighten the notation). In particular, near any point $\mathbf{s} \in \mathbf{j}(\mathbf{m})$,

$$v_{\mathbf{m}}(x) = C_{s,h}^{-1} \int_0^{\ell_s(x,h)} \zeta(r/\tau_0) e^{-r^2/(2h)} dr,$$

where ℓ_s is the function defined by Proposition 3.5 such that there exists a neighborhood V of \mathbf{s} such that $E(\mathbf{m}) \cap V \subset \{\eta(\mathbf{s}) \cdot (x - \mathbf{s}) > 0\}$ (see the lines below (4.6)). This choice of sign depends of course on \mathbf{m} and in order to avoid any confusion, we shall denote by $\ell_{s,\mathbf{m}}$ the function ℓ_s above.

Lemma 6.4. *Let $\mathbf{m} \neq \mathbf{m}' \in \mathcal{U}^{(0)}$ satisfy $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') \neq \emptyset$ and take $\mathbf{s} \in \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}')$. Then the functions $\ell_{s,\mathbf{m}}$ and $\ell_{s,\mathbf{m}'}$ can be chosen so that $\ell_{s,\mathbf{m}'} = -\ell_{s,\mathbf{m}}$ near \mathbf{s} .*

Proof. Note that the function $\ell_{s,\mathbf{n}}$ only makes sense when $\mathbf{n} \in \{\mathbf{m}, \mathbf{m}'\}$. We assume that $\ell_{s,\mathbf{m}}$ is given by Proposition 3.5 with the sign condition for \mathbf{m} below (4.6). As explained at the end of Section 3.3, $-\ell_{s,\mathbf{m}}$ also satisfies Proposition 3.5 with the opposite sign condition. Thus, this function satisfies the sign condition for \mathbf{m}' and can be chosen as the function $\ell_{s,\mathbf{m}'}$. ■

Proposition 6.5. *Assume that the hypotheses of Theorem 3 hold true and $\mathcal{U}^{(0),II} = \emptyset$. Then the conclusions of Proposition 5.1 are satisfied. Moreover, for all $\mathbf{m}, \mathbf{m}' \in \mathcal{U}^{(0)}$, one has*

$$\langle P\varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle \in \mathcal{E}_{\text{cl}} \left((-1)^{\delta_{\mathbf{m},\mathbf{m}'}-1} h e^{-(S(\mathbf{m})+S(\mathbf{m}'))/h} \sum_{\mathbf{s} \in \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}')} \frac{|\mu(\mathbf{s})|}{2\pi} \frac{\sqrt{D_{\mathbf{m}} D_{\mathbf{m}'}}}{D_{\mathbf{s}}} \right). \quad (6.1)$$

Finally, denoting $P^{\sharp} = P_2 + P_0$, there exists $\alpha > 0$ such that

$$\langle P\varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle = \langle P^{\sharp}\varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle (1 + \mathcal{O}(e^{-\alpha/h})). \quad (6.2)$$

Remark 6.6. (i) Note that relation (6.1) implies that

$$\langle P\varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle = 0 \quad (6.3)$$

for all $\mathbf{m}, \mathbf{m}' \in \mathcal{U}^{(0)}$ with $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset$. This is in particular the case when $\mathbf{m}' \notin \text{Cl}(\mathbf{m})$ (see Definition 6.2).

(ii) When $\mathcal{U}^{(0),II} \neq \emptyset$, the family $(\varphi_{\mathbf{m},h})_{\mathbf{m} \in \mathcal{U}^{(0)}}$ of quasimodes is not quasi-orthonormal and thus does not satisfy (i) of Proposition 5.1 (see indeed Remarks 5.2 and 6.3).

Proof of Proposition 6.5. Thanks to Remark 6.3 and $\mathcal{U}^{(0),II} = \emptyset$, the assumptions of Proposition 5.1 are satisfied.

Let us now prove (6.1) and take $\mathbf{m}, \mathbf{m}' \in \mathcal{U}^{(0)}$. When $\mathbf{m}' = \mathbf{m}$, this is exactly (i) of Proposition 5.1 and so we can assume that $\mathbf{m}' \neq \mathbf{m}$.

Assume moreover that $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset$. If $\sigma(\mathbf{m}) > \sigma(\mathbf{m}')$, then we have $\varphi_{\mathbf{m}} = C_{\mathbf{m},h} e^{-(f-f(\mathbf{m}))/h}$ on $\text{supp}(\varphi_{\mathbf{m}'})$ for some constant $C_{\mathbf{m},h}$. Hence $P\varphi_{\mathbf{m}} = 0$ on $\text{supp}(\varphi_{\mathbf{m}'})$, which implies $\langle P\varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle = 0$. If $\sigma(\mathbf{m}) < \sigma(\mathbf{m}')$, the same argument works since $P^*(e^{-f/h}) = 0$. Assume now that $\sigma(\mathbf{m}) = \sigma(\mathbf{m}')$. According to Lemma 4.4 (iv), it follows that $\text{supp}(\varphi_{\mathbf{m}}) \cap \text{supp}(\varphi_{\mathbf{m}'}) = \emptyset$ and so $\langle P\varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle = 0$. Summing up, if $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset$, we always have $\langle P\varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle = 0$, which is precisely (6.1) in that case.

Assume finally that $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') \neq \emptyset$ and so $\sigma(\mathbf{m}) = \sigma(\mathbf{m}')$. As in the proof of Proposition 5.1, we denote $\psi_{\mathbf{m}} = \tilde{v}_{\mathbf{m}} \tilde{\psi}_{\mathbf{m}}$ with $\tilde{v}_{\mathbf{m}} = \theta_{\mathbf{m}}(1 + v_{\mathbf{m}})$ and $\tilde{\psi}_{\mathbf{m}} = e^{-(f-f(\mathbf{m}))/h}$. Using the identity $\tilde{\psi}_{\mathbf{m}} \nabla \tilde{\psi}_{\mathbf{m}'} = \tilde{\psi}_{\mathbf{m}'} \nabla \tilde{\psi}_{\mathbf{m}}$ and working as in the proof of Proposition 5.1, we get

$$\langle (P_2 + P_0)\psi_{\mathbf{m}}, \psi_{\mathbf{m}'} \rangle = h^2 \langle \tilde{\psi}_{\mathbf{m}} A \nabla \tilde{v}_{\mathbf{m}}, \tilde{\psi}_{\mathbf{m}'} A \nabla \tilde{v}_{\mathbf{m}'} \rangle. \quad (6.4)$$

Observe now that $\theta_{\mathbf{m}} = \theta_{\mathbf{m}'}$ (see (4.8)) and

$$\text{supp}(\theta_{\mathbf{m}}(v_{\mathbf{m}} + 1)(v_{\mathbf{m}'} + 1)) \subset \bigcup_{s \in \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}')} \mathcal{C}_{s, 3\tau_0, 3\delta_0} \quad (6.5)$$

by (4.5). Since $\ell_{s,\mathbf{m}'} = -\ell_{s,\mathbf{m}}$ (see Lemma 6.4) and hence $v_{\mathbf{m}'} = -v_{\mathbf{m}}$ on the support of $\theta_{\mathbf{m}}(v_{\mathbf{m}} + 1)(v_{\mathbf{m}'} + 1)$ (see (4.6)), we have

$$\langle P_1 \psi_{\mathbf{m}}, \psi_{\mathbf{m}'} \rangle = \langle P_1(\theta_{\mathbf{m}}(v_{\mathbf{m}} + 1)\tilde{\psi}_{\mathbf{m}}), \theta_{\mathbf{m}}(-v_{\mathbf{m}} + 1)\tilde{\psi}_{\mathbf{m}'} \rangle.$$

Since in addition P_1 is formally anti-self-adjoint and $P_1(e^{-f/h}) = 0$, and $f - \delta_0 > \sigma(\mathbf{m}) = \sigma(\mathbf{m}')$ on the compact set $\text{supp}(\nabla \theta_{\mathbf{m}})$ (see (4.8)), we have

$$\begin{aligned} \langle P_1 \psi_{\mathbf{m}}, \psi_{\mathbf{m}'} \rangle &= -\langle P_1(\theta_{\mathbf{m}} v_{\mathbf{m}} \tilde{\psi}_{\mathbf{m}}), \theta_{\mathbf{m}} v_{\mathbf{m}} \tilde{\psi}_{\mathbf{m}'} \rangle + \mathcal{O}(e^{-(S(\mathbf{m})+S(\mathbf{m}')+2\delta_0)/h}) \\ &= -e^{(f(\mathbf{m}')-f(\mathbf{m}))/h} \langle P_1(\theta_{\mathbf{m}} v_{\mathbf{m}} \tilde{\psi}_{\mathbf{m}}), \theta_{\mathbf{m}} v_{\mathbf{m}} \tilde{\psi}_{\mathbf{m}} \rangle + \mathcal{O}(e^{-(S(\mathbf{m})+S(\mathbf{m}')+2\delta_0)/h}) \\ &= \mathcal{O}(e^{-(S(\mathbf{m})+S(\mathbf{m}')+2\delta_0)/h}). \end{aligned} \quad (6.6)$$

Combining (6.4) and (6.6), we get

$$\langle P \psi_{\mathbf{m}}, \psi_{\mathbf{m}'} \rangle = h^2 \langle \tilde{\psi}_{\mathbf{m}} A \nabla \tilde{v}_{\mathbf{m}}, \tilde{\psi}_{\mathbf{m}'} \nabla \tilde{v}_{\mathbf{m}'} \rangle + \mathcal{O}(e^{-(S(\mathbf{m})+S(\mathbf{m}')+2\delta_0)/h}). \quad (6.7)$$

Using (5.3), (6.5), and $f - \delta_0 > \sigma(\mathbf{m}) = \sigma(\mathbf{m}')$ on $\text{supp}(\nabla \theta_{\mathbf{m}})$, this implies

$$\begin{aligned} \langle P \psi_{\mathbf{m}}, \psi_{\mathbf{m}'} \rangle &= h^2 \sum_{s \in \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}')} \frac{1}{C_{s,h}^2} \int_{\mathcal{C}_{s, 3\tau_0, 3\delta_0}} \theta_{\mathbf{m}}^2 \zeta(\ell_{s,\mathbf{m}}/\tau_0) \zeta(\ell_{s,\mathbf{m}'}/\tau_0) \\ &\quad \times A \nabla \ell_{s,\mathbf{m}} \cdot \nabla \ell_{s,\mathbf{m}'} e^{-(2f + \ell_{s,\mathbf{m}}^2/2 + \ell_{s,\mathbf{m}'}^2/2 - f(\mathbf{m}) - f(\mathbf{m}'))/h} dx \\ &\quad + \mathcal{O}(e^{-(S(\mathbf{m})+S(\mathbf{m}')+2\delta_0)/h}). \end{aligned}$$

Since $\ell_{s,\mathbf{m}'} = -\ell_{s,\mathbf{m}}$ on $\mathcal{C}_{s,3\tau_0,3\delta_0}$ and ζ is even, it follows that

$$\begin{aligned} \langle P\psi_{\mathbf{m}}, \psi_{\mathbf{m}'} \rangle &= h^2 \sum_{s \in \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}')} \frac{-1}{C_{s,h}^2} \int \theta_{\mathbf{m}}^2 \zeta(\ell_{s,\mathbf{m}}/\tau_0)^2 A \nabla \ell_{s,\mathbf{m}} \cdot \nabla \ell_{s,\mathbf{m}} \\ &\quad \times e^{-(2f + \ell_{s,\mathbf{m}}^2 - f(\mathbf{m}) - f(\mathbf{m}'))/h} d\mathbf{x} \\ &\quad + \mathcal{O}(e^{-(S(\mathbf{m}) + S(\mathbf{m}') + 2\delta_0)/h}). \end{aligned}$$

Since $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') \neq \emptyset$, this quantity can be computed by the Laplace method as in the proof of Proposition 5.1. We obtain

$$\langle P\psi_{\mathbf{m}}, \psi_{\mathbf{m}'} \rangle \in \mathcal{E}_{\text{cl}} \left(\frac{-2h}{\pi} (\pi h)^{d/2} e^{-(S(\mathbf{m}) + S(\mathbf{m}'))/h} \sum_{s \in \mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}')} |\mu(s)| D_s^{-1} \right). \quad (6.8)$$

Relation (6.1) follows, using also (5.1).

Finally, (6.2) follows from (6.4), (6.6), (6.7), and (6.8) when $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') \neq \emptyset$, and when $\mathbf{j}(\mathbf{m}) \cap \mathbf{j}(\mathbf{m}') = \emptyset$, from $\langle P\varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle = 0 = \langle P^\# \varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle$, where the last equality can be proved as was the first one in the third paragraph, using $(P_2 + P_0)(e^{-f/h}) = (P_2 + P_0)^*(e^{-f/h}) = 0$. ■

6.2. Graded structure of the interaction matrix

Suppose from now on that the minimum points $\mathbf{m} \in \mathcal{U}^{(0)}$ of f are labeled as in (5.7). Let $(\tilde{e}_j)_{j=1}^{n_0}$ denote the basis of $\text{Ran } \Pi_h$ obtained from the sequence $(\Pi_h \varphi_{\mathbf{m}_{n_0-j+1}})_{j=1}^{n_0}$ by the Gram–Schmidt process, and let $e_j = \tilde{e}_{n_0-j+1}$ for $j = 1, \dots, n_0$. We recall that, taking the ε_* of Proposition 1.2,

$$\Pi_h = \frac{1}{2i\pi} \int_{\partial D(0, \varepsilon_* h/2)} (z - P)^{-1} dz$$

denotes the spectral projector associated with the n_0 eigenvalues of P of order $\mathcal{O}(h^{1+\alpha})$, $\alpha > 0$.

Let $\Upsilon = (v_{i,j})_{i,j=1}^{n_0-1}$ denote the matrix with coefficients

$$v_{i,j} = \langle P e_i, e_j \rangle. \quad (6.9)$$

Introduce also the matrix $\Upsilon^\# = (v_{i,j}^\#)_{i,j=1}^{n_0-1}$ defined by

$$v_{i,j}^\# = \langle P^\# \varphi_{\mathbf{m}_i}, \varphi_{\mathbf{m}_j} \rangle = \langle (P_2 + P_0) \varphi_{\mathbf{m}_i}, \varphi_{\mathbf{m}_j} \rangle. \quad (6.10)$$

Note that, in these definitions, we do not consider the contribution of the vectors e_{n_0} and $\varphi_{\mathbf{m}_{n_0}}$ which are collinear to $e^{-f/h}$ and so belong to the kernels of P , P^* , and $P^\#$. If we had added these last vectors in the definitions of Υ and of $\Upsilon^\#$, the last line and column of these matrices would have consisted of zeros. In particular, $\sigma(P|_{\text{Ran } \Pi_h}) = \sigma(\Upsilon) \cup \{0\}$.

In order to compute the spectrum of the matrix Υ and then the spectrum of $P|_{\text{Ran } \Pi_h}$, we recall some tools from [24, Section 5B]. We denote by $\mathcal{S}^+(E)$ the set of symmetric positive definite matrices on a vector space E , and by $\mathcal{S}_{\text{cl}}^+(E)$ the set of

h -depending matrices $M(h) \in \mathcal{S}^+(E)$ admitting a classical expansion $M(h) \sim \sum_j h^j M_j$ with $M_0 \in \mathcal{S}^+(E)$. We will sometimes forget E and write for short $\mathcal{S}^+, \mathcal{S}_{\text{cl}}^+$.

Definition 6.7. Let $\mathcal{E} = (E_j)_{j=1}^p$ be a sequence of finite-dimensional vector spaces, $E = \bigoplus_{j=1}^p E_j$, and let $\tau = (\tau_2, \dots, \tau_p) \in (\mathbb{R}_+^*)^{p-1}$. Let $\tau \mapsto \mathcal{M}(\tau)$ be a map from $(\mathbb{R}_+^*)^{p-1}$ into the set of matrices $\mathcal{M}(E)$.

- ★ We say that $\mathcal{M}(\tau)$ is an (\mathcal{E}, τ) -graded symmetric matrix if there exists $M \in \mathcal{S}^+(E)$ independent of τ such that

$$\mathcal{M}(\tau) = \Omega(\tau) M \Omega(\tau), \quad (6.11)$$

with $\Omega(\tau) = \text{diag}(\varepsilon_1(\tau) \text{Id}_{E_1}, \dots, \varepsilon_p(\tau) \text{Id}_{E_p})$, $\varepsilon_1(\tau) = 1$ and $\varepsilon_j(\tau) = \prod_{k=2}^j \tau_k$ for $j \geq 2$.

- ★ We say that a family of (\mathcal{E}, τ) -graded symmetric matrices $\mathcal{M}_h(\tau)$, $h \in]0, h_0]$, is classical if $\mathcal{M}_h(\tau) = \Omega(\tau) M_h \Omega(\tau)$ for some matrix $M_h \in \mathcal{S}_{\text{cl}}^+(E)$.
- ★ We say that $\mathcal{M}_h(\tau)$ is an (\mathcal{E}, τ) -graded almost symmetric matrix if there exists $M_h \in \mathcal{M}(E)$ such that

$$\mathcal{M}_h(\tau) = \Omega(\tau) M_h \Omega(\tau), \quad M_h + M_h^* \in \mathcal{S}^+(E), \quad M_h - M_h^* = \mathcal{O}(h^\infty). \quad (6.12)$$

- ★ We say that a family of (\mathcal{E}, τ) -graded almost symmetric matrices $\mathcal{M}_h(\tau)$, $h \in]0, h_0]$, is classical if the matrix M_h in (6.12) satisfies $M_h + M_h^* \in \mathcal{S}_{\text{cl}}^+(E)$.

Throughout, we denote by $\text{GS}(\mathcal{E}, \tau)$ (resp. $\text{GS}_{\text{cl}}(\mathcal{E}, \tau)$) the set of (resp. classical) (\mathcal{E}, τ) -graded symmetric matrices, and by $\text{GAS}(\mathcal{E}, \tau)$ (resp. $\text{GAS}_{\text{cl}}(\mathcal{E}, \tau)$) the set of (resp. classical) (\mathcal{E}, τ) -graded almost symmetric matrices.

For any $\mathcal{M} \in \mathcal{M}(E)$, we denote $\mathcal{M}^s = \frac{1}{2}(M + M^*)$ its real part. Obviously, the real part of a matrix in $\text{GAS}(\mathcal{E}, \tau)$ (resp. $\text{GAS}_{\text{cl}}(\mathcal{E}, \tau)$) belongs to $\text{GS}(\mathcal{E}, \tau)$ (resp. $\text{GS}_{\text{cl}}(\mathcal{E}, \tau)$).

Let $\{S_1 \leq \dots \leq S_p\}$ denote the set $\{S(\mathbf{m}_j); j = 1, \dots, n_0 - 1\}$ and, for all $k = 1, \dots, p$, let E_k denote the vector space generated by $\{e_r; S(\mathbf{m}_r) = S_k\}$. We also set $\tau_k = e^{-(S_k - S_{k-1})/h}$ for any $k = 2, \dots, p$.

Proposition 6.8. Let $\mathcal{E} = (E_1, \dots, E_p)$ and $\tau = (\tau_2, \dots, \tau_p)$ be as above. Then

$$h^{-1} e^{2S_1/h} \Upsilon^\# \in \text{GS}_{\text{cl}}(\mathcal{E}, \tau) \quad \text{and} \quad h^{-1} e^{2S_1/h} \Upsilon \in \text{GAS}_{\text{cl}}(\mathcal{E}, \tau).$$

Moreover,

$$e^{2S_1/h} \Upsilon = e^{2S_1/h} \Upsilon^\# + \Omega(\tau) \mathcal{O}(h^\infty) \Omega(\tau). \quad (6.13)$$

Proof. Mimicking the proof of [20, Proposition 4.12], we get

$$v_{i,j} = \langle P \varphi_{\mathbf{m}_i}, \varphi_{\mathbf{m}_j} \rangle + \mathcal{O}(h^\infty \langle P \varphi_{\mathbf{m}_i}, \varphi_{\mathbf{m}_i} \rangle^{1/2} \langle P \varphi_{\mathbf{m}_j}, \varphi_{\mathbf{m}_j} \rangle^{1/2}). \quad (6.14)$$

Then one deduces from Proposition 6.5 the existence of some $\alpha > 0$ such that

$$\begin{aligned} e^{2S_1/h} v_{i,j} &= e^{2S_1/h} (v_{i,j}^\# (1 + \mathcal{O}(e^{-\alpha/h})) + \mathcal{O}(h^\infty \langle P \varphi_{\mathbf{m}_i}, \varphi_{\mathbf{m}_i} \rangle^{1/2} \langle P \varphi_{\mathbf{m}_j}, \varphi_{\mathbf{m}_j} \rangle^{1/2})) \\ &= e^{2S_1/h} v_{i,j}^\# + \mathcal{O}(e^{(2S_1 - S_i - S_j)/h} h^\infty). \end{aligned}$$

This establishes (6.13).

It just remains to show that $h^{-1}e^{2S_1/h}\Upsilon^\# \in \text{GS}_{\text{cl}}(\mathcal{E}, \tau)$. Indeed, the fact that $h^{-1}e^{2S_1/h}\Upsilon \in \text{GAS}_{\text{cl}}(\mathcal{E}, \tau)$ will then follow from (6.13).

Using (6.1), (6.2) and the fact that $P^\#$ is symmetric, we deduce that $h^{-1}e^{2S_1/h}\Upsilon^\#$ is a graded matrix, say $h^{-1}e^{2S_1/h}\Upsilon^\# = \Omega(\tau)M_h\Omega(\tau)$, where M_h is a symmetric matrix having a classical expansion with leading term

$$(M_0)_{i,j} = (-1)^{\delta_{\mathbf{m}_i, \mathbf{m}_j} - 1} \sum_{s \in \mathbf{j}(\mathbf{m}_i) \cap \mathbf{j}(\mathbf{m}_j)} \frac{|\mu(s)|}{2\pi} \frac{\sqrt{D_{\mathbf{m}_i} D_{\mathbf{m}_j}}}{D_s}.$$

In order to show that M_0 is positive definite, it is sufficient to show that it has the form $M_0 = L^*L$, where L is an injective matrix from \mathbb{R}^{n_0-1} to $\mathbb{R}^{\#\mathcal{V}^{(1)}}$. To this end, let us define, for every $s_k \in \mathcal{V}^{(1)}$, $G(s_k) := \{\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}; s_k \in \mathbf{j}(\mathbf{m})\}$. For any $s_k \in \mathcal{V}^{(1)}$, the set $G(s_k)$ is nonempty. Moreover, from the structure of a Morse function near a (separating) saddle point, this set has at most two elements, and has only one element \mathbf{m} if and only if $s_k \in \partial\widehat{E}(\mathbf{m})$. If there is only one minimum point $\mathbf{m}_i \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$ in $G(s_k)$, we set

$$L_{k,i} = \sqrt{\frac{|\mu(s)|}{2\pi} \frac{D_{\mathbf{m}_i}}{D_{s_k}}}.$$

If there are two minimum points $\mathbf{m}_i \neq \mathbf{m}_j \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$ in $G(s_k)$, we define

$$L_{k,i} = \sqrt{\frac{|\mu(s)|}{2\pi} \frac{D_{\mathbf{m}_i}}{D_{s_k}}} \quad \text{and} \quad L_{k,j} = -\sqrt{\frac{|\mu(s)|}{2\pi} \frac{D_{\mathbf{m}_j}}{D_{s_k}}}.$$

We do not specify which index (i or j) receives a “−”, since this choice is irrelevant in what follows. The other coefficients of L are set to zero:

$$L_{k,i} = 0 \quad \text{whenever } \mathbf{m}_i \notin G(s_k), \text{ i.e. whenever } s_k \notin \mathbf{j}(\mathbf{m}_i).$$

A direct computation gives $M_0 = L^*L$. Moreover, it follows from [24, Lemma 5.1] that L is injective. Let us briefly explain the argument of [24, Lemma 5.1]. Let $X = (X_{\mathbf{m}})_{\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}} \in \mathbb{R}^{n_0-1}$ be such that $LX = 0$. For any $s_k \in \mathcal{V}^{(1)}$ such that $G(s_k)$ contains a unique element $\mathbf{m}(s_k) \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$, we have $X_{\mathbf{m}(s_k)} = 0$. It follows from the structure of L that $X_{\mathbf{m}} = 0$ for every $\mathbf{m} \in \text{Cl}(\mathbf{m}(s_k))$. But, for every $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$, there exist $s_k \in \mathcal{V}^{(1)}$ and $\mathbf{m}(s_k) \in \text{Cl}(\mathbf{m})$ such that $G(s_k) = \{\mathbf{m}(s_k)\}$. Thus $X_{\mathbf{m}} = 0$ for every $\mathbf{m} \in \mathcal{U}^{(0)} \setminus \{\underline{\mathbf{m}}\}$ and L is injective. Summing up, M_0 is positive definite and thus $h^{-1}e^{2S_1/h}\Upsilon^\# \in \text{GS}_{\text{cl}}(\mathcal{E}, \tau)$, which concludes the proof of Proposition 6.8. ■

6.3. The spectrum of GAS matrices

The results stated here are variants of those of [24, Section 5] and of [20, Appendix]. For $p \in \mathbb{N}^*$, a finite-dimensional vector space $E = E_1 \oplus \cdots \oplus E_p$, and $j \in \{1, \dots, p\}$, let us write a general matrix $M \in \mathcal{M}(E)$ in block form

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \text{ on the direct sum } (E_1 \oplus \cdots \oplus E_{j-1}) \oplus (E_j \oplus \cdots \oplus E_p). \quad (6.15)$$

If $A \in \mathcal{M}(E_1 \oplus \cdots \oplus E_{j-1})$ is invertible, the Schur matrix of M (with respect to the vector space $E_1 \oplus \cdots \oplus E_{j-1}$) is the matrix on $E_j \oplus \cdots \oplus E_p$ defined by

$$\mathcal{R}_j(M) = D - CA^{-1}B,$$

where by convention $\mathcal{R}_1(M) = M$. By the Schur complement method, M is invertible if and only if $\mathcal{R}_j(M)$ is invertible. Moreover, the map \mathcal{R}_j sends $\mathcal{S}^+(E)$ into $\mathcal{S}^+(E_j \oplus \cdots \oplus E_p)$ and $\mathcal{S}_{\text{cl}}^+(E)$ into $\mathcal{S}_{\text{cl}}^+(E_j \oplus \cdots \oplus E_p)$. We will also denote by $\mathcal{J} : \mathcal{M}(\bigoplus_{k=j}^p E_k) \rightarrow \mathcal{M}(E_j)$ the restriction map to the first vector space E_j of $\bigoplus_{k=j}^p E_k$. More precisely, with the notation of (6.15), we will write $\mathcal{J}(M) = A$ when $j = 1$. Of course, the map \mathcal{J} depends on $j \in \{1, \dots, p\}$, but we omit this dependence since the set on which \mathcal{J} is acting will be obvious in what follows.

Let E be a finite-dimensional vector space and $M_h \in \mathcal{S}_{\text{cl}}^+(E)$. By a standard result of perturbation theory of symmetric matrices (see [17, Theorem 6.1 in Chapter II]), the eigenvalues of M_h , after an appropriate labeling that we assume, have an asymptotic expansion in powers of h with a nonzero leading term. This justifies the following definition.

Definition 6.9. For $M_h \in \mathcal{S}_{\text{cl}}^+(E)$, we denote by $\sigma_{\equiv}(M_h)$ the set of asymptotic expansions (that is, formal power series in h) of the eigenvalues of M_h . Moreover, $m_{\equiv}(\Lambda, M_h)$ is defined as the multiplicity of $\Lambda \in \sigma_{\equiv}(M_h)$.

By an abuse of notation, if $\lambda \in \sigma(M_h)$ has an asymptotic expansion $\Lambda \in \sigma_{\equiv}(M_h)$, we will sometimes write $m_{\equiv}(\lambda, M_h)$ instead of $m_{\equiv}(\Lambda, M_h)$. Roughly speaking, $m_{\equiv}(\lambda, M_h)$ can be seen as the multiplicity modulo $\mathcal{O}(h^\infty)$ of the eigenvalue $\lambda \in \sigma(M_h)$. Note that if $\lambda, \nu \in \sigma(M_h)$ do not have the same asymptotic expansion, there exists $K_0 > 0$ such that $|\lambda - \nu| \geq h^{K_0}$ for $h > 0$ small enough.

Theorem 4. Let $\mathcal{M}_h = \Omega(\tau)M_h\Omega(\tau) \in \text{GAS}_{\text{cl}}(\mathcal{E}, \tau)$ and assume that $\tau_j = \tau_j(h) = \mathcal{O}(h^\infty)$ for every $j = 2, \dots, p$. Then, for $h > 0$ small enough,

$$\sigma(\mathcal{M}_h) \subset \bigcup_{j=1}^p \varepsilon_j(\tau)^2 (\sigma(\mathcal{J} \circ \mathcal{R}_j(M_h^s)) + D(0, \mathcal{O}(h^\infty))), \quad (6.16)$$

where we recall that $M_h^s = \frac{1}{2}(M_h + M_h^*)$ denotes the real part of M_h . Moreover, for all $j = 1, \dots, p$, $K > 0$ large enough and $\lambda \in \sigma(\mathcal{J} \circ \mathcal{R}_j(M_h^s))$, one has

$$n(\mathcal{M}_h, D_{j,\lambda}^K) = m_{\equiv}(\lambda, \mathcal{J} \circ \mathcal{R}_j(M_h^s)), \quad (6.17)$$

which does not depend on K , where $n(\mathcal{M}_h, D_{j,\lambda}^K)$ is the number of eigenvalues of \mathcal{M}_h in $D_{j,\lambda}^K = D(\varepsilon_j^2 \lambda, \varepsilon_j^2 h^K)$ counted with multiplicity. Finally, for all $K > 0$ large enough, there exists $C > 0$ such that

$$\forall z \in \mathbb{C} \setminus \bigcup_{j,\lambda} D_{j,\lambda}^K, \quad \|(\mathcal{M}_h - z)^{-1}\| \leq C \text{dist}(z, \sigma(\mathcal{M}_h))^{-1} \quad (6.18)$$

for $h > 0$ small enough.

Proof. For $j = 1, \dots, p$, we assume that the spectral parameter z belongs to the ring $\mathcal{C}_j = \{z \in \mathbb{C}; \varepsilon_j^2/R < |z| < R\varepsilon_j^2\}$, where $R > 1$ is large enough so that $\bigcup_{\lambda \in \sigma(\mathcal{J} \circ \mathcal{R}_j(M_h^s))} D_{j,\lambda}^K \subset \mathcal{C}_j$. We write \mathcal{M}_h as

$$\mathcal{M}_h = \Omega(\tau)M_h\Omega(\tau) = \begin{pmatrix} L_+ & 0 \\ 0 & L_- \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} L_+ & 0 \\ 0 & L_- \end{pmatrix} = \begin{pmatrix} L_+AL_+ & L_+BL_- \\ L_-CL_+ & L_-DL_- \end{pmatrix},$$

with $L_+ = \text{diag}(\varepsilon_1 \text{Id}_{E_1}, \dots, \varepsilon_{j-1} \text{Id}_{E_{j-1}})$ and $L_- = \text{diag}(\varepsilon_j \text{Id}_{E_j}, \dots, \varepsilon_p \text{Id}_{E_p})$. For $z \in \mathcal{C}_j$, we have $\mathcal{E} := L_+AL_+ - z = L_+(A - L_+^{-1}zL_+^{-1})L_+$ with $\|L_+^{-1}zL_+^{-1}\| = \mathcal{O}(\varepsilon_j^2\varepsilon_{j-1}^{-2}) = \mathcal{O}(\tau_j^2) = \mathcal{O}(h^\infty)$. Since $A \in \mathbb{S}_{\text{cl}}^+(E_1 \oplus \dots \oplus E_{j-1})$ modulo $\mathcal{O}(h^\infty)$, the matrix \mathcal{E} is invertible and

$$\mathcal{E}^{-1} = L_+^{-1}(A^{-1} + \mathcal{O}(h^\infty))L_+^{-1} = \mathcal{O}(\varepsilon_{j-1}^{-2}), \quad (6.19)$$

uniformly for $z \in \mathcal{C}_j$. By the Schur complement method, $\mathcal{M}_h - z$ is invertible if and only if

$$\mathcal{F} = L_-DL_- - z - L_-CL_+\mathcal{E}^{-1}L_+BL_- \text{ is invertible.} \quad (6.20)$$

In that case,

$$(\mathcal{M}_h - z)^{-1} = \begin{pmatrix} \mathcal{E}^{-1} + \mathcal{E}^{-1}L_+BL_-\mathcal{F}^{-1}L_-CL_+\mathcal{E}^{-1} & -\mathcal{E}^{-1}L_+BL_-\mathcal{F}^{-1} \\ -\mathcal{F}^{-1}L_-CL_+\mathcal{E}^{-1} & \mathcal{F}^{-1} \end{pmatrix}. \quad (6.21)$$

From (6.19), the matrix \mathcal{F} can be written, uniformly for $z \in \mathcal{C}_j$,

$$\mathcal{F} = L_-(D - CA^{-1}B)L_- - z + \mathcal{O}(\varepsilon_j^2h^\infty) = \varepsilon_j^2\mathcal{J}(D - CA^{-1}B) \oplus 0 - z + \mathcal{O}(\varepsilon_j^2h^\infty)$$

with $\mathcal{J}(D - CA^{-1}B) : E_j \rightarrow E_j$ and the shorthand

$$T \oplus 0 = \begin{pmatrix} T & 0 \\ 0 & 0 \end{pmatrix} : E_j \oplus \dots \oplus E_p \rightarrow E_j \oplus \dots \oplus E_p.$$

Thus,

$$\begin{aligned} \mathcal{F} &= \varepsilon_j^2\mathcal{J} \circ \mathcal{R}_j(M_h) \oplus 0 - z + \mathcal{O}(\varepsilon_j^2h^\infty) \\ &= \varepsilon_j^2\mathcal{J} \circ \mathcal{R}_j(M_h^s) \oplus 0 - z + \mathcal{O}(\varepsilon_j^2h^\infty). \end{aligned} \quad (6.22)$$

We obtain an upper bound on the resolvent of \mathcal{M}_h away from its expected spectrum. Let $K_0 > 0$ be such that $|\lambda - \nu| \geq h^{K_0}$ for all $\lambda, \nu \in \sigma(\mathcal{J} \circ \mathcal{R}_j(M_h^s))$ having different asymptotic expansions and let $K > K_0$. Define $\tilde{\mathcal{C}}_j = \mathcal{C}_j \setminus \bigcup_{\lambda \in \sigma(\mathcal{J} \circ \mathcal{R}_j(M_h^s))} D_{j,\lambda}^K$. Since $\mathcal{J} \circ \mathcal{R}_j(M_h^s)$ is symmetric, we have $(\varepsilon_j^2\mathcal{J} \circ \mathcal{R}_j(M_h^s) - z)^{-1} = \mathcal{O}(\varepsilon_j^{-2}h^{-K})$ for $z \in \tilde{\mathcal{C}}_j$, and

$$\begin{aligned} \mathcal{F} &= (\varepsilon_j^2\mathcal{J} \circ \mathcal{R}_j(M_h^s) \oplus 0 - z)(1 + (\varepsilon_j^2\mathcal{J} \circ \mathcal{R}_j(M_h^s) \oplus 0 - z)^{-1}\mathcal{O}(\varepsilon_j^2h^\infty)) \\ &= (\varepsilon_j^2\mathcal{J} \circ \mathcal{R}_j(M_h^s) \oplus 0 - z)(1 + \mathcal{O}(h^\infty)), \end{aligned}$$

and so \mathcal{F} is invertible and

$$\mathcal{F}^{-1} = (\varepsilon_j^2\mathcal{J} \circ \mathcal{R}_j(M_h^s) \oplus 0 - z)^{-1} + \mathcal{O}(\varepsilon_j^{-2}h^\infty) = \mathcal{O}(\varepsilon_j^{-2}h^{-K}). \quad (6.23)$$

Combining the first equality in (6.19), (6.21), (6.23) and using $\varepsilon_k = \mathcal{O}(\varepsilon_\ell h^\infty)$ for $k > \ell$, we get, for $z \in \tilde{\mathcal{C}}_j$,

$$(\mathcal{M}_h - z)^{-1} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & (\varepsilon_j^2 \mathcal{J} \circ \mathcal{R}_j(M_h^s) - z)^{-1} & 0 \\ 0 & 0 & -z^{-1} \end{pmatrix} + \mathcal{O}(\varepsilon_j^{-2} h^\infty), \quad (6.24)$$

which implies that there exists $C > 0$ such that for $z \in \tilde{\mathcal{C}}_j$,

$$\|(\mathcal{M}_h - z)^{-1}\| \leq C \operatorname{dist}(z, \varepsilon_j^2 \sigma(\mathcal{J} \circ \mathcal{R}_j(M_h^s)))^{-1}. \quad (6.25)$$

We now compute the eigenvalues of \mathcal{M}_h . For $\lambda \in \sigma(\mathcal{J} \circ \mathcal{R}_j(M_h^s))$, we introduce the spectral projectors

$$\Pi_\lambda = -\frac{1}{2i\pi} \int_{\partial D_{j,\lambda}^K} (\mathcal{M}_h - z)^{-1} dz \quad \text{and} \quad \pi_\lambda = -\frac{1}{2i\pi} \int_{\partial D(\lambda, h^K)} (\mathcal{J} \circ \mathcal{R}_j(M_h^s) - z)^{-1} dz.$$

Since the circumference of $\partial D_{j,\lambda}^K$ is $2\pi \varepsilon_j^2 h^K$, (6.24) implies

$$\Pi_\lambda = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \pi_\lambda & 0 \\ 0 & 0 & 0 \end{pmatrix} + \mathcal{O}(h^\infty).$$

Since E is a finite-dimensional space and the rank of a projector is an integer equal to its trace, we deduce that

$$\operatorname{rk}(\Pi_\lambda) = \operatorname{tr}(\Pi_\lambda) = \operatorname{tr}(\pi_\lambda) + \mathcal{O}(h^\infty) = \operatorname{rk}(\pi_\lambda).$$

Thus, the number of eigenvalues of \mathcal{M}_h in $D_{j,\lambda}^K$ counted with multiplicity is equal to $m_\equiv(\lambda, \mathcal{J} \circ \mathcal{R}^{j-1}(M_h^s))$. Since

$$\sum_{j=1}^p \sum_{\lambda \in \sigma_\equiv(\mathcal{J} \circ \mathcal{R}_j(M_h^s))} m_\equiv(\lambda, \mathcal{J} \circ \mathcal{R}_j(M_h^s)) = \sum_{j=1}^p \dim E_j = \dim E,$$

relations (6.16) and (6.17) of Theorem 4 follow.

To finish the proof of Theorem 4, we have to obtain the resolvent estimate (6.18). Since the estimate follows from (6.25) in $\tilde{\mathcal{C}}_j$, $j = 1, \dots, p$, it remains to prove it in $\mathcal{D}_1 = \{z \in \mathbb{C}; R\varepsilon_1^2 \leq |z|\}$, $\mathcal{D}_j = \{z \in \mathbb{C}; R\varepsilon_j^2 \leq |z| \leq \varepsilon_{j-1}^2/R\}$ for $2 \leq j \leq p$ and $\mathcal{D}_{p+1} = \{z \in \mathbb{C}; |z| \leq \varepsilon_p^2/R\}$. We only show it in \mathcal{D}_j when $2 \leq j \leq p$, since the two remaining situations can be treated in the same way. Mimicking the proof of (6.19), we have $\mathcal{E}^{-1} = L_+^{-1}(A^{-1} + \mathcal{O}(R^{-1}))L_+^{-1}$ for R large enough and $z \in \mathcal{D}_j$. Then \mathcal{F} defined in (6.20) satisfies, instead of (6.22),

$$\mathcal{F} = \varepsilon_j^2 \mathcal{J} \circ \mathcal{R}_j(M_h^s) \oplus 0 - z + \mathcal{O}(\varepsilon_j^2 R^{-1}).$$

Since $|z| \geq R\varepsilon_j^2$, this implies $\|\mathcal{F}^{-1}\| = \mathcal{O}(|z|^{-1})$ and $\|(\mathcal{M}_h - z)^{-1}\| = \mathcal{O}(|z|^{-1})$ from (6.21). The last estimate gives (6.18). ■

6.4. The spectrum of the interaction matrix

Combining Proposition 6.8 and Theorem 4, we obtain the following result.

Theorem 5. *Suppose that the assumptions of Theorem 1 are satisfied. Assume also that (Gibbs), (Confin), and (Morse) hold true and $\mathcal{U}^{(0),II} = \emptyset$. Let Υ^\sharp be defined by (6.10). Then*

$$\sigma(P) \cap \{\operatorname{Re} z < \varepsilon_* h\} \subset \{0\} \cup \bigcup_{j=1}^p e^{-2S_j/h} (\sigma(\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon^\sharp)) + D(0, \mathcal{O}(h^\infty))).$$

Moreover, for all $j = 1, \dots, p$, $K > 0$ large enough and $\lambda \in \sigma(\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon^\sharp))$, one has

$$n(P, D_{j,\lambda}^K) = m_\equiv(\lambda, \mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon^\sharp)),$$

where $n(P, D_{j,\lambda}^K)$ is the number of eigenvalues of P in $D_{j,\lambda}^K = D(e^{-2S_j/h} \lambda, e^{-2S_j/h} h^K)$ counted with multiplicity.

Note here that the matrix $h^{-1} \mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon^\sharp)$ belongs to $\mathcal{S}_{\text{cl}}^+$, which almost gives the statement of Theorem 3. In order to obtain an explicit version of Theorem 3, let us use the specific structure of the matrix Υ^\sharp . We say that a matrix M is \mathcal{A} -compatible if $M_{\mathbf{m},\mathbf{m}'} = 0$ for all \mathbf{m}, \mathbf{m}' with $\text{Cl}(\mathbf{m}) \neq \text{Cl}(\mathbf{m}')$. In that case, for all $\alpha \in \mathcal{A} \setminus \{\text{Cl}(\underline{\mathbf{m}})\}$, M_α denotes the matrix M restricted to the lines and columns whose label belongs to $\mathcal{U}_\alpha^{(0)}$. Note that M is \mathcal{A} -compatible iff M is block diagonal after a permutation of lines and columns which gathers the labels of each element of \mathcal{A} . Thus, if A is a square \mathcal{A} -compatible matrix,

$$\sigma(A) = \bigcup_{\alpha \in \mathcal{A} \setminus \{\text{Cl}(\underline{\mathbf{m}})\}} \sigma(A_\alpha),$$

counting the eigenvalues with multiplicity. Moreover, if B, C are two \mathcal{A} -compatible matrices which can be multiplied, then BC is \mathcal{A} -compatible and $(BC)_\alpha = B_\alpha C_\alpha$. The same way, if D is \mathcal{A} -compatible and invertible, then D^{-1} is \mathcal{A} -compatible and $(D^{-1})_\alpha = (D_\alpha)^{-1}$.

It follows from (6.2) and (6.3) that $\langle P^\sharp \varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle = 0$ when $\text{Cl}(\mathbf{m}) \neq \text{Cl}(\mathbf{m}')$ and hence the matrix Υ^\sharp defined by (6.10) is \mathcal{A} -compatible with

$$\Upsilon_\alpha^\sharp = (\langle P^\sharp \varphi_{\mathbf{m}}, \varphi_{\mathbf{m}'} \rangle)_{\mathbf{m}, \mathbf{m}' \in \mathcal{U}_\alpha^{(0)}} \quad (6.26)$$

for all $\alpha \in \mathcal{A} \setminus \{\text{Cl}(\underline{\mathbf{m}})\}$. The previous paragraph shows that $\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon^\sharp)$ is also \mathcal{A} -compatible and

$$(\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon^\sharp))_\alpha = \mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon_\alpha^\sharp),$$

for all $\alpha \in \mathcal{A} \setminus \{\text{Cl}(\underline{\mathbf{m}})\}$ and $j = 1, \dots, p$. This implies

$$\sigma(\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon^\sharp)) = \bigcup_{\alpha \in \mathcal{A} \setminus \{\text{Cl}(\underline{\mathbf{m}})\}} \sigma(\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon_\alpha^\sharp)), \quad (6.27)$$

counting the eigenvalues with multiplicity, and we obtain the following variant of Theorem 5, whose formulation is close to that of [24, Theorem 5.8].

Theorem 6. *Suppose that the assumptions of Theorem 1 are satisfied. Assume also that (Gibbs), (Confin), and (Morse) hold true and $\mathcal{U}^{(0),II} = \emptyset$. Let Υ_α^\sharp be defined by (6.26). Then*

$$\sigma(P) \cap \{\operatorname{Re} z < \varepsilon_* h\} \subset \{0\} \cup \bigcup_{\substack{\alpha \in \mathcal{A} \setminus \{\operatorname{Cl}(\underline{\mathbf{m}})\} \\ 1 \leq j \leq p}} e^{-2S_j/h} (\sigma(\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon_\alpha^\sharp)) + D(0, \mathcal{O}(h^\infty))).$$

Moreover, for all $\alpha \in \mathcal{A}$, $j = 1, \dots, p$, $K > 0$ large enough and $\lambda \in \sigma(\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon_\alpha^\sharp))$, one has

$$n(P, D_{j,\lambda}^K) = \sum_{\beta \in \mathcal{A}} m_\equiv(\lambda, \mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon_\beta^\sharp)),$$

where $n(P, D_{j,\lambda}^K)$ is the number of eigenvalues of P in $D_{j,\lambda}^K = D(e^{-2S_j/h} \lambda, e^{-2S_j/h} h^K)$ counted with multiplicity and with the convention that $m_\equiv(\lambda, M) = 0$ if $\lambda \notin \sigma_\equiv(M)$.

In this result, if $e^{-S_j/h}$ does not appear in the graded expression of Υ_α^\sharp , the matrix $\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon_\alpha^\sharp)$ acts on the trivial vector space $\{0\}$ and its spectrum is empty.

In order to emphasize the connection with the formulation of [24, Theorem 5.8], let us set, with a slight abuse of notation, $\mathcal{R}^k = \mathcal{R}_2 \circ \dots \circ \mathcal{R}_2$ (k times), with the convention $\mathcal{R}^0 = \operatorname{Id}$. It then follows from Appendix C that $\mathcal{R}^{j-1} = \mathcal{R}_j$ for every $j \in \{1, \dots, p\}$. Thus, for every $j \in \{1, \dots, p\}$, the term $\mathcal{J} \circ \mathcal{R}_j(e^{2S_j/h} \Upsilon_\alpha^\sharp)$ is nothing but $\mathcal{J} \circ \mathcal{R}^{j-1}(e^{2S_j/h} \Upsilon_\alpha^\sharp)$, which is the expression appearing in [24, Theorem 5.8].

Remark 6.10. A similar result holds true without the assumption $\mathcal{U}^{(0),II} = \emptyset$. This requires constructing adapted quasimodes as in [24, Section 3] (see for example formula (3.14) there) with cut-off functions as above.

Appendix A. Spectrum of transport operators

We begin the appendix with a result used to solve some transport equations.

Lemma A.1. *For $m \in \mathbb{N}^*$, let $\mathcal{P}_{\operatorname{hom}}^m$ denote the set of complex polynomials in d variables which are homogeneous of degree m . Let $A \in \mathcal{M}_d(\mathbb{C})$ and let $\mathcal{L}_A : \mathcal{P}_{\operatorname{hom}}^m \rightarrow \mathcal{P}_{\operatorname{hom}}^m$ be given by*

$$\mathcal{L}_A p = Ax \cdot \nabla p.$$

If $\sigma(A) \subset \{\operatorname{Re} z > 0\}$, then $\sigma(\mathcal{L}_A) \subset \{\operatorname{Re} z > 0\}$.

Proof. Assume first that A is diagonalizable and denote by $\lambda_1, \dots, \lambda_d$ its eigenvalues. After a linear change of variable in \mathbb{C}^d (leaving $\mathcal{P}_{\operatorname{hom}}^m$ invariant), we can assume that A is diagonal, that is, $A = \operatorname{diag}(\lambda_1, \dots, \lambda_d)$. For any $\gamma = (\gamma_1, \dots, \gamma_d) \in \mathbb{N}^d$ satisfying $\sum_{j=1}^d \gamma_j = m$, we have

$$\mathcal{L}_A x^\gamma = \mathcal{L}_A x_1^{\gamma_1} \cdots x_d^{\gamma_d} = |\gamma|_\lambda x^\gamma$$

with $|\gamma|_\lambda := \sum_{j=1}^d \lambda_j \gamma_j$. Thus, the monomials of degree m form a basis of eigenvectors of \mathcal{L}_A on $\mathcal{P}_{\text{hom}}^m$ and the corresponding eigenvalues are $|\gamma|_\lambda$.

In the general case, we note that $A \mapsto \mathcal{L}_A$ is continuous and recall that the spectrum is a continuous function of the matrix (see [17, Theorem II.5.14]). By density of the diagonalizable matrices in $\mathcal{M}_d(\mathbb{C})$, the above discussion shows that the eigenvalues of \mathcal{L}_A on $\mathcal{P}_{\text{hom}}^m$ are still the $|\gamma|_\lambda$'s where $\lambda_1, \dots, \lambda_d$ are the eigenvalues of A . Since $\operatorname{Re} \lambda_j > 0$ for all j implies $\operatorname{Re} |\lambda|_\gamma > 0$ for all $\gamma \in \mathbb{N}^d$ such that $\sum_{j=1}^d \gamma_j > 0$, the lemma follows. ■

Appendix B. The supersymmetric structure

We now give an example showing that the operators considered in this paper do not have a nice supersymmetric structure. We refer the reader to Hérau, Hitrik, and Sjöstrand's paper [15] and to the last author's work [23] for general discussions of supersymmetric structure for differential operators of second order. We say that P as in (1.8) admits a temperate supersymmetric structure if there exist a smooth $d \times d$ matrix $G(x, h)$ and $M > 0$ such that

$$P = - \sum_{i,j=1}^d (h \partial_{x_i} - \partial_{x_i} f(x)) \circ G_{i,j}(x, h) \circ (h \partial_{x_j} + \partial_{x_j} f(x)), \quad (\text{B.1})$$

and $\|G(x, h)\| \lesssim h^{-M}$ locally in x . Note that (B.1) implies $P(e^{-f/h}) = P^*(e^{-f/h}) = 0$ and that the present definition of temperate supersymmetric structure is weaker than that of [23].

Proposition B.1. *In dimension $d = 2$, there exists an operator P satisfying the assumptions of Theorem 2 and having no temperate supersymmetric structure.*

The counterexample constructed in the proof also shows that the determination of the supersymmetric structure (that is, the matrix G) of an operator having a temperate supersymmetric structure is an instable question. On the contrary, since all the closed forms on \mathbb{R}^d are exact, all the operators P satisfying the assumptions of Theorem 2 have a supersymmetric structure which may not be temperate (see Theorem 1.2 of [15]).

Proof of Proposition B.1. First, we consider an operator P_0 satisfying all the assumptions of Theorem 2 and having a temperate supersymmetric structure (for instance, the




Fig. B.1. The geometric setting in the proof of Proposition B.1.

Witten Laplacian described in (1.25)). Of course, P_0 is of the form (1.8). We assume in addition that the Morse function f is such that there exist two points $\rho_1, \rho_2 \in \mathbb{R}^2$, a simple smooth loop γ around ρ_1 but not ρ_2 , and $C_0 > 0$ such that

$$\max_{f|_\gamma} < C_0 < \min(f(\rho_1), f(\rho_2)), \quad (\text{B.2})$$

(see Fig. B.1). Let $\chi \in C_c^\infty(\mathbb{R}^2; [0, 1])$ be such that $\chi(\rho_1) = 1$ and $\text{supp}(\nabla \chi)$ is sufficiently close to γ (so, in particular, $\chi(\rho_2) = 0$). We define $P = P_0 + P_{\text{per}}$ with the perturbation operator

$$P_{\text{per}} = \sum_{j=1}^2 b_j^{\text{per}}(x, h) \circ h \partial_{x_j} + h \partial_{x_j} \circ b_j^{\text{per}}(x, h),$$

and the smooth and compactly supported vector field

$$b^{\text{per}}(x, h) = e^{2f/h} e^{-2C_0/h} d^* \chi \quad \text{where} \quad d^* = \begin{pmatrix} \partial_{x_2} \\ -\partial_{x_1} \end{pmatrix}. \quad (\text{B.3})$$

If the support of $\nabla \chi$ is close enough to γ , the function b^{per} and all its derivatives are exponentially small from (B.2). In particular, b^{per} satisfies (1.9) with a null classical expansion and P_{per} does not change the principal symbol of P_0 . Moreover, a direct computation gives

$$P_{\text{per}}(e^{-f/h}) = -(P_{\text{per}})^*(e^{-f/h}) = e^{f/h} h \operatorname{div}(b^{\text{per}} e^{-2f/h}) = 0. \quad (\text{B.4})$$

Thus, P (as P_0) satisfies the assumptions of Theorem 2. It remains to show that P has no temperate supersymmetric structure. Since P_0 has such a structure, it is equivalent to show that P_{per} has no temperate supersymmetric structure. Towards a contradiction, assume that P_{per} can be written as in (B.1) for some polynomially locally bounded matrix $G(x, h)$. Since $P_{\text{per}} = -(P_{\text{per}})^*$, the matrix G must be antisymmetric, say

$$G(x, h) = \begin{pmatrix} 0 & g(x, h) \\ -g(x, h) & 0 \end{pmatrix}$$

for some smooth function g with $|g(x, h)| \lesssim h^{-M}$ locally. Expanding (B.1) gives

$$\begin{aligned} P_{\text{per}} &= -(h\nabla - \nabla f) \cdot G(h\nabla + \nabla f) \\ &= h(d^* g) \cdot h\nabla - h\nabla \cdot (G\nabla f) - (G\nabla f) \cdot h\nabla \\ &= \left(\frac{h}{2} d^* g - G\nabla f \right) \cdot h\nabla + h\nabla \cdot \left(\frac{h}{2} d^* g - G\nabla f \right). \end{aligned}$$

Then G satisfies the relation

$$b^{\text{per}} = \frac{h}{2} d^* g - G\nabla f = \frac{h}{2} e^{2f/h} d^* (e^{-2f/h} g), \quad (\text{B.5})$$

which is similar to [23, (2.4)]. Comparing with (B.3), this equation is equivalent to

$$d^* (e^{-2f/h} g) = \frac{2}{h} e^{-2C_0/h} d^* \chi.$$

Since $d^*\psi = 0$ implies that ψ is constant in dimension 2, this gives

$$g = \frac{2}{h} e^{2(f-C_0)/h} (\chi + C(h)) \quad (\text{B.6})$$

for some constant $C(h) \in \mathbb{R}$. Computing g at $x = \rho_2$ where $\chi = 0$, (B.2), (B.6) and $|g(\rho_2, h)| \lesssim h^{-M}$ imply that $C(h)$ must be exponentially small. On the other hand, computing g at $x = \rho_1$ where $\chi = 1$ leads to $g(\rho_1, h) \geq h^{-1} e^{2(f(\rho_1)-C_0)/h}$ for h small enough. Then (B.2) shows that $g(\rho_1, h)$ is exponentially large, in contradiction with $|g(\rho_1, h)| \lesssim h^{-M}$. Summing up, P_{per} and hence P have no temperate supersymmetric structure and the proposition follows. ■

Appendix C. Iteration of the Schur complement

Let us conclude the appendix with a lemma about the Schur complement. We recall that, for a matrix $M \in \mathcal{M}_{d+d'}(\mathbb{C})$ with $d, d' \in \mathbb{N}^*$ written in block form

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \quad \text{with} \quad A \in \mathcal{M}_d(\mathbb{C}) \text{ invertible,}$$

the Schur complement of the block $D \in \mathcal{M}_{d'}(\mathbb{C})$ of M is the matrix defined by

$$\mathcal{R}(M) = D - CA^{-1}B \in \mathcal{M}_{d'}(\mathbb{C}).$$

Moreover, by the Schur complement method, M is invertible if and only if $\mathcal{R}(M)$ is invertible.

Lemma C.1. *For $d_1, d_2, d_3 \in \mathbb{N}^*$ and matrices $M \in \mathcal{M}_{d_1+d_2+d_3}(\mathbb{C})$ and $M' \in \mathcal{M}_{d_2+d_3}(\mathbb{C})$ written in block form*

$$M = \begin{pmatrix} A & B & C \\ D & E & F \\ G & H & I \end{pmatrix} \quad \text{and} \quad M' = \begin{pmatrix} A' & B' \\ C' & D' \end{pmatrix},$$

we denote respectively, when they make sense, by $\mathcal{R}_1(M)$, $\mathcal{R}_{1,2}(M)$ and $\mathcal{R}_2(M')$ the Schur complements of the blocks $\begin{pmatrix} E & F \\ H & I \end{pmatrix} \in \mathcal{M}_{d_2+d_3}(\mathbb{C})$ of M , $I \in \mathcal{M}_{d_3}(\mathbb{C})$ of M and $D' \in \mathcal{M}_{d_3}(\mathbb{C})$ of M' .

If M has the previous form with A and $\begin{pmatrix} A & B \\ D & E \end{pmatrix}$ invertible, then the Schur complements $\mathcal{R}_{1,2}(M)$, $\mathcal{R}_1(M)$ and $\mathcal{R}_2(\mathcal{R}_1(M))$ make sense and satisfy

$$\mathcal{R}_2(\mathcal{R}_1(M)) = \mathcal{R}_{1,2}(M).$$

Proof. First, since A and $\begin{pmatrix} A & B \\ D & E \end{pmatrix}$ are invertible, the respective Schur complements $\mathcal{R}_1(M)$ and $\mathcal{R}_{1,2}(M)$ of the blocks $\begin{pmatrix} E & F \\ H & I \end{pmatrix}$ and I of M make sense. Moreover, by the Schur complement method, the invertibility of A and $\begin{pmatrix} A & B \\ D & E \end{pmatrix}$ implies that the Schur complement

$E - DA^{-1}B$ of the block E of $\begin{pmatrix} A & B \\ D & E \end{pmatrix}$ is invertible, and a straightforward computation shows that

$$\begin{pmatrix} A & B \\ D & E \end{pmatrix}^{-1} = \begin{pmatrix} A^{-1} + A^{-1}B(E - DA^{-1}B)^{-1}DA^{-1} & -A^{-1}B(E - DA^{-1}B)^{-1} \\ -(E - DA^{-1}B)^{-1}DA^{-1} & (E - DA^{-1}B)^{-1} \end{pmatrix}.$$

It follows that

$$\begin{aligned} \mathcal{R}_{1,2}(M) &= I - (G \ H) \begin{pmatrix} A & B \\ D & E \end{pmatrix}^{-1} \begin{pmatrix} C \\ F \end{pmatrix} \\ &= I - GA^{-1}C + (H - GA^{-1}B)(E - DA^{-1}B)^{-1}(DA^{-1}C - F). \end{aligned} \quad (\text{C.1})$$

Moreover,

$$\mathcal{R}_1(M) = \begin{pmatrix} E & F \\ H & I \end{pmatrix} - \begin{pmatrix} D \\ G \end{pmatrix} A^{-1} (B \ C) = \begin{pmatrix} E - DA^{-1}B & F - DA^{-1}C \\ H - GA^{-1}B & I - GA^{-1}C \end{pmatrix}.$$

Since $E - DA^{-1}B$ is invertible, the Schur complement $\mathcal{R}_2(\mathcal{R}_1(M))$ of the block $I - GA^{-1}C$ of $\mathcal{R}_1(M)$ makes sense and satisfies

$$\mathcal{R}_2(\mathcal{R}_1(M)) = I - GA^{-1}C - (H - GA^{-1}B)(E - DA^{-1}B)^{-1}(F - DA^{-1}C). \quad (\text{C.2})$$

The statement of Lemma C.1 then follows from (C.1) and (C.2). ■

Funding. The authors are supported by the ANR project QuAMProcs 19-CE40-0010-01.

References

- [1] Berglund, N.: Kramers' law: validity, derivations and generalisations. *Markov Process. Related Fields* **19**, 459–490 (2013) Zbl [1321.58035](#) MR [3156961](#)
- [2] Bouchet, F., Reygner, J.: Generalisation of the Eyring–Kramers transition rate formula to irreversible diffusion processes. *Ann. Henri Poincaré* **17**, 3499–3532 (2016) Zbl [1375.82081](#) MR [3568024](#)
- [3] Bovier, A., Eckhoff, M., Gaynard, V., Klein, M.: Metastability in reversible diffusion processes. I. Sharp asymptotics for capacities and exit times. *J. Eur. Math. Soc.* **6**, 399–424 (2004) Zbl [1076.82045](#) MR [2094397](#)
- [4] Bovier, A., Gaynard, V., Klein, M.: Metastability in reversible diffusion processes. II. Precise asymptotics for small eigenvalues. *J. Eur. Math. Soc.* **7**, 69–99 (2005) Zbl [1105.82025](#) MR [2120991](#)
- [5] Coron, J.-M.: *Control and nonlinearity*. Math. Surveys Monogr. 136, American Mathematical Society, Providence, RI (2007) Zbl [1140.93002](#) MR [2302744](#)
- [6] Di Gesù, G., Le Peutrec, D.: Small noise spectral gap asymptotics for a large system of nonlinear diffusions. *J. Spectr. Theory* **7**, 939–984 (2017) Zbl [1395.60095](#) MR [3737885](#)
- [7] Dimassi, M., Sjöstrand, J.: *Spectral asymptotics in the semi-classical limit*. London Math. Soc. Lecture Note Ser. 268, Cambridge University Press, Cambridge (1999) Zbl [0926.35002](#) MR [1735654](#)
- [8] Eyring, H.: The activated complex in chemical reactions. *J. Chem. Phys.* **3**, 107–115 (1935)

- [9] Helffer, B., Klein, M., Nier, F.: Quantitative analysis of metastability in reversible diffusion processes via a Witten complex approach. *Mat. Contemp.* **26**, 41–85 (2004) Zbl [1079.58025](#) MR [2111815](#)
- [10] Helffer, B., Sjöstrand, J.: [Puits multiples en mécanique semi-classique. IV. Étude du complexe de Witten](#). *Comm. Partial Differential Equations* **10**, 245–340 (1985) Zbl [0597.35024](#) MR [0780068](#)
- [11] Helffer, B., Sjöstrand, J.: From resolvent bounds to semigroup bounds. *arXiv*:[1001.4171v1](#) (2010)
- [12] Helffer, B., Sjöstrand, J.: [Improving semigroup bounds with resolvent estimates](#). *Integral Equations Operator Theory* **93**, article no. 36, 41 pp. (2021) Zbl [1496.47067](#) MR [4271979](#)
- [13] Hérau, F., Hitrik, M., Sjöstrand, J.: [Tunnel effect for Kramers–Fokker–Planck type operators](#). *Ann. Henri Poincaré* **9**, 209–274 (2008) Zbl [1141.82011](#) MR [2399189](#)
- [14] Hérau, F., Hitrik, M., Sjöstrand, J.: [Tunnel effect and symmetries for Kramers–Fokker–Planck type operators](#). *J. Inst. Math. Jussieu* **10**, 567–634 (2011) Zbl [1223.35246](#) MR [2806463](#)
- [15] Hérau, F., Hitrik, M., Sjöstrand, J.: [Supersymmetric structures for second order differential operators](#). *Algebra i Analiz* **25**, 125–154 (2013) Zbl [1303.81086](#) MR [3114853](#)
- [16] Hérau, F., Nier, F.: [Isotropic hypoellipticity and trend to equilibrium for the Fokker–Planck equation with a high-degree potential](#). *Arch. Ration. Mech. Anal.* **171**, 151–218 (2004) Zbl [1139.82323](#) MR [2034753](#)
- [17] Kato, T.: *Perturbation theory for linear operators*. Grundlehren Math. Wiss. 132, Springer, New York (1966) Zbl [0148.12601](#) MR [0203473](#)
- [18] Kramers, H. A.: [Brownian motion in a field of force and the diffusion model of chemical reactions](#). *Physica* **7**, 284–304 (1940) Zbl [0061.46405](#) MR [0002962](#)
- [19] Landim, C., Mariani, M., Seo, I.: [Dirichlet’s and Thomson’s principles for non-selfadjoint elliptic operators with application to non-reversible metastable diffusion processes](#). *Arch. Ration. Mech. Anal.* **231**, 887–938 (2019) Zbl [1412.35093](#) MR [3900816](#)
- [20] Le Peutrec, D., Michel, L.: [Sharp spectral asymptotics for nonreversible metastable diffusion processes](#). *Probab. Math. Phys.* **1**, 3–53 (2020) Zbl [1484.60086](#) MR [4408002](#)
- [21] Lee, J., Seo, I.: [Non-reversible metastable diffusions with Gibbs invariant measure I: Eyring–Kramers formula](#). *Probab. Theory Related Fields* **182**, 849–903 (2022) Zbl [1489.82059](#) MR [4408505](#)
- [22] Lelièvre, T., Rousset, M., Stoltz, G.: [Free energy computations](#). Imperial College Press, London (2010) Zbl [1227.82002](#) MR [2681239](#)
- [23] Michel, L.: [Around supersymmetry for semiclassical second order differential operators](#). *Proc. Amer. Math. Soc.* **144**, 4487–4500 (2016) Zbl [1345.81043](#) MR [3531196](#)
- [24] Michel, L.: [About small eigenvalues of the Witten Laplacian](#). *Pure Appl. Anal.* **1**, 149–206 (2019) Zbl [1447.35236](#) MR [3949372](#)
- [25] Pavliotis, G. A., Stoltz, G., Vaes, U.: [Scaling limits for the generalized Langevin equation](#). *J. Nonlinear Sci.* **31**, article no. 8, 58 pp. (2021) Zbl [1473.35057](#) MR [4195749](#)
- [26] Talay, D.: Approximation of invariant measures of nonlinear hamiltonians and dissipative stochastic differential equations. In: *Progress in stochastic structural dynamics*, Publication du L.M.A.-C.N.R.S. 152, 139–168 (1999)
- [27] Villani, C.: [Hypocoercivity](#). *Mem. Amer. Math. Soc.* **202**, no. 950, iv+141 pp. (2009) Zbl [1197.35004](#) MR [2562709](#)
- [28] Witten, E.: [Supersymmetry and Morse theory](#). *J. Differential Geometry* **17**, 661–692 (1982) Zbl [0499.53056](#) MR [0683171](#)